



AGRICULTURAL RESEARCH INSTITUTE
PUSA

JOURNAL OF THE CHEMICAL SOCIETY

ABSTRACTS OF PAPERS ON ORGANICAL, INORGANIC, MINERALOGICAL, AND ANALYTICAL CHEMISTRY.

Committee of Publication:

A. E. H. MANN, M.C., D.Sc.	T. M. LOWRY, C.B.E., D.Sc., F.R.S.
C. L. DRAKE, M.A.	J. W. MCBAIN, M.A., Ph.D.
A. W. CHURCH, O.M.G., C.B.E.	J. I. O. MASSON, M.B.E., D.Sc.
C. L. DRAKE, M.A., Ph.D.	J. C. PHILIP, O.B.E., D.Sc., Ph.D., F.R.S.
R. H. PICKARD, D.Sc., Ph.D., F.R.S.	R. H. PICKARD, D.Sc., Ph.D., F.R.S.
N. V. SIDGWICK, M.A., Sc.D., F.R.S.	N. V. SIDGWICK, M.A., Sc.D., F.R.S.
J. F. THORPE, C.B.E., D.Sc., F.R.S.	J. F. THORPE, C.B.E., D.Sc., F.R.S.
Sir JAMES WALKER, D.Sc., LL.D., F.R.S.	Sir JAMES WALKER, D.Sc., LL.D., F.R.S.

Editor:

J. GREENAWAY.

Assistant Editor:

CLARENCE SMITH, D.Sc.

Assistant:

A. A. ELDRIDGE, B.Sc.

Indexer:

MARGARET LE PLA, B.Sc.

Abstractors:

J. BARGER, M.A., D.Sc., F.R.S.	G. W. ROBINSON, M.A.
H. J. EVANS, B.Sc.	E. H. RODD, D.Sc.
W. E. GARNER, M.A.	F. M. ROWE, D.Sc.
W. GODDEN, B.Sc.	W. P. SKERTCHLY.
C. R. HARRINGTON, B.A.	J. F. SPENCER, D.Sc., Ph.D.
C. K. INGOLD, D.Sc.	L. J. SPENCER, M.A., Sc.D.
K. KASHIMA, Rg.S.	E. STEDMAN, B.Sc.
J. KENNER, D.Sc., Ph.D.	J. S. G. THOMAS, D.Sc.
W. O. KERMAK, M.A., B.Sc.	W. THOMAS, M.A., M.Sc.
H. KING, D.Sc.	R. TRUSZKOWSKI, B.Sc.
S. I. LEVY, B.A., B.Sc.	D. F. TWISS, D.Sc.
S. S. MIHOLIC, Ph.D.	J. C. WITHERS, Ph.D.
G. F. MORRELL, D.Sc., Ph.D.	H. WREN, M.A., D.Sc., Ph.D.
J. R. PARTINGTON, M.B.E., D.Sc.	S. S. ZILVA, D.Sc., Ph.D.
T. H. POPE, B.Sc.	

1922. Vol. CXXII. Part II

LONDON:

GURNEY & JACKSON, 33, PATERNOSTER ROW, E.C. 4.
1922.

Abstractors of the *Journal of the Society of Chemical Engineers*
who have contributed to this volume.

J. F. BRIGGS.
T. H. BURNHAM.
A. COUSEN.
J. B. FIRTH.
A. GROUNDS, M.Sc.
A. J. HALL.
C. IRWIN.

J. H. LANE.
H. MOORE.
A. G. POLLARD.
A. R. POWELL.
J. REILLY.
H. C. REYNARD.
A. DE WAELE.

JOURNAL

OF

THE CHEMICAL SOCIETY.

ABSTRACTS OF CHEMICAL PAPERS PUBLISHED IN
BRITISH AND FOREIGN JOURNALS.

PART II.

General and Physical Chemistry.

Refraction of Light and Atomic Structure. Refraction Equivalents of Ions. JARL A. WASASTJERNA (*Öfvers. Finska Vet.-Soc.*, 1921, **63**, [A], No. 4, 18 pp.; from *Chem. Zentr.*, 1921, iii, 759).—Cuthbertson's rule for the relation between the refraction equivalents of atoms and their position in the periodic system is applicable to the refractⁿ and alkaline-earth metals. The next sheath of an alkali metal formation passes over to negative. In the case of contains two electrons in group. The next sheath in the alkali and alkaline-earth metals is identical with that in the preceding member of the horizontal series which is nearer to the nucleus on account of increased attraction. Numerical values are given for the refraction equivalent of positive and negative ions.

1482
■■■■■■■■
IARI

one of the alkali
outer electron
on which in salt
hereby becomes
the outer sheath
negative atom or

G. W. R.

The Molecular Refraction of Substances of Higher Melting Point and the Calculation of the Corresponding Refractive Exponents to the Temperature of Comparison, 20°. FRITZ EISENLOHR (*Ber.*, 1921, **54**, [B], 2857—2867).—Difficulties are experienced in calculating the refractive values of substances of higher melting point above 20°, since values calculated from the

Lorentz-Lorenz expression increase perceptibly with increasing temperature, whereas those calculated according to Gladstone-Dale decrease to about a corresponding extent. The only expression for the molecular refraction which is nearly independent of the temperature is that of Eykman, $(n^2-1)M/(n+0.4)d$, but this has never become popular with chemists. If a and b are the factors for the conversion of the Lorentz-Lorenz and Gladstone-Dale to the Eykman values, the expressions $n^2-1/n+0.4=a(n^2-1)/(n^2+2)$ and $n^2-1/n+0.4=b(n-1)$ are obtained, whence $a=n^2+2/n+0.4$ and $b=n+1/n+0.4$. For a given interval of temperature, for example 10° , it is found that the alteration in $\log a \times \log b$ is approximately constant and independent of the value of the refractive index; for the given temperature interval, $\log a = -0.00033$ and $\log b = +0.00024$. The latter value cannot be regarded as completely independent of the magnitude of the refractive index if this varies greatly from the normal value ($n=1.45$) and, for this case, a modified table of corrections is given, but, in general, the values quoted are sufficiently accurate. A number of examples are quoted showing the method of calculating the refractive index from one temperature to another over considerable intervals of temperature; in general, the observed and calculated values are in excellent agreement, but cymene affords an exception. A small number of other substances also do not behave in accordance with Eykman's formula. The substances appear to have a very appreciable vapour tension at the atmospheric temperature and to pass at a comparatively low temperature into a region in which there is no longer a perfectly uniform change of refractive index with the temperature. For safety, it is advisable to restrict observations to a region which does not embrace more than one-third of the temperature of ebullition as usually expressed. H. W. .

Optical Properties of Solutions. A Theory of the Structure of the Molecules of Electrolytes. JARL A. WASASTJERNA (*Acta Soc. Sci. Fennicae*, 1920, 50, No. 2, 129 pp.; from *Chem. Zentr.*, 1921, iii, 758—759).—The refractive indices and densities of aqueous solutions of a number of organic salts, sodium chloride, potassium chloride, and oxalic acid were determined for two different temperatures and for three different wave-lengths at each temperature. From the results obtained certain conclusions were drawn as to the influence of dissociation on the optical properties of electrolytes. The number of valency electrons in each atom may be directly calculated from Eisenlohr's atomic refraction constant and the results are in agreement with Drude's valency theory. A new interpretation of the Lorenz-Planck dispersion formula is given leading to a working hypothesis whereby the selective photoelectric effect can be calculated as well as, in certain cases, the alteration in refraction and dispersion due to ionisation. It is shown that the temperature coefficient of molecular refraction for dissolved salts is always negative by the Newton-Laplace formula, variably positive and negative by the Gladstone-Dale formula, and negative by the Lorentz-Lorenz formula. It is

further shown that the old n_D^2 specific refractive power is satisfied by the n_D^2 Dale formula. For the determination of the molecular refraction of dissolved salts, strong solutions should be used and the values extrapolated for 100%. The optical properties of salt solutions can be simply explained by assuming that the ions occur ready formed in the molecules of strong electrolytes whilst the molecules of weak electrolytes cannot be thus polarised.

G. W. R.

Notes of Spectrography. E. VON ANGERER (*Physikal. Z.*, 1921, 22, 521—523).—A description of various modifications of apparatus and technique used by the author in spectroscopic work. The points included refer to (a) the iron arc, (b) the mercury reference spectrum, (c) the reproduction of a wave-length scale, and (d) an ultra-violet monochromatic light filter.

J. F. S.

The Origin of Band Spectra. YUTAKA TAKAHASHI (*Proc. Phys. Math. Soc. Japan*, 1921, [3], 3, 20—28, 30—33).—It is possible to deduce a formula representing a system of band spectra from the oscillation of the atoms in a molecule if, with Bohr and Sommerfeld, the quantum relations $A_1 - A_2 = h\nu$ and $\int_L p_i dq_i = nh$ are accepted for the conditions of the radiation and the stationary state. A simple Deslandres formula $\nu = A + B(n_1^2 - n_2^2)$ where $B = h/8\pi^2 I_0$ is arrived at. If the angular momentum of the molecular rotation is not a complete multiple of $h/2\pi$ the formula becomes $\nu = A \pm B(n + \epsilon)^2$, where $\epsilon < 1$ as observed in many cases. The theoretical values of A and B in the last equation are found to be of the same order of magnitude as the observed ones for band spectra of nitrogen and oxygen. In the low potential discharge in hydrogen, Fulcher isolated two triplet bands. It is shown that these bands can be represented by the author's formula by taking suitable values for the constants.

CHEMICAL ABSTRACTS.

The Changeable Fine Structure of the Balmer Series. E. GEHROCKE and E. LAU (*Physikal. Z.*, 1921, 22, 556—557).—A change in pressure brings about a change in the intensity relationship of the two components of the Balmer series, which is a series of doublets. The intensity change is most noticeable in the case of H_α and least in the case of H_γ . Using a long tube of 2 cm. bore the ratio I_k/I_l , on changing the pressure from 0.1 mm. to 0.01 mm., increases by 50% for H_α , 11% for H_β , and 5% for H_γ . The ratio I_k/I_l is that of the intensity of the short wave-length component to the intensity of the long wave-length component. The change of the intensity relationship is nearly inversely proportional to the square of the series number. The bore of the tube has a controlling influence on the intensity relationship, thus, in a 5 cm. long tube of 1.5 mm. bore the longer wave-length component is stronger, but both components are faint. The appearance of the lines is unchanged on changing the bore from 5 mm. to 0.1 mm.

J. F. S.

Spectrum of Bromine Vapour ; Zeeman Effect. G. RIBAUD (*J. phys.*, 1917, 7, 205—208).—The emission spectrum of bromine

vapour was studied with the aid of a Geissler tube discharge. If a Geissler tube is placed perpendicularly to the field of an electromagnet, and strong fields are used, the apparent resistance of the tube becomes extremely high, and the discharge requires, therefore, a very high potential; moreover, the lines obtained are much broadened and difficult to measure. The tube is consequently introduced along the axis of the pole-pieces. Twenty-six lines in the region λ 4014 to λ 5332 were studied in a field of 21800 gauss. The values of $\Delta\lambda/H\lambda^2$ for the majority of the lines are grouped about the values 1.05×10^{-4} and 1.23×10^{-4} . Normal separation was not observed for any of the lines, but the lines λ 4766, 4785, and 4816 have a separation very nearly double the normal (cf. A., 1912, ii, 1114; Kimura, A., 1921, ii, 140, 141).

CHEMICAL ABSTRACTS.

Excitation of the Enhanced Spectrum of Magnesium in a Low Voltage Arc. PAUL D. FOOTE, W. F. MEGGERS, and F. L. MOHLER (*Phil. Mag.*, 1921, [vi], 42, 1002—1015).—A vacuum arc is described in which the electronic-atomic collisions take place at any desired definite velocity of the electrons. The neutral magnesium atom is shown to absorb quanta of the following values, which may be emitted as equivalent quanta of radiation: 46.9 volts, producing *L*-radiation; 22.8 volts (probably), producing double ionisation and resulting in the simple enhanced and the arc spectra; 7.61 volts, producing simple ionisation and resulting in the arc spectrum; 2.70 volts resulting in the single line spectrum. The simply ionised magnesium atom absorbs quanta as follows: 14.97 volts, producing double ionisation and resulting in the simple enhanced spectrum; 4.4 volts resulting in the single-line enhanced spectrum. With high current density other quanta may be absorbed corresponding with fundamental lines of the subordinate series.

J. F. S.

Band Spectra of Isotopes. L. GREBE and H. KONEN (*Physikal. Z.*, 1921, 22, 546—549).—The wave-lengths of the higher members of the band spectrum of uranium-lead and ordinary lead have been compared. The chosen wave-lengths lie between λ 4257.690 and 4281.458 Å.U. Eighteen lines have been compared, and it is shown that the wave-length of the line corresponding with uranium-lead is on the average 0.055 Å.U. shorter than that for ordinary lead. This figure agrees so far as its order is concerned with the assumption that the diatomic molecules are the carriers of the band spectrum.

J. F. S.

Wave-lengths of Lines in the Iron Arc from Grating and Interferometer Measurements, λ 3370—6750. CHARLES E. ST. JOHN and HAROLD D. BABCOCK (*Astrophys. J.*, 1921, 53, 260—299).—One thousand and twenty-six lines were measured with an accuracy of ± 0.001 Å. Manipulative details receive special consideration in this paper.

CHEMICAL ABSTRACTS.

Determination of the Terms of the Cyanogen Bands. A. KRATZER (*Physikal. Z.*, 1921, 22, 552—555).—A theoretical

paper in which the relationships of the violet cyanogen bands are investigated and the values obtained theoretically are compared with the experimental and empirical values of Heurlinger (*Z. Physik.*, 1920, i, 82). It is shown that the zero lines of the violet cyanogen bands are expressed by the formula $\nu = 25797.83 - n_1(2143.88 - n_1 \times 20.25) - n_2(2055.64 - n_2 \times 13.25)$, in which n_1 and n_2 are the quantum numbers. A comparison of the zero values calculated by this equation with the empirically deduced values of Heurlinger shows in most cases an astonishingly good agreement between the two sets of wave-lengths. The whole of the line systems of the violet bands of cyanogen may be calculated by the theoretically deduced formula. $\nu = \nu_e + B_1 n_1 + n_1 \nu_1^\circ (1 - n_1 u_1 x_1) - n_2 \nu_2^\circ (1 - n_2 u_2 x_2) \pm 2mB_2 n_1 + m^2(B_1 n_1 - B_2 n_1)$, where $\nu_e + B_1^\circ = 25797.83$; $\nu_e = 25795.91$; $\nu_1^\circ = 2143.88$; $\nu_1^\circ u_1 x_1 = 20.25$; $2B_1 n_1 = 3.841 - 0.0044n_1$; $\nu_2^\circ = 2055.64$; $\nu_2^\circ u_2 x_2 = 13.25$ and $2B_2 n_1 = 3.705 - 0.035n_2$. It now becomes possible to differentiate between the red cyanogen bands and the so-called nitrogen bands, since, as was shown by Heurlinger, the red cyanogen bands and the violet cyanogen bands have a common end term.

J. F. S.

The Absorption Spectrum of Hydrogen Chloride. WALTER F. COLBY and CHARLES F. MEYER (*Astrophys. J.*, 1921, 53, 300—309).—The hydrogen chloride absorption band extending from 3.16 to 3.70 μ was observed by the use of apparatus similar to that described by Imes (*Astrophys. J.*, 1919, 50, 251—276). The compensation chamber was found to be superfluous. The absorption chambers, 16 to 60 cm. in length, could be heated to incipient redness. The experiments confirmed results of Paton showing that heating increases the number of observable lines but does not change the wave-length. The wave numbers for 28 lines were tabulated and an equation was derived to represent them. The law of spacing is not parabolic, as has been supposed, but a cubic term is found to be necessary.

CHEMICAL ABSTRACTS.

An Absorption Band Spectrum for Water in the Region of Wave-lengths of Several Decimetres. RICHARD WEICHMANN (*Physikal. Z.*, 1921, 22, 535—544).—The method of the parallel wire system is so modified that it may be used for the determination of the refractive indices of liquids and gives results with an error of only 0.5%. An exciter has been constructed by means of which a practically monochromatic radiation up to $\lambda = 12$ cm. and of a sufficient intensity can be produced. The resonance curve has a perfectly smooth course and a decrement which in general does not exceed 0.04. The width of the spectrum lines is therefore not much greater than $\Delta\lambda = 1$ mm. In the spectrum of water over the range $\lambda = 65$ cm. to $\lambda = 27$ cm., three bands of anomalous dispersion are found, which at the same time are three absorption bands. The width of the two outside bands is approximately $\Delta\lambda = 3$ cm. The middle band is sharper and has a width of 0.5 cm. These results indicate that it is probable that water possesses a true absorption spectrum in the region of long wave-lengths, which is

a continuation of the usual long wave-length absorption spectrum of water. J. F. S.

Action of Electrical Fields on Absorption Lines (*D*-Lines of Sodium Vapour). R. LADENBURG (*Physikal. Z.*, 1921, 22, 549—552).—The effect of electric fields of 100,000 volt/cm. on the *D* lines of sodium has been investigated. The sodium light was produced in a quartz capillary filled with a mixture of neon and helium, and was examined in a Lummer-Gehrcke spectroscope. It is shown that no resolution of the absorption lines occurs, but that probably the *p* and *s* components are displaced unequally in the same sense. It is shown that two fundamentally different kinds of electrical action on spectrum lines must be differentiated, (1) the symmetrical, relatively strong Stark effect, which is proportional to the field strength, and (2) the unsymmetrical effect, which is proportional to the square of the field strength and occurs when the influenced electron together with the Coulomb's force act, and in comparison with which the external field is to be regarded as small. J. F. S.

Effect of a Strong Electrical Field on the Absorption Lines of Sodium Vapour. R. LADENBURG (*Naturwiss.*, 1921, 9, 667; from *Chem. Zentr.*, 1921, iii, 993).—The Stark effect has only been observed with emission spectra of gases under the influence of canal rays. Using a strong electrical field (150,000—200,000 volts/cm.) and a Lummer-Gehrcke interference spectrometer, a marked asymmetrical effect on the absorption lines of sodium vapour in the yellow was observed, and a displacement in the red of 0.02 Å. G. W. R.

Width of the Absorption Bands of the Rare Earths. KARL F. HERZFELD (*Physikal. Z.*, 1921, 22, 544—546).—A mathematical paper, in which the width and the influence of temperature on the width of the absorption bands of the rare earths is considered. The width is shown to be due to a Stark effect, which is brought about by the penetration, due to the vibration, into the electric fields of the neighbouring ions. This view is in keeping with the dependence of the width on the temperature and leads to widths for the bands which are of the correct dimensions. It is shown that observations on the width of the bands at very low temperatures will furnish information on the existence of an energy zero point. The strength of the electric field in water is estimated as of the order 5×10^7 volts/cm. J. F. S.

Spectroscopy of Uranium and some Rare Earths. G. MEYER and GREULICH (*Physikal. Z.*, 1921, 22, 583—585).—The minimum concentration of solutions of salts of uranium, cerium, lanthanum, didymium, and thorium which may be detected spectroscopically has been ascertained. The illumination was produced in three ways: (1) sparks from a transformer of 10,000 volts were passed from a platinum wire to a bundle of platinum wires moistened with the solution, (2) a syrupy mixture of the salt solution with

phosphoric acid was placed in the platinum electrodes of a Miethe extra current apparatus, and (3) hollow carbon arc carbons were filled with the residue obtained on evaporating the solution to dryness. The following gives the minimum concentrations (c) in mg./c.c. and the wave-lengths of the lines visible at this concentration: cerium, λ 3940.89, 4166.75; $c=0.01$; lanthanum, λ 3949.22; $c=0.006$; didymium, λ 4061.27, 4325.80; Pr λ 4206.81, 4429.41; $c=0.01$; thorium, λ 4391.30, 4382.10; $c=0.01$; uranium, λ 4090.28, 5919.61, $c=5.0$. It is thus seen that the minimum concentration of uranium is approximately 500 times greater than that of the other metals. Experiments were made to ascertain the influence of other salts on the amount of the elements in question which could be detected. The value of $G=U/(U+M) \times 100$ has been determined, in which U and M represent the amounts of uranium and the other element. It is shown that the presence of cerium, lanthanum, didymium, and thorium has very little effect on the visibility of the uranium lines; in all cases the value of G approximates to unity. In the case of fifteen other elements examined with uranium, it is shown that G has a considerable value. It is also shown that a knowledge of the value of $G=U/(U+Fe) \times 100=7$ gives a means of estimating the amount of uranium in a solution of a uranium salt. The method consists in adding an iron salt to the solution until the persistent uranium lines just vanish. In the case of a solution of uranium sulphate containing 15.8 mg./c.c., there was found, using this method, 15.4 mg./c.c. The influence of the addition of salts of iron, nickel, chromium, manganese, potassium, sodium, calcium, cobalt, lithium, aluminium, magnesium, uranium, cerium, didymium, and thorium on the persistence of the spectrum lines obtained from solutions of cerium, lanthanum, didymium, and thorium salts was also investigated. The values $100Ce/(Ce+M)$; $100La/(La+M)$; $100Di/(Di+M)$, and $100Th/(Th+M)$ vary between 0.5 and 1.0 for all cases except those in which sodium to cerium, of the above list, have been added. It therefore follows that the rare earths are detectable in much smaller quantities in mixtures than is uranium.

J. F. S.

Absorption of Ultra-violet Light. THOS. H. DURRANS (*Perf. Essent. Oil Rec.*, 1921, 12, 370—371).—The ultra-violet absorption spectra of eugenol, isoeugenol, methylisoeugenol, anethole, methylchavicol, and anisole were examined. *isoEugenol* absorbs more strongly than eugenol, and methylation of the phenolic group still further increases the absorption, an $N/10,000$ solution being opaque in thicknesses greater than 20 cm. The absorptive power of anethole is remarkable, 10 cm. of $N/10,000$ solution or 0.25 mm. of pure anethole completely stopping ultra-violet light of a wave-length of less than about 27 Å.U. The substitution of the propenyl side-chain by the allyl radicle, or its complete removal, causes an enormous diminution of the absorptive power. The paper is illustrated by photographs of the absorption spectra of the substances examined.

G. F. M.

Absorption of Light by Solutions. HARALD LUNELUND (*Öfvers. Finska Vet.-Soc.*, 1916—1917, 59, No. 21, 21 pp.; from *Chem. Zentr.*, 1921, iii, 810).—Working with aqueous solutions of "brilliant safranine," "tartrazine," "crystal-ponceau," and "rose Bengal" with and without addition of sulphuric acid, it was shown that the extinction coefficient, ϵ , calculated from the formula $I' = I \cdot 10^{-\epsilon d}$, agrees with Beer's law, according to which the extinction coefficient is proportional to the concentration. Using the König-Martens spectral photometer, it is shown that in aqueous solutions of safranine a marked maximum absorption occurs for $\lambda 519\mu\mu$. By addition of sulphuric acid the colour changes to green through violet, and the absorption maximum changes to $\lambda 622\mu\mu$. G. W. R.

A Theory of Chromo-isomerism of Solid Compounds. HOWARD J. LUCAS and ARCHIE R. KEMP (*J. Amer. Chem. Soc.*, 1921, 43, 1654—1665).—See this vol., i, 30.

Hydrated Boric Acid as the Basis of Systems Capable of [Showing] a High Phosphorescence. E. TIEDE, P. WULFF, and A. RAGOSS (*Physikal. Z.*, 1921, 22, 563).—When boric acid is mixed with certain organic substances, particularly aromatic and heterocyclic substances, and partly dehydrated by melting, it is found on cooling that the mixtures are strongly phosphorescent. These substances after illumination by ultra-violet light emit a phosphorescent glow of a colour which varies with the organic substance and in the best cases persists for two to three minutes after the exciting light is removed. J. F. S.

Destruction of Phosphorescent Zinc Sulphides by Ultra-violet Light. LEONARD B. LOEB and LLOYD SCHMIEDESKAMP (*Proc. Nat. Acad. Sci.*, 1921, 7, 202—207).—Three different samples of phosphorescent zinc sulphides were exposed in very thin layers to the action of ultra-violet light from a mercury arc. In each case there was a marked decrease in phosphorescent intensity (measured by a phosphoroscope) with increase of time of ultra-violet illumination. The reduction of intensity was accompanied in each case by a darkening in colour. Exposure to chlorine restored to some extent the original colour and ability to phosphoresce. Some points of similarity with the destruction of phosphorescence by the bombardment of α -particles are discussed. G. W. R.

Linear Polarisation of the Blue Light from the Focus of the Lilienfeld Röntgen Tube. H. SEEMAN (*Physikal. Z.*, 1921, 22, 581—582).—It has been shown by Lilienfeld and Rother (*ibid.*, 1920, 21, 249, 360) that the light from the point of convergence of the Lilienfeld tube is almost completely linear polarised in the sense that the vibrations of the electric vector are at right angles to the plane of the anti-cathode which is inclined generally at 45° to the direction of the cathode rays. The present paper shows that this phenomenon can be observed without special apparatus

in full daylight by means of a Nicol prism held before the eye. The phenomenon is not observed in Coolidge or Müller electron tubes because of the very intense white light. J. F. S.

Studies on the Dependence of Optical Rotatory Power on Chemical Constitution. IV. Aryl Derivatives of Bisimino-camphor. BAWA KARTAR SINGH, MAHAN SINGH, and JIWAN LAL (T., 1921, 119, 1971—1976).

Negative Optical Anomalies. GERVAISE LE BAS (*Chem. News*, 1921, 123, 271—272).—From a study of certain thiophen derivatives, it is shown that substitution α 2 in position 3 can reduce the negative anomaly of sulphur to zero. Substitution on both sides of the sulphur atom practically neutralises the negative anomaly. The full substitution of one ethenoid group or substitution on one side of the molecule reduces the negative anomaly to half its value. Chlorine and bromine have lost their usual positive anomalies for substitution in these cases. W. G.

Mechanism of the Photochemical Chlorine-Hydrogen Gas Reaction and the Question of the Damping of the Velocity [of Reaction] of Chlorine Activated by Light. RUDOLF GÖHRING (*Z. Elektrochem.*, 1921, 27, 511—518).—The paper opens with an account of the principal investigations on the photochemical reactions between hydrogen and chlorine which have been carried out in the last twenty years. The possible reactions which may occur photochemically in a mixture of chlorine, hydrogen, and a little oxygen after the reaction $\text{Cl}_2 + L = 2\text{Cl}$ are considered. From a large number of possible combinations of reactions two sets are picked out, which both conform to Bodenstein's empirical reaction equation. The Bodenstein equations are theoretically deduced. The light absorption by chlorine has been measured for several wave-lengths and the fraction of light absorbed from the radiation of an Osram lamp has been calculated. It is shown that the number of quanta absorbed by 1 c.c. per sec. is approximately 2×10^{11} . The number of collisions calculated for the gas theory and the number of absorbed quanta are compared with reaction velocity calculated by Bodenstein. It is shown that for $h\nu$ the number of molecules of hydrogen chloride formed is 5×10^5 . Inserting the collision number and the measured reaction velocity in the equations deduced does not lead to any contradiction of the initial assumptions. The time required for damping of the activity of chlorine activated by light is calculated, and it is shown that theoretically no effect was to be expected in the experiments of Bodenstein and Taylor (A., 1916, ii, 463), although under other definite conditions an effect might have been observed. J. F. S.

The Decomposition of Ozone by Light of the Visible Spectrum. ROBERT OWEN GRIFFITH and WILLIAM JAMES SHUTT (T., 1921, 119, 1948—1959).

Verification of the Photochemical Equivalent Law with Photographic Dry Plates. J. EGGERT and W. NODDACK (*Sitzungsber. Preuss. Akad. Wiss. Berlin*, 1921, 631—635).—Dry plates

of various kinds have been illuminated with light of wave-length $407.8\mu\mu$ for measured periods of time and the amount of metallic silver formed by the absorption of a measured amount of light energy has been ascertained. It is shown that for a given type of dry plate (Agfa reproduction) the ratio N/Q , where N is the number of silver atoms formed by an amount of light energy Q quanta is constant and equal to 3.0%, that is, the production of one atom of silver requires 3% of the light energy striking the plate. The light absorption by the non-illuminated plate has been directly measured and is shown to be about 12—15%. A further experiment shows that about one hundred times as many quanta are absorbed as are silver nuclei formed. This indicates that about 100 quanta must act on a single particle before it can be reduced by the developer. The results point definitely to the conclusion that not every absorbed quantum produces a silver particle, but, in the case of moderate illumination, every silver particle corresponds with one and only one absorbed quantum. This is in keeping with the experimental result that the number of developed nuclei is proportional to the amount of absorbed light.

J. F. S.

Photochemistry of Silver Compounds. FRITZ WEIGERT and W. SCHÖLLER (*Sitzungsber. Preuss. Akad. Wiss. Berlin*, 1921, 641—650).—Experiments have been carried out in silver chloride emulsions with the object of ascertaining the mechanism of the photographic copying process. It is shown that silver chloride is, of itself, neither light sensitive, nor does it furnish a noticeable quantity of the silver which constitutes the photographic positive. The colloidal metallic silver, which is present as an impurity in very small amounts in fresh unilluminated emulsions, is the only light sensitive substance present. The colouring of the emulsion layers therefore commences very slowly and after a while accelerates itself for a short time and then the rate of colouring decreases since the photographic yield in such solid systems is smaller with increasing quantities of silver. This behaviour is represented by an S-shaped curve which is obtained by plotting the amount of silver formed as ordinates against the time of illumination. The strong colouring in systems rich in silver acts in the same sense, since the silver behaves as a harmful light filter. The fact, that in this case the substance, which is light sensitive, is produced by the process itself, shows that the photochemical process takes place in molecules other than those which actually absorb the light. Since the system is solid the transmission of energy from the absorbing molecules to the reacting molecules cannot take place through collisions as in gaseous systems. It is therefore probable the energy transmission occurs by means of electrons. In a single atom, as, for example, the absorbing silver atom, no process other than the separation of an electron can occur through the absorption of an energy quantum. The electron is absorbed by the surrounding silver salts with the production of a photo-electric effect.

J. F. S.

Action of Ultra-violet Light on Colloidal Platinum. ELLWOOD B. SPEAR, P. F. JONES, A. S. NEAVE, and M. SHLAGER (*J. Amer. Chem. Soc.*, 1921, **43**, 1385—1391).—A number of experiments on the coagulating action of ultra-violet light on platinum sols of various concentrations have been carried out at 20—22°. It is shown that ultra-violet light will precipitate colloidal platinum from solutions which contain no electrolytes other than carbon dioxide or constituents of the hard glass containers. Thus a solution containing 0.038 gram of platinum per litre was half coagulated in ten hours by ultra-violet light, whilst in the absence of ultra-violet light coagulation was not complete in two years. Coagulation by electrolytes is greatly accelerated by ultra-violet light. Thus a solution containing 0.038 gram of platinum was half coagulated in seven minutes by a solution of sodium chloride containing 4.25 millimols per litre when the action took place in ultra-violet light, but when no ultra-violet rays were employed sixty-four minutes were necessary to accomplish the same change. The action of ultra-violet light is greater in dilute solutions than in concentrated solutions of the colloids. J. F. S.

The Action of Light of Short Wave-lengths on some Organic Acids and their Salts. FRANS MAURITS JAEGER (*T.*, 1921, **119**, 2070—2076).

Photochemistry of the Retina. FRITZ WEIGERT (*Z. Elektrochem.*, 1921, **27**, 481—487).—The processes occurring in the retina are considered and an hypothesis of the action is advanced on the basis of experiments on the velocity of bleaching of layers of collodion containing dyes by polarised light. The velocity of bleaching of layers of collodion containing cyanine by linear polarised light has been examined spectro-photometrically and dichrometrically. It is shown that the velocity of bleaching of a freshly-prepared layer is much greater than that of an already partly bleached layer of the same extinction. It is also shown that layers of dyes resemble very strongly the photochlorides. J. F. S.

Phototropy and Photoelectric Effect. PATRICK H. GALLAGHER (*Bull. Soc. chim.*, 1921, [iv], **29**, 961—976; cf. A., 1921, i, 715).—The author has applied the method of Padoa and Amaduzzi (cf. A., 1912, ii, 227) in a modified form to a study of photo-electric effects in relationship to phototropy in the case of a number of aldehyde-amines. Phototropic compounds show photo-electric fatigue on exposure to the light from a mercury lamp much more slowly than other substances. The results seem to indicate that in phototropic transformations there is a liberation of electrons. An electron so liberated will take up a new position in the molecule or will remain at some distance from the molecule in a state of forced equilibrium. Its intervention in the photo-electric effect will be secondary. The relationship between phototropy and phosphorescence and fluorescence is considered. W. G.

Passage of α -Rays through Materials. H. RAUSCH VON TRAUBENBERG and K. PHILIPP (*Physikal. Z.*, 1921, **22**, 587—588).—The range of α -rays in liquids and gases has been investigated. The range of the α -ray in water is found to be 60μ . In the case of gases, carbon monoxide, carbon dioxide, methyl bromide, methyl iodide, chlorine, hydrogen chloride, and ammonia have been investigated and results obtained which agree with the results obtained by Bragg by the ionisation method. In the case of liquids, the range was measured by immersing a Sidot screen in the liquid until it just fluoresced and measuring the thickness of the liquid layer with a horizontal microscope. The range in gases was obtained by passing the beam of rays parallel to the horizontal side of a glass wedge containing the gas and placing in the path of the rays a Sidot screen at a small angle to the horizontal. By observing the boundary of dark and light on the screen, the range may be deduced. Taking oxygen as standard, it is shown that the stopping power of the gases varies as \sqrt{Z} . It is shown that the stopping power of compounds is not strictly additive as Bragg has stated, and it is now shown that the stopping power of hydrogen varies in different compounds, thus H_2 , 0.200 per atom; but in the compounds C_2H_2 , C_2H_4 , C_2H_6 , CH_4 , 0.187; NH_3 , 0.173; and HCl , 0.16. J. F. S.

Collisions of α -Particles with Hydrogen Nuclei. J. CHADWICK and E. S. BIELER (*Phil. Mag.*, 1921, [vi], **42**, 923—940).—The relationships which hold in the collisions between α -particles and hydrogen nuclei have been investigated. The angular distribution of the hydrogen particles projected by α -particles of mean range 6.6 cm. has been determined up to an angle of 66° . The distribution for α -rays of mean ranges 8.2, 4.3, and 2.9 cm. has been obtained over a smaller range of angle. It is shown that the number of hydrogen particles projected within these angles by α -rays of high velocity is greatly in excess of that given by forces varying as the inverse square of the distance between the centres of the two nuclei. The variation in the number of hydrogen particles projected within a given angle with the velocity of α -rays has been observed over a wide range. It is shown that for α -rays of high velocity the variation is in the opposite direction to that given by the inverse square law; for α -rays of range less than 2 cm. and velocity less than 1.26×10^9 cm. per sec., however, the collision relation is about the same as that given by the inverse square law. The experimental collision relation is compared with those calculated by Darwin for various models of the α -particle, and the conclusion is drawn that the α -particle behaves in these collisions as an elastic oblate spheroid of semi-axes about 8×10^{-13} and 4×10^{-13} cm., moving in the direction of its minor axis. Outside this surface, the force varies approximately as the inverse square of the distance from the centre of the spheroid. J. F. S.

Recoil of Hydrogen Nuclei from Swift α -Particles. A. L. MCAULAY (*Phil. Mag.*, 1921, [vi], **42**, 892—904).—An account of

work undertaken to ascertain the number of atoms recoiling at various angles from an homogeneous α -ray beam, from thorium-*C* and radium-*C* respectively with the object of obtaining data bearing on the nature of the collisions, the size of the colliding particles, and similar quantities. The experimental method was similar in principle to that already described (*ibid.*, 1920). An experiment made to determine the relation between the α - and γ -ray activity of thorium-*C* is described.

J. F. S.

The Oxidising Properties of certain Radioactive Elements.

PIERRE LEMAY and LÉON JALoustRE (*Compt. rend.*, 1921, 173, 916—918).—The bromides of mesothorium, radiothorium, thorium-*X*, and radium were found to be catalysts of certain oxidation reactions. Their action is not due to the intermediate formation of ozone but to α -radiation, and, for the small amounts of material used, it is the same for the four elements.

W. G.

Scattering of β -Rays by Thin Metal Sheets. H. GEIGER and W. BOTHE (*Physikal. Z.*, 1921, 22, 585—587).—The scattering of β -rays by thin metal sheets has been measured (a) in the region of small scattering angle ($\phi=15^\circ$ or less) and (b) in regions of large scattering angle ($\phi=60^\circ$ or more). It is shown that there is a fundamental difference between the two types of scattering. In the region of smaller angles, the observed angle is produced by the superposition of many individual small scattering angles through which the β -ray is bent as it passes the individual atoms (multiple scattering), whilst in the region of larger angles the superposition plays a subordinate rôle; each scattering angle is produced by a single collision of a β -ray with a single atom. This is relatively rare, for it is only brought about when the path of the electron lies very close to the nucleus of the atom (individual scattering). In the case of β -rays from radium-*(B+C)*, it is shown that for the very thinnest layers (under 0.01 mm.) the scattering is less than that demanded by the square root law, but for thicker layers this law is confirmed. There is an inverse proportionality between mv^2 and the probable scattering angle, if the uncertain measurements for β -rays of high velocity are left out of account.

J. F. S.

Excitation of γ -Radiation by α -Particles from Radium Emanation. F. P. SLATER (*Phil. Mag.*, 1921, [vi], 42, 904—923).—A hard γ -radiation is emitted when α -particles from radium emanation impinge on metals such as lead and tin. The radiation differs but little in quality when the radiator is changed from one of high atomic number to one of low atomic number. The coefficients of absorption in lead are 1.8 cm.^{-1} for lead and 2.1 cm.^{-1} for tin. The intensity obtained is very small, and only a small fraction of the impinging α -particles can be effective. For the same absorption conditions the intensity is about 50% greater for the lead radiation than for the tin radiation. It appears to be emitted fairly uniformly in all directions, but differs in all other respects from the characteristic radiations, and is probably emitted from

the nuclei of the atoms in the radiator after direct collision with the α -particles. No hard radiation is emitted by the disintegration of the radium emanation nucleus in the ordinary course of radioactive transformation. Elements of high atomic number when bombarded by α -particles emit two well-defined types of radiation, corresponding roughly with their characteristic K and L radiations. Tin, an element of medium atomic number, emits one type, within the range of the experiments, of quality roughly the same as its L radiation. The quantity of radiation emitted by any element is so small that a very small fraction of the impinging α -particles can cause radiation. Radium emanation emits, under the conditions of the experiments, no soft radiation on disintegration.

J. F. S.

Excitation of Soft Characteristic X-Rays. O. W. RICHARDSON and C. B. BAZZONI (*Phil. Mag.*, 1921, [vi], **42**, 1015—1019).—A development of work previously published by the authors (A., 1917, ii, 521). The thermionic radiation from a tungsten wire has been allowed to fall on targets of carbon, molybdenum, copper, and tungsten and in all cases an emission of radiation has been detected. In the case of carbon and molybdenum, the critical potential at which radiation sets in has been determined with considerable exactitude. In the case of carbon, the target was prepared on a copper disk from india-ink, and the thermionic current was kept constant by reducing the temperature of the filament as the operating voltage was increased. Although the pressure was of the order 10^{-8} — 10^{-9} mm., a radiation effect was present at 220 volts which may have been due to residual gas as it was independent of the voltage above 220 volts. A radiation occurs at 286 volts. If it be admitted that the radiations are soft X-rays excited according to the same laws as hard X-rays and the quantum relation is applied, it is found that the wave-length of the shortest member of the group lies between 42.7 and 44.3 Å.U. and is probably close to 43.4 Å.U. Extrapolating from K_{β} for aluminium and using Moseley's relation, it is found that K_{β} for carbon should have the wave-length 45.5 Å.U. This makes it practically certain that the radiation is the K series of carbon. In the case of molybdenum, radiation is obtained at 356 volts, which corresponds with the wave-length 34.8 Å.U. Calculating from Vegard's empirical relation, the M_{α} line of molybdenum should be 35.2 Å.U., which indicates that the radiation is the M series of molybdenum.

J. F. S.

Scattering of Röntgen Rays by Anisotropic Liquids. E. HÜCKEL (*Physikal. Z.*, 1921, **22**, 561—563).—The scattering of Röntgen rays by *p*-azoxyanisole, *p*-azoxyphenetole, dianisylidene-azine, cholesteryl propionate and cholesteryl benzoate as solid, clear-liquid, and turbid-liquid crystals has been photographed with the object of ascertaining whether or no the liquid crystals possess a space lattice. Several of the photographs are reproduced in the paper, and show that the turbid liquids do not consist of

regions built up on a space lattice, and, further, that in the suspended liquids there is no noticeable amount of material built up on a space lattice. It is not impossible, however, that small quantities of crystals are suspended in the liquid, since the interference bands which they would produce are too weak to be visible. It is shown that the term anisotropic liquids must be used in place of that of liquid crystals, for the word crystal implies a space lattice. If this is not adopted it will be logical to term as crystals a gas with molecules which carry electric dipoles in an electric field, also paramagnetic gases in a magnetic field. Consequently, it is to be taken that anisotropic liquids possess a more or less ordered arrangement of their molecules on account of their electrical and magnetic characteristics. J. F. S.

A Precision Röntgen Spectrograph. H. SEEMANN (*Physikal. Z.*, 1921, 22, 580—581).—A Röntgen spectrograph is described which is designed to measure lines of the hardest spectrum region. Using rock salt, reflecting crystal measurements may be made beyond 1 Å.U., and also in the softest technical Röntgen spectrum by using a gypsum reflecting crystal, measurements may be obtained beyond 3 Å.U. The instrument is designed for both the camera and window methods of measurement. J. F. S.

Fine Structure of Röntgen Spectra. ADOLF SMEKAL (*Physikal. Z.*, 1921, 22, 559—561).—A theoretical discussion of the fine structure of Röntgen spectra in which it is shown that if the sum of the quanta of an electron layer is k , that is, the layer is k -quantal, then the number of energy levels belonging to the layer is $2k-1$. According to Bohr, the O -layer of the heaviest elements is tri-quantal and the P -layer is di-quantal; it follows that there must be $5O$ and $3P$ energy levels. Wentzel has shown by means of his principle of selection that the O -layer is actually tri-quantal. J. F. S.

The Existence of a New Radioactive Emanation in the Springs of Bagnoles-de-l'Orne and its Surroundings. P. LOISEL (*Compt. rend.*, 1921, 173, 1098—1101).—In studying the gases dissolved in the water of a large number of springs in the region of Bagnoles-de-l'Orne, a curve of activity was obtained which could not be explained by the presence of any known emanation. The curve showed at first a decreasing activity, then a rise to a maximum, followed by a decrease. The results indicate the presence of a new, simple, radioactive substance, to which the author provisionally gives the name *emilium*. W. G.

Radioactivity and Atomic Constitution. LISE MEITNER (*Naturwiss.*, 1921, 9, 423—427; from *Chem. Zentr.*, 1921, iii, 847).—A discussion of the subject of isotopes in relation to atomic transformations. From a formula connecting the number of nuclear constituents with the number of helium nuclei and the number of electrons, four possible types of atomic disintegration of radioactive elements are deduced. G. W. R.

Adsorption of Radium by Barium Sulphate. FRANK E. E. GERMANN (*J. Amer. Chem. Soc.*, 1921, 43, 1615—1621).—From experiments on the adsorption of radium, from a radium barium chloride solution, containing 22.79×10^{-10} gram of radium per c.c., by barium sulphate, it is shown that Kroeker's adsorption law is true in the case of the adsorption of a radioactive substance. The adsorption law of Freundlich is found to be equally true in the case of a radioactive substance. Consequently, it follows that the laws which apply to the adsorption of easily weighable masses also apply with an equal degree of accuracy to masses of the order 5×10^{-6} gram adsorbed by 1 gram of adsorbent. The relationship existing between adsorption taking place during precipitation and that taking place on a solid precipitate of definite surface area has been discussed. J. F. S.

The Radioactivity of Mineral Springs. II. ERICH EBLER and A. J. VAN RHYN (*Z. anorg. Chem.*, 1921, 119, 135—144; cf. A., 1911, ii, 1049).—Borings for thermal water undertaken by the town of Heidelberg in 1913 provided an opportunity for studying the relationship of radium to radium emanation content and the influence of depth and geological considerations on the radioactivity of the water. The waters from these borings were also compared with those from other springs. In the Heidelberg waters the radium emanation content was constant at about 10^{-10} "Curie" per litre, and was independent of the depth of the boring down to 600 m. The granite springs of Löwenbrunnen show a higher and the New Red Sandstone springs of Felsenmeer a lower emanation content, in agreement with the known relatively high radium content of granite and the low radium content of sandstones. The radium salt content of the water from the Heidelberg borings was of the order 10^{-12} gram per litre, but rose at a depth of 590 m. to $0.97 \cdot 10^{-10}$, of the same order as that in the Max spring at Bad-Dürkheim (*loc. cit.*). This observation supports the view that the radium content increases with the depth of the spring. The granitic water of Löwenbrunnen does not show a high radium salt content corresponding with its high emanation content. It is important when comparing different waters to measure both radium and radium emanation content, as these may vary independently of each other. E. H. R.

The Adsorption of Radioactive Substances. II. ERICH EBLER and A. J. VAN RHYN (*Ber.*, 1921, 54, [B], 2896—2912; cf. A., 1911, ii, 957).—The view has been expressed previously (*loc. cit.*) that the colloidal condition does not exercise a directive influence on the behaviour of colloids towards radio-elements, but that this depends on the chemical nature of the substances, the reactions which occur being expedited by the increased surface. Thus, the adsorption of radioactive material by silica or, more precisely, by its hydrate, is due to the formation of sparingly soluble silicates which owe their origin to the acidic nature of the hydrate. The discordant results obtained by many observers

are due to the fact that the acidity of the solution plays an important part, since acid has a de-adsorbent action probably in accordance with the scheme: $\text{RaSiO}_3 + 2\text{HCl} = \text{RaCl}_2 + \text{H}_2\text{SiO}_3$. In addition, the silicic acid and its gels which have been used are very ill-defined products which, owing to the number of factors affecting their condition, are frequently not reproducible. In this connexion it is pointed out that the statement of Horovitz and Paneth (A., 1915, ii, 305) that the separation of silica during the estimation of radium in minerals is of no importance is only true within certain limits, since if free acid is not present in considerable quantity, the precipitate carries down radium.

The adsorptive power of charcoal for radioactive material appears difficult to explain in the light of the hypothesis outlined above. It is pointed out, however, that the varieties of charcoal employed never consist of pure carbon. In a series of experiments the adsorptive power of blood and animal charcoal towards uranium-X has been compared with that of pure charcoal in differing degrees of fineness (prepared by the carbonisation of pure sugar, extraction of the residue with water, and ignition of the product at bright redness). It is found that the selective adsorption of uranium-X by pure carbon is very much less than that of the impure varieties. An active preparation is obtained, however, when pure charcoal is mixed with the residues prepared by extracting blood or animal charcoal with water or dilute acid and subsequent evaporation of the extracts to dryness. The adsorptive power of impure charcoal, therefore, appears to be due to the impurities contained in it.

The adsorption of radium by barium sulphate has been examined further. This so-called "precipitation" of radium sulphate is not, in reality, a precipitation in the usual sense since, on account of the small concentration of radium, the solubility product $L = K.C_{\text{Ra}}.C_{\text{SO}}$, is seldom attained. It is shown that co-precipitation of radium sulphate with barium sulphate from solutions containing radium and barium is only complete when the barium is completely precipitated. An exact distribution of radium between the solid and liquid phase does not appear to occur.

H. W.

The Adsorption of Thorium-B and Thorium-C by Ferric Hydroxide. JOHN ARNOLD CRANSTON and ROBERT ALEXANDER BURNETT (T., 1921, 119, 2036—2044).

Excitation of Atoms to Emission of Light by Electronic Collisions. IV. Behaviour of Individual Members of a Series. R. SEELIGER (*Physikal. Z.*, 1921, 22, 610—613; cf. A., 1921, ii, 143, *Z. Physik.*, 1920, i, 355; 1921, 5, 182, 7, 93).—An examination of the spectra of helium, hydrogen, mercury, and neon produced by collisions with electrons shows that two kinds of series are produced, the one, which shows the regular behaviour, namely, that the optimal velocity increases with increasing member number, and the intensity maximum is displaced toward

the cathode with increasing member number, and the other, where the opposite behaviour is observed. To the first belong the Balmer series of hydrogen and both subsidiary triplet series of mercury, whilst to the second belong, the series of neon and helium.

J. F. S.

Entropy of Electron Gas. RICHARD C. TOLMAN (*J. Amer. Chem. Soc.*, 1921, **43**, 1592—1601).—It is shown that considerable simplification in expression and thought can be achieved by treating electrons as atoms of a chemical substance. This substance, which may be regarded as a gas "*plus que parfait*," owing to the very considerable force of repulsion between electrons, will not deviate appreciably in behaviour from a perfect monatomic gas if the concentration is low enough and the temperature high enough so that the effects produced by the repulsive force between the electrons can be neglected in comparison with the effect produced by their thermal agitation. It is shown with a reasonable certainty that the entropy of electron gas may be calculated from the theoretical equation, $S = 5/2R \log_e T - R \log_e p + 3/2R \log_e M + S_1$, which is found to apply to the monatomic gases. Using this equation, the entropies of 1 mol. of electron gas in equilibrium with metal filaments at 2000° are found to be: tungsten 53.3, tantalum 52.3, molybdenum 51.0. The entropy may also be calculated from the heat of vaporisation of the electron gas by the equation $S = \Delta H/T$, in which ΔH is the heat absorbed when 1 mol. of electrons is reversibly evaporated at T . The following values are obtained: tungsten, $\Delta H = 107270$ cal., $S = 53.6$; tantalum, $\Delta H = 107970$ cal., $S = 54.0$; molybdenum, $\Delta H = 109770$, $S = 54.9$.

J. F. S.

Thermal Ionisation of Metallic Vapours. RICHARD C. TOLMAN (*J. Amer. Chem. Soc.*, 1921, **43**, 1630—1632).—Making use of work recently published on the entropy of electron gas (see preceding abstract) it is shown that further information on the thermal ionisation of metallic vapours may be obtained. Thus assuming that a metallic vapour can ionise into a positive ion and an electron thus: $R = R^+ + E^-$, it is shown that $\Delta F = EQ - 5/2RT \log_e T + (5/2R - 3/2R \log_e M_E - S_1)T$ which connects the free energy with the heat content and the entropy and is of the same form as the well known free energy equation $\Delta F = \Delta H_0 - \Delta C_p T \log_e T + IT$. The ionisation constant is given by the equation $\log_e K_p = -EQ/RT + 5/2 \log_e T - (5/2 - 3/2 \log_e M_E - S_1/R)$, or, putting in the numerical values of the constants $Q = 96540$ coulombs, M_E (the molecular weight of electron gas) $= 5.44 \times 10^{-4}$ gram. $S = -3.2$ cal./degree and $R = 1.985$ cal./degree the equation becomes: $\log_e K_p = -5040E/T + 2.5 \log T - 6.69$, where E is the ionisation potential in volts. Using the above formulæ, the author calculates the ionisation of calcium in the sun. Taking the ionisation potential as 6.1 volts, the partial pressure of un-ionised calcium vapour as 4 atms., and the temperature as 5500° abs., it is found that the partial pressure of the calcium ion or electron gas is 0.07 atm., corresponding with 1.7% ionisation.

J. F. S.

New Detectors of High Frequency Vibrations. W. HEIKE (*Z. anorg. Chem.*, 1921, 118, 255—263).—An account of experiments made with the object of discovering an artificial detector for use as a substitute for pyrites in wireless telegraphy. Preparations of zinc arsenide, ZnAs_2 , containing crystals of zinc antimonide, ZnSb , proved specially sensitive. The zinc-arsenic alloys are discussed in the succeeding paper (this vol., ii, 60).

E. H. R.

Conductivity of Solutions of Ternary Electrolytes in Propyl Alcohol. CHARLES A. KRAUS and JOHN EGBERT BISHOP (*J. Amer. Chem. Soc.*, 1921, 43, 1568—1575).—The electrical conductivity of anhydrous calcium nitrate and magnesium nitrate hexahydrate in anhydrous propyl alcohol and magnesium nitrate hexahydrate in mixtures of propyl alcohol and water has been determined at 18° for a series of concentrations. The present results are compared with those of Schlamp (*A.*, 1894, ii, 376) on lithium and calcium chlorides and it is shown that his results indicate that the calcium chloride must have been hydrated. Neither the present work nor that of Schlamp supports the view that ternary salts ionise in this solvent according to a binary process. The specific conductivity of anhydrous propyl alcohol at 18° is 1.7×10^{-7} .

J. F. S.

The Theory of Electrolytic Ions. XXVI. How is the Limiting Value of the Molecular Conductivity of Strong Electrolytes Determined? RICHARD LORENZ (*Z. anorg. Chem.*, 1921, 118, 209—222; cf. *A.*, 1921, ii, 158, 481, 482, 483).—A graphic method for determining the limiting molecular conductivity μ_0 of univalent electrolytes is described, based on Herz's theory of ionic conductivity (*A.*, 1921, ii, 482). According to this theory, the conductivity of a single ion is expressed by the equation $B(\lambda_0 - \lambda) = \psi(\omega)$, where B is a constant and $\omega = A[S]^{\frac{1}{2}}$, where A is another constant and S is the concentration. The ionic conductivity λ of the theory is identical with the molecular conductivity, μ . The graphic method described depends on the fact that, by means of the constants A and B , the $[S]^{\frac{1}{2}}$ -conductivity curve of any substance can be transformed to coincide with a "universal conductivity curve." The molecular conductivities as ordinates are plotted against $[S]^{\frac{1}{2}}$ as abscissæ on a prescribed scale, and by a simple geometric process the curve obtained is transformed into a form of the universal curve. By comparing this derived curve with a series of standard curves printed on transparent paper on the same scale, the values of μ_0 and of the constant B can be read off on the ordinate axis.

E. H. R.

Concentration and Potential Fall at Boundary Surfaces. H. FREUNDLICH (*Z. Elektrochem.*, 1921, 27, 505—511).—A theoretical paper in which the change in concentration and the *E.M.F.* at the boundary surfaces of two liquids is considered. It is shown that the concentration of saponin or amyl alcohol in the foam produced at the air-water interface is due to the lowering of the

surface energy at the interface. All cases of adsorption are shown to be fundamentally due to the same cause. The case of adsorption on crystal faces is particularly considered, and although in this case the adsorption is held to be due to residual affinity, it is shown that here the phenomenon is molecular-kinetic, whilst with the surface tension a thermodynamic function is employed. In the case of surface tension, the mean value of a large number of molecules is considered, whilst with crystal surfaces when considering the residual valencies individual molecules only are concerned. The *E.M.F.* existing at interfaces and the changes effected by the introduction of foreign non-electrolytes is considered.

J. F. S.

The Normal Potential of Silver. I. M. KOLTHOFF (*Z. anorg. Chem.*, 1921, 119, 202—212).—The electrolytic potential of silver was measured in silver nitrate solutions varying in strength from 0.001 to 0.1*N* at 18° against a normal calomel electrode. The mean value found for the silver potential was -0.7887 volt against the normal hydrogen electrode. Potassium nitrate, up to a concentration of 1.0*N*, has no measurable effect on the potential of the silver electrode. From measurements with a silver-silver bromide electrode in solutions of potassium bromide from 0.001 to 0.5*N*, the solubility product of silver bromide at 18° was found to be 4.1×10^{-13} . The concentration of bromine ions can be calculated from the equation $\log [\text{Br}] = (-\pi - 0.34852)/0.0576$. In presence of sulphates in solution, the silver electrode becomes "nobler"; the same is true of the mercury electrode. E. H. R.,

Some Phenomena in the Electromotive Behaviour of certain Metals. NIL RATAN DHAR (*Z. anorg. Chem.*, 1921, 118, 75—80).—If the potential difference between a metal and a solution of one of its salts increases with increasing dilution, it is to be expected, by extrapolation of Nernst's formula, that the potential difference would be still greater when the metal is in contact with water or a solution of a neutral electrolyte. Conversely, if the *E.M.F.* decreases as the dilution increases, it should be still less in water or a solution of a neutral electrolyte. A large number of measurements were made to test these theoretical deductions. The metals magnesium, zinc, and aluminium gave each a smaller *E.M.F.* in water or a neutral electrolyte than in a solution of its own salt, instead of greater, as was to be expected. Similarly, nickel and tin, by giving greater *E.M.F.* in water or neutral electrolytes, behaved in a manner contrary to what was expected. Only the metals antimony, copper, and silver, which have small solution tensions, behaved in the way indicated by theoretical considerations. A large number of measurements were made of the *E.M.F.* given by pairs of the above metals in water and potassium chloride solution respectively.

E. H. R.

Electromotive Behaviour of Aluminium. A. SMITS (*Z. Elektrochem.*, 1921, 27, 523—526).—Polemical. An answer to the criticism of Gunther-Schulze (*A.*, 1921, ii, 535) to the author's

paper (A., 1920, ii, 579) in connexion with anodic polarisation and passivity. The author maintains his former position. J. F. S.

Electromotive Behaviour of Aluminium. III. A. SMITS and C. J. DE GRUYTER (*Proc. K. Akad. Wetensch. Amsterdam*, 1921, **24**, 86—91; cf. A., 1920, ii, 579; 1921, ii, 371).—A continuation of previous work (*loc. cit.*). The influence of mercury on the potential of aluminium immersed in a solution of aluminium sulphate has been investigated. The aluminium electrode was placed in a solution of aluminium sulphate and small quantities of a mercury salt were added at intervals, and the potential was measured repeatedly. The influence of oxygen on the process was prevented by a stream of nitrogen which was continuously bubbled through the solution. It was found that the potential of the aluminium at first became less negative, reached a minimum, after which it rose to a strongly negative value which increased to a maximum. The maximum value was maintained for some time and then fell to a value which was close to the mercury potential. Up to the point where the maximum was observed, the aluminium electrode was grey in colour, but during the last stage it became lustrous, due to a coherent film of mercury. The changes are considered in connexion with the three-phase equilibrium of the four component system Al^{3+} — Hg^{2+} —anion— H_2O .

J. F. S.

Electrochemical Investigations of Gold-Copper Mixed Crystals. RICHARD LORENZ, W. FRAENKEL, and M. WORMSER (*Z. anorg. Chem.*, 1921, **118**, 231—253).—In connexion with the study of discharge and decomposition potentials, experiments were made in which a series of gold-copper alloys were used as anodes in an electrolytic cell. For studying the discharge potentials, $N/10$ -sodium hydroxide was used as electrolyte, with a copper cathode, and the cell was charged for a short time, about two minutes, with an *E.M.F.* of about 4 volts. The discharge potential was then plotted against time. With pure copper and with alloys containing less than 0.2 atom of gold, a distinct arrest occurred in the curve, whilst alloys with 0.2 atoms of gold and upwards behaved like pure gold, giving a smooth discharge curve. The charging, or decomposition, potentials were studied, using as electrolytes first a solution containing sodium hydroxide and ammonia, and then copper sulphate. In the former case, copper or an alloy with 0.1 atom of gold showed an early rise of current with *E.M.F.*, followed by a fall to zero and then a steady rise starting at about 0.7 volt. With 0.2 atom or more of gold, no current passed until the *E.M.F.* reached 1.05 volts, and the current increased steadily with the *E.M.F.* Using copper sulphate as electrolyte, the decomposition potential rose steadily with the gold content of the alloy up to about 0.25 atom of gold, and from 0.35 atom of gold upwards was indistinguishable from that of pure gold. This is in agreement with Tammann's observations on the limiting composition in gold-copper alloys (A., 1920, ii, 672). It is shown experimentally that

changes in the resistance of the cell during the experiments have no appreciable influence on the results obtained, and it follows that the maximum and minimum in the decomposition potential curves and the arrest point in the discharge potential curves for copper and the alloys less rich in gold have a common cause.

E. H. R.

Metallographic Investigations on the Cathodic Deposition of Metals on Aluminium and Chromium. S. KYROPOULOS (*Z. anorg. Chem.*, 1921, **119**, 299—304).—The cathodic deposition of copper, silver, chromium, and nickel on aluminium and chromium was studied. Copper and silver from nitrate solution, with a current density of 0.01 to 0.1 amp. per sq. cm., are deposited on tempered aluminium principally on the intercrystalline boundaries and to a very limited extent on the crystal faces, whilst chromium and nickel are deposited more freely in isolated spots on the crystal faces. With chromium and nickel, a higher current intensity favours deposition on the crystal faces of the aluminium. From cyanide solution copper may be deposited on the crystal faces when the current intensity is high; on pressed aluminium it is deposited along the gliding lines. It is noteworthy that deposition on the crystal faces is favoured under conditions such that production of hydrogen at the cathode is possible. Resistance to copper deposition is most clearly shown by passive chromium, deposition occurring only on isolated spots of non-passive metal or impurity. Under conditions which destroy the passivity of the chromium, such as hydrogen evolution at the cathode, deposition occurs on the crystal faces of the chromium.

E. H. R.

Electrolysis of Aqueous Solution of Cerous Salts. A. B. SCHIØTZ (*Z. Elektrochem.*, 1921, **27**, 521—523; cf. A., 1921, ii, 589).—After a short account of the work which has previously been carried out on the electrolysis of solutions of the rare earth salts, and of that on the cerium alloys, experiments are described on the electrolysis of concentrated solutions of mixtures of cerous and ferrous chlorides to which various additions had been made. The electrolysis took place between a platinum spiral which served as anode and a platinum gauze covered with lead dioxide which served as cathode. The solutions, which were used warm, contained 30 grams of lactic acid neutralised by sodium or potassium hydroxide, and 5—10 grams of sodium chloride in 400 c.c. of water to which was added cerous chloride (7.0—3.5 grams) and ferrous chloride (4.5 grams). The electrolysis was effected with a current density of 0.4—2.0 amp./sq. cm. and yielded in all cases a black, powdery deposit which was metallic and contained about 62% of cerium. Whether or no it contained oxygen could not be ascertained because of the difficulty of drying the substance without oxidation. The deposit corresponds with the iron-cerium alloy, Ce_2Fe_3 , prepared by Vogel (A., 1912, ii, 557). The electrolysis of cerous chloride in the presence of chloride of mercury, nickel, platinum, zinc, or aluminium gave deposits which contained no

cerium. The addition of potassium lactate to solutions of cerous chloride raises the overvoltage by 0.15 volt. J. F. S.

Biochemical and Electrochemical Oxidation of Organic Compounds. FR. FICHTER (*Z. Elektrochem.*, 1921, 27, 487—494).—A general discussion of electrolytic and biochemical oxidation of organic substances in which a number of similarities in the two processes are indicated. J. F. S.

Magnetism and the Constitution of Atoms. PIERRE WEISS (*Rev. sci.*, 1920, 58, 645—659).—A review of work already published (see A., 1911, ii, 91, 183, 250, 367, 694; 1915, ii, 737).

CHEMICAL ABSTRACTS.

Behaviour of Substances near the Absolute Zero. WILLIAM R. FIELDING (*Chem. News*, 1921, 123, 97—99).—The results previously obtained by the author (A., 1920, ii, 732; 1921, ii, 487) are applied in the present paper to several substances in the neighbourhood of the absolute zero. Making use of the specific heat of lead determined by Kammerlingh Onnes and Keesom over the range 46°Abs.—14°Abs., the author has determined the specific heat over the range 14°Abs.—0°Abs. by three methods and finds that by plotting the specific heat curve the specific heat appears to be zero at 6°Abs., by plotting the specific heats against the absolute temperatures the values 10°Abs., $s=0.0053$, 2°Abs., $s=0.0012$ and 1°Abs., $s=0.00005$ are obtained, whilst applying Debye's law (atomic heat= kT^3) leads to values which demand that the molecule of lead must change from Pb_8 to $Pb_{(3000-4000)}$. Similar calculations are made for water, copper, and carbon, and the absolute molecular weight is calculated for the absolute zero. The values recorded are, water 540, lead 1456, and copper 9450. If the values of p , the number of single molecules in the complex molecule, are plotted, it becomes possible to obtain by extrapolation the values for other elements. The values of p are shown to be periodic, thus, iodine 6.3, bromine 6.6, chlorine 14, and fluorine 24 all at 0°Abs. In the case of mercury, p is slightly greater than 2. The number of atoms in the benzene, naphthalene, and anthracene polymerides at the absolute zero are respectively 84, 100, and 132. J. F. S.

The Properties of Organic Liquids. W. HERZ (*Z. anorg. Chem.*, 1920, 118, 202—206).—The relation $T_c/d_c\sqrt{n}=\text{const.}$, where n is the number of atoms in the molecule, has been found to hold for 27 non-associated organic liquids and to have a value approximately equal to 450. When this is combined with the formula $n=T_c/0.9p_c$ (A., 1920, ii, 285), the relation $T_cp_c/d_c^2=\text{const.}$ is obtained, which should have a value of about 225×10^3 . This relation was verified for the above 27 liquids, with the exception that acetylene and carbon dioxide give respectively high and low values. It was shown previously (A., 1920, ii, 736) that the relation $d_c=M/9.24z$, where z is the total of the valencies of the atoms in the molecule ($C=4$, $N=3$, $O=2$, $H=1$), also holds for

non-associated liquids. With the first equation above, this gives $T_c z / M \sqrt{n} = 48.7$, which has been verified for a number of aliphatic liquids. In the case of aromatic liquids, the relation holds if the valency of carbon is taken as 3. From Trouton's rule $M.L/T_c = 14$, it follows that $Lz/\sqrt{n} = 686$, and this is also found to hold for a number of fluids, but aniline and dimethylcyclohexane give high values. Finally, from the relation $L = 0.666 p_c / d_c$, it follows that $p_c z / d_c \sqrt{n} = 1030$, and this relation is also satisfactorily verified in most cases, but acetylene gives a high value. It is pointed out that these relations can only be tested for relatively simple substances, of which the critical data do not differ widely. Not improbably complications may arise when data are available for other types of substance.

E. H. R.

General Improvement of the Equation of Condition : Special Equation of Condition for Hydrogen. LÉON SCHAMES (*Physikal. Z.*, 1921, 22, 630—634).—A theoretical paper in which the constants a and b of van der Waals's equation are considered on the basis of a perfectly rigid atom. A general equation of condition is evolved, and taking account of the compressibility of the atoms the equation $pv(1-b/v) + a_0/v(\alpha - v_0/v) = RT$ is deduced for hydrogen.

J. F. S.

Joule-Thomson Effect for Air. FREDERICK G. KEYES (*J. Amer. Chem. Soc.*, 1921, 43, 1452—1470).—The Joule-Thomson coefficients of air have been calculated from the author's equation of state for air based on the measurements of pressure, volume, and temperature by Amagat (*Proc. Nat. Acad. Sci.*, 1917, 3, 323), and have been compared with the coefficients as measured by Joule and Thomson, by Noell, and by Hoxton. The measurements of Joule and Thomson accord well with the calculations except at zero degrees, whilst the recent careful measurements of Hoxton (*Physical Rev.*, 1919, 13, 438) exceed the calculated values between 0° and 100° by about 15%. The value of the ice-point absolute temperature (273.36) deduced from the Chappuis 1000 mm. constant-pressure expansion coefficient for air, corrected by means of the Hoxton Joule-Thomson coefficients is too large, due to the fact that the Chappuis expansion coefficient is too small and the Joule-Thomson values too large. The calculated expansion coefficient and the Hoxton and Joule-Thomson coefficients lead to $T_0 = 273.197$. The value $T_0 = 273.135$ is, on the other hand, shown to be substantially correct in that this value is the mean value obtained by graphically extrapolating the measured constant volume and constant pressure expansion coefficients of air, nitrogen, hydrogen, and helium measured at various ice-point pressures. The value 273.135 is also the mean of T_0 as deduced from the same data by means of the equations of state for the respective gases. The pressure coefficient at zero of the Joule-Thomson coefficient as obtained by Hoxton is shown to be much larger than that obtained by other observers. The pressure coefficient given by Vogel and Noell (8.8×10^{-4}) is twice that calculated from the equation

(4.5×10^{-4}) for low pressures, but agrees approximately with that calculated for high pressures, by means of the equation of state. The Bradley and Hale average pressure coefficient to 200 atmospheres at 0° is shown to be 7.3×10^{-4} , whilst the average value over this range calculated by the equation of state is 7.1×10^{-4} . The Joule-Thomson coefficient itself at one atmosphere, as given by Vogel and Noell, is 0.277° , and from Bradley's work 0.268° . The equation gives 0.261° . An explanation of the discrepancies in the observed Joule-Thomson numbers is suggested which assumes that at the low pressure surface of the plug a temperature drop is superposed on the Joule-Thomson temperature change due to the momentary persistence of the fine gas streams issuing from the pores of the plug and their subsequent expansion. It is shown that this effect would produce a large apparent pressure coefficient, since the specific effect of the velocity-persistence-effect is shown to vary inversely as the pressure. The value of the measured Joule-Thomson effect at 0° and one atmosphere pressure is given as 0.303 by Hoxton, whilst that deduced from the high pressure measurements of Bradley and Hale is 0.268 , as compared with 0.261 calculated by the equation of state. The Joly constant-volume specific heats of air are considered, and it is shown that a correction is required to compensate for the increased specific heat of the copper sphere used by Joly to contain the air. The corrected Joly values are shown to be independent of pressure, which fact confirms the validity of the form of the air equation of state employed. As a further consequence, it is pointed out that at low pressures the Joule-Thomson effect must vary inversely with the absolute temperature. Another important consequence is that the constant volume air thermometer scale reads directly on the absolute scale and therefore requires no correction. J. F. S.

Simple Methods for the Determination of Melting Points and Critical Temperatures. The Melting Point of Arsenic. W. HEIKE (*Z. anorg. Chem.*, 1921, 118, 254).—In a paper under the above title, Rassow (A., 1921, ii, 164) quotes the author's determination of the melting point of arsenic, 830° . The author points out that later work gave a value 814.5° , which is more trustworthy (A., 1916, ii, 248). E. H. R.

Theory of Concentrated Solutions. Separation of Solids from Organic Mixtures. J. TIMMERMANS (*Bull. Soc. chim. Belg.*, 1921, 30, 276—290; cf. A., 1909, ii, 388).—From a study of experimental data in the literature relating to the freezing points of binary organic mixtures, it is concluded that deviations from the behaviour of ideal mixtures cannot in all cases be accounted for by assuming combination between the components or polymerisation, as suggested by Dolezalek and others (A., 1909, ii, 22; also Ikeda, A., 1908, ii, 932). It is necessary to take into account also certain physical factors, in the sense of van der Waals's theory, in particular for systems of limited mutual solubility. A provisional scheme of classification of binary organic liquids is based on these chemical and physical factors.

Freezing-point data respecting numerous binary mixtures containing benzene as the crystallising component were compared with the ideal curve obtained by plotting T and C from the thermodynamic equation: $\log_e C = Q(T_0 - T)/2T_0T$, in which Q is the molecular heat of fusion of the component which separates, C is the number of molecules of the latter per 100 molecules of the mixture, T is the temperature at which this component separates, and T_0 is the solidifying point of the same substance in the pure state (cf. Washburn, "Introduction to the Principles of Physical Chemistry," 1915, ch. 14). Experimental data for many of the mixtures agreed well with the curve, especially where the two components were closely related and not of widely different chemical character. Deviations from the curve in the direction of the axis of composition might reasonably be attributed to combination between the components, but deviations in the other direction could not in all cases be due entirely to polymerisation, and it is shown that they are to be expected if mixed crystals separate or if the components are not completely miscible in all proportions (cf. Kohnstamm, A., 1911, ii, 93). Similar considerations apply also to boiling-point curves, but abnormalities are here more frequent than with freezing-point curves. In many cases where the boiling-point curve is convex towards the axis of composition (cf. Dolezalek, *loc. cit.*), the assumption of polymerisation sufficient to account for the abnormality appears inadmissible (for example, for carbon tetrachloride, benzene, toluene, etc.), and here again the results may be due to the existence of a range of immiscibility. This explanation was verified in a number of cases, the suspected mixtures separating into two liquid phases when their temperature was lowered sufficiently, by means of liquid air or solid carbon dioxide.

J. H. L.

The Temperature of the Vapour Arising from Boiling Saline Solutions. GEORGE HARKER (*J. Roy. Soc. New South Wales*, 1920, 54, 218—226; cf. Sakurai, T., 1892, 61, 495).—Experiments with solutions of calcium chloride boiled in a hypsometer either by direct flame or by steam blown in, the only jacketing of the vapour column being by the slightly cooled issuing vapour, indicate that the vapour coming from a boiling solution has a higher temperature than the vapour from the boiling pure solvent.

W. G.

A Simple Receiver for Fractional Distillation under Diminished Pressure. E. J. WILLIAMS (*Chem. News*, 1921, 123, 265—266).—A simple receiver for fractional distillation under reduced pressure, working on the principle of that of Lothar Meyer, is figured and described. It is constructed with a small distillation flask, a small wide-necked tap funnel, and an adapter. W. G.

The Separation of Miscible Liquids by Distillation. ARTHUR FELIX DUFTON (*T.*, 1921, 119, 1988—1994).

Temperature Correction in Bomb Calorimetry. N. H. ENGSTRÖM (*Tekn. Tidskr. Kemi Berg.*, 1921, 51, 18—21).—A formula is evolved for the correction of temperature in bomb calorimetry. This formula is for general use where the tedious calculations by the application of the classical equation of Regnault-Pfaundler is not absolutely necessary. It is given as follows: $k = (1-n)v_1 - v$. The order and significance of the letters are the same as in the Regnault-Pfaundler equation. This new formula was tested by a large number of determinations and its error falls within the limits ± 0.001 — 0.003° . The error was seldom as high as $\pm 0.003^\circ$.
CHEMICAL ABSTRACTS.

Constitution and Heats of Combustion. F. OTTO H. BINDER (*Chem. Zeit.*, 1921, 45, 1114—1116; cf. A., 1921, ii, 241, 435).—In continuation of the method previously described (*loc. cit.*) for calculating the heat of combustion of a substance from its constitutional formula, the author has worked out from a consideration of a large number of hydrocarbons the average value of the following: Hydrogen atom attached to carbon 55560, carbon-carbon linking 16258, one carbon valency 24495, compared with 24240, the value obtained by calorimetric experiments on wood charcoal. The heat of combustion of a compound accordingly equals $cH \times 55560 + y \times 24240 + z \times 34170$, where cH is the number of carbon valencies bound to hydrogen, y the number not bound to either hydrogen or oxygen, and z the number of free carbon valencies. From consideration of hydrogen cyanide, cyanogen, and acetonitrile, a mean value for one carbon-nitrogen valency of 35301 cal. is obtained.
G. F. M.

Absolute Value of the Energy of the Linkings between the Atoms of Compounds. M. PADOA (*Gazzetta*, 1921, 51, ii, 239—245).—The method of calculation used by Fajans (A., 1920, ii, 354) is substantially that employed by the author (A., 1919, ii, 96), who, however, assumed the value 42 Cal. for the heat of sublimation of diamond and graphite indicated by Thomsen's results, whereas Fajans, on the basis of relationships established by Grüneisen between the heat of sublimation of a solid monatomic element and the ratio of atomic heat to the coefficient of thermal expansion, concluded that the heat of sublimation of carbon is at least 275 Cal. per gram-atom. In spite of the large difference between these two values, both authors conclude that the energy of the linkings between the carbon atoms is almost the same in diamond and in aliphatic hydrocarbons. It seems probable that neither of the above two values is the correct one, and the author considers 150 Cal. to be more nearly the truth; Kohn (A., 1921, ii, 302) arrives at values similar to this. Steiger's results (A., 1920, ii, 355, 735) are also discussed.

Taking the value 150 Cal. for the heat of sublimation of 1 gram atom of carbon and 81.3 Cal. for the heat of dissociation of 1 gram molecule of hydrogen, the energy of the C-C linking is calculated for the aromatic and aliphatic hydrocarbons previously considered.

The results show that the energy of the linking increases from simpler to more complex compounds. The differences of the values referred to graphite and to diamond vary from 5 to about 7% and are small considering that the side of the hexagon is 6.2×10^{-8} cm. in benzene and 1.45×10^{-8} cm. in graphite.

The energy of the simple linking in ethane is 69.9 Cal., that of the double linking in ethylene, 2×59.65 Cal., and that of the triple linking in acetylene, 3×53.33 Cal. The heat of combustion of acetylene (3 mols.) being 938.4 Cal. and that of benzene vapour 783.3 Cal., the difference between the three triple linkings and the nine simple linkings is 155.1 and that between one triple linking and three simple linkings hence 51.7 Cal.; the numbers just given for ethane and acetylene lead to the value 49.7 Cal. The absolute values of the double and triple linkings are, therefore, positive and unsaturated hydrocarbons all highly exothermic when formed from the free atoms. These considerations have been extended to various endothermic reactions, such as the formation of hydrogen telluride, the formation of cyanogen from carbon and nitrogen, the formation of nitrous and nitric oxides, the results indicating that, in all probability, all compounds must be formed from free atoms with generation of heat.

T. H. P.

Compressibility of Benzene, Liquid and Solid. THEODORE W. RICHARDS, EDWARD P. BARTLETT, and JAMES H. HODGES (*J. Amer. Chem. Soc.*, 1921, **43**, 1538—1542).—The compressibility of liquid benzene has been determined at 20°, solid benzene at 0°, and toluene at 0° and 20°, using a steel piezometer previously described (Richards and Bartlett, A., 1915, ii, 229). It is found that the compressibility of solid benzene at 0°, $(\Delta v / \Delta P v_0)_T$, is 0.0000305 over a pressure range 100 to 500 megabars. Incidentally, the compressibility of liquid toluene at 0° over the same pressure range was found to be 0.0000618. A new determination of the compressibility of liquid benzene at 20° confirmed satisfactorily the previous determination made in a glass piezometer (Richards and Shipley, A., 1916, ii, 376), giving the value 0.00007207 for the same range. The compressibility of solid benzene is shown to be less than half that of liquid benzene at the same temperature. Thus another case, supporting the majority of those investigated, is found in which the solid shows a much smaller compressibility than the liquid (A., 1915, ii, 518). No interpretation of this result is offered at present.

J. F. S.

Relative Volumes of the Chemical Elements. HAWKSWORTH COLLINS (*Chem. News*, 1921, **122**, 76—77, and **123**, 95—97).—A continuation of previous work (A., 1921, ii, 168). In the first part, a table is given of the relative volumes and the densities of a number of organic compounds, the latter quantity is compared with the experimentally determined values. The calculated results are based on the hypothesis previously put forward by the author. In the second part, a further long list is given. It is shown that the experimental facts obtained from the examination of 76 mole-

oules demonstrates that one relative volume of hydrogen is 15.25 at 15° whether, Cl, Br, I, NH_2 , NO_2 , SH, C_2H_3 , C_2H , CO_2Me , CO_2Et , CO_2Pr , CO_2H or $-\text{OH}$ be displaced; in this case $-\text{OH}=13.29$. In a further series of 24 compounds, the relative volume of hydrogen is 12.22 at 15° for the same replacements, $\text{OH}=10.21$. The methyl group has a volume 31.67 at one end of a chain and 28.64 at the other end. When an atom of hydrogen the volume of which is 15.25 is displaced by methyl, the volume of the latter is 31.67, and when the displacement occurs with hydrogen 12.22, the volume of the methyl is 28.64. The two valencies of carbon concerned are distinguished by the fact that one (first position) has hydrogen with a volume 15.25 and the other (second position) has hydrogen with a volume 12.22.

J. F. S.

Atomic and Molecular Volumes at the Absolute Zero. II.

W. HERZ (*Z. anorg. Chem.*, 1921, 119, 221—224).—In a previous paper it was shown by Lorenz and Herz (*A.*, 1921, ii, 536) that the molecular volumes of alkali salts reduced to absolute zero are smaller than the sums of the volumes of the constituent atoms, similarly reduced according to the law of corresponding states. The inquiry has been extended to a number of simple inorganic compounds other than salts. In the hydrogen compounds the molecular volume is smaller than the sum of the atomic volumes, but the difference becomes smaller with increasing molecular weight. In the halogen acids, for example, these differences are, HF, 7.52; HCl, 3.09; HBr, 1.09; HI, 0.01. In the trihaloids of the elements of the fifth group, however, the molecular volume is greater than the sum of the atomic volumes. The molecular volumes are also calculated from the dielectric constant by the Clausius-Mossotti equation $\gamma = D - 1/D + 2.M/d$. The values obtained are sometimes greater and sometimes smaller than the actual molecular volumes. The divergences may be due to association in the liquids.

E. H. R.

Surface Tensions and Densities of Liquid Mercury, Cadmium, Zinc, Lead, Tin, and Bismuth. THORFIN R. HOGNESS (*J. Amer. Chem. Soc.*, 1921, 43, 1621—1628).—The densities of liquid zinc, cadmium, tin, lead, and bismuth have been determined at various temperatures and the following values obtained: zinc $d=6.59-0.00097(t-419)$, lead $d=10.71-0.00139(t-327)$, cadmium $d=8.02-0.00110(t-320)$, bismuth $d=10.07-0.00125(t-269)$, and tin $d=7.01-0.00074(t-232)$. The surface tension for mercury and the above-named molten metals has been determined by the Cantor drop pressure method for a series of temperatures. The values of the surface tension may be calculated by the following equations: mercury $r=467-0.043(t+39)-0.000386(t+39)^2$, bismuth $r=378-0.063(t-269)$, cadmium $r=630-0.065(t-320)$, lead $r=444-0.077(t-327)$, tin $r=531-0.080(t-232)$, and zinc $r=758-0.090(t-419)$. It is shown that there is no appreciable difference between the surface tensions of mercury in dry air, hydrogen, and a vacuum.

J. F. S.

Relationship of the Internal Friction of Organic Liquids to other Properties. W. HERZ (*Z. Elektrochem.*, 1921, 27, 518—521).—In the case of a large number of simple non-associated organic liquids an approximately constant relationship exists between the surface tension and the critical pressure. This relationship is $\gamma_s : p_c \triangleq 1.6 \times 10^4$, in which γ_s is the surface tension at the boiling point and p_c the critical pressure. Between the surface tension and the internal friction (η_s), both at the boiling point, the approximate relationship $\gamma_s : \eta_s \triangleq 6 \times 10^3$ exists. It is also shown that the two expressions, $M\gamma_s/T_b d_s = \text{const.}$ and $Ld_s/\gamma_s = \text{const.}$, both hold for non-associated organic liquids, where M is the molecular weight, L the latent heat of vaporisation, and d_s the density at the boiling point T_b . In the case of the former equation, the constant value varies between 6.7×10^4 and 11.3×10^4 , whilst in the latter the value is between 2.0×10^4 and 3.2×10^4 . Two relationships connecting the internal friction and the critical temperature respectively with the molecular refractivity are deduced. These have the form $\eta_s MR/T_c = \text{const.}$ and $\eta_s MR/T_b = \text{const.}$, where MR is the molecular refractivity. The constant in the first case varies between 1.1×10^{-4} and 1.7×10^{-4} and in the second case between 1.6×10^{-4} and 3.2×10^{-4} . If n is the number of atoms in a non-associated liquid, the quantity $\eta_s n/T_b$ is constant and has a value between 0.86×10^{-4} and 1.39×10^{-4} . Finally, it is shown that $V_c \eta_s/T_b$ and $V_c \eta_s/T_c$ are both constant for the same class of organic liquid. In the first case the value varies between 9.0×10^{-8} and 12.6×10^{-8} , and in the second case between 5.8×10^{-8} and 8.0×10^{-8} . J. F. S.

Theory of Brownian Movement. GUSTAV JÄGER (*Sitzungsber. K. Akad. Wiss. Wien*, 1919, [2a], 128, 1271, 1298; from *Chem. Zentr.*, 1921, iii, 833).—Stokes's law is held to be inapplicable in the case of Brownian movement. It is shown from consideration of mechanical examples that if this law holds, osmotic pressure must be exerted as a continuous force and the consequences of Einstein's equation for Brownian movement and the calculation of molecular size from diffusion are justified. The distribution of energy for particles of any number of molecules may be obtained from Maxwell's law directly by gradual building up of groups from molecules. The velocity of particles relative to the liquid medium, including their Brownian movement, is only to a small degree dependent on their mass. G. W. R.

Influence of Water-content on the Adsorption-capacity of an Active Charcoal. HEINRICH HERBST (*Biochem. Z.*, 1921, 118, 103—119).—The highest adsorption is obtained with dry charcoal. For practical purposes 10% of water is not harmful, since the presence of water has a catalytic effect. H. K.

Absorption of Gases by Crystals. R. SEELIGER and K. LAPKAMP (*Physikal. Z.*, 1921, 22, 563—567).—Partly or completely dehydrated zeolites have the power of absorbing gases in large quantities. The absorption of a mixture of neon and helium,

nitrogen, oxygen, methane, nitric oxide, carbon dioxide, acetylene, hydrogen, and ammonia by dehydrated chabasite ($\text{CaAl}_2\text{Si}_4\text{O}_{16}\cdot 6\text{H}_2\text{O}$) has been examined at 0° under a series of pressures from 0° to 760 mm. It is shown that, with the exception of hydrogen, the amount of gas absorbed is greater the higher the critical temperature of the gas. The following volumes of gases in c.c./gram of chabasite at 0° and 760 mm are absorbed. Neon 25% + helium 75%, 4.10; nitrogen, 6.89; oxygen, 8.32; methane, 15.18; nitric oxide, 25.67; carbon dioxide, 39.60; acetylene, 56.46; hydrogen, 82.22, and ammonia, 180.59. The adsorption curves can be approximately reproduced by the formula $v = \alpha p^{1/n}$, in which α and $1/n$ are constants. All gases except nitric oxide, ammonia, and impure acetylene can be entirely removed from the crystals by lowering the pressure. The results do not indicate whether the absorbed gases are adsorbed according to the usual conception of the term, or whether they enter into the structure of the crystal lattice. In favour of the pure adsorption are the qualitative character of the adsorption isotherm, the connexion between the critical temperature and the amount of absorption, and the effect of reducing the pressure. Further, the connexion between $1/n$ and the critical temperature also indicates a pure adsorption. On the other hand, the large absorption of hydrogen and the difficulty experienced in removing nitric oxide, ammonia, and impure acetylene from the crystals point to the view that the absorbed gases enter into the crystal lattice in the position occupied by the water before dehydration.

J. F. S.

Graphic Deductions with regard to the Dissociation of Weak Electrolytes. N. SCHOORL (*Rec. trav. chim.*, 1921, 40, 616—620).—An expression is deduced from Ostwald's dilution law which gives approximately correct values for the degree of ionisation of weak electrolytes. Such approximation is depicted graphically in contrast with the calculated values.

H. J. E.

The Redissolution of a Salt during the Isothermal Evaporation of a Solution. ETIENNE RENGADE (*Compt. rend.*, 1921, 173, 1080—1081; cf. A., 1921, ii, 93).—A reply to Raveau (cf. A., 1921, ii, 682).

W. G.

Formulae for the Solubility of certain Salts in Aqueous Ethyl Alcohol and Water. W. D. TREADWELL (*Helv. Chim. Acta*, 1921, 4, 982—991).—Attempts are described to devise a mathematical expression for the diminution of the solubility of a salt in water by the addition of alcohol on the assumption that the solvent action is exerted by the water alone, and that its extent is lessened by its dilution with the alcohol. Consideration is confined for the present to substances which are almost insoluble in absolute alcohol. In these circumstances, the solubility is expressed by the equation $h = L_0 C_w^k$, in which L_0 is the solubility of the salt in pure water, C_w the concentration of the water in the solvent, and k a proportionality factor of the approximate order, 1. The results calculated from this formula ($k = 0.88$) are in harmony with the experimental data for the solubility of potassium chlorate in

mixtures of alcohol and water at the atmospheric temperature, at 30°, and at 40°. The influence of temperature, therefore, is manifest essentially in the alteration of the value, L_0 . Within certain limits of alcoholic concentration, similar observations are made with potassium nitrate, but, in this case, the value of k does not appear to be so independent of the temperature. At 14.5°, the solubility of potassium chloride is represented satisfactorily by the expression when $k=0.93$, but at 30° and 40° a distinct specific action of the alcohol is observed which causes a variation in the value of k with varying alcoholic content of the solution. This specific action is also in evidence in the cases of sodium chloride, etc., more noticeably, of ammonium sulphate.

The remarkable independence of the value of k of the temperature in the case of potassium chlorate and, to a more limited extent, in that of potassium nitrate points to the conclusion that alcohol behaves merely as an inert diluent. L_0 may therefore be expressed as a function of the temperature and replaced by the expression aK^n in which a and n are empirical constants and K is the ionisation product of water $\times 10^{14}$. The solubility of potassium chlorate in any mixture of alcohol and water between 12° and 40° is given by the expression: $L=8.4 \times K^{0.482} \cdot C_w^2 \times 0.88 = 7.4 \cdot K^{0.482} \times C_w^2$. That of potassium nitrate, sodium chloride, potassium chloride, potassium sulphate, and lead nitrate can be calculated satisfactorily if a constant value is given to n , but, in the case of sodium chloride a distinct linear variation of the exponent is noticeable which, if taken into consideration, permits a very exact calculation of the observed values. The solubility of non-ionised mercuric chloride cannot be calculated accurately if n is assumed to be constant.

H. W.

Application of Ideal Solution Equations to Dilute Aqueous Solutions. JAMES KENDALL (*J. Amer. Chem. Soc.*, 1921, 43, 1391—1396).—A theoretical paper in which it is shown that the present method of applying the fundamental ideal solution equation to dilute aqueous solutions, particularly in the vapour pressure and freezing-point depression equations, leads to certain incompatibilities, which have been pointed out and discussed. It is also shown that the discrepancy disappears when due notice is taken of the fact that the vapour pressure of water at the ordinary temperature must be ascribed practically entirely to its uni-molecular fraction, the shift in the equilibrium $(H_2O)_g \rightleftharpoons qH_2O$ on addition of a solute counterbalancing the effect of the association on the molecular fraction. Some fundamental errors in the present application of ideal osmotic pressure and freezing-point depression equations to dilute aqueous solutions have also been pointed out and corrected.

J. F. S.

Factors Affecting the Stability of Additive Compounds in Solution and their Influence on Ionisation Equilibria. I. JAMES KENDALL and PAUL M. GROSS (*J. Amer. Chem. Soc.*, 1921, 43, 1416—1426).—In an earlier paper (Kendall, Booge, and Andrews, *A.*, 1918, ii, 36, 37), it is postulated that the formation of solvent-

solute complexes is a pre-requisite to ionisation in solutions. On this assumption, it should be possible to predict the degree of dissociation in any given solution from (a) the stability of the complexes with respect to their components, and (b) the instability of the complexes with respect to their ionisation products. The first of these points has been dealt with in detail in previous papers (*loc. cit.*), and the rules formulated have here been extended to the association of pure liquids. In the present communication the study of the second point—the factors affecting the ionic disintegration of solvent-solute complexes in solution—has been commenced. These generalisations are shown to be in excellent qualitative agreement with the experimental data of earlier investigators. A more stringent test of their validity is furnished by careful conductivity determinations on specially selected systems (*cf.* following abstract). The connexion between the ionisation hypothesis presented here and the views of Abegg and Bodländer, on the one hand, and of Werner, on the other, has been shortly examined and some points of difference have been indicated.

J. F. S.

Compound Formation and Specific Conductivity in Solutions of the Types, Acid-Ester, Acid-Ketone, and Acid-Acid. JAMES KENDALL and PAUL M. GROSS (*J. Amer. Chem. Soc.*, 1921, 43, 1426—1439).—Complete specific conductivity-composition curves have been produced for fourteen systems of the types, acid-ester, acid-ketone, and acid-acid, at 25°. The substances employed are all liquids, and the specific conductivity of these was taken as the test of purity. It is shown that this property is far more sensitive to small amounts of impurities than the melting point or boiling point; the specific conductivity at 25° is found for the substances named to be: ethyl acetate, $<1 \times 10^{-9}$; ethyl benzoate, $<1 \times 10^{-9}$; benzyl benzoate, $<1 \times 10^{-9}$; acetone, 5.8×10^{-8} ; acetophenone, 5.5×10^{-8} ; acetic acid, 2.4×10^{-8} ; propionic acid, $<1 \times 10^{-9}$; monochloroacetic acid, 1.4×10^{-6} (60°); trichloroacetic acid, 6.2×10^{-9} (60°), and formamide, 4×10^{-6} . These results are from one-third to one-six hundred and twentieth of the previous best values. The conductivities of solutions of the above-named types are, in general, considerably in excess of those of the pure components, and increase uniformly with increasing diversity in character (that is, in the positive or negative nature of the constituent radicles) of their components. The results obtained have been correlated with those derived from freezing-point measurements on similar systems, and the validity of the fundamental relationships between compound formation and ionisation in solutions (see preceding abstract) has been confirmed. The maximum specific conductivity in no case corresponds with a composition of the system which is the same as that of an isolated compound. The following compounds have been isolated: $\text{CCl}_3 \cdot \text{CO}_2\text{H}$, $\text{CH}_3 \cdot \text{CO}_2\text{Et}$, m. p. -27° ; $\text{CCl}_3 \cdot \text{CO}_2\text{H}$, $\text{C}_6\text{H}_5 \cdot \text{CO}_2\text{Et}$, m. p. -23° ; $\text{CCl}_3 \cdot \text{CO}_2\text{H}$, $\text{C}_6\text{H}_5 \cdot \text{CO}_2 \cdot \text{C}_7\text{H}_7$, m. p. 11.9° ; $\text{CCl}_3 \cdot \text{CO}_2\text{H}$, COMePh , m. p. 26° ; $\text{H} \cdot \text{CO} \cdot \text{NH}_2$, $2\text{CH}_3 \cdot \text{CO}_2\text{H}$, m. p. -8° .

J. F. S.

Prediction of Solubility in Polar Solutions. JAMES KENDALL, ARTHUR W. DAVIDSON, and HOWARD ADLER (*J. Amer. Chem. Soc.*, 1921, **43**, 1481—1502; cf. preceding abstracts).—The various factors inducing deviations from the ideal solubility curve in polar solutions have been critically discussed, and it has been shown that the predominant disturbing factor is additive compound formation between the components of the solution. In previous papers (*loc. cit.*) it has been shown that the extent of such compound formation in any series is dependent on the diversity in character of the constituent radicles of the components, the basis of the diversity being the relative position of the variable radicles in the *E.M.F.* series. Ionisation, furthermore, has been found to increase uniformly with compound formation. These generalisations have been employed to predict solubility relationships in systems of various types. It has been shown that: (a) for a given solute in a series of different solvents, increasing solubility and increasing compound formation proceed in parallel, (b) for a series of different solutes of high melting point in a given solvent, increasing solubility and increasing compound formation also proceed in parallel at low temperatures. Illustrations of these rules for non-aqueous and aqueous solutions have been presented which indicate their agreement with the facts of experiment. A general discussion of solubility relationships in systems of increasingly complex nature (for example, $RX-HX$; $RX-H_2O$; $RX-R'X-H_2O$) has been made, and the rules deduced for the simpler types of systems have been found to be applicable, to a limited extent, to the more complex types. For example, (a) salts of a very weak base exhibit increasing hydrate formation and increasing solubility in water as the acid radicle X diverges from OH' ; salts of a very weak acid show the same behaviour as R diverges from H . (b) The increase in the solubility of a sparingly soluble salt in water on addition of a second salt containing a common ion, due to complex salt formation, is dependent on the diversity of the variable radicles.
J. F. S.

Solubilities of Acids in Aqueous Solutions of other Acids. JAMES KENDALL and JAMES C. ANDREWS (*J. Amer. Chem. Soc.*, 1921, **43**, 1545—1560; cf. preceding abstracts).—Solubility curves have been produced for hydrogen sulphide, boric acid, benzoic acid, and salicylic acid in aqueous solutions of hydrogen chloride of a series of concentrations and also for the same acids, except hydrogen sulphide in aqueous solutions of nitric acid. The changes of the specific conductivity of the solvent acids, due to the presence of the weaker acids in solution, has also been investigated. The results obtained, together with those of previous investigators, have been critically discussed in connexion with the rule established previously (*loc. cit.*), that compound formation between acids in pairs increases in extent with increasing diversity of acidic strength. The predicted and observed solubility curves for systems of various types have been compared, and satisfactory agreement, in general, has been obtained. The conductivity results also fall into line in

most cases. Certain abnormalities, however, do appear in particular systems, which are probably to be explained by the water entering into the equilibria in some as yet undefined way. J. F. S.

The Influence of Pressure on Spontaneous Crystallisation. MEINHARD HASSELBLATT (*Z. anorg. Chem.*, 1921, **119**, 353—364).—Experiments were made to determine the influence of pressure on the number of nuclei developed during spontaneous crystallisation at different temperatures (cf. Tammann, A., 1898, ii, 330). Experiments were made with betol, piperine, and papaverine, and it was found that increased pressure, up to 1000 kilos. per sq. cm., had the same effect on nuclei formation as on melting point. At 1000 kilos. the m. p. of betol is raised from 93° to 122·9°, and the temperature of spontaneous crystallisation was raised by a closely corresponding amount. In the case of piperine, the temperature of maximum formation of nuclei was raised from 35° to 70° by increasing the pressure from 1 to 1000 kilos. per sq. cm., and in the case of papaverine from 40° to 60°. The form of the curve showing the variation of number of nuclei with temperature was similar at both pressures. The maximum number of nuclei is doubled at the higher pressure in the case of piperine, but in the case of papaverine it is about the same at 1 kilo. and 1000 kilos. per sq. cm. E. H. R.

Crystallisation Velocity under High Pressure. MEINHARD HASSELBLATT (*Z. anorg. Chem.*, 1921, **119**, 325—352).—The conditions determining crystallisation velocity at normal pressure have been recorded by Tammann (A., 1898, ii, 425; 1899, ii, 272, 548). For examining the effect of pressure on crystallisation velocity, a modified form of an apparatus described by Tammann was used ("Kristallisieren und Schmelzen," p. 165). The substances studied were thymol, phenyl salicylate, benzophenone, apiole, calcium nitrate tri- and tetra-hydrates, cadmium nitrate tetrahydrate, mixed crystals of calcium and cadmium nitrates, guaiacol, form-anilide, benzoic anhydride, *m*-iodonitrobenzene, and benzy laniline. In general, the effect of pressure, up to 2000 kilos. per sq. cm., was remarkably small. In two cases there was an increase in crystallisation velocity of 43% and 18% respectively, but in most cases a slight fall, the greatest being 49% at 1000 kilos. per sq. cm. As the increased pressure has also the effect of raising the melting point considerably, for instance by 27·4° at 1000 kilos. in the case of phenyl salicylate, and the crystallisation temperature limits are raised proportionately, it can be concluded that the real effect of pressure alone is always to decrease the maximum crystallisation velocity. The influence of other substances on the crystallisation velocity of benzophenone was not appreciably affected by pressure. E. H. R.

Recrystallisation Produced by Annealing. PAUL GAUBERT (*Compt. rend.*, 1921, **173**, 1089—1092).—From a study of the behaviour of crystals of vanillin at a temperature slightly below their melting point and of substances like paraffin wax or cetin, which

give malleable crystals in the neighbourhood of their melting point, it is shown that recrystallisation is, as a rule, only possible if the crystals are sufficiently malleable for certain mechanical actions to modify their crystalline system. There is then produced, as it were, a slow polymorphic transformation, with, however, this difference, that in hammer-hardened substances there may remain some intact crystals which will act as nuclei to start the recrystallisation.

W. G.

Crystal Structure and Atomic Constitution. I. P. NIGGLI (*Z. Kryst. Min.*, 1921, **56**, 12—45; from *Chem. Zentr.*, 1921, iii, 833—834).—A discussion of the bearing of crystal structure on atomic constitution in the light of existing crystallographic data. Crystal symmetry with a known arrangement of atoms in the lattice implies also a certain minimal symmetry in the structure of the component atoms. A volume effect must follow from the actual effective size of atoms, as, for example, in the series formed by the carbonates of calcium, magnesium, ferrous iron, nickel, and cobalt, all of which have the calcite structure. From a consideration of published crystallographic data, the univalent metallic elements may be arranged in two series, namely, a principal series, lithium, sodium, potassium, rubidium, and caesium and a subsidiary series, copper (univalent), silver, and gold (univalent). Molecular volume does not increase continuously with electron number but shows periodicity. Similar results are given by bivalent and trivalent metallic elements. With higher valencies the periodicity in molecular volumes tends to disappear (cf. Bragg, A., 1920, ii, 537).

G. W. R.

Crystal Structure and Atomic Constitution. II. P. NIGGLI (*Z. Kryst. Min.*, 1921, **56**, 167—190; from *Chem. Zentr.*, 1921, iii, 834—835).—A continuation of an earlier discussion (see preceding abstract) of the relations between crystal structure and atomic constitution. The theories of Kossel, Vegard, Lewis, Langmuir, Lacombe, and Kohlweiler imply certain assumptions as to atomic structure. The periodicity in atomic volumes already noted is related to the arrangement of electrons in the atom. Whilst crystal form is connected with chemical constitution, isomorphism in its widest sense is a geometrical rather than a chemical effect. A comparison of the volume relationships in series of crystalline "isosteric" compounds permits a comparison of the effective sizes of atoms of differing valency, whereby it appears that atomic size diminishes with increase of valency.

G. W. R.

Investigation of Crystal Space Lattices by means of Röntgen Rays: a Simple Röntgen Tube. WALTER GERLACH (*Physikal. Z.*, 1921, **22**, 557—559).—The lattice constant, a , has been determined for magnesium oxide, calcium oxide, strontium oxide, and barium oxide. In all cases the substances are shown to be regular and belong to the sodium chloride type. The values of a recorded are MgO, 4.204; CaO, 4.762; SrO, 5.100; BaO, 5.47. Making use of the Debye-Scherrer method, it is shown that

the ions of magnesia are both bivalent. The lattice of fluorspar has been redetermined and the structure put forward by Bragg confirmed. The lattice constant is found to be $a = 5.452 \text{ \AA.U.} \pm 0.2\%$. Various forms of silicon have been examined and shown to be crystallographically similar. The lattice is of the diamond type and has the constant $a = 5.400 \times 10^{-8} \text{ cm.} \pm 0.2\%$. The transport of anti-cathode material towards the cathode has been considered. It is shown that after using a Siegbahn-Hadding tube for a prolonged period a deposit of the anti-cathode material is found on the cathode in the form of small crystals. In the case of aluminium cathodes, the crystals were well-formed cubes of 1 mm. edge. In the case of copper cathodes the transport of material takes place in the ionic form, and is not in any way to be regarded as a distillation. The amount of material transported is in accordance with the dimensions of the ionic stream.

A Röntgen tube is described in which a portion of the outside wall serves as the metallic hemisphere constituting the cathode, whilst the anti-cathode is situated at the geometrical centre of the hemisphere. The Röntgen rays pass through a number of holes in the cathode which is rendered air-tight by suitable cements.

J. F. S.

The Theory of Resistance Limits in Mixed Crystals. G. MASING (*Z. anorg. Chem.*, 1921, 118, 293—308).—It has been held by Tammann (*A.*, 1919, ii, 398) that the existence in binary mixed crystals of resistance limits to the attack of chemical agents which react with one constituent but not with the other provides a proof that the two kinds of atom in the mixed crystals are regularly distributed on a space lattice. The author shows that, making certain assumptions, it can be proved by the mathematical laws of probability that resistance limits will occur if the atoms are irregularly distributed on the space lattice. The limits calculated are not the same as those found experimentally, but they depend on certain simple arbitrary assumptions made with respect to the manner in which the solvent attacks the mixed crystals, and these may be insufficient. The work is only intended to show that the mere existence of resistance limits is no proof of the regular distribution of the atoms in the mixed crystals.

E. H. R.

Coagulation of Colloids by Electrolytes. HUMPHREY D. MURRAY (*Chem. News*, 1921, 123, 277—279).—A general review of the work within recent years on the coagulation of colloids by electrolytes.

J. F. S.

Kinetic and Static Coagulation Measurements of Suspensoids. FRIEDRICH VINCENZ VON HAHN (*Z. Elektrochem.*, 1921, 27, 501—505).—A general discussion of the various kinetic and static methods of measuring the coagulation of suspensoids with the object of ascertaining the suitability or otherwise of these methods for determining the relative stability of the sols. A new method of determining the stability by coagulation is described.

The apparatus consists of a long U-tube consisting of a narrow tube and a wide tube; the wide tube terminates at its lower end in a tap and extends slightly beyond the cross-piece joining the two tubes which is closed by a tap. In making a determination, the tap is closed and the wide limb filled with water and the narrow tube with the sol, both to the same height. The tap is now opened, and the levels adjust themselves; if now a portion of the disperse phase owing to coagulation reaches the continuation of the longer tube, the density of the sol is changed and the meniscus commences to move. The time which must elapse before this movement of the meniscus begins is characteristic of the stability of the sol. Two other, rather more sensitive, methods based on the same principles are described. J. F. S.

The Equilibrium in the Reduction of Nickel Chloride by Hydrogen. E. BERGER and G. CRUT (*Compt. rend.*, 1921, 173, 977—979).—Within wide limits the equilibrium of the reversible reaction $\text{NiCl}_2 + \text{H}_2 \rightleftharpoons \text{Ni} + 2\text{HCl}$, is determined accurately by the formula $29642/4.571T - 1.75 \log T - 0.006T^2/4.571 + \log p + \log [\alpha^2/(100 - \alpha)] = K$, taking the heat of reaction from left to right as -30700 cal. at 15° and the value of K as 5.93. The equilibrium concentrations are the same in whichever direction the reaction proceeds. W. G.

An Explanation of Liesegang's Rings. HUGH MCGUIGAN (*Science*, 1921, 54, 78—80).—When a minute drop of silver nitrate solution is placed on a gelatin gel containing potassium dichromate, there is an immediate formation of a circle of silver chromate about which is a sharply demarcated zone of influence from which the chromate has been removed. The Liesegang ring formation when a larger drop of silver nitrate is used is explained as follows: Silver chromate is formed and a clear zone results in the gelatin by the attraction of the chromate to the silver. Beyond this zone of influence, the chromate is fixed and remains so unless an attraction force is exerted. The silver nitrate now wanders on through the ring into the clear zone until it is sufficiently close to the chromate gelatin to exert an attraction which again draws the chromate and forms another ring and clear zone. At the same time the chromate exerts a "pull" on the silver and the ring is formed where the forces are balanced. Again it may be presumed that to start the chromate moving will require a greater force than to keep it moving after the movement has commenced; consequently the second ring is separated from the first. With each succeeding ring the concentration of the silver is less and this also will operate to remove the succeeding rings farther apart. As a requisite for ring formation the precipitate formed must be permeable to the liquid solution used, in this case silver nitrate.

CHEMICAL ABSTRACTS.

Preparation of Nitroanisole from Nitrochlorobenzene. A. V. BLOM (*Helv. Chim. Acta*, 1921, 4, 1029—1035).—The action of *o*- or *p*-nitrochlorobenzene on sodium methoxide in the presence

of methyl alcohol has been studied and by a combination of different titrations it has been possible to determine the rate of formation of each component of the change (cf. A., 1921, ii, 497). Below 70°, the reducing action of intermediately formed alkoxide is so slight as to be practically negligible. The yields of nitrophenol ethers and the rate of reaction depend, not only on temperature and time, but also on the quantity and composition of the solvent. Accelerating factors increase the yield of ethers if they do not also accelerate the subsidiary reactions. High alcoholic concentration and a minimum amount of solvent are favourable in this respect. The use of temperatures higher than 70° or of a large excess of alkali is disadvantageous on account of the increased production of dichloroazoxybenzene or nitrophenol. The possibility of the hydrolysis of the ether formed need not be taken into account, and the whole of the alkali may therefore be added at the commencement of the operation. H. W.

The Theory of Reaction Velocities. N. R. DHAR (*Z. anorg. Chem.*, 1921, **119**, 177—178).—In criticising Lewis's theory, Lindemann (A., 1920, ii, 743) states that the inversion of sucrose by acids, which should proceed 10^{13} times as fast in sunlight as in the dark, is not accelerated at all by light. The author has found that a solution of sucrose in tropical sunlight is completely inverted even in absence of acids, whilst in presence of hydrochloric acid the inversion is distinctly accelerated by sunlight. These observations therefore confirm the theory of Lewis in a qualitative manner. Reactions which have a large temperature coefficient are those most sensitive to light. E. H. R.

Rates of Hydrolysis of the Rare Earth Carbonates and the Serial Order of the Rare Earth Elements. PAUL H. M.-P. BRINTON and C. JAMES (*J. Amer. Chem. Soc.*, 1921, **43**, 1446—1451).—The relative rates of hydrolysis of the carbonates of ytterbium, thulium, yttrium, dysprosium, terbium, gadolinium, europium, samarium, neodymium, praseodymium, lanthanum, and cerium have been determined. An equivalent quantity of the carbonate of each of the earths was dissolved in a definite quantity of water and treated with the theoretical quantity of sodium carbonate and boiled. The carbon dioxide evolved was collected during half-hour periods and measured, and from the results the percentage hydrolysis calculated. In the case of lanthanum and cerous carbonate anomalous behaviour is displayed inasmuch as the whole of the carbon dioxide is evolved in the first half-hour and consequently on plotting percentage hydrolysis against the time of boiling they give horizontal line curves, whilst the curves for the other elements are slowly rising curves, slightly convex to the time axis. The curves place the elements, with the exception of the two named, in the order of their basicity. The position thus assigned to yttrium agrees with the solubilities of its compounds, whilst previous determinations of basicity have not done so. The increasing order of basicity of the ten elements concerned

is given as: ytterbium, thulium, yttrium, dysprosium, terbium, gadolinium, europium, samarium, neodymium, and praseodymium. A short discussion on the serial order of the rare earth elements is given. J. F. S.

The Explosion of Acetylene and Nitrogen. I. WILLIAM EDWARD GARNER and KICHIMATSU MATSUNO (T., 1921, 119, 1903—1914).

Velocity of Decomposition of Mono- and Di-substituted Malonic Acids. AUGUST L. BERNOULLI and HEINRICH JAKUBOWICZ (*Helv. Chim. Acta*, 1921, 4, 1018—1029).—In extension of the work of Bernoulli and Wege (A., 1919, ii, 503), the rate of decomposition of malonic, diallylmalonic, diethylmalonic, phenylmalonic, benzylmalonic, ethylmalonic, tartronic, mono- and dichloromalonic, and dibromomalonic acid has been measured at a variety of temperatures.

The relative effect of the different substituents as established by Bernoulli and Wege (*loc. cit.*) at 100° is found to be maintained at all temperatures investigated. As mono-substituents, bromine, chlorine, and hydroxyl accelerate the decomposition of malonic acid in order of decreasing effect. The temperature curves of the mono-substituted acids diverge from one another with increasing temperature, without, however, intersecting. The temperature curves of the di-substituted malonic acids (from 80° upwards) remain parallel to one another and to that of malonic acid so that the latter behaves as "symmetrically di-substituted." Two similar substituents depress the rate of decomposition greatly, even when the latter is accelerated by the same substituent acting singly. The retarding action becomes more marked with increase in mass of the substituent; this is particularly noticeable with dibromomalonic acid. The effect of substitution in the case of di-substitution is the result of two component actions, (a) a specific effect which is obvious in the case of mono-substitution, and (b) a symmetry effect which depends on the mass and not on the electrochemical character of the radicle or atom. H. W.

Influence of Neutral Salts on the Hydrolysis of Ethyl Formate. ALEXANDER BERNARD MANNING (T., 1921, 119, 2079—2087).

The Maximum Stability of Esters of Carboxylic Acids. KARL GUSTAV KARLSSON (*Z. anorg. Chem.*, 1921, 119, 69—96).—Experiments were made to determine the influence of hydrogen-ion concentration on the stability of methyl acetate and ethyl acetate. The experiments were carried out at 85.55° and 75.40°. The ester was heated in aqueous solution in presence of a "buffer" salt (sodium acetate) in a sealed quartz tube for a known time. The hydrogen-ion concentration was determined by direct measurement before and after the experiment and the amount of hydrolysis was determined by titration with barium hydroxide solution. The velocity constant was calculated from the usual unimolecular equation. In the case of methyl acetate, the lowest velocity con-

stant, that is, the greatest stability, was found at 85.55° with $P_H=4.70$, and in the case of ethyl acetate, $P_H=5.1$. The presence of neutral salts, sodium chloride, or potassium nitrate had no effect on the position of the point of maximum stability. The curves obtained by plotting the velocity constants as ordinates against the P_H values descend to a minimum at the P_H representing maximum stability, but they are not symmetrical about this point, rising more steeply on the alkaline side. It follows that H- and OH-ions cannot have the same hydrolytic activity (cf. Wijs, A., 1893, ii, 364; 1894, ii, 82). What function the reaction velocity is of the H- and OH-ions cannot yet be determined. Neutral salts increase the reaction velocity in alkaline solution, that is, when P_H is high, by increasing the acidity, but on the acid side of the point of maximum stability they have much less effect.

E. H. R.

Preparation and Hydrolysis of Benzyl Esters. E. H. VOLWILER and E. B. VLIET (*J. Amer. Chem. Soc.*, 1921, 43, 1672—1676).—The comparative rates of hydrolysis of certain benzyl esters were determined in order to obtain a basis for the correlation of chemical properties and physiological action. The rates of hydrolysis of these benzyl esters increase in the following order: salicylate, benzoate, stearate, cinnamate, acetate, succinate, and fumarate. The rate of hydrolysis of the benzyl group in benzyl aceto-oxybenzoate is of the same order as in benzyl salicylate.

W. G.

The Influencing of Catalysts and Specifically Active Catalysts. KARL W. ROSENMOND and FRITZ ZETZSCHE (*Ber.*, 1921, 54, [B], 2885—2888; cf. A., 1921, ii, 320, 392, 393).—A reply to the criticism of Abel (A., 1921, ii, 542).

H. W.

Catalytic Action at Solid Surfaces. VII. Influence of Pressure on the Rate of Hydrogenation of Liquids in the Presence of Nickel. E. F. ARMSTRONG and T. P. HILDITCH (*Proc. Roy. Soc.*, 1921, [A], 100, 240—252; cf. A., 1921, ii, 582; 1920, ii, 608).—The influence of pressure on the rate of absorption of hydrogen by pinene, ethyl cinnamate, ethyl linoleate, linolein, linolenin, geraniol, terpineol, oleic acid, citral, and carvone in the presence of nickel has been investigated. The results fall into three definite groups: (i) *Normal*. In the absence of substituent groups of the type mentioned below and in the presence of sufficient nickel (generally as long as at least 0.1% of nickel is present) the ethylenic union is hydrogenated at a rate which is in almost exact proportion to the absolute pressure of the hydrogen. To this group belong ethyl cinnamate, ethyl linoleate, and pinene. (ii) *Subnormal*. At very low concentrations of catalyst (for example, 0.01—0.02% of nickel calculated on the organic compound) the increase in the rate of hydrogenation becomes less than proportional to the increase in pressure. The compounds which absorb hydrogen most readily are in general more prone to show the sub-normal effect when the concentration of the catalyst is diminished, and it

is especially marked in the case of multi-ethylenic compounds such as derivatives of linoleic or linolenic acid or citral, although at low concentrations of catalyst it is also shown by ethyl cinnamate. (iii) *Abnormal*. If the unsaturated compound contains another group which has affinity towards nickel (but is not open to hydrogenation), it is found that increase in hydrogen pressure causes an increase in the rate of hydrogen absorption in greater than simple proportion to the altered concentration of the hydrogen. This is observed with unsaturated alcohols and carboxylic acids; the carboxyl group tends to produce the nickel salt and hydrogen, although at 180° the equilibrium is moving in the direction of reduction, whilst with the alcohol there is a tendency toward dehydration. On the other hand, unsaturated aldehydes and ketones show a normal reduction with increase in the hydrogen concentration. In general, an increase of the hydrogen concentration causes a directly proportionate increase in the rate of hydrogenation, provided that there are no disturbing factors; the increase of hydrogenation becomes abnormally large if other groups active toward nickel, but not open to hydrogenation, are also present. The nature of the unsaturated organic compound has a determining influence on the effect of the hydrogen concentration.

J. F. S.

Catalysis of the Mutarotation of Dextrose by Metals.

WILLIAM EDWARD GARNER and DOUGLAS NORMAN JACKMAN (T., 1921, 119, 1936—1948).

Oxidation Catalysis. I. L. KARCZAG (*Biochem. Z.*, 1921, 117, 69—86).—A group of colouring matters was subjected to the action of hydrogen peroxide under various experimental conditions in the presence of iron, copper, cobalt, manganese, nickel, and platinum salts. The conclusion is drawn that the first four metals named have a two-fold action, like that of catalase and oxydase respectively, whilst the two latter have only an oxydase-like action.

H. K.

Magnitude of Atoms. THEODORE W. RICHARDS (*J. Amer. Chem. Soc.*, 1921, 43, 1584—1591).—A theoretical paper in which it is pointed out that the contraction suffered during combination, taken in connexion with the compressibilities of the elements combined, makes possible an approximate estimate of the bulk occupied by the several constituents in a compound. Calculations are made of the diameters of the atoms of the halogens and the alkali metals in the alkali haloids when the following values in Ångström units are obtained: (a) in chlorides, chlorine, 2·8; lithium, 2·3; sodium, 2·9; potassium, 3·5; rubidium, 3·8; caesium, 3·8; (b) in bromides, bromine, 3·1; lithium, 2·4; sodium, 2·9; potassium, 3·5; rubidium, 3·8; caesium, 3·7; (c) in iodides, iodine, 3·4; lithium, 2·6; sodium, 3·1; potassium, 3·7; rubidium, 3·9; caesium, 3·8. These values lie about half-way between the values obtained by Bragg and Landé. Emphasis is placed on the importance of considering the compressing effect of the affinities concerned in

making any comparison of atomic volume. A series of volumes of the individual halogen and alkali metal atoms in alkali haloids is given in the paper.

J. F. S.

Analysis of the Atomic Structure. A. DAUVILLIER (*Compt. rend.*, 1921, 173, 1077—1079; cf. A., 1921, ii, 669).—A table is given showing the number of internal electrons and the number of quanta for each layer in the heavy atoms, and these data are discussed. A periodic table is given illustrating the author's theory of electronic structure of the elements, and from it, it is shown that the chemical analogy of the different elements depends, not only on the number of superficial corpuscles, but also on the subjacent electronic layer. The table enables the transformations that will be observed in the X-spectrum of the light elements to be foretold.

W. G.

Langmuir's Theory of the Arrangement of Electrons in Atoms and Molecules. CHARLES R. BURY (*J. Amer. Chem. Soc.*, 1921, 43, 1602—1609).—A theoretical paper in which an alternative has been suggested to the fourth postulate in Langmuir's arrangement of electrons in atoms and molecules (A., 1919, ii, 328). This avoids the necessity of postulating cells in the structure of the atom, and leads to different structures for the elements of higher atomic weights than that of neon. It is claimed that the application of Lewis and Langmuir's octet hypothesis of valency to these structures gives a better explanation of the chemical properties of the higher elements than can be obtained from Langmuir's structures.

J. F. S.

Motion of Electrons in Gases. J. S. TOWNSEND and V. A. BAILEY (*Phil. Mag.*, 1921, [vi], 42, 873—891).—An account is given of the method of finding the velocities of electrons in gases, and the application of the results to determine some properties of gaseous molecules. The paper also contains the results of a large number of experiments which have been made on the motion of electrons in hydrogen, nitrogen, and oxygen with ranges of forces and pressures similar to those used in experiments with air.

J. F. S.

New Type of Sodium Lamp for Polarimetry. HAROLD A. FALES and JACQUES C. MORRELL (*J. Amer. Chem. Soc.*, 1921, 43, 1629—1630).—A sodium lamp for use in all cases where a continuous sodium flame is required is described. The lamp consists of a small alundum thimble supported by nickel or nichrome wire in the flame of a Méker burner. The molten sodium chloride contained in the thimble slowly percolates through the pores of the thimble, so that the flame is continuously fed by sodium chloride vapour.

J. F. S.

Rotary Burner with Fixed Gas Connexion. GEORG LOCKEMANN (*Z. angew. Chem.*, 1921, 34, 594—595).—A modification of the rotary burner previously described (A., 1921, ii, 447) enables an ordinary Teclu burner with a non-flexible gas connexion to be

used. The burner tube is fitted into the hub of the horizontal pulley, and is continued on the upper side by a tube bent four times at right angles and jointed in the central vertical limb, so that the upper portion can be twisted round to bring the gas outlet to any desired distance from the axis of rotation of the pulley. A fitting is also provided to fix on the top of the burner tube to give an oblique instead of a vertical flame. G. F. M.

Inorganic Chemistry.

Molecular Volumes, Physical Properties, and Molecular Models of the Halogens. F. A. HENGLEIN (*Z. anorg. Chem.*, 1921, **118**, 165—171).—It has been shown by Biltz (A., 1921, ii, 437, 487) that there is a linear relation between the molecular volumes of the halogens and of their compounds. It is now shown that there is likewise a linear relation between the atomic volumes of the halogens at their boiling points and others of their physical properties including melting point, boiling point, critical point, latent heat of fusion or of evaporation, and normal potential; also between the molecular volumes of the halogen acids at the boiling point and many other of their physical constants. Owing to the close similarity in structure of the different halogen atoms, their properties are determined principally by the atomic radius. In contrast with most groups of the periodic system, the molecule of a halogen element has an especially simple structure; it can be represented diagrammatically by two cubes having a common edge, with the valence electrons situated at the corners.

E. H. R.

The Catalytic Formation of Hydrogen Chloride from Hydrogen and Chlorine without Explosion. BERNHARD NEUMANN [with BERGDAHL, BROY, and KARWAT] (*Z. angew. Chem.*, 1921, **34**, 613—620).—All non-explosion methods for the synthesis of hydrogen chloride gas yield a product contaminated with chlorine. Hoppe, who first suggested the use of metallic chlorides as catalysts to this reaction employed the chlorides of aluminium, tin, and zinc in concentrated solution at a temperature of 130°. A repetition of his work has shown that, under the most favourable conditions and with a very slow current of gas not more than 70% conversion is obtained. The reaction is favoured by an increase of concentration and of temperature, but a practical limit is set to the latter by the point at which evaporation becomes rapid. The author has used solid chlorides and higher temperatures. The hydrogen and chlorine were generated electrolytically in separate cells in the same electric circuit, passing into a mixing flask containing water and thence to a quartz tube in the shape of a pipette filled

with granules of quartz impregnated with the selected chloride and heated. The emerging gases were absorbed in a 10-bulb potash tube. The mixed gases were shielded from light and drawn through the apparatus by uniform slight suction, fluctuation of pressure being a source of explosions. With a low gas velocity, a complete conversion was effected by magnesium chloride at 300°, by calcium chloride at 305°, by aluminium chloride (which at this temperature is almost entirely decomposed) at 350°, and by quartz unimpregnated by any salt at 380°. Conversion was improved by warming the mixing flask to 50°, with the effect of adding 1 mol. of water to each mol. of hydrogen chloride, this temperature being an optimum. The dilution of the gases with oxygen had at 380° no influence on the reaction. The reactions $\text{Cl}_2 + \text{H}_2\text{O} = \text{ClOH} + \text{HCl}$, $\text{ClOH} + \text{H}_2 = \text{H}_2\text{O} + \text{HCl}$ are considered to occur, removal of the hydrogen chloride generated being facilitated by the partly dissociated chlorides. C. I.

Sulphate-free Sulphites for Standard Sulphur Dioxide Solutions. S. LANTZ SHENEFIELD, FRANK C. VILBRANDT, and JAMES R. WITHROW (*Chem. and Met. Eng.*, 1921, 25, 953—955).—Pure sodium sulphite heptahydrate was prepared by passing carefully purified sulphur dioxide into a solution of sodium carbonate to saturation, adding the requisite amount of sodium carbonate to transform the resultant sodium hydrogen sulphite into the normal sulphite, and cooling the solution to 0°, whereby a crystalline meal of the heptahydrate was obtained. All the operations were carried out in the absence of oxygen and the damp crystals were found to be free from sulphate. They were placed in a desiccator over sulphuric acid and sodium pyrogallate was used as oxygen absorbent. Although every precaution was taken to exclude air leakage, the crystals after one week contained 7.52% of sodium sulphate decahydrate. Similar oxidation occurred in the preparation of pure dry calcium sulphite, and the opinion is expressed that in both cases this is due to autoxidation of the salt. It is concluded that the validity of all investigations in the literature based on the preparation of sulphate-free sulphite for use as a sulphur dioxide standard is doubtful. A. R. P.

Polythionic Acids and Polythionates. E. H. RIESENFELD and G. W. FELD (*Z. anorg. Chem.*, 1921, 119, 225—270).—A method has been devised for estimating the proportions of tri-, tetra-, and penta-thionates together in solution with sulphite, thiosulphate, and sulphate, and the method has been applied to the study of the formation and stability of the polythionates. The hexathionic acid described by Debus (T., 1888, 53, 278) does not exist. The total polythionic acids can be determined by treatment in neutral solution with a mercuric salt, when four equivalents of sulphuric acid are produced for each molecule of polythionate, for instance: $2\text{S}_5\text{O}_8^{--} + 2\text{Hg}^{++} + 4\text{H}_2\text{O} = 2\text{HgS} + 4\text{SO}_4^{--} + 8\text{H}^+ + 4\text{S}$. The acid can then be titrated. Alternatively, the polythionates can be oxidised with bromine in alkaline solution; the whole of the sulphur is oxidised to sulphate and is precipitated and weighed

as barium sulphate. The trithionate is estimated by boiling the solution with an excess of copper sulphate, when the following reaction occurs: $\text{S}_3\text{O}_6'' + \text{Cu}'' + 2\text{H}_2\text{O} = \text{CuS} + 2\text{SO}_4'' + 4\text{H}^+$. The copper sulphide is filtered, ignited, and weighed as copper oxide. When boiled with excess of alkali, the polythionates form thiosulphate and sulphite thus: $2\text{S}_3\text{O}_6'' + 6\text{OH}' = \text{S}_2\text{O}_3'' + 4\text{SO}_3'' + 3\text{H}_2\text{O}$; $2\text{S}_4\text{O}_6'' + 6\text{OH}' = 3\text{S}_2\text{O}_3'' + 2\text{SO}_3'' + 3\text{H}_2\text{O}$; $2\text{S}_5\text{O}_6'' + 6\text{OH}' = 5\text{S}_2\text{O}_3'' + 3\text{H}_2\text{O}$. The sulphite and thiosulphate formed are estimated by titration with iodine. By applying three of these reactions, data can be obtained from which to calculate the proportions of tri-, tetra-, and penta-thionate present.

The interaction between hydrogen sulphide and sulphur dioxide was studied by leading a current of the former, as gas, into an aqueous solution of the latter at 0° , until a definite ratio of the reacting substances was present in solution. The total polythionate was estimated after sixty hours. The optimum ratio for polythionate formation was $2\text{SO}_2 : 1\text{H}_2\text{S}$. On the other hand, with the ratio $2\text{H}_2\text{S} : \text{SO}_2$, all the sulphur was precipitated in the elementary form. Evidence was obtained that, immediately after the preparation of a solution in the ratio $2\text{SO}_2 : \text{H}_2\text{S}$, an intermediate compound is formed which can be precipitated at a low temperature as the barium salt; in solution, this changes to barium thiosulphate. The proportion of tri- and tetra-thionic acids formed (determined after fourteen days) depends on the sulphur dioxide concentration, the formation of tetrathionic acid being favoured by low concentration. The proportion of pentathionic acid is, however, practically constant. These results are contrary to those obtained by Heinze (A, 1919, ii, 334). The sulphuric acid which is always formed reaches a maximum near the commencement of the reaction; it must therefore be formed from the intermediate compound, not by oxidation of the polythionic acids.

Of the three polythionic acids, the tetrathionic acid is the least stable and decomposes relatively quickly into tri- and pentathionic acids. The trithionic acid decomposes more slowly with formation of sulphur dioxide, whilst pentathionic acid decomposes only in the course of months with separation of sulphur. The order of stability is the same in neutral as in acid solution; in alkaline solution all the polythionates decompose quickly into thiosulphate and sulphite.

The phenomena observed are explained on the assumption that the above intermediate compound is a hydrate of the unknown sulphur monoxide, SO . This is stable in acid solution for a time, but in neutral or alkaline solution quickly forms thiosulphate. In acid solution, it slowly polymerises to pentathionic acid. By combination with sulphurous acid it forms tri- and tetra-thionic acid: $3\text{SO} + \text{H}_2\text{SO}_3 = \text{H}_2\text{S}_4\text{O}_6$; $\text{SO} + 2\text{SO}_2 + \text{H}_2\text{O} = \text{H}_2\text{S}_3\text{O}_6$. By hydrogen sulphide, it is reduced to sulphur.
E. H. R.

The Preparation of Hydrogen Selenide from Metallic Selenides. L. MOSER and E. DOCTOR (*Z. anorg. Chem.*, 1921, 118, 284—292).—The selenides of magnesium, aluminium, iron,

and zinc were prepared in a similar manner to the corresponding tellurides (this vol., ii, 48) by passing selenium vapour over the heated metal in a vacuum. The aluminium and magnesium compounds were also prepared by direct combination of the metal with selenium in a crucible, starting the reaction between the mixed components with a burning magnesium wire. All the products were considerably contaminated with metal except aluminium selenide, which, prepared by the latter process, was practically pure. Aluminium selenide, Al_2Se_3 , forms a light brown powder, unstable in air, and magnesium selenide, MgSe , is very similar. Zinc selenide, ZnSe , is citron-yellow and iron selenide, FeSe , is black and metallic; both are stable in air.

Hydrogen selenide was prepared in an apparatus similar to that used for hydrogen telluride, by dropping the metallic selenide slowly into acid. The best results were obtained using aluminium or magnesium selenide. The gas was liquefied at the temperature of a solid carbon dioxide-ether mixture and by revaporisation was obtained pure. It is not decomposed by daylight in the liquid or gaseous form, but is sensitive to ultra-violet light. Dry oxygen has no action on the dry gas, but in presence of moisture decomposition is rapid.


E. H. R.

Selenious Acid and Heteropolyselenites. ARTHUR ROSENHEIM and LEONHARD KRAUSE (*Z. anorg. Chem.*, 1921, **118**, 177—192).—A large number of heteropolyselenites with vanadates and molybdates have been described by Prandtl and others (A., 1907, ii, 477; 1912, ii, 167; 1916, ii, 333), who described numerous well-crystallised salts which, however, had variable compositions according to the conditions of preparation. These compounds have been re-examined with a view to determine whether they may not have a semi-colloidal constitution similar to that of the periodates (A., 1919, ii, 508). Experiments were first made to determine the state of aggregation of selenious acid in aqueous solution. Depression of the freezing point of water indicated slight association, which has a tendency to decrease with time. The dissociation was determined by the hydrogen-ion concentration method and from the electrical conductivity, the results obtained being $4.85 \cdot 10^{-3}$ and $3.45 \cdot 10^{-3}$ respectively. These results are taken to be consistent with the presence of associated $(\text{H}_2\text{SeO}_3)_2$ molecules in solution.


The method of estimating selenious acid by heating with potassium iodide and hydrochloric acid and distilling the iodine over into potassium iodide was improved by the addition of phosphoric acid to the hydrochloric acid. This prevents the formation of selenium iodide, which may be the cause of low results. It was also found that selenious acid may be accurately titrated with sodium hydroxide; the best indicators are, for the formation of NaHSO_3 , *p*-nitrophenol, and for complete neutralisation to Na_2SeO_3 , thymolphthalein.

Lithium selenite forms the hydrate $4\text{Li}_2\text{SeO}_4 \cdot 3\text{H}_2\text{O}$; its solubility has a negative temperature coefficient.

Molybdoselenites.—The composition of molybdoselenites was found to depend on the ratio of molybdate to selenious acid in the solution from which they were precipitated. When less than 1 mol. of selenium dioxide was present to 1 mol. of molybdate, the potassium and barium salts corresponded with $2R_2O, 2SeO_2, 5MoO_3, xH_2O$ and the ammonium salt with $3(NH_4)_2O, 2SeO_2, 8MoO_3, 6H_2O$. With more than 1 mol. of selenium dioxide per mol. of molybdate in solution, salts were obtained in which the proportion of base was variable, but the ratio $SeO_2 : MoO_3$ was always very nearly 1 : 1. The ammonium salt, $2(NH_4)_2O, 5SeO_2, 5MoO_3, 8H_2O$, white, microscopic prisms, was obtained by adding 15 mols. of selenium dioxide to a saturated solution of ammonium paramolybdate. The potassium salt,



forms microscopic prisms, and the barium salt,



a white, crystalline precipitate.

Vanadoselenites.—Vanadoselenious acid has the composition $4SeO_2, 3V_2O_5, 10H_2O$. An extensive series of experiments showed that, as the concentration of selenious acid in the mother-liquor increased from zero to 5*N*, the ratio of SeO_2 to V_2O_5 in the solid phase increased from 4 : 3 to 5·5 : 3. Similar behaviour was shown by the ammonium vanadoselenites having the approximate formula $3(NH_4)_2O, 12SeO_2, 8V_2O_5$. It is concluded that the variable composition of the vanadic acid compounds is due to the formation of adsorption compounds, although in the case of the molybdic acid compounds this is not so clearly demonstrated. E. H. R.

The Preparation of Hydrogen Telluride from Metal Tellurides. L. MOSER and K. ERTL (*Z. anorg. Chem.*, 1921, **118**, 269—283).—A new method for preparing metal tellurides was devised which consisted in distilling tellurium at a low pressure (8 mm.) over the hot, finely divided metal. In this way the tellurides of magnesium, $MgTe$, aluminium, Al_2Te_3 , iron, $FeTe$, and zinc, $ZnTe$, were prepared. The aluminium compound was obtained in an almost pure state; it is a blackish-brown, lustrous, amorphous substance, decomposing in the air with formation of tellurium hydride. The other tellurides were all more or less contaminated with excess of the respective metal. Magnesium telluride forms a brown, sintered mass, iron telluride is grey and metallic, and zinc telluride is pale brown. The last two are stable in air. For the preparation of hydrogen telluride a special apparatus was designed, in which the powdered metallic telluride was dropped very gradually into acid in an atmosphere of nitrogen. Aluminium telluride proved the most suitable substance from which to generate the gas, and hydrochloric acid the best acid to use. The yield of gas obtained under the best conditions was more than 80% of the theoretical. The gas was liquefied by passing through a tube immersed in a mixture of solid carbon dioxide and ether. In the liquid state hydrogen telluride is very sensitive both to daylight and ultra-violet light, but the dry gas is quite stable in light. The dry gas is, however, immediately oxidised by oxygen. E. H. R.

The Preparation of Telluric Acid. JULIUS MEYER and HANNS MOLDENHAUER (*Z. anorg. Chem.*, 1921, **119**, 132—134).—Telluric acid can be prepared in a pure state and in almost theoretical yield by oxidation of tellurium tetrachloride with chloric acid. Tellurium (10 grams) is boiled with 10 c.c. of nitric acid and 3 c.c. of hydrochloric acid until completely dissolved. To the hot solution is added gradually a concentrated solution of 9 grams of chloric acid and the solution is boiled until no more chlorine is evolved. A slight excess of chloric acid is added to avoid formation of any explosible chlorine oxide. The solution is filtered through asbestos and concentrated by distillation in a vacuum on the water-bath, thereby removing chlorine. The telluric acid can be crystallised out by the addition of concentrated nitric acid, collected, and finally freed from chlorine and nitrogen oxides by drying in a vacuum. It is obtained as a crystalline, snow-white powder, readily soluble in water, having the composition H_6TeO_6 . E. H. R.

Vapour Pressures of Ammonia-Salt Solutions. R. O. E. DAVIS, L. B. OLMSTEAD, and F. O. LUNDSTRUM (*J. Amer. Chem. Soc.*, 1921, **43**, 1580—1583; cf. this vol., ii, 56).—Vapour pressure curves have been constructed for the following solutions: $\text{Ca}(\text{NO}_3)_2$ 22.48%, NH_3 19.18%, H_2O 58.34%; NaI 32.34%, NH_3 16.06%, H_2O 51.60%; NH_4CNS 77.84%, NH_3 22.16%; NH_4NO_3 33.7%, NH_3 18.82%, H_2O 47.48%; CaCl_2 12.9%, NH_3 22.9%, H_2O 64.2%; $\text{Ca}(\text{NO}_3)_2$ 55.8%, NH_3 25.77%, H_2O 18.43%; NH_3 28.15%, H_2O 71.85%; NaI 64.88%, NH_3 26.02%, H_2O 8.20% over the temperature range -16° to 40° . It is shown that solutions of ammonium nitrate in ammonia and ammonium thiocyanate are very corrosive to iron and steel, the calcium chloride-ammonia solution is less corrosive, and calcium nitrate-ammonia, and sodium iodide-ammonia solutions show no immediate corrosive action. Calcium nitrate-ammonia solutions seem to be the most promising of these solutions for practical use as an absorbent for ammonia in the synthetic ammonia process. J. F. S.

The Action of Metals, such as Copper and Zinc, on an Aqueous Solution of Ammonium Nitrite. N. R. DHAR (*Z. anorg. Chem.*, 1921, **119**, 174—176).—The action of a solution containing ammonium nitrite on copper was attributed by Hofmann and Buhk (*A.*, 1921, ii, 43) to the hydrolysis of the nitrite with formation of free nitrous acid. The observation that the metal is attacked even in presence of urea, however, renders this explanation improbable. Further, solutions of other nitrites such as zinc nitrite, which are hydrolysed as much as ammonium nitrite, do not attack copper. The activity of the ammonium nitrite is probably related to its instability and ready decomposition into nitrous oxide and water. E. H. R.

The Structure of Pyrophosphoric Acid. III. D. BALAREFF (*Z. anorg. Chem.*, 1921, **118**, 123—130; cf. *A.*, 1915, ii, 446; 1917, ii, 467).—In previous papers it has been shown that there is a good

deal of evidence in favour of the unsymmetrical structure of pyrophosphoric acid. The synthesis of the pyro-acid by combination of the ortho- and meta-acids in sulphuric acid would give added support to the hypothesis of an unsymmetrical structure, but attempts in this direction were not successful, even in acid containing 15% of free sulphuric anhydride.

The change of colour of the salt $\text{NaAg}_3\text{P}_2\text{O}_7$ from white to yellow on heating has been attributed to its decomposition into NaPO_3 and Ag_3PO_4 , but it is now shown that this does not occur, the colour change being probably due to some physical change in the salt.

No evidence as to the structure of the pyro-acid could be gained from a study of the dehydration of dihydrogen phosphates of alkali metals. The potassium salt heated at 244° loses water and changes to the acid pyrophosphate, $\text{K}_2\text{H}_2\text{P}_2\text{O}_7$; the rubidium salt behaves in the same way. The sodium salt loses water very slowly at 180° and in the course of about one hundred and seventy-eight hours becomes completely converted into $\text{Na}_2\text{H}_2\text{P}_2\text{O}_7$. The products of further dehydration at a higher temperature depend on the water vapour pressure. In moist air at 305° , only soluble metaphosphate is formed, whilst in dry air at 330° about 75% of the metaphosphate formed is insoluble.

Phosphoryl bromide dehydrates orthophosphoric acid to pyrophosphoric acid, but not to the meta-acid. The action is a complex one, and depends on the temperature and proportions of the interacting substances.

A dilute solution of an alkali pyrophosphate, after prolonged boiling, shows the presence of orthophosphate, proving that hydration occurs slowly.

E. H. R.

Iso- and Heteropoly-acids. XVII. Polyborates in Aqueous Solution. ARTHUR ROSENHEIM and FELIX LEYSER (*Z. anorg. Chem.*, 1921, 119, 1—38).—An attempt was made to prepare simple and complex polyborates with the object of comparing these with salts of other acids such as telluric, antimonie, periodic, plumbic, and stannic acids, which show semi-colloidal properties. Methods for the quantitative estimation of boric acid were examined. The polarimetric method, depending on the influence of boric acid on the optical rotation of tartaric acid, is of limited application on account of the disturbing influence of salts or other substances present in solution. Titration with sodium hydroxide in presence of mannitol, using phenolphthalein as indicator, gives trustworthy results. Free boric acid in the presence of borate can be detected by boiling a sample of the substance for some minutes with dry acetone, filtering, evaporating the filtrate on a watch glass, moistening with a few drops of methyl alcohol, and igniting, when the characteristic flame coloration is given if free boric acid is present.

An investigation of the equilibrium in the system $\text{Na}_2\text{O}-\text{B}_2\text{O}_3-\text{H}_2\text{O}$ at 0° confirmed the existence of the three salts, $\text{Na}_2\text{O}, \text{B}_2\text{O}_3, 8\text{H}_2\text{O}$, $\text{Na}_2\text{O}, 2\text{B}_2\text{O}_3, 10\text{H}_2\text{O}$, and $\text{Na}_2\text{O}, 5\text{B}_2\text{O}_3, 10\text{H}_2\text{O}$. Sodium pentaborate can readily be crystallised from solutions containing Na_2O

and B_2O_3 in the ratio 1 : 5, but sometimes only crystals of borax are obtained, probably because the pentaborate is metastable at ordinary temperatures and borax is the less soluble salt. Potassium pentaborate, $K_2O, 5B_2O_3, 8H_2O$, is a well-defined, characteristic salt, separating from solutions in which the ratio $B_2O_3 : KOH$ is 3 : 1 or higher. Its solubility is very low, not much greater than that of potassium perchlorate. Potassium monoborate crystallises with $8H_2O$ at 0° , with $2.5H_2O$ at 30° . Rubidium pentaborate is very similar to the potassium salt, but crystallises with $10H_2O$. Thallium pentaborate, like the potassium salt, crystallises with $8H_2O$, but is more soluble than the latter. Guanidine forms a diborate crystallising in elongated prisms with $4H_2O$ and a pentaborate with $8H_2O$.

Experiments on the dehydration of pentaborates showed that, in the general formula $R_2O, 5B_2O_3, xH_2O$, two molecules of water are probably constitutively combined. Conductivity experiments indicated that in dilute aqueous solution the pentaborate anion is hydrolysed into the diborate anion and boric acid. In presence of great excess of boric acid this hydrolysis is prevented, and the specific conductivity of sodium pentaborate at 0° appears to be 85% of that of sodium diborate. Experiments on the hydrogen-ion concentration of solutions containing varying ratios of $NaOH$ to B_2O_3 confirm the existence of a pentaborate ion in concentrated solution.

The pentaborate ion appears to form complex anions with a number of metals. Whilst borax solution immediately precipitates zinc or cadmium hydroxide from a solution of a salt of the metal, sodium pentaborate does not. Cobaltous hydroxide dissolves in sodium pentaborate solution to form a red solution in which, although alkalis do not readily precipitate it, the cobalt is in the kation. When this solution is oxidised with hydrogen peroxide, however, some cobaltic oxide is precipitated and a yellow solution is formed containing a complex cobalt anion. Nickel, chromium (Cr'''), manganese (Mn''), and copper also appear to form complex anions. A very small quantity of a copper compound was isolated, having approximately the composition $2Na_2O, 4CuO, 12B_2O_3, 50H_2O$.

E. H. R.

The Atomic Weight of Carbon. E. MOLES (*Anal. Fís. Quím.*, 1921, 19, 255—259).—The value 12.005 for the atomic weight of carbon given by Richards and Hoover (A., 1915, ii, 96) is held to be based on an erroneous value for the atomic weight of sodium. The value 12.000 is claimed to be more exact. G. W. R.

The Oxidising Properties of Carbon Suspensions. F. FEIGL (*Z. anorg. Chem.*, 1921, 119, 305—309).—The oxidising effect of blood charcoal was studied in a qualitative manner by boiling solutions of different oxidisable substances with a suspension of the charcoal. In acid solution, hydrogen sulphide was oxidised to sulphuric acid, potassium iodide to iodine, mercurous salts to mercuric, oxalic acid to carbon dioxide. In alkaline solu-

tion, potassium iodide was oxidised to iodate, alkaline sulphides and sulphites to sulphate, cuprous and cupric sulphides to copper sulphate, cobalt sulphide to sulphate, potassium chromite to chromate. Sodium thiosulphate was unacted on in alkaline or neutral solution, and sodium nitrite was unaffected in alkaline solution. A quantitative study of the oxidation of trivalent chromium to chromate was made, after a method had been devised for removing from the solution a product, formed by the interaction of the charcoal and potassium hydroxide, which liberates iodine from potassium iodide. This was accomplished by boiling with potassium permanganate and removing the excess with hydrogen peroxide. The experiments showed that the proportion of chromate formed increased with the proportion of charcoal used, but that, with a constant quantity of charcoal, the amount of chromate formed increased with the quantity of chromium salt taken. Different charcoals varied widely in their oxidising power, but the differences seemed to bear no relation to the ash content.

E. H. R.

Aqueous Carbonic Acid Solutions. E. WILKE (*Z. anorg. Chem.*, 1921, **119**, 365—379).—The dissociation constant of carbon dioxide solutions was measured by the conductivity method, using a solution through which the gas was being continuously circulated. When an ordinary saturated solution was used, without circulation, the conductivity was found to increase during measurement, probably through electrolytic changes caused by the current. Even with the greatest precautions, variable results were obtained, confirming the observations of earlier workers. It was observed that, by contact with the metal electrodes (gold), even without passage of current, the conductivity gradually increased. In three hours the dissociation constant $K \cdot 10^7$ increased from 3.07 to 4.5. Light seemed to have an effect in the same direction. For measuring the hydrogen-ion concentration, a special hydrogen electrode was used consisting of a palladium capillary into which hydrogen was forced at a pressure of 20 atm. The hydrogen-ion concentration was determined in presence of sodium, potassium, and barium chloride. In these solutions carbon dioxide has the character of a strong acid, increasing with the concentration of salt. The hydrogen-ion concentration increases more rapidly than the total carbon dioxide concentration. The observations can be explained on the assumption that a solution of carbon dioxide in water contains orthocarbonic acid, H_4CO_4 , which, containing no ketonic oxygen, is a very weak acid. In concentrated salt solutions it is dehydrated to form the strong acid $CO(OH)_2$.

E. H. R.

Behaviour of Amorphous Carbon and Sulphur at High Temperatures. Carbon Sulphides. J. P. WIBAUT (*Proc. K. Akad. Wetensch. Amsterdam*, 1921, **24**, 92—101).—The action of sulphur on amorphous carbon at high temperatures has been investigated. Pure sugar charcoal has been heated with sulphur at temperatures from 400° to 1000° under reduced pressure for

prolonged periods of time. A slow evolution of hydrogen sulphide, due to the small amount of hydrogen present in the carbon, is observed and a carbon-like substance containing 1.98% of sulphur obtained. This substance yields no sulphur to toluene, even after prolonged boiling, and the residue after this treatment contained 2.03% of sulphur. Prolonged heating in a vacuum at temperatures up to 1010° did not reduce the sulphur content nor was any volatile compound obtained. Prolonged shaking with bromine water oxidised 9% of the sulphur to sulphuric acid, and heating in a current of hydrogen at temperatures up to 750° removes 77% of the sulphur as hydrogen sulphide; this reaction is exceedingly slow and must be regarded as an action between a sulphur compound and hydrogen and not as an action between hydrogen and sulphur vapour. This was further proved by the fact that heating in nitrogen did not reduce the sulphur content. The author considers that a solid carbon sulphide is formed which bears a strong resemblance to coal coke (cf. Stock and Praetorius, A., 1913, ii, 46). A further sulphide containing 3.5% of sulphur has been obtained by heating carbon purified by chlorine with sulphur. This substance has similar properties to the compound containing 2.0% of sulphur.

J. F. S.

. The Deviations from the Gas Laws of Carbon Disulphide.

ALFRED SCHULZE (*Z. anorg. Chem.*, 1921, **118**, 223—230).—A number of observations on the properties of carbon disulphide vapour indicate that it is associated to a small extent. The increase of pressure observed when carbon disulphide and ether vapours are mixed at constant volume at 80° under atmospheric pressure indicates association of the former to the extent of 0.14%, whilst vapour density determinations by Dumas's method give results corresponding with 2% association. Compressibility experiments at 80° showed 0.5% more association at 2 atmospheres than at 1 atmosphere pressure. The PV curves at 78.82° and 130.48° show that the amount of association decreases with increasing temperature, but at constant temperature increases with increasing pressure. It is probable that in the liquid phase association is more considerable.

E. H. R.

New Theory of the Constitution of Hydroxides, particularly those of the Basic Metallic Oxides. FR. TIEMANN (*Chem. Zeit.*, 1921, **45**, 1125).—To furnish an explanation for a number of phenomena in organic, inorganic, and electrolytic processes which are not in consonance with existing ideas, the author proposes a new theory of the constitution of the hydroxides of the pronounced electropositive metals. It is suggested that these compounds do not contain hydroxyl groups, but are to be regarded merely as true hydrates of the corresponding oxides, that is to say that sodium and calcium hydroxides, for example, are not correctly represented by the formulæ NaOH and Ca(OH)_2 but are actually $\text{Na}_2\text{O}\cdot\text{H}_2\text{O}$, and $\text{CaO}\cdot\text{H}_2\text{O}$, a molecule of water being closely associated with the metallic oxide similarly to the "water of

crystallisation" of salts. This applies to all the elements of groups I and II of the periodic system, whilst the constitution of the hydroxides of those of groups III and IV (aluminium, zinc) will depend on the electrochemical conditions under which they are produced. Only the hydroxides of the metalloids and non-metals are to be regarded invariably as true hydroxyl derivatives. With increasing basicity of the oxides the associated water molecules become increasingly firmly bound exactly as in the case of the increasingly basic character of salts containing associated water. So the dehydration of the hydroxides of calcium, strontium, and barium is effected with increasing difficulty in the order named.

The sucrosates are cited to illustrate the application of the theory. If calcium hydroxide is regarded as a hydroxyl compound, the chemical character of sucrose or dextrose is quite incompatible with the idea of a "neutralisation" of hydroxide, looked on as a generator of hydroxyl ions. There can therefore only be a question of the displacement of the associated water by the sugar, and the sucrosates must be formulated $C_{12}H_{22}O_{11} \cdot 2CaO$, $C_6H_{12}O_6 \cdot CaO$, etc. The isomerism of the hydroxides of tin and aluminium is also explained by reference to the theory, *e.g.* $Al(OH)_3$ and $Al_2O_3 \cdot H_2O + 2H_2O$ can both exist as individual substances, and either one or the other will be produced according to the conditions of the reaction. The non-appearance of hydrogen peroxide, derived from the union of two hydroxyl groups, during electrolysis of a metallic hydroxide is explained by the new theory as due to the absence of hydroxyl groups. The electrolysis actually is that of, say, $Na_2O \cdot H_2O$, the associated water taking no part in the process. There takes place simply a direct fission into metal and oxygen which are liberated at their respective poles. An assumption of the appearance of hydroxyl ions in any electrolytic process is quite unjustified.

G. F. M.

Preparation of Alcoholic Potassium Hydroxide Volumetric Solution. S. T. MCCALLUM (*J. Ind. Eng. Chem.*, 1921, **13**, 943).—A solution which does not darken in colour when kept is prepared by dissolving potassium hydroxide in methyl alcohol (purified wood spirit); the solution must be filtered through glass-wool to remove insoluble potassium carbonate, etc., before it is used.

W. P. S.

Existence of Tetra-hydrated Sodium Sulphate in Mixed Crystals with Sodium Chromate. THEODORE W. RICHARDS and W. BUELL MELDRUM (*J. Amer. Chem. Soc.*, 1921, **43**, 1543—1545).—It is shown that crystals of the tetrahydrate of sodium chromate, $Na_2CrO_4 \cdot 4H_2O$, dissolve sodium sulphate as $Na_2SO_4 \cdot 4H_2O$, a form of sodium sulphate otherwise unknown, to the extent of somewhat less than half the quantity corresponding with the same weight of sodium chromate in the supernatant liquid. When sodium sulphate was in large excess, no crystallisation could be induced by "seeding" the saturated mixture with the crystals of the mixed tetrahydrate above the transition temperature of sodium

sulphate, and below this temperature only crystals of the decahydrate could be formed. Thus under these conditions the tetrahydrate is so much more soluble than the phases containing more sodium sulphate as to be incapable even of metastable existence.

J. F. S.

Ammonium Radicles. III. Ammonium. HANS HEINRICH SCHLUBACH and FRITZ BALLAUF (*Ber.*, 1921, 54, [B], 2825—2834; cf. A., 1920, i, 822 and this vol., i, 15).—The authors' experience with tetraethylammonium leads them to expect that the ammonium radicle would be extremely sensitive to rise in temperature and that there is no hope of isolating it by the electrolyses of solutions of ammonium salts in liquid ammonia on account of the impossibility of avoiding the thermal effect of the current, and that the only prospect of success lies in displacement reactions effected at a low temperature. They find that when well-cooled ammonium chloride is added to a solution of potassium in liquid ammonia at -70° in the apparatus described previously for the preparation of tetraethylammonium, decolorisation of the solution takes place before the calculated volume of hydrogen has been evolved, the deficit amounting to as much as 65%; according to Moissan, the whole of the hydrogen is evolved by the time the solution becomes colourless. The deficit cannot be attributed to the solubility of hydrogen in liquid ammonia since this is shown to be too small to account for the observed effect, and it appears therefore that colourless ammonium is actually present in the solution. This conclusion is supported by the observation that the remainder of the hydrogen is evolved rapidly when the solution is cautiously warmed at about -40° . Repetition of Moissan's experiment shows that the non-observation of the production of ammonium is due to operation in too concentrated solution and consequent decomposition of the radicle by the heat liberated during the reaction. When a solution of potassium (1.8%) is added to a solution of ammonium chloride (1%) in liquid ammonia at -70° , a violent reaction is observed and each drop of added solution is immediately decolorised, formation of ammonium and its conversion into the colourless form appearing to occur instantaneously; tetraethylammonium and ammonium, therefore, stand to one another in the same relationship as triphenylmethyl to methyl. In spite of the violence of the reaction, the yields of ammonium by this method are good and readily reach 50%; the influence of concentration is, however, again apparent, and it is to be expected that an improvement in yield would be observed with more dilute solutions.

The behaviour of ammonium towards the reagents used with tetraethylammonium is described. Corresponding with the rapid isomerisation to the colourless form, the equilibrium is here greatly displaced in the direction of the latter, and it is probable that dissociation and consequent reaction only occur in close proximity to the temperature of decomposition. A reaction with dimethylpyrone could not be observed. Iodine, on the other hand, appears

to react immediately with ammonium, but the quantitative examination of this change could not be completed on account of the experimental difficulties involved. H. W.

Vapour Pressure of the System, Lithium Nitrate : Ammonia.

R. O. E. DAVIS, L. B. OLMSTEAD, and F. O. LUNDSTRUM (*J. Amer. Chem. Soc.*, 1921, **43**, 1575—1580).—The solution of ammonia in lithium nitrate has been studied with the object of finding an absorbent for ammonia in the synthetic production of this gas. The use of ammonium nitrate (Kurilov, A., 1898, ii, 156) and ammonium thiocyanate (Foote and Hunter, A., 1920, ii, 246) suffers from the serious drawback that the liquids produced, when these salts adsorb ammonia, attack metals rapidly. A large number of salts have been tested as to their suitability for this purpose, and of these lithium nitrate alone forms a liquid with ammonia in the absence of water, whilst calcium nitrate tetrahydrate liquefies in the presence of a little water. The ammonia contained in 1 c.c. of the lithium nitrate solution saturated at 24° is equivalent to 26.0 c.c. of 0.95*N* sulphuric acid, whilst that for the calcium nitrate solution under identical conditions is 18.5 c.c. of 0.95*N* acid. Vapour pressure measurements have been made for the solution 36.34% ammonia, 63.66% lithium nitrate, and for several other mixtures containing 6.06—58.66% of water. The solutions of ammonia in lithium nitrate have no action on machine steel, iron wire, and nichrome wire after several months' contact, but nickel steel shows a slight action after several months. The results show that a solution of lithium nitrate in ammonia with a small percentage of water should be a good absorbent for the removal of ammonia from mixtures of nitrogen, hydrogen, and ammonia. The absorption could be effected at 0° and a large proportion of the ammonia released either by a small increase of temperature or by reduction of the pressure. J. F. S.

The Reaction between Persulphates and Silver. GEOFFREY ISHERWOOD HIGSON (*T.*, 1921, **119**, 2048—2055).

Silver Bromate. J. H. REEDY (*J. Amer. Chem. Soc.*, 1921, **43**, 1440—1445).—In an earlier paper (A., 1915, ii, 733) it was shown that the electrode $\text{Ag}|\text{AgBrO}_3|0.1\text{N KBrO}_3$ only reached a steady value (0.631 volt) after being kept for five days, if the bromate was prepared by the action of bromine on silver nitrate solution, but if it was obtained by double decomposition of silver nitrate and potassium bromate, the correct value was at once obtained. Investigation now shows that silver bromate is dimorphous, existing as tetragonal bi-pyramids and as hair-like crystals. The tetragonal crystals are stable at temperatures below 98.5° (the transition point) and the fine, hair-like crystals are stable above this temperature. Difference of solubility of the two forms explains the irregular behaviour of the electrode mentioned above. The solubility of silver bromate has been determined at temperatures from 25° to 90° and the following values have been found: 25°, 0.196; 35°, 0.269; 45°, 0.371; 55°, 0.497; 65°, 0.648; 75°, 0.832; 85°, 1.055

and 90° , 1.325, the solubilities being expressed in grams per 100 grams of water. The solubility curve indicates 98.5° as the transition temperature, a value which is confirmed by a dilatometric measurement of this quantity. Dry silver bromate melts at $308\text{--}310^\circ$ and is stable toward heat and light, but in the presence of water it darkens slowly at the ordinary temperature and rapidly at high temperatures. If a little impurity, such as dust, is introduced into heated silver bromate decomposition occurs with explosive violence. Silver bromate crystals absorb a considerable quantity of air which is slowly evolved at high temperatures.


It is shown that silver bromate may be used as a standard in iodometry. The method of use consists in placing 1 gram of bromate with an excess of potassium iodide in 150 c.c. of water in a 250 c.c. flask; this is heated on a water-bath to effect double decomposition. The contents of the flask are cooled and made up to 250 c.c. Samples of 25 c.c. are withdrawn, acidified with dilute hydrochloric acid, and titrated with sodium thiosulphate. Arsenious oxide gives a result about 0.2% higher than silver bromate, but after recrystallising the arsenious oxide from hydrochloric acid this figure was reduced to 0.05%. This indicates that whilst silver bromate may have a somewhat higher oxygen equivalent than arsenious oxide, this defect is fully compensated by its greater definiteness. J. F. S.

Alkali Silver Thiosulphates and their Ability to Add Ammonia. ERIK JONSSON (*Ber.*, 1921, **54**, [B], 2556—2564).—Additive compounds of alkali silver thiosulphate and ammonia have been described previously by Schwicker (A., 1889, 942) and by Meyer and Eggeling (A., 1907, ii, 347), who, however, do not record analyses of their products. A repetition of their work has given somewhat different results. The ability to form additive compounds seems to depend on the presence of unused subsidiary valencies of the silver atom and is most marked in compounds of the type $\text{K}_2\text{S}_2\text{O}_3\cdot\text{Ag}_2\text{S}_2\text{O}_3$; it is scarcely noticeable in the case of the salts $2\text{M}_2\text{S}_2\text{O}_3\cdot\text{Ag}_2\text{S}_2\text{O}_3$ and $5\text{M}_2\text{S}_2\text{O}_3\cdot 3\text{Ag}_2\text{S}_2\text{O}_3$. The existence of colourless and yellow alkali silver thiosulphates (cf. Meyer and Eggeling, *loc. cit.*) is confirmed, but it appears doubtful whether their isomerism is explicable by assigning the respective formulæ $\text{AgS}\cdot\text{SO}_2\cdot\text{OK}$ and $\text{KS}\cdot\text{SO}_2\cdot\text{OAg}$, since their behaviour towards ethyl iodide indicates that the silver is attached to the sulphur atom in each case. Conversion of the colourless into the yellow modification can be effected frequently by cautious warming with water, but too drastic treatment leads to the formation of silver sulphide, sulphur dioxide, and sulphate. It appears, therefore, that the yellow compounds are intermediate products in the decomposition of the colourless salts, and the transformation is possibly explained by such a scheme as: $\text{KO}\cdot\text{SO}_2\cdot\text{SAg} \rightarrow \text{KO}\cdot\text{S}\cdot\text{SO}_2\cdot\text{Ag}$.


The following individual substances are described: the salt, $2\text{K}_2\text{S}_2\text{O}_3\cdot\text{Ag}_2\text{S}_2\text{O}_3$, colourless prisms, from silver nitrate and potassium thiosulphate in the presence of ammonia; the salt, $5\text{K}_2\text{S}_2\text{O}_3\cdot 3\text{Ag}_2\text{S}_2\text{O}_3$, long, colourless prisms; the compound,

$$3\text{KAgS}_2\text{O}_3\cdot\text{NH}_3\cdot 2\text{H}_2\text{O}$$

(cf. Schwicker, *loc. cit.*, who regards it as $\text{KAgS}_2\text{O}_3 \cdot \text{NH}_3$), colourless plates which are converted by warm dilute ammonia and into a yellow salt of the same composition and are transformed by warm water into the compound, $\text{KAgS}_2\text{O}_3 \cdot 1.5\text{H}_2\text{O}$, colourless needles, and KAgS_2O_3 , yellow, hexagonal pyramids; the salt, $2\text{NaAgS}_2\text{O}_3 \cdot 3\text{H}_2\text{O}$, small, irregular plates; the salt,



long prisms, and the compound, $(\text{NH}_4)_2\text{AgS}_2\text{O}_3$, prismatic crystals (by the action of ammonium thiosulphate on a solution of silver oxide in ammonia); the salt, $5\text{Rb}_2\text{S}_2\text{O}_3 \cdot 3\text{Ag}_2\text{S}_2\text{O}_3$, colourless prisms (cf. Meyer and Eggeling, *loc. cit.*) which is transformed by warm water into the salt, $3\text{Rb}_2\text{S}_2\text{O}_3 \cdot 4\text{Ag}_2\text{S}_2\text{O}_3$, yellow prisms; the salt, $3\text{RbAgS}_2\text{O}_3 \cdot \text{NH}_3 \cdot 2\text{H}_2\text{O}$ (Meyer and Eggeling record



the unstable yellow salt, $\text{Ag}_2\text{S}_2\text{O}_3 \cdot 3\text{NH}_3 \cdot \text{H}_2\text{O}$.

H. W.

Metallic Hydrides. II. Hydrides of the Alkaline-earth Metals and of Lithium. FRITZ EPHRAIM and EDUARD MICHEL (*Helv. Chim. Acta*, 1921, 4, 900—924; cf. A., 1921, ii, 638).—The preparation of the hydrides and the measurement of their dissociation tensions is recorded.

When attempts are made to compare the tensions of the different hydrides with one another, it becomes apparent that all measurement of dissociation pressure of the alkali and alkaline-earth hydrides are vitiated by the use of impure material containing a greater or less proportion of dissolved metal which tends to depress the tension. Within each group, the effect of the metal increases with its atomic weight, and the influence of sublimation lies in the same direction. In the cases of caesium and barium hydrides, these actions render the measurement of dissociation pressures at high temperatures almost impossible. The influence of the atomic weight of the metal on the stability of the alkali hydride cannot be regarded as elucidated completely, but the authors consider from their own experiments that a slight diminution of stability with increasing atomic weight of the metal is probable. The tension curves of lithium hydride could not be measured, since a material which would withstand the chemical action of the hydride and metal could not be found. It is, however, established that it is the most stable of all the alkali or alkaline-earth hydrides which is in accordance with its great heat of formation. Calcium hydride appears to be more stable than barium hydride, whilst the strontium compound occupies an intermediate position.

The behaviour of the alkali and alkaline-earth metals towards hydrogen is not confined to the formation of hydrides, XH and XH_2 , but extends also to the production of solutions, the phenomenon being more marked with the hydrides of the alkaline earths than with those of the alkalis. The absorption of hydrogen occurs previously to, and, in part, simultaneously with, the formation of the hydrides; this occurs to a greater extent with the alkaline earth than with the alkali metals.

The formation of hydrides occurs slowly with the alkali metals,

rapidly and with incandescence in the cases of the alkaline-earth metals. This appears to be due to the greater solubility of the hydride in the metal. The same explanation applies to the observation that calcium hydride, for example, can be formed at a temperature which is certainly considerably higher than the temperature of dissociation of the pure hydride. The liquid nature of the alkali hydrides at the temperature of their formation contributes also to the slowness of absorption of hydrogen, since the eutectic mass protects the metal from further action.

Investigation of the hydrides of lanthanum and cerium (Mathmann and Baur, A., 1903, ii, 213) and of neodymium and praseodymium (Muthmann and Beck, A., 1904, ii, 409) have given results similar to those now observed with the alkaline-earth metals except that the displacement of the tension due to the presence of an excess of metal is even more considerable. The increase in the action of an excess of metal with increasing atomic weight is therefore apparent not only within a group in the periodic system, but also from left to right with increasing valency of the metal.

H. W.

The Discovery of an Equilibrium between Cement and Lime-water. RICHARD LORENZ and GUSTAV HAEGERMANN (*Z. anorg. Chem.*, 1921, **118**, 193—201).—When finely-ground Portland cement, which has been previously treated with water and dried, is stirred with a fixed quantity of water in absence of air, the quantity of lime taken up by the water eventually reaches a maximum value. This maximum is much less than the solubility of lime in water, and depends on the quantity of cement present in proportion to the water, and also, to some extent, on the fineness of the particles. The existence of this maximum is shown to depend on the partition coefficient of lime between the water and the silica-alumina gel formed by the decomposition of the cement constituents such as monocalcium silicate and tricalcium aluminate. This partition coefficient was determined by repeatedly treating the cement with fresh water until the whole of it had decomposed. The ratio of lime in the solid phase to lime in the water was then found to be about 7.0. The existence of the partition coefficient shows that no definite compound is formed between the lime and the constituents of the gel. [See also *J. Soc. Chem. Ind.*, 1922, 15A.]

E. H. R.

The Solubility of Glucinum Sulphate in Water and Sulphuric Acid at 25°. HUBERT THOMAS STANLEY BRITTON (*T.*, 1921, **119**, 1967—1971).

Ammoniates of Magnesium Haloids. WILHELM BILTZ and GUSTAV F. HÜTTIG (*Z. anorg. Chem.*, 1921, **119**, 115—131).—For the investigation of the ammoniates of magnesium haloids special precautions were taken in the preparation of pure magnesium chloride, bromide, and iodide. Ephraim's results (A., 1912, ii, 546), which were not confirmed, were probably due to the fact that his magnesium chloride contained basic salts. The

hexammoniate of magnesium chloride is formed in about fifteen hours at room temperature, when pure magnesium chloride is saturated with ammonia, and the product is extraordinarily voluminous. Magnesium bromide behaves similarly, but in the case of the iodide the increase in volume when the hexammoniate is formed is relatively slight. Observations were made on the time taken for the vapour pressure to become steady with different proportions of ammonia in the solid phase. From these observations definite rules can be formulated regarding heterogeneous equilibrium in a solid-gas system. When two non-miscible substances are present in the solid phase, equilibrium is reached with gradually diminishing velocity, usually in the course of a few hours, depending on the temperature and the nature of the substances. When one constituent just disappears, for instance from a mixture containing principally a lower ammoniate and a small quantity of a higher ammoniate, equilibrium is attained very rapidly. When unsaturated mixed crystals are present, for example of two ammoniates, equilibrium is reached very slowly, often requiring several days. This case occurs with the magnesium haloids containing 5.5 to 5.8 mols. of ammonia. When the ammonia content is very nearly 6 mols., addition of a very small quantity of ammonia, even a few hundredths of a mol., causes a very rapid rise in vapour pressure, equilibrium being rapidly attained. On account of this, the dissociation pressures of the hexammoniates could not be accurately determined. These ammoniates may be compared with the zeolites, which, unlike normal hydrates, lose water very rapidly. The phenomenon has not been observed with other ammoniates.

Magnesium chloride and bromide both form a diammoniate and a monoammoniate; the iodide forms only a diammoniate. The table gives the heats of formation in Cals. and the absolute temperatures at which the dissociation pressure is 100 mm.

	Saturated mixed crystals.	2NH ₃ .	1NH ₃ .
MgCl ₂	13.3; 367°	17.9; 495°	20.8; 573°
MgBr ₂	15.2; 420°	20.1; 559°	21.7; 606°
MgI ₂	17.2; 475°	22.7; 636°	—

E. H. R.

The Solidification Diagram of the Zinc-Arsenic Alloys. W. HEIKE (*Z. anorg. Chem.*, 1921, **118**, 264—268).—Thermal examination was made of alloys containing from 6.6 to 92% of arsenic and with the aid of the results the equilibrium diagram was constructed. Two compounds, both melting without decomposition, are formed, Zn₃As₂, m. p. 1015°, and ZnAs₂, m. p. 771°; the former has a transition point at 671°. With excess of zinc, pure zinc appears with Zn₃As₂, no solid solutions being formed. Arsenic dissolves little zinc, but on the other hand is soluble to a considerable extent in the compound ZnAs₂. Two eutectics are formed, at 62% and 81.5% As respectively, the corresponding temperatures being 730° and 723°. Both compounds are very brittle,

and ZnAs_2 shows a well-marked cleavage. This compound always shows super-cooling during crystallisation of the alloys.

E. H. R.

The Fusion Diagram of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O} + \text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ at Pressures of 1 to 3000 kilo./cm.². MEINHARD HASSELBLATT (*Z. anorg. Chem.*, 1921, 119, 313—324).—The fusion diagram of the system $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O} + \text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ at the normal pressure has been previously determined (A., 1913, ii, 484). It was shown that the stable form of calcium nitrate forms a eutectic with cadmium nitrate at 91% $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ and 40.6° , whilst the unstable form of the calcium salt forms a continuous series of mixed crystals with the cadmium salt. The effect of increased pressure up to 3000 kilos. per sq. cm. on the diagram has now been investigated. The general form of the diagram is unchanged. The p - t curve for mixed crystals containing a high proportion of the calcium salt could not be followed at higher pressures on account of the rapid change of the calcium salt into the stable form. Excess of the cadmium salt inhibits this change, but as the pressure increases, more cadmium salt is needed to produce this effect. With increasing pressure, the m. p. of the stable calcium salt rises much more rapidly than that of the unstable. The latter does not form mixed crystals with cadmium nitrate. The lowering of the m. p. of the stable calcium salt by the cadmium salt is independent of the pressure. With increasing pressure, the eutectic point moves towards the cadmium side; at 1000 kilo./cm.² it is at 79% calcium nitrate, 47.8° ; at 2000 kilo./cm.², 74% and 55° , and at 3000 kilo./cm.², 71% calcium nitrate and 61.5° .

E. H. R.

Light Reactions of the Oxides of Titanium, Cerium, and the Earth Acids. CARL RENZ (*Helv. Chim. Acta*, 1921, 4, 961—968; cf. A., 1921, ii, 316).—Titanium dioxide, cerium dioxide, niobium pentoxide, and tantalum pentoxide are in themselves stable towards light, but become markedly photosensitive in the presence of suitable media. Reaction is due to reduction; this is the more remarkable since the oxides are reducible by purely chemical means with considerable difficulty.

Titanium dioxide, niobium pentoxide, and, to a less degree, cerium dioxide undergo reduction when exposed to light in the presence of certain organic liquids and reducing solutions, particularly glycerol. A lower oxide appears to be formed (zirconium dioxide is not photosensitive and does not form a lower oxide) which on exposure to air or on being heated regenerates the original material. In the case of niobium pentoxide, the process depends to some extent on the presence of impurities, notably stannic and tungstic acids, zirconium compounds, and titanous acid or its anhydrides. Brown vanadium pentoxide becomes black with greater or less rapidity when exposed to light beneath glycerol, benzaldehyde, cinnamaldehyde, cuminol, or aqueous mannitol solution; a lower oxide, initially vanadium tetroxide, is produced. Solutions of citric or tartaric acid in absolute alcohol become green

and ultimately blue when illuminated in the presence of vanadium pentoxide; carbon dioxide is evolved freely. Similar decomposition is observed with mandelic acid, but, in this instance, the vanadium pentoxide is blackened. Brown neodymium oxide containing praseodymium does not alter in appearance when illuminated under glycerol or phenylhydrazine; it becomes bluish-grey when warmed with the latter, owing to conversion of the brown to the yellow oxide of praseodymium. When exposed to sunlight in the presence of glycerol, benzaldehyde or tartaric acid dissolved in alcohol, bismuth oxide becomes grey and ultimately dark black. Reduction to the lower oxide and, possibly, to the metal takes place. In similar circumstances, antimony trioxide is also photosensitive.

H. W.

Concentration of the Erbium Earths. PAUL H. M.-P. BRINTON and C. JAMES (*J. Amer. Chem. Soc.*, 1921, **43**, 1397—1401).—Four methods for the concentration of the less basic of the rare earths have been investigated; the methods examined are: (1) formation of basic nitrates, (2) crystallisation of chlorides from 1:1-hydrochloric acid, (3) formation of basic chlorides, and (4) formation of basic thiosulphates. The authors highly recommend the first method for the separation of erbium, holmium, dysprosium, and the less basic earths from yttrium, and the second method for the separation of holmium and dysprosium from yttrium. The basic nitrate formation was carried out with (a) a solution of yttrium nitrate containing a little erbium, and (b) a yttrium-erbium-holmium mixture. In the former case, the solution was boiled and treated with a fairly concentrated solution of sodium hydroxide and thoroughly boiled. The yttrium hydroxide which first precipitated soon dissolved. The addition of sodium hydroxide was continued until minute crystals of the basic nitrate were observed swirling through the liquid. The whole was then set aside to cool, when a mass of needle-like crystals of the basic nitrate was obtained. These were collected, dissolved in the least amount of nitric acid, and treated with sodium hydroxide solution as before. The basic nitrate crystals from this when dissolved in nitric acid gave a rose-red solution which exhibited intense absorption bands of erbium, thus showing that the erbium was rapidly collecting in this fraction. The original filtrate was treated several times with sodium hydroxide, and although the concentration of nitrates was kept high the erbium absorption bands soon began to fade. The results obtained with the second mixture were equally good. The crystallisation of the chlorides was effected with a solution containing yttrium, holmium, and dysprosium. The solution in hydrochloric acid was boiled down until acid of constant boiling point was obtained. The solution was then evaporated until a scum appeared on the surface, when it was set aside for fifteen to twenty hours. The crystals were separated by decantation and the crystallisation proceeded with; by the time the tail fraction had become No. 9, and the head fraction No. 4 owing to the combination of small

head fractions, it was found that the atomic weights had become 92.5 and 124.0 respectively. After four further fractionations the atomic weight of the tail fraction No. 12 was 91.5 whilst that of the head fraction No. 4 was 133.70. The order of separation in order of decreasing solubility of the chlorides is erbium, yttrium, holmium, dysprosium. J. F. S.

Disglomeration and Formation of the Autogenous Lead Tree. A. THIEL (*Ber.*, 1921, 54, [B], 2755—2758; cf. A., 1920, ii, 622).—Disglomeration, which has been observed previously in the cases of lead and copper, is also exhibited by tin when the latter is preserved for some time under stannous chloride solution in a loosely stoppered bottle.

Large uniform crystals of lead become strongly corroded when preserved for some weeks beneath Heller's solution; a considerable quantity of lead powder is formed but, as expected, there is no evidence of disglomeration, that is, formation of deep fissures at the boundaries of the crystallites. Unexpectedly in the light of the previous theory, the large crystallites readily exhibit the formation of the lead tree when preserved beneath a solution of lead nitrate acidified with nitric acid. The phenomenon is observed only after the formation of a white skin of basic salt on the metal and is explained in the following manner. The presence of the skin inhibits the contact of dissolved lead salt and metal by convection and greatly impedes the diffusion of the lead ion. Beneath the skin, therefore, the solution soon contains practically only lead nitrate and is poor in lead ions, whereas the external solution still contains lead nitrate and therefore has a much higher lead ion concentration. The possibility of the formation of a short-circuited concentration cell is thus provided. H. W.

The Chemical Behaviour of Crystallised Binary Compounds with one Component Nobler than Hydrogen. G. TAMMANN (*Z. anorg. Chem.*, 1921, 118, 93—104).—The author draws a comparison between metallic mixed crystal series and binary compounds. In the former case, the members of a mixed crystal series behave chemically and electrically in a similar manner to one or other component, according to the proportion of each present, with sharply defined limits at molecular fractions which are generally a simple multiple of 1/8. This behaviour is correlated with the lattice structure of the mixed crystals, and may be expected also in crystallised binary compounds which have a similar lattice structure, the difference being that in the latter case the proportions of the two kinds of atom are fixed. It is to be expected that binary compounds will show similarity chemically or electrically to one or other component, and when two or more compounds of the two elements are formed, some will resemble one component and some the other. As an example confirming this view, the compounds of lead and palladium are cited. In this series the compounds Pb_3Pd , $PbPd$, $PbPd_2$, and $PbPd_3$ have been identified. In a solution of lead nitrate against

lead these all give a potential equal to that of palladium, but as soon as any free lead is present the potential drops to zero. Chemically, all the palladium-lead compounds are as resistant as palladium. Binary compounds can be conveniently classed as resistant or non-resistant, the former showing the properties of the nobler, the latter those of the baser component.

The above principles are applied to the discussion of a large number of binary compounds, principally metallic, such as those present in alloys of gold, silver, copper, and platinum, besides sulphides, silicides, and carbides, and it is shown that the compounds can be classified as resistant or non-resistant. The more base the inactive component is, the greater is the number of atoms of the nobler component necessary to protect it. In general, a smaller number of gold than of silver atoms are needed to protect a given atom of a baser metal. These considerations apply to compounds in which one component is nobler than hydrogen. When both are less noble than hydrogen, the classification into resistant and non-resistant does not apply, since both constituents and their compounds decompose water. Apart from compounds of this type, it is possible, in a series of binary crystallised conglomerates, to determine from a few measurements of their galvanic potential which of the components they will resemble in their chemical character.

E. H. R.

The Chemical and Electrical Behaviour of some Series of Alloys. WILHELM JENGE (*Z. anorg. Chem.*, 1921, **118**, 105—122).—With a view to test the theory put forward by Tammann (preceding abstract) that crystallised binary compounds, when no diffusion of the atoms is possible, may be expected to behave chemically and electrically as one or other of the constituent elements a number of series of alloys, in which binary compounds are formed, were examined. The alloys were used as anodes and subjected to the action of halogen, sulphate, or nitrate ions, and were tested against acids and alkalis. In the cobalt-silicon series, in which the compounds formed are Co_3Si , Co_3Si_2 , CoSi , CoSi_2 , CoSi_3 , those having less than 32% of silicon were readily attacked by acidic ions, but those with higher silicon content were resistant. That is to say, CoSi and the higher silicides behave as silicon, the other compounds as cobalt. Towards cold acids, the same compounds were respectively resistant and non-resistant and towards cold alkalis all were resistant except CoSi_3 , the behaviour of which resembled that of silicon. No sharp demarcation of properties was found in the behaviour of the alloys towards hot acids and alkalis because the cobalt loses its passivity and decomposes water. In the series of nickel-silicon alloys somewhat similar results were obtained, but the compound NiSi , unlike CoSi , was not resistant to halogen ions. In the manganese-silicon series only Mn_2Si was non-resistant to acids and all, that is, Mn_2Si , Mn_2Si_3 , and MnSi were resistant to sodium hydroxide.

Alloys of antimony with cadmium and tin and of bismuth with thallium were also examined. The compounds Cd_3Sb_2 and Zn_3Sb_2

have the potentials of cadmium and tin respectively, whilst CdSb and ZnSb soon approximate to the hydrogen potential. The cadmium alloys precipitate antimony and lead from solution, and Zn_3Sb_2 precipitates not only lead, but also cadmium. A bismuth-thallium alloy with the composition Bi_5Tl_3 gives the hydrogen potential, but after etching with hydrochloric acid, the bismuth potential, indicating that the surface thallium atoms have been removed, leaving only bismuth exposed.

Of the lead-thallium alloys, those with over 0.49 mol. of lead show the lead potential, those with 0 to 0.475 mol. of lead show the thallium potential.

Great difficulty was experienced in measuring the potentials of the alloys of magnesium with copper, lead, cadmium, and tin, but it appears that at the moment of contact with the electrolyte they have the magnesium potential, which, however, rapidly falls.

E. H. R.

Physical Chemistry of the Oxides of Lead. II. The Supposed Enantiotropy of Lead Monoxide. SAMUEL GLASTONE (T., 1921, 119, 1914—1927).

Binary Systems of the Sulphates, Chromates, Molybdates, and Tungstates of Lead. F. M. JAEGER and H. C. GERMS (*Z. anorg. Chem.*, 1921, 119, 145—173).—The paper comprises an account of thermal investigations of the binary systems of PbO with SO_3 , CrO_3 , MoO_3 , and WO_3 respectively, and of the different systems formed by pairs of the compounds PbSO_4 , PbCrO_4 , PbMoO_4 , and PbWO_4 . By an optical method the transition temperature of PbO from the red, tetragonal, low temperature form to the yellow, rhombic, high temperature modification was found to be 587° . The melting point of pure lead oxide is 879° . The following melting points were also freshly determined, and differ slightly from accepted values: chromium trioxide, 198° ; molybdenum trioxide, 795° ; tungsten trioxide, 1473° . The examination of the binary systems formed by lead monoxide with the acid anhydrides was limited in each case to the partial system $\text{PbO-PbM}'''\text{O}_4$. In the system PbO-PbSO_4 , the existence of the following compounds was recognised: Pb_4SO_7 , Pb_3SO_6 , Pb_2SO_5 , PbSO_4 . The first has no real melting point, but decomposes at 897° , forming Pb_3SO_6 , m. p. 961° , which has a transition point at 450° . Pb_2SO_5 (lanarkite) has m. p. 977° . PbSO_4 decomposes markedly at 1135° and its m. p. is estimated by extrapolation to be 1170° . The transition point of PbSO_4 from the β to the low temperature α form is $864 \pm 1^\circ$. Three eutectics are formed at 89 mols. % PbO and 835° , 60 mols. % PbO and 950° , and 34 mols. % PbO and 960° .

In the system PbO-PbCrO_4 , the compounds Pb_5CrO_8 , $\text{Pb}_7\text{Cr}_2\text{O}_{13}$, Pb_2CrO_5 , and PbCrO_4 were recognised. The first has no real melting point and exists only below 815° . $\text{Pb}_7\text{Cr}_2\text{O}_{13}$, m. p. 854° , has a reversible transition point at 744° , and forms with Pb_2CrO_5 , m. p. 920° , a eutectic at 68 mols. % PbO and 841° . Lead chromate, PbCrO_4 , is found to be trimorphous; the α -form

is stable below 707° ; the β -form between 707° and 783° , and the γ -form above 783° , melting at about 844° with evolution of oxygen. The heat effect of the change $\alpha \rightleftharpoons \beta$ is small and is sharper in binary mixtures with lead oxide than in the pure substance. The eutectic between PbCrO_4 and Pb_2CrO_5 occurs at 820° , but this part of the equilibrium diagram could not be determined accurately on account of decomposition.

The system PbO-PbMoO_4 shows only two compounds, Pb_2MoO_5 , m. p. 951° , and PbMoO_4 , m. p. 1065° . There are two eutectics, at 87.5 mols. % PbO , 762° , and at 40 mols. % PbO , 933° . The tungstates correspond with the molybdates, Pb_2WO_5 , m. p. 899° , and PbWO_4 , m. p. 1123° , with a transition point at 877° .

The equilibrium diagram for the system $\text{PbCrO}_4\text{-PbSO}_4$ is largely conjectural owing to the considerable amount of decomposition occurring at higher temperatures. There is a gap in the mixed crystal series between about 40% and 30% PbCrO_4 . The mixed crystals have transition points at 934° , 874° , and 748° . In the $\text{PbSO}_4\text{-PbMoO}_4$ system mixed crystals are formed containing up to 6 mols. % sulphate or 2 mols. % molybdate. There is a eutectic at 57 mols. % molybdate and 962° . At 879° , the mixed crystals have a transition point. In the $\text{PbSO}_4\text{-PbWO}_4$ system the mixed crystals of the β -type separating at the eutectic temperature contain respectively 37 mols. % sulphate and 7 mols. % tungstate. The eutectic is at 51 mols. % tungstate and 995° . At 875° occurs the $\beta \rightleftharpoons \alpha$ -sulphate transformation and at 859° the corresponding tungstate change. The $\text{PbCrO}_4\text{-PbMoO}_4$ diagram is largely hypothetical. The composition of the limiting mixed crystals on the molybdate side is 48 mols. % PbCrO_4 at 838° , the eutectic temperature. Transitions occur at 799° of $\gamma \rightarrow \beta$ -chromate mixed crystals, and at 697° $\beta \rightarrow \alpha$. In the $\text{PbCrO}_4\text{-PbWO}_4$ system the eutectic temperature is 837° and the limiting mixed crystals on the tungstate side contain 41 mols. % PbCrO_4 . Lead molybdate and lead tungstate form an isodimorphous mixed crystal series with a transition temperature at 1082° . A mixture containing 75 mols. % PbMoO_4 is in equilibrium at this temperature with both kinds of mixed crystal. E. H. R.

Photochemistry of Thallous Chloride. II. CARL RENZ (*Helv. Chim. Acta*, 1921, 4, 950—960).—A continuation of previous work (A., 1920, ii, 71). Thoroughly illuminated, blackish-brown, thallous chloride, in consequence of photolysis, contains as primary product more or less grey to slate-grey photothallous chlorides in addition to yellow intermediate thallous-thallic chlorides formed in accordance with the scheme: $6\text{TlCl} + \text{Light} = \text{photochloride} + \text{TlCl}_3, 3\text{TlCl}$. Thallic hydroxide, formed by subsidiary actions, is also present. These phases characterise the photo-processes in those cases in which the change of colour extends over the scale, greyish-brown, dark greyish-brown, blackish-brown, and hence occur when thallous chloride is illuminated in the dry condition, under water, and in the presence of solutions of many neutral salts. In the presence of reducing agents or of organic hydroxy-

acids, the action of light on thallous chloride only leads to the production of the photochloride. The formation of thallous-thallic chlorides is not observed in the presence of alkalis or alkali carbonates, which decompose these compounds immediately. The production of photothallous chlorides and of thallic hydroxide does not occur in the presence of hydrochloric acid, even without the addition of organic substances.

Photothallous chloride can be prepared by purely chemical methods if ferrous sulphate is added to a boiling, saturated aqueous solution of thallous chloride and the mixture is treated with an excess of ammonia. The black precipitate of photothallous chloride and iron hydroxides is allowed to settle, and is subsequently washed with hydrochloric acid until the iron compounds are dissolved; the slate-grey photothallous chloride so obtained behaves in exactly the same manner as the photosynthetic product.
H. W.

Ammoniates of Cupro- and Thallo-haloids. WILHELM BILTZ and WILHELM STOLLENWERK (*Z. anorg. Chem.*, 1921, **119**, 97—114).—The formation and vapour pressures at different temperatures of ammoniates of cuprous and thallous chloride, bromide, and iodide were investigated, using apparatus similar to that employed in experiments on the ammoniates of silver haloids (A., 1921, ii, 201). When saturated with ammonia gas, cuprous chloride first shrinks to a yellow mass, then swells and becomes greyish-white. Saturation at -70° to -30° requires at least a day. When the excess of ammonia is allowed to evaporate at room temperature and atmospheric pressure, cuprous chloride triammoniate remains. In damp air, it quickly turns green. The pressure isotherms also indicate the existence of a sesquiammoniate and a monammoniate. Cuprous bromide behaves similarly, forming a white triammoniate, a sesquiammoniate, and a monammoniate. Cuprous iodide absorbs ammonia quickly at room temperature. It forms four compounds, containing respectively 3, 2, 1, and $\frac{1}{2}$ mol. of ammonia. In the following table are given the heats of formation Q in Cals. and the temperatures in absolute degrees at which the dissociation pressures of all these compounds are equal to 100 mm.

	3NH_3 .	2NH_3 .	$1\frac{1}{2}\text{NH}_3$.	1NH_3 .	$\frac{1}{2}\text{NH}_3$.
CuCl	9.48; 283°	—	12.61; 326°	16.73; 417.5°	—
CuBr	9.50; 283°	—	13.15; 339°	14.64; 369.0°	—
CuI	10.37; 286.5°	11.30; 298°	—	14.70; 371.0°	15.22; 390°

Thallous haloids do not absorb ammonia at the ordinary temperature, but in liquid ammonia they all form triammoniates. The vapour pressures are all very close to those of ammonia itself. The triammoniates are soluble to a certain extent in liquid ammonia, the solubility increasing with rising temperature and with the atomic weight of the halogen. The heat of formation is about 7.1 Cal. for the ammonia compound of each of the three haloids. No lower ammoniates are formed.

E. H. R.

The Action of Molten Alkali Chlorides on Copper Oxide.

J. ARVID HEDVALL and GUNNAR BOOBERG (*Z. anorg. Chem.*, 1921, **119**, 213—216).—It was shown in a former paper (Hedvall and Heuberger, A., 1921, ii, 508) that potassium chloride could not be used as a flux in the fusion of cupric oxide with aluminium oxide on account of a reaction taking place between the potassium chloride and copper aluminate. It is now shown that when copper oxide is heated with potassium chloride, cuprous oxide is formed and oxygen evolved. This is best demonstrated by adding cupric oxide in small quantities to a mixture of potassium and sodium chlorides at 1000° and continuing the heating for one and a half hours. At the same time, a basic cupric chloride is formed which, by prolonged heating with sodium or potassium chloride solution, is obtained as the compound $3\text{CuO}, \text{CuCl}_2, 4\text{H}_2\text{O}$. E. H. R.

Phenomena of Diffusion in Metals in the Solid State and Cementation of Non-ferrous Metals. I. Cementation of Copper by means of Ferro-manganese.

G. SIROVICH and A. CARTOCETI (*Gazzetta*, 1921, **51**, ii, 245—261).—A bar of copper was arranged centrally in a porcelain tube glazed internally and the tube then packed with ferro-manganese containing 5% of wood charcoal, both these materials being capable of passing through a sieve with 64 meshes per sq. cm. and of being retained by one of 324 meshes per sq. cm. The tube was closed by means of rubber stoppers luted with sodium silicate, one of the stoppers having two holes to admit a thermo-couple for measuring the temperature and a glass tube bent at right angles and with its end dipping into mercury. After the tube had been heated for some hours at 900° in a Heraeus furnace, considerable proportions of the manganese were found to have penetrated the copper (cf. *J. Soc. Chem. Ind.*, 1922, 17A). T. H. P.

Tervalent Copper. G. SCAGLIARINI and G. TORELLI (*Gazzetta*, 1921, **51**, ii, 225—228).—Contrary to Moser's statement (A., 1907, ii, 549), the action of potassium persulphate on cupric hydroxide in presence of barium hydroxide at temperatures obtained by cooling with ice and salt results in various changes in the colour of the solution and in the deposition of a tenuous amaranth-red precipitate, which may be purified by repeated washing with ice-water by decantation. The compound thus obtained yields oxygen when treated with sulphuric acid, oxidises hydrochloric acid with liberation of chlorine, oxidises ammonia in the cold with production of nitrogen, nitrous acid, and traces of nitric acid, decolorises permanganate, and decomposes potassium iodide with liberation of iodine in quantity greater than that corresponding with the proportion of copper present. Since it does not yield hydrogen peroxide when treated with dilute acid, the compound lacks the grouping characteristic of peroxides and is thus different from the orange-yellow copper peroxide obtained by means of hydrogen peroxide. The ratio between the percentages of copper and active oxygen present is in agreement with the formula Cu_2O_3 . T. H. P.

Production of Single Crystals of Aluminium and their Tensile Properties. H. C. H. CARPENTER and CONSTANCE F. ELAM (*Proc. Roy. Soc.*, 1921, [A], 100, 329—353; cf. A., 1921, ii, 641).—

A continuation of work previously published (*loc. cit.*) on the production of large crystals of aluminium. The metal used in the present work had a purity of 99.6%, the impurity being 0.19% silicon and 0.14% iron. The test-pieces used were 70 mm. with a parallel portion 103 mm. long, 26 mm. broad, and 3 mm. thick, and were estimated to contain 1,687,000 small crystals in the parallel portion ($103 \times 26 \times 3$ mm.). The authors first describe the treatment necessary to convert the whole of the crystals into a single crystal. Three separate processes are shown to be necessary: (i) the aluminium strip is heated at 550° for six hours, (ii) the strip after cooling is subjected to a stress which is equivalent to 378 kilos. per sq. cm., and gives an average elongation of 1.6% on 76 mm., (iii) the test-piece is finally placed in a furnace at 450° and the temperature raised $15\text{--}20^\circ$ per day up to 550° and then for 1 hour at 600° . Applying this treatment to thirty-eight test pieces showed that nine pieces consisted of a single crystal, fourteen of two crystals, nine of three crystals, four of four crystals, and two of six crystals. The tensile strength of aluminium strips consisting of known numbers of crystals has been determined. It is shown that for strips consisting of 150 crystals per 25 mm. it is 708—740 kilos. per sq. cm., and these give an elongation of 36—38% on 76 mm. The tensile strength of strips consisting of a single crystal varies between 598 and 642 kilos. per sq. cm. and these strips suffer an elongation of 34—86% on 76 mm. The varying tensile strength and elongation was accompanied by differences in the type of stretching and fracture. Strips consisting of two crystals have a tensile strength of 441—550 kilos. per sq. cm. and suffer an elongation of 29—70% on 76 mm., whilst strips consisting of three crystals have a tensile strength of 456—567 kilos. per sq. cm. and suffer an elongation of 36—55% on 76 mm. A further series of experiments on the production of single crystals in bars is described. J. F. S.

The Thermal Treatment of certain Complex Aluminium Alloys. LÉON GUILLET (*Compt. rend.*, 1921, 173, 979—982).—In order to determine the effect of each constituent on the behaviour of duralumin under thermal treatment (cf. *ibid.*, 1919, 169, 508), the author has studied alloys of aluminium and copper, aluminium and silicon, aluminium, silicon, and copper, aluminium, magnesium, and silicon, and quaternary alloys containing all four elements. Measurements of hardness have been made on annealed samples, and on samples tempered at different temperatures, the measurements being made in the latter case immediately after tempering and also after the alloy had been kept for forty-eight hours at 20° . From the results of these measurements and from micrographic examinations of the alloys it is shown that the simultaneous presence of silicon, magnesium, and copper is indispensable to obtain the interesting results given by tempering high resistance aluminium alloys. W. G.

Solubility Limits of Carbon in Ternary Steels. I. The System Chromium-Iron-Carbon. KARL DAEVES (*Z. anorg. Chem.*, 1921, **118**, 55—66).—Experiments were made to determine the influence of chromium on the solubility of carbon in iron and to determine the position of the corresponding solubility line in the ternary chromium-iron-carbon diagram. The solubility falls off rapidly at first as the chromium content increases, then more slowly, the general form of the curve being hyperbolic. Points on the curve were determined by observing what chromium content was necessary, with a given carbon content, to cause the appearance of a eutectic in the structure of the metal. To make the hard alloys workable, for the preparation of polished surfaces, it was necessary to heat for several hours at 800°, just below the Ac_1 point, by which treatment the solid solution was broken up and the metal softened. Etching was accomplished by electrolysis in ammonium persulphate solution. In eutectoid alloys, the cementite is practically unattacked by hot sodium picrate solution. Cold alkaline potassium ferricyanide turns the hard constituent of the eutectic brown to yellow, leaving the mixed crystals untouched. The solubility curve explains many of the known properties of chromium steels. The melting point of steel and the arrest points are little affected by chromium up to 10%. [Cf. *J. Soc. Chem. Ind.*, 1922, 16A.] E. H. R.

Solubility Limits of Carbon in Ternary Steels. II. The System Tungsten-Iron-Carbon. KARL DAEVES (*Z. anorg. Chem.*, 1921, **118**, 67—74).—The effect of tungsten on the solubility of carbon in iron was studied in the same way as that of chromium (preceding abstract), and a solubility curve of similar form was obtained, separating eutectic from non-eutectic steels in the ternary diagram. Sudden changes in the physical properties of tungsten steels are correlated with changes of composition involving the passage from one side to the other of this limiting curve. The appearance of so-called double carbides of iron and chromium or of iron and tungsten, observed by different workers, is attributed to the same cause. Small amounts of tungsten in steel raise the melting point, but larger amounts depress it. E. H. R.

The Colour of Iron Alum. JANE BONNELL and EDGAR PHILIP PERMAN (*T.*, 1921, **119**, 1994—1997).

Complex Selenates. JULIUS MEYER (*Z. anorg. Chem.*, 1921, **118**, 1—47).—A large number of new complex selenates and incidentally some simpler compounds which have not hitherto been described, were prepared for comparison with the corresponding sulphates. The new selenates described belong to the chromi- and cobalti-series, and show the closest resemblance to the sulphates, differing from these occasionally only in their water of crystallisation. On account of the ease with which selenic acid is reduced, difficulties were at times encountered in the preparation of certain of the compounds.

[With LEONHARD SPEICH.]—Chromiselenates. Violet chromic selenate, $[\text{Cr}(\text{H}_2\text{O})_6]_2(\text{SeO}_4)_3 \cdot 3(\text{or } 4)\text{H}_2\text{O}$, forms a crystalline powder, readily soluble in water, from which it is precipitated by alcohol or acetic acid. Its aqueous solution dissolves chromic hydroxide with formation of green basic salts. When the violet salt is heated in solution or in the solid state at 90° it changes irreversibly into a green chromiselenate. The green salt prepared in the solid state has the composition $\text{Cr}_2(\text{SeO}_4)_3 \cdot 10\text{H}_2\text{O}$ and dissolves very slowly in water, probably only after addition of water. The green salt may have a constitution of the type $[\text{Cr}(\text{SeO}_4)(\text{H}_2\text{O})_5]_2\text{SeO}_4$. When a solution of the violet salt is boiled for some time, a green compound is formed which is precipitated by alcohol as a green oil and dries to an amorphous, green solid. It is very soluble in water and gives no precipitate with barium salts or with ammonia. It is probably a triselenatochromic acid, $[\text{Cr}(\text{SeO}_4)_3]\text{H}_3$.

Chloropentaquochromiselenate, $[\text{CrCl}(\text{H}_2\text{O})_5]\text{SeO}_4 \cdot 3\text{H}_2\text{O}$, was prepared from chloropentaquochromichloride and sodium selenate; it forms a bright green powder very soluble in water and alcohol. Attempts to obtain other chloro-selenates, corresponding with known chloro-sulphates, were not successful.

Dichlorotetraquochromihexaquochromiselenate,
 $[\text{CrCl}_2(\text{H}_2\text{O})_4](\text{SeO}_4)_2[\text{Cr}(\text{H}_2\text{O})_6]$,

forms a green, crystalline powder, readily soluble in water, slightly so in alcohol. An attempt to prepare a corresponding double chromi-aluminium selenate failed, although sulphates of the type $[\text{CrCl}_2(\text{H}_2\text{O})_4](\text{SO}_4)_2[\text{M}(\text{H}_2\text{O})_6]$, where $\text{M} = \text{Cr}, \text{Fe}, \text{Al}$, or V , were prepared by Werner and Huber (A., 1906, ii, 170).

Hexamminechromiselenate, $[\text{Cr}(\text{NH}_3)_6]_2(\text{SeO}_4)_3$, was prepared from the corresponding nitrate and selenic acid. It is precipitated from aqueous solution by alcohol as a heavy, yellow, finely crystalline powder. The salt is amorphous whilst the corresponding sulphate has $5\text{H}_2\text{O}$.

Chloropentamminechromiselenate, $[\text{CrCl}(\text{NH}_3)_5]\text{SeO}_4$, was prepared from purpureochromichloride and silver selenate. It forms a heavy, red, amorphous powder sparingly soluble in water. The corresponding sulphate is much more soluble, and crystallises with $2\text{H}_2\text{O}$.

Hexacarbamidechromiselenate, $[\text{Cr}(\text{NH}_2\text{CO}\cdot\text{NH}_2)_6]_2(\text{SeO}_4)_3$, prepared from hexacarbamidechromichloride and silver selenate, was obtained as a bright green, finely crystalline powder, moderately soluble in water, from which alcohol precipitates it.

Triethylenediaminechromiselenate, $[\text{Cr en}_3]_2(\text{SeO}_4)_3$, from the corresponding chloride and silver selenate, is a reddish-yellow, heavy, crystalline powder, soluble in water and precipitated by alcohol. When the dry salt is heated at 100° , the colour changes to reddish-violet.

Aluminium selenate, which has not before been described, forms a white, crystalline powder, easily soluble in water and precipitated by alcohol. It appears to contain less than $18\text{H}_2\text{O}$, but the analysis did not distinguish between 15 and $17\text{H}_2\text{O}$.

[With HANNS MOLDENHAUER.]—Complex cobaltiselenates. The

complex cobaltiselenates prepared were confined to those containing only one cobalt complex and to those with 6, 5, or 4 molecules of ammonia or 4 molecules of pyridine.


Hexammine(luteo)cobaltiselenate, $[\text{Co}(\text{NH}_3)_6]_2(\text{SeO}_4)_3 \cdot 5\text{H}_2\text{O}$, corresponds in every respect with luteocobaltisulphate.

Aquopentamminecobaltiselenate, $[\text{Co}(\text{H}_2\text{O})(\text{NH}_3)_5]_2(\text{SeO}_4)_3 \cdot 3\text{H}_2\text{O}$, was prepared both from the corresponding cobaltchloride and from selenatopentamminecobaltiselenate. The salt is similar in physical and chemical properties to roseocobaltisulphate.

Diaquotetramminecobaltiselenate, $[\text{Co}(\text{H}_2\text{O})_2(\text{NH}_3)_4]_2(\text{SeO}_4)_3 \cdot 3\text{H}_2\text{O}$, was prepared from carbonatotetramminecobaltiselenate and selenic acid. It dissolves in water to a deep red solution from which alcohol precipitates it as a bright red, crystalline powder. It loses its water of crystallisation on exposure to air.

Chloropentamminecobaltiselenate, $[\text{CoCl}(\text{NH}_3)_5]\text{SeO}_4$, was prepared from purpureocobaltchloride and silver selenate; it corresponds in its properties with purpureocobaltisulphate.

Chloroaquotetramminecobaltichloride selenate,




was obtained when dichlorotetramminecobaltichloride was treated with silver selenate, through hydration of one of the nuclear chlorine atoms. It forms a violet, crystalline powder, giving a violet aqueous solution.

Nitropentamminecobaltiselenate, $[\text{Co}(\text{NO}_2)(\text{NH}_3)_5]\text{SeO}_4$, from the corresponding chloride and silver selenate, forms bright yellow, microscopic crystals, giving a yellowish-brown, aqueous solution. It forms a periodide, as does the corresponding sulphate.


Sulphatopentamminecobaltiselenate, $[\text{Co}(\text{SO}_4)(\text{NH}_3)_5]_2\text{SeO}_4 \cdot 2\text{H}_2\text{O}$, was prepared from the corresponding sulphato-bromide and silver selenate. It is precipitated from aqueous solution by alcohol in rose-coloured leaflets consisting of microscopic, rhombic tablets. The corresponding sulphatosulphate contains only $1\text{H}_2\text{O}$.

Acid selenatopentamminecobaltiselenate,



was prepared by treating chloropentamminecobaltichloride with concentrated selenic acid. From the diluted solution the acid salt crystallised in reddish-violet crystal aggregates. It closely resembles the sulphato-sulphate and forms the starting material for the preparation of a series of selenatopentamminecobalti-salts including several of the following.

Normal selenatopentamminecobaltiselenate,



was obtained by treating the above acid selenate with alcohol; it has a brighter red colour than the acid salt.

Selenatopentamminecobaltisulphate, $[\text{Co}(\text{SeO}_4)(\text{NH}_3)_5]_2\text{SO}_4 \cdot \text{H}_2\text{O}$, was obtained from the selenato-bromide and silver selenate. It is precipitated by alcohol from aqueous solution in bright red, lustrous tablets. This salt is metameric with the above sulphatopentamminecobaltiselenate, but the two are not isomorphous, as the latter crystallises with $2\text{H}_2\text{O}$.

Selenatopentamminecobaltinitrate, $[\text{Co}(\text{SeO}_4)(\text{NH}_3)_5]\text{NO}_3$, was pre-

pared from the above acid selenate and ammonium nitrate. It separates in well-formed, bright red, sparingly soluble crystals.

Selenatopentamminecobaltibromide, $[\text{Co}(\text{SeO}_4)(\text{NH}_3)_5]\text{Br}$, was prepared from the above acid selenate and hydrobromic acid. It is thrown down by alcohol from aqueous solution as a bluish-red precipitate.

Selenatopentamminecobaltihexachloroplatinate,
 $[\text{Co}(\text{SeO}_4)(\text{NH}_3)_5]_2\text{PtCl}_6 \cdot 2\text{H}_2\text{O}$,
 forms lustrous, orange-red tablets, sparingly soluble in water.

In the tetramminecobalt-series only carbonic acid of the bi-valent acids could be introduced into the complex. With two univalent acid radicles, stereoisomerism becomes possible, and it was found possible to prepare the 1 : 2- and 1 : 6-dinitrotetramminecobaltiselenates.

Carbonatotetramminecobaltiselenates, $[\text{Co}(\text{CO}_3)(\text{NH}_3)_4]_2\text{SeO}_4 \cdot 3\text{H}_2\text{O}$, is similar to the corresponding sulphate, crystallising in dark red leaflets which lose their water of crystallisation on exposure to air.

Acid dichlorotetramminecobaltiselenate, $[\text{CoCl}_2(\text{NH}_3)_4]\text{SeO}_4\text{H}$, crystallises in dark green, well-formed needles but is unstable and readily changes to the chloroaquotetrammine salt described above.

Acid dichlorotetrapyridinecobaltiselenate, $[\text{CoCl}_2\text{Py}_4]\text{SeO}_4\text{H} \cdot 2\text{H}_2\text{O}$, is more stable than the dichlorotetrammine salt; it crystallises in lustrous green leaflets. The salt corresponds with the sulphate described by Werner and Feenstra (A., 1906, i, 450).

1 : 2-Dinitrotetramminecobaltiselenate, $[\text{Co}(\text{NO}_2)_2(\text{NH}_3)_4]_2\text{SeO}_4$, was prepared from flavocobaltinitrate (Jørgensen, A., 1898, ii, 592) and ammonium selenate; it forms dark brown crystals.

1 : 6-Dinitrotetramminecobaltiselenate, stereoisomeric with the last, was prepared from croceocobaltichloride (Jørgensen, *loc. cit.*) and silver selenate; it is precipitated from aqueous solution by alcohol in the form of minute, bright yellow crystals.

The electrical conductivities of many of the above salts in aqueous solution were measured at 25° and their magnitudes were found to agree with the constitutions ascribed to the different salts.

E. H. R.

The Green Colour of Tungsten Trioxide. J. A. M. VAN LIEMPT (*Z. anorg. Chem.*, 1921, 119, 310—312).—Tungsten trioxide generally has a yellow colour, but is sometimes green. A number of explanations of this phenomenon have been offered, but it is now shown experimentally that the green colour is due to reduction at ordinary temperatures by traces of organic matter to lower oxides. Provided the green oxide has not been ignited, the yellow colour may be restored by heating it in a current of oxygen.

E. H. R.

Chlorination by Mixed Carbon Monoxide and Chlorine. F. P. VENABLE and D. H. JACKSON (*J. Elisha Mitchell Sci. Soc.*, 1920, 35, 87—89).—Chlorination is successfully accomplished with a mixture of carbon monoxide and chlorine, containing the former in excess, in the following cases : zirconium dioxide at 480°, stannic

oxide at 400°, magnesium oxide at 475°, aluminium oxide at 450°, ferric oxide at 460°, chromic oxide at 625°, manganese dioxide at 460°, uranoso-uranic oxide at 500°. With chlorine in excess, the requisite temperature for zirconium dioxide is 425° and for ferric oxide 370°.

CHEMICAL ABSTRACTS.

Antimonic Acid and the Use of Sodium Antimonate in Analysis. E. S. TOMULA (*Z. anorg. Chem.*, 1921, **118**, 81—92).—The constitution of antimononic acid and of the salts derived from it has never been satisfactorily settled, and an attempt has now been made to solve the problem by the application of physico-chemical methods. The conductivity of the potassium salt was measured at 25° at dilutions from $V=32$ to 1024, and the basicity of the acid, by the Oswald-Walden rule, was found to be 1. This rules out the possibility that the salt is a pyroantimonate, $K_2H_2Sb_2O_7$, and since it gives a solution having an acid reaction, it cannot be the metantimonate, $KSbO_3$. It must therefore be the orthoantimonate, KH_2SbO_4 . Hydrogen-ion determinations in a $1/1024N$ -solution by the calorimetric method confirmed this view. The dissociation constant at this dilution was found to be $\alpha=0.957$ and the hydrogen-ion concentration $C_H=10^{-6.3}$. The equivalent conductivity of the sodium salt was, on account of its low solubility, determined only at dilutions $V=512$ and 1024, and was found to be of the same order as, although slightly lower than, that of the potassium salt. The hydrogen-ion concentration at $V=1024$ was $C_H=10^{-6.4}$, and it is concluded that the two salts have the same constitution. Delacroix (A., 1898, ii, 340; 1900, ii, 145) and Senderens (A., 1899, ii, 557) both isolated a soluble and an insoluble form of antimononic acid, which they called ortho- and pyro-acids, but they differed as to which was which. Conductivity experiments on the potassium salts show that the soluble acid is the ortho-acid, whether prepared by Senderens's or Delacroix's method. It is concluded, however, that a concentrated solution of antimononic acid is not a true solution but a supersaturated colloidal pseudo-solution, from which the acid soon separates in the insoluble form.

Determinations were made at 18°, 25°, and 33.5° of the solubility of sodium antimonate in water, in aqueous sodium acetate, and in aqueous methyl and ethyl alcohols. Expressed in mg. of $Na_2O.Sb_2O_5.6H_2O$ per 100 c.c. of solution, the solubility at 18° is, in water 56.4, in equal volumes of water and ethyl alcohol 0.1, and in 2.5% sodium acetate 3.1.

The following method is recommended for the estimation of antimony as sodium antimonate. The antimony must be in alkaline solution as sodium sulphantimonate, Na_3SbS_4 , and must be free from potassium, since in presence of potassium salts precipitation is incomplete. The solution is warmed at 80° and stirred while a solution of 30% hydrogen peroxide is run in drop by drop until vigorous evolution of oxygen commences, and it is then boiled until all oxygen evolution ceases. The alkaline solution is neutralised with acetic acid until it is acid to phenolphthalein, but still weakly alkaline to litmus. It is stirred a further quarter

of an hour and then one-half its volume of 96% alcohol is added, after which stirring is continued for ten minutes. After twelve hours, the crystalline sodium antimonate is filtered, washed on the filter with a solution containing 3 grams of sodium acetate, 3 grams of acetic acid, and 400 c.c. of ethyl alcohol per litre, and finally with 50% alcohol. The dried precipitate is separated from the filter-paper, which is burnt separately, is ignited for fifteen minutes in a porcelain crucible, and weighed as sodium metantimonate, NaSbO_3 . Special directions are given for procedure when tin is present, as it is then necessary to redissolve and reprecipitate the sodium antimonate. [See also *J. Soc. Chem. Ind.*, 1922, 12A.]

E. H. R.

The Reaction Limit of Chemical Agents on Copper-Gold Alloys and their Galvanic Tension. G. TAMMANN (*Z. anorg. Chem.*, 1921, 118, 48—54).—The reaction limit in different copper-gold alloys is reached when the molecular fraction of gold present is $1/8$, $2/8$, or $4/8$, according to the chemical reagent used. The reactivity of the mixed crystals may be regarded as due to the loosening of the copper atoms from their lattice combination by the chemical agent or, from another point of view, to the action of the chemical agent on copper atoms which have become detached from the lattice on account of their solution tension. From the latter point of view it was important to determine how the solution tension of the alloys varied with the composition. Measurements were made against a gold electrode in a number of electrolytes, and against silver with silver sulphate as electrolyte. The results showed that the limiting composition beyond which no copper ions appear in the solution and the alloy behaves electrically as pure gold, is at $2/8$ mol. fraction of gold. This method does not give such sharp limiting values as the chemical method, however, owing to the sensitiveness of the galvanic tension to impurities on the surface of the metal. The case of the cell silver | saturated silver sulphate | copper-gold is specially interesting, since, when the proportion of gold in the alloy does not exceed 0.145 mol., silver is visibly precipitated and the metal becomes negatively charged, whilst the alloys richer in gold do not precipitate silver and assume a weak positive charge. This weak positive charge indicates a superficial deposit of silver, so that the surface acts as a silver-gold alloy of corresponding composition. It is shown, from consideration of the mixed crystal lattice, that when $1/8$ mol. of gold or less is present in the copper alloy, conditions are favourable for the formation of silver crystals. It is suggested that those agents which find their active limit at $2/8$ mol. of gold, corresponding with the solution tension limit for copper ions, act first on the copper ions in solution, but as soon as the osmotic pressure of the copper ions exceeds the solution tension, the agent attacks the mixed crystal surface.

E. H. R.

Ruthenium Tetroxide. F. KRAUSS (*Z. anorg. Chem.*, 1921, 119, 217—220).—An aqueous solution of ruthenium tetroxide has

apparently a weak acid reaction, although this is difficult to demonstrate on account of the rapid decomposition of dyes by the solution. The solution behaves as an electrolyte, and is decomposed by the current with formation of a green colour. With alkali hydroxides it forms salts, but only the ammonium salt could be obtained in the pure state. It was prepared by adding concentrated ammonia to a concentrated solution of ruthenium tetroxide in water until the colour changed from yellow to greyish-brown. By evaporating, a salt of the composition $(\text{NH}_4)_2\text{RuO}_5$ was obtained. Under certain conditions, which could not be accurately determined, a mono- and a di-hydrate of this salt were obtained. In the preparation of ruthenium tetroxide, besides the yellow compound, a brownish-red substance was observed, less soluble in water than the tetroxide. This has not been identified. Ruthenium tetroxide can be estimated by distilling it in a current of dry air at 15° into a specially constructed weighed flask, dissolving in a little water, reducing with alcohol, evaporating with dilute hydrochloric acid, igniting in a stream of hydrogen, and weighing the ruthenium. E. H. R.

Mineralogical Chemistry.

Native Antimony from Kern County, California. C. H. BEHRE, JUN. (*Amer. J. Sci.*, 1921, [v], 2, 330—333).—A statement of the results of an examination of nodular masses of antimony with a crust of oxidation products. L. J. S.

Identity of Flagstaffite with Terpin Hydrate. F. N. GUILD (*Amer. Min.*, 1921, 6, 133—135).—A comparison of the crystal constants of flagstaffite (A., 1921, ii, 51) with those of terpin hydrate suggests the identity of these; and this is confirmed by comparative tests made on the natural and artificial materials. Variable results for the m. p. are obtained, owing to loss of water before melting. When heated very slowly the crystals soften at about 100° , and finally melt near 116° . Anhydrous terpin from flagstaffite has m. p. 105° . The terpin hydrate formula, $\text{C}_{10}\text{H}_{20}\text{O}_2 \cdot \text{H}_2\text{O}$, is adopted, since the material analysed had been partly dehydrated by remaining over sulphuric acid. L. J. S.

The Natural Iron Hydroxides. KARL WILLMANN (*Centr. Min.*, 1921, 673—678).—A review is given of the colloidal and crystalline forms of iron hydroxides. The scaly (Rubinglimmer) and acicular (Samtblende or needle-iron-ore) forms of goethite differ in optical characters and are regarded as dimorphous forms of $\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$. New analyses of Rubinglimmer from the Eleonore mine near Giessen gave Fe_2O_3 89.90, H_2O 10.77 = 100.67, and Fe_2O_3 88.11, H_2O 11.97 = 100.08, agreeing with this formula. L. J. S.

Curite, a New Radioactive Mineral. ALFRED SCHÖPF (*Compt. rend.*, 1921, 173, 1186—1187).—The new mineral is found at Kasolo, Katanga, Belgian Congo, as translucent, reddish-brown, acicular crystals on torbernite or as compact or earthy masses, and consists of minute needles with straight optical extinction. It is readily soluble in cold nitric acid and in hot hydrochloric acid; when heated, it turns dark brown. Analysis gives the formula $\text{PbO}, 5\text{UO}_3, 4\text{H}_2\text{O}$.

PbO.	UO ₃ .	H ₂ O.	Fe ₂ O ₃ .	Total.	d^{17}
21.32	74.22	4.00	0.17	99.71	7.192

L. J. S.

Analytical Chemistry.

The Conditions for the Maximum Precipitation of an Amphoteric Electrolyte. ADA PRINS (*Chem. Weekblad*, 1921, 18, 657—658).—The minimum solubility occurs at a definite hydroxyl-ion concentration depending for each amphoteric electrolyte on its solubility product as base and as acid. The concentrations of the positive and negative ions are inversely proportional to their charges.

S. I. L.

Use of Cæsium Chloride in Microchemistry. ENRIQUE HERRERO DUCLOUX (*Anal. Asoc. Quím. Argentina*, 1921, 9, 215—227).—Cæsium chloride may be used as a reagent in microchemistry by reason of the well-defined double chlorides which it forms with different metals. Crystallographic descriptions, with photomicrographs, are given of the double salts thus formed with silver, mercury, lead, platinum, gold, palladium, arsenic, antimony, cadmium, tin, copper, aluminium, iron, zinc, nickel, cobalt, manganese, calcium, magnesium, thallium, cerium, and indium.

G. W. R.

Use of the Zeiss Water Interferometer (Rayleigh-Löwe) for the Analysis of Non-aqueous Solutions. ERNST COHEN and H. R. BRUINS (*Proc. K. Akad. Wetensch. Amsterdam*, 1921, 24, 114—122).—A description of the Rayleigh-Löwe water interferometer and the method of use for estimating the concentration of aqueous solutions are given. The accuracy obtainable with this instrument is about 99.9998% for aqueous solutions. To use this instrument for non-aqueous solutions greater precautions in the fixing of the temperature are required. The thermostat liquid should also be chosen so as to have a refractive index of the same order as that of the solvent used. When these precautions are taken an accuracy equal to that obtained for water solutions may be obtained with solutions in organic solvents.

J. F. S.

A Buffer Solution for Colorimetric Comparison. T. C. McILVAINE (*J. Biol. Chem.*, 1921, **49**, 183—186).—The author covers the whole range from P_H 2.2 to P_H 8.0 by mixing two solutions only, viz. a 0.2*M*-disodium phosphate solution with 0.1*M*-citric acid. A table to obtain any desired P_H is given and also a graph.
G. B.

A Simplified Form of Apparatus for Air Analysis. CHARLES CLAUDE GUTHRIE (*J. Biol. Chem.*, 1921, **48**, 365—371).—The apparatus, which is illustrated in the original, “differs from the well known forms in dimensions rather than in principles or in design.”
E. S.

Estimation of the Gases of the Blood. DONALD D. VAN SLYKE and WILLIAM C. STADIE (*J. Biol. Chem.*, 1921, **49**, 1—42).—Improvements in the technique of using the apparatus previously described for the extraction from the blood and measuring of the carbon dioxide (A., 1917, ii, 422—423), oxygen (A., 1918, ii, 82), and carbon monoxide (A., 1920, ii, 53). All the gases can now be estimated in 1 c.c. of blood by a modified form of the original apparatus with narrow measuring tube enclosed in a water-jacket. In the oxygen estimation, the blood is now laked with water, and the amount of ferricyanide has been greatly reduced; the results are about 5% higher than those obtained by Haldane's method. Particularly in the case of the carbon dioxide estimation, the errors are fully analysed, and examples of calculations are given.
G. B.

Mechanical Shaker and other Devices for Use with the Van Slyke Blood Gas Apparatus. WILLIAM C. STADIE (*J. Biol. Chem.*, 1921, **49**, 43—46; cf. preceding abstract).—A motor-driven shaker, a levelling scale, and tonometer rotator are described with the help of figures.
G. B.

Estimation of Chlorine in Benzaldehyde. SCHIMMEL & Co. (*Ber. Schimmel & Co.*, 1921. 56—61; from *Chem. Zentr.*, 1921, iv, 771).—Benzaldehyde is burnt in a small lamp so constructed that the amount of liquid burnt may be found by weighing at the beginning and end of the experiment. The products of combustion are passed through two U-tubes containing glass beads moistened with 0.02*N*-potassium hydroxide solution, foaming being prevented by the use of a few drops of petroleum. The washings from the absorption apparatus are united and, after addition of an equivalent amount of 0.02*N*-sulphuric acid, concentrated in a closed flask. The solution is then made alkaline to phenolphthalein and the colour discharged by one or two drops of 0.02*N*-sulphuric acid. Five drops of 10% potassium chromate solution are added and the chlorine is estimated in the usual way with 0.02*N*-silver nitrate solution.
G. W. R.

Action of Potassium Ferrocyanide on Silver Haloids. G. B. BONINO (*Gazzetta*, 1921, **51**, ii, 261—265).—Potassium ferrocyanide does not appear to interact with silver bromide or iodide,

but with the chloride it reacts according to the equation : $3\text{AgCl} + \text{K}_4\text{Fe}(\text{CN})_6 = \text{Ag}_3\text{KFe}(\text{CN})_6 + 3\text{KCl}$. The chlorine ion may therefore be estimated in presence of the iodine ion by precipitating and washing the mixed silver haloids, treating the latter with a known volume of 0.1N-potassium ferrocyanide solution, removing the insoluble silver potassium ferrocyanide by filtration, and determining the amount of the residual potassium ferrocyanide by titration with standard permanganate solution. The reaction has not yet been studied quantitatively in presence of bromides.

T. H. P.

Estimation of Small Quantities of Bromides and Chlorides in Iodides. I. M. KOLTHOFF (*Pharm. Weekblad*, 1921, 58, 1568—1569).—The iodide in solution is treated with excess of sodium nitrite in presence of sulphuric acid. After filtering and boiling, to remove iodine and excess of nitrous-acid, chlorides and bromides are estimated by Volhard's method. The method is quantitative, but does not distinguish between chlorides and bromides.

S. I. L.

Bromine Normally Present in Animal Tissues. A. DAMIENS (*Bull. Sci. Pharmacol.*, 1921, 28, 85—93; from *Chem. Zentr.*, 1921, iv, 847).—Thirty grams of finely-divided tissue are extracted by heating with dilute potassium hydroxide solution. The dried residue is incinerated with a mixture of potassium nitrate and sodium carbonate, and the ash dissolved in water. Iodine is estimated in a portion of the solution as follows: the silver haloid precipitate, obtained by addition of silver nitrate solution in the presence of nitric acid, is suspended in water and a stream of chlorine is passed, first in the cold and then in the presence of 1 c.c. of sulphuric acid with warming. Air is passed through and the precipitate removed by centrifuging. The iodine is then estimated by a colorimetric method. Another portion of the solution is precipitated with silver nitrate and nitric acid. The precipitate is removed and the filtrate is treated with zinc and sulphuric acid. When less than a milligram of iodine is present, bromine may be estimated colorimetrically in the filtrate. In the presence of larger amounts of iodine, the filtrate, after the reduction of the silver haloid precipitate, is neutralised with ammonia, diluted to 40 c.c. and after addition of 1 gram of iron ammonium sulphate, concentrated to 10 c.c. Bromine may then be estimated as before (cf. A., 1921, i, 476).

G. W. R.

Estimation of Sulphur in Pyrites. L. GADAIS (*Ann. Chim. Analyt.*, 1921, 3, 330—335).—A critical review of Lunge's method and certain suggested modifications. For accurate work, the original method without alteration is to be preferred, care being taken to adhere strictly to all details of the operations. A variation of this method, in which the insoluble gangue is not filtered off before the precipitation of the iron with ammonia, is not recommended, as this gangue contains substances such as the sulphates of barium, strontium, calcium, and lead which may be partly

dissolved by ammonia and reprecipitated on subsequent acidification and addition of barium chloride. A second modification of Lunge's procedure, which obviates the washing of the gangue, consists in making up the solution in aqua regia to 100 c.c., filtering off 50 c.c., and proceeding with this aliquot portion as in the original method. This is free from serious objection provided the insoluble residue is comparatively small in amount. Finally, a rapid control method, not suitable for accurate work, consists in diluting the original solution to about 800 c.c., adding ammonia directly to this, digesting for two hours at a moderate temperature, cooling, making up the volume to 1000 c.c., and filtering off 500 c.c. in which sulphate is estimated in the usual way. G. F. M.

Volumetric Estimation of Sulphide by Oxidation to Sulphate. H. H. WILLARD and W. E. CAKE (*J. Amer. Chem. Soc.*, 1921, **43**, 1610—1614).—Sulphide is rapidly and quantitatively oxidised to sulphate by an excess of hypobromite in 2.5*N*-sodium hydroxide or by hypochlorite in 4*N*-sodium hydroxide. The excess of oxidising agent is determined iodometrically. The method gives accurate results for the estimation of sulphur in steels and in sulphides if care is taken to exclude all other reducing agents. In the case of steels, the method is used as follows: 5 grams of steel are placed in a flask, through which hydrogen may be passed, which is connected with a 10-bulb tube for absorbing the hydrogen sulphide. Air is removed from the apparatus by hydrogen and 100 c.c. of hydrochloric acid (*d* 1.1) are added. After the reaction has moderated, the solution is heated just to boiling for five minutes after the steel has dissolved. The solution in the absorption tube should contain 6—7 grams of sodium hydroxide in 50 c.c. of solution. The contents of the bulbs are washed with as little water as possible into a flask containing 10 c.c. of 0.3*N*-hypobromite solution and kept for three or four minutes, then 2 or 3 grams of potassium iodide are added, and the solution is diluted to 150 c.c. It is then exactly neutralised with concentrated hydrochloric acid and 5 c.c. excess added and titrated with 0.1*N*-sodium thiosulphate. If hypochlorite has been used instead of hypobromite, more sodium hydroxide must be used. Sulphides which are soluble in hydrochloric acid are estimated in the same way as steels. Insoluble sulphides are ignited with powdered iron in an atmosphere of hydrogen or carbon dioxide, thus producing ferrous sulphide. The estimation is then carried to completion as above. The error of the method is about 0.1%. J. F. S.

Estimation of Sulphurous Acid. VICTOR COPPETTI (*Ann. Chim. Analyt.*, 1921, **3**, 327—330).—The gravimetric method of Haas for the estimation of sulphurous acid, which consists in expelling the sulphur dioxide from the solution under examination by distillation in an atmosphere of carbon dioxide, absorbing the gas in a solution of iodine, and weighing the resulting sulphuric acid as barium sulphate, gives accurate results volumetrically if means are taken to prevent loss of iodine by volatilisation in the

current of carbon dioxide. For this purpose, an apparatus is described consisting essentially of a 300 c.c. flask to contain the iodine solution, to the bottom of which extends the gas delivery tube from the distillation flask. Surmounting the flask is a spherical absorption vessel containing *N*/10-thiosulphate solution, through which the carbon dioxide and iodine vapours leaving the flask must pass. When distillation is complete, the thiosulphate solution containing all the volatilised iodine is allowed to run back into the flask and the excess of iodine in the latter is titrated back with standard thiosulphate solution. G. F. M.

The Estimation of Sulphates by means of a Suspension of Barium Chromate. I. M. KOLTHOFF (*Rec. trav. chim.*, 1921, 40, 686—699; cf. Andrews, A., 1890, 414).—Although it would appear on theoretical grounds that an estimation of sulphate by means of barium chromate was not practicable, the method is valid, since the reaction $\text{BaSO}_4 + \text{CrO}_4^{2-} \rightleftharpoons \text{BaCrO}_4 + \text{SO}_4^{2-}$ proceeds very slowly from left to right. The solution, acidified with hydrochloric acid, should be heated during neutralisation with ammonia to avoid the loss of chromic acid which accompanies the precipitation of barium chromate at ordinary temperatures, and, having regard to the hydrolysis of ammonium chromate, a slight excess of ammonia should be added. Barium chromate is adsorbed by ferric, aluminium, and zinc hydroxides, so that when these metals are present the method gives results which are too low, but trustworthy results can be obtained when ions which affect the reacting substances are absent. In presence of calcium, the results are too low, the error becoming smaller with increasing acidity of the solution and decreasing concentration of calcium. Practical details are given. H. J. E.

Catalytic Action of Copper in the Oxidation of Ammonia by means of Persulphate. G. SCAGLIARINI and G. TORELLI (*Gazzetta*, 1921, 51, ii, 277—280).—Quantitative investigation of the oxidation of ammonia by means of either potassium or ammonium persulphate in presence of copper sulphate shows that the oxygen of the persulphate first oxidises the ammonia to nitrous acid, which is converted into ammonium nitrite, this being decomposed, with liberation of nitrogen, by the heat developed: $2\text{NH}_3 + 3\text{O}_2 = 2\text{HNO}_2 + 2\text{H}_2\text{O}$, $2\text{NH}_3 + 2\text{HNO}_2 = 2\text{NH}_4\cdot\text{NO}_2$, and $2\text{NH}_4\cdot\text{NO}_2 = 4\text{H}_2\text{O} + 2\text{N}_2$. The catalytic effect of copper salts appears to be due to oxidation of the copper to a more highly oxidised compound, which passes on its surplus oxygen to the ammonia.

The estimation of persulphate by means of ferrous salts may be replaced advantageously by the following method, in which use is made of a Schultze and Tiemann's apparatus furnished with a mercury valve (*Z. anal. Chem.*, 1870, 9, 401; *Ber.*, 1873, 6, 1041): 40 c.c. of water and 0.4—0.6 gram of copper sulphate are boiled in the flask for about ten minutes, the caoutchouc tube being then clipped and the boiling continued for five minutes to expel the air. The apparatus is allowed to cool, the mercury rising in

the barometer tube and a vacuum becoming established in the flask. A known volume of the persulphate solution and afterwards about 100 c.c. of ammonia solution are drawn into the cold flask through the clipped tube, the flask being then heated and the evolved gas collected over water and measured. The results obtained in this way agree exactly with those yielded by the ferrous sulphate method.

T. H. P.

[Method for the Estimation of Tri-, Tetra-, and Pentathionates present together in Solution with Sulphite, Thio-sulphate, and Sulphate.] E. H. RIESENFELD and G. W. FELD (*Z. anorg. Chem.*, 119, 225—270). See this vol., ii, 45.

The Accuracy of Dumas's Method for the Estimation of Nitrogen in the Cases of Substances rich in Nitrogen. ERNST MOHR (*Ber.*, 1921, 54, [B], 2758—2767).—A mathematical treatment of the influence of the errors in measuring weight of substance, volume of nitrogen, temperature and pressure on the accuracy of the process.

The usual procedure of estimating the volume accurately to within 0.05 or 0.1 c.c., and the temperature and pressure to degrees Centigrade and millimetres of mercury is sufficient for substances containing 20—25% of nitrogen, but involves considerable error when more than this amount is present. The errors due to inaccurate reading of pressure and temperature cannot be minimised by increasing the weight of substance taken. On the other hand, the errors due to volume and weight of substance become considerable when a small quantity of substance is taken and can be diminished by increasing the amount. The practice of using small weights of material when dealing with substances rich in nitrogen by Dumas's method is to be deprecated; Pregl's method should be used in preference. The error involved in the measurement of pressure does not depend to an appreciable extent on whether the gas is moist or dry (above 50% potassium hydroxide solution), but the error involved in measurement of temperature is lower in the latter case. In spite of this fact, the measurement is generally made in preference over water by reason of the customary large diameter of the Schiff's nitrometer and the formation of foam over the potassium hydroxide solution.

A simple and accurate method of calculation is as follows. The temperature is first brought to whole degrees by addition or subtraction of x° in the direction of smallest change and the pressure is then changed by $3x$ mm. in the same sense as the alteration of temperature.

A plea is entered for the more uniform recording of analytical data in the literature and more precise statement of the exact condition of the nitrogen with respect to moisture as also for the reduction of pressures to 0° .

H. W.

The Kjeldahl Nitrogen Method and its Modifications. A. E. PAUL and E. H. BERRY (*J. Assoc. Off. Agric. Chem.*, 1921, 5, 108—132).—Investigations are described into the most suitable

apparatus and method for the nitrogen estimation, with particular reference to the case of cotton-seed meal as presenting special difficulty. The most suitable type of bulb-trap for use in the distillation is the one in which both inlet and outlet tubes enter the bulb and are bent in opposite directions. It is advisable, although not absolutely necessary, to have enough acid in the receiver to neutralise all the ammonia distilled. Practically all the ammonia appears in the first 75 c.c. of the distillate, and all is in the first 100 c.c. During the initial digestion of the material with acid, the flame should never touch the flask above the surface of the liquid; the flask should be protected by a ring of asbestos. The volume of the digesting liquid should at no time be less than 10 c.c. If mercury is used to aid the digestion there is a loss of from 2 to 15% of ammonia unless enough potassium sulphide is added to precipitate all the mercury before distillation. The use of copper sulphate during digestion does not necessitate the subsequent addition of potassium sulphide. The use of permanganate is unnecessary. Digestion with sulphuric acid alone never gives maximum results. Mercury gives a much more rapid digestion than copper salts, and potassium sulphate is more efficient than sodium sulphate. The most rapid and efficient digestion is given by the use of 0.7 gram of mercuric oxide and 10 grams of potassium sulphate. In this way, the liquid becomes clear in one to one and a half hours and further heating for three hours completes the digestion. The amount of copper used in the digestion has little effect on the result. A 2 gram sample of cotton-seed meal and similar substances is preferable. Digestion should not be carried out in an atmosphere containing nitrous fumes. Nitrates in the digestion mixture are not only entirely lost, but they also bring about a considerable reduction in the amount of ammonia recovered.

A. G. P.

Modification in the Kossel-Neumann Method for the Estimation of Phosphorus in Organic Substances. MARIO A. MANCINI (*Biochem. ter. sper.*, 1921, 8, 4—7; cf. Falk and Sugiura, A., 1915, ii, 577).—0.5 Gram of the substance is heated in a 200 c.c. Kjeldahl flask with 6—8 c.c. of sulphuric acid (d 1.184), the boiling being continued for one hour. Nitric acid (d 1.4) having been carefully added drop by drop, boiling is continued until the evolution of oxides of nitrogen ceases. The operation is repeated four or five times. The clear, light yellow liquid is washed into a beaker with hot water, and ammonium hydroxide solution is added in slight excess, followed by 30—40 c.c. of 40% ammonium nitrate solution and 20 c.c. of 25% nitric acid. The liquid is mixed, at 100°, with 120—140 c.c. of a boiling 3% solution of ammonium molybdate. After being kept for at least two hours, it is filtered and the precipitate washed with a hot solution containing 50 c.c. of nitric acid and 50 grams of ammonium nitrate per litre. Hot 25% ammonium hydroxide solution is poured over the filter, which is washed with hot water until the washings give no reaction with Nessler's reagent. To the filtrate is added

50 c.c. of ammoniacal "magnesia mixture," and, after being kept for twenty-four hours, the precipitate is filtered, washed, and ignited in the usual manner.

CHEMICAL ABSTRACTS.

The Composition and Preparation of a Neutral Solution of Ammonium Citrate. C. S. ROBINSON (*J. Assoc. Off. Agric. Chem.*, 1921, 5, 93—97).—To overcome the confusion attached to the term "neutral ammonium citrate solution," it is recommended that such a solution should be one showing a p_H value of 7.0. The solution contains 45.33 grams of ammonia and 172.00 grams of anhydrous citric acid per litre at 20° and has d 1.09. To prepare a litre of solution 172.00 grams of anhydrous citric acid are dissolved in 700 c.c. of water, nearly neutralised, cooled, and made up to a convenient volume, maintaining the density above 1.09. Of this, 5 c.c. are diluted to about 20 c.c., and standard ammonia solution is added until the colour produced with phenol-red indicator matches that produced with the same quantity of indicator by an equal volume of neutral standard phosphate solution (50 c.c. of $M/5$ -dihydrogen potassium phosphate + 29.63 c.c. of $M/5$ sodium hydroxide in 200 c.c.). The calculated amount of ammonia is then added to the bulk of the solution.

A. G. P.

A Modified Method for the Estimation of Phosphoric Acid.

A. W. CLARK and R. F. KEELER (*J. Assoc. Off. Agric. Chem.*, 1921, 5, 103—105).—Two grams of the sample are dissolved in 30 c.c. of concentrated nitric acid and 10 c.c. of hydrochloric acid. The solution is diluted to 200 c.c. and filtered through a dry filter. A portion equivalent to about 0.25 gram is neutralised with ammonia and acidified with nitric acid. Fifty c.c. of 20% ammonium nitrate solution are added and then sufficient ammonium molybdate solution. After remaining over-night, the precipitate is collected on a Gooch crucible, washed eight times with 2% nitric acid (12—13 c.c. each time), then twice with cold water, and dried for two hours at 120°. Precipitation in the cold and drying at 120° give a less variable precipitate than the usual method. The conversion factor of ammonium phosphomolybdate to phosphoric acid is 0.03723.

A. G. P.

Nephelometric Method for the Estimation of Phosphoric Acid and its Compounds in Small Quantities of Blood.

W. R. BLOOR (*Bull. Soc. Chim. Biol.*, 1921, 3, 451—475).—A detailed description of the author's adaptation (A., 1918, ii, 452) of Kober and Egerer's method (A., 1915, ii, 794).

E. S.

Toxicology of Arsenic. N. TARUGI (*Boll. Chim. Farm.*, 1921, 60, 569—576).—The results of experiments with rabbits and guinea-pigs show that arsenic, either organically combined or mixed mechanically with organic matter, always yields gases containing arsenic when putrefaction occurs. Such evolution of gas may occur quickly and must be borne in mind in cases of suspected arsenical poisoning. In the putrefaction of animal matter, the mechanism by which gaseous arsenic compounds are formed

is more complicated than the action of the "arsenic moulds," since such gases are not formed in the initial stages of the putrefaction where aerobiosis appears to predominate, and since also the gases contain not only alkylarsines but hydrogen arsenide as well.

T. H. P.

The Use of Silica Crucibles for the Estimation of Potassium in Soils. J. S. JONES and J. C. REEDER (*Soil Sci.*, 1921, **12**, 419—432).—For the estimation of potassium in soils by the fusion method, silica crucibles may be used instead of platinum, provided that certain limits of temperature are observed. To ensure perfect fusion, the muffle must reach a temperature of 812° , and to avoid loss of potassium by volatilisation the temperature must not exceed 855° . For this purpose an electrically heated muffle is preferable. An electrical arrangement for heating a silica crucible of the J. L. Smith type, 10 cm. long, 2 cm. in diameter at the top, and 1.8 cm. at the bottom is described and is very satisfactory for this type of work.

W. G.

Detection of Magnesium in Presence of Manganese and Phosphoric Acid. A. PURGOTT (*Gazzetta*, 1921, **51**, ii, 265—266).—In presence of phosphoric acid, manganese is precipitated almost completely as tertiary manganous phosphate, even from solutions containing large proportions of ammonium chloride; the precipitate undergoes gradual or, in the hot, rapid transformation into pale pink crystals of manganous ammonium phosphate, analogous in composition and properties to magnesium ammonium phosphate. The formation of this manganese precipitate and hence the danger of mistaking it for the magnesium precipitate may be avoided by treating the hydrochloric acid solution, not only with ammonia solution to remove the kations accompanying magnesium, but also at the same time with ammonium sulphide, which eliminates the whole of the manganese as sulphide; the filtrate is then tested for magnesium.

If the metals of the third group are precipitated in the usual way with ammonia solution, manganese phosphate is precipitated in considerable quantity in addition to certain amounts of calcium, barium, and strontium phosphates, a little manganese and much calcium, barium, and strontium passing into solution. Similar behaviour is shown by a mixture of magnesium phosphate with a manganese salt, but less magnesium goes into solution and still less manganese is precipitated as phosphate. If the quantity of manganese salt is very considerably greater than that of the calcium, barium, and strontium phosphates, the equilibrium, $\text{Ca}_3(\text{PO}_4)_2 + 3\text{Mn}(\text{OH})_2 \rightleftharpoons \text{Mn}_3(\text{PO}_4)_2 + 3\text{Ca}(\text{OH})_2$, is displaced towards the right hand side and the calcium, barium, and strontium pass into solution almost completely.

T. H. P.

Apparatus for the Gasometric Estimation of Zinc in Zinc Powder. EDGAR BEYNE (*Ann. Chim. Analyt.*, 1921, **3**, 360).—The apparatus consists of a gas burette, the upper part of which below the glass stop-cock is expanded into a bulb of 300 c.c.

capacity, and the lower part is graduated in 0.5 c.c. from 300 c.c. to 375 c.c. The lower end of the burette is connected with a rubber tube to a bulb for adjusting the liquid level and gas pressure in the burette, and the upper end is connected through a small refrigerating spiral to a Koninck apparatus which is charged with the zinc dust under examination, and a mineral acid. In a simpler form of apparatus the latter is replaced by an ordinary flask connected by means of a glass tube with the stop-cock of the burette.

G. F. M.

The Direct Iodometric Estimation of Lead Peroxide.
SAMUEL GLASSSTONE (T., 1921, 119, 1997—2001).

A Very Sensitive Reagent for Copper: the Kastle-Meyer Reagent. PIERRE THOMAS and GEORGES CARPENTIER (*Compt. rend.*, 1921, 173, 1082—1085).—The Kastle-Meyer reagent, which is a 2% solution of phenolphthalein in 20% potassium hydroxide solution decolorised by boiling with zinc powder, gives a pink coloration with copper salts and is capable of detecting 1 part of copper in 100,000,000 parts of water. Four drops of the reagent are added to 10 c.c. of the solution to be tested and then one drop of hydrogen peroxide (5—6 vols.).

W. G.

The Iodometric Estimation of Copper and Arsenic present together, especially in Paris and Schweinfürth Green. I. M. KOLTHOFF and C. J. CREMER (*Pharm. Weekblad*, 1921, 58, 1620—1624).—Arsenic trioxide can be estimated by means of iodine if the cupric-ion concentration is very much reduced, which can be effected by addition of excess of pyrophosphate or tartrate, with which the copper forms complex salts. 0.6—0.8 Gram of the pigment is boiled with 25 c.c. of water and 5 grams of sodium pyrophosphate to a clear solution. After cooling, *N*/10-iodine solution is added until the deep blue solution becomes green; the end-point is very sharp. The iodine required is equivalent to the arsenious oxide present. Ten c.c. of 4*N*-sulphuric acid and 2 grams of potassium iodide are then added, and after ten minutes the iodine liberated by reduction of the cupric compound is titrated with *N*/10-thiosulphate, starch being added towards the end.

The results were checked by analysis of the pigments by the standard Lunge-Berl method. This method was found unsatisfactory for copper, the results being always high. Addition of hydrazine sulphate before the sodium hydroxide reduces the copper compound, the element being precipitated and weighed as metal; this modification is much more accurate than the accepted method.

The iodine titration is rapid and simple, and gives accurate results for both elements.

S. I. L.

Separation and Estimation of Copper, Lead, Antimony, and Tin. Analysis of White Metals. A. KLING and A. LASSIEUR (*Compt. rend.*, 1921, 173, 1081—1082).—0.5—1.0 Gram of the alloy is dissolved in 10 c.c. of hydrochloric acid in the presence of potassium chlorate. The solution is diluted to 100 c.c.,

and neutralised with sodium hydroxide. Any precipitate formed is redissolved by the addition of 4—5 grams of tartaric acid. The liquid is transferred to a conical flask coated inside with wax, and to it are added 10 c.c. of concentrated hydrofluoric acid and then, after half an hour, 10 grams of sodium acetate, 1 c.c. of glacial acetic acid and water to bring the volume to 300 c.c. A white precipitate of lead fluoride is formed, but to the liquid 20 c.c. of a 10% solution of sodium sulphide are added and after a time the precipitate of the sulphides of copper, lead, and antimony is filtered off. In the filtrate the tin may be estimated either by precipitation with cupferron (cf. A., 1920, ii, 452), or electrolytically after decomposition of the complex fluoro-compound by the addition of boric acid and redissolving the tin sulphide by boiling with hydrogen peroxide. The precipitate of the mixed sulphides is extracted with 80 c.c. of sodium sulphide solution (*d* 1.14) and the antimony estimated electrolytically in the extract after the addition of potassium cyanide. The copper and lead sulphides are dissolved in nitric acid and the two metals estimated simultaneously by electrolysis. W. G.

Estimation of Mercury in the Mercurial Pills of the [French] Codex. MAURICE FRANÇOIS (*J. Pharm. Chim.*, 1921, 24, 369—379; *Ann. Falsif.*, 1921, 14, 340—347).—The pills are heated with concentrated nitric acid for two hours on a water-bath, and after a further two hours the solution is filtered, and the destruction of the organic matter is completed by adding a small quantity of a 10% solution of bromine. After ten minutes the liquid is rendered strongly alkaline with sodium hydroxide and after adding potassium iodide the mercury is precipitated in the form of a grey powder by means of formaldehyde solution. After two hours the supernatant liquid is decanted through a filter, the mercury is washed with dilute sodium hydroxide solution, and is then treated, together with the filter-paper and its contents, with diluted acetic acid and *N*/10-iodine solution. After ten minutes' agitation, the mercury is completely transformed into mercuric iodide, which dissolves in the potassium iodide present, and the excess of iodine is titrated back with thiosulphate. The method is of general application to all the mercurial pills of the French Codex without modification except in the case of opiated mercurous iodide pills, with which, after treatment with nitric acid, a crystalline precipitate of mercuric iodonitrate, $\text{HgI}_2 \cdot \text{Hg}(\text{NO}_3)_2$, is formed. All that is necessary in this case, however, is to dissolve the crystals in the liquor by addition of potassium iodide, and omitting the bromine treatment, to proceed with the precipitation of the mercury as above. In pills containing soap, although the fat acids are not destroyed by the nitric acid treatment, they do not retain any mercury in combination, and form a layer on the surface of the liquor, which solidifies on cooling, and is therefore easily separated. G. F. M.

Volumetric Estimation of Aluminium. ERNST JOSEF KRAUS (*Chem. Zeit.*, 1921, 45, 1173).—The neutral or faintly acid solution containing the aluminium in the form of sulphate, and free from

other interfering metals, is titrated with standard disodium hydrogen phosphate solution, a few drops of silver nitrate solution being used as indicator, as yellow silver phosphate only commences to form after all the aluminium has been precipitated as phosphate according to the equation: $\text{Al}_2(\text{SO}_4)_3 + 2\text{Na}_2\text{HPO}_4 = 2\text{AlPO}_4 + 2\text{Na}_2\text{SO}_4 + \text{H}_2\text{SO}_4$. The titration is preferably carried out in a boiling solution as the silver phosphate is more pronouncedly yellow in colour and therefore more easily noticeable under these conditions. In presence of other metals, such as iron, etc., the aluminium should first be separated, by addition of excess of sodium hydroxide, for example, and finally precipitated as hydroxide with ammonia. The precipitate after washing is dissolved in a slight excess of dilute sulphuric acid to form a solution suitable for the titration. The method gives good results even when only very small quantities of the metal are to be estimated. G. F. M.

Improved Method for the Separation of Iron and Manganese. M. CARUS (*Chem. Zeit.*, 1921, 45, 1194).—In the usual method for the separation of manganese from the metals of the iron group by precipitating the latter as basic acetates, the contamination of the precipitate with manganese is not due to the co-precipitation of the basic acetate of this metal, but to the formation of insoluble higher oxidation products owing to the action of dissolved oxygen. A perfect separation of the manganese in one operation, even when a large excess is present, is obtained by causing the precipitation with sodium acetate to occur in presence of a small quantity of hydrogen peroxide, in which case no oxidation of the manganese salts can occur. The precipitate obtained, after washing with dilute acetic acid containing a small quantity of sodium acetate and hydrogen peroxide, and finally with hot water, is then completely free from manganese. G. F. M.

The Acidimetric Estimation of Dichromate. I. M. KOLTHOFF and E. H. VOGELENZANG (*Rec. trav. chim.*, 1921, 40, 681—685).—By reason of the hydrolysis of potassium chromate solution, neither chromic acid nor dichromate can be neutralised using phenolphthalein as indicator. A study of the neutralisation of dichromate with sodium hydroxide leads to the following conclusions:—Thymolphthalein should be used as indicator; if phenolphthalein is used the solution must have been saturated previously with sodium chloride, or barium chloride must be used to precipitate the chromate formed. The barium chloride may only be added when the solution is already yellow, otherwise the precipitate of barium chromate contains chromic acid. It is claimed that the results are correct to within 0.2%, but the method is not recommended owing to difficulties with the indicator.

H. J. E.

Modified Method for the Estimation of Iron and Vanadium after Reduction by Hydrogen Sulphide. G. E. F. LUNDELL and H. B. KNOWLES (*J. Amer. Chem. Soc.*, 1921, 43, 1560—1566).—The estimation of iron or of vanadium by reduction with

hydrogen sulphide followed by titration with potassium permanganate or dichromate ordinarily leads to high values. These high values are not due to the presence of sulphur, and persist in spite of the complete expulsion of hydrogen sulphide and the avoidance of organic matter extracted from filter-paper. They may be ascribed to polythionic acids which are not destroyed by boiling in moderately concentrated sulphuric acid solution and are volatilised slowly from dilute and more rapidly from concentrated solutions. Reasonably accurate estimations may be made by restricting the volume of the original solution to 100 c.c. and proceeding by the following modified process. The solution is acidified with sulphuric acid until it contains 2.5% sulphuric acid by volume, reduced by hydrogen sulphide for thirty minutes in cold solution, and for a further fifteen minutes, during which the solution is raised to the boiling point. The hot solution is treated with 15 c.c. of 1:1-sulphuric acid and boiled for thirty to sixty minutes, during which a stream of carbon dioxide is passed through, until the total volume of liquid is about 50 c.c. The solution is cooled, the current of carbon dioxide being maintained, diluted to 200 c.c., and titrated with standard permanganate solution. In some cases it is advisable to filter off the separated sulphur after the hydrogen sulphide treatment. The filtrate is treated for a further ten minutes with hydrogen sulphide and the process completed as above. The average errors found by this method are for iron 0.5% and for vanadium 0.1%. For accurate estimations of iron and vanadium in solutions containing platinum (as in rock analysis), a preliminary separation of the hydrogen sulphide group, followed by expulsion of the gas, complete oxidation with permanganate, and reduction with sulphur dioxide is recommended. J. F. S.

Estimation of Vanadium and Chromium in Ferrovandium by Electrometric Titration. G. L. KELLEY, J. A. WILEY, R. T. BOHN, and W. C. WRIGHT (*J. Ind. Eng. Chem.*, 1921, **13**, 939—941).—Three grams of the ferrovandium are dissolved in a mixture of nitric acid and hydrochloric acid, sulphuric acid is then added, the mixture evaporated to expel all nitric acid and hydrochloric acid, cooled, and the solution diluted to 1000 c.c. One hundred c.c. of this solution are treated with 25 c.c. of sulphuric acid (d 1.58), diluted to 300 c.c., boiled, and 20 c.c. of 10% ammonium persulphate solution and 10 c.c. of 0.25% silver nitrate solution are added; the mixture is boiled for ten minutes, 5 c.c. of hydrochloric acid (1:3), are then added, the boiling is continued for a further ten minutes, the mixture treated with 26 c.c. of sulphuric acid, cooled at 5°, and titrated with ferrous ammonium sulphate solution, the end-point of the titration being determined electrometrically. This titration is a measure of the vanadium and chromium together. The vanadium is estimated by boiling 100 c.c. of the original solution with a few c.c. of ferrous sulphate solution, adding 20 c.c. of sulphuric acid, d 1.58, and 40 c.c. of nitric acid (d 1.40), diluting the mixture to 200 c.c., and boiling it at such a rate that the volume is reduced to 100 c.c. in one hour.

The solution is then cooled and titrated as before. The difference between the two titrations is a measure of the amount of chromium present.

W. P. S.

[**Estimation of Antimony as Sodium Antimonate.**] E. S. TOMULA (*Z. anorg. Chem.*, 1921, **118**, 81—92).—See this vol., ii, 74.

Estimation of Bismuth. O. A. CRITCHETT (*Eng. and Min. J.*, 1921, **112**, 58).—A solution of the ore in a mixture of nitric and hydrochloric acids, having been evaporated with sulphuric acid until fumes appear, is diluted, treated with a drop of hydrochloric acid, filtered, and the filtrate boiled with sodium thiosulphate and aluminium foil. The precipitate is collected, returned to the beaker with hot water, potassium hydroxide added, and the liquid, after having been boiled, is filtered through the original paper. The latter, with the precipitate, is repeatedly evaporated with nitric and sulphuric acids until the paper is completely destroyed. After dilution, boiling, filtering, addition of a slight excess of ammonium hydroxide, and again boiling, the precipitate is collected, washed, and dissolved in hot dilute nitric acid. The bismuth may then be estimated (a) as oxide after treatment with ammonium carbonate, (b) as oxychloride after neutralising with ammonium hydroxide, adding a little hydrochloric acid, diluting, and boiling, or (c) by titration with permanganate after dilution, treatment with ammonium oxalate, boiling, and separation, with subsequent washing by decantation with hot water, of the precipitate of bismuth oxalate.

CHEMICAL ABSTRACTS.

Analytical Chemistry of Tantalum, Columbium, and their Mineral Associates. I. The Use of Tartaric Acid in the Analysis of Natural Tantalocolumbates. II. The Separation of Zirconium from Tantalum and from Columbium. WALTER RAYMOND SCHOELLER and ALAN RICHARD POWELL (*T.*, 1921, **119**, 1927—1935).

Evaluation of the Degree of Unsaturation of Mineral Oils in the Bergius Process. H. I. WATERMAN and J. N. J. PERQUIN (*Rec. trav. chim.*, 1921, **40**, 677—680; cf. Dean and Hill, Technical Paper 181, *Bureau of Mines*, 1917).—Determination of the iodine number of an unrefined mineral oil, before and after treatment by the Bergius process, shows that it is a little greater after treatment.

H. J. E.

Estimation of Phenanthrene. ARTHUR G. WILLIAMS (*J. Amer. Chem. Soc.*, 1921, **43**, 1911—1919).—The phenanthrene is oxidised by iodic acid to phenanthraquinone, which is precipitated as toluphenanthrazine by means of 3:4-tolylenediamine and weighed as such, the procedure being as follows:

For materials containing 30% or more of phenanthrene, 0.25 gram is weighed into a 50 c.c. conical flask, 0.75 gram of iodic acid and 20 c.c. of glacial acetic acid are added and the mixture is boiled for two and a half hours under an air condenser. After cooling for

several hours, any anthraquinone formed from anthracene present in the original material is filtered off on a Gooch crucible and washed with the minimum amount of glacial acetic acid. The filtrate and washings are evaporated to slightly less than 25 c.c. and then the volume is made exactly to 25 c.c., the mixture is cooled and 1 gram of 3:4-tolylenediamine is added and the flask left in running water at 20° over-night. The toluphenanthrazine is collected in a Gooch crucible and washed first with 25 c.c. of 50% acetic acid saturated with the phenanthrazine and then with 200 c.c. of cold water. The precipitate is dried and weighed and to the weight is added 0.053 gram to allow for the toluphenanthrazine remaining in solution in the 25 c.c. of glacial acetic acid. The factor for conversion into phenanthrene is 0.6052. Carbazole, if present in amounts exceeding 10%, interferes in the determination of phenanthrene, and some of the high-boiling coal tar constituents interfere; a crude anthracene cannot be directly analysed.

For the detection of phenanthrene, the material is oxidised as described above, the mixture being cooled and filtered. The filtrate is poured into water and the precipitate collected and washed with water. The precipitate is warmed with concentrated sodium hydrogen sulphite solution and any residue is filtered off. The filtrate is washed in a separating funnel with one or two portions of carbon tetrachloride and then, after the addition of a fresh portion of carbon tetrachloride, is acidified with hydrochloric acid containing ferric chloride. The carbon tetrachloride layer, which contains the phenanthraquinone, is separated and tested by Hilpert and Wolf's reaction (cf. A., 1913, ii, 733), using a solution of antimony pentachloride in carbon tetrachloride, a purplish-red precipitate being obtained on boiling if phenanthrene was originally present.

W. G.

Chemical Analysis of Caoutchouc Articles. ANDRÉ DUBOSC (*Ann. Chim. Analyt.*, 1921, 3, 335—344).—A résumé of the methods employed and the estimations necessary for a complete chemical analysis of caoutchouc and ebonite articles. (1) The acetone extract contains the natural resins, added resins, free sulphur, oils, and waxes, and is evaporated to dryness and quantitatively examined for these substances by the usual methods. (2) The chloroform extract should be almost colourless. Brown coloration indicates the presence of tar or asphaltic adulterants. (3) The extract in alcoholic potash of the insoluble residue of the chloroform extraction should not exceed 15% from a material containing about 50% of caoutchouc, a larger proportion indicating added oils or fats. (4) The aqueous extract reveals the presence of starch or dextrans. (5) The estimation of total sulphur is best carried out by Henriquez's method as modified by the Bureau of Standards. (6) The ash is determined on the residue from the acetone extraction, and a rough estimate of the caoutchouc content can be obtained by subtracting ash and total sulphur from 100. (7) None of the various methods which have been proposed for the estimation of caoutchouc itself give entire satisfaction. The

total foreign matter may be obtained, however, by solution in boiling nitrobenzene, diluting with chloroform, filtering through a tared filter, washing with acetone, drying, and weighing. G. F. M.

Estimation of Oxalic Acid in Urine. E. SALKOWSKI (*Biochem. Z.*, 1921, **118**, 259—266).—A reply to Bau's criticism of the author's method (A., 1921, ii, 356). H. K.

Estimation of the Amino-acids of Feeding Stuffs. T. S. HAMILTON, W. B. NEVENS, and H. S. GRINDLEY (*J. Biol. Chem.*, 1921, **48**, 249—272).—Further improvements are made in the application of Van Slyke's method to the estimation of amino-acids in feeding stuffs (cf. Eckstein and Grindley, A., 1919, ii, 204). Non-protein nitrogen is first removed from the material by successive extractions with anhydrous ether, cold absolute alcohol, and cold 1% trichloroacetic acid, any protein removed by the latter being recovered by precipitation with colloidal ferric hydroxide. The main portion of the protein is then extracted with dilute (0.2%) sodium hydroxide. Starch is removed from the residue by treatment with hot 2% trichloroacetic acid and the remaining protein extracted by treatment first with boiling 20% hydrochloric acid and then with cold 5% sodium hydroxide. A small quantity of protein extracted with the starch is recovered by precipitation of the latter by addition of alcohol. The various fractions of protein thus obtained are hydrolysed with concentrated hydrochloric acid, united, and submitted to the Van Slyke analysis. The method is applied to oats, corn, cotton-seed meal, and lucerne. E. S.

Estimation of Hippuric Acid in Urine. J. SNAPPER and E. LAQUEUR (*Arch. Néerl. Physiol.*, 1921, **6**, 48—57).—To 100 c.c. of urine 25 grams of sodium chloride are added, and a little concentrated hydrochloric acid. An aliquot portion is extracted six times with ethyl acetate, and the total extract is washed once with one-quarter of its volume of water. The wash water is washed with an equal volume of ethyl acetate, which is added to the original extract. After evaporation of the ethyl acetate, the urea is decomposed by sodium hypobromite, and the residual hippuric acid estimated by Kjeldahl's method. G. B.

The Rotation of Dextrose in Solutions of Trisodium Phosphate. Mutarotation as an Analytical Method. HANS MURSCHHAUSER (*Biochem. Z.*, 1921, **117**, 215—225).—The mutarotation of dextrose is accelerated by trisodium phosphate. It follows a unimolecular law, the velocity constants being also linear functions of the concentration of sodium phosphate. As the mutarotation is a function of the hydroxyl ion, its use is indicated for distinguishing salts of different alkalinity. H. K.

Conditions Affecting the Quantitative Estimation of Reducing Sugars by Fehling's Solution. Elimination of certain Errors Involved in Current Methods. F. A. QUISUMBING and A. W. THOMAS (*J. Amer. Chem. Soc.*, 1921, **43**, 1503—1526).—The various sources of error in the current methods of

using Fehling's solution are discussed. The inter-relationship of temperature, 60—110°, and time of heating, ten minutes to two hours, to determine the best time and optimum temperature for reduction has been investigated. Fehling's solution has been studied from the point of view of the nature and concentration of the alkali, concentration of copper sulphate and potassium sodium tartrate, to find the maximum and minimum concentrations of these constituents necessary to give the greatest yield of cuprous oxide and to ensure the formation of the complex cupric tartrate ion. Auto-reduction of Fehling's solution at different times and temperatures of heating has been measured, showing those conditions under which absolutely no "blank" reduction is obtained. Conditions affecting the physical properties of the precipitated cuprous oxide, the photosensitiveness, and the keeping quality of Fehling's solution are recorded. A study of surface oxidation involving different methods of heating and vessels of various sizes has been carried out in order to determine the loss of copper due to surface oxidation and how to avoid it. The catalytic effect of the walls of the container has been demonstrated. A modified method of procedure for the estimation of sugars by means of Fehling's solution is described. In this method the solutions required are, (1) copper sulphate solution containing 82.4 grams of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ per litre, (2) alkaline tartrate solution; 376 grams of crystallised potassium sodium tartrate dissolved in water in a litre flask, and the calculated amount of sodium hydroxide solution added to make 1 litre of this solution containing 130 grams of sodium hydroxide. The sodium hydroxide is made from material purified by alcohol which is kept in concentrated solution for several days to allow carbonates and other insoluble impurities to separate. To make an estimation, 25 c.c. of the copper sulphate solution, 25 c.c. of the alkaline tartrate solution, and 50 c.c. of the sugar solution are placed in a beaker and warmed on a water-bath at 80°. After exactly thirty minutes, the cuprous oxide is filtered on a Gooch crucible and washed. This may either be dried and weighed as cuprous oxide, or dissolved in nitric acid and estimated electrolytically. From the weight of copper, the amount of sugar may be obtained from tables which are given in the paper or by means of the following equations, in which x is the amount in milligrams of copper obtained from y mg. of sugar: dextrose, $y = 0.474x + 0.000115x^2$; levulose, $y = 0.526x + 0.000078x^2$; lactose, $y = 0.813x + 0.000003x^2$; maltose, $y = 0.990x + 0.0000005x^2$; and invert-sugar, $y = 0.504x + 0.0000870x^2$. This method is designed for use in the analysis of saccharine materials containing sucrose, dextrose, levulose, invert-sugar, lactose, and maltose. It is not claimed that the method can be used for the accurate quantitative estimation of 0.1% or less of invert-sugar in approximately pure sucrose.

J. F. S.

Errors in the Estimation of Sugar by Fermentation with Yeast. CARL LANGE (*Berlin Klin. Woch.*, 1921, 58, 957—959; from *Chem. Zentr.*, 1921, iv, 848).—The estimation of sugar in

urine by fermentation with yeast is subject to certain errors. The evolved gas is not entirely carbon dioxide, but contains also nitrogen from the reaction of carbamide, ammonia, or amino-acids with nitrites. The latter may be formed by reduction of nitrates in the urine, or may occur in the yeast. There may also be an evolution of carbon dioxide from carbonates present in the urine. Errors may be diminished to some extent by sterilisation, and precipitation of carbonates with calcium chloride. The estimation of sugar in urine, however, gives only an approximate value.

G. W. R.

Limitations of the Modified Lewis-Benedict Method of Blood Sugar Estimation. RUTH A. GUY (*Biochem. J.*, 1921, **15**, 575—576).—0.0001 Gram of acetone added to 2 c.c. of blood produces an observable colour change in Lewis and Benedict's picric acid method of sugar estimation (*A.*, 1915, ii, 111), but no change is noted with similar quantities of acetoacetic acid. G. B.

The Catalytic Power of Flour. O. FERNÁNDEZ and A. PIZARROSO (*Anal. Fís. Quím.*, 1921, **19**, 265—268).—The catalytic activity of flour, measured by its decomposition of hydrogen peroxide, has been proposed as a measure of its grade of extraction. The results obtained by the authors, using samples of flour of different origin, showed no correlation between catalytic activity and total- or phytin-phosphorus. The test is held to be of little value.

G. W. R.

Identification of Oxycellulose by means of the Barium Compound. ERNST BECKER (*Zellstoff u. Papier*, 1921, **1**, 5—7).—When 2 grams of oxycellulose are shaken with 50—60 c.c. of barium hydroxide solution for four hours, and the insoluble portion is washed with water until the filtrate is free from barium compounds, a barium compound of oxycellulose remains, of which the content of barium varies according to the origin of the oxycellulose.

CHEMICAL ABSTRACTS.

Benzidine Hydrochloride as a Reagent for Wood Cells. C. VAN ZIJF (*Pharm. Weekblad*, 1921, **58**, 1539—1542).—A solution of 0.2 gram of benzidine in 19 c.c. of water, to which 1 c.c. of 25% hydrochloric acid has been added, gives an orange stain to ligneous matter in microscope sections. Sections of material containing tannins or acids which act on iron should be washed free from iron with a 1% solution of hydrochloric acid in alcohol after cutting. Starch can be stained with iodine in the same section without interfering. The reagent also detects diseased tissues in Hevea bark, and for this reason, as well as because of its cheapness as compared with phloroglucinol, is of great use for control on rubber plantations.

S. I. L.

Detection of Formaldehyde with Phenols. B. PFYL, G. REIF, and A. HANNER (*Chem. Zeit.*, 1921, **45**, 1220—1221).—Phenol reactions for the detection of formaldehyde as previously carried out (cf. *A.*, 1921, ii, 663) are not sufficiently trustworthy

when the test is to be adapted to the detection of methyl alcohol in potable spirits and tinctures after distillation with permanganate, as colour reactions are often simultaneously given by the traces of other aldehydes, alcohols, etc., which pass over into the distillate, and the mixed colours produced allow of no certain conclusions being drawn. A solution of guaiacol, or of *apomorphine* hydrochloride in concentrated sulphuric acid (0.02 gram in 10 c.c.) gives, however, a reagent with which a sharp distinction can always be observed if methyl alcohol is present in the original tincture. The reaction is best carried out by adding a few drops of the distillate to 0.5 c.c. of the reagent in a watch glass. With the guaiacol reagent a clear dark red colour is produced when formaldehyde is present which is readily distinguished from the pale yellow coloration produced in its absence, whilst with the *apomorphine* reagent a characteristic precipitate is formed in presence of formaldehyde. As little as 0.25% of methyl alcohol in tinctures, etc., can be detected with certainty by the new method. G. F. M.

Orcinol Reaction of Furfuraldehyde. ED. JUSTIN-MUELLER (*J. Pharm. Chim.*, 1921, [vii], 24, 334—336).—As little as 1 part of furfuraldehyde in 600,000 parts of solution may be detected by the blue coloration which is obtained when 5 c.c. of the solution, 5 c.c. of hydrochloric acid, and 0.02 gram of orcinol are heated together, cooled, and shaken with a few drops of amyl alcohol.

W. P. S.

Source of Error in Tests for Acetone. E. PITTARELLI (*Policlinico*, 1921, 28, 621; *J. Am. Med. Assoc.*, 76, 1803).—Caoutchouc under the influence of steam or boiling water yields a volatile substance which responds to all the tests most characteristic of acetone; therefore conclusions based on technique involving distillation in the presence of caoutchouc may be misleading.

CHEMICAL ABSTRACTS.

Microchemical Reactions of "Dulcin" [*p*-Ethoxyphenyl-carbamide]. G. DENIGÈS and R. TOURROU (*Compt. rend.*, 1921, 173, 1184—1186).—When a few particles of "dulcin" on a microscope slide are moistened with one drop of nitric acid (*d* 1.39) they dissolve and on the addition of a drop of water microscopic orange or brick-red crystals of *p*-ethoxynitrophenylcarbamide are obtained. These crystals are soluble in chloroform and, on evaporation after the addition of a drop of acetic acid, characteristic crystals are obtained. If dilute nitric acid is used in place of the concentrated acid the "dulcin" does not dissolve, but the mass gradually becomes permeated with crystals of the nitro-compound and under the microscope there is the appearance of a marked effervescence. "Dulcin" is soluble in concentrated sulphuric acid or glacial acetic acid and is deposited as a micro-crystalline precipitate from such solutions on the addition of water or alkali.

W. G.

The Vitali Reaction for Cocaine. PAUL HARDY (*J. Pharm. Chim.*, 1921, [vii], 24, 325—330).—Pure cocaine, and most sub-

stituted cocaines, yield a yellow coloration with the Vitali test in the cold; *isotropylocoaine*, however, yields a violet coloration, and the presence of a small quantity of this substance in certain specimens of cocaine causes these to give a violet coloration with the test. When the reaction mixture is heated, pure cocaine produces at most a slight yellow coloration. W. P. S.

Morphine, Codeine, and Narcotine in Indian Opium. JITENDRA NATH RAKSHIT (*Analyst*, 1921, 46, 481—488).—The morphine content of the bulk of Indian opium is between 8.5 and 10.5%. The B.P. process for estimating morphine is inefficient in India owing to the high temperature which prevails. The U.S.P. process is better and compares favourably with the author's polarimetric process (*Analyst*, 1918, 43, 321). The solubility of codeine and narcotine has been determined in various solvents at temperatures between 20° and 100°. Methods have also been worked out for the estimation of codeine and narcotine, for which the original should be consulted. H. K.

Estimation of Tannin and Colouring Matters in Wines. W. FRESSENIUS and L. GRUNHÜT (*Z. anal. Chem.*, 1921, 60, 406—417).—One hundred c.c. of the wine is evaporated to expel alcohol, cooled, diluted to 100 c.c. and 50 c.c. of this solution are treated with a few grams of pure animal charcoal; after some hours, the mixture (a further quantity of charcoal must be added if all the colour has not been removed) is diluted to 1 litre, filtered, and 400 c.c. are diluted to 1 litre, treated with 10 c.c. of sulphuric acid, (*d* 1.11), and 30 c.c. of indigo solution (3 grams of indigo dissolved in 20 c.c. of sulphuric acid and diluted to 1 litre) and the mixture is titrated with permanganate solution. Twenty c.c. of the de-alcoholised wine not treated with charcoal are titrated in a similar manner. The difference between the two titrations is a measure of the tannin and colouring substances in the wine. The permanganate solution should contain 1.33 grams of potassium permanganate per litre and be standardised against 10 c.c. of *N*/10-oxalic acid solution; the volume of permanganate solution used is divided into 0.0416 to obtain the tannin value of 1 c.c. of permanganate solution. W. P. S.

Determination of the Size of Particles. Attempts to Explain the Formation of Layers in Clay Turbidities and their Use in Soil Analysis. ERNST UNGERER (*Kolloid Chem. Beihefte*, 1921, 14, 63—95).—The layers in clay suspensions are brought about solely by the size and weight of the suspended particles, so that each layer corresponds with particles of a definite size. The turbidity between two layers contains particles uniformly dispersed. Layers are formed both in solutions containing electrolytes and in solutions which do not contain electrolytes, but a high concentration of a coagulating electrolyte will probably prevent the formation of layers. The individual layers either rise or fall with a uniform velocity, and from the rate of falling or rising the size of the particles in a given layer may be calculated by means of Stokes's law. J. F. S.

General and Physical Chemistry.

Refraction of Light at Corresponding Temperatures. W. HERZ (*Z. physikal. Chem.*, 1921, 98, 175—180).—A theoretical paper in which from the examination of the data for a large number of substances it is shown that the refractive index of all substances has the same value 1.126 at the critical temperature. The formulæ of Gladstone and Dale, and of Lorentz and Lorenz yield approximately the same specific refraction values for this temperature. It is also shown that at other comparable temperatures the refractive indices approximate to the same value, although in these cases considerable differences do occur. Thus at the boiling point, ten liquid substances give a mean value of 1.335 for the refractive index, the extreme values being 1.348 and 1.320. J. F. S.

The Spectrochemistry of Aliphatic Dienes with Conjugated Double Bonds. K. VON AUWERS and H. WESTERMANN (*Ber.*, 1921, 54, [B], 2993—2999).—The physical constants of a number of carbinols and the corresponding dienes are recorded. The specific exaltations in the cases of the latter substances with “undisturbed, doubly disturbed, and singly disturbed” conjugation are $E\Sigma_{\text{Retr.}}$ +2.0, +1.4, and +0.9 and $E\Sigma_{\text{Disp.}}$ +50%, +43%, and +35%, respectively. The boiling points of the dienes are depressed by branching of the chain, but raised by the approximation of the methyl group to the double bond; this regularity is not always very definitely marked. The density of isomeric compounds increases relatively greatly when the methyl group is attached to a doubly-bound carbon atom. Similarly, the index of refraction increases with similar alteration in structure; these changes are consonant with those observed with cyclic compounds.

The following constants are recorded: Δ^{β} -pentene- δ -ol, b. p. 64°/62 mm., $d_4^{17.95}$ 0.8382, $n_a^{17.95}$ 1.42558, $n_D^{17.95}$ 1.42821, $n_B^{17.95}$ 1.43502, $n_{\gamma}^{17.95}$ 1.44065. Δ^{β} -Hexen- δ -ol, b. p. 59°/27 mm., $d_4^{23.3}$ 0.8370, $n_a^{23.3}$ 1.43025, $n_D^{23.3}$ 1.43286, $n_B^{23.3}$ 1.43962, $n_{\gamma}^{23.3}$ 1.44510. Δ^{β} -Heptene- δ -ol, b. p. 63°/11 mm., $d_4^{14.4}$ 0.8422, $n_a^{14.4}$ 1.43698, $n_D^{14.4}$ 1.43965, $n_B^{14.4}$ 1.44620, $n_{\gamma}^{14.4}$ 1.45176. ϵ -Methyl- Δ^{β} -hexen- δ -ol, b. p. 75°/45 mm., d_4^{19} 0.8411, n_a^{19} 1.43539, n_D^{19} 1.43788, n_B^{19} 1.44478, n_{γ}^{19} 1.45045. ζ -Methyl- Δ^{β} -hepten- δ -ol, b. p. 67°/11 mm., $d_4^{15.2}$ 0.8354, $n_a^{15.2}$ 1.43679, $n_D^{15.2}$ 1.43926, $n_B^{15.2}$ 1.44585, $n_{\gamma}^{15.2}$ 1.45127. η -Methyl- Δ^{β} -octen- δ -ol, b. p. 85°/12 mm., $d_4^{14.5}$ 0.8402, $n_a^{14.5}$ 1.44202, $n_D^{14.5}$ 1.44448, $n_B^{14.5}$ 1.45114, $n_{\gamma}^{14.5}$ 1.45651. δ -Methyl- Δ^{γ} -heptene- ϵ -ol, b. p. 66°/17 mm., $d_4^{17.9}$ 0.8525, $n_a^{17.9}$ 1.44525, $n_D^{17.9}$ 1.44792, $n_B^{17.9}$ 1.45454, $n_{\gamma}^{17.9}$ 1.46018. δ -Methyl- Δ^{γ} -octen- ϵ -ol, b. p. 89°/16 mm., $d_4^{19.5}$ 0.8495, $n_a^{19.5}$ 1.44576, $n_D^{19.5}$ 1.44838, $n_B^{19.5}$ 1.45490, $n_{\gamma}^{19.5}$ 1.46059.

$\Delta^{\beta\delta}$ -Pentadiene, b. p. 43°, $d_4^{13.6}$ 0.6887, $d_{20}^{13.6}$ 0.685, $n_a^{13.6}$ 1.42991, $n_D^{13.6}$ 1.43443, $n_B^{13.6}$ 1.44655, $n_{\gamma}^{13.6}$ 1.45694, n_D^{30} 1.4309. $\Delta^{\beta\delta}$ -Hexadiene, b. p.

80°, $d_4^{15.05}$ 0.7237, $d_{20}^{15.05}$ 0.720, $n_a^{15.05}$ 1.44962, $n_D^{15.05}$ 1.45420, $n_B^{15.05}$ 1.46651, $n_\gamma^{15.05}$ 1.47764, n_D^{20} 1.4514. $\Delta^{\delta\delta}$ -Heptadiene, b. p. 107°, $d_4^{16.3}$ 0.7341, $d_{20}^{16.3}$ 0.731, $n_a^{16.3}$ 1.45101, $n_D^{16.3}$ 1.45543, $n_B^{16.3}$ 1.46695, $n_\gamma^{16.3}$ 1.47735, n_D^{20} 1.4534. ζ -Methyl- $\Delta^{\delta\delta}$ -heptadiene, b. p. 117°, $d_4^{15.5}$ 0.7361, $d_{20}^{15.5}$ 0.733, $n_a^{11.5}$ 1.44887, $n_D^{15.5}$ 1.45302, $n_B^{15.5}$ 1.46397, $n_\gamma^{15.5}$ 1.47379, n_D^{20} 1.4505. η -Methyl- $\Delta^{\delta\delta}$ -octadiene, b. p. 149°, $d_4^{11.6}$ 0.7515, $d_{20}^{11.6}$ 0.751, $n_a^{11.6}$ 1.45427, $n_D^{11.6}$ 1.45831, $n_B^{11.6}$ 1.46903, $n_\gamma^{11.6}$ 1.47812, n_D^{20} 1.4553. ϵ -Methyl- $\Delta^{\delta\delta}$ -hexadiene, b. p. 104°, $d_{20}^{19.7}$ 0.745, n_D^{20} 1.4606, (i) $d_4^{19.7}$ 0.7439, $n_a^{19.7}$ 1.45659, $n_D^{19.7}$ 1.46117, $n_B^{19.7}$ 1.47340, $n_\gamma^{19.7}$ 1.48433, (ii) $d_4^{19.65}$ 0.7473, $n_a^{19.65}$ 1.45589, $n_D^{19.65}$ 1.46037, $n_B^{19.65}$ 1.47262, $n_\gamma^{19.65}$ 1.48316. β -Methyl- $\Delta^{\alpha\gamma}$ -butadiene, b. p. 34°, $d_4^{18.8}$ 0.6826, $d_{20}^{18.8}$ 0.682, $n_a^{18.8}$ 1.41902, $n_D^{18.8}$ 1.42309, $n_B^{18.8}$ 1.43422, $n_\gamma^{18.8}$ 1.44405, n_D^{20} 1.4224. δ -Methyl- $\Delta^{\gamma\epsilon}$ -heptadiene, b. p. 131°, $d_4^{24.4}$ 0.7598, $d_{20}^{24.4}$ 0.763, $n_a^{24.4}$ 1.45610, $n_D^{24.4}$ 1.46003, $n_B^{24.4}$ 1.47125, $n_\gamma^{24.4}$ 1.48103, n_D^{20} 1.4625. δ -Methyl- $\Delta^{\gamma\epsilon}$ -octadiene, b. p. 150°, $d_4^{20.3}$ 0.7708, $d_{20}^{20.3}$ 0.771, $n_a^{20.3}$ 1.46206, $n_D^{20.3}$ 1.46617, $d_{20}^{20.3}$ 1.47724, $n_D^{20.3}$ 1.48708, n_D^{20} 1.4663. $\beta\gamma$ -Dimethyl- $\Delta^{\alpha\gamma}$ -butadiene, b. p. 70°, $d_4^{22.1}$ 0.7239, $d_{20}^{22.1}$ 0.725, $n_a^{22.1}$ 1.43307, $n_D^{22.1}$ 1.43703, $n_B^{22.1}$ 1.44758, $n_\gamma^{22.1}$ 1.45704, n_D^{20} 1.4382. H. W.

[Substitution Processes]. K. VON AUWERS (*Ber.*, 1921, 54, [B], 3000—3003).—A reply to K. H. Meyer (*A.*, 1921, i, 853). The view that the methyl group causes optical exaltation cannot be maintained. In certain of the examples quoted by Meyer the differences lie within the limits of experimental error, whereas in other examples readily polymerised substances, for instance, acrylic acid and its esters, have been chosen and the data obtained are not due to the unimolecular forms. H. W.

The Molecular Refraction of Aromatic Hydrocarbons and "Aromatic" Carbon. K. VON AUWERS (*Ber.*, 1921, 54, [B], 3188—3194).—In a recent communication (*A.*, 1921, ii, 473), von Steiger has endeavoured to show that the molecular refraction of hydrocarbons can be calculated more accurately from linking refractions than from atomic refractions and he has postulated that all C—C bonds and all C—H bonds in aromatic hydrocarbons are equivalent among themselves but different from the corresponding values in the aliphatic series. This mode of calculation appears to the author to be unsound theoretically, since the atomic refraction of aromatic hydrogen as deduced from the molecular refraction of benzene, diphenyl, and naphthalene, in these circumstances is negative, whilst also the "linking refraction" for (C—H)_{ar} falls with decreasing wave-lengths. Von Steiger has illustrated his arguments by citing the cases of benzene, diphenyl, and naphthalene, and the agreement between the observed and calculated values is excellent. Unfortunately, the data quoted for naphthalene are incorrect, and the amended figures show that the presumption of additivity is not fulfilled. Further, the data for the three hydrocarbons are not directly comparable, since they are obtained under widely-differing conditions; when due allowance is made for this factor, it is found to be impossible to deduce constant atomic or linking refractions. The molecular refractions

of homologues of benzene, and other aromatic hydrocarbons cannot be calculated from von Steiger's aromatic and aliphatic linking refractions, since the differences between the observed and calculated values increase more and more with increase in the number of side-chains.

The fundamental idea in von Steiger's refractometric hypothesis and thermochemical theories (A., 1920, ii, 355) is the existence of a practically tervalent "aromatic" carbon atom of the graphite type which differs from the quadrivalent "diamond" carbon atom of the paraffins (cf. Debye and Scherrer, A., 1917, ii, 437). To the author, this view of the varying valency appears to be misleading. The supposed ter- and quadri-valency of the carbon atom does not, as in the case of other elements, correspond with different stages of oxidation or different electric charges; the atom is invariably quadrivalent and the apparent variation is caused by alteration in the distribution or compensation of its affinity. H. W.

The Mechanism of Continuous Luminous Radiation. J. DUCLAUX (*Compt. rend.*, 1921, **173**, 1355—1357).—The author has previously put forward the hypothesis (cf. *ibid.*, 1914, **158**, 1879) that the emission and absorption of the continuous spectrum depend on the establishment or rupture of chemical valencies. Evidence in support of this hypothesis is now given. W. G.

The Spectrum of Fluorine. WILLIAM R. SMYTHE (*Astrophys. J.*, 1921, **54**, 133—139).—With a discharge passing between gold electrodes, the ultra-violet region of the fluorine spectrum was photographed through a fluorite window, the gas having been prepared by the electrolysis of potassium hydrogen fluoride and purified by being passed through sodium fluoride and a freezing trap. Ten fluorine lines, all in the red, were observed and tabulated with an accuracy of ± 0.1 Å. An impurity which was supposed to have the composition CF_4 was observed following a flare back from the charcoal chamber. The approximate positions of nine heads of bands between λ 4829 and λ 6525 were observed in this spectrum.

CHEMICAL ABSTRACTS.

Arc-cathode Spectra. ARTHUR ST. C. DUNSTAN and BENJAMIN A. WOOTEN (*Astrophys. J.*, 1921, **54**, 65—75).—The spectra of strontium, barium, lithium, copper, and lead were observed when the vapours of the metals were introduced into a direct current arc by means of an alternating current arc placed below it in the same enclosure. It was shown that by cooling the electrodes alternately, the intensity of the metallic lines remained, as before, greater at the cathode than at the anode when the metal was introduced into the arc symmetrically; neither was the effect due to electrolytic or electrostatic separation. Since the phenomenon is fully developed in $1/120$ second, the transference of vapour from one electrode to the other must be of little importance. The light appears to be due chiefly to bombardment of the metallic vapour by electrons from the cathode. Whilst the lines of all

metals are stronger at the cathode, the difference decreases with increasing atomic weight.

CHEMICAL ABSTRACTS.

The Electric Furnace Spectrum of Scandium. ARTHUR S. KING (*Astrophys. J.*, 1921, 54, 28—44).—The spectrum of a pure sample of scandium oxide was observed in the carbon arc and in the electric furnace through the range λ 3015—6559 Å. The furnace was operated at the usual low gas pressure, and temperatures of 2200°, 2250°, and 2600° were used. A table gives the classification of 257 lines with respect to their behaviour in the arc and at various furnace temperatures. Some 25 lines are indicated as being enhanced in the arc; 150 are enhanced in the furnace. The 29 arc-flame lines belong to the latter class, some of them showing moderate strength at all temperatures, whilst others increase in intensity more or less rapidly with increasing temperature. These lines are not due to oxidation, but the bands, on the other hand, are quite probably due to the oxide. Lines which appear at a low temperature and those enhanced in the furnace are weak or lacking in the general solar spectrum but prominent in that of sun-spots. The Zeeman effect for scandium lines is large and apparently uniform in sunspot spectra, although laboratory observations have not as yet been made. Scandium oxide fused in the furnace seemed to form a carbide with the graphite of the boat. The glossy black residue became a greyish-brown powder on exposure to the air, presumably reabsorbing oxygen.

CHEMICAL ABSTRACTS.

Wave-length Measurements in Arc Spectra Photographed in the Yellow, Red, and Infra-red. F. M. WALTERS, jun., (*Bureau of Standards, Sci. Papers*, 1921, 17, No. 411, 161—177).—Observations were made with a grating spectrograph of 640 cm. radius and 299 lines per mm. The photographic plates were sensitised to yellow, red, or infra-red light with pinacyanol and dicyanin. The wave-lengths of silver, aluminium, gold, bismuth, cadmium, mercury, lead, antimony, tin, and zinc are given in international units and to 0.01 Å. The longest wave-length measured is that at 10395 Å. in the spectrum of cadmium. The work is compared with previous results of others and discrepancies are accounted for as arising from impurities, ghosts, or overlapping spectra of higher orders.

CHEMICAL ABSTRACTS.

Extension of the Ultra-violet Spectrum and the Progression with Atomic Number of the Spectra of Light Elements. R. A. MILLIKAN (*Proc. Nat. Acad. Sci.*, 1921, 7, 289—294).—The results of an investigation of the extreme ultra-violet radiations emitted by the second ring or shell of electrons in the atoms of atomic number 2 to 13 (helium to aluminium) are recorded. The ultra-violet spectrum has been photographed down to $\lambda=136.6$ Å. in the case of aluminium and to $\lambda=149.5$ Å. in the case of copper. The L_{α} lines of aluminium, magnesium, and sodium are found at 144.3 Å., 232.2 Å., and 372.2 Å., respectively. Aluminium emits no radiations when excited by condensed sparks in a vacuum between

144.3 Å. and 1200 Å., where the M spectrum due to the three outer electrons begins. The chief lines below 2000 Å. due to the three outer electrons of aluminium are 1379.7, 1384.5, 1605.9, 1612.0, 1671.0, 1854.7, and 1862.7 Å. Magnesium is like aluminium, there is a blank between its L_a line at 232.2 Å. and the lines due to the two outer electrons which begin at 1700 Å. The lines measured below 2000 Å. are 1735.2, 1737.9, 1751.0, and 1753.7 Å. Sodium emits no lines between the L line 376.5 Å. and the lines due to the single M electron which have their convergence wave-length at 2412.63 Å. The spectrum due to the six L ring electrons of oxygen begins at 230 Å. and extends with much complexity and strength up to 834.0 Å., where the strongest line is found; above this point the lines are few in number and relatively faint. The ratio of the K/L frequency for oxygen is about 35. The strongest oxygen lines are 321.2, 374.3, 507.8, 525.7, 554.2, 599.5, 610.1, 616.7, 625.2, 629.6, 644.0, 703.1, 718.5, and 834.0 Å. The spectrum of the four L -ring electrons of carbon begins at 360.5 Å. and extends with much complexity and strength up to 1335.0 Å., where the strongest line lies; above this point the lines are widely scattered and relatively weak. The ratio of the K/L frequency in carbon is about 30. The spectrum due to the five L -ring electrons of nitrogen is simple; it begins at 685.6 Å. and reaches a maximum at 1085.3 Å. The only other strong lines of nitrogen are 685.6 Å., 916.2 Å., and 991.1 Å. The ratio of the K/L frequency is 34.8. In the case of fluorine, a strong line appears at 657.2 Å. and a second at 607.2 Å.; these are the only lines thus far identified as coming from the seven L -ring electrons of fluorine. The spectrum from the three L -ring electrons of boron is very simple. It consists of 676.8, 760.0, 1624.4, two doublets, 2164.2, 2166.2, and 2496.9, 2497.8, and a single spark line, 3451.5 Å. The ratio of the frequency K/L is 37. The spectrum of the two L -ring electrons of glucinum begins at 2175 Å. and finishes in its L_a doublet at 3130.6 and 3131.2 Å. Similar experiments with lithium reveal no lines between the shortest wave-lengths measurable by the present method and the familiar series due to its single L -ring electron the L_a line of which is at 6708 Å., and its convergence wave-length at 2299 Å. The progression shown in these optical spectra is simple and very like that exhibited by X-ray spectra.

J. F. S.

The Structure of the Elements of Mean Atomic Number.

A. DAUVILLIER (*Compt. rend.*, 1921, **173**, 1458—1461).—An extension of previous work (cf. A., 1921, ii, 669), further results being given for gold and new measurements of the L series for cerium and antimony.

W. G.

The Structure of the Bismuth Lines. H. NAGAOKA and Y. SUGIURA (*Astrophys. J.*, 1921, **53**, 339—348).—The light resulting from the bombarding of bismuth by an electronic current was observed by means of a glass Lummer-Gehrcke plate crossed with one of quartz or crossed with an echelon grating. For the line λ 4722 the five known positive components were confirmed and

seven weak negative satellites were discovered, showing the line to have the general structure common to lines of heavy metals. The principal component is probably multiple, its constituents varying in relative intensity. The line λ 4122 has four strong components. The line λ 4308 consists of two principal lines separated by 0.350 Å., each being a narrow doublet; there are also two faint satellites. The authors point out some constant frequency differences and suggest that these may be interpreted in terms of the quantum theory. CHEMICAL ABSTRACTS.

The Zeeman Effect. H. R. WOLTJER (*Chem. Weekblad*, 1921, 18, 677—682).—A paper written in connexion with the twenty-fifth anniversary (October 31st, 1921) of the first announcement of Zeeman's discovery. The importance and subsequent developments of his work are discussed at some length. S. I. L.

Anisotropy of Molecules. C. V. RAMAN (*Nature*, 1922, 109, 75—76).—An extension to direct visual observations is described of the method whereby it has been shown photographically (Lord Rayleigh, A., 1920, ii, 574) that the light scattered by molecules is, in general, not completely polarised when observed in a direction transverse to the pencil of light traversing the gas. With carbon dioxide, the effect is conspicuous, and visual determinations of its magnitude have been made. Similar evidence that the molecules of gases are not spherically symmetrical and are anisotropic in their properties is furnished by observations on the polarisation of light of the sky; liquids also show an imperfect polarisation attributable to anisotropy. A. A. E.

Sols with Non-spherical Particles. H. ZOCHER (*Z. physikal. Chem.*, 1921, 98, 293—337).—The double refraction discovered by Diesselhorst, Freundlich, and Leonhardt (A., 1916, ii, 65) with flowing vanadium pentoxide sols has been found in the following sols: Soap solution, clay suspensions, sols of silver cyanate, benzo-purpurin, "benzo-brown," primulin, sodium alizarinsulphonate, alizarin, *p*-azoxyphenetole, *p*-azoxyanisole, anthracene, cerasin-orange, and aniline-blue. Some observations on the spacial partition of the double refraction in agitated sols showed that in the eddies phenomena were observed which exhibited many analogies to the behaviour of uniaxial crystals in convergent light. These observations serve as methods for determining the flowing-double refraction and pleochroism and for determining the character of the sols. The rotation of the dark cross in eddies in opposition to the polarisation direction shows the direction of flow does not always need to correspond with the main vibration direction of the light. The coincidence of the dark cross with the direction of polarisation in old vanadium pentoxide sols and soap solutions is attributed to the bending of the doubly refracting particles. The sign of the double refraction of soap solutions varies with the age and concentration of the sol. The change in the sign of the double refraction is explained as follows: Colloidal particles in the form of short rods must arrange themselves with their symmetry axis

in the direction of flow, particles in the form of leaflets must arrange themselves with the axis of symmetry at right angles to this direction. If now, in the process of ageing, the rods pass into leaflets, then the optical character of the direction of flow must at first be the same as that of the colloidal particles and afterwards of the opposite sign. Aniline-blue sols, prepared by pouring an alcoholic solution of the dye into water, show on flowing a strong negative double refraction in the red, negative pleochroism in the region orange to green, and a positive double refraction in the blue. Anomalous interference bands are produced by this anomalous behaviour. The optical anisotropy produced by a magnetic field has the opposite sign, that is, the colloidal particles arrange themselves at right angles to the lines of force. On the other hand, they arrange themselves parallel to the electrical lines of force. The negative double refraction of benzopurpurin can be made to disappear by heating. The addition of electrolytes causes the double refraction to reappear on cooling. The coagulum obtained by the addition of an excess of electrolyte gives a doubly refracting sol on peptisation if the addition of the electrolyte was slow or if the original sol was doubly refracting, but if the coagulation took place rapidly or if the original sol was isotropic, there will be no double refraction. When acidified, double refracting red sols may be changed into double refracting blue sols. The strong pleochroism has always a negative sign, and the particles of the sols are diamagnetic. The double refraction of vanadium pentoxide sols disappears more slowly the older and more concentrated the sol. Very concentrated old vanadium pentoxide sols and concentrated, electrolyte-poor benzopurpurin solutions show a persistent double refraction similar to that of the crystalline liquids. The view of Diesselhorst and Freundlich (*loc. cit.*), that the growth of non-spherical particles is not a crystallisation but an aggregation of non-spherical primary particles in parallel layers, is confirmed, with the addition that the primary particles may be crystalline and that eventually they may undergo crystalline processes.

J. F. S.

Recent Advances in Stereochemistry. B. K. SINGH (*J. Proc. Asiatic Soc. Bengal*, 1921, **17**, 213—230).—A general account of the historical development of stereochemistry and a discussion of the relation between optical activity and chemical constitution. Particular consideration is given to optical activity in homologous series, the effect of conjugated unsaturation, and of position isomerism on optical activity. The Walden inversion is also discussed.

H. W.

Investigations on the Fundamental Law of Photochemistry. P. LASAREV (*Z. physikal. Chem.*, 1921, **98**, 94—97).—A theoretical discussion of work previously published on the bleaching of dyes by light (A., 1912, ii, 219; *Ann. Physik*, 1907, [iv], **24**, 661). The experiments on the bleaching in the presence of oxygen under increased pressures (up to 150 atm.) are particularly considered. The fundamental law for the bleaching of dyes has the form

$-dC/dt = \alpha_0(1 - e^{-K_1 C})(1 - e^{-K_2 C})$, in which C is the concentration of the dye, K a constant proportional to the absorption constant, C_1 and K_1 similar values for the substance non-sensitive to light, and α the photochemical constant. It is assumed that in the reaction, during the absorption of light, an electron from the first Bohr orbit springs over to the furthest orbit. It is shown that the total absorption of light is proportional to the absorption of the largest (ionised) atoms. The increase in volume during photochemical reactions is explained by the assumption that the light, inasmuch as it occasions an increase in the volume of the atoms by effecting the electron spring, brings about an increase in the gaseous volume.

J. F. S.

New Measurements of Precision in the X-Ray Spectrum. M. SIEGBAHN (*Compt. rend.*, 1921, 173, 1350—1352).—By the use of three spectrographs, each adapted to a particular spectral region, it was possible to measure a wave-length with an accuracy of 0.01—0.005%, and by this means the wave-length of the copper line $K\alpha_1$ was found to be 1537.36×10^{-11} cm. in a vacuum. A more precise instrument is described the readings of which are accurate to within 0.002%, and with this the value of λ for $K\alpha_1$ was found to be 1537.302×10^{-11} cm.

W. G.

Spectrographic Study of the De-intensifying of Barium Platinocyanide in the Villard Effect. A. ZIMMERN and E. SALLES (*Compt. rend.*, 1922, 174, 80).—Fluorescent screens which have undergone the Villard effect can be regenerated by exposure to diffused light. It is shown that the radiations capable of destroying the Villard effect are found in four principal bands of almost equal breadth. The first is entirely in the infra-red, the second in the greenish-yellow, the third in the blue, and the fourth in the ultra-violet.

W. G.

The Theory of Absorption of X-Rays by Matter and the Principle of Correspondence. LOUIS DE BROGLIE (*Compt. rend.*, 1921, 173, 1456—1458).—An expression for the atomic coefficient of absorption of a substance for a radiation of wave-length λ has previously been given (cf. A., 1920, ii, 208) in which the constant, α , was defined by the hypothesis that, for the possible changes of internal configuration of an atom placed in a system in thermodynamic equilibrium at the temperature T , the probability A_{12} of the return of the atom from a configuration of energy ϵ_1 to a configuration of less energy, ϵ_2 , is proportional to $\epsilon_1 - \epsilon_2$ and also to the absolute temperature in such a way that $A_{12} = \alpha(\epsilon_1 - \epsilon_2)T$. A mathematical proof of the truth of this hypothesis and a means of calculating the constant α are given, and it is shown that the coefficient of the Bragg-Pierce law is a universal constant capable of being expressed as a function of the constants of the electron and of the radiation.

W. G.

Calculation of the X-Ray Absorption Frequencies of the Chemical Elements. I and II. WILLIAM DUANE (*Proc. Nat. Acad. Sci.*, 1921, 7, 260—267; 267—273).—In the first paper,

the author has calculated the K critical absorption frequencies of the elements magnesium, sulphur, calcium, iron, selenium, molybdenum, tin, cerium, dysprosium, tungsten, lead, and uranium on the basis of the Rutherford-Bohr theory of the structure of atoms and the mechanism of radiation. The assumption is made that the electrons are distributed in circular orbits, which do not lie in planes passing through the nucleus of the atom. In order to estimate the forces exerted on an electron in one orbit, A , due to the electrons in the parallel orbit, B , it is assumed that they are the same as if the charges are concentrated, half at the nearest point to the orbit A and half at the point furthest from A . A series of values is obtained which are in fair agreement with those observed. In the second paper, it is assumed that the forces are the same as if the electricity of the electrons in the orbit B is uniformly distributed along the orbit. The values calculated on the second assumption are somewhat smaller than those of the first calculations.

J. F. S.

The Laws of Absorption of X-rays. F. K. RICHTMYER (*Physical Rev.*, 1921, 18, 13—30).—An experimental study of the absorption of X-rays by water (oxygen); aluminium, copper, molybdenum, silver, and lead, each over a great part of the range of wave-lengths 0.093 Å. to 0.95 Å. The longest of these waves is shorter than the critical K absorption wave-length of oxygen, aluminium, or copper, and falls between K and L for molybdenum, silver, and lead. Rays from a Coolidge tube were analysed by a spectrometer with a sodium chloride crystal, and the intensity was measured by ionisation of methyl bromide. Scattered and fluorescent radiation from the absorbing screen were suppressed by placing the screen in front of the spectrometer slit nearest the tube. Detailed tables and curves of the results are given. The mass-absorption coefficients, μ/ρ , of these substances is given by $\mu/\rho = F\lambda^3 + (\sigma/\rho)$ where λ is wave-length and σ/ρ is the mass-scattering coefficient, except in the region close to λ_K on the short-wave-length side, where the equation gives values which are somewhat too large, the error being greater as the difference between λ and λ_K is smaller. For each substance F has a constant value for $\lambda < \lambda_K$ and another much smaller constant value for $\lambda > \lambda_K$. The mass-scattering coefficient is independent of wave-length, and is, to a first approximation, the same for all elements. The atomic-absorption coefficient is given by $\mu_a = 2.29 \times 10^{-27} N^4 \lambda^3 + \sigma_a$, where N is the atomic number and σ_a the atomic-scattering coefficient, for $\lambda < \lambda_K$. No trace of the suggested "J" absorption discontinuity was found. At 0.093 Å., μ/ρ for water is still much greater than for the hard rays of radium-C.

CHEMICAL ABSTRACTS.

The Luminescence of certain Oxides Sublimed in the Electric Arc. E. L. NICHOLS and D. T. WILBER (*Physical Rev.*, 1921, 17, 707—717).—Thin films of metallic oxides were obtained by subliming the metal or one of its salts from the crater of a direct-current carbon arc on to a metal disk. Oxides prepared by sublimation are found to respond to excitation by cathode rays.

The shift in colour with increasing temperature in luminescence is in general to the violet. The effect of pressure on luminescence is discussed. The most active oxides were those of calcium, magnesium, zinc, zirconium, silicon, and aluminium.

CHEMICAL ABSTRACTS.

Anode Rays of Glucinum. G. P. THOMSON (*Nature*, 1921, 107, 395).—The method of positive ray analysis applied to lithium (Aston, A., 1920, ii, 344) has been extended to glucinum and found to yield a well-marked parabola corresponding with a single charge and an atomic weight 9.0 ± 0.1 ($\text{Na} = 23$). No second parabola was observed which could be ascribed with certainty to glucinum, but it is doubtful if a line of intensity one-tenth that of the first could have been detected. No indication was found which would suggest that the atom of glucinum can lose two electrons under the prevailing experimental conditions. A. A. E.

Reactions in Penetrating Radium Radiation and in Ultra-violet Radiation Filtered by Quartz Glass. II. The Hydrogen Peroxide Equilibrium set up in Radium Radiation. ANTON KAILAN (*Z. physikal. Chem.*, 1921, 98, 474—497; cf. A., 1912, ii, 10, 522; 1920, ii, 576).—Radiation from 0.1 gram of radium metal after passing through 1 mm. of glass into 100 c.c. of water in contact with air produces a solution containing 6×10^{-6} gram-equivalent of hydrogen peroxide; in the presence of 0.01*N*-sulphuric acid the amount of hydrogen peroxide becomes 10^{-4} , and in 1.0*N*-acid the amount is 5×10^{-4} gram-equivalent. In the last-named solution 6×10^{12} and in the first 3×10^{12} molecules of hydrogen peroxide are formed per second. These values are considerably less than those obtained nine or ten years ago, and reasons for the discrepancy are advanced. A comparison of the values obtained for the velocity of formation of hydrogen peroxide then and now with the rate of formation by other reactions which have been obtained in a similar manner to the present shows that both in aqueous and in non-aqueous solutions, when the decomposition of hydrogen peroxide is neglected, the number of molecules of hydrogen peroxide formed is of the same order as that of the ion pairs absorbed from the radiation by the medium. In ultra-violet light from a quartz glass lamp under similar conditions, the reduction of the peroxide concentration by increase in the hydrogen-ion concentration is less than in the radium rays, whilst the decomposition of hydrogen peroxide is more than one hundred times greater than in the radium rays. J. F. S.

Stationary Electron Vibrations without Radiation Resistance. A. D. FOKKER (*Physica*, 1921, 1, 107—109).—It is theoretically shown that a Bohr atom with electrons revolving in closed paths may be free from radiation resistance without violating the principles of the classical theory. CHEMICAL ABSTRACTS.

The Radioactivity of the Oxides of Uranium. CHARLES STAEBLING (*Compt. rend.*, 1921, 173, 1468—1471; cf. A., 1920, ii, 5).—In continuation of previous work (*loc. cit.*), it is shown that

the radioactivity of the green oxide, which has been restored to its original value, commences to diminish as before. The restoration of activity may be brought about by simple calcination without passage through ammonium uranate. During this process of restoration, the oxide undergoes loss in weight, due, for the most part, to the removal of moisture. The black oxide which shows little loss in activity also shows little loss in weight or alteration in activity on calcination. The author considers that the green oxide exposed to the air in thin layers undergoes hydration as well as loss in activity, the hydration diminishing the superficial density of the uranium atoms. The black oxides obtained by calcination at high temperatures do not hydrate and show little or no decrease in radioactivity. These facts tend to confirm the existence of a definite black oxide, which is probably an allotropic modification of the green oxide.

W. G.

Isotopy of the Radio-elements and Meitner's Nucleus Model. MAXIMILIAN CAMILLO NEUBURGER (*Z. physikal. Chem.*, 1921, 99, 161—167).—A theoretical paper in which, on the basis of Meitner's nuclear hypothesis (A., 1921, ii, 293), the author has subdivided isotopes into four groups. These are termed isotopes of the first, second, third, and fourth order. Isotopes of the first order have the same nuclear charge and arrangement of the outside electrons, but different nuclear mass, total number of nuclear constituents, arrangement of the nuclear constituents, number of each kind of nuclear constituent, and probability of disintegration; radium and meso-thorium-I illustrate this group. Isotopes of the second order have the same nuclear charge, arrangement of the outside electrons, nuclear mass, and total number of nuclear constituents, but different numbers of each kind of nuclear constituent, arrangement of the nuclear constituents, and probability of disintegration. An example of this class is furnished by ionium and uranium-Y. Isotopes of the third order differ only in the arrangement of the nuclear constituents and the probability of disintegration, all other properties being identical; this is the case with radium-D and actinium-B. Isotopes of the fourth order are identical in all the properties mentioned; no examples of this class are known.

J. F. S.

Calculation of the Branching Relationships for Dual α -Disintegration and the Meitner Nucleus Model. MAXIMILIAN CAMILLO NEUBURGER (*Z. physikal. Chem.*, 1921, 99, 168—171; cf. Meitner, A., 1921, ii, 293).—A theoretical paper in which it is shown that the method of calculation adopted by Smekal (A., 1921, ii, 149) for determining the branching relationship is so closely dependent on the assumptions made as to the structure of the radioactive nucleus, that it cannot possibly be of general use. The branching relationship cannot be calculated by this method, since it leads to inaccurate values which lead to results opposed to the experimental data. The agreement found for the branching relationship by this method for the uranium, radium, thorium,

and actinium families with the experimentally determined values for this relationship is accidental. J. F. S.

Emissivity of Iron and Copper. K. LUBOWSKY (*Elektro-techn. Z.*, 1921, 42, 79—81).—The emissivity of both bright and blackened iron and copper was determined. Curves are given showing the effect of convection currents obtained from a flat plate of the material, first perpendicular to, and second in the direction of the air stream. Expressed in watts per sq. cm. per 1° of temperature excess, the emissivity varies from 0.75×10^{-3} to 1.5×10^{-3} under the various conditions. CHEMICAL ABSTRACTS.

The Effects of Impurities on the Ionisation Potentials Measured in Thermionic Valves. L. S. PALMER (*Radio Rev.*, 1921, 2, 113—125).—The method of investigation used is based on a study of the current-voltage curves measured in hard and soft 3-electrode tubes. The details of the method have been given in a previous paper. Certain bends in these curves are interpreted as being due to the bombardment of the filament by positive ions. The ionisation potentials of helium, mercury, and argon have thus been determined. There are variations in the values for the ionisation potentials obtained in helium. These are ascribed to impurities, and the amount of impurity active accords with the values found by Horton in helium (A., 1919, ii, 210; 1920, ii, 660; 1921, ii, 672). This effect of the impurity also explains the low values of the potentials obtained by Stead and Gossling (A., 1920, ii, 659). The characteristic of the soft 3-electrode tube can be used to determine the resonance and ionising potentials of the contained gas. The nature of the gas and presence of the impurities can be determined from the critical points of the characteristics. Methods of measuring ionisation potentials are liable to error if impurities are present. Horton's previous work is confirmed and the result has been extended to argon with traces of mercury. The value of the ionisation potential decreases slightly with increase of gas pressure, causing changes in the characteristics similar to those for impurities. A number of critical potentials obtained for helium and argon are given. CHEMICAL ABSTRACTS.

Estimation of the Radium Content of Radioactive Luminous Compounds. E. A. OWEN and WINIFRED E. PAGE (*Proc. Physical Soc.*, 1921, 34, 27—32).—The absorption of γ -radiation in zinc sulphide has been measured by different methods, and its value found to be approximately the same by all. No indication was observed of the occurrence of an abnormal absorption when the radioactive salt was mixed with the sulphide. A table has been drawn up which gives the values of the ratio of the true to the apparent radium content for tubes of different diameters filled with a radioactive luminous compound. This table applies to a compound the apparent density of which is 2.03 grams per c.c. The value of the absorption of γ -radiation from radium has been measured for zinc sulphide (0.101), barium sulphate (0.079), barium chloride (0.100), barium carbonate (0.046), lead monoxide (0.294),

bismuth sesquioxide (0.206), and uranium pentoxide (0.315). The absorption coefficients are given in brackets in cm^{-1} . The mass absorption coefficient has been calculated in each case, and it is shown that the barium salts have nearly the same value, the mean value being 0.053. J. F. S.

Electrical Conductivity of Salts and Mixtures of Salts. A. BENRATH and K. DREKOFF (*Z. physikal. Chem.*, 1921, 99, 57—70).—The specific conductivity of molten mixtures of potassium sulphate with sodium sulphate, magnesium sulphate, potassium fluoride, and lithium sulphate respectively has been determined over the whole range of compositions and from the results relationships to the diagram of condition are drawn. It is shown that the conductivity isotherm can be deduced only to a small extent from the diagram of condition. It is also shown that the point at which the eutectic appears is a well defined conductivity point. Pure salts as well as salt compounds change their conductivity with temperature according to the formula $\log k = a + bT$. This formula is not generally applicable to mixtures of salts and mixed crystals; attempts to obtain a theoretical basis for the formula gave the relationship: $\log k'' + bT'' = \text{const.}$ The formula of Tubandt, $\log K = c - d/T$, is not generally applicable. For the complete explanation of the phenomena, other factors, particularly the viscosity, must be known and taken account of in the calculation. J. F. S.

The Electrical Conductivity of Anhydrides of the Higher Fatty Acids. D. HOLDE and IDA TACKE (*Chem. Ztg.*, 1921, 45, 1246—1247; cf. A., 1921, i, 842).—The values previously given for the molecular conductivity of oleic anhydride and oleic acid in acetone, on being adjusted to a dilution of 50 and 8 respectively gave 2.015×10^{-4} and 0.53×10^{-4} . If it is assumed that these two compounds are only slightly dissociated, that is, that the molecular conductivity increases proportionally to the square root of the dilution, the calculated figure for λ_{50} for oleic acid becomes 1.33×10^{-4} , a figure not differing very greatly from that of the anhydride; the difference may be due to impurities in the acetone used. A. R. P.

Measurement of Dielectric Constants. J. F. KING and W. A. PATRICK (*J. Amer. Chem. Soc.*, 1921, 43, 1835—1843).—A bridge method for the measurement of dielectric constants is described in which an arrangement consisting of an electron tube, a condenser, and an induction coil connected in a circuit serve as the source of alternating current of symmetrical wave form. Preliminary measurements of the dielectric constant of mixtures of ethyl alcohol with benzene, ethyl ether, and carbon tetrachloride respectively up to 100% by weight of alcohol, are recorded. The dielectric constant increases with each addition of alcohol to the other constituent in a fairly regular manner. J. F. S.

Electrical Moments of Carbon Monoxide and Dioxide Molecules. HANS WEIGT (*Physikal. Z.*, 1921, 22, 643).—The

dielectric constants of carbon monoxide and carbon dioxide and their dependence on temperature have been determined with the object of ascertaining by means of the Debye theory the electrical moment of the two gases. The values obtained are $\mu_{\text{CO}_2} = (0.1420 \pm 0.0017) \times 10^{-18}$, $\mu_{\text{CO}} = (0.1180 \pm 0.0016) \times 10^{-18}$, both of which are in good agreement with that calculated from the Lorentz-Lorenz formula. J. F. S.

Application of the Theory of Allotropy to Electromotive Equilibria. IV. A. SMITS (*Z. physikal. Chem.*, 1921, **98**, 455—459; cf. A., 1915, ii, 217; 1916, ii, 77; 1917, ii, 232).—The author offers a corrected interpretation of that published ("Theorie der Allotropie," Leipzig, 1921, 403—405) for the electromotive behaviour of metals as soon as they are attacked by acids. The *E.M.F.* of *M*/2-zinc sulphate dissolved in water, 0.05*N*-sulphuric acid, 0.1*N*-, 0.5*N*- and 1.0*N*-sulphuric acid, and *N*-potassium sulphate has been determined at 18° in an atmosphere of nitrogen. The results show that the potential of zinc is not effected by the presence of acid below 0.5*N*, and from this point upwards the potential is only changed by 2 millivolts. Similar experiments are described with zinc chloride in hydrochloric acid solutions; here it is shown that a concentration of hydrochloric acid of 0.1*N* is without influence on the potential, but that 0.5*N* and 1.0*N* cause a change in the *E.M.F.* of 0.021 and 0.038 volt, respectively. J. F. S.

Electromotive Behaviour of Aluminium. A. GÜNTHER-SCHULZE (*Z. Elektrochem.*, 1921, **27**, 579—582).—A reply to Smits, in which the author maintains his previous assertions (cf. this vol., ii, 20). J. F. S.

Electromotive Behaviour of Metallic Compounds with Electron Conductivity. G. TRUMPLER (*Z. physikal. Chem.*, 1921, **99**, 9—56).—Metallic or mixed conducting binary compounds combined with the metallic component as conductor and immersed in a solution of a salt of the metal which is saturated with the compound, have a zero potential when measured against the metallic component in the same solution. The formula put forward previously by Haber for such cases has been confirmed in the case of cuprous iodide, and cuprous, lead, and silver sulphides. It is also shown that metallic and mixed conducting compounds combined with the metallic component as conductor possess the same potential as an indifferent electrode (platinum) when immersed in a solution saturated with these substances and containing the negative component both as ion and in the free condition. A necessary condition in this case is the stability of the compound towards the metallic conductor and the solution. This has been investigated in the case of lead, cupric, cuprous, silver, and ferrous sulphide against sodium sulphide and polysulphide solutions. If in the last case pure electrolytes are used, the free negative component has no influence on the potential against the solution; this is much more dependent on the concentration of the components in the solution. This has been shown in the cells $\text{Ag}|\text{AgBr}|\text{KBr}, \text{Br}_2(\text{sat.})$; $\text{Ag}|\text{AgI}|\text{KI}, \text{I}_2(\text{sat.})$. In the case of the

mixed conducting silver sulphide in the presence of free sulphur the potential lies between that of the pure components. This potential is a characteristic of the mixed conductor. The three forms of conductors, metallic, mixed, and electrolytic, are characterised by the potential relationship obtained by the above-named measurements. From the different behaviour of the metallic and electrolytic conductors (compounds) with respect to the influence of the negative component on the potential relations as obtained by the above method, a fundamental difference of the internal structure is deduced, according to which in the pure metallic conducting compounds the space lattice points are occupied by atoms or molecules and not by ions. They appear, in opposition to electrolytes, to be non-polar. It is shown that pure cuprous sulphide is an electrolyte, or a mixed conductor in which the metallic component of the conductivity is very small. The passivity of lead sulphide in solutions of lead salt has been investigated and a process for activating the substance is put forward. The data in the literature on the solubility of lead sulphide and cupric sulphide are corrected. The electro-metric determination of the solubility of lead sulphide is not possible on account of the complete passivity of the lead electrode in sulphide solutions. The solubility product of lead sulphide has been determined approximately from the solubility in hydrochloric acid and the value $[Pb^{++}] \cdot [S^{--}] = 5 \times 10^{-29}$ obtained. It is shown that cupric sulphide is not stable in sodium sulphide and that in the presence of copper only cuprous sulphide can exist in sodium sulphide solution, and consequently the measurement of the copper potential in sodium sulphide solution does not give the solubility product of cupric sulphide (cf. Knox, A., 1908, ii, 830). The solubility product of cuprous sulphide is found to be $[Cu^+]^2 \cdot [S^{--}] = 2 \times 10^{-47}$. J. F. S.

The "Salt Error" of the Quinhydrone Electrode. S. P. L. SØRENSEN, M. SØRENSEN, and K. LINDERSTRØM-LANG (*Compt. rend. Trav. Lab. Carlsberg*, 1921, **14**, No. 14, pp. 31).—Biilmann's quinhydrone electrode (A., 1921, ii, 372) gives, in the presence of sodium chloride, too small a potential (with 3.99*N*-sodium chloride + 0.01*N*-hydrochloric acid 0.6931 volt instead of 0.7044 volt at 18°). This is due to a lowering of the solubility of quinhydrone and can be overcome by saturating the salt solution with quinhydrone and with one of its constituents, for example, with quinol. This and various other questions are discussed theoretically in detail, and the theory is verified by experiments. G. B.

The Quinhydrone Electrode. EINAR BIILMANN and HAKON LUND (*Ann. Chim.*, 1921, [ix], **16**, 321–340).—It has previously been shown (A., 1921, ii, 372) that aqueous solutions of quinhydrone may be used for the preparation of reversible electrodes with very constant potential, which may serve to determine hydrogen-ion concentrations in solutions which are one-tenth molar. Concordant results are not obtained with stronger solutions. It is possible to construct electrodes, however, in which the electromotive

reaction is a transformation of one solid body into another solid body. The preparation of such electrodes with benzoquinone-quinhydrone and with quinol-quinhydrone is described. Such electrodes have hydrogenation potentials $\pi_{18}=0.7562$ and 0.6179 , respectively. They may be used for measurements over a wide range of hydrogen-ion concentration. W. G.

Electromotive Force produced by the Relative Displacement of an Electrode and an Electrolyte. STÉFAN PROCOPIU (*J. Chim. Physique*, 1921, **19**, 121—134).—It is found that the movement of an electrode in a liquid gives rise to an *E.M.F.* of movement which is general for all metallic electrodes. The dimensions of this *E.M.F.* are determined only by the solution pressure of the metal and the osmotic pressure of the metallic ion in the solution. The effects produced when a series of metals is moved in water, nitric acid, sulphuric acid, salts of the metals, and potassium hydroxide have been measured. The phenomenon is explained by the existence of a layer of solution round the electrode of composition different from that of the rest of the solution. This leads to a relationship between the photoelectric *E.M.F.* and that of movement, and makes it possible to find the effect of movement on the electrical resistance of liquids. J. F. S.

The Overvoltage of the Mercury Cathode. EDGAR NEWBERY (*T.*, 1922, **121**, 7—17).

Theory of the Electrolytic Ionic Condition and the Calculation of the Electrolytic Solution Constants, and of the Related Quantities from the Chemical Relationships. KARL FREDENHAGEN (*Z. physikal. Chem.*, 1921, **98**, 38—69).—A theoretical paper in which it is shown that neither Arrhenius's electrolytic dissociation theory nor Nernst's theory of solution tension furnishes any information on the nature of the ionic condition or the nature of ionic dissociation. Neither of these theories can be directly deduced from the facts of inorganic chemistry, and they do not bring these facts into relationship with one another. The author has put forward an hypothesis of the ionic condition and the electrolytic solution constants which removes the above-named defects and shows how ordinary thermal dissociation passes into electrolytic dissociation. The hypothesis also shows how the solution constants of the elements, and the partition and dissociation constants of chemical compounds may be calculated from the chemical relationships of the compounds and elements and from a factor which expresses the condition of the solvent. The hypothesis permits a qualitative deduction of the electrical and dielectrical behaviour of conductors of the second class.

J. F. S.

An Electrolytic Current Intensification Effect, a New Electrolytic Displacement Effect, and the Connexion between Electrolysis and the Emission of Electrons in a Vacuum. II. D. REICHINSTEIN and F. KLEMENT (*Z. physikal. Chem.*, 1921, **99**, 275—289; cf. *A.*, 1921, ii, 729).—A continuation of work previously

published (*loc. cit.*). It is shown that in a closed circuit by means of superimposed alternating current a stronger direct current can be produced, the work from which can be greater than that of the alternating current used to generate it. The action of a high frequency current on an electrolytic cell is not a specific high frequency action; qualitatively, it is the same as that of a low frequency current. All passivity phenomena, both anodic and cathodic, show on superposing an alternating current on the direct current a decrease of the direct current polarisation, and thereby allow themselves to be used as detectors or current intensifiers. Using the commutator method, an intensification of two hundred times has been obtained and the experiments indicate that this may be still further increased. A new experiment with a highly evacuated electron tube is described, which indicates that after removal of the polarisation it is possible to generate cathode rays with low potentials and cold electrodes. J. F. S.

Applicability of the Gas Laws to Strong Electrolytes. II. J. N. BRØNSTED (*K. Danske Videnskab. Selskab., Math.-fys. Medd.*, 1920, 3, 1—21; cf. A., 1920, ii, 78).—By a modification of the original electrometric method, consisting in the use of a streaming electrode, *E.M.F.* determinations have been carried out with one electrolyte dissolved in a concentrated solution of another. Cells of the type: $\text{Ag, AgCl} \left| \begin{array}{c} \text{solvent } (n-c_1) \\ \text{chloride } (c_1) \end{array} \right| \begin{array}{c} \text{solvent } (n-c_2) \\ \text{chloride } (c_2) \end{array} \left| \text{AgCl, Ag and} \right.$
 $\text{Ag} \left| \begin{array}{c} \text{solvent } (n-c_1) \\ \text{silver salt } (c_1) \end{array} \right| \begin{array}{c} \text{solvent } (n-c_2) \\ \text{silver salt } (c_2) \end{array} \left| \text{Ag} \right.$ were employed. Support is afforded to the theory that the simple gas laws are valid for ions present in salt solutions the concentrations of which are large in comparison with that of the ions concerned. Conditions under which the law fails to apply are discussed, and the results of Loomis, Essex, and Meacham (A., 1917, ii, 353), that slight changes in the activity coefficient occur with varying concentrations, are confirmed. CHEMICAL ABSTRACTS.

Newer Investigations on the Anomaly of the Strong Electrolytes. LUDWIG EBERT (*Jahrb. Radioaktiv. Elektronik*, 1921, 18, 134—196).—An alphabetical bibliography of the work on strong electrolytes is given. The whole subject of the behaviour of strong electrolytes is discussed under the headings (i) chief anomalies, and (ii) the hypothesis of complete dissociation of the strong electrolytes and the activity of electrical forces between the ions. J. F. S.

Transport Numbers of Sulphuric Acid by the Concentration Cell Method. ALFRED L. FERGUSON and WESLEY G. FRANCE (*J. Amer. Chem. Soc.*, 1921, 43, 2150—2160).—A method is described for the determination of the transport numbers of a uni-bivalent electrolyte by measurement of the potentials of concentration cells. The transport number of the anion of sulphuric acid for concentrations between 0.1*M* and 0.01*M* has been measured and found to be 0.1868 ± 0.0007 at 25°. This value is compared,

with the values obtained by other investigators. It is shown that the dissociation values determined from freezing-point data are more satisfactory for calculating the potentials of concentration cells than those obtained from conductivity data. A correction to the formula for the potential of a concentration cell has been developed which takes into account the undissociated part of the acid. It is shown that the concentration cell method is entirely satisfactory for the determination of the transport numbers of sulphuric acid.

J. F. S.

Influence of Gelatin on the Transport Numbers of Sulphuric Acid. ALFRED L. FERGUSON and WESLEY G. FRANCE (*J. Amer. Chem. Soc.*, 1921, **43**, 2161—2171).—The effect of gelatin on the transport numbers of gelatin has been investigated. It is shown that the addition of sulphuric acid up to 20% increases the transport number of the anion of sulphuric acid (0.1—0.01*M*) from 0.187 in the absence of gelatin to 0.685 in the presence of 20% gelatin. The effective concentration of 0.1*M* and 0.01*M* sulphuric acid solution is reduced by the addition of gelatin. The conductivities of sulphuric acid are reduced by the addition of gelatin. An hypothesis is put forward to account for the action of gelatin in the presence of electrolytes.

J. F. S.

Electro-osmosis. A. H. W. ATEN (*Chem. Weekblad*, 1921, **18**, 690—692).—The theory of electro-osmosis of liquids through porous solids is discussed, and the volume passing in unit time shown to be directly proportional to the current density, the dielectric constant of the liquid and the potential difference between the liquid and the porous material, and inversely proportional to the viscosity and conductivity of the liquid. The various attempted technical applications of the phenomenon, and of kataphoresis (which term is restricted to the motion of solid particles suspended in a liquid under the influence of the electric field) are described.

S. I. L.

Some Electro-osmotic Experiments with de Haen's Membrane Filters. P. H. PRAUSNITZ (*Kolloid Z.*, 1921, **29**, 293—309).—The electro-osmosis of water, solutions of ammonia, sulphuric acid, ammonium sulphate, sodium sulphate, potassium ferricyanide, and aluminium sulphate, through de Haen filters for varying size of pores, has been investigated at 25° and under a pressure of 43 cm. of water. It is shown that the de Haen filters are particularly well adapted to the study of the influence of electrolytes on electro-endosmosis. Filters of medium-sized pores (Nos. 20 and 320) when 5 cm. thick allow 100—500 c.c. of water to pass per hour under the pressure used in the experiments. Distilled water ($\lambda = 6 \times 10^{-6}$) travels toward the cathode through the membrane under a current of 100 volts and about 0.1 ampere at the rate of 1—2 litres per hour. The addition of sulphuric acid to the anode compartment, or any addition of thorium chloride or aluminium sulphate, reduces the electro-endosmosis to values below that of water and in cases may produce even a weak anodic endosmosis. In

other cases, the addition of electrolytes brings about an increased cathodic endosmosis, in which the amount of water transported varies with the time the current is flowing and the nature and concentration of the electrolyte. It is shown that, in general, with an electrical conductivity of $200\text{--}400 \times 10^{-6}$ and a consumption of energy of $0.01\text{--}0.06$ KWH/litre of transported liquid, 8—16 litres/hour of liquid can be made to pass through a diaphragm of 100 cm^2 . It is not always necessary that the electrolyte should pass through the membrane as such; particularly in the case of ammonia it is shown that the addition of the base to the cathode compartment can bring about a very considerable endosmosis of the water, which is to be attributed to the wandering in the opposite direction of those ions which are responsible for the discharging of the membrane. The transport of the SO_4 -ion through a de Haen membrane from negative to positive compartments brings about a considerable passage of water from the positive to the negative compartment. In every case there is an optimum electrolyte concentration, and if this is exceeded anomalies occur which are found to be connected closely with the electrical conductivity. The discharge of the membrane is held to be due in all probability to the adsorption of hydroxyl ions, for all processes which effect a reduction of the hydroxyl-ion concentration also effect a reduction of the electro-endosmosis (cf. *D.R.-P.* 333575, 1919). J. F. S.

Attainment of Constant High Temperatures. E. MOSER (*Z. angew. Chem.*, 1921, **34**, 625).—The apparatus or object to be maintained at constant temperature is immersed in paraffin contained in a cylindrical glass vessel surrounded by an outer glass vessel of similar shape but enlarged to a spherical form at the bottom to accommodate a liquid of suitable boiling point, which is kept in ebullition so that its vapour surrounds the inner vessel. The two vessels are sealed together round the top except where a tube serves to connect the vapour jacket with a reflux condenser. The lower end of the condenser tube projects downwards for some distance into the vapour jacket and the vapour enters it through a lateral orifice. The end of the tube is drawn out to a jet and disposed vertically above the boiling liquid. By this arrangement the condensed liquid is warmed again before it falls back into the jacket and is prevented from falling on the hot walls of the vessels. Liquids of high boiling point, such as ethyl benzoate, b. p. 212.5° , have been successfully used in the outer vessel. J. H. L.

The Expansions of some Refractory Materials at High Temperatures. B. BOGITCH (*Compt. rend.*, 1921, **173**, 1358—1360).—The expansion curves for bauxite, clay, silica, chromite, and magnesia over the temperature range 0° to 1600° are given. The smallest expansion was shown by the bauxite, which is suitable for use in furnaces subject to sudden changes in temperature. Silica showed the most irregular expansion; it expanded rapidly up to 600° and then only very slowly, and above 1000° showed a slight contraction. The curve shows two singular points at 210°

and 570° respectively, which are the transformation points of cristobalite and quartz. Furnaces constructed with such bricks must only be heated up very slowly, the temperature rise being less than 50° per hour. The curve for the clay bricks is intermediate between the curves for bauxite and silica. Magnesia and chromite showed the biggest expansions, and such bricks are only suitable for the construction of furnaces in continuous work. W. G.

Coefficient of Expansion of Molten Salts. W. HERZ (*Z. physikal. Chem.*, 1921, 98, 98—102).—In previous papers (A., 1914, ii, 25, 245; 1921, ii, 381) it has been shown that Mendeléev's equation is true for the expansion of normal organic liquids. In the present paper, the equation has been applied to the molten inorganic substances, sodium molybdate, lithium nitrate, sodium nitrate, rubidium bromide, caesium nitrate, potassium tungstate, caesium iodide, potassium molybdate, sodium tungstate, rubidium fluoride, caesium sulphate, and thallous nitrate. The data employed are taken from the work of Jaeger (A., 1918, ii, 33). The densities calculated by the Mendeléev formula agree in most cases to about 1 per 1800 with the experimental values. The only cases where divergence is shown are thallous nitrate and sodium tungstate, and here it is not much greater than in the other cases. The density values calculated by the Mendeléev equation are compared with those calculated by Jaeger (*loc. cit.*) by a single constant formula for caesium iodide and rubidium fluoride, and by a two constant formula for caesium sulphate and the value of the single constant Mendeléev formula demonstrated. J. F. S.

General Methods for the Determination of the True Molecular Constitution of Pure Substances and their Mixtures. MARIO BASTO WAGNER (*Z. physikal. Chem.*, 1921, 98, 81—93).—A theoretical paper in which the author differentiates between the thermodynamics of mixtures treated in connexion with their independent components and the thermodynamics of mixtures treated in connexion with their true components. The advantages and drawbacks of both methods of treatment are explained and the nature of the results obtainable is pointed out in each case. The various properties which may be used in the determination of constitution are enumerated and their application is indicated. These include volume, heat change, vapour pressure, solubility, freezing point, boiling point, specific heat, compressibility, thermal coefficient of expansion, and osmotic pressure. The properties, dielectric constant, viscosity, refraction, and surface tension do not permit conclusions being drawn which may be used in the second method of treatment. J. F. S.

Thermodynamics of Mixtures. IX. MARIO BASTO WAGNER (*Z. physikal. Chem.*, 1921, 98, 151—156; cf. A., 1921, ii, 375).—A theoretical paper in which the theory of the heat of mixtures is developed on the basis of the thermodynamics of mixtures treated in connexion with the true components (cf. preceding abstract). J. F. S.

Theory of Equations of State. II. MARIO BASTO WAGNER (*Z. physikal. Chem.*, 1921, **98**, 244—251; cf. preceding abstract).—The hypothesis developed previously is extended. The present paper deals with the theory of the heat of vaporisation and the internal pressure.
J. F. S.

Equation of Condition of Nitrogen at Small Pressures and Low Temperatures. R. BARTELS and A. EUCKEN (*Z. physikal. Chem.*, 1921, **98**, 70—79).—The equation of condition of nitrogen has been measured by means of a constant pressure gas thermometer between the temperatures 90° and 76° Abs. and between the pressures 0.5 and 1.0 atm. The results are represented by means of an empirical formula for the correction quantity B' of the form $v = RT/p - RTB'$. This formula has been used to calculate the molecular heat of nitrogen for the ideal gas conditions, and the value 6.873 obtained for C_p . This value is 0.07 cal., or 1% lower than that (6.944 cal.) demanded by the classical theory, whilst the Berthelot equation of condition leads to a value 0.2 cal. lower than the classical value for the temperature 92° Abs. A comparison of the reduced B' values for a range of temperature shows that the curves for hydrogen and nitrogen cannot be brought into coincidence, but that at low temperatures they diverge.
J. F. S.

Equation of Condition. II. Principal Equation of Condition. III. Principal Equation of Condition and the Equation of Condition for Individual Substances. IV. Compressibility Equations of Liquids. A. WOHL (*Z. physikal. Chem.*, 1921, **99**, 207—225, 226—233, 234—241; cf. A., 1914, ii, 430).—A series of theoretical papers in the first of which the author further develops the equation of condition previously published (*loc. cit.*). The various relationships brought out by this equation are considered and compared with the equations of van der Waals and Kamerlingh Onnes. In the second paper, equations of condition are developed and considered for helium, hydrogen, water, and ethyl alcohol. The third paper deals with compressibility equations for liquids, particularly highly compressed substances. The substances ethyl ether, carbon dioxide, and hydrogen are considered.
J. F. S.

Extension of Ebullioscopy and its Application to Binary Mixtures. ERNST BECKMANN and OTTO LIESCHE (*Z. physikal. Chem.*, 1921, **98**, 438—454).—The authors have combined the Raoult equation $P = P_0[N/(N+n)]$ with Dalton's law $P = B[N'/(N'+n')]$ to form the equation $P_0[N/(N+n)] = B[N'/(N'+n')]$ and shown that this equation may be used generally in the ebullioscopy of binary mixtures, and may be extended to meet complicated cases, since the determination of the absolute boiling point fixes the value of P_0 with the help of vapour pressure tables or formulæ. The introduction of the boiling point difference Δ and the ebullioscopic constant E leads to a special ebullioscopic equation

$\Delta = E(n - n')$, which may only be used for small Δ values. If in either of the above equations n or n' is given the value zero, the two limiting cases are obtained, in which the second component (represented by small letters) is non-volatile or is not soluble in the first component.

J. F. S.

A Vapour Pressure Formula with a General Integration Constant. F. A. HENGLEIN (*Z. physikal. Chem.*, 1921, **98**, 1—13).—A theoretical paper in which the constants of a previously published rule (A., 1920, ii, 732) for the comparison of the vapour pressure curves of two substances, $\log T_1 = a \log T_2 + b$, have been brought into relationship with the Trouton constants. From this relationship a vapour pressure formula has been deduced which contains two specific constants and has the form $\log p_{(\text{atm.})} = -k/T^n + 4.6222$. The formula holds, in comparison with other two constant vapour pressure formulæ, over greater temperature ranges and is characterised by the fact that the integration constant is the same for all substances. The index n is shown to be equal to $\lambda/(v - v')p \cdot K$ where K has the same value for all substances, and consequently $\lambda/(v - v')p$ may be calculated from the above formula; on the other hand, n may be determined from $\lambda/(v - v')p$ or λ/RT . To calculate the actual formula, two vapour pressure determinations are necessary, or the knowledge of one vapour pressure point and the molecular latent heat of vaporisation at the same pressure. The formula is shown to be equally true for sublimation pressures, and it may also be applied to decomposition pressures.

J. F. S.

Theory of Binary Mixtures. VII. Mixtures of Ethyl Ether and Bromoform. F. DOLEZALEK and M. SCHULZE (*Z. physikal. Chem.*, 1921, **98**, 395—429; cf. A., 1920, ii, 32).—The vapour pressure of a series of mixtures of ethyl ether and bromoform has been determined at 25° and 75°; the contraction on mixing, the coefficient of expansion, and the density of the mixtures have been determined at 25°. The results show that on mixing the components a mutual partial complex formation occurs. The resulting compound consists of one molecule of ether and one molecule of bromoform; it occurs in the solution in a dissociated condition and on heating it decomposes. Bromoform itself is strongly associated, but with decreasing concentration the association decreases. The vapour pressure was determined by a new static method. A theory is developed whereby the molecular constitution may be calculated from the vapour pressure measurements. From two measured vapour pressures the combination constant K and the association constant k were determined, and from these the molecular constitution of the mixture was calculated for 25°. The results show that at 25° bromoform consists of 65% of bimolecules and that in the mixture of 1 mol. of ether and 1 mol. of bromoform, 23% of the equi-molecular compound is present. The vapour pressure for all mixtures has been calculated from the molecular composition, and found to be in good agreement with the experimental values. A marked contraction occurs on mixing

ether and bromoform; the contraction curve is strongly depressed on account of the association of the bromoform. The contraction has been calculated on the basis of the theory and found to agree with the observed contraction. From the coefficient of expansion the contraction at 15°, 20°, and 30° has been calculated. From the contraction at 25°, the density of the mixtures was calculated, and the results show that the measurement of volume change by the differential method is the most exact for the measurement of easily volatile liquid mixtures. J. F. S.

Vapour Pressure of Binary Mixtures. GERHARDT C. SCHMIDT (*Z. physikal. Chem.*, 1921, 99, 71—86).—A new simple static method of measuring the vapour pressure of liquids is described. It is shown that the dynamic method of measuring the vapour pressure of mixtures gives only moderate results and then only when the experiments are made under definite conditions, and it is inferior in every way to the new static method. The vapour pressures of the following mixtures, benzene-ethyl ether, benzene-chloroform, benzene-methyl acetate, methyl acetate-ethyl acetate, benzene-carbon disulphide, benzene-toluene, toluene-carbon disulphide, ethyl alcohol-methyl alcohol, benzene-methyl alcohol, chloroform-carbon disulphide, and acetone-chloroform have been measured up to a pressure of 4 atmospheres. By plotting the molecular compositions as abscissæ and the vapour pressures as ordinates, it is found that at all temperatures (20—100°) mixtures of ethyl alcohol and methyl alcohol give a straight line. Mixtures of benzene and ether and of benzene and chloroform give approximately straight lines up to 30°, but at higher temperature curves which are convex to the abscissa axis, and the higher the temperature the greater the convexity. Mixtures of acetone and chloroform at 0° give a straight line, and at higher temperatures curves convex to the abscissa axis, which become more convex as the temperature is raised. Mixtures of benzene and toluene, ethyl acetate and methyl acetate, and benzene and methyl acetate at low temperatures give straight lines, but at higher temperatures the curves are concave to the abscissa axis and the concavity becomes greater as the temperature is raised. Mixtures of benzene and carbon disulphide, toluene and carbon disulphide, chloroform and carbon disulphide, and benzene and methyl alcohol, give at 0° curves which are concave to the abscissa axis, and the higher the temperature the greater the concavity, this being particularly noticeable in the case of benzene-methyl alcohol. From the fact that the higher the temperature the greater the concavity or convexity, the conclusion can be drawn that for every liquid mixture there is a temperature at which the relationship between the vapour pressure and the molecular composition may be represented by a straight line, generally speaking this occurs at low temperatures. It is shown that a series of deductions from Dolezalek's theory of binary solutions, which refer to the influence of temperature on the course of the isotherms, are not confirmed by the experiments. Consequently the theory cannot be maintained. J. F. S.

Intersecting Vapour Pressure Curves and Deductions therefrom. CARL VON RECHENBERG (*Z. physikal. Chem.*, 1921, 99, 87—104).—A large number of pairs of substances are recorded for which the vapour pressure curves intersect. From a comparison of the intersecting vapour pressure curves the following conclusions are drawn. All liquid and solid substances consist of large molecular complexes of undetermined variable composition of different stability, and present as many classes of compounds as they have different molecular types. Every individual compound behaves in a characteristic manner. The division of liquids into associated and non-associated groups from surface tension measurements is incorrect. All compounds are associated, but the so-called associated compounds are more loosely combined. The sub-division only separates the most divergent compounds, but does not take account of the intermediate and transition compounds. A table is drawn up of the various aggregates according to the increase of vapour pressure with increase of temperature. The table is very similar to Rothmund's solubility table, from which it follows that vapour formation and solubility are parallel phenomena and must be due to the same cause. J. F. S.

Vaporisation of Solutions of Liquid Pairs possessing Intersecting Vapour Pressure Curves. CARL VON RECHENBERG (*Z. physikal. Chem.*, 1921, 99, 105—115; cf. preceding abstract).—The Bancroft rule, that liquids with intersecting vapour pressure curves form solutions with a characteristic point, has been confirmed, when notice is taken of the fact that the property of the characteristic solution only obtains inside a limited pressure region. A large number of homogeneous mixtures with intersecting vapour pressure curves are tabulated with the minimum boiling point and the concentration of the characteristic fraction at 760 mm.

J. F. S.

Distillation and Rectification. L. GAY (*Chim. et Ind.*, 1921, 6, 567—578; cf. this vol., ii, 85).—In order to obtain the maximum efficiency from a column, *ceteris paribus*, the higher the temperature of the initial mixture the lower should be the level of its introduction into the distillation apparatus. This level should be such that an initial liquid or gaseous mixture meets a liquid or gaseous mixture of approximately the same composition in the column. If the liquid mixture is at its boiling point, or rather if the gaseous mixture is at its condensation point, then the composition of the two mixtures is identical. The determination of the minimum intensity of heat and the minimum number of compartments of the column necessary for an initial gaseous mixture has been carried out in a similar manner to that used for an initial liquid mixture (*loc. cit.*), and applied to the separation of practically pure water and alcohol from any gaseous mixture of water and alcohol at 95°. The determination of the minimum intensity of heat necessary to remove a trace of impurity from a liquid has also been carried out. F. M. R.

Heat of Formation of Silver Iodide. HUGH STOTT TAYLOR and WILLIAM THEODORE ANDERSON, jun. (*J. Amer. Chem. Soc.*, 1921, **43**, 2014—2017).—Braune and Koref's calorimetric measurement of the heat of formation of silver iodide (A., 1914, ii, 536), using *N*-potassium cyanide instead of 3*N*-solutions as solvent for silver iodide, have been repeated. The result 15,100 cal. per mol. of silver iodide has been obtained, which confirms Braune and Koref's value.
J. F. S.

Relationship between Heat of Solution and Heat of Fusion of Organic Substances. GEORG GEHLHOFF (*Z. physikal. Chem.*, 1921, **98**, 252—259).—The heat of solution of naphthalene, diphenyl, diphenylamine, azobenzene, α -naphthylamine, β -naphthylamine, urethane, paraffin (wax), nitronaphthalene, nitrophenol, *m*-dinitrobenzene, phenol, menthol, thymol, pyrogallol, resorcinol, citric acid, and chloral hydrate has been determined in 1—3% solutions in the solvents water, benzene, ethyl ether, alcohol, carbon disulphide, acetone, amyl acetate, aniline, chloroform, and acetic acid. The results show that for a series of substances, particularly hydrocarbons, aromatic amines, urethane, and azobenzene, the heat of solution in various solvents is practically identical with the heat of fusion. Organic substances containing the hydroxyl group give very different values for the heat of solution and the heat of fusion, whilst substances containing the nitro-group take up an intermediate position between the other substances. The heat of solution of the sodium, potassium, and ammonium haloids has been determined in aqueous solution, and the results are discussed in connexion with the structures and space lattices of these salts.
J. F. S.

Heats of Solution and of Transformation of the Acido- and Aquo-cobalt Pentammines. ARTHUR B. LAMB and JOHN P. SIMMONS (*J. Amer. Chem. Soc.*, 1921, **43**, 2188—2199).—A method of calorimetry has been developed in which only observations on a silver coulometer and of time and electrical resistance are required. It is shown that the same cobalt sulphide, Co_2S_3 , is produced by the action of sodium sulphide on chloro- and aquo-pentammine-cobalt chloride in aqueous solution. The solubility of chloro-pentamminecobalt chloride in water at 0°, 30°, and 50° has been determined, and the values 2.244, 6.047, and 10.01, respectively, have been obtained in grams per litre. The heat of solution has been calculated from these data and the value 13,440 cal. obtained, which is in fair agreement with the experimental value, 12,400 cal. The heats of solution of a number of acido- and aquo-pentammine-cobalt salts in water have been measured. The following values are found at 25°, in cal., chloro-chloride, —12,420; aquo-chloride, —6,460; bromo-bromide, —12,660; aquo-bromide, —9,220; nitrate-nitrate, —14,900; and aquo-nitrate, —15,300. The heat of reaction of solutions of pentammine salts with solutions of sodium sulphide has been measured at 25° and that of the solid salts with sodium sulphide solutions calculated. The

following values in cal. at 25° are obtained, chloro-chloride, 15,070; aquo-chloride, 12,930; bromo-bromide, 13,290; aquo-bromide, 11,600; nitrate-nitrate, 12,340, and aquo-nitrate, 11,860. From these results the heat of transformation of the acido-salt to the aquo-salt in each case has been calculated, and the values chloride, 2,140 cal.; bromide, 1,690 cal., and nitrate, 480 cal. have been obtained. J. F. S.

Heat of Wetting of Silica Gel. W. A. PATRICK and F. V. GRIM (*J. Amer. Chem. Soc.*, 1921, **43**, 2144—2150).—The heat change occasioned by wetting silica gels with water, ethyl alcohol, benzene, carbon tetrachloride, and aniline, respectively, has been determined at 25°. The quantity of liquid used in each experiment was 50·07 c.c. and the amount of gel 2·3—2·5 grams. The following values in cal./gram of gel were obtained: water, 19·22; alcohol, 22·63; benzene, 11·13; carbon tetrachloride, 8·42, and aniline, 17·54. The heats of wetting have been discussed and found to be in keeping with the changes of surface energy. J. F. S.

Viscosity of Gelatin Sols. ROBERT H. BOGUE (*J. Amer. Chem. Soc.*, 1921, **43**, 1764—1773).—A number of experiments have been effected with gelatin sols to determine the relation between viscosity and concentration. The data obtained have been applied to Hatschek's formula for the viscosity of emulsoids, and it is shown that the value of A'/A , representing the volume occupied per unit weight of dispersed phase, is not a constant with varying concentration, but rises regularly to a maximum and thereafter regularly diminishes with increasing concentration. This behaviour is noticed also in the case of other colloids. A tentative explanation is presented based on the effect which increasing concentrations of dispersed phase will have on the surface tension of the dispersion medium. Assuming the equilibrium, surface tension \rightleftharpoons solvation potential, an empirical relation is developed which, with a fair degree of accuracy, defines the equilibrium. At very high concentrations, a reversal of phase probably occurs, at which no expressions yet developed adequately represent the relations involved. A high degree of solvation is shown to be indicated by a high coefficient of viscosity, that is, solvation and viscosity are parallel functions. Isoelectric gelatin at a hydrogen-ion concentration 2×10^{-5} is shown to have the lowest viscosity and the lowest degree of solvation, gelatin chloride at a hydrogen-ion concentration $3 \cdot 1 \times 10^{-4}$ the highest, and calcium gelatinate at a hydrogen-ion concentration $2 \cdot 5 \times 10^{-6}$ is intermediate. If an excess of acid is allowed to remain in the gelatin solution, even although the acid be of very low concentration, the viscosity and degree of solvation will be reduced. J. F. S.

A New Viscostalagmometer for the Estimation of Surface Tension and Viscosity of Liquids of Very Different Limpidity. I. TRAUBE (*Biochem. Z.*, 1921, **120**, 106—107).—The essential feature consists in the adaptation to the ordinary form of stalagmo-

meter of five interchangeable ground-in mouth-pieces of differing capillary bore, thus allowing measurements to be performed with the same apparatus on liquids of a great range of fluidities.

H. K.

Ionisation and Surface Activity of Aqueous Solutions of Aliphatic Acids. WILH. WINDISCH and PHIL. OSSWALD (*Z. physikal. Chem.*, 1921, 99, 172—188).—The connexion between ionisation and surface activity of organic acids has been investigated. It is shown that of *N*/10-acid or alkali when added to an aqueous solution acts only as a diluent, that is, the surface activity is independent of the hydrogen-ion concentration. Experiments have been made to determine the concentration of acids of different strengths, such as hydrochloric, formic, and acetic acids, which must be added to solutions of nonoic acid so that the same size of drops may be obtained, and also the concentrations of these acids which will produce the same colour with methyl-orange solution. It is shown that free nonoic acid in aqueous solution is 10% dissociated, and therefore the addition of acid increases its surface activity. The simultaneous presence of sodium chloride is without action on the activity. With continued addition of acid, the size of the drops sinks to a constant minimum. A solution of sodium nonoate exhibits a small activity, because through hydrolysis a small quantity of acid exists in the free condition. By the addition of a small quantity of alkali, the free molecules are ionised and the size of the drop sinks to that of water. An addition of hydrochloric acid produces fresh acid molecules, and the size of the drop sinks to a minimum, which is reached when all the aliphatic acid ions have disappeared. Consequently it follows that the surface activity of aqueous solutions of aliphatic acids is due to ionisation processes, and the possibility of calculating the size of the drops due to the action of various additions is pointed out.

J. F. S.

High Pressure due to Adsorption, and the Density and Volume Relations of Charcoal. WILLIAM D. HARKINS and D. T. EWING (*J. Amer. Chem. Soc.*, 1921, 43, 1787—1802).—The present paper furnishes evidence in favour of the hypothesis previously expressed (*A.*, 1921, ii, 87) by the authors, which states that the liquids which penetrate into charcoal are compressed by a force due to molecular attraction, which acts as a pressure of many thousand atmospheres (20,000 or more). The present paper shows that it is the liquid in the micro-pores and not the macro-pores which is compressed, and consequently the pressure would probably be much greater than the figure mentioned. Not only charcoal but other porous substances and fine powders should exert this compressive effect, but to a much smaller extent. A number of experiments are described in which the compression is demonstrated by volume changes, but the values measured have not exceeded 72 atms. Ethyl ether, which is much more compressible than water, occupies a volume in charcoal which is

about 10% less than that occupied by the amount of water which outside the charcoal is equal in volume to the ether. It is probable that the water in the micro-pores is compressed by about 25% or more, whilst the ether is compressed by 40%. The liquids in the macro-pores of the diameter 1.2×10^{-3} cm. are not compressed sufficiently to produce a noticeable effect on the volume. The true volume relations in 1 c.c. of a characteristic coconut shell charcoal are 0.28 c.c. micro-pores, 0.18 c.c. macro-pores, 0.54 c.c. carbon, which give a density of 1.60 for the carbon. The density of the lumps of such a charcoal is 0.866. Charcoals which are inactive as adsorbents of gases do not exert a compressive effect on the adsorbed liquids of a sufficiently high magnitude to be very evident, although there appears to be a slight effect of this nature. The densities of the carbon in the two inactive wood charcoals investigated are 1.65 and 1.50. When coconut-shell charcoals alone are considered, it is found that the lower the apparent density in an organic liquid, the less is the adsorptive action on vapours, and it is indicated that this relation may be expected to hold better the more compressible the liquid which is adsorbed, so that either ethyl ether or pentane or another highly compressible liquid should be used in such tests. Simple thermodynamic equations are given for the heat of immersion or adsorption of a plane surface. Whilst there is probably no definite area of surface inside a lump of charcoal, a definition for an apparent area may be given. The one chosen here is that the apparent area with respect to the heat of immersion is the area of the plane surface of carbon which will develop the same amount of heat on immersion as is equal to that developed by the immersion of 1 gram of the charcoal in the same liquid. Since the film in the charcoal is probably a number of molecules thick, this apparent area is probably larger than corresponds with the carbon surface. This method indicates that the area of 1 gram of charcoal is 120 sq. metres. The magnitude of the heat of immersion of liquids on mercury is 3.3×10^{-6} cal. for octane and 3.25×10^{-6} cal. for isobutyl alcohol.

J. F. S.

The Origin of the Potential Differences Responsible for Anomalous Osmosis. JACQUES LOEB (*J. Gen. Physiol.*, 1921, 4, 213—226).—If a collodion bag, coated on the inside with protein (gelatin) separate two identical acid solutions and a neutral salt be added to the inside, the rate of diffusion inwards through the membrane is a minimum at the isoelectric point of the protein, increases with the acidity to a maximum, and then decreases. The potential difference between the two solutions varies similarly. In the absence of protein, these characteristic variations are not obtained. The anomalous variation of the potential difference is apparently a consequence of the Donnan equilibrium between solution and protein, and this potential difference causes electrical endosmosis, and hence the irregular rates of diffusion. The potential difference is also influenced by the diffusion potential, but this exists although no protein is present.

W. O. K.

Penetration of Electrolytes into Gels. III. Influence of the Concentration of the Gel on the Coefficient of Diffusion of Sodium Chloride. WALTER STILES and GILBERT SMITHSON ADAIR (*Biochem. J.*, 1921, 15, 620—626).—The coefficient of diffusion from normal solution into gelatin and agar gels was measured by the indicator method previously described (*A.*, 1920, ii, 235). With increasing concentration of gel, the coefficient of diffusion at first decreases rapidly according to an exponential relation, but with concentrations above 2% the curve connecting coefficient of diffusion and concentration is approximately a straight line. It is shown how this result can be harmonised with the relation between diffusion and viscosity found by Einstein, Sutherland, and von Smolukowski. The diffusion coefficients found when extrapolated for diffusion in water are about 7.5% higher than those obtained from the results of Öholm (*A.*, 1905, ii, 147). This is attributed to the fact that in Öholm's experiments the salt concentration refers to ionised+un-ionised salt, whereas the authors measure only the rate of penetration of the ions, which diffuse about twice as fast as the un-ionised salt.

The resistance which even a stiff jelly offers to salt diffusion is quite small; the rate is reduced by less than 2% for each gram of gelatin added to 100 c.c. when the concentration exceeds 2%.

G. B.

Penetration of Electrolytes into Gels. IV. Diffusion of Sulphates. WALTER STILES (*Biochem. J.*, 1921, 15, 629—635).—In 0.5% agar, ammonium, potassium, sodium, and magnesium sulphates diffuse almost as fast as in pure water. In 10% gelatin, the rate is decreased by about 25%. The diffusion coefficient of copper sulphate in agar is considerably greater than in water, and in 10% gelatin less than in water, but still greater than the values for other sulphates indicate. The abnormal results of von Fürth and Bubanović (*A.*, 1920, ii, 94) have not been confirmed. The comparatively small uptake of sulphates by plant tissue is not to be explained on the basis of retarded diffusion through a gel.

G. B.

A General Theory of Solutions of Electrolytes. JNANENDRA CHANDRA GHOSH (*Z. physikal. Chem.*, 1921, 98, 211—238).—A theoretical paper in which the author develops a new theory of electrolyte solutions on the basis (i) that in solutions of strong electrolytes only ions exist, and (ii) in solutions of weak electrolytes or those of medium strength there exists an equilibrium between a non-polar form (non-electrolyte) and a polar form (which behaves as a strong electrolyte). This hypothesis is examined in connexion with the data obtained in recent years on electrical conductivity, activity coefficients, conductivity in non-aqueous solvents, influence of salts on solubility relations, absorption of light by salt solutions, and the relation between osmotic pressure and activity coefficients.

J. F. S.

Theory of Strong Electrolytes. H. KALLMANN (*Z. physikal. Chem.*, 1921, 98, 433—437).—A number of criticisms of Ghosh's

hypothesis of solutions of electrolytes (T., 1918, **113**, 449, 627, 707, and preceding abstract) are put forward. J. F. S.

Solubilities in Mixtures of Two Solvents. GEORGE PUCHER and WILLIAM M. DEHN (*J. Amer. Chem. Soc.*, 1921, **43**, 1753—1758; cf. A., 1917, ii, 445).—The solubility of forty-five organic substances has been determined at 20—25°, in ethyl alcohol, quinoline, and equimolecular mixtures of the two solvents. In some cases, the solubility in the mixed solvent lies either above or below that in the individual solvents; this is most marked in the case of carbamide, "saccharin," *p*-nitrophenol, acetanilide, resorcinol, piperine, cinnamic acid, and chloral hydrate and in all these cases compound formation is indicated. The solubility of carbamide, "saccharin," brucine, and *o*-nitrophenol in a series of mixtures of alcohol and quinoline has been determined at 25°, and the solubility of lactose at 1° and 25° and raffinose at 25° in a series of mixtures of water and pyridine has also been determined. It is shown that the solubility of carbamide decreases quite regularly as the concentration of quinoline increases, but at 60% quinoline a break occurs and the compound of one molecule of quinoline and three molecules of carbamide is formed, m. p. 154°. Compound formation is indicated in the case of "saccharin," brucine, lactose, and raffinose, but no definite compound was isolated. J. F. S.

Compound Formation and Viscosity in Solutions of the Types Acid-Ester, Acid-Ketone, and Acid-Acid. JAMES KENDALL and ELIZABETH BRAKELEY (*J. Amer. Chem. Soc.*, 1921, **43**, 1826—1834; cf. Kendall and Gross, this vol., ii, 32, 33).—The viscosity-composition curves at 25° have been determined for the systems trichloroacetic acid-ethyl acetate, acetic acid-ethyl acetate, trichloroacetic acid-ethyl benzoate, acetic acid-ethyl benzoate, trichloroacetic acid-acetone, acetic acid-acetone, trichloroacetic acid-acetophenone, acetic acid-acetophenone and trichloroacetic acid-acetic acid. The results obtained indicate extensive compound formation in solutions of the esters and ketones with a strong acid (trichloroacetic). The same substances with a weak acid (acetic) give viscosity curves which are much less abnormal, although compound formation is still evident. The curve for the system acetic acid-trichloroacetic acid exhibits considerable compound formation and an examination of the results of previous workers on systems of the types acid-acid and acid-base shows that this behaviour is general except when the components are of similar acidic strength. The rules formulated in previous papers relating to the extent of compound formation with diversity in character of the components are confirmed by the present work. J. F. S.

Compound Formation and Conductivity in Systems of the Types Formic Acid-Metal Formate and Sulphuric Acid-Metal Sulphate. JAMES KENDALL, HOWARD ADLER, and ARTHUR W. DAVIDSON (*J. Amer. Chem. Soc.*, 1921, **43**, 1846—1853; cf. preceding abstract).—Electrical conductivity determinations

have been made of solutions of various concentrations of barium, magnesium, and lead formates in anhydrous formic acid at 25°, and of solutions of sodium, barium, calcium, magnesium, zinc, lead, and silver sulphates in anhydrous sulphuric acid. In this case, the solutions contained 0.04 equivalent of solid sulphate. Comparing the results for the formates with those obtained for other formates in formic acid by Schlesinger (A., 1919, ii, 91; 1920, ii, 72, 73), it is shown that a definite parallelism exists between the degree of ionisation and the extent of compound formation. In the case of the sulphates, however, the uncertainty of the correction to be applied for the solvent is so great as to obscure the results. Variations in compound formation, solubility, and ionisation in systems of the general type $HX-RX$ are all more extreme the weaker the acidic radicle X . Hydroxides in aqueous solution differ more widely than formates in formic acid, and these are more diverse than sulphates in sulphuric acid solution.

J. F. S.

The Photography of Opaque Crystals. M. FRANÇOIS and CH. LORMAND (*Bull. Soc. chim.*, 1921, [iv], 29, 1056—1059).—The application of the apparatus previously described for the photography of transparent crystals (cf. A., 1921, ii, 626) to the photography of opaque crystals is given in detail.

W. G.

Stereoscopic Photography of Crystals. M. FRANÇOIS and CH. LORMAND (*Bull. Soc. chim.*, 1921, [iv], 29, 1059—1063).—For this purpose, a microscope, the tube of which can be inclined successively to the left and right, is used. It is fitted with the apparatus previously described for the photomicrography of crystals (cf. A., 1921, ii, 626, and preceding abstract). A photograph is taken with the tube inclined to the left and then one with it inclined equally to the right. The examination of the two photographs in a stereoscope gives a stereoscopic picture of the crystals in relief.

W. G.

Ionic Properties and Crystallo-chemical Relationships. I. The Properties of the Ions which appear in Crystals. H. GRIMM (*Z. physikal. Chem.*, 1921, 98, 353—394).—A theoretical paper in which it is shown that the electrostatic action of the ions in crystals is due to the following ionic properties, (i) the charge of the ions, (ii) the radius of the ions, (iii) the structure of the ions, particularly the surface and number of exterior electrons. The ions are divided into five principal groups depending on the number of exterior electrons. (i) Ions of the helium type (2 exterior electrons), (ii) ions of the Ne, Ar, Kr, Xe, and emanation type (8 exterior electrons), (iii) ions of the Cu^+ , Ag^+ , Au^+ type (probably with 18 exterior electrons), (iv) ions of the Te^+ and Pb^{++} type, (v) ions of the type Mn^{++} , Fe^{++} , Co^{++} , Ni^{++} (transition ions). It is shown from many compounds and elements that the expression $\phi_A - \phi_{Ne} > \phi_{Xe} - \phi_{Kr} > \phi_{Kr} - \phi_A$ holds for many physical properties where ϕ is the numerical value of the property and the index is that of the inactive gas the structure of which comes nearest to that of the ion concerned. This inequality is due to an analogous inequality of the ionic radii. The ionic radii of O^{--} , S^{--} , Se^{--} , Te^{--} ; Mg^{++} , Ca^{++} , Sr^{++} , Ba^{++} have been calculated and that

of Cs^+ has been approximated. The series order of the other ionic radii has been deduced from the molecular volumes of isomorphous compounds in which the interchangeable ions have the same number of exterior electrons. Ions with different structures and different exterior sheaths show a similar electrostatic action when the difference in the ionic radius compensates the other properties.

J. F. S.

Structure of Organic Crystals. SIR W. H. BRAGG (*Proc. Physical Soc.*, 1921, 34, 33—50).—The author has determined the X-ray spectra of naphthalene, α -naphthylamine, benzoic acid, acenaphthene, β -naphthol, and α -naphthol by the powder method. To interpret the results, he has made the assumption that the benzene and naphthalene rings are actual structures having definite size and form, and that they are built as a whole into the organic substances in which they occur. It is shown from crystallographic and other data that this hypothesis is an extremely probable one, and it leads to results which are in excellent agreement with crystallographic data. It is shown that the unit cell of naphthalene contains two molecules and has a mass of 213 Å.U., where 1 Å.U. = 10^{-24} grams, the linear dimensions of the cell are $a=8.34$, $b=6.05$, and $c=8.69$ Å.U. Two molecules of anthracene occupy the unit cell, and this has the linear dimensions $a=8.7$, $b=6.1$, and $c=11.6$ Å.U. The unit cell of acenaphthene contains four molecules, and has linear dimensions 8.32, 14.15, and 7.26 Å.U., respectively, whilst α -naphthol has a unit cell which also contains four molecules and has linear dimensions $a=13.1$, $b=4.9$, and $c=13.4$ Å.U.; the same number of molecules occupy the unit cell of β -naphthol and here the linear dimensions are $a=5.85$, $b=4.28$, and $c=8.7$ Å.U. The unit cell of α -naphthylamine also contains four molecules, and has the dimensions $a=8.62$, $b=14.08$, and $c=7.04$ Å.U. Benzoic acid has a wide spacing between the planes, the unit cell contains four molecules and has the dimensions $a=5.44$, $b=5.18$, and $c=21.8$ Å.U. In a note the author stated, "It is convenient in this work to extend the Ångstrom system of units so that an Å.U. of area is 10^{-16} cm.², of volume 10^{-24} cm.³ and of mass 10^{-24} grams." J. F. S.

Röntgen Spectroscopic Investigation of Organic Compounds. I. and II. K. BECKER and W. JANCKE (*Z. physikal. Chem.*, 1921, 99, 242—266, 267—274).—In the first part an account is given of the X-ray examination of the structure of a number of organic substances. The substances were examined in the form of compressed pastilles by a method which is a combination of that due to Bragg and Debye-Scherrer. In the case of indigotin it is found that the unit cell is an hexagonal parallelepiped with a rhombus as base of edge 19.55 Å.U. and height 11.80 Å.U. This contains 12 indigotin molecules. The figures refer to indigotin which has been subjected to a pressure of 4000—5000 atoms. Indigotin which has not been so compressed has a unit crystal cell with the linear dimensions $a=b=20.20$ Å.U., $c=12.15$ Å.U., which means that there is a volume contraction of 9.8%, the base decreases 3.2% and the height 2.9%. Carbamide has a unit cell

of dimensions $a=b=8.75$ Å.U., $c=7.24$ Å.U., and contains 8 molecules. The unit cell of succinic acid contains 2 molecules and has dimensions $a=5.00$ Å.U., $b=8.20$ Å.U., $c=5.40$ Å.U., $\beta=94.70^\circ$. Compression reduces all values, the c value relatively most, the values being: volume 10.3%, a 2.2%, b 4.8%, and c 5.4%. Measurements are recorded for quinol, quinone, anthraquinone, anthracene, naphthalene, phenanthrene, resorcinol, phthalic anhydride, phthalic acid, cinnamic acid (*trans*), β -phenylpropionic acid, azobenzene, hydrazobenzene, lithium oxalate, maleic acid, succinic acid, maleic acid, *d*- and *l*-tartaric acids, pentaerythritol, α -methylglucoside, and acetylenedicarboxylic acid. The results show that the number of molecules in the unit crystal cell is always greater than one, and is not the same for nearly related substances such as maleic acid and succinic acid. In the second paper, the results of X-ray examination of a large number of aliphatic acids are given. The results are arranged in four groups. I. Acids with an odd number of carbon atoms, formic, propionic, valeric, heptioic, nonoic, undecoic. II. Acids with an even number of carbon atoms, acetic, butyric, octoic, lauric, myristic, palmitic, stearic (also $\alpha\alpha'$ -distearin). III. Isomeric acids, *isobutyric*, *isovaleric*, trimethylacetic. IV. Unsaturated acids, crotonic, undecenoic, oleic, and elaidic. In many cases, the acids were measured as their lithium salts. It is shown in the case of the even-number saturated acids apart from the first two members that they possess an hexagon symmetry with 72 molecules in the unit crystal, or possibly rhombohedral with 24 molecules; the uneven-number acids are tetragonal with 24 or 48 molecules in the unit cell. There is no fundamental difference between the free acids and the lithium salts.

J. F. S.

Mixtures of Anisotropic Liquids and the Identity of Grandjean's Stratified Liquids with Liquids of the Azoxyphenetole Type. G. FRIEDEL and L. ROYER (*Compt. rend.*, 1921, 173, 1320—1322; cf. Grandjean, A., 1921, ii, 91).—In the case of mixtures of substances capable of furnishing anisotropic liquids, the two phases, the conic and thread-like (*liquides à coniques et liquides à fils*), are perfectly distinct and separated from one another by an absolute discontinuity. Mixtures of Grandjean's stratified liquids with other anisotropic liquids always show an absolute discontinuity between the stratified liquid and the conic phase. On the other hand, these mixtures do not show any discontinuity between the thread-like phase, and the stratified liquid of Grandjean. It is suggested, therefore, that Grandjean's stratified liquids are only a special aspect of the thread-like liquids of the type of azoxyphenetole.

W. G.

Some Fundamental Conceptions of Colloidal Chemistry. RICHARD ZSIGMONDY (*Z. physikal. Chem.*, 1921, 98, 14—37).—A theoretical paper in which the nature of matter in the colloidal condition is considered. The views put forward by Nägeli ("Gärung," Munich, 1879) on the nature of micellæ are considered, and it is shown that in general these views are in keeping with

experimental data. The conceptions to be generally adopted are expressed as follows: by micellæ in its broadest sense is to be understood a molecular complex of the dispersed material, which is not permeated by the dispersion medium. In the narrower sense, micellæ are crystalline, ultramicroscopic particles of the dispersed material. Micellar complexes are formed by the association of micellæ, and are ultramicroscopic particles generally, but in isolated cases may be microscopic. According to the form of the association, micellar complexes form, on the one hand, microscopic or ultramicroscopic particles which are permeated by the dispersion medium, or, on the other, ultramicroscopic or microscopic swelling crystals. J. F. S.

An Interesting Colloid Gel. ROSS AIKEN GORTNER and WALTER F. HOFFMAN (*J. Amer. Chem. Soc.*, 1921, 43, 2199—2202).—Dibenzoylcystine is insoluble in water and in the crystalline state has no hydrophilic properties, but it can give rise to rigid gels which do not contain more than 0.2% of dibenzoyl-L-cystine. The 0.2% gel is prepared by dissolving 0.2 gram of dibenzoyl-L-cystine in 5 c.c. of 95% alcohol. The solution is heated and water added to make the solution up to 100 c.c. After cooling for two to three hours, the solution sets to a gel comparable with that of a 5% gelatin gel. The gel is transparent, but after several days opaque nuclei are formed, due to crystallisation, which is accompanied by syneresis, and after several weeks most of the dibenzoyl-L-cystine has separated in definite crystals. On applying strong suction, the liquid may be drained from the gel, thus showing the coarse dispersion. The gel has a fibrillar structure. J. F. S.

Coagulation of Colloidal Solutions of Arsenious Sulphide by Electrolytes. E. F. BURTON and E. D. MACINNES (*J. Physical Chem.*, 1921, 25, 517—525).—The coagulation of solutions of arsenic sulphide sols of concentrations 0.027 gram per c.c. to 0.00337 gram per c.c. by means of potassium, lithium, magnesium, barium, aluminium, and zirconium chlorides, lanthanum sulphate, and cerium nitrate of various concentrations has been investigated. It is shown in the case of aluminium chloride that the volume of the electrolyte required for coagulation of a given amount of the disperse phase varies inversely as the concentration of the electrolyte solution, provided the final concentration of the colloid is kept constant. For univalent ions, it is shown that the concentration of ion necessary to produce coagulation increases with decreasing concentration of the colloid, whilst for bivalent ions the concentration of ion required to produce coagulation is almost constant and independent of the concentration of the colloid. With trivalent ions, the concentration of ion required to produce coagulation varies almost directly with the concentration of the colloid, whilst quadrivalent ions necessary for coagulation decrease much more rapidly than the concentration of the colloid. These results are in keeping with earlier results of Burton and Bishop (*A.*, 1921, ii, 176). It is also shown that the coagulation curve for barium chloride shows

a tendency to resemble the curve for trivalent ions as given by aluminium chloride, whilst the curve for the quadrivalent zirconium ion differs from that of the ceric ion, which is similar to that for trivalent ions. J. F. S.

Acidity and Basicity. RUDOLF KELLER (*Z. physikal. Chem.*, 1921, 98, 338—351).—A theoretical paper in which the terms acidity and basicity are considered. It is shown for the determination of the acidity of a combination solute-solvent, or the sense of the electric charge, not only the inner chemical constitution of the dissolved substance is to be considered, but also the chemical constitution of the solvent, which may, in the case of substances which are nearly amphoteric, be determinative. Further, the difference in the dielectric constants of the solvent and solute must be considered; this quantity makes its influence noticeable in the case of amphoteric, non-ionised, and associated solutes. The so-called basic dyes in aqueous solution usually migrate toward the anode, that is, they are acid according to the ruling terminology; this is particularly so in the case of the slightly dispersed and non-dissociated colour colloids; the so-called acid dyes are for the greater part basic or amphoteric. Certain acid dyes, such as picric acid, Martius-yellow, aurantia, are really acid, that is, negative, and wander to the anode. The capillary rule of Fichter-Sahlbom is generally correctly expressed in the literature, but incorrectly defined, since it places the dyes in the incorrect position in the electro-polarity series. All the experimental capillarity experiments show that immediately on dipping the paper strips, the neutral and positive dyes rise with the solvent. In this correct form, the rule holds, not only for colloids, but also for ion-dispersoids. Gobbelsröder ("Kapillaranalyse," Dresden, 1910) has actually found the strongest acids below in the anode region. The strongest acids and bases, such as hydrochloric and sulphuric acids and potassium hydroxide in suitable solvents, migrate in directions which are opposed to their nature. Concentration cells may be built up in which the strongest acids behave as bases. J. F. S.

Homogeneity and Dispersity. WA. OSTWALD (*Z. physikal. Chem.*, 1921, 99, 155—159).—A theoretical paper in which the conception specific surface is investigated on the basis of the phase rule. J. F. S.

Reactions in Heterogeneous Systems. The Systems $K_2CO_3|BaSO_4-K_2SO_4|BaCO_3$; $K_2CO_3|CaC_2O_4-K_2C_2O_4|CaCO_3$ and $K_2CrO_4|AgIO_3-KIO_3|AgCrO_4$. E. RAMANN and H. SALLINGER (*Z. physikal. Chem.*, 1921, 98, 103—150).—The systems named in the title have been experimentally investigated. By the addition by barium chloride solution to an excess of a mixture of potassium carbonate and potassium sulphate solution, the relation in which barium divides itself between the carbonate and the sulphate ions has been approximately determined. This ratio, $[CO_3^{''}]:[SO_4^{''}]$, is termed the precipitation equilibrium. In the

case of the addition of silver nitrate solution to an excess of the mixed solutions of potassium iodate and chromate, the original equilibrium partition of the silver takes place according to the anion ratio, $[\text{IO}_3'] : [\text{CrO}_4''/2]$, as in the previous case, but the two equilibria differ to an extraordinary degree in stability. Whilst the first can be approximately fixed by the rapid addition of alcohol, the second can only be fixed by a special experimental procedure. In the system $x\text{K}_2\text{CO}_3 + y\text{K}_2\text{C}_2\text{O}_4 + z\text{CaCl}_2 (z < x + y)$, the precipitation equilibrium could not be fixed by the addition of alcohol. The precipitation equilibrium in general is labile, inasmuch as immediately after the precipitation a reaction is set up which brings about a change in the relationship of the salts in the solid phase and leads to the formation of one of the salts (solution equilibrium). The solution equilibrium has been determined for various concentrations for the system $\text{K}_2\text{CO}_3|\text{BaSO}_4\text{--K}_2\text{SO}_4|\text{BaCO}_3$; for the system $\text{K}_2\text{CO}_3|\text{CaC}_2\text{O}_4\text{--K}_2\text{C}_2\text{O}_4|\text{CaCO}_3$, at various total potassium salt concentrations and various temperatures and for the system $\text{K}_2\text{CrO}_4|\text{AgIO}_3\text{--KIO}_3|\text{Ag}_2\text{CrO}_4$ at various total salt concentrations and one temperature (20°). In the system $\text{K}_2\text{CO}_3|\text{BaSO}_4\text{--K}_2\text{SO}_4|\text{BaCO}_3$, the partition relationship $\text{K}_2\text{CO}_3|\text{K}_2\text{SO}_4$ shows a marked dependence on the total salt concentration. No measurements were made on the temperature influence on this ratio. In the system $\text{K}_2\text{CO}_3|\text{CaC}_2\text{O}_4\text{--K}_2\text{C}_2\text{O}_4|\text{CaCO}_3$, the partition relationship $\text{K}_2\text{CO}_3|\text{K}_2\text{C}_2\text{O}_4$ is but slightly influenced by the total salt concentration but very strongly influenced by temperature. The ratio $\text{K}_2\text{CO}_3|\text{K}_2\text{C}_2\text{O}_4$ is shown to be a strict linear function of the temperature. The data obtained for the systems investigated and those obtained by Knüpfner (A., 1898, ii, 420) for the system $\text{KCNS}|\text{TlCl--KCl}|\text{TlCNS}$ and by Noyes and Kohr (A., 1903, ii, 201) for the system $\text{KOH}|\text{AgCl--KCl}|\text{AgOH}$ have been examined by means of the formula put forward by Nernst and Noyes (A., 1891, ii, 142).

J. F. S.

Studies on Chemical Affinity. XIII. Reaction Affinity in Systems of Solid Salts. J. N. BRÖNSTED (*Z. physikal. Chem.*, 1921, 98, 239—243).—In earlier papers (A., 1920, ii, 290, 298), a method was described whereby the reaction affinity between very soluble solid salts might be determined. This method does not yield the required results when applied to sparingly soluble salts. In the present paper, a method is described for the determination of the reaction affinity in systems of sparingly soluble solid salts. The method has been applied to the reaction $[\text{CoCl}(\text{NH}_3)_5]\text{Br}_2 + [\text{CoBr}(\text{NH}_3)_5]\text{Cl}_2 \rightarrow [\text{CoCl}(\text{NH}_3)_5]\text{Cl}_2 + [\text{CoBr}(\text{NH}_3)_5]\text{Br}_2$ and consists in determining the solubility of the four cobalt complex salts in mixtures of (i) potassium formate and chloride solution, (ii) potassium formate and bromide, (iii) sodium chlorate and chloride, and (iv) sodium chlorate and potassium bromide, where the total alkali concentration is $0.5N$ in each case. Then from the solubility products of the four cobalt complex salts by means of the gas laws the reaction affinity may be calculated. The results of the experiments lead to the value $A_0 = 621$ cal. for $T = 273^\circ$, a value

which is probably accurate to 1%. The system is characterised by the fact that the lighter ions are combined with the lighter ions and the heavier ions with the heavier.

J. F. S.

The Velocity of Decomposition of High Explosives in a Vacuum. III. Mercuric Fulminate. ROBERT CROSBIE FARMER (T., 1922, 121, 174—187).

Sensitiveness of Very Sensitive Explosives. J. EGGERT (*Z. Elektrochem.*, 1921, 27, 547—558).—The sensitiveness of a number of explosives, including the iodides of nitrogen, has been experimentally investigated. The results show that the intermolecular decomposition of nitrogen iodide takes place according to the equation $8\text{NH}_3\text{NI}_3 = 5\text{N}_2 + 6\text{NH}_4\text{I} + 9\text{I}_2$, whether the decomposition takes place in the dark or in light or is occasioned by detonation. Nitrogen iodide is not sensitive to shock, and many effects which were previously attributed to detonation are now shown to be secondary mechanical effects. The sensitiveness of nitrogen iodide and silver amide is not materially changed by lowering the temperature to -190° . Gradual isothermal increase of pressure to 5000 atm. brings about a decomposition in 70% of the cases of nitrogen iodide or silver amide, but all other explosives examined are unchanged by this treatment. An attempt is made to find an explanation of the sensitiveness of explosives; since earlier experiments have shown that the mechanical effects are not to be explained as secondary temperature effects, there only remains the possibility that local increases of pressure occur with all mechanical influences and these are to be regarded as causing the detonation. This view is supported by the present experiments on the behaviour of the sensitiveness with increase of pressure. A consideration of the processes occurring during the decomposition of explosives from the point of view of Herzfeld's theory of velocity of chemical reaction leads to a possible explanation of some of the properties of these substances. In this connexion, the primary, generally very simple, unimolecular decomposition of the explosive, which depends on the persistence of the molecule alone, is to be differentiated from the accompanying, at least bimolecular reaction, due to the collision between the products of reaction of the primary process. If in the primary reaction products are formed which possess more energy and a greater persistence than the original substances, or if, following on the primary process, other processes which are subjected to several negative accelerations occur, then there will be no marked tendency to detonation. But if, on the other hand, the primary reaction leads to the stable end products of the reaction, or if these products undergo other reactions with no retardations or at most only one retardation, then the explosive is one which has a high sensitiveness and a great tendency to detonation. Pressure is the factor which removes the retardation to the reactions, since it causes the atoms, molecules, and atomic groupings which have affinities for one another to come closer together.

J. F. S.

The Autoracemisation of Potassium Chromioxalate. ERIC KEIGHTLEY RIDEAL and WILLIAM THOMAS (T., 1922, 121, 196—202).

Neutral Salt Action, at Higher Salt Concentrations; the Velocity of Hydrolysis of Ethyl Acetate, and the Hydrogen-ion Activity of the Catalyst. GÖSTA ÅKERLÖF (*Z. physikal. Chem.*, 1921, 98, 260—292).—The velocity of hydrolysis of ethyl acetate by 0.05*N*-solutions of hydrochloric, sulphuric, and nitric acid at 25° has been determined in the presence of various concentrations of the chlorides of ammonium, potassium, sodium, magnesium, calcium, and barium, the nitrates of ammonium, potassium, sodium, magnesium, and calcium, and the sulphates of ammonium, potassium, sodium, and magnesium. The hydrogen-ion activity of the catalyst in the various reaction mixtures has been determined by means of *E.M.F.* measurements. An attempt is made to ascertain the cause of the neutral salt action on the velocity of hydrolysis at constant acid concentration by combining the velocity measurements with the hydrogen-ion activity. A relationship between these two quantities is furnished by the formula $K = C\alpha \cdot f(n)\sqrt[3]{\alpha}$, in which K is the reaction velocity, n the acid titre, and α the hydrogen-ion activity. The formula holds for all concentrations of catalyst, and is not influenced by the nature or concentration of the added salt, as far as the present experiments go. This formula is probably also true for the inversion of sucrose and allied reactions. The foregoing shows that the hypothesis of the catalytic activity of the non-ionised molecule is incorrect. The salt molecules have no activity of their own, but they are the cause of the change of activity of the catalyst. Exactly how the neutral salt changes the activity of the catalyst is not known with certainty, but the most probable hypothesis is that the water sheath of the hydrogen ion is changed by the addition of neutral salts. The concentration of the hydrogen ion remains constant whilst the activity changes. According to this view, the hydration numbers do not represent definite quantities, but the ions control all the water molecules in the solution. The various neutral salt actions are controlled by the various forces with which the different ions attract the water molecules. J. F. S.

Velocity of Hydrolysis of Acetals. ANTON SKRABAL and ANTON SCHIFFRER (*Z. physikal. Chem.*, 1921, 99, 290—313).—The velocity of hydrolysis of methylene dimethyl ether, methylene diethyl ether, ethylidene diethyl ether, methylene diacetate, and ethylidene diacetate has been investigated. The reactions were carried out with hydrochloric acid of various concentrations, (0.001—0.3*N*), and also in alkaline solution. It is shown that the measured velocity constants represent the velocity of hydrolysis of the first ether group; the removal of the second group is much more rapid, and is regarded as a water hydrolysis. The velocity constants vary very much for the different cases, thus for the acid hydrolysis the value of k_a is 70,000 for methylene diethyl

ether, whilst for ethylidene diacetate it is 0.00650, whilst in the case of alkaline hydrolysis the value of k is 1,100 for methylene diacetate and 96 for ethylidene diacetate. In the case of the hydrolysis of esters it is shown that the velocity depends largely on the nature of the acid but very little on the nature of the alcohol. These rules are established from the published data of a number of authors.

J. F. S.

Velocity of Hydrolysis of Sucrose. R. H. CLARK (*J. Amer. Chem. Soc.*, 1921, 43, 1759—1764).—The velocity of hydrolysis of sucrose by hydrochloric, hydrobromic, sulphuric, or nitric acid has been determined at 25°. The experiments were effected in the presence of acids of varying concentration and in the presence of potassium nitrate and dextrose. It is shown that the addition of a neutral substance, to keep the concentration of the water constant while varying the concentration of the acid, has no appreciable effect in producing a numerical proportionality between the quantity of hydrogen ions present and the inversion velocity of sucrose. The increase in the velocity of hydrolysis of sucrose produced by a strong acid in the presence of a neutral salt of that acid over the velocity produced by the acid alone can only to a small extent be attributed to the fact that the volume of the reaction is reduced by the presence of the salt. On the assumption that both the dissociated and undissociated forms of an acid are catalytically active in sucrose inversion, the values of K_i for the hydrogen ion from hydrochloric, hydrobromic, and nitric acids have the same value ($K_i=0.234$); the corresponding value for sulphuric acid is considerably lower ($K_i=0.144$).

J. F. S.

Affinity Dimensions of Weak Acids and Bases in Alcoholic Solution and the Alcoholysis of their Salts. HEINRICH GOLDSCHMIDT, CARL GÖRBITZ, HAAKON HOUGEN, and KRISTIAN PAHLE (*Z. physikal. Chem.*, 1921, 99, 116—154).—The theory of the alcoholysis of salts of weak acids with bases of the aniline type in anhydrous alcohol and in alcohol containing water is developed. A method of determining the degree of alcoholysis based on electrical conductivity measurements is described. The electrical conductivity of a series of sodium, aniline, and ammonium salts in pure and in aqueous alcohol has been determined. The affinity dimension of several organic acids in alcohol solution has been determined. The alcoholytic constant of eleven aniline salts has been measured and from the results the affinity constant K_a , corresponding with the equilibrium $(\text{Aniline} \times \text{H}^+)/\text{Aniline ion}$, has been calculated. The same quantity has been determined for the three toluidines and mono- and di-methyl anilines. The influence of water and the alcoholytic constants has been investigated.

J. F. S.

The Rate of Solution of Iron in Dilute Sulphuric Acid both when Stationary and under Rotation. JOHN ALBERT NEWTON FRIEND and JOHN HORACE DENNETT (*T.*, 1922, 121, 41—44).

Theory of Catalysis in Homogeneous Gas Reactions. KARL F. HERZFELD (*Z. physikal. Chem.*, 1921, 98, 161—174).—A theoretical paper in which an attempt is made to answer the question, "Why does a reaction proceed more rapidly through an intermediate product than by the direct action?" It is shown that since at equal concentration the slowest of all the partial reactions is determinative of the velocity of the whole process, this slowest reaction must proceed more rapidly than the original reaction when catalysis occurs. After a short review of the previously published theoretical work on gas reactions, the author considers such reactions in a way which assumes the existence of a few free atoms. Then for catalysis it is necessary that at not too small concentrations of the catalyst the heat of activation of the intermediate product (that is, practically its heat of formation from atoms) is markedly smaller than that of the starting material; in the best case about one-half. This gives then an upper limit for the acceleration which at not too high concentrations may be reached by a given catalyst. Examples on which to test the hypothesis are not known.

J. F. S.

Catalytic Oxidation of Carbon Monoxide. T. H. ROGERS, C. S. PIGGOT, W. H. BAHLKE, and J. M. JENNINGS (*J. Amer. Chem. Soc.*, 1921, 43, 1973—1982).—The present work was undertaken with the object of finding a catalyst which will completely oxidise carbon monoxide at ordinary temperatures. A number of such catalysts have been prepared which cause rapid and complete oxidation. The chief of the catalysts obtained is a specially prepared manganese dioxide on which is precipitated the oxide of silver or copper or both. The silver or copper is best precipitated as the carbonate and subsequently hydrolysed to the hydroxide. The presence of water vapour limits the life of these catalysts. The decomposition temperature of silver oxide when simultaneously precipitated with calcium hydroxide is considerably lower than that of silver oxide alone.

J. F. S.

Catalytic Oxidation of Carbon Monoxide at Ordinary Temperatures. DAVID R. MERRILL and CHARLES C. SCALIONE (*J. Amer. Chem. Soc.*, 1921, 43, 1982—2002).—A number of catalysts for use in protective masks against carbon monoxide are described. The most successful are mixtures containing manganese dioxide and basic copper carbonate and manganese dioxide, copper oxide, cobaltic oxide, and silver oxide, respectively. These mixtures oxidise carbon monoxide rapidly and at ordinary temperatures, and, when protected by a drying agent such as calcium chloride, are suitable for use in protective masks against carbon monoxide of all concentrations below those in which an oxygen helmet would be necessary. The factors influencing activity, such as the constituents of the mixtures, the conditions of precipitation, the mechanical treatment of the precipitate, and method of drying, are discussed and a description is given of the efficiency of the mixtures under various conditions.

J. F. S.

Oxidation Catalysis. II. L. KARCZAG (*Biochem. Z.*, 1921, 119, 16—22).—Experiments with dyes at high dilutions show that the ferrous ion catalyses oxidation with hydrogen peroxide much faster than the ferric ion. The order of mixing is of influence, the controlling factor being whether the ferrous ion is first oxidised to ferric or not.

H. K.

Electronic Theory of the Interior of the Atom. A. D. FOKKER (*Arch. Néerland.*, 1921, 5, [iiia], 193—242).—A mathematical paper in which it is shown that the application of the electronic theory to the mutual actions of the electrons inside the atom gives results which are in accordance with experimental observations.

J. F. S.

Theory of Allotropy. MAURICE COPISAROW (*J. Amer. Chem. Soc.*, 1921, 43, 1870—1888).—Allotropy is defined as the capacity of an element to exist in forms differing in the mode of their intramolecular linking. It is therefore a function of the valency, but it does not imply isomerism or polymerism, since allotropes need not contain the same or a multiple number of atoms in their respective molecules. The number of possible forms in which an element can exist on this basis is indicated for the elements of all valencies. It is theoretically possible for an element to have in certain cases more than one molecular form corresponding with each mode of linking. Regarding allotropes as the most chemically and physically distinct forms of an element, it follows that several molecular forms, each containing a different number of atoms, but all having the power of free rotation, will differ among themselves to a less extent than when compared with a molecular structure of the same element in which all atoms are rigidly fixed. Thus it follows that valency and the saturation or fixation of atoms, and not the actual number of atoms, play the predominant part in the determination of allotropes. In this light, allotropy becomes the capacity of an element to exist in forms differing in the mode of their intramolecular linking. Molecular forms differing in the distribution of their intramolecular linking and in the number of atoms, but all belonging to one type of linking can be termed *allotropoids*. The allotropy of carbon, phosphorus, iron, and nickel are considered from the point of view of the hypothesis.

J. F. S.

Types of Valency. IRVING LANGMUIR (*Science*, 1921, 54, 59—67; cf. *ibid.*, 1921, 53, 290; *J. Ind. Eng. Chem.*, 1920, 12, 386, and A., 1919, ii, 328; also Rydberg, A., 1915, ii, 94).—Recognition of three distinct types of valency, (a) positive valency (the number of electrons an atom can give up), (b) negative valency (the number of electrons an atom can take up), and (c) co-valency (the number of pairs of electrons an atom can share with its neighbours), as well as the numerical values, for most of the elements, of each type are shown by a simpler method than that hitherto employed to be derived from certain postulates regarding the structure of atoms, and the relationships between the different types of valency are thereby further clarified. On the basis of the

Rutherford atom, only three postulates, consistent with those previously proposed, are necessary: (a) The electrons in atoms tend to surround the nucleus in successive layers containing 2, 8, 8, 18, 18, and 32 electrons respectively (the extra electrons, if any, remaining in the outside layer as an incomplete layer or "sheath"). (b) Two atoms may be coupled together by one or more duplets (stable pairs of electrons forming the first complete layers) held in common by the completed sheaths of the atoms. A group of neutral atoms interacting by transfer of electrons gives rise to the conception of electrovalency, including positive and negative valency differing only in algebraic sign, whilst interaction by the sharing of duplets corresponds with co-valency. It is shown that the sum of the electrovalencies and co-valencies for all the atoms in any complete compound is zero, a complete compound being defined as one in which all the atoms possess complete layers of electrons. (c) The residual charge on each atom and on each group of atoms tends to a minimum. The term "residual charge" is employed to express the total charge of an atom or aggregate of proximate atoms regardless of sign. It is pointed out, however, that the first and third postulates are often in conflict, and that in such cases the tendency of the first may prevail against that of the third.

Chemical compounds are therefore classified according to the types of valency exhibited by their atoms as follows: (1) complete compounds, including (a) compounds without co-valency, such as certain simple metallic salts, volatile halogen compounds, silicates, and most minerals, and (b) compounds without electropositive atoms, such as organic compounds; (2) incomplete compounds, including metallic substances, and compounds such as zinc oxide, ferrous-ferric oxide, lead sulphide, and cupric oxide, which contain both electropositive and electronegative atoms; (3) exceptional cases, including nitrogen, carbon monoxide, nitric oxide, and the cyanogen radicle, the structure of which is not accounted for by this theory. These substances may have a single octet structure or possibly a triple bond structure. Other exceptions are boron hydride and certain other compounds forming double molecules.

A. A. E.

The Expression of the Octet Theory of Valence in Structural Formulæ. GRANVILLE A. PERKINS (*Philippine J. Sci.*, 1921, 19, 1-22).—A system for writing structural formulæ is developed, based on the octet theory of valency as presented by Langmuir.

Electronegative valence, the physical interpretation of which is vacancy for one more electron in the outer shell of the atom, is represented by a line in the usual way. Electropositive valence, on the other hand, is variable and its maximum is represented by the total number of electrons in the shell. It cannot be represented by "unsatisfied bonds" as it never causes direct union between atoms, and is expressed as follows, for example, Na^1 , Ca^{II} . The neutral atoms having electronegative valence may satisfy that valence by acquiring an electron, that is, becoming

a negative ion; this electron may be obtained from the shell of an atom showing positive valence tendencies, and it is then left as a positive ion. These two oppositely charged ions may then form a stable compound by electrostatic attraction. Such a union is termed a salt-forming union and is expressed thus: $\text{Na}^+ \dots \text{Cl}^-$, or $\text{Ca}^{++} :::: \text{O}^{--}$. On the other hand, two atoms may be held together by the shell-completing forces of both atoms acting on one or more pairs of electrons, which are then shared between them. Each shell thus fills one or more electron vacancies by sharing electrons, and thus "satisfies" one or more "bonds." This is expressed in the usual way: $\text{H}-\text{H}$, $\text{H}-\text{O}-\text{H}$, etc., and is termed a direct union. All valency relations cannot, however, be expressed in terms of either of the above unions, and a special feature of the system is the introduction of the conception of a "borrowing direct union," for cases in which both the shared electrons are supplied by the same atom. In this case, the borrowing atom fills two vacancies in its shell, and the lending atom neither gains nor loses electrons; but the union is essentially polar, since the originally neutral borrowing atom has acquired at least an interest in two negative electrons and therefore becomes a negative pole. This relationship is expressed thus: $\overset{\text{H}}{\text{H}} > \text{O} \propto \text{O}, \underset{\text{O}}{\text{O}} = \text{S} \propto \text{O}$,
8

indicating that the O atom is bound to S, or to O, by virtue of having its two electron vacancies filled by borrowing an interest in two of the shell electrons of the other oxygen atom in hydrogen peroxide, and in the sulphur atom of sulphur trioxide. The mode of application of the system to all the known elements is indicated in the original paper.

G. F. M.

The Thermos Flask in the Chemical Laboratory. CLAUDE W. BOURLET and WALTER THOMAS (*Chem. News*, **123**, 336).—The thermos flask can be utilised advantageously in the laboratory in numerous instances where it is desired to maintain substances above the normal temperature, as, for example, in reactions between liquids, or liquids and solids at elevated temperatures such as hot oxidation by permanganate and other similar reactions which are not strongly exothermic. Oil and water emulsions and similar mixtures are conveniently "split" by warming and keeping overnight in a thermos flask, and flocculent or colloidal precipitates can be dealt with in a similar manner. In carrying out certain colour tests, for example, the Halphen reaction, and reactions in media which gelatinise or become viscous at ordinary temperatures, a water-bath can with great advantage be replaced by the vacuum flask.

G. F. M.

Water Pump. JOHANNES WETZEL (*Chem. Ztg.*, 1921, **45**, 1122).—A water pump of improved efficiency is described; particular attention is directed to the relative diameters of the injector tube and outlet tube and to the central position of the jet as regards the outlet tube.

W. P. S.

Inorganic Chemistry.

Separation of the Element Chlorine into Isotopes (Isotopic Elements). The Heavy Fraction from the Diffusion.

WILLIAM D. HARKINS and ANSON HAYES (*J. Amer. Chem. Soc.*, 1921, **43**, 1803—1825).—By means of elaborate diffusion a definite increase in the atomic weight of chlorine has been obtained. The increase in atomic weight amounts in different experiments to from slightly less than to considerably more than one part in a thousand (1 in 645). The diffusion experiments were effected with hydrogen chloride, and a considerable amount of the isotopic acid has been obtained. The separation of isotopes by diffusion is discussed in terms of the Rayleigh diffusion equation, which applies to the diffusion into a vacuum. It is shown that for a high efficiency the pressure of the gas on both sides of the diffusion wall should be low, first to secure good mixing, and secondly to insure that the passage through the porous partition shall be entirely molecular. A rapid and precise method for the determination of the atomic weight of isotopic chlorine has been devised. It is found that in the separation of isotopes the percentages of the different isotopes present are as important factors as the atomic weight differences. Thus, contrary to what has been supposed, it is shown, even aside from the greater difficulties involved in obtaining and handling neon, that it is easier to produce a small increase in the atomic weight of chlorine (of the magnitude of 0.05 unit) than to produce the same increase in the atomic weight of neon. J. F. S.

Viscosities of the Hydrogen Haloids. H. HARLE (*Proc. Roy. Soc.*, 1922, [A], **100**, 429—440).—The viscosity of hydrogen chloride, bromide, and iodide has been determined by the method of transpiration through a capillary tube. Measurements were made at temperatures in the neighbourhood of 20° and 100°, and from the results the viscosity is calculated for 0° and 100° respectively. The following values in *C.G.S.* units are recorded: hydrogen chloride, $\eta_0 = 1.332 \times 10^{-4}$; $\eta_{100} = 1.837 \times 10^{-4}$; hydrogen bromide, $\eta_0 = 1.710 \times 10^{-4}$; $\eta_{100} = 2.365 \times 10^{-4}$; hydrogen iodide, $\eta_0 = 1.731 \times 10^{-4}$, $\eta_{100} = 2.403 \times 10^{-4}$. The values of *C*, the Sutherland constant, are 357, 375, and 390 for the three gases respectively. J. F. S.

Numerical Revision of the Data referring to the Density of Gaseous Hydrogen Bromide: Atomic Weight of Bromine.

E. MOLES (*J. Chim. Physique*, 1921, **19**, 135—138).—The molecular weight of gaseous hydrogen bromide with reference to oxygen has been calculated to be 80.944, and from this the atomic weight of bromine is 79.936. If, however, the value for the weight of the normal litre of oxygen recently published (cf. following abstract) is used the atomic weight of bromine becomes 79.927. J. F. S.

Critical Study of the Modern Value of the Density of Gaseous Oxygen. E. MOLES (*J. Chim. Physique*, 1921, 19, 100—120).—A discussion of the modern results for the density of oxygen leads to the value $L_0 = 1.42891 \pm 0.00003$ as the most probable value for the weight of the normal litre of oxygen. The value is deduced from 162 measurements made by nine workers using oxygen obtained from three different sources and by three essentially different methods. If only the most recent measurements by weighing a globe of the gas are considered the mean is $L_0 = 1.42894$. Up to this the value accepted was 1.42905, which can no longer be maintained, and although the difference is only 1 in 10,000, yet it cannot be neglected in the calculation of molecular weights. It is proposed that the value 1.42891 be accepted as the standard, and this in some cases may be rounded to 1.4289. If this value is calculated to the conventional value of g , $g = 980.665$ C.G.S. units then the value of L_{0N} becomes 1.42897, which may be rounded to 1.4290. J. F. S.

Solubility. VII. Solubility Relations of Rhombic Sulphur. JOEL H. HILDEBRAND and CLARENCE A. JENKS (*J. Amer. Chem. Soc.*, 1921, 43, 2172—2177).—The solubility of rhombic sulphur has been determined in carbon tetrachloride, benzene, toluene, *m*-xylene, heptane, and ethylene dichloride at 0°, 25°, 35°, 45°, and 54°. The results are expressed in molecules of S_8 per 100 molecules of solvent and in grams per 100 grams of solvent. The following values in grams per 100 grams of solvent are recorded: Carbon tetrachloride, 0°, 0.339; 25°, 0.831; 35°, 1.155; 45°, 1.564; and 54°, 2.008; heptane, 0°, 0.124; 25°, 0.362; 35°, 0.512; 45°, 0.698; and 54°, 0.926; toluene, 0°, 0.897; 25°, 2.018; 35°, 2.722; 45°, 3.620; and 54°, 4.85; *m*-xylene, 25°, 1.969; 45°, 3.604; benzene, 25°, 2.074; 54°, 5.165; ethylene dichloride, 25°, 0.826; 40°, 1.380; 79°, 5.43; and 97.5°, 9.97. It is shown that, with the exception of certain minor discrepancies, the solubilities of sulphur accord well with the internal pressure relations of the substances involved. J. F. S.

The Constitution of Selenium. H. PÉLABON (*Compt. rend.*, 1921, 173, 1466—1468).—The grey selenium previously described (cf. A., 1921, ii, 533), the specific resistance of which may vary from a few ohms to several million ohms, is now shown to consist of two modifications. The α -modification has a very high resistance and is prepared by heating selenium at a temperature only just above its melting point and then allowing it to cool slowly. The specific resistance of this modification diminishes at first very rapidly with rise in temperature and then more slowly. At 200°, the resistance is about 70,000 ohms, and at 218° the substance melts, the melting being accompanied by a sudden rise in resistance. The β -modification has a very small specific resistance. It is obtained by maintaining the molten selenium at a temperature near its boiling point for some time and then allowing it to cool. The β -modification is not stable at low temperatures, and it is

readily changed into the α -form by oscillations of its temperature between 15° and 200°.

W. G.

Influence of Freezing on Colloidal Selenium. II. A. GUTBIER, FR. HEINRICH, and J. HUBER (*Kolloid Z.*, 1921, **29**, 287—293; cf. A., 1921, ii, 693).—A continuation of work previously described (*loc. cit.*) on the effect of freezing on selenium sols. Three new series of experiments are now described in which it is shown that the sensitiveness of undialysed selenium sols, prepared by means of sulphur dioxide, towards cold increases with the time during which they are frozen. The system may be frozen for short periods and will then melt to form a typical colloid, but more prolonged freezing increases the tendency to coagulate to irreversible gels on melting, and very prolonged freezing will even cause coagulation to take place in the solid. In this respect, there is no fundamental difference between dialysed and undialysed sols, but the dialysed system is somewhat more sensitive. The reducing action of sulphur dioxide on solutions of selenious acid is retarded by freezing the mixture. If the mass is kept frozen for a long time the colloid is precipitated in the ice and thereby the reduction process is accelerated. The precipitation appears to exercise a nucleus action, by which the reduction in the solid mass is accelerated, so that eventually more selenium is precipitated in the frozen mass than would be produced in a solution in the same time. In general, the coagulation in the frozen sol commences where the sol is in contact with the air, as is shown by the formation of a coloured ring on the upper surface of the ice. When test-tubes are used for the experiments, the curved lower portion of the tube always shows an increased concentration of selenium. The form of the vessel in which the freezing takes place is shown to exert an influence on the coagulation. When a sol is poured on ice and frozen it is found that the disperse phase diffuses into the ice. It has been found scarcely possible to freeze selenium sols in capillary tubes.

J. F. S.

Metallurgy of Tellurium by the Wet Way. PIERRE HULOT (*Bull. Soc. chim.*, 1921, [iv], **29**, 1070—1071).—The method of reduction of potassium anhydrotellurate by zinc and hydrochloric acid with the precipitation of metallic tellurium (cf. A., 1920, ii, 174) is modified by using aluminium and sodium or potassium hydroxide for the production of nascent hydrogen, with the result that the reduction is completed in one hour instead of ten. The aluminium must be pure and free from copper.

W. G.

Manganese in the Catalytic Oxidation of Ammonia. CHARLES SNOWDEN PIGGOT (*J. Amer. Chem. Soc.*, 1921, **43**, 2034—2045).—The action of manganese dioxide alone and when mixed with copper oxide or silver oxide, and various alloys of manganese, copper, silver, iron, and silica as catalysts in the oxidation of ammonia have been investigated. A mixture of manganese dioxide with 40% of copper oxide at 800° is shown to have an efficiency of more than 90%. All the substances mentioned catalyse the

oxidation of ammonia with various efficiencies which with suitable conditions are about 50%. The physical structure of the oxide catalyst is of great importance, extremely fine subdivision giving a large adsorbing surface with very small pores being conducive to greater activity.

J. F. S.

Attempt to Prepare Nitro-nitrogen Trichloride. II. Behaviour of Mixtures of Nitrogen and Chlorine in a Flaming Arc. WILLIAM ALBERT NOYES (*J. Amer. Chem. Soc.*, 1921, 43, 1774—1782; cf. A., 1913, ii, 584).—Attempts to prepare nitro-nitrogen trichloride or nitrogen trichloride by passing mixtures of nitrogen and chlorine through a flaming arc were entirely unsuccessful. Passing the same gases through a powerful ozoniser had the same result and no action at all occurred with Strutt's active nitrogen. The results, although negative, confirm Lewis's hypothesis that electrons are held jointly by two atoms rather than the view that electrons are transferred from one atom to another when atoms combine.

J. F. S.

Vapour Pressures of Aqueous Solutions of Nitric Acid. WILLIAM C. SPROESSER and GUY B. TAYLOR (*J. Amer. Chem. Soc.*, 1921, 43, 1782—1787).—The total and partial vapour pressures of nitric acid solutions of the concentrations 20%, 40%, 56%, 68%, and 80% by weight have been measured at temperatures 0°, 35°, 50°, 65°, and 80°. From the data the vapour pressure for each 10% increase in concentration and each 10° rise in temperature has been calculated and tables drawn up. Attempts to measure the vapour pressure of 90% acid were unsuccessful owing to the decomposition of the acid at all temperatures except 0°.

J. F. S.

Reducing Actions of Arsenious Acid. MORITZ KOHN (*Monatsh.*, 1921, 42, 221—226).—When copper sulphate is heated with ammonia solution and arsenious anhydride in a sealed tube in a boiling water-bath, it undergoes reduction to cuprous salt with formation of arsenic acid. After the resulting cuprous solution has been oxidised to the cupric condition by atmospheric oxygen, estimation of the arsenic acid reveals more of the latter than corresponds with the equation, $2\text{Cu}^{++} + \text{AsO}_3''' + 2\text{OH}' = \text{H}_2\text{O} + 2\text{Cu}' + \text{AsO}_4'''$. Evidently oxidation of the ammoniacal cuprous solution to the cupric stage activates the atmospheric oxygen for the oxidation of the residual unchanged arsenious acid. Depression of the concentration of hydroxyl ions by addition of ammonium salts retards the reduction of cupric to cuprous salt.

T. H. P.

Catalysis in the Interaction of Carbon with Steam and with Carbon Dioxide. HUGH STOTT TAYLOR and HARVEY A. NEVILLE (*J. Amer. Chem. Soc.*, 1921, 43, 2055—2071).—The effect of potassium carbonate, sodium carbonate, lithium carbonate, barium carbonate, calcium carbonate, sodium chloride, ferric oxide, copper, sodium silicate, borax, and nickel as catalysts on the interaction of steam on carbon has been investigated, using various forms of

carbon. It has been discovered that the interaction of carbon and carbon dioxide is likewise accelerated by the presence of the same materials, and a striking parallelism between the catalysts for the two reactions has been shown. The probable mechanism of the steam-carbon reaction has been investigated. It has been shown that good catalysts for the water-gas reaction, for example, iron oxide, are ineffective in the steam-carbon and carbon dioxide-carbon reactions. Catalysis of the water-gas reaction thus being excluded from consideration, it has been shown that the acceleration of the reaction $C + CO_2 = 2CO$, and therefore the acceleration of the reaction $C + 2H_2O = CO_2 + 2H_2$, may be ascribed to increased adsorption of carbon dioxide by the carbon surfaces in presence of active catalytic agents. Adsorption measurements confirm this view. The results have been considered in connexion with the suggested existence of a surface complex, C_xO_y , and have been found to agree with this conception. J. F. S.

A Modification of Silicon Soluble in Hydrofluoric Acid.

WILHELM MANCHOT (*Ber.*, 1921, **54**, [B], 3107—3111).—Moissan and Siemens (*A.*, 1904, ii, 560) have isolated specimens of silicon from a silver regulus which suffer loss of weight (up to 99%) when treated with hydrofluoric acid, and subsequently Lebeau (*A.*, 1906, ii, 168) has brought forward evidence to show that a similar modification is present in copper-silicon alloys rich in the latter. The substance, however, does not appear to have been investigated fully.

Considerable difficulty is experienced in preparing silicon which is completely resistant towards hydrofluoric acid; the product obtained by Wöhler's method requires frequently repeated treatment with the acid before a stable specimen is secured. It is shown that an appreciable gain in weight occurs when the latter is melted with silver in an electric furnace in an atmosphere of air, oxygen, or carbon monoxide, but this is in itself insufficient to account for the subsequent solubility of the silicon. The latter, however, is only produced in the soluble condition when the regulus is suddenly cooled. The silver may be replaced by aluminium. It dissolves in hydrofluoric acid with evolution of hydrogen.

It is remarkable that Moissan and Siemens describe their product as exactly resembling crystalline silicon, whereas the author's specimens are dark to pale brown and amorphous; they leave a pale brown powder after being treated with hydrofluoric acid.

H. W.

Organogels of Silicic Acid. B. S. NEUHAUSEN and W. A. PATRICK (*J. Amer. Chem. Soc.*, 1921, **43**, 1844—1846).—Hydrogels of silicic acid have been repeatedly soaked in alcohol, acetone, and benzene for long periods, dried in a vacuum of 5 mm. at 80°, 120°, and 270° successively for periods of one hour, and analysed. The dried products contain: alcogel, 4.23% water, 3.90% alcohol; acetone gel, 4.77% water, 3.75% acetone; benzene gel, 4.31% water, 3.82% benzene. Hence it follows, in opposition to Graham's

statement, that the water of a hydrogel of silicic acid cannot be entirely replaced by a second solvent. It has been previously shown that heating silicic acid gels in a vacuum at 300° for six hours does not reduce the water content below 4·8%, a quantity which is very near the amount of water left in the organo-gel above, and indicates that this water is very firmly combined in the gel.

J. F. S.

Relative Densities of Alkali Metal Amalgams and Mercury.

II. EDGAR C. BAIN and JAMES R. WITHROW (*J. Physical Chem.*, 1921, 25, 535—544; cf. A., 1916, ii, 431).—The methods of formation and the relative densities of liquid and solid amalgams of sodium, potassium, ammonium, and calcium have been investigated. It is shown that solid amalgams float on the mother-liquor and do not sink as stated by Kerp (A., 1898, ii, 516). The method used by Kerp (*loc. cit.*) produces sodium and potassium amalgams, which are like the amalgams produced by other methods inasmuch as the solid is lighter than the liquid. Oily and pasty amalgams are due to fine crystals incorporated in the liquid. By electrolysis of solutions of potassium chloride, amalgams with concentrations up to 2·31% of potassium were obtained. No difficulties, except those of collection and preservation, were met with in the production of ammonium amalgams by the electrolysis of solutions of ammonium chloride, but crystalline amalgams cannot be obtained in this way. Electrolysis of an acidified solution of calcium acetate is not a satisfactory method for the production of calcium amalgams, but it does produce a dilute calcium amalgam.

J. F. S.

The Production of Potassium Hydrogen Sulphate from Ammonium Hydrogen Sulphate and Potassium Sulphate. W. DOMINIK (*Przemysl Chem.*, 1921, 5, 10—15, 37—40, 63—67).—Reaction in the presence of steam superheated at 200—400° takes place according to the equation $M_2SO_4 + NH_4HSO_4 = 2MHSO_4 + NH_3$; lack of steam results in the production of pyrosulphates. In the above reaction, $K = [MHSO_4]^2 / [M_2SO_4] \cdot [NH_4HSO_4]$ increases with the temperature according to a logarithmic function. Theoretical consideration is given to the use of a mixture of sodium and potassium sulphates in the reaction. CHEMICAL ABSTRACTS.

Crystal Structures of Sodium Chlorate and Sodium Bromate. ROSCOE G. DICKENSON and ELBRIDGE A. GOODHUE (*J. Amer. Chem. Soc.*, 1921, 43, 2045—2055).—A large amount of X-ray spectrometer data for sodium chlorate and bromate has been obtained and tabulated. In agreement with other observers, it has been found that the nature of the crystal surface has a considerable effect on both the absolute and relative intensities of reflection. The ease with which reflections from one face may be mistaken for those from another and the necessary precautions to avoid this have been pointed out. It is shown that the atoms in sodium chlorate and sodium bromate are very

probably arranged with the symmetry of the Schönflies space group T^4 , all oxygen atoms being equivalent. Sets of positions of the atoms in the unit structure, only slightly different for the chlorate and bromate, have been suggested. J. F. S.

Preparation and Study of the Rarer Alkali Bromates. Rubidium Bromate. HAROLD D. BUELL and C. R. McCROSKY (*J. Amer. Chem. Soc.*, 1921, **43**, 2031—2034).—Rubidium bromate was prepared by treating the pure carbonate with an excess of bromic acid and recrystallising the product (cf. A., 1920, ii, 688). The following figures are the solubilities in 100 grams of water at various temperatures: 25°, 2.93; 30°, 3.55; 35°, 4.28, and 40°, 5.08. The corresponding figures for caesium bromate are 25°, 3.66; 30°, 4.53; 35°, 5.32. The following melting points are recorded: potassium bromate, 405°; caesium bromate, 420°; rubidium bromate, 430°. Both rubidium and caesium bromates form small, cube-like crystals, which, however, belong to the hexagonal system; they have a refractive index between 2.144 and 2.22. J. F. S.

Vapour Pressure of some Salts. II. H. VON WARTENBERG and H. SCHULZ (*Z. Elektrochem.*, 1921, **27**, 568—573; cf. Albrecht and Wartenberg, *ibid.*, 162).—Using the method previously employed, the authors have determined the vapour pressures of lithium chloride, caesium chloride, rubidium chloride, lithium bromide, caesium bromide, rubidium bromide, sodium fluoride, potassium fluoride, lithium fluoride, caesium fluoride, rubidium fluoride, sodium iodide, caesium iodide, and rubidium iodide. The measurements were carried in most cases up to the boiling point of the salt in question. The following data are recorded, pressures being in atmospheres: lithium chloride, b. p. 1382°, m. p. 606°, $\log p = -37200/4.57T + 4.923$; caesium chloride, b. p. 1303°, m. p. 626°, $\log p = -37400/4.57T + 5.190$; rubidium chloride, b. p. 1383°, m. p. 717°, $\log p = -37800/4.57T + 4.998$; lithium bromide, b. p. 1310°, m. p. 549°, $\log p = -35600/4.57T + 5.109$; caesium bromide, b. p. 1300°, m. p. 627°, $\log p = -36750/4.57T + 5.113$; rubidium bromide, b. p. 1350°, m. p. 681°, $\log p = -36980/4.57T + 4.964$; sodium fluoride, b. p. 1695°, m. p. 988°, $\log p = -56600/4.57T + 6.299$; potassium fluoride, b. p. 1505°, m. p. 846°, $\log p = -41900/4.57T + 5.138$; lithium fluoride, b. p. 1676°, m. p. 842°, $\log p = -55100/4.57T + 6.190$; caesium fluoride, b. p. 1251°, m. p. 684°, $\log p = -34700/4.57T + 4.982$; rubidium fluoride, b. p. 1410°, m. p. 775°, $\log p = -40000/4.57T + 5.243$; sodium iodide, b. p. 1300°, $\log p = -37000/4.57T + 5.130$; lithium iodide, b. p. 1170°, $\log p = -40300/4.57T + 6.105$; caesium iodide, b. p. 1280°, $\log p = -36600/4.57T + 5.165$; and rubidium iodide, b. p. 1305°, $\log p = -37000/4.57T + 5.148$. The results show that the heats of formation of the haloids of potassium, rubidium, and caesium lie very close together, whilst that of the lithium salts is much higher and that of the sodium salts much lower. J. F. S.

Catalytic Influence of Foreign Oxides on the Decomposition of Silver Oxide, Mercuric Oxide, and Barium Peroxide.

JAMES KENDALL and FRANCIS J. FUCHS (*J. Amer. Chem. Soc.*, 1921, **43**, 2017—2031).—The effect of foreign oxides on the temperature and rate of decomposition of silver oxide, mercuric oxide, and barium peroxide under an oxygen pressure of one atmosphere has been experimentally investigated. In almost all the systems examined the added oxides (CuO , MnO_2 , Fe_2O_3 , CeO_2 , SiO_2 , CrO_3) induce a considerable change in the decomposition temperature. Most commonly, there is a marked lowering in this point; thus, quoting an extreme case, an equimolecular mixture of barium dioxide and cupric oxide has an oxygen equilibrium pressure of 1 atm. at 322° , a temperature which is approximately 500° below the decomposition temperature of pure barium dioxide. In a few systems a comparatively small rise in the decomposition temperature is indicated. In all cases, however, the rate of oxygen evolution is markedly increased. The decomposition temperatures recorded refer to true equilibrium conditions, concordant values being obtained with rising and falling temperature. The results are tentatively ascribed to the formation of unstable intermediate compounds between the two oxides present. The increased rates of decomposition may be referred to adsorption effects, but the large temperature changes point strongly to the actual participation of the added oxide in the reaction. Direct evidence was given in certain mixtures of the production of stable complexes.

J. F. S.

Physical Chemistry of the Oxides of Lead. III. Hydrated Lead Monoxide. SAMUEL GLASSTONE (*T.*, 1922, **121**, 58—66).

Reactions in Fused Salt Media. I. Basic Lead Chromates. J. F. G. HICKS (*J. Physical Chem.*, 1921, **25**, 545—560).—The reaction between lead monoxide and sodium chromate in fused sodium chloride and a 50% mixture of sodium and potassium nitrates has been investigated and the equilibrium diagram of the system $\text{PbO}-\text{PbCrO}_4$ constructed. It is shown that reactions in fused salts can be brought about in such a way as to yield products analogous to those prepared from the same initial substances in aqueous solution. The chief difference between these reactions and their analogues in aqueous solution lies in the smaller velocity of the former, probably due to the relative insolubility of the reacting substances in the fluxes as compared with water. There is in several cases a reaction between the flux and the dissolved substance, similar to hydrolysis. This analogy would appear to indicate that these solvolytic reactions are ionic, but the second phase of the reaction forming basic lead chromates points to a non-ionic reaction. It may be a purely molecular (additive) reaction, resulting in the formation of compounds of higher orders. Such a conclusion explains satisfactorily the slowness of the reactions, aside from the low solubility of the solutes in the fused salt medium. Whilst at least one basic lead chromate,

$\text{PbO}, \text{PbCrO}_4$, may be prepared by alkaline hydrolysis, it does not follow that the basic lead chromates formed in fused salt media are formed by an entirely analogous process. Two new basic chromates have been prepared by the present method which have not been obtained by the wet process; these have the formulæ, $2\text{PbO}, \text{PbCrO}_4$ and $3\text{PbO}, \text{PbCrO}_4$ respectively. A fourth compound, $\text{PbO}, 2\text{PbCrO}_4$ is readily prepared by the fusion process and has been stated to be formed by the wet process, but the author has been unable to prepare it by this method. The existence of the four basic lead chromates named is confirmed by the equilibrium diagram of the system $\text{PbO}-\text{PbCrO}_4$. Other basic lead chromates described in the literature are shown to be mixtures or solutions of the compounds named above in one another. Salts of ortho-chromic acid are shown not to exist; whilst the salt Pb_2CrO_5 may be the salt of the monohydrate of ordinary chromic acid, it could equally well be a true basic salt or a compound of a higher order, so far as the present work is concerned. At the temperature ($225-800^\circ$) the stable form of lead monoxide is yellow in colour. This, combining with yellow lead chromate yields red compounds, all of which point to compounds of a higher order. The red compound formed by alkaline hydrolysis of normal lead chromate could well be considered a basic salt, but the addition of lead oxide and lead chromate molecules to form a compound of higher order might as easily take place in aqueous solution as in the nitrate flux used to form a compound of very closely the same composition. The reddening of lead monoxide in the nitrate flux has been shown to be due to change in crystalline form, and not, as might be supposed, to the formation of red lead.

J. F. S.

Reduction of Copper Oxide by Hydrogen. ROBERT N. PEASE and HUGH STOTT TAYLOR (*J. Amer. Chem. Soc.*, 1921, **43**, 2179—2188).—An investigation of the characteristics of the reduction of cupric oxide by hydrogen and the effect on the reaction of adding metallic copper to the oxide and water vapour and oxygen to the hydrogen has been carried out. It is pointed out that the reaction is auto-catalytic, copper being the auto-catalyst. The reaction appears to take place at the copper-copper oxide interface. This is shown by the character of the reduction curve and the fact that the addition of metallic copper accelerates the reaction. It is shown that the presence of water vapour in the hydrogen markedly interferes with the formation of the original copper nuclei from which the reaction zone, that is the copper-copper oxide interface, spreads out; it does not markedly affect the subsequent reaction at the interface, however. The presence of oxygen in the hydrogen strongly inhibits the reaction at the interface, but in all probability has no marked effect on the primary reaction, that is, the formation of the original copper nuclei.

J. F. S.

The Oxidising Properties of Sulphur Dioxide. III. Copper Chlorides. WILLIAM WARDLAW and FREDERICK WILLIAM PINKARD (*T.*, 1922, **121**, 210—221).

Behaviour of Ammoniacal and Alkaline Copper Solutions.

MORITZ KOHN (*Monatsh.*, 1921, **42**, 83—87).—When arsenic is heated in a sealed tube with ammoniacal copper solution, the cupric compound undergoes reduction, first to cuprous salt and subsequently to copper, $3\text{Cu}^{++} + \text{As} + 3\text{OH}' = \text{AsO}_3''' + 3\text{Cu}' + 3\text{H}'$ and $3\text{Cu}' + \text{As} + 3\text{OH}' = \text{AsO}_3''' + 3\text{Cu} + 3\text{H}'$; the copper liberated unites with the excess of arsenic to form greyish-black copper arsenide. Exactly similar changes occur when antimony is heated with ammoniacal copper solutions containing tartaric acid, the solid deposited being then reddish-black and containing the copper and the excess of antimony. With bismuth and ammoniacal copper solutions containing tartaric acid, rapid action occurs, but the reduction proceeds only as far as the cuprous compound. Alkaline solutions of copper salts, such as Fehling's solution or an aqueous solution containing copper sulphate, glycerol and potassium hydroxide, are rapidly reduced by arsenic, antimony, or bismuth, with separation of copper, $2\text{As}(2\text{Sb}) + 3\text{Cu}^{++} + 6\text{OH}' = 2\text{AsO}_3(\text{SbO}_3)''' + 6\text{H}' + 3\text{Cu}$ or $2\text{Bi} + 3\text{Cu}^{++} = 2\text{Bi}''' + 3\text{Cu}$. T. H. P.

Separation of the Isotopes of Mercury. J. N. BRÖNSTED and G. VON HEVESY (*Z. physikal. Chem.*, 1921, **99**, 189—206, and *Phil. Mag.*, 1922, **43**, [vi], 31—49).—A partial separation of the isotopes of mercury has been achieved by two processes. (1) Evaporation method (ideal distillation) based on the difference in the velocities of evaporation of the isotopes. The distillate was found to be richer and the remainder poorer in the lighter isotope than the original substance. (2) Effusion method. A fraction of the mercury vapour penetrates through narrow openings into a condensation chamber where the lighter isotope is found in a relatively larger amount than in ordinary mercury. The results of the experiments agree with the hypothesis, according to which the evaporation, as well as the effusion velocity of the isotopes, is inversely proportional to the square root of their molecular weights; they are further in conformity with Aston's results, obtained by means of mass spectrographic observations. The partial separation achieved was proved by measurements of density. The density difference found between the heaviest and lightest mercury amounts to 0.49%, corresponding with a difference of 0.1 unit in the atomic weight of mercury. J. F. S.

Physico-chemical Analysis of Aluminium Oxy-salts and Aluminium Oxide Sols. MONA ADOLF, WOLFGANG PAULI, [with FRANZ JANDRASCHITSCH] (*Kolloid Z.*, 1921, **29**, 281—287; cf. A., 1917, ii, 563; 1921, ii, 700).—The composition and nature of aluminium oxy-chloride sols have been investigated by means of measurements of the concentration of hydrogen and chlorine ions, the total chlorine concentration, and the electrical conductivity. A number of transport determinations have also been made. It is shown that it is impossible to remove all the chlorine from the products of hydrolysis of aluminium chloride by washing. A quantity of chlorine, which is greatly in excess of that contained

in the hydrochloric acid required for peptisation, always remains. On boiling well washed aluminium hydroxide with dilute hydrochloric acid, sols of various compositions are obtained up to a limiting composition represented by the formula $[\text{Al}(\text{OH})_3]_2\text{AlOCl}$. Of these two have been investigated. Aluminium oxy-dichloride $\text{AlCl}_2\cdot\text{OH}$ behaves as a ternary electrolyte and on dilution is strongly dissociated, the process being complete at a dilution $0\cdot00106N$. This compound shows a remarkably small hydrolysis, which amounts to $0\cdot1\%$ at $0\cdot068N$, and in a $0\cdot00106N$ solution is only $0\cdot25\%$. The compound aluminyl monochloride, $\text{Al}(\text{OH})_2\text{Cl}$ or AlOCl , behaves as a binary electrolyte. The conductivity data indicate that a complex ionisation occurs in this case of the type $\text{Al}(\text{OH})_2\text{Cl}|\text{AlO}$, indicating a compound in which one aluminium atom of the complex acts as the central atom of a negative complex, whilst the other furnishes a stable univalent positive ion. The peptisation of aluminium hydroxide leads to a sol of the composition $2[\text{Al}(\text{OH})_3]\text{Al}(\text{OH})_2\text{Cl}$, and this on dilution undergoes complex ionisation represented by the formulæ (1) $10\text{Al}(\text{OH})_3, 4\text{AlOCl}, \text{AlO}|\text{Cl}$; (2) $12\text{Al}(\text{OH})_3, 5\text{AlOCl}, \text{AlO}|\text{Cl}$; (3) $16\text{Al}(\text{OH})_3, 7\text{AlOCl}, \text{AlO}|\text{Cl}$. On the other hand, no complex ionisation of the form $\text{Al}(\text{OH})_4|\text{AlO}$ has been observed in the case of aluminium hydroxide. J. F. S.

Germanium. I. Extraction from Germanium-bearing Zinc Oxide. Non-occurrence in Samarskite. L. M. DENNIS and JACOB PAPISH (*J. Amer. Chem. Soc.*; 1921, **43**, 2131—2144).—A method of extracting germanium residues obtained in the smelting of certain American zinc ores has been investigated and is described. The residue contains zinc oxide, considerable quantities of lead, arsenic, and cadmium, and small quantities of indium, tin, and antimony, in addition to the germanium. A kilogram of the crude oxide is placed in a 5-litre Pyrex flask, which is fitted with a two-holed rubber stopper carrying a bent glass tube for connexion to a Liebig condenser, and a second short glass tube for introduction of the acid. The condenser leads under the surface of water contained in a 4-litre bottle which acts as receiver. Two and a half litres of hydrochloric acid ($d\ 1\cdot18$) are added to the ore and the flask is heated until 2 litres of distillate have been collected. The distillate at this point contains all the germanium and a great deal of the arsenic from the ore. The distillate is poured into a 15-litre bottle until 10 litres have been accumulated, carefully acidified with sulphuric acid until it is $6N$ (this must be carefully done to prevent loss of germanium chloride), and treated with washed hydrogen sulphide. The precipitate is at first yellow, due to arsenic, but later becomes whiter, due to the germanium. When precipitation is complete, the bottle is stoppered and kept for twenty-four hours. The solution is filtered by suction and washed with $3N$ -sulphuric acid which has previously been saturated with hydrogen sulphide. The filtrate is kept for forty-eight hours, when usually a small quantity of a white precipitate (corresponding with 2 mg. of germanium) separates, the majority of the supernatant liquid is siphoned off, and the small amount of liquid and

precipitate worked up with the filtrate from the next lot of precipitate. At this point two different methods of procedure are possible.

(1) The moist sulphides are added to hot 50% sodium hydroxide until a small quantity remains undissolved; this is then just dissolved by the addition of a little more sodium hydroxide. The solution is made strongly alkaline by the addition of 8 grams of solid sodium hydroxide and placed in a large Pyrex flask fitted with a rubber stopper carrying a delivery tube, a separating funnel, and a glass tube reaching almost to the bottom of the flask. The flask is connected to a condenser and receiver as before. Washed chlorine is passed in to oxidise the arsenic to the quinquevalent condition. When the solution is saturated with chlorine, the rate of entry of the chlorine is reduced and concentrated hydrochloric acid is added in large excess from the funnel. The flask is heated until half the liquid has distilled. The germanium chloride passes over and is hydrolysed by the water in the receiver, forming white, hydrated germanium dioxide. Should oily drops form in the receiver, more water is added to reduce the acid concentration and so allow the hydrolysis to proceed. The receiver is now replaced by a second one, the distilling flask is filled up again with concentrated hydrochloric acid, and the distillation continued as before. Most of the germanium chloride passes over in the first distillation, but for a complete separation the distillation must be repeated several times. The hydrated oxide is filtered, washed first with dilute sulphuric acid and then with water, and dried at 110° . It is pure white, and contains traces of sodium, calcium, and iron, but no arsenic. The filtrates from the hydrated oxide are treated with hydrogen sulphide and the germanium sulphide is recovered. The impurities mentioned are removed by dissolving in a slight excess of sodium hydroxide, saturating with chlorine, and distilling with hydrochloric acid, hydrolysing the distillate as before.

(2) This process is generally superior to the former in its greater economy of reagents. The wet sulphides are washed with 3*N*-sulphuric acid until free from chlorine and dried at 110° . They are then roasted in shallow iron dishes at temperatures not exceeding 500° . This removes a great deal of the arsenic. The roasted material is dissolved in sodium hydroxide (50%), chlorinated, and distilled with twice its weight of concentrated hydrochloric acid. A repetition of the distillation removes the last trace of arsenic. Either process gives a very pure germanium dioxide; the yield is better by the first, but the second is more rapid and economical.

Germanium in ores is estimated by grinding 20–100 grams of the finely ground, dried, and weighed ore into a paste with water and pouring into a solution of sodium hydroxide in a hard flask, the proportions being 2 of ore : 1 sodium hydroxide : 5 of water. The flask is fitted with a delivery tube for leading in chlorine, a fractionating column, and a small tap funnel. It is connected with a Liebig's condenser which leads to two Erlenmeyer flasks in series containing water to the depths of 3 cm. and half full respectively. The receivers are cooled with ice. The distilling flask is surrounded

by ice and the contents are saturated with chlorine; the solution is then neutralised by hydrochloric acid and an excess equal to twice the weight of the ore added. The ice is removed and a slow distillation in chlorine carried out until one-half the liquid in the flask has passed over. An equal volume of hydrochloric acid is added to the distilling flask and the distillation continued until its volume is again reduced by one-half. This is then repeated once more. The receivers are disconnected, acidified with sulphuric acid to make the solutions 6*N*, saturated with hydrogen sulphide, and kept for twenty-four hours. In a successful experiment there will be no germanium in the second flask. The precipitate is brought on to an ashless paper and washed with 3*N*-sulphuric acid, saturated with hydrogen sulphide until free from chloride, then washed with alcohol to remove the acid, and dried. The bulk of the precipitate is placed in a porcelain crucible and dried, moistened with 1 : 1 nitric acid, and warmed to drive off all the liquid, allowed to cool, treated with concentrated nitric acid, dried, and ignited. The filter-paper is incinerated in a second crucible, treated with concentrated nitric acid, and ignited. The filtrate from the germanium sulphide is kept forty-eight hours and the small precipitate formed filtered and treated as above. The weight of the germanium dioxide in the crucibles represents the germanium in the ore. Using this method, the amount of germanium in the material used for the extraction of germanium was found to be: sample I 0.247%, sample II 0.19%. Samarskite has been analysed by this method and found not to contain germanium.

J. F. S.

Crystalline Structure of Iron and Steel. ARNE WESTGREN and AXEL E. LINDH (*Z. physikal. Chem.*, 1921, **98**, 181—210).—Various types of iron and steel have been subjected to Röntgen ray spectrographic examination. It is shown that the results of Hull (*Phys. Review*, 1917, **9**, 84, **10**, 661), that pure iron at ordinary temperatures (α -iron) possesses a space-centred cubic lattice, is in keeping with the present work. At 800—836°, that is, within the so-called β -iron region, the atoms are grouped in exactly the same way as in α -iron. Since in the authors' opinion allotropy and polymorphy are synonymous, β -iron can only be regarded as a particular modification of α -iron. In austenite and in pure iron stable at 1000°, the crystals possess a face-centred cubic lattice. This is also characteristic of γ -iron, and thereby a fundamental difference is established between α -iron and γ -iron. In martensite, the iron occurs in its α -modification. This is also the case with high speed tool steel which has been hardened at 1275°. Photomicrographs of some of the preparations examined are included in the paper.

J. F. S.

Alloys of Iron and Uranium. E. P. POLUSHKIN (*Iron and Steel Inst., Carnegie Schol. Mem.*, 1920, **10**, 129—150; cf. *Rev. Metal.*, 1920, **17**, 421).—Alloys of iron with uranium are pyrophoric, this property varying directly with the content of uranium. Alloys containing carbon are decomposed by water, but this effect is not

exhibited by silicon, except when present in very large amount, or by vanadium. A polished sample placed on the emulsion of a photographic plate leaves an impression which, after development by the usual process, represents the structure of the specimen. Iron-uranium alloys contain the uranium carbides UC , U_2C_3 , and U_3C_8 , the compound Fe_3C, U_2C_3 , as well as the compounds V_2C , Fe_6U , and $FeSi$. Neither uranium nor any of its compounds already mentioned forms solid solutions with iron. Vanadium has a greater affinity for uranium than for carbon. CHEMICAL ABSTRACTS.

Equilibria in the System Fe-C-O: the Equilibrium Fe_β -Martensite-Ferrous Oxide-Gas. W. REINDERS and P. VAN GRONINGEN (*Rec. trav. chim.*, 1921, 40, 701-706).—A considerable number of determinations of points of univariant equilibrium for this system have been carried out. The conclusion is drawn that the transition temperature for $Fe_\beta \rightarrow Fe_\gamma$ is 905° ; this is in accord with previous results obtained by different methods. By extrapolation of the graphic results, the quintuple point of the system is found at 740° and 2300 mm. H. J. E.

Researches on the Metallic Carbonyls. ROBERT LUDWIG MOND and ALBERT EDWARD WALLIS (T., 1922, 121, 29-32).

The Action of Nitric Oxide on the Metallic Carbonyls. ROBERT LUDWIG MOND and ALBERT EDWARD WALLIS (T., 1922, 121, 32-35).

Expansion of Chromium and of Nickel-Chromium Alloys over a Wide Range of Temperatures. P. CHEVENARD (*Compt. rend.*, 1922, 174, 109-112).—Between 0° and 100° the expansion of chromium is exactly reversible, the curve showing no singular point. The true coefficient of expansion, which is 6.8×10^{-6} at 0° , increases rapidly with the temperature, but the curve shows a slight concavity towards the increasing temperature.

Nickel-chromium alloys containing up to 16% of chromium and from 0.5 to 2.5% of manganese were examined over the temperature range 0° to 1000° . The addition of chromium leads to a very rapid weakening of the anomaly of dilatation of nickel, and when the chromium content reaches 5% the anomaly disappears. The addition of chromium to nickel affects the expansion of this metal very little at the ordinary temperature, but tends to increase it at higher temperatures; this effect being probably due to the presence of the compound Ni_2Cr_3 . W. G.

Zirconium. J. W. MARDEN and M. N. RICH (*Bur. Mines Bull.*, 1921, 186, 146 pp.; cf. A., 1920, ii, 547).—The work is divided into four parts, dealing successively with an historical review of zirconium minerals, the salts of zirconium, and the metal; experimental work on zirconium: the furnaces used; and a bibliography of zirconium and its compounds. Analytical methods are given for the estimation of zirconium in ferrozirconium, steel, alloys such as nickel-zirconium, and a method of separation of titanium, columbium, tantalum, and zirconium. The physical and chemical

properties of amorphous and coherent zirconium are described. The cupferron method is the only one effecting complete separation of zirconium from aluminium.

CHEMICAL ABSTRACTS.

Polymorphic Transformations of Antimony Trisulphide. SAMUEL WILSON and C. R. McCROSKY (*J. Amer. Chem. Soc.*, 1921, **43**, 2178—2179).—The rate of transformation of the red, amorphous variety of antimony trisulphide into the black variety has been determined in the presence of *N*-, 7*N*-, 12*N*-hydrochloric acid, 7*N*-phosphoric acid, 7*N*-sulphuric acid, glacial and *N*-acetic acids, and a saturated ethereal solution of hydrochloric acid. Of these acid solutions, only the aqueous solution of hydrochloric acid gave any appreciable transformation at 18—22°, even after two months. With the aqueous hydrochloric acid, the time necessary for complete transformation was 0.5 day, 1 day, and 10.5 days for 12*N*, 7*N*, and *N*, respectively. The action is due to a solution of the red variety and a reprecipitation of the less soluble black variety. The effect of temperature was investigated with a 20% aqueous hydrochloric acid. Complete conversion at 26.5° required forty-four hours; at 30°, twenty-nine hours; 35°, sixteen hours; 40°, nine hours; 68.5°, sixty-two minutes, and 75°, thirty-two minutes. A 20% solution of hydrobromic acid gave no change after twenty hours at 75°.

J. F. S.

Mineralogical Chemistry.

The Ultimate Composition of British Coals. THOMAS JAMES DRAKELEY [with FREDERICK WILLIAM SMITH] (*T.*, 1922, **121**, 221—238).

Kasolite, a New Radioactive Mineral. ALFRED SCHOEP (*Compt. rend.*, 1921, **173**, 1476—1477).—This occurs together with curite (this vol., ii, 77) and torbernite at Kasolo, Katanga, Belgian Congo. It forms compact, crystalline aggregates, with sometimes tufts and radiating groups of prismatic crystals on the surface. The colour is ochre-yellow to brownish-yellow and the streak ochre-yellow. The crystals are monoclinic, with the optic axial plane perpendicular to the plane of symmetry, and the acute bisectrix of the optic axes nearly perpendicular to a perfect cleavage. $d^{17} 5.962$, *H.* 4—5. Analysis I is of translucent crystals showing no sign of alteration, II of less fresh massive material, and III of clear crystals:

	SiO ₂	PbO.	UO ₂	H ₂ O.	Fe ₂ O ₃	CaO.	MgO.	CO ₂	Total.
I.	9.42	36.20	49.28	3.59	0.41	0.06	0.03	0.85	99.84
II.	9.14	34.44	49.00	3.77	0.58	—	—	0.53	—
III.	9.00	32.16	48.26	3.28	0.40	—	—	—	—

The simplest formula corresponding with the mean of these analyses is $3\text{PbO}, 3\text{UO}_3, 3\text{SiO}_2, 4\text{H}_2\text{O}$. The mineral is decomposed by acids with the separation of gelatinous silica and of lead chloride or sulphate. Heated in the reducing flame on charcoal it fuses to a black glass with beads of lead. Its radioactivity is less than that of curite.

L. J. S.

Melanovanadite, a New Mineral from Peru. WALDEMAR LINDGREN (*Proc. Nat. Acad. Sci. U.S.A.*, 1921, 7, 249—251).—This occurs as bunches of black needles on a black shale from Minasragra, Cerro de Pasco. The crystals are monoclinic with a perfect cleavage parallel to the plane of symmetry. The streak is dark reddish-brown, d^{15} 3.477, H . $2\frac{1}{2}$. Only the thinnest splinters are brown and translucent under the microscope, and the optical characters are masked by the strong absorption. Analysis by L. F. HAMILTON gives the formula $2\text{CaO}, 3\text{V}_2\text{O}_5, 2\text{V}_2\text{O}_4$.

V_2O_5 .	V_2O_4 .	CaO .	MgO .	$\text{Al}_2\text{O}_3, \text{Fe}_2\text{O}_3$.	SiO_2 .	Total.
52.61	33.34	9.89	0.27	1.89	1.66	99.66

Before the blowpipe the mineral readily fuses to a brown liquid. It is readily soluble in acids, giving an apple-green solution, and in potassium hydroxide with a brown colour.

L. J. S.

Some Natural and Synthetic Melilites. A. F. BUDDINGTON (*Amer. J. Sci.*, 1922, [v], 3, 35—87).—To test Schaller's hypothesis regarding the composition of the melilite group of minerals, comprising akermanite, gehlenite, humboldtilite, sarcolite, fuggelite, and other varieties (A., 1916, ii, 632) more than one hundred synthetic crystalline mixtures of $2\text{CaO}, \text{MgO}, 2\text{SiO}_2$ (akermanite), $2\text{CaO}, \text{Al}_2\text{O}_3, \text{SiO}_2$ (gehlenite), and $3\text{R}'\text{O}, \text{R}_2\text{O}_3, 3\text{SiO}_2$ ($\text{R}' = \text{Ca}$ or Na , $\text{R} = \text{Fe}$ or Al) were prepared from appropriate glasses at temperatures above 1000° . The quenching method was used (cf. Ferguson and Buddington, A., 1920, ii, 621), and the glasses were crystallised by annealing at a temperature just below the melting point or the dissociation point. The homogeneity, optical characters, and melting points were determined, and the synthetic materials compared with the natural minerals. The gehlenite of Velardeña consists approximately of 76% of gehlenite, 17% of akermanite, and 7% of ferric and ferrous compounds, and agrees very closely with artificial gehlenite, having the same ratio of akermanite to gehlenite, but free from iron compounds. Other natural gehlenites examined confirm the opinion that the artificial series of solid solutions of $2\text{CaO}, \text{MgO}, 2\text{SiO}_2$ and $2\text{CaO}, \text{Al}_2\text{O}_3, \text{SiO}_2$ are pure synthetic analogues of the akermanite-gehlenite series of minerals. Mixtures of akermanite and gehlenite form a complete series of solid solutions with $3\text{CaO}, \text{Al}_2\text{O}_3, 3\text{SiO}_2$ (grossularite) plus 10% of $3\text{Na}_2\text{O}, \text{Al}_2\text{O}_3, 3\text{SiO}_2$, except for a trace of inhomogeneity in some preparations high in akermanite. These mixtures when crystallised correspond in their properties with the humboldtilites, which are interpreted as isomorphous mixtures of positive uniaxial akermanite and a negative, uniaxial, tetragonal, moderately birefringent form of grossularite, with minor amounts of gehlenite, a

ferrous compound, and $3R'O, R_2O_3, 3SiO_2$ compounds. The compositions of the humboldtilites lie in a zone which exhibits the lowest temperatures of complete melting for the components involved. Artificial crystalline mixtures containing ferric iron ($3CaO, Fe_2O_3, 3SiO_2$) were found to be quite different from the natural melilites of similar composition rich in ferric iron. The latter probably were formed at temperatures lower than those of the present experiments, since some of them were found to decompose or invert at temperatures as low as 850° . A new specimen of melilite rich in ferric iron, from Capo di Bove, differs in composition from any hitherto known. It contains :

SiO_2	Al_2O_3	Fe_2O_3	FeO	MgO	CaO	Na_2O	K_2O	Total
40.03	5.66	7.76	0.4	9.43	32.17	2.83	1.72	100.0

The crystals have a tabular or pseudo-cubic habit and are intimately associated with nephelite and pyroxene.

E. H. R.

Analytical Chemistry.

The Graphical Representation of the Composition of Chemical Compounds. JULIUS HÜBSCHER (*Chem. Ztg.*, 1922, 46, 19—20).—The repeated calculation of the percentage of the various constituents of chemical compounds in mixtures can be avoided by calculating once and for all the percentage of the required constituent (for example, anhydrous sodium carbonate in the decahydrate) and dividing a vertical line in these proportions. A square is then described with this line as its right-hand side and the points marked off joined to the left-hand top corner. From this co-ordinate system the weight of any constituent corresponding with a given weight of any other constituent can be rapidly read off with sufficient accuracy for practical purposes. Should the given weights not lie between 0 and 100 they can be divided by a suitable factor and the corresponding weight read off the diagram and subsequently multiplied by the same factor.

H. C. R.


The Sensitiveness of Coloured Indicators at Temperatures above the Ordinary. I. M. KOLTHOFF (*Rec. trav. chim.*, 1921, 40, 775—785; cf. Schoorl, A., 1907, ii, 388).—Indicators which are themselves weak acids are almost all as sensitive to hydrogen ions at higher temperatures as at ordinary temperatures. Those which are weak bases become less sensitive to hydrogen ions, but retain the same sensitiveness to hydroxyl ions. Theoretical deduction of these facts is given in addition to experimental evidence. The suggestion is made that the results obtained may be of service in providing a colorimetric method of studying variations in dissocia-

tion constant and of hydrolysis constant with change of temperature and may also be of use in analytical work. H. J. E.

Further Elaboration of the Indicator Method without Buffers. L. MICHAELIS and R. KRÜGER (*Biochem. Z.*, 1921, **119**, 306—327).—The salt error and temperature coefficient of *m*-nitrophenol have been determined, and a new one-colour indicator, 2 : 5-dinitrophenol, described. A theoretical and practical treatment of the effect on the P_H of a solution of the addition of an indicator is given, and instructions for the colorimetric estimation of P_H in solutions weak in buffers, for example, sea and river waters. A discussion of the theory of the salt-error and of the activity theory of ions is also given. H. K.

A Stable Single Buffer Solution, p_H 1 to p_H 12. S. F. ACREE, R. R. MELLON, PAULINE M. AVERY, and E. A. SLAGLE (*J. Infect. Dis.*, 1921, **29**, 7—10).—The components of the buffer solution are: (1) One mol. of potassium dihydrogen phosphate, with dissociation constant, K_a 1.1×10^{-2} ; (2) 0.625 mol. of sodium formate, K_a 2×10^{-4} ; (3) 0.375 mol. of sodium acetate, K_a 2×10^{-5} ; (4) the second group of dipotassium hydrogen phosphate, K_a 2×10^{-7} ; (5) 1 mol. of sodium phenolsulphonate, K_a 10^{-10} (approx.); (6) 0.005 molar thymol to saturation (for water, 0.08), K_a 0.5×10^{-10} (approx.); (7) the third group of phosphoric acid, K_a 10^{-12} . A curve is given from which the amounts of 0.5 molar hydrogen chloride or sodium hydroxide necessary to produce a given p_H may be directly determined. CHEMICAL ABSTRACTS.

Colour Standards for the Colorimetric Measurement of Hydrogen-ion Concentration. LOUIS J. GILLESPIE (*J. Bact.*, 1921, **6**, 399—405).—The recently published studies of Medalia are in disagreement with other published data (cf. *ibid.*, 1920, **5**, 441—468). A colorimeter for two-coloured indicators is described for the measurement of the hydrogen-ion exponent of indicators. The optical assumptions underlying its use are practically the same



as those on which ordinary colorimetry are based. The instrument is used as follows. The glass vessels A and C are fixed in position, and B can be moved up and down, the motion being measured by a pointer (not shown) fixed to B and moving on a scale divided into 100 parts. The pointer moves from 0 to 100 when B moves from contact with C to contact with A. The acidified indicator solution may be placed in B and the alkaline indicator solution of the same strength in C. A is left empty. If the scale reads 70, the path of light along the left dotted line passes through the alkaline form of the indicator for 10% of its path and the acid form for 30%. The light along the right-hand dotted line traverses an indicator solution in tube E, again of the same strength, and over a path equal in length to the total path on the left. The

solution, the p_H of which is unknown, is placed in tube E , in which titrations may be made.

CHEMICAL ABSTRACTS.

A New Apparatus for Colorimetric Estimations. O. MANNEBACH (*Chem. Ztg.*, 1922, 46, 20).—The apparatus consists of a wooden box open at the top the inside of which is painted black. Into this fits a glass vessel filled with water. The bottom of the box has a hole cut in it allowing light to come up through the glass vessel, and below is a rotating frame carrying a dead-white porcelain plate. The solutions to be compared are placed in glass tubes 16 mm. in diameter and 300 mm. long, which are placed in the glass vessel in a slanting position. The apparatus is suitable for the colorimetric estimation of carbon in iron. H. C. R.

A Gas Receiver of Convenient and Practical Form for Sampling Expired Air for Analysis. CHARLES CLAUDE GUTHRIE (*J. Biol. Chem.*, 1921, 48, 373—378).—Whilst less efficient than the mercury receiver, the apparatus described has the advantage of being inexpensive. Expired air, after storage in it for some hours, gave results, on analysis, with an error of about 1% for carbon dioxide and considerably less for oxygen. E. S.

Apparatus for Estimation of the Gases in Blood and Other Solutions. DONALD D. VAN SLYKE (*Proc. Nat. Acad. Sci.*, 1921, 7, 229—231).—Essentially a much simplified form of the apparatus described in A., 1917, ii, 422, and this vol., ii, 78. The upper part of a large pipette (for instance, 50 c.c.) ends in a tap funnel. The stem above the wide portion has a mark, indicating a definite volume a from there to the tap (for instance, 2 c.c.), and below the wide portion another mark indicating a volume A (for instance, 50 c.c.) from it to the tap. The pipette is joined below to a tube of 800 mm. connected with an open manometer and through a tap, with a mercury reservoir. A definite volume of solution (for instance, 1 c.c. of blood) is sucked in through the tap funnel and then the necessary reagent (for instance, acid for a carbonate solution), making a total solution of S c.c. (for instance, 2.5 c.c.) and the mercury is allowed to fall to the lower mark. The lower tap is closed, and the pipette is shaken for one to two minutes to establish equilibrium. Mercury is then let in through the lower tap until the gas volume is a c.c. and the pressure is read (m mm.). The zero point is then determined by expelling the gases or after absorbing one or more of them by introducing small, measured volumes of gas-free absorbent solutions. The pressure is then lowered until the space above the solution is again a c.c. and read (n mm.). The volume at $N.T.P.$ of the gas given off is $V = a(m-n)/760 \cdot \{273/T + S\alpha/(A-S)\}$. The term $S\alpha/(A-S)$, in which α is the volume of the gas dissolved in 1 c.c. of the solution at $N.T.P.$ corrects for the portion of the gas remaining dissolved when equilibrium is reached. It is negligible for oxygen and nitrogen, but not for carbon dioxide. The solubility of the latter gas also imposes an empirical correction for reabsorption of the gas while undergoing reduction from $50-S$ to a c.c. With $S=50$ c.c. $a=2$ c.c. the factor is 1.020, that is 2% of the carbon dioxide is reabsorbed. G. B.

The Use of Mercuric Nitrate instead of Silver Nitrate in the Estimation of the Halogens. I. M. KOLTHOFF and ADA BAK (*Chem. Weekblad*, 1922, 19, 14—16).—The method proposed by Votoček (A., 1918, ii, 238, 272, 330), in which mercuric nitrate is used with sodium nitroprusside as indicator, gives very accurate results for chlorides (and for bromides, cyanides, and thiocyanates) if a correction is applied for the excess of mercuric salt necessary to produce a precipitate under given conditions of composition and volume of solution. Tables of corrections are given. Neither dilute acids nor the common metals (except copper, cobalt, nickel, and cadmium) interfere. The method estimates accurately chlorides in conductivity water, and gives good results for as little as 9 mg. of chlorine per litre; it is suitable for the estimation of chlorides in urine. S. I. L.

Use of Perchloric Acid as an Aid to Digestion in the Kjeldahl Nitrogen Estimation. BRAINERD MEARS and ROBERT E. HUSSEY (*J. Ind. Eng. Chem.*, 1921, 13, 1054—1056).—In the estimation of nitrogen in such substances as milk, urine, casein, gelatin, dried blood, etc., by the Kjeldahl method, the time required for the digestion with sulphuric acid is reduced to about twenty minutes if perchloric acid is added to the mixture. For each gram of sample, 25 c.c. of sulphuric acid, 1 gram of copper sulphate, and 2 c.c. of 60% perchloric acid should be used, but the presence of an excess of perchloric acid causes loss of nitrogen. W. P. S.

Micro-Kipp Apparatus for the Preparation of Air-free Carbon Dioxide for Use in the Micro-estimation of Nitrogen by Pregl's Method. A. SCHOELLER (*Z. angew. Chem.*, 1921, 34, 586).—The apparatus consists of two small cylindrical bulbs, one above the other; the upper one contains fused potassium-sodium carbonate and is provided with a tapped delivery tube, whilst a side tube on the lower bulb connects with an upper acid reservoir. The apparatus is made all in one piece. W. P. S.

Estimation of Very Small Quantities of Arsenic in Silicate Rocks. O. HACKL (*Chem. Ztg.*, 1921, 45, 1169).—Ten grams of the finely-powdered sample are heated at 250° in a tube through which a current of dry carbon dioxide saturated with bromine vapour is passed; the outlet end of the tube is connected with a receiver containing 10 c.c. of dilute nitric acid and the receiver may be connected with a vessel containing sodium hydroxide solution to absorb the excess of bromine. The contents of the receiver are subsequently evaporated with the addition of a small quantity of sulphuric acid, the residue is dissolved in water, and the arsenic in this solution estimated by the Gutzeit method. W. P. S.

Method for Direct Estimation of Carbon Dioxide and Oxygen in the Berthelot Bomb and its Importance for the Metabolic Balance of Herbivora. W. KLEIN and MARIA STEUBER (*Biochem. Z.*, 1921, 120, 81—89).—An extension of the

work of Zuntz and co-workers on the use of the Berthelot bomb calorimeter for the analysis of organic substances, the main feature now introduced being the determination of the oxygen used, by weighing. It may prove useful for the analysis of the metabolic excreta of herbivora. H. K.

Micro-elementary Analysis by Pregl's Method. A. SCHOELLER (*Z. angew. Chem.*, 1921, **34**, 581—583).—A short review of the method, with a detailed description of the apparatus used. W. P. S.

Simplified Construction of the Metal Parts of the Apparatus used in Pregl's Micro-analytical Method. A. SCHOELLER (*Z. angew. Chem.*, 1921, **34**, 587).—Convenient methods of attaching the heating apparatus for the lead peroxide, the micro-burner, the drying chamber, etc., to the stand are described, the parts being provided with brass rods which fit into a boss on the stand. W. P. S.

Solid Sodium Hydroxide as an Absorbent for Carbon Dioxide in Steel Analysis. G. L. KELLEY and E. W. EVERS (*J. Ind. Eng. Chem.*, 1921, **13**, 1052).—Powdered sodium hydroxide, which will pass through a 5-mesh sieve but be retained on a 20-mesh sieve yields satisfactory results when used to absorb carbon dioxide in the estimation of carbon. W. P. S.

The Separation of Silver from Mercurous Salts. I. M. KOLTHOFF (*Pharm. Weekblad*, 1921, **58**, 1680—1683).—The treatment of the mixed chlorides precipitated in Group I with ammonia will not separate small quantities of silver from mercurous chloride. A better method is to treat the mixed chlorides, after exhaustive washing with boiling water to remove lead, with 2% potassium cyanide solution. Silver and mercuric cyanides dissolve in the reagent, whilst black metallic mercury is precipitated. After filtering, silver is again precipitated by means of hydrochloric acid; mercuric chloride remains in solution, and may be detected, after filtering, by means of sodium sulphide. The test detects 0.05 mg. of silver in presence of 50 mg. of mercurous mercury, and will also detect 0.5 part of mercury in presence of 100 parts of silver. S. I. L.

The Titration of Zinc. E. MONASCH (*Pharm. Weekblad*, 1921, **58**, 1652—1656).—The thiocyanate method of Kolthoff and van Dijk (*ibid.*, 538) has been applied to the estimation of zinc in alloys. The potassium mercuric thiocyanate solution is prepared by dissolving 23.7 grams of mercuric thiocyanate in a concentrated aqueous solution of 14.4 grams of potassium thiocyanate, and is stable for many months. Compounds of all the common metals interfere, but ferric and aluminium salts do not affect the reaction. Since the author uses aluminium in the separation of zinc from alloys, the method is suitable for the estimations, but iron salts must first be oxidised by means of peroxide. S. I. L.

Estimation of Minute Amounts of Lead in Water, with Notes on certain Causes of Error. D. AVERY, A. J. HEMINGWAY, V. G. ANDERSON, and T. A. READ (*Proc. Austral. Inst. Min. Met.*, 1921, 173—199).—By the following method it is possible to detect and estimate with considerable accuracy 1 part of lead in 100,000,000 parts of water. A known volume of the water is filtered, and 2.5 to 5 litres of the filtrate are evaporated to about 100 c.c., just neutralised with hydrochloric acid, and 2 c.c. excess of acid are added, the liquid is filtered and the filtrate cooled and made up to 250 c.c. Hydrogen sulphide is passed into the solution for one hour and, after remaining over-night, the precipitate, which is usually barely visible, is collected and washed with cold water containing hydrogen sulphide in solution. Two portions of 2 c.c. of hot nitric acid (d 1.2) are then successively poured over the filter to dissolve the sulphides, and the filter is washed with hot water. The filtrate is evaporated with 1 c.c. of sulphuric acid until the latter fumes strongly, and the liquid, after cooling, is treated with 20 c.c. of cold water and 10 c.c. of absolute alcohol, and the mixture set aside over-night. The precipitated lead sulphate is collected and washed with a mixture of 65 vols. of water, 32 vols. of absolute alcohol, and 3 vols. of sulphuric acid. It is dissolved off the paper by dropping successively two portions of 5 c.c. of hot 33% ammonium acetate solution round the edges and then washing the paper thoroughly with hot water. The solution is transferred to a 50 c.c. Nessler tube and treated with 1 c.c. of 10% potassium cyanide solution, 1 c.c. of ammonia solution, and six drops of freshly-prepared ammonium sulphide solution. Into the standard tube are placed the same reagents in the same quantities; the liquid, which must be absolutely colourless, is diluted to 45 c.c., and a standard solution of lead acetate (1 c.c. = 0.00001 gram of lead) is added until the tint matches that of the assay. If more than 8 c.c. of the standard are required, a proportionately smaller quantity of the sample should be used. To estimate the lead in the sediment, it is evaporated to dryness with hydrochloric acid, the residue is taken up with 2 c.c. of the same acid, the solution filtered, and the assay finished in a similar way to that of the water.

The estimation of lead in urine is carried out by evaporating 1 litre of the sample with 50 c.c. of nitric acid to dryness, first on a water-bath, then on the hot-plate. The dish is then placed in a cool, electrically heated muffle and gradually heated to 450—500° to destroy organic matter and nitrates. The residue is dissolved in water, the liquid just neutralised with hydrochloric acid, 2 c.c. more acid are added, and the solution is filtered. The filtrate is treated for lead as described above.

All the materials used in the above work should be redistilled from lead-free glass apparatus, the ammonium acetate should be made by neutralising freshly distilled ammonia with acetic acid, and the ammonium sulphide must be made immediately before use. All filter-papers before use must be washed with hot dilute hydrochloric acid, hot ammonium acetate solution, and hot water

successively to remove the minute amounts of lead introduced by the acid washing process of rendering them ashless. A blank test must be made, using the same quantity of redistilled water as that used for the assay and putting it through the whole of the operations.

A. R. P.

New Method for the Volumetric Estimation of Copper. S. MINOVIĆI and AL. IONESCU (*Bul. Soc. Chim. România*, 1921, 3, 89—93).—The salt $\text{CuSO}_4 \cdot 4\text{NH}_3$ is quite stable at the ordinary temperature, and is quantitatively precipitated from aqueous solution by addition of eight volumes of 98% alcohol. The precipitate, after thorough washing with alcohol, is redissolved in water, and titrated with *N*/10-sulphuric or oxalic acid, with methyl-red as indicator.

J. K.

Rapid Estimation of Mercury in Ores. ALFRED HEINZELMANN (*Chem. Ztg.*, 1921, 45, 1226—1227).—The author has made comparative tests on the estimation of mercury in ores, using a modification of Whitton's method (*U.S. Bureau of Mines, Bull.* 78, 1918) which consists in heating the ore with a mixture of 3 grams of fine iron filings and 3 grams of good lime and collecting the mercury on a cooled silver plate previously weighed, and his own method (*A.*, 1921, ii, 521) and finds that both methods give equally satisfactory results. [Cf. *J. Soc. Chem. Ind.*, 1922, 61A.]

A. R. P.

The Analysis of Aluminium Alloys. H. MENDE (*Chem. Ztg.*, 1922, 46, 49—50).—One gram of the alloy is heated with 5 c.c. of water and 12 c.c. of strong potassium hydroxide solution (the latter added gradually) on a water-bath until nothing further dissolves. The liquid is diluted with water, the clear liquor decanted through a small filter-paper, and the precipitate washed by decantation, first with dilute potassium hydroxide solution, then with hot water. The filter-paper is burnt in a platinum spiral, the ash added to the metallic residue in the beaker, and the whole dissolved in 5 c.c. of strong nitric acid and 15 c.c. of water; 75 c.c. of water are added, the liquid is boiled and the precipitate collected, washed, ignited, and weighed as tin dioxide. It should be tested for traces of silica. The filtrate is evaporated with 5 c.c. of sulphuric acid until the latter fumes strongly, the mass is treated with 150 c.c. of water, and the precipitated lead sulphate estimated in the usual way. The filtrate is electrolysed with a current of 0.5 ampere at 2.0—2.2 volts for one and a half hours at 75°, using a spiral anode and a gauze cathode. The gain in weight of the latter represents copper. The solution is neutralised with sodium hydroxide after adding any zinc found by treating the potassium hydroxide solution of the alloy with sodium sulphide, and 50 c.c. of a 50% solution of sodium hydroxide are added in excess. The solution is again electrolysed, using the same anode and cathode (the latter being coppered or silvered) for two to three hours at 70°, using a current of 1—1.5 amperes at 4 volts. The gain in weight of the cathode represents zinc. Silicon is estimated in a separate

trial in which 3 grams of the alloy are evaporated with 50 c.c. of a mixture of 2 parts of sulphuric acid, 3 parts of hydrochloric acid, 1 part of nitric acid, and 4 parts of water. After the metal has dissolved, the solution is evaporated with 30 c.c. of strong sulphuric acid until the latter fumes strongly, the liquid is treated with 300 c.c. of water, and the silica, which separates in an easily filterable, flocculent form, is collected, washed, ignited, and weighed as usual. Iron is determined by treating 3 grams of the alloy in potassium hydroxide solution, collecting the insoluble residue, and dissolving it, after thorough washing, in nitric acid. The solution is evaporated to dryness, the nitric acid destroyed by evaporation with hydrochloric acid, the copper and heavy metals are removed by hydrogen sulphide, the iron is precipitated from the oxidised solution by ammonia, redissolved in acid, reprecipitated as before, and the final precipitate either weighed as Fe_2O_3 after ignition, or dissolved in sulphuric acid, reduced, and titrated in the usual way.

For the estimation of traces of iron in pure aluminium, 3 grams of the metal are heated with 50 c.c. of strong potassium hydroxide solution, 200 c.c. of 50% sulphuric acid are added, the solution is heated until everything has dissolved, then quickly cooled, and titrated with $N/10$ -potassium permanganate. Copper and silicon are estimated as described above for alloys. A. R. P.

Germanium. II. Identification of Germanium by its Visible Arc Spectrum. JACOB PAPISH (*Chem. News*, 1922, 124, 3).—Photographs of the arc spectrum of germanium were obtained by means of a Hilger constant deviation spectrograph fitted with a flint glass prism ($n_D=1.7537$), the arc being made between carbon electrodes, the lower positive one carrying the material to be tested. Under these conditions, the little-known germanium line in the blue, λ 4686, is very sharp and intense, and very suitable for the identification of the metal. The line overlaps the prominent zirconium line of wave-length 4688.

G. F. M.

The Carrying Down of Calcium Oxide by Precipitates of Ferric Oxide. A. CHARRIOU (*Compt. rend.*, 1921, 173, 1360—1362; cf. Toporescu, A., 1920, ii, 450).—In order to get the minimum co-precipitation of calcium hydroxide with ferric hydroxide, the concentration of the calcium salt should be as small as possible and the minimum amount of ammonium hydroxide requisite for the precipitation of the ferric hydroxide should be used. W. G.

Estimation of Nickel in Steels. H. RUBRICIUS (*Chem. Ztg.*, 1922, 46, 26).—From 2 to 5 grams of the borings are dissolved in 40–80 c.c. of nitric acid (d 1.2), and the solution is cooled and treated with 250 c.c. of ammonia (d 0.91). It is transferred to a graduated flask and diluted to 500 c.c. of which 250 c.c. are filtered through a dry paper, diluted to 500 c.c., and the solution, heated at 40° , is treated with 20–30 c.c. of a 1% alcoholic solution of dimethylglyoxime. After half an hour the precipitate is filtered

on a 15 cm. paper, washed with hot water, dried, and ignited to nickelous oxide in a platinum crucible and weighed. $0.7858 \times \text{NiO} = \text{Ni}$.
A. R. P.

Estimation of Chromium in Ferrochromium by Electro-metric Titration. G. L. KELLEY and J. A. WILEY (*J. Ind. Eng. Chem.*, 1921, **13**, 1053—1054).—Twenty grams of sodium carbonate are fused in a nickel crucible and then cooled, the crucible being rotated during the cooling so that the carbonate forms a lining. A mixture of 16 grams of sodium peroxide and 1 gram of the sample is then fused for three minutes in this crucible, the heat being so regulated that the sodium carbonate lining is not fused; when cold, the contents of the crucible are dissolved in 300 c.c. of water, the solution is boiled for thirty minutes, cooled, 80 c.c. of sulphuric acid (d 1.58) are added, the solution is boiled for a further five minutes, cooled, filtered, and the filtrate diluted to 1 litre. One hundred c.c. of this solution are treated with 25 c.c. of sulphuric acid and titrated with ferrous ammonium sulphate solution, using the apparatus described previously by the authors and Adams (A., 1917, ii, 512).
W. P. S.

Tungsten. HERBERT LAVERS (*Proc. Austral. Inst. Min. Met.*, 1921, 101—152).—The paper contains a short description of the chemistry and metallurgy of tungsten together with a summary of the various processes that have been described for the estimation of tungsten in low grade ores. A volumetric method is recommended in which the tungsten is precipitated with cinchonine hydrochloride from acid solutions, and the precipitate dissolved in ammonium acetate, an excess of lead acetate added, and the excess determined by titration with ammonium molybdate. [Cf. *J. Soc. Chem. Ind.*, 1922, Feb.]
A. R. P.

Estimation of Thorium in Monazite Sand by an Emanation Method. HOMER H. HELMICK (*J. Amer. Chem. Soc.*, 1921, **43**, 2003—2014).—A method of estimating thorium in monazite sand is described. The sample is sieved through a 40 per cm. mesh and dried at 115—120°. Two grams are well mixed with 5 grams of potassium hydrogen fluoride and 5 grams of anhydrous, recently fused, metaphosphoric acid, and slowly brought to the highest temperature obtainable with a Méker burner in a 35 c.c. platinum crucible. A further 5 grams of metaphosphoric acid are slowly added during the heating; when the mass is clear, it is allowed to cool. After cooling, 20 c.c. of 80% orthophosphoric acid are added, and the crucible is heated in an air-bath at 250—255° for three hours, the solution being automatically stirred with a platinum wire; in this way a viscous solution is obtained. A small vessel made of glass and fitted with a ground stopper and inlet and outlet tubes is suspended in a vessel of concentrated sulphuric acid at 190—200° and the liquid from the crucible poured in. The solution vessel is removed from the acid, allowed to cool, and the remaining contents of the crucible are washed in with two quantities of 20 c.c. of water, the total volume made up to 75 c.c.,

and the vessel closed in an air-tight manner. The electroscope measurements are made, (i) with a blank, prepared in exactly the same way as the sample except that the monazite is omitted, (ii) with a standard solution containing a known amount of thorium, and (iii) with the sample solution. The percentage of thorium is calculated by means of the expression $X = AT_s(T_b - T_u)/T_s(T_b - T_s)$ in which X is the percentage of thorium in the sample, A the percentage in the standard, T_s the time of discharge of the electroscope by the standard, T_b by the blank, and T_u by the sample under investigation. The main sources of error of Cartledge's emanation method (A., 1919, ii, 120) were adsorption of thorium X by suspended matter in final solutions, by filters, and by the walls of the vessels used. These errors are avoided in the present method by producing a complete solution of the sand without filtration and in a single vessel. Other improvements made use of are a highly efficient form of vessel to contain the solution during de-emanation; the use of an automatic regulator to maintain constant pressure in the ionisation chamber during measurements; maintenance of optimum pressure gradient along the air current line through the measuring apparatus, thus assuring a minimum error on account of pressure variations, and maximum speed of measurement; protection of the insulation of the electroscope by means of a current of dry air. Analyses by this method gave results agreeing very well with results obtained by gravimetric methods, and required much less time for each determination.

J. F. S.

Detection of Bismuth in Urine. PIERRE AUBRY (*J. Pharm. Chim.*, 1922, [vii], 25, 15—18).—Following the administration of bismuth salts the metal is at least to some extent eliminated in the urine. In certain cases, it manifested its presence as a black precipitate of bismuth sulphide, but in other cases no precipitation occurred, and the metal was detected by evaporating to dryness, calcining the residue, dissolving the ash in hot dilute nitric acid, and adding a reagent containing 1% of quinine sulphate and 2% of potassium iodide dissolved in slightly acidified water, which produces an orange-red coloration or precipitate, according to the amount of bismuth present. This reagent is sufficiently sensitive to detect 1 part of Bi_2O_3 in 600000 parts of water. G. F. M.

Estimation of Alcohols by Acetylation. H. WOLFF (*Chem. Umschau*, 1922, 29, 2—3).—The acetylation is carried out by weighing out 0.5 gram of the sample in a test-tube 0.6—0.8 cm. wide and 10 cm. long. One c.c. of acetic anhydride is added and the tube sealed, the lower end being kept in cold water the while. The tube is heated for one hour in a boiling water-bath, removed, and allowed to cool. It is then placed in a well-stoppered, thick-walled flask, and broken by vigorous shaking. The stopper is bound on and the flask heated at about 50° for half an hour on the water-bath with constant shaking, cooled, and neutralised to phenolphthalein. Twenty-five c.c. of $N/2$ -alcoholic potash are

added and enough alcohol to make a clear solution. The flask is either left over-night or warmed for a quarter to half an hour at 50–60°, cooled, and titrated back with *N*/2-acid. The results obtained are within less than 1% of the theoretical. Solvents such as light petroleum or benzene have no effect on the result. If esters are present, their saponification value must be allowed for.

H. C. R.

New Methods of Blood Sugar Estimation. Estimation of the True Sugar Content of Urine. D. G. COHEN-TERVAERT (*Nederl. Tijdschr. v. Geneesk.*, 1921, 65, ii, 857–864, 3065–3069).—I. The methods of Schaffer and Hartmann (A., 1921, ii, 417), Folin and Wu (A., 1919, ii, 308), and of Ponder and Howie (A., 1921, ii, 417) deserve full recommendation. The first-named has been worked out for 0.1–0.2 c.c. of blood.

II. Sumner's method (A., 1921, ii, 564) has been compared with a fermentation method due to Nagasaki (*Nederl. Tijdschr. v. Geneesk.*, 1915, ii, 1478) and found to be very accurate. A few small modifications were introduced.

G. B.

A Source of Error in Testing Urine for Dextrose with *o*-Nitrophenylpropionic Acid. GEORGES RODILLON (*J. Pharm. Chim.*, 1922, 25, 56–57).—Dextrose is not the only reducing agent likely to be found in urine which will reduce *o*-nitrophenylpropionic acid to indigotin. The presence of hydrogen sulphide will both show this reaction and give a positive result in the sodium nitroprusside reaction for acetone. It does not reduce Fehling's solution. If the tests are carried out after clarifying the sample with lead acetate, these misleading results are avoided.

H. C. R.

The Identification of Lævulose in Presence of Aldoses. I. M. KOLTHOFF (*Chem. Weekblad*, 1922, 19, 1–2).—To 2 c.c. of the 1% sugar solution are added in succession 4 c.c. of *N*/10-iodine solution and 5 c.c. of 2*N*-sodium hydroxide. This order of addition must not be reversed. The mixture after shaking is left for one to one and a half hours and the excess of iodine removed by addition of a few drops of *N*-thiosulphate. Two c.c. of Fehling's solution No. II and 2 c.c. of Fehling's solution No. I are added, the tube is shaken, and warmed in a boiling water-bath for not more than five minutes. A red coloration in one minute shows 5% fructose in presence of glucose; in two minutes, 2½%, and in four minutes 1% of fructose. After five minutes, glucose alone gives the red colour.

The test detects 0.2 mg. of lævulose in presence of 10 mg. of dextrose, 10 mg. of sucrose, and 10 mg. of lactose.

S. I. L.

Estimation of Sugar by Titration with Alkali of the Cuprous Oxide Precipitated from Fehling's Solution. A. HANAK (*Z. Unters. Nahr. Genussm.*, 1921, 42, 248–250).—The cuprous oxide obtained from the inverted sugar solution containing not more than 0.5% of invert-sugar, and 50 c.c. of Fehling's solution is washed, dissolved in aqua regia, diluted to 250–300 c.c. with water free from carbon dioxide, and carefully neutralised with

sodium hydroxide so that it gives a pale greenish-yellow colour with methyl-orange. Phenolphthalein is added and $N/2$ -sodium hydroxide run in until the red colour remains for three minutes in the boiling solution. One c.c. of $N/2$ -sodium hydroxide = 0.0159 gram of copper. H. C. R.

Detection of Traces of Lactose in Urine by Formation of Formaldehyde. KURT HERZBERG (*Biochem. Z.*, 1921, 119, 81—92).—The injection of lactose for testing the functioning capacity of the kidney necessitates a ready method for its recognition in urine. Concentrations of 0.2% can be recognised by isolation of the osazone, oxidation of the latter with permanganate to formaldehyde which can be detected by a colour test, for example, the reaction with peptone and ferric chloride. H. K.

Polarisation of Normal Sucrose Solution. VLAD. STANĚK (*Z. Zuckerind. čechoslov.*, 1921, 45, 417—423; 425—431).—After correction for water, ash, invert-sugar, error of the polarimeter, etc., a normal solution of sucrose (recrystallised) polarised 99.9° Ventzke, using a dichromate filter, or 100.1° Ventzke, when a filter was not used. W. P. S.

Use of Invertase for Sucrose Estimation. T. SWANN HARDING (*Sugar*, 1921, 23, 546—547).—Errors in acid hydrolysis of sucrose by the Clerget method include the hydrolysis of other sugars present and the effect of the acidity on optical non-sugars. Using invertase, no effect is found on compounds present other than sucrose. Invertase can be obtained of sufficient strength to invert a 10% sucrose solution in two hours. The activity of invertase may be retained for a year and a half.

CHEMICAL ABSTRACTS.

Detection of Fatty Acids by the Formation of their Sodium Uranyl Salts. J. BARLOT and (Mlle) M. T. BRENET (*Compt. rend.*, 1922, 174, 114—116).—Streng's reaction for the micro-chemical detection of sodium (cf. *Ber. oberhess. Ges. Nat. Heilkunde*, 1883, 22) based on the formation of a characteristic crystalline precipitate of sodium uranyl acetate with uranyl acetate in the presence of acetic acid, gives positive results if acetic acid is replaced by its homologues in which there is an even number of carbon atoms in the straight chain. In the case of derivatives of acetic acid, the reaction depends on the nature of the substituent. The chloroacetic acids do not give the reaction, but sodium phenylacetate and uranyl nitrate give, at once, crystals of the double salt, $\text{CH}_2\text{Ph}\cdot\text{CO}_2\text{Na}, (\text{CH}_2\text{Ph}\cdot\text{CO}_2)_2\text{UO}_2$. W. G.

[**Estimation of Arachidic Acid.**] J. PRITZKER and R. JUNG-KUNZ (*Z. Unters. Nahr. Genussm.*, 1921, 42, 232—241).—See this vol., i, 208.

A rapid Method for Determining the Acetyl Value of Oils and Fats. ALEXANDRE LEYS (*J. Pharm. Chim.*, 1922, 25, 49—56).—The following values are determined: the saponification

value S of the oil or fat, the saponification value S' of the acetylated oil or fat and the ratio K between the weight of the acetylated matter and that of the oil or fat from which it has been prepared. The acetyl value A is given by $A = S' - S/K$. To determine K , a weighed quantity of the oil or fat is boiled with ten times its volume of acetic anhydride under a reflux condenser for two hours, cooled, diluted with benzene, and transferred to a tared dish. The benzene and excess of acetic anhydride are evaporated on a water-bath and the acetylated oil or fat is weighed. The acetyl value of an oil or fat dissolved in a neutral solvent having no acetyl value may be obtained without evaporating off the solvent as follows. One weighed portion (P') of the mixture is acetylated and the excess of acetic anhydride removed as above. The saponification value S' of this is then obtained. A further portion is saponified without acetylating, giving a saponification value S . $S' - S = A_0$ represents the quantity of potassium hydroxide required to neutralise the acetic acid which has combined with 1 gram of the mixture. This has increased its weight by $42 \times A_0 / 56 = 0.75 A_0$. Therefore P' grams of fat has become $P'' = P'(1 + 0.75 A_0)$ grams of acetylated fat. A saponification value S'' can therefore be calculated, based on the weight of acetylated oil or fat. The acetyl value $A = S'' - S/K$.

H. C. R.

Detection of Vegetable Oils in Animal Fats. The Phytosteryl Acetate Test. C. F. MUTTELET (*Ann. Falsif.*, 1921, 14, 327—333).—The cholesteryl acetate obtained from butter, lard, or beef fat (by treating the fatty acids at 70° with alcoholic digitonin solution, collecting the precipitated compound, converting it into the acetate and recrystallising the latter twice from alcohol) has m. p. 114.0° to 114.3° , whilst the phytosteryl acetate obtained in a similar way from coconut oil or earthnut (arachis) oil has m. p. 124.5° to 126.5° . This method will detect the presence of 10% of vegetable oil in an animal fat.

W. P. S.

Presence and Estimation in the Total Lipoid Ether-soluble Phosphorus of Phosphorus Compounds other than Phosphatides. (MLLE) ELIANE LE BRETON (*Bull. Soc. Chim. Biol.*, 1921, 3, 539—546).—The ordinary indirect method of estimating lecithins as ether-soluble phosphorus is subject to an error, unless the material is purified by precipitation with acetone, according to MacLean (A., 1914, i, 1197). About 20% of impurities remain in the acetone, which contains free and combined glycerol as well as phosphorus. These impurities are not formed by hydrolysis of lecithin during extraction.

G. B.

Titrimetric and Spectrometric Analysis of Keto-Enol Mixtures. KARL VON AUWERS and HELENE JACOBSEN (*Annalen*, 1922, 426, 161—236).—The general plan of the series of researches of which this paper forms a part has already been fully described (A., 1918, ii, 381), and it is now shown that by following the method of computation previously indicated spectrochemical data may be employed to estimate with moderate certainty the proportion

of enol in a keto-enol mixture. It is true that the method breaks down in certain cases owing to the difficulty of estimating the optical constants for one or other of the pure constituents; K. H. Meyer's titration method also gives somewhat uncertain values in particular cases. However, when the indications of both methods are definite they are in remarkably good agreement.

The following substances have been examined. The percentages represent the enol-content estimated spectrochemically, whilst the figures in brackets are the values determined by titration with bromine: ethyl acetylmalonate, 70% (69); ethyl propionylmalonate, 43% (44); ethyl *n*-butyrylmalonate, 55% (55); ethyl isobutyrylmalonate, 47% (47); ethyl isovalerylmalonate, 57% (55); ethyl hydroxymethylenemalonate, 100% (94); ethyl hydroxymethylene-ethyl ketone, ?% (<100); hydroxymethylenecyclohexanone, assumed 100% (100); 3-methyl-6-hydroxymethylenecyclohexanone, <100% (91%); 3:4-dimethyl-6-hydroxymethylenecyclohexanone, ?% (89); hydroxymethylenementhone, ?% (97); acetylacetone, 82% (84); propionylacetone, 76% (72); methylacetylacetone, 33% (36); ethylacetylacetone, 31% (29); propylacetylacetone, 31% (32); benzoylacetone, 100% (100); propionylacetophenone, 93% (94); butyrylacetophenone, 88% (90); isovalerylacetophenone, 89% (85); methylbenzoylacetone, 6% (9); methyl benzoylacetate, 20% (19); ethyl benzoylacetate, 24% (21); ethyl α -benzoyl-*n*-butyrate, 2% (4). C. K. I.

Colour Reactions of Phenacetin and Acetanilide. L. EKKERT (*Pharm. Zentr.-h.*, 1921, 62, 735—737).—Under definite conditions, phenacetin and acetanilide give different colorations when hydrolysed with sulphuric acid and then oxidised with potassium dichromate. If 0.1 gram of phenacetin is boiled for one minute with 5 c.c. of 10% sulphuric acid, the solution then cooled, diluted to 5 c.c., and treated with 2 drops of 1% potassium dichromate solution, a violet-red or red coloration is obtained. Acetanilide under similar treatment, but when boiled for two minutes, yields gradually a greenish-blue coloration. W. P. S.

Alkalimetric Estimation of Amino-acids and Peptides. RICHARD WILLSTÄTTER and ERNST WALDSCHMIDT-LEITZ (*Ber.*, 1921, 54, [B], 2988—2993).—The acid of ammonium salts can be estimated alkalimetrically with phenolphthalein as indicator if the aqueous solution of the salt is mixed with a sufficient amount of alcohol, since ammonia does not affect the indicator in alcoholic solution. It is essential that the solution should contain about 97% of alcohol and that relatively much indicator should be used. Amino-acids and polypeptides show a similar behaviour. Characteristic differences are shown, however, in the concentration of the alcohol which is necessary for the elimination of the action of the amino-groups or of hydroxyl ions. The polypeptides, peptones, and proteins behave in the same manner as the ordinary carboxylic acids in solutions containing 40% of alcohol, whereas amino-acids of the aliphatic series or of aliphatic character require an alcoholic

concentration of about 97% to produce this effect. Ethyl alcohol can be replaced by propyl alcohol, which appears to be even more effective, but not by methyl alcohol. The behaviour enables amino-acids and polypeptides to be estimated simply in mixtures of the substances by titrating with alkali hydroxide solution to neutrality towards phenolphthalein in 50% and 97% alcoholic solution. If a and b are the volumes of alkali solution used, the proportion, x , required by the amino-acids (since the majority of the latter and, in any case, those which predominate in the general mixtures neutralise in 50% alcoholic solution 28% of the amount required for complete neutralisation) is $100(b-a)/72$ and the proportion used by the polypeptides is $b-x$. H. W.

Estimation of Urea. F. MEZGER (*Pharm. Zentr.-h.*, 1921, 62, 719—721).—A simple gasometric method is described. One c.c. of the urine is placed in a small tube and this in turn is placed in a reaction bottle containing 30 c.c. of hypobromite solution and connected with the top of a burette; the latter is filled with water to the zero point by means of a levelling reservoir and serves as a measuring vessel for the nitrogen evolved when the urine and hypobromite solution are mixed. A definite volume (for example, 1 c.c.) of 2% urea solution is treated in the same way and at the same time in an exactly similar apparatus. This is taken as the standard and the amount of urea in the urine is calculated from the ratio of the two volumes of nitrogen obtained. W. P. S.

Quantitative Gravimetric Micro-analysis of Urea. Application to the Estimation of Urea in 1 c.c. of Blood. MAURICE NICLOUX and GEORGES WELTER (*Compt. rend.*, 1921, 173, 1490—1493).—The use of Fosse's xanthidrol reagent (cf. A., 1914, i, 859; ii, 506) is combined with Pregl's micro-methods of analysis (cf. *Die quantitative organische Mikroanalyse*, Berlin, 1917). To 1 c.c. of solution containing not more than 0.2 mg. of urea are added 1 c.c. of glacial acetic acid and 0.2 c.c. of 5% xanthidrol solution in methyl alcohol. The mixture is stirred, and after thirty minutes the precipitate is collected in a Neubauer micro-crucible and washed alternately with saturated solutions of xanthyl-carbamide in alcohol and water and finally with two drops of distilled water. The precipitate is dried at 105—110° and weighed on a micro-balance. For the estimation of urea in serum, 1.0 c.c. or 0.5 c.c. is used. It is diluted with exactly five times its volume of water, and a volume of Tanret's reagent equal to the original volume of the serum is added. After five minutes the precipitate is filtered off and, in 1 c.c. of the filtrate, the urea is precipitated as above. The weight of xanthylcarbamide obtained is thus just equal to the weight of urea present in 1 c.c. of the original serum. W. G.

Estimation of Uric Acid in Blood. CH. O. GUILLAUMIN (*J. Pharm. Chim.*, 1922, [vii], 25, 5—15).—The following *modus operandi* is recommended for the colorimetric estimation of uric

acid in blood: The blood or serum is first freed from albuminous substances by mixing 4 c.c. with 4 c.c. of 10% sodium tungstate solution, 28 c.c. of distilled water, and 4 c.c. of 2.7% sulphuric acid, shaking vigorously, and filtering until a clear filtrate is obtained. For ordinary clinical work, the uric acid may then be estimated directly in the filtrate by adding to 20 c.c. 1.5 c.c. of 40% solution of crystallised sodium carbonate, 0.5 c.c. of the sodium phosphotungstate reagent, and sufficient water to make 25 c.c.; and comparing in a colorimeter with a standard sulphite solution of uric acid prepared according to the directions of Folin and Wu (A., 1919, ii, 308). For a more exact estimation, it is preferable to separate the uric acid from the dealbuminised blood by means of silver lactate as described by Folin (*loc. cit.*). This separation of the uric acid is, however, always necessary if working with a solution containing red corpuscles, and by so doing it was demonstrated that blood usually contains a slightly lesser quantity of uric acid than serum. The author does not favour the dealbuminisation of the blood by Grigaut's method, using 20% trichloroacetic acid.
G. F. M.

Detection and Estimation of Morphine and Other Alkaloids in Animal Excreta and Organs. CURT WACHTEL (*Biochem. Z.*, 1921, 120, 265—283).—Previous methods for the estimation of morphine in animal tissues are very laborious and time-consuming. A new process is described which takes about one and a half days. Urine is submitted to a preliminary purification by basic lead acetate and the morphine, in the lead-free filtrate, precipitated by phosphotungstic acid in weakly acid solution. The washed precipitate is decomposed in alkaline solution by sodium potassium tartrate and the morphine, in solution, oxidised to ψ -morphine by an excess of a standard solution of potassium ferricyanide. The excess of the latter is estimated iodometrically. When the morphine is present in tissues a preliminary extraction is made by means of acidified alcohol. The extract from brain and muscle in addition to purification by basic lead acetate has to be submitted to a further purification by boiling with copper sulphate, the copper-free filtrate being then precipitated with phosphotungstic acid. The process is applicable to other alkaloids provided they are not adsorbed by the basic lead acetate precipitate. H. K.

The Bliss Method for the Separation of Strychnine from Quinine. L. E. WARREN and A. H. CLARK (*J. Amer. Pharm. Assoc.*, 1920, 10, 267—270; cf. A., 1920, ii, 276).—The method is shown to be untrustworthy. The quinine fractions invariably contained strychnine. In most cases the strychnine fraction was comparatively free from quinine so that the method may have a qualitative value in detecting the presence of strychnine in mixtures.
CHEMICAL ABSTRACTS.

The Estimation of Proteins in Blood—a Micro-method. PAUL E. HOWE (*J. Biol. Chem.*, 1921, 49, 109—113).—The author

uses 0.5 c.c. of blood and the micro-Kjeldahl method. For the fibrinogen and non-protein nitrogen, he follows the technique of Cullen and Van Slyke (A., 1920, ii, 398). For the rest, the method indicated in the following abstract. G. B.

The Use of Sodium Sulphate as the Globulin Precipitant in the Estimation of Proteins in Blood. PAUL E. HOWE (*J. Biol. Chem.*, 1921, 49, 93—107).—In order to apply the Kjeldahl method more readily, the author substitutes sodium sulphate for the ammonium salt. At 37°, critical zones in the precipitation are observed when 13.5, 17.4, and 21.5% of sodium sulphate are present in solution. These correspond respectively with the precipitation of euglobulin (as usually separated by carbon dioxide or saturated sodium chloride) and of that of two ψ -globulins I and II. G. B.

Reaction for Blood with Hydrogen Peroxide in statu nascendi. A. PATZAUER (*Chem. Ztg.*, 1921, 45, 1056).—A small quantity of magnesium peroxide is dissolved in very dilute acetic acid and, when violent evolution of gas ceases, guaiacum tincture or benzidine dissolved in acetic acid is added, followed by the ethereal extract of the blood. W. P. S.

Measuring Soil Toxicity, Acidity, and Basicity. R. H. CARR (*J. Ind. Eng. Chem.*, 1921, 13, 931—933).—The acidity or basicity of a soil and the amount of soluble iron and aluminium in the same may be estimated as follows: 50 grams of the soil are shaken for two minutes with 30 c.c. of a saturated solution of potassium thiocyanate in 95% alcohol; if, after settling, the liquid has a red colour, *N*/10-alcoholic potassium hydroxide solution is added until the colour disappears. Each c.c. of the alkali solution required is equivalent to 200 lbs. of calcium carbonate per acre. Should a red colour not appear on the addition of the thiocyanate, the mixture is titrated with *N*/10-alcoholic hydrochloric acid until a pink coloration is obtained. The relative amount of aluminium in solution is indicated by the depth of the blue colour produced when the thiocyanate extract of the soil is treated with a few drops of logwood tincture. Soluble iron and aluminium salts seem to be toxic to plants. W. P. S.

The Estimation of Tannin in Wines. PHILIPPE MALVEZIN (*Bull. Soc. chim.*, 1921, [iv], 29, 1087—1088; cf. A., 1912, ii, 612).—A reply to Clarens (cf. A., 1921, ii, 719). W. G.

General and Physical Chemistry.

Visible Refraction and Dispersion in Aqueous Solutions of Salts, particularly Coloured Salts. G. LIMANN (*Z. Physik*, 1921, 8, 13—19).—The refractive indices for the D , H_α , H_β , and H_γ lines have been measured for 0.3, 0.5, 1.0, 2.0, and 4.0*N* solutions of ferrous chloride, bromide and nitrate, copper nitrate and chlorate, ferric chloride, manganese chloride, bromide and nitrate; nickel chloride, bromide, nitrate and chlorate, cobalt chloride, bromide, nitrate and chlorate, aluminium chloride and nitrate; potassium chromate, dichromate, ferrocyanide and ferricyanide, chromic chloride and bromide, lithium sulphate, chromate and dichromate, magnesium chromate, caesium bromide and nitrate, glucinum chloride and nitrate, mercuric chloride, lead nitrate, and sodium chromate. Two constants are deduced from the measurements by means of which the refraction of a solution may be calculated if the equivalent conductivity for the particular concentration is known. The influence of temperature on the refraction and dispersion has been determined in the case of potassium bromide over the range 6—30°. The results show that the dispersion decreases markedly with decrease of temperature, but the refraction is not much influenced. Thus over the measured range of temperature the refraction changes only 1/1500 for a change of 1°.

J. F. S.

Eisenlohr's Refractometric Constants. WOJCIECH SWIĘTOSŁAWSKI (*Roczniki Chemji*, 1921, 1, 104—115).—A theoretical paper in which the author attempts to prove the constancy of atomic refraction of the atoms of organic substances. An analysis of the experimental data leads the author to the conclusion that the method employed by Brühl and Eisenlohr is not a rational method. This method of determination of the atomic refraction, based on the calculations of the mean value obtained from a large number of compounds, only serves to cancel the individual differences of the compounds investigated. The author has used a different method. For the calculation of the atomic refractions he takes the molecular refractions of the eight compounds, pentane, isopentane, hexane, octane, diisobutyl, diisomyl, cyclohexane, and cyclopentane, and has calculated the most probable mean value of the refractivity of the group CH_2 , ζ_{CH_2} , and that of hydrogen, ζ_{H} , by the method of least squares, the equations employed being of the type $M_D = n\zeta_{\text{CH}_2} + 2\zeta_{\text{H}} + \Sigma\Delta\zeta$, and $M'_D = n\zeta_{\text{CH}_2} + \Sigma\Delta\zeta$. In the solution of the equations, it is assumed that the increment $\Sigma\Delta\zeta = 0$. The values thus obtained are compared with those of Eisenlohr, $\zeta_{\text{C}} = (2.418) 2.490$, $\zeta_{\text{H}} = (1.100) 1.066$, $\zeta_{\text{CH}_2} = (4.618) 4.622$, Eisenlohr's values being bracketed. The results show an agreement for the value of ζ_{CH_2} , but a difference of about 3% for ζ_{H} . In the case of compounds containing oxygen, the equation $M_D = n\zeta_{\text{CH}_2} + \zeta_{\text{O}} + \Sigma\Delta\zeta$ is used. The figures obtained show considerable deviations of the

value $\Sigma\Delta\zeta$, that is to say, this increment cannot be regarded as zero. Similar results are obtained for both alcohols and ethers. The agreement obtained on comparing the author's value with those of Eisenlohr is explained by the dominant rôle of $n\zeta_{OH}$, and the insignificant influence of $\Sigma\zeta_x$, which corresponds with a residual refraction of the molecule in the equation $M_D = n\zeta_{OH} + \Sigma\zeta_x + \Sigma\Delta\zeta$. The author concludes that the methods employed by Brühl and Eisenlohr to determine the atomic refractivity mask the variability of this quantity and scarcely give any real knowledge of the refraction of the atoms.

J. F. S.

Molecular Coefficients of Refraction. K. VON AUWERS and H. KOLLIGS (*Ber.*, 1921, 55, [B], 21—45).—In recent publications, Eisenlohr (*A.*, 1920, ii, 717; 1921, ii, 1, 229) has pointed out that the "molecular coefficient of refraction" $M \times n_D^{20}$ is more sensitive to constitutive changes than the generally employed molecular refraction, and has calculated values for aliphatic, benzenoid, and polymethylene compounds; this work is now exhaustively criticised.

A protest is raised against Eisenlohr's method of deriving corrected values. Thus in the case of the *cyclohexane* series, he has assumed that the E values for the hydrocarbons may be derived from those of the corresponding, analogously constituted, alcohols and ketones by the deduction of 0.35. This rule is not fulfilled by 1-methyl*cyclohexan-4-ol* or 1-methyl*cyclohexan-4-one*, and the values for these substances have therefore been recalculated from the relationships that are supposed to exist among position-isomeric *cyclohexane* derivatives. It is shown, however, that these recalculated values lead to data for the refractive index which differ so widely from the experimental values that the latter could only have been obtained by the use of grossly impure material and that this assumption is devoid of experimental basis.

The work on the benzenoid hydrocarbons has been examined in detail. In correcting the observed values for the refractive index to a temperature of 20°, Eisenlohr has preferred the factor 0.00065 or 0.00060 for each degree to the generally employed factor 0.00045. An exhaustive survey shows that this is generally unjustifiable. In the cases of benzene, toluene, and, to a less extent, of ethylbenzene, the higher factor is to be preferred, but in the cases of seventeen other hydrocarbons it is shown experimentally that the lower factor is in very close agreement with the observations. Since the bulk of Eisenlohr's data for the constants of aromatic hydrocarbons are based on von Auwers's observations which have been regarded as erroneous when they do not harmonise with the theory, the values have been redetermined with more particular reference to *m*-ethyltoluene, *m*-propyltoluene, *p*-ethyltoluene, *p*-propyltoluene, and *p*-diethylbenzene. The previous specimens were obtained by the reduction of the corresponding ketones by Clemmensen's method. The latter are prepared by the Friedel-Crafts method and also according to the scheme $C_6H_4MeBr \rightarrow C_6H_4Me \cdot CHMe \cdot OH \rightarrow C_6H_4Me \cdot C(OMe)_2$, and the

physical constants of the products obtained in either manner are found to be identical. The constants of the hydrocarbons obtained by their reduction are in good harmony with those observed previously. A second method of preparation consists in replacing the hydroxyl group of the carbinols, $C_6H_4Me \cdot CHR \cdot OH$, by bromine and reduction of the bromides by sodium in the presence of moist ether; the indices of refraction of these preparations are mostly lower than those of the first group, practical identity being encountered only with *p*-propyltoluene. Thirdly, the Fittig synthesis is employed which gives specimens having refractive indices lower by 0.002—0.003 than those of the Clemmensen preparations. The discrepancies are far beyond the limits of experimental error. It appears impossible to decide which preparations have the greater claim to purity, since, on the one hand, the Clemmensen products are generally obtained smoothly and in good yield, whereas the "Fittig" specimens, although isolated from more complex mixtures, have constants which harmonise better with Eisenlohr's rules.

Examination of the new data obtained for the benzenoid hydrocarbons in accordance with Eisenlohr's principles leads the author to the following conclusions. Ortho-derivatives can be easily recognised by the markedly higher refractive coefficient, but this does not represent any advance, since it has been shown previously that they are differentiated so sharply from their position isomerides by density, refractive index, and specific exaltation that any of these factors is alone suitable for this purpose. The older methods do not allow the distinction between meta- and para-compounds and in this respect Eisenlohr's constant appears also to be useless; thus in the cases of *m*- and *p*-xylene the refractive indices do not differ by more than one part in a thousand parts and this position is not improved by multiplying them by their molecular weights. Position isomeric tri-derivatives are easily distinguished from one another by their different molecular refractive coefficients, but this was easily possible previously. The relationships of substances with four side chains cannot yet be elucidated.

The molecular coefficient of refraction cannot replace the spectrochemical constants, particularly the specific exaltation, but can only serve as an amplification of them. If it is desired to assign a substance to a particular group or to elucidate the general character of a group of substances, the most trustworthy data are to be derived from the specific exaltation, since in this the influence of homology or position isomerism is but little noticeable. When, on the other hand, it is desired to elucidate questions of structure within a group of substances, the molecular refractive coefficients are likely to be extremely valuable.

The following new data are recorded. Benzene, n_D^{20} 1.50092. Toluene, n_D^{20} 1.49653, n_D^{20} 1.49647. Ethylbenzene, n_D^{20} 1.49606. Propylbenzene, n_D^{20} 1.49241. *iso*Propylbenzene, n_D^{20} 1.49199. *m*-Xylene, n_D^{20} 1.49782. *p*-Xylene, n_D^{20} 1.49685, n_D^{20} 1.49680. 1-Methyl-2-ethylbenzene, n_D^{20} 1.50381. 1-Methyl-2-propylbenzene, n_D^{20} 1.49931. *o*-Cymene, n_D^{20} 1.501, n_D^{20} 1.50188. *p*-Cymene (from camphor), n_D^{20}

1-48885, whence n_D^{20} 1-48894 (from cymene-*p*-sulphonic acid), n_D^{24} 1-48942, whence n_D^{20} 1-49122. *p*-Diethylbenzene, n_D^{20} 1-49671. ψ -Cumene, n_D^{19} 1-50527, whence n_D^{20} 1-50523. Mesitylene, n_D^{20} 1-49981. Pehnitrol, n_D^{20} 1-51865. Ethyl- ψ -cumene, n_D^{15} 1-51047, whence n_D^{20} 1-50856. 1 : 3 : 5-Methyl-2-ethylbenzene, n_D^{20} 1-51167. *m*-Tolyl-methylcarbinol (from magnesium *m*-tolyl bromide and acetaldehyde), b. p. 112°/12 mm. 1-Methyl-3- α -bromoethylbenzene, $C_6H_4Me \cdot CHBrMe$, a colourless liquid, b. p. 101°/12 mm. ($\beta\gamma$ -*Di-m-tolylbutane*, $C_6H_4Me \cdot CHMe \cdot CHMe \cdot C_6H_4Me$, crystallises in slender, lustrous needles, m. p. 97°.) 1-Methyl-3-ethylbenzene (from *m*-methylacetophenone by Clemmensen's method), b. p. 159·7—160·5°, d_4^{19} 0·8669, d_4^{20} 0·867, n_a^{19} 1·49575, n_D^{19} 1·49966, n_D^{19} 1·51102, n_D^{19} 1·52056; (from 1-methyl-3- α -bromoethylbenzene), b. p. 159°, d_4^{19} 0·8622, d_4^{20} 0·862, n_a^{19} 1·49279, n_D^{19} 1·49650, n_D^{19} 1·50767, n_D^{19} 1·51678. *p*-Tolylmethylcarbinol, b. p. 108/12 mm., is oxidised to *p*-methylacetophenone, b. p. 105—106°/13 mm., d_4^{18} 1·0045, d_4^{20} 1·003, n_a^{18} 1·52880, n_D^{18} 1·53391, n_D^{18} 1·54835, n_D^{18} 1·56127, n_D^{20} 1·5332, which is identical with a product prepared by the Friedel-Crafts reaction. 1-Methyl-4- α -bromoethylbenzene, a colourless liquid, b. p. 105—106°/12 mm. 1-Methyl-4-ethylbenzene (from *p*-methylacetophenone by Clemmensen's method), b. p. 160·5°, n_D^{21} 1·49424, n_D^{20} 1·49490; from the Friedel-Crafts ketone), b. p. 161°, d_4^{18} 0·8650, whence d_4^{19} 0·8657, d_4^{20} 0·862, n_a^{19} 1·49372, n_D^{19} 1·49775, n_D^{19} 1·50889, n_D^{19} 1·51833; (from 1-methyl-4- α -bromoethylbenzene), b. p. 160°, n_D^{24} 1·49287, n_D^{20} 1·49481. *m*-Tolylethylcarbinol, b. p. 114°/12—13 mm. 1-Methyl-3- α -bromopropylbenzene, a colourless liquid, b. p. 114°/14 mm. 1-Methyl-3-propylbenzene (from the bromide), b. p. 180°, d_4^{17} 0·8646, d_4^{20} 0·863, n_a^{17} 1·48968, n_D^{17} 1·49340, n_D^{17} 1·5038; (from *m*-bromotoluene, propyl bromide, and sodium), b. p. 177—178·5°, d_4^{20} 0·8601, n_a^{20} 1·48978, n_D^{20} 1·49321, n_D^{20} 1·50392. *p*-Tolylethylcarbinol is a colourless liquid, b. p. 114°/12—13 mm. 1-Methyl-4- α -bromopropylbenzene, a colourless liquid, b. p. 108°/13 mm. *p*-Propionyltoluene, b. p. 114°/14 mm. 1-Methyl-4-propylbenzene (from the corresponding bromide), b. p. 182°, d_4^{18} 0·8620, d_4^{20} 0·861, n_a^{18} 1·49278, n_D^{18} 1·49655, n_D^{18} 1·50765, n_D^{18} 1·51687; (from *p*-propionyltoluene, according to Clemmensen), b. p. 183°, d_4^{20} 0·860, d_4^{18} 0·8617, n_a^{18} 1·49269, n_D^{18} 1·49641, n_D^{18} 1·50752; (by reduction of the ketone prepared by the Friedel-Crafts method), b. p. 182·5°, n_D^{20} 1·49542; (from *p*-bromotoluene, propyl bromide, and sodium), b. p. 181—181·5°, d_4^{23} 0·8554, d_4^{20} 0·858, n_a^{23} 1·48706, n_D^{23} 1·49065, n_D^{23} 1·50119. 1-Methylcyclohexan-4-ol, b. p. 172—173°, d_4^{18} 0·9192, d_4^{20} 0·916, n_a^{18} 1·45742, n_D^{18} 1·45959, n_D^{18} 1·46558, n_D^{18} 1·47025, n_D^{20} 1·4579. 1-Methylcyclohexan-4-one, b. p. 169·2°, d_4^{19} 0·91685, d_4^{20} 0·917, n_a^{19} 1·44285, n_D^{19} 1·44509, n_D^{19} 1·45110, n_D^{19} 1·45595, n_D^{20} 1·4450. H. W.

Occurrence of Spark Lines (Enhanced Lines) in the Arc.
I. Lead and Tin. G. A. HEMSALECH and A. DE GRAMONT (*Phil. Mag.*, 1922, 43, [vi], 287—306).—A convenient method is

described for obtaining the arc spectra of volatile metals. This consists in placing a piece of the metal on a sheet of copper (5 cm. \times 10 cm. \times 1 mm.) and connecting to the positive pole. The negative pole is a pointed carbon rod; the arc is struck by lowering the point until it touches the metal and then withdrawing it a little. The metal melts and forms a globule and the arc between it and the carbon point burns steadily. The heat generated is dissipated by the copper plate. Experiments are described which seem to indicate that spark lines are brought out in the arc when the degree of ionisation is reduced, as, for example, by blowing air through the arc. The effect of liquid media on the character of the arc spectrum of the metal has been studied by placing a drop of the liquid between the poles before striking the arc. The spark lines, or so-called high temperature lines, disappear from the arc when the cathode is formed by white-hot carbon, but they are brought out prominently when the electrodes are cooled down to about -190° by immersion in liquid air. A detailed account is given of the observations on the relative behaviour of various types of lead and tin lines in the arc under various conditions of discharge. It is definitely shown that the presence of hydrogen is not essential for the excitation of spark lines. Spark lines attain a high degree of development only in a medium which offers a comparatively high resistance to the flow of electricity. When hydrogen is present in the medium, either free or in combination, its spectrum resembles that which is only observed with high tension condenser discharges; namely, its lines are symmetrically broadened as though under the influence of a strong electric field (Stark effect).

J. F. S.

Röntgen Absorption Spectrum of Chlorine. AXEL E. LINDH (*Z. Physik*, 1921, 6, 303—310).—The Röntgen absorption spectrum of chlorine has been examined, using free chlorine and the compounds lithium, sodium, potassium, and thorium chlorides, potassium chlorate, potassium perchlorate, ammonium stannichloride and dichlorotetra-aquochromium chloride, $(\text{Cr}[\text{H}_2\text{O}]_4\text{Cl}_2)\text{Cl}$. It is shown that with univalent chlorine in all cases the absorption edge has the same wave-length, within the limits of the experimental error. In the case of quinque- and septa-valent chlorine, a displacement of the absorption edge toward shorter wave-lengths is observed which is greatest in the case of septavalent chlorine. A metallic Röntgen tube and new apparatus for photometric measurements both designed for measurements of the above-named type are described.

J. F. S.

Series Regularities in the Resonance Spectrum of Iodine. R. MECKE (*Z. Physik*, 1921, 7, 73—85).—A theoretical paper in which, on the basis of Wood's measurements of the resonance spectrum of iodine (A., 1911, ii, 82, 950; 1912, ii, 325, 1018; 1913, ii, 994; 1914, ii, 233; "Researches in Physical Optics," II, 1919), which is shown to consist of a series of doublets of the band type and is expressed by the series formula $\nu = A + Bm + Cm^2 + Dm^3$, the author has examined the series regularities of this spectrum.

It is shown that the first constant is given invariably by the exciting line, whilst the two final coefficients, C and D , maintain their values at all exciting lines. On the other hand, the second factor varies, but this may be represented in the form $B=b_0-b_1n$, in which n is a whole number, so that a number of partial series can be differentiated. In the case of the most accurately investigated resonance spectrum, that of the green mercury line, eight such series of doublets can be characterised, which can also, according to the position of the components of the doublets, be divided into two groups. In one group, b_1 has the value 0.655, and, in the other, 0.75, and n can have the values 0, 1, 2, and 3. For $n=0$ an apparent triplet is indicated. Further regularities are probable in the series, but they cannot be definitely characterised. It appears that the value of m represents an azimuth quantum number. In consequence of the very small moment of inertia, however, the rotation cannot be ascribed to the molecule itself; it is therefore assumed that the moment of inertia is due to a rotating electron ring. On this assumption, the series formula can be developed, and resolved into the two energy terms. Further, it is shown that resonance spectra, in the same way as line spectra, are brought into existence by electron springs.

J. F. S.

Divergence from Stokes's Law during the Excitation of the Fluorescence of Iodine Vapour. PETER PRINGSHEIM (*Z. Physik*, 1921, 7, 206—216).—The resonance spectrum of iodine vapour has been examined. It is shown that on heating the vapour from the ordinary temperature to 320°, the intensity of the anti-Stokes members increases many times, and at the same time also that of the members of positive order number some become stronger whilst the majority become less bright. It is thought probable that these oppositely behaving resonance lines belong to series which originate from some of the seven iodine absorption lines covered by the mercury line.

J. F. S.

Difference between the Absorption Spectrum and the Complete Fluorescence Spectrum of Iodine Vapour indicated by Lenz's Theory. PETER PRINGSHEIM (*Z. Physik*, 1921, 8, 126—131).—Theoretically, it is to be expected that the members of higher order number in the resonance spectrum of cold iodine vapour will not be markedly absorbed; this conclusion is fully confirmed by experiment. Correspondingly, the fluorescence spectrum of iodine vapour excited by white light is not a simple reversal of the absorption spectrum, but is considerably richer in lines, and so the colour change of the fluorescence of iodine vapour of high density is explained by a partial reabsorption of the fluorescence light. In iodine vapour at higher temperatures, resonance lines of higher order number are also strongly absorbed. The colour change in the emission spectrum of iodine vapour of constant density, occasioned by fluorescence or electric discharge at higher temperatures is considered, and it is shown that the cause cannot, at present, be indicated theoretically.

J. F. S.

Ultra-red Reflection Power of Silica. CLEMENS SCHAEFER and MARTHA SCHUBERT (*Z. Physik*, 1921, 7, 313—315).—The reflection spectra of various varieties of silica has been measured; the varieties include quartz, chrysoprase, hornstein, chalcedony, opal, and quartz-glass. It is shown that all varieties except the last two exhibit an identical quantitative reflection power. The reflection maximum at about 9μ exhibits about 90% of the metallic reflection. In the case of opal and quartz-glass, the reflection maximum lies in the same position as in the other varieties, but the reflection power has fallen to 40—50%. This result is attributed to a distortion of the space lattice in the amorphous varieties of silica. It is remarkable that the water maximum, which lies at 3.2μ , is not observed in the case of opal and it is suggested that this is to be attributed to the probable fact that the water is not arranged in the space lattice in opal. J. F. S.

The L Doublet of Neon. WALTER GROTRIAN (*Z. Physik*, 1921, 8, 116—125).—From a consideration of the recent work on the systematics of examination of Röntgen spectra, the view is put forward that only one permanent condition of unexcited atoms exists, and that the different absorption band edges are to be explained, not by different initial conditions, but by different final conditions. From this, the conclusion is drawn that of the inactive gases helium has one ionisation-potential, neon three, argon five, and krypton seven. In the case of neon, it is shown that the doublet difference, $L_2 - L_1$, is identical with the constant A found by Paschen for the neon spectrum, $A = 782 \text{ cm}^{-1}$. The absolute value of the ionisation potential of neon has as yet not been determined with certainty. J. F. S.

Characteristic Ultra-red Frequencies of Selenates and Chromates. CLEMENS SCHAEFER and MARTHA SCHUBERT (*Z. Physik*, 1921, 7, 297—308).—The reflection spectra of potassium, manganese, nickel, copper, zinc, and cadmium selenates, an isomorphous mixture of lithium and sodium selenate, magnesium ammonium selenate, manganese ammonium selenate, nickel ammonium selenate, cobalt ammonium selenate, nickel potassium selenate, and zinc potassium selenate have been determined in the region of the ultra-red, using ordinary light. It is shown that all selenates exhibit in the region 11μ a definite maximum of the reflection power, which is to be attributed to the vibrations of the selenate radicle. This maximum is analogous to the maximum at 9μ exhibited by the sulphates (cf. A., 1916, ii, 505, 506; 1918, ii, 282, 315). In those cases where the selenates contain water of crystallisation, the water bands previously recorded (*loc. cit.*) are also observed. The crystals have also been examined in polarised light, and it is shown that the reflection maximum at about 11μ can be resolved into two groups of characteristic vibrations in the case of uni-axial crystals and into three in the case of bi-axial crystals. Similar measurements were made with potassium chromate, sodium potassium chromate, magnesium ammonium chromate, and potassium dichromate. In the first three cases

using ordinary light, a single well-developed maximum is found, which lies respectively at 11.16μ , 11.2μ , and at complex maximum 11.3μ and 11.42μ . In the case of potassium dichromate, three maxima are observed, the first two of which are complex and lie at 10.44μ and 11.1μ , 12.14μ and 13.22μ , and 18.12μ . Magnesium ammonium chromate has been examined in polarised light and exhibits a maximum at 11μ which falls into three groups: 11.46μ parallel to the a axis, 11.28μ parallel to the b axis, and 10.36μ parallel to the c axis.

J. F. S.

Characteristic Ultra-red Frequencies of Chlorates, Bromates, and Iodates. CLEMENS SCHAEFER and MARTHA SCHUBERT (*Z. Physik*, 1921, 7, 309—312).—The reflection spectra of sodium, potassium, silver, and barium chlorates, sodium, silver, cadmium, and barium bromates and potassium iodate have been measured. It is shown that in all cases but that of sodium bromate, where a single maximum is observed, two reflection maxima are exhibited. The maxima have the following wave-lengths: sodium chlorate, 10.04μ , 16.04μ ; potassium chlorate, 10.12μ , 16.22μ ; silver chlorate, 10.76μ , 16.38μ ; barium chlorate, 10.24μ , 19.6μ ; sodium bromate, 12.24μ ; silver bromate, 12.62μ , 13.10μ ; cadmium bromate, 12.32μ , 13.20μ ; barium bromate, 12.3μ , 12.64μ , and potassium iodide, 12.48μ , 13.16μ . The values are discussed in connexion with those found for other acid radicles (cf. preceding abstract).

J. F. S.

Röntgen Spectra. ELIS HJALMAR (*Z. Physik*, 1921, 7, 341—350).—Röntgen spectra of the elements copper to tantalum have been measured. It is shown to be probable that the β_1 lines of the K series of the lighter elements are not simple. This has been established in the case of sulphur. The new line is regarded by the author as the continuation of the β_2 line of the next higher element, and the theoretical significance of this is discussed. The valency of sulphur appears to have no action on the emission spectra, but the experimental difficulties experienced in the work make it advisable to leave this question open. Exact measurements of a number of lines of the K -series are recorded, and the earlier measurements of lines of the L series are controlled. Two long tables of wave-lengths measured in the present work are included in the paper.

J. F. S.

Systematics of Röntgen Spectra. D. COSTER (*Z. Physik*, 1921, 6, 185—203).—The author has measured the Röntgen spectra of the elements tantalum, tungsten, osmium, iridium, platinum, gold, thallium, lead, bismuth, thorium, and uranium, using the method adopted by Hadding (*ibid.*, 1920, 3, 369). The object of the work was to obtain measurements of the very weak lines and from these to examine various relationships and regularities obtaining among X-ray spectra. Tables of the lines measured are given in the paper and also of the L doublets; a table of the constant wave-length differences $\beta_2 - \beta_5$, $\gamma_1 - \gamma_6$, $\gamma_2 - \gamma_3$, $\beta_4 - \beta_5$, $\beta_2 - \beta_8$, and $\beta_2 - \beta_7$ is drawn up. The relationship $(L_3 - L_1) - M\gamma = L\beta_3 - L\beta_2$, put forward

by Smekal (A., 1921, ii, 615), is evaluated and found to be only in moderate agreement with the experimental data; the divergences between the two sets of values are attributed to errors in the (L_2-L_1) values.

J. F. S.

Systematics of Röntgen Spectra. ADOLF SMEKAL (*Z. Physik*, 1921, 7, 410—412).—A reply to Coster (preceding abstract). The author points out that Coster's criticism is based on a misunderstanding of his method of treatment.

J. F. S.

Fluorescence of Cadmium Vapour. J. STEPH. VAN DER LINGEN (*Z. Physik*, 1921, 6, 403—404).—Light from cadmium sparks was allowed to pass through an evacuated quartz globe at 280° containing a scrap of cadmium. It is found that an absorption line 2288 Å.U. appears at this temperature, and with changing vapour density a band symmetrical with this line appears. When the band extends to 2307 Å.U. a second absorption line, 3260 Å.U. is observed. To ascertain whether the emitted light is due to resonance or fluorescence, the cadmium light around 2288 Å.U. was isolated and passed through the globe. It was found that at a density corresponding with that at which the line 2288 Å.U. is emitted, the vapour shows an ultra-violet fluorescence band which is sharply terminated at 2288 Å.U. and fades at 2314 Å.U. At higher densities the head disappears at 2288 Å.U., and the band is then terminated at 2313 Å.U. and extends to 3005 Å.U., where it also ends sharply. Between 2745 and 3005 Å.U. the band contains sixteen lines which are closer together the smaller the wave-length.

J. F. S.

Dependence of the Intensity of the Fluorescence of Dyes on the Wave-length of the Exciting Light. S. I. VAVILOV (*Phil. Mag.*, 1922, [vi], 43, 307—320).—It has been shown by Nichols and Merritt (*Physical Rev.*, 1910, 31, 376, 381) that the specific fluorescence of resorufin and eosin increases toward long wave-lengths. The present paper describes experiments with fluorescein, eosin-S extra, and rhodamine-B extra, designed to test how far this result is general. It is shown that within the limits of the errors of observation the specific fluorescence of the dyes mentioned is independent of the wave-lengths of their absorption bands. This result is equivalent to the conclusion that the absorption curves of the dyes are physically simple ones. Einstein's theory is not confirmed, but the deviations required by this theory are so small that they are only a little greater than the experimental errors, and consequently no very definite statement can be made. The intensity of fluorescence radiated by a definite molecular resonator depends only on the value of the absorbed energy and on the mechanism of the resonator. In the case of excitation by white light, the intensity of fluorescence F can therefore be expressed by the equation

$$F = K \int_{\lambda_1}^{\lambda_2} I_{\lambda} \cdot xc \cdot e^{-xod} \cdot d\lambda$$
, where I_{λ} is the energy of the exciting source in the wave-length interval $\lambda \dots \lambda + d\lambda$, xc the coefficient of absorption of the solution over the same wave-length interval, C the

concentration, and K the specific fluorescence. The result of Nichols and Merritt (*loc. cit.*) probably only shows the physical complexity of the bands of the dyes examined by them. J. F. S.

Ionisation brought about by Quinine Salts. GERHARD C. SCHMIDT (*Z. Physik*, 1921, 8, 160—164).—It is shown that quinine nitrate as well as quinine sulphate (Le Bon, *Compt. rend.*, 1900, 130, 891) emits light on warming and makes the surrounding air a conductor; other salts of quinine have no such action. All quinine salts when heated to just above the melting point change into a modification which gives intensely green fluorescent solutions instead of the usual violet fluorescent solutions. Quinine hydrogen sulphate in aqueous or chloroform solution gives a violet fluorescence, but when heated at 50° for an hour the fluorescence is light blue; when heated to the melting point this changes to an intense green which is maintained at very high dilutions. In the case of quinine valerate, the green colour changes back to violet on dilution when in water solution; in chloroform solution the heated salt gives a green fluorescence, but a violet fluorescence in benzene. The absorption spectrum of the green fluorescing solution is different from that of the violet fluorescing solution. The following hypothesis is advanced to explain the luminescence of the nitrate and sulphate when warmed, and its absence in other salts. On heating, these two salts pass into a new modification, which on stronger heating passes into the green, fluorescing variety; the change into the intermediate modification is accompanied by the emission of electrons or ions. In the case of the other salts, there is no intermediate compound formed, but a direct change into the green, fluorescing variety without emission of ions or electrons. J. F. S.

Action of Light on Silver Chloride, Bromide, and Iodide. PETER PAUL KOCH and FRITZ SCHRADER (*Z. Physik*, 1921, 6, 127—131).—The results of experiments, made to ascertain the loss of weight of particles of silver chloride, silver bromide, and silver iodide on illumination, previously published (*A.*, 1921, ii, 289) and a large number of other experiments made by the same method are collected and considered. The experiments were made with the suspended particles in air, dried air, dry nitrogen, moist nitrogen, moist argon, and dry argon, and a number of improvements were made in the technique of the method. The results show that silver chloride illuminated by intensities up to 3,000,000, silver bromide up to 12,000,000, and silver iodide up to 5,600,000 candle metre seconds (the light coming from the positive crater of an arc through 80 cm. of 1% copper sulphate solution) do not undergo a change in weight greater than 1—2%. The larger changes previously recorded are attributed to disturbances in the earlier experiments, among which the secondary reaction of silver bromide with the surrounding gas is mentioned. The rate of colouring of the silver haloids by light has also been determined; it is shown that silver bromide, after an illumination of 5 seconds by a light of intensity 3200 candle metre seconds, shows a definite coloration, silver chloride shows a coloration in three minutes when illuminated by

940 candle metre seconds, and silver iodide after ten minutes' illumination by the same light shows only a weak greyish-black coloration. In all cases the coloration increases with the duration of the illumination.

J. F. S.

Luminous Path of α -Rays in Crystals. H. GEIGER and A. WERNER (*Z. Physik*, 1921, 8, 191—192).—An experiment is described which answers the question, "Is the number of metallic centres, which are excited by an α -particle on its entry into a crystal, sufficiently great in every case to bring about a visible scintillation?" A thin, highly polished section of artificial willemite was arranged in the field of a microscope so that the α -rays from a polonium preparation should strike it at a small angle. At a magnification of 400, point scintillations were not seen, but luminous lines of 0.02 mm. length. These represent the path of the α -particle in the crystal, and show that the number of centres is extremely large, and in the case of perfect crystals sufficient to ensure a scintillation for every α -particle hitting the crystal.

J. F. S.

The Meitner Nuclear Model of the Radio-elements as the Basis of a Relationship between the Range and Total Number of Nuclear Particles of the α -Radiators. MAXIMILIAN CAMILLO NEUBURGER (*Z. physikal. Chem.*, 1921, 99, 327—331).—A theoretical paper in which, on the basis of the Meitner nuclear model, a relationship between the range and the total number of nuclear particles of the α -radiators is deduced. The range of most of the α -radiators of the uranium, thorium, and actinium families is calculated by means of this new relationship and found to be in good agreement. The divergence of the two sets of values lies between $\pm 0.8\%$ and $\pm 1.8\%$. Applying this relationship to the hypothetical elements uranium III and uranium IV, it is found that the former is practically inactive whilst the latter has a much shorter life than uranium I. This is further evidence against the existence of these hypothetical isotopes (cf. this vol., ii, 85).

J. F. S.

Beta Rays and Atomic Number. J. L. GLASSON (*Phil. Mag.*, 1922, [vi], 43, 393—396).—A theoretical paper in which the relationships between the atomic absorption, α , and the atomic scattering, β , of β -rays respectively and the atomic number are considered. The values of α , β , the atomic absorption $a = \alpha A/D$, the atomic scattering $b = \beta A/D$, the atomic weight A , the density D , and the atomic number N are tabulated for a number of elements from magnesium to bismuth. It is shown from the table that the values of a are approximately constant for elements of the same period of the periodic system; the elements examined fall into four groups which have values for a in the ratio 1 : 2 : 3 : 4. These groups are magnesium and aluminium, $a = 89$; iron, cobalt, nickel, copper, and zinc, $a = 172$ —196; palladium, silver, and tin, $a = 260$ —268; and platinum, gold, lead, and bismuth, $a = 348$ —372. It therefore seems likely that the value of a is a periodic function of the atomic number. The value of b increases with the atomic number of the element and on plotting $\log b$ against $\log N$ a curve is obtained

which is represented by $\log b = 2.09 \log N - 0.44$ and therefore $b = 0.36 \times N^{2.09}$. J. F. S.

Scattering of Röntgen Rays by the Atoms of a Crystal. W. L. BRAGG, R. W. JAMES, and C. H. BOSANQUET (*Z. Physik*, 1921, 8, 77—84).—A theoretical paper in which the experimental results previously published (A., 1912, ii, 477) on the scattering of Röntgen rays by the sodium and chlorine atoms of rock-salt crystals have been used in connexion with the arrangement of the electrons inside the atom. In the case of the sodium atom, the expected amplitude curve has been calculated for the case that of the eight exterior electrons four vibrate in di-quantic circular orbits and four in di-quantic elliptical orbits, and it is shown that such a model has a very satisfactory agreement with the experimental facts. J. F. S.

Characteristic X-Rays from Boron and Carbon. A. LL. HUGHES (*Phil. Mag.*, 1922, [vi], 43, 145—161).—The photoelectric effect of the radiation from carbon and boron bombarded by electrons has been measured as a function of the energy of the electrons. Two slight but definite breaks in the curves for both carbon and boron were obtained. These are considered to correspond with the *K*- and *L*-absorption wave-lengths. The breaks occur at 215 volts ($\lambda = 57.5$) and 34.5 volts ($\lambda = 358$) for carbon and 148 volts ($\lambda = 83.5$) and 24.5 volts ($\lambda = 505$) for boron. The *L* point for carbon is in good agreement with the measured values recently obtained by Millikan (*Astrophys. J.*, 1920, 52, 47) for the *L* emission lines of carbon vapour. The *K* points for carbon and boron are approximately where they would be expected on extrapolating the known values of the *K_a* critical absorption wave-lengths for elements heavier than magnesium. J. F. S.

Absorption of the *K* X-Rays of Silver in Gases and Gaseous Mixtures. P. W. BURBIDGE (*Phil. Mag.*, 1922, [vi], 43, 381—389).—The total absorption coefficient of silver *K* radiation has been measured in air, carbon dioxide, sulphur dioxide, and methyl iodide vapour. Using the same silver *K* radiation, it has been shown with mixtures of sulphur dioxide and air, carbon dioxide and air, sulphur dioxide and carbon dioxide, and methyl iodide and air that the absorption in gases is atomic (as in the case of solids and liquids) and therefore additive in mixtures and compounds. J. F. S.

Experimental Decision of the Question of the Radioactivity of all Elements. II. G. HOFFMANN (*Z. Physik*, 1921, 7, 254—259; cf. A., 1920, ii, 575).—Making use of the very sensitive method of measuring α -ray activity previously described (*loc. cit.*), the author has extended his measurements to platinum. The measuring instrument itself shows an α -ray activity represented by 0.3—0.5 α -particle per hour. Using a cleaned platinum dish of 39 sq. cm. surface, an activity of 14.6 per hour for the whole dish is recorded, or 0.39 per sq. cm. The measurements were made over a period of about fifty hours. The results are analysed for α -rays which

may come from known α -radiators of all ranges uniformly distributed through the mass of the platinum and for α -radiators situated on the surface of the platinum, and the analysis indicates a definite α -ray activity of platinum.

J. F. S.

Protactinium Content of Pitchblende Residues and the Branching Relationship of the Actinium Series. OTTO HAHN and LISE MEITNER (*Z. Physik*, 1921, 8, 202—204).—A criticism of the value put forward by St. Meyer (*Mitt. Inst. Rad. Forsch.*, 1920, 127) for the branching relationship of the actinium series. The value, 4.2%, is about 25% higher than that found by the present authors, $3 \pm 0.3\%$. It is shown to be probable that Meyer's preparation contained 1—2% of ionium, which would explain the difference in the two figures.

J. F. S.

Existence of Isotopes of Uranium and the Meitner Nuclear Model. MAXIMILIAN CAMILLO NEUBURGER (*Z. physikal. Chem.*, 1921, 99, 321—326).—The author discusses the position of uranium- Z_2 in the disintegration series from the point of view of Meitner's nuclear model. The position given by Hahn (*A.*, 1921, ii, 498) in the series $U_{III} \xrightarrow{\alpha} UZ_1 \xrightarrow{\beta} UZ_2 \xrightarrow{\beta} U_{IV}$ is criticised, and it is shown, for several reasons, that only two elements of the uranium type, uranium I and uranium II, can possibly exist. The elements uranium II and uranium III are shown to be identical and the

disintegration series: $U_I \begin{cases} \xrightarrow{\alpha'} UX_1 \xrightarrow{\beta} UX_2 \xrightarrow{\beta} U_{II} \\ \xrightarrow{\alpha} UZ_1 \xrightarrow{\beta} UZ_2 \xrightarrow{\beta} U_{II} \end{cases} \begin{cases} \nearrow Io \\ \searrow UY \end{cases}$ is

put forward to represent the disintegration of uranium. A method for the detection of isotopes of the fourth order (this vol., ii, 107) is described. It consists in determining the number of particles emitted by the disintegrating substance; should this number be in keeping with the disintegration constant, then the substance does not consist of isotopes of the fourth order, but of a single element.

J. F. S.

A New Radioactive Substance. A. PICCARD and E. STAHEL (*Physikal. Z.*, 1922, 23, 1).—A quantity of uranium-X, separated from uranyl nitrate by the ether-water method, has been repeatedly measured with respect to its β -radiation for a prolonged period. It is shown that in addition to the hard β -radiation of uranium-X, there is a soft radiation which with increasing age of the preparation decreases more and more slowly; this indicates the presence of a substance of longer life than uranium-X. This substance has been provisionally named uranium-V. It has an approximate half life of forty-eight days, or about twice as long as uranium- X_1 . The β -radiation due to this element is half absorbed by an aluminium sheet 0.003 mm. thick, that is, the absorption coefficient is $2300 \text{ cm.}^{-1} \text{ Al}$. It is shown that this substance may be a member of the actinium series, but it has not been decided whether it or uranium-Y is the first thorium isotope in the disintegration series put forward by one of the authors.

J. F. S.

Some Oxidising Properties of Thorium-X. PIERRE LEMAY and LÉON JALOUSTRE (*Compt. rend.*, 1922, 174, 171—172).—Thorium-X exerts a very marked catalytic action in the oxidation of adrenaline and morphine, the oxidation being much more rapid than in the presence of manganese salts. Oxidation could not, however, be proved in the case of the primary alcohols of the fatty series. W. G.

Ionisation Tension of the Halogen Hydrides. PAUL KNIPPING (*Z. Physik*, 1921, 7, 328—340).—The ionisation tension of hydrogen cyanide, chloride, bromide and iodide has been determined by the method previously described by Franck and Knipping for helium (A., 1920, ii, 72). The following values are recorded: hydrogen cyanide, 15.5 volts, 357 Cal.; hydrogen chloride, 14.4 volts, 331 Cal.; hydrogen bromide, 13.8 volts, 317 Cal.; and hydrogen iodide, 13.4 volts, 308 Cal. The electron affinity, in Cal. and volts, the limiting wave-length, and frequency have been calculated for chlorine, bromine, and iodine, and the values compared with those of other investigators. The following values are recorded: chlorine, 98 Cal., 4.26 volts, $\lambda=2890$, $\nu=34600$; bromine, 68 Cal., 2.96 volts, $\lambda=4150$, $\nu=24100$, and iodine, 61 Cal., 2.65 volts, $\lambda=4600$, $\nu=21500$. J. F. S.

Ionisation and Excitation Tension of Nitrogen. ERICH BRANDT (*Z. Physik*, 1921, 8, 32—44).—The ionisation tension of nitrogen has been found to be 17.75 ± 0.1 volt with reference to the resonance tension of helium, 20.5 volts. Higher ionisation stages have been found at 25.41 ± 0.1 volt and 30.72 ± 0.2 volt, both with reference to the value 17.75 volts. The limits of light excitation of the nitrogen molecule by electron collisions have been investigated, and an analogy between the light-electric curve and that of band emission is pointed out. J. F. S.

Electrical Conductivity of Zinc Sulphate Solutions in the Presence of Sulphuric Acid. HERMAN V. TARTAR and HARMON E. KEYES (*J. Ind. Eng. Chem.*, 1921, 13, 1127—1129).—The conductivity of zinc sulphate solutions increases with the sulphuric acid concentration, but the conductivity of sulphuric acid is decreased by the addition of zinc sulphate; when magnesium sulphate is added to a strongly acid zinc sulphate solution, an additional decrease in conductivity occurs. The addition of small quantities of gelatin does not affect the conductivity. The temperature coefficients of conductivity vary, and are a function of the acid and zinc concentration. During the electrolysis of zinc sulphate solutions there is an increase in the volume of the solution, the increase amounting to 1.5% for each 100 grams of zinc deposited. W. P. S.

Anomalous Dissociation in Aqueous Solutions. A. J. RABINOWITSCH (*Z. physikal. Chem.*, 1921, 99, 338—360).—By anomalous dissociation is understood the phenomenon which occasions a decrease in the molecular or equivalent conductivity

with increasing dilution, that is, the degree of dissociation, as expressed by $\alpha = \lambda/\lambda_\infty$, decreases with increasing dilution. With the object of finding a connexion between the viscosity and the electrical conductivity of solutions of salts in water, the author has determined the density, viscosity, and electrical conductivity of aqueous solutions of caesium chloride, cadmium chloride, cadmium bromide, cadmium iodide, zinc bromide, zinc iodide, and sodium iodide at 0°, 25°, and 50°. It is shown that in the case of caesium chloride the viscosity and the conductivity are very closely connected. A comparison of the corrected molecular conductivity curves and the relative viscosity curves at 25° and 50° for the halogen salts of cadmium with one another and with those of other salts shows that the form of the relative viscosity curve has a strong influence on the form of the $\lambda\eta_a$ curve. In those cases with a high value for η_a , the introduction of the usual viscosity correction of the $\lambda\eta_a$ curves can lead to an anomalous form of the curve because of over-correction. This anomaly is general in the case of highly concentrated aqueous solutions. The over-correction is explained as follows: In introducing the correction, it is assumed that the ionic mobility is inversely proportional to the viscosity of the medium. This assumption, however, does not hold because a reduction of the ionic diameter is brought about by increasing concentration, due largely to a dehydration of the ions. With increasing temperature, the anomaly extends over a considerable portion of the $\lambda\eta_a - V$ curve. This is in opposition to the view that the anomaly is due solely to the over-correction on account of a reduction of the ionic diameter. If the over-correction is due alone to the inapplicability of Stokes's law to all concentrations, then it is to be expected that the minimum, displaced on warming, should correspond with higher viscosity values; but since this is only observed in the case of caesium chloride, it follows that this cause of over-correction does not explain the anomaly in all cases. It is held that the anomalous form of the conductivity curve is not to be explained by the over-correction alone, but is due to several causes. It is suggested that the causes which bring about the "true anomalous dissociation" in non-aqueous solutions are operative in aqueous solutions also. This suggestion would explain the displacement of the minimum with increasing temperature, since with increasing temperature water in respect of its dielectric constant approaches in behaviour the other solvents. It is also shown that in concentrated aqueous solutions, in addition to the over-correction of the conductivity values, there is also a typical anomalous progression of the molecular and equivalent conductivity, which in all probability has its origin in the same causes as in non-aqueous solutions.

J. F. S.

Electrolytic Dissociation of Salts in Concentrated Solutions, and in the Fused and Solid States. A. J. RABINOWITSCH (*Z. physikal. Chem.*, 1921, 99, 417—433; cf. preceding abstract).—The conductivities, viscosities, and densities of aqueous solutions of caesium chloride, silver nitrate, ammonium nitrate, thallium

nitrate, and thallium silver nitrate were determined at high concentrations. In the case of thallium ammonium nitrate the results were extended to the fused salt. The molecular conductivities were extrapolated to the pure salts. After the application of the viscosity correction, all the molecular conductivity curves assumed an abnormal form, and on extrapolation to pure salt gave high values of $\mu\eta_\alpha$, where μ is the molecular conductivity and η_α the relative viscosity. The values of $\mu\eta_\alpha/\mu_\infty = \alpha$ were therefore abnormally high. The true values of α for the salts probably lie between the values calculated from the corrected and the uncorrected (μ/μ_∞) conductivities, and nearer the former. The course of the dissociation in very concentrated solutions is thus abnormal. The extrapolation was confirmed by the form of the curve for thallium silver nitrate, which continued smoothly up to the fused salt.

J. R. P.

Reversal of the Molecular Conductivity Curve and Abnormal Dissociation. A. J. RABINOWITSCH (*Z. physikal. Chem.*, 1921, 99, 434—453; cf. preceding abstract).—Abnormal dissociation occurs when the degree of dissociation (or the molecular conductivity) decreases with increasing dilution. It is most commonly observed with non-aqueous solutions, and is most marked when the dielectric constant of the solvent is small. The various hypotheses which have been advanced to account for the phenomenon are reviewed. In the ordinary method of representation of conductivities, the dilution V is taken as the variable volume of solution in which a constant weight of one gram equivalent of solute is dissolved. The author considers the results obtained when a constant weight of one gram equivalent of solvent is taken and the volume V_1 of solution containing it is taken as the dilution. If k is the specific conductivity, the product $kV_1 = \mu_1$ is the newly-defined molecular conductivity. The curves representing the values of μ_1 and V_1 are called "reversed conductivity curves." When only one constituent of the solution is capable of ionisation and the other is a strong ioniser (for example, acetic acid in water), the original molecular conductivity curve is normal but the reversed curve is abnormal, exhibiting a maximum but not a minimum, after correction for viscosity. When the ionising constituent is also a strong ioniser (for example, silver nitrate in aniline or water), the original curve is abnormal, without a maximum, or apparently abnormal (without correction). The reversed curve is normal (water in silver nitrate and thallium silver nitrate), or, in consequence of the uncorrected increase in viscosity is apparently abnormal (aniline in silver nitrate). When neither constituent is ionisable, but when its polymeride or solvate is ionisable, then in both cases abnormal curves result, in the latter case with a maximum, the position of which indicates approximately the maximum content of solvate in the solution. To the right and left of the maximum the conductivity decreases in consequence of the diminution of solvation due to insufficient quantity of solvent and to depolymerisation respectively. Examples of such curves are

furnished by acetic acid and aniline and by acetic acid and pyridine. A definition of abnormal dissociation is given. J. R. P.

Effective Potential Difference of Electro-osmosis and Allied Phenomena. H. FREUNDLICH (*Trans. Faraday Soc.*, 1921, 16, Appendix, 146—149).—A theoretical paper in which the author discusses the nature of the potential difference at work in electro-osmosis, cataphoresis, and allied phenomena. A great deal of experimental work is reviewed, and it is shown that the Nernst potential difference, ϵ , is in no sense identical with the electrokinetic potential difference, ζ ; the Nernst potential represents the total difference between the interior of the first phase and that of the second phase, the electrokinetic potential represents that portion of the difference which falls within the displaceable liquid layers. J. F. S.

Selenium [Cells]. WILHELM SPÄTH (*Z. Physik*, 1921, 8, 165—183).—The influence of illumination on the dielectric constant of selenium has been investigated in connexion with the fatigue and inertia of the element. It is shown in all selenium cells that the change in conductivity on illumination runs parallel with a change in capacity. The final capacity value and the final conductivity are reached with similar inertia phenomena. The conductivity of the cells increases two- or three-fold on illumination, but the capacity increases only 5%. A cell which, with respect to its conductivity, exhibits fatigue, also exhibits the same phenomenon with respect to its capacity. The dependence of the change in capacity on the intensity of illumination is much the same as that of the conductivity. With increasing intensity of illumination, the sensitiveness of the capacity increases in a parabolic manner. The inertia on darkening the cell is greater than the inertia on illumination as far as the capacity effect is concerned, and, further, it is considerably smaller in soft cells. The fatigue effect in soft cells on intense illumination is more marked, both with regard to the capacity change and the conductivity. The author has put forward an hypothesis of the mode of action of the selenium cell and has calculated the equilibrium of the cell for illuminated and dark conditions. J. F. S.

Fatigue of Alkali Metal Cells in the Neighbourhood of the Discharge Potential and the Influence of this on the Results of Photometric Measurements. H. ROSENBERG (*Z. Physik*, 1921, 7, 18—64).—Sodium, potassium, and rubidium photoelectric cells have been examined in connexion with the fatigue effects often noticed in these cells. It is shown that these cells, when filled as usual with an inactive gas, rapidly exhibit fatigue effects and also a rapid recovery. The effects are apparently not proportional to the intensity of the light or to the photoelectric current. In addition, slow changes of the sensitiveness of the cells are observed, which must be regarded as a type of fatigue effect. The cause of the fatigue is found to be due to the adsorption of a positively charged layer of gas by the alkali metal, but it is

not decided whether electron absorption or a reduction of the field is the ultimate cause of the fatigue effect. The process appears to be intimately connected with the appearance of a powerful collision ionisation. The pure photo-effect is shown to be strictly proportional to the amount of light which has entered the cell. Taking into account these phenomena, a photometric process has been tested which leads to intensity ratios which are accurate to 0.1%.

J. F. S.

Equilibrium Law of Electrolytes. K. JABLZYŃSKI and F. J. WIŚNIEWSKI (*Roczniki Chemji*, 1921, 1, 116—134).—The authors have deduced a dilution law of the form $n^{4/3}/n_0 = k$ and have used this to calculate the value of k for potassium, sodium, and lithium chlorides, using data obtained from cryoscopic measurements. It is shown that only in the case of potassium chloride is the value of k constant, whilst in the case of the other two salts it may have even negative values. These facts are explained by hydration of the ions. Corrections which take into account the hydration have been introduced into van't Hoff's equation, and lead to a remarkably constant value for k in all three cases; thus for potassium chloride $k = 3.09$, for sodium chloride 3.10, and for lithium chloride 2.93. Furthermore, the number of molecules of water combined with the various ions has been calculated as follows: potassium, 0; sodium, 3; lithium, 11; chlorine, 0. Hydrochloric acid behaves in the same way as the salts mentioned, and when the hydrogen ion is taken as combined with nine molecules of water a very constant value, 4.74, is obtained for k . The equation has been applied to those weak electrolytes which apparently follow Ostwald's dilution law. The value of k has been calculated for acetic acid from cryoscopic measurements and a remarkably constant value 0.00504 obtained, whereas the value of k calculated from Ostwald's equation increases considerably with increasing concentration. On applying the equation to ebullioscopic data of solutions of the chlorides mentioned above, it is shown that the value of k increases even more strongly than when cryoscopic data are used. This is explained by assuming that the ions are more heavily hydrated at 100° than at 0°. This is shown to be in keeping with facts. The value of k at 100° is smaller than that at 0°, which shows that the degree of ionisation decreases with increase of temperature. This the authors attribute to a fall in the dielectric constant of water with increase of temperature. A rule has been deduced, in keeping with facts, which shows that the equilibrium constant of salts in aqueous solution is proportional to the dielectric constant of the solvent. J. F. S.

Velocity of Sound in Air and Hydrogen at 0° and 1 Atm. E. GRÜNEISEN and E. MERKEL (*Ann. Physik*, 1921, [iv], 66, 344—364).—The velocity of sound in air is found to be 331.57 m./sec. and in hydrogen 1260.6 m./sec. The ratio of the specific heats for air is $C_p/C_v = 1.4034$ and for hydrogen 1.408, both at 0° and 760 mm.

J. F. S.

Specific Heats of Ammonia, Sulphur Dioxide, and Carbon Dioxide. J. R. PARTINGTON and H. J. CANT (*Phil. Mag.*, 1922, [vi], 43, 369—380).—The ratio of the specific heats $k=c_p/c_v$ for ammonia, sulphur dioxide, and carbon dioxide has been determined by a method depending on the relative velocities of sound in the gases and in air. The specific heats have been calculated using Berthelot's equation of state in all calculations. The following results were obtained at one atmosphere pressure: ammonia at 14.5° , c_p/c_v , 1.308, C_p 8.77 cal., C_v 6.70 cal.; sulphur dioxide at 13.2° , c_p/c_v , 1.290, C_p 9.47 cal., C_v 7.34 cal.; carbon dioxide at 20° , c_p/c_v , 1.303, C_p 8.76 cal., C_v 6.72 cal. J. F. S.

Numerical Values of the Gas Constants. F. HENNING (*Z. Physik*, 1921, 6, 69—72).—The data on which the gas constant is based have been critically discussed and, on the basis of the most trustworthy experimental data, the value of this constant, in different units, has been calculated. The following values are recommended for general use as the most accurate, $R=0.08204\pm0.00003$ litre-atm./degree mol.; $R=(8.313\pm0.003)10^7$ erg/degree mol.; and $R=1.986\pm0.001$ cal.₁₅/degree mol. J. F. S.

Theoretical Determination of the Chemical Constants of Monatomic Gases. E. BRODY (*Z. Physik*, 1921, 6, 79—83).—The author have deduced a mathematical expression for calculating the chemical constants of ideal monatomic gases on the basis of the first quantum theory and from statistical considerations. The expression obtained is the same as that deduced by Stern (A., 1919, ii, 219) and Planck (*Sitzungsber. Preuss. Akad. Wiss. Berlin*, 1916, 653). The expression has the form $C=\log_e (2\pi m)^{3/2} k^{5/2}/h^3$. J. F. S.

Chemical Constants of Sodium and Potassium. R. LADENBURG and R. MINKOWSKI (*Z. Physik*, 1921, 8, 137—141).—The chemical constants of sodium and potassium have been calculated from existing data on the specific heat, vapour pressure, and latent heat of fusion of these elements. In the case of sodium, the value of the chemical constant C is found to be 0.846, and on inserting this value in the equation $C=C_0+1.5 \log M$, where M is the molecular weight, the value of the constant C_0 is found to be -1.20 instead of -1.59 , as in the case of mercury, argon, hydrogen, cadmium, and zinc. The data on which the calculation is based are analysed with the object of finding the cause for the discrepancy. In the case of potassium, the value of C is found to be 1.016, which gives the value of C_0 as -1.37 . J. F. S.

Absolute Entropy and Chemical Constants. MAX PLANCK (*Ann. Physik*, 1921, [iv], 66, 365—372).—An answer to the criticism of Ehrenfest and Trkal (*ibid.*, 1921, 65, 609) in which the difficulties in calculation of entropy experienced in the author's book ("Theorie der Wärmestrahlung," 1921) are pointed out. It is also doubted whether the equation $s=k \log W$, in which W is the thermodynamic probability, will give the value of the entropy. The author answers the objections and criticisms and points out that they arise from a confusion of terms. J. F. S.

Equation of Condition. RUDOLF WEGSCHEIDER (*Z. physikal. Chem.*, 1921, 99, 361—382.)—A theoretical paper in which the equation of condition put forward by Wohl (this vol., ii, 117) is discussed and criticised. It is shown that this equation of condition leads to results which can scarcely be accepted; for example, the decrease of pressure at constant volume with increasing temperature in liquid systems, consequently this equation can have no theoretical significance. The critical point is always given by equations of condition which yield isotherms of the van der Waals form, by the coincidence of the three roots of the equation $v=f(p, T)$. The equality of four roots involves the introduction of a relationship between the constants; the equality of two roots leads to conditions which include, in addition to the critical point, also the maxima and minima.

J. F. S.

Physical Properties of Vapour-Liquid Systems. I. WOJCIECH SWIENTOSLAWSKI (*Roczniki Chemji*, 1921, 1, 276—296).—Starting from relationships based on van der Waals's hypothesis of corresponding conditions, $\tau=T/T_c$ and $d_c/d=K=f(\tau)$. Where T , d , T_c , and d_c are the temperature and corresponding density and T_c and d_c the same critical values, the author has deduced the following regularities from the experimental work of Young, Amagat, Mathias, Villard, Kamerlingh Onnes, and Crommelin. (1) The value of K or $\log K$ increases with increasing molecular weight of the liquid, and also when the liquid is associated. (2) Oxygen and argon have small values for K . In the case of associated liquids, it is shown that the concentration of non-associated molecules, y , is given by $y\% = 100K/K_A$, where $K=d_c/d_p$ for a non-associated liquid and K_A is the same ratio for an associated liquid at the same relative temperature ($\tau=\text{const.}$). The concentration of non-associated molecules in methyl alcohol and propyl alcohol increases proportionally to the temperature, but this is not true for ethyl alcohol. Ramsay's results for water and ethyl alcohol are satisfactory if it is assumed that the associated molecule of water contains three simple molecules and that of ethyl alcohol two simple molecules. The value of K plays an important rôle in all cases where the physical state of a system liquid-vapour is characterised. The relationship between the surface tension γ and the reduction of pressure below the meniscus in a capillary tube of radius r is given by $(2\gamma/r\Delta f)_{\tau=\text{const.}} = K = \text{const.}$, and in the case of osmotic pressure $(P/\Delta f)_{\tau=\text{const.}} = K = \text{const.}$ The constant of Trouton's rule may be calculated by the equation $W_{\text{mol.}} = bR \log_e K = 20.4$, in which $R = 2 \text{ cal.}$ $\log_e K = 2.303$, $\log_{10} K = 2.203 \times 2.38$ or 2.38 , depending on the ratio $K=d_c/d_p$ for the liquid at the relative boiling point $\tau=0.65$ under a pressure of one atmosphere.

J. F. S.

Physical Properties of Vapour-Liquid Systems. II. WOJCIECH SWIENTOSLAWSKI (*Roczniki Chemji*, 1921, 1, 297—304; cf. preceding abstract).—It is shown that over the range of relative temperatures $\tau=0.50$ to $\tau=0.95$ the expression $d_c^2/T \log_e K = B$ is constant, where d_c is the density of the liquid, and K the ratio d_c/d_p in

which d_p is the density of the vapour at the absolute temperature T . The deviations of B never exceed a mean amount of $\pm 2.3\%$. A minimum value of B is found between $\tau=0.85$ and 0.90 and a maximum about $\tau=0.60-0.65$. Associated liquids are exceptional; these show a continual increase in the value of B from $\tau=0.50$ to the critical temperature $\tau=1$.
J. F. S.

Vaporisation Coefficients of Solid and Liquid Mercury. M. VOLMER and I. ESTERMANN (*Z. Physik*, 1921, 7, 1-12).—The authors have determined the velocity of vaporisation of mercury at temperatures from 59° to -180° . This demands the determination of the fraction, (α), of the vaporised mercury which is condensed on a cooled surface, whilst the rest ($1-\alpha$) is reflected. It has been previously shown by Knudsen (*Ann. Physik*, 1915, [iv], 47, 697; 1916, 50, 472) that at 19.45° $\alpha=0.95$, but at -140° $\alpha=1.00$. For the temperature range $59-0^\circ$ the rate of vaporisation and consequently the value of α has been determined by measuring the rate at which a small globule of mercury deposited on the interior of the outside wall of a highly evacuated Dewar vessel filled with liquid air decreases. The Dewar vessel is kept in a thermostat at the required temperature during the experiment and the temperature of the bath up to 40° is taken as the temperature of the mercury, but for higher temperatures a correction is applied to the bath temperature to get the temperature of the mercury. This method is inapplicable at temperatures below 0° and above 60° . For these temperatures, 0° to -64° , the rate of vaporisation is determined by measuring the rate of change of the electrical resistance of the condensed film of mercury. A third method is also described which depends on the difference of readings in a graduated capillary tube due to vaporised mercury. The results show that for pure liquid mercury, irrespective of the temperature, the value of α is unity, but for solid mercury it is less than unity and decreases with decreasing temperature. It is shown that with sulphur, phosphorus, and benzophenone the value of α lies between 0.2 and 0.5 . It is also shown that there is a close connexion between α and the extent to which a liquid may be supercooled.
J. F. S.

Coefficients of Vaporisation and their Relationship to Ostwald's Step Rule. M. VOLMER and I. ESTERMANN (*Z. physikal. Chem.*, 1921, 99, 383-394).—The coefficient of vaporisation, α , has been determined for mercury over the temperature $+59^\circ$ to -64° , and found, for the liquid phase, to be equal to unity irrespective of the temperature. The known expression $G=1/\sqrt{2\pi R} \cdot \sqrt{M/T} \cdot p$ expresses actually the velocity of vaporisation for all temperatures: If this equation yields results for other substances similar to those obtained for mercury, then it becomes the first strictly true reaction velocity equation. In the case of solid mercury α is found to be approximately 0.9 . The ratio α (solid)/ α (liquid) has been determined for sulphur, phosphorus, and benzophenone and the values $0.3-0.4$, $0.4-0.5$, and 0.25 respectively have been obtained. The Ostwald step rule and the

limits of its applicability to the separation of the liquid and solid phases from the vapour condition are shown to be due to the different α -values of the solid and liquid phases. J. F. S.

The Separation of Miscible Liquids by Distillation. II. ARTHUR FELIX DUFTON (T., 1922, 121, 306—308).

Divergences of the Value of the Mechanical Equivalent of Heat. WOJCIECH SWIENTOSLAWSKI (*Roczniki Chemji*, 1921, 1, 171—177).—The author gives a short description of facts which show that the value of the mechanical equivalent of heat requires exact verification. It is shown to be probable that the values obtained by electrical and mechanical methods are not in agreement. The measurements of Jaeger and Steinwehr show that in bomb calorimetry conditions often exist which increase the amount of heat obtained by the introduction of electrical energy into a calorimeter. It is also pointed out that adiabatic calorimetry has not been employed in the measurement of this constant. J. F. S.

Heat of Vaporisation of Sodium and the Probability of the Transition of the Sodium Atom from the Resonance to the Normal Condition on the Basis of Optical Measurements. R. LADENBURG and R. MINKOWSKI (*Z. Physik*, 1921, 6, 153—164).—The authors have measured the magnetic rotation of the plane of polarisation in the immediate neighbourhood of the D line by the method described by Senftleben (*Ann. Physik*, 1915, [iv], 47, 949) at various temperatures of the emitting vapour. The sodium used was contained in an air-free glass tube in an electric furnace the temperature of which was constant to $\pm 0.1^\circ$. It is shown that the dependence of N (the number of electrons which vibrate with the atoms) on the temperature can be regarded as a relative vapour pressure curve, if the theory that N is proportional to the number of sodium atoms present is accepted. Calculation from the experimental results leads to the value for the heat of vaporisation of sodium $l_{T=0}=25.9$ Cal. This value is probably uncertain to about 2%. J. F. S.

Action of Forces between Separated Atoms in Diamond and Aliphatic Molecules. K. FAJANS (*Z. physikal. Chem.*, 1921, 99, 395—415).—A theoretical paper in which it is shown that if in analysing the results of heat of combustion experiments the assumption is made, as a first approximation, that only neighbouring atoms have a noticeable energetic action on one another, then it appears that the firmness of a C—C linking in the diamond is somewhat greater than that in aliphatic hydrocarbons. From this it follows that the heat of combustion of hydrocarbons with branched chains should be somewhat smaller than that of the straight chain isomerides, a conclusion which is in keeping with experimental data. To explain the above relationships it is assumed, (1) that the energy of the C—C linking in aliphatic substances and diamond is constant, irrespective of whether or no the carbon atom is bound to further carbon atoms or also with hydrogen. (2) That separated and distant atoms have a notice-

able energetic action on one another. As a second approximation in the quantitative analysis, only those linkings which lie on a common edge of the tetrahedron are considered. From a comparison of diamond with the hydrocarbons the relationship $y_m - 2x_m + h \sim 1 - 2 \text{ Cal.}$, between the energies of the three types of edge linkings (C-C, C-H, H-H), where y_m , x_m , and h are respectively the energies of formation of the three types of linking. This relationship indicates a method of explaining the difference in properties of the chain isomerides and also the exceptional position of the first member of homologous series. In the case of aliphatic hydrocarbons, alcohols, ketones, and esters of monobasic acids, the chain isomerides with low boiling point and heat of vaporisation possess a lower heat of combustion, a stronger saturation of the forces in the molecule corresponding with a weaker intermolecular force. The difference in the boiling points and the specific heats of the chain isomeride is therefore brought into relationship with the difference in the intramolecular energy, and the latter is attributed to the mutual action between separated and distant atoms.

J. F. S.

The Meaning of the Atomic Constants of Heat of Combustion and Molecular Refraction. WALTER HÜCKEL (*J. pr. Chem.*, 1921, [ii], 103, 241—248).—A theoretical paper, in which the author states the various relations that have been found by Steiger (A., 1921, ii, 473) and others between the various "atomic" and "linking" constants of organic compounds and of the elements forming them, and points out some relations that seem to follow.

W. O. K.

New Data in the Thermochemistry of Organic Substances. WOJCIECH SWIENTOSLAWSKI (*Roczniki Chemji*, 1921, 1, 305—315).—The heat of dissociation of hydrogen, chlorine, bromine, iodine, and solid carbon into individual atoms, determined during recent years, has been used to obtain the value of the constant ϵ of the equations $(C-I) = (H-I) + \epsilon - 3$; $(C-Br) = (H-Br) + \epsilon - 3$; $(C-Cl) = (H-Cl) + \epsilon$; $(C-C) = (H-C) + \epsilon$; $(C-N) = (H-N) + \epsilon$; $(C-S) = (H-S) + \epsilon + 2$; $(C-O) = (H-O) + \epsilon + 5$; $(C-H) = (H-H) + \epsilon + 15$. The symbols $(C-I)$, $(H-I)$ indicate the heat of formation of the single link between the atoms named. When the values 81.3 Cal., 106 Cal., 46 Cal., and 36 Cal. are taken as the heat of formation of the molecules of hydrogen, chlorine, bromine, and iodine, respectively, as represented by $2X = X_2 + Q \text{ Cal.}$, and the heat of formation of solid diamond as $-\overset{\cdot}{\underset{\cdot}{\text{C}}} = C_{\text{sol}} \times 287 \text{ Cal.}$, it is possible to calculate the heat of formation of the links $(C-C)$ and $(C-H)$. According to Fajans (A., 1920, ii, 354), $(C-C) = 137.5 \text{ Cal.}$ and $(C-H) = 117.0 \text{ Cal.}$ Using the above data in the equations, a mean value of 20.6 is obtained for ϵ . From the heat of formation of the linkings $(H-Cl) = 115.5 \text{ Cal.}$, $(H-Br) = 72.0 \text{ Cal.}$, $(H-I) = 52.7 \text{ Cal.}$, it is shown that $(C-Cl) = 136.3 \text{ Cal.}$, $(C-Br) = 89.7 \text{ Cal.}$, and $(C-I) = 70.3 \text{ Cal.}$ An analysis of the above results shows that the heat of formation of a link is an additive

quantity made up of parts due to the links of the atoms concerned. These parts have been calculated as follows: (C—)=68.8 Cal., (H—)=48.2 Cal., (Cl—)=67.4 Cal., (Br—)=23.8 Cal., (I—)=4.4 Cal. The heat of formation of a link between two atoms is the sum of the two atomic values. This regularity is probably existent in inorganic compounds also, for if the above atomic values of the halogens are subtracted respectively from the heat of formation of the corresponding haloids of potassium, the value of (K—) is found to be 91.2, 94.5, and 93.7 Cal., and by the same method the value for sodium (Na—) is 83.3, 84.0, and 82.7 Cal. J. F. S.

Heat of Coagulation of Ferric Oxide Hydrosol with Electrolytes. FREDERICK L. BROWNE and J. HOWARD MATHEWS (*J. Amer. Chem. Soc.*, 1921, 43, 2336—2352).—The heat of coagulation of ferric oxide hydrosol of various concentrations and purity by various concentrations of sodium oxalate, sulphate, chloride, and hydroxide, potassium ferricyanide, and sulphuric acid has been determined. It is shown that the dilution of ferric oxide hydrosols of varying purity is accompanied by measurable heat effects until the purity exceeds 21 or 22 (the ratio of gram equivalents of ferric oxide per litre to gram equivalents of chlorine per litre is termed the purity of the sol), beyond which point the heat of dilution is zero. If the heat of dilution per gram-equivalent of chlorine in the sol is plotted against the chlorine content, the curve is of the same nature as that for the heat of dilution of ferric chloride solutions of varying concentration. The thermal behaviour of ferric chloride solutions on dilution confirms the observations of Goodwin (*Physical Rev.*, 1896, 9, 251; 1900, 11, 193), to the effect that the hydrolysis of ferric chloride takes place in two steps, the first an instantaneous and the second a slow reaction. The mixing of ferric chloride solutions with common electrolytes involves very significant heat changes. The heats of coagulation have been reduced to curves showing the heat of coagulation as a function of the purity which vary widely in shape with the nature of the coagulant; the heat effect is usually positive, but with sodium sulphate and sodium chloride at low purities it is negative. As the purity increases, the heat of coagulation becomes smaller, and finally becomes negligible, except in the case of sulphuric acid. The peculiar behaviour of this last electrolyte is due to solution of the ferric oxide. When the heat of coagulation is plotted against the concentration of the coagulating electrolyte, smooth curves are obtained which show no break at the point of limiting concentration. The heat effects observed on coagulating ferric oxide sols with electrolytes are to be attributed to the action of the electrolytes in the coagulants on the electrolytes, particularly ferric chloride, present in the sol. The change in dispersity of the ferric oxide on coagulation does not involve a greater heat change than 1—2 cal./gram-equivalent of ferric oxide. In the case of ferric oxide hydrosols, there is no heat of peptisation analogous to the heat of solution of crystalloids. J. F. S.

Kation Volumes in Permutite. RICHARD LORENZ (*Z. Physik*, 1921, 6, 269—270).—A criticism of the work of Günther-Schulze

on the determination of the kation volumes in permutite (A., 1921, ii, 624). It is pointed out that the value found for the alkali ion is practically the same as that of the alkali atom. It is suggested that the alkali ions in permutite fill only a small amount of the space and have a large space to vibrate in. This great mobility is characteristic of permutite. The kation radii calculated by Günther-Schulze are probably too large, for the new determinations of Fajans and Herzfeld and also those of Landé are much smaller and lie very close to those calculated from the ionic mobility by Lorenz and Born.

J. F. S.

The Velocity of Extension of Thin Layers of Oils on the Surface of a Sheet of Water. PAUL WOOG (*Compt. rend.*, 1922, 174, 162—165).—For fatty oils, where all the molecules intervene, the velocity of extension is practically proportional to the ratio (the carboxyl groups of the glycerides + the double linkings + the acidity)/viscosity. In mineral oils, however, the velocity of extension depends only on the unsaturated molecules, which are few in number, and hence the viscosity plays an insignificant part. In mixtures of the two types of oil, the results depend on the relative proportions in which the active fatty oil and the comparatively inactive mineral oil are mixed.

W. G.

High Pressure Due to Adsorption. WILLIAM D. HARKINS and D. T. EWING (*Chem. News*, 1922, 124, 23—24, 31—35).—Cf. this vol., ii, 123.

J. F. S.

Electro-adsorption as a Purely Chemical Process. I. M. KOLTHOFF (*Kolloid Z.*, 1922, 30, 35—44).—The author discusses electro-adsorption (adsorption of ions) and from the experimental results of Freundlich, Odén, and others shows that the process is strictly a chemical one. The adsorption of copper, lead, morphine, codeine, papaverine, atropine, ferricyanide, oxalate, chromate, and iodate ions by norit carbon confirms this view. It is shown that the adsorption isotherm can be deduced directly from the stoichiometric relationships, if it is assumed that the electro-adsorption is a process in which a sparingly soluble substance, contained in the adsorbent, is converted by the adsorbed ion into another sparingly soluble substance. The coefficient $1/n$ of the adsorption isotherm bears a direct relationship to the valency of the reacting substances. If $1/n$ is equal to a for a univalent ion, then it has the value $a/2$ for a bivalent ion and $a/3$ for a trivalent ion. This relationship is confirmed by the present experiments and by the earlier work of Freundlich and Odén. The significance of the quantity of the adsorbent is explained by the characteristic constitution of its surface layer of particles. The active ion, which determines the charge in the surface layer, and in adsorption forms the insoluble salt, is present in this layer in far larger concentration than in the liquid. When equilibrium has been set up on the surface of a particle, then the concentration of the active ion, in contact with the excess of reacting ions, is small. In such a case, if fresh adsorbent is placed in the liquid, then the reacting ions

are again in contact with large concentrations of active ions, and the reaction commences anew. The general chemical theory in its relationships to colloid chemistry is shortly discussed. J. F. S.

Origin of the Charge of a Colloidal Particle and its Neutralisation by Electrolytes. JNANENDRA NATH MUKHERJEE (*Trans. Faraday Soc.*, 1921, 16, Appendix, 103—115).—The charge of a number of suspensoids has been shown to be due to the adsorption of a common ion and the nature of the chemical forces which cause this adsorption has been defined. The effect of the electrical charge of a surface (due to the disorption of ions) on ions of opposite sign has been theoretically investigated for the case when chemical action does not take place between the surface and the ion. It has been shown that the electrical forces are sufficiently strong to account for adsorption. The stability of the adsorption of an ion of opposite sign by electrical forces has been given a quantitative form, assuming that the charge of the surface consists of ions which can be treated as point charges widely separated from each other. Actual calculations justify this assumption. The reversal of the charge by electrical adsorption has been discussed. The following series of the adsorbability of kations by a negatively charged surface has been deduced from the theory: $\text{Th}^{+++} > \text{Al}^{+++} > \text{Ba}^{++} > \text{Sr}^{++} > \text{Ca}^{++} > \text{Mg}^{++} > \text{H}^+ > \text{Cs}^+ > \text{Rb}^+ > \text{K}^+ > \text{Na}^+ > \text{Li}^+$. This order is the same as that of the precipitating power of these ions for a number of suspensoids of widely differing chemical properties. Equations deduced from the theory agree satisfactorily with the experimental observations of Elissaffoff (A., 1912, ii, 419) on electro-endosmosis. The chemical affinity of the surface atoms is not always negligible, and for this reason the series given above cannot be expected to hold in all cases. J. F. S.

Electrical Theory of Adsorption. W. HARRISON (*Trans. Faraday Soc.*, 1921, 16, Appendix, 116—118).—A criticism of Mukherjee's theory of adsorption (cf. preceding abstract). The author is of the opinion that the double layer consists of a surface of rigidly fixed atoms under continuous bombardment of positively and negatively charged ions, any particular point on the rigid surface becoming in turn negative, neutral, and positive, these conditions arising in any order. The observed contact difference is the average effect of these conditions. When several kinds of atoms are present in the solution, the average number of any one of them at the surface will depend on their concentration, valency, and mobility. The variation of contact difference from negative to neutral and positive was observed with cotton and aluminium sulphate near the neutral point (*J. Soc. Dyers and Col.*, 1911, 27, 279; 1918, 34, 91). These variations occurred during the same experiment, the readings being direct measurements of *E.M.F.* developed by filtration under pressure. J. F. S.

Diffusion of Hydrogen through Iron and Platinum. GERHARD C. SCHMIDT and TH. LÜCKE (*Z. Physik*, 1921, 8, 152—159).—The diffusion of hydrogen through iron and platinum has

been investigated. A sheet of the metal was fixed between two glass vessels, hydrogen was electrolytically generated on one side of the sheet and the potential on the other side measured over a period of time up to about twenty-four hours. The metal sheets were of various thicknesses. It is shown in the case of iron that the passage of the hydrogen through the metal is always indicated by a fall of potential. In the case of thin sheets, the fall of potential is observed as soon as the electrolysis commences; the fall of potential is rapid at first, and gradually becomes slower until, when the liberation of hydrogen has proceeded for some time, it becomes constant. With sheets of medium thickness (0.5—1.0 mm.) the fall of potential is gradual, and with sheets thicker than 1.0 mm. there is no fall of potential. On breaking the polarising current, the potential increases, the more quickly the thinner the metal. From this fact it is deduced that the hydrogen diffuses into the liquid. With thicker sheets, the increase of potential is slower, which is explained by the longer time required for the relatively large volume of hydrogen to diffuse into the liquid. Similar results are obtained in the case of platinum. The results show that the passage of hydrogen across platinum plates is a diffusion and not, as stated by Nernst and Lessing (*Göttinger Nachr.*, 1902, 146), due to small pores in the metal. The Nernst-Lessing rule was shown by them to hold for palladium and thin sheets of platinum, but not for thicker sheets of platinum. It is shown that the reason for the rule holding with palladium is that on account of the great power of palladium for absorbing hydrogen practically none is liberated from the other side, and consequently the condition of the rule is obtained.

J. F. S.

Dispersoid Analysis of Non-aqueous Systems. WOLFGANG OSTWALD (*Trans. Faraday Soc.*, 1921, 16, Appendix, 89—93).—The author describes the preparation of membranes and jellies suitable for dispersoid analysis of solutions in alcohol of 97%, for both dialysis and diffusion experiments. The mercury sulphide alcosol, the preparation of which is explained, is an especially stable sol suitable for these tests. Solutions of dyes, resins, neutral salts, fatty acids, and soaps, as far as at present examined, produced by the spontaneous solution of these substances in 97% alcohol, are all molecular-dispersed, or contain, at any rate, molecular-dispersed particles in considerable proportions. Zein, the protein of maize which is soluble in alcohol, is also found to be dialysable.

J. F. S.

Solubility. III. Solubility of Metal Ammonia Salts in Salt Solutions. J. N. BRÖNSTED and AGNES PETERSEN (*J. Amer. Chem. Soc.*, 1921, 43, 2265—2292; cf. A., 1920, ii, 536).—A long list of the solubilities of eighty-six cobalt, chromium, and rhodium, complex ammine salts in water at 0° and 20° is given in the paper. The solubilities of a large number of complex ammine salts of cobalt and chromium in solutions of potassium chloride, formate, chlorate, trichloroacetate, dichloroacetate, monochloroacetate, thio-

cyanate, hydroxide, nitrate, and phthalate; sodium chloride, sulphate, chlorate, nitrate, formate, oxalate, and benzenesulphonate; magnesium sulphate and chloride; calcium formate, acetic acid; and mono-, di-, and trichloro-acetic acids of various concentrations have been determined at 0°. In the preceding papers (*loc. cit.*) it was shown that the expression $\log S/S_0 = a(C_i^{1/3} - S_0^{1/3})$ represents approximately the solubility changes of salts in hetero-ionic solvents up to about 0.1*N*-solutions, when a is a constant which for uni-univalent salt systems has the value 1/3, C_i is the concentration of the solvent salt, S_0 the solubility in water, and S that in the salt solution at the same temperature. The experimental data furnished for salts of the same type in the present paper agree approximately with the formula, and an equation of the same type is found for bivalent salts. The individual nature of the salts employed is more pronounced in the larger amount of material now presented and tends to obscure any numerical regularity in salt systems of promiscuous types. The experiments lead to the following rules. The influence of the solvent increases with increasing valency of the ion of the solvent salt, which, in the case of ter-bivalent salts, leads to an enormous increase of solubility, even on the addition of solvents of very low concentration. When both the dissolved and dissolving ions are of higher valency, the effect of the sign of the electric charge is very marked, producing the highest solubility values when the multivalent ions of solvent and solute are of different sign. The reason for the appearance of individuality in the solubility curves is to be sought partly in the hydration of the solvent and the solute, and partly in the fact that in these ions the mass is not to be regarded as a point, but as a particle of varying size bearing a number of electric charges of opposite signs, the algebraic sum of which gives the valency of the ion. It is probable that the size of the ion in the present cases will have an influence of considerable magnitude. J. F. S.

Probability of Spontaneous Crystallisation of Supercooled Liquids. C. N. HINSHELWOOD and HAROLD HARTLEY (*Phil. Mag.*, 1922, [vi], 43, 78—94).—The statistical investigation of the spontaneous crystallisation of supercooled salol, phenol, *p*-toluidine, and *o*-nitrophenol leads to the following conclusions: In all these cases crystallisation is provoked by colloidal organic dust particles, the activity of which diminishes in general as the result of heating or ageing. The effectiveness of the dust particles depends on their radius. If this is equal to the radius of a small particle of the solid which should thermodynamically be in equilibrium with the supercooled liquid, then crystallisation occurs at once. If the radius is less than this, the supercooled liquid has an average life depending on the discrepancy between the equilibrium radius and the radius of the particles present. The magnitudes of these quantities are discussed. The results of crystallisation experiments are given: 16 sets of 70 tubes of salol were kept at 32.3° in a thermostat and the number which had crystallised at various times noted; similar results are recorded for 12 sets of 41 tubes of phenol at 18—25°;

8 sets of 122 tubes of *p*-toluidine at 33.36—17.48° and 5 sets of 130 tubes of *o*-nitrophenol at 38.72—31.1°.

J. F. S.

Structure of [Hanging] Drops of Mixed Liquid Crystals. O. LEHMANN (*Ann. Physik*, 1921, [iv], 66, 323—343).—The structure of hanging drops of liquid crystals of *p*-hydroxyphenetole and cholesteryl benzoate has been examined between parallel Nicols by means of the dichroic effects produced. A number of diagrams and descriptions of the structure observed are included in the paper.

J. F. S.

Submicrons Visible in the Tyndall Cone to the Naked Eye. J. TRAUBE and P. KLEIN (*Kolloid Z.*, 1922, 30, 19—20).—When 30 c.c. of 1/30000 *N*-lead nitrate solution are treated with 2 drops of *N*/2-sodium carbonate solution in a rectangular vessel, stirred, and illuminated by a Tyndall cone, it is found that in forty to sixty seconds an extremely large number of small particles are visible by means of a hand lens. The particles are in violent and irregular motion. As the time of the existence of these particles increases more and more light is reflected from them, so that in about two minutes they are visible to the naked eye at a distance of 2 metres. It is held that the particles are crystals and are visible on account of their rotation, which causes an intermittent reflexion of light from the crystal faces. The phenomenon is not produced by all concentrations of lead nitrate, *N*/10000 is too concentrated and *N*/70000 is too dilute; the best concentration is that given above. The same phenomenon may be observed with very dilute solutions of calcium sulphate, calcium ammonium phosphate, barium sulphate, mercurous chloride, and lead chloride.

J. F. S.

Fibrous Structure of Hard Drawn Metal Wires. M. ETTISCH, M. POLANYI, and K. WEISSENBERG (*Z. physikal. Chem.*, 1921, 99, 332—337).—From the X-ray examination of hard and soft wires of copper, tungsten, iron, molybdenum, palladium, aluminium, silver, and zinc, it is shown that the crystallite in soft wires is arranged irregularly, whilst in hard wires it is arranged regularly. In the case of the metals with a space centred cubic lattice (tungsten, iron, molybdenum), it is found that only a single lattice plane lies in the section of the wire, namely, the plane (110), whilst in the case of the metals with a face centred cubic lattice (copper, palladium, and aluminium) two lattice planes lie in the section of the wire, namely, the planes (111) and (100). The crystallite arrangement in hard wires is termed a fibrous structure because this arrangement was first observed in natural fibres such as ramie and silk.

J. F. S.

Molecular Structure of Amorphous Solids. C. V. RAMAN (*Nature*, 1922, 109, 138—139).—In view of the fact that glass exhibits a very strong scattering of light, its magnitude being of the order that might be expected if the arrangement of the molecules

was irregular, it is assumed that such is the case. If the arrangement of the molecules approximated to the crystalline state, the scattering of light would be merely that due to the thermal movements of the molecules, and would be much smaller. Careful microscopic examination failed to disclose any inclusions, to which the effect has previously been ascribed (Strutt, *Proc. Roy. Soc.*, 1919, [A], 95, 476).
A. A. E.

Action of Penetrating Radium Radiation on Inorganic and Bio-colloids. III. A. FERNAU and WO. PAULI (*Kolloid Z.*, 1922, 30, 6—13; cf. A., 1915, ii, 722; 1917, ii, 189).—A continuation of previously published work (*loc. cit.*). In the present paper the influence of the radiation from a radium preparation equivalent to 78.6 mg. of radium on gelatin, albumin, casein, acid albumin, and lecithin has been investigated. Of the radiation emitted by the radium preparation, only 25% of the hardest β -radiation and 99% of the γ -radiation came in contact with the colloids. It is shown that the viscosity of a 0.99% solution of glutin from bone glue at 35° falls rapidly at first under the influence of the radiation and then successively less rapidly, but in the presence of 0.1*N*-sodium chloride the fall in viscosity is much slower. The addition of 0.01*N*-hydrochloric acid to 0.5% glutin solution causes at first a rapid fall in the viscosity, which becomes less rapid as the time increases. Under the action of the radiation, the rate of fall of the viscosity is still more rapid. The action of the radiation is analogous to the action of heat on glutin. Glutin jellies when exposed to the radiation liquefy, but the portion nearest the tube containing the radium sets to a solid, opaque mass which sticks to the tube and is insoluble in water, although soluble in alkali hydroxide. The coagulation of serum-albumin, which had been freed from globulin, by the radiation is found less rapid in the presence of sodium chloride, the retardation being greater the greater the concentration of the salt. The time required for coagulation of albumin depends on the concentration of the albumin; it decreases at first with increasing concentration, passes through a minimum at 0.5—1.0% of albumin, and then increases. The action of the radiation on solutions of the sodium salts of acid albumin and casein is to displace the equilibrium toward the acid side, that is, to decrease the concentration of the hydroxyl ion. Thus an acid albuminate, prepared from serum-albumin, had a hydroxyl-ion concentration $1.66 \times 10^{-5}N$; after subjection to the radiation for a week the value was $1.57 \times 10^{-6}N$, and after three weeks $3.62 \times 10^{-8}N$, whilst sodium caseinate solution had the value 8.64×10^{-5} , which became 1.19×10^{-6} after treatment with the radiation for a week. A solution of 3 grams of lecithin in 200 grams of water exhibited at 25°, under the action of the radiation, a gradual fall in viscosity, but scarcely any change in the hydrogen-ion concentration or in the specific conductivity. Similar results are also found with a 0.2% emulsion of cholesterol. The results show that whilst albumins undergo considerable change under the action of radium radiation, this is not the case with lipid emulsions.
J. F. S.

Protective Colloids. XI. Carrageen as a Protective Colloid. I. General Colloid-chemical Investigation of the Extract of Irish Moss. A. GUTBIER and J. HUBER (*Kolloid Z.*, 1922, 30, 20—31; cf. A., 1921, ii, 537).—The method of preparation and properties of colloidal solutions prepared from Irish moss are described. Purified carrageen (30 grams) is placed in a litre of water at the ordinary temperature and kept until the swelling process is complete, then it is vigorously shaken for five hours. The mucilage is separated by pressing through linen. Several other preparations are described; the amount of dry colloid varies between 0.4400 gram and 0.2100 gram per 100 c.c. of mucilage, and the ash between 0.0246 and 0.0600. Dialysis reduces the amount of ash and removes the turbidity and the foul odour and raises the degree of dispersion. Carrageen solutions do not produce fungoid growths when kept, and it is therefore unnecessary to add stabilising agents. On keeping, the viscosity of a 0.55% mucilage fell from 15.29 to an approximately constant value of 7.19 in thirty-three days. The rate of decrease in viscosity is rapid at first and subsequently much slower. The change in viscosity with ageing indicates the change of the mucilage to a more stable system. Boiling the mucilage reduces the viscosity by an amount which increases with the length of time of boiling. This indicates a change in the chemical constitution of the mucilage, but after filtration no further change occurs on prolonged keeping. Heating therefore may be regarded as a hastened ageing and gives rise to stable colloidal solutions. The viscosity increases with increasing concentration, but the amount of increase is greater the higher the temperature employed in the preparation. Electrolytes have no unfavourable action on the mucilage, no coagulation or visible change being brought about except in the case of sodium hydroxide, where it becomes light brown in colour. Increasing concentration of hydrochloric acid, sodium hydroxide, and sodium chloride decreases the viscosity to values which approach a constant value as the concentration of the electrolyte increases.

J. F. S.

Protective Colloids. XI. Carrageen as Protective Colloid. II. Colloidal Silver. A. GUTBIER, A. WOLF, and A. KRIESS (*Kolloid Z.*, 1922, 30, 31—35; cf. preceding abstract).—Extract of Irish moss prepared as indicated previously (*loc. cit.*) has been investigated as to its efficacy as a protecting colloid in the case of colloidal silver. Before use, the mucilage is aged for a considerable time and thereby stabilised. Colloidal silver prepared in the presence of carrageen by hydrazine hydrate has colours varying from dark reddish-brown to brownish-yellow in transmitted light, depending on the concentration of the silver nitrate used. It is exceedingly stable and remains unchanged for two months, and may be evaporated to dryness on a water-bath to an almost black residue which is completely reversible. On the other hand, evaporation in a vacuum over sulphuric acid yields a viscous paste which is not completely reversible, but yields a secondary system with water which is only stable in dilute solutions. Colloidal

solutions of silver have been prepared in the presence of carragheen with a concentration of 17% of silver, but they were not very stable. Carragheen itself has a reducing action on silver nitrate and produces colloidal silver. This action is not complete, but the percentage of reduced silver increases with the dilution of the silver nitrate; thus *N*/10-silver nitrate gives 8.98% of reduced silver, whilst *N*/640-silver nitrate gives 58.54% of reduced silver, the amount of carragheen being the same in all cases. J. F. S.

[Ultra-violet Spectroscopical Studies on the Antagonistic Action of Salts in Organic Colloidal Solution. TETSUTARO TADOKORO (*J. Coll. Agric. Hokkaido Imp. Univ.*, 1921, **10**, 37—49; cf. A., 1918, ii, 432).—The effect of different salts on the degree of dispersion of organic colloidal solutions (egg-albumin, taka-diastrase, blood-serum, and lettuce juice) was studied by observing the changes induced in their absorption spectra in the ultra-violet. Certain pairs of salts were shown to have antagonistic effects on dispersion. Ultramicroscopic observations confirmed the conclusions drawn from the spectroscopical observations. G. W. R.

The Importance of Imbibition for some Biochemical Problems. M. RICHTER-QUITTNER (*Biochem. Z.*, 1921, **121**, 273—292).—By means of the Zsigmondy-Haen ultra-filtration apparatus, the author has examined the influence of electrolytes and non-electrolytes on the velocity of ultra-filtration of various dispersoid systems, for example, sera, for substances which inhibit imbibition will accelerate filtration. All potassium salts, heavy metal salts, urea, and sugar inhibit, whilst other kations favour imbibition. For anions the Hofmeister series holds good. H. K.

[Reversal of Phases in Emulsions and Precipitation of Suspensoids by Electrolytes. SHANTI SWARUPA BHATNAGAR (*Trans. Faraday Soc.*, 1921, **16**, Appendix, 27—31).—A study of the reversal of phase in oil-water emulsions in the presence of pure soaps by electrolytes (cf. T., 1920, **117**, 542), and the precipitation of arsenic sulphide sols by lithium chloride (Mukherjee, *J. Amer. Chem. Soc.*, 1915, **37**, 2030). It is shown that the power of ions in reversing the phases in oil-water emulsions in the presence of soap follows the order: $Al^{+++} > Cr^{+++} > Ni^{++} > Pb^{++} > Ba^{++} > Sr^{++} > Ca^{++}$. The amount of electrolyte required varies with different soaps but the order of the different electrolytes remains the same. The greater the dilution, that is, the distance between the oil particles in an emulsion, the larger the amount of a multivalent ion required to bring about reversal of the phases. Some measurements are given of the effect of diluting milk on the rate of coagulation by dilute sulphuric acid. It is shown that the greater the dilution the larger is the concentration of acid required to bring about an immediate coagulation. Soaps are regarded as exerting a protective action on oil-water emulsions, and it is shown that the protective action follows the order: potassium stearate > sodium stearate > sodium and potassium oleate > potassium oleate > sodium oleate.

J. F. S.

Peptisation and Precipitation. N. G. CHATTERJI and N. R. DHAR (*Trans. Faraday Soc.*, 1921, 16, Appendix, 122—127).—Silver chloride, silver chromate, calcium silicate, cupric fluoride, and barium sulphate when precipitated in the presence of glycerol or concentrated sucrose solutions are not peptised. The hydroxides of iron, nickel, cobalt, thorium, and mercury may be peptised by adding an alkaline hydroxide to the solutions of salts of these metals in the presence of glycerol or concentrated sucrose solution. In the case of gold or uranium salts, ammonium hydroxide should be used instead of sodium hydroxide. In these cases, peptisation can only be brought about if the alkali is added to the mixture of the salt and glycerol, but precipitation takes place if the solution of the salt is added to the mixture of glycerol and alkali. These results are supported by conductivity measurements. The major portion of the hydroxides of aluminium, chromium, lead, mercury, and copper, when dissolved in sodium or potassium hydroxide, exists as a colloid, whilst zinc hydroxide mainly forms a zinc oxide, as shown by conductivity experiments. Similarly, the hydroxides of aluminium, iron, and chromium form colloidal solutions in acetic acid, whilst zinc hydroxide forms zinc acetate. The transformation of the blue hydroxide of copper to the black variety and that of the blue hydroxide of cobalt to the pink form are changes of the same type, and are almost equally effected by different catalysts. The hydroxides of aluminium, chromium, and zinc respectively are adsorbed by ferric hydroxide when the latter is precipitated in the presence of solutions of the hydroxides of these metals in potassium hydroxide.

J. F. S.

A Kinetic Precipitation Measurer. WO. OSTWALD and F.-V. VON HAHN (*Kolloid Z.*, 1922, 30, 62—70).—Two new precipitation measures are described, which permit of the precipitation process and the peptisation of colloidal solutions being studied kinetically. The one instrument, the two-limbed measurer, is based on the principle of level difference in communicating tubes, and the change of this difference due to decrease of density of one liquid by the sedimentation of the disperse phase contained in it. The second apparatus, the single-limbed measurer, depends on the rise of the seal liquid column of an enclosed gaseous volume, through change in the density of the seal liquid in consequence of the precipitation of the disperse phase which it contains. A number of examples of the use of these instruments in the kinetic measurement of the precipitation processes of typical sols are given.

J. F. S.

Distribution of certain Drugs between Immiscible Solvents. W. O. EMERY and C. D. WRIGHT (*J. Amer. Chem. Soc.*, 1921, 43, 2323—2335).—Distribution experiments between water and chloroform have been carried out with caffeine, antipyrine and *p*-acetoxyacetanilide. It is shown that the most suitable temperature for the extraction of caffeine from water by chloroform is the ordinary temperature, although a change of temperature from 12° to 40° only reduces the amount recovered from water by a single extraction from 96.3% to 93.2%. The distribution is the more favourable to

the chloroform the more dilute the aqueous solution. The effect of small concentrations of sodium hydroxide, sodium acetate, sulphuric acid, sodium salicylate, sucrose, alcohol, and citric acid on the distribution of caffeine between water and chloroform has been examined at 25°. The presence of sulphuric acid leads to a less efficient extraction of the caffeine, whilst sodium salicylate reduces the amount extracted from 96% to 20.7%, due in all probability to the formation of a molecular compound between caffeine and sodium salicylate. The solubility of caffeine in water, and aqueous solutions of sulphuric acid (*N*), citric acid (*N*), potassium bromide (*N*) and (2.5*N*), sodium salicylate (*N*) and (0.1*N*), sodium benzoate (*N*) and (0.1*N*) has been determined at 25°. The solubility in grams per 100 c.c. of solvent is 2.071 in water, 3.355 in sulphuric acid, 6.411 in citric acid, 2.136 and 2.035 respectively in potassium bromide, 22.22 and 4.921 respectively in sodium salicylate, and 15.27 and 3.422 respectively in sodium benzoate. The solubility determinations and cryoscopic measurements of mixtures of caffeine with sodium salicylate and sodium benzoate respectively confirm the complex salt formation with these salts. Distribution experiments of caffeine between absolute chloroform and water, *N*-sulphuric acid, *N*-potassium bromide, 0.1*N*-sodium salicylate, and 0.1*N*-sodium benzoate have been made at 25°. It is shown that potassium bromide has but little effect on the partition, whilst sodium salicylate reduces the amount of caffeine recovered from the water solutions by the greatest amount. The distribution of antipyrine shows that there is no tendency to the formation of double salts with sodium salicylate, but, on the other hand, there is a far greater depression in the percentage of antipyrine extracted from *N*-sulphuric acid, when compared with that from pure water solution, due presumably to difference in basicity. In the case of *p*-acetoxyacetanilide, somewhat lower distribution ratios are obtained than for either caffeine or antipyrine. The solubility of *p*-acetoxyacetanilide in water at 25° is found to be 0.239 gram per 100 c.c. of water, and in chloroform at the same temperature 3.250 grams in 100 c.c. of chloroform.

J. F. S.

Theory of the Liesegang Rings. MARTIN H. FISCHER and GEORG D. McLAUGHLIN (*Kolloid Z.*, 1922, 30, 13—16).—It is shown that Liesegang rings are only observed when a liquid phase is present; they may be formed when two dissolved substances, which react with one another, are brought together in the presence of a third insoluble substance; the formation of the rings occurs on the side of the solution of lower concentration, the rings tend to increase in thickness from the commencement of their formation and during the growth tend to pass from the originally liquid and amorphous condition to the solid and crystalline form. All substances which give satisfactory Liesegang rings are impermeable toward molecularly dissolved substances, and consequently it would seem unlikely that after the formation of the first ring a second and third ring could be formed. The view is put forward that true semi-permeable media are solvated and when these become less solvated

holes appear through which the dissolved substance can diffuse and so go to form a second ring.

J. F. S.

Thermodynamical Theory of Explosions. I and II. SIR J. B. HENDERSON and H. R. HASSÉ (*Proc. Roy. Soc.*, 1922, [A], 100, 461—482).—A theoretical paper in which calculations are made of the maximum temperature and pressure of an explosion in a closed vessel, also the calculation of the curve of reversible adiabatic expansion of the products, and incidentally the ideal indicator diagram. The calculations are applied to a gun, in which the variation with time of the volume of the chamber, of the composition of the products, and the temperature and pressure of explosion are worked out and the value of the maximum pressure determined. The maximum temperature obtained with Mark I cordite is 3210° , and with MD cordite 2870° ; whilst the maximum pressure of Mark I cordite is 8370 atmospheres or 55.1 tons/sq. in.

J. F. S.

Equation for the Velocity of Reaction of Hydrogen Peroxide and Potassium Permanganate. JAN ZAWIDZKI (*Roczniki Chemji*, 1921, 1, 135—139).—From the measurements of Baeyer and Villiger (*A.*, 1900, ii, 719) on the course of the reaction between hydrogen peroxide and potassium permanganate, the author calculates that the time law of this reaction is expressed by the differential equation $dx'/dt = k_2(1-x')/\sqrt{a}\sqrt{x'}$. The velocity coefficient $k' = k_2/\sqrt{a}$ is calculated for -16° and $+15^{\circ}$ and the values 0.119 and 0.379 respectively are obtained. From these, the temperature coefficient of the velocity constant is found to be 1.452.

J. F. S.

The Velocity of Reaction in Mixed Solvents. II. The Velocity of Saponification of the Ethyl Esters of some Mono-substituted Benzoic Acids. ALBERT ERIC CASHMORE, HAMILTON MCCOMBIE, and HAROLD ARCHIBALD SCARBOROUGH (*T.*, 1922, 121, 243—253).

Influence of Temperature on the Velocity of Interpenetration of Solids. H. WEISS and P. HENRY (*Compt. rend.*, 1922, 174, 292—294; cf. *A.*, 1920, ii, 545; 1921, ii, 551).—From a study of the velocity of solution of silver antimonide, Ag_3Sb , in solid silver at different temperatures in the case of a silver-antimony alloy containing 14% of antimony, it is shown that the velocity is proportional to the temperature, being expressed by the equation $v = Ka^T$, where T is the absolute temperature and, in this case, $K = 4.17 \times 10^{-12}$, and $a = 1.0324$.

W. G.

The Mechanism of Catalytic Hydrogenation. A. SKITA (*Ber.*, 1922, 55, [B], 139—143).—In a recent communication, Willstätter and Waldschmidt-Leitz (*A.*, 1921, ii, 185) have pointed out the necessity of priming the platinum catalyst with oxygen during the course of hydrogenations and have advanced the hypothesis that a platinum peroxide or oxide is intermediately formed. This suggestion is quite consonant with the author's repeated observation that the activity of the catalyst produced *in situ* is

superior to that of the pre-formed agent, since the experimental conditions do not guarantee the complete absence of oxygen. Comparative experiments with a platinum catalyst produced in situ do not show any difference in the rate of hydrogenation of pulegone or *as-p*-xylidine when every trace of oxygen is excluded and when special precautions to this end are not observed; the formation of a peroxide as catalyst cannot therefore be assumed in these cases. Further, if the platinum catalyst is in reality a peroxide, its oxygen must liberate iodine from potassium iodide, and hydrogenation must be impossible in the presence of the salt. It is found, however, that phenol is reduced smoothly to *cyclohexanol* at 40° in the presence of potassium iodide. On the other hand, the addition of potassium iodide completely inhibits the reduction of phenol or *as-p*-xylenol at the atmospheric temperature, whereas in its absence they are transformed readily into *cyclohexanol* and 1:4-dimethylcyclohexan-2-ol, b. p. 179°, respectively. Reduction occurs quantitatively but slowly if the mixtures are heated at 50°. At the higher temperature, it appears, therefore, impossible that hydrogenation should depend on the formation of a platinum peroxide. It has not yet been elucidated whether the failure of the action at the atmospheric temperature is due to the inactivation of a platinum peroxide or to poisoning of the catalyst.

H. W.

The Mechanism of Catalytic Action in the Hydrolysis of Fats. E. BRINER and (MLLE) A. TRAMPLER (*Helv. Chim. Acta*, 1922, 5, 18—20).—The product obtained by the sulphonation of phenylacetic acid is inferior to hydrochloric or sulphuric acid as a catalyst in the hydrolysis of ethyl acetate in homogeneous solution. It thus appears that the presence of a common group in catalyst and substrate does not confer any supplementary chemical activity on the former. In Twitchell's and similar catalysts, the activity of the sulphonic group is diminished by the presence of the aliphatic radicle, but this effect is more than compensated by the greater mutual solubility caused by the two similar groups.

H. W.

Conclusions from the Spherical Form of the Simplest Atom. L. ZEHNDER (*Physikal. Z.*, 1922, 23, 53—55).—A theoretical paper of a highly speculative character, in which, on the basis of a spherical primordial atom, an atomic ether, the atoms of which have a mass of about one-millionth that of hydrogen, elasticity, and gravitational forces, the author draws a number of conclusions respecting the atomic weight of helium, the structure of the carbon atom, and the subdivision of organic compounds into three classes, aliphatic, aromatic, and carbohydrates.

J. F. S.

A New Model of the Nucleus. MAXIMILIAN CAMILLO NEUBURGER (*Z. physikal. Chem.*, 1921, 99, 454—473).—Previous speculations (cf. A., 1921, ii, 479) are extended by the consideration of isohydrogen particles, of mass 1. The disintegration series of thorium, uranium, and actinium are considered on the basis of this hypothesis.

J. R. P.

Structure of the Atomic Nucleus. O. CHWOLSON (*Z. Physik*, 1921, 7, 268—284).—A theoretical paper in which an attempt is made to deduce the structure of the atomic nucleus. It is maintained that any hypothesis of the structure of the atomic nucleus must conform to the following: (1) the mass of the atomic nucleus is practically the same as that of the atom, (2) the positive charge, E , of the nucleus is eN , where $-e$ is the charge of the electron, and N the atomic number of the corresponding element, (3) the atomic weight is roughly $M=2N$, (4) in the case of hydrogen $M=2N$ is not true, and (5) the hypothesis must not be in conflict with Bohr's hypothesis. As the basis of the present hypothesis, the author assumes that the volume density, δ , of electricity of both signs is a constant and that a given charge, E , irrespective of its sign, has the same volume, $v=\delta/E$. From this it is shown that the atomic nucleus cannot easily be represented as a sphere built up of spherical particles, and the assumption is made that it is cylindrical and is built up of disks. This conception is strictly in keeping with the relatively large volume of the nucleus in comparison with that of the electron. It is further assumed that the positive elementary disk is identical with the helium nucleus and the α -particle. The elementary positive disk is termed the "pentalon." Negative electricity consists of spherical electrons and positive electricity of disk-shaped pentalons. If $-e$ is the charge of an electron, e' , the charge of the pentalon is $e'=2e$; the mass of the pentalon is $m'=4$. The order number of the pentalon N' is $N'=N+1$, where N is the atomic number of the element. The atomic nucleus is a disk or cylinder made up of pentalons, between each pair of which is a disk of negative electricity, of charge $-e$, equal to that of an electron. The disk of negative electricity is termed a "piezo-electron" and its order number is $N''=N-2$. The atomic nucleus of an element of atomic number N consists of $N-1$ pentalons and $N-2$ piezo-electrons; these $2N-3$ disks form a cylinder which has no interstices. The charge of the nucleus is $E=(N-1)2e-(N-2)e=Ne$. Calculating from the above, it is shown that the radius of the pentalon is $69.9s$, where s is the radius of the electron, and the thickness, D , is $0.0005450s$; the radius of the piezo-electron is also $69.9s$, and the thickness is half that of the pentalon. The pentalon has a mass which is 7340 times that of the electron. The thickness $D(N)$ of the nucleus of an element of atomic number N is $D(N)=(N-1)D+(N-2)D/2=(3N-4)D/2$ and the mass, $M_1=4(N-1)+2(N-2)=2N$. The hypothesis is compared with the Bohr hypothesis, and found not to conflict with it. The hydrogen nucleus on the basis of the hypothesis is shown to be built up of one pentalon and one piezo-electron, both of double thickness and of reduced radii in the ratio $1:\sqrt{2}$. That is, the hydrogen nucleus is three times as thick as the helium nucleus, and has a radius $49.5s$ and a thickness $0.001635s$.

J. F. S.

Attempt [to Find] a Physical Basis for Atomic Radiation. E. GEHROCKE (*Z. Physik*, 1921, 6, 400—402).—The author imagines

the electrons surrounding an atomic nucleus as rings round the nucleus and not as spheres rotating in an orbit round the nucleus. The ring form of the electron has the same mass, volume, and surface as the spherical electron, and is to be regarded as a spacial isotope of the spherical electron. If for any reason the ring electron changes its diameter, a change in the ether must also occur, and this change in the ether gives rise to emission or adsorption; emission when the ring reduces its diameter and absorption when the diameter increases. The change in the diameter of the ring consists in the displacement of the ether over the space between the initial and final positions of the ring. The mechanism of this process is visualised as follows. An atomic ether, E , the atoms of which are arranged in a space lattice, is assumed. The space lattice has spherical symmetry in the neighbourhood of a positive atomic nucleus. Between the atoms of E , a magnetic ether, M , is assumed, which in comparison with E can be regarded as a continuum. The atomic ether, E , is the carrier of electric forces and is named the electric ether. The magnetic ether can spread out in stream lines (magnetic field lines), whilst the electric ether cannot spread out in the same way, but may change the shape of its space lattice. The sudden change of the space lattice, due to a change in diameter of the ring electron, occasions vibrations in the ether atoms which are transmitted to neighbouring ether atoms, and when the transmission of the vibration extends outside the ring emission of radiation results. Absorption results when incoming vibrations increase the kinetic energy of the ring electron, that is, increase its velocity and so increase its diameter. It is therefore not the ring electron which makes the spring at each energy step, but the E ether which is displaced by the thickness of a ring electron. The generator of light and Röntgen radiation in the material atom is, therefore, not a vibrating electron, but the whole system of ether atoms between two radii, R and r , swinging in a pendulum-like manner. The frequency, ν , of the vibrations of the ether atoms is given by the ether between the radii R and r , and is proportional to the thickness and inversely proportional to the square of the length, that is, $\nu \sim R-r/Rr=1/r-1/R$.

J. F. S.

Condition of Motion of Molecules in Space. J. BÖESEKEN, CHR. VAN LOON, DERX, and P. HERMANS (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 24, 198—204).—A theoretical paper in which it is shown that atoms which are not directly bound to one another exercise a mutual action on other atoms in the same molecule. Saturated non-cyclic molecules execute, among others, movements in which the parts of the molecule revolve in opposite directions, or with different velocities round the single bonds as axes. In the case of non-uniformly loaded molecules, that is, practically all molecules except those of hydrogen, nitrogen, oxygen, ethane, hexachloroethane, and similar molecules, the movements are irregular, because the most stable position of the atoms will be passed most frequently. In the saturated cyclic molecules with six or seven atoms, the ring-forming atoms are not fixed in one

plane, but lie in a curved surface which travels through space in undulatory movements; the atoms in this surface are under no tension.

J. F. S.

Transition Elements and the Octet Theory: A New Arrangement of the Rare Earth Elements in the Periodic Classification. R. G. W. NORRISH (*Chem. News*, 1922, 124, 16—22).—A theoretical paper in which the author considers the fourth postulate of Langmuir's octet hypothesis from the point of view of valency and colour of the ions produced in solutions of salts of the metals. A new arrangement of the periodic system is drawn up, which has as its basis that the transition elements and the rare earth metals have a variable electronic structure as far as the outermost shell of electrons is concerned. It is shown that the elements which yield coloured kations can all be grouped together in a rectangle which extends from groups 4A to 1B and from series IV to VI. These elements are assumed to possess the variable electronic structure. The rare earth metals which give coloured kations are sharply cut off from those which give colourless kations by this rectangle. The existing data concerning the characteristic infra-red frequency, the latent heat of vaporisation, the compressibility, and the magnetic susceptibility are tabulated with the periodic arrangement, and it is shown that these properties maintain a more or less high value inside the rectangle but fall off sharply on either side of it. The elements inside the rectangle (excluding copper, silver, gold, and zirconium) are paramagnetic, whilst to the right of the rectangle all the elements except tin, which is feebly paramagnetic, are diamagnetic and those to the left of the rectangle are either feebly paramagnetic or diamagnetic. It appears that the magnetic susceptibility and colour are dependent on the presence of an incomplete subsidiary ring in the atom.

J. F. S.

Radii of the Alkali Metals and the Alkali Metal Ions. RICHARD LORENZ (*Z. Physik*, 1921, 6, 271—275).—A theoretical paper in which the author has discussed the values previously obtained for the atomic radii (A., 1921, ii, 191) and the space filling of the atoms concerned. A new table of the atomic radii, ionic radii, and the space filling of the alkali metals and alkali metal ions has been drawn up which is based on the present discussion and recent work of other investigators.

J. F. S.

A Lecture Experiment for Demonstrating the Dependence of the Antiseptic Action of Mercury Compounds on the Degree of Ionisation. G. JOACHIMOGLU (*Biochem. Z.*, 1921, 121, 259—261).—The fermentation of dextrose by yeast with the visible production of carbon dioxide in fermentation tubes affords a means of demonstrating the antiseptic influence of mercuric ions. Addition of mercuric chloride inhibits fermentation completely, but repression of the ionisation by addition of sodium chloride gives a moderate evolution of carbon dioxide. Mercuric cyanide has practically no inhibiting influence.

H. K.

Inorganic Chemistry.

The Steric Formula of the Molecule of Water. JEAN PICCARD (*Helv. Chim. Acta*, 1922, 5, 72—74).—It is only possible to explain certain physical properties of water if it is assumed that the centre of gravity of the positive charges of the molecule does not coincide with that of the negative charges. This asymmetry of the charges might result from the inequality in the distances of the two hydrogen atoms from the oxygen atom or by such a disposition of the hydrogen atoms that the valencies uniting them to the oxygen atom are disposed at an angle other than 180° . There is, however, no evidence in favour of the first supposition. A consideration of oxonium salts, on the other hand, shows that in them the co-ordinative valency of oxygen is three, and that the oxygen atom may be regarded as placed in the centre of an equilateral triangle with its valencies directed towards the apexes and therefore inclined to one another at an angle of 120° . An explanation is thus given for the observation that only those carbon-oxygen heterocyclic rings are stable in which the total number of atoms is five or six. Thus, for example, if the valencies of carbon form an angle of $109^\circ 28'$ with one another and those of oxygen an angle of 120° , the mean deviation in a cyclic system composed of three carbon and three oxygen atoms is only $2^\circ 38'$ whereas in cyclohexane the mean deviation is $5^\circ 16'$. H. W.

Electrolytic Concentration of Aqueous Solutions of Nitric Acid. I. HENRY JERMAIN MAUDE CREIGHTON (*J. Franklin Inst.*, 1922, 193, 89—95).—When a solution of nitric acid is electrolysed in a cell in which the anode and cathode are separated by a porous diaphragm, concentration of the acid occurs through electrolytic decomposition of water and at the same time the concentration of the acid in the anolyte is increased at the expense of that in the catholyte on account of the different migration velocities of the hydrogen and nitrate ions. The acid in the catholyte is further reduced by reduction to nitrogen oxides, hydroxylamine, or ammonia according to the nature of the cathode metal. Experiments were made in which 70—71% nitric acid was electrolysed in a diaphragm cell using platinum electrodes, and provision was made for returning the gaseous nitrogen oxides formed in the anolyte chamber back to the catholyte chamber. The current used was 6—8 amperes and the *E.M.F.* 3.0—6.5 volts. It was found possible thus to increase the concentration of acid in the anolyte chamber to 99.65% HNO_3 . The porous cylinder used underwent considerable disintegration in the process. [See further *J. Soc. Chem. Ind.*, 1922, 172A.]

E. H. R.

Preparation of Silicic Acid and Tungsten Hydroxide Sols by means of Hildebrand Cells. M. KRÖGER (*Kolloid Z.*, 1922, 30, 16—18).—By the electrolysis of a 1.5% solution of sodium silicate

between a mercury cathode and a platinum anode in a Hildebrand cell sols of silicic acid are obtained which do not gelatinise until they have been kept for four weeks. A 6% solution gelatinises as soon as the solution becomes neutral. The electrolysis is started with a current of 0.55 ampere, but this steadily falls as the process proceeds, and in ninety minutes has reached the value 0.12 ampere. After one hundred and ten minutes' electrolysis, the solution has a neutral reaction toward litmus. In the case of the 6% solution, the gelatinisation proceeds so rapidly that the waves occasioned by stirring are often reproduced in the gel, which is usually as clear as glass. Electrolysis of a 30% solution of water glass causes silica to separate on the anode. Electrolysis of a 2% solution of sodium tungstate, using the apparatus named above with a silver anode, rapidly produces the hydrosol of tungsten hydroxide. The removal of alkali may be hastened by the cautious addition from time to time of a little hydrochloric acid, but in no circumstances may the neutral point be passed. Should the solution become acid, blue tungsten compounds are produced. The tungsten hydroxide hydrosols are clear and transparent, but of a deep brown colour, which in dilute solutions is yellowish-brown. They are coagulated by potassium chloride to form a black powder which resembles the lower oxides of tungsten.

J. F. S.

Influence of Tungstic Acid on the Gelatinisation of Silicic Acid in Concentrated Hydrochloric Acid Solutions. M. KRÖGER (*Kolloid Z.*, 1922, 30, 18—19).—The time required for the gelatinisation of silicic acid in the presence of tungstic acid by hydrochloric acid has been investigated. The solution consisted in each case of 7 c.c. of a solution of sodium silicate containing 33.7% of silica, to which had been added volumes of a 10% solution of sodium tungstate varying from 0.5 c.c. to 15 c.c., the total volume in each case being made up to 22 c.c. Nine c.c. of 9.77*N*-hydrochloric acid were added in each case and the time required for complete gelatinisation was noted. The time-tungstic acid concentration curve passes through a minimum at 1 c.c. of tungstic acid and a maximum at 2.5 c.c. of tungstic acid and then falls continuously.

J. F. S.

Frequency of the Electrons in the Neon Atom. LAURENCE ST. C. BROUGHALL (*Phil. Mag.*, 1922, [vi], 43, 339—344).—A mathematical paper in which the frequency, angular velocity, and linear velocity of the electrons in the neon atom have been calculated on the assumption that the electrons do not radiate energy under normal conditions. The following numerical results are recorded: Frequency of the electrons about the axis XX' , $n_3 = 0.73 \times 10^{16}$; frequency about YY' and ZZ' , $n' = 1.00 \times 10^{16}$; angular velocity about XX' , $\omega_3 = 4.58 \times 10^{16}$ rad./sec.; angular velocity about YY' and ZZ' , $\omega' = 6.28 \times 10^{16}$ rad./sec.; instantaneous linear velocity of the outer electrons about XX' , $v_3 = 2.98 \times 10^8$ cm./sec.; instantaneous linear velocity of the outer electrons about YY' and ZZ' , $v' = 4.08 \times 10^8$ cm./sec.; instantaneous linear velocity of the inner electrons about YY' and ZZ' , $v = 3.83 \times 10^8$ cm./sec.,

The value of v is small when compared with the velocity of light, in consequence of which it follows that no appreciable error is committed in not correcting for the variation of mass with velocity according to the equation $m_v = m_0(1 - v^2/c^2)^{1/2}$, where c is the velocity of light. J. F. S.

Crystal Structures of the Alkali Haloids. I. RALPH W. G. WYCKOFF (*J. Washington Acad. Sci.*, 1921, 11, 429—434).—From data as to the crystalline structure of the alkali haloids obtained from powder photographs and on the assumption that the unit cell consists of four molecules, it is held by the author that there are only two possible arrangements of the molecules within the unit cell, namely, the sodium chloride arrangement and the zinc sulphide arrangement. The geometrical considerations involved and the method of calculating the nature of the diffraction effects to be expected are given elsewhere (Wyckoff and Posnjak, following abstract). A closer accord of the normal decline of intensity of reflection with the spacing of the reflecting planes as observed in experiments on sodium chloride and similar crystals is obtained by assuming the intensity to be proportional to the 2.35 power of the spacing instead of the simple square. Experimental determinations of the intensity for a few of the principal lines of the spectrum were compared with the intensities calculated with alternative assumption of the two groupings above mentioned. The results agreed with the assumption of the sodium chloride grouping in the case of sodium bromide, sodium iodide, potassium bromide, potassium iodide, and rubidium chloride. Cæsium bromide and cæsium iodide have a similar structure to cæsium chloride, which has been shown to be body-centred (Davey and Wick, *Physical Rev.*, 1921, 17, 403). G. W. R.

Crystal Structure of Ammonium Platinichloride. RALPH W. G. WYCKOFF and EUGEN POSNJAK (*J. Amer. Chem. Soc.*, 1921, 30, 2292—2309).—The crystal structure of ammonium platinichloride has been determined by a general method which is based on the theory of space groups. It is shown that crystals of ammonium platinichloride have a structure which is analogous to that commonly assumed for fluorspar, in which the PtCl_6 groups occupy the positions of the calcium in fluorspar or crystals and the NH_4 groups the positions of the fluorine. The unit crystal cell has a side 9.843×10^{-8} cm. The only assumption made which is not required in the ordinary determination of the wave-length of X-rays from a reflection spectrum is that the four hydrogen atoms of the ammonium radicle are exactly alike, with this exception, that in attempting to place the chlorine atoms with accuracy, it was assumed that atoms scatter X-rays in an amount which is roughly proportional to their atomic numbers and that in a lattice arrangement of atoms the intensities of reflection follow quantitatively the order of $1/(h^2 + k^2 + l^2)$. J. F. S.

Revision of the Atomic Weight of Glucinum. Analysis of Glucinum Chloride. O. HÖNIGSCHMID and L. BIRCKENBACH (*Ber.*, 1922, 55, [B], 4—12).—The ratios $\text{BeCl}_2 : 2\text{Ag}$ and

$\text{BeCl}_2 : 2\text{AgCl}$ have been determined in the same manner as used previously in the determination of the atomic weight of bismuth (Hönigschmid and Birckenbach, A., 1921, ii, 646). As mean result of all analyses, the value $G1=9.018$ is adopted, this figure being about 1% lower than that assigned by the International Commission.

Technical glucinum carbonate is converted into the basic acetate and purified from iron compounds by repeated crystallisation from glacial acetic acid. The purified acetate is sublimed and converted into the nitrate. The solution of the latter in water is treated with an excess of ammonium carbonate and filtered from any undissolved aluminium compounds; the glucinum carbonate is subsequently precipitated by boiling the filtrate, and is finally converted into the oxide by calcination in a platinum dish in an electrically heated furnace. The pure oxide is transformed into the chloride by ignition with carbon in a current of chlorine. The apparatus used is identical with that described previously (*loc. cit.*). Great caution is needed in the quantitative decomposition of glucinum chloride by water.

Glucinum chloride has d_4^{25} 1.8995.

H. W.

Inorganic Luminescence Phenomena. IV. Preparation of Pure Magnesium Sulphide and its Phosphorescence.
II. Phosphorescent Magnesium Sulphides. ERICH TIEDE and FRIEDRICH RICHTER (*Ber.*, 1922, 55, [B], 69—74).—The specimens of magnesium sulphide described previously (A., 1916, ii, 619) were not sufficiently pure to allow definite conclusions to be drawn with respect to the capacity of the substance to phosphoresce. Pure magnesium sulphide has now been prepared by the ignition of magnesium oxide or, preferably, of anhydrous magnesium sulphate in a current of nitrogen laden with carbon disulphide vapour. The compound is not phosphorescent, but becomes so by suitable additions of manganese, bismuth, or antimony, the optimal amount of metal for 1 gram of sulphide being 0.001—0.002 gram of manganese as sulphate or chloride, 0.0024 gram of bismuth as the basic nitrate, and 0.0013 gram of antimony as potassium antimonyl tartrate. The main band of emission of phosphorescent light in the case of magnesium sulphide containing manganese lies in the red, approximately between 615 and 765 $\mu\mu$ with a maximum at 720 $\mu\mu$. The intensity at the atmospheric temperature for metal content from 0.00023—0.004 gram is almost constant and not markedly dependent on wide variations of temperature and duration of ignition. The duration of the phosphorescence is small. Magnesium sulphide containing bismuth exhibits an intensely blue phosphorescence which is excited by daylight or arc or mercury-vapour light. The band lies between 430 and 550 $\mu\mu$ with a maximum at 465 $\mu\mu$. Specimens of magnesium sulphide containing antimony have a delicate, yellow colour and a persistent, intensely yellowish-green phosphorescence after excitation by daylight or arc or mercury-vapour light, or, particularly, by exposure to cathode rays. The band lies between 570 and 610 $\mu\mu$ with a maximum at 545 $\mu\mu$.

H. W.

g**

Lead in the Uranium Minerals of Madagascar. MUGUET (*Compt. rend.*, 1922, 174, 172—173).—The industrial treatment of several tons of betafite from Madagascar resulted in the isolation of lead to the extent of 0.6% of the mineral treated. The mineral being perfectly crystalline and practically free from impurities, the lead apparently occurs in the same chemical form as the uranium and is a disintegration product of uranium. The radioactivity of this lead has increased regularly for six months. W. G.

Abnormal Crystallisation of Lead Azide by Protective Colloids. A. G. LOWNDES (*Trans. Faraday Soc.*, 1921, 16, Appendix, 128—129).—Lead azide when formed in large crystals is liable to explode and is therefore not entirely suitable for detonators. It is generally held that the explosions are caused by the fracture of large crystals. It is shown in preparing lead azide that if instead of running sodium azide and lead acetate solutions into water the solutions are run into a 0.5% solution of gelatin or dextrin small crystals which are not liable to fracture are produced. The presence of ferric chloride also causes the formation of small crystals, but these are useless for the filling of detonators. A number of photomicrographs of lead azide prepared by the various methods are included in the paper. J. F. S.

Preparation of Catalytic Copper. JEAN PICCARD (*Helv. Chim. Acta*, 1922, 5, 147—148).—The following modified method (cf. Piccard and Larsen, A., 1917, i, 644) yields catalytic copper which is superior in its activity to Kahlbaum's "copper-bronze" or "naturkupfer C."

Granulated zinc (600 grams) is mixed with finely-powdered potassium dichromate (190 grams) or, preferably, with an equal weight of hydrated sodium dichromate, in a three-litre flask and concentrated hydrochloric acid (1300 c.c.) is added within five to ten minutes. A vigorous action ensues which leads to the formation of a clear, blue solution. The latter is filtered through glass wool into a flask filled with carbon dioxide, a current of the gas being passed over the filter during the process. The solution is treated with brisk agitation with copper sulphate (120 grams) dissolved in ice-cold water (700 c.c.). Reduction is instantaneous. The precipitated copper is washed five times by decantation with water, then filtered, and washed successively with ordinary and absolute alcohol and benzene. If the product is to be dried, the final washing should be effected with benzene containing a little vaselin. After desiccation in a vacuum, the copper is relatively stable towards air, but is preferably preserved in evacuated sealed tubes. It may also be stored as a paste beneath alcohol. H. W.

Colloidal Copper Hydroxide. C. PAAL and HERMANN STEYER (*Kolloid Z.*, 1922, 30, 1—5).—The preparation of solid colloidal copper hydroxide is described. To 50 c.c. of 2% sodium prot-albinate solution, which acts as protective colloid, 20 c.c. of *N*-sodium hydroxide and 20 c.c. of 1% copper sulphate solution are added alternately in small quantities at a time. A light blue

turbid sol is produced which is dialysed for four days, treated with 3 drops of *N*-sodium hydroxide, and evaporated to dryness at 60° in a vacuum. A blackish-blue, brittle substance is obtained in the form of lamellæ which dissolve in water to form the original sol. The solid colloid contains 14.02% of copper and 1.34% of sodium. By using sodium lysalbinat as protective colloid and varying the quantities of the other reagents, solid colloids, similar in appearance and properties to the above but varying in composition, may be obtained. The colloid richest in copper contains 35.47% of copper and 5.31% of sodium. If solutions of the sols are heated for some time on a water-bath black lamellæ are deposited which dissolve in water to give a dark brown, turbid hydrosol. This does not change in colour when treated with 4*N*-ammonia solution, even after keeping for four days. The product is regarded as colloidal cupric oxide, and has been obtained containing 28.58% of copper and 4.37% of sodium.

J. F. S.

The Peroxidic Compounds of Copper. (MISS) JOAN ALDRIDGE and MALCOLM PERCIVAL APPLEBEY (T., 1922, **121**, 238—243).

Reduction of Solutions of Ferric Salts with Mercury. LEROY W. MCCAY and WILLIAM T. ANDERSON, jun. (*J. Amer. Chem. Soc.*, 1921, **43**, 2372—2378).—Neutral and acid solutions of ferric chloride are completely and rapidly reduced when shaken with a little mercury. In the case of ferric sulphate, the reduction proceeds to an equilibrium which at 20° lies at about 53% of ferrous iron, but if a little free hydrochloric acid or sodium chloride is added the reduction becomes complete. Under similar conditions, solutions of titanous acid are not reduced. If the mercurous salt is filtered off, the ferrous iron in the solution may be estimated by titration with either potassium permanganate or dichromate. Experiments on the estimation of iron in ferric alum and other ferric compounds show that this constitutes a rapid, accurate, and convenient method. Solutions of potassium ferricyanide, potassium chromate, ammonium molybdate, sodium vanadate, and potassium antimonate when acidified with hydrochloric acid are all reduced when shaken with mercury.

J. F. S.

Strengths of Cobaltamine Bases and Werner's Theory of Bases. ARTHUR B. LAMB and VICTOR YNGVE (*J. Amer. Chem. Soc.*, 1921, **43**, 2352—2366).—The relative strengths of the following cobaltamine bases: hexamminecobaltic dibromide dihydroxide, hexamminecobaltic hydroxide, aquopentamminecobaltic hydroxide, diaquotetramminecobaltic hydroxide, triethylenediaminecobaltic hydroxide, diaquodiethylenediaminecobaltic hydroxide, carbonato-tetramminecobaltic hydroxide, 1 : 2-dinitrotetramminecobaltic hydroxide, 1 : 6-dinitrotetramminecobaltic hydroxide, and dinitroaquotriamminecobaltic hydroxide have been determined by measuring the electrical conductivity over a range of concentrations at 25° and comparing these with the conductivities at zero concentration, as calculated from conductivity measurements of salts derived

from these bases. The determinations show that these bases are very strong, some of them being as highly ionised as potassium hydroxide; the replacement of ammonia by ethylenediamine has no effect on the strength of the base. The replacement of ammonia by water molecules produces a marked and progressive decrease in the strength; the substitution of acid-groups has no marked effect on the strength, the electrostatic effect of a decrease in valency probably counteracting the chemical effect of the acid radicle; the stronger acid radicle produces the weaker base. There is no marked difference in the ionisation of the successive hydroxyl groups in hexamminecobalt hydroxide. The improbability of the distinctive features of Werner's theory of bases is pointed out, and a more probable interpretation suggested. Equations have been derived, giving an important correction of the conductivity of solutions of bases for the conductivity of the carbon dioxide dissolved in the water. These equations have been applied in the present work.

J. F. S.

Sub-salts of Bismuth. HENRY GEORGE DENHAM (*J. Amer. Chem. Soc.*, 1921, **43**, 2367—2371).—Making use of the method and apparatus formerly employed in the preparation of the sub-salts of lead (T., 1917, **111**, 29; 1918, **113**, 249; 1919, **115**, 109), the author has prepared sub-salts of bismuth from bismuth sub-oxide. The products obtained are a sub-oxyiodide, $2\text{BiI}_2 \cdot 3\text{BiO}$, the sub-iodide, BiI_2 , and bismuth dimethyl. The *sub-oxyiodide* is a non-volatile, brick-red substance which is stable in dry air; it commences to decompose at 350° . A saturated solution gives a faint darkening with hydrogen sulphide and a faint turbidity with silver nitrate. It is decomposed into the metal and a soluble trivalent bismuth salt by sulphuric, hydrochloric, and acetic acids. It is insoluble in alcohol and in aqueous potassium iodide solutions, and reduces acid solutions of potassium permanganate. *Bismuth sub-iodide* is a volatile substance which crystallises in red, orthorhombic needles. In aqueous solution, it gives stronger reactions for bismuth and iodine than the oxyiodide; it dissolves freely in potassium iodide solution, giving solutions of the colour of dichromate solutions. It speedily reduces aqueous solutions of iodine and acid permanganate, and is decomposed at 400° into bismuth tri-iodide and metallic bismuth. The distillate of excess methyl iodide from the preparation of the above compounds was yellow in colour, but on exposure to air it became colourless, and a white solid which turned yellow on keeping separated. This white compound is probably *bismuth dimethyl*, and is insoluble in alcohol; it is a strong reducing agent, and on oxidation is converted into *bismuth dimethoxide*, $\text{Bi}(\text{OCH}_3)_2$.

J. F. S.

Mineralogical Chemistry.

Chemical Aspects of Volcanism with a Collection of the Analyses of Volcanic Gases. E. T. ALLEN (*J. Franklin Inst.*, 1922, 193, 29—80).—The author has collected and tabulated all published analyses of volcanic gases of any importance for the purpose primarily of determining whether any relation exists between the composition of the gases and the temperature at which they issue from the ground. The analyses are made the basis of a critical inquiry into the nature of the original gases and into the extent to which they have become contaminated with atmospheric gases and water. Consideration of the gases contained in igneous rocks leads to the conclusion that these are the source of the original volcanic gases, of which the most important is water. The original gases are probably changed in composition by the time they reach the point of collection; in some cases they probably lose a portion of the strong acid gases by interaction of these with metallic oxides; in most cases they become diluted with steam from surface waters and by atmospheric gases. In some volcanic gases the relations between the inert gases, argon and nitrogen, point to an atmospheric origin for these constituents; in others, they must either be of deep-seated origin or their original atmospheric relation has been changed by selective solubility in the magma. The composition of the original gases may also be changed by interaction with mineral substances, by surface oxidation, or by the shifting of chemical equilibrium due to change of temperature. Such equilibria as the following may be affected: $\text{CO}_2 + \text{H}_2 \rightleftharpoons \text{H}_2\text{O} + \text{CO}$; $3\text{S}_2 + 4\text{H}_2\text{O} \rightleftharpoons 4\text{H}_2\text{S} + 2\text{SO}_2$; $\text{H}_2\text{S} + 2\text{H}_2\text{O} \rightleftharpoons 3\text{H}_2 + \text{SO}_2$; $2\text{H}_2\text{S} \rightleftharpoons 2\text{H}_2 + \text{S}$. These and other reactions are discussed from the thermal point of view. A more or less abrupt evolution of gas from the magma may occur during crystallisation. Energy derived from shifting chemical equilibria may have a considerable influence in prolonging the life of an eruption, but not in initiating volcanic activity. Surface combustion is sometimes an important factor in keeping up the temperature. Secondary explosions of great violence are undoubtedly produced by the access of surface water to hot volcanic ejecta. Primary explosions are probably the result of pressure from magmatic gases, not the result of chemical action. E. H. R.

Absence of Cobalt in Cornetite. ALFRED SCHOEP (*Min. Mag.*, 1922, 19, 301—302).—Cornetite from the original locality, Star of the Congo mine in Katanga, Belgian Congo (A., 1920, ii, 441), is intimately mixed with minute black specks of heterogenite (A., 1921, ii, 649). When freed from these, the mineral shows no cobalt when tested microchemically with mercury thiocyanate. It thus agrees completely with the cornetite from Bwana Mkubwa in Northern Rhodesia (A., 1921, ii, 701). L. J. S.

Chemical Investigation of Japanese Minerals containing Rarer Elements. II. Analysis of Columbite and Monazite of Ishikawa, Iwaki Province. YŪJI SHIBATA and KENJIRŌ KIMURA (*J. Chem. Soc. Japan*, 1921, 42, 957—964; cf. A., 1921, ii, 269).—Columbite (*d* 5.59) from Ishikawa, Iwaki Province, gave on analysis:

Nb ₂ O ₅ , Ta ₂ O ₅ .	SnO ₂ .	Al ₂ O ₃ .	MnO.	FeO.	H ₂ O.	Total.
78.94	0.38	0.36	3.56	15.99	0.90	100.13

with traces of SiO₂, TiO₂, WO₃, and CaO. The amount of Ta₂O₅ was about 10%. The ratio (Fe,Mn)O : (Nb,Ta)₂O₅ = 1 : 1.03.

Monazite (*d* 5.17) from the same locality gave on analysis :

P ₂ O ₅ .	SiO ₂ .	UO ₃ .	ThO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	Ce ₂ O ₃ .	(La,Nd) ₂ O ₃ , etc.	Y ₂ O ₃ , etc.	CaO.	MgO.	H ₂ O.	Total.
27.52	2.98	0.42	11.08	0.80	0.66	21.08	31.27	3.53	0.52	0.27	0.56	100.69

Various samples differed; some contained traces of ZrO₂, TiO₂, and SnO₂, whilst others did not contain UO₃, and the quantities of ThO₂ and SiO₂ were very irregular. On spectroscopic examination, lines of praseodymium, gadolinium, dysprosium, erbium, ytterbium, holmium, and terbium were observed. K. K.

Density, Refractivity, and Composition Relations of some Natural Glasses. C. E. TILLEY (*Min. Mag.*, 1922, 19, 275—294).—The natural glasses fall into two groups: (a) tektites, including moldavites and australites, which are perhaps of meteoric origin, and (b) volcanic glasses, including rhyolitic, trachytic, and basaltic obsidians. The refractive index and density were determined for a number of these, and their specific refractivity ($r_g = (n-1)/d$) compared with that of artificial glasses of definite composition (silica, feldspars, and CaSiO₃-MgSiO₃). When plotted on graphs, the different glasses fall in well-defined areas according to their composition. Values for the specific refractivity of various rock-constituents in a state of glass are calculated. An analysis is given of tachylite forming the selvage of a basic andesite at Kildonan, Eigg, Western Isles of Scotland. An estimation of water in the pitchstone of Newry, Ireland, gave 7.04%. The presence of water has a marked influence on the specific refractivity, as shown in the case of analysed rhyolite-obsidians from the Island of Lipari. L. J. S.

Augite from Hawaii. HENRY S. WASHINGTON and H. E. MERWIN (*Amer. J. Sci.*, 1922, [v], 3, 117—122).—Loose, jet-black crystals of augite from the volcano of Haleakala on the island of Maui, were powdered and pure material separated by heavy solutions and the electromagnet, *d* 3.358. In thin sections the material is pale grey with a darker surface film; extinction-angle 47° (red) to 49° (blue), α 1.700, β 1.706, γ 1.724. An analysis of the powder dried at 110° gave :

SiO ₂ .	TiO ₂ .	Al ₂ O ₃ .	Cr ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	MnO.	CaO.	MgO.	Na ₂ O.	K ₂ O.	H ₂ O.	Total.
47.70	1.89	6.82	0.23	3.36	4.43	0.16	21.35	13.34	0.65	0.03	0.16	100.11

Tschermak's (A., 1921, ii, 121), Boeke's (A., 1914, ii, 283), and Zambonini's (1914) views as to the constitution of aluminous augites are criticised; and the alumina and excess of ferric oxide are regarded as being in solid solution in the pyroxene constituents (diopside-hedenbergite, aegirite-jadeite, and clinoenstatite). The present analysis is interpreted as $\text{CaMgSi}_2\text{O}_6$, 69.12; $\text{CaFeSi}_2\text{O}_6$, 15.13; $\text{NaFeSi}_2\text{O}_6$, 5.08; MgSiO_3 , 1.90; FeSiO_3 , 0.40; $(\text{Al,Fe})_2\text{O}_3$, 8.65%. Assuming, with Zambonini, that the alumina is present as spinel (12.23%), there remains a residue of silica (4.20%) and CaSiO_3 (6.61%).

L. J. S.

Analytical Chemistry.

A Microvolumenometer. A. BOLLAND (*Roczniki Chemji*, 1921, 1, 147—156).—A volumenometer is described which consists of a thick-walled capillary tube which is closed at its lower end; the tube is 50 mm. long and has a capacity of 0.02 c.c., it is accurately graduated into forty divisions each corresponding with 0.0005 c.c. and widens at the top into the shape of a test-tube 40 mm. long. The method of weighing, filling, and emptying the apparatus is described. It is shown that it may be used for quantitative micro-analysis by measuring the volume of precipitates after centrifuging.

J. F. S.

Standard Dropping Pipette. HEDE HALPHEN (*Pharm. Zentr.-h.*, 1921, 62, 767—768).—A pipette which will deliver 20 drops of water per gram should have an orifice 3 mm. in diameter (external measurement); such a pipette can be made by drawing out a glass tube at one end and passing this narrow portion as far as it will go through a 3 mm. hole drilled in a brass plate. The tube is then cut exactly at the surface of the plate.

W. P. S.

Rapid Extraction Funnel. H. WOLFFRAM (*Chem. Ztg.*, 1922, 46, 93).—A very simple apparatus to replace a Soxhlet extractor consists of a cylindrical part somewhat wider than the extraction thimble to be employed, narrowing below to an outlet tube in the usual way. A small glass triangle prevents the thimble from closing up the way into the outlet tube, and three or four indentations in the sides of the glass cylinder prevent the thimble from leaning over against the side of the glass.

G. F. M.

Proposed Standard Method of Colorimetry. HERBERT E. IVES (*J. Opt. Soc. Amer.*, 1921, 5, 469—478).—For the purpose of simplifying spectrometric measurements for colour specification, the author has designed a method of colour measurement, which consists essentially of the spectrophotometry of adjacent patches of the spectrum, each patch of a width fixed by the hue scale, and

narrow enough so that there is no colour difference in the spectrophotometer field at each setting. The number of patches is determined by the kind of colour and the degree of accuracy required. An instrument is described by which the measurements can be made and in which any colour may be reproduced for comparison. J. F. S.

The Salt Error of Coloured Indicators. I. M. KOLTHOFF (*Rec. trav. chim.*, 1922, 41, 54—67; cf. Sørensen, A., 1909, i, 861; 1910, i, 147).—The corrections that should be made for the presence of solutions of sodium chloride and potassium chloride of concentrations ranging from decinormal to normal have been determined for a large number of indicators. The results are discussed in detail for each indicator and a table summarising the chief results is appended, together with notes as to the suitability of the different indicators for various conditions. H. J. E.

The Colorimetric Estimation of Hydrogen-ion Concentration without Buffer Solutions. I. M. KOLTHOFF (*Pharm. Weekblad*, 1922, 59, 104—118).—Hydrogen-ion concentration may be estimated by comparing the colour given with standard indicators with the colours given by mixtures of coloured salts in specified proportions. For many indicators, suitable comparison solutions may be made by mixing a solution of 11.262 grams $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ in 250 c.c. of 1% hydrochloric acid solution with a solution of 18.2 grams of crystallised cobalt nitrate in 250 c.c. of 1% hydrochloric acid solution; both these solutions may be employed after long keeping. Mixtures of these salt solutions in various proportions given in the tables show the same tints as the following: Neutral-red in solutions with range p_{H} from 7.0 to 8.0; methyl-orange, range 3.05—4.60; tropæolin-OO, range 1.80—3.00; methyl-red 5.2—6.0. For solutions having p_{H} between 4.2 and 6.0, where methyl-red is used as indicator, suitable comparison solutions may be made up with permanganate solution (0.004*N*) and dichromate solution (0.01*N*—potassium dichromate in 0.4*N*— H_2SO_4), but these solutions are not stable in colour, and must be prepared afresh as required. For the range 4.4—6.0 with this indicator, the ferric chloride solution may be mixed with one obtained by mixing 10 c.c. of 0.025% pure methyl-red solution with 10 c.c. of 4*N*-acetic acid solution, and diluting to 250 c.c.; the latter solution is stable for about a week.

For the range of p_{H} between 8.2 and 10.0, with phenolphthalein as indicator, the comparison solutions are best prepared by adding to 10 c.c. of 0.5*N*-sodium carbonate solution given volumes of a 0.004% phenolphthalein solution. From the figures prepared for this table, the dissociation constants of phenolphthalein have been calculated; the values agree approximately with the figures deduced from theory (assuming the indicator is a bibasic acid) by Rosenstein (A., 1912, ii, 893).

For the range of p_{H} between 6.0 and 7.0, *p*-nitrophenol is a suitable indicator, the colour comparison solutions being prepared in this case also from an alkaline solution of the indicator. With this

range completed, it is possible to use the method for all values of p_H between 2.0 and 10.0. S. I. L.

The Titration of Moderately Strong Acids or Bases in the Presence of Very Weak Ones. I. M. KOLTHOFF (*Pharm. Weekblad*, 1922, 59, 129—142).—The observation of Tizard and Boeree (T., 1921, 119, 132) that the hydrogen-ion concentration is equal to $\sqrt{K_1 K_2}$ in the case of two acids having the same concentration, in the presence of sufficient alkali to neutralise the stronger one, and where K_1 and K_2 are the dissociation constants of the two acids, is confirmed theoretically and by experiment. To obtain accurate results by titration, it is necessary that the ratio $K_1 : K_2$ should be not less than 1×10^4 , and the approximate concentrations must be determined before the final titration is made. For this purpose, the approximate concentration of both acids is first determined by titration with *N*-sodium hydroxide solution in presence of tropæolin-*O*, and the concentration of the stronger acid by titration in presence of neutral-red or phenol-red. The value of p_H can then be calculated from the above, and a comparison solution made up from the sodium salt of the strong acid and the necessary proportion of the weak acid, or a buffer solution of the same p_H can be selected. With this comparison solution to give the colour required for the particular indicator employed, the accurate titration may be made. Generally, neutral-red or an indicator of the same transition range is most suitable. S. I. L.

Formation of Nitrogen Oxides in the Slow Combustion and Explosion Methods in Gas Analysis. G. W. JONES and W. L. PARKER (*J. Ind. Eng. Chem.*, 1921, 13, 1154—1155).—When the time of burning is not more than three minutes and the platinum wire is not heated above bright yellow, the formation of nitrogen oxides in the slow combustion method does not exceed 0.003 c.c., and nitrogen oxides are not produced in the explosion method when air is used as the oxygen supply. If a mixture of air and oxygen is employed in the explosion method, appreciable quantities of nitrogen oxides are formed and the error thus introduced may amount to 2%; their formation is due to increase in the flame temperature produced by the addition of oxygen.

W. P. S.

Estimation of Water in Fuels. ANDRÉ MARINOT (*Ann. Chim. Analyt.*, 1922, [ii], 4, 7—8).—In the estimation of water in fuels or fuel oils, errors may be introduced by the absorption of oxygen or the distillation of anthracene concurrently with the evaporation of the water, even at temperatures below 100°. These errors are avoided by carrying out the operation in a current of dry hydrogen, and an apparatus is described for the purpose consisting of two 100 c.c. flasks connected together and heated in a constant level water-bath. From 10—15 grams of the material are placed in each flask and a slow current of hydrogen dried in a calcium chloride-sulphuric acid tower is passed through for one and a half hours. The vapours from the flasks pass through a

horizontal elongated bulb where any anthracene is deposited into a weighed U-tube containing calcium chloride, the exit of which is protected from atmospheric moisture by a guard tube.

G. F. M.

Potassium Ferricyanide as a Reagent in Iodometry.

I. M. KOLTHOFF (*Pharm. Weekblad*, 1922, 59, 66—68).—The accuracy of the author's method of standardising thiosulphate solutions by the use of potassium ferricyanide (*ibid.*, 1919, 56, 1618) having been questioned, a further examination has been made. The method is found extremely accurate and very rapid. The ferricyanide must be pure and dry, and if the zinc sulphate and hydrochloric acid used are free from iron, a perfectly white precipitate of zinc ferrocyanide is obtained, and the end-point with starch is very sharp.

S. I. L.

High Percentage Hydrogen Peroxide (Perhydrol) for the Estimation of the Total Sulphur in Illuminating Gas. ALOYS KLEMMER (*Chem. Ztg.*, 1922, 46, 79).—The sulphur compounds, including hydrogen sulphide, carbon oxysulphide, carbon disulphide, and mercaptan, are oxidised to sulphuric acid by passing the gas through a strongly alkaline solution of hydrogen peroxide. Ten c.c. of perhydrol are mixed with 80 c.c. of fairly concentrated sodium hydroxide, and the gas is led through the thick, crystalline paste consisting of sodium peroxide, $\text{Na}_2\text{O}_2 \cdot 8\text{H}_2\text{O}$, which is formed, at a rate not exceeding 100 litres per hour. At the end of the operation the liquid is acidified with hydrochloric acid, boiled to expel the excess of hydrogen peroxide, and the sulphuric acid precipitated as barium sulphate.

G. F. M.

Estimation of Sulphur in Cast Irons and Steels. ANDRÉ MARINOT (*Ann. Chim. Analyt.*, 1922, [ii], 4, 5—6).—Five grams of the metal are treated with a reagent consisting of 30 c.c. of 50% sulphuric acid and 60 c.c. of hydrochloric acid in an apparatus consisting of a conical reaction flask of 375 c.c. capacity surmounted by a vertical condenser into the top of which is ground a small pear-shaped gas washer from which a delivery tube leads into a flask containing 200 c.c. of a 2.5% solution of zinc acetate acidified with acetic acid. The gas washer consists of a vertical narrow tube surmounting the condenser and extending almost to touch the apex of a conical muff which surrounds it and is sealed to the tube at the base, where, however, it is perforated with five or six small holes which allow of the escape of the gas into the outer envelope of the pear and thence through the delivery tube into the zinc acetate flask. All the sulphur in the metal is evolved as hydrogen sulphide, and a slow stream of carbon dioxide is passed through the whole apparatus to displace the gas and prevent the formation of colloidal sulphur or organic sulphur compounds. The sulphur is finally estimated by oxidising the zinc sulphide formed in the flask by standard iodine solution and titrating back the excess.

G. F. M.

Estimation of Available Sulphur in Golden Sulphide of Antimony. B. D. W. LUFF and B. D. PORRITT (*J. Soc. Chem. Ind.*, 1921, 40, 275—278T).—It is suggested that the "available sulphur" be estimated by extraction with carbon disulphide after the pigment has been heated at 150° for five hours in an atmosphere rendered slightly alkaline with ammonia; this heating converts any amorphous sulphur which may be present into a form which is soluble in the solvent. The carbon disulphide used should be purified previously by distillation over sulphur. W. P. S.

Sulphates in Blood. W. DENIS (*J. Biol. Chem.*, 1921, 49, 311—317).—A nephelometric method for the estimation of inorganic sulphates in blood or plasma is described. Proteins are removed by means of mercuric chloride and the precipitate produced by the addition of acidified barium chloride solution is compared with a standard in a nephelometer. No evidence was obtained of the existence in deproteinised blood of other compounds of sulphur. E. S.

A Micro-method for Estimation of Nitrogen. D. ACÉL (*Biochem. Z.*, 1921, 121, 120—124).—After destruction of the organic matter with sulphuric acid in the usual manner, the ammonia formed is not distilled off, but is determined colorimetrically by Nessler's reagent. The control is treated with standard ammonium chloride solution until the colours match. The method is suitable for the estimation of nitrogen in as little as 0.001—0.003 c.c. of serum or urine if diluted for measurement. H. K.

A Rapid Method for the Estimation of Ammoniacal Nitrogen. R. MEURICE (*Ann. Chim. Analyt.*, 1922, [ii], 4, 9—10).—In the estimation of ammonium salts by converting them into hexamethylenetetramine and titrating the free acid thus produced, errors are likely to occur if phenolphthalein is used as indicator in the preliminary exact neutralisation of the ammonium salt owing to the uncertainty of this indicator in presence of ammonia. This error is eliminated if rosolic acid is used as indicator for the preliminary neutralisation, but as under ordinary conditions it is also sensitive to hexamethylenetetramine a special device is adopted to render it insensitive. After the mixture of ammonium salt and formaldehyde has been kept about thirty minutes, an equal volume of ether is added, and the whole is well shaken, whereby the rosolic acid passes into the ether and becomes insensitive to the amine, although still sensitive to a strong base such as sodium hydroxide. Titration of the free acid with standard alkali can now therefore be proceeded with until the appearance of a pale rose colour which persists on agitating the liquid with the ether. G. F. M.

Precipitation of Arsenic Sulphide from Arsenates. J. H. REEDY (*J. Amer. Chem. Soc.*, 1921, 43, 2419).—The time required for precipitation of arsenic sulphide from solutions of arsenates may be greatly reduced by the addition of a small quantity of a soluble iodide, such as ammonium iodide. The hastening of the

precipitation is due to the reduction of the arsenic acid to arsenious acid, according to the equations $\text{H}_3\text{AsO}_4 + 2\text{HI} \rightarrow \text{H}_3\text{AsO}_3 + \text{H}_2\text{O} + \text{I}_2$; $\text{H}_2\text{S} + \text{I}_2 \rightleftharpoons 2\text{HI} + \text{S}$. The reaction is applied by adding 1—2 c.c. of *N*-ammonium iodide solution to the hot solution which contains 4 c.c. of 6*N*-hydrochloric acid in 40 c.c. of solution, just before the hydrogen sulphide is passed in. Precipitation begins immediately and is usually complete in four to five minutes. Complications arise in the precipitation of mercury and copper as iodides and in the partial reduction of mercury to the mercurous condition. This difficulty is, however, removed during the digestion with yellow ammonium sulphide which oxidises both metals to the higher valency, precipitating them as mercuric and cupric sulphides. J. F. S.

Method for the Estimation of Free and Combined Carbon Dioxide. J. A. SHAW (*J. Ind. Eng. Chem.*, 1921, 13, 1151—1152).—A definite quantity of the solution under examination (for example, mine water) is drawn into a cylindrical bulb which is fitted at the top and bottom with three-way taps; the tap at the top connects with a small funnel for the introduction of the sample and with a measuring burette, whilst the tap at the bottom connects with a smaller cylindrical bulb and with a mercury reservoir. At the commencement of the operation the whole apparatus is filled with mercury. Any gas liberated from the sample is passed over into the burette, the upper tap is then closed and the mercury reservoir lowered so that the whole of the liquid is drawn into the lower cylinder, a low pressure being thus produced in the upper cylinder and above the surface of the liquid. The gas collecting in the upper cylinder is then forced over into the burette by raising the mercury reservoir, the tap between the two cylinders being meanwhile closed, and, by repeating these operations several times, the whole of the carbon dioxide may be collected and its volume measured in the burette. Concentrated sulphuric acid is added after the sample has been introduced into the cylinder when it is desired to estimate the total (free and combined) carbon dioxide. W. P. S.

Laboratory Notes. [Estimation of Silica, Phosphorus in Iron, etc.]. ERNST MURMANN (*Österr. Chem. Zeit.*, 1921, 24, 142).—In the estimation of silica, two evaporations with an intervening filtration are necessary in order to render the whole of the silica insoluble, care being taken that the evaporation is carried to complete dryness of the residue. To estimate phosphorus in iron, soil extracts, etc., the sample is evaporated with hydrochloric and nitric acid, and the residue heated in order to destroy any organic matter; by extracting the residue with nitric acid, the phosphoric acid is obtained in solution and it is not necessary to dissolve the ferric oxide resulting from the decomposition of the ferric nitrate during the heating. The grinding of ores is facilitated if the powder is moistened with ether, and graphitic carbon burns readily if it has been moistened previously with a small quantity of magnesium acetate solution. The use of chalk, mineral phos-

phate, or bone-ash is recommended in place of sand in sand-baths, since sand is liable to scratch glass vessels and cause them to break when heated.

W. P. S.

Volumetric Estimation of Potassium. MACHELEIDT (*Woch. Brau.*, 1922, 39, 23—24).—A standard solution of sodium hydrogen tartrate is prepared by dissolving 60 grams of tartaric acid and 16 grams of sodium hydroxide in water and diluting to 1 litre. Six grams of potassium hydrogen tartrate are added and the solution is shaken for several hours. Thirty c.c. are filtered off and titrated with *N*/10-barium hydroxide solution. A second 30 c.c. are shaken for one to two hours with 0.5—0.75 gram of the salt mixture to be tested, the solution is filtered into a tared basin and, without washing the filter, titrated with the barium hydroxide solution. The solution is weighed before and after filtering, and allowance is made for the loss. The difference between the two titrations is calculated to potassium oxide. [See further *J. Soc. Chem. Ind.*, 1922, 200.]

A. R. P.

Estimation of the Calcium Content of Blood. R. WEISS (*Deut. med. Woch.*, 1921, 47, 1298; from *Physiol. Abstr.*, 1922, 6, 568).—A small measured quantity of serum is placed in a specially graduated tube, ammonium oxalate is added, and the precipitate washed on the centrifuge with water. It is dissolved in sulphuric acid and titrated with permanganate. The figures given are: Normal 12, in tetany as low as 4, in rickets as high as 18 mg. per 100 c.c.

E. S.

Micro-estimation of Calcium in Whole Blood, Plasma, and Serum by Direct Precipitation. GUY W. CLARK (*J. Biol. Chem.*, 1921, 49, 487—517).—The method described is practically identical with that of Halverson and Bergeim (*A.*, 1916, ii, 270).

E. S.

Sensitiveness and Applicability of Qualitative Reactions.
III. Strontium Ions. O. LUTZ (*Z. anal. Chem.*, 1921, 60, 433—441; cf. *A.*, 1921, ii, 596).—The minimum quantities of strontium which can be detected by various reagents are as follows: in each case, 5 c.c. of the strontium salt solution were treated with 0.5 c.c. of the reagent in the cold and the observation made after five minutes. Sodium phosphate, 1:9400; sodium sulphite, 1:12000; ammonium oxalate, 1:50000; ammonium carbonate and ammonia (at 100°), 1:210000; sulphuric acid, 1:125000. The addition of 5 c.c. of alcohol increases the sensitiveness of the sulphuric acid test to 1:1400000.

W. P. S.

Estimation of Magnesium in Technical Nickel. K. CHALUPNY and K. BREISCH (*Chem. Ztg.*, 1922, 46, 91).—For the estimation of small quantities of magnesium in technical nickel 10—15 grams of the metal must be used, and the filtration and washing of the voluminous precipitate of nickel sulphide are very laborious and a frequent source of error. The advantage of a

method whereby the magnesium could be precipitated whilst the nickel remained in solution is apparent, and the non-precipitation of nickel phosphate from potassium nickelocyanide by sodium phosphate affords a means of attaining this object. The analysis is carried out as follows: 15–20 grams of the metal are dissolved in nitric acid, and the solution is twice evaporated to dryness with hydrochloric acid to precipitate silicic acid. To the filtered solution is then added 5–10 grams of ammonium chloride, followed by bromine water and ammonia to precipitate iron, manganese, etc. The filtrate is approximately neutralised with hydrochloric acid, and a concentrated solution of potassium cyanide is added until the precipitate of nickel cyanide is just redissolved, followed by 20 c.c. of 10% disodium hydrogen phosphate solution and a third of the volume of concentrated ammonia. After keeping overnight, the crystalline precipitate of magnesium ammonium phosphate is separated, and washed free from nickel. It is dissolved in dilute hydrochloric acid and reprecipitated with sodium phosphate and ammonia (this step is necessary on account of the large excess of alkali cyanide, present during the first precipitation, which causes somewhat too high results). The precipitate is collected in a Gooch crucible, ignited, and weighed as magnesium pyrophosphate in the usual way.

G. F. M.

Estimation of Zinc as Zinc Pyrophosphate. D. BALAREFF (*Z. anal. Chem.*, 1921, 60, 442–448).—The precipitation of the zinc phosphate should be made under the following conditions. The slightly acid solution, containing ammonium chloride and ammonium phosphate is treated with ammonia until it is slightly alkaline in reaction towards litmus; the mixture is kept at the ordinary temperature for eighteen hours, then heated on a water-bath for fifteen minutes, and the precipitate is collected, ignited, and weighed. The amorphous precipitate which first forms when zinc is precipitated as phosphate from a solution containing ammonium salts is not zinc phosphate but ammonium zinc phosphate; if the zinc is precipitated by treating the hot solution with ammonia, ammonium zinc phosphate and zinc phosphate are precipitated together, and the results obtained are too low (cf. Austin, A., 1900, ii, 49, and Dakin, A., 1900, ii, 624). W. P. S.

The Separation of Aluminium from Glucinum. III. HUBERT T. S. BRITTON (*Analyst*, 1922, 47, 50–60; cf. A., 1921, ii, 657, 712).—Berzelius's method, consisting in boiling the precipitated hydroxides with ammonium chloride solution whereby the glucinum hydroxide is dissolved, is unsatisfactory, as no means could be found by which the occlusion of glucinum hydroxide by aluminium hydroxide could be eliminated, the results being accordingly low for glucinum and correspondingly high for aluminium. Wünder and Wenger's sodium carbonate fusion method (A., 1912, ii, 687) is satisfactory, but the time required for an analysis is long, as two fusions are necessary for a complete separation. The thiosulphate method in which the neutral salt solutions are boiled

with an excess of sodium thiosulphate until evolution of sulphur dioxide has ceased does not give quantitative separations owing to adsorption of glucina by the aluminium hydroxide which is precipitated. Haven's ether-hydrochloric acid method (A., 1898, ii, 142) is quantitative and is one of the most satisfactory and easily manipulated of all the methods investigated. Of the remaining methods, none were investigated, but it is probable that only Kling and Gelin's basic acetate distillation method (A., 1914, ii, 867) and Renz's ethylamine method (A., 1903, ii, 729) are quantitative, and as the former requires considerable manipulation and time, and the latter involves the use of a large quantity of an expensive reagent, they have no particular feature to recommend them.

G. F. M.

Detection of Manganese with Benzidine and of Cobalt by means of the Thiocyanate Reaction. HUGO DIRZ (*Chem. Ztg.*, 1922, 46, 121—122).—The author claims priority for the method described by Feigl and Stern (A., 1921, ii, 278) of detecting traces of manganese by the blue coloration produced on adding an acetic acid solution of benzidine to the peroxidised manganese compound formed by autoxidation in an alkaline medium. As little as 0.000008 gram of manganese can be detected by this method. No interference with the reaction by iron salts occurs provided a considerable excess of acetic acid is present, and it can therefore be used for the detection of manganese in iron ores and slags. Vogel's thiocyanate reaction for cobalt (*Ber.*, 1879, 12, 2314) is rendered more sensitive by using acetone instead of either amyl or ethyl alcohols, and quantities of the order of 0.003 mg. of the metal can be detected by this means.

G. F. M.

Separation of Molybdenum and Tungsten by means of Selenium Oxychloride. HENRY BALDWIN MERRILL (*J. Amer. Chem. Soc.*, 1921, 43, 2383—2387).—Mixtures of molybdenum trioxide and tungsten trioxide may be quantitatively separated by boiling 1 gram of the mixture with 30 c.c. of a 1 : 1 mixture of selenium oxychloride and concentrated sulphuric acid for sixty minutes. The solution is decanted through a weighed Gooch crucible and the residue washed several times with small quantities of selenium oxychloride and finally brought on to the filter with a hot 10% solution of ammonium nitrate. The crucible is ignited and weighed, and gives the weight of tungsten trioxide, the molybdenum trioxide being obtained by difference. The above method, which is effective for mixtures made by mixing the two oxides by hand, does not work with mixtures of the two oxides precipitated together, if the amount of tungsten trioxide is greater than 10%. In such cases, it is impossible to dissolve all the molybdenum trioxide owing to the formation of solid solutions. This difficulty is overcome by dissolving the oxides in ammonia, adding sufficient nitric acid to precipitate most of the tungsten, evaporating to dryness, and proceeding as described above. The method gives good results.

J. F. S.

Technical Estimation of the Colloidal Part of Tungsten Powder. A. LOTTERMOSER (*Kolloid Z.*, 1922, 30, 53—61).—Two methods are described for the estimation of the percentage of colloidal tungsten in commercial tungsten powders. The methods are (1) a sedimentation method, and (2) an optical method, both of which lead to approximately the same results. Sedimentation method: the sample (20 grams) is shaken thoroughly with 100 c.c. of water in a tube and allowed to sediment for two days, 75 c.c. of the supernatant liquid are removed, and 75 c.c. of water added to the sediment. The mixture is shaken and allowed to sediment for a further two days. The process is repeated as long as a measurable quantity of tungsten remains in the supernatant liquid. The sediment is then dried and weighed and gives the non-colloidal portion. The addition of ammonia to the water shows that whilst most specimens of tungsten powder are very finely divided and give suspensions, some only are truly colloidal, inasmuch as they are peptised by ammonia. Optical method: this consists in estimating the quantity of tungsten in the solutions from which the powder has sedimented by means of its absorption of light. The light from a quartz mercury lamp is allowed to pass through the solution on to a potassium photo-electric cell and the absorption determined from the galvanometer deflection. In this connexion, it is shown that the de Beer-Fick law is applicable. Five specimens of tungsten powder have been examined and found to contain respectively 7.5%, 5.5%, 11.25%, 39.75%, and 7.0% of colloidal tungsten. The fourth and fifth samples are peptised by ammonia. J. F. S.

Separation of Columbium and Tantalum by means of Selenium Oxychloride. HARRY BALDWIN MERRILL (*J. Amer. Chem. Soc.*, 1921, 43, 2378—2383).—A method of analysis of mixtures of tantalum and columbium oxides, and for the preparation of the pure oxides is described. The mixed oxides together with titanium oxide if such be present are separated together from the mineral, ignited, and weighed. A sample (0.2—0.3 gram) of the mixed ignited oxides is boiled in an Erlenmeyer flask with 50 c.c. of a 1:1 mixture of selenium oxychloride and concentrated sulphuric acid for thirty minutes on a sand-bath, care being taken that clouds of vapour are not evolved. The solution after cooling is decanted on to an asbestos pad in a Gooch crucible and filtered by suction. The filtrate is poured into a large volume of water and boiled, when hydrated columbium pentoxide is precipitated. The residue is boiled with 20 c.c. of the 1:1 mixture for fifteen minutes, decanted, and treated as before and the process repeated until the filtrate on hydrolysis gives only a faint cloudiness due to traces of tantalum pentoxide. The residue from the flask is now washed into the Gooch crucible and without much washing the crucible is ignited and weighed; the gain in weight gives the amount of tantalum pentoxide, whilst the loss of weight gives the columbium pentoxide, with titanium if this is present. The method gives results which have a maximum error of 3% and

is therefore better than the Marignac method whilst at the same time being more rapid.

To prepare pure columbium oxide, the mixed oxides are extracted with sufficient solvent to dissolve all the columbium, but it should not be boiled until all the columbium is dissolved, since this would mean the solution of much tantalum oxide. The dissolved oxide is precipitated with water and ammonia, filtered, and ignited. It still contains some tantalum oxide which by repeating the treatment can be removed and very pure columbium oxide obtained. Pure tantalum oxide is prepared by boiling the mixed oxides with the reagent until all the columbium oxide is dissolved; thus whilst sacrificing a little tantalum oxide, a very pure product is obtained. The complete removal of the columbium oxide is hastened by adding a little more sulphuric acid to the 1 : 1 mixture of selenium oxychloride and sulphuric acid.

J. F. S.

The Estimation of Small Quantities of Antimony in Copper and Brass. B. S. EVANS (*Analyst*, 1922, 47, 1—9).—Five grams of the sample are dissolved in 60 c.c. of nitric acid (*d* 1.2) and 10 c.c. of sulphuric acid and the solution evaporated until it fumes strongly. When cold, the mass is dissolved in 100 c.c. of water, 14 grams of sodium hypophosphite are added, and the solution is heated nearly to boiling whereby the copper is precipitated. The solution is filtered and the precipitate washed with hot water. A further 2 grams of sodium hypophosphite and 100 c.c. of hydrochloric acid are added to the filtrate and the liquid is boiled for fifteen minutes to precipitate any arsenic present. After cooling slightly, 10 c.c. of benzene are added to the filtrate and the liquid is well shaken so that the colloidal arsenic becomes suspended in the benzene layer. The liquid is filtered through a wet filter (to retain the benzene and arsenic) and the latter is washed twice with warm water. The filtrate is heated to boiling and a spiral roll of clean copper foil is dropped in and the boiling continued for one and a half to two hours; the liquid is then poured off, the copper washed rapidly with cold water, covered with water, and treated with 1 gram of sodium peroxide. The liquid is warmed until the deposited antimony has dissolved, then decanted off and the strip washed with cold water. The solution is treated with 0.5 gram of zinc sulphide and, after one and a half to two hours, filtered. The filtrate is acidified with hydrochloric acid, treated with sulphur dioxide, and evaporated to 10 c.c. Five c.c. of standard antimony solution (1 c.c.=0.0001 gram Sb), 80 c.c. of water, and a few drops of hydrochloric acid are treated with sulphur dioxide, and the solution is evaporated to 10 c.c. Both assay and standard solutions are treated with 5 c.c. of 1% gum arabic solution, diluted to 100 c.c., treated with hydrogen sulphide for a few seconds, and transferred to Nessler tubes. The liquid having the greater depth of colour is poured from the glass until the colours match; the depth of the two liquids is measured and the result calculated from these figures. The standard antimony solution is made by dissolving 0.2764 gram

of potassium antimonyl tartrate in 100 c.c. of hydrochloric acid and diluting to 1 litre.

A. R. P.

I. Estimation of Methyl Alcohol in Remains for Forensic Purposes. II. Occurrence of Methyl Alcohol in the Human Body. H. JANSCH (*Vrtiljschr. ger. Med. öffentl. Sanitätsw.*, 1921, 62, 1—18; from *Chem. Zentr.*, 1921, iv, 983—984).—A weighed portion of the finely divided remains is acidified with tartaric acid and distilled in a current of steam. The distillate (500—1500 c.c.) is repeatedly redistilled, recovering about 60% each time, until a distillate of 5 to 10 c.c. is obtained. Fatty acids are removed by filtration. The later distillations are carried out with alkaline reaction. Acetaldehyde and glycerol which may be present are suitably oxidised. The density and refractive index of the end distillate are determined; from these the amount of methyl alcohol present is calculated, using appropriate tables. For the detection of methyl alcohol, the iodoform, benzoyl chloride, and morphine-sulphuric acid tests are used. Methyl alcohol is a normal constituent of human fæces and urine with mixed diets. It originates probably from pectins in the food.

G. W. R.

Apparatus for the Estimation of Methoxyl Groups. WILLIAM M. CUMMING (*J. Soc. Chem. Ind.*, 1922, 41, 20r).—An improved form of apparatus for the estimation of methoxyl groups by Hewitt and Jones's modification of the Zeisel method in which the methyl iodide is absorbed in pyridine, consists of a round-bottomed flask of 250 c.c. capacity with a neck 10 inches long to which a delivery tube is attached by a ground glass joint. A thermometer is provided with its bulb opposite the delivery exit, and a carbon dioxide inlet tube. The delivery tube leads to a narrow bored U-tube, one arm of which contains four convolutions, and each convolution a bulb. This absorber holds about 10—15 c.c. of pyridine, and by passing a very slow current of carbon dioxide the whole of the methyl iodide is carried out of the flask and completely absorbed. The thermometer should not register more than 40° (cf. T., 1919, 115, 1030).

G. F. M.

A Method for the Estimation of Trimethylene Glycol in Crude Glycerol. L. V. COCKS and A. H. SALWAY (*J. Soc. Chem. Ind.*, 1922, 41, 17—20r).—The method previously described (*ibid.*, 1918, i, 123, 158r) consisting in the distillation of the crude glycerol and the determination of the specific gravity and acetin value of the distillate, from which figures the trimethylene glycol content was calculated from the known specific gravity and acetin values of the pure substances, gives only approximate results, as no allowance is made for alterations in volume when glycerol, water, and trimethylene glycol are mixed. The specific gravity of a series of mixtures containing known proportions of these three substances has now been systematically determined, and tables and curves have been constructed by means of which the trimethylene glycol content of any glycerol distillate of known gravity and apparent

glycerol content can be deduced. The pure trimethylene glycol prepared for the purpose boiled at 210—211°/760 mm., or 171°/174 mm.; $d_{20}^{20}=1.0554$. The reduction in specific gravity with increasing trimethylene glycol content was fairly regular, and as a simple method for calculating the amount present, apart from reference to the curves, all that is necessary is to divide by the given factor the difference between the observed sp. gr. and the sp. gr. of pure glycerol of the strength indicated by the acetin figure of the mixture. The factor in question increases regularly with the acetin value from 0.00134 per 1% trimethylene glycol for 50% acetin to 0.00179 for 95% acetin value. An accuracy of ± 0.2 is claimed. G. F. M.

Test for Sucrose in the Presence of Dextrose. LEON A. CONGDON and CHARLES R. STEWART (*J. Ind. Eng. Chem.*, 1921, 13, 1143—1144).—When a dry mixture of sucrose and dextrose is extracted with hot ethyl acetate, the dextrose dissolves and the sucrose remains insoluble; on cooling the ethyl acetate solution, crystals of dextrose are obtained. W. P. S.

Clarification of Urines by Zinc Ferrocyanide. C. CARREZ (*Ann. Chim. Analyt.*, 1922, [vi], 4, 11—12).—Polemical. The author points out that his method involving the use of potassium ferrocyanide and an excess of zinc acetate (A., 1908, ii, 329) was never intended as a preliminary step for any estimation other than that of glucose in the urine, and Thiéry's criticism (A., 1921, ii, 527) therefore has no point. G. F. M.

Colour Reaction of Sucrose. FERDINAND KRYZ (*Oesterr. Chem. Ztg.*, 1921, 24, 141—142).—When a mixture of 1 c.c. of saturated ammonium nickel sulphate solution, 1 c.c. of sucrose solution, and a few drops of sulphuric acid or hydrochloric acid is boiled, the green colour of the mixture changes to yellow and then to red; this red coloration persists even when the mixture is cooled. The reaction cannot be obtained with less than 0.005 gram of sucrose, but other sugars do not interfere. The coloration is not obtained when nitric acid is used in place of the sulphuric acid or hydrochloric acid. W. P. S.

Re-testing the 100°-point of the Saccharimeter. II. Preparation of Chemically Pure Sucrose. ANTON KRAISY (*Z. Ver. deut. Zuckerind.*, 1921, 785—797).—A method of procedure for recrystallising refined sugar from alcohol for the preparation of pure sucrose for purposes of standardisation is described. The product obtained after several successive crystallisations is considered to be free from invert-sugar. Such a product yielded 0.002—0.005% of ash, and reduced 36—38 mg. of copper under the well-known Herzfeld conditions for estimating invert-sugar in sucrose, and 1.5—1.8 mg. under the conditions of the method recently proposed by the author (*ibid.*, 123). [See also *J. Soc. Chem. Ind.*, 1922, 151A.] J. H. L.

Detection of Formic Acid in Wine. W. FRESSENIUS and L. GRÜNHUT (*Z. anal. Chem.*, 1921, 60, 457—463).—The wine is acidified with sulphuric acid and extracted with ether; the ethereal extract is then shaken with dilute sodium hydroxide solution, the alkaline aqueous solution is separated, evaporated to dryness, and the residue heated at 130° for one hour to remove any traces of formaldehyde. The residue is dissolved in 10 c.c. of water and the solution treated with 5 c.c. of hydrochloric acid (*d* 1.12) and 0.4 gram of magnesium turnings. After two hours, the mixture is distilled, 5 c.c. of distillate being collected; this distillate is boiled for one minute with the addition of 2 c.c. of milk and 7 c.c. of hydrochloric acid (*d* 1.12) containing a trace of ferric chloride. If the wine contained formic acid or its salts, a violet coloration develops in the mixture. W. P. S.

Apparatus for Measuring the Hydrogen-ion Concentration of a Solution. Application to the Detection of Mineral Acids in Vinegar. ANDRÉ KLING, A. LASSIEUR, and (MME) A. LASSIEUR (*Compt. rend.*, 1922, 174, 165—168).—A compensation electro-metric method for measuring hydrogen-ion concentration is described in which a millivoltmeter capable of measuring 1200 millivolts with an accuracy of 1 millivolt is used. The hydrogen-ion concentration of vinegar measured with this apparatus or by the colorimetric method, using thymolsulphonphthalein as indicator, serves as a ready means of detecting the presence of mineral acids. The presence of 0.24% of sulphuric acid in vinegar alters the p_H value at 18° from 2.67 to 1.96. W. G.

Estimation of β -Hydroxybutyric Acid. Estimation of Acetone Substances in the Urine. Estimation of Acetone Substances in the Blood. ROGER S. HUBBARD (*J. Biol. Chem.*, 1921, 49, 351—357, 357—374, 375—384).—A modification of Shaffer's method (*A.*, 1914, ii, 77; 1916, ii, 352) for the estimation of β -hydroxybutyric acid is described, in which the time required for the oxidation is reduced to half an hour. The method is applied to urine after removal of interfering substances by precipitation with basic lead acetate, copper sulphate, and sodium hydroxide. The estimation of acetone plus acetoacetic acid in urine is made more accurate by adding to the technique of Shaffer a distillation from acid potassium permanganate solution. The same method is applied to blood after treatment of the latter with colloidal iron, basic lead acetate, and sodium hydroxide.

It is found that in normal urine the total acetone averages about 2 mg. per 100 c.c., the greater part representing β -hydroxybutyric acid; in blood, the total acetone varies normally between 0.1 and 1.0 mg. per 100 c.c. C. R. H.

Effect of the Presence of Filter-paper on Permanganate-Oxalate Titrations. STEPHEN G. SIMPSON (*J. Ind. Eng. Chem.*, 1921, 13, 1152—1154).—Filter-paper reduces permanganate rapidly, especially when the paper is highly disintegrated. In the titration of calcium oxalate precipitates with permanganate solution, the

precipitate should be washed off the filter-paper with hot water and the paper added only when the titration is nearly complete.

W. P. S.

The Estimation of *H*-Acid. HENRY R. LEE (*J. Ind. Eng. Chem.*, 1921, 13, 1049—1051).—The choice of a diazonium salt for use as a standard solution for the estimation of the hydroxyl group should depend on the stability of the salt in the solution in which it is prepared and in which it is to be used, the completeness of the coupling with the intermediate product in question, and the rate of coupling. Comparative experiments in which *H*-acid was titrated with benzenediazonium chloride and with *p*-toluenediazonium chloride respectively show that the rate of decomposition of diazobenzene is approximately eight times as rapid in acid solution, and one and a half times as rapid in alkaline solution, as that of *p*-diazotoluene. Moreover, the rate of coupling of *p*-diazotoluene is slightly more rapid than that of diazobenzene, whilst the secondary coupling, which is marked in the case of commercial samples of *H*-acid titrated with diazobenzene, is very slight when coupled with diazotoluene, and consequently, in the latter case, the end-point is more definite. The use of *p*-diazotoluene in the estimation of *H*-acid, γ -acid, *J*-acid, *S*-acid, and other naphthol- and aminonaphthol-sulphonic acids is recommended. A stock solution of *p*-toluidine hydrochloride is prepared by dissolving 10.7082 grams of pure *p*-toluidine in 40 c.c. of pure concentrated hydrochloric acid, and diluting with water to 1 litre. One hundred c.c. of this solution are placed in a 250 c.c. graduated flask, cooled in ice and salt until frozen, and diazotised with 102 c.c. of *N*/10-sodium nitrite solution. After thirty to forty minutes, this is made up to 250 c.c. with water, and shaken well. This solution should be kept at 0° and protected from light. In carrying out the estimation, 5 grams of dry *H*-acid or 10 grams of press-cake are dissolved in 400 c.c. of water and sufficient sodium hydrogen carbonate to produce a clear solution, and made up to 500 c.c. with water. Twenty-five c.c. of this solution are placed in a 600 c.c. beaker, 200 c.c. of ice-water added, and 2 grams of sodium hydrogen carbonate. This solution is cooled in ice and the diazonium solution is added from a burette, jacketed with ice-water, with good agitation. When the titration is almost finished, about 2 grams of sodium carbonate are added to increase the rate of coupling, and 10—15 grams of salt are added near the end-point. The titration is continued to the point when a spot on filter-paper gives a faint purple ring when spotted with *H*-acid solution. The titration is complete if this purple colour develops again when tested after the solution has been left for five minutes. F. M. R.

Chlorohydrocarbons and Carbon Chlorides. II. The Knowledge of the Saturation Character of the Di-, Tri-, and Tetra-chloroethylenes. B. M. MARGOSCHES and RICHARD BARU (*J. pr. Chem.*, 1921, [ii], 103, 216—226).—The iodine numbers of $\alpha\alpha$ - and $\alpha\beta$ -dichloroethylene, trichloroethylene, and tetrachloroethylene are practically zero as determined by the Hübl, Wijs, or other "iodine solution" in which the active agent is iodine

monochloride, and therefore these solvents can be used safely as solvents for fats in the determination of their iodine number. This is in accordance with previous work on the influence of chlorine on the degree of unsaturation of a double bond.

W. O. K.

Detection of Coconut Oil in Butter. C. F. MUTTELET (*Compt. rend.*, 1922, 174, 220—223).—The method is based on the fact that pure butter contains cholesterol but not phytosterol, whilst the vegetable fat contains phytosterol, and the melting points of the acetates of these two sterols differ. The sterols in the butter are precipitated by the addition of 20 c.c. of a 1% alcoholic solution of digitonin to the fatty acids from 50 grams of the butter. The precipitate is collected and washed free from fatty acids by chloroform and ether, and then boiled with 2—4 c.c. of acetic anhydride. The acetate is twice crystallised from alcohol and its melting point determined. The acetate from pure butter has m. p. 113·6—114·2°. With 10% of coconut oil present in the butter the resulting acetate has m. p. 114·5° and the m. p. increases as the percentage of coconut oil in the butter increases.

W. G.

A Simple and Exact Method for the Direct Estimation of Acetaldehyde in the Presence of Acetone. WILHELM STEPP and ROBERT FRICKE (*Z. physiol. Chem.*, 1921, 116, 293—301).—To estimate the acetaldehyde, the solution is treated with an excess of alkaline-ammoniacal silver of known strength, the reduced silver is filtered off, and the unreduced silver is titrated in the acidified filtrate with *N*/10-ammonium thiocyanate, using ammonium iron alum as indicator. An accuracy of 0·1 mg. can be obtained. In order to estimate the acetone, the acetaldehyde is removed by boiling with silver oxide or with Fehling's solution, the residual liquid is distilled, and the acetone estimated in the distillate by the Messinger-Huppert method. If acetaldehyde has to be estimated at the same time, a known suspension of silver oxide is used and the residual silver oxide is dissolved in ammonia and estimated volumetrically.

S. S. Z.

Spectrochemical Reaction of Methylfurfuraldehyde and Hydroxymethylfurfuraldehyde Phloroglucides. TETSUTARO TADOKORO (*J. Coll. Agric. Hokkaido Imp. Univ.*, 1921, 10, 52—56; cf. Oshima and Tadokoro, *A.*, 1918, ii, 255).—The difference in the colour reactions of methylfurfuraldehyde and hydroxymethylfurfuraldehyde as phloroglucides was observed by means of the ultra-violet spectroscope. The colour reaction when the two aldehydes are allowed to react with phloroglucin and hydrochloric acid changes with the time and is complete in five minutes. At a half to two minutes after the beginning of the reaction, the two phloroglucides give almost the same absorption band in the visible spectrum ($\lambda\lambda$ 4200—5000 and $\lambda\lambda$ 4100—5000), but the methylfurfuraldehyde phloroglucide shows an absorption band in the ultra-violet at $\lambda\lambda$ 2400—3800. Five minutes after the beginning of the reaction, methylfurfuraldehyde phloroglucide gives an

absorption at $\lambda\lambda$ 4300—4800, whilst hydroxyfurfuraldehyde phloroglucide shows an absorption band at $\lambda\lambda$ 5000—5500. G. W. R.

A Supposed Method for the Quantitative Separation of "Saccharin" from *p*-Sulphaminobenzoic Acid. WALTHER HERZOG and J. KREIDL (*Österr. Chem. Ztg.*, 1921, 24, 165—166).—A method for the estimation of *p*-sulphaminobenzoic acid in commercial "saccharin" has been described by O. Beyer ("Kontrolle und Herstellung von Saccharin," p. 97), which consists in dissolving the material in a slight excess of ammonia, adding a 50% excess of acetic acid, and keeping for twelve hours. The para-acid is said to be completely precipitated under these conditions, whilst the more strongly acidic "saccharin" remains in solution as undecomposed ammonium salt. Experimental investigation of the method with known mixtures of the two pure substances showed, however, that the results were inaccurate to the extent of 2—3%. For example, in mixtures containing 5 and 25% respectively of para-acid, only 3.33 and 23.11% were found. G. F. M.

Apparatus for Use in Titrating Intermediates with Unstable Diazo-solutions. C. P. ATKINSON (*J. Soc. Dyers and Col.*, 1922, 38, 15—16).—The apparatus is intended for the estimation of intermediates for azo-dyestuff manufacture by titration with standard diazonium solutions.

An iron tripod about thirty inches high supports a circular tin trough containing a supply of ice-water, and inside the trough is a circular glass vessel, with an outlet through the centre of the trough, to contain the supply of diazonium solution. The burette, jacketed with the outer tube of a condenser through which the ice-water flows, is supported by a triangle attached to the three legs of the tripod. In one of the three legs of this support a funnel holder is fitted to receive the waste water as it flows from the jacket and conduct it to the sink. F. M. R.

Mercury or Water Ureometer for the Estimation of Urea in Urine or in Blood. RENÉ CLOGNE (*J. Pharm. Chim.*, 1922, vii, 25, 99—100).—The method of graduation in the usual form of ureometer is modified so that the graduated tube immediately below the inlet tap is very narrow and is marked in twentieths of a c.c. as far as the 2 c.c. graduation, and then in fifths as far as the 25 c.c. graduation. This has the advantage of enabling the same apparatus to be utilised for estimating the small amount of urea usually present in blood, as, if the same quantity is taken as of urine for an estimation, namely, 2 c.c., the reading of the volume of liberated gas will fall within the range of the 0.05 c.c. graduations, and an amount as small as 2.5 grams per litre can be accurately estimated. G. F. M.

Silicotungstic Acid Applied to the Estimation of Caffeine. A. AZADIAN (*Bull. Soc. chim. Belg.*, 1922, 31, 15—18).—Silicotungstic acid gives with caffeine, in the presence of 5% hydrochloric acid, a precipitate having the composition

$$12\text{W}\text{O}_3 \cdot \text{SiO}_2 \cdot 2\text{H}_2\text{O} \cdot 3\text{C}_8\text{H}_{10}\text{O}_2\text{N}_4 \cdot 6\text{H}_2\text{O}.$$

On ignition, it leaves a residue having the composition $12\text{WO}_3, \text{SiO}_2$, and the factor for converting the weight of this residue into weight of caffeine is 0.2646. For the estimation, a known volume of an extract containing caffeine is evaporated to a syrup, which is extracted with hot water, the extract being made up to 50 c.c. Sufficient hydrochloric acid is added to bring the acid concentration up to 5% and then a solution of silicotungstic acid is added. The mixture is boiled and then left for twenty-four hours; the precipitate is collected, dried, ignited, and weighed. W. G.

New Method for the Detection of Thymine. OSKAR BAUDISCH and TREAT B. JOHNSON (*Ber.*, 1922, 55, [B], 18—21).—The method depends on the conversion of thymine into carbamide, acetylcarbinol, and pyruvic acid, the latter being identified as indigotin. Preliminary experiments show that the action is not influenced by the presence of uracil, cytosine, or sugar.

A solution of sodium hydrogen carbonate in water is treated successively with aqueous solutions of thymine and ferrous sulphate; the mixture is well-shaken with air, which causes the gradual conversion of the white, ferrous hydrogen carbonate into ferric hydroxide. The latter is removed and the filtrate concentrated on the water-bath, when the original odourless solution which does not reduce Fehling's solution acquires a characteristic odour and strong reducing properties, probably owing to a Cannizzarro reaction resulting in the formation of acetylcarbinol and pyruvic acid. The presence of the former is conveniently established by distillation of the liquid and treatment of the distillate with *o*-aminobenzaldehyde; the solution is boiled until the odour of the latter disappears, cooled, acidified with hydrochloric acid, and made alkaline again with sodium hydrogen carbonate. The presence of 3-hydroxy-2-methylquinoline is shown by the blue fluorescence of the solution; the reaction is unusually sensitive. The residue from the distillation contains pyruvic acid, the presence of which is detected by the formation of indigotin after addition of *o*-nitrobenzaldehyde and sodium hydroxide. The dye is extracted with chloroform. The presence of 2—5 mg. of thymine is established readily by the formation of the blue chloroform solution.

H. W.

Application of Folin and Denis's Phosphotungstic Reaction to the Estimation of Uric Acid in Urine. THIÉRY (*J. Pharm. Chim.*, 1922, [vii], 25, 87—92).—All the methods for the estimation of uric acid in urine hitherto proposed have the disadvantage of requiring at least 100 c.c. of urine for each estimation. The author considers that the direct application of Folin's phosphotungstic reagent to urine without any preliminary treatment gives results sufficiently accurate for all ordinary purposes, at least in such cases where the volume of urine at disposal is small. The reagents required are Folin's reagent, a solution containing 120 grams of anhydrous sodium carbonate per litre, and a standard solution of uric acid containing 0.2 part per 1000, prepared by means of mono- and di-sodium phosphate. Into a 100 c.c. graduated tube 1 c.c.

of urine and 2 c.c. of the phosphotungstic reagent are introduced, and into a further series of tubes 1, 2, 3, 4, etc., c.c. of standard uric acid solution are placed, each with 2 c.c. of the reagent. The contents of all the tubes are made up to 40 c.c. with the sodium carbonate solution and, after keeping fifteen minutes, when the blue colour has reached its maximum intensity, the volume is made up to 100 c.c. with distilled water, and the uric acid is estimated colorimetrically by comparison with the standard colours in a Duboscq colorimeter. The results furnished by this method are very close to those given by the present author's volumetric silver method after treatment of the urine with potassium zinc ferrocyanide (A., 1921, ii, 527). G. F. M.

The Estimation of Purine Bases in Urine. H. STEUDEL and SUNG-SHENG CHOU (*Z. physiol. Chem.*, 1921, **116**, 223—225).—It is suggested that the filtrate from the second copper sulphate precipitation obtained in Krüger's method for the estimation of the purine bases in urine should after its decomposition with hydrogen sulphide be boiled with magnesium oxide in order to eliminate all traces of ammonia the nitrogen of which might otherwise be ascribed to the purine bases. S. S. Z.

Electrometric Titration of Azo-dyestuffs. D. O. JONES and H. R. LEE (*J. Ind. Eng. Chem.*, 1922, **14**, 46—48).—The difficulties encountered in the analysis of azo-dyestuffs with titanous chloride by previous methods are enumerated, and an analytical procedure is described whereby the electrometric method of following oxidimetric reactions (cf. A., 1919, ii, 471, 480) is used in conjunction with titanous chloride for the analysis of azo-dyestuffs and nitro-compounds.

A sample of the finely powdered dye (0.5—1.0 gram), sufficient to require 30—45 c.c. of $N/4$ -titanous chloride for reduction, is placed in a reaction flask with 25 c.c. of distilled water, and heated on a steam-bath for ten minutes to dissolve or soften the particles. Twenty-five c.c. of 40% sulphuric acid are added, the flask is stoppered, and a current of carbon dioxide is passed through for five minutes; 35—50 c.c. of titanous chloride, being at least 5 c.c. of $N/4$ -titanous chloride in excess of that required for reduction, are added, the mixture is boiled for five minutes, and cooled to 30°. In the back titration, the potentiometer is adjusted, and the voltages read for each addition of $N/20$ -ferric alum solution. The latter is added in 5 c.c. portions at first, gradually decreasing to 0.1 c.c. or less. When passing over the end-point, the poles are reversed in the usual manner, and the voltages read as the additions of ferric alum become larger. Volts are plotted as ordinates and c.c. of ferric alum solution as abscissæ, and the end-point is determined from the curve. For routine analysis, almost all azo-dyestuffs can be analysed with sufficient accuracy without reading the voltmeter or plotting a curve. The potentiometer is adjusted at the beginning of the back titration, until, on closing the circuit, the galvanometer shows no deflection. A permanent large swing of the galvanometer is obtained at the end-point. F. M. R.

Estimation of the Percentage of Fibrin in Blood and Plasma. H. C. GRAM (*J. Biol. Chem.*, 1921, 49, 279—295).—Citrated plasma is recalcified and heated at 35° for one and a half hours; the clot is then washed and dried, treated with alcohol and ether to remove lipoids, and weighed. The whole estimation, including the platelet count and the cell volume, is performed on 4.5 c.c. of blood.

E. S.

Chemical Blood Analysis. III. The Importance of the Ultra-filtration Method for the Analysis of Blood. M. RICHTER-QUITNER (*Biochem. Z.*, 1921, 124, 106—113).—Examples are given of the use of the Zsigmondy-Haen ultra-filtration apparatus for the estimation in blood of residual nitrogen, uric acid, chloride, sodium, free potassium, and calcium. The residual nitrogen, the uric acid, the chloride, and sodium are completely dialysable, but only a portion of the potassium, calcium, or dextrose is free and dialysable.

H. K.

Colorimetric Estimation of the Concentration of Hydrogen Ions in Very Small Quantities of Blood by Dialysis. J. LINDHARD (*Compt. rend. Trav. Lab. Carlsberg*, 1921, 14, No. 13, pp. 13).—A modification of the method of Dale and Evans (*A.*, 1921, i, 142), in which the required quantity of blood is reduced to three drops, so that the process can be applied repeatedly by finger pricks. Hirudin is used, and phenolsulphonephthalein as indicator. The dialysate is not titrated but compared with a colour scale of phosphate-indicator mixtures. The error for the dialysis of phosphate mixtures compared with a separate scale is about P_H 0.02, but compared with the original mixture the P_H agrees in the second place of decimals. Bicarbonate solutions gave by the electrometric method a P_H 0.2 to 0.3 higher than by the colorimetric, but this discrepancy the author attributes to loss of carbon dioxide in dialysis and inapplicability of the electrometric control (cf., however, Evans, *A.*, 1921, i, 904).

G. B.

Preparation of Colloidal Gold Solution for Testing Spinal Fluid. A. O. GETTLER and J. W. JACKSON (*Arch. Neurol. Psychiatry*, 1921, 6, 70—71).—To one litre of water (distilled in copper vessels from potassium permanganate) are added in turn 10 c.c. of 1% auric chloride solution, 7 c.c. of 2% potassium carbonate solution, and 0.5 c.c. of 1% oxalic acid solution. The liquid is heated until it boils, then removed from the flame and vigorously shaken, 0.2 to 0.3 c.c. of concentrated formaldehyde solution being simultaneously added, and the shaking continued for one minute; after three or four minutes, the colour usually commences to develop. If it does not do so, an additional 0.1 to 0.2 c.c. of formaldehyde solution is added, with agitation during and after the addition. A deep red colour should rapidly develop. **CHEMICAL ABSTRACTS.**

General and Physical Chemistry.

Relation between Atomic Volumes [of Elements in Combination] and Optical Refractivities. GERVAISE LE BAS (*Nature*, 1921, 108, 272—274).—A periodic relationship is known to exist between the atomic volumes of elements in combination, the differences between the atomic volumes of successive members of the same series being of the order of that of hydrogen. Similar serial and group relations are observed with the atomic refractivities, the serial differences again being of the order of the atomic refractivity of hydrogen. When the atomic volumes of carbon, nitrogen, oxygen, fluorine, silicon, phosphorus, sulphur, chlorine, arsenic, selenium, bromine, and iodine are plotted against their atomic refractivities, the points obtained lie approximately on a straight line. A. A. E.

Molecular Refraction of some Molten Salts and their Degree of Dissociation. G. MEYER and HECK (*Z. Elektrochem.*, 1922, 28, 21—23).—The molecular refraction of molten sodium hydroxide and nitrate and potassium hydroxide and nitrate has been measured at a series of temperatures between 320° and 440°. The method adopted was to allow a ray of light (sodium) to fall into a prism of the molten substance and be reflected from a metallic mirror inside the prism. The densities of the molten salts were determined for all the temperatures used. The mean molecular refraction over the whole temperature range is found to be : sodium nitrate 11.54, potassium nitrate 14.09, sodium hydroxide 5.37, and potassium hydroxide 7.71. These values, which are accurate to one unit in the second decimal place, the refractive index being accurate to one unit in the third decimal place, are in good agreement with the values calculated from the atomic refractions. Using the values 2.80 and 2.59, respectively, for the atomic refractions of the sodium atom and the sodium ion, it is calculated that molten sodium nitrate is dissociated to 61.7%. The influence of temperature on the molecular refraction of all the substances examined is very small. J. F. S.

Some Problems of the Mass Spectrograph. F. W. ASTON and R. H. FOWLER (*Phil. Mag.*, 1922, [vi], 43, 514—528).—A mathematical discussion of some points raised by the performance and further design of the mass spectrograph. J. R. P.

The Broadening of Spectral Lines. J. FRANCK (*Festschrift Kaiser Wilh. Ges. Ford. Wiss. Zehnjährigen Jubiläum*, 1921, 77—81).—A summary of the manner in which the broadening of spectral lines has been explained, or is capable of explanation, both by classical methods and in terms of the quantum theory. Radiation damping has been fairly adequately explained on the quantum

theory by Stern and Volmer; the Döppler effect may be explained by the influence of radiation pressure, whilst collision damping is considered to be due to the electron transitions produced by a collision, together with a sudden warping of the electron orbits.

CHEMICAL ABSTRACTS.

Intensity and Broadening of Spectrum Lines. CHR. FÜCHTBAUER and G. JOOS (*Physikal. Z.*, 1922, **23**, 73—80).—The form of a spectrum line broadened by another gas depends on the latter, not on the absorbing gas. The constant representing the broadening is, with hydrogen and nitrogen, proportional to the density. In the first pair of the principal series of caesium, the number representing the dispersing electrons on the classical theory is equal to the number of caesium atoms. For the mercury line 2537, the number of resonators is one forty-fifth of the number of atoms. The maximum of 2537 is less displaced towards the red by hydrogen than by carbon dioxide and nitrogen.

J. R. P.

Excitation Stages in Open Arc-light Spectra. I. Sodium, Potassium, Calcium, Strontium, Barium, and Magnesium. II. Silver, Bismuth, Cadmium, Zinc, Air, and Copper. B. E. MOORE (*Astrophys. J.*, 1921, **54**, 191—216, 246—272).—An investigation of the variation of the spectrum with current, using a potential of 2000 volts and currents of 0.02 to 1 ampere, distinguishes between five stages of excitation; the features of each are described, and a number of intensity-current curves are given. The relation of the results to previous work, including the Bohr theory, is briefly discussed.

A. A. E.

Influence of the Pressure of Foreign Gases on the D-Lines in Saturated Sodium Vapour. R. MINKOWSKI (*Physikal. Z.*, 1922, **23**, 69—73).—Measurements of the magnetic rotation of the D-lines of sodium vapour in presence of nitrogen under various pressures showed that the constant representing, on the classical theory, the number of dispersion electrons in unit volume is affected by pressure. The difference in the values of this constant determined by absorption measurements and extrapolated from the magnetic rotations is largely, perhaps entirely, to be ascribed to the influence of pressure. The broadening of the lines under the influence of pressure is largely confined to the side near the red. The method may be used in the determination of small partial pressures of monatomic vapours in mixtures.

J. R. P.

The Production of Enhanced Line Spectra. R. A. SAWYER and A. L. BECKER (*Science*, 1921, **54**, 305—306).—When calcium wires are exploded by the Anderson method (*Astrophys. J.*, 1920, **51**, 37), it is found that, as the size of the wires employed is decreased, the energy of the stimulus remaining the same, the intensity of the enhanced lines is increased, indicating a more complete ionisation of the calcium atoms. A fine asbestos fibre about 3 cm. in length was therefore saturated with an aqueous

solution of a calcium salt, and the charge of the high-tension condensers thrown across it. The fibre remained uninjured, and the calcium spectrum thus produced showed a striking enhancement of the spark lines of calcium over the arc lines, indicating that a large proportion of the emitting atoms were ionised. A table giving the relative intensities, under various conditions, of prominent spark- and arc-lines of calcium shows that the new source of light, provisionally called the "super-spark," yields a degree of ionisation comparable with, or perhaps in excess of, that existing in the high chromosphere of the sun and in the early (or hot) type *B* stars. Very minute amounts of material suffice for the production of intense spectra by this method, and practically only metallic lines are produced; the spectra of hydrogen, oxygen, or of the acid radicle of the salt used do not appear, and only the strongest air lines could be observed.

A. A. E.

The Evolution of the Spectrum of Magnesium under the Influence of Increasing Electric Fields. Applications to Astrophysics. A. DE GRAMONT and G. A. HEMSALECH (*Compt. rend.*, 1922, 174, 356—361).—In a discussion of previous work (A., 1921, ii, 611), it is shown that the spectral effects of intense electric fields are particularly marked in the initial stage of a luminous phenomenon (arc or spark) when the temperature and the electrical conductivity of the vapours are not very high. As the temperature increases, the emission, which depends on rapid falls in potential, diminishes or disappears. In consequence, the authors consider it dangerous to conclude that a star has a high temperature because the spark rays predominate in its spectrum.

W. G.

The Lines K_α of the Light Elements. V. DOLEJSEK (*Compt. rend.*, 1922, 174, 441—443).—The author has again measured the K_α lines of the elements from zinc to chlorine. He has found for these elements the lines α_3 and α_4 which Hjalmar had measured for the elements sulphur to sodium. They are inseparable above scandium. The line α_7 , according to Siegbahn's notation, was also found for the elements calcium to chlorine. In addition, a new satellite of α_1 of shorter wave-length has been found. It appears to be an emission band, and is denoted by α_1' . The line α_3 found by Duane and Stenström for tungsten could not be observed.

W. G.

The Complexity of the *K* Series of the Light Elements and its Theoretical Interpretation. A. DAUVILLIER (*Compt. rend.*, 1922, 174, 443—445).—Ten components of the *K* series for copper have previously been reported. Their wave-lengths are now recorded. Working under the same conditions (cf. A., 1921, ii, 669), the *K* series of molybdenum has been studied without observing any rays other than those found by Duane. The line α_3 found by Duane for tungsten does not exist either for copper or molybdenum. An attempt is made to show the relationship

between the author's measurements of the rays α_7 , α_3 , and α_4 of copper and those of Hjalmar for the elements calcium to sodium.

W. G.

Wave-lengths Longer than 5500 Å. in the Arc Spectra of Yttrium, Lanthanum, and Cerium, and the Preparation of Pure Rare Earth Elements. C. C. KIESS, B. S. HOPKINS, and H. C. KREMERS (*U.S. Bureau of Standards, Sci. Papers*, 1921, No. 421, 318—351).—Photographic determinations were made in the yellow, red, and infra-red regions of the arc spectra, as follows : yttrium, 170 lines to 7881·868 Å., lanthanum, 410 lines to 9078·99 Å., cerium, 1700 lines to 9024·68 Å. A description is given of the methods used in the purification of the substances employed, compounds of cerium, samarium, lanthanum, neodymium, and gadolinium being obtained from a sample of "Welsbach" sodium rare-earth sulphate, and those of yttrium, dysprosium, and erbium from gadolinite and xenotime. The cerium was removed from the "Welsbach" residues by precipitation with potassium bromate as basic ceric bromate; the other elements were then separated by way of the fractional crystallisation of the double magnesium nitrate. The yttrium-group material was freed from the elements of the cerium group, and then fractionally crystallised as bromate.

CHEMICAL ABSTRACTS.

The L Series of the X-Ray Spectrum. D. COSTER (*Compt. rend.*, 1922, 174, 378—379).—The author has again measured the spectra of a large number of elements (tantalum to rubidium), and in general the new results confirm the conclusions of previous work and in addition lend support to Bohr's theory of the structure of the atom (cf. this vol., ii, 277). Certain of the author's results are more or less opposed to the results and conclusions of Dauvillier (cf. A., 1921, ii, 421, 475, 699) and these divergences are summarised.

W. G.

The Infra-red Absorption Spectra of Alkali Hydroxides. G. E. GRANTHAM (*Physical Rev.*, 1921, 18, 340; cf. Howe and Gibson, *ibid.*, 1917, 10, 767).—When the absorption of solutions of sodium, potassium, and lithium hydroxides (of various concentrations) and of ammonium and caesium hydroxides (one solution each) was determined with reference to that of water, all the curves showed a broad, intense absorption band with a maximum at about $2\cdot29\mu$, except in the case of the ammonium hydroxide solution, for which the maximum was found to be at $2\cdot20\mu$. The absorption was found to be proportional to the concentration, but not to vary systematically with the atomic weight of the metal used. It is suggested that the band is due to dissociated hydroxide ions, although in the case of other hydroxide solutions such a band has not previously been observed. Maximum absorption of water was observed at $1\cdot48$ and $1\cdot98\mu$. The effect of dissolving an alkali hydroxide is to decrease the absorption of the water in the band at $1\cdot48\mu$ by an amount proportional to the concentration of the solution and to the atomic weight of the metal of the base.

A. A. E.

The Ultra-violet Absorption Spectra and the Optical Rotation of the Proteins of Blood Sera. S. JUDD LEWIS (*Proc. Roy. Soc.*, 1922, [B], 93, 178—194; cf. A., 1917, ii, 62).—Eleven sera, six from the horse and five from man, were used for the separation and purification by modifications of well-known methods of the albumin, euglobulin, and ψ -globulin. The optical rotation and the ultra-violet absorption spectra of the individual proteins were examined in detail. ψ -Globulin, euglobulin, and albumin from horse sera had specific rotations of -52° , -43° , and -57.4° , respectively, the corresponding values for human being -46° and -48° for the globulins and varying values for the albumin. The absorption curves are similar in form and character to that of serum, well-developed bands being best found at concentrations of 0.08% for albumin and 0.04% for the globulins. The absorption curve of ψ -globulin is the same for the horse and man and differs from euglobulin in the extinction coefficients but not in general form. For horse and human albumin, the curves are the same except for a constant ratio in their magnitudes, due possibly to physical or chemical association of an aggregate of little or no absorptive power. H. K.

The Fluorescence of Mercury Vapour. J. S. VAN DER LINGEN and R. W. WOOD (*Astrophys. J.*, 1921, 54, 149—160).—Since the fluorescent spectrum of mercury vapour cannot be excited in quiescent vapour, but only in vapour which is being distilled from the metal at a temperature not less than 150° , the active molecules are presumably not neutral monatomic molecules, but others, possibly diatomic, present only during distillation. The spectrum consists of lines at 2536, 2539, and 2346 Å., and four structureless bands with maxima at 2346, 2540, 3300, and 4850 Å. The complete spectrum is excited by light from a zinc spark, but single lines excite only a part. The relation of the spectrum to the exciting light is examined in some detail. A. A. E.

Hydrates of Boron Trioxide as a Constituent of Systems Capable of Strong Phosphorescence and containing Organic Compounds. ERICH TIEDE and PETER WULFF (*Ber.*, 1922, 55, [B], 588—597).—The phosphorescence of partly dehydrated boric acid (A., 1921, ii, 75) originates from traces of organic compounds, is removable, not by recrystallisation, ignition in a stream of oxygen, boiling with nitric acid, or fusion with potassium nitrate, but by boiling with fuming nitric acid for twenty-four hours, subsequently recrystallising three times from water specially purified, and finally igniting the upper portion of the filtered mass (which had not been in contact with filter-paper) in a platinum crucible. Phosphorescence could not be induced in such a product by any means other than by introduction of organic impurities. For this purpose, even those present in ordinary distilled water suffice. A number of fluorescent boric acid compounds were prepared by the respective additions to purified boric acid of fluorescein, its sodium salt (uranin), phenolphthalein, benzoic acid, quinone,

aniline, dimethylaniline, diphenylamine, phenol, quinol, naphthol, terephthalic acid, pyridine, and quinoline—in short, aromatic or heterocyclic compounds, but not aliphatic compounds, unless these (for example, ethylene glycol or sugar) undergo some decomposition—followed by fusion of the mixtures to transparent glasses, or by heating for two hours in the vacuum of a water-pump until a partly sintered mass is produced. Inorganic compounds, even those of uranium or the platinocyanides, with the exception of boron nitride (Tiede and Büscher, A., 1921, ii, 74) do not produce phosphorescent materials, nor can the boric acid be replaced by silicic or phosphoric acids. Hydration to orthoboric acid consequent on exposure of the products to air largely destroys the phosphorescence, as does also too intense dehydration. Phosphorescence is excited by daylight, but usually best by the ultra-violet light of the quartz lamp, and does not usually persist for more than two minutes. The intensity, however, is frequently much greater than that from the phosphorescent alkaline earths and sufficient momentarily to illuminate comparatively large spaces. Blue tones predominate, but reddish-yellow and green effects have been observed. Röntgen, cathode, and radium rays have no effect. The new products show only slight tendency towards thermoluminescence, down to the temperature of liquid air. In common with analogous cases, they are produced when the medium (boric acid) suffers contraction. They offer no support for the view that luminescence of organic compounds is dependent on their slow decomposition (Perrin, A., 1918, ii, 418; 1919, ii, 177). Pure anthracene, in spite of its fluorescence, causes no fluorescence in boric acid, but compounds like ethyl dihydrocollidinedicarboxylate, which are excited by radium rays, give a product in which fluorescence is produced by ultra-violet illumination, but is indifferent to radium. It is presumed that the phosphorescence emanates from definite organic derivatives of boric acid (cf. Dimroth and Faust, A., 1922, i, 155; Pictet, A., 1903, i, 601). Thus orange crystals result from evaporation of a solution of boric acid containing uranin (1 : 3000), but these, on fusion, become light green and simultaneously acquire capacity for phosphorescence.

J. K.

Action of Red and Infra-red Rays on the Phosphorescent Sulphides. MAURICE CURIE (*Compt. rend.*, 1922, **174**, 550—553).—As an explanation of the extinguishing action of the less refrangible part of the spectrum on certain phosphorescent substances, it is suggested that the action of the extinguishing rays will consist in rendering the medium a conductor by the detachment of electrons from the atoms of sulphur, in the case of the sulphides. In support of this theory, experiments are described in which an increase of conductivity was found for finely-powdered sulphur and certain sulphides when they were exposed to the action of red rays. This hypothesis also gives an explanation of the fact that the extinguishing action is not limited to the red and infra-red portions of the spectrum.

W. G.

Preparation of Phosphorescent Magnesium Sulphide. ERICH TIEDE and FRIEDRICH RICHTER (*Z. Elektrochem.*, 1922, 28, 20—21).—Strongly phosphorescent preparations of magnesium sulphide containing small quantities of bismuth, antimony, or manganese may be prepared. These substances exhibit a violet, golden-yellow, or dark red phosphorescence and an afterglow which in every way corresponds with that of the well-known preparations of the alkaline earth sulphide. Details of the method of preparation of phosphorescent magnesium sulphide are not given in the paper.
J. F. S.

Experimental Proof of some Theories of Natural Rotatory Power of Optically Active Solutions. PAUL WETTERFORS (*Z. Physik*, 1922, 8, 229—242).—With the object of testing the hypotheses of optical rotation put forward by Lorentz and Livens ("Theorie elektr. opt. Erscheinungen," 1906; A., 1913, ii, 543; 1914, ii, 830), the author has measured the specific rotation, $[\alpha]$, the refractive index, n , and the rotatory dispersion of solutions of camphor in ethyl alcohol, propyl alcohol, acetone, benzene, and toluene, and of α -bromocamphor in the three last-named solvents. The measurements were made with solutions of various concentrations from 10.02 to 49.98 grams per 100 c.c. of solution at temperatures from 11.1° to 18.0°. Four series of measurements were made in each case with light of wave-lengths 589 $\mu\mu$, 546 $\mu\mu$, 436 $\mu\mu$, and 710 $\mu\mu$, the first being obtained from a sodium lamp and the three latter from a quartz mercury lamp by the use of suitable light filters. The specific rotation of α -bromocamphor in acetone solution in various concentrations, c , is given by the equations $[\alpha]_{\lambda=710}=86.87+0.0625c$; $[\alpha]_{\lambda=589}=140.08+0.1043c$; $[\alpha]_{\lambda=546}=172.01+0.1389c$; $[\alpha]_{\lambda=436}=350.25+0.3561c$. It is shown in general the rotation increases with increase of temperature, since the concentration of a solution decreases with increasing temperature the specific rotation is also greater. An exception is found in the case of acetone solutions of α -bromocamphor; here the rotation decreases with increasing temperature, but the specific rotation is independent of temperature. The rotatory dispersion, defined as the ratio $[\alpha]_{\lambda} : [\alpha]_{589}$ where $[\alpha]_{\lambda}$ is the specific rotation for the wave-length λ and $[\alpha]_{589}$ the same quantity for $\lambda=589$, changes in general with the concentration. Bromocamphor shows an increase, camphor a decrease for the wave-lengths $\lambda=546\mu\mu$ and 436 $\mu\mu$ ($\lambda>589\mu\mu$), but for $\lambda=710\mu\mu$ ($\lambda<589$) an increase when the concentrations are greater. Solutions of camphor in acetone are exceptional, for here the rotatory dispersion is constant. According to Livens's theory, the expression $[\alpha]=rv^2/2 \cdot (n^2-1)[a(n^2-1)+1]$ must hold; this has been rewritten in the form $[\alpha]=ay_0(n^2-1)^2+y_0(n^2-1)$, where $y_0=rv^2/2$. On plotting the values of $[\alpha]/(n^2-1)$ against (n^2-1) , a straight line should be obtained. Treating the present experimental values in this way actually leads to straight lines for solutions of camphor in benzene and toluene, and for bromocamphor in benzene; in all other cases a straight line passes only approximately through the points. The values of a and y_0

have been evaluated by the method of least squares, and it is shown that the value of a is always negative as shown by Peacock (T., 1914, 105, 2782; 1915, 107, 1547). The numerical values of a are largest for bromocamphor in toluene and benzene and smallest for camphor in ethyl alcohol. The a value for camphor in toluene is the same for $\lambda=546\mu\mu$ and $\lambda=436\mu\mu$. It is shown, therefore, that the Lorentz-Livens hypothesis can only claim to give qualitative representation of the present material. According to Oseen's hypothesis, the expression $k=[\alpha]:(n^2+2)^2$ must be constant. The experimental data show that this is not the case, for in the case of solutions of camphor in benzene, toluene, and the alcohols, the k value increases with increasing concentration, but decreases with acetone solutions. Further, except in the case of solutions of camphor in propyl alcohol, the value of k is a linear function of the concentration.

J. F. S.

Photochemical Studies. XIV. Fundamental Photochemical Laws. J. PLOTNIKOW (*Z. wiss. Photochem.*, 1922, 21, 134—140).—A theoretical paper in which Einstein's photochemical equivalent law and the Grotthus-van't Hoff photochemical absorption law are critically examined. It is shown that Einstein's law leads to the following improbabilities. On calculating the energy absorbed, by one gram molecule, from light of various wave-lengths, it is found that the extreme ultra-violet, $100\mu\mu$, gives up 300,000 cal. and Röntgen rays 3×10^9 cal., which indicates that with decreasing wave-length light becomes much more active, until, with Röntgen rays, all molecules will be simply destroyed. Further, all atoms in all reactions in all solvents must react with the same velocity at all temperatures and must take up the same quantity of energy. It is shown that the Grotthus-van't Hoff law is diametrically opposed to the Einstein law. Nine classes of photolytic reactions are quoted in which the Einstein law in no way represents the facts, and two further classes are noted in which deviations from the law of -8% to 22% are observed. The photolysis of hydrogen iodide and hydrogen bromide alone give an approximate agreement, but here there is a divergence varying between -3.3% and $+4\%$. Weigert's recent paper (*Z. Physik*, 1921, 5, 421), which shows the applicability of the Einstein law and characterises the Grotthus-van't Hoff law as untenable, is criticised. It is shown that the reactions studied and the method of operation is much too complicated to make it possible to draw conclusions of the type drawn by Weigert from the results. In the present author's opinion, the Einstein law in its present form cannot be maintained.

J. F. S.

Photosynthesis and the Electronic Theory. II. HENRY H. DIXON and NIGEL G. BALL (*Sci. Proc. Roy. Dubl. Soc.*, 1922, 16, 435—441; cf. Dixon and Poole, A., 1920, ii, 343).—In continuation of previous work, the photo-electric properties of chlorophyll were investigated. The electronic theory of sensitisation is supported by experiments in which sensitised photographic plates were affected by light at a temperature of -185° , chemical

action being thereby precluded. Using collodion films, it was shown that sensitisation by chlorophyll is also effective even at the temperature of liquid air. Since only light of visible wavelengths is absorbed by chlorophyll, and since earlier experiments have shown that electrons are not expelled from chlorophyll, it follows that the light absorbed is wholly used in the displacement of electrons within the molecule, thereby making some atomic group or groups reactive. In photosynthesis, these groups may react with carbon dioxide and water, according to the scheme $\text{chlorophyll-}a + \text{CO}_2 = \text{chlorophyll-}b + \text{CH}_2\text{O}$ and $\text{chlorophyll-}b + \text{H}_2\text{O} = \text{chlorophyll-}a + \text{O}_2$. It is assumed that these reactions only take place during illumination and depend on its intensity. The velocity of the first reaction would only increase with intensity of illumination so long as carbon dioxide is in excess, whilst the velocity of the second reaction would only be indirectly influenced by this factor.

G. W. R.

Absorption of Röntgen Rays. K. A. WINGÅRDH (*Z. Physik*, 1922, 8, 363—376).—A method for the determination of the absorption coefficients of substances in solution for the homogeneous Röntgen rays of the K_α line of molybdenum is described. The atomic absorption coefficients are calculated, and it is shown that the logarithms of these when plotted against the logarithms of the ordinal numbers of the elements give two straight lines. Oxygen showed a marked deviation.

J. R. P.

Spark Lines in the Röntgen Spectrum. GREGOR WENTZEL (*Ann. Physik*, 1921, [iv], 66, 437—461).—It is assumed that the inner shells of the atom may under certain conditions undergo repeated ionisation. By the spontaneous transition of an atom from such a state to a state of equal level, Röntgen lines will be emitted which correspond with spark lines in the optical spectrum. It is shown that the emission can have only one fine-structure of known Röntgen lines, which is extended usually towards the region of short wave-lengths. All the lines which previously caused difficulty in the systematic arrangement of Röntgen spectra can be regarded as spark lines; in particular the hard $K\alpha$ satellites, from $K\alpha_3$ to α_6 can be derived from two combination relations, and the complex structure of the K -absorption edges observed by Fricke is also elucidated. The magnitudes of the frequency differences and their linear change with ordinal number are satisfactorily explained.

J. R. P.

X-Rays and Thermodynamic Equilibrium. L. DE BROGLIE (*J. Phys. Radium*, 1922, [vi], 3, 33—45).—The conditions of thermodynamic equilibrium between atoms, electrons, and radiation in an enclosure maintained at a temperature T are considered. The statistical equilibrium between the atoms is described by the law of Maxwell and Boltzmann, which may be applied to systems in which the total energy may assume only a certain number of discontinuous values. The equilibrium density of

radiation in the region of frequencies considered may be represented by Wien's law. The free electrons are considered as forming an ideal gas, with density proportional to $T^{\frac{3}{2}}$. The atomic coefficient of absorption of a substance is then shown to be given by $\mu = C\lambda^3 \sum_p n_p E_p^2$, where C is a constant, λ the wave-length, n_p the number of electrons in the level p (in the sense of Bohr's theory), and $E_p = h\nu_p$ the energy emitted when the atom passes from the ionised to the normal state. This result is compared with experiment.

J. R. P.

Stopping Power and Atomic Number. J. L. GLASSON (*Phil. Mag.*, 1922, [vi], 43, 477—481).—By a discussion of known results it is shown that the stopping power of an atom is proportional to $N^{\frac{1}{2}}$, where N is the atomic number, rather than to $A^{\frac{1}{2}}$, where A is the atomic weight, as previously supposed. This agrees with the higher stopping power of hydrogen as compared with helium

J. R. P.

Isotopy of the Radio-elements. M. L. NEUBURGER (*Nature*, 1921, 108, 180).—Meitner's (this vol., ii, 15) nucleus model of the radio-elements permits of the division of the radioactive isotopes into four classes: (1) those which possess only the same nuclear charge and the same arrangement of their outer electrons, for example, radium and mesothorium-I, (2) those which have in addition the same atomic weight and the same total number of nuclear "building stones," for example, ionium and uranium-Y, (3) those having the same number of each nuclear building stone, but a different arrangement of these in the atomic nucleus, for example, radium-D and actinium-B, (4) those possessing the same arrangement of building stones, and thus the same probability of disintegration, for example, radium-G and actinium-D. Such elements cannot at present be designated isotopes since there is no available means of distinguishing between them.

A. A. E.

The Ionium Content of Radium Residues. ELIZABETH RONA (*Ber.*, 1922, 55, [B], 294—301).—Ionium has been estimated in various radium residues according to the "indicator" method with the aid of uranium-X and the content has been compared with that of pitchblende. The final residues obtained during the extraction of radium in Austria still contain approximately 16% of the ionium present originally in the pitchblende. Thorough extraction of the latter with nitric acid on a laboratory scale gives insoluble residues which are almost completely free from ionium.

The importance of the estimation of ionium for that of proto-actinium in uranium minerals and for the inter-relationships of the actinium series is discussed.

* H. W.

Artificial Disintegration of the Elements. A Lecture delivered before the Chemical Society on February 9th, 1922. SIR ERNEST RUTHERFORD (*T.*, 1922, 121, 400—415).

Luminescence of Flame Ions in the Air Spark. H. GREINACHER (*Physikal. Z.*, 1922, 23, 65—69).—When a flame coloured

with a salt vapour is brought into a strong electric field, the ions may be caught on two platinum wires, outside the flame, on which they may afterwards be detected spectroscopically by heating. When a spark is passed between a wire in the flame and a brass sphere a few centimetres from the flame, or two wires similarly placed, the spectrum of the spark shows the lines of hydrogen and the metal ions. Lithium, sodium, potassium, rubidium, caesium, thallium, and strontium were examined. The metals move principally to the negative electrode, but partly to the positive. In the case of strontium, all the lines leave the flame, but in the other cases only a limited number. In spite of the strong field in the spark, only the lines with a small exciting potential (small $h\nu$) appear. The ions are partly produced in the flame and partly in the spark. An apparent change in wave-length occurs.

J. R. P.

Ionic Equilibria on Metallic Surfaces. H. VON EULER [with ARVID HEDELIUS and ZIMMERLUND] (*Z. Electrochem.*, 1922, **28**, 2—6).—An account is given of the sorption of gold, silver, and mercury ions on the surfaces of the metals. The method consisted in determining analytically the change in the concentration of solutions of silver nitrate and silver sulphate after they had been in contact with metallic silver and gold respectively. Similar experiments are described for solutions of mercuric chloride in contact with mercury. It is shown that 1 sq. metre of silver surface absorbs 0.35 mg. ion from silver nitrate and 0.5 mg. ion from potassium chloride solution, whilst the same area of gold absorbs 0.063 mg. ion from silver nitrate. A sq. metre of mercury surface absorbs 0.004 mg. ion from mercuric chloride solution. The results are discussed in connexion with the charge of an electric double layer.

J. F. S.

Evidence for the Existence of Homogeneous Groups of Large Ions. P. J. NOLAN (*Physical Rev.*, 1921, **18**, 185—198).—Using, as did Blackwood (*ibid.*, 1920, **16**, 85), the Zeleny method for the measurement of the mobilities of ions, instead of that previously employed by McClelland and Nolan (*Proc. Roy. Irish Acad.*, 1919, **35**, [A], 1, and previous papers), evidence has been adduced to confirm the author's original statement that when large ions are produced in air in various ways, the ions can be divided into a number of groups, each group having a definite mobility. It is considered that Blackwood's negative results were due to ions changing from group to group when the time spent in the measuring chamber is relatively long.

A. A. E.

The Scattering of Electrons by Nickel. C. DAVISSON and C. H. KUNSMAN (*Science*, 1921, **54**, 522—524).—When a nickel target is bombarded by a stream of electrons, there is observed, besides the emission of slowly moving secondary electrons characteristic of all metals, an appreciable emission of electrons of higher speed. After experimental and mathematical examination of

the phenomenon, it is suggested that the sharp deflexions experienced by these scattered electrons must result from their penetration into the atomic structure and their deflexion by the strong field which they encounter.

A. A. E.

Dielectric Constants of some Esters at Low Temperatures.

L. C. JACKSON (*Phil. Mag.*, 1922, [vi], 43, 481—489).—A method is described for the determination of the dielectric constants of solid and liquid substances by the use of triode valve circuits. The results, with a frequency of 4.7×10^5 per second, were, at the absolute temperatures indicated, as follows: methyl formate, 2.56 at 78.65° ; ethyl formate, 2.40 at 81.3° ; *n*-propyl formate, 2.39 at 79.5° ; *n*-butyl formate, 2.43 at 78.7° ; methyl acetate, 2.58 at 77.3° ; ethyl acetate, 2.48 at 79.15° ; *n*-propyl acetate, 2.42 at 80.9° ; *n*-butyl acetate, 2.41 at 77.6° . The values are smaller than those found at the ordinary temperature (5 to 9). The acid radicle apparently contributes but little to the value of the dielectric constant. The dielectric constant of *n*-butyl acetate increases linearly with decrease of temperature to the melting point at which there is a sudden fall, followed by a very gradual rise as the temperature is still further lowered.

J. R. P.

Conductivity of Concentrated Solutions of Sodium and Potassium in Liquid Ammonia.

CHARLES A. KRAUS and WALTER W. LUCASSE (*J. Amer. Chem. Soc.*, 1921, 43, 2529—2539).—The specific conductivity of solutions of sodium and potassium in liquid ammonia at its boiling point (-33.5°) has been measured for concentrations from 0.6*N* to saturated solutions. It is shown that as the concentration increases, the specific conductivity increases enormously, reaching the values 0.5047×10^4 and 0.4569×10^4 for saturated solutions of sodium and potassium respectively. The specific conductivity of solutions of sodium and potassium at the same equivalent concentration is very nearly the same. The equivalent conductivity of saturated solutions of sodium and potassium in liquid ammonia is of the same order of magnitude as that of such metals as strontium and iron. The solubility of sodium and potassium in liquid ammonia at its boiling point has been found to be 5.367 and 4.866 mols. of ammonia per atom of sodium and potassium respectively. The general result of the work is the furnishing of further evidence in support of the hypothesis that the conduction process in metals consists in a motion of negative carriers of sub-atomic dimensions, which carriers are identical for all metals.

J. F. S.

Potentials at the Junctions of Univalent Chloride Solutions.

DUNCAN A. MACINNES and YU LIANG YEH (*J. Amer. Chem. Soc.*, 1921, 43, 2563—2573).—Using a slightly modified form of Lamb and Larson's flowing junction apparatus (*A.*, 1920, ii, 347) the contact potentials of the type, $MCl||M'Cl$, in which *M* and *M'* are either hydrogen or an alkali metal (lithium, sodium, potassium, caesium, or ammonium) have been determined. The measurements were made at 25° and were for solutions of equal concentrations.

Tables of results of all possible combinations are given for the concentrations 0.1*N* and 0.01*N*, and the values compared with those calculated by Lewis and Sargent's formula (A., 1909, ii, 369). In all cases reproducible values were obtained, within a few hundredths of a millivolt. The Lewis and Sargent form of Planck's original equation $E_c = RT/F \log_e \Delta_c/\Delta_c'$ expresses the results for most of the cases studied and gives a qualitative agreement in all cases. The experimentally determined potentials can, however, be obtained from the differences between numbers which are characteristic for each ion at each concentration. This is equivalent to the expression given above, except that for the lithium and potassium ions, values of Δ which are slightly different from those obtained from conductivity measurements would have to be used.

J. F. S.

A Quick Acting Hydrogen Electrode. P. J. MOLONEY (*J. Physical Chem.*, 1921, 25, 758—761).—A new form of hydrogen electrode is described. The electrode consists of a short length of platinum wire sealed into the end of a glass tube and projecting 4 mm. from the end; the tube should have an internal diameter of about 4 mm. A loop of 1 mm. diameter glass rod is sealed to the end of the tube so that the platinum wire occupies the centre of the loop. A glass tube, of slightly greater diameter than the electrode tube, blown into the form of a bell at one end and having a bulb a little higher up which is pierced near the top by a small hole, fits over the electrode tube and is kept in position by a piece of rubber tube. The platinum wire is platinised in the usual way. The object of the electrode is to ensure a rapid equilibrium between the hydrogen in the platinum and the hydrogen ion in the solution. To use the electrode it is drawn up the tube until the wire and glass loop are in the middle of the bulb in the jacket tube and the whole is immersed in the liquid to be measured until the tube connecting the bell and bulb is filled, then the electrode is pushed down until the wire and loop are in the bell, hydrogen is admitted under the bell, and the whole adjusted so that the wire is surrounded by hydrogen whilst the lower end of the loop just dips into the main bulk of the liquid. In this way only a small volume of liquid, namely, that held in the loop, has to be saturated with hydrogen and be in equilibrium with the wire. Equilibrium values are rapidly reached with this electrode.

J. F. S.

Behaviour of a Non-attackable Electrode in the Process Leading to the Equilibrium $3\text{HNO}_3 = 2\text{NO} + \text{HNO}_2 + \text{H}_2\text{O}$. ALFONS KLEMENC (*Z. Elektrochem.*, 1922, 28, 55, 56); H. PICK (*ibid.*, 56—57).—Polemical; cf. A., 1920, ii, 409; 1921, ii, 297.

J. F. S.

Overvoltage on Electrodes and its Connexion with the Hydration of Ions. N. ISGARISCHEV and SOPHIE BERKMANN (*Z. Elektrochem.*, 1922, 28, 40—47).—The decomposition voltage of *N*- and *N*/10-solutions of sulphuric acid and the sulphates of rubidium, aluminium, potassium, ammonium, sodium, lithium,

magnesium, zinc, copper, manganese, and nickel has been determined and the values have been compared with the degree of dissociation, the surface tension, the molecular hydration and the ionic hydration. It is shown that there is only one break in the anode current-voltage curve, and this is attributed to the discharge of the SO_4'' ion. The results show that the surface tension is in no way related to the discharge potential of the anion (E); the degree of dissociation also stands in no direct relationship to the value of E ; in some cases it is found that an increase in the value of the degree of dissociation is accompanied by an increase of E . It is concluded, therefore, that the discharge voltage is dependent on the velocity of the anode process, that is, a lower potential will be required for decomposition the greater the velocity of the anode process. To investigate the cause of the difference in the velocity of the anode process in different cases, the authors have compared the value of E with the hydration of the molecules and the ions. It is shown that there is no connexion between the value of E and the number of molecules of water combined with the molecules of the electrolyte. The more hydrated the cations the smaller is the decomposition voltage found to be, and since the more hydrated the cation the less hydrated is the anion, it follows that the lower is the potential required to discharge it. The anode process is expressed by the scheme, (a) $\text{SO}_4'' + 2\oplus = \text{SO}_4$; (b) $\text{SO}_4 = \text{SO}_3 + \text{O}$; (c) $\text{Na}_2\text{SO}_4(\text{H}_2\text{O})_r \rightleftharpoons 2\text{Na}(\text{H}_2\text{O})_{66} + \text{SO}_4''(\text{H}_2\text{O})_m$; (d) $\text{SO}_4''(\text{H}_2\text{O})_m \rightleftharpoons \text{SO}_4'' + m\text{H}_2\text{O}$. The cause, therefore, of the retardation of the anode process is the insufficiently rapid formation of SO_4'' ions from the hydrate. It is shown that there is a definite periodicity between the value of E and the atomic weight of the cation.

J. F. S.

Influence of Colloids on Overvoltage. N. ISGARISCHEV and SOPHIE BERKMANN (*Z. Elektrochem.*, 1922, 28, 47—50).—The discharge potential of the hydrogen ion on copper, silver, and platinum cathodes has been measured in 2*N*-sulphuric acid containing various concentrations of gelatin (0—6%). It is shown that the nature of the cathode has no influence on the discharge potential. The value of this quantity at 18° rises to a maximum with increase in the gelatin concentration and then commences to fall. There is a corresponding increase in the anode discharge voltage with increase in the gelatin concentration. The change is explained as due to the formation of a complex between the hydrogen ion and the gelatin which is slowly decomposed at the electrode (see preceding abstract). The maximum potential represents the point where all the ions have combined with gelatin and a calculation based on an approximate value of the molecular weight of gelatin shows that a complex consisting of one hydrogen ion and 1173 molecules of gelatin is formed.

J. F. S.

Free Energy of Dilution and the Activities of the Ions of Potassium Bromide in Aqueous Solutions. J. N. PEARCE and HARRY B. HART (*J. Amer. Chem. Soc.*, 1921, 43, 2483—2492).—Measurements of the *E.M.F.* of cells of the type $\text{Ag}|\text{AgBr}, \text{KBr}(c)|$

KHg_x have been made for various concentrations ($c=2.0-0.001N$) at 25° , 30° , and 35° . The decrease of free energy and the decrease in heat content attending the reaction in these cells have been calculated. The free energy decrease accompanying the transfer of 1 mol. of potassium bromide from various concentrations (c) to $0.10N$ have also been calculated. The *E.M.F.* of concentration cells with ion transference of the type $\text{Ag}|\text{AgBr}, \text{KBr}(c_1)|\text{KBr}(c_2), \text{AgBr}|\text{Ag}$ and of concentration cells without ion transference of the types $\text{Ag}|\text{AgBr}, \text{KBr}(c_1)|\text{KHg}_x-\text{Hg}_x\text{K}|\text{KBr}(c_2), \text{AgBr}|\text{Ag}$ and $\text{KHg}_x|\text{KBr}(c_1), \text{AgBr}|\text{Ag}-\text{Ag}|\text{AgBr}, \text{KBr}(c_2)|\text{Hg}_x\text{K}$ have been determined and from these values the transport number of the potassium ion has been calculated. The results show that dilution has very little influence on the transport number of potassium and there is only a very gradual increase with increasing dilution, whilst the effect of temperature is practically negligible between 25° and 35° . The decrease of free energy and heat content accompanying the transference of 1 mol. of potassium bromide from a concentration c_1 to a concentration c_2 have been calculated. The activity coefficients of the bromide ion have been calculated and a comparison made of the activity coefficients of the chloride and bromide ions. For all concentrations up to $0.5N$, the activity coefficients of these two ions are practically equal. J. F. S.

Electro-chemical Behaviour of Metallic Conducting Compounds. G. TAMMANN (*Z. Elektrochem.*, 1922, 28, 36-40).—A discussion on the electromotive behaviour of compounds which exhibit a metallic conductivity. It is shown that of the five cobalt silicides, only those are attacked by acids which contain more than one atom of cobalt to one atom of silicon. In the case of the lead palladium intermetallic compounds, it is found that they have the palladium potential, but if there is an excess of lead over that required for the formation of a compound in the lead palladium alloy the mixture shows the lead potential. A number of other cases are cited and discussed. J. F. S.

Activity Coefficients and Colligative Properties of Electrolytes. HERBERT S. HARNED (*J. Amer. Chem. Soc.*, 1922, 44, 252-267).—A theoretical paper in which on the basis of the empirical equation $\log F_a' = \alpha'c - \beta'c^m$ (A., 1920, ii, 664) which connects the activity coefficient of an electrolyte with the molecular concentration at a given temperature, and Duhem's equation, equations are derived for calculating the vapour pressures of solutions at this temperature. The validity of the above equation has been rigorously tested, and it is found to hold for all electrolytes over the whole range where data were examined with the exception of sulphuric and hydrochloric acids. In these cases, it holds up to a concentration $3M$. The activity coefficients of solutions of hydrochloric acid, sodium chloride, and potassium chloride have been investigated and the values of the parameters of the above equation for these electrolytes and others have been collected and tabulated, from which the vapour pressures and osmotic pressures of their solutions may be calculated. It is suggested that the

equation and method employed here, although empirical and an approximation, will prove useful in testing the consistency and accuracy of activity data and also will be of service in organising the colligative data of concentrated solutions. J. F. S.

Ghosh's Theory of Electrolytic Solutions. CHARLES A. KRAUS (*J. Amer. Chem. Soc.*, 1921, 43, 2514—2528).—A theoretical paper in which the hypothesis of electrolytic solutions put forward by Ghosh (*T.*, 1918, 113, 449, 627, 777, 790) is discussed. Ghosh's equation expressing the conductivity of solutions of electrolytes as a function of the concentration is compared with the experimental values in aqueous and non-aqueous solutions. On plotting the values of $\log A$ against those of the cube root of the concentration, the experimental points lie on a curve which is concave toward the axis of concentration. If Ghosh's equation were applicable, the points should lie on a straight line. The deviation of the points from a linear relation is consistent throughout, and cannot be accounted for by errors in the original observations. It is held that in its present form Ghosh's hypothesis does not satisfactorily account for the properties of solutions of electrolytes. The postulates and assumptions on which the hypothesis is based are discussed and several inconsistencies indicated. The use of Maxwell's law for the distribution of velocities and Clausius's virial theorem to systems of charged particles is open to serious objections. J. F. S.

Distribution of Thermal Energy in the Quadrivalent Chlorides of Carbon, Silicon, Titanium, and Tin. WENDELL M. LATIMER (*J. Amer. Chem. Soc.*, 1922, 44, 90—97).—The specific heats of carbon, silicon, titanium, and tin tetrachlorides have been measured from the temperature of liquid air to 298° Abs., and, in addition, the specific heat of carbon tetrachloride was measured at still lower temperatures down to 39.1° Abs., in order to determine the whole specific heat curve for these compounds. The following values of $C_p M/5$ are recorded: Carbon tetrachloride, solid I, 39.1°, 0.60; 40.7°, 0.65; 63.8°, 2.05; 79.6°, 2.81; 91.0°, 3.40; 99.5°, 3.60; 199.5°, 5.84; 208.0°, 5.90; solid II, 229.4°, 6.20; 235.6°, 6.25; 238.0°, 6.77; liquid, 253.8°, 6.40; 265.1°, 6.30; and 290°, 6.34. Silicon tetrachloride, solid, 77.4°, 3.60; 86.6°, 3.83; 94.8°, 4.14; 131.3°, 4.80; 168.6°, 5.68; 181.0°, 5.95; liquid, 208.8°, 6.74; 294.3°, 6.94. Titanium tetrachloride, solid, 86.7°, 4.26; 92.8°, 4.40; 194.6°, 6.19; 231.8°, 6.31; 247.7°, 6.21; liquid, 251.6°, 7.26; 294.3°, 7.32. Tin tetrachloride, solid, 89.0°, 4.59; 95.9°, 4.71; 161.2°, 5.89; 200.7°, 6.33; 227.3°, 6.62; liquid, 266.1°, 7.56; 294.0°, 7.86. All temperatures are in absolute degrees. The heat of transition of solid I to solid II in the case of carbon tetrachloride at 224.6° Abs. is 7.10—7.19 cal./gram. The heat of fusion has been determined as follows: carbon tetrachloride, 4.1—4.22 cal./gram at 249°; silicon tetrachloride, 10.85 cal./gram at 203.3°; titanium tetrachloride, 11.77 cal./gram at 248.0°, and tin tetrachloride 8.38—8.42 cal./gram at 239.9°. The entropy, S_{298} , of each of the compounds at 298° Abs. and the entropy change

in the formation, ΔS_{298} , of each compound from its elements have been calculated. The following values are recorded: carbon tetrachloride, S_{298} , 49.06, ΔS_{298} , -50.84; silicon tetrachloride, S_{298} , 56.43, ΔS_{298} , -48.27; titanium tetrachloride, S_{298} , 59.51, ΔS_{298} , -47.09; tin tetrachloride, S_{298} , 61.84, ΔS_{298} , -49.66.

J. F. S.

Relation between Statistical Mechanics and Thermodynamics. RICHARD C. TOLMAN (*J. Amer. Chem. Soc.*, 1922, 44, 75—90).—A theoretical paper in which the relationship between statistical mechanics and thermodynamics is investigated by a method which is based, like that of Boltzmann and Planck, on a consideration of the arrangement of the elements or molecules of a system. Instead, however, of arbitrarily defining entropy by the equation $S = k \log W$ as do Planck and Boltzmann, an attempt is made to discover a statistical mechanical quantity which has the same derivatives with respect to energy, volume, and number of molecules as does the thermodynamic quantity entropy. In this way, an expression for entropy is developed which differs from that of Planck by an additive term; it is further shown that the addition of such a term is necessary in order that the derivative of the entropy of the system with respect to number of molecules shall have its correct value. The equation thus developed has the form $S = (N\epsilon - N\psi)/T$, in which N is the number of molecules, ϵ has a significance defined below, and T is the temperature. In connexion with the derivation of the above expression, it is shown that the quantity ϵ occurring in the Maxwell-Boltzmann distribution law $dN = Ce^{-\epsilon/kT} dq_1 \dots dp_n$ must be taken, not as the energy of a molecule having co-ordinates and momenta falling in the region $dq_1 \dots dp_n$, but rather as the rate of increase in the total energy of the system per molecule added in the region $dq_1 \dots dp_n$, when the system is in its configuration of maximum probability. These two quantities are the same for dilute gases, but for concentrated systems may be quite different. Hence the new methods introduced may be of considerable importance in dealing with the properties of imperfect gas or rather concentrated systems. It is pointed out that the equations developed which connect the statistical mechanical quantities with thermodynamic quantities do not agree with the equations of Marcelin and Adams (*A.*, 1915, ii, 328; 1921, ii, 628).

J. F. S.

Relation between Entropy and Probability. Integration of the Entropy Equation. GEORGE A. LINHART (*J. Amer. Chem. Soc.*, 1922, 44, 140—142).—Two equations are deduced which express the change of specific heat of an element or compound with the entropy. The equations are based on the assumptions that the change depends on the probability of the "randomness" of the individual particles, the certainty that particles will have a fixed position at the point of zero kinetic energy, and the zero probability that this condition persists at high temperatures. The equations deduced are tested by means of recently published work on the specific heat of copper when a good agreement between the

observed and calculated values of the specific heat over the range 14.51—390.0° Abs. is found. J. F. S.

Latent Heats of Fusion. I. Benzophenone, Phenol, and Sulphur. (MRS.) K. STRATTON and J. R. PARTINGTON (*Phil. Mag.*, 1922, [vi], 43, 436—446).—The following values of the latent heats of fusion, in gram-calories per gram, were measured at the melting points by electrical heating: benzophenone, 21.70; phenol, 29.06; sulphur (monoclinic), 8.85. In a theoretical discussion, it is suggested that intramolecular vibrations as well as rotations are indicated in the case of some liquid molecules.

J. R. P.

The Freezing Points of Pure Organic Liquids as Thermometric Constants at Temperatures below 0°. JEAN TIMMERMANS, (MLLE) H. VAN DER HORST, and H. KAMERLINGH ONNES (*Compt. rend.*, 1922, 174, 365—367).—The following organic liquids have been carefully purified and their boiling points, densities, and freezing points accurately determined, so that these constants may be used as references by working with specimens kept in sealed tubes. All boiling points are given at 760 mm. pressure, and f. p. is the freezing point obtained under conditions where the thermodynamic equilibrium between the liquid and crystalline phases can be maintained for a long time to within 0.01°.

Carbon tetrachloride, b. p. 76.75°; d_4^{20} 1.63255; f. p. -22.9°; chlorobenzene, b. p. 132.00°; d_4^{20} 1.12795; f. p. -45.2°; chloroform, b. p. 61.2°; d_4^{20} 1.52635; f. p. -63.5°; ethyl acetate, b. p. 77.15°; d_4^{20} 0.92450; f. p. -83.6°; toluene, b. p. 110.70°; d_4^{20} 0.88445; f. p. -95.1°; carbon disulphide, b. p. 46.25°; d_4^{20} 1.29270; f. p. -111.6°; ethyl ether (stable form), b. p. 34.60°; d_4^{20} 0.73625; f. p. -116.3°; (unstable form), f. p. -123.3°; methylcyclohexane, b. p. 100.3°; d_4^{20} 0.78640; f. p. -126.4°; isopentane, b. p. 27.95°; d_4^{20} 0.63950, f. p. -159.6°.

W. G.

Hydration of Ions. S. ASCHKENASI (*Z. Elektrochem.*, 1922, 28, 58—59).—A theoretical paper in which Fajans's treatment of the hydration of ions (*Naturwiss.*, 1921, 729) is discussed and a possible connexion with the known irregularities in the vapour pressure curves of aqueous solutions of electrolytes suggested.

J. F. S.

A Receiver for Fractionation in a Current of Gas or under Reduced Pressure. T. S. WHEELER and E. W. BLAIR (*J. Soc. Chem. Ind.*, 1922, 41, 59—60T).—The device consists essentially of an upper and a lower receiver connected together through a two-way tap which makes the connexion alternatively with the top or bottom of the upper receiver. In the former case, the vacuum is maintained in, or the gas is passed through, the upper and lower receivers, whilst the distillate collects in the upper receiver. When it is desired to collect a new fraction, the two-way tap is turned into the second position and the fraction which has collected in the upper receiver passes into the lower one, which is then detached, another put in its place, and evacuated (if neces-

sary), and the distillation continued with the two-way tap in the first position as before. If it is desired to keep the distillate out of contact with air, the lower receiver is provided with inlet and outlet taps, which are turned off before detaching it from the apparatus, and the liquid thus remains enclosed in the inert atmosphere in which it was distilled.

G. F. M.

Comparison of the Linking of Carbon Atoms in Graphite and in Aromatic Hydrocarbons. J. P. WIBAUT (*Rec. trav. chim.*, 1922, 41, 96—102).—A theoretical paper based on the work of Fajans (A., 1920, ii, 354), and criticising the theory put forward by von Steiger (A., 1920, ii, 355) that the linking of the carbon atoms in aromatic hydrocarbons is, from the point of view of energy, equivalent to the linking in graphite.


H. J. E.

Atomic Linkings in the Carbon Compounds. I. and II. HANS BEUTLER (*Z. anorg. Chem.*, 1921, 120, 24—30, 31—47).—I. Assumptions that periodic oscillations of melting points and boiling points in homologous series are due to heteropolar linking of the carbon atoms in the chain (cf. Cuy, A., 1921, ii, 429), are shown to be untenable, on account of the non-dissociation of the carbon compounds. The heats of combustion also vary in an oscillatory manner in homologous series, and this may be explained by considering the lattice-energy of the crystal structure. Compounds with an even number of carbon atoms probably have a more highly symmetrical lattice than those with an odd number, and as a result they have higher melting points and lower heats of combustion. Even in liquids, a kind of potential lattice structure probably exists, and influences the periodically varying physical properties. As the critical temperature is approached, the influence of the lattice is weakened by the strong thermal forces and the differences between odd and even members of the series disappear.

The electron structure of stable ions such as the Cl^- or SO_4^{2-} ion is discussed, and it is shown that ion formation is due to the splitting of a compound into parts each of which has an electron structure similar to that of a noble gas, with an outer shell of eight electrons. (This is identical with the Lewis-Langmuir theory, to which, however, no reference is made.) The structure of a non-dissociating substance of the type of $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$ is probably such that the co-ordinated groups are attached tetrahedrally to the central atom; in the electron system, the central atom is surrounded by eight electrons forming the corners of a cube, four edges of which are common each to one edge of the electron cubes of one of the co-ordinated groups. The possibility of two isomerides is at once apparent in this arrangement.

II. In the preceding section it was shown that isomerism is possible in the compound of the formula $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$; this is due to the fact that free rotation about the central Pt atom is inhibited probably by directing forces from the inner layers of electrons in the Pt atom. In carbon compounds $\text{CR}_1\text{R}_2\text{R}_3\text{R}_4$ free rotation is possible, since only two electrons are contained within the shell of eight electrons and restraining forces are absent. The electron

structures of methane, its halogen substitution products, ethylene, acetylene, and aromatic hydrocarbons are discussed. The methane molecule is represented by a central carbon nucleus surrounded by four electrons and four hydrogen nuclei arranged tetrahedrally at the eight corners of a cube with the other four electrons between the



carbon and hydrogen nuclei. In CH_3Cl there are fourteen electrons, two of which are common to the eight surrounding the carbon atom and the eight surrounding the chlorine atom. Diagrammatically the electron structure can be represented thus, where the electrons of the outer shell are represented by small circles. In ethylene it is supposed that the eight electrons

which are not situated directly between the hydrogen and carbon nuclei, revolve in an elliptical path of which the major axis is parallel to the C—C axis. The observed paramagnetism attributed to the double bond confirms this view. The electron structural formula for benzene shows three electrons between each pair of adjacent carbon atoms. In naphthalene, three electrons are shown between each pair of adjacent carbon atoms except between the carbons common to both rings, where there are only two. Reduction to 1:4-dihydronaphthalene, however, converts one ring into a true benzene ring, and a true ethylenic bond appears between the carbon atoms 2 and 3 in the other ring.

E. H. R.

A Method for Measuring Vapour Densities for the Determination of Atomic Weights. A. MAGNUS and E. SCHMID (*Z. anorg. Chem.*, 1921, 120, 232—240).—An apparatus is described for the measurement of vapour densities, the distinguishing feature of which is the total enclosure of the manometer and measuring vessel in a double-walled chamber which can be kept constant at the temperature of boiling water. The apparatus was used for determining the relation between the pressure p and pv/mT , where v is the volume, m the weight of the vapour, and T the absolute temperature for benzene and chloroform. Contrary to the results of Ramsay and Steele, the relation was found to be truly linear. The molecular weights calculated from the results are 78.096 for benzene and 119.380 for chloroform, whilst the atomic weight of chlorine calculated from these is 35.455, showing that the apparatus gives results of a high degree of accuracy.

E. H. R.

Regularities in the Molecular Volumes of Inorganic Compounds. FR. A. HENGLEIN (*Z. anorg. Chem.*, 1921, 120, 77—84).—It was recently shown by Biltz (A., 1921, ii, 437) that there is a linear relationship between the molecular volumes of the halogens and of their compounds. This is now shown to be a particular case of a more general law that the molecular volumes of certain groups of inorganic compounds of like crystalline form bear a linear relation to constant magnitudes characteristic of their kations or anions. For the halogens, these magnitudes x may be expressed by the constants $F' : 0.8525$, $\text{Cl}' : 1.000$, $\text{Br}' : 1.066$, $\text{I}' : 1.179$, and

the magnitudes y for the alkali metals, $\text{Li}^+ : 1.000$, $\text{Na}^+ : 1.428$, $\text{K}^+ : 2.108$, $\text{Rb}^+ : 2.478$. The molecular volumes of the haloids of the alkali metals can be calculated from the general equation $\text{mol. vol.} = 14.67x \cdot y + 0.73x + 56.65y - 51.60$. Cesium is abnormal in this series, its constant being smaller than that of rubidium although in other series, for example in the sulphates, selenates, manganates, and chlorates, it is normal. The cubic haloids of copper, silver, and thallium form a good series in which the following constants appear, $\text{Cu}^+ : 1.00$, $\text{Ag}^+ : 0.920$, $\text{Tl}^+ : 1.40$. Hexagonal silver iodide does not fit in with the other cubic substances. The same law is shown to hold for numerous other isomorphous series containing metals of the second and fourth groups. The following density determinations were made: lithium fluoride, $d^{20}_4 2.597$; sodium fluoride, $d^{20}_4 2.726$; potassium fluoride, $d^{20}_4 2.369$.
E. H. R.

The Variation of the Mechanical Properties of Metals and Alloys at Low Temperatures. LÉON GUILLET and JEAN CURNOT (*Compt. rend.*, 1922, 174, 384—386).—The hardness and resilience of a number of metals and alloys have been measured at 20° , -20° , -80° , and -190° . The results show that there is, in general, an increase in the hardness with cooling, there being a fairly big jump from the value at -80° to that at -190° . Fragility at low temperatures is a characteristic of ferrite, the rapidity of the fall in resilience as a function of the temperature being greater as the ferrite content is higher. Nickel and copper, on the other hand, do not lead to fragility and aluminium, if present to any great extent, tends to produce a slight increase in the resilience. Pure austenite sufficiently rich in nickel does not show fragility at low temperatures. Special perlitic steels containing nickel have a high fragility in liquid air, but the addition of nickel retards the lowering of the fragility with decrease in temperature.

Measurements of hardness and resilience after a specimen has been kept for sixteen hours at -190° and then for twenty-four hours at $+20^\circ$ give evidence of a return to normal properties at the ordinary temperature.
W. G.

Surface Tension of Corresponding States. RICHARD LORENZ and W. HERZ (*Z. anorg. Chem.*, 1922, 120, 320—328).—The expression $\gamma_s/\gamma_e = (1 - T_s/T_k)/(1 - T_e/T_k) \cdot (V_e/V_k)^{2/3}$ is deduced. T_s/T_k and T_e/T_k have been shown by Lorenz (A., 1916, ii, 311) to be practically constant. According to the best average values, $T_s/T_k = 0.64$, $T_e/T_k = 0.44$, $(V_e/V_s) = 0.855$, and therefore $\gamma_s/\gamma_e = 0.578$. This was tested on results obtained by Jaeger (A., 1917, ii, 33) and figures are given for 89 organic compounds. The average value for γ_s/γ_e was found to be 0.52, 75% of compounds agreeing very closely. Twenty-one inorganic compounds of known boiling points gave an average for $\gamma_s/\gamma_e = 0.58$. Assuming $\gamma_s/\gamma_e = 0.578$, the authors have calculated the boiling points of several salts. The higher the critical temperature the further does γ_s/γ_e deviate from the mean.
W. T.

Surface Tension of Aqueous Solutions of Night-blue. R. DE IZAGUIRRE (*Anal. Fis. Quím.*, 1922, 20, 40—55; cf. Traube, A., 1912, ii, 740, 741).—The surface tension of aqueous solutions of night-blue of varying concentrations and with the addition of different electrolytes was investigated by means of the stalagmometer. The increase of surface tension with concentration is fairly rapid at first, but then slows down. In the experiments with addition of electrolytes, it was found that an abrupt fall in surface tension to that for pure water took place at the point where coagulation of the solution occurred. The coagulating power of various ions can thus be studied by means of the stalagmometer. Other physical properties, including the aspect under the ultramicroscope, do not show this abrupt change at the point of coagulation. Solutions purified by dialysis give results analogous to those obtained with unpurified material, but the stability of the solutions obtained is less, since smaller concentrations of salts are necessary for coagulation.
G. W. R.

Theory of Adsorption Processes. A. EUCKEN (*Z. Elektrochem.*, 1922, 28, 6—16).—A theoretical paper in which the process of physical absorption is considered. By physical adsorption is understood that adsorption, for example, of argon by charcoal, where it is unlikely that the adsorbed molecules are held by the attractions of chemical forces. It is shown that the adsorption forces are similar to the forces which are responsible for condensation phenomena. They are practically independent of the temperature and are operative over a relatively small distance which is of the same order as the molecular radius.
J. F. S.

Adsorption of Ammonia by Silica Gel. L. Y. DAVIDHEISER and W. A. PATRICK (*J. Amer. Chem. Soc.*, 1922, 44, 1—8).—A continuation of previous work (A., 1920, ii, 417). The adsorption of ammonia by silica gels containing respectively 4.98% and 0.33% of water has been determined for a series of pressures at 0°, 30°, 40°, and 100°. A smaller amount of ammonia is adsorbed by the less hydrated gel than by the more hydrated compound, although the less hydrated gel is still capable of adsorbing large amounts of ammonia. It is shown that the adsorption of ammonia may be satisfactorily explained on the basis of capillary condensation, provided corrections are made for the amount of gas which dissolves in the water.
J. F. S.

Adsorption by Precipitates. V. Adsorption during the Precipitation of Colloids by Mixtures of Electrolytes. HARRY B. WEISER (*J. Physical Chem.*, 1921, 25, 665—683; cf. A., 1919, ii, 269; 1920, ii, 228; 1921, ii, 625).—The precipitation of colloidal ferric hydroxide and colloidal arsenious sulphide by the simultaneous addition of pairs of electrolytes, and the adsorption of ions during the precipitation of the colloids by mixtures of electrolytes, have been investigated. The precipitation experiments were carried out with pairs of the electrolytes, potassium chloride,

barium chloride, strontium chloride, potassium oxalate, potassium sulphate, and potassium bromate. The adsorption of the oxalate ion in the presence of the chloride and sulphate ion respectively and the barium ion in the presence of the potassium ion was investigated. It is shown that the precipitating action of mixtures of pairs of electrolytes is approximately additive if the precipitating power of each is of the same order of magnitude, but may be far from additive if the electrolytes have widely varying precipitating power. In the latter case, the action of the electrolytes is antagonistic in the sense that relatively more of each is necessary than if the other were absent. In the simultaneous adsorption by solids from mixtures of two electrolytes having no common ion, the most strongly adsorbed kation and anion are taken up most and the other pair least readily; from mixtures having one common ion the oppositely charged ions are each adsorbed less than if the other were absent, but the most readily adsorbed ion is displaced the least. Factors which influence the precipitating action of mixtures of electrolytes are the effect of the presence of each precipitating ion on the adsorption of the other and the stabilising action of the ions with the same charge as the colloid. The second factor is of minor importance with mixtures of electrolytes that have similar precipitation values and have stabilising ions in common. The precipitating action of mixtures of electrolytes with similar precipitating power is additive, since the adsorption of the precipitating ions is similar and consequently the adsorption of each is affected but slightly by the presence of the other within the limits of the precipitation concentration. With mixtures of electrolytes of widely varying precipitating power, the precipitating action may be approximately additive when the influence of both factors, mentioned above, is small, but may be far from additive when the influence of both factors is large. These extreme cases are realised experimentally.

J. F. S.

Use of the Water Interferometer as a Pressure Gauge.
L. VAN DOREN, H. K. PARKER, and P. LOTZ (*J. Amer. Chem. Soc.*, 1921, **43**, 2497—2501).—The Zeiss water interferometer can be modified so as to serve as a pressure gauge. The modification consists in replacing the two water cells by two brass cells fitted with stout plate glass windows at the ends. One cell is filled with water at atmospheric pressure and the other with water under pressure. On calibrating the instrument, it was found that pressures up to sixty atmospheres could be measured. The range of the instrument may be increased by inserting a thin glass plate in the path of the beam passing through the non-pressure chamber of such a thickness that it will compensate for a pressure of sixty atmospheres in the other chamber. Thus by means of a series of glass plates of different thickness the range of the instrument can be extended to almost any pressure. In the calibration of the instrument for pressures up to 270 atmospheres at 20° and 30° it is found that at the higher temperature the effect of increased pressure on the refractive index diminishes.

J. F. S.

Osmotic Pressures of Concentrated Solutions of Sucrose as Determined by the Water Interferometer. PAUL LOTZ and J. C. W. FRAZER (*J. Amer. Chem. Soc.*, 1921, **43**, 2501—2507).—The authors describe a modification of the method used by Frazer and Myrick for measuring the osmotic pressure (*A.*, 1916, ii, 603). The principal feature of the new apparatus is the use of the water interferometer (preceding abstract) for determining the magnitude of the pressures developed. The apparatus is a decided improvement on that of Frazer and Myrick, being more rapid in action and more trustworthy. Pressures up to 273 atmospheres have been measured, but the instrument has been calibrated up to 330 atmospheres. Pressure measurements are recorded for sucrose solutions of concentrations up to saturated solutions at 30° and 55·7°. Attention is directed to the results of Woods as calculated for vapour-pressure measurements, and it is shown that his results (*Trans. Faraday Soc.*, 1915, **11**, 29) showing a decrease in osmotic pressure with rise in temperature in concentrated solutions are relatively correct, although his actual values for the osmotic pressure appear to be high when compared with these direct measurements. J. F. S.

System: Ammonia-Water as a Basis for a Theory of the Solution of Gases in Liquids. BENJAMIN S. NEUHAUSEN and W. A. PATRICK (*J. Physical Chem.*, 1921, **25**, 693—720).—A static method has been developed for measuring the partial pressure of a component of a liquid mixture which has a very small partial pressure compared with that of the second component. This method has been used to determine the partial pressures of water and ammonia solutions at 0°, 20°, and 40° at partial pressures of ammonia varying from 1000 mm. to 4000 mm. The measurements in the case of ammonia are made to 2—4 mm., and in the case of water to 0·08 mm. The solubility of ammonia in water was determined at 0°, 20°, and 40° at pressures from 750 mm. to 3600 mm., and the densities of the saturated solutions were also measured. A theory of the nature of solutions of gases in liquids first advanced by Graham has been amplified, and solutions of various gases in liquids classified on the basis of some of the physical and chemical properties of the gas. The formula $V = K(P\sigma/P_0)^{1/n}$ has been found to represent the solubility of ammonia, hydrogen chloride, sulphur dioxide, and carbon dioxide in water at various temperatures and pressures. In this formula V is the volume occupied by the liquefied gas dissolved per gram of water, P_0 is the vapour pressure, and σ the surface tension of the liquefied gas at the temperature, whilst P is the equilibrium gas pressure. The constant K for ammonia has value 0·49 and $1/n$ the value 0·69. The solubility data of hydrogen chloride, sulphur dioxide, and carbon dioxide have been plotted according to this formula. J. F. S.

Change of State of Aggregation and Polymorphism. KARL SCHAUM (*Z. anorg. Chem.*, 1921, **120**, 241—260).—The assumption of Ostwald that there is a relatively broad region of metastability, below the normal melting point, in which spontaneous crystallisation

is impossible, appears unnecessary. On the other hand, it is highly improbable that, in a small quantity of liquid in a limited time, the necessary number of molecules in the correct space-lattice orientation to form a nucleus in equilibrium with the liquid at a given temperature will ever come together. It is reasonable to assume with Ostwald the existence of "anisotropic" molecules, but these cannot alone have a "resonance" effect on other "isotropic" molecules and so cause crystallisation. The ageing of a liquid by keeping it some time above its melting point, whereby the tendency to supercooling is increased, is doubtless due to the change of anisotropic molecules into the isotropic form. Anisotropic molecules probably become associated with one another to form space-lattice fragments, and when a crowd of these come together they may cause a field of force strong enough to bring about the formation of a crystal nucleus. As the temperature falls, this process is facilitated through the slowing down of translatory and rotatory motion, but, on the other hand, the probability of the crowding together of anisotropic molecules and space-lattice fragments is lessened.

[With ELLI RIFFERT.]—The crystallisation process is investigated statistically by counting the number of nuclei formed when a thin film of liquid is allowed to crystallise between microscope slide and cover-glass under different conditions. A large number of organic substances were used for the purpose. There is a strong tendency for the nuclei to form in the outer zone of the cover-glass, particularly at the edge, and sometimes they will form again and again at a particular spot. The tendency to nuclei formation at boundary surfaces, especially a triple boundary such as air-glass-liquid, is probably a surface tension phenomenon. The number of nuclei formed increases as the temperature is lowered, finally reaching a maximum. In these experiments, the melted substance was brought suddenly down to the temperature at which the nuclei were to be formed. Pressure may cause the development of nuclei. The substances examined (about 200) fall into five different types, according to the manner in which they crystallise; the states of aggregation assumed by crystallisation from nuclei are described as radiating, bunching, star-like, tabular, and mosaic.

E. H. R.

The K_{α} Doublet, with New Determinations of the Lattice Constants of some Crystals. WALTHER GERLACH (*Physikal. Z.*, 1922, 23, 114—120).—When a somewhat coarse powder is used in the Debye-Scherrer method, the K_{α} lines appear double. A method for the measurement of the fine structure of Röntgen lines is described. The lattice constants, in 10^{-8} cm., of the following crystals were determined: silicon (identical, within the limits of error, with "amorphous," graphitoid, and crystalline silicon): 5.415 ($\pm 0.3\%$) with copper radiation; 5.410 ($\pm 0.3\%$) with nickel radiation; regular zinc blende 5.390 ($\pm 0.3\%$) with copper radiation; calcium fluoride 5.455 ($\pm 0.1\%$) with copper radiation; 5.478 ($\pm 0.2\%$) with nickel radiation. The doublet distance $K_{\alpha_1-\alpha_2}$

was determined with an accuracy of about 3% as 3.842 X.U. (1 "X-Unit" = 10^{-11} cm.) for copper, and 3.772 X.U. for nickel. Bragg's model for fluorite was confirmed. J. R. P.

Freezing of Hydrosols. A. LOTTERMOSER (*Kolloid Z.*, 1922, 30, 133).—A complaint that Gutbier (this vol., ii, 142) has used the author's statements in connexion with the influence of freezing on sols in a manner opposite to that which the author has intended. This has arisen because the author's statements have not been understood in the sense demanded by the context. J. F. S.

Influence of the Concentration of Colloids on their Precipitation by Electrolytes. HARRY B. WEISER and HENRY O. NICHOLAS (*J. Physical Chem.*, 1921, 25, 742—757).—The precipitation of colloidal chromic oxide, Prussian blue, ferric oxide, and arsenious sulphide of various concentrations by electrolytes with precipitating ions of different valency has been investigated. With colloidal chromic oxide, Prussian blue, and ferric oxide, the precipitating values of all electrolytes decrease as the concentration of the colloid decreases, irrespective of the valency of the precipitating ion; but the decrease is least marked with electrolytes having univalent precipitating ions. With colloidal arsenious sulphide, the precipitating value of electrolytes with univalent precipitating ions increases as the concentration of the colloid is decreased. The manner in which the precipitation value of an electrolyte varies with the concentration of colloid is determined to a large extent by the relative adsorbability of the precipitating ion and the stabilising ion. If the adsorption of the stabilising ion of an electrolyte is negligible and the adsorption of the precipitating ion is very large, the precipitation value varies almost directly with the concentration of the colloid. This condition may be realised experimentally with electrolytes which have tervalent and quadrivalent ions which precipitate in extremely low concentration. If the adsorption of the stabilising ion of an electrolyte is appreciable, the precipitating value is increased. This effect is more pronounced the greater the dilution of the colloid, since the decreased opportunity both for the collision and the coalescence of particles combines to render the colloid proportionately more stable so that relatively more of the precipitating ion must be added for complete coagulation. The influence of adsorption of the stabilising ion is more marked with electrolytes having univalent precipitating ions which precipitate only in very high concentration. Under these conditions, the precipitation value decreases much less sharply than under the conditions obtaining when the stabilising ion is only slightly adsorbed, and may even increase as the concentration of the colloid is decreased. The laws put forward by Burton and Bishop (*A.*, 1921, ii, 176) in connexion with the variation in the coagulative power of ions with the concentration of the disperse phase are shown to be quite limited in their applicability. The hypothesis of Krulyt and van der Spek (*A.*, 1919, ii, 498) on the nature of the process of coagul-

ation is inadequate, since it fails to take into account the influence of the stabilising ions of electrolytes. J. F. S.

Relation of Anomalous Osmosis to the Swelling of Colloidal Material. F. E. BARTELL and L. B. SIMS (*J. Amer. Chem. Soc.*, 1922, **44**, 289—299).—The authors discuss some seven hypotheses put forward previously to account for the swelling of colloids, and find none of them is applicable to all cases of swelling. The relationship of the swelling of gels to the osmotic effects is considered, and as a working hypothesis it is assumed that a gel may be considered as composed essentially of a framework of membranes and that the hypothesis which accounts for the flow of a liquid through membranes also accounts to a certain extent for the flow of a solution into a cellular gel. The view is held that swelling is the result of imbibition together with the operation of effects which produce anomalous osmosis. It would therefore follow that in swelling the tendency for normal osmosis is always operative, solution in the membrane may take place in some instances, forces of attraction and repulsion are operative, capillarity plays a part, negative or positive adsorption of solute may have an effect, electrical effects due to electrical orientation of the system may enter in, and swelling unaccompanied by electrical effects may occur, as with solutions of non-electrolytes with some gels. For the purpose of comparing anomalous osmotic effects with swelling effects, a series of osmotic pressures using parchment membranes and a series of swelling experiments using *Laminaria* have been carried out with solutions of potassium citrate (0.05*M*), sodium hydroxide (0.0001*N*), hydrochloric acid (0.0001*N*), and aluminium chloride (0.05*M*). The results show that those solutions which exhibit a negative osmotic tendency with parchment produce a swelling effect with *Laminaria*, and those which have a positive osmotic tendency produce a shrinking of the *Laminaria*. A further series of experiments with solutions of various electrolytes of several concentrations confirms the foregoing results. In all cases, the osmosis-concentration curves have the exactly opposite form to those of the concentration-swelling curves. J. F. S.

Lecture Experiment. Protective Colloids. J. NEWTON FRIEND (*Nature*, 1922, **109**, 341).—It appears to be a general law that protective colloids or organic emulsoids tend to retard the velocities of such reactions, whether chemical or physical, as involve a change of state in one or more of the components. Examples of such retardation, the rate of which conforms in many cases to the requirements of the adsorption law, are to be found in the solution of metals in acids, corrosion in neutral media, solution and precipitation of salts, and replacement of one metal by another. An illustration is afforded by the precipitation of mercuric iodide from solutions of mercuric chloride and potassium iodide, the unstable yellow form first precipitated ordinarily turning rapidly into the more stable red variety. In the presence, however, of 1% of gelatin, after a momentary formation of colloidal mercuric iodide, a fine, canary-yellow precipitate is produced, and remains

practically unchanged for half an hour or more. The slow change to the red variety is markedly accelerated by sunlight. A. A. E.

Pure Aniline and Water Emulsions. SHANTI SWARUPA BHATNAGAR (*J. Physical Chem.*, 1921, 25, 735—741).—The rate of coagulation of aniline-water emulsions by potassium chloride, sodium chloride and sulphate, barium, strontium, chromium, and aluminium sulphates has been investigated at 46°. The results show that in general the precipitating power of electrolytes is the same as in the case of many sols, tervalent ions being more effective than bivalent ions and these more so than univalent ions. The order of the precipitating efficiency of the kations is given as $\text{Al} > \text{Cr} > \text{Ba} > \text{Sr} > \text{K} > \text{Na}$. The effect of dilution on the emulsions is to increase the time required for precipitation; thus an emulsion which is normally precipitated in ten minutes by barium chloride requires eighteen minutes seven seconds when diluted by an equal volume of water saturated with aniline, and when again diluted by the same volume of saturated water forty minutes. J. F. S.

The Detergent Power of Soap Solutions. R. T. A. MEES (*Chem. Weekblad*, 1922, 19, 82—85).—The conception of Martin Fischer of a soap solution as a two-phase system is inadequate, since it does not exclude other gels, for example, gelatin solutions. Similarly, the orientation theory of Langmuir and Harkins does not exclude other substances possessing polarity within the molecule. The unique power of soap solutions to stabilise emulsions can be explained by combining the two theories, since only soaps have the necessary polar character within the molecule without at the same time having molecules soluble as such in oil and water. A soap such as sodium palmitate has the reactive sodium ion, which tends to confer solubility in water, opposed within the molecule to the inert, saturated carbon-hydrogen chain, which tends to confer solubility in oily substances. When such a soap is intimately admixed with an oily substance and water, it forms a bounding layer between the particles of each, its molecules being orientated with the sodium ion towards the water, the carbon-hydrogen chain towards the oil, acting as a layer possessing polar solubilities; in such a system, surface tension disappears.

Earthy impurities, such as clays, resemble soaps in stabilising water-oil emulsions, being lyophil towards both; in presence of soap solution, they are adsorbed by the latter in the bounding surfaces between the particles of oily matter and water.

The detergent power of soaps is therefore primarily a chemical property due to the polarity of molecules insoluble as such in oil and water. S. I. L.

Cellulose Nitrate as an Emulsifying Agent. HARRY N. HOLMES and DON H. CAMERON (*J. Amer. Chem. Soc.*, 1922, 44, 66—70).—The use of cellulose nitrate as an emulsifying agent has been studied. It is shown that this substance serves as an excellent emulsifying agent in dispersing water or glycerol throughout

amyl acetate, acetone, or other solvents for the nitrate. Visible concentration films of cellulose nitrate were observed round large drops of water emulsified in amyl acetate-benzene mixtures. An increase in concentration of cellulose nitrate at the acetone-glycerol interface was observed and studied quantitatively. The properties of an ideal emulsifying film are discussed. J. F. S.

Chromatic Emulsions. HARRY N. HOLMES and DON H. CAMERON (*J. Amer. Chem. Soc.*, 1922, **44**, 71—74).—Two immiscible liquids can be emulsified with suitable emulsifying agents, such as cellulose nitrate, to produce a transparent emulsion when the refractive index of the two phases is the same, and when the dispersive power ($n_F - n_C$) of the two phases is the same. A chromatic (structural colour) emulsion is produced when the refractive index of both phases is the same and the dispersive power of one phase is much greater than that of the other phase. The greater the difference the more intense the resulting colour.

J. F. S.

Application of Statistics to Chemical Equilibria. K. F. HERZFELD (*Physikal. Z.*, 1922, **23**, 95—100; cf. A., 1921, ii, 313).—Statistical principles have been applied to chemical equilibria, in connexion with polymorphic substances and equilibria in gaseous systems at low temperatures and at high temperatures. A reason is advanced for the good agreement found between experimental results and those calculated by the Nernst approximation formula, and it is shown to be probable that the conventional chemical constant is always about 3.

J. F. S.

Equilibrium in Mixtures of Binary Electrolytes. CHARLES A. KRAUS (*J. Amer. Chem. Soc.*, 1921, **43**, 2507—2514).—A theoretical paper in which it is shown that the isohydric principle, when applied to mixtures of electrolytes, leads to a functional relationship of the form $(C_i)^2/C_u = F(\Sigma C_i)$ in which C_u is the concentration of the undissociated fraction, C_i the ion concentration, ΣC_i the sum of the concentrations of all the negative and positive ions in the solution, and $F(\Sigma C_i)$ any explicit function of the total ion concentration of the solution. As the concentration of the second electrolyte diminishes, the function $P_i/C_u = F(P_i)$, in which P_i is the ion product, reduces, in the limit, to the same form as the function resulting from the isohydric principle. The values of the specific conductivity of mixtures of sodium chloride and hydrochloric acid as calculated according to this principle are shown to furnish a better agreement with the measured values than are those on the isohydric principle.

J. F. S.

Partition of Chlorine between Water and a Gaseous Phase. W. S. TROV (*Nachr. Physik.-chem. Lab. Semsoinsés*, 1917, 102—110; from *Chem. Zentr.*, 1921, iii, 1222).—The partition of chlorine at 20° between water and air containing this gas is given by the formula $(y - 1.748)x^2 = (72.52)^3$, where x is the volume of chlorine per thousand volumes of air and y is the volume dissolved in the water under the corresponding partial pressure. For example,

when $x=10$ parts per thousand, $y=74.27$ under a partial pressure of 7.6 mm. G. W. R.

Nature of Secondary Valency. III. Partition Coefficients in the System Water-Ethyl Ether. HOMER W. SMITH (*J. Physical Chem.*, 1921, **25**, 605—615, 616—627; cf. A., 1921, ii, 315, 324).—In the first paper, details are given of the determination of the partition coefficient of forty-nine organic acids and fifteen organic bases at 25° in the system water-ethyl ether. The results obtained from a comparative treatment of these partition coefficients in terms of molecular volume are in good agreement with the results obtained previously (*loc. cit.*) in the systems water-xylene, and water-chloroform. The results are discussed in connexion with the principles laid down in previous papers. The second paper is a supplementary note on the method of correction employed in the earlier papers. Corrected values for the partition of chloroacetic acid in the system water-chloroform and for a number of organic acids and bases in the systems water-ether are given.

J. F. S.

Nature of Secondary Valency. IV. Partition Coefficients in the System Glycerol-Acetone. HOMER W. SMITH (*J. Physical Chem.*, 1921, **25**, 721—734; cf. preceding abstract).—The partition coefficients of fifty acids, amines, and amine acid phthalates have been determined in the system acetone-glycerol. The present results are in excellent agreement with those obtained in the systems previously investigated (*loc. cit.*). In the present series of experiments, a fact, which has not been observed previously, is noted, namely, that under certain conditions, concerning the nature of the solvents and the solutes, a solute is not distributed, but is concentrated entirely in one or other of the solvents. The series constants for the acids and amines have the same absolute value in this system, as contrasted with the different values found in systems containing water as one solvent. Of the substances examined in the system glycerol-acetone, benzoic acid, benzoic acid, $\alpha\beta$ -dibromopropionic acid, 2:4-dinitrobenzoic acid, diisobutylamine, salicylic acid, 2:4:6-trinitrobenzoic acid, and *n*-valeric acid pass completely into the acetone layer.

J. F. S.

Equilibrium in the Reaction between Sulphur Dioxide and Water. F. RUSSELL BICHOWSKY (*J. Amer. Chem. Soc.*, 1922, **44**, 116—132).—The reactions between liquid sulphur and aqueous sulphuric acid have been investigated over a wide range of conditions. At low concentrations of sulphuric acid, two reactions occur simultaneously, namely, (1) $S_{(\lambda\mu)} + 2H_2SO_4(aq) = 2H_2O(liq) + 3SO_2(gas)$ and (2) $4S_{(\lambda\mu)} + 4H_2O(liq) = 3H_2S(gas) + H_2SO_4(aq)$. Both reactions are reversible. In the acids of a greater molecular fraction than 0.05, and at temperatures up to the critical temperature of water, the second reaction becomes negligible. Equilibrium pressures for the first reaction have been measured for acids of molecular fractions ranging from 0.041 to

0.363 and at temperatures from 527° Abs. to 613° Abs., using an apparatus of a new type. The vapour pressures of sulphuric acid of various concentrations have been measured. Using these and other existing data, the free energy for the reaction $S_{(A)} + 2H_2SO_4(M) = 2H_2O(gas) + 3SO_2(gas)$ has been calculated to be 31500 cal. at 25° and standard conditions. From the data the free energy of both aqueous sulphuric acid and sulphur dioxide may be calculated. A new type of apparatus for measuring equilibrium or vapour pressures of corrosive liquids up to high pressures and temperatures is described. This consists of a quartz tube contained in a mercury-filled steel bomb which communicates with a pressure gauge. The quartz tube is constructed so that the material is contained in a chamber at the top; this communicates through a quartz capillary with a second chamber filled with mercury which is in contact with the mercury in the steel jacket. The upper part of the bomb only is heated.

J. F. S.

Equilibria in the Systems: Carbon Disulphide-Methyl Alcohol and Carbon Disulphide-Ethyl Alcohol. E. C. McKELVY and D. H. SIMPSON (*J. Amer. Chem. Soc.*, 1922, **44**, 105—115).—The temperature of complete mutual miscibility of mixtures of carbon disulphide with ethyl and methyl alcohols respectively have been determined. The critical solution temperature of the system carbon disulphide-methyl alcohol is 35.7° and the critical concentration is about 85% of carbon disulphide; solid methyl alcohol separates from the layer rich in alcohol at -99.6°. The critical solution temperature of the system carbon disulphide-ethyl alcohol is -24.4° and the critical concentration about 83% by weight of carbon disulphide. All mixtures freeze at -111.7°. It is shown that the critical solution temperature offers an extremely sensitive criterion of the purity or changes of purity of either component, provided that the other component is either pure or of constant composition. Distillation of carbon disulphide from ceresin is shown to remove the impurities and yield a perfectly pure product.

J. F. S.

The Quaternary System: Ammonium Chloride-Sodium Sulphate-Ammonium Sulphate-Sodium Chloride-Water. ALBERT CHERBURY DAVID RIVETT (*T.*, 1922, **121**, 379—393).

Induction Periods in Reactions between Thiosulphate and Arsenite or Arsenate. A Useful Clock Reaction. GEORGE SHANNON FORBES, HOWARD WILMOT ESTILL, and OSMAN JAMES WALKER (*J. Amer. Chem. Soc.*, 1922, **44**, 97—102).—The precipitation of arsenious sulphide from an acidified solution (acetic acid) containing sodium thiosulphate and arsenite is preceded by an exceedingly sharp and reproducible induction period. The duration of the induction period is strictly in inverse proportion to the concentration of thiosulphate and is practically independent of the concentration of the arsenite or the acetic acid. Hydrochloric acid prolongs the induction periods and is much better for experimental demonstration of induction periods, of the relationship

between concentration and reaction velocity, and of the temperature coefficient of the reaction velocity, than the familiar sulphurous acid and iodate reaction. The prolongation of the periods by strong acids, the limiting concentrations yielding precipitates when arsenates are used, and the speed of precipitation were also investigated. J. F. S.

The Combustion of Complex Gaseous Mixtures. WILLIAM PAYMAN and RICHARD VERNON WHEELER (T., 1922, 121, 363—379).

Oxidation of Nitric Oxide and its Catalysis. CHARLES LALOR BURDICK (*J. Amer. Chem. Soc.*, 1922, 44, 244—251).—The rate of oxidation of nitric oxide by oxygen in the presence of glass, pumice, pumice impregnated with tungstic acid, nickel oxide, cobalt oxide, nickel, cobalt, and manganese dioxide, platinised pumice, platinised asbestos, coke, charcoal, and coconut charcoal, has been investigated at temperatures from 0° to 100°. It is shown that the reaction $2\text{NO} + \text{O}_2 = 2\text{NO}_2$ is not catalysed by ordinary porous material either impregnated or unimpregnated, but that in the presence of the highly absorptive forms of carbon or charcoal the rate of the reaction may be accelerated as much as five hundred times. The presence of water vapour in the gaseous mixtures greatly decreases the activity of the charcoal catalysts. Increase of temperature above the point at which the aqueous vapour condenses counteracts this effect to some extent. The temperature coefficient of the velocity of reaction for both the catalysed and uncatalysed change is generally negative. In the presence of water vapour the temperature coefficient of the catalysed reaction is apparently positive. This is probably due to the decreased absorption of water by the catalyst at elevated temperatures. J. F. S.

The Transformation of Ammonia into Carbamide. C. MATIGNON and M. FRÉJACQUES (*Compt. rend.*, 1922, 174, 455—457; cf. A., 1921, ii, 33).—In continuation of previous work (*loc. cit.*), the yields of carbamide from ammonium carbamate at the equilibrium point at different temperatures are given. Applying the law of mass action to the system in equilibrium the authors calculate the heat of the reaction to be -6 cal., the experimental value being -7.7 . The progress of the reaction with time at different temperatures from 130° to 145° has been studied. The curves indicate that the water formed has an effect on the velocity of the reaction. The effect of certain catalysts such as thorium oxide, aluminium oxide, kaolin, etc., has been examined. The results indicate that whilst they may accelerate the reaction at low temperatures, they are without benefit at about 150°. W. G.

The Processes $\text{NH}_2\cdot\text{CO}\cdot\text{ONH}_4 + \text{H}_2\text{O} \rightleftharpoons (\text{NH}_4)_2\text{CO}_3$ and $\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3$. CARL FAURHOLT (*Z. anorg. Chem.*, 1921, 120, 85—102).—Experiments on the velocity of the conversion of ammonium carbamate into carbonate indicated that the reaction proceeded first with liberation of carbon dioxide, and this con-

clusion was confirmed by an investigation of the velocity of the reaction $\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3$. A new method was devised for estimating the proportion of hydrated carbon dioxide, that is, of carbonic acid, in a solution of the gas. When an aqueous solution of carbon dioxide is added to a large excess of ammonia, the anhydrous carbon dioxide immediately forms carbamate, which is so stable in the concentrated ammonia that there is time to precipitate the carbonate as barium carbonate, filter, wash, and titrate it. By this method the dissociation constant of carbonic acid to $\text{H}^+ + \text{HCO}_3^-$ was found to be $10^{-6.3}$, against $10^{-6.7}$ to $10^{-6.4}$ found by Thiel and Strohecker (A., 1914, ii, 361). This method was applied to making approximate determinations of the velocity of hydration of carbon dioxide in solutions of sodium hydroxide, sodium carbonate, and mixtures of carbonate and hydrogen carbonate. At 0° , hydration is immediate in sodium hydroxide solution, but in a solution containing $\text{Na}_2\text{CO}_3 : \text{NaHCO}_3$ in the ratio 6 : 4 only 77% of the carbon dioxide added is hydrated in the first eighty seconds. Hydration is thirteen times as rapid at 18° as at 0° . Velocity of hydration is inversely proportional to hydrogen-ion concentration. By strong acids, carbonic acid is immediately dehydrated, but by weak acids much more slowly.

The same method of analysis was used for investigating the decomposition of ammonium carbamate in acid, aqueous, ammoniacal, and alkaline solution. Before analysis, the solutions were made alkaline with 0.1*N* sodium hydroxide to stabilise the carbamate. In aqueous solutions containing ammonia and ammonium chloride, the proportion of carbamate present at equilibrium increases with the concentration of either. The equilibrium constant for the reaction $\text{NH}_3 \cdot \text{CO} \cdot \text{O}^- + \text{H}_2\text{O} \rightleftharpoons \text{HCO}_3^- + \text{NH}_3$ at 0° had an average value of about 0.34. By even very weak acids, carbamate is immediately decomposed. On adding alkali again, however, some carbamate is re-formed, since two reactions proceed simultaneously with nearly the same velocity, carbamate being formed from ammonia and carbon dioxide, and carbonate from carbonic acid, which is formed at once by hydration of part of the carbon dioxide by the alkali added.

In a 0.05 mol. solution of carbamate in water at 0° , equilibrium is reached in about a day; in a 0.1*N*-ammonia solution, three days are required, in a 0.4*N*-ammonia solution, ten days; at 18° the reaction is twenty times as fast. In sodium hydroxide solution, the reaction is much slower, requiring about three days in a 0.1*N*-solution and a month in a 1.0*N*-solution, at 18° . E. H. R.

The Velocity of Formation of Barium Peroxide. NOBUJI SASAKI (*Mem. Coll. Sci. Kyoto*, 1921, 5, 9—96).—For the investigation of the velocity of formation of barium peroxide from barium oxide and oxygen at high temperatures, 400—625°, a special balance was devised, so constructed that the specimen under examination could be weighed at any time, being suspended from one arm of the balance, which was totally enclosed and operated from outside by means of magnets. A new type of gas washer was also devised.

It consists of a U-tube of glass, one arm being narrow and the other expanded into an elongated bulb. The two arms are bridged by a narrow glass spiral in such a manner that the gases, entering by the narrow arm, bubble through the liquid in the spiral and escape at the top of the wide arm, which acts as a reservoir for the washing liquid. The velocity curves for the formation of barium peroxide were of an unusual form. The greater part of the paper is devoted to a mathematical analysis of these curves, and it is shown that they may be explained on the assumption that the oxide consists of a great number of very small, equal spherical particles which begin to react one after another, the number of spheres becoming active in unit time being proportional to the quantity of peroxide already formed. E. H. R.

The Influence of Protective Colloids on the Corrosion of Metals and on the Velocity of Chemical and Physical Change. JOHN ALBERT NEWTON FRIEND and REECE HENRY VALLANCE (T., 1922, 121, 466—474).

Influence of Temperature on Two Alternative Modes of Decomposition of Formic Acid. C. N. HINSHELWOOD, H. HARTLEY, and B. TOPLEY (*Proc. Roy. Soc.*, 1922, [A], 100, 575—581).—The thermal decomposition of formic acid has been investigated at 283°, and from the results an hypothesis has been formulated in connexion with the mechanism of the reaction. Sabatier and Mailhe (A., 1911, i, 515) have shown that formic acid may decompose according to the equations (a) $\text{H}\cdot\text{CO}_2\text{H}=\text{CO}_2+\text{H}_2$, (b) $\text{H}\cdot\text{CO}_2\text{H}=\text{CO}+\text{H}_2\text{O}$, and (c) $2\text{H}\cdot\text{CO}_2\text{H}=\text{H}\cdot\text{CHO}+\text{CO}_2+\text{H}_2\text{O}$. Assuming that in the decomposition certain molecules are activated, either by collision with other molecules or by the absorption of quanta of radiation, the question arises whether every molecule which possesses the necessary critical energy reacts at once or whether, in addition to possessing this, it must be in a certain phase. Under the conditions of the experiments described, only reactions a and b come into consideration. If E_1 and E_2 are the critical energies for the two alternative reactions and N is the total number of molecules, it follows from the kinetic theory that the number of molecules possessing these respective energies are $N_1=N\epsilon^{-E_1/RT}$ and $N_2=N\epsilon^{-E_2/RT}$, taking the system as of two degrees of freedom. Then $N_1/N_2=\epsilon^{-(E_1-E_2)/RT}$, so that if one mode of reaction is associated with a slightly lower critical energy it would predominate very largely if the number of active molecules were the only factor. Actually it is found that although the two alternative modes of decomposition of formic acid proceed with the same velocity, yet the critical increments are so different that one reaction should predominate almost entirely unless a phase restriction is introduced, and an interpretation of the phase factor is suggested for this case, as follows. The critical increments for the two reactions are calculated to $E_{\text{CO}}=16000$ cal./mol. and $E_{\text{CO}_2}=28000$ cal./mol. The numbers of molecules activated for the two reactions are proportional to $\epsilon^{-28000/RT}$ and $\epsilon^{-16000/RT}$, respectively, that is about ϵ^{10} molecules should yield carbon monoxide for one yielding the

dioxide, but since the rates of both reactions are identical a phase restriction must govern the reaction yielding the monoxide. If the formic acid molecule is assumed to be adsorbed on the glass surface by the residual affinity of the carbonyl group, then in the case of a sufficiently activated molecule a violent collision between the hydrogen- and hydroxyl-groups may occur, and a molecule of hydrogen be split off. Although the requisite degree of activation may be attained but rarely, nevertheless a high percentage of these collisions in activated molecules may result in this type of decomposition. On the other hand, for water to be eliminated and carbon monoxide left, a high degree of activation is not necessary, possibly since the removal of a hydroxyl group, as a whole, from an organic compound is a much less profound change than the fission of the hydrogen and oxygen, but very few of the activated molecules may decompose, because it is necessary to await the contingency that the forces holding the hydroxyl and hydrogen to the carbon should be weak enough to allow the simultaneous "uncoupling" of the molecule. J. F. S.

The Replacement of Halogen in 4-Chloro-3-nitrobenzonitrile and in 4-Bromo-3-nitrobenzonitrile. II. T. J. F. MATTAAR (*Rec. trav. chim.*, 1922, 41, 103—111; cf. this vol., i, 251; also Conrad, Hecht, and Brückner, A., 1890, 4, 327, and 1046; A., 1891, 796).—In order to measure the effect of the cyanogen group on the replacement of halogen the velocity of reaction of 1-chloro-2:4-dinitrobenzene, 4-chloro-3-nitrobenzonitrile, 1-bromo-2:4-dinitrobenzene and 4-bromo-3-nitrobenzonitrile with sodium methoxide, with sodium ethoxide, and with sodium phenoxide in both ethyl alcohol and methyl alcohol solution has been investigated. The following conclusions are drawn: (1) the halogen atom is more active in the substituted dinitrobenzene than in the substituted nitrobenzonitrile, (2) chlorine is more reactive than bromine in corresponding compounds, (3) sodium ethoxide reacts more quickly than sodium methoxide, (4) the reaction in ethyl alcohol takes place about twice as quickly as that in methyl alcohol. H. J. E.

The Effect of Temperature and the Concentration of Hydrogen Ions on the Rate of Destruction of Antiscorbutic Vitamin (Vitamin-C). V. K. LAMER, H. L. CAMPBELL, and H. C. SHERMAN (*J. Amer. Chem. Soc.*, 1922, 44, 172—181; cf. *ibid.*, 165).—Using the quantitative method previously described (*loc. cit.*), the authors claim to have shown that, under the experimental conditions, the velocity of destruction of vitamin-C by heat decreases with the time and in greater degree than would be expected if the reaction followed the unimolecular law, or the square root rule of Schutz. The temperature coefficient is of low order, namely, $Q_{10}(60-80^{\circ})=1.23$; $Q_{10}(80-100^{\circ})=1.12$. The effect of reducing the hydrogen-ion concentration from $p_H=4.3$ to $p_H=5.2-4.9$ is to increase the destruction during one hour at 100° from 50% to about 58%. When the material was made alkaline, $p_H=10.9-8.3$, the destruction was 61—65%. If re-acidification after heating in

alkaline solution was omitted, there was still further destruction, if the material was allowed to remain alkaline for five days at 10°. The material used as the source of vitamin-*C* was tomato juice.

W. G.

The Influencing of Catalysts and Specifically Active Catalysts. E. ABEL (*Ber.*, 1922, 55, [B], 322—323).—A final reply to Rosenmund and Zetzsche (cf. Abel, A., 1921, ii, 542; Rosenmund and Zetzsche, A., 1921, ii, 320, 392, 393; this vol., ii, 41).

H. W.

The Oxygen-Hydrogen Catalysis by the Platinum Metals and the Contact Potentials in Presence of Aqueous Electrolytes. K. A. HOFMANN (*Ber.*, 1922, 55, [B], 573—588).—Since previous experiments (A., 1920, ii, 240) were performed with platinised porous earthenware tubes, whilst Willstätter and Waldschmidt (A., 1921, ii, 185) employed platinum black, these investigators are not justified in concluding that their results are incompatible with the views of Hofmann and Zipfel (*loc. cit.*). In the present work, no indication could be discovered of the peroxide-hydride, to which Willstätter and Waldschmidt ascribe the favourable effect of oxygen on the catalytic activity of hydrogen (D.R.-P. 301364, Willstätter and Waldschmidt, *loc. cit.*). The volume of oxygen occluded by platinised earthenware is shown to be so small that any peroxide produced cannot bear any relation to the actual effect of oxygen on catalysis, which is rather due to its influence on the catalyst surface. At the same time, the small amounts of gas adsorbed by porous surfaces render them much more suitable than platinum for determinations of variations of contact potential during catalysis. Measurements of the potential difference between such an electrode and one of platinum in contact with hydrogen at atmospheric pressure show that, whether the electrolyte be 17% sulphuric acid or 4% sodium hydrogen carbonate solution, contacts impregnated with platinum, palladium, or iridium in contact with a mixture of hydrogen and oxygen are electrically modified in the opposite sense to that of their previous treatment (with one or other of the gases), and the more rapidly this occurs the more rapidly is water-formation catalysed. The activity diminishes on prolonged treatment with hydrogen or, especially in the case of palladium, with oxygen. It is improved by repeated variation of this treatment, by repeated use with the stoichiometrical mixture of the gases, and by varying galvanic polarisation. Light and moderate temperature changes are much less effective. Hydrogen exerts a greater effect on the catalysis potentials with increased hydrogen-ion concentration of the electrolyte, whilst the reverse is true of oxygen. These results indicate that neither pure hydrogen nor hydrogen-oxygen mixtures produce any special oxidising or reducing compound on the catalyst surface of the kind suggested by Willstätter which determines the potential. Direct instantaneous determinations of the effect on the same electrode potentials of the addition of hydrogen peroxide show that this always acts as a strong oxidising agent, and renders it

highly improbable that any peroxide-hydride could exert a strong reducing influence, apart from the extreme rapidity with which the known peroxides are decomposed by platinum metals. Although it is not improbable that hydrogen peroxide is an intermediate product in the formation of water, its existence is so transient as to elude all attempts to detect it.

J. K.

Atomic Structure. N. BOHR (*Nature*, 1921, 108, 208—209; cf. *ibid.*, 1921, 107, 104, 170).—Further examination of the orbits of the electrons postulated in the previous communication has shown that the conception of the atom there outlined cannot be reconciled with that which assumes the atom to be composed of a number of well-defined spherical shells of electrons moving in sharply separated regions. The electrons in the atom are still regarded as being arranged in groups in such a way that the orbit of every electron within a particular group is characterised by the same total number of quanta. In proceeding outwards from the nucleus, the number of quanta characterising a certain group of orbits is larger by one unit than that of the preceding group. The electrons of certain sub-groups will also penetrate into the region of the orbits of electrons of inner groups, giving rise to group-coupling, which is reflected in the stability of the atom. These considerations are applied to the cases of the iron group and the rare earths.

A. A. E.

A Magnetic Model of Atomic Constitution. J. K. MARSH and A. W. STEWART (*Nature*, 1921, 109, 340).—If the central magnetic pole in a tank is maintained at a strength always exactly equal and opposite to the combined powers of a number of floating magnets, and if the number of the latter corresponds with units of atomic weight in various elements, the magnets arrange themselves in two clearly defined and sharply separated groups, termed for convenience the nucleus and ring, respectively. It is a remarkable coincidence that the series of groupings obtained corresponds exactly with Aston's suggestions regarding the atomic structure of a number of elements and isotopes. In the case of boron-11 and carbon, both nuclei contain six magnets arranged in two pairs of concentric triangles, a peculiarity which seems to have some connexion with certain chemical and physical similarities.

A. A. E.

Correlation of Atomic Structure and Spectra. HAROLD S. KING (*J. Amer. Chem. Soc.*, 1922, 44, 323—328).—A theoretical paper in which it is shown that Bury's modification (this vol., ii, 43) of Langmuir's hypothesis of the structure of elements is confirmed in addition to the relationship between the atomic diameter and the number of electron shells, also by the relationship between the atomic number and the number of lines in the arc and spark spectra, respectively.

J. F. S.

Motion of Electrons in Atoms. J. S. TOWNSEND and V. A. BAILEY (*Phil. Mag.*, 1922, [vi], 43, 593—600).—It is shown that when an electron moving with a velocity of the order of 10^8 cm. per second collides with a molecule of argon it loses only about

one ten-thousandth of its energy, whereas previous experiments showed that in collision with oxygen, nitrogen, or hydrogen, it loses more than 1% of its energy. The mean free path of an electron moving with these velocities in argon is about ten times that in the other gases at the same pressure. J. R. P.

Relationship between the Internal Friction and the Chemical Constitution of Gases. HARRY SCHMIDT (*Z. Elektrochem.*, 1922, 28, 50—55).—A theoretical paper in which the author has deduced equations by means of which the atomic radius of the elements and the mean cross-section of the simpler molecules may be calculated. Using the formula $r = \sqrt{4k\sqrt{M}/7\pi\eta}$, in which r is the atomic radius, k a constant, M the molecular weight, and η the coefficient of viscosity, the following values of the radii have been obtained: oxygen, 1.000; nitrogen, 1.035; hydrogen, 0.752; chlorine, 1.491; sulphur, 1.477; bromine, 1.841; helium, 0.795; neon, 0.946; argon, 1.336; krypton, 1.553; xenon, 1.794. The mean cross-section of a number of simple compound molecules are also calculated and tabulated. J. F. S.

Atomic Radius and Ionisation Potential. MEGH NAD SAHA (*Nature*, 1921, 107, 682—683; cf. Basu, *Phil. Mag.*, 1920, [vi], 40, 619).—The radius, a_x , of an atom (the distance from the nucleus to the outermost electron of the Rutherford-Bohr model) may be calculated from the ionisation potential, V_x , of an element x , from the equation: $a_x = 10^{-8} \times (0.532 \times 13.56) / V_x$ cm. The atomic radii calculated in this way are smaller than those calculated either from crystal data or from the kinetic theory. The following values, expressed as $a \times 10^8$, were obtained, those derived from other methods being placed within brackets: hydrogen, 0.530; helium, 0.28 (1.08); neon, 0.33 (0.65, 1.01); lithium, 1.34 (1.50); sodium, 1.41 (1.77); potassium, 1.67 (2.07); rubidium, 1.73 (2.25); caesium, 1.86 (2.37); copper, 0.94 (1.37); silver, 0.95 (1.77); gold, 0.83; magnesium, 0.95 (1.42); calcium, 1.18 (1.70); strontium, 1.27 (1.95); barium, 1.39 (2.10); zinc, 0.77 (1.32); cadmium, 0.81 (1.60); mercury, 0.69; thallium, 0.99 (2.25); manganese, 0.98 (1.47). A. A. E.

New Method of Determining Molecular Diameters by the Electromagnetic Rotation of the Discharge in Gases. C. E. GUYE and R. RÜDY (*Compt. rend.*, 1922, 174, 382—384).—By applying the theory of ionisation by shock to the phenomenon of rotation of the electric discharge in gases, the velocity of rotation is given by $V = eH/12\pi\sigma^2mM$. Thus with a constant magnetic field and for the same gas, the product of the pressure and the velocity observed must be a constant. A new series of measurements has been made on a number of carefully purified gases. The molecular diameters obtained by this method show a close agreement in value with those deduced from viscosity measurements. Contrary to the general opinion, the observed velocity of rotation is not inversely proportional to the density of the gas,

but to the product of the square of the molecular diameter and the density. W. G.

Atomic Weights and Frequencies. A. REYCHLER (*Bull. Soc. chim. Belg.*, 1922, **31**, 67—72).—Warrington's formula for calculating atomic weights (A., 1915, ii, 160) gives too low a value for elements of the two short series. The author's formula, $A = 2N + 0.007N^2$, gives better values, but is not applicable to hydrogen and is not very satisfactory for elements of the first short series. Attention is directed to the relation of this formula to the equation $\nu = \gamma(N - c)^2$, which expresses frequency of radiation of an element when used as an anticathode. H. J. E.

Qualities of Valency. R. M. CAVEN (*Nature*, 1921, **108**, 210—211).—Polemical. Langmuir's (*ibid.*, 1921, **108**, 101) view that the sodium and chlorine atoms in sodium chloride are not united by a chemical bond is attacked, and criticism offered regarding the postulation of covalency and electrovalency in certain cases. A. A. E.

A Theoretical Derivation of the Principle of Induced Alternate Polarities. ARTHUR LAPWORTH (T., 1922, **121**, 416—427).

An Explanation of the Property of Induced Polarity of Atoms and an Interpretation of the Theory of Partial Valencies on an Electronic Basis. WILLIAM OGILVY KERMACK and ROBERT ROBINSON (T., 1922, **121**, 427—440).

Friend's Theory of Valency. J. D. MAIN SMITH (*Chem. News*, 1922, **124**, 84—86).—Friend's theory of valency as extended in a recent paper (T., 1921, **119**, 1040) is subjected to a detailed criticism. In particular, the direct translation of Thomson's physical conception of "ionised" and "non-ionised" valency into chemical terms is criticised, on the ground that residual valency may partake of the properties of both. Friend's criticisms of some aspects of Werner's theory are discussed. E. H. R.

Space-filling and the Equation of Condition. J. J. VAN LAAR (*Z. anorg. Chem.*, 1921, **120**, 203—208).—A theoretical discussion of the theory of space-filling numbers in relation to van der Waals's equation. E. H. R.

Inorganic Chemistry.

Active Modifications of Hydrogen and Nitrogen produced by α -Rays. F. H. NEWMAN (*Phil. Mag.*, 1922, [vi], **43**, 455—462).—The α -rays from polonium were allowed to act on nitrogen at different pressures in the presence of sodium, potassium, sulphur, phosphorus, iodine, magnesium, arsenic, mercury, and an alloy

of sodium and potassium. Some gas was absorbed. Similar experiments with hydrogen gave absorptions with sulphur, phosphorus, and iodine. The absorption was shown to be due, at least in some cases, to the formation of nitrides and hydrides. The chemical activity of the gases is due to active modifications, probably consisting of neutral atoms and triatomic molecules, and not to ions. The α -rays are the only ones effective in the rays from radioactive substances.

J. R. P.

Constitution of Hydrogen Peroxides. B. ODDO and R. BINAGHI (*Gazzetta*, 1921, 51, ii, 343—348).—See this vol., i, 314.

Separation of the Isotopes of Chlorine. J. N. BRÖNSTED and G. HEVESY (*Nature*, 1921, 107, 619).—When a strong aqueous solution of hydrogen chloride is cooled to -50° and evaporated in a high vacuum, the mixture of water and hydrogen chloride being condensed on a surface cooled with liquid air, the condensed portion of the acid is found to be richer, and the remaining portion poorer, as regards the lighter constituent of chlorine, than ordinary hydrogen chloride. After repeated separations, the two portions were converted into sodium chloride, of which saturated aqueous solutions at 20° had d 1.20222 and 1.20235, respectively, corresponding with a difference of 0.024 unit in the atomic weights of the chlorine isotopes, if it is assumed that the latter have equal atomic volumes. Equal quantities of the isotopic sodium chlorides were also treated with a slight excess of silver nitrate solution, and the excess of silver salt accurately estimated. The results indicated a difference in the atomic weight of 0.021 unit.

A. A. E.

The Separation of Chlorine into Isotopes. WILLIAM D. HARKINS (*Nature*, 1921, 108, 209; cf. Harkins and Broeker, *ibid.*, 1920, 105, 230).—See Harkins and Hayes, this vol., ii, 140.

A. A. E.

Attempt to Separate the Isotopes of Chlorine. H. HARTLEY, A. O. PONDER, E. J. BOWEN, and T. R. MERTON (*Phil. Mag.*, 1922, [vi], 43, 430—435).—According to Aston, ordinary chlorine consists essentially of a mixture of three parts of Cl^{35} and one part of Cl^{37} , and the molecules should be present in the proportions $\text{Cl}^{35}\text{Cl}^{35} : \text{Cl}^{35}\text{Cl}^{37} : \text{Cl}^{37}\text{Cl}^{37} = 9 : 6 : 1$. If light which has passed through a column of such chlorine enters a mixture of chlorine and hydrogen, the initial reaction should use up the three types of molecules in the proportions $1 : 10^9 : 10^{24}$, and the hydrogen chloride formed should be almost entirely HCl^{37} . The gas, after exposure, was treated with mercury to remove unchanged chlorine and the hydrogen chloride, absorbed in water, was converted into sodium chloride. The equivalent of the latter, determined by conversion to silver chloride, showed that no separation had occurred within the limits of experimental error. The circumstances which would prevent a separation of isotopes in the experiment are discussed.

J. R. P.

The Atomic Weight of the Chlorine in an Ancient Mineral, Apatite, from Balme. (MILLER) ELLEN GLEDITSCH and B. SAMDAHL (*Compt. rend.*, 1922, 174, 746—748).—The chlorine in a sample of apatite from Balme (cf. Curie, A., 1921, ii, 396) was converted into sodium chloride which was carefully purified without possible contamination by chlorine from other sources. Atomic weight determinations gave the value for chlorine as 35.49 and 35.45. Thus taking into consideration the age and origin of the mineral, it may be concluded that the chlorine at the time of the formation of the minerals of the primary magma contained the two isotopes in the same proportion as it does to-day or that the two isotopes were then formed in constant proportions. W. G.

The Reaction between Iodine and Sulphurous Acid. ROBERT MILROY MACAULAY (T., 1922, 121, 552—556).

Strength of Hypoiodous Acid. ADOLF FÜRTH (*Z. Elektrochem.*, 1922, 28, 57—58).—Making use of the constants $[I']/[HIO][H^+]/[I_2] = 3 \times 10^{-13}$ (Bray, *Abegg's Handb.*, 4, 2, 456) and $[I_2][I']/[I_3] = 1.3 \times 10^{-3}$ (Jakowkin, A., 1899, ii, 593) and a series of determinations of the concentration of I_3 ions in a sulphuric acid solution of various concentrations of iodine in 0.1N-potassium iodide by a colorimetric method, the author has calculated the dissociation constant of hypoiodous acid. The values found lie between 1.8×10^{-11} and 4.4×10^{-11} , which are in good agreement with a value 4.8×10^{-11} calculated from experiments of Deventer (*Chem. Weekblad*, 1905, 2, 135). This value shows that hypoiodous acid is about a thousand times weaker than hypochlorous acid.

J. F. S.

Preparation and Properties of the Persulphides of Hydrogen. JAMES H. WALTON and LLEWELLYN B. PARSONS (*J. Amer. Chem. Soc.*, 1921, 43, 2539—2548).—The preparation and properties of hydrogen disulphide and hydrogen trisulphide are described. The two compounds are prepared together by the action of hydrochloric acid on sodium polysulphide. Other acids decomposed the polysulphide completely into hydrogen sulphide and sulphur. Two kilos. of crystallised sodium sulphide and 300 grams of flowers of sulphur are placed in a 3-litre round-bottomed flask and mixed, 800 c.c. of water are added, and the flask is closed with a Bunsen valve. After all the sodium sulphide has dissolved, the flask is heated on a water-bath for three to four hours with occasional shaking, when the whole of the sulphur dissolves and a clear solution of polysulphide is obtained. The polysulphide solution is run into hydrochloric acid (d 1.19) which is cooled to -4° to -10° . The addition is made under the surface of the acid at the rate of 3 litres an hour, the solution being rapidly stirred. After a quantity of polysulphide has been added, a brown scum rises to the surface and decomposes with evolution of hydrogen sulphide. At this point, the addition is stopped and the emulsion allowed to settle when a crude oil separates. The oil is a mixture of the two sulphides and about 400—500 c.c. are obtained from

2 kilos. of polysulphide. The oil is separated and dried with phosphoric oxide. The dried mixture is distilled in a quartz glass apparatus fitted with two receivers in series. The first is cooled by cold water and the second by ice and salt. The distillation is carried out at 20—25 mm. pressure and the flask heated in a glycerol bath at 120°. From two volumes of the freshly prepared mixture $\frac{2}{3}$ volume of hydrogen trisulphide is obtained in the first receiver and $\frac{1}{3}$ volume of hydrogen disulphide in the second receiver, whilst a residue of sulphur dissolved in the trisulphide remains in the distilling flask. It is essential that quartz apparatus is used in the distillation, because the alkali in glass decomposes both sulphides very rapidly.

The analysis of the persulphides is effected as follows. A weighed quantity (2—3 grams) is placed in a tared quartz test-tube (75 mm.), corked, and weighed. The tube is almost filled with carbon disulphide and the contents are poured into a tared quartz dish (75 mm. diam.), and the tube washed with carbon disulphide. About 20—30 c.c. of carbon disulphide are sufficient. About 10 c.c. of pure acetone are added to the solution, and the dish is covered by a watch-glass. After the evolution of hydrogen sulphide has ceased, the cover-glass is rinsed with carbon disulphide and the contents of the dish are allowed to evaporate spontaneously and the residue of rhombic sulphur crystals is dried at 90° and weighed. This method is rapid and accurate.

Hydrogen trisulphide, H_2S_3 , is a mobile, yellow, oily liquid with an odour similar to that of camphor and sulphur monochloride. Its vapours have an irritating action on the eyes and nose. On cooling strongly, the yellow colour is lost and at the temperature obtained with solid carbon dioxide and ether it is quite colourless. It is soluble in benzene, toluene, chloroform, carbon disulphide, ether, and heptane, and is catalytically decomposed by alcohols, ketones, nitrobenzene, aniline, and pyridine. Hydrogen trisulphide behaves like glass on cooling, becoming more and more viscous as the temperature is reduced to -75° , but on warming again there is a short delay in the rise of temperature at -52° to -53° . It may be distilled in a vacuum, but at atmospheric pressure it foams, evolves hydrogen sulphide, and leaves a gummy residue of sulphur in the mixed sulphides.

Hydrogen disulphide, H_2S_2 , has not been obtained quite free from trisulphide, the amount of impurity is not large and is represented by a deficiency of 1—2% of hydrogen sulphide on decomposition. It is a colourless, mobile oil with a much more severe and irritating odour than the trisulphide, causing tears and a smarting sensation in the nostrils. It is soluble in the same solvents as the trisulphide, but is much more sensitive to impurities. Acetone catalyses the decomposition very violently, so that great care is required in the analysis. It boils at 74.5° and melts between -88° and -90° , but shows no sharp point of solidification. Hydrogen trisulphide has the same action on solutions of copper oleate, ferric chloride, stannic iodide, silver nitrate, and mercuric bromide in ether as hydrogen sulphide. Silver oxide, cupric

oxide, lead dioxide, and mercuric oxide decompose hydrogen trisulphide so violently and with such development of heat that it is ignited; lead oxide, stannic oxide, and magnetite decompose it violently, whilst arsenious, arsenic, ferric, and zinc oxides, barium dioxide, and manganese dioxide bring about only a slow decomposition. Potassium permanganate and potassium dichromate decompose it violently. Most metallic salts decompose the trisulphide, but massive metals have a very slow action. It dissolves readily in liquid hydrogen sulphide and one molecule of the trisulphide dissolves eight atoms of sulphur. J. F. S.

Influence of Freezing on Colloidal Selenium. III. A. GUTBIER and R. EMSLANDER (*Kolloid Z.*, 1922, 30, 97—110; cf. A., 1921, ii, 693; this vol., ii, 142).—Further experiments on the effect of freezing on selenium sols are described. It is shown that selenium sols prepared by means of hydrazine hydrate are stable toward freezing when of suitable concentration, even in the presence of large concentrations of selenious acid. The stability of such sols is dependent on the temperature of formation to a marked extent. A particularly stable system is formed when a 0.002*M* solution of selenium dioxide is reduced by a 1:2000 solution of hydrazine hydrate at 60°, ten parts of selenium dioxide solution being reduced by one part of hydrazine hydrate. The coagulation of selenium sols is greatly accelerated by freezing. Selenium sols may be prepared by the solution of selenium in hydrazine hydrate at ordinary temperature and pouring this molecular disperse system into a large volume of water. With this system, it is shown that the stability of colloidal selenium depends mainly on the degree of dispersion. It is also shown that an optimum concentration of electrolyte is necessary for the stability of selenium hydrosols. In the absence of electrolytes, this system is quite unstable toward freezing. The protective action of the sodium salts of protalbic and lysalbic acids on selenium sols consists in a retardation of the spontaneous coagulation of the suspension. J. F. S.

Hydrates of Selenium Dioxide. WILHELM MANCHOT and KARL ORTNER (*Z. anorg. Chem.*, 1922, 120, 300—309).—Crystallisation of solutions of selenium dioxide did not give trustworthy results, because the crystals obtained were very hygroscopic and could not be dried in a desiccator without undergoing decomposition. The freezing curve of hydrated selenium dioxide showed only one eutectic point at -23° ; the eutectic mixture containing 57% SeO_2 , and consisting of ice and the hydrate $\text{SeO}_2 \cdot \text{H}_2\text{O}$. A maximum was found on the solubility curve corresponding with a composition of 86.07% SeO_2 , that is, $\text{SeO}_2 \cdot \text{H}_2\text{O}$. The vapour pressure curve indicated that $\text{SeO}_2 \cdot \text{H}_2\text{O}$ was the only hydrate present. Molecular weight was found by the freezing-point method. The degree of dissociation was found to be approximately the same as for tartaric acid. No evidence could be obtained of the existence of the associated selenious acid $(\text{H}_2\text{SeO}_3)_2$ in a

freshly prepared solution as claimed by Rosenheim and Krause (this vol., ii, 47).
W. T.

The Elimination of the Heat of Reaction in the Synthesis of Ammonia by High Pressures. GEORGES CLAUDE (*Compt. rend.*, 1922, 174, 681—683).—The device adopted to remove the heat of reaction consists in having the catalyst in a thin-walled inner tube and allowing the oncoming cold gases to circulate through the annular space. The inner tube is so constructed that the amount of heat removed at any one point is just equal to the heat generated at that point. This is effected by varying the thickness of the inner tube along its length. With this new arrangement it is claimed that the life of the reaction tube is considerably increased and the catalyst can very easily be replaced when necessary.
W. G.

The Rôle of Gaseous Impurities in the Catalytic Oxidation of Ammonia. Influence of Hydrogen Phosphide. E. DECARRIÈRE (*Compt. rend.*, 1922, 174, 460—461; cf. A., 1921, ii, 503, 546).—Hydrogen phosphide when present in the ammonia-air mixture as the sole gaseous impurity and only to the extent of 0.00002% exercises a harmful effect and reduces the oxidation by nearly 30%. If the gas is present to the extent of 0.02%, the yield falls from 93.8% to 3.9%. In every case, however, the activity of the catalyst is regenerated when the impurity is removed, the regeneration being slower the greater the percentage of the impurity previously present. In some cases, however, industrial catalysts of the type of those of Ostwald may be destroyed by the momentary presence of traces of hydrogen phosphide in certain favourable circumstances.
W. G.

The Rôle of Gaseous Impurities in the Catalytic Oxidation of Ammonia Gas. EUGÈNE DECARRIÈRE (*Compt. rend.*, 1922, 174, 756—758; cf. preceding abstract).—It has previously been shown that hydrogen sulphide is capable of partly neutralising the injurious influence of acetylene in the catalytic oxidation of ammonia in the presence of finely divided platinum. It is now shown that it can also partly neutralise the injurious influence of hydrogen phosphide. This beneficial effect is not due to the dissociation of the sulphide with the consequent liberation of hydrogen, since hydrogen itself is incapable of neutralising the effect of the phosphide.
W. G.

Nitrous Anhydride. F. FOERSTER (*Ber.*, 1922, 55, [B], 490—491).—In a recent communication (A., 1921, i, 778) Wieland has shown that nitrous fumes behave in many respects as if constituted entirely as nitrous anhydride, N_2O_3 , and has assumed that, in spite of density determinations, a small proportion of this substance must be present in equilibrium with the mixture of nitric oxide and nitrogen peroxide. The experimental evidence on this point has been provided by Le Blanc (*Z. Elektrochem.*, 1906, 12, 544) and the author (A., 1910, ii, 1059).
H. W.

Magneto-chemical Investigation of the Constitutions in Mineral Chemistry. The Acids of Phosphorus. PAUL PASCAL (*Compt. rend.*, 1922, 174, 457—460).—The method of magnetic analysis previously applied to the compounds of sulphur (A., 1921, ii, 692) has now been applied to compounds of phosphorus. The molecular susceptibilities of a number of phosphorus compounds are given. In all its incompletely saturated derivatives the diamagnetism of phosphorus or its oxy-groups is greatly weakened, and phosphorus and its oxy-radicles possess a constant magnetic individuality. In its saturated derivatives, phosphorus has a different but constant magnetic susceptibility. The results accord perfectly with the rational formulæ $\text{PO}(\text{OH})_3$, and $\text{RPO}(\text{OH})_2$ for the phosphoric and phosphinic acids. W. G.

The Use as a Drying Agent of Phosphoric Oxide Treated with Ozone. JOHN JOB MANLEY (T., 1922, 121, 331—337).

Atomic Weight of Boron. G. P. BAXTER and A. F. SCOTT (*Science*, 1921, 54, 524—525).—Boron was obtained by reduction of boric oxide with excess of magnesium, and extraction with either hydrochloric or hydrobromic acid. It was then converted into the haloid by passing over it at 700° either dry chlorine or helium saturated with bromine nearly at the boiling point of the latter. The excess of halogen having been removed, the haloids were repeatedly distilled with the use of Hempel fractionating columns in sealed, all-glass vessels, with the exclusion of air. Analysis was effected by comparison with silver in the usual way, and by each method gave a result of 10.83 ± 0.01 . This indicates the proportion of the heavier isotope to be about five times that of the lighter, a result which is more in accord with the observations of Aston (A., 1920, ii, 718) than the previous determination (10.900) by Smith and van Haagen (A., 1920, ii, 247). A. A. E.

Researches on the Chemistry of Coal. II. The Resinic Constituents and Coking Propensities of Coals. W. A. BONE, A. R. PEARSON, E. SINKINSON, and W. E. STOCKINGS (*Proc. Roy. Soc.*, 1922, [A], 100, 582—598; cf. *ibid.*, 1917, [A], 96, 119).—Strongly coking bituminous coals may be extracted for prolonged periods by organic resin solvents without impairing their coking propensities. It is shown that the pyridine-chloroform method of extracting coals does not, as stated by Clark and Wheeler (T., 1913, 103, 1706), effect a complete, or nearly complete, separation between the resinous constituents and the degradation products of the celluloses of which coal is conglomerated; but that, on the contrary, it yields an admixture of resins with a predominance of non-resinous substances, which latter are chiefly of cellulosic origin, and have been provisionally designated "humic" substances in the present paper. It is shown that resins may be extracted and isolated in a pure condition from such coals by treatment with pyridine and amyl alcohol, in equal proportions, in an atmosphere of nitrogen, followed by successive treatments with ethyl ether and light petroleum. These resins do not normally much exceed

1% of the coal substance, and, although they may be a contributory, are not usually the chief cause of the coking propensities of coals. A series of non-resinous substances, amounting in many cases to as much as 4%, may be extracted from strong coking coals by a special pyridine or pyridine-amyl alcohol treatment. These substances for the most part are insoluble in ether but soluble in chloroform. On heating these substances out of contact with air, a strong exothermic reaction takes place between 275° and 375° which is accompanied by the elimination of water. This reaction is reminiscent of the behaviour of cellulose and shows these substances to be chiefly cellulosic in type and origin. The coking propensities of coals are principally due to the presence, or the formation in them by heat, of such non-resinous substances of cellulosic origin, the fusion temperatures of which are below those at which they undergo rapid decomposition. The still more complex substances, also of cellulosic origin, which constitute the main portion of the coal substance, but which decompose without fusion, have little or no direct influence on its coking propensities.

J. F. S.

Modifications of Silicon. Solubility of Silicon in Hydrofluoric Acid. WILHELM MANCHOT and HERBERT FUNK (*Z. anorg. Chem.*, 1922, 120, 277—299).—Specimens of silicon obtained from an aluminium regulus have been examined and the influences of (1) temperature of the fusion, (2) concentration of silicon, and (3) rate of cooling investigated. The temperature of the fused mass was varied from 900° to 1650° and was found to have but little effect. The concentration of silicon could be varied from 0.25 to 10% without much effect, but a higher concentration (>10%) favoured the formation of crystals. Rapid cooling of the regulus gave a greyish-black, amorphous (no crystalline form could be detected at 960 magnification) silicon (d 2.23) which reacted briskly with hydrofluoric acid with evolution of hydrogen leaving a brown, amorphous residue (d 2.20) apparently insoluble in hydrofluoric acid. This brown variety was found to be very active—reacting violently with fuming nitric acid, sodium hydroxide, chlorine, and bromine at ordinary temperatures. This reactivity was found to be due to adsorbed hydrogen; after removing the hydrogen it reacted with hydrofluoric acid and behaved in the same way as the greyish-black, amorphous form. On cooling the regulus slowly, crystalline silicon was obtained (d 2.30); this form reacted but slowly with hydrofluoric acid. The aluminium could be replaced by a silver regulus. Silicon of 99% solubility in hydrofluoric acid as claimed by Moissan and Siemens (*A.*, 1904, ii, 560) could not be prepared. Silicon in the mixture (silicon and silica) was estimated by the volume of hydrogen liberated from potassium hydroxide solution. Silicon completely resistant towards hydrofluoric acid could not be prepared. Ordinary silicon heated at 2000° and suddenly cooled behaves in the same way as when slowly cooled. This leads the author to believe that the various forms obtained from the reguli are not allotropic modifications. The

sudden cooling of the solution and solidification of the metal solvent gives extremely fine particles ($d=1\mu$); the reactivity is ascribed to this fineness and not to a new form. Such a degree of fineness could not be obtained mechanically.

W. T.

Formation of Potassium Perchlorate from Potassium Chlorate. VICTOR LENHER, HOSMER W. STONE, and HELEN H. SKINNER (*J. Amer. Chem. Soc.*, 1922, **44**, 143—144).—The formation of potassium perchlorate from potassium chlorate by the action of acids has been investigated. On treating potassium chlorate with sulphuric acid, taking care to add the acid slowly and keeping the mixture cold until the yellow colour has disappeared, a yield of 11% of perchlorate is obtained. Evaporation to dryness of chlorate with nitric acid of various concentrations on a steam-bath gives a yield of 30% of perchlorate; fuming nitric acid yields no perchlorate under the same conditions. A 15% yield of perchlorate is obtained when potassium chlorate is boiled with 85% phosphoric acid. Potassium chlorate and chromium trioxide boiled with just sufficient water to maintain a solution give 11% of perchlorate. Chloric acid gives no perchlorate with potassium perchlorate. The following acids have no action of potassium chlorate even at the boiling point: saturated oxalic acid solution, 25% tartaric acid, acetic acid both glacial and dilute, 50% chloroacetic acid, 25% lactic acid, 50% arsenic acid, 20% permanganic and persulphuric acid. Formic, trichloroacetic, hydrofluoric, and hydrochloric acids decompose potassium chlorate without producing perchlorate.

J. F. S.

Preparation and Constitution of a Double Potassium Ammonium Orthophosphate. R. M. CORELLI (*Gazzetta*, 1921, **51**, ii, 380—385).—Attempts to prepare a salt analogous to sodium ammonium hydrogen orthophosphate by the interaction of dipotassium hydrogen phosphate and ammonium chloride or phosphate result in the elimination of ammonia and formation of potassium dihydrogen phosphate. This extreme instability indicates the compound formed to be a tertiary phosphate, and *potassium di-ammonium phosphate*, $K(NH_4)_2PO_4 \cdot 4H_2O$, may be obtained by passing ammonia into aqueous potassium dihydrogen phosphate solution cooled in ice and salt, filtering the solution quickly in an atmosphere of ammonia and pressing the crystals between filter-paper. The compound deliquesces and loses ammonia in the air, but may be preserved in sealed tubes.

T. H. P.

Crystal Structures of Potassium and Ammonium Stannichlorides. ROSCOE G. DICKINSON (*J. Amer. Chem. Soc.*, 1922, **44**, 276—288).—The crystal structures of potassium and ammonium stannichlorides have been determined from measurements of the X-ray reflection spectra and from unsymmetrical Laue photographs. The structure is shown to be similar for the two salts and of the calcium fluoride type. It has been found possible to carry out the analyses without making any quantitative assumption concerning "normal decline" of intensity, and with only

rough assumptions as to the relative reflecting powers of the atoms. The results lead to the conclusion that six equivalent chlorine atoms are grouped about each tin atom; and the structure is accordingly regarded as built up of SnCl_6'' ions and potassium or ammonium ions. The SnCl_6'' ion occupies the positions filled by calcium in fluorspar and the potassium or ammonium ions occupy the positions of the fluorine. It is shown that the general structure and cleavage of the crystals investigated are similar to those of fluorspar. The following distances between the atoms are recorded: potassium stannichloride, between tin and chlorine, 2.44 Å.U., tin and potassium, 4.31 Å.U., potassium and chlorine, 3.52 Å.U., ammonium stannichloride, tin and chlorine, 2.46 Å.U., nitrogen and tin, 4.35 Å.U., and nitrogen and chlorine, 3.55 Å.U. J. F. S.

Simple Method for the Preparation of Sodium Hydroxide Free from Carbon Dioxide. JACOB CORNOG (*J. Amer. Chem. Soc.*, 1921, 43, 2573—2574).—Solutions of sodium hydroxide free from carbonate may be prepared as follows. Distilled water contained in an Erlenmeyer flask is boiled to remove carbon dioxide, after which, when the water has cooled sufficiently, a layer of ethyl ether 3—4 cm. deep is placed on the water. Pieces of metallic sodium, not exceeding 1 cm. in diameter, are dropped into the flask. They fall no further than the ether, where they remain suspended and are slowly attacked by the water dissolved in the ether and the sodium hydroxide passes into the water. After the desired quantity of sodium has reacted, the larger portion of the ether is pipetted off and the last traces are removed by boiling the solution. It is shown that there is no danger of fire if the depth of the ether is great enough to prevent the suspended sodium from being simultaneously in contact with the air and water. With the care usually employed in working with ether and by keeping the ether layer from three to four times as thick as the diameter of the pieces of sodium added, the reaction may be performed easily and without danger. A further advantage of using ether is that the oil adhering to the sodium is removed with the ether. The product gives no precipitate with solutions of barium hydroxide. J. F. S.

Sodium Hyposulphite. FREDERICK W. HEYL and FRANK E. GREER (*Amer. J. Pharm.*, 1922, 94, 80—92).—The most satisfactory laboratory method for the production of sodium hyposulphite is by the action of sodium formaldehydesulphoxylate on sodium hydrogen sulphite. The former substance may be prepared by the method of D.R.-P. 256460, or more conveniently by the reduction of commercial "hydrosulphite" with zinc dust and zinc oxide in presence of formaldehyde solution, and recrystallising from water at a temperature not exceeding 70° the crystals first obtained. The purity of the product, which approximates to 100%, is best determined by direct titration of a hot solution with standard solution of methylene-blue. Sodium formaldehydesulphoxylate is soluble in glycerol to the extent of about 74 grams in

100 c.c. Administered intravenously, it apparently has no toxic action on white rats. It was not found possible to prepare analytically pure anhydrous sodium hyposulphite even by the method from the sulphoxylate indicated above, and salting out the product with strong brine, although a purity of 97.7% is claimed in U.S. Pat. 990457. The yields obtained amounted to 55–60% of the theoretical with a purity of 80–85%, and neither by recrystallisation nor salting out from air-free aqueous solutions in an inert atmosphere could the salt be further purified. Unlike the sulphoxylate, sodium hyposulphite and also its decomposition products are toxic in doses of about 200 mg. upwards per kilo. of body weight.

G. F. M.

The Density of Cæsium at the Absolute Zero. W. HERZ (*Z. anorg. Chem.*, 1921, **120**, 159).—In a previous paper (A., 1919, ii, 220), it was overlooked that, in calculating the density of a number of elements at the absolute zero, the value arrived at for cæsium, 1.791, is lower than the density at higher temperatures. The equation used cannot therefore be applicable to cæsium. A value harmonising better with that of the other alkali metals is obtained by taking the density at 27° as the density at the melting point d_m , and applying the relation $d_0 : d_m = 1.21$, when the value 2.222 is obtained, and the corresponding atomic volume 59.77. A correction must be made in a subsequent paper in which the atomic volume at absolute zero is used for calculating molecular volumes of cæsium salts (Lorenz and Herz, A., 1921, ii, 536).

E. H. R.

Highly Basic Lithium Salts of Weak Inorganic Acids. ARTHUR ROSENHEIM and WERNER REGLIN (*Z. anorg. Chem.*, 1921, **120**, 103–119).—It has been shown that some highly basic salts of lithium, such as those of antimonie and periodic acids, have semi-colloidal properties (A., 1918, ii, 194 and 1919, ii, 508). Those lithium salts which are characterised by low solubility and negative temperature coefficient of solubility are generally those of weak acids which may be expected to form polymerised or associated molecules. A number of such lithium salts, including phosphite, hypophosphate, phosphate, vanadate, molybdate, tungstate, borate, and carbonate, have been examined with respect to their state of aggregation.

Dilithium hydrogen phosphite, $\text{Li}_2\text{HPO}_3 \cdot \text{H}_2\text{O}$, has a negative temperature coefficient of solubility in water, which ranges from 9.07 grams at 0° to 4.24 grams at 98° of anhydrous salt per 100 grams of solution, the solid phase being the monohydrate throughout. The hydrogen-ion concentration at 20° in a normal solution is $7.34 \cdot 10^{-9}$ ($P_H = 8.14$). The equivalent conductivity at dilutions from $v = 32$ to $v = 1024$ was measured, and also the freezing-point depression. The values for the latter do not indicate association in solution.

Lithium hypophosphate, $\text{Li}_2\text{PO}_3 \cdot 3.5\text{H}_2\text{O}$, does not form a clear, aqueous solution and its solubility would not be determined directly with accuracy. An estimate of the solubility was made from the conductivity, making use of known data to calculate the migration

velocities of the ions. The value obtained for the solubility was 0.1267 gram of Li_3PO_3 per litre at 25° , a value much lower than the apparent solubility determined directly. Whether the apparently colloidal solution is really colloidal or a non-colloidal suspension cannot be stated.

Lithium phosphate was obtained in the form of a dihydrate, $\text{Li}_3\text{PO}_4 \cdot 2\text{H}_2\text{O}$, which at 60° is slowly converted into $\text{Li}_3\text{PO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$. It also forms a cloudy solution; the solubility, determined by the conductivity method, is 0.297 gram of Li_3PO_4 per litre at 25° . Two compounds of lithium with vanadic acid were obtained. The first, $4\text{Li}_2\text{O} \cdot \text{V}_2\text{O}_5 \cdot 4\text{H}_2\text{O}$, is precipitated when a solution of vanadium pentoxide in a large excess of a saturated solution of lithium hydroxide is heated. It appears to have a negative temperature coefficient of solubility, but at low temperatures when stirred with water it changes into $3\text{Li}_2\text{O} \cdot \text{V}_2\text{O}_5 \cdot 18\text{H}_2\text{O}$ or $\text{Li}_3\text{VO}_4 \cdot 9\text{H}_2\text{O}$, transparent, rhombohedral needles. This appears to be identical with Ditte's $4\text{Li}_2\text{O} \cdot \text{V}_2\text{O}_5 \cdot 14\text{H}_2\text{O}$ (A., 1887, 705). The solubility coefficient of $\text{Li}_3\text{VO}_4 \cdot 9\text{H}_2\text{O}$ is positive, the maximum solubility being 6.25 gram of Li_3VO_4 per 100 grams of solution at 35.2° ; above this temperature, the monohydrate is stable and the temperature coefficient is negative.

Lithium molybdenate was found to have the composition $4\text{Li}_2\text{MoO}_4 \cdot 3\text{H}_2\text{O}$, not $5\text{Li}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ (Wempe, A., 1913, ii, 59). Its solubility at 25° is 44.81 grams of Li_2MoO_4 per 100 grams of solution and the temperature coefficient is small and negative. The tungstate has a similar composition, $4\text{Li}_2\text{WO}_4 \cdot 3\text{H}_2\text{O}$, and its solution is rapidly decomposed by atmospheric carbon dioxide.

Lithium borate, $\text{LiBO}_2 \cdot 8\text{H}_2\text{O}$, has a positive solubility coefficient and shows normal depression of freezing point in solution.

Lithium carbonate when boiled in aqueous solution loses carbon dioxide, and finally a solution is obtained containing $\text{LiOH} : \text{Li}_2\text{CO}_3 = 1 : 1$ (approx.). It follows that the solubility determinations of Bewad (*J. Russ. Phys. Chem. Soc.*, 1884, 16, 591) are not trustworthy at higher temperatures.

E. H. R

Crystallographic and Atomic Symmetries of Ammonium Chloride. RALPH W. G. WYCKOFF (*Amer. J. Sci.*, 1922, [v], 3, 177—183).—The symmetry of ammonium chloride as indicated by the development of the external faces and by etch-figures on the faces is that of the enantiomorphous hemihedral class, whilst from the X-ray data, both with the spectrometer and the powder method, the internal structure is tetrahedral cubic with one chemical molecule to the unit cell. The latter is based on the ratio $n^3/m=1$, where n is the order of the reflection and m the number of molecules in the unit cell. Several more complex enantiomorphous groupings with several molecules in the unit cell are considered; but from a chemical point of view these are highly improbable, since they collected all the ammonium groups about one point in the unit and all the chlorine atoms about another point. The crystallographic observations are therefore regarded as requiring revision.

L. J. S.

Decomposition and Stabilisation of Ammonium Nitrate in Presence of Oxidisable Material. ALEXANDER FINDLAY and CYRIL ROSEBOURNE (*J. Soc. Chem. Ind.*, 1922, **41**, 58—59r).—The stability of mixtures of ammonium nitrate with small quantities of woodmeal and with starch was investigated by heating the mixture contained in a test-tube, provided with a capillary delivery tube leading below the surface of mercury to a graduated gas burette. The tube was exhausted and placed in a boiling water-bath. A mixture of 23.75 grams of ammonium nitrate and 1.25 grams of woodmeal gave 1.5 c.c. of a mixture of carbon dioxide (33.1%) and nitrogen (66.9%) in twenty-four hours, and 47.1 c.c. in fifty days. A mixture of the same amount of nitrate and 2 grams of soluble starch gave in one day 46.5 c.c. of a gas having the composition CO_2 27.04%, N_2 72.4%, and CO 0.56%. After some days, the rate of evolution slowed down considerably, and eventually the gas evolved appeared to consist of nitrogen only. In both cases, the addition of 0.25 gram of carbamide to the mixtures proved a most effective stabiliser, no appreciable evolution of gas occurring with the woodmeal mixture in a period of thirty-five days. Diphenylamine and phenyl benzyl ether also act as stabilisers, but they are not suitable for the purpose, owing to subsidiary reactions occurring between these substances and the ammonium nitrate. G. F. M.

Solubility of Silver Chloride in Dilute Chloride Solutions and the Existence of Complex Argentichloride Ions. II. GEORGE SHANNON FORBES and HARRIET ISABELLE COLE (*J. Amer. Chem. Soc.*, 1921, **43**, 2492—2497; cf. A., 1912, ii, 49).—To solutions of sodium, ammonium, hydrogen, rubidium, calcium, barium, strontium, and potassium chloride of concentrations not greater than 0.1*N* a 0.001*N*-solution of silver chloride was added at 25° until an incipient precipitation was observed. The total silver then in solution was invariably given by $2 \times 10^{-10}(\text{MCl}) + 3.4 \times 10^{-5}(\text{MCl}) + b$, where 2×10^{-10} is the solubility product of silver chloride, 3.4×10^{-5} a constant possibly connected with a complex of the type AgCl_2 , and b is apparently the constant sum of silver chloride in precipitate, in dispersion, and in dissolved molecules. The great difference between this observed total and the minute silver content of filtrates from large precipitates of silver chloride is discussed. It is shown that silver chloride should be most insoluble at 25° in 0.0025*N*-chloride solutions. J. F. S.

Crystal Structure of Silver Oxide. RALPH W. G. WYCKOFF (*Amer. J. Sci.*, 1922, [v], **3**, 184—188).—Silver oxide (Ag_2O), which crystallises as small, isotropic octahedra, was examined by the X-ray powder method, and the type of cubic structure deduced is the same as that assigned to cuprous oxide. The unit cube, containing two molecules of Ag_2O , has an edge of 4.768 Å.U.

L. J. S.

Vapour Pressure of Metallic Calcium. NORMAN B. PILLING (*Physical Rev.*, 1921, **18**, 362—368; cf. Langmuir, *ibid.*, 1913, **2**, 329; 1914, **4**, 377).—The vapour pressure of calcium was computed

with the aid of Langmuir's expression, which requires measurement of the rate of evaporation of a certain weight of metal of known surface area at constant temperature and in a vacuum. When the coefficient of reflection, s , of the calcium molecules from the enveloping walls is taken into consideration, the expression becomes $m = \sqrt{M/2\pi RT} \cdot p(1-s)$, where M is the molecular weight of the vapour, assumed to be monatomic, R the gas constant, T the absolute temperature, p the pressure in dynes per sq. cm., and m the rate of evaporation in grams per sq. cm. per second. The value of s was determined by control experiments with zinc and cadmium, the vapour pressures of these metals in the solid state being known (Egerton, *Phil. Mag.*, 1917, [vi], **33**, 33). Vapour-pressure curves are given for liquid and solid calcium, the vapour pressure at the melting point, 805° , being 2.0 mm. and the b. p./760 mm. being computed to be 1240° . The calcium employed contained 1.62% of magnesium, 0.009% of iron, and 1.25% of calcium chloride. A. A. E.

A Simple Process for Obtaining Crystallised Gypsum.

L. BOURGEOIS (*Bull. Soc. chim.*, 1922, [iv], **31**, 160—161).—Three volumes of nitric acid are diluted with one volume of water and this solution is saturated at just below its boiling point with calcium sulphate. The clear liquid is decanted off and allowed to cool. After eight days, crystals of gypsum begin to appear. W. G.

Some Compounds in the System $\text{CaO-P}_2\text{O}_5$ and their Relation to Basic Slag. TH. DIECKMANN and ED. HOUDRE-MONT (*Z. anorg. Chem.*, 1921, **120**, 129—149).—A number of compounds of calcium and phosphoric acid have been prepared and examined with respect to their physical properties and solubility in citric acid with the object of throwing some light on the constituents of basic slag. The solubility of tricalcium phosphate, $\text{Ca}_3(\text{PO}_4)_2$, in 2% citric acid was found to be affected by the temperature to which it was heated. Starting at 94.6%, it fell to 60% as the temperature was raised to 540° , and then rose steadily to 96.2% at the melting point. The composition of the insoluble residue varied in a similar manner, and at the point of minimum solubility had the composition of oxyapatite, $3(\text{Ca}_3\text{P}_2\text{O}_8)\cdot\text{CaO}$. Further heating to higher temperatures appeared to reconvert this into the normal phosphate. The melting point of tricalcium phosphate is 1670° . Oxyapatite was prepared from a mixture of tricalcium phosphate and calcium carbonate. Formation of the compound was complete at about 1450° , and its low solubility in citric acid, about 55%, confirmed the opinion that this compound is formed during the heating of tricalcium phosphate alone. By melting the oxyapatite with the calculated quantity of insoluble pyrophosphate according to the equation $3(\text{Ca}_3\text{P}_2\text{O}_8)\cdot\text{CaO} + 2\text{CaOP}_2\text{O}_5 = 4\text{Ca}_3(\text{PO}_4)_2$, a highly soluble tricalcium phosphate was obtained. The melting point of oxyapatite is 1540° , d 2.99; it crystallises in doubly refracting needles.

Tetracalcium phosphate, $4\text{CaO}\cdot\text{P}_2\text{O}_5$, has a high solubility in

citric acid, but if heated for some time at 1000° the solubility falls to about 25% and the insoluble residue has the composition of oxyapatite. This observation explains earlier statements that tetracalcium phosphate has a low solubility. Its m. p. is 1630° . It is decomposed by boiling water into lime and oxyapatite. Calcium pyrophosphate is practically insoluble in citric acid; m. p. 1230° , d 3.09. Calcium metaphosphate is also insoluble in citric acid. The molten substance when cooled forms a glass and is difficult to crystallise. The melting point is between 970° and 980° , d 2.82 (crystals) and 2.65 (amorphous). Calcium silicophosphate, $5\text{CaO} \cdot \text{P}_2\text{O}_5 \cdot \text{SiO}_2$, has the same solubility in citric acid as tricalcium phosphate, and is probably therefore a compound of the latter with the normal orthosilicate, $\text{Ca}_3(\text{P}_2\text{O}_4)_2 \cdot 2\text{CaO} \cdot \text{SiO}_2$. Its melting point is $1760\text{--}1780^{\circ}$. Synthetic fluorapatite, $\text{CaF}_2 \cdot 3\text{Ca}_3(\text{PO}_4)_2$, has a low solubility, about 10%, in citric acid; m. p. $1630\text{--}1650^{\circ}$, d 3.18. The above observations show the importance of adding sufficient silica to basic slag to neutralise any excess of lime if a soluble phosphate is to be obtained, since the silicophosphate, unlike tetracalcium phosphate, does not decompose during slow cooling to form insoluble oxyapatite. [See *J. Soc. Chem. Ind.*, 1922, April.]

E. H. R.

Calcium Silicides. LOTHAR WÖHLER and F. MÜLLER (*Z. anorg. Chem.*, 1921, **120**, 49—70).—It has been shown by a number of workers that, besides the calcium silicide, CaSi_2 , corresponding with ordinary calcium carbide, a second calcium silicide exists, but various formulæ have been ascribed to it, for example, Ca_3Si_4 (Hackspill, A., 1908, ii, 589) and $\text{Ca}_{11}\text{Si}_{10}$ (Kolb and Formhals, A., 1910, ii, 35). It is now shown that the second compound is calcium monosilicide, CaSi or Ca_2Si_2 . The compound is formed free from disilicide when a mixture of calcium and silicon in atomic proportions or with excess of silicon up to 100% is heated in a magnesia boat in an atmosphere of carbon dioxide at 1050° . After a quarter of a minute, a violent reaction takes place and the mass becomes incandescent. It must then be cooled rapidly. The mass breaks up readily into small, lustrous, metallic leaflets and larger crystals. The density of the substance, containing a small proportion of uncombined silicon, is 2.346. By dilute acids it is attacked readily with evolution of a spontaneously inflammable silicon hydride and formation of hydrated silica. It is only slowly attacked by concentrated acids, with evolution of hydrogen, and by cold water, more readily by warm water and by dilute ammonia. By contrast, the disilicide dissolves in hydrochloric acid with formation of yellow silicone and without production of spontaneously inflammable silicon hydride. When, in the preparation of the substance, so large an excess of silicon is used as to prevent the necessary rise in temperature during the reaction, a mixture of mono- and di-silicides is formed. From this it was concluded that the monosilicide is formed endothermically from the disilicide. This was confirmed by an experimental determination of their heats of combustion and calculation of their heats of

formation, which were found to be, for Ca_2Si_2 , +166.3 cal. and for CaSi_2 , +208.7 cal.

Pure calcium disilicide can be prepared by heating the monosilicide at $1000\text{--}1010^\circ$ in a current of hydrogen. If excess of silicon is present, this reacts with the calcium hydride formed to give a further quantity of the disilicide. The pure disilicide evolves only hydrogen and no silicon hydride with dilute acid. The disilicide is stable at least up to 1050° , for at this temperature the monosilicide, in absence of air, dissociates with development of heat into disilicide and calcium.

The silicon hydride formed by the action of dilute acids on the monosilicide is probably silicoethylene formed according to the equation $\text{Ca}:\text{Si}:\text{Si}:\text{Ca} + 4\text{HCl} = 2\text{CaCl}_2 + \text{H}_2:\text{Si}:\text{Si}:\text{H}_2$. E. H. R.

System Ammonia-Magnesium-Mercury. Formation of Magnesium Hexammoniate. ALBERT G. LOOMIS (*J. Amer. Chem. Soc.*, 1922, 44, 8—19).—When dilute magnesium amalgam is brought into contact with ammonia, a solid phase separates slowly which has a bright metallic lustre and is decomposed on exposure to air or by reducing the ammonia pressure to below two atmospheres. If the magnesium amalgam contains more than 0.15% of magnesium, the whole becomes completely solid with excess of ammonia, whilst richer magnesium amalgams become solid if more ammonia than 0.035 mol. per 100 grams of amalgam is added. The determination of the composition of the solid phase has been attempted. It is shown that magnesium dissolves in mercury to the extent of 0.323% at 23° . The vapour pressure of the system at 0° has been studied by removing measured volumes of ammonia. The vapour-pressure curve shows a gradually decreasing ammonia pressure as ammonia is removed from which it is concluded that in addition to the vapour phase there are only two phases present, one at least of which is of variable composition. The ratio of ammonia to magnesium has been determined in the crystals and the formula $\text{Mg}(\text{NH}_3)_6\text{Hg}_2$ given to the compound. The solubility of the crystals at 22.4° in pure mercury has been determined and is represented by 0.00593 gram of magnesium per 100 grams of mercury. The ammonia pressure of the crystals has been measured at 22.4° . From the experimental results the value of x in the above formula is calculated as 17.5—18.4. From the large value thus obtained for the amount of mercury, it is concluded that solid solutions are formed by the molecular compound, magnesium hexammoniate, with the excess of mercury present. J. F. S.

Variously Coloured Modifications of Colloidal Copper. C. PAAL and HERMANN STEYER (*Kolloid Z.*, 1922, 30, 88—97).—Colloidal copper exists in the following differently coloured modifications: reddish-brown (Lottermoser, A., 1899, ii, 558), blue (Gutbier, A., 1903, ii, 81; 1905, ii, 327), ruby-red (Paal and Leuze, A., 1906, ii, 356), brown (Billitzer, A., 1902, ii, 454), olive (Ehrenhaft, *Anz. Wiener Akad.*, 1902, 39, 241), and green (Paal and Leuze, *loc. cit.*). The ruby-red variety exists in two forms known as the α - and β -modification, respectively. The

present paper gives a description of the preparation and properties of the α - and β -ruby-red and the blue modifications of copper hydrosol by the use of sodium lysalbate and protalbate and the free acids. The method of converting the ruby-red variety into the blue variety is described. J. F. S.

German Silver. WALTER VOIGT (*Z. anorg. Chem.*, 1922, 120, 309—319).—The equilibrium diagram of the copper-nickel-zinc alloys was investigated by Tafel (A., 1908, ii, 846). All alloys except those rich in zinc were easily worked. The electric resistance was determined after heating the wires to the temperatures of minimum resistance (cf. Credner, A., 1913, ii, 280). If the ratio of copper to nickel remains constant, then increasing amounts of zinc have but little effect. The resistance increases much more rapidly on increasing the nickel content. As regards the thermoelectric effect, if the ratio copper: nickel is constant then increasing amounts of zinc increase it. Increase in the ratio of zinc to copper alone decrease it; a considerable decrease was observed on increasing the nickel content, the copper-zinc ratio being kept constant. In general, the surface of the thermoelectric effect (on diagram) runs parallel to that of the reciprocal of resistance, that is, conductivity. All the alloys showed a solution tension approximately equal to that of copper. They all precipitated copper gradually from a solution of copper sulphate. The passivity of nickels protects the alloy only when its content is 31 mol.% and 43 mol.%. W. T.

Crystal Structures of the Cuprous Haloids. RALPH W. G. WYCKOFF and EUGEN POSNJAK (*J. Amer. Chem. Soc.*, 1922, 44, 30—36).—The crystal structure of cuprous chloride, bromide, and iodide have been deduced from the examination of X-ray reflection spectra using films of powder. They are shown to possess the zinc sulphide arrangement and the length of the side of the unit cube is 5.49, 5.82, and 6.10 Å.U. respectively. J. F. S.

Separation of Isotopes. Theory of Resolution of Isotopic Mixtures by Diffusion and Similar Processes. Experimental Separation of Mercury by Evaporation in a Vacuum. ROBERT S. MULLIKEN and WILLIAM D. HARKINS (*J. Amer. Chem. Soc.*, 1922, 44, 37—65).—The various phenomena of diffusion and effusion in gases are discussed in connexion with the separation of isotopes. It is shown that for the processes of molecular diffusion through a porous membrane, molecular effusion, and non-equilibrium evaporation, the relative rates of escape of isotopes are proportional to their respective molecular fractions and inversely proportional to the square roots of their molecular weights. In all the above processes, the molecules move independently. As collisions between molecules become more frequent, molecular diffusion passes into capillary transpiration, or mass motion, and non-equilibrium evaporation passes into ordinary distillation. In either case practically no separation of isotopes occurs. If a gaseous mixture of isotopes diffuses into another gas, with no convective mixing, the diffusion coefficients of any two isotopes

are probably in the ratio $x_1/x_2 \sqrt{M_2/M_1}$, where c is a variable quantity usually in the neighbourhood of 1 or 2. The theory is complicated, but indicates a higher rate of separation than for molecular diffusion and evaporation, for which c is always 2. A further possible method of separation is that of "initial diffusion," which would take advantage of the fact that the front layer of molecules diffusing into a gas or a vacuum should be enriched in the lighter isotopes. The theory of non-equilibrium evaporation is discussed. If the vapour pressures of isotopes when pure are equal, a separation must occur when an isotopic liquid is evaporated in a vacuum in such a way that practically all the evaporating molecules are condensed. For an efficient separation, good mixing must take place in the surface and body of the liquid. This consideration makes the method inapplicable to solids, but solutions or liquid alloys might be used. Simple and closely approximate equations are developed for the change in the molecular fraction of any isotope and for the change of atomic weight in a mixture of any number of isotopes, when the latter is separated into fractions by a diffusion or evaporation process. A number of equations are evolved which apply to various sets of conditions. The rate of separation of two isotopes as measured by the change of atomic weight for a given operation is proportional to the square of the interval between the molecular weights of the two isotopes, inversely proportional to the ordinary molecular weight and proportional to the product of the molecular fractions of the isotopes. Similar relations also hold for a mixture of any number of isotopes. For a given element, the separation coefficient (equal to the decrease of atomic weight of the isotopic mixture for the first small portion of diffusate or condensate) is inversely proportional to the molecular weight of the compound in which that element is combined. A table of separation coefficients is given, calculated from atomic weights and the results of positive ray analysis. The value given for mercury has been calculated from the present experimental data. Most of the equations developed apply to the separation by diffusion of any gaseous mixture, whether it consists of isotopes or not. In systematic fractionation the diffusate or condensate, being formed at any time has an atomic weight less than that of the corresponding residue, which becomes denser as the diffusion proceeds, by a constant amount. Thus the enrichment of the light fraction is a maximum at the beginning. The atomic weight of the residue increases indefinitely, however, in proportion as the logarithm of its quantity decreases, whilst at the same time the atomic weight of the total diffusate approaches that of the original material. The use of cuts of two gives equal and opposite enrichments for the two fractions, equal in magnitude to 0.693 times the separation coefficient, or initial enrichment for the light fraction. Formulæ are given for calculating the proportions and numbers of isotopic species in compounds containing several isotopic atomic species. Zinc chloride contains twelve molecular isotopes, and if, for example, tin has six isotopes, the compound SnCl_4 is a mixture of thirty and SnCl_2Br_2 of fifty-four different molecular isotopes.

The existence of many new isomerides due to isotopism is pointed out. Experimental work on the partial separation of mercury into isotopes by non-equilibrium evaporation is described. Calculation based on data for a very efficient evaporation gives the value 0.0057 for the separation coefficient of mercury. The results agree with those of Brönsted and Hevesy (this vol., ii, 149), but the efficiency of the present separation is better. By making four successive cuts of approximately two, on both light and heavy fractions, a much larger decrease, of 64 parts per million, or 0.013 unit of atomic weight, has been obtained on the lightest fraction and a corresponding increase of 69 parts per million or 0.014 unit on the extreme heavy fraction. The results are in complete agreement with the theory developed. The total difference in density between the extreme fractions is 133 parts per million and a difference of 0.027 unit in the atomic weight. Data are given which show that a slight separation of isotopes occurs during an ordinary distillation under reduced pressure. A classification of the possible methods of separating isotopes is given in outline.

J. F. S.

Chromates of Thallium. G. CANNERI (*Gazzetta*, 1922, 52, i, 33—36).—Gröger's method of obtaining basic chromates (A., 1920, ii, 313) yields with thallous salts only normal thallous chromate. The action of dilute acid on thallous chromate yields thallous dichromate, but if a quantity of sulphuric acid insufficient to dissolve the chromate is employed and the boiling solution is filtered and concentrated, the double salt, $\text{TiHCrO}_4 \cdot \text{Ti}_2\text{CrO}_4$, is obtained in cinnabar-red crystals, which often form cruciform twins.

Thallic chromate may be obtained pure as a golden yellow, crystalline powder by dissolving freshly precipitated thallic oxide in excess of chromic anhydride solution.

The *E.M.F.* of the electrode, mercury amalgam|saturated thallous chromate solution varies continuously with the temperature, the curve exhibiting no characteristic point corresponding with the change in colour which thallous chromate crystals undergo when heated at 60°. Similarly the conductivity of saturated thallous chromate shows no sudden variation between 25° and 90°.

T. H. P.

Molecular Condition of Metals Dissolved in Mercury. FRANZ SKAUPY (*Z. Elektrochem.*, 1922, 28, 23—27).—A theoretical discussion on the views which are at present held with regard to the condition of metals dissolved in mercury.

J. F. S.

Atomic Weight of Yttrium. H. C. FOGG and C. JAMES (*J. Amer. Chem. Soc.*, 1922, 44, 307—316).—The atomic weight of yttrium has been determined from the ratio $\text{YtCl}_3 : 3\text{Ag}$, and as a mean of twenty-one analyses the value 89.03 is obtained, the extreme values being 88.97 and 89.08. It is shown that to obtain very pure yttrium it is essential to use, finally, such a method as the cacodylate separation, in which the yttrium is precipitated whilst the more basic cerium earths remain in the form of quite

soluble salts. The material employed in the present work was obtained from Norwegian gadolinite and was submitted to a long process of purification which consisted in converting the oxalates of the rare earths into sulphates and these into bromates which were subjected to fractional crystallisation. A fraction containing yttrium and erbium bromates and practically no holmium was further purified by fractional precipitation as basic nitrate, which was continued until the erbium bands had practically disappeared. Further fractional precipitations by means of sodium nitrite and potassium ferricyanide were also carried out. The oxide obtained at the end of this series of purifications had a pale cream colour, due to a minute trace of praseodymium oxide, and this was removed by converting into acetate and precipitating as cacodylate, which effectually removed the last trace of impurity. The final purification was attempted by precipitation as the dimethyl phosphate, or *p*-sulphobenzoate and also by crystallisation of the acetate. The two former methods effected no purification, but the crystallisation of the acetates concentrated the coloured oxide in the mother-liquors. It was found, however, to be too wasteful and tedious.

J. F. S.

[Revision of the Atomic Weight of Lanthanum.] Correction.

G. P. BAXTER (*J. Amer. Chem. Soc.*, 1922, 44, 328).—A correction of some eight figures in a table contained in a paper by Baxter, Tani, and Chapin on the atomic weight of lanthanum (*A.*, 1921, ii, 454). The following corrected values are given: $\text{LaCl}_3 : 3\text{Ag} = 0.757895$; atomic weight of lanthanum = 138.914 (mean); $\text{LaCl}_3 : 3\text{AgCl} = 0.570413$; atomic weight of lanthanum = 138.913 (mean).

J. F. S.

The Separation of the Rare Earths by Basic Precipitation.

WILHELM PRANDTL and JOHANNA RAUCHENBERGER (*Z. anorg. Chem.*, 1921, 120, 120—128).—In a previous paper (*A.*, 1920, ii, 434) a method was given by which lanthanum could be separated from neodymium and praseodymium by regulated precipitation of the basic chlorides by ammonia in presence of ammonium chloride. The investigation has now been extended to samarium and it is found that the solubility of samaria in ammonia-ammonium chloride solutions at different concentrations and temperatures approximates closely to that of the didymia earths, being generally somewhat lower. Attempts to discover an unknown element between neodymium (atomic number 60) and samarium (62) by fractional precipitation of neodymia and samaria were unsuccessful.

With the object of applying the basic precipitation method more widely, the solubilities of lanthana, praseodymia, neodymia, and samaria were examined in presence of ammonia and ammonium nitrate at different temperatures, by adding the calculated quantity of ammonia, according to the equation $\text{M}'''(\text{NO}_3)_3 + 3\text{NH}_3 + 3\text{H}_2\text{O} \rightleftharpoons \text{M}'''(\text{OH})_3 + 3\text{NH}_4\text{NO}_3$ to the neutral nitrates in 1, 2, 3, 4, and 5*N*-ammonium nitrate solutions, and shaking for a day in a thermostat to attain equilibrium. The biggest separation of the solubility curves was found in 4—5*N*-ammonium nitrate solu-

tion at 100° , instead of $2-3N$ at 50° as in the case of the chlorides. The separation therefore depends on the nature of the anion, showing that it is due to the formation of basic salts of varying composition and solubility. The solubilities were then further examined in presence of an equivalent quantity of magnesium or zinc nitrate under similar conditions. The presence of magnesium offered no advantage for the separation of lanthanum from the other earths, but the solubilities of the last differed more among themselves at high concentrations of ammonium nitrate in presence of magnesium nitrate. The presence of zinc nitrate, however, increased the solubilities markedly, especially that of lanthana, which became four times that of the other earths. The practical application of this result will be reported later. E. H. R.

The Minimum Solubility of Aluminium Hydroxide in Water. A. MASSINK (*Chem. Weekblad*, 1922, 19, 66).—Since the minimum solubility of an amphoteric electrolyte lies at the isoelectric point, which for aluminium hydroxide requires $p_H \pm 7$, waters which are treated with aluminium compounds as coagulants should have a hydrogen-ion concentration close to that value.

S. I. L.

Tempering. A. POUCHOLLE (*Compt. rend.*, 1922, 174, 611—613).—Curves are given showing the variation in length of steel wires as a function of the time of cooling after different treatments. Tempering is characterised by the absence of the transformation point Ar_1 at low temperature. The transformation of γ -iron into α -iron only takes place at the point Ar_2 , that is, at about 200° . Over the temperature range $650-200^{\circ}$, from the point Ar_1 to the point Ar_2 , the curves do not show any angular point. However, invariably in this region, and only in this region, mechanical tensions appear, being manifested by sharp cracks and accompanied by the projection of the thin skin of oxide. The transformation point Ar_1 is lowered if the annealing observed follows a tempering. Similarly, the temperature at which tempering is obtained is lowered by successive temperings. The amplitude of the inflection Ar_1 diminishes (α) by rise in temperature until it becomes nil, (β) by the duration of the heating if the temperature remains constant. W. G.

Reduction of Ferric Chloride. A. PICKLES (*Chem. News*, 1922, 124, 93—94).—When hydrogen is passed through ferric chloride solution containing pieces of fine copper gauze, the ferric chloride is reduced rapidly; varying quantities of cuprous chloride are produced at the same time, and the reaction, which appears to be one of adsorption, is accelerated when the gauze is so arranged that it comes into intimate contact with the hydrogen. W. P. S.

The System Ferric Oxide-Sulphuric Acid-Water. MALCOLM PERCIVAL APPLEBEY and SIDNEY HERBERT WILKES (*T.*, 1922, 121, 337—348).

Examination by the X-Ray Spectrum of Metallic Oxides which are Stable at Red Heat (Prepared by Different Methods and having Different Properties). J. A. HEDVAL (*Z. anorg. Chem.*, 1922, **120**, 327—340).—That the properties of certain oxides prepared by different methods vary as regards colour, density, etc., is generally explained by assuming the existence of allotropic modifications. As this seems highly improbable, these oxides have been examined by the X-ray spectrum. The author shows that identical spectra can only be obtained from the same modification, and vice versa. The following oxides were examined. Ferric oxide (27), ferrosiferic oxide (2), alumina (5), cobaltous oxide (2), cobaltocobaltic oxide (3), nickel oxide (6), magnesia (7), zinc oxide (6), cupric oxide (11), stannic oxide (3). The numbers in brackets give the number of different methods employed in their preparation. No oxide was found to exist in the amorphous form; they were all crystalline or cryptocrystalline. The different preparations of the same oxide gave identical spectra, which were the same as for the minerals. Many oxides were heated at 1150° for some time; this caused a change in some of their properties, but the X-ray spectrum remained unchanged. These changes in properties are not caused by changes in structure. W. T.

Properties of Subsidiary Valency Groups. III. The Preparation, Properties, and Molecular Volume Relationships of the Hydrates and Ammines of Cobalt Fluoride, Bromide, Iodide, Nitrate, Carbonate, and Citrate. GEORGE L. CLARK and HENRY K. BUCKNER (*J. Amer. Chem. Soc.*, 1922, **44**, 230—244; cf. A., 1921, ii, 116).—The preparation and properties of a number of hydrates and ammines of several salts of cobalt are described. *Cobaltous fluoride trihydrate* is prepared by the action of hydrofluoric acid on cobalt carbonate; it forms rose-red crystals which only lose the last of the water at 300° in a current of nitrogen; $d=2.583$. *Cobaltous fluoride hexammine*, prepared by passing dry ammonia over the anhydrous fluoride placed on a balance pan until the requisite weight has been absorbed, which required about twenty-four hours, is a reddish-brown liquid at 25° which solidifies in ice and salt. It is easily soluble in water without hydrolysis, and has $d=1.744$. *Cobaltous fluoride triammine* is produced as a light pinkish-brown compound by keeping the hexammine in dry air; at 40° it loses all its ammonia. *Cobalt citrate tetrammine* is prepared by dissolving cobalt citrate in concentrated ammonia and evaporating spontaneously in a vacuum. It is a pink, very stable, crystalline compound, $d=1.686$. The preparation of many other compounds which have previously been prepared is also described. The molecular volume of all the above-named compounds is tabulated and the apparent volume of the subsidiary group calculated. The present work furnishes further evidence of the existence of cavities in the space lattice of solids, which may hold secondary valency groups, the sizes of the cavities depending on the relative volumes of kation and anion and varying in a parallel fashion with the stability of secondary

valency compounds. The constant relationship of physical properties among the halogens, free and combined, is shown by the linearity of the molecular volumes of the cobalt haloids, as well as by those of numerous other metal haloids, when plotted against the atomic volume of the halogens at the boiling point. Such linearity is not maintained after the formation of hydrates or ammines. The great importance of the percentage contraction in the formation of hydrates, ammines, and polyhaloids is shown by a straight line proportionality to stability. J. F. S.

Electrometric Study of the Hydrolysis of some Complex Cobaltammines under the Action of Barium Hydroxide. PAUL JOB (*Compt. rend.*, 1922, 174, 613—616).—Electrometric titration of complex cobaltammines with barium hydroxide gives a clear indication when the whole of the water of constitution of the cobalt complex has been replaced by the hydroxyl group. This method is capable of much wider application than the electrical conductivity method previously employed (cf. A., 1920, ii, 320). The curve obtained with the sulphatopentammine nitrate is quite regular and shows no inflexion, whereas the curves obtained with roseopentammine chloride and diroseotetrammine chloride both show marked inflexions and indicate when the replacement of the water of constitution is complete. W. G.

The Equilibrium of Tungsten and its Oxides with Hydrogen and Water Vapour; Carbon Monoxide and Carbon Dioxide and Oxygen. J. A. M. v. LIEMPT (*Z. anorg. Chem.*, 1921, 120, 267—276).—The measurements of Chaudron (A., 1920, ii, 379) are used to calculate the equilibrium constants for the reactions $\text{WO}_2 + 2\text{H}_2 \rightleftharpoons \text{W} + 2\text{H}_2\text{O}$; $\text{W}_2\text{O}_5 + \text{H}_2 \rightleftharpoons 2\text{WO}_2 + \text{H}_2\text{O}$; and $2\text{WO}_3 + \text{H}_2 \rightleftharpoons \text{W}_2\text{O}_5 + \text{H}_2\text{O}$; and the equilibrium curves are constructed. Making use of the water-gas constant $p_{\text{CO}} \cdot p_{\text{H}_2\text{O}} / p_{\text{CO}_2} \cdot p_{\text{H}_2}$, the constants for the three reactions $\text{WO}_2 + 2\text{CO} \rightleftharpoons \text{W} + 2\text{CO}_2$; $\text{W}_2\text{O}_5 + \text{CO} \rightleftharpoons 2\text{WO}_2 + \text{CO}_2$; $2\text{WO}_3 + \text{CO} \rightleftharpoons \text{W}_2\text{O}_5 + \text{CO}_2$ are also calculated. Further, from the known thermal dissociation of water, the vapour pressures and heats of formation of the three oxides of tungsten are calculated. The vapour pressures at 1773° are: $\text{WO}_2 \rightleftharpoons \text{W} + \text{O}_2$, $p = 10^{-8.2}$; $2\text{W}_2\text{O}_5 \rightleftharpoons 4\text{WO}_2 + \text{O}_2$, $p = 10^{-8.0}$; $4\text{WO}_3 \rightleftharpoons 2\text{W}_2\text{O}_5 + \text{O}_2$, $p = 10^{-6.7}$ atm. The heats of reaction are: $\text{W} + \text{O}_2 = \text{WO}_2 + 122,800$ cal.; $4\text{WO}_2 + \text{O}_2 = 2\text{W}_2\text{O}_5 + 121,400$ cal.; $2\text{W}_2\text{O}_5 + \text{O}_2 = 4\text{WO}_3 + 123,000$ cal. E. H. R.

Relationships between the Different Oxides of Uranium. PIERRE JOLIBOIS and ROBERT BOSSUET (*Compt. rend.*, 1922, 174, 386—388).—The decomposition of uranium trioxide, when heated in a vacuum is irreversible and yields the oxide U_3O_8 at 502°. The dioxide when heated in oxygen is oxidised very rapidly, the action commencing at about 185°. The only product is the oxide U_3O_8 . When heated in a current of hydrogen, the oxide U_3O_8 shows signs of reduction at 625° and the reduction can be completed at 650°. The only product is the dioxide UO_2 . In three hours at 1000° in a vacuum the oxide U_3O_8 only loses a very small

fraction of its oxygen. To obtain the dioxide by simple dissociation, the oxide U_3O_8 must be calcined at 2000° in a vacuum. W. G.

The Oxides of Uranium. P. LEBEAU (*Compt. rend.*, 1922, 174, 388—391; cf. preceding abstract).—From a consideration of the work done on the oxides of uranium the author concludes that the only oxides which have a definite existence are UO_3 , U_3O_8 , and UO_2 . The so-called black oxides have the composition U_3O_8 . They are stable in air and can be heated at 1000° under atmospheric pressure without decomposition. The green oxides prepared at temperatures below 800° contain varying amounts of uranium trioxide and can undergo change when exposed to moist air, the uranium trioxide present undergoing hydration (cf. Staehling, this vol., ii, 106). W. G.

The Amphoteric Character of Stannic Hydroxide and its Bearing on the Isomerism of the Stannic Acids. GEORGE ERNEST COLLINS and JOHN KERFOOT WOOD (*T.*, 1922, 121, 441—449).

Germanium. III. Germanium Tetrabromide and Germanium Tetrachloride. L. M. DENNIS and F. E. HANCE (*Chem. News*, 1922, 124, 66—69, 82—84; *J. Amer. Chem. Soc.*, 1922, 44, 299; cf. this vol., ii, 150).—When bromine vapour is passed over germanium, prepared by reduction of the dioxide with hydrogen, superficial reaction takes place in the cold, but the most favourable temperature for the reaction is about 220° . The germanium tetrabromide which distils over is best purified from bromine by fractional distillation. It has b. p. 185.9° (corr.) and m. p. 26.1° . It crystallises in small, white, flattened octahedra belonging to the cubic system. The liquid substance can be supercooled considerably, as low as -18° , without crystallising; n_D^{25} , 1.6269; d_{25}^{25} = 3.1315; specific conductivity < 0.000078 reciprocal ohm. It is very sensitive to water and fumes in air. With dry ammonia gas, it forms a white, solid compound.

Germanium tetrachloride was prepared by passing chlorine over germanium. Reaction started at 80° , was rapid at 180° , and at 360° the metal became incandescent. The crude product could not be freed from chlorine by fractional distillation, but the chlorine was removed by passing dry air through the liquid in a suitable apparatus with a condenser to keep back the germanium tetrachloride. The tetrachloride is a colourless, mobile liquid which fumes in air, b. p. 86.5° (corr.), m. p. -49.5° , n_D^{25} , 1.3606, d_{25}^{25} , 1.874. When the liquid is placed in water, reaction takes place slowly with a peculiar crackling sound. E. H. R.

Are Tantalum and Columbium Pentachlorides Conductors of Electricity in the Fused State? WILHELM BILTZ and ARTHUR VOIGT (*Z. anorg. Chem.*, 1921, 120, 71—76).—According to HAMPE (A., 1888, 211), tantalum pentachloride is a good conductor, whilst columbium pentachloride is an insulator. To test the accuracy of these observations the two pentachlorides have been prepared in such a manner that moisture was altogether excluded and the

substance was distilled out of contact with air directly into the conductivity apparatus. The tantalum pentachloride was prepared by passing chlorine over a heated mixture of tantalum pentoxide and sugar carbon and distilling the pentachloride. It formed a snow-white mass of crystals which, when melted at 230—240°, had a conductivity $\kappa=0.30.10^{-6}$. The platinum electrodes were strongly attacked, but the liquid appeared clear and unchanged. Columbium pentachloride was obtained by passing a current of chlorine over columbium sulphide, and, after removal of the sulphur chloride, distilling the columbium pentachloride. It crystallised in long, yellow needles, and its conductivity at 220—235° was $\kappa=0.22.10^{-6}$. The electrodes were attacked also in this case. The two compounds are therefore both insulators of about the same order as the best conductivity water. E. H. R.

Action of Selenium on Gold. H. PÉLABON (*Compt. rend.*, 1922, 174, 391—392).—It is now shown that gold is attacked by selenium if immersed in the latter at a temperature just below its boiling point. The gold fixes some of the selenium, which is only removed with difficulty on heating, and at the same time some of the gold passes into the selenium. The β -selenium previously recorded (this vol., ii, 141) is, therefore, not a pure compound, its low electrical resistance being explained by the presence in it of gold dust held in suspension and by the crystalline configuration of the selenium. W. G.

Preparation of Chloroplatinic Acid by Means of Hydrogen Peroxide. PAUL RUDNICK (*J. Amer. Chem. Soc.*, 1921, 43, 2575—2577).—Details of a method for working up platinum residues are given. The residues from the determination of potassium are extracted with the smallest possible quantity of hot water and filtered hot to remove asbestos and paper fibre. On cooling, pure potassium chloroplatinate is obtained as crystals. The filtrate from the crystals and all other aqueous filtrates are reduced with zinc and hydrochloric acid, and after removing the platinum black the filtrates are exposed to sunlight for several weeks, whereby further quantities of platinum black separate. Alcohol filtrates are treated as above after the alcohol has been expelled. The granular platinum black is washed and suspended in hydrochloric acid, treated with concentrated hydrogen peroxide, and stirred by passing in a stream of hydrogen chloride. The solution of the metal is complete in a few hours. The solution is treated with the necessary amount of potassium chloride and the potassium platinichloride added to the quantity obtained above. The material is stocked in this form. The reagent is prepared by reducing a weighed amount of potassium platinichloride with alkaline sodium formate, washing the platinum black, and dissolving as above. The solution is then made up so that 10 c.c. of the reagent contains 1 gram of metallic platinum. J. F. S.

Mineralogical Chemistry.

The Relation of Chalcedony to other Forms of Silica.

EDWARD W. WASHBURN and LOUIS NAVIAS (*Proc. Nat. Acad. Sci.*, 1922, 8, 1—5).—Chalcedony from Yellowstone Park and black pebble flint from France were examined in comparison with quartz, cristobalite, tridymite, and silica-glass, and determinations are given of d , n , inversion-temperature, coefficient of cubical expansion on either side of the inversion-temperature, and change in volume on inversion. The results suggest that the raw materials consist of quartz, but that after calcination the material is cristobalite. This is fully confirmed by the X-ray spectra. After calcination at 1450° for two hours, the chalcedony gave 99.87% SiO_2 and the flint 99.6% SiO_2 .
L. J. S.

Camsellite, a New Borate Mineral from British Columbia.

H. V. ELLSWORTH and E. POITEVIN (*Trans. R. Soc. Canada*, 1921, Sect. IV, [iii], 15, 1—8).—This is found as white, fibrous masses, intermixed with chrysotile and dolomite, filling shear zones in serpentine near Douglas Lake in the Nicola mining division. The blades give straight extinction with negative elongation and are probably orthorhombic; refractive indices α 1.575, γ 1.649. Analyses I, II, and III are of the camsellite, dolomite, and chrysotile respectively, separated from one another as far as possible; after deducting considerable amounts of impurities these are reduced to Ia, IIa, IIIa.

	SiO_2 .	CO_2 .	B_2O_3 .	Al_2O_3 .	Fe_2O_3 .	FeO.	MnO.	CaO.	MgO.	H_2O (at 110°).	H_2O ($>110^\circ$).	Total.
I.	7.65	5.64	29.07	0.26	0.86	0.95	0.85	3.69	41.72	0.52	9.88	100.12*
II.	1.15	48.10	trace	0.09	—	0.29	0.68	29.05	20.98	0.03	0.69	100.12*
III.	39.95	1.15	trace	0.32	1.44	0.13	0.06	0.75	41.43	1.80	13.04	100.08§
Ia.	—	—	40.40	0.29	0.85	1.28	1.09	—	45.24	0.26	10.55	100.00
IIa.	—	47.25	—	0.09	—	0.30	0.68	29.77	21.50	—	0.41	100.00
IIIa.	40.80	—	—	0.33	1.48	0.13	0.06	—	41.90	1.84	13.35	100.00

* Also Na_2O , K_2O 0.03, NiO trace. † Also MgO (insol.) 1.01. § Also Cr_2O_3 0.01, NiO , B_2O_3 , Na_2O , K_2O traces. ‡ Also Na_2O , K_2O 0.04. || Also Cr_2O_3 0.01.

The formula $2\text{MgO} \cdot \text{B}_2\text{O}_3 \cdot \text{H}_2\text{O}$ given by Ia, corresponds with that of the manganese borate sussexite. Up to 550° only 2.95% H_2O is lost. The mineral readily fuses to a black, opaque globule; and it is soluble in acids.
L. J. S.

Tschermigite (Ammonium-Alum) from Wyoming.

THEODORE ERICKSON (*J. Washington Acad. Sci.*, 1922, 12, 49—54).—Tschermigite occurs as colourless to white, columnar masses and imperfect crystals in brown bituminous shale near Wamsutter, Wyoming. It is optically isotropic, n 1.457, d 1.645. Associated are a pale yellow ammoniacal jarosite (containing $(\text{NH}_4)_2\text{O}$ 1.25%) and crystals of gypsum. Analysis gave (also traces of Fe_2O_3 , CaO , K_2O , Cl):

Al_2O_3 .	$(\text{NH}_4)_2\text{O}$.	Na_2O .	MgO .	SO_3 .	H_2O .	Insol.	Total.
11.57	5.23	0.21	0.13	35.11	47.82	0.06	100.13

At 105°, the mineral fuses in its water of crystallisation and loses three-fourths of its water; nearly all the water is lost at 200°. The ammonium sulphate commences to decompose at 360°, and on ignition the total loss is 88.06%. Sodium alum when ignited gives $\text{Na}_2\text{O} + \text{Al}_2\text{O}_3$ presumably as sodium aluminate.

L. J. S.

Melanovanadite, a New Mineral from Peru. WALDEMAR LINDGREN, L. F. HAMILTON, and CHARLES PALACHE (*Amer. J. Sci.*, 1922, [v], 3, 195—203).—A repetition of the earlier paper (this vol., ii, 155) giving some additional details. The mineral is hygroscopic; material which had been exposed to a moist atmosphere contained up to 16.6% H_2O , of which about 10.5% is lost over sulphuric acid or at 105°. A new analysis of such material gave:

V_2O_5	V_2O_5	CaO	$\text{Fe}_2\text{O}_3 + \text{Al}_2\text{O}_3$	H_2O (above 105°)	Total
33.48	49.38	10.65	1.39	5.90	100.80

The melanovanadite has probably been derived from a vanadium sulphide which is present in the black shale; and by its own alteration it gives rise to pascoite ($\text{Ca}_2\text{V}_6\text{O}_{17}, 11\text{H}_2\text{O}$). A microscopical examination of polished sections of patronite shows that this is a mixture of three minerals which are probably vanadium sulphides. Measurements of the monoclinic crystals of melanovanadite give the axial ratios $a : b : c = 0.4737 : 1 : 0.5815$, $\beta = 88^\circ 37\frac{1}{2}'$.

L. J. S.

Dewindtite, a New Radioactive Mineral. ALFRED SCHOEP (*Compt. rend.*, 1922, 174, 623—625).—This occurs at Kasola, Katanga, Belgian Congo, as a canary-yellow powder intimately mixed with torbernite and a white, powdery material (which resembles impure talc, but contains much alumina—a partial analysis is given). Under the microscope, it is seen to consist of minute scales of rectangular or square outline, $d\ 4.8$. In the closed tube it yields water and becomes brown, returning to yellow on cooling. Before the blowpipe in the oxidising flame it fuses to a black globule, and in the reducing flame yields beads of lead. It is soluble in nitric acid, and decomposed by hydrochloric or sulphuric acid. The mean of several analyses is given under I;

P_2O_5	UO_3	PbO	$\text{Al}_2\text{O}_3(\text{Fe}_2\text{O}_3)$	CaO	MgO	H_2O	Insol.	Total
I. 10.01	55.50	21.74	2.06	1.32	2.75	5.82	0.40	99.60
II. 10.84	60.13	23.55	—	—	—	5.46	—	[99.98]

deducting impurities (represented by the white powdery material), II corresponds with $4\text{PbO}, 8\text{UO}_3, 3\text{P}_2\text{O}_5, 12\text{H}_2\text{O}$.

L. J. S.

Japanese Minerals containing Rare Elements. III. Analyses of Beryl of Naegi, Mino Province. YŪJI SHIBATA and TAKU UEMURA (*J. Chem. Soc. Japan*, 1922, 43, 48—62).—Beryl ($d\ 2.6$ and having no radioactivity) from Naegi in Mino Province gave on analysis:

SiO_2	Fe_2O_3	Al_2O_3	GLO	CaO	MgO	Na_2O	Ignition	Total
60.69	0.77	20.79	11.10	1.00	0.99	1.57	2.75	99.66

No alkali metals other than sodium were detected. The authors deduce the formula, $5\text{GfO} \cdot 2\text{Al}_2\text{O}_3 \cdot 10\text{SiO}_2$, the constitution $\text{Gf}[\text{SiO}_3 \cdot \text{Al}(\text{OH}) \cdot \text{SiO}_3 \cdot \text{Gf} \cdot \text{SiO}_3 \cdot \text{Al} < \frac{\text{SiO}_3}{\text{SiO}_3} > \text{Gf}]_2$ being suggested.

K. K.

Thortveitite, a Silicate of Scandium. JAKOB SCHETELIG (*Norsk Geol. Tidsskr.*, 1922, 6, 233—244).—Thortveitite (A., 1912, ii, 56; 1920, ii, 627) is found in felspar quarries in Setersdalen, Norway, as large, greyish-green, prismatic crystals resembling epidote in appearance. These are monoclinic ($a:b:c=0.7674:1:0.5569$, $\beta=77^\circ 28'$) and usually twinned on $m(110)$; H 6—7, optically negative, α 1.7561, β 1.7926, γ 1.8093 (Na). Analysis I by J. SCHETELIG, II by J. ŠTĚRBA-BÖHM, and III by F. TAUCHERT.

	SiO ₂ .	Sc ₂ O ₃ .	Yt ₂ O ₃ .	(Di,Er) ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	GfO.	Ign.	Total.	d.
I.	42.9	37.6	17.7*		2.1	0.8	—	0.4	100.9	3.57
II.	45.45	42.06	8.89†		2.83	—	0.51	0.54	100.28	3.566
III.	45.55	38.61	10.47‡	4.26§	3.13	—	—	—	100.02	—

* Mol. wt. 270. † Mol. wt. 320. ‡ Mol. wt. 226. § Mol. wt. 380.

These lead to the diorthosilicate formula $(\text{Sc},\text{Y})_2\text{Si}_2\text{O}_7$, analogous to that of thalenite (A., 1899, ii, 766). Spectroscopic analysis shows the presence of several other rare-earths, but an absence of cerium, zirconium, etc.

L. J. S.

The Annaheim Meteorite. R. A. A. JOHNSTON and H. V. ELLSWORTH (*Trans. R. Soc. Canada*, 1921, [iii], 15, sect. IV, 69—92).—This iron, weighing 11.84 kilos., was found in 1916 and probably represents the meteor observed on January 21, 1914. The structure is that of a coarse octahedrite, and the different constituents were studied in detail by the metallographic method. I is the bulk analysis of the iron (also insol. in nitric acid 0.003; Ti, V, Al, Sn absent); II of the kamacite groundmass readily dissolved in 10% hydrochloric acid. The constituents insoluble in this dilute acid were separated into: III, bright, tin-white lamellæ of tænite rich in nickel; IV, tin-white, uncrystallised phosphide (schreibersite) giving ratios near $\text{Fe}_5(\text{Ni},\text{Co})_2\text{P}_2$; V, long, slender crystals of phosphide (rhabdite?), $(\text{Fe},\text{Ni},\text{CO})_3\text{P}$. VI, central portion of a nodule of chromiferous troilite, which in the outer portion is intergrown with 38.71% graphite.

	Fe.	Ni.	Co.	Cu.	Mn.	Cr.	Si.	P.	C.	Total.	d.
I.	91.51	7.84	0.46	0.08	trace	0.001	0.002	0.219	0.012	0.01	100.12
II.	93.10	6.39	0.48	0.03	—	—	—	—	—	—	100.00
III.	60.74	37.38	0.67	0.64	nil	—	nil	0.65	—	trace	100.08
IV.	61.28	25.62	0.47	0.09	—	—	—	13.06	—	—	100.52
V.	40.28	41.36	0.23	0.77	—	—	—	15.35	—	n. d.	98.68*
VI.	62.91	0.28	trace	0.16	—	0.96	—	—	35.47	—	99.76

* Also SiO₂ 0.69.

L. J. S.

The Pitts Meteorite. S. W. MCCALLIE (*Amer. J. Sci.*, 1922, [v], 3, 211—215).—This meteorite was observed to fall on April 20, 1921, at Pitts in Wilcox Co., Georgia. Four fragments, weighing

57, 42½, 30, and 2 ounces, were picked up; *d* 7-23. Analyses by E. EVERHART gave I for the metallic portion (forming about 90% of the whole); and II for the stony portion, of which 77-62% consists of pyrrhotite (FeS), the remainder being near to hypersthene in composition.

	Fe.	Ni.	Co.	Sn.	Cu.	Si.	S.	Mn.	
I.	0.09	0.32	trace	0.20	0.00	0.00	28.30	0.00	
II.	91.50	6.67	0.45	0.04	trace	1.40	0.02	0.05	
	MnO.	Na ₂ O.	K ₂ O.	MgO.	Al ₂ O ₃ .	FeO [<i>sic</i>].	P ₂ O ₅ .	SiO ₂ .	C [diff.].
II.	0.27	1.17	0.32	5.96	0.30	65.52	0.07	8.46	2.32
								L.	J. S.

L. J. S.

Analytical Chemistry.

Phenol-red as an Indicator for Acidity of Media. A. MASSINK (*Pharm. Weekblad*, 1921, 58, 1133—1136).—The colorimetric determination of p_H in the region 7—8 by means of phenol-red is affected by the electrolyte-content of the solution, the values being too low with low salt-content, and too high with high salt-content.

S. I. L.

Continuous Reading Electro-titration Apparatus. KENNETH H. GOODE (*J. Amer. Chem. Soc.*, 1922, 44, 26—29).—An electro-titration apparatus for the determination of hydrogen-ion concentration is described by means of which the *E.M.F.* between a calomel electrode and a hydrogen electrode may be read continuously. The voltmeter consists of a three electrode vacuum valve ("audion"). The three electrode valve consists of a highly exhausted glass bulb containing an incandescent filament surrounded by a grid of fine wire, which is itself surrounded by a metallic plate. A battery of 20—100 volts connected between the plate and filament, through a d'Arsonval galvanometer, produces a current through the plate circuit, the magnitude of which is proportional to the potential of the grid. The filament is connected to a 6-volt circuit containing a resistance which adjusts the current to 1-06 amperes. The current in the plate circuit, I_p , may be considered as the sum of a constant current, I_0 , which is independent of the grid potential and a current ($I_p - I_0$) which is a linear function of the grid potential. The current I_0 is balanced by an equal current in the opposite direction and ($I_p - I_0$) is measured with the galvanometer which is calibrated to read either volts or Sørensen units directly. The calomel cell is connected to the negative pole of the filament circuit and the hydrogen electrode to the grid. The apparatus as described with a d'Arsonval galvanometer of sensitivity of $10 \cdot 1 \times 10^{-6}$ amperes for one scale division is sensitive to 0-1 Sørensen unit or 0-006 volt.

J. F. S.

Titration in Ethyl Alcohol as Solvent. EDNA R. BISHOP, ESTHER B. KITTRIDGE, and JOEL H. HILDEBRAND (*J. Amer. Chem. Soc.*, 1922, **44**, 135—140).—Alcoholic solutions of acetic, maleic, palmitic, and sulphuric acids, *p*-chlorophenol, and phenol have been titrated with an alcoholic solution of sodium ethoxide, using the hydrogen electrode as indicator. The hydrogen electrode was measured against the electrode HgHg_2Br_2 | $\text{N}/10\text{KBr}$ in alcohol. Similar titrations of alcoholic solutions of ammonia and aniline by alcoholic hydrogen chloride were also carried out. Sharp and accurate end-points were obtained, but the titrations take much longer than titrations in aqueous solutions by the same method, since the hydrogen electrode requires ten minutes before it reaches a constant value. The hydrogen potentials at which a series of indicators changes colour was determined by titrating pairs of the above-named substances and using the indicators and then measuring the potential of the mixture which just gives the colour change. The potential range 0.14—0.45 was measured with aniline and hydrogen chloride, 0.45—0.70 and 0.90—0.97 with acetic acid and sodium ethoxide and 0.75—0.98 volt with *p*-chlorophenol and sodium ethoxide. The positions of most of the indicators in alcohol are close to their positions in water; some exceptions occur, for example, cyanine is displaced very much to the acid side, and some indicators which are useful in water cannot be used in alcoholic solution. The following colour changes and potentials are recorded: Benzaldehyde-green, green 0.69 colourless. Bromophenol-blue, yellow 0.34, green 0.47 blue. Cresol-red, pink 0.20 orange 0.30. Curcumin, greenish-yellow 0.66 red 0.85 orange 0.91 golden. Cyanine, colourless 0.24 blue; gallein, rose 0.68 violet-blue. Iodeosin, golden brown 0.20 pink; methyl-green, blue 0.66 lavender. Methyl-orange, pink 0.20 orange 0.23 yellow. Methyl-violet, violet 0.35 colourless. Methyl-red, red 0.54 orange 0.62 yellow. Naphthol-benzoin, light brown 0.70 blue. *p*-Nitrophenol, colourless 0.61 yellow green. Phenolphthalein, colourless 0.68 red. Resorcin-blue, red 0.39 blue. Rosolic acid, golden 0.58 orange 0.67 pink. Sodium alizarin-sulphonate, greenish-yellow 0.50 orange 0.55 rose 0.82 violet. Thymol blue, red 0.30 golden. Thymolphthalein, colourless 0.82 blue. Trinitrobenzene, colourless 0.68 golden-orange. Tropæolin salmon-pink 0.20 orange 0.23 golden, and tropæolin OO, pink 0.15 orange 0.20 yellow. Palmitic acid in the presence of tripalmitin may be titrated in alcoholic solution with sodium ethoxide, using thymolphthalein as indicator. J. F. S.

Simple Theory of the Nephelometer. P. V. WELLS (*J. Amer. Chem. Soc.*, 1922, **44**, 267—276).—The theory of the nephelometer and its use are considered. It is shown that although the depth ratios by reflection and by transmission are equal, both methods cannot be used in nephelometry except for an intermediate range of concentrations. For extremely dilute suspensions, for example, less than 10^{-5} gram/c.c., the transmissions are quite insensitive, whilst the reflection measurement remains sensitive down to the limit of vision. On the other hand, for very turbid suspensions

the transmission probably follows less complicated laws. Although masses in suspension much too small to be detected by the most delicate balance can be easily measured in a Tyndall beam, the precision of such a measurement can never exceed that of the best photometry, that is, about 0.2%. For sensitive and rapid work nephelometry takes its place with other volumetric methods. The phenomena of diffuse reflection and transmission are much more complicated than is represented in the present considerations, but the simple theory presented gives a general idea of the phenomena and may serve as a starting point of a future investigation.

J. F. S.

Nephelometry. A Nephelometer with Constant Standard.

A. A. WEINBERG (*Biochem. Z.*, 1921, **125**, 292—310).—A new nephelometer and point-source of light are described which embody the best features of the Kober (*A.*, 1917, ii, 266) and Kleinmann (*A.*, 1920, ii, 634) nephelometers, the most striking feature being the introduction of the Lummer-Brodhun prism whereby the two fields are rendered concentric. Nephelometric estimations with such an instrument are comparative and a modified nephelometer is described with a permanent standard. This is attained by replacing one of the tube systems by two Nicols, the lower of which can be rotated relatively to the upper, thus reducing the light to any degree as recorded by a scale on the Nicol prism. For coloured solutions, a slip of coloured glass can be inserted.

H. K.

Quantitative Analysis by Centrifuge. OLOF ARRHENIUS (*J. Amer. Chem. Soc.*, 1922, **44**, 132—134).—The estimation of calcium, magnesium, and phosphoric, sulphuric and nitric acids may be effected by precipitating the various substances in the usual way and transferring the precipitate and mother-liquor to tubes 2.5 cm. diameter at the top and provided with a steep funnel below leading to a capillary tube 4 cm. long and 1 mm. diameter. The precipitate is allowed to settle and by means of gentle shaking is, for the most part, caused to settle in the capillary stem. The tube is then centrifuged until the height of the column of precipitate is constant; this usually requires thirty minutes. The height of the column is then read and reduced to the weight standard by comparison with a precipitate similarly obtained from a known weight of the same substance. The method claims no great degree of accuracy, and is suggested as a suitable method of analysis for soil and similar substances.

J. F. S.

The Penetrability of Filter-paper. R. C. GRIFFIN and H. C. PARISH (*J. Ind. Eng. Chem.*, 1922, **14**, 199—200).—An apparatus is described for testing the penetrability of filter-paper by observing the time required to pass 100 c.c. of distilled water at 20° through a 2-inch disk of the paper under a constant head of 9 inches of water. The apparatus, which is constructed mostly of lead, consists of an overflow cup connected by a pipe to the under side of a

wire gauze which supports the disk of filter-paper, and is placed 9 inches below the top of the overflow cup. The distilled water is fed into the connecting pipe at such a rate that it overflows at the cup as well as through the outlet pipe of the apparatus after passing through the filter-paper, and the number of seconds required to collect 100 c.c. at the outlet is timed with a stop watch. It was found that the temperature of the water had a marked bearing on the speed of filtration, water at 30°, for example, passing through more than twice as rapidly as water at 0° under otherwise similar conditions. The time factor is also important, and even when distilled water is used the filtration slows down after a time owing to the hydration and expansion of the fibres. A paper which initially had a penetrability of twenty-five seconds had slowed down after two hours to one thousand seconds. G. F. M.

Estimation of Perchlorate by Rothmund's Method. FRITZ KÖNIG (*Z. anorg. Chem.*, 1921, 120, 48).—In the estimation of perchlorate by reduction with titanous sulphate by Rothmund's method (A., 1909, ii, 434), it is unnecessary to pass a current of hydrogen or carbon dioxide through the flask if a long, narrow glass tube is adapted to the reflux condenser to prevent circulation of air. For the oxidation of excess of titanous sulphate, ferric ammonium sulphate is better than permanganate in view of the fact that the latter may attack the hydrochloric acid, and that in the back-titration of silver nitrate in the chlorine determination by the Volhard process a ferric salt must be present. Soluble titanic sulphate is now a commercial product and can be reduced electrolytically, using lead electrodes, to titanous sulphate.

E. H. R.

Comparative Values of Different Specimens of Iodine for Use in Chemical Measurements. C. W. FOULK and SAMUEL MORRIS (*J. Amer. Chem. Soc.*, 1922, 44, 221—229).—Iodine purified by several methods, including wet and dry sublimation and drying in the presence of sulphuric acid and phosphoric oxide, has been compared by direct titration with sodium thiosulphate with iodine prepared by Baxter's atomic weight method (A., 1905, ii, 81, 579). Using some new forms of titration flasks and other pieces of apparatus for handling iodine, it is shown that the maximum difference between the various specimens and the highly purified iodine is 0.024%. It is shown that a rubber stopper may be used in place of a glass stopper for closing the flask in which the iodine is titrated. The usual method, as given in text-books of analytical chemistry, of drying iodine by exposing it in a desiccator with a drying agent, is open to criticism if the iodine has previously solidified in the presence of water. Powdered iodine when exposed under a bell-jar with water takes up 0.09% of its weight in forty-eight hours, whilst crystals take up only 0.05% under the same conditions in five days. An exposure to sulphuric acid for ten days removes the whole of the water in both cases. It is suggested that solids like iodine, which have a measurable vapour pressure at ordinary

temperatures, possess peculiar adsorptive properties, due to the fact that a fresh surface is being continually exposed. J. F. S.

Method of Determining Traces of Oxygen in Hydrogen.

ALFRED T. LARSON and ERNEST C. WHITE (*J. Amer. Chem. Soc.*, 1922, **44**, 20—25).—A rapid and accurate method of estimating traces of oxygen in nitrogen-hydrogen mixtures such as are used in the synthetic manufacture of ammonia is described. The method consists, essentially, in passing the gas mixture through a platinised platinum catalyst which is heated at 305° in a carefully regulated diphenylamine vapour-bath, and measuring the rise in temperature due to the combustion of the hydrogen by means of a thermo-element. The deflexions of a sensitive galvanometer attached to the thermo-element give a measure of the oxygen content of the gas mixture. Concentrations of oxygen as low as 0.001% may easily be determined with an error of 3%. The apparatus was calibrated for a rate of flow of 560 c.c. per minute by adding electrolytic oxygen from an alkali electrolysis to the oxygen-free mixture of nitrogen and hydrogen. By the use of a less sensitive galvanometer, the apparatus may be used for concentrations of oxygen up to about 1%. Precautions necessary for efficient working of the apparatus are indicated. J. F. S.

The Estimation of Sulphur in Iron Pyrites. G. CHAUDRON

and G. JUGE-BOIRARD (*Compt. rend.*, 1922, **174**, 683—685).—During the dissolution of pyrites in nitric acid or aqua regia, there is always a separation of free sulphur if the temperature exceeds 60°, in the case of marcasite or pyrites containing other metallic sulphides. If, however, the acid is allowed to act at the room temperature, there is no separation of sulphur, but the time required is much longer. W. G.

A New Method of Estimating Sulphur in Organic Compounds. H. TER MEULEN (*Rec. trav. chim.*, 1922, **41**, 112—

120).—The substance is vaporised or decomposed by heating in a current of hydrogen and the mixture of vapours and hydrogen then passed through a heated quartz tube containing platinised asbestos. All the sulphur is thus converted into hydrogen sulphide, which is led into an alkaline solution and estimated iodometrically. In the case of very small quantities, colorimetric estimation by means of sodium plumbite is suitable. Experimental errors are avoided by using small quantities of material, heating gently, and not allowing the hydrogen to pass too quickly. It is occasionally necessary to burn off a deposit of carbon from the catalyst.

The results obtained by this method are in good agreement with the theoretical values. H. J. E.

Volumetric Method for the Estimation of Hyposulphurous and Sulphoxylic Acids. FERRUCCIO DE BACHO (*Giorn. Chim. Ind. Appl.*, 1921, **3**, 501—502).—This method is based on the property possessed by formaldehyde of forming with hyposulphite a solution highly resistant to oxidation by means of atmospheric oxygen [cf. *J. Soc. Chem. Ind.*, 1922, 250A.] T. H. P.

Microchemical Estimation of Nitrogen. C. VALLÉE and M. POLONOWSKI (*Compt. rend. Soc. Biol.*, 1921, **84**, 900—901; from *Chem. Zentr.*, 1921, iv, 1080).—For microchemical estimation of nitrogen, the substance is heated with 1 c.c. of sulphuric acid, 1 gram of potassium sulphate, and a small piece of quartz, and diluted with 6 c.c. of water. After addition of 3 c.c. of sodium hydroxide solution, the ammonia is carried over by a current of air into 0.02*N*-sulphuric acid. G. W. R.

Estimation of Small Quantities of Nitrogen by Kjeldahl's Method. J. K. PARNAS and RICHARD WAGNER (*Biochem. Z.*, 1921, **125**, 253—256).—A description of a slight modification of Pregl's micro-Kjeldahl method of analysis. H. K.

Nesslerisation of Ammonia Solutions. CLARENCE E. MAY and HARRY P. ROSS (*J. Amer. Chem. Soc.*, 1921, **43**, 2574—2575).—In the estimation of ammonia in urine by the Folin-Bell method (A., 1917, ii, 268) and of urea in urine by the Folin-Youngburg method (A., 1919, ii, 304), the authors were troubled by the formation of clouds and precipitates during the time (twenty minutes) which the solution is kept after the addition of the reagent. An investigation of the reactions shows that such disturbing occurrences may be avoided if the following points are observed. Distilled water only must be used for dilutions; the flask must be rinsed with nitric acid before each determination to remove the mercury film which may have been formed in a previous experiment; the Nessler solution should be the one recommended by Folin and Wu (A., 1919, ii, 308); the whole of the Nessler solution must be added rapidly; after the addition, the solution must not be shaken or stirred and must be left unmoved for at least twenty minutes and any dilution must be made only after this time has elapsed; the solution to be Nesslerised should not contain more than 1.0 mg. of ammonia nitrogen in 150 c.c. On Nesslerisation and dilution to 200 c.c., the brown colour is very intense, and with such concentrated solutions part of the product tends to deposit on the walls of the flask and interfere with the colorimetric work. J. F. S.

The Volumetric Estimation of Hydroxylamine and Hydrazine. ALBIN KURTENACKER and JOSEF WAGNER (*Z. anorg. Chem.*, 1921, **120**, 261—266).—It was shown by Rupp and Mäder (A., 1913, ii, 618) that hydroxylamine can be oxidised quantitatively by a bromate-bromide mixture in sulphuric acid solution to nitric acid, according to the equation $\text{NH}_2\cdot\text{OH} + 6\text{Br} + 2\text{H}_2\text{O} = \text{HNO}_3 + 6\text{HBr}$. As a method of analysis, however, the reaction has serious limitations, since, unless only a very small quantity of hydroxylamine is taken and a large excess of bromine used, oxidation is incomplete and nitrous acid is formed. It is found, however, that if bromate alone is used in strong hydrochloric acid solution, nascent chlorine being then the oxidising agent, complete oxidation is assured, and the reaction can be used for the estimation of hydroxylamine. The hydroxylamine solution (containing about 1.14 gram of $\text{NH}_2\cdot\text{OH}\cdot\text{HCl}$ per litre) is mixed with excess of

0.1*N*-potassium bromate solution and acidified with hydrochloric acid. After a quarter of an hour (the reaction is not instantaneous), potassium iodide solution is added and the liberated iodine is titrated with thiosulphate. Sulphuric acid cannot be used instead of hydrochloric, and bromide must not be present.

Hydrazine is oxidised by bromate or a bromide-bromate mixture in hydrochloric acid solution instantaneously to nitrogen. The estimation can be carried out exactly as for hydroxylamine, estimating excess of bromate by means of iodide and thiosulphate, or a direct bromate titration can be made using indigo as indicator. The hydrazine solution is mixed with potassium bromide and hydrochloric acid and titrated with 0.1*N*-bromate solution at about 60°. Towards the end of the titration, a few drops of indigo solution are added and the titration is continued until the colour becomes yellow.

Hydroxylamine and hydrazine may be estimated together. A sample is first titrated as for hydroxylamine, and the bromate equivalent for the sum of the two found. A second sample is then oxidised with bromate in an atmosphere of carbon dioxide and the nitrogen evolved is collected and measured. This gives the amount of hydrazine present and that of the hydroxylamine can be calculated.

E. H. R.

The Action of Nitrous Acid on Iodides in the Presence of Oxygen. MAURICE LOMBARD (*Bull. Soc. chim.*, 1922, [iv], 31, 161—169).—When nitrous acid acts on potassium iodide in the presence of oxygen, the nitrous acid is regenerated as fast as it disappears. Without taking slight losses into account, two causes limit the phenomenon. One of these causes is independent of the analyst, and is small but relatively constant. It is the production of a small amount of nitrogen. The other cause, which is of variable importance, depends almost entirely on the method of working. It is the production of nitrate (which may be nil); and it is this which, coupled with the losses by diffusion, leads to such variable results. Thus the estimation of nitrites by the liberation of iodine is impossible in the presence of oxygen. Comparable results might be obtained by working under carefully controlled conditions which would, however, be difficult to realise, and different analysts would probably obtain widely differing results.

W. G.

Estimation of Oxides of Nitrogen [in Air]. V. C. ALLISON, W. L. PARKER, and G. W. JONES (*U.S. Bureau of Mines, Tech. Paper* 249).—The phenoldisulphonic acid method for the estimation of nitrates in water analysis is adapted to the estimation of small quantities of oxides of nitrogen in air. The sample is taken in an evacuated tube of about 250 c.c. capacity, and after the usual analysis has been made for carbon dioxide, carbon monoxide, oxygen, etc., 5 c.c. of 10% sodium hydroxide solution and 5 c.c. of hydrogen peroxide are introduced into the tube, which is then closed with a rubber stopper and rotated to coat the inside with a film of liquid, and is then left for thirty minutes. The contents are then washed through a filter-paper into a 150 c.c. beaker and

evaporated just to dryness. The residue is treated with 2 c.c. of the phenoldisulphonic acid reagent, diluted to 10 c.c., filtered into a Nessler tube, 15 c.c. of ammonia are added, and the whole made up to 100 c.c. and compared with standards similarly prepared from a standard potassium nitrate solution. The method is sensitive to 10 parts of oxides of nitrogen in 1,000,000 parts of air with an accuracy of about 5 parts per million. G. F. M.

The Argentometric Titration of Phosphoric Acid. I. M. KOLTHOFF (*Pharm. Weekblad*, 1922, 59, 205—215).—The solubility product of silver phosphate in water is of the order 1×10^{-21} ; accurate determinations are impossible, since decomposition occurs, the salt becoming black. Accurate results are easily obtained in estimating phosphoric acid by means of excess of silver nitrate if the solution is made neutral to phenol-red by addition of 0.1N-sodium hydroxide; phenolphthalein is less suitable. In place of sodium hydroxide, sodium acetate may be added; about 2 grams are required for 10 c.c. of 0.1 molar phosphate solution, and in this case ammonium, calcium, and magnesium salts do not interfere.

In estimating phosphoric acid in urine, the ammonium magnesium salt is precipitated, washed and dissolved in nitric acid, the solution being neutralised with sodium hydroxide, using dimethyl-yellow as indicator, and titrated with silver nitrate, the excess of which is estimated by Volhard's method. S. I. L.

Errors caused by Nitrates and Nitrites in the Estimation of Arsenic by the Distillation Method and a Means for their Prevention. J. J. T. GRAHAM and C. M. SMITH (*J. Ind. Eng. Chem.*, 1922, 14, 207—209).—In the presence of nitrates or nitrites the distillation method for the estimation of arsenic as usually carried out, using cuprous chloride or cuprous chloride and ferrous sulphate as the reducing agents, gives low results owing to some volatile substance, probably nitrosyl chloride, carried over with the arsenic trichloride into the distillate oxidising it to quinquivalent arsenic. The extent of the oxidation depends largely on the length of time the distillate is kept before titration. The error may be avoided by using hydrazine sulphate in conjunction preferably with sodium bromide as the reducing agent, whereby the nitrates, etc., are reduced to nitrogen. For a sample containing the equivalent of not more than 0.6 gram of As_2O_3 , 50 c.c. of a solution containing 2% each of hydrazine sulphate and sodium bromide in dilute (1 : 4) hydrochloric acid will be required for the analysis. The mixture is boiled for two to three minutes, 100 c.c. of concentrated hydrochloric acid are added, and the distillation is proceeded with in the usual way. G. F. M.

The Separation of Arsenic from Tungsten, Vanadium, and Molybdenum by Means of Methyl Alcohol in a Current of Air. L. MOSER and J. EHRLICH (*Ber.*, 1922, 55, [B], 430—437).—It has been shown previously (A., 1912, ii, 866) that tervalent arsenic can be separated quantitatively from antimony and other metals at the temperature of boiling water by volatilising it partly as

methyl arsenite and partly as arsenic trichloride in a current of air. In the presence of tungstic acid, however, the removal of arsenic is incomplete, owing to the adsorption of a portion of the arsenic trichloride by colloidal tungstic acid. The difficulty can be removed by bringing the latter into highly disperse solution, which is readily effected by means of pyrogallol; the procedure is described fully in the original. Accurate results are obtained thereby in the estimation of arsenic, but the method suffers under the disadvantage that the pyrogallol must be destroyed before the tungsten can be estimated in the residue. The action of a number of other substances (oxalic, tartaric, and citric acids) is considered and the following process is recommended. The solution containing the tungstate and arsenious acid (arsenic acid must be reduced by one of the customary methods) is evaporated to small bulk and treated with glacial acetic acid (20 c.c.) and concentrated hydrochloric acid (120—150 c.c.); after addition of methyl alcohol (30 c.c.), distillation is effected as described previously in a current of air. Arsenic is estimated, preferably iodometrically, in the distillate; the residue in the distillation flask is evaporated completely to dryness on the water-bath, dissolved in dilute sodium hydroxide solution, and the tungstic acid precipitated as mercurous tungstate.

The separation of arsenic from molybdenum and vanadium is effected without difficulty by the ester method; it is only necessary to make certain that a sufficient amount of the reducing agent is added (even when the arsenic is present in the tervalent state), since molybdic and vanadic acids themselves attack the latter.

H. W.

The Theory of the Distillation of Arsenic and a New Separation of Arsenic from all Metals in a Current of Air. L. MOSER and J. EHRLICH (*Ber*, 1922, 55, [B], 437—447; cf. Moser and Perjatel, A., 1912, ii, 866; Moser and Ehrlich, preceding abstract).—A study of the rate of distillation of arsenic chloride from a solution of arsenious oxide in concentrated hydrochloric acid shows that the process is influenced greatly by the hydrolysis of the chloride and that this effect becomes more marked as the distillation proceeds towards completion. The difficulty has been overcome largely by suitable addition of potassium bromide to which a catalytic action has been commonly ascribed. Its function, however, is shown to depend on its ability to repress hydrolysis of arsenic chloride, since it may be regarded as attracting the water from the mixture for purposes of solution and thus rendering the solutions of arsenic chloride in hydrochloric acid more concentrated with regard to the latter. The effect is exhibited by a number of other substances, for example, barium chloride, barium acetate, sodium pyroborate, sodium benzoate, sodium nitroprusside, crystalline lactose, etc., which are freely soluble in water, but insoluble or very sparingly soluble in concentrated hydrochloric acid.

The following method of estimating arsenic is recommended.

It does not involve the use of methyl alcohol. It requires no supervision and is complete within about forty minutes (or thirty minutes after addition of potassium bromide). Since the liquid never comes to its boiling point, it is impossible for compounds of antimony to distil. The apparatus consists of a wide-mouthed flask of 300 c.c. capacity provided with a rubber stopper carrying an inlet tube for air, a stoppered dropping funnel, and a bulb tube connected with a long glass tube dipping into water (250 c.c.) contained in a beaker which is cooled by running water. The arsenious oxide (0.15–0.25 gram) is dissolved in concentrated hydrochloric acid (d 1.19, 50 c.c.) in the flask which is immediately immersed up to the neck in boiling water, whilst a brisk current of air is passed through the solution. At intervals of ten minutes, further additions of concentrated hydrochloric acid (20 c.c.) are made. After forty to sixty minutes the distillation is interrupted and the arsenic titrated in the distillate with $N/10$ -potassium bromate solution. The procedure is similar when potassium bromide (about 1.5 grams) is used, but two or at most three additions of hydrochloric acid only are necessary. Arsenic acid must be reduced in the usual manner, for example, with ferrous sulphate, hydrazine sulphate, or even with potassium bromide alone.

H. W.

Microchemical Estimation of Carbon and Hydrogen. FRITZ WREDE (*Ber.*, 1922, **55**, [B], 557–563).—A valuable detailed discussion, unsuitable for abstraction, of apparatus, absorbent and oxidising materials, and procedure necessary for the microchemical estimation of carbon and hydrogen in organic compounds by Pregl's method.

J. K.

Estimation of the Carbon Dioxide Content of Air, with Special Reference to the "Aeronom." HANS RAUCH (*Z. Hyg. Infektionskrankh.*, 1920, **91**, 1–26).—A critical comparison was made between various methods for the estimation of carbon dioxide as used in analysing the air of schools, dwellings, etc. Results obtained with the "aeronom," an instrument in which the difference of pressure, measured with an oil manometer, of an enclosed volume of air before and after its content of carbon dioxide has been absorbed with sodium hydroxide, differ by as much as 20% when compared with the more exact Pettenkofer's baryta method. These larger errors were found also when the carbon dioxide was absorbed by sodium carbonate or sodium hydroxide as in the methods of Lung-Zeekendorf and Wolput. The author concludes that of the methods studied in determining the ventilating conditions in rooms, Pettenkofer's method is preferable.

CHEMICAL ABSTRACTS.

Estimation of Carbon Dioxide in Mineral Carbonates. L. A. SAYCE and A. CRAWFORD (*J. Soc. Chem. Ind.*, 1922, **41**, 57–58T).—The ignition method in a Teclu furnace until the substance shows no further loss in weight gives consistent results where the decomposition temperature is low enough for it to be

applied without complications arising, as, for example, reduction to metallic lead with cerussite, and oxidation of the ferrous oxide to an unknown extent with chalybite. The Schrötter method is unsuitable for estimating the carbonates of metals having comparatively insoluble sulphates. Hydrochloric acid can be used successfully for the decomposition, but this introduces the danger of loss of acid by volatilisation. Armstrong's simple modification of the Schrötter apparatus is capable of fair accuracy. Garrett's apparatus, consisting of a stout test-tube to the top of which is attached a Y-tube, one branch of which bears a tap funnel containing dilute acid, and the other a combined condenser and drying tube leading to potash bulbs or soda-lime tubes, gave good results with witherite, but for some unexplained reason very inconsistent results were obtained with calcite. Collin's calcimeter, by reason of the ease of operation, and the short time needed for an estimation seems well adapted for dealing with readily soluble carbonates when an accuracy of 0.1 or 0.2% is sufficient. G. F. M.

The Volumetric and Gravimetric Estimation of Zinc. STEFAN URBASCH (*Chem. Ztg.*, 1922, 46, 6—7, 29—30, 53—55, 97—99, 101—103, 125—127, 133—134, 138—139).—The separation of zinc from the common metals and its estimation by means of potassium ferrocyanide and sodium sulphide and as zinc oxide have been studied in detail. In all cases the ore is dissolved in aqua regia, the solution evaporated with sulphuric acid, the heavy metals removed with hydrogen sulphide and the iron and aluminium with ammonia in the presence of ammonium chloride if the ferrocyanide or gravimetric method is to be used, or of magnesium chloride if the sulphide titration is adopted. The ferrocyanide titration is carried out in a bulk of 150 c.c. containing 2—3 drops of hydrochloric acid and 3 c.c. of a 0.03% ferric chloride solution. The ferrocyanide is added to the boiling zinc solution until the blue colour just fades to white, a standard zinc chloride solution is then added, drop by drop, until the blue colour just reappears, and this amount of zinc is deducted from that found in the first titration. In the sulphide method, ammonia and ammonium salts should be kept as low as possible. The separation of cadmium from zinc by hydrogen sulphide is not complete in one operation if much zinc is present, but a satisfactory separation may be obtained by boiling the solution of the mixed metals with aluminium foil and finally adding hydrogen sulphide water to prevent re-solution of the cadmium. The separation of zinc from nickel and cobalt by the "salting out" method is discussed with regard to the necessary acidity of the solution; the author recommends saturating a solution containing less than 0.15 gram of zinc and about 1 c.c. of normal acid per 100 c.c. with hydrogen sulphide at 50°. [*Cf. J. Soc. Chem. Ind.*, 1922, 218A.] A. R. P.

The Detection of Small Quantities of Lead in Urine. O. SCHUMM (*Z. physiol. Chem.*, 1922, 118, 189—214).—Copper sulphate is added to the urine and the lead precipitated with the copper as the sulphides by hydrogen sulphide; by then separating the

mixture by electrolysis, it is possible to detect small quantities of lead of the order of a few decimilligrams in a litre and a half.

S. S. Z.

Studies on Thallium Compounds. I. Analytical. ARTHUR JOHN BERRY (T., 1922, 121, 394—399).

A New Iodometric Method for the Estimation of Copper. RUDOLF LANG (*Z. anorg. Chem.*, 1921, 120, 181—202).—The usual iodometric method for estimating copper depends on the reduction of a cupric salt by an iodide to cuprous iodide followed by titration of the free iodine formed. The reaction $\text{Cu}^{++} + 2\text{I}^- \rightleftharpoons \text{CuI} + \text{I}_2$ is, however, reversible, and in the present paper methods are developed for estimating copper which consist in oxidising a cuprous salt with a known quantity of iodine and titrating the excess. The most suitable salt for oxidation is cuprous thiocyanate, the reaction being carried out in presence of ammonium oxalate or tartrate. The reaction is complicated, however, by the possibility of a reaction between thiocyanate and iodine such as $\text{KCNS} + 8\text{I} + 4\text{H}_2\text{O} = \text{H}_2\text{SO}_4 + \text{ICN} + \text{KI} + 6\text{HI}$. A lengthy investigation was carried out to study the extent to which this reaction proceeds in presence of thiocyanates of potassium, sodium, and ammonium, oxalic acid, ammonium oxalate, ammonium sulphate, nitrate, and chloride in presence and absence of copper salts, and it was found that the reaction could be altogether inhibited in presence of oxalic acid and the ammonium salts of strong acids, including ammonium thiocyanate in high concentration. The following method was finally adopted for the estimation of copper. A solution of cupric salt, weakly acidified with a mineral-acid, and containing at most 0.28 gram of copper is reduced with excess of sulphurous acid and diluted to 0.1*N*. Ammonium thiocyanate solution is added to precipitate the copper as cuprous thiocyanate and the solution is boiled to expel sulphur dioxide. After cooling, a mixture of 5 volumes of ammonium oxalate solution containing 45 grams of hydrated salt per litre and 7 volumes of oxalic acid solution containing 120 grams of hydrated acid per litre is added, and the whole is diluted to 400 c.c. Next 0.1*N*-iodine solution is run in until a clear solution is obtained, the mixture being well shaken, and excess of iodine is titrated with thiosulphate solution.

The cupric salt can be reduced with cyanide instead of sulphurous acid, the advantage of this method being that the reduction is selective for copper and that the reducing agent need not be removed. Conditions were determined experimentally under which reaction between cyanide and iodine is prevented, and it was found necessary, after reduction, to acidify before adding iodine since only cyanogen ions, not hydrocyanic acid, reduce iodine. The following method was finally adopted. The cupric salt solution containing at most 0.28 gram of copper is made ammoniacal in a long-necked flask and reduced with 0.5*N*-potassium cyanide solution. After addition of 1 gram of ammonium thiocyanate, the solution is acidified with concentrated oxalic acid solution, keeping

cold, so that all the copper is precipitated as thiocyanate. The oxidation with iodine and titration are then carried out as above.

In place of the expensive potassium iodide solution of iodine, a potassium cyanide solution can be used, prepared by dissolving 3 grams of potassium cyanide in a little water, dissolving in this 12.7 grams of iodine, and making up to a litre. This solution is quite stable.

The above methods are applicable in presence of all the commoner metals. Notes are given of slight precautions and modifications necessary in presence of silver, mercury, lead, bismuth, arsenic, antimony, cobalt, manganese, iron, barium, strontium, calcium, and magnesium. E. H. R.

A Rapid Iodometric Estimation of Copper and Iron in Mixtures of their Salts. IAN WILLIAM WARK (T., 1922, 121, 358—363).

Estimation of Metallic Aluminium and Aluminium Oxide in Commercial Metal. H. V. CHURCHILL (*J. Ind. Eng. Chem.*, 1922, 14, 81); JULIAN H. CAPPS (*ibid.*, 81—82).—The first author criticises a recent paper by Capps (A., 1921, ii, 657), whose results indicate an atomic weight of 26.81 for aluminium, and concludes that the method described is untrustworthy. Capps replies, admitting an error pointed out by Churchill, but considers that his figures after recalculation still support his contention that the accepted atomic weight for aluminium is wrong. W. P. S.

The Analysis of Aluminium Alloys. ANTON BRENNER (*Chem. Ztg.*, 1922, 46, 188).—The solution of aluminium and zinc hydroxides in alkali hydroxide solution is treated with sodium sulphide, the precipitate is collected on a filter, washed with hot water, and dissolved in dilute hydrochloric acid. The cold solution is treated with sodium carbonate and the precipitated zinc carbonate collected, washed, ignited, and weighed as zinc oxide. The method is more rapid than that involving separation of the zinc sulphide from organic acid solutions. A. R. P.

The Separation of the Oxides of Iron and Aluminium from Admixture with Calcium Oxide by the Nitrate Method. CHARRIOU (*Compt. rend.*, 1922, 174, 751—754).—In St. Claire Deville's method for the separation of the oxides of iron and aluminium from calcium oxide by means of ammonium nitrate, it is preferable to carry out the drying, after the precipitation of the hydroxides with ammonia, at a temperature not exceeding 150° and in the presence of ammonium nitrate. The residue is then extracted three times with a boiling 10% solution of ammonium nitrate and subsequently washed with boiling water. Under these conditions the separation is complete. W. G.

A New Quantitative Method for the Estimation of Iron in the Blood. ANSON L. BROWN (*J. Amer. Chem. Soc.*, 1922, 44, 423—425).—To 0.5 c.c. of blood, laked in 4 c.c. of water, 1 c.c. of concentrated hydrochloric acid and 0.01 gram of potassium chlorate

are added, and the mixture is heated in a boiling water-bath until it becomes white or light yellow and the whole of the proteins are precipitated. The mixture is allowed to cool and is filtered, and the residue is washed with water until 15 c.c. of filtrate are obtained. The iron is estimated colorimetrically in an aliquot portion of the filtrate against a standard iron solution, using potassium thiocyanate as indicator.

W. G.

Separation of Germanium and Arsenic. JOHN H. MÜLLER (*J. Amer. Chem. Soc.*, 1921, **43**, 2549—2552).—A method of separating minute traces of arsenic from germanium is described. The method consists in adding a large excess of hydrofluoric acid to the mixed oxides and precipitating cold with hydrogen sulphide. By these means arsenic is precipitated quantitatively. Hydrogen sulphide is without action on solutions of fluorogermanic acid and its salts. The separation is sufficiently accurate to estimate as little as 0.01% of arsenic in germanium compounds. The method is especially useful in connexion with the preparation of pure germanium compounds, as the fractional crystallisation of the double fluoride can thus be avoided and the operation carried out in small volumes of solution. Germanium dioxide is non-toxic, and solutions of it when injected subcutaneously show a marked erythropoietic action.

J. F. S.

Co-precipitation of Vanadic Acid with Ammonium Phosphomolybdate. J. R. CAIN and J. C. HOSTETTER (*J. Amer. Chem. Soc.*, 1921, **43**, 2552—2562).—The authors have investigated the conditions under which the simultaneous precipitation of vanadic acid and ammonium phosphomolybdate is best carried out in connexion with the method of estimation of vanadium in steel (*A.*, 1912, ii, 1101). The ratio of vanadium to phosphorus in the precipitate has been determined for mixtures in which the phosphorus is constant and the vanadium varies, and for mixtures in which the vanadium is constant and the phosphorus varies. The influence of dilution, temperature, and acidity of the solutions has also been investigated. The results show that the co-precipitation of vanadic acid with ammonium phosphomolybdate may be explained as a phenomenon resulting from a partition of the vanadic acid in some undissociated form between the solution and the solid phase. The maximum absorption by the solid phase occurs at a temperature of 40—50° and in a nitric acid concentration of 4*N*. The effect of dilution in lowering the amount of occlusion is a direct result of the partition law and can be minimised by the presence of ammonium nitrate, the latter probably tending to repress the dissociation of the vanadium complex which is occluded. The ammonium vanadophosphomolybdates are probably a series of solid solutions the end members of which may be ammonium phosphomolybdate and ammonium phosphovanadate.

J. F. S.

Estimation of Methyl Alcohol. KARL H. A. MELANDER (*Svensk. Pappers Tid.*, 1921, **24**, 277—278).—A mixture of 22 grams

of iodine and 5 c.c. of the solution containing about 2% of methyl alcohol is heated for half an hour at 70° in a slow current of hydrogen, the flask having previously been connected with a condenser, a wash-bottle containing a little red phosphorus in water at 50°, and an absorption apparatus containing a few c.c. of pyridine. For a further period of half an hour a higher temperature is employed, and the current of hydrogen increased. The pyridine solution is concentrated on a water-bath, dissolved in water, and titrated with 0.1N-silver nitrate solution.

CHEMICAL ABSTRACTS.

Vapour Pressures of Dilute Alcohol Solutions. R. THOMAS (*J. Soc. Chem. Ind.*, 1922, 41, 33—34r).—A rapid method of determining the partial pressures of alcohol and water in aqueous solutions of alcohol is described. This method also constitutes a method of analysing air containing water and alcohol vapour. It depends on the facts that water is retained by a layer of calcium carbide which has no action on alcohol, and that alcohol is completely absorbed by a layer of ignited alumina or by passing it through 98% sulphuric acid. The process is carried out by passing dry air, free from carbon dioxide, through a bubbler containing the alcohol solution at a rate of 4—5 litres an hour and thence over calcium carbide contained in a U-tube and finally through 98% sulphuric acid or over ignited alumina in a U-tube. The loss of weight of the alcohol bubbler gives the total weight of vapour, and the increase in weight of the sulphuric acid or alumina gives the weight of alcohol. It is essential that fresh carbide be used in each experiment, for calcium hydroxide would retain some of the alcohol. It is shown for solutions containing 5% and 10% of alcohol that the partial pressure of the alcohol is approximately proportional to its concentration. From the data obtained, a table has been constructed showing the ratio of the concentration of alcohol in the gaseous phase to its concentration in the liquid phase for the temperature range 15—26°. J. F. S.

Estimation of Glycerol in the Presence of Sugars. L. F. HOYT and H. V. PEMBERTON (*J. Ind. Eng. Chem.*, 1922, 14, 54—56).—A method is described for the estimation of glycerol in transparent soaps, which almost invariably contain added sucrose. The soap is dissolved in water, the solution acidified with sulphuric acid, and boiled to remove any alcohol which may be present. The mixture is filtered to separate insoluble fatty acids, the filtrate treated with a small quantity of silver sulphate, diluted to a definite volume, and again filtered. A portion of this filtrate is oxidised in the usual way with potassium dichromate-sulphuric acid mixture, using a larger excess of the oxidising mixture than is customary in glycerol analysis; this estimation gives the sum of the glycerol and sugar, and the latter is then estimated separately in another portion of the filtrate. W. P. S.

Sensitive Test for Phenols. JAMES MOIR (*J. S. African Chem. Inst.*, 1922, 5, 8—9).—Five c.c. of *p*-nitroaniline hydrochloride

solution (1.5 grams of *p*-nitroaniline dissolved in a mixture of 40 c.c. of hydrochloric acid and 500 c.c. of water) are treated with dilute sodium nitrite solution until decolorised, and the mixture is then added to the solution to be tested; an orange-coloured precipitate is formed if much phenol is present. After one minute, an excess of sodium hydroxide is added; a coloration develops varying in depth from salmon-pink to ruby-red, according to the quantity of phenol present. The salmon-pink coloured solution exhibits a broad absorption band at λ 494 and this serves to distinguish phenol from the cresols, etc. The test will detect 1 part of phenol in 1,000,000 parts of solution. W. P. S.

Proteinogenous Amines. XIV. A Microchemical Colorimetric Method for Estimating Tyrosine, Tyramine, and other Phenols. MILTON T. HANKE and KARL K. KOESSLER (*J. Biol. Chem.*, 1922, **50**, 235—270).—The colour reaction given by phenol, *o*-, *m*-, and *p*-cresols, and *p*-hydroxyphenylacetic, *p*-hydroxyphenylpropionic and *p*-hydroxyphenyl-lactic acids with sodium *p*-diazobenzene sulphonate may be used for the colorimetric estimation of these substances. The details of the method are practically identical with those previously described for the estimation of iminazole derivatives (A., 1920, ii, 67). Tyrosine and tyramine give with a solution of diazobenzenesulphonic acid in sodium carbonate a pink coloration which rapidly changes to yellow. Addition of sodium hydroxide produces an intensification of the yellow and a restoration, to a small extent, of the pink coloration, but at no stage is the intensity proportional to the concentration of the phenol. If, however, sodium hydroxide and then hydroxylamine hydrochloride are added to the solution, an intense bluish-red colour is obtained which is proportional to the amount of tyrosine or tyramine present. The coloration so produced may consequently be employed for the estimation of small quantities of these two compounds. Ammonium salts, leucine, glycine, hydrogen peroxide, formaldehyde, acetaldehyde, acetone, acetoacetic acid, dextrose, and alcohols interfere with the colour. An explanation based on a tautomeric change of the keto-enol type is tentatively advanced to account for the colour changes described in the case of tyrosine and tyramine. E. S.

Proteinogenous Amines. XV. A Quantitative Method for the Separation and Estimation of Phenols including Phenol, *o*-, *m*-, and *p*-Cresols, and *p*-Hydroxyphenylacetic, *p*-Hydroxyphenylpropionic and *p*-Hydroxyphenyl-lactic Acids, Tyrosine, and Tyramine. MILTON T. HANKE and KARL K. KOESSLER (*J. Biol. Chem.*, 1922, **50**, 271—288).—The method of separation described was designed to render possible the application of the author's method for the estimation of certain phenolic substances (preceding abstract) to mixtures of these substances. It is intended primarily for use in bacterial metabolism studies on tyrosine and is not applicable to more complex liquids such as urine or blood. The liquid containing the mixture of phenols is acidified and distilled, whereby phenol and the cresols pass over.

Non-aminophenolic acids are removed from the residual liquid by extraction with ether, after which it is made alkaline with sodium carbonate and extracted with amyl alcohol. Tyramine is thus removed, whilst tyrosine remains in the alkaline solution. Separation is thus effected into four fractions. If, as will normally be the case in bacterial metabolism experiments, only one member of each fraction is present, the estimation is proceeded with after appropriate treatment of the various solutions. The method fails, however, in those cases where more than one member of each fraction is present since further separation is impossible.

E. S.

Estimation of Sugar in Small Quantities of Blood by the Pavy-Sahli Method. SVEND' HUBERT REIST (*Schweiz. med. Woch.*, 51, 419—423; from *Chem. Zentr.*, 1921, iv, 1056).—0.1 c.c. of blood is washed into a test-tube with 3 c.c. of alcohol. The mixture is shaken for half an hour, filtered through hardened filter-paper, and the residue washed twice with 2 c.c. of alcohol. The alcohol is then removed over a water-bath and 0.3 c.c. of Pavy's solutions and 1.8 c.c. of water are added. The liquid is heated to boiling and while boiling titrated with standard (approximately 0.1%) dextrose solution.

G. W. R.

Estimation of Lævulose (Fructose) in Straw. S. H. COLLINS (*J. Soc. Chem. Ind.*, 1922, 41, 56—57r).—The colouring matters present in this case bear a high ratio to the sugar and attempts to obtain concentrated solutions sufficiently translucent for the polarimeter failed. It was found in practice that a solution obtained by continuous extraction, of which 10 parts of liquid represented 1 part of straw, was as dark as could be used even with a 200 mm. tube and a quartz wedge saccharimeter illuminated by a 100 c.p. electric lamp. By means of hot and cold water circulating through the jacketed polarimeter tube, temperatures of about 70° and 10°, constant to a fraction of a degree, could be obtained, and ten readings were taken in order to reduce chance errors to unimportance. A serious error was found to be due to the deflexion of the zero by temperature alterations. This was traced to the end glasses of the polarimeter tube and was overcome by annealing them for three hours at 300°. The least trace of alkalinity must be avoided, otherwise the lævulose solutions become almost black at the higher temperature. Faint acidity is not of so much consequence. With straw containing 5—7% of lævulose the error in the estimation is about 0.05% with carefully annealed end glasses. The results of the investigation have shown that with straws containing much sugar three-fourths of the total is lævulose, but when the total sugar is low in amount, lævulose is absent.

G. F. M.

Micro-estimation of Lactose by means of Potassium Permanganate. Application to Milk. G. FONTÈS and L. THIVOLLE (*Bull. Soc. Chim. Biol.*, 1922, 4, 23—42).—The author's method for the estimation of dextrose in blood (*A.*, 1921, ii, 563)

can, with slight modifications, be used for the estimation of lactose in milk. E. S.

A Simpler Method of Determining Acetyl Values. LEON W. COOK (*J. Amer. Chem. Soc.*, 1922, **44**, 392—394).—A modification is given of André's formula (cf. A., 1921, ii, 419) for calculating the acetyl value of an oil from its saponification values before and after acetylation. The new formula is $A = (S' - S)/(1 - 0.00075S)$, where A is the acetyl value, S the saponification value before acetylation, and S' the value after acetylation. Similarly, the percentage of alcohol in the original sample may be calculated by the formula: percentage of alcohol = $M(S' - S)/(560 - 0.42S')$ provided that the molecular weight, M , of the alcohol is known. W. G.

The Estimation of Fatty Acids Based on their Volatility in Steam. W. ARNOLD (*Z. Unters. Nahr. Genussm.*, 1921, **42**, 345—372).—From a large number of determinations of the Reichert-Meissl and Polenske values of pure fatty acids, using the Polenske apparatus and weights of fatty acid varying from 0.01 gram to 0.05 gram, it appears that butyric and hexoic acids only give a Reichert-Meissl value; octoic and decaoic acids give both Reichert-Meissl and Polenske values, the Reichert-Meissl value of decaoic acid being distinct but very small. Acids from lauric upwards only give Polenske values. Of the insoluble acids decaoic and lauric are easily volatile, palmitic and stearic volatile with difficulty, myristic standing midway between the two groups. In the case of palmitic and stearic acids, the amount that passes over is so small that the Polenske value is practically independent of the quantity of acid originally present. With myristic acid, this is only the case if more than 0.06 gram is present. Lauric acid gives a maximum Polenske value with 0.3 gram present. In the case of the easily volatile acids, there is a rough proportionality between the Reichert-Meissl and Polenske values and the weight of acid present. The later fractions of the higher acids are only very slowly volatile in steam, but even fractions of a milligram are easily visible in the condensate. The fact that the Polenske value of the higher fatty acids is independent of the weight present is useful in the analysis of mixtures of fatty acids. The work of Dons (A., 1908, ii, 238; 1909, ii, 75) on the estimation of decaoic, lauric, and myristic acids in butter by distillation of the fatty acids in steam is described. If 1 gram of mixed fatty acids, is distilled in 100 c.c. of steam (measured as water), the partition coefficient of the volatile acids between distillate and condensate is given by $K = (m_1 \times 100)/m_2$, where m_1 is the weight of acids remaining in the flask and m_2 the weight distilling over. The partition of the different acids is unaffected by the other components of the mixture and depends on their volatility under the conditions of distillation and on the proportion present in the mixture. Dons found that if c_1 is the quantity reckoned as c.c. $N/10$ alkali hydroxide volatilising with the first 100 c.c. of water, $c_1 = \alpha r$, where r is the "volatility" and is equal to 0.244, 0.060, 0.016 for decaoic, lauric, and myristic acids respectively, and α

equals the percentage of the respective acid in the mixture. The partition coefficient is given by $K = (100 \times 1 - r)/r$. For decoic, lauric, and myristic acids K has the values 138.1, 733.3, and 2640, respectively. The general formula for r is $r = l_2/(Kl_1 + l_2)$, where l_1 is the weight of the acid phase and l_2 that of the condensed steam. From this it follows that the volatility of a fatty acid becomes greater with decrease in l_1 . This is borne out by Arnold's experiments. If a series of distillations of a given weight of fatty acid is undertaken, using each time 100 c.c. of aqueous condensate, the weight of a given acid distilling over at the n th distillation is given by $c_n = c_1(1 - r)^{n-1}$.

In practice, Dons's method is applied as follows. Using 0.5 gram of fatty acids, the Polenske value (p_{16}) of the sixteenth distillation is determined. It is considered that decoic and lauric acids have by this time been completely eliminated from the mixture, so that this value is due only to myristic and the higher acids. The latter have a constant figure of 0.2. Hence c_1 (myristic acid) $= p_{16} - 0.2$. Hence c_1 (myristic acid) $= c_{16}/(1 - r)^{15}$, where $r = 0.0704$. The percentage of myristic acid is given by $c_1/0.016$. Using 1 gram of fatty acids, 700 c.c. of distillate are then collected and the sum of the Reichert-Meissl and Polenske values for the first six distillations, and the Polenske value (p_7) for the seventh distillation obtained. The seventh distillate is free from decoic acid, as it no longer gives a Reichert-Meissl value; $p_7 - 0.2$ is therefore due entirely to lauric and myristic acids, c_1 for myristic acid is known, so $c_7 = c_1(1 - r)^6$. Hence the lauric acid in the seventh distillate is given by $p_7 - 0.2 - c_7$ and from this c_1 (lauric) and the percentage of lauric acid in the mixture can be calculated. The percentage of decoic acid can be obtained in a similar manner from the results of the first six distillations. Dons's method is interesting from a theoretical point of view, but cannot be considered established as a practical method of analysing fatty acid mixtures.

H. C. R.

A Micro-extraction Apparatus. FRITZ LAQUER (*Z. physiol. Chem.*, 1922, 118, 215—217).—An adaptation of an extraction apparatus for the detection of small quantities of lactic acid of the order 2—10 mg. in tissues.

S. S. Z.

Estimation of Oxalic Acid in Urine. E. MISLOWITZER (*Biochem. Z.*, 1921, 126, 77—81).—A comparison has been made of the old standard method of Salkowski and the new calcium acetate process of Bau (A., 1921, ii, 356) for the estimation of oxalic acid in urine. Bau's process was found inferior and especially so if more ether be used for the extraction of the oxalic acid and the extraction be carried out mechanically.

H. K.

The Relation between the Refractive Index and the Chemical Characteristics of Oils and Fats (Glycerides). G. F. PICKERING and G. E. COWLISHAW (*J. Soc. Chem. Ind.*, 1922, 41, 74—77).—An attempt has been made to discover a quantitative relation between the refractive index and the molecular weight, free fatty

acid content, and iodine value of the following oils : linseed, soja bean, cotton-seed, ground nut, palm-kernel, and coco-nut. The refractive index of each of these oils can be calculated with a high degree of accuracy from the formula $n_D^{20} = 1.4643 - 0.000066(S.V.) - 0.0096(A.V./S.V.) + 0.000117(I.V.)$ where $S.V.$, $A.V.$, and $I.V.$ are the saponification, acid, and iodine values respectively. The equation has a useful application in the commercial analysis of the above oils.

E. H. R.

Use of the "Silver Method" in the Estimation of Acetaldehyde. Its Application in the Estimation of other Aldehydes. A Convenient Method of Accumulation of Aldehyde and other Volatile Substances from Body Fluids. ROBERT FRICKE (*Z. physiol. Chem.*, 1922, **118**, 241—246).—Further details regarding the method described by Stepp and Fricke (this vol., ii, 236). Acetaldehyde can be removed from body-fluids by steam distillation.

S. S. Z.

The Detection of Aldol in the Urine of Diabetic Patients. ROBERT FRICKE (*Z. physiol. Chem.*, 1922, **118**, 218—223).—By distilling acid urine from diabetic patients small quantities of crotonaldehyde have been demonstrated in the distillate; possibly this is derived from the aldol present in the urine. The "dime-don" method of precipitation was employed.

S. S. Z.

Micro-estimation of Total Acetone in Urine. HEINRICH LAX (*Biochem. Z.*, 1921, **125**, 262—264).—An apparatus is pictured and a method described, suitable for clinical use, for the estimation of acetone in 2 c.c. of urine in ten minutes. Twenty-five c.c. of water, 2 c.c. of urine, and 3 drops of 80% acetic acid are distilled for four minutes into a second flask containing 2 c.c. of water and 2 drops of concentrated hydrochloric acid. The ammonia is thus retained and the acetone passes on and is absorbed by 30 c.c. of water kept at 0°. Ten c.c. of $N/20$ -iodine solution are added and 1 c.c. of 33% sodium hydroxide solution and after three minutes the excess of iodine after acidification is titrated with sodium thiosulphate.

H. K.

Estimation of Monobromocamphor. ELGAR O. EATON (*J. Ind. Eng. Chem.*, 1922, **14**, 24).—A weighed quantity of about 0.2 gram of monobromocamphor is dissolved in alcohol, 50 c.c. of $N/2$ alcoholic potash solution, and 25 c.c. of 0.4% alcoholic silver nitrate solution are added and the mixture is boiled under a reflux apparatus for ninety minutes; during the boiling, further quantities (25 c.c. in all) of the alcoholic silver nitrate solution are added through the reflux apparatus. After cooling, the mixture is diluted with water to 200 c.c., the solution decanted into a beaker, and the insoluble portion washed by decantation. This dilute alcohol solution is boiled for five minutes with the addition of 1 gram of zinc dust, filtered, the filtrate acidified with nitric acid, and treated with an excess of silver nitrate solution; the silver bromide formed is collected and weighed. The weight of silver bromide found

multiplied by 1.23 gives the corresponding amount of monobromocamphor. Potassium bromide is formed in the first part of the process, and not silver bromide.

W. P. S.

Extraction and Characterisation of Alkaloids. W. C. COLLEDGE (*J. S. African Chem. Inst.*, 1922, 5, 5—8).—A description of methods of purifying extracted alkaloids, methods of testing, general alkaloidal reagents, certain colour reactions, etc. Physiological tests may be applied when any doubt exists as to the nature of a supposed alkaloid.

W. P. S.

Extraction of Alkaloids from Viscera. W. C. COLLEDGE (*J. S. African Chem. Inst.*, 1922, 5, 3—5).—The Stas-Otto method is recommended; substances produced by putrefactive changes and extracted together with the alkaloids do not interfere with the identification of the latter, but if desired, the alkaloidal residue may be purified by dissolving it in a mixture of benzene and amyl alcohol, shaking the solution with dilute hydrochloric acid, separating the acid aqueous layer, extracting this with light petroleum, and then rendering the solution alkaline and extracting it with ether. For the detection of morphine, the alkaloidal residue is dissolved in acetic acid, the solution treated with lead acetate to precipitate meconic acid, and the morphine is extracted subsequently from the ammoniacal solution by means of amyl alcohol. In some cases, dialysis of the original material (stomach contents, etc.) affords a means of separating certain alkaloids such as strychnine and morphine.

W. P. S.

Iodic Acid as a Microchemical Reagent for the Detection of Organic Bases. L. ROSENTHALER (*Schweitz. Apoth. Zeit.*, 1921, 59, 477—479; from *Chem. Zentr.*, 1921, iv, 1055—1056).—The precipitates obtained with iodic acid and solutions of organic bases have in many cases characteristic forms, suitable for use in microchemical work. Strychnine, cinchonine, morphine, codeine, hydrastinine, pyridine, quinoline, and aniline give characteristic crystalline precipitates. Other alkaloids and organic bases may be distinguished by colorations.

G. W. R.

Identification of Alkaloids under the Microscope from the Form of their Picrate Crystals. BURT E. NELSON and HELEN A. LEONARD (*J. Amer. Chem. Soc.*, 1922, 44, 369—373).—The more commonly occurring vegetable alkaloids may be tentatively identified under the microscope by the form or habit of their picrate crystals prepared under standard conditions. A chart is given showing the crystalline structure of the picrates of twenty-five of the alkaloids. For the test, the aqueous solution of the separated alkaloid is slightly acidified with hydrochloric acid and a slight excess of a saturated solution of picric acid is added. The precipitated picrate is washed in a centrifuge and recrystallised from the smallest possible amount of warm 95% alcohol. The crystals are separated centrifugally and examined under a microscope without a cover-slip.

W. G.

Estimation of Carnosine in Muscle Extract. GEORGE HUNTER (*Biochem. J.*, 1921, 15, 689—694).—Consistent results can be obtained in the estimation of carnosine in muscle by the diazo-method of Koessler and Hanke (*A.*, 1920, ii, 67). Details are given of the method recommended by the author for preparing the protein-free extract, together with a few analytical results. Ox muscle is found to contain from 0.4% to 0.6% of carnosine.
W. O. K.

Pyrimidines. XCII. New Methods of Identifying the Pyrimidine, Thymine. TREAT B. JOHNSON and OSKAR BAUDISCH (*J. Amer. Chem. Soc.*, 1921, 43, 2670—2674).—The thymine molecule is completely destroyed at the ordinary temperature when subjected in aqueous solution to the oxidising action of the system, ferrous sulphate plus sodium hydrogen carbonate plus air. The products formed are pyruvic acid, acetylcarbinol, urea, and formic acid. The urea may be identified by its xanthidrol derivative and the formic acid by its reducing action. The pyruvic acid is best identified by its interaction with *o*-nitrobenzaldehyde in alkaline solution to form indigotin. For the detection of acetylcarbinol Baudisch's specific reaction (*cf.* *A.*, 1918, ii, 412) with *o*-amino-benzaldehyde is best used. Uracil and cytosine are also oxidised by the above reagent, but in neither case is any pyruvic acid obtained and hence the detection of this acid among the products of the oxidation is sufficient proof of the presence of thymine in the original material.
W. G.

Colorimetric Estimation of Uric Acid. Estimation of 0.03 to 0.5 mg. quantities by a New Method. J. LUCIEN MORRIS and A. GARRARD MACLEOD (*J. Biol. Chem.*, 1922, 50, 55—63).—The phosphotungstic acid reagent used by Folin and Denis (*A.*, 1913, ii, 162) is replaced by an arsenotungstic acid reagent, which produces a greater depth of colour. Uric acid is precipitated from urine or deproteinised blood by the addition of zinc chloride and sodium carbonate (*cf.* *A.*, 1916, ii, 456). After separation by centrifuging, it is dissolved in hydrochloric acid and the colour developed in the presence of sodium cyanide by the addition of the arsenotungstic acid reagent compared with a standard. The formation of precipitates is avoided by the use of sodium cyanide as the only alkali.
E. S.

A Modification of Folin's Colorimetric Method for the Estimation of Uric Acid. HENRY JACKSON, jun., and WALTER W. PALMER (*J. Biol. Chem.*, 1922, 50, 89—101).—The estimation is made by the method of Folin and Denis (*A.*, 1913, ii, 162), using a modified phosphotungstic acid reagent. The addition of sodium carbonate is omitted, the necessary alkalinity being furnished by the use of sodium cyanide. By these modifications a greater intensity of colour is obtained and the formation of precipitates avoided.
E. S.

General and Physical Chemistry.

A Modified Form of Double Slit Spectrophotometer. A. L. NARAYAN and G. SUBRAHMANYAM (*Phil. Mag.*, 1922, [vi], 43, 662—663).—This consists essentially of an electromagnetically maintained pendulum carrying a double slit, which is mounted in front of the collimator slit of the spectrograph. This form of spectrophotometer is free from the defects of Vierordt's type, giving a better method of regulating the brightness of the spectrum. It also possesses many of the advantages of the sector photometer.
W. E. G.

Molecular Refraction of some Molten Salts. G. MEYER and ADOLF HECK (*Z. physikal. Chem.*, 1922, 100, 316—333; cf. this vol., ii, 241).—The index of refraction of molten sodium nitrate, potassium nitrate, sodium hydroxide, and potassium hydroxide has been determined by the method of autocollimation at a series of temperatures between 320° and 440°. The following values of the refractive index are recorded: sodium nitrate, $643.9\mu\mu$, $n=1.479-1.75 \times 10^{-4}t$; $589.3\mu\mu$, $n=1.499-2.12 \times 10^{-4}t$; $579.1\mu\mu$, $n=1.487-1.75 \times 10^{-4}t$; $546.1\mu\mu$, $n=1.483-1.50 \times 10^{-4}t$; potassium nitrate, $643.9\mu\mu$, $n=1.525-3.12 \times 10^{-4}t$; $589.3\mu\mu$, $n=1.664-4.00 \times 10^{-4}t$; $579.1\mu\mu$, $n=1.556-5.75 \times 10^{-4}t$; $546.1\mu\mu$, $n=1.538-3.12 \times 10^{-4}t$; sodium hydroxide, $643.9\mu\mu$, $n=1.467-1.25 \times 10^{-4}t$; $589.3\mu\mu$, $n=1.458-8.75 \times 10^{-5}t$; $585.7\mu\mu$, $n=1.460-8.75 \times 10^{-5}t$; $558.9\mu\mu$, $n=1.471-1.12 \times 10^{-4}t$; potassium hydroxide, $643.9\mu\mu$, $n=1.453-1.0 \times 10^{-4}t$; $589.3\mu\mu$, $n=1.479-1.5 \times 10^{-4}t$; $585.7\mu\mu$, $n=1.479-1.33 \times 10^{-4}t$; $558.9\mu\mu$, $n=1.475-1.33 \times 10^{-4}t$. The density of molten sodium hydroxide and potassium hydroxide has been determined and the following values have been found: sodium hydroxide, 340°, 1.89; 400°, 1.86; 440°, 1.84; potassium hydroxide, 380°, 1.87; 420°, 1.83, and 440°, 1.81. The molecular refractivity calculated according to the formula $(n^2-1) \cdot M / (n^2+2) \cdot d$ is in all cases in keeping with the theoretical value. In the case of sodium nitrate, the degree of dissociation has been calculated from the molecular refraction and shown to be 64.7%.
J. F. S.

A New Band Spectrum of Oxygen. C. RUNGE (*Physica*, 1921, 1, 254—261).—A direct current, high voltage arc was operated in a cylinder through which a stream of oxygen was passed, and a group of bands photographed in the region λ 2200—4900 Å. Although closely crowded towards the ends of the region, the separate lines could be observed from λ 2980 to λ 3900. The vibration frequencies are given by an expression of the form $a+bn+cn^2$; for the first series, n is a whole number, and for the related series, negative fractional numbers differing by whole numbers. Formulæ representing seven bands are given.

CHEMICAL ABSTRACTS.

Structure of the Band Spectrum of Helium. W. E. CURTIS (*Proc. Roy. Soc.*, 1922, [A], 101, 38—64). An investigation of the structure of the red band near λ 6400, the green band near λ 5730, and a blue band near λ 4550 in the helium spectrum. The principal features of the structure are shown to be accounted for by the quantum theory of band spectra, which are considered to originate in the passage of molecules between stationary states characterised by constant values of angular momentum, such passage being possibly accompanied by a change in the configuration of the molecule, as assumed by Schwarzschild. A brief résumé of the theory is given. In each of the three bands a new type of series is found, which, although closely related to the others, has not yet received a theoretical explanation. Certain other departures from the theory are also noted. It is concluded that the spectrum is due to an unstable helium molecule having a moment of inertia equal to about 1.8×10^{-40} gram cm^2 . J. S. G. T.

The Spectra of Helium, Hydrogen, and Carbon in the Extreme Ultra-violet. J. C. McLENNAN and P. A. PETRIE (*Trans. Roy. Soc. Canada*, 1921, 15, iii, 15—25).—In identifying the wave-lengths obtained in vacuum grating spectra of helium, and possibly of hydrogen, regard must be paid to the possibility of certain of the recorded wave-lengths originating in carbon or mercury introduced into the discharge tube. The authors support the contention of Millikan (*Astrophys. J.*, 1920, 52, 47) that certain of the lines obtained by Lyman (A., 1920, ii, 207) in the Schumann spectrum of helium are due to carbon. In particular, the wave-lengths $\lambda\lambda$ 1931, 1657, and 1561 Å., together with others of less intensity, originate in the atoms of carbon. The series of wave-lengths the frequencies of which are given by $\nu = 4N(1/2^2 - 1/n^2)$ exists for the spark spectrum in helium. J. S. G. T.

The Corpuscular Spectra of the Elements. MAURICE DE BROGLIE (*Compt. rend.*, 1922, 174, 939—941; cf. A., 1921, ii, 232, 292, 615).—The spectra of the four elements silver, tin, gold, and uranium are given, showing the position of the lines which appear on the photographic plate. To identify the origin of the rays, a selective screen which will absorb certain radiations more strongly than others may be interposed between the source of the X-rays and the apparatus. Thus a screen containing the oxides of the rarer earths causes the almost complete disappearance of the β - and γ -rays of the K spectrum of tungsten. W. G.

A New Method of Absorption Spectroscopy. WALTER GERLACH and ERICH KOCH (*Ber.*, 1922, 55, [B], 695—697).—The sources of light usual in spectroscopy suffer under the drawback that they are either insufficiently intense or not so constant as is desirable. The defect can be overcome by using as source of light a wire which suffers disintegration by a high-tension condenser discharge. A battery of Leyden jars (capacity about 30,000 cm.) is connected with a spark gap of constant dimensions and the iron wire which is to be disintegrated (about 2 cm. long and 0.031 mm.

diameter). The battery is slowly charged from a small machine until an arc is struck across the gap, when the wire is disintegrated with a blinding light and the current is broken; a single discharge is invariably sufficient for spectroscopic purposes. The constant brightness of the source of light is guaranteed by the constant energy expended in producing it, which depends only on the dimensions of the spark gap and the thickness of the wire.

The disposition of the apparatus and the methods for its adjustment are fully described and figured in the original. H. W.

The Absorption Spectrum of Liquid and Gaseous Oxygen. W. W. SHAVER (*Trans. Roy. Soc. Canada*, 1921, **15**, iii, 7—14).—The absorption spectra of oxygen, liquid and gaseous, were determined by means of a quartz spectrograph, employing a spark between aluminium electrodes under water as the light source. In the visible region, eight absorption bands were observed in the case of liquid oxygen, and seven with gaseous oxygen. The mean wave-lengths of the bands in the former case were 6285, 5800, 5350, 4816, 4458, 3828, 3631, and 3461 Å. In the case of gaseous oxygen at 140 atmospheres pressure, the band at 4458 Å. was absent. At 107 atmospheres pressure, the bands at 6285, 5800, and 4816 Å. alone were observed. In the ultra-violet, four broad bands, about 30 Å. wide, were observed in the case of both liquid and gaseous oxygen. In the former case, the centres of the bands were located at 2795, 2731, 2681, and 2631 Å., and each band consisted of a fine set of symmetrical triplet bands. In the gaseous spectrum, the similar bands observed were slightly displaced towards the ultra-violet. It was shown that the bands were not due to the presence of nitrogen or ozone. J. S. G. T.

An Absorption Band Spectrum for Water in the Region of Wave-lengths of Several Decimetres. RICHARD WEICHMANN (*Ann. Physik*, [iv], **66**, 501—545).—A more detailed account of work previously published (this vol., ii, 5).

Studies in Catalysis. XV. Absorption Spectra of Triethylsulphonium Bromide in Various Solvents, in the Short Infra-red Region. HENRY AUSTIN TAYLOR and WILLIAM CUDMORE McCULLAGH LEWIS (T., 1922, **121**, 665—675).

Absorption Spectrum of Benzene Vapour and the Fundamental Magnitudes of the Benzene Molecule. VICTOR HENRI (*Compt. rend.*, 1922, **174**, 809—812).—The ultra-violet absorption spectrum of benzene vapour is composed of four series of superposed bands, which obey the formulæ, $1/\lambda = 37703 + 921.4n - (159p + 2p^2) + 2(m^2 - q^2)$; $1/\lambda = 37494 + 921.4n - (159p + 2p^2) + 2(m^2 - q^2)$; $1/\lambda = 37613 + 921.4n - (159p + 2p^2) + 2(m^2 - q^2)$; $1/\lambda = 37426 + 921.4n - 166p + 2(m^2 - q^2)$, where n has the values 0 to 7, p the values 0 to 6, and $q = m$ or $q = m \pm 1$ and $m = 1, 2 \dots 10$. Eight groups of bands correspond with the eight values of n . The positions of the heads of the bands in each group correspond with the different values of p , and the values of m give the fine structure of

each band. The intensity of the bands diminishes rapidly as p increases. The constants 921.4 and 159 represent the number of vibrations of the atomic nuclei or of the groups of atoms in the molecule. Hence the infra-red absorption spectrum is given by $1/\lambda = 921.4n - 159p$, and this is confirmed by the results of Coblenz and Pucciati. The moment of inertia of the benzene molecule is deduced as being 1.45×10^{-38} . From this the distance between the carbon atoms is calculated as being 1.85×10^{-8} cm. and the diameter of the molecule as 2.6×10^{-8} . Thus the benzene molecule is a very symmetrical edifice the movements of which obey the simple laws deduced for diatomic molecules, and it may be considered as being formed of two halves, $C_3H_3-C_3H_3$, which vibrate with respect to one another.

W. G.

Ultra-violet Absorption Spectrum of Phenol in Different Solvents. F. W. KLINGSTEDT (*Compt. rend.*, 1922, 174, 812—815).—A quantitative study of the ultra-violet absorption spectrum of phenol in different solvents shows that the spectrum of phenol dissolved in pentane or hexane, which the author calls the normal spectrum, comprises two regions in the ultra-violet. In the first, between $\lambda = 2860$ and 2400 there are three intense narrow bands, and in the second, between $\lambda = 2325$ and the extreme ultra-violet there are two broad bands. The absorption spectrum changes with the nature of the solvent, and from this point of view the solvents may be divided into two groups. The solvents of the first group, such as carbon tetrachloride, chloroform, ether, do not modify the general aspect of the absorption spectrum, only producing a displacement and a broadening of the bands. The solvents of the second group, such as methyl or ethyl alcohol or water, completely modify the absorption spectrum of phenol. In these solvents there is only a single very broad and uniform band. The absorption spectrum of pure liquid and solid phenol occupies a position intermediate between the two preceding types. W. G.

Fundamental Laws of Photochemistry. II. Influence of Cooling on the Absorption of Light by Dyes. P. LASAREV (*Z. physikal. Chem.*, 1922, 100, 266—270; cf. this vol., ii, 103).—The absorption of light of wave-lengths $\lambda = 514, 543, 559, 576, 595, 617$, and 644 by layers of collodion containing cyanin and pinacyanol has been measured at ordinary temperatures and at the temperature of liquid air. In the case of cyanin, it is shown that the simple absorption band becomes narrower, whilst in the case of pinacyanol the single band is resolved into two absorption bands at the temperature of liquid air.

J. F. S.

Application of Photo-electric Cells to the Measurement of the Light Absorption in Solutions. II. H. VON HALBAN and K. SIEDENTOPF (*Z. physikal. Chem.*, 1922, 100, 208—230; cf. A., 1921, ii, 145).—An arrangement for measuring the absorption of light by solutions is described, which depends on the use of photo-electric cells and in which variations in the intensity of illumination of a mercury lamp are eliminated by compensation

with two photo-electric cells. The cells are used only as a zero instrument. A rotating sector and a grey wedge are used to measure the weakening of the light. The greatest variation of an individual measurement of the absolute value of the extinction coefficient from the mean value does not exceed 0.5%. Measurements with the wedge permit of the identity of two objects in respect of their light absorption being established to 0.1% and differences can also be determined with the same accuracy. Measurements are recorded for solutions of potassium chromate in potassium hydroxide, equimolecular quantities of potassium chromate and copper sulphate in ammonia, azobenzene in alcohol, and anthracene in alcohol for the mercury lines 579 $\mu\mu$ and 254 $\mu\mu$.

J. F. S.

Colour and Chemical Constitution. XIII. Calculation of the Colour of Monocyclic Dyes. JAMES MOIR (*Trans. Roy. Soc. Sth. Africa*, 1921, 10, 35—39; cf. A., 1921, ii, 475).—Dyes which contain only one active colour ring do not fit into the scheme previously described (A., 1921, ii, 6) for those containing two such rings, but their colours are deducible by a similar factorial scheme, based on *p*-hydroxybenzyl alcohol (for which, and probably also for *p*-cresol, in faintly alkaline aqueous solution, $\lambda=290$). The factors in parenthesis are employed in conjunction with this value: replacement of carbinol hydrogen by phenyl (1.135); conversion of phenyl carbinol into a phthalein (1.060); replacement of ring hydroxyl- by amino-group (1.140); replacement of amino-hydrogen atoms by methyl (1.035). Values are thus predicted for certain compounds for which measurements are at present lacking. It appears that colour cannot be traced to lower terms than the presence of one benzene ring, with one ionisable group, usually with another active atom, and that the physical cause of colour is the periodic motion of an electron round a molecule. Although the sinuous orbit previously conceived (A., 1921, ii, 475) is now considered not to be very likely, the tautomeric pauses then assumed are a probable feature of the orbit. It is suggested that the colours of dyes containing two active colour rings (*loc. cit.*) may be based on 4:4'-dihydroxybenzhydrol ($\lambda=539$). The pink colours attributed to phenylphenolphthalein and to its analogue from salicylic acid (A., 1917, ii, 349; 1919, ii, 78) were due to impurity.

J. K.

Constitution and Colour. VIII. F. KEHRMANN (*Helv. Chim. Acta*, 1922, 5, 158—163; cf. A., 1921, ii, 476).—The effect of salt formation on the colour of basic nitrogen compounds is to lighten it if the unsaturated condition is destroyed, and to intensify it if the unsaturated condition persists (for example, in the cases of auramine base, azo-compounds, azomethines, and members of the quinoline, pyridine, acridine, phenazine, etc., series). If no change occurs in the degree of saturation (for example, ammonium, imonium, cyclonium, and diazonium compounds), the colour is also unchanged. Similar generalisations apply to basic sulphur and oxygen compounds, with the reservation that knowledge of the colour changes accompanying destruction of the unsaturated con-

dition in these cases is at present insufficiently specific. The behaviour of the group, $\cdot\text{CH}:\text{CH}\cdot$, is apparently analogous to that of the azo-group, since intensely coloured sulphates of diphenylcyclopentadiene (Borsche and Menz, A., 1908, i, 147) and of unsaturated ketones (Kehrmann and Effront, A., 1921, i, 348) have been described. The effect of salt formation on the colour of iodonium compounds is so far unknown. The above rules also apply to all those cases of salt formation from pseudo-bases (for example, carbinol bases) in which the nature of the changes is properly understood. J. K.

Fluorescence and Photochemistry. R. W. WOOD (*Phil. Mag.*, 1922, [vi], 43, 757—765).—The theory of Perrin (cf. A., 1918, ii, 418) is tested by exposing aqueous solutions of eosin, rhodamine, and fluorescein to a very intense beam of sunlight. The products, "photo compounds," are usually coloured, non-fluorescent substances, which are bleached by further action of the light. Their absorption bands have a totally different form from those of the original substances. Rhodamine is almost non-fluorescent at 100°, but decomposes under the action of light as rapidly as at ordinary temperatures. This is contrary to Perrin's theory. The relation between the rate of breakdown of eosin and the intensity of the light is at variance with the Bunsen-Roscoe law, whereas unstable non-fluorescent substances behave normally. The proportionality between the emission of fluorescence and the intensity of the exciting light holds over a wide range of intensity. W. E. G.

The Quantitative Determination of the Fluorescent Powers (the Spectro-fluorescometry) of Cellulose, Sugars, and other Substances. SAMUEL JUDD LEWIS (*J. Soc. Dyers and Col.*, 1922, 38, 68—76, 99—108).—An instrument has been designed by which the fluorescent powers of different substances can be quantitatively compared. It consists of a Hilger quartz spectrograph with the back of its camera modified so as to accommodate the lens of an auxiliary camera of the ordinary type directed towards the position usually occupied by the photographic plate of the spectrograph. The usual spectrum falls on to the paper, fabric, or other substance placed in the position of the photographic plate of the spectrograph. The auxiliary camera is directed on to the ultra-violet region only, and since the lens is of glass, any light transmitted by it must consist of visible light produced from the ultra-violet light by the fluorescent substance under examination. As a standard for comparison, Whatman No. 44 filter-papers were used and quantitative comparison was obtained by comparing the times required by the standard and the substance under examination to produce equal photographic effects in a given part of the spectrum. Great difficulties were met in obtaining suitably sensitive photographic plates, and it is suggested that the properties of gelatin may vary with the season of the year at which it is prepared. This method of investigation was applied to determine the fluorescent properties of many different textiles and woods, of papers made from pure cotton, flax, hemp, esparto, and ramie, of cellulose

derivatives, and of a number of carbohydrates, including several sugars, starch, and dextrin. The results are presented in the form of curves in which the fluorescence, expressed as a percentage of that of the arbitrary standard, is plotted against the wave-length of the incident light. Cellulose acetate was the most strongly fluorescent of the cellulose derivatives examined, the maximum strength being at λ 2500—2800, where it was seven times that of the standard. Cellulose nitrate is almost devoid of fluorescent properties. The sugars examined included xylose, dextrose, galactose, levulose, sucrose, maltose, and lactose, and indications were found of some relation between structure and fluorescent power. There is some evidence to suggest that the cellulose complex may contain groups more nearly related to dextrose than to *d*-fructose.

E. H. R.

The Spectra Structure of the Luminescence excited by the Hydrogen Flame. HORACE L. HOWES (*Physical Rev.*, 1921, 17, 469—474).—An investigation, with determinations of wave-lengths, of the spectra of the luminescence of air-slaked lime, and of certain phosphorescent sulphides ("strontium sulphide-bismuth sodium sulphate," "strontium sulphide-bismuth potassium phosphate," and a calcium sulphide compound containing both bismuth and fluorine as active elements), when partly bathed in a hydrogen flame, but at a temperature below red heat.

A. A. E.

Scattering of Light by Dust Free Liquids. II. W. H. MARTIN and S. LEHRMAN (*J. Physical Chem.*, 1922, 26, 75—88).—A continuation of previously published work (A., 1920, ii, 573). In the present paper, the relative intensity and polarisation of the light scattered by dust-free benzene, toluene, xylene, chlorobenzene, methyl alcohol, ethyl alcohol, *n*-propyl alcohol, *isobutyl* alcohol, *isoamyl* alcohol, water, and mixtures of carbon disulphide with ether, benzene with hexane, and hexane with *cyclohexane* have been measured with an accuracy greater than that obtained in earlier work. The increased accuracy was made possible by the use of cross-shaped containers with sealed in, flat, glass end plates in place of the bulbs used previously. Measurements of the light scattered by two-component liquid solutions show that the relative intensity of the scattered light is always somewhat greater than that calculated on the assumption that the scattered light is an additive property for the two liquids. Liquids which polarise the scattered light very far from completely show, on dilution, much more nearly complete polarisation. Measurements of the ratio of the intensity of incident light to that of scattered light have been made for liquids. The results show that ether and water scatter about one-tenth as much light as do the same weights of these liquids in the gaseous state. For benzene and its homologues, the ratio is about one-fifth. Measurements of the intensity of the scattered light for various wave-lengths show that this value varies inversely as the fourth power of the wave-length.

J. F. S.

Radiation and Chemical Action. T. W. J. TAYLOR (*Nature*, 1921, 108, 210; cf. following abstract).—An experimental refutation of Lewis's explanation that the velocity of inversion of sucrose by dilute acid is not increased by sunlight because the activating rays lie in the region of 1μ , and at this wave-length water would absorb the radiation almost totally in the first thin layer, so that the bulk of the liquid would remain unaffected (cf. Lindemann, A., 1920, ii, 743). A solution of sucrose containing hydrogen chloride was forced upwards in bright sunlight through extremely fine capillary jets; the liquid formed fine columns about 8 cm. high, which then broke up into clouds of small drops (about 0.015 cm. in diameter), the drops rising a further 40 cm. After falling during an average time of 0.68 second, the drops were collected. The rotation was only 0.24° less than that of an unexposed control portion of the solution. Even assuming that the radiation density at 1μ had been reduced, by absorption, inside each drop to 10^{-8} of its value, which is considered to be unlikely, a difference in polarimeter readings of about 13° would have been anticipated. A. A. E.

Radiation and Chemical Action. W. C. MCC. LEWIS (*Nature*, 1921, 108, 241; cf. preceding abstract).—An alternative reply to Lindemann's criticism (A., 1920, ii, 743) of the radiation hypothesis of chemical reactions, based on the relatively small absorption capacity of the reactant solutes (cf. Lewis and McKeown, A., 1921, ii, 623). A clear distinction must be drawn between photochemical and thermal conditions, the former involving an absorption coefficient term. It is argued mathematically that the photochemical fractional decomposition and the thermal value (about 0.01 of the magnitude of the former) would be inappreciably small in Taylor's experiments. Further, Taylor carried out no determination of the amount of radiation absorbed by the sugar in the solution; if this were small, no chemical change in excess of the thermal change would be anticipated. A. A. E.

Dependence of Velocity of Reaction on the Concentration in Photochemical Processes. M. VOLMER and K. RIGGERT (*Z. physikal. Chem.*, 1922, 100, 502—511).—The velocity of the dissociation of anthracene in hexane solution into a positive residue and an electron has been determined with the object of testing Luther and Weigert's (A., 1905, ii, 785) expression for the dependence of the velocity of photo-decomposition on the concentration. The expression has the form $v = dC/dt = kJ_0(1 - e^{-\epsilon C d})/d$, where C is the concentration, ϵ the extinction coefficient, d the thickness of the illuminated layer, J_0 the intensity of the light, and k a constant. The amount of change was measured by the saturation current. It is shown that the current experimentally observed and that calculated from the above-mentioned equation are in excellent agreement, thus proving the correctness of the formula.


J. F. S.

Theory of Induced Reactions. CHR. WINTHER (*Z. physikal. Chem.*, 1922, 100, 566—571).—A theory of induced reactions is

put forward which is based on the conversion of oxygen into ozone in the presence of zinc oxide by ultra-violet light and other induced reactions. In the case mentioned, it is assumed that the light is absorbed by the zinc oxide and the light energy is given out again in the form of very short wave-length radiation which is photochemically absorbed by the oxygen. J. F. S.


The Action of Light on Silver Bromide. ERNST JOHANNES HARTUNG (T., 1922, 121, 682—691).

Action of Light on Uranium Salts. J. ALOY and E. RODIER (*Bull. Soc. chim.*, 1922, iv, 31, 246—249).—Under the influence of light, uranyl salts are converted into uranous salts in the presence of the acid entering into the constitution of the salt and a readily oxidisable substance such as alcohol. The reaction which occurs is as follows:




Two molecules of acid are necessary for each molecule of uranyl salt. In this way, the fluoride, chloride, bromide, iodide, and sulphate have been prepared, and an aqueous solution of the nitrate. If an insufficient amount of acid is present, basic salts are obtained. If the reaction is allowed to proceed further, a black precipitate of uranous hydroxide is obtained. W. G.

Photolysis of Uranyl Oxalate and Uranyl Acetate. EMIL BAUR and A. REBMANN (*Helv. Chim. Acta*, 1922, 5, 221—239).—If photolysis consists of two processes respectively akin to anodic oxidation and cathodic reduction, the presence of a suitable third substance should affect the quantitative, and possibly the qualitative, results of the process. Thus, the photolysis of oxalic acid by uranyl salts accords with the scheme:



which is supported by the decomposition of glyoxylic acid under similar conditions (A., 1919, ii, 264):



In the latter reaction, no evidence could be obtained of the formation of glycollic acid, or of formaldehyde, derived from its decomposition in the photolysis of uranyl oxalate. On the other hand, the influences of mercuric chloride, potassium iodide, ferrous sulphate, quinol, uranyl chloride, and sodium uranyl oxalate respectively in increasing the proportion of carbon dioxide formed in comparison with carbon monoxide are explicable in accordance with the theory. Sodium sulphite has no effect (compare Hatt, A., 1918, ii, 143; Trümpler, A., 1916, ii, 9), and formic acid only gives rise to excess of carbon dioxide in the later stages of the reaction, probably because the formate- takes place independently of the oxalate-photolysis. The gas produced in the photolysis of uranyl acetate (A., 1918, ii, 143), and considered to be hydrogen, is now shown to have been carbon monoxide originating from glyoxylic acid, frequently present as an impurity in acetic acid. Two reactions occur

during the photolysis of uranyl acetate: (1) true photocatalytic decomposition into methane and carbon dioxide (Fay, A., 1896, i, 464), (2) non-catalytic oxidation of acetate-ions to ethane and carbon dioxide by uranyl salts, which only occurs in acetic acid solution with a sufficient concentration of acetate-ions. The photolysis is favoured by the presence of free acetic acid, but restricted by chloridions and mercuric-ions to some extent, almost entirely by ferric salts or formic acid. Such a reaction cannot be looked on as, for example, a unimolecular decomposition of the uranyl acetate complex by light, but as an action of the activated uranyl complex on surrounding molecules by interchange of electrons. In other words, it is always an oxidation-reduction process, and so closely related to electrolysis. Probably Fay's reaction is connected with Kolbe's electrolytic decomposition of acetates, acetyl peroxide and atomic hydrogen being the respective anodic and cathodic products, which then interact and form methane and carbon dioxide. Since the photolytic oxidation of acetic acid to ethane and carbon dioxide, and of glyoxylic acid to carbon monoxide and carbon dioxide, can be reproduced electrochemically only by the use of a considerable potential difference, it is concluded that the energy quantum absorbed by the uranyl complex must also have such a value as would suffice, not only for the production of oxygen, but also for the formation of peroxides. The formation of uranyl peroxide, observed by Usher and Priestley (A., 1906, ii, 881), is recalled in this connection. J. K.

Photolysis of Uranyl Oxalate. EMIL BAUR and HAGGEN-MACHER (*Z. physikal. Chem.*, 1922, 100, 36—41).—The action of sunlight on aqueous solutions of uranyl sulphate and oxalic acid has been investigated. The products of the reaction are found to be formic acid, carbon dioxide, and carbon monoxide. The amount of formic acid formed is very small, and it is held that it must remain small because it is used up as fast as it is produced in reducing the uranyl-ion to uranous-ion. The gas evolved contains a slight excess of carbon dioxide above that required for the simple stoichiometric relationship $\text{CO}_2:\text{CO}$. The formation of formic acid is regarded as a cathodic reduction of carbon dioxide, thus: $\text{CO}_2 + 2\text{H}^+ + \ominus\ominus = \text{H}\cdot\text{CO}_2\text{H}$. J. F. S.

Action of Ultra-violet Light on Gels. EDWARD O. HOLMES, jun., and WALTER A. PATRICK (*J. Physical Chem.*, 1922, 26, 25—41).—The action of ultra-violet light from an iron arc on silica gels, which had been impregnated with acetone, acetic acid, and nitric acid, respectively, and also on celluloid, has been investigated by a tensimetric method. The gel was placed in both bulbs of the tensimeter, and one was exposed to ultra-violet light whilst the other was kept in the dark and the difference in pressure measured. It is shown that the gels give off gaseous products, provided the liquid contained in the gel is decomposed photochemically into gaseous products which are not adsorbed by the gel. Thus increase of pressure was observed in the case of gels containing acetic acid and acetone, but not in the case of nitric acid, for the decomposition

products of nitric acid are strongly adsorbed by silicic acid gel. The gaseous products consist of a mixture of the vapour of the organic liquid itself along with those gases resulting from the photo-chemical decomposition of the liquid. Celluloid behaves in a similar manner under the influence of ultra-violet light. The mechanism of the liberation of the adsorbed liquid is explained as follows. In the gel there are a large number of pores partly filled with the adsorbed substance under great negative pressure and therefore having a very low vapour pressure. The negative pressure in the film is caused by the surface tension acting along the sides of the pores and round their circumference. The force of the surface tension is reduced by the action of ultra-violet light owing to its decomposition of the liquid of the film with the formation of gas bubbles, which prevent the film from completely wetting the surface of the pores, resulting in a decrease of the negative pressure and subsequently an increase in the vapour pressure of the adsorbed substance, and consequent evaporation of some of the liquid. The same mechanism explains the liberation of gases by celluloid under the influence of ultra-violet light, only here the case is complicated by the fact that light decomposes the structure of the gel itself (cellulose nitrate) as well as the solvent in the pores. Hence the reason for celluloid turning brown and becoming brittle under the action of light is evident.

J. F. S.

β -Ray Spectra and their Meaning. C. D. ELLIS (*Proc. Roy. Soc.*, 1922, [A], 101, 1—17).—The shortest wave-length that has hitherto been measured by the crystal method is 0.07 Å.U., which is greater than the wave-lengths of γ -rays emitted by many radioactive substances. A method based on the quantum theory has been developed for the measurement of the wave-lengths of such rays, and has been applied to the cases of the γ -rays of radium-*B*, radium-*C*, and thorium-*D*. The method involves the measurement of the energies of the different lines in the natural β -ray spectrum of the element in question, and the energy of the corresponding line in the excited spectra of a substance of neighbouring atomic number. The numerical results obtained support the view that γ -rays are emitted from the nucleus. The quantum theory is probably applicable to the nucleus, and a part, at least, of the structure of the nucleus is expressible in terms of stationary states.

J. S. G. T.

Coloration and Luminescence produced by the Action of Becquerel Rays. STEFAN MEYER and KARL PRZIBRAM (*Z. physikal. Chem.*, 1922, 100, 334—336).—A number of examples are recorded which show that the coloration, brought about in glass, quartz, and similar materials by exposure to radium rays, may be removed entirely or in part by prolonged heating at comparatively low temperatures. It is also shown that thermoluminescence is exhibited on heating such coloured material at comparatively low temperatures. Thus a piece of colourless glass which has been turned brown by the rays and has remained unchanged for years in daylight on heating at 100° for twenty-four hours became con-

siderably lighter in colour, after several days at 130—140° it showed a violet tinge, and at 150—200° for several days it was definitely violet. Violet coloured glass is more resistant. Quartz which had been coloured brown became colourless after several days' heating at 150—200°. Sapphire which was originally light blue and had become topaz coloured by exposure to radium rays, after heating at 60° for a day became yellow, after a further two days at 80—90° it was pale yellow with a blue tinge, and after several days at 150—200° it regained its blue colour. A piece of kunzite which had become green in colour when heated for nine hours at 90° showed a feeble thermoluminescence, which disappeared in three hours. By heating at 170—180°, thermoluminescence was again shown, and the green colour had become much paler in half an hour. The same piece of material was then slowly heated at 140° without luminescence appearing, but at 160° weak luminescence appeared which became stronger as the temperature was raised to 185°, and after two hours' heating at 190° the luminescence failed and the kunzite had regained its original lilac colour. J. F. S.

The Existence of the New Radioactive Element Uranium-V described by Piccard and Stahel. OTTO HAHN (*Physikal. Z.*, 1922, 23, 146—150).—The decrease in activity of a number of uranium-X preparations has been determined to test the observations of Piccard and Stahel (*A.*, 1922, ii, 185). The observed activity was the sum of the exponentially decreasing activity of pure uranium-X and a constant α activity due to the presence of ionium. No evidence of the existence of uranium-V was found. W. E. G.

Existence of Isotopes of the Disintegration Products of Actinium and the Meitner Nuclear Model. MAXIMILIAN CAMILLO NEUBURGER (*Z. anorg. Chem.*, 1921, 120, 150—158).—The actinium family shows many irregularities when compared with the radium and thorium families, which it is suggested are due to the existence of unknown isotopes. The production of isotopes by branching at radioactinium is probable since this element gives α -particles with two ranges. A scheme is given for the disintegration of radioactinium in which the disintegration process, $\alpha'-\beta-\alpha-\beta$, is assumed (cf. Meitner, *A.*, 1921, ii, 293). W. E. G.

Electrochemical Behaviour of Liquid Sodium Amalgams. THEODORE W. RICHARDS and JAMES BRYANT CONANT (*J. Amer. Chem. Soc.*, 1922, 44, 601—611).—A number of improvements are described in the electrolytic preparation and the subsequent manipulation of pure liquid sodium amalgam, as well as in the apparatus for the measurement of the *E.M.F.* of sodium amalgam concentration cells with aqueous electrolytes. The *E.M.F.* of many concentration cells of liquid sodium amalgam has been measured at 25°, and a few cells also at 15° and 35°. The deviations of these potentials from the simple concentration law were found to be greater in sodium amalgam than with any other mercurial solution hitherto investigated in detail. The heats of transference

of sodium from one amalgam to another calculated by the Helmholtz equation are found to be unusually large. J. F. S.

Solid Thallium Amalgams and the Electrode Potential of Pure Thallium. THEODORE W. RICHARDS and CHARLES P. SMYTH (*J. Amer. Chem. Soc.*, 1922, **44**, 524—545).—Pure thallium in compact form is definitely shown to possess at 20° an electrode potential 2.1 millivolts higher than saturated thallium amalgam. When immersed in a *N*-solution of thallous ions and connected with a normal calomel electrode, the total potential 0.6192 is indicated at 25°. The potential of pure thallium is not influenced by quenching the metal, which is so soft that no important strain can exist in it. α -Thallium is the only phase of the pure metal which is stable at ordinary temperatures. Finely divided electrolytic thallium sponge gives a potential 0.6 millivolt higher than the compact fused form. The difference is probably due to the fine state of division, and disappears on long keeping. The difference of potential shown by β -thallium, if it could exist at ordinary temperatures, would probably be nearly three times as great. Solid thallium amalgams may be made having potentials anywhere between that of pure thallium and the 2-phase amalgam. As mercury is added, the potential decreases at first, then remains constant, between 4% and 10% of mercury, and later decreases again until at about 15% of mercury the liquid phase becomes permanent at 20°. Crystals of solid amalgam separated centrifugally from the liquid containing more mercury showed as much as 20% of mercury, but some of this was undoubtedly adhering mother-liquor. The densities of solid thallium amalgams indicate an increase in volume of thallium on amalgamation. The corresponding curve shows a slight inflection at about 5% of mercury. The hardness of solid thallium amalgams increases with added mercury until about 5% is present, when it begins slowly to diminish. From these phenomena it is inferred that α -thallium dissolves mercury, increasing in hardness and volume up to about 5% of mercury. When more mercury is added, another solid phase of about the same hardness and volume, but containing more mercury, appears. These two phases appear to exist mixed together in equilibrium, over the range from about 5% of mercury to 10% at 20°. With more than about 10% of mercury the α -solid solution ceases to exist and the other phase continues to dissolve mercury in solid solution until it is saturated, with perhaps 15% of mercury. The other solid phase is probably a solid solution of mercury in β -thallium, which appears in definite crystalline form, making the amalgam friable after the admixture of α -thallium has been eliminated. β -Thallium can exist at ordinary temperatures only in the presence of much mercury, but it is always the form present in equilibrium with the liquid amalgam. Pure thallium melts at 303.5° and has a transition point at 235.3°. J. F. S.

Electromotive Behaviour of Aluminium. ROBERT MÜLLER and FRANZ HÖLZL (*Z. anorg. Chem.*, 1921, **121**, 103—109).—The potential of aluminium and amalgamated aluminium has been

measured in solutions of aluminium chloride, potassium chloride, hydrochloric acid, and sodium hydroxide of various concentrations and in mixtures of hydrochloric acid and aluminium chloride of various concentrations. The values of the potential are found to be in qualitative agreement with the Nernst theory. It is shown, from a comparison of the potentials of pure and amalgamated aluminium in acid and alkaline electrolytes, to be probable that the activation by means of mercury and the apparent anomaly of the potential of pure aluminium is occasioned by a protecting layer of hydroxide. In concentrated hydrochloric acid (6.8 *N*), pure aluminium is more electropositive than the amalgam; cf. Dhar (*ibid.*, 1921, 118, 75). J. F. S.

Acidifying Action of Hydroxy-organic Compounds on Boric Acid and Molybdic Acid. E. RIMBACH and P. LEY (*Z. physikal. Chem.*, 1922, 100, 393—407).—The authors have measured at 18° the hydrogen potential of solutions of boric acid, molybdic acid, glycollic acid, lactic acid, malic acid, citric acid, tartaric acid, propylene glycol, glycerol, mannitol, and dextrose and also mixtures of the two inorganic acids with various concentrations of each of the organic compounds. The hydrogen-ion concentration has been calculated in each case, and the increase in hydrogen-ion concentration occasioned by the mixing. The results are represented by curves and tabulated. The increase in the acidity is attributed to complex formation in which the hydroxyl group is operative. In the case of boric acid, tartaric acid produces a much greater increase than the monohydroxy-acids, whilst in the mixtures of molybdic acid the reverse is the case. Similar results are obtained with the non-acidic hydroxy-compounds. With molybdic acid, glycerol increases the acidity most, whilst mannitol has a smaller action than propylene glycol, but with boric acid, mannitol and dextrose have the most pronounced action, whilst propylene glycol has a greater action than glycerol. J. F. S.

Thermodynamic and Electrokinetic Potential Difference at the Surface of Two Liquids. H. FREUNDLICH and A. GYEMANT (*Z. physikal. Chem.*, 1922, 100, 182—196).—The thermodynamic potential, ϵ , and the electrokinetic potential, ζ , of the liquids phenol, guaiacol, benzonitrile, and aniline against aqueous solutions of a number of electrolytes have been measured. The ϵ -potential was measured by Beutner's method and the ζ -potential determined from the microscopic observations on the velocity of kataphoresis of the liquid drops. The two potentials are shown to be very largely independent of one another, a fact in keeping with the views of Freundlich (*Kolloid Z.*, 1921, 28, 240). The negatively charged drops are generally discharged and recharged oppositely by the kations, particularly strongly by multivalent kations such as calcium and aluminium, whereas such an effect does not occur in connexion with the ϵ -potential. A certain similarity of action is observed in the case of some organic ions; the kations of basic dyes and the alkaloids have a positive effect on

both potentials, whilst the anions of organic acids have a negative action. Exceptions and great variations in the dimensions of the action are, however, to be observed. The regularities previously published (*loc. cit.*) are confirmed. The partition relationship of the ions is determinative of the ϵ -potential, the organic liquid phase being more strongly positive or negative the more soluble the kation or anion is in the liquid. In the case of the ζ -potential, the adsorption and valency of the kations are responsible for the discharge and positive recharging of the drops; thus, for example, the kations of the basic dyes and the multivalent kations have a particularly strong action. All four liquids are negatively charged towards water in respect of their ζ -potential, even aniline, despite its strongly basic character. J. F. S.

Measurement of Overvoltage. HERMAN V. TARTAR and HARMON E. KEYES (*J. Amer. Chem. Soc.*, 1922, **44**, 557—567).—The conclusions of earlier investigators that overvoltage increases with time and current density and varies with the nature of the electrode surface have been confirmed. A comparison, based on the transition point for hydrogen-zinc deposition from solutions containing sulphuric acid and zinc sulphate, has been made by the two methods used for overvoltage measurement. The data obtained show that the commutator method gives erroneous results; when gas is being evolved at the experimental electrode, the error is large and varies with the speed of rotation of the commutator. The direct method has been found to give more consistent results and to be the more accurate method; the criticism that it is subject to variation due to electrode spacing and size and shape of the containing vessel is shown to be invalid. The existence of an appreciable transfer resistance at the surface of the electrodes examined is doubtful. While gas is being evolved, it is evidently not of the magnitude which has been stated. Evidence is advanced which indicates that the ordinary hydrogen overvoltage tables, obtained by using pure acid solutions as electrolyte, do not show the current densities necessary for the deposition from acid solutions of metals having a greater *E.M.F.* than hydrogen. J. F. S.

Low-voltage Standard Cells. J. OBLATA (*Proc. Phys. Math. Soc. Japan*, 1921, [iii], **3**, 136—149; cf. *ibid.*, 1921, [iii], **3**, 64).—The properties of various cell combinations were studied in order to obtain a satisfactory low-voltage standard. The following two were the most reproducible and constant: Cd amalgam | CdI_2 | PbI_2 | Pb amalgam, and Cd amalgam (10%) | $\text{CdSO}_4 \cdot 8/3\text{H}_2\text{O}$ | Cd amalgam (dilute). The *E.M.F.* of the former is 0.09838 at 20° and that of the latter is of the order of 0.01 volt, giving different values according to the concentration of the dilute amalgam. The temperature coefficients were determined between 15° and 30°. The heats of formation of cadmium chloride, iodide, and bromide were computed by applying the Gibbs-Helmholz equation. They were found to be 94090, 49340, and 74700 cals., respectively.

CHEMICAL ABSTRACTS.

Modifications of Kohlrausch's Method for the Measurement of the Conductivity of Electrolytes. MARCEL ABRIBAT (*Bull. Soc. chim.*, 1922, iv, 31, 241–245).—The ordinary telephone receiver used for determining the zero point on the Wheatstone bridge is replaced by a galvanometer. In order to permit of this, the current passing through the galvanometer is converted from alternating to direct by means of a thermionic valve. The arrangement of the apparatus is sketched. To make the readings still more exact, an amplifier may be used. W. G.


Questions Relating to the Constitution and Function of Electrolytes. W. MADELUNG (*Annalen*, 1922, 427, 35–109).—A theoretical enquiry into the constitutional causes which determine the formation and stability of electrically charged atomic aggregates (ions) in crystals and in solution. The form of the paper does not admit of satisfactory abstraction, and the original should be consulted in order to obtain a just appreciation of the comprehensive character of the author's views.

Hantzsch's theory of ψ -isomerism is discussed and extended. The cyanogen compounds described in the preceding paper (cf. this vol., i, 438) are ψ -acids in their non-ionisable forms, and true acids in the forms corresponding with their salts: disodium cyanamide is $\text{Na}\cdot\text{N}\cdot\text{C}\cdot\text{N}\cdot\text{Na}$, sodium dicyanamide is $\text{Na}\cdot\text{N}\cdot\text{C}\cdot\text{N}\cdot\text{CN}$, and sodium cyanoform is $\text{Na}\cdot\text{N}\cdot\text{C}\cdot\text{C}(\text{CN})_2$. Corresponding with ψ -acids and ψ -bases, a class of ψ -salts can be recognised as falling within the same view of ψ -isomerism. Aluminium and stannic chlorides are not ionised as such; they are therefore ψ -salts. In general, the univalent and some bivalent metals tend to form salts, whilst most of the ter- and quadri-valent metals form ψ -salts. The sodium compound of acetyl acetone is an electrolyte and non-volatile, like sodium chloride, whilst the aluminium derivative is a non-electrolyte and is volatile, like aluminium chloride: both the sodium compounds are salts, and both the aluminium compounds ψ -salts. In order to form ions, the ter- and quadri-valent metals have to combine with other ions or molecules to form centrically orientated complexes: $[\text{Fe}(\text{H}_2\text{O})_6]^{+++}$, $[\text{Fe}(\text{CN})_6]^{----}$, $[\text{Cr}(\text{NH}_3)_6]^{+++}$, $[\text{SnCl}_6]^{--}$.

Ionisation therefore is largely a matter of the formation of centrically orientated complexes. In acids, it is assisted by the accumulation of a number of electro-negative peripheral groups around the nuclear atoms, for example, $[\text{Fe}(\text{CN})_6]^{----}$ and $[\text{SnCl}_6]^{--}$. In bases, the accumulation of hydrogen atoms either as such, as amino- or hydroxyl-groups, or as water or ammonia molecules, has a similar effect; cf. $[\text{NH}_4]^+$, $[\text{PH}_4]^+$, $[\text{Cr}(\text{NH}_3)_6]^{+++}$, etc. Even carbon may become the central atom of an ion if sufficiently loaded with electrically powerful groups; the ion of guanidine is represented by the formula $[\text{C}(\text{NH}_2)_3]^+$, which expresses its strongly basic character much better than the customary formula based on the integral valencies of the organic chemist. *tert.*-Butyl iodide, $(\text{CH}_3)_3\text{CI}$, is better termed trimethylcarbonium iodide, as it is ionised in solution in liquid sulphur dioxide, a remarkable instance

of the tendency of a large accumulation of hydrogen atoms to promote ion-formation; the ions $[\text{C}(\text{CH}_3)_3]^+$, $[\text{C}(\text{NH}_2)_3]^+$, and $[\text{CO}_3]^-$ are strictly analogous.

Apparently simple ions, H^+ , OH^- , NO_3^- are subject to the same laws regarding the tendency to centric orientation and the influence of the peripheral groups. Hydrogen- and hydroxyl-ions are invariably combined with at least one molecule of water: $[\text{OH}_3]^+$, $[\text{H}(\text{OH})_2]^+$, and often with more. By assuming a further degree of hydration, chains may be produced which are strongly reminiscent of the structure of the crystalline compound :



The electrical influence of peripheral groups is often felt, not only in the circumjacent atoms, but also in more remote parts of the molecule. This phenomenon is termed induction; it is fully discussed and illustrated by many examples in the original paper.

C. K. I.

[Theory of Strong Electrolytes.] SVANTE ARRHENIUS (*Z. physikal. Chem.*, 1922, 100, 9—35).—A theoretical paper in which the author discusses the theory of strong electrolytes as put forward by Ghosh (T., 1918, 117, 449, 627, 707; 1920, 117, 823, 1390; this vol., ii, 125). The various calculations made by Ghosh in connexion with the dissociation of strong electrolytes and the more complete calculations of Partington (A., 1920, ii, 151) are considered, and in connexion with those of the former author it is shown that they are not always consistent, nor have they always the deep significance which Ghosh places on them. For example, in a comparison of the Arrhenius formula, $i=1+(n-1)\alpha$, with Ghosh's formula, $i=n(1+\frac{1}{2}\log_e \alpha)$, calculations are put forward by Ghosh which indicate the superiority of the second formula. The present author points out that in obtaining values from the second formula values which had been corrected for viscosity are used in the case of salts of the potassium chloride type, but not for those of the calcium chloride type, whilst with the former formula no correction was applied anywhere. The author concludes that could the Ghosh hypothesis be applied it would be a distinct advance on the previous hypotheses, but, unfortunately, it cannot be applied without great modifications.

J. F. S.

The λ -V Curves of Non-aqueous Solutions of Binary Electrolytes. P. WALDEN (*Z. physikal. Chem.*, 1922, 100, 512—531).—A theoretical paper in which the influence of the viscosity, dielectric constant, and mobility on the form of the λ -V curves is discussed.

J. F. S.

Dissociation and Hydrolysis Equilibria in Solutions of Salts of Carbonic and Boric Acids. HEINRICH MENZEL (*Z. physikal. Chem.*, 1922, 100, 276—317).—From electrometric measurements of mixtures of sodium carbonate and hydrogen carbonate in various concentrations at 18°, the second dissociation constant of carbonic acid has been determined and the value $K_2=6.2 \times 10^{-11}$ found, which agrees well with the value found by Auerbach and

Pick (*Arbt. Kais. Gesundh.*, 1911, **38**, 2). The hydrolysis of sodium carbonate has been calculated from the results, and the following values are recorded: $0.2M$, 1.12%, $0.1M$, 1.76%, $0.05M$, 3.0%, $0.02M$, 5.5%, $0.01M$, 8.3%, and $0.005M$, 10.65%. The dissociation constant for boric acid has been determined at 18° by *E.M.F.* measurements of solutions of borax and sodium metaborate, and the value $K_B = 5.7 \times 10^{-10}$ obtained. This value is much smaller than that recorded by Walker (T., 1900, **77**, 17), but is in keeping with the values recorded by Lundberg and Lundén at other temperatures. Concentrated solutions of borax and metaborate show, in consequence of the formation of polyborate ions, a somewhat stronger alkalinity than would be expected from the dissociation and hydrolysis. The degree of hydrolysis of sodium metaborate has been determined, and the following values have been obtained: $0.4M$, 0.50%; $0.2M$, 0.57%; $0.1M$, 0.76%; $0.04M$, 1.22%; $0.02M$, 1.7%, and $0.01M$, 2.3%. The equilibrium sodium carbonate + borax \rightleftharpoons metaborate + bicarbonate has been experimentally examined in dilute solutions, and also in two cases for concentrated solutions where the polyborate formation was taken into account.

J. F. S.

Calculation of the Second Dissociation Constant of Dibasic Acids from the Hydrogen-ion Concentration. MARCEL DUBOIX (*J. Chim. physique*, 1921, **19**, 179—187).—Two methods are given for calculating the second degree of dissociation of dibasic acids, which are applicable to solutions containing 0.01—0.005 gram-mol. per litre. At these concentrations, the determination of the hydrogen-ion concentration presents no difficulties. It is shown that Noyes's formula can be rigorously applied only to solutions less concentrated than 0.001 gram-mol./litre. Making use of known hydrogen-ion concentrations, the following values have been calculated: tartaric acid, $K_2 = 32 \times 10^{-6}$ at 25° , $K_2 = 28.4 \times 10^{-6}$ at 76° ; malic acid, $K_2 = 4.1 \times 10^{-6}$ at 100° ; succinic acid, $K_2 = 2.5 \times 10^{-6}$ at 25° , and $K_2 = 1.5 \times 10^{-6}$ at 100° . J. F. S.

Behaviour of the Photo-haloids in a Direct Electric Current. C. TUBANDT and GERHARD ESCHENHAGEN (*Z. physikal. Chem.*, 1922, **100**, 489—501).—The effect of passing a direct electric current through photo-haloids has been investigated. It is shown that both the synthetic photo-haloids and those which have been formed by the action of light are simple electrolytic conductors; the presence of colloidal metals, whether adsorbed or mechanically mixed, does not affect the character of the electrical conductivity. Metallic silver may be separated electrolytically from strongly illuminated dry silver chloride, and also from synthetic photo-chlorides and bromides, but not from photo-iodides even when these are mixed with large quantities of colloidal silver. The colloidal silver in synthetic photo-haloids and also that in illuminated silver haloids is unaffected by a direct current, but the silver-ions, as in the case of pure silver haloids, wander uninfluenced by the silver particles toward the electrode; a displacement of the silver

particles is not observed even when the current is allowed to flow for prolonged periods.

J. F. S.

Cathodic Deposits from Mixed Solutions of Two Simple Metallic Salts. WALTHER HILDEBRANDT CREUTZFELDT (*Z. anorg. Chem.*, 1921, **121**, 25—52).—Mixtures of salts of the following pairs of metals, copper-cadmium, copper-lead, copper-iron, silver-copper, silver-cadmium, and silver-zinc, have been electrolysed by various currents and voltages and binary, fine-grained, crystalline deposits obtained. It is shown that the composition of the deposit changes continuously with the current density and the composition of the mixture. By changing these factors, a series of mixed crystals may be deposited which varies from one pure constituent to the other constituent. Measurements of the *E.M.F.* of the deposit show that they are readily oxidised, and when placed in water they lose the more electropositive constituent in a few hours. In order that a deposit of a given concentration should be obtained, it is necessary that the current density should be increased with increasing metal concentration in the solution. A more vigorous agitation of the solution acts in the same sense as an increased current density, whilst an increased temperature acts in the opposite sense. The dependence of the *E.M.F.* of the deposit on the composition is considered in connexion with Tammann's theory. The partition of both atom types in the electrolytic deposit is not the normal one observed in mixed crystals from molten mixtures, but a completely unordered one. The probability for the unprotected atoms of the electropositive metal is given by $\Sigma w = p^{s-1} \cdot q + p^{2s-4} \cdot q^2 + p^{3s-7} \cdot q + \dots$, in which p is the molecular fraction of the less electropositive metal, q that of the electropositive metal, and $s-1$ the number of atoms of the less electropositive metal necessary to protect one atom of the more electropositive metal from the action of reagents. The fraction α of the unprotected atoms of the more electropositive metal is given by the equation $\alpha = (q - \Sigma w)/q$. The *E.M.F.* is assumed to be proportional to the number of unprotected atoms. It is shown that the potential-composition curve is determined by a whole number s . Four copper atoms are necessary to protect a single cadmium or lead atom, seven silver atoms are required for a single copper atom, and twenty-two copper atoms are necessary to protect a single iron atom.

J. F. S.

Determination of the Specific Heat of Solid Substances by the Adiabatic Electrical Method. ERNST COHEN and A. L. TH. MOESVELD (*Z. physikal. Chem.*, 1922, **100**, 151—158).—The specific heat of cadmium sulphate has been determined by an adiabatic electrical method previously described (this vol., ii, 156). The calorimeter liquid used was bromoform (specific heat 0.1241) and three determinations, using 823 and 676.5 grams respectively of $\text{CdSO}_4 \cdot 8/3\text{H}_2\text{O}$, were carried out. The value 0.200 between 17° and 21° was obtained, which agrees well with 0.2008 found by Cohen, Kruisheer, and Moesveld (*A.*, 1921, ii, 156). It is shown

that the values found by Holsboer (0.2225) (A., 1901, ii, 226) and by Seibert, Hulett, and Taylor (A., 1917, ii, 231) are untrustworthy.
J. F. S.

Determination of the Higher Point of Inversion of the Specific Heat of the Saturated Vapour of Benzene. G. BRUHAT and A. DELAYGUE (*Compt. rend.*, 1922, 174, 937—939).—Using a Cailletet apparatus with the tube jacketed with glycerol electrically heated, the authors have determined the two points of inversion of the specific heat of the saturated vapour of benzene. The values obtained are 122° and 258° respectively. From a theoretical study of the specific heat of the saturated vapour of benzene, using the characteristic equation for benzene, they obtain the values 121° and 254°.
W. G.

Specific Heats. II. M. PADOA (*Gazzetta*, 1922, 52, i, 25—29; cf. this vol., ii, 27).—The views expressed in the former paper lead to the conclusion that the low specific heats of elements deviating markedly from Dulong and Petit's law are due to the high number of valencies used by the atoms to ensure the stability of the crystalline edifice and to the exceptional energy of the corresponding linkings. Thus, if it were possible to separate the atoms of the abnormal element to such an extent that, although participating in the formation of the crystalline network, such atoms were no longer able to exert mutually disturbing actions, their specific heats should increase to the values exhibited by the other elements. Such a condition should be attained in solid solutions of an element of abnormal specific heat in excess of another of normal specific heat. Experimental confirmation of these views is furnished by measurements of the specific heats of silicon-iron and sulphur-selenium mixtures. In the former case, the atomic heat of silicon is 4.74—4.83 in alloys containing 95—50% of silicon, whilst the values for 33.6 (FeSi), 14 and 10% of silicon are, respectively, 5.27, 6.04, and 6.42. Heating of the 10% silicon mixture at above 1000°, followed by rapid cooling, fails to alter the specific heat. In sulphur-selenium mixtures containing 4, 9, 28.77, 90.35 (rhombic), and 90.35 (monoclinic)% of sulphur, the values of the atomic heat of sulphur are respectively 7.03, 6.00, 5.87, 5.50, and 5.78.
T. H. P.

Method of Formation of the Atoms in Carbon Compounds; HANS BEUTLER (*Z. anorg. Chem.*, 1921, 121, 67).—In a previous paper (this vol., ii, 259) it was shown that the lattice energy of carbon compounds alternated from member to member. This fact was deduced from the behaviour of the melting point and the heat of combustion of the homologues of aliphatic series. It is now pointed out that the recent work of Becker and Janke (this vol., ii, 128) experimentally confirms the author's hypothesis, since this work shows that the even members of the monocarboxylic aliphatic acids crystallise in the hexagonal and the odd members in the tetragonal system.
J. F. S.

Moment of Inertia and Structure of the Carbon Dioxide Molecule. A. EUCKEN (*Z. physikal. Chem.*, 1922, **100**, 159—170).—A theoretical paper in which the moment of inertia of the carbon dioxide molecule and its structure are considered. It is shown to be probable that in the molecule of carbon dioxide the carbon atom is situated practically on the line which joins the two oxygen atoms and performs small elastic circular vibrations round these atoms. The moment of inertia vertical to the line joining the oxygen atoms has the value 50×10^{-40} cm². gram, and the mutual distance of the oxygen atoms 2.0×10^{-8} cm. Carbon dioxide and presumably sulphur dioxide are to be treated thermally as diatomic gases; on this basis the chemical constant i for carbon dioxide has the value 0.80, and the molecular heat at high temperatures, in consequence of the linear vibration of the oxygen atoms and the circular vibrations of the carbon atom, approaches the limiting value $C_{\infty} = 5/2R + 4/2R + 4/2R = 13/2R$. J. F. S.

Some Relationships between the Critical Temperature and the Melting and Boiling Points. MAURICE PRUD'HOMME (*J. Chim. physique*, 1921, **19**, 188—189).—The following two relationships are found to hold for the critical temperature (T_c), the freezing point (T_f), and the boiling point (T_b) of members of the zero group: $T_b(T_c - T_f)/T_f(T_c - T_b) = 1 + \epsilon$, where ϵ is a fraction having a mean value 0.05, and $T_c = (T_f - T_b)(1 + \epsilon')$ where ϵ' has very nearly the same value as ϵ . These formulæ hold for other substances and have been confirmed in sixty different cases. J. F. S.

Factors Determining the Hygroscopic Properties of Soluble Substances. I. Vapour Pressures of Saturated Solutions. GRAHAM EDGAR and W. O. SWAN (*J. Amer. Chem. Soc.*, 1922, **44**, 570—577).—The vapour pressures of saturated aqueous solutions of potassium chloride, potassium nitrate, ammonium chloride, ammonium sulphate, ammonium nitrate, ammonium dihydrogen orthophosphate, sodium nitrate, carbamide, and a mixture of ammonium chloride and potassium nitrate have been measured at temperatures between 20° and 30°. The relationship of the vapour pressures of saturated aqueous solutions to the hygroscopic properties of soluble substances is discussed. The theory relating the vapour pressure of a saturated solution to the temperature and solubility is considered, and on the basis of the Clausius-Clapeyron equation and van't Hoff's equation the expression $\log P_1 - \log P_2 = i(S_2 - S_1) - Q_1/R(1/T_1 - 1/T_2)$ is deduced. This equation gives calculated vapour pressures which are in good agreement with those found. J. F. S.

The 25°-Isotherms of the Systems Magnesium Nitrate-Sodium Nitrate-Water and Magnesium Sulphate-Magnesium Nitrate-Water. DOUGLAS NORMAN JACKMAN and AGNES BROWNE (*T.*, 1922, **121**, 694—697).

Heat of Formation of Compounds of Metallic Magnesium with Metallic Zinc, Cadmium, Aluminium, and Calcium.

WILHELM BILTZ and GEORG HOHORST (*Z. anorg. Chem.*, 1921, 121, 1—21).—The heat of formation of the intermetallic compounds of magnesium with zinc, cadmium, aluminium, and calcium has been calculated from a series of determinations of the heat of solution of the individual metals and the compounds in hydrochloric acid of various concentrations. The metals were used in the form of fine powders or thin foil, and the solution was hastened by the use of small quantities (about 8 mg.) of a platinum catalyst. The following values of the molecular heat of solution in $\text{HCl}, 8\text{H}_2\text{O}$ at 18° are recorded: magnesium, 113.1 Cal.; zinc, 33.55 Cal.; aluminium, 125.1 Cal.; calcium, 130.1 Cal.; cadmium, 19.77 Cal.; MgZn_2 , 167.6 Cal.; MgCd , 123.7 Cal.; Mg_4Al_3 , 779 Cal.; and Mg_4Ca_3 , 800 Cal. The difference between the sum of the heats of solution of the constituent metals and that of the compounds gives the heat of formation of the compounds from the solid metals thus: $\text{Mg} + 2\text{Zn} = \text{MgZn}_2 + 12.6$ Cal.; $\text{Mg} + \text{Cd} = \text{MgCd} + 9.2$ Cal.; $4\text{Mg} + 3\text{Al} = \text{Mg}_4\text{Al}_3 + 49$ Cal.; and $4\text{Mg} + 3\text{Ca} = \text{Mg}_4\text{Ca}_3 + 43$ Cal. The densities of the compounds and that of calcium have been determined and the following values of d_4^{25} obtained, MgZn_2 , 5.126; MgCd , 5.383; Mg_4Al_3 , 2.151; Mg_4Ca_3 , 1.701; Ca, 1.542. The molecular volumes are calculated and compared with the sum of the atomic volumes, and it is shown that in all cases the compound is formed with a contraction in volume. J. F. S.

Heat of Reaction of Oxygen with Hæmoglobin. EDWARD F. ADOLPH and LAWRENCE J. HENDERSON (*J. Biol. Chem.*, 1922, 50, 463—490).—Estimations of the heats of reaction of oxygen and carbon monoxide with hæmoglobin at 22° gave average values of 6,950 and 14,700 cal., respectively. The individual values, however, showed large variations, the extreme values for oxygen being 1,500 and 10,900 cal. and those for carbon monoxide 8,300 and 25,200 cal., although the same apparatus gave consistent results when used for inorganic reactions. The authors conclude from their own results and from those of others that K and Q for oxyhæmoglobin are not related by the van't Hoff isochore, and discuss various factors which may account for this. E. S.

Adhesion Forces in Solutions. II. Adhesion Series. NIKOLAI SCHILOV, NAD. BULYGINA, NIK. ZEREVTINOV, LIDIE LEPIN, OLGA MICHAJLOVA, MICH. DUBININ, SERG. WOSNESSENSKI, AGNESS IVANITZKAJA, LIDIE ORLOVA, and JOH. SASLAVSKY (*Z. physikal. Chem.*, 1922, 100, 425—462; cf. A., 1920, ii, 591).—In an earlier paper (*loc. cit.*) it was shown that the adsorption of a series of salts of the same anion by charcoal increases with increasing valency of the kation; such a series is termed "convalent." A series of acids under the same treatment gave decreased adsorption with increasing valency of the anion; such a series is termed "antivalent." A series where the univalent ions come at one end, the bivalent at the other, and the trivalent between or irregu-

larly among the uni- and bi-valent ions is termed a mixed series. A number of adsorption experiments have been carried out on the adsorption of electrolytes by alumina, ferric oxide, manganese dioxide, peat, wool, and finely divided metals. The results show in the case of alumina that the adsorption series for acids and for copper salts are both convalent, with ferric oxide and acids the adsorption series is also convalent; manganese dioxide, acids, and copper salts all give convalent series. Peat gives an antivalent adsorption series with acids. In the case of finely divided metals, the adsorption of copper salts by tungsten and antimony is antivalent but by nickel salts convalent. The work with finely-divided metals was rendered difficult owing to chemical action occurring between the two phases; it is shown that an adsorption always precedes a chemical action. Thus iron powder was kept in contact with silver nitrate and it was found that for one hundred and fifty minutes silver was removed from solution to the extent of 3—8% without iron passing into solution, but once the chemical action commenced it proceeded very rapidly to completion. The adsorption of a series of chlorides by wool is convalent. Coagulation series have been investigated for the action of electrolytes on mastic sols, colloidal ferric hydroxide, arsenic sulphide, antimony sulphide, manganese dioxide, and for suspensions of charcoal in alkalis, and in tannic acid, and iron oxide in *N*/50-hydrochloric acid. The coagulation series of mastic sols is antivalent for acids and for copper salts, but for sulphates it is convalent, although the hydrogen-ion falls among the multivalent ions. The nitrates and chlorides both formed mixed series. With positively charged ferric hydroxide sols, the anion series is convalent, in the case of the acids, copper salts, and potassium salts, whilst the kations all formed mixed series. Arsenic sulphide and antimony sulphide both give an antivalent series for the anions and a convalent series for the kations, whilst manganese dioxide gives an antivalent series for anions and a mixed series for kations. The electrode potential of metals in acids has been divided into similar series, and from the measurements of Ostwald (*ibid.*, 1887, i, 583) it is shown that zinc, cadmium, tin, lead, iron, and antimony give convalent series, mercury and copper antivalent series, and silver and bismuth mixed series. From the work of other investigators it is shown that the influence of anions on the solubility, compressibility of solutions, and the abnormal elevation of the boiling point is convalent, whilst the influence of anions in acid solution on the rate of hydrolysis of esters is antivalent, but in alkaline solution convalent. A calculation of the molecular contraction in the formation of solid chlorides and sulphates gives rise in both cases to convalent series. The experimental results are discussed, and it is shown that the series is due to the processes being controlled by molecular forces rather than by ionic forces. Such molecular forces are termed "cohesion forces" where similar molecules are concerned, and "adhesion forces" where dissimilar molecules participate.

J. F. S.

Validity of the Laws of the Internal Friction of Liquids. W. HERZ (*Z. anorg. Chem.*, 1921, **121**, 95—98).—A theoretical paper in which the author examines the formulæ put forward by Bingham and Harrison (A., 1918, ii, 382) and Batschinski (A., 1913, ii, 928) to represent the fluidity of the benzene haloids. Extrapolating to the absolute zero leads to negative values for the fluidity, and from this the author concludes that the fluidity is not a strict linear function of either the temperature or the specific volume. J. F. S.

The Viscosities of some Hydrocarbons. G. CHAVANNE and (Mlle) H. VAN RISSEGHEM (*Bull. Soc. chim. Belg.*, 1922, **31**, 87—94; cf. Bartoli and Stracciati, A., 1886, 215; also Thorpe and Rodger, T., 1897, 360).—Viscosity measurements of some hydrocarbons lead to the following conclusions: Viscosity increases directly with the number of side chains, for example, 2:2:3:3-tetramethylbutane is a solid; high viscosity is thus conditioned by the presence of tertiary butyl groups, which also determine high melting and low boiling points. Cyclic compounds have greater viscosity than the corresponding aliphatic compounds, but in this case the introduction of a methyl group diminishes viscosity; this is also the case with polymethylenes, the viscosities of which are considerably greater than those of the corresponding aromatic substances. H. J. E.

Thermal Evolution of Gases Absorbed by Charcoals and Carbonised Lignites. STUART McLEAN (*Trans. Roy. Soc. Canada*, 1921, **15**, iii, 73—84).—An investigation of the heat developed when air, oxygen, nitrogen, or carbon dioxide is absorbed by wood or cocoanut charcoal, or lignite carbonised at various temperatures between 350° and 550°. Oxygen develops the greatest amount of heat per unit volume of gas adsorbed. During the adsorption, both carbon dioxide and carbon monoxide are formed. The greatest evolution of heat per unit mass of adsorbent occurs with carbon dioxide. The thermal effect for air is much smaller than for oxygen, and less carbon dioxide and carbon monoxide are produced. In the case of oxygen adsorption, the amount of gas adsorbed diminishes until a constant value is attained on repeating the experiment. The same holds for the heat developed per unit mass of adsorbent. The heat developed by the adsorption of carbon dioxide is not accounted for by the latent heat of evaporation. The coarse kinds of charcoal adsorb more oxygen and form carbon dioxide more readily than other kinds. J. S. G. T.

Influence of the Hydrogen-ion Concentration on the Permeability of Dead Membranes, on Adsorption by Protein Sols, and on the Metabolism of Cells and Tissues. ALBRECHT BETHE (*Biochem. Z.*, 1922, **127**, 18—33).—The diffusion of acid dyes through parchment is accelerated in acid solution and depressed in alkaline solution, whilst the reverse is true for basic dyes. If the dyes are allowed to diffuse into a protein solution, for instance, milk, gelatin, or serum, there is preferential distribution of the

dye in the protein solution if the acid dye is dissolved in acid solution and the basic dye in alkaline solution. The process can then be reversed by addition of alkali or acid. The analogy is extended to vital staining, where the hydrogen-ion concentration is assumed to play an essential rôle. H. K.

Substantive Cotton Dyeing. RUDOLF AUERBACH (*Kolloid Z.*, 1922, 30, 166—168).—The influence of various salts on the dyeing of cotton by substantive dyes was investigated. In each case an optimum concentration of the salt was found, and this was almost inversely proportional to its precipitating power on the colloid systems. The amount of dye taken up by the threads also decreases with increasing precipitating power of the salt. The strong mineral acids do not follow the same rule, no optimum concentration being found. W. T.

Pressure of Oxygen in Equilibrium with Silver Oxide. FREDERICK G. KEYES and H. HARA (*J. Amer. Chem. Soc.*, 1922, 44, 479—485).—The equilibrium pressure of oxygen over silver and silver oxide has been measured up to about 800°; the present results confirm and extend the previously published results of Lewis (*Z. physikal. Chem.*, 1906, 55, 449). The following equation, ordinarily derived by the use of the perfect gas law and by assuming the heat constant, is found empirically to represent the data exactly: $\log p = -2859/T + 6.2853$, where the pressure is in atmospheres, and the temperature in degrees absolute. The Nernst heat theorem as given by Pollitzer to fit the data of Lewis entirely fails to represent the present data. Neglecting only the difference in the coefficient of expansion between silver and silver oxide, the following equation is derived from the Clausius equation for the external heat of the reaction, in calories per molecule of oxygen, $Q = 6582(v - 20.7)p/T$, where v calculated from the equation of state of oxygen is given in c.c./mol. The value for the internal heat at 25°, calculated with the aid of Kopp's rule, is 12642 cal., in agreement with the mean of the measurements and computations (12640) considered by Lewis. The internal heats, although they pass through a minimum at about 400—500°, increase again with rising temperature. The analogy between this chemical reaction and a simple evaporation, therefore, breaks down, since no sign is shown of a critical point, with the heat equal to zero, although Δv appears to be diminishing progressively. J. F. S.

A Simple Dialyser. A. GUTBIER and A. MAYER (*Z. anorg. Chem.*, 1922, 121, 215—216).—A simple and efficient dialyser can be constructed as follows. A polished ring of wood, bone, or horn is grooved on the outer side, a piece of parchment paper is well soaked in distilled water and then folded over a test-tube or beaker (according to the size of dialyser required) to form a kind of bag, the glass vessel is then removed and the parchment bag tied around the ring. If necessary (for sensitive systems), in place of the ring a disk of the same material with a groove along its edge and a hole in the plate to fill and empty the bag can be used. W. T.

Temperature Dependence of the Solubility, the Activity, and Osmotic Coefficients of Salts. J. N. BRÖNSTED (*Z. physikal. Chem.*, 1922, 100, 139—150).—Formulæ are developed to represent the dependence of the activity and osmotic coefficients on the temperature. These have the form $\delta \log_e f / \delta T = \alpha c / RT^2$ and $\delta \phi / \delta T = \alpha c / 2RT^2$, in which f is the activity coefficient and ϕ the osmotic coefficient, c the concentration, and $\alpha = \delta U_1 / \delta c$. In the case of potassium chloride in 0.1*N*-potassium chloride the value $\delta \log_e f / \delta T$ is calculated to 0.000092 and $\delta \phi / \delta T = 0.000046$. Further formulæ are developed for the dependence on temperature of the solubility of sparingly soluble salts in pure water and in salt solutions. These formulæ have been verified by means of solubility determinations of β -cobalt croceonitrate in water, *N*/10-sodium nitrate, and potassium nitrate at various temperatures. J. F. S.

Physical Chemistry of Lipoids. Diffusion of Methylene-blue through Organic Solvents. S. LOEWE (*Biochem. Z.*, 1922, 127, 231—240; cf. A., 1912, ii, 741, 742).—The author has examined qualitatively the distribution and diffusion of methylene-blue from a gelatin gel in one limb of an inverted Y-shaped apparatus through a supernatant organic solvent (carbon tetrachloride, benzene, toluene, ether), containing in solution one of a great variety of substances belonging to the groups of plant resins, wool-fat and its products, and various lipoids, into a gelatin gel in the other limb. With a few exceptions, the amount of methylene-blue taken up from the gelatin gel by the organic solvent and its dissolved substance and by the pure gelatin gel is small. H. K.

Electrical Charges of Colloidal Particles and Anomalous Osmosis. JACQUES LOEB (*J. Gen. Physiol.*, 1922, 4, 463—486).—In a system consisting of water and salt solution separated by a collodion-gelatin membrane, the whole being on the acid side of the isoelectric point of the membrane, the transport curve of water to salt solution shows a rise up to a concentration of the salt of $M/32$, then a fall up to a concentration of $M/4$, and subsequently a continuous rise. The preliminary rise is the greater the greater the valency of the kation. On the alkaline side of the isoelectric point, curves of a similar nature are obtained, but in this case it is the valency of the anion which influences the preliminary rise.

The fluctuations in those curves ("anomalous osmosis") are explained as due to the effect of the salts on the potential difference across the membrane and on the potential difference between the solution in the pores of the membrane and the membrane itself.

Both these effects should disappear when the whole system is at the isoelectric point of the membrane; and it is, in fact, found, under these conditions, that solutions of salts composed of uni- and bi-valent ions behave like sucrose solutions. Salts containing ter- or quadri-valent kations or quadrivalent anions still behave anomalously, probably owing to the fact that the isoelectric gelatin forms loose compounds with the multivalent ions, thus setting up a potential difference between itself and the solution. C. R. H.

An Optical Method for the Determination of the Reciprocal Solubility of Slightly Miscible Liquids. C. CHÉNEVEAU (*Compt. rend.*, 1922, 174, 815—817).—The process employed consists in using the method of autocollimation, whilst making it differential. The refractive indices of the liquid considered as the solvent and of the solution are measured in a specially constructed prism divided into two cells. A formula is given by means of which, from the readings taken, it is possible to calculate the solubility of the one liquid in the other. The results obtained are in very close agreement with those obtained by gravimetric methods.
W. G.

Application of the Electron Theory of Chemistry to Solids. (SIR) J. J. THOMSON (*Phil. Mag.*, 1922, [vi], 43, 721—757; cf. A., 1921, ii, 252).—The part played by electrons in determining the structure of solids is considered. Each kind of atom is associated with a definite number of electrons with which it holds other atoms in chemical combination. The simplest case is that of a chemical element. In the crystalline state each unit cell will contain one atom and a definite number of electrons; one for a univalent atom, two for a bivalent atom, and so on. The arrangement of the electrons in the unit cell determines the shape of the cell and the crystalline structure. Possible unit cells are limited to (1) Parallelepiped; (2) Hexagonal prisms; (3) Rhombic Dodecahedra; (4) Cubo-octahedra.

The univalent element gives the simplest type of cell. The atom is situated at the centre of a parallelepipedon (or cube) with an electron at each corner and each electron held in common by the atoms in eight such parallelepipeda. For a bivalent element the electrons may be arranged at the corners of an hexagonal prism or in cubes with additional electrons at the centres of one set of parallel faces. The cells of tervalent elements may be hexagonal prisms with an electron at the centre of each end, cubes with electrons at the centres of two sets of parallel faces, or as rhombic dodecahedra. Similarly, the four electrons of quadrivalent elements may be arranged as in the face-centred cube or as in the hexagonal prism with two electrons along the axis at equal distances from the centre, and so on.

The symmetries represented by the triclinic, monoclinic, rhombic, and tetragonal systems in crystallography can be obtained if an assemblage of regular cells be strained homogeneously in any way without fracture. The cells would still maintain an atom in each cell and the original number of electrons.

Making the assumptions with regard to the forces between the atoms and electrons given in the previous paper, the stability of the above electronic arrangements and their effect on some physical properties is discussed. The values for the bulk modulus, the critical frequency, and the dielectric constant calculated for elements of different valencies, agree closely with those found by experiment. Expressions are given for the potential energy per unit volume of solid, the compressibility, the latent heat of fusion, etc., for a

number of types of unit cells. It is shown that the potential energy of an atom on the surface will be greater than that in the interior of the crystal and this excess will be proportional to the surface tension.

The constitution of salts may be inferred from their physical properties. From the magnitude of the compressibility of potassium chloride and sodium chloride it is deduced that the chlorine and sodium or potassium atoms are present as singly positively charged units with one electron per unit cell. The other six electrons are arranged as a regular octahedron about the chlorine atom at a distance considerably smaller than the length of the cube. The number of unit cells occupied by a chemical molecule and the presence of chemical groups (for example, CN or NH_4) in crystals can also be indicated by compressibility measurements. It is shown that Richard's value for the compressibility of the diamond can be explained on the assumption that the carbon atoms are arranged in groups in the crystal. The stability of chains of carbon atoms is discussed. A long chain of carbon atoms is unstable. A stable arrangement for chains of CH_2 is possible if this group be regarded as equivalent to an atom of a bivalent element with two disposable electrons.

W. E. G.

Effect of Electrolytes on the Constitution of Soap Solutions, as Deduced from Electromotive Force. CYRIL SEBASTIAN SALMON (T., 1922, 121, 711—715).

Hysteresis Phenomena and Coagulation of Sols with Ultramicros of Rod-like Structure, in particular Vanadium Pentoxide compared with Fibrin Sols. G. WIEGNER, J. MAGASANIK, and H. GESSNER (*Kolloid Z.*, 1922, 30, 145—165).—Vanadium pentoxide sols contain ultramicros of elongated rod-like structures similar to fibrin sols; in course of time these ultramicros grow, this being accelerated by an increase in concentration of the sol. The tendency to coagulate is much increased after this growth, and can be brought about by filter-paper, bores, carbon, silk threads, and silicic acid and gelatin gels. This gelatinisation is favoured or perhaps conditioned by vanadyl salts formed by the reduction of the vanadium pentoxide. The serum obtained from the gel coagulates more sol; this is accounted for by its high content of vanadyl salts. Salts (contained in filter-paper) favour the reduction to vanadyl salts and therefore accelerate the coagulation. Vanadic acid stabilises the sol. Dilution of sol and consequently decrease of concentration of the active vanadic acid ions brings about gelatinisation. These results show a close analogy to fibrin sols. Quantitative investigations are now being carried out.

W. T.

The Protective and Flocculating Action of Hydrophilic Colloids and Hydrophobic Sols. H. FREUNDLICH and E. LOENING (*Festschrift Kaiser Wilhelm Ges. Förderung Wiss. Zehnjährigen Jubiläum*, 1921, 82—89).—It was found by Brossa and Freundlich (A., 1915, ii, 241) that the flocculating value of electro-

lytes for ferric oxide sol decreased with increase in the percentage of albumin present. In addition Gann (A., 1917, ii, 21) found that egg-white had a coagulating action on acidic gold sol in the absence of electrolytes. An attempt is made to correlate these facts. By experiments with both gold and silver sols it was shown that the sensitisation of a hydrophobic sol by a hydrophilic colloid is merely a coagulation which is too slight to become immediately observable. It can be detected only by the fact that when the sol is thus sensitised, smaller amounts of electrolytes are required for flocculation. Measurements of the flocculation value of electrolytes showed that with increasing gelatin content the value diminished to a low figure. In this way the sensitisation of the ferric oxide sol by albumin is not essentially a different phenomenon from the coagulation of gold sol by egg-white. The transition from coagulating to protective action observed by Gann is a general phenomenon. It probably depends on the fact that the oppositely charged hydrophilic sol in low concentration coagulates the particles of hydrophobic sol, whereas in higher concentration the former actually envelops and thus protects them. The properties of the hydrophobic sol are thus completely masked. Silver sol protected by gelatin has the same charge as pure silver sol and is therefore not reversed in charge by addition of gelatin. The action in this case is considered due to the amphoteric nature of the protective hydrophilic colloid. CHEMICAL ABSTRACTS.

Capillary-electric Phenomena in Lyophile Sols. H. R. KRUYT and H. G. DE JONG (*Z. physikal. Chem.*, 1922, 100, 250—265).—The influence of the electrolytes potassium chloride, barium chloride, potassium sulphate, potassium thiocyanate, potassium ferri cyanide, sodium chloride, lithium chloride, strontium chloride, magnesium sulphate, cadmium sulphate, and lanthanum nitrate on the viscosity of agar sols of various concentrations has been measured at 50°. It is shown that above the temperature at which gelatinisation occurs (40°) agar sols follow the Poiseuille law exactly. Agar sols exhibit the quasi-viscous effect noted by Smoluchowski, and the particles are discharged by kations at concentrations corresponding with the valency of the added kation. The electrical behaviour of the agar sol is very similar to that of a suspensoid, and the electrical charge has a capillary-electric character. An emulsoid sol possesses two stabilising factors: the electric charge and the hydration, both of which must be removed before a coagulation can be effected. If the charge only is removed, a lyophile sol remains, but if the hydration is removed a suspensoid remains. Salting-out is a combined discharge and dehydration. Albuminous substances do not form ion-disperse solutions, but colloid disperse sols. The amino-acidic character exhibits itself only in the molecules which form the boundary layer of the particles and in those which form the double layer. J. F. S.

Convertibility of Chemical Energy and the Conception of Affinity. R. PLANK (*Z. physikal. Chem.*, 1922, 100, 372—392).—A mathematical paper in which the maximum work of reversible

reactions is considered and from the consideration it is shown that the conception of affinity at present adopted is unsatisfactory. Suggestions are made for changing the present conception.

J. F. S.

The Le Chatelier-Braun Principle. CARL BENEDICKS (*Z. physikal. Chem.*, 1922, **100**, 42—51).—A discussion of the Le Chatelier-Braun theorem as formulated by Chwolson (*Lehrb. d. Physik*, 1905, Braunschweig). A number of cases are brought forward, chiefly metallographic, for which the theorem does not hold, and it is shown that the theorem is not generally applicable. It is always true for a system in equilibrium, and is to be regarded as a general criterion of stable equilibria, for only when the theorem holds can there be a stable equilibrium. At extremely high temperatures and pressures, the theorem holds more often.

J. F. S.

An Explanation of Liesegang's Rings. S. C. BRADFORD (*Science*, 1921, **54**, 463—464; cf. *A.*, 1916, ii, 474; 1917, ii, 366; 1920, ii, 235, 593).—Objection is raised to the views of McGuigan (*ibid.*, 1921, **54**, 78), and particularly to the statement that the chromate of itself is unable to diffuse in the gelatin. Further, bands of lead chromate can be obtained in gelatin under suitably chosen conditions. It is maintained that the attractive force described by McGuigan is that of adsorption.

A. A. E.

Influence of Intensive Drying on Internal Change. A. SMITS (*Z. physikal. Chem.*, 1922, **100**, 477—478).—The author directs attention to the experiments of Baker (*T.*, 1912, **51**, 2339) on the intensive drying of nitrous anhydride and nitrogen tetroxide. He points out that the large change in boiling point brought about in these substances by excessive drying probably indicates that the internal change has been stopped and that fractional distillation of such an extremely dry unary system will probably result in the separation of the pseudo-components [cf. also Baker, *T.*, 1922, **121**, 568].

J. F. S.

Change of Properties of Substances on Drying. HERBERT BRERETON BAKER (*T.*, 1922, **121**, 568—574).

The Dushman Equation for the Velocity of a Unimolecular Reaction. W. E. GARNER (*Nature*, 1921, **108**, 211).—It is suggested that the term ν in Dushman's equation (*A.*, 1921, ii, 315) $\kappa = \nu e^{(-N h \nu)/(RT)}$ is a frequency characteristic of some degree of freedom in the decomposing molecule, and is not related to the frequencies of the activating radiation. The "period of existence" of the molecule (the time corresponding with one molecular vibration) will then be $1/\nu$. It is argued, with the support of Tolman's data (*A.*, 1921, ii, 248), that the exact agreement obtained by Dushman for observed and calculated values of κ in the case of phosphine is accidental.

A. A. E.

Ignition Point of Detonating Gas [Hydrogen and Oxygen]. ALEXANDER MITSCHERLICH (*Z. anorg. Chem.*, 1921, **121**, 53—66).—The ignition temperature of detonating gas has been determined for

a variety of conditions in porcelain and glass tubes. The conditions which were varied, included the rate of flow of the gas, the diameter of the tube, the pressure and the dilution of the gas. It is shown that the ignition temperature of detonating gas at 150 mm. is highly dependent on the rate of flow in the sense that it increases with decreasing velocity, but beyond a given velocity further increase has but a slight influence on the temperature of ignition. The following values illustrate this fact, $v=280$, $t=592^\circ$; $v=187$, $t=592^\circ$; $v=130$, $t=593^\circ$; $v=93$, $t=594^\circ$; $v=37$, $t=601^\circ$, where v is the number of c.c. passing a cross-section of 1 sq. cm. per minute. When the value of v is kept constant, it is shown that the ignition temperature increases with increasing pressure, thus, $p=150$ mm., $t=592^\circ$; $p=200$, $t=607.5^\circ$; $p=250$, $t=622^\circ$; $p=300$, $t=630.5^\circ$; $p=400$, $t=644^\circ$; $p=500$, $t=651.5^\circ$; $p=600$, $t=659.5^\circ$; $p=650$, $t=663.0^\circ$. The diameter of the porcelain or glass tube is without influence on the explosion temperature for the diameters 3.6, 5.5, 9.5, and 11 mm., but this is not true for narrow tubes; thus with tubes less than 0.5 mm. diameter no definite ignition temperature could be observed.

J. F. S.

Chemical Kinetics of Perchloric Acid and its Salts.

G. BREDIG and J. MICHEL (*Z. physikal. Chem.*, 1922, **100**, 124—138).—The velocity of reaction in the reduction of aqueous perchloric acid by tervalent titanium, tervalent molybdenum, and bivalent chromium salts in hydrochloric acid and sulphuric acid solutions has been measured for various concentrations at 40° . In the case of titanium, which is most completely examined, the velocity is proportional to the first power of the concentration of the tervalent titanium, the perchloric acid, or the perchlorate. The concentration of sulphuric acid or hydrochloric acid has only very slight influence on the velocity in dilute solutions of the reactants. The influence of the acids increases from a definite acid concentration (4.45*N*-sulphuric acid, 4.69*N*-hydrochloric acid) linearly with the increasing concentration of the acids. The form of the kinetic equation is of the first order and of much lower order than is demanded by the equation representing the reaction, namely, $4\text{Ti}(\text{SO}_4)_3 + 4\text{H}_2\text{SO}_4 + \text{HClO}_4 = 8\text{Ti}(\text{SO}_4)_2 + 4\text{H}_2\text{O} + \text{HCl}$. As in most similar cases, an intermediate reaction of a lower order must be assumed, the nature of which has not been ascertained. The reaction velocity of the titanium reaction increases three times for an increase in temperature of 10° . In the present experiments, the times required for an equal amount of reduction of perchloric acid in the presence of the same excess of sulphuric and hydrochloric acids at the same temperature are in the ratio: $\text{Ti}^{\text{III}} : \text{Mo}^{\text{III}}$ (olive-green) : $\text{Cr}^{\text{II}} : \text{Mo}^{\text{III}}$ (orange) = 1 : 18 : 1300 : 40000. J. F. S.

Kinetic Study of Alkaline Solutions of Iodine. O. LIÉVIN (*Compt. rend.*, 1922, **174**, 868—870).—In alkaline medium iodine is converted into iodate by processes which differ according to the degree of alkalinity. The alkali, the concentration, and the iodide formed all influence the reaction. As the amount of alkali hydroxide used for a given weight of iodine increases the free iodine

disappears more rapidly but the hypiodite disappears more slowly. The presence of the iodide accelerates the reaction. Thus the iodide is a product of the reaction which it catalyses and the sodium hydroxide retards a reaction in which it is used up. Dilution causes a diminution in the velocity of the reaction. If sodium carbonate or the tribasic phosphate are used as the alkali then the reaction is accelerated by the addition of alkali, in so far as the formation of iodate is concerned. Further with sodium carbonate as the alkali, dilution of the system causes an increase in the velocity of reaction.

W. G.

Velocity of Decomposition of Silver Permanganate. A. SIEVERTS and H. THEBERATH (*Z. physikal. Chem.*, 1922, **100**, 463—475).—The thermal decomposition of silver permanganate takes place according to the equation $\text{AgMnO}_4 = \text{AgMnO}_3 + \text{O}$, and in the presence of water the residue combines with half a molecule of water. The process of the decomposition at 30° may be approximately expressed by the equation for an autocatalytically accelerated reaction, $dx/dt = Kx(a-x)$, where a is the initial amount of silver permanganate and $(a-x)$ the amount of silver permanganate undecomposed at time t . The decomposition product, the nature of which has not been established, accelerates the reaction proportionally to its mass. The constants of the decomposition velocity of the various preparations are different. The presence of water has a very strong influence on this reaction. Dry silver permanganate decomposes very slowly even at 50° , but small traces of water increase the decomposition velocity very much, whilst larger additions of water have little further influence. The addition of powdered pumice and the decomposition product of the reaction changes only the initial velocity. Temperature has a very marked influence on the reaction. At 0° , even in the presence of water and the decomposition product, the decomposition does not take place. At 50° , the decomposition process is no longer represented by the same equation as at 30° ; the tx curves for 50° are characterised by a long, almost straight line. Comparison with the experiments at 30° shows that an increase of 20° in the temperature of reaction causes the initial velocity to increase from 12 to 20 times, the mean velocity between 19% and 38% decomposition, 6.1—7.6 times, and between 38% and 57% decomposition, 3.4—4.3 times.

J. F. S.

Separation of Halogens from Organic Compounds. H. VON EULER and A. FAHLANDER (*Z. physikal. Chem.*, 1922, **100**, 171—181).—Determinations have been made of the velocity and extent of the separation of halogens from the sodium salts of chloroacetic, bromoacetic, and bromopropionic acids in water to which finely-divided silver, silver oxide, copper oxide, lead oxide, and platinum sponge have been added. The results show that the hydrolysis of a 0.1N-solution of sodium chloroacetate is not noticeably accelerated either at 75° or 100° by the presence of finely-divided silver of 1μ diameter in an hour. At 100° , the hydrolysis of the chloroacetate solution amounted to 8.8% in ninety minutes, but in the

presence of 5 grams of copper oxide in 25 c.c. of the 0.1*N*-solution it amounted to 9.6% in the same time. At 75°, in five hours 2.6% of the same solution was hydrolysed, but with 1 gram of finely-divided silver there was no increase in the amount decomposed, with 2 grams of silver oxide the amount was 7.2%, and with 4 grams of copper oxide, 3.8%, all in 25 c.c. of *N*/10-solution. It is shown that the accelerating action cannot be attributed to the amount of the oxides dissolved. In the case of the most concentrated bromoacetate solution employed, the velocity increases proportionally with the amount of silver oxide added, but more slowly with more dilute solutions. The velocity of bromine separation is roughly one hundredth of that occasioned by the presence of the same amount of silver oxide (cf. Donnan, T., 1904, **85**, 555; 1910, **97**, 1882). J. F. S.

Reduction of Nitro-compounds by Stannous Chloride. III. HEINRICH GOLDSCHMIDT, EINAR STORM, and ODD HASSEL (*Z. physikal. Chem.*, 1922, **100**, 197—207; cf. A., 1904, ii, 607; 1906, i, 734).—A number of experiments are described on the reduction of *m*-nitroaniline, *m*-nitrobenzenesulphonic acid, *o*-nitrobenzenesulphonic acid, and *o*-nitroaniline by stannous chloride and bromide in the presence of hydrochloric, hydrobromic, sulphuric, and benzenesulphonic acids and lithium, ammonium, potassium, calcium, and cadmium chlorides. It is shown that in the reduction by stannous chloride and hydrochloric acid a portion of the acid may be replaced by a metallic chloride without any reduction of the velocity of reaction ensuing. Cadmium chloride behaves differently from the other metallic chlorides, and reduces the velocity constant from 10.35 to 5.14. In the reduction with stannous bromide and hydrobromic acid, a portion of the hydrobromic acid may be replaced by metallic bromides without reduction of the velocity of reaction, but cadmium bromide brings a marked reduction in the velocity; here the constant is reduced from 33.37 to 8.47, an action which is attributed to the formation of complex salts. The addition of strong acids such as sulphuric and benzenesulphonic acids has but little action on the velocity of reaction. Reduction with stannous chloride and sulphuric acid in the absence of hydrochloric acid proceeds very slowly. Thus, 0.1*N*-stannous chloride and *N*-hydrochloric acid has a velocity constant 9.96 with 0.033*N*-*m*-nitroaniline, but if the hydrochloric acid is entirely replaced by *N*-sulphuric acid, the value falls to 0.44. J. F. S.

The Influence of Temperature and Constitution on the Decomposition Velocity of Substituted Malonic Acid. HEINRICH JAKUBOWICZ (*Z. anorg. Chem.*, 1922, **121**, 113—127).—The temperature-velocity curves are given for malonic acid and some of its derivatives up to 110°. For malonic acid, the decomposition becomes measurable at 66°, and the temperature coefficient was found to be 2—3 for 10° increase. The temperature-velocity curves for diethyl- and diallyl-malonic acids run parallel with those of malonic acid. The introduction of a hydroxyl group slightly increases the velocity of decomposition. The monochloro-acid

decomposes much more rapidly and occupies a position between the allyl and phenyl derivatives. The introduction of another chlorine atom reduces the velocity considerably; that is, the more symmetrical compound is the more stable. The dichloro-derivative decomposes more rapidly than its dibromo-analogue. The values of K at 100° are phenylmalonic acid 0.1970, chloromalonic acid 0.1855, allylmalonic acid 0.1321, hydroxymalonic acid 0.0820, malonic acid 0.0755, methylmalonic acid 0.0598, dichloromalonic acid 0.0585, dibromomalonic acid 0.0224.

W. T.

Slow Hydrolysis of Salts. A. TIAN (*J. Chim. physique*, 1921, **19**, 190—216; cf. A., 1921, ii, 439).—The author discusses the hypothesis put forward by Wagner (A., 1913, ii, 200, 765) to explain the slow hydrolysis of salts and finds it at variance with facts, and puts forward a new hypothesis, which has the following form. The slow hydrolysis of salts is always preceded by a normal hydrolysis, limited by the reverse reaction which always occurs between ions and is extremely rapid. As soon as the solubility product of one of the substances (acid or base) produced by the hydrolysis has been reached, a new phase appears and at the same time a new cause of retrogradation of the hydrolysis. This supplementary retrogradation is due to a reaction between the two phases (colloid and aqueous solution) present; it is realised with a velocity which increases with the surface of contact. This cause of retrogradation is very great at first, when the colloidal particles are very small, but as a consequence of their union the surface of the particles decreases and with it the speed of the supplementary retrogradation, and from this point the hydrolysis increases. The progression of the hydrolysis is therefore as slow as the evolution of the colloid. This hypothesis has been tested by means of conductivity measurements on solutions of ferric chloride in the presence of silica gels and gelose, and found to be in keeping with the experimental data.

J. F. S.

Catalysis. XIII. Temperature Coefficient of Catalysed and Non-catalysed Reactions. R. M. PURKAYOSTHA and N. R. DHAR (*Z. anorg. Chem.*, 1922, **121**, 156—166).—The reaction $\text{H}_2\text{C}_2\text{O}_4 + \text{KMnO}_4 + \text{MnSO}_4 + \text{H}_2\text{SO}_4$ has a temperature coefficient 2.96 between 10° and 20° , almost the same value as that found by Dhar for a similar reaction with chromic acid as the oxidising agent (T., 1917, **111**, 727). The reaction with permanganate is unimolecular with respect to the oxidising agent; in the case of chromic acid reaction the rate was found to be independent of the concentration of the oxidising agent. The reaction between mercuric chloride and phosphorous acid was found to be bimolecular for phosphorous acid and semimolecular for mercuric chloride. The temperature coefficient is 3.4 at 30 – 40° and 3.26 at 40 – 50° . The photochemical reactions $\text{Na}_2\text{SO}_3 + \text{O} = \text{Na}_2\text{SO}_4$ and $\text{NaClO} = \text{NaCl} + \text{O}$ are discussed and are regarded as semimolecular reactions. All the reactions with a high temperature coefficient were found to be sensitive to light rays.

W. T.

Peculiar Catalytic Action in Chemical Synthesis by Glow Discharge. FRITZ PANETH (*Z. physikal. Chem.*, 1922, 100, 367—371).—When a glow discharge is allowed to pass through a tube containing tin electrodes and a low pressure of hydrogen, there is no combination between the hydrogen and the tin, but if the hydrogen contains a small amount of methane, tin tetrahydride is the product. The action is explained by the assumption that the excited hydrogen atom contains a higher quantic electron, and when this comes in contact with an atom of tin, vapour combination occurs. The present action is discussed in connexion with the formation of active nitrogen and triatomic hydrogen.

J. F. S.

Reactions on Metallic Surfaces. OTTO RUFF (*Z. physikal. Chem.*, 1922, 100, 419—424).—It is shown that metallic calcium when alloyed with more electro-positive metals such as strontium, barium, sodium, or potassium, or with calcium nitride, combines more readily with nitrogen than pure calcium. From this observation it is concluded that the heteropolar nature of a metallic surface favours the combination with a homopolar gas, and from this it is assumed that it acquires influence charges from the surface valencies of the metal which determine its position and behaviour towards the surface.

J. F. S.

The Structure of the Atom and the Physical and Chemical Properties of the Elements. NIELS BOHR (*Z. Physik*, 1922, 9, 1—67).—An address delivered before the Physical and Chemical Societies of Copenhagen. A résumé is given of the recent work on the series spectra of the elements (cf. A., 1921, ii, 137) and their dependence on the orbit of the electrons within the atom. On the basis of his previous theories, the author develops an interpretation of the periodic table of the elements. In order to arrive at the relative stability of the electronic configurations in the atoms of the elements, the process of formation of an atom from the positive nucleus is followed step by step. Thus an element with an atomic number N is built up by the attachment of N successive electrons, each electron giving rise to a spectrum. The actual increase in charge of the nucleus, which occurs throughout the process, is taken into account. For the first two elements, hydrogen and helium, the knowledge of the spectra produced during the combining process is fairly exhaustive. For the remaining elements, the arc and spark spectra may be represented with good approximation by formulæ of the Rydberg type, $\nu = K/(n'' + \alpha_k)^2 - K/(n' + \alpha_k)^2$.

The orbit of the first electron is circular and the final stationary condition is attained when n and $k=1$. The combination of the second electron, as the spectrum of helium shows, takes place in two different ways. In the normal condition, the second electron follows the same type of path as the first, but in a different plane and in the metastable condition it moves in a 2_1 orbit ($n=2$ and $k=1$). The third electron, as in the lithium atom, is more loosely bound than the first two electrons and describes an eccentric orbit

2₁. For a small part of its path it approaches the nucleus to within a distance comparable with that of the innermost electrons. The fourth, fifth, and sixth electrons follow similar orbits. In the carbon atom, the configuration of the paths of the four loosely bound electrons possesses a tetrahedral symmetry. The seventh electron, as in the nitrogen atom, and the eighth, ninth, and tenth electrons follow circular paths of the type 2₂, lying within the region of the type 2₁ and being tightly bound. For neon, there will be two electrons with a path 1₁, four electrons with elliptical paths 2₁, and four electrons with a circular path 2₂. The paths of the electrons of this atom will possess a high degree of symmetry. In a similar manner, the stability of the electronic configurations of the elements with higher atomic numbers is considered. Particular attention is directed to sodium, where it is shown that the first terms of the *P* and *S* series spectra correspond with electronic orbits 3₁ and 3₂, so that the last bound electron is less tightly held than the last electron in the lithium atom. With the beginning of the fourth period orbits of the type 4₁ appear, but these are comparatively unstable, the outer electron of potassium being very loosely held—being only twice as fast as the corresponding 3₃ orbit. In the case of scandium and the elements with higher atomic numbers, the 3₃ orbits become more stable, and unsymmetrical electronic arrangements are produced which are accompanied by the appearance of colour, of paramagnetism, variable valency, and an increase in the complexity of the spectra. Throughout the first long period these irregularities occur, but when there are six each of electrons in the 3₁, 3₂, and 3₃ orbits the electronic configuration again becomes symmetrical. Thus the spectra of copper becomes less complex than that of iron, and at the end of the period is found the element krypton with a very symmetrical electronic arrangement. Colour is explained as due to the transference of an electron from one orbit to another, and occurs only with an unsymmetrical arrangement of the inner electrons. In the last period of the table electrons with orbits of the type 7₁ will be present. The Röntgen spectra also point to the same stability relationships as were found with the series spectra.

W. E. G.

Ionisation Potential and the Size of the Atom. A. S. EVE (*Trans. Roy. Soc. Canada*, 1921, 15, iii, 37—40).—The author tabulates the values of the respective ionisation potentials, the atomic diameters as determined by Bragg (A., 1920, ii, 537), and the cube roots of the atomic volumes, so far as these constants are all known for the elements of groups 1—7 and for the inert gases. In the first group, the products of the ionisation potential by either the atomic diameter or the cube root of the atomic volume are approximately constant and fairly concordant, the respective mean values being 18.3 and 15.5. It is concluded that the work done in removing an exterior electron is inversely proportional to the atomic radius. The same holds for the elements Mg, Ca, Sr, and Ba of group II, whilst the values for Zn, Cd, and Hg differ


considerably from the mean values for these four. This difference is possibly related to the fact that whilst the former have only a single ring of external electrons, the latter, according to Urbach, have a double ring of electrons in the outer zone. In the case of neon and argon, the diameters as deduced by Bragg afford values of the products in far better agreement with theory than those found by using the cube roots of the respective atomic volumes.

J. S. G. T.

The Atomic Weight of isoHelium. MAXIMILIAN CAMILLO NEUBURGER (*Physikal. Z.*, 1922, **23**, 145—146).—A theoretical paper in which the author applies his theory of the nucleus (this vol., ii, 208) to calculate the atomic weight of Rutherford's *iso*-helium. The assumption is made that oxygen is composed of one α -particle, four *isohelium* nuclei, and two electrons, nitrogen of one α -particle, three *isohelium* nuclei, one positively charged hydrogen atom, and two β -particles, and carbon of four *isohelium* nuclei and two β -particles. The maximum change in mass which can occur during the collision of an α -particle with atoms of oxygen, nitrogen, or carbon is taken as 0.009 unit, and from the known atomic weights of oxygen, nitrogen, carbon, helium, and hydrogen it is found that the atomic weight of *isohelium* is 3.00113 ± 0.006 .

W. E. G.

The Genesis of the Elements. MAXIMILIAN CAMILLO NEUBURGER (*Physikal. Z.*, 1922, **23**, 133—136).—A theoretical paper which deals with the structure of the atomic nucleus. The author proposes a modification of the Meitner nuclear model which embraces all the elements and not merely the radio-elements. Taking into account the recent work of Rutherford and Aston, he is able to give definite nuclear formulæ to the lighter elements, for example, nitrogen = $\alpha + 3\alpha_1 + H^+ + 2\beta$, oxygen = $\alpha + 4\alpha_1 + 2\beta$ (α = Helium nucleus, α_1 = *isohelium* nucleus, H^+ = hydrogen nucleus and β = nuclear electrons). By making certain assumptions as to the existence of isotopes, the genetic relationship of the lighter elements is given, for example,




Suggestions for the Construction of a Periodic Table on a Wider Basis. D. BALAREFF (*Z. anorg. Chem.*, 1921, **121**, 22—24).—A method of arrangement of the elements, which takes account of the various valency states of the elements, is described. The method consists in taking nine equal rectangular sheets of glass; these are placed parallel to one another; on the first sheet all elements of zero valency are marked in the positions they would occupy in the Mendeléeve system, and on the second sheet all elements which exhibit univalency are marked, and so on. Hence each sheet of glass has those elements marked on it which exhibit a given valency. On bringing the sheets close together, the Mendeléeve system is obtained, but at the same time relationships depending on the valency are brought out and emphasised.

J. F. S.

Twenty-eighth Annual Report of the Committee on Atomic Weights. Determinations Published during 1921. GREGORY PAUL BAXTER (*J. Amer. Chem. Soc.*, 1922, **44**, 427—437).—The report of the Atomic Weight Committee is discussed, and much of the numerical data concerned with recent determinations of the atomic weight of oxygen, fluorine, chlorine, nickel, zinc, germanium, cadmium, antimony, lanthanum, and bismuth is reproduced. J. F. S.

An Electrical Doublet Theory of the Nature of the Molecular Forces of Chemical and Physical Interaction. R. D. KLEEMAN (*Physical Rev.*, 1921, **18**, 303—312; cf. A., 1910, ii, 492).—It has already been shown that the attraction between two atoms is proportional to the product of the square roots of their atomic weights; the suggestion is now made that a neutral atom with a positive nucleus surrounded by electrons is an electrical doublet. Since the force between two such doublets would on the average be an attraction proportional to the product of the moments of the two doublets, it follows that the electrical moment of an atom is proportional to the square root of its atomic weight. A. A. E.

Nature of the Ionogen Linking. HUGO KAUFFMANN (*Z. physikal. Chem.*, 1922, **100**, 238—249).—It is shown that the introduction of methoxyl groups into the ortho- and para-position of triphenyl carbinol, fuchsines, and similar substances increases the basic character of these substances, whilst in the meta-position the basic character is unchanged. It is shown that the basicity is not centralised in one carbon atom, but distributed over all the ortho- and para-positions and that every methoxyl group introduced in those positions has an active share of the total basicity of the compound. In salts with a radicle as kation, the basic character is not centralised, but is spread over the whole kation, and consequently the positive charge of the kation is not situated on a single atom but is spread over the whole ion. The valency of an anion in union with such an ion is therefore a divided valency which acts in parts between the anion and the positions on the kation where the positive charge is located. Hence the ionogen linking is a divided linking. In the case of acetic acid it is represented as $\text{CH}_3 \cdot \text{C}^{\text{O}} \text{O} \cdot \text{H}$ and in trichloroacetic acid as in



the annexed formula. The author then discusses homo- and hetero-polar linkings, and shows that these are only limiting cases and that most linkings are intermediate between these. From the discussion, the following results are drawn. The valency has its seat in the atomic nucleus. Valency is measured by the number of lines of electric force which lead from the nucleus of an atom to the connecting electrons. To bring this statement into line with the usual chemical valency numbers, it is necessary to divide the above-mentioned number by $4\pi e$, where e is the absolute value of the charge of an electron. A result of this statement is that all valencies

are positive; negative valencies are impossible and that which has hitherto been termed negative valency is the other end of the line of force from the electron. The characteristics of the homo- and hetero-polar linkings are illustrated by means of the electric moment. If M_A represents the electric moment of the valency of an atom A , M_B that of an atom B , and M_{AB} that of the linking $A-B$, then $M_{AB} = M_A - M_B$. If the atom nucleus of A is similar to that of B , then M_{AB} is zero. It follows that the electric moment of a pure homopolar valency is always zero. When M_{AB} is equal to M_A , the linking is heteropolar and the electric moment is the greater the more pronouncedly heteropolar is the linking. J. F. S.

Double and Triple Bonds, and Electron Structures in Unsaturated Molecules. E. D. EASTMAN (*J. Amer. Chem. Soc.*, 1922, 44, 438—451).—A theoretical paper dealing with electron structure of unsaturated molecules of compounds of carbon, nitrogen, and boron. It is suggested that multiple linking involves the transfer of two electrons from the inner to the outer shell in one or both of two atomic nuclei, the group of eight being preserved about each, with two jointly held electrons constituting the bond. This hypothesis is sufficient to explain the restricted region of the periodic system in which elements characterised by multiple bond formation occur. To meet requirements imposed by the facts relating to stability, reactivity, and free rotation, the electrons of unsaturated octets are assumed to be held in equilibrium positions at greater distances from the nucleus than in the ordinary case, the transfer from the extended position to the usual one liberating energy. No attractive force between electrons is assumed, and the electron arrangement is taken as cubic, although subject to distortion. On this basis, a mechanism is provided for addition and condensation reactions. Many interesting points of view are afforded in the study of conjugation, the structure and substitution reactions of benzene, and tautomeric displacement of double bonds. Formulæ are illustrated and discussed which represent, more satisfactorily in several cases than other current hypotheses, many of the properties of chain hydrocarbons, the oxides of carbon, the carbonyl and carboxyl groups, nitrogen, oxygen, and the hydrides and oxygen-containing ions of boron.

J. F. S.

Mesohydry. II. GIUSEPPE ODDO (*Gazzetta*, 1922, 52, i, 42—56).—A claim for priority (cf. A., 1907, ii, 15) principally over Kauffmann (*Ahrens Sammlung*, 1907, 11, 1—102; 1918, 12, 1—112), Gebhard (A., 1912, ii, 242), and Hantzsch (A., 1908, ii, 462; A., 1910, i, 200).

T. H. P.

Mesohydry. III. Relation between the Affinity of Acids and the Atomic Ratio O : H in their Functional Groups, and the Constitutional Formulæ of the Acids. GIUSEPPE ODDO (*Gazzetta*, 1922, 52, i, 56—79; cf. A., 1907, ii, 15, and preceding abstract).—Starting with the conception that tetragonal meso-

hydric nuclei, $M \begin{array}{c} \diagup O \\ \diagdown O \end{array} H$, are not capable of dissociation and that

they become so in solution only in so far as they undergo hydrolytic transformation into the group, $O:M-OH$, the author shows that an intimate relation exists, for all acids, between the number of oxygen and hydrogen atoms existing in the functional acid group and the force of affinity.

With the monobasic acids, there are the following five different types, that in which $nO:nH$ has the value 2:1 required by the non-ionisable mesohydric nucleus, being taken as a basis: (1) $nO:nH=4:1$, acids even more energetic than HCl , such as $HClO_4$; (2) $nO:nH=3:1$, acids as energetic as HCl , such as $HClO_3$; (3) $nO:nH=2:1$, very weak acids, such as $R\cdot CO_2H$; (4) $nO:nH=2:2$ (or more), acids somewhat more energetic than the 2:1-acids, such as H_2CO_2 ; (5) $nO:nH=1:1$ (or more), very feeble acids, such as $HClO$.

With the dibasic acids, if the oxygen and hydrogen atoms constituting the dibasicity are distributed between two oxidisable atoms, the relations for each of the acid functional groups thus resulting are the same as for the monobasic acids, except for the mutual influence of the two groups, this being marked in the 1:2-position, slight in the 1:3-position, and almost zero in the 1:4-position. Thus, succinic acid is almost equivalent to acetic acid in this respect, and dithionic acid to nitric acid or to $R\cdot SO_3H$. If, however, the oxygen and hydrogen atoms causing the dibasicity are united to the same oxidisable atom, only two types result: (1) $nO:nH=4:2$, acids of the average energy of monobasic acids of types (1) and (2), for instance, sulphuric acid; (2) $nO:nH=3:2$, very weak acids, such as H_2SO_3 and (3) $nO:nH=2:2$, still weaker acids, such as H_2SO_2 . With tri- and poly-basic acids, the relations are the same as with dibasic acids. T. H. P.

Space Formulæ. FRIEDRICH RINNE (*Z. physikal. Chem.*, 1922, **100**, 408—418).—A theoretical paper in which the spacial configuration of crystalline substances and the atomic diameter are considered on the bases of the Barlow and Pope hypothesis of close packing, the molecular volume, and the X-ray measurements of crystal structure. J. F. S.

Still-head for the Prevention of Spray. FRIEDRICH L. HAHN (*Z. anal. Chem.*, 1922, **61**, 52—53).—The lower end of the delivery tube, below the cork closing the distillation flask, is blown into the form of an elongated bulb of slightly smaller diameter than that of the neck of the flask. An internal tube extends from an opening at the shoulder of the bulb to nearly the bottom of the latter, and the bulb itself is filled with pieces of broken glass. The vapours from the flask pass through the tube into the bulb, and then escape through the delivery tube to the condenser. Since

the bulb is contained in the neck of the flask, reflux action is but slight and distillation is rapid. W. P. S.

A Sodium Press for the Laboratory. ERNST BECKMANN (*Ber.*, 1922, 55, [B], 766—768).—The instrument is figured and fully described in the original. The essential features are the construction of the press cap in two separable portions whereby cleaning is greatly facilitated, the arrangement of the piston in such a manner that it has only a vertical and not a rotatory motion, the provision of a powerful wheel in place of the handle usual in copying presses, and a calibration device whereby the weight of the sodium wire can be ascertained from the readings of the rotations of the wheel. H. W.

Apparatus for Hydrogenation at Ordinary Pressures. J. KLIMONT (*Chem. Zeit.*, 1922, 46, 275).—An ordinary distillation flask of about 150 c.c. capacity has a glass tube sealed horizontally into the neck and bent at right angles inside the flask so that it passes vertically downwards, and ends in a small depression blown in the bottom of the flask, which is also provided with a rubber stopper carrying a short glass tube bent at right angles, which may be connected with a manometer, and a thermometer dipping into the liquid in the flask. The flask is filled by means of a long-stemmed funnel with a mixture of the substance to be hydrogenated and the catalyst, which is heated by means of a Bunsen burner and kept in constant agitation by a stream of hydrogen, which enters through the tube dipping into the liquid and leaves through the other side tube. H. C. R.

A Reversible Turbidity Phenomenon. RAPHAEL ED. LIESE-GANG (*Kolloid Z.*, 1922, 30, 165—166).—The double salts mercuric silver iodide and copper silver iodide undergo a colour change at 70° and 45°, respectively, and have found application as colour thermoscopes. A similar change at a lower temperature is produced by a mixture of 18 c.c. of 10% gelatin solution, 1 c.c. of 20% potassium ferricyanide, and 1 c.c. of nitric acid (*d* 1.148). This is completely clear, but becomes opaque on solidification and again clear on liquefaction. After hydrolysis of the gelatin, the turbidity is fainter. Nitric acid can be replaced by hydrochloric, citric, and other acids. W. T.

Inorganic Chemistry.

Triatomic Hydrogen. II. GERALD L. WENDT and ROBERT S. LANDAUER (*J. Amer. Chem. Soc.*, 1922, 44, 510—521; cf. A., 1920, ii, 425).—Triatomic hydrogen has now been prepared by three new methods, the Siemens glass tube ozoniser, the high frequency Tesla discharge, and the thermionic emission methods. Attempts to show its presence in hydrogen produced as nascent hydrogen from

solutions of hydrogen-ions were unsuccessful. This hydrogen is catalytically decomposed by finely divided platinum, nickel, copper, lead, antimony, and cadmium. It is unaffected by contact with silver, mercury, tin, bismuth, molybdenum, zinc, or aluminium. Contraction in volume occurs when this hydrogen is produced by the electrical discharge at 3 mm. pressure. This is particularly marked at the temperature of liquid ammonia and solid carbon dioxide. Triatomic hydrogen is condensed to the liquid form by exposure to the temperature of liquid oxygen. The spectrum of hydrogen at the temperature of liquid oxygen shows a progressive intensification of the secondary line spectrum at the expense of the primary series spectrum, which is probably due to the gradual formation of the triatomic form. The present work is held to confirm the mechanism of the formation previously advanced, and also it shows that the new gas is probably H_3 and not *iso*- H_2 , an atomic species proposed by Harkins as a constituent of heavier atoms (cf. Harkins, A., 1920, ii, 479, 745). J. F. S.

Constitution of Haloids. WILHELM BILTZ (*Z. physikal. Chem.*, 1922, **100**, 52—87).—A theoretical paper in which the author discusses the constitution of the haloids from considerations of the melting point, temperature of decomposition, and electrolytic conductivity of the molten compounds. In the case of the platinum, iridium, and vanadium compounds, it is shown that the heat of formation of the highest chloride is much less than that of the remaining lower chlorides; he therefore considers that the higher chlorides are to be considered as co-ordination complexes of the same type as the amines. In the case of the melting points of the chlorides, it is shown that the lower chlorides have much lower values than the higher chlorides, and in the same way the lower chlorides have a much smaller conductivity in the molten condition than the higher chlorides. All haloids melting between 960° and 450° are good conductors, those melting between 450° and 200° are partly good conductors, partly poor conductors, and some do not conduct at all, and those melting below 200° , with the exception of the gallium compounds, do not conduct at all. The lower chlorides are therefore regarded as true salts, whilst the higher compounds are regarded as complexes. A comparison of the electro-valency towards chloride with the atomic number indicates periodicity as shown by a curve made by plotting the two values. J. F. S.

Significance of the Density of Hydrogen Bromide with Reference to the Atomic Weight of Bromine. GREGORY PAUL BAXTER (*J. Amer. Chem. Soc.*, 1922, **44**, 595—601; cf. Guye, A., 1919, ii, 333).—A discussion on the determination of atomic weight from gas densities, particularly in the case of the atomic weight of bromine from the gas density of hydrogen bromide. The pitfalls of such determinations are carefully described. It is pointed out that no method of extrapolation can be satisfactory or convincing where, as in the case of hydrogen bromide, the accuracy of the numerical data diminishes as the region to be

extrapolated approaches. The conclusion cannot be avoided that whilst in the case of hydrogen bromide the gas density method of determining molecular weights may be regarded as an interesting and satisfactory corroboration of gravimetric evidence, its value ceases at that point. To use such as a criterion on which to judge the best gravimetric evidence cannot be justified on scientific grounds. On the other hand, from the point of view of the general application of Avogadro's hypothesis to gases at low pressures, the results of the very careful density determinations, published from Guye's laboratory, are particularly significant and convincing.

J. F. S.

Adsorption of Iodine by Silver Iodide. FRANK E. E. GERMANN and RALPH N. TRAXLER (*J. Amer. Chem. Soc.*, 1922, **44**, 460—464).—Carey Lea's experiments indicate that iodine is strongly adsorbed from solution by silver iodide (A., 1888, 1). This point has been investigated; dry thoroughly washed silver iodide precipitated from *N*-solutions of silver nitrate by potassium iodide was added to aqueous and alcoholic solutions of iodine, but there was no appreciable adsorption. In the same way, moist silver iodide precipitated from *N*- and 0.1*N*-solutions of silver nitrate gave negative results. Decolorisation of the iodine solution was obtained with unwashed silver iodide both from *N*- and 0.1*N*-solutions, using an excess of silver nitrate. The decolorisation was due to a chemical reaction between the iodine in solution and the silver nitrate occluded in the silver iodide and adsorbed on its surface. Unwashed silver iodide precipitated from dilute solutions of silver nitrate and potassium iodide decolorised the iodine solutions most easily, due to the greater adsorption of silver nitrate by the silver iodide and the larger quantity of solution held because of the increased porosity. A sherry-red solution of iodine in alcohol contains about 0.008 millimol. per c.c. From the above, it is concluded that there is no adsorption of iodine from solution by silver iodide above a maximum limit of 0.02 millimol. per gram of silver iodide. Hence if Carey Lea worked with 25 c.c. of sherry-red solution of iodine, he must have added more than 10 grams of silver iodide to obtain complete decolorisation.

J. F. S.

Vapour Pressure of Selenium Oxychloride. VICTOR LENHER, G. B. L. SMITH, and G. G. TOWN (*J. Physical Chem.*, 1922, **26**, 156—160).—The vapour pressure of selenium oxychloride has been determined at temperatures from 84.3° to 117.2° by means of a modified Ramsay and Young apparatus and from 123.5° to 177.2° by the distillation method. The results are recorded for every degree over the whole range of temperature. The vapour pressure is expressed by the formula $\log P = 5.8503 + 0.000219T - 830.9/(T - 178)$, where *P* is the pressure in millimetres of mercury and *T* the temperature in absolute degrees. Selenium has no influence on the vapour pressure between 90° and 120°, selenium monochloride has no influence on the vapour pressure between 91° and 135°, neither does selenium tetrachloride up to its saturation point show any effect between 89° and 130°. Selenium oxychloride,

when distilled under reduced pressures, has a light straw colour, m. p. $10\cdot8^\circ$, b. p. $177\cdot2^\circ/744$ mm.; when distilled under atmospheric pressure, it decomposes slightly and becomes reddish-brown; this decomposition reverses to some extent on cooling. There is a small amount of dissociation between 265° and 290° , since the mean value of the molecular weight over this range of temperature is $151\cdot4$. The following minor decompositions occur at or near the boiling point under atmospheric pressure. $2\text{SeOCl}_2 \rightleftharpoons \text{SeO}_2 + \text{SeCl}_4$; $2\text{SeCl}_4 \rightleftharpoons \text{Se}_2\text{Cl}_2 + 3\text{Cl}_2$; $2\text{Se}_2\text{Cl}_2 \rightleftharpoons \text{SeCl}_4 + 3\text{Se}$; and $4\text{SeOCl}_2 \rightleftharpoons \text{Se}_2\text{Cl}_2 + 2\text{SeO}_2 + 3\text{Cl}_2$. J. F. S.

Nitrosylselenic Acid. JULIUS MEYER and WALTER WAGNER (*Ber.*, 1922, 55, [B], 690—692).—*Nitrosylselenic acid*, $\text{NO}\cdot\text{O}\cdot\text{SeO}_2\cdot\text{OH}$, a colourless, crystalline, snow-like mass, is obtained by the action of liquid nitrogen trioxide on ice-cold, anhydrous selenic acid, a large excess of the former being used initially and the greater part being subsequently allowed to evaporate. It has m. p. 80° (decomp.). It is somewhat unstable even at the atmospheric temperature, and is immediately decomposed by water. H. W.

Cathodic Reduction of Elementary Nitrogen. FR. FICHTER and RICHARD SUTER (*Helv. Chim. Acta*, 1922, 5, 246—255).—Tschirch's view that absorption of atmospheric nitrogen by leguminous plants is due to electronic action receives support from the work of Hiedemann (A., 1921, ii, 694). Since, further, biochemical oxidation can be reproduced electrochemically by use of high potential differences (Fichter, this vol., ii, 23), the attempt has been made to reduce nitrogen to ammonia by similar means. Rayleigh's contradiction (T., 1897, 71, 181) of Davy's original positive result can be disregarded, since he employed platinum electrodes and a high current density instead of gold electrodes and a low density. Experiments are now described in which, after making full allowance for the effects of impurities by blank experiments, on an average 0.3 mg. of ammonia per 160 amp. min. was consistently obtained from nitrogen under 200 atm. pressure by the use of platinum electrodes of large surface, freshly and thickly platinised, the electrolyte being 1% sulphuric acid (cf. Fischer and Priess, A., 1913, ii, 285; Tiede and Schleede, A., 1921, ii, 328). The cathode surface quickly loses its efficiency, and has to be renewed. Its activity cannot be restored by oxidation (Willstätter and Waldschmidt-Leitz, A., 1921, ii, 185). No definite evidence of ammonia formation could be obtained when a mercury cathode was employed. J. K.

Formation and Decomposition of the Higher Nitrogen Oxides. MAX BODENSTEIN, FRIEDRICH BOES, (FRL.) LINDNER, and RAMSTETTER (*Z. physikal. Chem.*, 1922, 100, 68—123).—The equilibria $\text{N}_2\text{O}_4 \rightleftharpoons 2\text{NO}_2$ and $2\text{NO}_2 \rightleftharpoons 2\text{NO} + \text{O}_2$ have been investigated over the temperature range $281\cdot7$ to $403\cdot8^\circ$ Abs., and $498\cdot9^\circ$ to $825\cdot3^\circ$ Abs., respectively. The velocity of the reaction $2\text{NO} + \text{O}_2 = 2\text{NO}_2$ has been determined at a series of temperatures. It is shown that the equilibrium $2\text{NO}_2 \rightleftharpoons \text{N}_2\text{O}_4$ is represented by the formula

$\log p_{\text{NO}_2}^2/p_{\text{N}_2\text{O}_4} = \log k_p = -2692/T + 1.75 \log T + 0.00483T - 7.144 \times 10^{-5}T^2 + 3.062$. This equation embodies the measurements of Wourtsel (A., 1920, ii, 108) between 0° and 84° , and the corrections made by the authors on measurements of the rate of oxidation of nitric oxide. The dissociation $2\text{NO}_2 \rightleftharpoons 2\text{NO} + \text{O}_2$ is to be represented by the equation $\log p_{\text{NO}}^2 \cdot p_{\text{O}_2}/p_{\text{NO}_2}^2 = \log k_p = -5749/5 + 1.75 \log T - 0.00050T + 2.839$. Both constants are true for values of p measured in atmospheres. The experimental measurements in both the above cases were effected by introducing suitable quantities of pure nitrogen tetroxide into a quartz flask and measuring the pressure with a quartz glass manometer at different temperatures. The older measurements on the velocity of combination of oxygen with nitric oxide at lower temperatures ($0-90^\circ$) and low pressures were extended to higher temperatures ($140-390^\circ$), at which the reverse reaction is considerable. The measurements were made, as before, by observing pressure changes as indicated by a bromonaphthalene manometer. The reaction takes place, as expected, according to the equation $dx/dt = kC_{\text{O}_2} \cdot C_{\text{NO}}^2 - k'C_{\text{NO}_2}^2$. The velocity increases with increasing temperature, but more slowly the higher the temperature, until eventually it is practically independent of the temperature. According to the present observations, the decomposition of nitrogen dioxide must also be measurable, and measurements were made at temperatures of $319-383^\circ$ and the velocity shown to be given by $dx/dt = k'C_{\text{NO}_2}^2 - kC_{\text{NO}}^2 \cdot C_{\text{O}_2}$. The value of k' increases with temperature 1.5 times for an increase of 10° ; k/k' gave a good agreement with the value of K deduced from the measurements. The theory of gas reactions of the third order is considered, and it is shown that simultaneous collisions between three molecules are not too few to give true trimolecular reactions, as is stated by Trautz (A., 1916, ii, 422). In the present work at 0° , where the greatest velocity is observed, it is shown that one hundred thousand times as many such collisions occur as lead to reaction, and that, from the variation of the interval between collisions of two molecules with the temperature and the probability of triple collisions, the unusual dependence of this reaction on the temperature may be deduced. J. F. S.

Purification of Phosphoric Oxide. GEORGE INGLE FINCH and RAYMOND HAROLD KELSALL PETO (T., 1922, 121, 692-693).

The Behaviour of Amorphous Carbon and Sulphur when Heated : Sulphides of Carbon. J. P. WIBAUT (*Rec. trav. chim.*, 1922, 41, 153-171; cf. A., 1919, ii, 190; also Mixter, A., 1893, ii, 571).—The sulphur which is retained by carbon after a mixture of the two elements has been heated has now been further investigated and the author considers that it is chemically combined, although no definite compound has been isolated. The product is probably a mixture of carbon with a sulphide of carbon which is practically non-volatile at 1000° , although the quantity of sulphur retained depends on the nature of the carbon and on the extent of its surface (cf. Langmuir, A., 1915, ii, 467). It is pointed out that these results are analogous to those obtained by Lowry

and Hulett (A., 1920, ii, 536) and by Rhead and Wheeler (T., 1912, 101, 831, and 1913, 103, 461) in the case of carbon and oxygen. It is suggested that the sulphur present in coal as iron pyrites is fixed by the carbon in the process of coking. H. J. E.

Processes in Gas Generators and Blast Furnaces. H. VON JÜPTNER (*Z. physikal. Chem.*, 1922, 100, 231—237).—A theoretical paper in which relationships between carbon monoxide and carbon dioxide in gas generators and blast furnaces is discussed. It is shown that the direct reduction of iron oxide by carbon in the blast furnace is advantageous, for it involves a reduction in the amount of fuel used, but the process is slow. In the electric blast furnace, better results are obtained with porous wood charcoal than with the denser coke, and also the liquid condition of the ore accelerates the process. J. F. S.

The State of Carbonic Acid in Aqueous Solution. A. THIEL (*Z. anorg. Chem.*, 1922, 121, 211—214).—A theoretical paper. Wilke (this vol., ii, 52) assumed that the increase in the strength of carbonic acid caused by the addition of neutral salts is due to the formation of orthocarbonic acid. This is shown to be incorrect. No other explanation is offered. W. T.

Solubility of Potassium Permanganate in Solutions of Potassium Sulphate and Sodium Sulphate. H. M. TRIMBLE (*J. Amer. Chem. Soc.*, 1922, 44, 451—460).—The solubility of potassium permanganate in solutions of sodium sulphate and potassium sulphate of concentrations varying between 0 and 10.75% and 0 and 21.80%, respectively, have been determined at 25°. The densities of the mixed solutions and of solutions of the single salts have also been determined for a series of concentrations. It is shown that the solubility of potassium permanganate in solutions of potassium sulphate decreases with increasing concentration of the sulphate. In solutions of sodium sulphate the solubility increases to a maximum with increasing concentration of sodium sulphate. The maximum solubility lies at about 6% of sodium sulphate, after which there is a slow decrease in solubility to a minimum in solutions which contain the maximum amount of sodium sulphate soluble in the presence of permanganate crystals. For the lower concentrations of each sulphate, potassium sulphate acts powerfully to decrease the solubility of potassium permanganate in solutions of sodium sulphate, whilst sodium sulphate increases its solubility in solutions of potassium sulphate to a much smaller degree. J. F. S.

Thermal Decomposition of Potassium Permanganate. E. MOLES and M. CRESPI (*Z. physikal. Chem.*, 1922, 100, 337—345).—The thermal decomposition of potassium permanganate has been investigated at temperatures up to 542°. It is shown that pure dry potassium permanganate shows the first signs of decomposition at 200°. The pressures observed when permanganate is heated at lower temperatures than this are due to the presence of carbonates and substances containing water. The

decomposition is complete at 240° , and the oxygen pressure of the residue up to 485° corresponds with that of pure manganese dioxide. From 485° upwards the two dissociation curves diverge from one another, and this divergence indicates the formation of the complex assumed by Askenasy and Solberg (A., 1912, ii, 1167). The absorption equilibrium does not coincide with the liberation equilibrium, a behaviour which is also observed with manganese dioxide. From the dissociation values of potassium permanganate, the heat of dissociation has been calculated and the value 60,000 cal. obtained. The absence of potassium hydroxide in the residue after heating has been established from conductivity measurements, and it is therefore proved that potassium oxide is not one of the decomposition products of potassium permanganate as indicated by Rudolf (A., 1901, ii, 388) in the equation $10\text{KMnO}_4 = 3\text{K}_2\text{MnO}_4 + 7\text{MnO}_2 + 2\text{K}_2\text{O} + 6\text{O}_2$; consequently it is assumed that manganite must be formed according to the equation $10\text{KMnO}_4 = 2\text{K}_2\text{MnO}_4 + 3\text{K}_2\text{MnO}_3 + 5\text{MnO}_2 + 6\text{O}_2$. J. F. S.

Electrolytic Production of Sodium Perborate. PEDER CHR. ALSGAARD (*J. Physical Chem.*, 1922, 26, 137—155).—A long account of experimental work on the electrolytic preparation of sodium perborate is given (*Tidskr. kem. Farm. Terap. Kristiania*, 1916, Nos. 17, 18), which is followed by an account of the author's own experiments. It is shown that by electrolysing a solution containing 45 grams of borax, 130 grams of sodium carbonate, 45 grams of sodium hydrogen carbonate, 2 grams of potassium dichromate, and 2 grams of sodium silicate per litre at 10° between a suitably shaped copper pipe as cathode and a platinum anode by means of a current of 6 amperes at 7—8 volts a yield of 11.55 grams of sodium perborate per litre may be obtained. This corresponds with a 40% current efficiency. [Cf. *J. Soc. Chem. Ind.*, 1922, 326A.] J. F. S.

The Preparation of Sodium Hydrogen Carbonate. E. TOPORESCU (*Compt. rend.*, 1922, 174, 870—873).—A study of the equilibrium of the four salts, sodium chloride, sodium hydrogen carbonate, ammonium chloride, and ammonium hydrogen carbonate, with their saturated solutions at 15° , together with measurements on certain mixtures of two or three of them together. From the results, a solubility diagram has been constructed by Le Chatelier's method (cf. A., 1921, ii, 248). From this diagram it is possible to calculate the amount of the different salts which will crystallise when a solution of known initial composition is progressively evaporated. W. G.

The Manufacture of Sodium Carbonate by the Ammonia Process. H. LE CHATELIER (*Compt. rend.*, 1922, 174, 836—841).—Using the results of Toporescu (cf. preceding abstract), diagrams are constructed by means of which it is possible to calculate the theoretical yields in the ammonia-soda process, under different conditions, and starting with mixtures of any initial composition. The theoretical maximum yield may be considerably reduced by

very slight changes in the initial composition. These results, obtained by studying the concentration of solutions initially very dilute, are applied to manufacturing conditions in which salts are added to a fixed amount of water, which is kept constant throughout the separation of the sodium hydrogen carbonate. It is possible to calculate the amount of water, on the one hand, or salt, on the other, which it may be necessary, under the manufacturing conditions, to add in order to get the maximum yield of pure sodium hydrogen carbonate.

W. G.

Density of Aqueous Solutions of Ammonium Perchlorate.

A. MAZZUCHELLI and S. ANSELMi (*Gazzetta*, 1922, **52**, i, 147—152).—The densities of aqueous ammonium perchlorate solutions of $p\%$ concentration or n -normality are given by the equations, $d_{15}=0.99913+4.6826 \cdot 10^{-3} \cdot p+1.425 \cdot 10^{-5} \cdot p^2+2 \cdot 10^{-7} \cdot p^3$; $d_{25}=0.99907+4.7898 \cdot 10^{-3} \cdot p+1.920 \cdot 10^{-5} \cdot p^2+1.33 \cdot 10^{-8} \cdot p^3$, and $d_{15}=0.99913+5.632 \cdot 10^{-2} \cdot n-5.24 \cdot 10^{-4} \cdot n^2-5.2 \cdot 10^{-5} \cdot n^3$. The results show that ammonium perchlorate dissolves in water with increase of volume at 25° and with decrease of volume at 15° , and thus support de Boisbaudran's view (cf. Roozeboom, "Heterogene Gleichgewichte," II, i, 402) that no fundamental distinction can be drawn between salts dissolving with contraction and those dissolving with expansion of volume.

T. H. P.

The Decomposition of Ammonium Nitrate by Heat.

HORACE LEONARD SAUNDERS (*T.*, 1922, **121**, 698—711).

The Ammonium Carbonate-Carbamate Equilibrium.

RUDOLF WEGSCHEIDER (*Z. anorg. Chem.*, 1921, **121**, 110—112).—A theoretical paper in which the author makes an attempt to explain the divergences of the views of Faurholt (this vol., ii, 272) and the author (*A.*, 1916, ii, 617) in connexion with the equilibrium in a solution of ammonium carbamate. It is shown that the equilibrium in such solutions cannot be regarded as definitely settled until some hypothesis is put forward which will explain the older results and those of Faurholt, or until it is shown that the results of Fenton and of Burrows and Lewis, despite their agreement, are erroneous.

J. F. S.

Purity of Atomic Weight Silver. I. Gases in Pure Silver and Iodine. GREGORY PAUL BAXTER and LEON WOODMAN PARSONS (*J. Amer. Chem. Soc.*, 1922, **44**, 577—591).—The amount of gases contained in silver and iodine which has been purified for atomic weight work has been determined. It is shown that iodine sublimed in air yields small quantities of gases when resublimed in a vacuum. The proportions varied from 0.0005% in the case of crude iodine to 0.00015% in the case of the purest material. These proportions are far smaller than those found by Guye and Germann (*A.*, 1914, ii, 727). Crude silver, when converted into iodide in a vacuum, was found to yield 0.006% of gas, whilst the purest material when treated in the same way yielded only 0.00063%. In this case also, the proportion of gas found in the pure metal

is only one-seventh that found by Guye and Germann. The foregoing figures are based on the assumption that the iodine and the silver are responsible for all the gas evolved, and that the gas is as heavy as air. It has been shown to be likely that a portion of the gas obtained was either liberated from the walls of the apparatus or diffused through the reaction tube. Furthermore, in the case of silver, at least, a large portion of the gas was found to be hydrogen. There is therefore good reason to believe that the real percentages are smaller than those given above. When the maximum corrections are applied, the atomic weights of silver and of elements referred to silver are affected in the most unfavourable cases by only 0.002 unit.

J. F. S.

Purity of Atomic Weight Silver. II. Solid Impurities.

GREGORY PAUL BAXTER (*J. Amer. Chem. Soc.*, 1922, **44**, 591—594; cf. preceding abstract).—Atomic weight silver when examined spectroscopically shows the presence of only calcium as impurity, and this in an amount not greater than 0.00004%. It is therefore concluded that in the present state of the most accurate chemical analysis the purity of atomic weight silver is fully adequate.

J. F. S.

The Absorption of Nitrogen by Calcium and its Alloys.

OTTO RUFF and HELLMUTH HARTMANN (*Z. anorg. Chem.*, 1922, **121**, 167—177).—Methods of preparing calcium alloys are given. The rate at which alloys rich in calcium absorb nitrogen depends on volume relationship, temperature, potential of the added metal, and the calcium nitride content of the alloy. The contraction of calcium in the formation of calcium nitride keeps the exposed surface porous, but pure calcium is almost passive towards nitrogen. Metals more strongly positive than calcium accelerate the adsorption (K, Ba), of the others, some have no effect (Mg, Pb, Sn), others retard it (As, Sb), and some inhibit it (Bi, Cu, Zn). The addition of more positive metals has a loosening effect on the valency electrons of calcium. Calcium nitride acts as a catalytic agent in all cases. With calcium alloys containing 5% of the nitride, pure argon can be obtained from atmospheric nitrogen in a few minutes even at a temperature below 320°.

W. T.

Zinc Borate. T. C. N. BROEKSMIT (*Pharm. Weekblad*, 1922, **59**, 265—268).—The admixture of solutions containing equivalent weights of zinc sulphate and borax causes the precipitation of a gelatinous mass, which is very difficult to filter and wash. When dry, it contains boric acid, even after prolonged washing, but zinc oxide is also present. The precipitate is soluble in boric acid solution if it is kept moist, but becomes insoluble when dried.

S. I. L.

Thallous Nitrite in certain Complex and Double Nitrites.

V. CUTTICA and A. PACIELLO (*Gazzetta*, 1922, **52**, i, 141—147).—Although thallium closely resembles in physical properties the heavy metals, thallous nitrite exhibits a chemical and physico-chemical behaviour quite different from that shown by the nitrites

of these metals. In aqueous solution, thallous nitrite, like the alkali nitrites, is highly dissociated, and the salt shows no tendency to form either complex or double nitrites. In the compounds described below, this nitrite enters with the character of the alkali nitrites.

Cupric thallous nitrite [*thallous cuprinitrite*], $\text{TL}_3[\text{Cu}(\text{NO}_2)_5]$, forms opaque, black, rounded crystals, yields pale green, aqueous solutions, and is decomposed by dilute acid, with formation of a blue solution and liberation of nitrous acid. At the ordinary temperature, both the solid salt and its aqueous solutions are stable.

In order to trace the displacement of the equilibria produced in solution of cupri-nitrous complexes by variation of the relative concentrations of the components, conductivity measurements have been made on solutions containing cupric chloride and barium nitrite in different proportions. The conductivity-composition curve shows two inflexions, due to the appearance of two complexes corresponding with the ions $[\text{Cu}(\text{NO}_2)]$ and $[\text{Cu}(\text{NO}_2)_2]$.

Thallous nickelonitrite, $\text{TL}_4[\text{Ni}(\text{NO}_2)_6]$, forms stable, flesh-red crystals and yields stable, green, aqueous solutions.

Barium thallous nitrite, $\text{TLNO}_2 \cdot 2\text{Ba}(\text{NO}_2)_2$, forms golden-yellow, prismatic crystals and is stable in the air and very readily soluble in water.

Lead thallous nitrite, $\text{Pb}(\text{NO}_2)_2 \cdot 2\text{TLNO}_2 \cdot \text{H}_2\text{O}$, forms orange-yellow crystals and in aqueous solution, especially when heated, undergoes partial hydrolysis with separation of lead hydroxide. T. H. P.

Thallium Bismutho-, Stibio-, and Arseno-thiosulphates.

G. CANNERY (*Gazzetta*, 1922, 52, i, 37—41; cf. Carnot, A., 1877, i, 50; Hauser, A., 1903, ii, 487; Szilágyi, A., 1921, ii, 199, 207).—*Thallous bismutho-thiosulphate*, $\text{TL}_3\text{Bi}(\text{S}_2\text{O}_3)_3$, prepared either from the double compound, $\text{TL}_2\text{S}_2\text{O}_3 \cdot 2\text{Na}_2\text{S}_2\text{O}_3 \cdot 8\text{H}_2\text{O}$ (cf. Vortmann and Padberg, A., 1890, 12), and bismuth chloride or from potassium bismutho-thiosulphate and a thallous salt, forms a sparingly soluble, microcrystalline, yellow powder. It is moderately stable in neutral solution at a low temperature, but decomposes with formation of sulphur dioxide and bismuth sulphide when gently heated, the decomposition being retarded by alcohol and accelerated by a trace of acid. The salt is soluble in excess of potassium bis-mutho-thiosulphate solution, yielding a clear liquid, with which alcohol forms a microcrystalline, orange-yellow precipitate containing both thallium and potassium.

Thallous stibio-thiosulphate, $\text{TL}_3\text{Sb}(\text{S}_2\text{O}_3)_3$, similarly prepared, forms a white, microcrystalline powder, and is stable in a dry atmosphere, but in presence of moisture rapidly reddens, owing to formation of antimony oxysulphide: $2\text{TL}_3\text{Sb}(\text{S}_2\text{O}_3)_3 \rightleftharpoons \text{SbOS}_2 + 4\text{SO}_2 + 3\text{TL}_2\text{S}_2\text{O}_3$.

Thallous arseno-thiosulphate, $\text{TL}_3\text{As}(\text{S}_2\text{O}_3)_3$, similarly obtained, forms a white powder and decomposes more readily than the preceding compounds, in accordance with the equations: $2\text{TL}_3\text{As}(\text{S}_2\text{O}_3)_3 = \text{As}_2\text{S}_3 + 3\text{TL}_2\text{S}_3\text{O}_6$, $\text{TL}_2\text{S}_3\text{O}_6 + 2\text{H}_2\text{O} = \text{TL}_2\text{S} + 2\text{H}_2\text{SO}_4$, and $\text{TL}_2\text{S} + \text{H}_2\text{SO}_4 = \text{TL}_2\text{SO}_4 + \text{H}_2\text{S}$. T. H. P.

Density and Molecular Volume of the Oxides of Lanthanum, Praseodymium, Neodymium, Samarium, and Europium. WILHELM PRANDTL (*Ber.*, 1922, 55, [B], 692—694).—The densities of the earth oxides depend greatly on their method of preparation and on the temperature to which they have been subjected. The nitrates when ignited give very voluminous, porous oxides which cannot be freed from adsorbed gases and consequently have low densities. The oxides obtained from sulphates are only deprived of the last traces of sulphur trioxide with great difficulty. Compact oxides are most conveniently prepared from the oxalates; the present series is therefore obtained by ignition of the latter in an electrically heated furnace at about 900° until constant in weight. Praseodymium sesquioxide is prepared by the reduction of the black oxide, Pr_4O_7 , by hydrogen at 900°. The following data are recorded :

Oxide.	Colour.	d_4^{25} .	Molecular volume.
La_2O_3	Pure white	6.51	50.08
Pr_2O_3	Greenish-yellow	6.87	48.01
Nd_2O_3	Reddish-bluish-grey	7.24	46.49
Sm_2O_3	White with yellow tinge	7.43	46.95
Eu_2O_3	Almost white with reddish-yellow tinge	7.42	47.44
Gd_2O_3	Colourless	7.407	48.95
Pr_4O_7	Black	6.71	
Eu_2O_3 (from the nitrate)		6.55	

H. W.

Corrosion of certain Aluminium Alloys. L. ROLLA (*Gazzetta*, 1922, 52, i, 79—87).—The author applies Desch's method of investigating corrosion, in which the metal is made the positive pole in the electrolysis of sodium chloride solution (A., 1911, ii, 381) to various aluminium-zinc and aluminium-magnesium alloys, which were prepared in a current of hydrogen in the electric furnace.

[With MARIO FRASSINETI].—The results obtained with the aluminium-zinc alloys show that solid solutions of zinc in aluminium undergo corrosion in such a way that neither of the two components exerts a true protecting action on the other. This result indicates that, although the solution pressure of the aluminium is greater than that of the zinc, the phenomenon is not electrochemical, but purely chemical in character. Corrosion effected by means of electrolytic chlorine does not modify substantially the ratio between the weights of the disintegrated metals remaining adherent and those dissolved or precipitated when the alloys are treated with either sea-water or sodium chloride solution. The results obtained are fully confirmed by micrographic investigation of the corroded alloys.

[With MARIO BULLI].—Two magnesium-aluminium alloys containing (1) 5.68% Mg and 93.3% Al and (2) 6.9% Mg and 92.18% Al, with small proportions of iron, silica, and carbon, were examined, the latter undergoing the more rapid corrosion. In either case, a voluminous precipitate, mostly aluminium hydroxide, was formed, and the proportion of magnesium removed was greater than that

of the aluminium. The protective action of the adherent stratum of aluminium hydroxide exerts an important influence on the course of the corrosion, rendering it discontinuous. T. H. P.

Solubility of Aluminium Nitrate Crystals in Solutions of Nitric Acid of Various Strengths at Various Temperatures.

LOWELL H. MILLIGAN (*J. Amer. Chem. Soc.*, 1922, **44**, 567—570).—The solubility of the ennea-hydrate of aluminium nitrate has been determined in nitric acid of concentrations varying from 4.98% to 72.4% at 0.4°, 20°, 40°, and 60°. The solubility increases with increasing temperature for constant nitric acid concentration, and for the same temperature decreases rapidly with increasing nitric acid concentration. J. F. S.

The Electrochemical Behaviour of Alloys of Manganese with Copper, Nickel, Cobalt, and Iron. G. TAMMANN and E. VADERS (*Z. anorg. Chem.*, 1922, **121**, 193—208).—The mixed crystals of manganese with other metals show the following limiting concentrations for the action of chemical reagents and in their electrochemical behaviour: Au-Mn 0.50 mol. Au. Ag-Mn 0.75 mol. Ag. Cu-Mn 0.50 mol. Cu. Ni-Mn 0.50 mol. Ni. Co-Mn 0.50 mol. Co. Fe-Mn none. The manganese employed was not pure, and therefore these limits are only approximate. More silver (0.75 mol.) is necessary to protect the manganese than the more positive metal copper (0.5 mol.). This limit in the case of silver holds for precipitation of metals as well as for the potential of the alloys.

In the case of copper, however, a sharp limit for the action of chemical reagents could not be obtained, but the potential showed a sudden change at the concentration 0.5 mol. Cu, and the decomposition potential of the element $\text{Cu}[\text{MnCl}_2]_x[\text{Mn}_x\text{Cu}_{1-x}]$ is independent of the composition of the alloy when $x < 5$; with $x > 5$ it decreases with decreasing amount of copper to zero. The same applies to the nickel and cobalt alloys. In the case of nickel-manganese alloys a sharp limit (0.5 mol. Ni) is found for the evolution of hydrogen from hydrochloric acid, and for the precipitation of cadmium from cadmium chloride solutions; this limit does not exist in the case of the cobalt-manganese alloys. Iron is the least able to protect the manganese in these alloys. W. T.

Preparation of Green Manganous Sulphide. FRIEDRICH L. HAHN (*Z. anorg. Chem.*, 1922, **121**, 209—210).—For the preparation of the green sulphide the following procedure is necessary. One hundred grams of manganese sulphate are dissolved in 300 c.c. of water. One hundred c.c. of 20% ammonia solution are saturated with hydrogen sulphide, and another 100 c.c. of the ammonia added (colourless ammonium sulphide). Thirty c.c. of this are saturated with sulphur at the boiling temperature and then made up to 100 c.c. with the colourless solution (yellow ammonium sulphide). To a boiling mixture of 5 c.c. of the manganese solution, 20 c.c. of 20% ammonia, and 100 c.c. of water, are added 30 c.c. of the yellow ammonium sulphide. The precipitated manganese sulphide

is dark green. To this is immediately added the colourless and yellow ammonium sulphide and the warm manganese salt is added in portions, the mixture being kept almost boiling and well stirred. If the manganese salt is added too rapidly, the flesh-coloured sulphide separates. W. T.

Structure and Simple Displacement of Iron. O. MÜGGZ (*Z. anorg. Chem.*, 1921, **121**, 68—72).—In previous papers (*Jahrb. Min.*, 1899, **2**, 63; 1901, **14**, 314), it was shown that ordinary (α) iron could not possess a plane-centred cubic structure. It is now shown, in keeping with the measurements of Westgren and Lindh (this vol., ii, 152), that the simple displacement of iron is in keeping with a space-centred cubic structure. The simple displacement which accompanies the working of α -iron is impossible in γ -iron above the transition point. Consequently, the behaviour of α -iron on working is quite different from that of γ -iron. A characteristic feature of the simple displacement in the working of α -iron is a relatively large increase in volume, which may amount to two-thirds. This is due to the formation of relatively wider concave channels. J. F. S.

The Colour of Ferric Oxide. J. ARVID HEDVALL (*Z. anorg. Chem.*, 1922, **121**, 217—224).—The colour of ferric oxide may vary from bright yellow to bluish-black. The light-coloured variety is obtained by the cautious heating of any of the sulphates of iron; the darker varieties are obtained by heating any of the other compounds of iron. The sulphate also gives the darker variety on heating with a flux or alone at 650°. When heated at 650—1000°, all the varieties become brown or dark violet; above 1000°, they become black or bluish-black (the colour of specular iron ore). Heating at above 650° causes a permanent change in colour. These varieties are explained either by the assumption of the existence of several stable modifications of ferric oxide or by that of different crystalline forms of the same modification. The author examined specimens of ferric oxide prepared by twenty-seven different methods; they were all crystalline. The bright yellow variety consisted of thin plates, the others were small grains or short prisms. They all gave the same X-ray spectrum, and therefore belong to the same system. The change from the light yellow to the darker variety was studied; it took place at 700°, the leaflets losing their form and becoming granular. W. T.

Cobalt-Tungsten Alloys. KARL KREITZ (*Metall u. Erz*, 1922, **19**, 137—140).—In the cobalt-tungsten series there is evidence of the formation of two compounds, Co_6W , melting at 1500° and CoW , melting at 1650°. Between 0 and 40% of tungsten, the alloys consist of a series of mixed crystals; from 40—70% of tungsten, of a mixture of eutectic and mixed crystals or of eutectic and the alloy CoW , according as the tungsten is less or more than 45%. The eutectic contains 44.5% of tungsten and melts at 1480°. Addition of tungsten to cobalt rapidly increases the hardness and brittleness and reduces the specific electrical

conductivity. Up to 3% of tungsten the alloys are more readily attacked by sulphuric acid than is cobalt; with more than 3% the resistance of the alloys to attack by acids rapidly increases. [Cf. *J. Soc. Chem. Ind.*, 1922, May.] A. R. P.

The Hydrolysis of the Roseocobaltic Salts. P. JOB (*Compt. rend.*, 1922, 174, 943—946).—By a quantitative application of the method previously described (this vol., ii, 301), using an oxygen electrode, the author has shown that, under the influence of hydroxyl ions, the roseocobaltic salts are converted into hydroxo-salts and that the equilibrium constant of this reaction at 16° is 6×10^{-10} . The constant of hydrolysis of the roseo-ion is calculated as being 1.2×10^5 . W. G.

Electrolytic Reduction of Chromic Chloride to the Bivalent State. M. C. TAYLOR, W. A. GERSDORFF, and E. J. TOVREA (*J. Amer. Chem. Soc.*, 1922, 44, 612—614).—When chromic chloride solution is electrolysed in a two-compartment cell between a spiral spongy lead cathode and five graphite anodes, it is reduced to chromous chloride. With a total cathode surface of 1.24 (dcm.)² and a current of 1.6 amperes, a current efficiency of 96% may be obtained over the period required to reduce 87% of the chromium, if the solution is kept stirred and if the current is reduced when hydrogen commences to be evolved. With a steady and unchanged current strength the efficiency is only 53% for the same amount of reduction. J. F. S.

Normal Chromium Azide and the Formation of Complex Salts. E. OLIVERI-MANDALÀ and G. COMELLA (*Gazzetta*, 1922, 52, i, 112—115).—The compound $\text{CrN}_9 \cdot 3\text{C}_6\text{H}_5\text{N}$, the behaviour of which indicates it to be a complex salt of the formula $\left[\text{Cr} \begin{smallmatrix} (\text{N}_3)_3 \\ (\text{C}_6\text{H}_5\text{N})_3 \end{smallmatrix} \right]$ (cf. A., 1919, ii, 468), may be obtained in better yield and purer condition by concentrating in a vacuum an alcoholic solution of chromium azide containing pyridine in solution.

Chromium azide, $\text{Cr}(\text{N}_3)_3$, prepared by prolonged treatment of an absolute alcoholic solution of crystallised chromium nitrate with anhydrous sodium sulphate and evaporation of the filtered liquid with the calculated proportion of sodium azide in a vacuum, forms a highly hygroscopic, dark green, amorphous mass. If the alcoholic solution of the chromium azide is not thoroughly dry, partial hydrolysis occurs, with formation of the basic azides, $\text{OH} \cdot \text{Cr}(\text{N}_3)_2$ and $\text{CrN}_3(\text{OH})_2$.

The compound $\text{CrN}_9 \cdot 3\text{NaN}_3$ forms green crystals, with ammonia yields a blue coloration and with silver nitrate, not silver azide, but a highly explosive complex salt, so that it evidently contains a complex anion, $[\text{Cr}(\text{N}_3)_6]'''$, analogous to $[\text{Cr}(\text{CN})_6]'''$ and $[\text{Cr}(\text{SCN})_6]'''$. The corresponding free acid, $\text{H}_3\text{CrN}_{18}$, could not be isolated owing to its ready decomposability. T. H. P.

New Views on the Constitution of the Chromic Acid. N. R. DHAR (*Z. anorg. Chem.*, 1921, 121, 99—102).—A theoretical paper in which the author considers the behaviour of chromic

acid in solution. It is suggested that chromic acid is an acid of medium strength, and that it dissociates in the two stages $\text{H}_2\text{CrO}_4 \rightleftharpoons \text{H}^+ + \text{HCrO}_4'$, $\text{HCrO}_4' \rightleftharpoons \text{H}^+ + \text{CrO}_4''$, but the dissociation constant of the first stage is extremely large in comparison with that of the second stage. Similar behaviour is found in the case of maleic and dipropyl malonic acid.

J. F. S.

The Preparation of Gaseous Metallic Hydrides from Alloys and Solutions. FRITZ PANETH, ADOLF JOHANNSEN, and MAX MATTHIES (*Ber.*, 1922, 55, [B], 769—775).—The gaseous hydrides of bismuth (A., 1919, ii, 67—68) and tin (A., 1920, ii, 41) have been obtained by melting the respective metals with magnesium and decomposing the products so formed with dilute acids. The yields, however, are very small and uncertain. Numerous attempts have therefore been made to find more favourable conditions for the formation of the alloys, but these have not resulted in any improvement in the preparation of the hydrides; and have led the authors to the conclusion that the process cannot be regarded as a simple decomposition of the magnesium bismuthide or stannide. Evidence in favour of this view is found in the observation that the relatively best yields of hydride are obtained from those portions of the mixtures in which the magnesium and tin or bismuth have not been melted together completely, that bismuth hydride is obtainable from magnesium which is merely coated with fused metallic bismuth, that the well-defined compounds, Na_3Bi and Na_4Sn , do not give a trace of metallic hydride, and, in particular, that thorium-C deposited on magnesium foil continues to evolve the metallic hydride for a long time after removal of the thin film is complete. It therefore appears possible to obtain metallic hydrides from suitable solutions of the metals instead of from their alloys and, in this connexion, the production of tin hydride by the action of magnesium on a solution of tin sulphate is described in detail; the operation proceeds with such uniformity and gives such good yields that it is suitable for lecture demonstration purposes. A solution of stannous chloride in hydrochloric acid may replace that of tin sulphate, but, in this case, the deposit formed when the gas is led through the heated Marsh tube consists of colourless stannous chloride (due to the hydrogen chloride carried forward in the gas) instead of metallic tin. This possibly accounts for the previous non-observance of the formation of tin hydride when tin is used for evolving hydrogen in Marsh's test for arsenic.

H. W.

The Preparation of Gaseous Metallic Hydrides by the Spark Discharge. FRITZ PANETH, MAX MATTHIES, and EDGAR SCHMIDT-HEBBEL (*Ber.*, 1922, 55, [B], 775—789).—Attempts are described to produce bismuth hydride by passing active hydrogen over bismuth powder, but the gas is not formed in identifiable amount. Small quantities of it are produced when rods of bismuth are used as electrodes in the formation of active hydrogen. Better results are obtained by the analogous use of lead, and the further experiments are mainly performed with this metal. The

yields with the disposition of apparatus outlined above are small and very irregular, in spite of the apparent uniformity in the conditions, but some improvement can be effected by using finely divided lead or, preferably, granulated lead packed round the platinum electrode and arranging the tube (it is figured and described fully in the original) in such a manner that the hydride produced is removed rapidly from the neighbourhood of the discharge. Even under these conditions, the experiments are very uncertain, and the yields of hydride are found to diminish with increasing purity of the reagents, thus indicating that some impurity is functioning as catalytic accelerator. Systematic search has shown that methane and the vapours of ethyl alcohol, ethyl ether, glycerol, light petroleum, paraffin, and the pyrogenic vapours from caoutchouc and paper have a very marked positive action. On the other hand, mercury, oxygen, iron, iodine, sulphur, sulphur dioxide, hydrogen sulphide, silicon hydride, ammonia, arsine, and stibine have no appreciable effect. Subsequent experiments are therefore performed in general with the addition of coal gas which is purified by successive treatment with concentrated sulphuric acid, alkaline sodium hyposulphite solution, potassium hydroxide (50%), calcium chloride, and phosphorus pentoxide. The arrangement of the apparatus is sketched and described in detail in the original. The product, which is condensed by liquid air, contains the metallic hydride mixed with a relatively large excess of hydrocarbons or the products derived from them under the action of the electric discharge, and the mixtures cannot be approximately separated by any available process of fractionation. It is therefore necessary to elaborate a method for the analysis of minute amounts of metallic hydride in the presence of an indefinite mixture of hydrocarbons. This is effected by decomposing the hydride by heat and measurement of the pressure of the resultant gases; the latter are cooled by liquid air, whereby only the hydrogen remains in the gaseous condition, and this is then absorbed by a heated palladium capillary. The pressure of residual gas is then estimated under the same conditions as previously, the difference in the readings giving the amount of hydrogen. The deposit of metal is weighed. The method gives satisfactory results with stibine and with tin hydride, to which the formula, SnH_4 , is assigned. With lead hydride it is unsatisfactory, since this substance cannot be separated by fractionation from unsaturated hydrocarbons, which subsequently become hydrogenated by the hydrogen derived from the hydride.

Under the experimental conditions adopted, volatile hydrides are formed from lead, bismuth, tin, antimony, tellurium, germanium, arsenic, and selenium, but not from aluminium, zinc, or mercury.

H. W.

Mineralogical Chemistry.

New Hypothesis of the Origin of Natural Fuels. G. CALCAGNI (*Gazzetta*, 1922, 52, i, 87—93).—The author advances and discusses the view that coal, natural graphite, petroleum, etc., may have had their origin in organic compounds, even of great complexity, existing on the earth long prior to the appearance of life.

T. H. P.

The Constitution of Anthracite. ARTHUR GROUNDS (*J. Soc. Chem. Ind.*, 1922, 41, 88—93r).—Three constituents can be recognised in anthracitic coals, namely fusain, occurring to the extent of about 1% or less, a glossy constituent similar in appearance to the vitrain of bituminous coals and designated ψ -vitrain, which forms the major portion of the seams, and a constituent similar to the clarain of bituminous seams present to the extent of 5—10% at most in the anthracites examined. The fusain was generally soft, porous, and pulverulent, and could be separated from the other constituents by fractional sieving. In certain cases, a harder variety was found associated with the soft fusain, and this contained a higher percentage both of ash and of volatile organic matter than the latter. Whilst the fusain in bituminous seams usually contains less volatile organic matter than the associated coal, the anthracitic fusain sometimes contains more and sometimes less. In one case, the dense fusain contained 23.22% compared with 11.60% for the adjacent ψ -vitrain, calculated on a moisture- and ash-free basis. The ash of fusain varies widely from about 3.0—15.0% and is much higher than the ash of ψ -vitrain, which usually varies from 0.1 to 1.5%. The water-soluble constituent of the ash of ψ -vitrain amounts to 1.7 to 6.9% of the total, and thereby differs widely both from that of vitrain, which usually contains about 70% of water-soluble substance, and of fusain, which varies from 10 to 50%. This variation with fusain is doubtless due to its porous nature, which allows the percolation of saline solutions which deposit their solid matter in the fusain on being subjected to concentration by heat. In all the seams examined, the ψ -vitrain gives an almost constant $\text{Al}_2\text{O}_3 : \text{SiO}_2$ ratio of 1.10, very similar to that of the fusain examined by the author (1.17), but very different from that of the ordinary vitrain of bituminous seams (2.55).

G. F. M.

Some Constituents of Lignites. II. R. CIUSA and M. CROCE (*Gazzetta*, 1922, 52, i, 125—128; cf. A., 1921, ii, 343).—The various natural organic compounds found in lignites may be grouped, according to their melting points, in the following classes: (1) fictelite, m. p. 46°; dinite, m. p. 35°; and scheerite, m. p. 44°; (2) simonellite, m. p. 62°; (3) konleinite, m. p. 108—114°; (4) branchite, m. p. 75—81°; bombicite, m. p. 75°; hartite, m. p. 75°; the hydrocarbon of Terni lignite (*loc. cit.*), m. p. 75°, and hofmannite, m. p. 71°.

The authors find that, when purified, all the substances of group (4), as well as mixtures of them, have m. p. 74—75°, and all have compositions and molecular weights corresponding with the formula $C_{20}H_{34}$.

The dihydrocamphene, $C_{20}H_{34}$, m. p. 74—75°, obtained by Étard and Méker (A., 1898, i, 443), and by Houben (A., 1906, i, 21), has been prepared by various methods and is found to have m. p. 85—86° (cf. Hesse, A., 1906, i, 375), its mixture with hartite having m. p. 52°. T. H. P.

Fossil Wax of Monte Falò. R. CIUSA and R. VOIS (*Gazzetta*, 1922, 52, i, 135—136).—This wax, m. p. 47—49° (crude), 50—52° (purified), contains the paraffin hydrocarbons $C_{23}H_{48}$, $C_{24}H_{50}$, and $C_{26}H_{54}$, together with higher members of the series, and is free from compounds containing sulphur or oxygen. T. H. P.

Stasite, a New Mineral Dimorphous with Dewindtite. ALFRED SCHOEP (*Compt. rend.*, 1922, 174, 875—877).—As a dirty yellow material, this occurs mixed with torbernite at Kasolo, Katanga, Belgian Congo. Under the microscope, it is seen to consist of minute, flattened prisms of a golden-yellow colour and with straight extinction, d 5.03. In the oxidising flame it fuses to a black globule, and it is soluble in nitric acid to a yellow solution. The extremes of several analyses made on two lots of material dried at 100° are given under I (also CaO 0.30, MgO trace).

	PbO.	UO ₃ .	P ₂ O ₅ .	Ign.	H ₂ O.	Insol.
I.	25.53—26.20	55.77—56.28	10.32—10.60	6.24—6.71	5.60	0.40—1.17
II.	26.40	57.19	10.68	—	5.70	—

The mean values calculated to 100% given under II correspond with the formula $4PbO, 8UO_3, 3P_2O_5, 12H_2O$, identical with that for dewindtite (this vol., ii, 305). The new mineral is thus dimorphous with dewindtite, differing from it in colour, density, and form of the crystals. L. J. S.

Aphthitalite from Kilauea. H. S. WASHINGTON and H. E. MERWIN (*Amer. Min.*, 1921, 6, 121—125).—On the walls of hot (about 800°) crevices in new lava are bright blue encrustations of small, hexagonal plates of aphthitalite, which on cooling become nearly white. In cooler parts of the crevices the encrustation consists of thenardite. The aphthitalite is optically uniaxial with ω 1.487, ϵ 1.492; analysis gave

SO ₃ .	Cl.	K ₂ O.	Na ₂ O.	CaO.	CuO.	H ₂ O.	Insol.	Total.
51.50	0.03	23.72	22.76	0.39	0.46	0.25	0.76	99.87

corresponding with K_2SO_4 43.88, Na_2SO_4 52.12, or $K_2SO_4 : Na_2SO_4$ about 1:1.5. The variable composition of aphthitalite and its relation to arcanite (orthorhombic K_2SO_4) are discussed. Anhydrous copper sulphate crystallised from hot sulphuric acid is orthorhombic (pseudo-hexagonal) with β 1.72 and $\gamma - \alpha$ 0.02, and is thus related crystallographically to the alkali sulphates. The alkalis and copper were probably volatilised as sulphides, which in contact with the air are deposited as sulphates. L. J. S.

A Tantalate and Columbite from South Dakota. WILLIAM P. HEADDEN (*Amer. J. Sci.*, 1922, [v], 3, 293—299).—Analyses of fragments of brown or greyish-black columbite: I—III, from the Old Mike mica mine in Custer Co.; IV, from Harney City, Pennington Co.; V and VI, from Tin Mountain, Custer Co. Fragments of a black tantalate (presumably tapiolite) gave VII—XIII. Crystals determined to be tapiolite from Prospect (near the Old Mike mica mine) gave XIV.

	Cb ₂ O ₅ .	Ta ₂ O ₅ .	TiO ₂ .	SnO ₂ *.	FeO.	MnO.	Total.	d.
I.	68.00	9.88	0.53	0.88	5.45	14.79	99.53	5.201
II.	67.20	10.10	0.92	0.96	5.96	15.08	100.22	5.421
III.	63.90	13.74	0.80	0.54	5.92	14.95	99.85	5.496
IV.	34.60	46.02	1.52	0.38	13.32	4.31	100.15	6.444
V.	27.22	53.47	1.30	0.44	11.91	5.66	100.00	6.725
VI.	28.81	53.67	1.63	0.38	12.00	5.20	100.69	6.845
VII.	8.63	69.55	1.50	5.29	10.84	4.19	100.00	6.954
VIII.	5.49	76.08	2.84	0.22	14.16	1.21	100.00	7.019
IX.	6.97	77.24	0.81	1.18	13.60	1.02	100.82	7.180
X.	5.56	78.28	1.33	0.28	13.35	1.22	100.62	7.468
XI.	4.32	79.50	0.92	0.32	13.42	1.56	100.04	7.794
XII.	3.69	81.40	0.58	0.12	12.55	1.73	100.07	7.878
XIII.	1.97	83.57	trace	0.10	13.28	1.19	100.10	7.975
XIV.	5.18	77.23	1.38	0.32	14.84	0.42	100.00	7.190

* Including some WO₃; small amounts of cassiterite have been deducted.

From these analyses it is seen that with increasing tantalic acid, not only is there an increase in density, but also iron replaces manganese.

L. J. S.

Analytical Chemistry.

Xylenol-blue and its Proposed Use as a New and Improved Indicator in Chemical and Biochemical Work. ABRAHAM COHEN (*Biochem. J.*, 1922, 16, 31—34).—The author recommends as an indicator 5-hydroxy-1:4-dimethylbenzenesulphonaphthalein (xylenol-blue), which has an acid range from P_H 1.2 (red) to P_H 2.8 (yellow), and an alkaline range from P_H 8.0 (yellow) to P_H 9.6 (blue). This new indicator possesses several advantages over thymol-blue, in place of which it can be successfully employed.

W. O. K.

Influence of Ethyl Alcohol on the Colour Change of Phenolphthalein. RUDOLF WEGSCHEIDER (*Z. physikal. Chem.*, 1922, 100, 532—536).—It is shown that the presence of alcohol in acidimetric titrations increases the amount of alkali necessary to produce a colour change with phenolphthalein as indicator. The cause of the slight increase is discussed and it is shown to be probable that a combination of the alcohol with the phenolphthalein explains the experimental results. (Cf. Hildebrand, A., 1909, ii, 25).

J. F. S.

The Application of the Iodine Electrode in Potentiometric Titrations. I. M. KOLTHOFF (*Rec. trav. chim.*, 1922, **41**, 172—191).—When an iodine electrode is used for the titration of an iodide by means of silver nitrate, it is found that in a neutral solution an error of 0.8% occurs, probably due to adsorption. In a solution containing free sulphuric acid the error is negligible, although the potential requires a considerable time to reach constant value. By means of the iodine electrode, an iodide may be titrated with silver nitrate in presence of an equivalent quantity of bromide or twenty times the equivalent of chloride. Mercuric perchlorate is a suitable reagent for accurate titration of iodides; mercuric nitrate in nitric acid solution is useless, as the acid oxidises a portion of the iodide. By using the perchlorate, a quantity of iodide as small as 0.6 mg. may be estimated with accuracy; good results are obtained in presence of an equivalent quantity of bromide or a large excess of hydrochloric acid.

Mercuric chloride may be estimated accurately by the potentiometric method by using the iodine electrode. Other metals have been investigated from this point of view, and it is found that the method is applicable to thallous salts, probably to palladium and some other rare metals, but not to lead or bismuth. H. J. E.

Hydrated Oxalic Acid as an Oxidimetric Standard. ARTHUR E. HILL and THOMAS M. SMITH (*J. Amer. Chem. Soc.*, 1922, **44**, 546—557).—The trustworthiness of crystallised oxalic acid as a standard in oxidation processes has been investigated. It is shown that crystals of hydrated oxalic acid as usually prepared from aqueous solution contain several tenths per cent. of included water, which is not entirely lost by four months' exposure to atmospheres of the same aqueous tension as the crystals or higher. Crystals superficially dry will lose this included water in about twenty-four hours if ground to pass a 100-mesh sieve and set in an atmosphere of an aqueous tension in equilibrium with the hydrate. A mixture of the hydrated and anhydrous oxalic acid is the only desiccating agent giving an aqueous tension in equilibrium with the hydrate at all temperatures. The powdered hydrate can be dried in about an hour in a current of air passed over this desiccating agent, so that its oxidimetric value agrees with that of sodium oxalate within 0.025%. J. F. S.

Pipettes. VERNEY STOTT (*J. Soc. Glass Tech.*, 1921, **5**, 307—325).—Pipettes should be adjusted for a particular time of delivery, and when testing pipettes it is important to record both the time of delivery and the capacity. The time required for the delivery of the contents of the pipette should be within certain limits, since pipettes are untrustworthy if the time of delivery is too short and also if it is too long. A definite time should be allowed for drainage when graduating, testing, or using pipettes. W. P. S.

Device for the Removal of Fumes. M. FISCHLER (*Chem. Zeit.*, 1922, **46**, 224).—A device for the removal of fumes during the oxidation of organic substances with sulphuric and nitric acids

consists of a wide earthenware or sheet-iron tube which is bent so as to enter the flue of a fume chamber; the lower end of the tube is provided with a small siphon trap whilst a wide side-tube receives the neck of the flask in which the material is being treated. A tubulure on the side tube admits the bent stem of a tapped funnel so that further quantities of acid may be added to the flask when necessary. The upper end of the tube may be slotted to receive a burner which aids the draught.

W. P. S.

Detection of Oxygen in Organic Compounds. JEAN PICCARD (*Helv. Chim. Acta*, 1922, 5, 243—244).—It is suggested that the fact that solutions of iodine in solvents free from oxygen are blue, but brown in alcohol, ether, or acetone, can form the basis of a method for the qualitative detection of oxygen in organic compounds. Since the brown colour depends on the formation of an additive compound with the oxygen atom, iodine in excess of this requirement must be avoided. Thus, although benzene containing 2% of ether dissolves iodine with a colour scarcely distinguishable from that obtained with pure benzene, a distinct brown colour is observable on the addition of the same proportion of ether to a solution (almost colourless) of iodine in one million parts of benzene if a layer 90 cm. deep be examined.

J. K.

Clinical Method for the Estimation of Chlorides in Blood. HERMAN FRIEND (*J. Biol. Chem.*, 1922, 51, 115—119).—For clinical purposes, plasma and not whole blood should be used for chloride estimations. In the method described, the plasma is separated immediately to prevent alterations in the distribution, precipitated by shaking with alumina cream, and chlorides estimated in the filtrate by titration with silver nitrate.

E. S.

Catalytic Analyses. I. Estimation of Chloric Acid. SHIN-ICHIRO HAKOMORI (*J. Chem. Soc. Japan*, 1922, 43, 62—70).—This method is based on the fact that the ferrous-ion acts as a catalyst in the reaction between chloric acid and hydriodic acid (Green, *J. Physical Chem.*, 1908, 12, 389). To a solution containing approximately potassium iodide 10%, ferrous ammonium sulphate 3%, and 0.2% of the sample, about 3*N*-hydrochloric acid is added, and after ten to fifteen minutes the mixture is titrated with sodium thiosulphate solution.

K. K.

The Estimation of Total Sulphur in Urine. ROBERT ROBISON (*Biochem. J.*, 1922, 16, 134—136).—Sulphur can be best estimated in the urine by a modification of Benedict's method. 2.5 c.c. of a solution of copper nitrate (40 grams) and copper chloride (15 grams) in water (100 grams) are added to the urine (10 c.c.) and the mixture is evaporated to dryness, and then heated over a plate or small flame to oxidise the sulphur. After being heated for a further twenty minutes, the residue is dissolved in 2*N*-hydrochloric acid (10 c.c.) and diluted with distilled water (300 c.c.). The oxidised sulphur is then estimated as barium sulphate.

W. O. K.

Analysis of Chlorosulphonic Acid. G. WEISSENBERGER and A. ZODER (*Z. anal. Chem.*, 1922, **61**, 41—48).—Volumetric methods, which depend on the estimation of the total acidity of the sample (after suitable dilution) and subsequent titration of the hydrochloric acid, yield untrustworthy results owing to the fact that the large quantity of sulphate ions present interferes with the titration of the chloride by Mohr's or Volhard's methods. The quantity of chlorosulphonic acid and of free sulphuric acid found is too high and the sulphur trioxide is too low. Gravimetric methods yield correct results for the hydrochloric acid and the sulphuric acid. Fairly correct results are also obtained by distilling the sample and collecting the portion which has b. p. 154—156°.

W. P. S.

Estimation of very small Quantities of Injurious Acids in Air. G. LAMBRIS (*Z. anal. Chem.*, 1922, **61**, 20—40).—Flue gases may contain sulphuric, sulphurous, and hydrochloric acids, which, when allowed to escape into the atmosphere, may have an injurious effect on vegetation. A method for the estimation of these acids consists in passing the air at the rate of about 100 litres per hour for two or three hours through an absorption vessel packed with cotton wool moistened with water and then through a second vessel containing cotton wool moistened with hydrogen peroxide. The sulphuric and hydrochloric acids are absorbed in the first vessel, whilst the sulphurous acid is absorbed and oxidised to sulphuric acid in the second vessel. The contents of the vessels are subsequently titrated and the quantities of the different acids estimated in the usual way. About 10% of the sulphurous acid is retained in the first vessel; this may be removed and conveyed over into the second vessel by means of a current of air free from acidity, or its quantity may be estimated by neutralising the contents of the vessel, using methyl-orange as indicator, then adding hydrogen peroxide, and titrating the sulphuric acid formed by the oxidation of the sulphurous acid. The method is suitable for the estimation of 1 part of acid per 500,000 parts of air.

W. P. S.

Estimation of Sulphate-ion as Barium Sulphate I. K. P. CHATTERJEE (*Z. anorg. Chem.*, 1922, **121**, 128—134).—The author carried out the precipitation under various conditions. The amount of hydrochloric acid added to make the precipitate granular and easily filtered should not exceed 0.1% of the volume of the liquid. Excess of barium chloride is not as detrimental as an excess of hydrochloric acid provided the precipitate is well washed. Rapid precipitation results in a stronger adsorption of the mother-liquor than slow precipitation. Dry barium sulphate gives up the adsorbed chloride more readily than the wet salt.

W. T.

Cathodic Deposition of Tellurium and Selenium from their Oxy-acids and their Electroanalytical Estimation. ERICH MÜLLER [with MENZEL and SCHUBERT] (*Z. physikal. Chem.*,

1922, 100, 346—366).—The detection of selenium and tellurium may be effected as follows: (a) selenium:—2 c.c. of the solution containing selenium is treated with 3 drops of concentrated sulphuric acid and a few small crystals of hydrazine sulphate and heated. A red coloration or precipitate indicates the presence of selenium. This reaction is sensitive to 5 mg. of selenium per litre; (b) tellurium:—1 c.c. of solution with 1 c.c. of concentrated ammonia and a few crystals of hydrazine sulphate when boiled give a brown coloration or precipitate if tellurium is present. This reaction is sensitive to 10 mg. of tellurium per litre; (c) selenium and tellurium in the same solution:—3 c.c. of solution are treated with 3 drops of concentrated sulphuric acid and a little hydrazine sulphate and boiled, a red coloration of selenium results; this is filtered, the filtrate made alkaline with ammonia, more hydrazine sulphate added, and the solution again boiled; a brown coloration indicates tellurium; (d) selenious acid in the presence of selenic acid:—3 c.c. of the solution are treated with 5 c.c. of concentrated sulphuric acid and a little sodium sulphite and boiled; a red precipitate or coloration indicates selenious acid. The reaction is sensitive to 5 mg. of selenium per litre. After filtration, 3 c.c. of concentrated hydrochloric acid and more solid sodium sulphite are added and the solution is boiled; a red precipitate or coloration indicates selenic acid; the sensitiveness is the same as above; (e) tellurous acid in the presence of telluric acid: these two acids cannot be detected in the presence of one another by simple reduction, since both acids are reduced, although at different rates. On electrolysis in sulphuric acid solution, the tellurous acid alone is reduced, and when this is complete the solution is concentrated and boiled with hydrochloric acid and sodium sulphite, when a brown precipitate or coloration indicates telluric acid. The cathodic decomposition potential of tellurous acid in 2*N*-sulphuric acid is -0.08 volt. No deposition potential could be obtained for tellurium from a 2*N*-sulphuric acid solution of telluric acid, from which it follows that tellurium cannot be deposited from telluric acid. A simple decomposition potential could not be observed for selenium from selenious acid, although selenium was seen to be deposited at about 0.05 volt, but the first selenium deposited appears to act as an insulating diaphragm and prevents further deposition; when the voltage is raised so that hydrogen is liberated, the selenium leaves the electrode in flakes. On adding copper sulphate to the sulphuric acid solution of selenious acid, the selenium and copper are deposited together in a conducting form at $+0.15$ volt, so that all the selenium may be deposited at this potential. Selenium cannot be electrolytically deposited from selenic acid on platinum either at 20° or 80° . Tellurium in tellurous acid may be quantitatively estimated as follows. A maximum weight of 0.25 gram of tellurous acid is dissolved in 175 c.c. of 2*N*-sulphuric acid and electrolysed for two and a half hours between two platinum Winkler electrodes which are connected directly with a single lead accumulator. The solution must be rapidly stirred during the electrolysis. The deposit, which is uniformly dense and grey, is washed with water and alcohol

and dried over sulphuric acid in a desiccator. The average error of the method is $\pm 0.1\%$. Tellurous acid may be estimated in the presence of telluric acid by this method, and after the tellurium from the tellurous acid has been removed the solution is boiled with hydrochloric acid to reduce the telluric acid to tellurous acid, and the estimation of this then carried out as before. Selenium may be estimated in selenious acid as follows. A quantity of selenious acid containing not more than 0.07 gram of selenium is dissolved in 2*N*-sulphuric acid and mixed with a quantity of copper sulphate in 2*N*-sulphuric acid such that the concentration of copper is four times that of the selenium, and electrolysed for two hours at the ordinary temperature with rapid stirring between two Winkler platinum electrodes which are connected directly with a single lead accumulator. The deposit of copper and selenium is washed with water and alcohol and carefully dried. The weight of the copper is known from the amount of copper sulphate solution used. The method is good, but suffers in accuracy on account of the small amount of selenium which may be used in a determination. Several attempts are described which were designed for the estimation of selenious acid in the presence of tellurous acid, but all failed to give the desired result. J. F. S.

The Estimation of Nitrogen in Nitric Esters. H. KESSELER, R. RÖHM, and G. LUTZ (*Z. angew. Chem.*, 1922, **35**, 145).—The esters are saponified at 40—50° with aqueous potassium hydroxide (1:1), the nitrates and nitrites so obtained reduced to ammonia with Devarda's alloy (50% copper, 45% aluminium, and 5% zinc), and the ammonia distilled into *N*/10-sulphuric acid and titrated back with *N*/10-sodium hydroxide with methyl-red as indicator. The method is specially recommended for nitro-starches, to which other rapid methods cannot be applied. H. C. R.

Colorimetric Estimation of Phosphorus. LUIGI LOSANA (*Giorn. Chim. Ind. Appl.*, 1922, **4**, 60—62).—The method developed by the author (cf. Namias, *Stahl und Eisen*, 1890) is based on the observation that treatment of ammonium phosphomolybdate with hot sodium thiosulphate solution yields a liquid coloured intensely blue owing to the formation of molybdenum compounds which have not been identified. The optimum conditions have been fixed and two colorimeters designed, one for low and the other for moderately high percentages of phosphorus. [Cf. *J. Soc. Chem. Ind.*, 1922, May.] T. H. P.

A Rapid Colorimetric Method for the Estimation of the Inorganic Phosphorus in Small Amounts of Serum. FREDERICK F. TISDALL (*J. Biol. Chem.*, 1922, **50**, 329—337).—The estimation is performed on 1 c.c. of serum. Proteins are removed by means of trichloroacetic acid, and the inorganic phosphates then precipitated as strychnine phosphomolybdate by the addition of a strychnine molybdate reagent (cf. Embden, A., 1921, ii, 462). After centrifuging and washing with the minimum amount of water, the precipitate is dissolved in 1% sodium hydroxide,

diluted, and the colour produced on reduction with 20% potassium ferrocyanide and concentrated hydrochloric acid compared with a standard. The error does not exceed 5%. E. S.

Gravimetric Estimation of Hydrogen Phosphide and a New Apparatus for Gas Analysis. L. MOSER and A. BRUKL (*Z. anorg. Chem.*, 1921, **121**, 73—94).—The reactions of phosphine with solutions of iodic acid, silver nitrate, mercuric chloride, gold chloride, and copper sulphate have been investigated with the object of finding a gravimetric method for the estimation of high concentrations of phosphine. The various methods for estimating small quantities of phosphine are discussed. Phosphine containing only a few per cent. of hydrogen is best prepared by the action of 1 : 10-sulphuric acid on aluminium phosphide, and gas obtained by this method was used in all analyses described. When phosphine is led into a 2*N*-solution of iodic acid, the gas is completely absorbed, but occasionally a small amount of fog, due to the inflammable hydride, is formed; this, however, is completely absorbed in half an hour. The solution is then heated and a current of carbon dioxide passed through until all the iodine has been distilled into a solution of potassium iodide. The amount of iodine is determined by titration with thiosulphate, and the phosphoric acid left in the flask estimated as magnesium pyrophosphate. The results obtained in both cases are identical. The reaction is represented by the equation $8\text{HIO}_3 + 5\text{PH}_3 = 5\text{H}_3\text{PO}_4 + 4\text{I}_2 + 4\text{H}_2\text{O}$. The error is a little more than 0.1%. Phosphine is shown to react with excess of silver nitrate in stages according to the equations: (1) $3\text{AgNO}_3 + \text{PH}_3 = \text{PAg}_3 + 3\text{HNO}_3$; (2) $\text{PAg}_3 + \text{AgNO}_3 + 2\text{H}_2\text{O} = \text{H}_3\text{PO}_2 + 4\text{Ag} + \text{HNO}_3$; (3) $\text{H}_3\text{PO}_2 + 2\text{AgNO}_3 + \text{H}_2\text{O} = \text{H}_3\text{PO}_3 + 2\text{Ag} + 2\text{HNO}_3$; (4) $\text{H}_3\text{PO}_3 + 2\text{AgNO}_3 + \text{H}_2\text{O} = \text{H}_3\text{PO}_4 + 2\text{Ag} + 2\text{HNO}_3$. The total action is therefore represented by the equation $8\text{AgNO}_3 + \text{PH}_3 + 4\text{H}_2\text{O} = \text{H}_3\text{PO}_4 + 8\text{Ag} + 8\text{HNO}_3$. By the action of a cold saturated solution of silver acetate added drop by drop to a vessel through which a steady stream of phosphine is added, the authors have been able to prepare *silver phosphide*, PAg_3 . This substance is an easily oxidisable, velvety, black powder which cannot be dried without decomposition; it is not decomposed by air-free cold water, but in water containing air it decomposes into metallic silver. It is rapidly decomposed by boiling water, concentrated nitric acid, warm hydrochloric acid with the liberation of phosphine, and concentrated sulphuric acid with the liberation of sulphur dioxide. The estimation of phosphine by means of silver nitrate takes place in a closed absorptiometer which consists of a bulb (120 c.c. capacity) to the bottom of which a glass tube 10 cm. long and 3 mm. bore is attached, and to the top a capillary tube which is bent twice at right angles for connecting with a gas burette. The capillary carries a tap which, in addition to the ordinary bore, is also bored along its axis. The apparatus is placed with its lower tube in a beaker containing *N*/10-silver nitrate solution and filled with the liquid and the gas drawn in and the apparatus shaken. After fifteen minutes the liquid is all run into the beaker and the apparatus

washed with a little dilute nitric acid and water, and the washings added to the beaker. The solution is then treated with hydrochloric acid, filtered, and the phosphoric acid in the filtrate estimated as magnesium pyrophosphate. The results obtained by this method are in excellent agreement with those obtained by the iodic acid method. The substitution of an ammoniacal solution of silver nitrate for aqueous silver nitrate has no advantage, and attempts to estimate the phosphine by weighing the precipitated silver failed. A *N/5*-solution of mercuric chloride can be used as absorbent for phosphine. The method of work is the same as described for silver nitrate. The reactions taking place are given by the equations (1) $6\text{HgCl}_2 + 2\text{PH}_3 = 3\text{HgCl}_2 \cdot \text{P}_2\text{H}_3 + 6\text{HCl}$; (2) $6\text{HgCl}_2 + \text{PH}_3 + 3\text{H}_2\text{O} = \text{H}_3\text{PO}_3 + 6\text{HCl} + 6\text{HgCl}$; (3) $8\text{HgCl}_2 + \text{PH}_3 + 4\text{H}_2\text{O} = \text{H}_3\text{PO}_4 + 8\text{HCl} + 8\text{HgCl}$. When the absorption is complete, the liquid and precipitate are boiled with bromine water, the mercury is removed by hydrogen sulphide, and the phosphoric acid determined as magnesium pyrophosphate. The results are uniformly good. A *N/5*-solution of gold chloride is a good absorbent for phosphine; the method employed is the same as described above. The following reactions occur during the absorption: (1) $3\text{AuCl}_3 + \text{PH}_3 + 3\text{H}_2\text{O} = 3\text{AuCl} + \text{H}_3\text{PO}_3 + 6\text{HCl}$; (2) $4\text{AuCl}_3 + \text{PH}_3 + 4\text{H}_2\text{O} = \text{H}_3\text{PO}_4 + 4\text{AuCl} + 8\text{HCl}$; (3) $3\text{AuCl} + \text{PH}_3 = \text{Au}_3\text{P} + 3\text{HCl}$; (4) $3\text{Au}_3\text{P} + \text{AuCl}_3 + 6\text{H}_2\text{O} = 10\text{Au} + 3\text{H}_3\text{PO}_2 + 3\text{HCl}$; (5) $3\text{H}_3\text{PO}_2 + 2\text{AuCl}_3 + 3\text{H}_2\text{O} = 2\text{Au} + 3\text{H}_3\text{PO}_3 + 6\text{HCl}$; (6) $3\text{H}_3\text{PO}_3 + 2\text{AuCl}_3 + 3\text{H}_2\text{O} = 2\text{Au} + 3\text{H}_3\text{PO}_4 + 6\text{HCl}$. The total and completed action is represented by the equation $8\text{AuCl}_3 + 3\text{PH}_3 + 12\text{H}_2\text{O} = 8\text{Au} + 24\text{HCl} + 3\text{H}_3\text{PO}_4$. After the action is completed, the gold in solution is precipitated by hydrogen sulphide, filtered, and the phosphoric acid in the filtrate estimated as above after oxidation by bromine water. A *N/5*-solution of copper sulphate is a good but much slower absorbent for phosphine than the other salts mentioned. The analysis, using this salt, is carried out as in the case of gold. The absorption occurs according to the equations: (1) $3\text{CuSO}_4 + \text{PH}_3 + 3\text{H}_2\text{O} = \text{H}_3\text{PO}_3 + 3\text{Cu} + 3\text{H}_2\text{SO}_4$; (2) $4\text{CuSO}_4 + \text{PH}_3 + 4\text{H}_2\text{O} = 4\text{Cu} + \text{H}_3\text{PO}_4 + 4\text{H}_2\text{SO}_4$; (3) $3\text{CuSO}_4 + 2\text{PH}_3 = \text{Cu}_3\text{P}_2 + 3\text{H}_2\text{SO}_4$; (4) $\text{Cu}_3\text{P}_2 + 3\text{Cu} = 2\text{Cu}_3\text{P}$. J. F. S.

Estimation of Krypton and Xenon in Absolute Values by Spectrophotometry. CHARLES MOUREU and ADOLPHE LEPAPE (*Compt. rend.*, 1922, 174, 908—913).—A modification of previous work (A., 1911, ii, 439, 1134). The authors have prepared mixtures of pure argon with differing amounts of pure krypton on the one hand and pure xenon on the other. For the mixtures containing krypton, the pressures at which the intensity of the line $\lambda=5870\cdot91$ of the krypton is just equal to that of the argon lines $\lambda=5860\cdot54$ and $5912\cdot31$ respectively have been determined for each mixture. For the xenon mixtures, the xenon line $\lambda=4671\cdot22$ and the argon line $\lambda=4702\cdot5$ have been used. These figures thus give a means of calculating the amounts of krypton or xenon in argon-krypton and argon-xenon mixtures respectively. The limits of dilution at which measurements can be made are for krypton $0\cdot5-4\cdot5 \times 10^{-3}$

and for xenon $2-8 \times 10^{-4}$. If the mixture to be examined contains too much krypton or xenon, it must first be diluted with pure argon, but if it is too dilute it must be concentrated by fractionation, using coconut charcoal.

W. G.

Estimation of Carbon in Cast Iron and Steel by Corleis's Apparatus. G. BATTA and H. THYSSEN (*Bull. Soc. chim. Belg.*, 1922, 31, 112—117).—By means of a number of blank experiments, errors in the estimation have been traced to incomplete drying of the gas before absorption of the carbon dioxide by alkali. Detailed experimental methods based on a considerable number of analyses have been worked out and are recommended.

H. J. E.

Rapid Analysis of Potassium Perchlorate. VICTOR LENHER and MARTIN TOSTERUD (*J. Amer. Chem. Soc.*, 1922, 44, 611—612).—The following method is described as the most satisfactory for the estimation of perchlorate. A 0.5 gram sample is mixed in an agate mortar with 1.0 gram of manganese dioxide, transferred to a porcelain crucible, and heated for fifteen minutes at $600-700^{\circ}$. After cooling, the mass is extracted with hot water, filtered, and the chlorine determined by either Mohr's or Volhard's method. The results are about 0.2—0.3% low, due to loss by volatilisation. A blank experiment with the manganese dioxide should always be carried out. Greater accuracy is obtained by Lamb and Marden's method (*A.*, 1912, ii, 681) of heating the perchlorate alone in a glass tube and retaining the fumes evolved with a plug of asbestos.

J. F. S.

Estimation of Sodium in Serum without the Use of Platinum Dishes. S. J. WILSON (*J. Biol. Chem.*, 1922, 50, 301—302).—In the estimation of sodium as pyroantimonate by Kramer and Tisdall's method (*A.*, 1921, ii, 463), the precipitation may be carried out in tinned dishes, in place of platinum ones, without loss of accuracy.

E. S.

Effect of Hydrogen-ion Concentration on the Estimation of Calcium. ALFRED T. SHOHL (*J. Biol. Chem.*, 1922, 50, 527—536).—In the estimation of calcium in the presence of magnesium and phosphates by McCrudden's method (*A.*, 1910, ii, 243; 1911, ii, 1136) the hydrogen-ion concentration must be kept within the limits P_H 4.0 and P_H 5.6. With a more acid solution, calcium oxalate dissolves, whilst with a less acid one magnesium ammonium phosphate is precipitated.

E. S.

A Rapid and Accurate Method for estimating Calcium in Urine. ALFRED T. SHOHL and FRANK G. PEDLEY (*J. Biol. Chem.*, 1922, 50, 537—544).—Uric acid and other substances present in the urine which would interfere with the titration by permanganate are oxidised by boiling with ammonium persulphate after acidification with nitric or sulphuric acid. The calcium is then precipitated by addition first of oxalic acid and then of ammonia until neutral to methyl-red, which brings the solution to P_H 4.8—5.3 (cf. preceding abstract). After remaining over-night, the precipitate is filtered

on to a hardened filter-paper, washed, transferred to the original flask, and the calcium oxalate estimated by titration with permanganate. E. S.

The Analytical Chemistry of Colloidal Disperse Systems. I. Estimation of Silver Ion in the Presence of Colloidal Silver. A. GUTBIER, J. HUBER, and O. KUPFINGER (*Ber.*, 1922, 55, [B], 748—752).—The method depends on the precipitation of the protected metallic colloid in reversible form by the addition of methyl alcohol and titration of the silver remaining in solution by Volhard's method. To avoid adsorption of silver ions by the precipitate, it is necessary, before precipitation, to convert the dissolved silver into the complex form, which can be most conveniently effected by means of solid ammonium carbonate.

A measured volume of the protected colloidal system, prepared at the atmospheric temperature and in the dark, is agitated with a corresponding excess of pure solid ammonium carbonate until the salt is completely dissolved and the precipitated silver carbonate has again passed entirely into solution. The mixture is slowly poured into methyl alcohol which has been freshly distilled over lime, after which it is allowed to remain at rest for about twenty-four hours until the precipitate has completely subsided. The supernatant liquid should now be colourless. The precipitate is carefully washed with alcohol by decantation and finally on the filter. The filtrate and washings are united, freed from alcohol by evaporation on the water-bath, cooled, acidified with nitric acid, and titrated according to Volhard's method. H. W.

The Preparation of Lead Test-papers and Notes on the Titration of Zinc with Sodium Sulphide. E. OLIVIER (*Bull. Soc. chim. Belg.*, 1922, 31, 102—109).—Experimental details of the method used in the author's laboratory for titrating zinc solutions are described and instructions for making and testing the lead papers used as indicators given. The presence of free ammonia, owing to its solvent action on zinc sulphide, introduces an error; the substance should be removed by boiling or by neutralisation. Further, ammonium salts should not be present in excess, otherwise the clearness of the end-point is impaired. H. J. E.

Estimation of Zinc as Sulphate. A. GUTBIER and K. STAIB (*Z. anal. Chem.*, 1922, 61, 97—103).—Zinc salts may be converted into sulphate by evaporation with sulphuric acid in a platinum basin, heating the residue at about 500° for fifteen minutes, cooling it, adding a small quantity of water, and repeating the evaporation and heating; the residue then consists of anhydrous zinc sulphate, and may be weighed as such; zinc sulphate does not dissociate below 675°. If desired, the sulphate may be heated over a blast-flame until no further loss in weight occurs, and the resulting zinc oxide weighed. W. P. S.

Rapid Analysis of White Metal and Similar Alloys. L. BERTIAUX (*Ann. Chim. Analyt.*, 1922, 4, 77—79).—Antimony is

estimated in one portion by solution of the sample in sulphuric acid followed by titration with permanganate. A second portion of 10 grams is mixed with fine tin filings equal to ten times the antimony content of the alloy and the mixture is treated with sufficient nitric acid to dissolve everything. The resulting mass is boiled with a large quantity of water to precipitate completely the tin and antimony, the liquid is cooled, made up to bulk, and an aliquot part filtered through a dry paper and used for the estimation of the other metals by any convenient process. The tin is determined by difference. [Cf. *J. Soc. Chem. Ind.*, 1922,.]

A. R. P.

Estimation of Aluminium in Tungsten. VICTOR FROBOESE and KÄTHE FROBOESE (*Z. anal. Chem.*, 1922, **61**, 107—110).—About 3 grams of the finely divided metal are ignited in a platinum crucible for one hour, then fused with the addition of a mixture of potassium carbonate and sodium carbonate, and cooled; the mass is dissolved in water, and the solution filtered. The filtrate, which may contain a trace of aluminium, is evaporated nearly to dryness, acidified with hydrochloric acid, evaporated, the dry residue treated with dilute hydrochloric acid and water, and the solution filtered; this filtrate is boiled, rendered alkaline with sodium hydroxide, again acidified, the aluminium is precipitated as hydroxide by the addition of ammonia, and the precipitate is collected, ignited, and weighed. The main quantity of the aluminium remains in the insoluble portion from the carbonate fusion; this insoluble portion is ignited in a platinum crucible, heated with hydrofluoric acid and a drop of sulphuric acid, fused with potassium pyrosulphate, the mass dissolved in water and sulphuric acid, and the solution filtered. The filtrate is treated with "cupferron" reagent to precipitate the iron, and the aluminium is then precipitated in the usual way in the filtrate from the iron precipitate. W. P. S.

Estimation of Free Acid in Acid Aluminium Sulphate Solutions. H. ZSCHOKKE and L. HÄUSELMANN (*Chem. Ztg.*, 1922, **46**, 302).—The following modification of Ivanov's method (*A.*, 1913, ii, 343, 1078) is recommended. Ten c.c. of the aluminium sulphate solution, 10 c.c. of barium chloride solution (1 : 10), 5 c.c. of potassium ferrocyanide solution (1 : 10) which must not be more than six days old, and 60 c.c. of boiling water are poured in the above order into a 100 c.c. flask. A gelatin solution (1 : 50) is added drop by drop with agitation until the precipitate that is formed becomes flocculent and settles easily. The mixture is cooled and diluted to 100 c.c., allowed to settle, and filtered. Fifty c.c. of the clear, colourless filtrate are diluted with 50 c.c. of water and titrated with *N*/10-sodium hydroxide until neutral to methyl-orange. The temperature of the solution must never exceed 85°, and the excess of potassium ferrocyanide must not be too great, or low results will be obtained. If the quantity of acid present be more than 6 grams per litre, the filtrate remains cloudy, when a few drops of *N*/10-sodium hydroxide should be added before precipitation. H. C. R.

Estimation of Manganese as Sulphate. J. HUBER (*Z. anal. Chem.*, 1922, 61, 103—107).—Manganese sulphate is rendered anhydrous by heating it at 360—400°; dissociation commences only at 650°. W. P. S.

Estimation of Manganese by Permanganate and Investigation of some Manganites. P. B. SARKAR and N. R. DHAR (*Z. anorg. Chem.*, 1922, 121, 135—155).—The authors find that manganese can be accurately estimated by permanganate if one of the following salts is present: magnesium sulphate, potassium nitrate, potassium sulphate, cadmium sulphate, sodium nitrate, potassium fluoride, sodium acetate and lithium, sodium, calcium, barium, and strontium chlorides. Some of the metals of these salts form manganites, others simply coagulate the hydrated manganese dioxide formed and thus make possible a sharp end-point. Coloured salts cannot be used, neither did the titrations succeed in the presence of such sparingly soluble salts as calcium phosphate or sulphate, etc., because the concentration of electrolyte is too small. The preparation, composition, and properties of several manganites are described, and the formation of manganites and the position of the metal in the periodic system discussed. Absolutely pure hydrated manganese dioxide can be prepared by heating a solution of manganous sulphate and sodium nitrate and gradually adding potassium permanganate to the well-stirred mixture. The precipitate is filtered and well washed with hot water. W. T.

The Estimation of Manganese by Knorre's Persulphate Method. P. NICOLARDOT, M. GELOSO, and E. RÉGLADE (*Ann. Chim. Analyt.*, 1922, 4, 69—77, 102—110).—Manganese dioxide, precipitated from sulphate solutions by boiling with ammonium persulphate, contains less than the theoretical amount of oxygen unless at least nine times as much iron is present, and if the precipitate is dissolved in ferrous sulphate solution and the excess of the latter titrated with permanganate, the manganese factor of the permanganate varies from $0.498 \times \text{iron factor}$ with pure solutions of manganese sulphate to $0.49176 \times \text{iron factor}$ when the original solution contains a large excess of iron. Variations in the acidity of the solution have very little effect, especially in the presence of iron, as long as the free acid does not exceed 5%; in any case, traces of manganese remain in the mother-liquor and should be estimated by boiling the solution with a further quantity of persulphate and a little silver nitrate and titrating the permanganate so formed with standard sodium arsenite. Of the common metals likely to be present in the solution, cobalt gives very, and nickel slightly, high results. Instructions are given for estimating manganese in ferromanganese and spiegeleisen. [Cf. *J. Soc. Chem. Ind.*, 1922, May.] A. R. P.

Titration of Tin with Ferric Chloride. L. SMITH (*Z. anal. Chem.*, 1922, 61, 113—120).—The hydrochloric acid solution of the tin salt is boiled, pieces of zinc foil are added, and, after about

ten minutes, the mixture is again boiled, and a further quantity of zinc is added, followed by hydrochloric acid. When all the metal is in solution, hot water is added slowly so that the boiling is not interrupted, and the solution is then titrated with ferric chloride solution. The end-point of the titration is denoted by the appearance of a greenish-yellow coloration. About 5 grams of zinc are required for each gram of tin to be reduced, and zinc is preferable to aluminium for the purpose.

W. P. S.

Micro-analysis of Organic Substances. G. WELTER (*Ann. Chim. Analyt.*, 1922, 4, 33—37).—Micro-chemical methods may be used for the estimation of carbon, hydrogen, nitrogen, halogens, sulphur, methoxyl groups, phosphorus, etc., in organic substances; these methods are described briefly. Only a few mg. of the substance are required for an estimation, but all weighings must be made on a balance sensitive to 0.001 mg. The error in the results obtained is about 0.5%.

W. P. S.

The Estimation of Aromatic Hydrocarbons in Mineral Oil Fractions. H. I. WATERMAN and J. N. J. PERQUIN (*Rec. trav. chim.*, 1922, 41, 192—198).—An attempt to apply the "aniline point" method of Tizard and Marshall (*A.*, 1921, ii, 280) to the estimation of aromatic hydrocarbons in various mineral oils and their products shows that the nature of the fraction influences the result considerably. Experiments carried out with oils containing added benzene, naphthalene, and phenanthrene show that addition of equal weights of these substances does not produce equal effects on the "aniline point."

H. J. E.

Estimation of Glycerol by the Dichromate Method. J. KELLNER (*Z. deut. Oel-Fett-Ind.*, 1921, 14, 751—752; cf. Steinfels, *Seifensieder Ztg.*, 1915, 42, 721; Tortelli and Ceccherelli, *A.*, 1914, ii, 750).—Analysis of glycerol by Steinfels's (*loc. cit.*) dichromate method gives low results unless sulphuric acid, *d* 1.230, is used instead of that of *d* 1.175.

CHEMICAL ABSTRACTS.

Physiology of the Phenols. I. The Estimation of Phenols in the Blood. K. F. PELKAN (*J. Biol. Chem.*, 1922, 50, 491—497).—The method of Folin and Denis (*A.*, 1912, ii, 1011) is modified and made applicable to blood. No heat is used in the precipitation of proteins and the concentration of filtrates, since this would cause a loss of volatile phenols. The estimation is performed on 10 c.c. of blood. Proteins are removed by shaking with tungstic acid and alumina cream, and the uric acid is precipitated by the addition of a solution of silver lactate in lactic acid. Free phenols are then estimated colorimetrically in one portion of the filtrate, using a phosphotungstic-phosphomolybdic reagent, whilst in a second portion the estimation is performed after heating for ten minutes on a water-bath with a few drops of concentrated hydrochloric acid. If the last operation is carried out in a narrow tube no loss of volatile phenols occurs and the difference between the two results then represents conjugated phenols. Since lactic acid gives a coloration with the phosphotungstic-phosphomolybdic

reagent, the standard must be treated with a corresponding amount of the silver lactate reagent (cf. Benedict and Theis, A., 1918, ii, 481). E. S.

Detection of β -Naphthol in Foods, Spices, and Beverages. YUTAKA KINUGASA and HISAJIRÔ TATSUNO (*J. Pharm. Soc. Japan*, 1922, 18—24).—The method is based on Riegler's method for the detection of nitrous acid (A., 1897, ii, 464). When a solution (1 c.c.) suspected to contain β -naphthol is treated with 2 drops of 0.1% sodium 1:4-naphthylaminesulphonate solution, 1 drop of 0.1% nitrous acid, and 1 drop of hydrochloric acid, shaken, and ammonia added, a red coloration is produced. The colour occurs with 1/1000 mg. of β -naphthol per litre. Colorations are also given by this reagent with the following substances in concentration of 1/100,000: α -naphthol, violet-red; resorcinol, quinol, phloroglucinol, pyrogallol and thymol, all yellow; phenol, orange-red; benzoic and cinnamic acids, almost colourless, and salicylic acid, slight yellow. [Cf. *J. Soc. Chem. Ind.*, 1922, May.] K. K.

Distillation Method for the Estimation of Santalol in Santal Oil. C. W. HARRISON (*J. Assoc. off. Agric. Chem.*, 1921, 5, 166—171).—In the U.S. Pharmacopœia method for estimating santalol in santal oil, the latter is acetylated and a known quantity of the dry product saponified. From the amount of alkali required to saponify the oil, the percentage of santalol present in the original substance is calculated by means of a formula given. When the santal oil is adulterated with a saponifiable oil, the method is not applicable. A modified method, which, however, is not yet completely standardised, is suggested. It consists in acidifying the product obtained after the saponification of the acetylated oil, distilling in a current of steam under specified conditions, and estimating by titration the acidity of the distillate. From this figure the percentage of santalol in the original oil can be calculated. [Cf. *J. Soc. Chem. Ind.*, 1922, 346A.] J. R.

The Estimation of Small Quantities of Dextrose by Bertrand's Process. IRENE GREINER (*Biochem. Z.*, 1922, 128, 274—278).—Accurate results for the estimation of quantities of dextrose less than 10 mg. are obtained by Bertrand's process, if 10 c.c. of the sugar solution are taken, mixed with 10 c.c. of the copper sulphate solution and 10 c.c. of a solution containing 150 grams of sodium carbonate and 30 grams of sodium hydrogen carbonate per litre are added, followed by 10 c.c. of the sodium potassium tartrate solution, and the process carried out in the standard manner, except that, after boiling the solution for three minutes, it is allowed to cool for fifteen minutes. For the titration the use of a 5 c.c. burette graduated in 1/50 or 1/100 c.c. is recommended. A table showing mg. of copper equivalent to 1 to 10 mg. of dextrose is given. H. K.

A Colorimetric Method for the Estimation of Sugars in Normal Human Urine. OTTO FOLIN and HILDING BERGLUND (*J. Biol. Chem.*, 1922, 51, 209—211).—Reducing substances are

removed from the urine by treatment with "Lloyd's alkaloidal reagent," which is described as a concentrated fullers' earth. The sugars are then estimated in the filtrate by the method of Folin and Wu (A., 1920, ii, 337). For total sugar, a separate portion of the filtrate is first hydrolysed by boiling with hydrochloric acid.

E. S.

Inhibition Phenomena in Amylases. URBAN OLSSON (*Z. physiol. Chem.*, 1922, 119, 1—3; cf. this vol., i, 390).—A method for measuring the liquefaction of starch is based on the principle of recording the time taken by a glass ball to drop in the starch solution contained in an evacuated tube.

S. S. Z.

The Estimation of Pectin as Calcium Pectate and the Application of this Method to the Estimation of the Soluble Pectin in Apples. MARJORY HARRIOTTE CARRÉ and DOROTHY HAYNES (*Biochem. J.*, 1922, 16, 60—69).—Pectin can be estimated as calcium pectate, empirical formula $C_{17}H_{23}O_{16}Ca$, by precipitation with calcium chloride and hydrochloric acid under determined conditions. A method is described for the extraction of the soluble pectin from apples, and some analytical results are given.

W. O. K.

A Reaction of Wood and some Observations on Anethole. OSKAR ADLER (*Biochem. Z.*, 1922, 128, 32—34).—All varieties of wood when warmed with a glacial acetic acid solution of phenylhydrazine hydrochloride become coloured green, adhering fragments of bark becoming reddish-brown. The colours fade on prolonged warming. Furfuraldehyde (but not pentoses), oil of anise, and oil of fennel give this green coloration. Anethole is a constituent of these oils and gives the coloration when of commercial purity, but when pure does not. The constituent responsible for the reaction has not been traced, but it distils over with anethole and is produced from anethole by oxidising agents, by the action of ultra-violet light, and by prolonged heating. Known oxidation products of anethole and allied substances do not give the reaction.

H. K.

The Analysis of Partly Hydrolysed Fats. II. W. FAHRION (*Chem. Umschau*, 1922, 29, 54—55, 60—61, 66—67, 75—76, 88—89).—In using the factor $(100 \times \text{acid number})/(\text{saponification number})$ for calculating the percentage of free fatty acids in a sample of partly hydrolysed fat, it is assumed (1) that the saponification value of the neutral fat is the same as the acid value of the fatty acids obtained from it, and (2) that the free fatty acids have no ester value, their acid value being the same as their saponification value. Neither of these assumptions is justified. The former assumption overlooks the fact that the weight of fatty acids obtained is only about 95% of the weight of fat treated, and involves an error in the percentage of free fatty acids amounting to a maximum of +1.1% when there are 50% present. That the latter assumption is not justified is shown by a list of acid values and saponification values of the fatty acids of various oils and fats. In almost every

case the saponification value is higher than the acid value, the difference being more than 10 units in the case of the fatty acids from cotton seed, apricot kernel, cherry kernel, walnut and linseed oils, and palmitic and oleic acids, especially when the oils or fatty acids have been stored for long periods in the light. This difference is shown to an even greater degree by the fatty acids of kapok and baobab oils and by those of certain marine oils. It has been explained by supposing a linking up of carbonyl groups with ethylenic linkings, but the case of palmitic acid cannot be explained in this way. A third assumption made in using the above formula is that the various glycerides in a fat are all split at the same rate. This is approximately true in the case of alkaline saponification, but it is doubtful if this is so in that of hydrolysis with steam, castor seed lipase, or hydrochloric acid. For example, castor seed lipase scarcely attacks triacetin and hydrolyses tributyrin incompletely, and the neutral fat from a sample of palm kernel oil partly hydrolysed by steam under pressure showed a markedly lower saponification number than the free fatty acids. Therefore the quantity of free fatty acids in partly hydrolysed fats cannot be calculated from the acid number, because the acid number of the fatty acids can appreciably fall both during the process and afterwards.

H. C. R.

The Analysis of Sour Milk. ANDRÉ KLING and ARNOLD LASSIEUR (*Ann. Falsif.*, 1922, 15, 95—101).—If the state of the milk is such that the sample cannot be made homogeneous, the estimation must be carried out on the whole sample. The "extract at 100°" of a milk no longer fresh must always be suspect, and may be very different from that of the fresh milk. It bears no relation to the appearance of the sample when analysed. The estimation of butter fat is more trustworthy, and varies little with time. The acidity of the fat obtained should, however, be determined to ensure that the glycerides are not partly hydrolysed. The estimation of casein precipitated by acetic acid is quite untrustworthy in the case of sour samples. The estimation of lactose is of doubtful value, but is best done by Hildt's method. The ash is affected by the partial volatilisation of chlorides to the extent of about 2%. The total nitrogen is quite unaffected. It is therefore recommended that decisions on samples of milk which have become sour should be based on estimations of butter fat and total nitrogen.

H. C. R.

A Rapid Method for the Estimation of Acetaldehyde. N. K. SMITT (*Bull. Bur. Bio-Tech.*, 1922, No. 5, 117—118).—Benzidine hydrochloride gives a yellow coloration with acetaldehyde, which increases in intensity with time and finally becomes brown or orange. Formaldehyde gives a similar colour with the reagent, but in this case it develops only slowly. For the estimation of acetaldehyde in solution, the solution is diluted until it contains from 0.1 to 5% and then to 5 c.c. of the diluted solution 5 c.c. of the benzidine hydrochloride solution are added and after thirty

minutes the colour is matched against a set of standard tubes of acetaldehyde to which the reagent was added at the same time. The method is not, however, sufficiently sensitive to be used for the estimation of very small amounts of acetaldehyde. W. G.

Detection of Acetaldehyde in Urine. WILHELM STEPP (*Biochem. Z.*, 1922, 127, 13—17).—Five % of a diluted urine is distilled through a fractionating column and the vapour absorbed in ice-cold water. The presence of acetaldehyde is best recognised by combination with dimethylhydroresorcinol. H. K.

[Estimation of] Vanillin. E. TSCHIRSCH (*Seifensieder Ztg.*, 1921, 48, 856).—Two grams of vanillin (m. p. 80—81°) are mixed with 100 c.c. of water and 30 c.c. of *N*/2-potassium hydroxide solution, the mixture being kept cold, and frequently shaken, for two hours; it is then titrated with *N*/2-sulphuric acid. The difference multiplied by 3.8 gives the percentage of vanillin.

CHEMICAL ABSTRACTS.

Estimation of Digitoxin in Digitan. L. E. WARREN (*J. Amer. Pharm. Assoc.*, 1922, 11, 8—12).—No satisfactory method of estimating digitoxin has been described, but the Keller method as modified by Fromme is considered the best. Digitan is a mixture of the tannates of the glucosides of digitalis diluted with lactose. The Keller method modified by omitting clarification with lead acetate yielded a digitoxin fraction which, when tested pharmacologically, had only about 17% of the activity of pure digitoxin.

CHEMICAL ABSTRACTS.

The Hypobromite Reaction on Urea. PAUL MENAUL (*J. Biol. Chem.*, 1922, 51, 87—88).—The author was unable to obtain accurate results in the estimation of urea by Stehle's modification (*A.*, 1921, ii, 605) of the hypobromite method.

E. S.

The Gasometric Estimation of Urea. RAYMOND L. STEHLE (*J. Biol. Chem.*, 1922, 51, 89—92).—A reply to Menaul (cf. preceding abstract).

E. S.

The Electrometric Estimation of Cyanides in the Presence of Haloids. ERICH MÜLLER and HANS LAUTERBACH (*Z. anorg. Chem.*, 1922, 121, 178—192).—The electrometric estimation of cyanides was carried out by Treadwell (*A.*, 1911, ii, 827). When silver nitrate solution is run into a solution of a cyanide containing a silver electrode a sudden increase of potential is observed when $\text{CN} : \text{Ag} = 2 : 1$ corresponding with the completion of the reaction $2\text{CN}' + \text{Ag}^+ = \text{Ag}(\text{CN})_2'$ (1). This sudden change is due to the commencement of the reaction $\text{Ag}(\text{CN})_2' + \text{Ag}^+ = 2\text{AgCN}$ (2). The authors found another sharp change in the potential when reaction (2) is completed and the concentration of silver-ion in the solution increases. This second maximum rate of change of potential is more accurate than the first, which gives a slightly low result. This second point is also of importance in the estimation of a

mixture of cyanide and haloids. The solubility products of the silver salts are $\text{AgI}=10^{-16}$; $\text{AgBr}=6.4 \times 10^{-13}$; $\text{AgCN}=4.8 \times 10^{-12}$; $\text{AgCl}=10^{-10}$. Saturated solutions of these salts contain a higher concentration of silver ion than a solution of potassium silver cyanide, because they are all soluble in a solution of potassium cyanide. Hence a solution containing cyanide and haloids should give several sharp changes of potential in an electrometric titration which would correspond with the end of the several precipitations. The solubilities of the bromide, cyanide, and chloride of silver are near to one another and sudden changes of potential corresponding with these three were not observed.

The results obtained were as follows :

(I+CN) three sudden changes giving	(1) $\frac{1}{2}\text{CN}$ (2) $\frac{1}{2}\text{CN}+\text{I}$ (3) $\text{CN}+\text{I}$
(Br+CN) two " " "	(1) $\frac{1}{2}\text{CN}$ (2) $\text{CN}+\text{Br}$
(Cl+CN) " " "	(1) $\frac{1}{2}\text{CN}$ (2) $\text{CN}+\text{Cl}$
(I+Br(or Cl)+CN) three " " "	(1) $\frac{1}{2}\text{CN}$ (2) $\frac{1}{2}\text{CN}+\text{I}$ (3) $\text{CN}+\text{I}+\text{Br(or Cl)}$
(Br+Cl+CN) two " " "	(1) $\frac{1}{2}\text{CN}$ (2) $\text{CN}+\text{Br}+\text{Cl}$

The apparatus is described.

W. T.

Sensitiveness of certain Tests for Hydrocyanic Acid.

THURE SUNDBERG (*Z. anal. Chem.*, 1922, **61**, 110—112).—The author agrees with the statement of Kolthoff (A., 1918, ii, 128) that the ferrocyanide reaction is the most trustworthy for the identification of hydrocyanic acid; the test will detect as little as 0.023 mg. of hydrogen cyanide in 10 c.c. of solution. Other sensitive tests are the guaiacum—copper sulphate test (0.001 mg. HCN) and the copper—benzidine acetate test (0.005 mg. HCN; A., 1921, ii, 224).

W. P. S.

Modification in the Process of Extraction of Alkaloids.

DOMENICO LIOTTA (*Arch. Farm. speriment. Sci. aff.*, 1921, **32**, 27—30; cf. A., 1920, i, 587).—The estimation of nicotine in tobacco may be simplified as follows. The dried, powdered material (1.5—3 grams) is completely decolorised by treatment with chlorine and then steeped in water at the ordinary temperature for twenty-four to forty-eight hours. From the resulting solution the nicotine is precipitated by means of silicotungstic acid, the precipitate being calcined; multiplication of the weight of the calcined residue by 0.1139 gives the weight of nicotine present in the sample taken. The method gives results in good agreement with those furnished by the ordinary method, and is doubtless applicable to the estimation of other alkaloids.

T. H. P.

Estimation of the Alkaloids in Extract of Aconite.

ASTROUC, E. CANALS, and R. BORDIER (*J. Pharm. Chim.*, 1922, **25**, 161—164).—The official method of the French Codex for the estimation of the alkaloids in extract of aconite gives low results owing to their incomplete extraction by the specified quantity of ether. The following method of operation is recommended in preference : 5 grams of extract are diluted to 25 c.c. with water, 10 c.c. of 10% nitric acid are added, followed by 5 c.c. of ammonia, and the solution is extracted three times with quantities of 100 c.c. of ether with vigorous and repeated agitation during ten minutes

for each extraction. It is then extracted four times with 50 c.c. of ether, and the fourth extract should be free from alkaloid when tested with Valser and Mayer's reagent. The alkaloids in the united ethereal extracts are then transferred to aqueous solution by shaking with dilute nitric acid followed by four washings with water, and they are then precipitated in the usual way from the filtered aqueous solution by adding 15 c.c. of 5% silicotungstic acid and 20 c.c. of 10% nitric acid. The precipitate is collected on a filter and ignited, and the weight of the residue multiplied by the factor 0.793 gives the weight of alkaloid in the 5 c.c. of extract taken.

G. F. M.

Method for the Estimation of Procaine [Novocaine]. A. W. HANSON (*J. Assoc. Off. Agric. Chem.*, 1921, 5, 163—166).—When novocaine is heated with 0.1N-sodium hydroxide solution, it is hydrolysed with a quantitative formation of sodium *p*-aminobenzoate. This compound without extraction can be titrated with a mixed potassium bromide and potassium bromate reagent under specified conditions. The estimation depends on the fact that each molecule of novocaine reacts with 3 molecules (6 atoms) of bromine. [Cf. *J. Soc. Chem. Ind.*, 1922, 345A.]

J. R.

The Estimation of the Purine Bases in Urine. E. SALKOWSKI (*Z. physiol. Chem.*, 1922, 119, 121—124).—Estimations of the purine bases carried out by Fridericia showed that the silver method of the author gives lower results than the Krüger-Schmid method. This is interesting in view of Steudel and Chon's recent results (this vol., ii, 239).

S. S. Z.

Estimation of Uric Acid in Blood. A. GRIGAUT (*Bull. Soc. Chim. Biol.*, 1922, 4, 11—22).—Proteins are precipitated by metaphosphoric acid and the colorimetric method of Folin and Denis (A., 1913, ii, 162) is then applied directly to the protein-free filtrate.

E. S.

Estimation of Uric Acid in Blood. STANLEY R. BENEDICT (*J. Biol. Chem.*, 1922, 51, 187—207).—In the colorimetric method described, a new arsenic phosphotungstic acid reagent is used which, with uric acid, produces a much greater depth of colour than the phosphotungstic acid reagent. The new reagent is prepared by adding 50 grams of arsenic pentoxide, 25 c.c. of 85% phosphoric acid, and 20 c.c. of concentrated hydrochloric acid to a solution of 100 grams of sodium tungstate in 600 c.c. of water, boiling for twenty minutes, and finally diluting to 1 litre. For the estimation, proteins are removed from the blood by the tungstic acid method and 5 c.c. of the filtrate are taken. After dilution to 10 c.c., 4 c.c. of a 5% sodium cyanide solution containing 2 c.c. of concentrated ammonia per litre and 1 c.c. of the arsenic phosphotungstic acid reagent are added and the mixture is immersed in boiling water for three minutes. After cooling, the colour developed is compared with a standard. Results obtained by this method are usually somewhat higher than those obtained by the precipitation method of Folin and Wu (A., 1919,

ii, 308). The possible causes of this and the relative merits of the two methods are discussed at some length in the original.

E. S.

A New Colorimetric Method for the Estimation of Plasma Proteins. HSIEN WU (*J. Biol. Chem.*, 1922, **51**, 33—39).—The method depends on the colour reaction which the three proteins give with phospho-18-molybdotungstic acid (phenol reagent) in the presence of sodium carbonate. A tyrosine standard is used, estimations by the Kjeldahl method having shown that, for human plasma, 1 mg. of tyrosine is equivalent to 16.4 mg. of fibrin, 25.2 mg. of globulin, or 27.5 mg. of albumin. Separation of the proteins is effected, in the main, by the method of Cullen and Van Slyke (*A.*, 1920, ii, 398). Fibrin and albumin, after purification, are estimated directly, whilst the albumin and globulin contained in the filtrate from the fibrin are precipitated and estimated together. The value for the globulin is then obtained by difference. E. S.

The Tyrosine Content of Proteins. OTTO FÜRTH and WALTER FLEISCHMANN (*Biochem. Z.*, 1922, **127**, 137—149).—A comparison of the various processes for the estimation of tyrosine in proteins shows that the quantity of tyrosine which can be isolated gravimetrically is far below that estimated. The method of Folin and Denis (*A.*, 1912, ii, 1012), and the colorimetric estimation by the diazo-reaction and by Millon's reagent give maximum values, the most suitable method being the absorption of bromine in acid solution by the protein hydrolysate after removal of substances precipitable by phosphotungstic acid. H. K.

Micro-estimation of Albumin. C. VALLÉE and M. POLONOWSKI (*Compt. rend. Soc. Biol.*, 1921, **84**, 901—903; from *Chem. Zentr.*, 1921, iv, 1080).—The total nitrogen is estimated in 1 c.c. of the liquid under examination by the method previously described (this vol., ii, 312). In a further 2—3 c.c. of the liquid the albumin is precipitated by addition of two drops of glacial acetic acid and a small quantity of sodium chloride, and heating on a water-bath at 90°. The liquid is made up to the original volume, centrifuged, and the nitrogen estimated in an aliquot portion of the clear liquid. The albumin nitrogen is thus obtained by difference. G. W. R.

Estimation of Pepsin. H. W. VALTEICH and C. C. GLOVER (*J. Amer. Pharm. Assoc.*, 1921, **10**, 595—606).—Comparative examination of methods for the estimation of pepsin showed that whilst Northrup's electrolytic method is trustworthy, the best method is that of U.S.P. IX, although the use of storage eggs instead of fresh eggs is recommended. CHEMICAL ABSTRACTS.

A New Method for the Estimation of Pepsin. KARL GLÄSSNER (*Biochem. Z.*, 1922, **127**, 312—315).—This method is a precipitation method and depends on the observation that globin (readily prepared from hæmoglobin by a slight modification of Strauss and Grützner's method; *A.*, 1921, i, 200) is precipitated from hydrochloric acid solution by ammonia and is insoluble in

excess of ammonia, especially if a few drops of ammonium chloride are added. A series of dilutions of pepsin or gastric juice and acid in geometrical progression are made and 1% globin solution is added to each. After digestion at 40° for fifteen minutes, addition of 3—5 drops of ammonium chloride solution (10%) and 1—2 c.c. of 1% ammonia solution gives a flocculent precipitate in the tubes containing undigested globin. H. K.

Tannase. DONALD RHIND and FRANCIS EDWARD SMITH (*Biochem. J.*, 1922, 16, 1—2).—A method has been elaborated for estimating the hydrolysing power of tannase by measuring the tannin present in a solution before and after action by the enzyme for varying periods. The tannin is estimated by titrating the whole solution with potassium permanganate in presence of indigo-carmin, the tannin is then removed by Nierenstein's caseinogen method (A., 1911, ii, 236), and the solution is again titrated with potassium permanganate. The difference between the two titrations represents the amount of gallotannin present.

W. O. K.

Estimation of Peroxydase in Milk. FRANCIS E. RICE and TORATORO HANZAWA (*J. Ind. Eng. Chem.*, 1922, 14, 201—202).—The method of Bach and Chodat for the estimation of peroxydase in plant juices (A., 1904, i, 542) is adapted for milk. It depends on the oxidation of pyrogallol by hydrogen peroxide, the reaction being catalysed by peroxydase. The number of mg. of purpurogallin precipitated by the action of 10 c.c. of milk is called the "peroxydase number." The reaction takes seven days, and air must be excluded. The residue after filtration is washed with light petroleum to remove fat. Whole and skim milk are about equal in peroxydase activity. Heating milk below 68.3° for thirty minutes reduces, but does not destroy peroxydase activity, which is also reduced slightly by keeping on ice for two days. Samples of milk for the estimation of peroxydase cannot be preserved with mercuric chloride or formaldehyde.

H. C. R.

Estimation of the Antiscorbutic Vitamin (Vitamin-C). H. C. SHERMAN, V. K. LAMER, and H. L. CAMPBELL (*J. Amer. Chem. Soc.*, 1922, 44, 165—172).—A modified diet free from vitamin-C and containing an adequate amount of vitamin-A, namely, ground oats 59%, skim milk powder, heated on open trays at 110°, 30%, freshly prepared butter fat 10%, and sodium chloride 1% is recommended. The method of measurement consists in a series of observations on animals receiving no vitamin-C and different measured amounts up to the amount which affords complete protection, and permits optimum growth. The symptoms and autopsy findings are interpreted in terms of the percentage of the required amount of antiscorbutic vitamin which was actually received by the animal in any individual case.

W. G.

Blood Analysis. IV. Ashing Methods. M. RICHTER-QUITTNER (*Biochem. Z.*, 1921, 126, 97—104).—For the estimation of total alkali metals, phosphorus and iron, for example, in blood,

ashing methods can be employed successfully. Corpuscles contain fewer mineral constituents than serum and still less if the flow of blood be stemmed in a vein.

H. K.

Analysis of Blood and Urine. I. LUDWIG PINCUSSEN and ARISTOMENIS FLOROS (*Biochem. Z.*, 1921, **125**, 42—45).—To avoid the expensive torsion balance in Bang's micro-methods, the authors recommend a fine graduated pipette holding 0.1 gram of blood. By working quickly the same pipette can be used for measuring the two drops of blood, for duplicate analysis, without use of an anti-clotting agent. Examples are given of the estimation of the reducing sugar of blood. Bang's method for the analysis of sugar in blood is applicable to sugar in urine if the coagulable protein is removed by addition of 12 c.c. of 0.05% copper sulphate solution to 1 c.c. of diluted urine (1 : 100). The results agree with the macro-method. Other substances which produce a high reduction figure are creatine and uric acid, the urine of persons with an increased uric acid and creatine content showing increased reduction.

H. K.

Analysis of Blood and Urine. II. LUDWIG PINCUSSEN and KATE MOMFERRATOS-FLOROS (*Biochem. Z.*, 1921, **125**, 46—48).—For the micro-estimation of acetone in urine, a brisk current of air is drawn for twenty-five minutes through 1—2 c.c. of urine, mixed with 20—30 mg. of oxalic acid, 0.5 gram of sodium chloride and 5 c.c. of water, and the free acetone absorbed by 10—15 c.c. *N*/100-iodine solution mixed with 1 c.c. of 33% sodium hydroxide solution. The excess of iodine is estimated volumetrically after acidifying. The combined acetone as acetoacetic acid is obtained by changing the receiver and passing the current of air through the boiling solution. For the estimation of ammonia in blood, 5 c.c. mixed with 0.2% potassium oxalate were treated with 0.1 c.c. of sodium hydroxide solution and the volume made up to 25 c.c. exactly with alcohol, whereby protein is precipitated. The solution is filtered rapidly into a vessel containing a drop of dilute acid to fix the ammonia. Twenty c.c. of filtrate were made alkaline with sodium carbonate solution and a current of air was passed through for fifteen minutes at 40° and the ammonia absorbed by 2 c.c. of *N*/50-sulphuric acid.

H. K.

Method for the Estimation of Total Base in Urine. CYRUS H. FISKE (*J. Biol. Chem.*, 1922, **51**, 55—61).—Organic matter is removed from the urine by evaporation with a mixture of sulphuric and nitric acids. Phosphates are removed from the residue by the ferric chloride-ammonium acetate method, after which the filtrate is evaporated to dryness and ignited first with sulphuric acid and then with ammonium carbonate. The residue contains the sulphates of sodium, potassium, calcium, and magnesium. Estimation of sulphate by the benzidine method (*A.*, 1921, ii, 556) gives the equivalent of the bases present.

E. S.

General and Physical Chemistry.

The Rôle of the Physicist in the Development of Chemical Theory. Presidential Address. Delivered at the Annual General Meeting, March 30th, 1922. SIR JAMES WALKER (T., 1922, **121**, 735—745).

Relationships between Molecular Refraction and Other Properties. W. HERZ (*Z. physikal. Chem.*, 1922, **101**, 54—62).—From Berthelot's equation, $M=11.4 d \{T_k/p_k(2-T/T_k)\}$ (A., 1899, ii, 404), and Guye's expression, $MR=1.8T_k/p_k$, in which MR is the molecular refraction and the other symbols have their usual significance, the author has derived the expression $M/d_s=5MR$, where d_s is the density at the boiling point. This expression says that the molecular volume for non-associated liquids at the boiling point has a value equal to five times the molecular refraction. This statement has been tested for thirty-three substances and shown to be fairly in keeping with facts in the case of the simpler aliphatic liquids, but in the case of halogen-substituted compounds, aromatic, heterocyclic, and inorganic substances, large divergences are observed. The expression $MR=M/5d_s=0.9c/d_kT_s$ is deduced, from which $M=11.97c/T_s$ is obtained, where c is the molecular boiling-point elevation constant. This equation is shown to be in keeping with fact to the extent of 10—20%. Further expressions are derived by which from the molecular volume at the boiling point and the molecular refraction the molecular diameter and the internal pressure may be calculated. In many cases it is shown that $ML/MR=8p_k$, where L is the latent heat of vaporisation at the boiling point. A constant relationship is shown to exist between the coefficient of expansion, the molecular refraction, and the critical pressure, which has the form $MRp_k/(1/k_{20}+293)=\text{constant}$ (0.9). Evaluation of this constant for seventeen substances gives figures which lie between the extremes 1.38 and 0.84.

J. F. S.

Regularities in Systems of Bands. A. KRATZER (*Ann. Physik*, 1922, [iv], **67**, 127—153).—The work of Heurlinger (*Diss. Lund.*, 1918), and Lenz (*Verh. Deut. physikal. Ges.*, 1919, **21**, 632) on series lines in band spectra is extended by the application of the quantum theory to the consideration of the anharmonic vibrations of a molecular oscillator. A formula, similar to that deduced empirically by Deslandres (A., 1920, ii, 69) is obtained for the zero lines in a series of bands, and is found to represent very exactly the lines in the band spectrum of oxygen and the lines in the violet band spectrum of cyanogen. The significance of the constants in Deslandres's formula is interpreted on the basis of the quantum theory, and a formula involving nine constants is obtained, to represent the whole of the lines in the violet band spectrum of

cyanogen, exclusive of the fine structure. The application of the theory to the lines in the red band spectrum of cyanogen and the band spectrum of nitrogen is briefly indicated. J. S. G. T.

The Structure of the Balmer Series. OTTO OLDENBERG (*Ann. Physik.*, 1922, 67, 69—72).—The new measurements of McLennan and Lowe (A., 1921, ii, 666) for the H_{α} , H_{β} , H_{γ} , and H_{δ} lines are not considered to be in opposition to the Sommerfeld theory of the structure of the Balmer series, the differences being explained by the neglect of the small shift which is caused by the steep rise in intensity of a neighbouring line. Experiments with the discharge tube used by Wood (A., 1920, ii, 569) indicate that the third members of the triplets found by McLennan and Lowe are not produced in the absence of nitrogen. The new lines which are observed on the admission of nitrogen are found to be somewhat sharper than the Balmer doublet. W. E. G.

Extension of the X-Ray Spectrum to the Ultra-violet. E. H. KURTH (*Physical Rev.*, 1921, 18, 461—477).—Certain improvements in technique made it possible to obtain the following values for the convergence wave-lengths in Å.U.: *K* series, carbon 42·6, oxygen 23·8; *L* series: carbon 375, oxygen 248, aluminium 100, silicon 82·5, titanium 24·5, iron 16·3, copper 12·3; *M* series: aluminium 326, titanium 85·3, iron 54·3, copper 41·6; *N* series: iron 247, copper 116. It is suggested that the radiation from solid targets may differ from the radiation from gaseous atoms, especially for the lighter elements. A. A. E.

The Significance of the Half Terms in Spectral Series Formulæ. PAUL D. FOOTE and F. H. MOHLER (*J. Opt. Soc. Amer.*, 1922, 6, [i], 54—56).—The ratio of the constant a^* for the enhanced spectrum of an alkali earth to the constant a for the arc spectrum of the alkali with the next lower atomic number has been computed on the basis of the quantum theory and compared with the observed values. W. E. G.

The Occurrence of Spark Lines (Enhanced Lines) in the Arc. II. Magnesium, Zinc, and Cadmium. G. A. HEMSALCH and A. DE GRAMONT (*Phil. Mag.*, 1922, [vi], 43, 834—871; cf. this vol., ii, 176).—Further experimental observations have been made on the spectra observed in liquid and vacuous film arcs and in arcs surrounded by gaseous media. The results point to the existence of four distinct stages in the spectral development of the arc flash (1) the film stage during which the spark lines are enhanced and symmetrically broadened, (2) a short region marked by the strong emission of spark lines, (3) a stage during which the spark lines are but feebly emitted and at the end of which their emission ceases abruptly, and (4) the last period of the flash during which arc lines are alone emitted. The flashing up of the spark lines and their symmetrical broadening at the instant of striking the arc is shown to be due to the momentary establishment of a powerful electric field between the electrodes. The symmetrical broadening of the lines is believed to be due to the Stark effect.

The disappearance of the magnesium spark line λ 4481 at a definite point during the second phase occurs when the electric field is reduced below 500 volts/cm. The spark lines are brought out prominently in liquid air and disappear when the temperature is raised. It is thus impossible to regard these as high temperature lines. The only types of magnesium lines which are emitted in the absence of electrical or special chemical actions are the flame line λ 4571 and the series triplets. A summary of the relative sensitiveness of the various magnesium lines to electrical and chemical actions is given.

W. E. G.

The Spectrum of Glucinum Fluoride. SNEHAMOY DATTA (*Proc. Roy. Soc.*, 1921, [A], **101**, 187—194).—The previous investigation on the band spectra of the fluorides of magnesium, calcium, strontium, and barium (cf. A., 1921, ii, 529) has been extended to a study of the spectrum of potassium glucinum fluoride. The presence of potassium does not interfere with the band spectrum of glucinum fluoride, neither does glucinum oxide present spectra in the vicinity of the fluoride bands. There are six groups of bands all in the ultra-violet between λ 2800 and λ 3400 and all fading towards the red. The strongest band at 3009 was selected for a detailed investigation, and it was shown to consist of three series of lines which depart from the ordinary formulæ. Each group of glucinum fluoride bands is composed of alternating double and single heads, and this is the only similarity they present to the spectra of the other alkaline-earth fluorides. The formula, given in the previous paper for the E_1 group of magnesium fluoride gives satisfactory agreement for the glucinum fluoride bands.

W. E. G.

Emission Bands of Erbium Oxide. E. L. NICHOLS and H. L. HOWES (*Science*, 1922, **55**, 53).—A confirmation of the work of Mallory (*Physical Rev.*, 1919, **14**, 54). At temperatures at, or slightly below, 1000° , there are three regions, in which the principal emission bands of the spectrum of erbium oxide are situated, which are brighter than for an ideal black body at the same temperature.

A. A. E.

N-Series in X-Ray Spectra. V. DOLEJŠEK (*Nature*, 1922, **109**, 582).—By using Siegbahn's apparatus, lines which most probably must be referred to the *N*-series of uranium and thorium have been detected. Calculations show that for these elements, some of the lines must have wave-lengths of about the measured values ($8\cdot6$ — $12\cdot0$ Å.U. and $9\cdot4$ — $13\cdot5$ Å.U., respectively), whilst for bismuth and the elements in its neighbourhood all the *N*-lines must have a wave-length of more than 13 Å.U.

A. A. E.

Spectrophotometric Study of Solutions of Cupric Chloride. FREDERICK H. GETMAN (*J. Physical Chem.*, 1922, **26**, 217—246).—The extinction coefficients of a series of solutions of cupric chloride ($2\cdot622$ — $0\cdot8443M$) have been determined at $28\cdot3^\circ$ for various wave-lengths lying between $460\text{ }\mu\mu$ and $560\text{ }\mu\mu$. A further series of extinction coefficients of a number of binary mixtures of cupric

chloride with hydrochloric acid, potassium chloride, lithium chloride, calcium chloride, aluminium chloride, and zinc chloride, respectively, have also been determined over the same region of the spectrum; in each case the concentration of the copper salt was kept constant ($0.8443M$), whilst the concentration of the added chloride was progressively increased. It is shown that in solutions of cupric chloride, Beer's law holds only in the region of longer wave-lengths, and that in this region both ions and undissociated molecules may be assumed to possess nearly equal absorbing powers. No apparent relation has been found to exist between absorption and dissociation; in fact, no satisfactory interpretation of the mechanism of the absorption of light by electrolytes seems to be possible, unless the view of Hantzsch be adopted, that absorption is independent of dissociation. The absorption of light by solutions of cupric chloride has been attributed to the presence of complex ions, the composition of which is conditioned, primarily, by the concentration of the solution. In extremely dilute solutions, absorption is assumed to be due to the presence of copper atoms holding in combination as many molecules of solvent as correspond with the co-ordination number of the metal, which, in this case, is four. In extremely concentrated solutions, complex anions containing copper have been shown to be present, and to these the hypothetical formula $CuCl_4''$ has been assigned. Other complex ions, in which one, two, or three of the water molecules of the tetrahydrated copper-ion have been replaced by chlorine atoms, are assumed to be present in solutions of intermediate concentrations. A study of the absorption spectra of the metal amines has shown that substitution of a chlorine atom for a water molecule in a complex ion causes a displacement of the minimum in the absorption curve toward the red end of the spectrum. Increase in concentration, or the addition of colourless chlorides, is shown to displace the absorption minimum of $0.8443M$ cupric chloride toward the red end of the spectrum. The cause of the observed displacement of the absorption minimum is believed to be the same as in the case of the metal amines, namely, the successive replacement of water molecules by chlorine atoms in complex ions.

J. F. S.

Absorption of Light by Solutions of Cupric Lithium Chloride. FREDERICK H. GETMAN (*J. Physical Chem.*, 1922, 26, 377—383).—The absorption of light by solutions of cupric lithium chloride ($CuCl_2 \cdot LiCl \cdot 2H_2O$) has been examined spectrophotometrically, and the resulting absorption curves have been compared with corresponding curves for cupric chloride. The minima in the absorption curves were found to lie nearer the red end of the spectrum than the corresponding minima in the curves obtained with solutions of cupric chloride. In dilute solutions, the absorption curves of cupric lithium chloride and cupric chloride were found to be identical. From a comparison of the conductivities of solutions of cupric lithium chloride with the sum of the conductivities of cupric chloride and lithium chloride at the

same dilutions, it is found that the double salt, when dissolved in water, undergoes almost complete dissociation into the two component salts. The possibility suggested by Donnan that the solid salt may undergo dissociation as indicated by the equation $\text{CuCl}_2, \text{LiCl}, 2\text{H}_2\text{O} \rightleftharpoons (\text{Li}_2\text{H}_2\text{O})^+ + (\text{Cu}_3\text{Cl})^-$ (A., 1905, ii, 806) is emphasised as offering a satisfactory explanation of the fact that the colour of the compound is widely divergent from that characteristic of copper salts. J. F. S.

Absorption Spectra of Glasses of Various Composition containing Didymium. F. WEIDERT (*Z. wiss. Photochem.*, 1922, 21, 254—264).—A spectrographic comparison of the absorption spectrum of didymium present as a solid solution in various glasses with that of aqueous solutions of the nitrate of didymium and other rare earths, and an investigation of the effect of the composition of the glass on the absorption spectrum. Eight of the glasses investigated contained various amounts of four or five of the oxides of silicon, boron, arsenic, lead, aluminium, calcium, potassium, and sodium, together with a mixture of the oxides of neodymium, praseodymium, samarium, cerium, and lanthanum. The remaining glass contained pure neodymium oxide. The aqueous solutions employed for comparison included, not only a solution of the mixture referred to, but also solutions containing respectively the equivalent of 1.37 gram of neodymium oxide, 0.76 gram of praseodidymium oxide, and 0.42 gram of samarium oxide per litre. The absorption bands characteristic of the glasses were broader and less sharp than the corresponding bands afforded by the aqueous solutions. The presence of boron in the glass increased the want of sharpness of the bands. Lead-potash glasses afforded the sharpest bands. Reproductions of certain of the absorption spectra are given, and the paper discusses the shift of the bands in various cases. J. S. G. T.

The Absorption of Colouring Matters of the Flavone Group.
II. Absorption Spectra of Phenyl Styryl Ketone and its Derivatives. YÛJI SHIBATA and WILLY NAGAI (*J. Chem. Soc. Japan*, 1922, 43, 101—122).—To determine the effect of the hydroxyl group on the absorption of light in compounds intimately related to the flavone group, the authors have studied the absorption spectrum of phenyl styryl ketone and its hydroxyl derivatives. Phenyl styryl ketone in 0.0001*N*-alcoholic solution shows a large absorption band in the ultra-violet (the centre being λ 3.300). The absorption of flavone, therefore, is more hypsochromic than that of phenyl styryl ketone in the same concentration (cf. Shibata and Kimotsuke, *J. Chem. Soc. Tokyo*, 1918, 39, 771). The effect on the absorption of light of the introduction of the hydroxyl group into phenyl styryl ketone is more complex than in the case of flavone. The effect of the introduction of the hydroxyl group into the phenyl group is very small; whilst in the styryl group it is distinct. In the latter case, it is bathochromic and accompanies the appearance of one or two new absorption bands. By introducing the hydroxyl group into both the phenyl and styryl groups

at the same time, the band is enlarged, and is bathochromic. When two hydroxyl groups are introduced into the group, the effect is additive, except in some cases of ortho-replacement. From the point of view of light absorption, natural naringenin is not shown to be a derivative of phenyl styryl ketone. K. K.

Spectrographic Detection of Change in Blood Pigment due to certain Organic and Inorganic Poisons, and the Characteristic Spectra of Yolk of Egg and Urobilin. L. LEWIN and E. STENGER (*Z. wiss. Photochem.*, 1922, **21**, 221—226; cf. A., 1908, ii, 1048).—In continuation of previous work, the authors have investigated photographically the effect of hydrocyanic acid, phenylhydrazine, acetylene, hydrogen phosphide, and sulphurous acid on the absorption bands of blood, both when the poison was added to blood withdrawn from an animal and when the poison was injected into the living animal. Only in the case of addition of phenylhydrazine did any appreciable shift of the absorption bands occur. In the case of blood derived from a rabbit after injection of phenylhydrazine, absorption bands occurred at $\mu\mu$ 629, 609, 581, 557, and 402, compared with absorption bands at $\mu\mu$ 577, 537, and 415 for normal blood. In the case of hydrocyanic acid, whilst no appreciable shift of the bands occurs, the sharpness of the bands diminishes, at first slowly, as the percentage of hydrocyanic acid increases. The absorption spectrum of yolk of egg showed bands at $\mu\mu$ 482, 453, 429, 401, and 380. The absorption spectrum of preparations of urobilin showed a band in the region $\mu\mu$ 494. By addition of an alcoholic solution of zinc chloride and ammonia, the maximum intensity in the band moves to $\mu\mu$ 509 and the solution fluoresces. J. S. G. T.

Optical Rotatory Dispersion. II. Tartaric Acid and the Tartarates. THOMAS MARTIN LOWRY and PERCY CORLETT AUSTIN (*Phil. Trans.*, 1922, [A], **222**, 249—308).—Bakerian lecture. The rotatory power of tartaric acid for a series of nine wave-lengths has been determined in aqueous solutions of eleven different concentrations ranging from 5 to 55% by weight, and also for twenty-one and twenty-six wave-lengths, respectively, at two other concentrations. The optical rotatory power of tartaric acid, like those of its methyl and ethyl esters, is expressed to a close degree of approximation by the formula $\alpha = k_1/(\lambda^2 - \lambda_1^2) - k_2/(\lambda^2 - \lambda_2^2)$. The rotatory power of sodium tartrate agrees very closely with Biot's law, $\alpha = k/\lambda^2$, but requires for its exact expression a two-term formula similar to that used for tartaric acid. Potassium and ammonium tartrate and Rochelle salt give similar dispersion curves, but deviate more widely from Biot's law. A number of solutions which give negative rotations have been examined for a range of wave-lengths and the corresponding dispersion curves have been plotted. In the presence of an excess of boric acid, the rotatory dispersion of tartaric acid is no longer complex but simple, and may be expressed over a wide range of wave-lengths by the equation $\alpha = k/\lambda^2 - \lambda_0^2$. Tartar emetic also gives a simple dispersion curve. When tartar emetic is dissolved in an excess of potassium hydroxide,

or when a corresponding solution is prepared with bismuth in place of antimony, strongly levorotatory solutions are obtained, but these again are characterised by a simple rotatory dispersion. It is suggested that tartaric acid, like nitrocamphor, exists in solution in two labile isomeric forms, and that the anomalous dispersion of the acid and of many of its derivatives is due to the presence of two isomeric compounds of opposite rotatory power and unequal dispersion. Derivatives which give simple rotatory dispersion are assumed (like salts of nitrocamphor) to be fixed in one of these forms. J. F. S.

The Inversion of the Rotatory Power in Anisotropic Liquids.

L. ROYER (*Compt. rend.*, 1922, **174**, 1182—1185).—Cholesteryl cinnamate shows inversion of its rotatory power. The wave-length at which this inversion occurs varies with the temperature, the wave-length increasing as the temperature decreases. The wave-length of the reflected light varies with the temperature in the same manner, and it would appear that these two wave-lengths are identical. W. G.

Phototropy of Inorganic Salts. Cuprous Chloride and Bromide. GOPAL SINGH (T., 1922, **122**, 782—785).

Conversion of Energy in Photochemical Processes.

A. BYK (*Z. wiss. Photochem.*, 1922, **21**, 197—199).—The contributions of Grothaus and of Draper to the theory of photochemical processes are briefly reviewed, and attention is directed to outstanding ambiguities in the application of Einstein's quantum photochemical law to such processes. J. S. G. T.

Velocity of Reaction in the Photochemical Dissociation of Ferric Chloride. E. PUXEDDU and F. L. VODRET (*Gazzetta*, 1922, **52**, i, 229—234; cf. A., 1920, ii, 406).—The velocity of decomposition of ferric chloride in ethereal solution according to the equation $\text{FeCl}_3 = \text{FeCl}_2 + \text{Cl}$ has been measured under various conditions, both sunlight and electric arc light being used to effect the reaction. The course of the reaction was followed by titrating the ferrous salt with permanganate in presence of a few c.c. of Zimmermann's solution. As far as a certain stage in the reaction, the latter follows the equation for unimolecular reactions. The change proceeds more slowly in arc light than in sunlight, and yields a dense, highly transparent precipitate of ferrous chloride. T. H. P.

An Attempt to Detect Induced Radioactivity resulting from α -Ray Bombardment. A. G. SHENSTONE (*Phil. Mag.*, 1922, [vi], **43**, 938—943).—It is shown that the violent dismemberment of a molecule by an α -particle does not give rise to unstable nuclei, capable of emitting mass particles of a range greater than 2.0 mm. Also no cumulative effect is observable after a heavy bombardment by the α -particles. These negative results do not preclude the possibility of radioactive disintegrations taking place which involve the emission of β -particles or γ -radiation. W. E. G.

The Production of β -Ray Spectra by Radioactive Substances. LISE MEITNER (*Z. Physik*, 1922, 9, 131—144).—Experiments on the β -ray spectra of thorium-*B* have shown that it is possible to distinguish between a primary β -ray emitted from the nucleus and a secondary β -spectrum from the *K* and *L* rings of electrons. The primary β -rays are supposed to be converted into γ -rays in the nucleus and these set free electrons from the envelope of the atom. The source of the secondary β -rays was ascertained by allowing the γ -rays from thorium-*B* to set free electrons from an isotopic element, namely, lead, and comparing the velocities of these electrons with those emitted from the radioactive element. Two groups of β -rays were emitted identical with those from thorium-*B*. The energy of the γ -rays calculated from the energy of the secondary electrons, and the work required to separate these electrons from the nucleus, were found to agree with a wavelength $\lambda_\gamma = 5.2 \times 10^{-10}$ cm. and with a velocity for the primary β -ray of 73.9%. This β -ray was found to be present in the spectra emitted from thorium-*B*. The intensity was, however, very small, so that the primary β -ray emission is almost entirely converted into γ -rays. Evidence of the presence of electrons from the *K*, *L*₁, and *L*₂ rings was found. Similar conclusions are drawn for other elements, but the β -spectra of radium and radiothorium are entirely secondary in origin. The velocity of the primary β -ray, the atomic weight, and the life period of the isotopes of lead, radium-*B*, thorium-*B*, and radium-*D* are shown to be intimately related.

W. E. G.

The Relationship between β - and γ -Rays. LISE MEITNER (*Z. Physik*, 1922, 9, 145—152; see preceding abstract).—Arguments are adduced for the conclusion that γ -rays from radioactive substances have their origin in the nucleus. From the behaviour of uranium-*X* and radium-*C*, it is concluded that the primary β -ray is intense only when the γ -ray emission is small. The results of Ellis (*A.*, 1921, ii, 422) on radium-*B* preparations are explained on the assumption that there are three γ -ray frequencies emitted from the nucleus, two of which correspond with two different primary β -rays. The middle γ -ray could be accounted for if the most rapid β -ray under certain conditions gave up part of its energy and left the nucleus with the remainder.

W. E. G.

Reflection of X-Rays from Imperfect Crystals. C. G. DARWIN (*Phil. Mag.*, 1922, [vi], 43, 800—829).—In the recent work of Bragg, James, and Bosanquet (*A.*, 1921, ii, 477) on the reflection of X-rays from crystals, use was made of the theoretical formulæ of the author. The position of the electrons in a crystal can be inferred from a study of the intensity with which the various faces of a crystal reflect a given wave-length. The problem falls into two stages, the first from crystal to molecule and the second from molecule to electron. The formulæ are, however, falsified by the phenomenon of extinction. Extinction is shown to lead to two effects, primary and secondary. The primary diminishes the reflection from a perfect crystal below the amount given by the

simpler theory. This is, however, negligible for crystals invisible under a high-power microscope. The reflection from a small, perfect crystal, a conglomerate composed of a large number of small crystals oriented in nearly the same direction, and crystal powders is examined, and formulæ are worked out for Q , a quantity which is, approximately, the amplitude of the wave scattered by all the electrons in a single atom in the direction of the reflected beam.

W. E. G.

Positive Ray Analysis of Lithium and Magnesium. A. J. DEMPSTER (*Physical Rev.*, 1921, **18**, 415—422; cf. *ibid.*, 1918, **11**, 316; *Science*, 1920, **52**, 559).—Lithium has two isotopes with atomic weights 6 and 7, respectively, the relative proportions of which vary with the conditions. A suggested possible explanation is that the rates of evaporation of the two isotopes vary with the surface conditions of the metal. Magnesium has three isotopes with atomic weights 24, 25, and 26, the relative proportions being 7 : 1 : 1. These ratios give a mean atomic weight of 24.336.

A. A. E.

The Disappearance of Gas in the Electric Discharge. IV. THE RESEARCH STAFF OF THE GENERAL ELECTRIC CO., LONDON (*Phil. Mag.*, 1922, [vi], **43**, 914—937; cf. A., 1920, ii, 730; 1921, ii, 369, 533).—A further account of experiments on the disappearance of gas in the presence of phosphorus vapour. The decrease in pressure was measured by use of a lamp filament as the hot wire of a Pirani gauge and by determining whether the pressure is above or below the value at which the glow potential is equal to the applied potential. The amounts of hydrogen adsorbed increase with increase in the phosphorus introduced. A marked step in the adsorption, however, occurs for 0.09 to 0.27 mg. of phosphorus. Sodium fluoride and phosphorus together exert a greater effect than either substance singly; spluttered tungsten is also effective. The adsorption of gases takes place in two stages; the first is practically instantaneous and is impossible to control.

The solid films on the walls are possibly in an electrically polar condition, and it is suggested that gaseous ions enter into combination with the surface by virtue of their charges. The very small adsorption of inactive gases, which do not form negative ions, suggests that it is the negative ions which are adsorbed by this layer.

W. E. G.

The Potential of the Iodine Electrode and the Activity of the Iodide Ion at 25°. A. McKEOWN (*Trans. Faraday Soc.*, 1922, **17**, 517—521).—*E.M.F.* measurements have been made for cells of the type $\text{Pt} + \text{I}_2 | \text{KI solution sat. with I}_2 | \text{sat. KCl} | \text{N calomel electrode}$, the concentration of potassium iodide being varied between wide limits. From the results, the activities of the iodide-ion and the tri-iodide-ion in the various solutions have been calculated. It was found that the activities of both ions increased much less rapidly than their concentrations. The potential of the normal

(saturated) iodine electrode has been deduced in terms of the activity of the iodide-ion. The value found is +0.2454 volt, the normal calomel electrode being taken as zero. W. T.

The Effect of Telluric Acid on the Potential of a Silver Anode. FRANZ JIRSA (*Z. Elektrochem.*, 1922, 28, 186—191).—The author has determined the effect of halogen ions on the potential of a silver anode, and has investigated the nature of the depolarisation effected by telluric acid. In general, the halogens depolarise a silver anode in an alkaline electrolyte and reduce the value of the decomposition voltage. Apart from the fluorine-ion, which exerts no depolarising effect in an alkaline electrolyte, the maximum depolarisation is effected by the iodine-ion, the minimum by the chlorine-ion. Fluorine-ions accelerate the appearance and evolution of oxygen at the anode. When the polarisation of a silver anode in an alkaline electrolyte containing halogen ions is a maximum, the dioxide, Ag_2O_2 , alone is formed. The monoxide, Ag_2O , is produced when a silver anode is polarised to +1.40 volts in an alkaline electrolyte containing telluric acid. The following reactions occur: $\text{Ag}_2\text{O} + \text{K}_2\text{TeO}_4 + \text{H}_2\text{O} \rightleftharpoons \text{Ag}_2\text{TeO}_4 + 2\text{KOH}$; $2\text{Ag}_2\text{O}_2 + 2\text{K}_2\text{TeO}_4 + 2\text{H}_2\text{O} \rightleftharpoons 2\text{Ag}_2\text{TeO}_4 + 4\text{KOH} + \text{O}_2$. The potential of the electrode $\text{Pt}/\text{Ag}_2\text{O}_3$ in an alkaline electrolyte containing telluric acid is constant and equal to 1.57 volts. J. S. G. T.

Effect of Changing the Hydrogen-ion Concentration on the Potential of the Zinc Electrode. THEODORE W. RICHARDS and THEODORE DUNHAM, jun. (*J. Amer. Chem. Soc.*, 1922, 44, 678—683).—Measurements of the potential of zinc in acid solutions of zinc sulphate, of various concentrations, measured at 18° against the calomel electrode, through a sodium sulphate bridge, and also directly against a zinc electrode containing pure *N*-zinc sulphate solution, showed marked rise in potential as the acid concentration was increased. The magnitude of this rise in potential indicated that it must be due partly to the liquid junction potential, owing to the presence of the hydrogen-ion. Equal concentrations of the sulphate-ion in the form of alkali sulphates added to the zinc sulphate solution gave smaller elevations in the potential, which elevations were in the order of the transport numbers of the kations concerned. It is assumed that the change in liquid junction potential produced by sodium sulphate is small and therefore it may be inferred that about two-thirds of the change produced by sulphuric acid is due to solution junction potential and the remaining third to a real increase in the single electrode potential of zinc, due, presumably, to the depression of the electrolytic dissociation of zinc sulphate. J. F. S.

Variations in the Conductivity of Solid Electrolytes. P. VAILLANT (*J. Phys. Radium*, 1922, 3, 87—100; cf. A., 1912, ii, 419; 1920, ii, 722).—The changes in conductivity of solid electrolytes with temperature are due to two independently varying phenomena; the first of these occurs instantaneously, whilst the second is a slow change, frequently in an opposite direction to the first. These

progressive changes in the conductivity at constant temperature are not related to phosphorescence, being shown by many types of salts. The polarisation which occurs on passing the current is maintained indefinitely unless the crystals are short-circuited. The conductivity during the initial period is given by the relation : $\log C_T = a + bT$ or $\log C_T = \alpha - \beta/T$. In the second phase, the changes in conductivity follow different laws, being manifested by a series of oscillations of decreasing amplitude. W. E. G.

Abnormality of Strong Electrolytes and the Ionisation Theory of Ghosh. JAMES KENDALL (*J. Amer. Chem. Soc.*, 1922, **44**, 717—738).—The ionisation hypothesis put forward by Ghosh to explain the abnormality of strong electrolytes (T., 1918, **113**, 449, 627, 707, 790; 1920, **117**, 823, 1390) has been critically analysed and the following conclusions have been drawn. The second and third postulates are absolutely incompatible, for if the particles of a salt in solution possess a definite space-lattice arrangement, as in the crystalline state, then the combination of two oppositely charged particles to form a salt molecule or a completely saturated electrical doublet is impossible. Ghosh's calculation of the electrical work required to separate the component radicles of a gram-molecule of a dissolved salt is erroneous. The value actually derived, assuming a cubic space lattice arrangement, exceeds that obtained by Ghosh by 75%. The experimental conductivity data for potassium chloride solutions are not in agreement with the equation which Ghosh obtains from his postulates, or with various corrected forms of this equation. The agreement claimed by Ghosh to exist between his equations and the experimental results of previous investigators is largely fictitious. The remarkable coincidence of calculated and observed values in many tables is due to far too frequent errors of calculation and transcription. In other cases, the only legitimate deduction is that the equations tested serve very well as interpolation equations over a restricted range. No confirmation of the theory of Ghosh is afforded by the experimental data for (a) the variation of equivalent conductivity with dilution for salts in aqueous solution, (b) the temperature coefficient of the ratio μ_v/μ_∞ , (c) the electrical conductivity of non-aqueous solutions, (d) the molecular number i and the Clausius theorem, (e) the ionisation of strong acids, (f) the ionisation of weak acids, (g) the ionisation of transition acids, (h) the electrical conductivity of pure salts in the solid and fused states, (i) the conductivity of salts in mixtures of pyridine and water, (j) the ionisation of salts in solvents of low dielectric constant, (k) the partition of a salt between an ionising and a non-ionising solvent, and (l) the vapour pressure of the hydrogen haloids in aqueous solution. The rôle of the solvent in ionisation, which is ignored altogether by Ghosh, must be taken into account in any complete hypothesis of conducting solutions. In its present form, the ionisation hypothesis of Ghosh is unacceptable, and certain of the postulates on which it is based must be either modified or rejected.

J. F. S.

Ionic Equilibria of Strong Electrolytes. ALLEN E. STEARN (*J. Amer. Chem. Soc.*, 1922, **44**, 670—678).—The equivalent conductivities of the haloids of sodium and potassium of varying weight normal concentrations (0.1—4.0N) have been determined at 25°. A series of similar determinations has also been made for mixtures of a sodium and a potassium haloid. The observed values for the mixed salt solutions have been compared with those calculated from the values for the pure salt solutions at equivalent total concentrations on the basis of the isohydric principle. The differences between the calculated and observed values were found to be positive and to increase with the total salt concentration and with the numerical value of the ratio of sodium to potassium salt in the mixture, but to decrease generally from chloride to iodide. In general, the results are to be explained on the basis of complex formation, although at present no explanation is offered for the seeming decrease in complex formation on proceeding from chlorides to iodides. Tables of the specific gravities and the relative viscosities of all solutions measured are given in the paper. J. F. S.

Influence of the Solvent on Ionisation and the Accompanying Heat Effect. SIDNEY MAURICE NEALE (*Trans. Faraday Soc.*, 1922, **17**, 505—516).—In passing from pure water as solvent to 93.8% acetone, the ionisation constant of picric acid falls from about 0.2 to 0.002, but even in the latter case the deviation from Ostwald's dilution law still manifests itself, and, to a similar extent, the value of k decreasing by about 50% on increasing the dilution from 32 to 1024 litres. In anhydrous acetone, the ionisation constant for picric acid has fallen to the order of 10^{-6} , and Ostwald's dilution law is obeyed. This indicates that the anomaly of strong electrolytes is to be referred to some property of the system solute plus solvent, rather than to high degree of dissociation. The heat of ionisation of picric acid varies largely with the nature of the solvent, passing through a minimum at about 70% acetone. In the case of *p*-nitrobenzoic acid, the heat of ionisation is practically zero. In 44.35% acetone, the ionisation constant has fallen one-twentieth of its value in pure water, but the heat of ionisation is still zero. W. T.

The Hydrolysis of a Salt formed from a Weak Acid and a Weak Base. R. O. GRIFFITH (*Trans. Faraday Soc.*, 1922, **17**, 525—527).—A theoretical paper. The author points out that it is necessary to state more exactly what is meant by the "degree of hydrolysis" in the case of a salt of a weak acid and a weak base at great dilution. He shows in the case of aniline acetate that at a dilution of 5000 litres the concentration of undissociated acid is not equal to the concentration of undissociated base, that is, the degree of hydrolysis differs for each ion. The calculation also shows that the hydrogen-ion concentration, whilst practically constant down to $v=500$ (the limit of Tizard's measurements (*T.*, 1910, **97**, 2494)), should not be independent of the concentration between $v=500$ and $v=5000$, but should fall with increasing dilution. W. T.

[**Biochemical and Electrochemical Oxidation of Organic Compounds.**] A. NATHANSOHN (*Z. Elektrochem.*, 1922, **28**, 129—130).—A criticism of the views put forward and the interpretation of the author's work (*Koll. Chem. Beihefte*, 1919, **11**, 261) by Fichter in a paper under the same title (this vol., ii, 23).

J. F. S.

Simple Formula for the Calculation of the Specific Heats of Solids. H. J. KRASE (*J. Amer. Chem. Soc.*, 1922, **44**, 784—.786).—A simple form of the specific heat equation of Lewis and Gibson (A., 1918, ii, 29), $C_p = f(T/\theta)$ has been obtained which fits the experimental facts. This equation has the form $C_p = 2.91 + 2.89 \tanh 2.95 \log T/\theta$. Since the entropy equation is obtained by integration of the specific heat equation $S_p = \int (2.91 + 2.89 \tanh 2.95 \log T/\theta) d \log_e T/\theta$, and on evaluating the integration constant by putting the entropy equal to the Lewis-Gibson value 1.62 when $\log T/\theta = 0$, $S_p = 6.70 \log T/\theta + 5.18 \log \cosh 2.95 \log T/\theta + 1.62$.

J. F. S.

Polymerisation at the Critical Temperature. WILLIAM R. FIELDING (*Chem. News*, 1922, **124**, 182—184, 198—199, 215—217; cf. A., 1920, ii, 732; A., 1921, ii, 487).—Calculations are made of the polymerisation factor at the critical temperature for a large number of volatile organic and inorganic compounds.

W. E. G.

Cinnamic Acid in Cryoscopy. P. FALCIOLA (*Gazzetta*, 1922, **52**, i, 175—179).—The results furnished by anthracene, naphthalene, *o*-, *m*-, and *p*-nitrobenzaldehydes, and phenylpropionic and phenylpropionic acids differ somewhat from those obtained by Mathews (A., 1917, ii, 356) and show that the value of the cryoscopic constant for cinnamic acid is about 100. The fact that phenylpropionic acid, when dissolved in cinnamic acid, produces almost normal depressions of the freezing point confirms Bruni's conclusion (A., 1899, ii, 731; 1900, ii, 714) that there is no tendency to the formation of solid solutions when saturated compounds are dissolved in unsaturated ones, although such tendency is exhibited in the inverse case (cf. also Mascarelli and Pestalozza, A., 1907, ii, 936; 1908, i, 527).

T. H. P.

A Micro-method for the Determination of Molecular Weight in a Melting-point Apparatus. KARL RAST (*Ber.*, 1922, **55**, [B], 1051—1054).—The freezing point of camphor is depressed to such an unusual extent by the presence of dissolved substances that the effects can be measured with sufficient accuracy for the determination of molecular weight by means of an ordinary thermometer and with minute amounts of material.

The substance under investigation and the camphor are weighed into a small tube which is subsequently closed with a cork carrying a pointed needle. The substances are melted together, well mixed, allowed to solidify, and withdrawn by means of the needle. A portion of the mixture is transferred to a melting-point tube the

bottom of which must be semi-spherical; it is pressed firmly together, giving a compact column not exceeding 1 mm. in height. The tube is heated in the usual melting-point apparatus. The mixture rapidly assumes the appearance of melting ice and is later converted into a cloudy liquid in which the presence of a delicate crystalline skeleton can be sharply detected by means of a lens. As the temperature rises, the latter disappears from above downwards, and the point at which the last trace vanishes is recorded as the melting point. Since the readings required are differential, the use of normal thermometers and the correction for the exposed part of the thread are unnecessary. H. W.

Orientation of Molecules in Surfaces. VII. Vaporisation in Steps as Related to Surface Formation. WILLIAM D. HARKINS and LATHROP E. ROBERTS (*J. Amer. Chem. Soc.*, 1922, **44**, 653—670; cf. A., 1921, ii, 242).—An account is given of the amount of energy involved in the various steps in the process of vaporisation, based on the assumption that the surface through which the vaporisation occurs is not highly curved. The results are expressed in *micri-ergs* as unit which is defined as 10^{-14} ergs. The relationships found at a corresponding temperature of 0.7° are tabulated for oxygen, nitrogen, ethyl ether, ethyl acetate, carbon tetrachloride, benzene, chlorobenzene, methyl alcohol, and ethyl alcohol, and the data indicate that, at a definite corresponding temperature, in the case of non-associated liquids the molecules of which are symmetrical, the molecular values for the latent heat of surface formation, l , the total surface energy, e , the energy of thermal emission, j , and the internal heat of vaporisation, λ , are nearly proportional to the critical temperatures of the liquids. The same relation seems to hold for the free surface energy, γ , provided that the temperature range is not too great. Thus the free surface energy of ethyl ether at a corresponding temperature of 0.7 is 4.0 as calculated from the value for carbon tetrachloride, and 3.9 as calculated from the value for chlorobenzene, whilst the experimental value is 4.0 . This statement, as applied to the latent heat of vaporisation alone, is somewhat similar to Trouton's law, which is known to be not entirely exact. Since the principle expressed above is much more general in its application, it is to be expected that it will prove to be somewhat less exact. The energy values for ethyl ether are lower than those for carbon tetrachloride, and this is a consequence of the lower critical temperature of the ether. The effect of a lack of symmetry in the molecule, especially when marked, is to lower the molecular free surface energy, latent heat of surface formation, and total surface energy and to increase the energy of thermal emission. Values are calculated for ethyl alcohol from the critical temperature on the assumption of a symmetrical molecule, using the values of carbon tetrachloride as a basis, and from the actual results it is evident that the molecular free surface energy, the total surface energy, and more markedly the latent heat of surface formation are considerably lowered by the dissymmetry of the molecule. The most striking

effect is, however, the very great increase in the energy of thermal emission. The symmetry considered is that of the electro-magnetic forces round the molecule rather than a symmetry with respect to mass. The substitution of the slightly polar chlorine atom for hydrogen in benzene gives almost exactly the calculated value for a symmetrical molecule except in the case of the thermal emission, which is considerably increased, since it is the most sensitive of all the quantities to changes of molecular symmetry. Since e is decreased and j increased by increasing dissymmetry of the molecule, the ratio e/j serves as a remarkably sensitive index of molecular symmetry. The ratio e/λ , which is equal to $e/(e+j)$, varies in the same way, but not to so large an extent. According to Stefan's law, the ratio of the total energy necessary to carry a molecule from the interior of a liquid into the surface to its total heat or energy of vaporisation, e/λ , is equal to $1/2$. This is shown not to be the case. Not only is this an increasing function with increasing symmetry of the molecule, but also with increasing corresponding temperature. Its value appears to approach unity as the corresponding temperature approaches unity. Thus a molecule in the surface at a high corresponding temperature is, in terms of relative energy, very much more nearly in the vapour phase than when the corresponding temperature is low. The relationships noted above are just those indicated by the hypothesis that molecules in the surface are oriented, the orientation increasing with increasing dissymmetry, and decreasing with increasing thermal agitation. The effect of thermal agitation is illustrated in the case of the alcohols; compounds of the polar-non-polar type. It is shown for these compounds that the molecular surface energy increases with the temperature. The effect of increased agitation is to overcome the orientation partly, and to throw the more polar groups into the outer surface, thus increasing the surface energy.

J. F. S.

Vapour Pressure of Sulphur Dioxide and Ammonia. F. W. BERGSTROM (*J. Physical Chem.*, 1922, **26**, 358—376).—The vapour pressure of ammonia and sulphur dioxide has been determined at temperatures from slightly above the normal boiling point to the freezing point. An improved form of apparatus is described which is capable of giving vapour pressure measurements below 900 mm. at low temperatures. The measurements are compared with those of Burrell and Robertson (*A.*, 1916, i, 6; ii, 81) and of Brill (*A.*, 1906, 847), and it is found that the author's results are in keeping with the Ramsay-Young relationship, whilst those of Brill and of Burrell are not. It has been shown experimentally that Burrell's vapour pressure measurements for ammonia are erroneous. The triple point of ammonia has been found to lie at -77.9° at a pressure of 45.5 mm. The value found by Cragoe, Meyers, and Taylor (*A.*, 1920, ii, 370) is -77.70 at 44.9 mm.

J. F. S.

Binary Liquid Mixtures. O. FAUST (*Z. physikal. Chem.*, 1922, **101**, 94—103).—A theoretical paper in which, for the purpose

of comparing the relative efficiency and accuracy of the static and dynamic methods of determining the vapour pressure of liquid mixtures, the author has recalculated the results obtained by Schmidt (this vol., ii, 119), Beckmann and Faust (A., 1915, ii, 143), and von Zawidzki (A., 1901, ii, 6) for the vapour pressure of mixtures of acetone and chloroform. Curves are constructed and the shape of the curves discussed in connexion with Dolezalek's theory of liquid mixtures. J. F. S.

Vapour Pressures of Binary Mixtures. H. CASSEL (*Z. physikal. Chem.*, 1922, **101**, 104—108).—A theoretical paper in which the author criticises mathematically the assertion of Schmidt (this vol., ii, 119) that Dolezalek's theory of liquid mixtures cannot be maintained. The author shows that the measurements of Schmidt for mixtures of benzene and toluene show divergences from a straight line curve which vary between 1.0 and 2.4%. It is shown that the simplicity of Dolezalek's theory in comparison with the complicated expressions of van der Waals and Jahn has justified itself in numerous cases, so that its heuristic significance as a usable working hypothesis cannot be denied (cf. preceding abstract). J. F. S.

Binary Liquid Mixtures. ALFRED SCHULZE (*Z. physikal. Chem.*, 1922, **101**, 109—116).—A theoretical paper in which the author criticises the assertions of Schmidt (this vol., ii, 119) and shows that his statements in connexion with Dolezalek's theory of liquid mixtures cannot be maintained. Further, the results on which Schmidt bases his assertions are a confirmation of the theory (cf. preceding abstracts). J. F. S.

A Continuous Flow Calorimeter, and the Determination of the Heat of Neutralisation of a Solution of Hydrochloric Acid by One of Sodium Hydroxide. FREDERICK G. KEYES, LOUIS J. GILLESPIE, and SHINROKU MITSUKURI (*J. Amer. Chem. Soc.*, 1922, **44**, 707—717).—A continuous flow calorimeter has been so developed that measurement of the rise of temperature for ten different rates of flow leads to a value of the heat of neutralisation, which is associated with an error, as calculated by the method of least squares, which is less than 0.1% of the heat change itself. It is shown that a few slight modifications in the calorimeter will probably increase the precision. For the heat of neutralisation resulting in the formation of $\text{NaCl}, 521\text{H}_2\text{O}$ (about 0.107*N*-sodium chloride) at 32.3°, the value 13280 cal./mol. is found. In the calculation, the specific heat data of Richards and Rowe were used, so that the unit is the 18° calorie (equivalent to 4.18 joules). It is held that this value is accurate within about 0.3%, although it is 2.6% higher than the corresponding point on Wörmann's curve for the same neutralisation (*Ann. Physik*, 1905, [iv], **18**, 775). The present value leads to a value for the heat of ionisation which is about 0.5% lower than the corresponding point on Noyes, Kato, and Sosman's curve (A., 1910, ii, 257). In the opinion of the authors, the 1% difference in the heat change when potassium

hydroxide is substituted for sodium hydroxide in the neutralisation is questionable, because no correction was applied for the carbonate present.
J. F. S.

Heats of Neutralisation of Potassium, Sodium, and Lithium Hydroxides with Hydrochloric, Hydrobromic, Hydriodic, and Nitric Acids at various Dilutions. THEODORE W. RICHARDS and ALLAN W. ROWE (*J. Amer. Chem. Soc.*, 1922, **44**, 684—707).—The heat of neutralisation of the four acids by the three bases mentioned in the title have been determined at a series of dilutions. It is shown that the most serious errors in work of this kind are due to irregularity of mixing, presence of carbonates in the hydroxides, and inadequate avoidance of loss of heat by cooling. The solutions neutralised were molecular quantities in 100 gram-molecules of water, and they were combined in all possible pairs over two ranges of temperature not far apart, and the results were interpolated to exactly 20°. The values of the heat of neutralisation obtained ranged from 137500 to 14085, sodium hydroxide giving the lowest values among the bases and hydriodic acid among the acids. With the help of the heats of dilution, of factors and products, previously determined, the heats of neutralisation at other concentrations have been determined, and these have been extrapolated in two ways through a short range to infinite dilution. The heat of formation of water from its ions at 20° is found by this short extrapolation to be probably not greater than 13·69 Cal._{20°} or 57·22 kilojoules and possibly not smaller than 13·62 Cal._{20°} or 56·93 kilojoules.

J. F. S.

Pressure Volume Relation of Superheated Liquids. K. L. WISMER (*J. Physical Chem.*, 1922, **26**, 301—315).—A number of experiments are described, designed to obtain actual *PV* curves at high temperatures to see whether the curve shows any tendency to bend more sharply as the limit of superheating is approached, and to find what degree of superheating may be attained. Experiments were made with ethyl ether, ethyl chloride, and isopentane, which were subjected to a pressure of 20 atmospheres, heated to various temperatures, and the pressure then reduced to 1 atmosphere. The time before explosion occurred was then noted. In this way, the highest temperature to which the liquids may be superheated at ordinary pressures was obtained. The values found are ethyl ether 143°, isopentane 136°, and ethyl chloride 126°. The minimum pressures which can be reached with liquid ether at temperatures above 143° increase with the temperature, the points lying approximately on a straight line joining the critical point with the —72 atmospheres' point of J. Meyer ("Kenntnis des negativen Druckes in Flüssigkeiten," 1911). The pressure-volume relations have been determined experimentally for liquid ethyl ether at 121·5°, 127·9°, and 133·8°, and for liquid ethyl chloride at 99·6°, 109·7°, and 117·4°, at pressures down to 1 atmosphere. The pressure-volume relation under these conditions is almost linear. There is no tendency shown by the *PV* curves to bend more sharply at low pressures as

the limit of superheating is reached. The actual experiments showed a marked departure from van der Waals's curve.

J. F. S.

The Corresponding States : The Halogenated Derivatives of Benzene. MAURICE PRUD'HOMME (*Bull. Soc. chim.*, 1922, [iv], 31, 295—299).—In the system of measurements of relativity constituted by the reduced values of the temperature, pressure, and density, the values obtained for the four monohalogenated derivatives of benzene are identical. The properties of the halogens are thus completely masked in such compounds.

For this series of compounds, the relation $M = AD_cT_c/P_c$ holds good independently of the halogen present. M is the molecular weight and A is a constant of value 21.65 for this series. W. G.

The Determination of Surface Tension from the Maximum Pressure in Bubbles. SAMUEL SUGDEN (*T.*, 1922, 121, 858—866).

Method of Calculating Fluidity, Surface Tension, and Reaction (Equilibrium) Pressure. HENRY JERMAIN MAUDE CREIGHTON (*J. Franklin Inst.*, 1922, 193, 647—655).—A theoretical paper in which it is shown that by means of a formula similar to the Ramsay and Young vapour-pressure equation, it is possible to calculate the fluidity (viscosity) of liquids, the equilibrium pressure of chemical reactions, and the surface tension of liquids. The expression has the form $R' = R + c(T_B - T_B')$, where R and R' are the ratios of the absolute temperatures at which two substances have the same value for any of the properties mentioned, T_B' and T_B the temperatures of one of the substances corresponding with the two values of the property chosen, and c is a small constant. The equations have been tested and shown to be in very good agreement with facts. J. F. S.

The Calculation of Adsorption of Vapours at Different Temperatures. L. BERÉNYI (*Z. angew. Chem.*, 1922, 35, 237—238).—On the basis of Polányi's theory of adsorption, a formula has been evolved, by which the equation of state of the adsorbed substance within certain limits of pressure may be approximately calculated. The results for a range of temperatures which extends, for substances boiling between 170° abs. and 400° abs., from 0.06 to 1.4 times the boiling point in degrees abs. show no greater error than that of experimental determination. The formula applies to the reversible and physical adsorption of chemically homogeneous vapours, alone, or from a mixture of comparatively inert gases. From the observed data, p_1 , or the pressure of the gas, and x_1 , the quantity adsorbed at a temperature T_1 , the amount of gas adsorbed at the temperature T_2 , or x_2 , may be calculated by the formula : $x_2 = x_1 \times (1 + F_2 - F_1/100)$. $F_2 - F_1 = F$ is a correction factor which takes into account the thermal expansion and the compression of the adsorbed layer. The means by which the value F is determined, with relation to absolute temperature and the variation of the vapour from the law of gases, is given. H. M.

Electro-adsorption as a Purely Chemical Process. WOLFGANG OSTWALD (*Kolloid Z.*, 1922, 30, 254—260).—A theoretical paper in which the author discusses and criticises a paper under the same title by Kolthoff (this vol., ii, 197). It is shown that chemical adsorption, that is, the concentration of a dissolved substance on the surface in consequence of the chemical energy potential existing there, is quite possible. Therefore the hypothesis put forward by Kolthoff (*loc. cit.*) is not a purely chemical hypothesis, but a combination of chemical and electrical hypotheses. The deduction of the well-known adsorption formula by means of the law of mass action leads to concrete practically applicable formulæ only when instead of the stoichiometric weight concentration of the adsorbing substance the actual reacting mass of this substance is employed. The active quantity corresponds with the sum of the chemically reacting surface layers of the adsorbing substance. The stoichiometric weight concentration must therefore be multiplied by a factor which connects the actual reacting mass with the total mass. The demand of Kolthoff's hypothesis, that the value of the index must be the same for equivalent adsorbents whilst for electrolytes of different valencies the values stand in a simple relationship to one another, is not in keeping with experimental results. The cases quoted by Kolthoff are accidental or exceptions. In other cases, the hypothesis is in no way confirmed.

J. F. S.

Diffusion in Porous Vessels. A. L. HERRERA (*Mem. rev. soc. cient. "Antonio Alzate,"* 1921, 39, 343—347).—When solutions of reactive salts are allowed to diffuse very slowly together, the precipitates or crystals formed show analogies to natural cells. A solution of potassium silicate, d 1.5 or more, containing potassium carbonate not in excess of normality, is placed inside an unglazed porcelain cup, and the latter placed in calcium chloride of d 1.0068. After twenty-four hours, numerous very fine tubes of calcium silicate are formed on the outside of the porous cup, bearing on their surface microscopic cells, and in some cases are formed entirely of the latter.

CHEMICAL ABSTRACTS.

An Application of the Optical Method of Determining the Solubility of One Liquid in Another. C. CHÉNEVEAU (*Compt. rend.*, 1922, 174, 1019—1021; cf. this vol., ii, 355).—In the case of inorganic or organic salts which are without action on and insoluble in aniline, when their aqueous solutions are in contact with aniline the ratio of the lowering of the refractive index of the aniline to the quantity of water dissolved is constant and the same for equimolecular solutions. The lowering of the refractive index by unit mass is the same for all salts which fulfil the above conditions.

W. G.

Le Chatelier's "New" Geometrical Representation. ERNST JÄNECKE (*Chem. Ztg.*, 1922, 46, 361).—A claim for priority against Le Chatelier with regard to the method of geometrical representation of solutions of reciprocal pairs of salts (cf. Jänecke, A., 1906, ii, 833; Le Chatelier, A., 1921, ii, 248).

H. W.

The Partial Miscibility of Liquid Couples. Applications to Reactions between Dissolved Substances. RENÉ DUBRISAY (*Ann. Chim.*, 1922, [ix], 17, 222—256).—A more detailed account of work already published (*A.*, 1919, i, 73; ii, 78; 1920, ii, 508, 756; 1921, i, 535; ii, 282, 344). W. G.

Sizes of Atoms in Crystals. ROBERT N. PEASE (*J. Amer. Chem. Soc.*, 1922, 44, 769—774).—A theoretical paper in which, as an alternative to Bragg's view that the distances between atoms in crystals can be expressed in terms of atomic radii characteristic only of the element concerned, it is suggested that the particular element is of secondary importance for elements the atoms of which are of the same rare-gas type, and that the contribution of an atom to the distance between it and another atom in a crystal depends on the number and arrangement of electrons about its positive nucleus, on the type of lattice at least as far as this affects the latter, and, ordinarily, on the magnitude of the charge carried by the atom. The interatomic distances in substances crystallising in the diamond type of lattice have been analysed from this point of view. The following values are recorded: diamond 1.54 (1.54), silicon 2.30 (2.35), grey tin 2.80 (2.80), silicon carbide 1.92 (1.95), zinc sulphide 2.41 (2.35), cuprous chloride 2.41 (2.43), cuprous bromide 2.52 (2.57), cuprous iodide 2.66 (2.78), and silver iodide 2.80 (3.18). The values in brackets are due to Bragg, and the present values generally agree better with the observed values than do those of Bragg. J. F. S.

Transition from Crystalloid to Colloid Properties within Homologous Series. EDGAR J. WITZEMANN (*J. Physical Chem.*, 1922, 26, 201—216).—A theoretical paper in which an attempt is made to answer the question "How do such systems as constitute living organisms come into existence?" It is shown that the emulsoid structure of biocolloids (lipoids, carbohydrates, and proteins) constitutes a gross heterogeneity which is also present in a much less conspicuous form in the lower crystalloid members of the same homologous series. In this way, it follows that the particular properties of biocolloids arise from the cumulative effect of certain groups in the molecule. This effect of chemical constitution may be easily traced with soaps, but less easily with the carbohydrates and proteins. Associated with the material heterogeneity there is a well-known heterogeneity of energy. It is shown that such heterogeneities may arise spontaneously on releasing one system into another. The best known energy differentiation under these conditions is the surface energy. It appears that the surface energy phenomena in solutions are energy differentiations definitely associated with material differentiations and these in turn are due to chemical constitution, and that changes in the latter necessarily involve the whole series of effects. In connexion with biological phenomena, these two heterogeneities have been regarded as associated with a vital force, and even now they are frequently thought to be maintained in organisms by a constant expenditure of energy obtained from chemical reactions in the

organism. It is now shown that certainly part of the heterogeneity occurring in the structure of organisms arises from a predisposition on the part of the component substances themselves to give rise to these effects. Such colloidal systems as those prepared from simple biocolloids are self-reproducing from the components; they arise spontaneously and require no unusual outlay of energy for their construction and maintenance. It is shown that the quantitative statement of Le Chatelier's principle may be easily discovered in the phenomena of true and colloidal solutions so fundamentally involved in biological phenomena. J. F. S.

Cataphoresis of Colloidal Carbon. STEPHANIE GOLDBERG (*Kolloid Z.*, 1922, 30, 230—234).—Colloidal carbon prepared by Sabbatani's method (A., 1914, i, 357; ii, 198) has been thoroughly dialysed to remove sulphuric acid and the influence of various factors on the rate of cataphoresis examined. The sols examined had a specific conductivity at 18° of 22.5×10^{-6} reciprocal ohms, and had various migration velocities toward the anode from 18.3×10^{-5} to 13.5×10^{-5} cm./volt sec. Dilution of the solution increases the velocity slightly to a maximum value, whilst filtration also causes an increase from 18.8×10^{-5} to 22.2×10^{-5} cm./volt sec. The addition of acids, irrespective of their nature, causes a decrease in the velocity; thus 0.001*M*-sulphuric acid decreases the velocity from 13×10^{-5} to 11×10^{-5} , and 0.002*M* reduces it to 8×10^{-5} cm./volt sec. Bases have a similar action, and in one case with ammonia the velocity was reduced from 19×10^{-5} to 13×10^{-5} cm./volt sec. Acids and bases change the colour of colloidal carbon, but the original colour may be regained by neutralisation. In the case of salts, only those of aluminium have any action on the velocity of migration, and in this case the velocity increases to a maximum with 1/500000*M*-aluminium sulphate and then falls with greater concentrations. The addition of colloidal ferric hydroxide generally coagulates the colloidal carbon, but if so much colloidal ferric hydroxide is added that no coagulation takes place, then the velocity is reduced about 20%. Dyes, such as crystal-violet, auramine, and methylene-blue, in small concentrations, reduce the velocity to zero and with increasing concentrations then increase it. J. F. S.

Ionic Synergism. II. Investigation of Mastic Sols. L. MICHAELIS and N. HIRABAYASHI (*Kolloid Z.*, 1922, 30, 209—215; cf. A., 1921, ii, 682).—The limiting concentrations of hydrochloric, acetic, sulphuric, nitric, and sulphasalicylic acids, by which precipitation of mastic sols may be brought about, have been determined and the P_H value of the solutions has been measured. The effect of adding lithium chloride, rubidium chloride, sodium carbonate, sodium citrate, calcium chloride, sodium chloride, and potassium sulphate to mastic sols in the presence of measured concentrations of hydrogen-ions has also been investigated. The results show that the anion is not entirely without influence on the precipitation of electro-negative mastic sols; certain anions (sulphasalicylic acid) are strongly antagonistic to the kations, but

the more common anions have so small a specific action that as a first approximation they may be neglected in the investigation of the synergism of the hydrogen-ion and the metallic kations. The results therefore furnish an approximate confirmation of the rules established in the case of congo-rubin (*loc. cit.*) both for the univalent alkali-ions and for the bivalent calcium-ion. From the results, it follows that the relationship of the activity of a uni- and a bi-valent ion cannot be given in absolute terms, but only with reference to the P_H value, and only with increasing value of P_H does the relationship approach an absolute limiting value. Changes in the concentration of the hydrogen- and hydroxyl-ions are only without effect on the condition of a mastic sol when P_H is greater than about seven or eight.

J. F. S.

Soaps and Proteins. I. Colloidal Chemistry of Soaps and the Manufacture of Soap. MARTIN H. FISCHER, GEORGE D. McLAUGHLIN, and MARIAN O. HOOKER (*Koll. Chem. Beihefte*, 1922, 15, 3—102).—A general account of the preparation of soaps, and a consideration, with experimental results of the systems soap-water, soap-alcohol and soap- X , where X is one of a long series of organic liquids. Definitions of a number of terms in their colloid-chemical significance are given; these include hysteresis, swelling, liquefaction, power of gelatinisation, and solvation, syneresis, and sol. The behaviour of soaps towards indicators and many other properties of soaps are considered. Reproductions of twenty-nine photographs are included in the paper, illustrating the behaviour and properties treated.

J. F. S.

Non-, Uni-, and Bi-variant Equilibria. XXI. F. A. H. SCHREINEMAKERS (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 23, 1151—1160; cf. A., 1920, ii, 361).—A theoretical paper in which the author considers the equilibria of n components in n phases, where the quantity of one of the phases approaches zero. The influence of a new substance on an invariant P or T equilibrium is discussed.

J. F. S.

Heterogeneous Equilibria: The Ternary System Sodium Sulphate-Sodium Carbonate-Water. ALFRED ERNEST DAWKINS (*T.*, 1922, 122, 776—781).

Nature of Secondary Valency. V. Partition Coefficients in Systems containing Water as one Component, with Special Reference to the Absolute Values of the Series Constants. HOMER W. SMITH (*J. Physical Chem.*, 1922, 26, 256—271; cf. this vol., ii, 270).—A continuation of previous work in which the series constants of the partition of organic acids and bases between water and various organic liquids have been derived from data largely taken from the literature. It is shown that in each of these systems the series constants for the various dissolved substances are related to each other in a simple proportion. The series constants for twelve systems in which water is one component have been tentatively formulated by the study of diethylamine. In all twenty of the systems studied, the series constants

for the various systems are related to each other in simple proportion. In every system in which water is one component, the amines differ from the acids in the value of their series constants by a constant amount. It is concluded that the intermolecular forces concerned in the solvent powers of various molecular species differ in intensity by discrete and simply-related amounts.

J. F. S.

Nature of Secondary Valency. VI. HOMER W. SMITH (*J. Physical Chem.*, 1922, **26**, 349—357; cf. A., 1921, i, 324; ii, 315; this vol., ii, 270).—The results of the preceding papers are summarised and discussed. It is shown from the examination of twelve hundred determinations of the partition coefficient in twenty-one systems of two immiscible liquids that all may be covered by an equation of the form $100P = V_m/60.00 \pm a$, where V_m is the molecular volume at the boiling point and a is a constant. It is found that a limited number of values of a are required to cover the behaviour of all substances in any one system. All substances covered by any one value of a in any system have been termed a series, and the value of a a series constant. The various values of a in any one system are related to one another in simple proportion.

J. F. S.

Kinetics of Reactions in Heterogeneous Solutions. The Reduction and Oxidation Actions of Alternating Currents. OSCAR COLLENBERG and SVEN BODFORSS (*Z. physikal. Chem.*, 1922, **101**, 117—149).—The action of metallic iron, copper, zinc, tin, and cadmium on a 4% solution of ferric alum has been investigated by studying the rate of solution of the metals. The results show that the solution of the metals takes place in accordance with Nernst's hypothesis, and the velocity of solution is in keeping with the laws of unimolecular reactions in heterogeneous systems. Apart from zinc, the reaction constant for the individual metals, despite their electrochemical dissimilarity, are very similar. The similarity becomes more apparent when the quantities of the metals dissolved in a given time are compared in terms of equivalents. Using the above metals as electrodes, the electrolysis of ferric alum has been carried out by means of alternating currents of various periodicity; it is shown that the current brings about no fundamental difference in the course of the reaction beyond an increase in the total yield. The influence of the periodicity is very small.

J. F. S.

Thermal Dissociation of Ammonia with Special Reference to Coke Oven Conditions. G. E. FOXWELL (*J. Soc. Chem. Ind.*, 1922, **41**T, 114—125).—The thermal decomposition of ammonia diluted with coal gas in silica tubes packed with porous material such as coke or firebrick has been investigated at temperatures between 520° and 850°. It is shown that when an excess solid surface is present the decomposition is bimolecular. The velocity of decomposition, when the tube is filled with coke, increases slowly with increasing temperature, the temperature coefficient being

1.136. The value of the velocity constant, k , may be expressed by the Arrhenius formula $k_2 = k_1 e^{[A(T_1 - 1/T_2)]}$, where A is 13300. For tubes with a radius less than 0.5 cm., $k \propto 1/r^2$, but it is probable that for wider tubes $k \propto 1/r$. When gases containing ammonia are passed through porous material, the amount of decomposition in unit time varies directly as the square of the radius of the pores; hence it is the size of the pores rather than the total porosity that is of importance. Silica bricks have considerably less decomposing effect than siliceous (80% SiO_2) bricks, whilst the presence of iron is very deleterious. The composition of the ash of the coke has an important bearing on the rate of decomposition. Rutile and orthoclase have very little, if any, action; iron pyrites is converted into ferrous sulphide during carbonisation, and this increases the velocity of decomposition to some extent. Iron oxide, which is converted into metallic iron, enormously increases the rate of decomposition and the same is true, although to a less extent, of lime. The value of k for coke is found to be 0.00200 at 755° .
J. F. S.

Decomposition of Nitrous Acid. ALFONS KLEMENC and FRIEDRICH POLLAK (*Z. physikal. Chem.*, 1922, **101**, 150—171).—The velocity of decomposition of nitrous acid in dilute acid solutions (acetic, nitric, sulphuric) has been determined. The reaction mixture was stirred continuously by a stream of nitrogen to remove the nitric oxide from the solution. It is shown that the velocity of decomposition, according to the equation $3\text{HNO}_2 = \text{HNO}_3 + 2\text{NO} + \text{H}_2\text{O}$, as measured by the decreasing permanganate titre, depends on the velocity with which the nitric oxide is removed from the solution. This process is determinative of the velocity. The velocity also depends on the pressure of nitric oxide above the solution. It is assumed that undissociated nitrous acid possesses a considerable pressure of nitric oxide and in consequence the velocity of decomposition is proportional to the concentration of undissociated nitrous acid. This fact has been confirmed under the most widely varying conditions. On the basis of the "invasion" and "evasion" coefficients for carbonic acid measured by Bohr, it has been shown that the concentration of free nitric oxide in aqueous solution may be regarded as approximately the same as that of the undissociated nitrous acid. When nitrogen is led through a solution of nitrous acid the velocity of decomposition is increased, and the velocity constant increases with the increasing velocity at which the nitrogen is passed through. The velocity of the direct decomposition of nitrous acid cannot be determined by the method adopted, and is in all probability immeasurably rapid. Nitrous acid decomposes spontaneously into nitric oxide and the hydroxyl radicle and its transitory existence in aqueous solution is due to a mutual action between it and the solvent. The temperature coefficient of the reaction is in keeping with the quantities which define the solubility of a gas. The presence of mineral acids increases the velocity of decomposition.
J. F. S.

Thermal Decomposition of Nitrogen Pentoxide in Solution.

ROGER H. LUECK (*J. Amer. Chem. Soc.*, 1922, **44**, 757—769).—The velocity of decomposition of nitrogen pentoxide in carbon tetrachloride and in chloroform solutions has been determined gasometrically at temperatures between 25° and 55°. It is shown that such systems are of special value in considering the newer hypotheses concerning the relationships of radiant energy to chemical action. The fact that the velocity constants obtained in solution so closely approximate to those obtained in the gas phase by Daniels and Johnston (*A.*, 1921, ii, 249) is of particular importance because of the wide variation of conditions under which the measurement has been made. The experimental facts show the energy of activation or the critical increment to be practically independent of the temperature. It has also been shown that, in accordance with a necessary deduction from the Perrin-Lewis radiation theory, a change in reaction velocity brought about by temperature effects or the influence of solvent diminishes as the magnitude of the velocity constant increases. A comparison has been made between the values of the velocity constant experimentally obtained and those calculated from the critical increment, by means of the equation recently put forward by Dushman (*A.*, 1921, ii, 315), and although the agreement is not exact, it is far better than can be obtained by means of the older equations developed by Trautz and Lewis. The nitrogen peroxide formed in the reaction has been shown to exert an autocatalytic effect on the thermal decomposition of the nitrogen pentoxide.

J. F. S.

The Velocity of Absorption of Carbon Dioxide by Alkaline Solutions.

PAUL RIOU (*Compt. rend.*, 1922, **174**, 1017—1019).—A study of the absorption of carbon dioxide by solutions of sodium carbonate. For a given area of absorbing surface the velocity of absorption of carbon dioxide diminishes as the concentration of the sodium carbonate increases. It is also diminished by the presence of increasing amounts of sodium chloride or hydrogen carbonate. The velocity of absorption is markedly increased by agitation or rise in temperature of the liquid. It is diminished by diluting the carbon dioxide with air.

W. G.

The Speed of Sulphonation of Phenols. I. The Effect of Temperature and the Methyl Group.

ARTHUR FRED CAMPBELL (*T.*, 1922, **121**, 847—857).

Formation and Hydrolysis of Lactones. II. ANTON KAILAN and EMANUEL FRANZ NEUMANN (*Z. physikal. Chem.*, 1922, **101**, 63—93; cf. *A.*, 1920, ii, 606).—The velocity of lactone formation in aqueous and ethyl-alcoholic solutions in the case of hydroxybutyric acid and hydroxyvaleric acid has been determined both in the absence of catalysts and in the presence of hydrochloric, hydriodic, sulphuric, chloroacetic, and *n*-butyric acids. All experiments were carried out at 25°, and in some cases an addition of potassium chloride or sodium chloroacetate was made to the

reaction mixtures. The rate of hydrolysis of butyrolactone and valerolactone in the presence of hydrochloric acid has also been determined. It is shown that the autocatalysis of the lactone formation in aqueous solution is practically proportional to the hydrogen-ion concentration alone, and that consequently the results of Henry (A., 1892, ii, 1303), which differ from this result, must be attributed to the presence of the silver salt of the hydroxy-acid in his solutions. Lactone formation in the presence of chloroacetic acid also depends on the concentration of the hydrogen-ion. When hydrochloric acid is used as catalyst in concentrations up to $N/5$, the velocity of lactone formation is proportional to the total concentration of this acid, and if the view be held that the undissociated hydrochloric acid exerts a specific catalytic action, then in this case it must be the same as that of the hydrogen-ion ($k_m = k_H$). With larger concentrations of hydrochloric acid, the velocity of reaction increases much more rapidly than the hydrogen-ion concentration, but more slowly than the total acid concentration, so that in these concentrations the action of the undissociated hydrochloric acid molecule must be decidedly less than that of the hydrogen-ion. This difficulty is removed if the hydrogen-ion is regarded as the only independent catalyst in this reaction, and the divergences from the proportionality between concentration and velocity as due to an increase in the activity of the hydrogen-ion brought about by the addition of hydrochloric acid exactly as is the case on the addition of neutral salts. The increase in the velocity of reaction which occurs on the addition of potassium chloride to hydrochloric acid is not to be attributed to a driving back of the dissociation alone, as was assumed by Taylor and Close (A., 1917, ii, 253), and consequently the deduction of the ratio k_m/k_H from this is not permissible. No connexion could be found between the velocity of reaction and Noyes's activity constants. The velocity of lactone formation in 99% alcohol, by volume, in the case of γ -hydroxyvaleric acid, both in the presence and absence of hydrochloric acid or other catalysts, takes place much more rapidly than in water. Water exerts a greater retardation on the velocity of lactone formation by hydrochloric acid than it does in the case of esterification by the same acid. The gradual addition of water causes the velocity of lactone formation to fall gradually to a point below that obtained in pure water, after which it gradually rises to that obtained in pure water. In the autocatalytic lactone formation in 99% alcohol, only a part of the observed velocity is to be explained by the concentration of the hydrogen-ions.

J. F. S.

The Alkaline Hydrolysis of the Esters of the Two Ethylenedicarboxylic Acids. ANTON SKRABAL and EMMY RAITH (*Monatsh.*, 1921, 42, 245—249).—The study of the hydrolysis of tartaric esters (A., 1920, ii, 239) and succinic esters and their homologues (A., 1921, ii, 34) has been extended to the investigation of the alkaline hydrolysis of the methyl esters of fumaric and maleic acids. The experiments were conducted at 25° in aqueous

solution containing a molecular proportion each of sodium carbonate and sodium hydrogen carbonate, the dilution being necessarily high on account of the low solubility in water of both esters. Referred to $[\text{OH}'] = 1$, the constants for the two stages of the hydrolysis are, for the fumaric ester, $k_1 = 414$, $k_2 = 19$, ratio $n = 22$; for the maleic ester, $k_1 = 48$, $k_2 = 0.72$, $n = 67$. Comparing these values with the first and second dissociation constants of the acids, which are: fumaric acid 1.0×10^{-3} and 3.2×10^{-5} ; maleic acid, 1.4×10^{-2} and 2.6×10^{-7} , fumaric acid falls into line with the saturated, straight chain dicarboxylic acids in that the acid strength and rate of hydrolysis of the esters are symbatic. Maleic acid, however, is abnormal, the first hydrolysis constant being unexpectedly low compared with the high value of the first dissociation constant. The observation may have an important bearing on the question whether there may not be a structural as well as a spatial difference between the two acids. E. H. R.

Kinetics of the Reaction between Ammonia and Aromatic Aldehydes. FRIEDRICH DOBLER (*Z. physikal. Chem.*, 1922, 101, 1—33).—The velocity of the reaction between ammonia and alcoholic solutions of benzaldehyde, *p*-tolualdehyde, *m*-xylyl-aldehyde, anisaldehyde, cinnamaldehyde, *p*-chlorobenzaldehyde, and *o*-, *m*-, and *p*-nitrobenzaldehydes has been investigated. The method of investigation depends on the fact that ammonia may be estimated in the presence of hydrobenzamide by titration with hydrochloric acid, using hæmatoxylin as indicator. The formation of hydrobenzamide at 20° is shown to be strictly bimolecular, and the corresponding hydramides formed from the other aldehydes are also in most cases formed according to the bimolecular formula. In the case of benzaldehyde and *p*-tolualdehyde, the dependence of the velocity constant on temperature has been determined; it is found that k is nearly doubled by raising the temperature 10° and trebled by raising the temperature 20°, thus: for benzaldehyde, $k_{20} = 0.233$, $k_{30} = 0.400$ and $k_{40} = 0.627$, the two latter values are not very constant. It is found that, on the one hand, an equilibrium is set up between aldehyde and ammonia, and, on the other, between hydramide and water. The influence of temperature on the equilibrium has been investigated, and it is shown that with increasing temperature the equilibrium is displaced so that less hydramide is formed; thus in the case of benzaldehyde at 20°, 48.4% of the ammonia has been converted into hydramide, at 30°, 43.8%; 40°, 36.3% and 50°, 23.2%. An addition of 0.01*N*-benzoic acid practically doubles the reaction velocity in the case of benzaldehyde; small additions of ammonium chloride effect a large increase in the velocity of hydramide formation, in one case this amounted to an increase in the k value from 0.233 to 2.325. Ammonium sulphate has no effect on the specific reaction velocity, whilst sodium hydroxide reduces the k value from 0.233 to 0.028. These changes in velocity are shown to be due to the same cause, namely, changes in the dissociation of ammonium hydroxide. A decrease in the hydroxyl-ion con-

centration increases the specific reaction velocity, and an increase decreases the velocity. Double bonds and chlorine in the para-position in the nucleus of the aldehyde increase the reaction velocity respectively six and three times. The introduction of a methyl group in the para-position, for example, *p*-tolualdehyde, increases the velocity, whilst a second methyl group in the ortho-position, *m*-xylyl-aldehyde, reduces the value somewhat, but it is still greater than that of the original benzaldehyde. The methoxyl group in anisaldehyde reduces the velocity to about one-half. The substitution of a nitro-group in the meta-position increases the velocity, but in the ortho- and para-positions it reduces the velocity, the para-substitution causing the greater reduction. The ionic concentration of the reaction mixtures does not change during the hydramide formation at 0° and 20°, as is shown by the constant value of the electrical conductivity.

J. F. S.

Catalysts and Chemical Equilibrium. J. CLARENS (*Bull. Soc. chim.*, 1922, [iv], **31**, 299—307).—From a study of the action of glass wool and cupric chloride respectively as catalysts of the reversible reaction $4\text{HCl} + \text{O}_2 \rightleftharpoons 2\text{H}_2\text{O} + 2\text{Cl}_2$, it is shown that the action of the catalyst is different for each of the inverse reactions which determine the equilibrium. In general, it is considered that there is no reason to suppose that a catalyst modifies equally the velocity of two inverse reactions.

W. G.

The Influencing of the Catalysis of Hydrogen Peroxide with Platinum by Röntgen Rays. ROBERT SCHWARZ and WALTER FRIEDRICH [with H. WUNNERLICH] (*Ber.*, 1922, **55**, [B], 1040—1051).—The catalytic decomposition of hydrogen peroxide in the presence of colloidal platinum is retarded by Röntgen rays, the effect being observed when the mixture of the substances is exposed to the radiation or when the platinum sol is treated in this manner separately and added shortly afterwards to the hydrogen peroxide; the latter alone does not undergo an appreciable change under the influence of Röntgen rays. The maximal value of the retardation is 77% (calculated on the percentage of hydrogen peroxide) and the highest values are observed in the sixth and seventh hours of the experiments. The cause of the phenomenon does not appear to lie in coagulation, diminution in the degree of dispersivity, or alteration in the charge of the particles, as is shown by special experiments, and is rendered improbable by the further observation that the illuminated sol recovers its activity completely within about sixteen hours. It appears probable, therefore, that the platinum sol suffers a reversible change at its surface which must be attributed to the presence of water. Since the action of the Röntgen rays is accompanied by an emission of β -rays from the platinum particles which are able to decompose water, it is possible that an oxide of platinum is formed or that hydrogen is adsorbed by the sol. If, however, it is assumed with Haber that the platinum catalysis of hydrogen peroxide is conditioned by the presence of dissolved or combined oxygen in the catalyst, the formation of an oxide would in all probability accelerate the

change. It is therefore more likely that the first stage of the catalysis is delayed because the platinum charged with hydrogen is not immediately able to form the intermediate product, Pt_2O_y . The first stage of the action therefore consists in the combustion of the dissolved hydrogen by oxygen derived from the peroxide after which catalysis can take place in the sense of Haber's theory. The recovery of the platinum sol within sixteen hours is attributed to the gradual decomposition of the platinum hydrogen alloy, and this hypothesis is supported by the known complete instability of the product at 100° . It is also possible that the removal of hydrogen is effected by dissolved atmospheric oxygen, but direct experiments in the absence of oxygen do not lend support to this view, although their interpretation is hampered by the extreme difficulty of excluding oxygen completely in such observations. Pure water is not altered by exposure to Röntgen rays in such a manner as to give rise to contact poisons.

Catalase, also, is altered in such a manner by Röntgen rays that its power to catalyse the decomposition of hydrogen peroxide is lessened; as with the platinum sol, the enzyme recovers its activity completely after several hours. A satisfactory explanation of the phenomena cannot yet be given. H. W.

The Catalytic Activity of Copper. W. G. PALMER (*Proc. Roy. Soc.*, 1922, [A], 101, 175—186; cf. A., 1920, ii, 609; 1921, ii, 542).—This communication describes the effect on the dehydrogenating activity of copper of the addition of magnesium oxide, ferric oxide, zinc oxide, manganous oxide, and sodium carbonate. The presence of small amounts of sodium carbonate completely destroys the activity. The addition of 1% or less of an oxide reduces the activity considerably, but with higher concentrations, owing to the separation of two phases, an increase in the catalytic effect is observed. The pure oxides are usually less active than pure copper, and hence must be deficient in the density of radiation of about $1.0\ \mu$ wave-length (cf. previous papers). They, however, adsorb water and probably alcohol to a greater extent than metals, but this increase may be counteracted by the opacity of the oxide film to radiation from the copper core. Increased adsorption thus increases the activity, provided the oxide films are transparent to the radiation of the main catalyst.

W. E. G.

A Defined Organic Catalyst with a Hydrogen-ion Optimum. ERIK WIDMARK and CARL ALEX JEPSSON (*Skand. Arch. Physiol.*, 1922, 42, 43—61; from *Physiol. Abstr.*, 1922, 7, 6; cf. A., 1921, ii, 183).—The kinetics of the catalysis of acetoacetic acid into acetone and carbon dioxide is investigated in detail. It is found to have a reaction optimum at $P_H\ 4.09$, and this corresponds exactly with the optimum reaction calculated on the assumption that only undissociated molecules of catalyst and of substrate enter into the reaction. For this calculation from the dissociation constants, the original must be consulted. The conclusion is drawn that only the undissociated aniline molecule is effective,

and acts only on the undissociated molecule of acetoacetic acid. The conclusion cannot be generalised: for example, alanine has its optimum at P_H 8, at which reaction acetoacetic acid is almost completely ionised. The suggestive comparison with the behaviour of enzymes is indicated. W. O. K.

The Bôhr Atom. The Lagrange Theorem applied to Electronic Orbits. MARCEL BRILLOUIN (*J. Phys. Radium*, 1922, 3, 65—73).—A mathematical paper emphasising the exclusively spectroscopic character of Bôhr's theory. W. E. G.



Atoms and Molecules. II. ALBERT C. CREHORE (*Phil. Mag.*, [vi], 43, 886—914; cf. A., 1921, ii, 632).—A theory of the structure of atoms is presented which is in conformity with the Saha form of the electromagnetic theory. In the models given, the negative electron is assumed to be an oblate spheroid with a ratio of axes about three to one and the hydrogen atom consists of a positive charge of two units with very small dimensions situated in between two negative electrons. A structure is suggested for nitrogen and oxygen, which indicates that the heavier products of disintegration found by Rutherford may possess a mass 12 and a charge of 5.


It is also sought to compute the known distances between the atoms forming a molecule. The shape of the negative electron is a most important factor in determining the equilibrium distance, and hence it is a universal distance. Only electrostatic forces are considered, and it is shown that equilibrium is possible between the attractive and repulsive forces of the atoms and electrons when the eccentricity of the electron lies between the limits 0.9 and 1.0. This agrees with the value found in the previous paper. It is shown that helium and neon atoms will not form compounds and must be monatomic. W. E. G.

Unsymmetrical Addition to the Double Bond. I. A Theory of the Reaction Mechanism of the Direct Union. GRANVILLE A. PERKINS (*Philippine J. Sci.*, 1921, 19, 645—660; cf. this vol., A., ii, 138).—A theory of the mechanism of chemical reaction based on the Langmuir octet theory of valence. Diagrams are given showing the relationships between the forces of attraction and repulsion for two types of reaction, (a) a direct addition reaction, where an interpenetration of the outer shells of electrons occurs, and (b) a dissociation addition reaction where only secondary valence comes into play. These two types are illustrated by double bond reactions. W. E. G.

The Dissociation of Hexaphenylethane from the Point of View of the Octet Theory of Valence. HOWARD IRVING COLE (*Philippine J. Sci.*, 1921, 19, 681—690).—A historical survey of the previous investigations on the dissociation of hexaphenylethane is given and the Lewis-Langmuir octet theory of valence applied to this problem on the lines suggested by Perkins (this vol., A., ii, 138). On this theory hexaphenylethane cannot possibly dissociate without a rearrangement of one or more of the phenyl

groups. Symmetrical dissociation occurs in non-ionising solvents such as benzene, giving the quinoid modification as in Gomberg's

formula, $(C_6H_5)_2=C=C$ , which contains a para-carbon atom in a tervalent condition. In an ionising solvent such as sulphur dioxide, which contains unshared electrons, the dissociation occurs unsymmetrically. An unstable union with a solvent molecule takes place, which supplies two electrons to the shell of one of the carbon atoms in a phenyl group. In this manner is obtained a colourless negative ion, $(C_6H_5)_3\equiv C^-$, and a coloured positive ion, $(C_6H_5)_2=C=C$ ; the latter exists only in combination

with a solvent molecule. These two ions can combine to give $(C_6H_5)_3\equiv C$  $=C=(C_6H_5)_2$. These modifications are essentially

the same as those of Gomberg, except that, on this view, colourless triphenylmethyl can exist only in the form of a negative ion. The physical and chemical properties appear to support this theory. The oxidation of triphenylmethyl to form a peroxide is explained by the formation of intermediate positive and negative ions. An oxygen molecule is added to the positively charged ion, which then combines with the negative ion to give the peroxide.

W. E. G.

A Proposed Laboratory Test of the Theory of Relativity.

HAROLD S. KING (*Nature*, 1922, 109, 582—583).—Since it is impossible to conceive of the annihilation of mass, such as appears in the synthesis of an atom of helium from four hydrogen atoms, it must be supposed that mass is electromagnetic in origin, loss of mass being ascribed to overlapping of fields. According to the theory of relativity, 1 gram of matter is equivalent to 9×10^{20} ergs, or 2.1×10^{13} calories; moreover, 1 gram of radium in disintegrating to lead evolves 3.7×10^9 cal. Thus, it is possible to calculate the atomic weight of radium (226.038) by addition of 1 gram atom of radium-*G* (206.000), 5 gram atoms of helium (20.000), 4 gram electrons (0.0005) [?], and 3.7×10^9 cal. (0.038). If Hönigschmid's value (225.97) for the atomic weight of radium is accepted, it would therefore appear that radium-*G* has an atomic weight less than 206 (the lowest experimental figures being 206.08 and 206.05, due respectively to Richards and Hönigschmid), or that energy is derived from outside sources as suggested by Perrin, or that the equation connecting mass and energy is not correct. It is therefore suggested that an extension of the method of positive ray analysis to the determination of the atomic weights of radium, radium-*G*, and helium to an accuracy of 1 part in 10,000 not only would be of intrinsic value, but also would show whether the relation of mass and energy, based on the theory of relativity, holds.

A. A. E.

Inorganic Chemistry.

A Simple Hydrogen Generator for Use in making Hydrogen-ion Determinations. P. H. CATHCART (*J. Ind. Eng. Chem.*, 1922, **14**, 278).—The hydrogen is generated electrolytically in a bell jar inverted in a cylinder containing 10% sodium hydroxide. The outer vessel consists of a large glass precipitating jar, the hydrogen reservoir being a large inverted percolator with its mouth resting on a support about 1 inch from the bottom of the precipitating jar. A rubber stopper in the top of the hydrogen reservoir carries the cathode (which consists of an iron disk suspended from an iron wire) and a glass tube with stop-cock. The inner vessel is held down by four rubber-covered wires. The anode is similar to the cathode, of about 4 sq. in. area, and so adjusted in the annular space between the two vessels that it just touches the liquid when all gas has been withdrawn from the reservoir. The cathode just projects below the lower rim of the reservoir. A 110 volt D.C. circuit was used for generating the hydrogen, with a switch so arranged that the current is automatically cut off when the reservoir becomes full. A diagram of the apparatus is given. H. C. R.

Rectilinear Diameter of Hydrogen. E. MATHIAS, C. A. CROMMELIN, and H. KAMERLINGH ONNES (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, **23**, 1175—1184; cf. A., 1921, ii, 256).—The density of liquid and gaseous hydrogen has been determined in grams/c.c. at temperatures from -240.57° to -249.89° . The critical density is calculated from the results and the value 0.03102 obtained, a figure which is considerably smaller than the corresponding figure for any other substance. The equation of the diameter of the form $y=a+bt$ is considered, and the values of the coefficient are evaluated from the experimental data. The values obtained are $a=-0.063510$ and $b=-0.00039402$. The value of the slope of the diameter b is the smallest observed for any substance.

J. F. S.

The Crystal Structure of Ice. (SIR) W. H. BRAGG (*Proc. Physical Soc.*, 1922, **34**, 98—103).—This is derived independently of direct X-ray analysis on certain suppositions connected with the low density of ice. St. John (*Proc. Nat. Acad. Sci.*, 1918, **4**, 193) and Dennison (*Physical Rev.*, 1921, **17**, 20) have applied the methods of X-ray analysis and both refer the structure to a lattice composed of four right triangular prisms but disagree as to the dimensions. In the present paper it is assumed that ice belongs to that class of crystals in which the molecules are broken up into positive and negative ions, the hydrogens having given up their valency electrons to the oxygens. The density of ice is so low that the most economical spacing is adopted; the atoms are arranged in a hexagonal lattice with each oxygen surrounded by four hydrogens and each hydrogen by two oxygens. The distance between the

centres of two oxygen atoms is 2.76, the distance between consecutive basal planes is 3.67, and the distance between two atoms in the same plane is 4.52. The structure is extremely empty and a loose arrangement of water molecules would occupy less space. The hydrogen atom has apparently a larger diameter than the oxygen atom if oxygen has the value 1.30. These results agree exactly with the X-ray analysis of Dennison and the conclusions are supported by a comparison between the calculated and observed intensities of reflection.

W. E. G.

The Variations in the Chemical Composition of Sea-water and the Evaluation of the Saline Content. GABRIEL BERTRAND, FREUNDLER, and (Mlle) MÉNAGER (*Compt. rend.*, 1922, **174**, 1251—1253).—The authors have determined the total halogens and the calcium and magnesium in the water from the Atlantic ocean and the Mediterranean sea. The halogens being calculated as chlorine, the ratio of chlorine to calcium and magnesium respectively in the two samples of water was by no means the same. Thus the method of evaluating the saline content of sea-water from the density or chlorine content by means of Knudsen's tables can only be considered as at the best an approximation.

W. G.


The Dissociation of Chlorine. H. VON WARTENBERG and F. A. HENGLEIN (*Ber.*, 1922, **55**, [B], 1003—1006).—Difficulties are experienced in attempting to measure the dissociation of active gases such as chlorine by observation of the abnormal increase of pressure when they are heated owing to the impossibility of finding a completely resistant material for the construction of the apparatus. Since, however, the dissociation of binary compounds is dependent on the total pressure, it is possible to effect such measurements at temperatures which silica vessels will withstand if an extremely low pressure is used. In this manner, preliminary measurements of the dissociation of chlorine at 10^{-3} mm. pressure have been made. The apparatus consists of a quartz bulb surrounded by a quartz mantle, the space between the two being continuously evacuated. The bulb is connected with a 2 mm. wide quartz capillary which leads to a quartz thread manometer and mercury seal cooled to -30° and thence to a Volmer pump. The chlorine is generated by heating auric chloride in a glass capillary which is subsequently sealed off. The bulb is surrounded by platinum foil and heated in a Heraeus furnace, the temperature measurements being accurate to $\pm 2^{\circ}$. The temperature interval ranges from 985° to 1151° absolute.

It is certain that the heat of dissociation of chlorine is considerably lower than the value 106,000 cal. usually adopted and for $T=0$ is about 70,000 cal.

H. W.

The Structure of Chlorine Dioxide and Related Compounds. GRANVILLE A. PERKINS (*Philippine J. Sci.*, 1921, **19**, 729—740; cf. this vol., ii, 138).—A discussion of the constitutional formulæ of some of the oxides of chlorine and nitrogen. In the Langmuir theory no place is given to bonds consisting of

one, three, or five electrons. The suggestion is made that in exceptional cases an odd number of electrons may form a chemical bond. In chlorine dioxide, five electrons are shared between the three atoms as in the formula :



On this view chlorine dioxide contains a three electron bond. An odd electron is also shared in nitric oxide and nitrogen peroxide. The electron binding strength of atoms and its variation with the covalence is discussed.

W. E. G.

Hydroxylamine. I. Simple Method of Preparation of Free Hydroxylamine. HANS LECHER and JOSEF HOFMANN (*Ber.*, 1922, 55, [B], 912—919).—The method adopted is essentially a modification of that of Lobry de Bruyn, its most important feature being the avoidance of the final distillation of the product.

The apparatus consists of a Brühl receiver in which the reaction vessel is placed. The lower tubulus (generally used for evacuation) is provided with a calcium chloride tube whilst the upper tubulus is fitted with a stopper carrying two dropping funnels, the ends of which are placed above the reaction vessel. The tubulus of the lid serves for the introduction of a very efficient stirrer, which is provided with the customary mercury seal. The finely divided hydroxylamine hydrochloride is suspended in absolute ethyl alcohol in the reaction vessel and a small quantity of solid phenolphthalein is added. A solution of sodium ethoxide in absolute ethyl alcohol is introduced slowly from one of the dropping funnels into the well-stirred suspension, the rate of addition being so controlled that the red coloration which occurs as each drop enters the suspension disappears instantaneously. Should ultimately a pale pink coloration persist, it is removed by cautious addition of a solution of hydroxylamine hydrochloride in ethyl alcohol from the second dropping funnel. The precipitated sodium chloride is removed and the filtrate allowed to remain undisturbed in a freezing mixture of ice and salt until it attains a temperature of -18° . The large crystals of hydroxylamine thus produced are filtered, washed with absolute ether, and brought into a desiccator which is immediately exhausted. In this manner, there is no loss of hydroxylamine by decomposition during the preparation, but about 5.4% of it is retained by the precipitated sodium chloride. The yield of solid hydroxylamine is about 40% of that theoretically possible, but the remainder can be readily recovered as the hydrochloride from the alcoholic mother-liquor. The product contains about 97% of hydroxylamine, the remainder being water. It has m. p. $32-35^{\circ}$, d_{4}^{10} 1.335, d_{4}^{14} 1.334. It is distinguished from Lobry de Bruyn's purest specimens by its greater instability; after twenty-four hours it smells of ammonia and after four days it is completely liquefied and contains only 42.55% of hydroxylamine. Hydroxylamine crystallises in the rhombic system; the crystals obtained from ethyl alcohol and by the solidification of the molten substance are identical.

H. W.

Behaviour of Carbon at High Temperatures. F. SAUERWALD (*Z. Elektrochem.*, 1922, **28**, 183—185).—Continuing the work of Ryschkewitsch (A., 1921, ii, 258, 586, 696) and Münch (A., 1921, ii, 586), the author heated rods of carbon and graphite to various temperatures, attaining finally the temperature of the positive crater of the arc. The heating was effected by passing a current through the rods in an atmosphere of hydrogen. The ends of the rods were copper-plated. Small globules appeared on the surface of the Acheson graphite employed, and these are attributed to the distillation of impurities from the hotter interior of the rod to the surface. This was confirmed by a determination of the respective ash contents of the rod and of the globules. The conclusion is not in agreement with that of Ryschkewitsch, who found similar globules to be composed of pure graphite. J. S. G. T.

Behaviour of Carbon at High Temperatures. EUGEN RYSCHKEWITSCH (*Z. Elektrochem.*, 1922, **28**, 185—186).—The author contends that the experimental conditions employed by Sauerwald (see preceding abstract), differ so materially from his own, that the results obtained in the two cases are not comparable. He remarks more particularly on the different atmospheres and current densities employed in the two cases, and on the possibility of impurities being derived, in Sauerwald's experiments, from the heated metallic connexions employed. J. S. G. T.

Preferential Catalytic Combustion of Carbon Monoxide in Hydrogen. ARTHUR B. LAMB, CHARLES C. SCALIONE, and GRAHAM EDGAR (*J. Amer. Chem. Soc.*, 1922, **44**, 738—757).—With the object of finding a method of removing small quantities of carbon monoxide from hydrogen before its use in the synthetic production of ammonia, a number of experiments have been carried out on the preferential combustion of carbon monoxide in mixtures containing air, hydrogen, carbon monoxide, and carbon dioxide by means of a catalyst termed "hopcalite," which is a mixture of metallic oxides and in the present experiments was composed of 60% of manganese dioxide and 40% of cupric oxide. It is shown that "hopcalite," which oxidises carbon monoxide in air rapidly and catalytically at the ordinary temperature, does not affect hydrogen in air under the same conditions, and that by its means carbon monoxide can be completely and continuously removed from a dry mixture containing, for example, 0.5% of carbon monoxide, 89.50% of hydrogen, and 10% of air without any noticeable action on the hydrogen. A similar mixture containing 1% of carbon monoxide, on the other hand, liberates so much heat that under these conditions oxidation of the hydrogen sets in, and the catalyst is soon heated to incandescence and is destroyed. It is also shown that moist mixtures of carbon monoxide and air and of hydrogen and air require much higher temperatures for oxidation. In the presence of water vapour at a pressure of 50 mm. of mercury, carbon monoxide in a 0.5% mixture with air is completely oxidised at a temperature of 90° or lower. Hydrogen, on the other hand, mixed with air, does not begin to be oxidised under these conditions

until a temperature of 120—125° is reached. It is possible, therefore, to remove carbon monoxide completely and continuously from a mixture having, for example, the composition 0·5% of carbon monoxide, 0·5% of oxygen, 24·75% of nitrogen, and 74·25% of hydrogen and containing water vapour at a pressure of 50 mm. of mercury without any considerable oxidation of hydrogen. On the other hand, in a similar 1% carbon monoxide mixture, the hydrogen soon takes fire and the catalyst becomes incandescent. The effect of space velocity, carbon dioxide, hydrogen sulphide, and the concentration of carbon monoxide on the efficiency of the catalyst have been investigated. Formulæ for the temperature rise in the catalyst, and the factor limiting the permissible concentration of carbon monoxide have been derived and verified at different space velocities in tubes of various diameters, and on the basis of these results the proper conditions for the technical application of this method of removing carbon monoxide from hydrogen have been outlined. The present results are in agreement with those of Rideal (T., 1919, 115, 993). J. F. S.

Preparation of Bright Metallic Sodium and Potassium, or their Alloys, in Nitrogen. G. BORNEMANN (*Z. angew. Chem.*, 1922, 35, 227).—A glass tube of 15—20 mm. diameter is drawn out into a short constriction of 3—4 mm. diameter about 20 cm. from one end which is sealed up. A fine wire gauze thimble is fitted tightly into the tube above the constriction and the required amount of metal introduced into the open end of the tube, which is then sealed. The tube is then placed in a horizontal position and the metal melted, whereby the oxygen, moisture, and carbon dioxide are removed, leaving an atmosphere of nitrogen. On now placing the tube vertically and carefully remelting the metal it flows through the gauze and the constriction into the lower portion of the tube as perfectly bright untarnished metal, all impurities remaining behind on the gauze filter. The tube is then drawn off and sealed at the constriction. G. F. M.

Sodium Silicate. A. ERDENBRECHER (*Mikrokosmos*, 1921, 15, 55—60).—The addition of not more than 40 grams of sodium hydroxide to 100 c.c. of the solution obtained when 20—25 grams of the salt $\text{Na}_2\text{SiO}_3 \cdot 9\text{—}10\text{H}_2\text{O}$ are dissolved in 40 c.c. of water yields the hydrate $\text{Na}_2\text{SiO}_3 \cdot 9\text{H}_2\text{O}$, rhombic, m. p. 47°. Sixty grams of sodium hydroxide similarly yield the hydrate $\text{Na}_2\text{SiO}_3 \cdot 6\text{H}_2\text{O}$, monoclinic, m. p. 63·5°, whilst larger amounts yield the hydrate $\text{Na}_2\text{SiO}_3 \cdot 4\text{H}_2\text{O}$, hexagonal, m. p. 83—85°. Other hydrates of sodium metasilicate mentioned in the literature are probably mixtures of these, as changes in the composition of the mother-liquor cause changes in the degree of hydration.

CHEMICAL ABSTRACTS.

Preparation of Ammonium Chloride. P. MONDAIN MONVAL (*Compt. rend.*, 1922, 174, 1014—1017).—A study of the conditions governing the crystallisation of ammonium chloride from solution at 15° when in solution alone or in the presence of one or more of

the salts, sodium chloride, sodium carbonate, and ammonium carbonate. A Le Chatelier diagram showing the surfaces of saturation is given. Commercially in the ammonia-soda process the salt is not crystallised out from the liquid from which the sodium hydrogen carbonate has been extracted, but ammonia is first added to convert the hydrogen carbonates into normal carbonates. W. G.

The Melting Point of Normal Ammonium Sulphate. R. KATTWINKEL (*Ber.*, 1922, 55, [B], 874).—Contrary to the statement of Caspar (A., 1920, ii, 431), a definite melting point cannot be assigned to normal ammonium sulphate; with increasing temperature it suffers loss of ammonia, decomposition being complete at 365°. H. W.

Conditions of Formation and Stability of Ammonium Carbamate. C. MATIGNON and M. FRÉJACQUES (*Bull. Soc. chim.*, 1922, [iv], 31, 307—316).—A more detailed account of work already published (A., 1920, ii, 250). W. G.

Liquid Crystals of Calcium Phosphate. P. GAUBERT (*Compt. rend.*, 1922, 174, 1115—1117).—When the rhombohedra obtained by crushing calcite are ignited, they retain their shape and if then they are submitted to the action of phosphoric acid certain optical phenomena are observed due to the formation of liquid or soft crystals of calcium phosphate. These are described in detail. W. G.

Action of Nitrogen on Mixtures of Barium Oxide and Carbon at High Temperatures. PAUL ASKENASY and FRITHJOF GRUDE (*Z. Elektrochem.*, 1922, 28, 130—151).—The rate of combination of compressed mixtures of barium carbonate and wood charcoal or graphite with nitrogen has been investigated up to 1600°. It is shown that the reaction proceeds with a practicable velocity at 1400°, and in fifteen minutes 60% of the barium has been converted into a mixture of cyanide and cyanamide when an excess of nitrogen is present. In agreement with Caro and in opposition to Ewan and Napier, the primary reaction is the formation of barium carbide, and from this much cyanide is formed which then undergoes a secondary reaction to form barium cyanamide. The formation of cyanide at comparatively low temperatures (1300—1400°) reaches a maximum of 65% in fifteen to thirty minutes, which is not increased at 1600°. With increasing temperature, the formation of the cyanamide rapidly increases. There is no evidence that a basic cyanide is first formed, nor is there any evidence of a sublimation of barium cyanide at the lower temperatures. The reproducibility of the combination of nitrogen is best achieved in the presence of an excess of carbon. When there is a shortage of carbon it is the cyanamide formation, and not that of the cyanide, which is effected. It is shown that the reaction proceeds more rapidly when compressed materials are used. Impurities in the carbon, such as silica or alumina, retard the reaction, whilst iron, in opposition to previous statements, accelerates it, but disturbs the reproducibility. A mixture of barium oxide and carbon takes up the same amount of

nitrogen as an equivalent quantity of 55% barium carbide, and the relative amounts of barium cyanide and cyanamide formed are the same in both cases. A very long list of references on this subject, with comments, precedes the paper. J. F. S.

Electro-deposition of Lead from Mathers's Perchlorate Bath. I. Structure of the Deposit. W. E. HUGHES (*J. Physical Chem.*, 1922, **26**, 316—323).—The author has made a macroscopic and microscopic examination of the structure of lead deposited from Mathers's perchlorate bath (U.S. Pat. 931944, 1909; A., 1911, ii, 113). The specimens were etched by using them as anodes in the electrolysis of 30 c.c. of 30% perchloric acid in 270 c.c. of water for various periods with a current density of 9—10 amperes/sq. dm. and a voltage of 1.5. The etched surface was much worn, cindery, and dull and light grey in colour with no crystalline appearance. Where the outside surface of the lead was directed toward the cathode during etching, a thin, skin-like substance is formed, which after a while detaches itself and comes away in flakes; no such action occurs when the inside surface is directed to the cathode. The term inside surface indicates the lead surface which was in contact with the cathode during the deposition. When examined microscopically (125 diameters) the deposit is seen to consist of irregular cells with lustrous, slightly yellow walls. The interior of each cell appeared to be made up of a mosaic of small bright and dark particles. The author deduces from the observations that the structure of the lead is ordered, and not amorphous; the size of the grain is reduced by the colloid (peptone) contained in the bath and that the colloid separates from solution with the lead, not in any haphazard fashion, but in continuous layers in the deposit. J. F. S.

Amalgams. III. Colloidal Copper Amalgam. C. PAAL and HERMANN STEYER (*Kolloid Z.*, 1922, **30**, 215—228; cf. A., 1919, ii, 69, 516).—Hydrosols of copper amalgam may be prepared by (a) shaking copper hydrosol with mercury, (b) keeping copper hydrosol in contact with mercury without shaking, (c) mixing either red or blue copper hydrosol with mercury hydrosol. The preparations were all effected in an atmosphere of hydrogen, and the amalgam sols possess colours which differ markedly from those of the original copper sols. In the first preparation, sols of an amalgam represented by the formula $\text{CuHg}_{0.02}$ may be obtained, in the second preparation the composition of the amalgam is given by the formula $\text{CuHg}_{0.34}$, and in the third preparation, using the red copper hydrosol, the amalgam has the composition $\text{CuHg}_{1.35}$. The copper amalgam sols are less stable and more easily oxidised than the gold amalgam hydrosols previously described (*loc. cit.*), and require a larger concentration of protecting colloid (sodium lysalbate or sodium protalbate) to stabilise them than the corresponding gold amalgam sols. J. F. S.

Copper Sulphide. W. GLUUD (*Ber.*, 1922, **55**, [B], 952—953).—The oxidation of copper sulphide by air at the atmospheric

pressure proceeds rapidly in ammoniacal suspension with the production of a mixture of sulphate and thiosulphate; cuprous sulphide is similarly but more slowly oxidised. In neutral or acid solution, the change is slower and necessitates the use of compressed air at temperatures up to 160° ; copper sulphate is formed. In certain circumstances the sulphur is deposited in the elementary state; the chief conditions are that the copper solution should not be precipitated completely, that the oxidation should be effected immediately, and that the solution should contain, in addition to ammonia, considerable amounts of dissolved salts, preferably ammonium compounds. H. W.

Mechanism of the Dehydration of Crystalline Aluminium Hydroxide and of the Adsorption of Water by the Resulting Alumina. LOWELL H. MULLIGAN (*J. Physical Chem.*, 1922, 26, 247—255).—Crystalline aluminium hydroxide in air at atmospheric pressure and humidity is not affected by temperatures below 145° , and remains constant in composition as the normal hydroxide, $\text{Al}(\text{OH})_3$, up to this temperature. The decomposition and evolution of water commences just above 145° , and as far as the evolution of the chemically combined water of the hydroxide is concerned, is practically complete at 200° . Slight irregularities are observed in the drying curve between 145° and 196° , which are attributed to the very slow rate at which equilibrium is reached at these particular temperatures and the resultant experimental errors. All the water is not driven off at 200° , but an amount equal to about 8% of the original aluminium hydroxide is retained at this temperature, and is driven off slowly as the ignition temperature is increased, very much higher temperatures being required for complete dehydration. Above 200° , the curve has the general form of an adsorption curve, and the 8% of water, which is retained at 200° , appears to be practically adsorbed by the highly porous aluminium oxide. There is no evidence of the existence of hydrates formed by step-wise dehydration. When alumina, produced by drying aluminium hydroxide at temperatures as low as 275° , is allowed to take up water, this water is simply adsorbed, and does not recombine chemically with the alumina. This is established by the fact that the drying curve after "rehydration" bears no resemblance to the original dehydration curve of aluminium hydroxide. The higher the ignition temperature at which alumina is prepared the smaller is the amount of water which it is capable of adsorbing under a given set of conditions.

[With W. J. MEAD.]—The original aluminium hydroxide, the alumina prepared by its ignition, and the alumina after "rehydration" have been subjected to an X-ray examination. It is shown that the X-ray pattern produced by the original crystalline aluminium hydroxide is identical with that of the mineral gibbsite, which is a definitely crystalline chemical compound. Alumina prepared by calcining the crystalline hydroxide at 325° gives no trace of the hydroxide structure, neither is it similar to the mineral diaspore, $\text{Al}_2\text{O}_3 \cdot \text{H}_2\text{O}$, nor to corundum, Al_2O_3 . It shows a set of

lines which indicate a crystalline structure. The adsorption of water does not alter its structure. Alumina prepared by calcination at 600° shows a faint pattern similar to the preceding specimen, but the bulk of the material is probably amorphous. When calcination temperatures somewhat above 1000° are employed, the product gives a faint corundum pattern, and still higher temperatures increase the intensity of this pattern until it becomes that of pure corundum.

J. F. S.

Mechanism of the Reduction of Permanganate and its Physico-chemical Basis. I. Reaction between [Potassium] Permanganate and Formic Acid in Slightly Acid Solution.

JOSEF HOLLUTA (*Z. physikal. Chem.*, 1922, **101**, 34—53).—The velocity of the reduction of potassium permanganate by formic acid in faintly acid solutions has been investigated at 22°, 25.1°, 24.7°, and 15.3°. The results are discussed in connexion with previously published work of Schilov (A., 1903, ii, 720) and of Skrabal (A., 1905, ii, 17, 18; 1906, ii, 658), and it is shown that the reduction takes place in the following stages: (1) $2(\text{MnO}_4' + \text{HCO}_2' = \text{MnO}_4''' + \text{H}^* + \text{CO}_2)$ (measurably slow); (2) $2\text{MnO}_4''' + 8\text{H}^* + 5\text{HCO}_2' = \text{MnO}_4' + \text{Mn}(\text{HCO}_2)_5'' + 4\text{H}_2\text{O}$ (instantaneous); (3) $\text{Mn}(\text{HCO}_2)_5'' = \text{Mn}^{***} + 5\text{HCO}_2'$ (equilibrium); (4) $2\text{Mn}^{***} + \text{HCO}_2' = 2\text{Mn}^{**} + \text{H}^* + \text{CO}_2$ (more rapid than 1); (5) $2(2\text{Mn}^{***} + 4\text{H}_2\text{O} = \text{Mn}^{**} + \text{Mn}(\text{OH})_4 + 4\text{H}^*)$ (immeasurably fast); (6) $\text{MnO}_4' + 4\text{Mn}^{**} + 8\text{H}^* = 5\text{Mn}^{***} + 4\text{H}_2\text{O}$ (immeasurably fast); (7) $2(\text{Mn}[\text{OH}]_4 + 4\text{H}^* = \text{Mn}^{***} + 4\text{H}_2\text{O})$ (equilibrium), and (8) $2\text{Mn}^{***} + \text{HCO}_2' = \text{Mn}^{**} + \text{H}^* + \text{CO}_2$ (very slow). These partial equations give a total equation for the reaction which has the form $2\text{MnO}_4' + 11\text{H}^* + 5\text{HCO}_2' = 2\text{Mn}^{**} + 5\text{CO}_2 + 8\text{H}_2\text{O}$. The measured temperature coefficient for 10° is 1.86.

J. F. S.

Mixed Crystal Formation in Ternary Systems containing Water, Ammonium Chloride, and Ferrous, Cobaltous, or Nickel Chloride. FREDERICK WILLIAM JEFFREY CLENDINNEN (T., 1922, **121**, 801—805).

Complex Nitrites of Nickel. V. CUTTICA (*Gazzetta*, 1922, **52**, i, 210—215).—In consequence of the slight tendency to dissociate exhibited by nickel nitrite, the latter is able to form complexes, double nickel nitrites occupying a position between true complex compounds and double salts and possessing the constant co-ordination number of the former and the great mobility of the component simple ions in solution shown by the latter. Cryoscopic measurements of aqueous solutions of nickel thallium nitrite, $4\text{TlNO}_2 \cdot \text{Ni}(\text{NO}_2)_2$, give a molecular weight of 107—111, the calculated value being 1050; hence rather more than nine of the eleven simple ions are manifest in solution. Since thallium nitrite is completely dissociated in the experimental conditions employed, the nickel nitrite is only partly dissociated.

Owing to the markedly "imperfect" nature of double nickel nitrites it is necessary, in the preparation of triple nickel nitrites, that the active masses of the component ions be high in order

that undissociated molecules of the complex may be formed and that the solubility product may be exceeded. The failure of Przibylla (A., 1898, ii, 162; 1899, ii, 222) and of Reichard (A., 1904, ii, 488, 741) to obtain triple nickel cadmium nitrites was probably due to neglect of this condition. The author has prepared *nickel cadmium potassium nitrite*, $\text{Ni}(\text{NO}_2)_2 \cdot 2\text{Cd}(\text{NO}_2)_2 \cdot 4\text{KNO}_2$, which is pale yellow; *nickel cadmium thallium nitrite*, of analogous formula, which is brick red, and *nickel cadmium ammonium nitrite*, $\text{Ni}(\text{NO}_2)_2 \cdot \text{Cd}(\text{NO}_2)_2 \cdot 2\text{NH}_4\text{NO}_2$, which is pale flesh-coloured. These three compounds are profoundly disgregated by the dissociating action of water, the nickel and cadmium nitrites undergoing only limited dissociation.

Many of the triple nickel nitrites known may be represented by the general scheme for the double nickel nitrites, for example, $\left[\text{Ni}(\text{NO}_2)_6\right]_{\text{K}_2}^{\text{Ba}(\text{Ca}, \text{Sr}, \text{Pb})}$, $\left[\text{Ni}(\text{NO}_2)_6\right]_{(\text{NH}_4)_2}^{\text{Cd}}$, the nickel in these exhibiting analogy to iron and cobalt. In the case of nickel cadmium potassium and nickel cadmium thallium nitrites, which do not correspond with this scheme, if the fundamental co-ordinating action capable of functioning in two concentric spheres is attributed to the nickel, it may be assumed that in the inner sphere are co-ordinated six nitrous residues in the same ion as occurs in the other nickel nitrites and that in the second sphere two whole cadmium nitrite molecules remain bound. This supposition, expressed by the formula $\left[\text{Ni}(\text{NO}_2)_6\right]_{[\text{Cd}(\text{NO}_2)_2]_2}^{\text{K}(\text{Tl})_4}$, is in accord with the similarity in colour between these compounds and the double and certain other triple nickel nitrites. T. H. P.

Complex Chlorides containing Gold. I. Pollard's Ammonium Silver Auric Chloride. HORACE L. WELLS (*Amer. J. Sci.*, 1922, 3, [v], (16), 257—259).—The triple chloride examined by Pollard (T., 1920, 117, 99) was investigated. It was found that when solutions somewhat similar to the one recommended by Pollard were diluted with an equal volume or more of 1:1, or stronger, hydrochloric acid better products for analysis were obtained. The results of analysis suggest the formula $(\text{NH}_4)_6\text{Ag}_2\text{Au}_3\text{Cl}_{17}$ for the triple salt against $(\text{NH}_4)_8\text{Ag}_3\text{Au}_4\text{Cl}_{23}$ put forward by Pollard. The corresponding potassium salt could not be obtained, but a triple salt of caesium was prepared and is being investigated. W. T.

Complex Chlorides containing Gold. II. Caesium Triple Salts. HORACE L. WELLS (*Amer. J. Sci.*, 1922, 3, [v], 315—326).—Five new triple chlorides of caesium have been prepared. $\text{Cs}_4\text{Ag}_2\text{Au}'''_2\text{Cl}_{12}$, very black, opaque, powder black. $\text{Cs}_4\text{ZnAu}'''_2\text{Cl}_{12}$, yellow (sometimes red), transparent, powder pale yellow. $\text{Cs}_4\text{HgAu}'''_2\text{Cl}_{12}$, orange, transparent, powder yellow. $\text{Cs}_4\text{CuAu}'''_2\text{Cl}_{12}$, crystals black, powder pale brown. $\text{Cs}_4\text{Au}_2'\text{Au}'''_2\text{Cl}_{12}$, very black, opaque, powder black. These triple salts are apparently isomorphous, an interesting case of isomorphous replacement of two univalent atoms by one bivalent atom. Caesium calcium auric

chloride could not be prepared. These caesium triple salts do not correspond with the ammonium triple chloride (Wells, preceding abstract). In preparing these triple chlorides a large excess of caesium chloride over the theoretical quantity is desirable, but the solutions should be very dilute in respect to gold in order to avoid the deposition of the sparingly soluble caesium auric chloride. Presence of concentrated hydrochloric acid is favourable. W. T.

Mineralogical Chemistry.

Cristobalite in Basalt from Washington. EARL V. SHANNON (*J. Washington Acad. Sci.*, 1922, 12, 195—196).—Minute, white crystals of cristobalite (SiO_2) occur with plagioclase, magnetite, opal, etc., in the steam-cavities of basalt at Spokane; they have the form of cubo-octahedra. L. J. S.

Becquerelite, a New Radioactive Mineral. ALFRED SCHOEP (*Compt. rend.*, 1922, 174, 1240—1242).—This mineral occurs as canary- to orange-yellow, crystalline crusts on pitchblende from the Kasolo mine, Katanga, Belgian Congo. Minute crystals are orthorhombic with perfect (001) and (110) cleavages. The acute negative bisectrix emerges through (001). The mean of two analyses on material dried at 100° gave :

UO_3 .	Fe_2O_3 .	PbO .	SiO_2 .	SO_3 .	H_2O .	Total.
86.51	0.54	5.25	0.83	1.01	5.82	99.96

Needles of anglesite and crystals of curite (this vol., ii, 77) and soddite (this vol., ii, 451) are present; and microchemical tests of the canary-yellow material show only traces of lead. Deducting lead, etc., and adding water (4.21%) lost at 100° , the formula is deduced as $\text{UO}_3, 2\text{H}_2\text{O}$. L. J. S.

Sincosite, a New Mineral. WALDEMAR T. SCHALLER (*J. Washington Acad. Sci.*, 1922, 12, 195).—This forms green, rectangular plates, which are optically uniaxial (sometimes biaxial) and negative, in black carbonaceous shale near Sincos, Peru. Analysis :

CaO .	V_2O_4	P_2O_5 .	H_2O .	Insol.	Total.
12.1	36.3	31.7	19.9	0.3	100.3

gives the formula $\text{CaO}, \text{V}_2\text{O}_4, \text{P}_2\text{O}_5, 5\text{H}_2\text{O}$. Since the mineral belongs to the uranite group (autunite, torbernite, etc.), the "equivalent valency" of quadrivalent vanadyl-vanadium with hexavalent uranic-uranium is suggested. L. J. S.

Optical-crystallographic Properties of Calcium Oxalate Monohydrate. EDGAR T. WHERRY (*J. Washington Acad. Sci.*, 1922, 12, 196—200).—The crystallographic and optical constants

hitherto determined for the mineral whewellite are tabulated, and the relation of the optical orientation to the various habits of the crystals is pointed out. Fragments of the mineral, crystals from plant tissues, and crystalline precipitates (prepared by boiling together solutions of the constituent ions) were examined microscopically. Refractive indices, α 1.490, β 1.555, γ 1.650.

L. J. S.

The Chemical Constitution of Felspars. Analysis of Two Microclines. M. E. DENAEYER (*Bull. Soc. chim. Belg.*, 1922, **31**, 131—147).—Theories of the constitution of complex silicates (W. and D. Asch, "Die Silicate," Berlin, 1911; P. Erculisse, "Classification chimique des silicates naturels," Brussels, 1920; J. Jakob, A., 1920, ii, 754) are discussed, and an attempt is made to apply Werner's co-ordination theory to the problem. Some formulæ based on the latter theory are given and a pseudo-phase rule diagram based on these is used to suggest the mutual relationship of quartz, orthose, plagioclases, micas, and spinels. The connexion between the views put forward by Jakob and by F. W. Clarke is briefly discussed.

The analysis of two microclines (triclinic potash-soda felspars) is considered to indicate that they are built up of three separate components in solid solution, and a study of results of analyses carried out by other workers tends to confirm this view.

H. J. E.

Soddite, a New Radioactive Mineral. ALFRED SCHOEP (*Compt. rend.*, 1922, **174**, 1066—1067).—This forms with curite (A., 1922, ii, 77) fine-grained crystalline aggregates at Kasolo, Katanga, Belgian Congo. In veins in this material minute prismatic crystals were found which from their optical characters are orthorhombic. H 3—4, d^{17} 4.627. Heated in a bulb-tube the mineral gives off water and oxygen and turns black, and it is infusible before the blow-pipe. The mean of several partial analyses is :

SiO ₂ .	UO ₃ .	Fe ₂ O ₃ .	H ₂ O.
7.83	85.33	0.40	6.23

agreeing with the formula $12\text{UO}_3, 5\text{SiO}_2, 14\text{H}_2\text{O}$.

L. J. S.

Ore Deposits and their Genesis in Relation to Geographical Distribution. A Lecture Delivered before the Chemical Society on December 8th, 1921. JOHN WALTER GREGORY (T., 1922, **121**, 750—772).

Meteoric Iron from Odessa, Ector Co., Texas. GEORGE P. MERRILL (*Amer. J. Sci.*, 1922, [v], **3**, 335—337).—The much rusted fragment of 1120 grams as received was stated to be from a larger mass found near Odessa. The etched surface shows a coarse octahedral structure. Analysis by E. V. SHANNON gave :

Fe.	Ni.	Co.	Cu.	Pt.	Cr.	Mn.	C.	P.	S.	Total.
90.69	7.25	0.74	0.02	nil	trace	nil	0.35	0.23	0.03	99.31

L. J. S.

Meteoric Irons from Alpine, Texas, and Signal Mountain, Lower California. GEORGE P. MERRILL (*Proc. U.S. Nat. Mus.*, 1922, 61, art. 4, 1—4).—A fragment of a mass, said to weigh two tons, from Alpine shows the finely granular structure of an ataxite, and gave anal. I by J. E. WHITFIELD. A fragment from a mass, said to have been observed to fall at Signal Mountain several years ago, gave II. The medium octahedrite structure grades into a finely granular structure at the exterior of the mass, the latter structure having evidently resulted by the heating during flight through the earth's atmosphere.

	Fe.	Ni.	Co.	Cu.	S.	P.	C.	Si.	Total.
I	93.60	5.62	0.43	0.016	0.012	0.328	0.008	0.015	100.029
II	91.47	7.86	0.60	0.015	0.002	0.041	—	0.004	99.992

A pallasite from Cold Bay, Alaska, is also briefly described.

L. J. S.

Analytical Chemistry.

The Application of Conductometric Methods to Precipitation Analysis. I. M. KOLTHOFF (*Z. anal. Chem.*, 1922, 61, 171—180).—The author discusses the conditions necessary for obtaining good results by conductometric methods in precipitation analysis and details the possible sources of error under three headings: errors in the determination of the conductivity, errors due to the solubility of the precipitate being too great, and errors due to the composition of the precipitate not being constant. In the first case, the analysis is more exact the slower the rate of migration of the ions taking the place of those precipitated, whilst the end-point is sharper the greater the speed of the anion which is precipitated and the greater the degree of dissociation of the salt. The speed of the ions present which take no part in the reaction is immaterial, but, as the presence of indifferent electrolytes decreases the accuracy of the method, they should be kept as low as possible and the reagent used for precipitation must be a strong electrolyte. The solubility of the precipitate must, at the most, not exceed a normality corresponding with one-twentieth of the normality of the solution from which it is produced. Errors may arise owing to the slow rate at which the precipitate forms or to adsorption of salts by the precipitate; the former may be overcome by the addition of alcohol to the solution.

A. R. P.

Estimation of Hydrogen-ion Concentration in Urine with Indicators. ADOLF SILBERSTEIN (*Biochem. Z.*, 1922, 128, 534—539).—The colorimetric estimation of the P_H of coloured liquids like urine is simplified by removing the natural colouring matter by shaking with charcoal, for instance, "Carbo medic," Merck. There is no change of P_H after this treatment of urine.

H. K.

Reaction between Thiosulphuric and Nitrous Ions. P. FALCIOLO (*Gazzetta*, 1922, 52, i, 179—182).—If sodium thiosulphate, even in very dilute solution, is treated with dilute sodium or potassium nitrite solution and the liquid is then acidified with either an inorganic or organic acid or a salt, such as alum, giving an acid solution, more or less marked effervescence occurs and the solution assumes a yellow colour which, according to the amounts of the substances used, may at first be green or orange-brown. The reaction is equally sensitive in aqueous alcoholic solution and is shown distinctly by 0.0001N-sodium thiosulphate solution, which does not readily yield sulphur when treated with a mineral acid or give a coloration with ferric chloride. Only excessive proportions and concentrations of sulphurous acid prevent the reaction with traces of thiosulphate. The nitrous ion in presence of the nitric ion may be detected by means of the reaction (cf. Bodnár, A., 1914, ii, 67).

T. H. P.

The Volumetric Estimation of Dithionates. A. FISCHER and W. CLASSEN (*Z. angew. Chem.*, 1922, 35, 198—199).—Sodium dithionate is not attacked by cold alkaline or neutral oxidising agents and but little in acid solution. On heating with acid, evolution of sulphur dioxide commences and oxidation follows. Attempts to devise a direct titration with an acid oxidising agent were unsuccessful either through loss of sulphur dioxide or through secondary reactions. A method was therefore adopted of distilling the sulphur dioxide into a known volume of an oxidising agent. Potassium permanganate and dichromate proving unsuitable, iodine solution was employed. The sodium dithionate is weighed into a flask, distilled hydrochloric acid free from chlorine added, and a stream of carbon dioxide passed through the apparatus. The flask is then heated and at the end of the reaction carbon dioxide is again passed through the apparatus. The results are in close agreement with those of gravimetric analysis. The method may be applied to mixtures of dithionates with other sulphur salts, most of which are decomposed by cold hydrochloric acid or oxidised by alkaline hydrogen peroxide.

C. I.

Estimation of Ammonia in Urine by Schloesing's Bell-glass Method. W. MESTREZAT and (MLLE) M.-P. JANET (*Bull. Soc. Chim. Biol.*, 1922, 4, 154—164).—Schloesing's method is modified and used for the estimation of ammonia in urine. To 20 c.c. of urine contained in a desiccator are added a few drops of sulphuric acid to dissolve any ammonium magnesium phosphate, and five drops of a 10% solution of mercuric cyanide to prevent bacterial decomposition of urea. Ten c.c. of 20% milk of lime are then added, after which a basin containing 10 c.c. of 0.1N-sulphuric acid is rapidly placed on a tripod in the desiccator and the latter closed. After remaining for three days at a room temperature of 15—20°, the excess of acid is titrated. No appreciable quantity of ammonia is evolved from nitrogenous organic substances present in the urine.

E. S.

The Estimation of Ammoniacal Nitrogen in Nitrogenous Organic Substances, and particularly in Proteins and their Products of Decomposition. J. FROIDEVAUX (*Compt. rend.*, 1922, 174, 1238—1240).—To 15 c.c. of the solution under examination 35 c.c. of 60% aqueous sodium hydroxide are added and air, carefully freed from ammonia, is bubbled through at the ordinary temperature at the rate of 150—200 bubbles per minute. The issuing air is passed through a known volume of standard acid. The acid is replaced from time to time and titrated. The ammonia absorbed is plotted against the time. The resulting curve consists of a line sharply inclined to the time axis and a line slightly inclined to the same axis, these being joined by a curved portion. The first line corresponds with ammoniacal nitrogen and the second line with ammoniacal nitrogen coming from the slow decomposition of protein or amino-acids. The curved portion is a combination of the two. The two lines are produced to meet and their junction represents the total ammoniacal nitrogen originally present as such in the sample. W. G.

The Analysis of Liquid Nitrogen Peroxide. A. SANFOURCHE (*Bull. Soc. chim.*, 1922, [iv], 31, 316—319).—The sampling is done by means of a Durand washing flask, the central tube of which is drawn out to a fine orifice, the flask being kept immersed in ice.

For the estimation of nitric acid in the sample, 10 c.c. of the liquid are measured out into a cylindrical gas-drying tube, which is then surrounded with ice, and air is bubbled through the liquid until the whole of the nitrogen peroxide has evaporated. The residual nitric acid, which should not evolve nitrous vapours when warmed by the hand, is diluted with water and titrated with *N*-sodium hydroxide.

For the estimation of nitrogen peroxide and nitrous anhydride, a known volume of the sample is dissolved in 20 c.c. of concentrated sulphuric acid cooled in ice, and aliquot portions of the solution are titrated with *N*/10-permanganate, and analysed in a Lunge nitrometer, respectively. From these results it is possible to calculate the percentages of nitrogen peroxide and nitrous anhydride respectively in the sample. W. G.

Estimation of Nitrates in Drinking Water by Mayrhofer's Method. A. REUSS (*Z. Unters. Nahr. Genussm.*, 1922, 43, 174—183).—The consumption of indigotin solution is considerably increased in the presence of sodium chloride, the reaction between small quantities of nitrates and the indigotin being facilitated by this substance. If only small quantities of nitrates and small quantities of chlorides are present in the water, it is advisable to add sufficient sodium chloride to bring the concentration up to 1 gram per litre. The indigotin solution must then be standardised against a nitrate solution containing 1 gram of sodium chloride per litre. Mayrhofer's tables can then be used. The indigotin solution must not have the slightest sediment or its titre will be variable. It is best filtered through asbestos without suction until no suspended particles are visible with a lens. The solution should

be added at a rate of 2—3 drops per second, the last drops being added at a somewhat slower rate. Care must be taken that the whole of the 5 c.c. of sulphuric acid actually reach the liquid in the flask. The acid used must be of the same composition as that used in standardising the indigotin solution which should be prepared as described by Lehmann in an appendix to the paper.

H. C. R.

The Estimation of Inorganic Phosphorus in Blood Plasma by Bell and Doisy's Method. BURTON A. MYERS and MARIAN C. SHEVKY (*J. Lab. Clin. Med.*, 1921, 7, 176—180; cf. A., 1920, ii, 769).—Phosphorus was not recovered quantitatively from plasma by Bell and Doisy's method, when added in known quantity. The only solution correctly measured was one which was weaker than the standard; all stronger than the standard showed considerable minus errors. This difficulty was obviated by using a series of standards of such strength that one might be chosen containing approximately 0.25 mg. per 100 c.c. more phosphorus than the unknown. A small series of estimations thus made gave an error ranging from 2.3 to 10% with an average of 5.8%. In many rabbit plasmas and some human plasmas larger amounts of molybdic acid and quinol solutions than prescribed in the original directions must be used to develop the proper colour. 1.5 c.c. of molybdic acid and 3 c.c. of quinol for each 5 c.c. of plasma filtrate (1 : 5 dilution) are recommended.

CHEMICAL ABSTRACTS.

Qualitative Reactions for Arsenic. I. M. KOLTHOFF (*Pharm. Weekblad*, 1922, 59, 334—350).—The reactions of Mayençon and Bergeret (this Journal, 1874, 1008; reduction to arsine and detection with mercuric chloride paper), Bougault (reduction to free arsenic by use of hypophosphite), Bettendorf (reduction to arsenic, using stannous chloride), and Reinsch have been examined. The most suitable is the first, which under the conditions recommended will detect one-millionth of a gram (1 μ .g.) in 1 c.c. To 1 c.c. of the neutral solution are added 1 c.c. of 22% hydrochloric acid containing 1% of stannous chloride, and 0.1 gram of finely divided aluminium foil. The gas is led through cotton-wool on to a strip of paper 4 mm. wide prepared by saturating filter-paper in 5% aqueous mercuric chloride solution. After one hour the colour of the paper is examined. Antimony must be removed by treating the paper with hydrochloric acid or potassium iodide.

Mercury salts interfere strongly, copper salts moderately; salts of other metals do not seriously affect the test.

Bougault's reaction will detect 2 mg. of arsenic per litre. Two c.c. of the solution, admixed with 1 c.c. of the reagent, and 4 c.c. of strong hydrochloric acid, are heated in the water-bath for thirty minutes. A brown colour shows presence of arsenic. The solution may be filtered, a stain remaining on the paper. S. I. L.

Aluminium for the Arsenic Reaction. G. ROMIJN (*Chem. Weekblad*, 1922, 19, 177—179).—The author agrees with Kolthoff

(preceding abstract) that aluminium is to be preferred to zinc, which he had formerly recommended (Romijn, *Pharm. Weekblad*, 1917, **54**, 1216), in the Mayençon and Bergeret reaction. He details some improvements in the apparatus which allow of the detection of 0.5 μ .g. of arsenic with a reaction time of two to five minutes. He suggests that in presence of formaldehyde hexamethylene-tetrarsine may be formed. S. I. L.

Estimation of Volatile Matter in a Coal. A. DESSEMOND (*Rev. ind. min.*, 1921, **1**, 451—456).—The essential analytical operations for estimating the volatile substances and ash of a coal by desiccation, distillation, and incineration are discussed. In the formula $V = v/(100 - c)$, where $c = \%$ ash, $v = \%$ volatile substances in the crude coal, and $V = \%$ volatile substances exclusive of the ash, calculations show that the percentage of V increases as the percentage of ash increases and becomes infinite when the ash becomes 100. A general investigation of the formula and of the errors usual in the procedures has led to the proposal of a simple formula expressing the content of volatile substances in a coal exclusive of the ash: $V = [v - (c/n)]/(100 - c)$, where n is a variable coefficient depending on the impurities. Numerous samples of coal are given, and it is suggested that the value $n = 10$ be adopted as a sufficiently accurate approximation for most coals. CHEMICAL ABSTRACTS.

The Separation of Zinc by Means of Ammonium Phosphate. G. LUFF (*Chem. Zeit.*, 1922, **46**, 365—366).—Magnesium and manganese may be separated quantitatively from zinc by precipitation with ammonium phosphate in solutions containing at least one-fifth of their bulk of ammonia (d 0.923), whilst iron and aluminium may be precipitated as phosphates free from zinc by addition of ammonium phosphate to the solution containing at least one-fifth of its bulk of glacial acetic acid. The zinc is recovered from the filtrate by rendering it neutral to litmus, in the first case with hydrochloric acid, and in the second case with ammonia. The separation of manganese and zinc must be carried out in an atmosphere of hydrogen to prevent precipitation of manganese dioxide. [Cf. *J. Soc. Chem. Ind.*, 1922, June.] A. R. P.

An Electro-volumetric Method for Lead. D. A. MACINNES and ERIC B. TOWNSEND (*J. Ind. Eng. Chem.*, 1922, **14**, 420—421).—The solution containing the lead as nitrate together with free nitric acid is electrolysed with a current of 13 amperes for thirty to forty-five minutes in a platinum dish, 9 cm. in diameter and roughened on the inside, which acts as anode whilst the cathode consists of a small platinum disk rotated at 600 revolutions per minute. The deposit is washed with cold water and dissolved in an excess of standard oxalic acid and 5 c.c. of nitric acid heated at 80°. The solution is rinsed into a beaker, a few c.c. of sulphuric acid are added, and the excess of oxalic acid is estimated by titration with permanganate. The results prove that the deposit consists entirely of lead dioxide with variable amounts of water, and that, contrary to previous statements, it contains no higher oxides.

The method described is more exact than that of weighing the deposit and calculating the lead by using an empirical factor, as the amount of water in the deposit varies with its weight.

A. R. P.

Red Coloration of Sodium Hypochlorite Solutions. TEALDI MARIO (*Boll. Chim. Farm.*, 1922, **61**, 165—166).—Sodium hypochlorite prepared from chemically pure calcium hypochlorite and sodium hydrogen carbonate develops no red coloration, and the presence of manganese in the glass of the containing vessel neither induces the coloration nor enhances the intensity of solutions already reddened. If commercial reagents are used in its preparation the salt at once turns red, irrespective of the presence of manganese in the reaction vessel. The coloration is not modified by addition of an iron salt, but is destroyed by sodium thiosulphate. The conclusion is drawn that the red colour is due to salts of permanganic acid derived from the manganese in the reagents used.

T. H. P.

Chemical Analysis with Membrane Filters. III. Their Application to Volumetric Analysis. GERHART JANDER (*Z. anal. Chem.*, 1922, **61**, 145—171).—Membrane filters are particularly suited to volumetric work involving precipitation of a definite compound followed by solution of the precipitate and titration, owing to the ease with which the precipitate can be removed from the filter simply by spraying water over it. The estimation of calcium by precipitation and titration of the oxalate with permanganate, of manganese by precipitation of the sulphide followed by solution of the latter in standard acid and titration of the excess acid with alkali, and of chromic acid in the presence of other oxidising acids is described in detail.

The estimation of manganese is carried out by treating the solution, 70 c.c., containing not more than about 0.25 gram of metal with 30 c.c. of 10% ammonium chloride solution, a few c.c. of ammonia and, after boiling, a large excess of freshly prepared ammonium sulphide solution. Boiling is continued for ten minutes to change the precipitate into the green modification, more ammonium sulphide is added, and the solution is cooled and filtered on a "30—60 second" membrane filter. The precipitate is washed first with water containing ammonium sulphide, then with neutral 2% sodium sulphate solution, rinsed off the filter with cold water, and dissolved in 60 c.c. of *N*/5-sulphuric acid. The solution is boiled to expel hydrogen sulphide, cooled, and titrated with *N*/5-sodium carbonate solution, using methyl-orange as indicator.

The volumetric estimation of chromium as chromate in solutions containing other oxidising substances, for example, chlorates or bromates derived from the oxidation of trivalent chromium to the hexavalent form in alkaline solution by means of chlorine or bromine, is carried out as follows: the solution is just acidified with acetic acid, 20 c.c. of *N*-sodium acetate solution are added, and the chromic acid is precipitated from the boiling solution by the addition, drop by drop, of a slight excess of *N*/10-lead acetate

solution. After cooling, the precipitate is collected on a membrane filter, washed with cold water, rinsed back into the beaker, and dissolved in 100 c.c. of *N*-hydrochloric acid. An excess of a standard solution of ferrous sulphate is added and the excess is estimated by titration with potassium dichromate as usual.

A. R. P.

The Estimation of Cobalt in Steel. ALOIS EDER (*Chem. Ztg.*, 1922, 46, 430).—The method depends on the separation of the iron from a weakly acid sulphate solution by means of an emulsion of zinc oxide and precipitation of the cobalt in an aliquot part of the filtrate after acidification with hydrochloric acid by means of an excess of a solution of nitroso- β -naphthol in 50% acetic acid, followed by ignition of the precipitate mixed with oxalic acid to the oxide Co_3O_4 , which is weighed. [Cf. *J. Soc. Chem. Ind.*, 1922, June.]

A. R. P.

Detection and Estimation of Small Quantities of Nickel and Cobalt in Silicate Rocks. O. HACKL (*Chem. Ztg.*, 1922, 46, 385—386).—A large quantity of the sample is digested with aqua regia, the filtered solution is evaporated to expel nitric acid, the residue digested with hydrochloric acid, the solution diluted with water, treated with hydrogen sulphide, filtered, and the filtrate made ammoniacal and saturated again with hydrogen sulphide. A large excess of 5% hydrochloric acid is added, the solution set aside until the precipitate has settled, and then filtered, the precipitate being washed with dilute hydrochloric acid solution saturated with hydrogen sulphide. It is dissolved in aqua regia and the solution divided into two parts, one of which is tested for nickel by means of α -dimethylglyoxime in weakly ammoniacal tartaric acid solution, and the other for cobalt, after removing the trace of iron present by any suitable method, by Vogel's thiocyanate reaction or by means of nitroso- β -naphthol. The quantitative estimation of the metals is made in the same way, except that the sulphide precipitate after dissolving in aqua regia is freed from iron by triple precipitation of the solution with ammonia or by precipitation with barium carbonate. Nickel and cobalt are again precipitated as sulphides in the filtrate from the iron, the sulphides are ignited to oxides, and these reduced to and weighed as metals. Either the nickel or the cobalt is then estimated and the other found by difference. In all cases, the insoluble residue from the original aqua regia treatment of the ore must be tested for the presence of nickel and cobalt by dissolving it in hydrofluoric acid and testing the solution as described above.

A. R. P.

A Modified Reaction of Tin. HANS HELLER (*Z. anal. Chem.*, 1922, 61, 180—182).—The following modification of the iodide method for the detection of tin gives more trustworthy results than that recommended by Mazuir (A., 1920, ii, 197). One c.c. of the solution containing only a little hydrochloric acid is mixed with 0.5 c.c. of 5% potassium iodide solution and 0.5 c.c. of strong

sulphuric acid is introduced, by means of a pipette, below the surface of the liquid. If tin is present an immediate precipitate of yellow iodide forms at the surface between the two liquids. The precipitate dissolves readily on addition of hydrochloric acid and careful shaking. Arsenic and antimony interfere with the test.

A. R. P.

The Volumetric Estimation of Titanium Dioxide in Bauxite.

H. J. WINCH and V. L. CHANDRATREYA (*Chem. News*, 1922, 124, 231—232).—0.3 Gram of the sample is dissolved by fusion with 3 grams of potassium pyrosulphate and the fused mass is dissolved in hydrochloric acid. The solution is boiled with 0.15 gram of tin powder and 5 grams of ammonium chloride until all the tin has dissolved. Cold water is added to the liquid followed by 50 c.c. of 4% mercuric chloride solution to destroy excess of stannous chloride (titaneous chloride is not oxidised by mercuric chloride in the cold) and 5 c.c. of ferric chloride solution containing 10 grams of iron per litre. The liquid is then titrated with potassium dichromate as usual. The iron is determined in a separate sample by reduction with stannous chloride followed by titration with the same dichromate solution. The difference in the two readings corresponds with the titanium dioxide present.

A. R. P.

Detection and Estimation of Vanadium in Steels. GEORGES MISSON (*Bull. Soc. chim. Belg.*, 1922, 31, 123—126).—A colorimetric method of estimating vanadium is described, for which considerable accuracy is claimed. The steel is dissolved in nitric acid, any organic matter oxidised by means of permanganate, and, after oxidising any precipitated oxide of manganese with a solution of sodium peroxide in dilute nitric acid, a further quantity of the latter reagent is added in order to produce the colour for comparison. It is essential that all the reagents should be free from chlorine. Modifications are described which render the method applicable to steels containing tungsten, chromium, and nickel.

H. J. E.

Detection of Traces of Osmium by Means of Potassium Thiocyanate. MAX HIRSCH (*Chem. Ztg.*, 1922, 46, 390).—The metal to be tested for osmium is dissolved in fuming nitric acid or by fusion with sodium hydroxide and potassium nitrate or chlorate, followed by solution of the fused mass in water and acidification of the solution with nitric acid. In either case, the solution is distilled and the distillate collected in cold water. This solution is acidified, concentrated potassium thiocyanate solution added, and the whole shaken with ether or amyl alcohol. A blue colour in the ethereal layer indicates osmium. The colour is just perceptible at a dilution of 1 part of osmium per million parts of solution.

A. R. P.

The Clarification of Solutions containing Reducing Sugars by Basic Lead Acetate. The Effect of Different Deleading Agents. DUANE T. ENGLIS and CHUK YEE TSANG (*J. Amer. Chem. Soc.*, 1922, 44, 865—867).—The removal of excess of basic

lead acetate, used as a clarifying agent from solutions of dextrose or l  vulose results in the loss of sugar. Of the agents tried, namely, potassium oxalate, disodium hydrogen phosphate, potassium sulphate, potassium sodium tartrate, and sodium carbonate, the least loss of sugar occurred when disodium phosphate was used. In general, the loss of l  vulose is much greater than that of dextrose. If the precipitate is washed, a much smaller loss is observed. W. G.

Detection of Lactic Acid in Organic Liquids. F. D'ARBELA (*Rend. accad. med.-fis. fiorentina; Sperim.*, 1921, 75, 415—416).—Pittarelli's method (A., 1921, ii, 418) is not specific, as the coloration is given by many other substances. Uffelmann's reaction is preferred.

CHEMICAL ABSTRACTS.

The Analysis of Solutions of Ammonium Citrate. C. S. ROBINSON and SELMA L. BANDEMAR (*J. Ind. Eng. Chem.*, 1922, 14, 429—432).—The methods proposed for the determination of the ratio of ammonia to citric acid in ammonium citrate solutions have been examined and the results obtained are tabulated. The formaldehyde method invariably yields accurate results provided sufficient of the reagent is added to combine with all the ammonia and the addition of alkali is carried to the first permanent pink colour of the phenolphthalein. The only other method that gives accurate results is as follows: 50 c.c. of the solution are diluted to 500 c.c. and 10 c.c. trials are distilled with an excess of standard alkali to obtain the ammonia. The residue in the distillation flask is titrated with standard acid until the solution is distinctly acid to methyl-red; it is then boiled to expel carbon dioxide, a few drops of phenolphthalein are added, and the solution is titrated with standard alkali hydroxide until a permanent pink colour is obtained. The difference between the total alkali and acid used gives the amount of alkali used in neutralising the citric acid. A. R. P.

Golodetz's Reaction (The Benzoyl Peroxide Reaction). H. C. J. H. GELISSEN (*Rec. trav. chim.*, 1922, 41, 224—227).—Golodetz's reaction (A., 1908, ii, 330) is not specific for benzoyl peroxide, but is given by all peroxides which, on treatment with concentrated sulphuric acid, yield phenolsulphonic acid. It may therefore be regarded as based on Hehner's reaction. Liebermann's test and Millon's test for phenol apply equally to benzoyl peroxide under the conditions indicated for Golodetz's reaction. A modification of Golodetz's procedure is recommended, as the test is considerably more sensitive if the sulphuric acid is cooled to -20° instead of being gently heated; the tests for phenol are also clearer at the lower temperature. H. J. E.

Indican Reactions for the Detection of Urine in Stains. GIUSEPPE JEMMA (*Arch. Farm. speriment. Sci. aff.*, 1921, 32, 193—194).—Replacement of thymol by α -naphthol in the application of Jolles's test (A., 1915, ii, 593, 655) to the detection of urine in stains (Latta, A., 1920, ii, 339) renders the reaction from two to four times more sensitive. The use of α -naphthol-*p*-sulphonic acid in this test leads, however, to unsatisfactory results. T. H. P.

General and Physical Chemistry.

Spectrum of Hydrogen. T. R. MERTON and S. BARRATT (Bakerian Lecture, *Phil. Trans.*, 1922, [A], 222, 369—400).—The wave-lengths of about 1200 lines in the secondary spectrum have been measured and classified into physically related groups according to the effects of the pressure of the gas, electrical excitation, and the presence of helium. This method of classification has been compared with the results of other investigators relating to the Stark and Zeeman effects and with the regularities observed by Fulcher. By means of a new method of measuring the widths of spectrum lines, it is shown that the secondary spectrum is due to hydrogen molecules. No evidence of the presence of secondary hydrogen has been found in the spectra of the sun. The effect of impurities and changes in the methods of electrical excitation on the relative intensities of the Balmer and secondary series is discussed, and it is shown that the greater the purity of the hydrogen the more prominent the secondary spectrum. The light from different portions of the capillary of the discharge tube varies in character with the nature of the discharge. With a condenser and spark gap in the electrical circuit, a mixture containing hydrogen and helium gives the hydrogen line at the two ends of the tube. Mercury lines and the lines of the heavier elements appear in the middle of the tube. On cutting out the condenser, this unequal distribution slowly changes. A partial separation of the gases in the discharge tube would explain these phenomena and the observation of Wood (cf. A., 1921, ii, 665) with long vacuum tubes, that the Balmer series are strongest in the middle of the tubes.

W. E. G.

Excitation of Gas-spectra during Chemical Reactions. F. HABER and W. ZISCH (*Z. Physik*, 1922, 9, 302—326).—A continuation of the work of Haber and Just (cf. A., 1911, ii, 954) on the emission of electrons in chemical reactions. The nature of the light emitted from flames of sodium and mercury burning in the halogens is investigated. Sodium vapour, diluted by nitrogen, is burnt in chlorine, bromine, iodine, or oxygen, the concentrations being so adjusted that the temperature of the flame is maintained below the limit at which temperature radiation becomes visible. With chlorine, a luminosity, greyish-green in appearance, is first observed at 350—360°. At higher temperatures, it becomes yellow, and at 473° is of sufficient intensity to make it possible to photograph the sodium doublet. The source of the *D*-line is the free sodium atom which is activated by collision with a freshly formed molecule, NaCl or NaCl₂, which has not yet dissipated its energy of combination. It is considered that the conductivity and luminosity of the inner cone of the Bunsen flame are partly chemical in origin and this view is supported by the intense colour of the inner cone when an alkali metal is added to the flame.

Mercury, burnt under similar conditions, gives a green cone with a band showing maxima between $595\ \mu\mu$ and $300\ \mu\mu$, whereas the wave-length, corresponding with the heat of formation of mercuric chloride, is $546\ \mu\mu$.

The conductivity of oxygen and potassium vapour at low pressures is considerable, so that electrically charged particles, possibly electrons, are liberated. New determinations of the vapour pressure of sodium vapour between 473° and 563° have been made. The relationship between light and electron emission during chemical reaction is discussed. In light emission, the excitation of the electron takes place from the normal state to the first quantum orbit, whereas in electron emission the electron is raised to an infinitely large orbit.

W. E. G.

Doublets in the Visible Spectrum. S. GOUDSMIT (*Arch. Néerland.*, 1922, [iii], 6, 116—126).—Diverse empirical relationships have been found between the doublets in the visible spectrum. Sommerfeld gives $\Delta\nu = 0.365 (Z - z)^4$; where $\Delta\nu$ is the breadth of the doublet in cm^{-1} , Z is the nuclear charge or atomic number, and z is the number which indicates the "screening effect" of the electrons; Sommerfeld calls $(Z - z)$ the effective nuclear charge (Z_{eff}). From this equation he has deduced the effective nuclear charge from the L -doublets, the difference between this and the true nuclear charge gives the screening effect of the L -series, and the mean value 3.63 was obtained. It is here assumed that Z_{eff} has the same value at all points on the orbit of the electron, but whilst this may be true in the case of a circular orbit, it cannot hold when the orbit is an ellipse. Lithium gives a spectrum closely resembling that of hydrogen, the doublet $2p$ being almost of the same magnitude. The present author investigated whether this doublet and as a result all the doublets in the spectra have not the same origin as that of hydrogen, that is, that they were due to relativity. This point of view is contradictory to the well-established theory of Sommerfeld concerning the origin of the spectral series, but the results obtained were in good agreement. The objections to this method of treatment are discussed.

W. T.

The L -Series of the Elements Barium to Rubidium. D. COSTER (*Arch. Néerland.*, 1922, [iii], 6, 76—91).—The author gives the wave-lengths and intensities of the L -series of these elements found by Hjalmar (A., 1921, ii, 145) and himself, using the method of Siegbahn. The values found differ somewhat from those found by Siegbahn. The results are discussed in detail on the basis of Bohr's new theory (this vol., ii, 277).

W. T.

Observations on the Rare Earths. XI. The Arc Spectrum of Yttrium. L. F. YNTEMA and B. S. HOPKINS (*J. Opt. Soc. Amer.*, 1922, 6, 121—134).—A determination of the emission spectrum of yttrium. The yttrium material which contained 0.005% of holmium was obtained in the work on atomic weights at the University of Illinois. The spectrum, however, showed

that other rare earth elements, erbium and dysprosium, and also magnesium and silicon, were present in small amounts. The yttrium material has a few lines in common with the eurosamarium of Eder. A comparison is made with the values previously obtained by Eder. W. E. G.

The L-Series of Lutecium and Ytterbium and the Identification of Celtium with the Element of Atomic Number 72. A. DAUVILLIER (*Compt. rend.*, 1922, 174, 1347—1349).—By a modification of his previous technique (cf. A., 1921, ii, 421, 475, 669), the author has been able to make a detailed study of the L-series of lutecium and ytterbium in the form of their oxides. Twenty-six rays have been identified for lutecium, and the wave-lengths of these together with those for ytterbium are tabulated. In addition, the existence of some feeble lines indicated the presence of a trace of thulium. These lines were $\gamma_7=1272.1$; $\gamma_1=1311.3$; and $\beta_2=1456.3$. Further the existence of two very feeble lines, $\beta_2=1319.4$ and $\alpha_1=1561.8$, show the existence of a trace of celtium and assign to it the atomic number 72. This element had been discovered in the same preparation by Urbain (A., 1911, ii, 115) by the appearance of a group of unknown lines in the arc spectrum. W. G.

The N-Series of Röntgen Spectra. V. DOLEJŠEK (*Z. Physik*, 1922, 10, 129—136).—With the aid of the spectrograph of Siegbahn (*Z. Physik*, 1922, 9, 68), determinations have been made of the N-series of uranium, thorium, and bismuth. All these lines were severely tested, to eliminate the M lines of higher order, by absorption through different thicknesses of aluminium and black paper and by a comparison of the spectra reflected from different crystals. A scheme of the N lines of the gas niton (86) shows the presence of seven N, five O, and three P levels in the atom. The various modes of transference of the electrons between these levels are also indicated. All the lines which have been found (λ 8,594— λ 13,255) agree with the theoretical expectations, and all possible lines, except that due to the transference between N_7 and O_4 , have been observed. The P_1-P_2 and O_1-O_2 doublets have been separated for the first time. The hardest lines of the N-spectra for the three metals follow the Moseley frequency relation. W. E. G.

X-Ray Spectra. WILLIAM DUANE and R. A. PATTERSON (*Proc. Nat. Acad. Sci.*, 1922, 8, 85—90).—In view of certain criticisms of their previous work, the measurements of the L-absorption limits of platinum, gold, and bismuth (cf. A., 1920, ii, 407) have been repeated with increased accuracy. The absorption limits are found to possess somewhat shorter wave-lengths than the shortest emission lines associated with them. The authors have also measured the absorption spectrum for the K-series of molybdenum, and there are considerable discrepancies between their measurements and those of Overn (*Physical Rev.*, 1921, 17, 350). The ratio of the intensities of α_1 to α_2 is given as 1.97 as

compared with 2.00 predicted by Bohr. The ratio of the intensities of β to γ in the first order is 6.3, and in the second order is 5.46.

W. E. G.

A Chromophore Grouping of Atoms in Inorganic Triple Salts and a General Theory for the Cause of the Colours of Substances. HORACE L. WELLS (*Amer. J. Sci.*, 1922, [v], 3, 417—422).—The author finds that complex salts containing the same metal in different states of valency are highly coloured, and he suggests that there is a constant exchange of electrons between the atoms of different valency and that this activity of electrons affects the passage of light, producing colours or opacity. An attempt is also made to extend this theory to explain the colours of substances in general by assuming spontaneous exchanges of electrons which affect the passage of light.

W. T.

The Labile Nature of the Halogen Atom in Organic Compounds. III. The Absorption Spectra of Bromomalonic Derivatives and Nitroparaffins, and their Bearing on the Question of an Oxygen-Halogen Linking. HUGH GRAHAM and ALEXANDER KILLEN MACBETH (*T.*, 1922, 121, 1109—1115).

The Relationship between Chemi- and Photo-luminescence in Unsaturated Silicon Compounds. H. KAUTSKY and H. ZOCHER (*Z. Physik*, 1922, 9, 267—284).—The compounds obtained by the action of acids on calcium silicide, for example, oxydisilin, silical hydroxide, and leucone (cf. A., 1921, ii, 505) have been further investigated. Chemiluminescence occurs when silicones, mixtures of oxydisilin, and silical hydroxide, are oxidised by chromic acid or potassium permanganate in acid solution. The colour of the silicones varies from pale yellow to deep red as the concentration of silical hydroxide increases, and the chemiluminescence follows identical colour changes. Oxydisilin gives only a feeble green light, and the dark red silical hydroxide a dark red light. Similar changes in colour occur for mixtures of silical hydroxide and leucone, and the latter, a white substance, shows no chemiluminescence. The photochemical oxidation of oxydisilin with ethyl iodide and water by short wave-lengths of light is accompanied by the emission of light of long wave-length. The intensity of the photoluminescence is dependent on the silical hydroxide concentration, and the change is apparently autocatalytic. Both chemi- and photo-luminescence are strengthened in liquid air, although the velocity of reaction is reduced. The substances in liquid air also exhibit phosphorescence with the same colour as that of the chemiluminescence. The emitted light is plane polarised.

W. E. G.

The Theory of Klein and Rosseland applied to Fluorescence, Photochemical Processes, and the Electron emission from Hot Substances. J. FRANCK (*Z. Physik*, 1922, 9, 259—266).—The theory proposed by Klein and Rosseland (cf. 1921, ii, 291) is extended to collisions between atoms and molecules. It is sought to determine the percentage of excited molecules which

undergo a change from the active into the normal condition after collision with a slow unexcited atom. The fluorescence of dilute iodine gas is shifted towards the red end of the spectrum by the addition of another gas. This result suggests that some of the excited molecules may transfer their rotation and vibration quanta to the colliding atoms without the emission of light. Observations of Wood and of the author on the resonance spectrum of the mercury line 2536.7 lead to the conclusion that the number of rayless transformations in the collision of excited atoms with slow moving atoms is not small compared with unity. In solids at high temperatures, the number of rayless quantum changes with liberation of electrons is large compared with the number of quantum changes taking place with the emission of light. W. E. G.

Photocatalysis. II. The Photosynthesis of Nitrogen Compounds from Nitrates and Carbon Dioxide. EDWARD CHARLES CYRIL BALY, ISIDOR MORRIS HEILBRON, and DONALD PRYCE HUDSON (T., 1922, 121, 1078—1088).

Photochemical Activity of the Triphenylmethanesulphonic Acids. EDWARD O. HOLMES, jun. (*J. Amer. Chem. Soc.*, 1922, 44, 1002—1008).—Solutions of magenta, malachite-green, methyl-violet, and crystal-violet which have been decolorised by sulphur dioxide when exposed to ultra-violet light of wave-length between 2200 Å.U. and 3300 Å.U. develop their original colours. Comparison of the absorption spectra of the original dyes and those which had regained the colour showed that the regained colour is due to the original dye, since the absorption spectrum is the same in both cases. It is shown that the reaction with sulphurous acid is reversible and consists of a decolorising action and a colorising action, the latter of which is photochemical. The large quanta of energy supplied by the ultra-violet light cause the photochemical action to increase and consequently a displacement of the position of equilibrium occurs. The equilibrium position can also be displaced by chemical and thermal means.

J. F. S.

A Determination of the Number of α -Particles per Second Emitted by Thorium-C of known γ -Ray Activity. ALLEN G. SHENSTONE and HERMAN SCHLUNDT (*Phil. Mag.*, 1922, [vi], 43, 1038—1047).—A direct comparison is made between the number of α -particles and the γ -ray activities of radium-C and thorium-C. α -Particles of ranges 8.6 cm. (thorium-C) and 6.96 cm. (radium-C) were counted by the wheel method devised by Rutherford, and accurate γ -ray measurements were made at the same time as the counts. The ratio $\alpha\text{Th-C}/\alpha\text{Ra-C}$ for equal γ -ray activities is not independent of the thickness of the wall of the γ -ray electroscope. Values of this ratio have been determined for thicknesses of lead ranging from 3.3 to 14.3 mm. For 3.3 mm., the ratio = 0.75. The γ -ray activities of thorium-C, radium-C, and a radium standard are given for the same range of thicknesses of lead. W. E. G.

The Interpretation of β -Ray and γ -Ray Spectra. C. D. ELLIS (*Proc. Camb. Phil. Soc.*, 1922, **21**, 121—128).—The researches of Rutherford and Chadwick have shown that the β -rays from a radioactive atom may be separated into a general and a magnetic line spectrum. The latter (cf. this vol., ii, 339) may be accounted for, if the γ -rays are the primary phenomena in the disintegration of the atom. The details of five different β -ray spectra (radium-*B*, radium-*C*, radium-*D*, thorium-*B*, thorium-*D*) lend support to this theory. The γ -rays are emitted during the movement of an electron from one stationary state in the nucleus to another. Some of these rays are absorbed in the electronic structure of the same atom and eject electrons with characteristic energies from the *K*, *L*, and *M* levels. When an electron arrives in a stationary state, in which it is not permanently stable, it is ejected from the nucleus. This electron, which has a variable kinetic energy, gives rise to the continuous or general spectrum. The numerical results obtained in this paper do not support the theory of Meitner (this vol., ii, 416), which affirms that the emission of a β -particle is the primary phenomenon in the disintegration of the atom.

W. E. G.

The Chemical Action of Penetrating Radium Rays. XIII. The Velocity of Formation and Equilibrium of Hydrogen Peroxide. ANTON KAILAN (*Monatsh.*, 1921, **42**, 387—398).—It is known that penetrating radium rays accelerate both the formation and decomposition of hydrogen peroxide (cf. A., 1912, ii, 10). Experiments have now been made to determine the equilibrium concentration of hydrogen peroxide in neutral solution and in solutions of varying degrees of acidity, since the stability of hydrogen peroxide is known to depend on the hydrogen-ion concentration. Experiments were made using the same samples of radium as in the previous work, at temperatures from 6° to 13°. No difference was found between Merck's perhydrol and ordinary hydrogen peroxide. At equilibrium the concentration of hydrogen peroxide, in gram equivalents per litre, was 0.005 and 0.001 when the hydrogen-ion concentration was 0.5 and 0.008 gram-ion per litre, respectively, whilst in neutral solution the equilibrium concentration was only 0.0006. Impurities present in ordinary distilled water lowered the equilibrium concentration in acid solution. The reaction velocity in either direction agrees with the equation $k=1/t \log a-b/a-b-y$, where *b* is the concentration at equilibrium, *a* the original concentration and *a-y* the concentration after time *t*. The values of *k* found at hydrogen-ion concentrations 10^{-7} , 0.008, and 0.5 were 7×10^{-3} , 0.7×10^{-3} , and 0.14×10^{-3} , respectively. The higher equilibrium concentration in acid solutions than in neutral solutions is due mainly to the retardation of decomposition of hydrogen peroxide in acid solution. It is calculated that for each primary β -particle, in neutral solution $3 \cdot 10^3$ and in acid solutions from 0.1 to 1.0*N*, $6 \cdot 10^3$ molecules of hydrogen peroxide are formed per second. A discrepancy between the velocity of formation found now and in 1911—1912 (*loc. cit.*) is discussed at length;

it may be due to physical changes in the glass apparatus brought about by prolonged exposure to radium rays.

The number of molecules of hydrogen peroxide formed is of the same order as the number of ion pairs from the total absorbed rays. This supports the view that there is a relation between ionisation and chemical action for β - and γ -rays as well as for α -rays. On the other hand, the ratio of hydrogen peroxide molecules decomposed to number of ion pairs is 1500:1 and the mechanism of decomposition appears to be different from that of formation. Some experiments on the decomposition of hydrogen peroxide in the light of a quartz mercury vapour lamp are also recorded. The velocity coefficient is much greater than with radium rays.

E. H. R.

Electrical Charges of Colloidal Particles and Anomalous Osmosis. II. Influence of the Radius of the Ion. JACQUES LOEB (*J. gen. Physiol.*, 1922, 4, 621—627; cf. this vol., ii, 354).—The rate of transport of water and the potential difference across a collodion-gelatin membrane separating solutions of the chlorides of potassium, sodium, and lithium from pure water, when the P_H is on the acid side of the isoelectric point of gelatin, vary inversely with the radius of the kations. At P_H 3.0 the influence of the three salts on the potential difference between the liquid and the membrane inside the pores of the gelatin is identical. The influence of the three salts on the potential difference across the membrane varies inversely as the relative mobilities of the kations, which suggests that this influence may be due to a diffusion potential.

C. R. H.

Effect of Hydrogen Pressure on the Electromotive Force of a Hydrogen-Calomel Cell. I. WILLIAM R. HAINSWORTH and DUNCAN A. MACINNES (*J. Amer. Chem. Soc.*, 1922, 44, 1021—1032).—The effect of pressure of hydrogen on the cell $H_2|HCl(0.1N)|HgCl|Hg$ has been investigated at 25° for pressures up to 400 atmospheres, and the change of *E.M.F.* by pressure calculated. The thermodynamic theory of the change of *E.M.F.* has been investigated and the expression $\Delta E = 0.02958 \log p + 6.56 \times 10^{-6}(p-1) + 3.7 \times 10^{-10}(p^2-1)$ obtained for the change of *E.M.F.* The calculated and the observed values have been compared and are as follows: 50 atmos. $\Delta E = 0.0504$ volt (0.0506), 100 atmos. $\Delta E = 0.0594$ (0.0598), 200 atmos. $\Delta E = 0.0688$ (0.0694), 300 atmos. $\Delta E = 0.0745$ (0.0753), and 400 atmos. $\Delta E = 0.0787$ (0.0797). The values in brackets are the calculated values. The values indicate that the thermodynamic theory as developed does not quite represent the measurements at higher pressures. This is probably due to the solubility of hydrogen, and it would appear that a correction to account for the solubility must be introduced.

J. F. S.

Carbon Monoxide-Oxygen Cell with Glass as Electrolyte. HARTMUT KALLMANN (*Z. Elektrochem.*, 1922, 28, 81—85).—The *E.M.F.* of cells containing mixtures of carbon monoxide, carbon dioxide, and oxygen on one side of a glass wall and air on the other

side has been determined at 717° under various pressures from 920–800 mm. It is shown that the *E.M.F.* for such a cell is expressed by the equation $E = 1.118 - 0.0707 \log p_{\text{CO}}/p_{\text{CO}} \cdot p_{\text{O}_2}^{1/2}$. The experimental values agree with the values calculated by the equation to within 0.9%.
J. F. S.

Use of Phthalate Solutions for Hydrogen Electrode Standards. EARLE T. OAKES and HENRY M. SALISBURY (*J. Amer. Chem. Soc.*, 1922, **44**, 948–951).—The proposal recently made to replace the calomel electrode as a standard of *E.M.F.* measurement by a hydrogen electrode made up in alkaline solutions of phthalates has been investigated; and it is found that when such a phthalate solution, of P_{H} value 6.0 as measured by indicators, is measured against a calomel electrode, the *E.M.F.* slowly drifts in forty-eight and a half hours from about 0.57 to 0.66, that is, to a P_{H} value of 7.01. This change in the P_{H} value was confirmed by indicators. It is held that the change in the P_{H} value is due to a change in the phthalate itself, probably a reduction, and not to reduction of impurities, for the purification had been such as to exclude the large quantities of impurities necessary to bring about the observed change. If impurities are responsible for the change in *E.M.F.* their action must be that of catalysts reducing the phthalate. Hence, unless specially pure, phthalate solutions are unsuitable as standards for measuring hydrogen-ion concentrations.
J. F. S.

Instability of Phthalate Potentials. WM. MANSFIELD CLARK (*J. Amer. Chem. Soc.*, 1922, **44**, 1072–1073; cf. preceding abstract).—The author has re-examined some phthalate hydrogen electrodes over a period of twenty-four hours and has been unable to find a drift of more than 1.0 millivolt, and in the last fifteen hours of the observation the drift was less than 0.00005 volt. There was also no change in the Sørensen value. The author is therefore unable to explain the very large changes observed by Oakes and Salisbury (*loc. cit.*).
J. F. S.

New Conceptions of Electrolytes. III. The Hydration of the Hydrogen Ion. ERLING SCHREINER (*Z. anorg. Chem.*, 1922, **121**, 321–334; cf. A., 1921, ii, 425, 498).—The author measured the *E.M.F.* of the hydrogen ion in 0.001, 0.002, and 0.005 molar hydrochloric acid in potassium chloride solutions the concentration of which varied from 0.001 to 3.0 molar. The quinhydrone electrode was employed (cf. Büllmann, A., 1921, ii, 372). These measurements do not give the concentration of hydrogen ions, but the activity of these ions which is associated only with the non-hydrated ions. The author has given the method of calculation in a previous communication (A., 1921, ii, 425). The results obtained were fairly constant and showed that 8–9 molecules of water are on an average associated with each hydrogen ion, a result in good agreement with that found by Bjerrum (*Z. anorg. Chem.*, 1920, **109**, 275).
W. T.

Electrochemical Oxidation of Organic Compounds.

ERICH MÜLLER (*Z. Elektrochem.*, 1922, **28**, 101—106).—A theoretical paper in which the author discusses the views put forward by Fichter (this vol., ii, 23) and makes his own views clear. It is shown that the assertion of Fichter, that his views on electrochemical oxidation and those of the author are in agreement, is not true. Whilst the author assumes the discharge of anions, Fichter supports the view of a primary formation of oxygen and the accompanying purely chemical oxidation. Fichter assumes the intermediate formation of peroxides or peracids, the method of formation and decomposition of which is not explained, and consequently furnishes nothing toward the explanation of electrochemical processes. Such per-compounds may be prepared chemically, but their decomposition is to some extent different from what would be expected from the view that they are formed at the anode and from the products of electrolysis. The author accepts the formation of intermediate hydroxy-compounds, formed by the discharge of anions, which decompose in a manner in keeping with facts, and present a mechanism for both the chemical and electrochemical oxidation. The view of Fichter that the substitution of discharge processes at the anode by purely chemical oxidations contributes to the understanding of the course of electrolytic oxidations is incorrect.

J. F. S.

Magnetic Properties and Atomic Structure. B. CABRERA (*Anal. Fis. Quím.*, 1922, **20**, 92—97).—A theoretical discussion of the relation between magnetic properties and atomic structure in the metals of the iron group. It is suggested that the N -electrons may be divided into two concentric subdivisions, the more superficial, N_2 , being the valency electrons and those beneath, N_1 , those with which magnetic properties are associated. It is supposed that the relation between magnetic moment and the number of N_1 electrons is periodic. The magnetic moment reaches a maximum for bivalent manganese and trivalent iron with five N_1 -electrons and diminishes to zero with increasing number of N_1 -electrons.

G. W. R.

Magnetism and Atomic Structure. II. The Constitution of the Hydrogen-Palladium System and other Similar Systems. A. E. OXLEY (*Proc. Roy. Soc.*, 1922, [A], **101**, 264—279; cf. A., 1921, ii, 82).—The specific magnetic susceptibility of palladium black, the value of which is $+64.6 \times 10^{-7}$, is considerably reduced by occluded hydrogen, the greatest reduction observed being about 75% to $+14.7 \times 10^{-7}$ when great care was taken to prevent loss of hydrogen after charging. The specific susceptibility gradually returns to its original value as the occluded hydrogen escapes. If the occluded hydrogen were in the atomic state, it would be expected to increase the specific susceptibility. On the other hand, molecular hydrogen, liquid or gaseous, being diamagnetic, should reduce the specific susceptibility by a calculable amount. The maximum reduction on this assumption would be 0.19×10^{-7} , an amount insufficient to account for the observed

reduction. The experimental results may be accounted for on the assumption that a loose compound is formed, possibly having the composition Pd_2H . That the presence of such a loose compound should lower the magnetic susceptibility is not surprising when it is remembered that the carbonyls of iron and of nickel are both diamagnetic. It is noteworthy that a loose compound of the composition PdH would form a system of two nuclei with an electronic system resembling that of silver, the atomic number of which exceeds that of palladium by one. Such a system would be expected to have a magnetic susceptibility very nearly equal to that of silver, -1.8×10^{-7} . If half the palladium atoms were in this form, its susceptibility would be reduced by about 50%. The difference between the occluding powers of amorphous and crystalline palladium is discussed. The slow diffusion of hydrogen into crystalline palladium can be understood when its face-centred cubic lattice is taken into account.

When manganese is fused in an atmosphere of hydrogen, or is deposited electrolytically, it becomes ferromagnetic with a specific susceptibility $+2000 \times 10^{-6}$, its normal value being $+11.0 \times 10^{-6}$. It may be supposed that the presence of hydrogen produces with the manganese, atomic number 25, a certain number of electron systems similar to that of iron, atomic number 26, and that these are the cause of the observed ferromagnetic properties. This suggestion is discussed in connexion with the Lewis-Langmuir theory.

E. H. R.

Determination of High Temperatures by Effusion of Gases.

YOHEI YAMAGUCHI (*J. Chem. Soc. Japan*, 1922, **43**, 1—21).—The author has studied the relations between pressure and temperature on the passage of a gas through a capillary tube of quartz. (1) The increase of pressure necessary to pass the same quantity of air through a quartz capillary tube at various temperatures was investigated. A linear relation, $\log p = a \log T + b$, holds between the pressure (p) and the absolute temperature (T) within the range 17° to 1205° . This, however, cannot be applied to the measurement of temperature, owing to the complexity of the apparatus necessary, and the difficulty of regulating the pressure. (2) Using the same capillary, the relation between the effluent velocity of the transpired air and its temperature has been investigated, no simple linear relation being found for the temperature interval 20.5° to 1046° . (3) The relation was investigated between temperature and the effluent velocity of air as measured by its effusion through a small hole in the round bottom of a quartz tube. A linear relation, $t = a + b\sqrt{T}$, holds between the effluent time (t) and the absolute temperature (T) of the tube heated by an electric furnace at temperatures from 17° to 1125° which can be applied to the measurement of high temperatures.

K. K.

Considerations on Cooling and Heating Curves. R. ARIANO (*Gazzetta*, 1922, **52**, i, 246—261).—The author considers the courses followed normally, that is, when no transformation occurs, during

cooling or heating, by: (1) the temperature-time curve, and (2) the curve representing the difference between the temperature of the substance examined, T , and that of a second substance, chosen for comparison, T_1 , as a function of the latter temperature, $T - T_1 = f(T_1)$.

As regards the temperature-time curve, it is first shown theoretically that Newton's law expresses, without appreciable error, the transference of heat between furnace and sample in the case of the cooling of metallic samples of small dimensions. Analogous reasoning applies in the case of heating curves. Cooling curves are exponential in form and are defined by equations of the type $T = k + Ae^{-a\theta/Pc}$, whilst heating curves are probably represented by equations of the form $T = k + Ae^{-a\theta/Pc} + f(\theta)$, θ being the time, a the product of the coefficient of external thermal conductivity and the surface, P the weight of the sample, and c its mean specific heat. The results of cooling and heating experiments with silver show that A in the former of the two above equations is sensibly identical with the temperature at the beginning of the cooling (t_1), k being consequently nearly zero in magnitude.

As regards the differential curve, $T - T_1 = f(T_1)$, it is shown theoretically and confirmed experimentally that, if the sample chosen for comparison undergoes no transformation in the interval of temperature chosen, this curve exhibits a maximum almost at the beginning of the cooling. In order that the normal course of the curve may be as nearly rectilinear as possible, the specific heats of the two samples and their mode of variation with temperature should be almost coincident; the same should hold for the values of the ratio, weight : surface, for the coefficients of linear expansion and their variation with temperature, and for the coefficients of external conductivity.

T. H. P.

Revision of the Entropies of the Elements. GILBERT N. LEWIS and W. M. LATIMER (*J. Amer. Chem. Soc.*, 1922, **44**, 1008—1017).—A theoretical paper in which on the basis of new determinations of the specific heats of many elements the entropy has been recalculated. By means of the equation $S = 3/2R \log_e W + 25.70$ the entropies of nitrogen, oxygen, fluorine, sodium, chlorine, potassium, calcium, zinc, bromine, cadmium, iodine, caesium, and mercury have been calculated in the state of monatomic gas. In four cases where the experimental values are of sufficient accuracy a comparison is made between the experimental values and the calculated values for 25° and 1 atm. The following are recorded where the values in brackets are those calculated: helium 29.2 (29.8), argon 36.4 (36.7), cadmium 40.0 (39.8), and mercury 41.3 (41.5). A table of entropies at 1 atm. and 25° has been drawn up and is appended to the paper.

J. F. S.

Electrically Heated Apparatus for the Determination of Melting Points. UYTENBOGAART, jun. (*Chem. Ztg.*, 1922, **46**, 493).—The apparatus consists of a circular glass tube at the bottom of which a platinum resistance is placed; it is filled with water, glycerol, or sulphuric acid (d 1.84). The liquid, warmed by the

platinum resistance, rises uniformly in the two limbs and the currents, joining at the top, pass down a central tube which contains the thermometer and substance, with a kind of whirlpool motion. For temperatures up to 75°, an ordinary 4-volt accumulator is sufficiently powerful, but for higher temperatures it is preferable to use the lighting current with a suitable resistance in the circuit. H. W.

Binary Liquid Mixtures. GERHARD C. SCHMIDT (*Z. physikal. Chem.*, 1922, **101**, 286—291).—An answer to the criticisms of Faust (this vol., ii, 423), Cassel (this vol., ii, 424), and Schulze (this vol., ii, 424) on the views expressed by the author in connexion with Dolezalek's theory of binary liquid mixtures (this vol., ii, 119).

J. F. S.

Heat of Vaporisation and the Difference, $m'-m$, of the Specific Heats at the State of Saturation for Argon, Oxygen, Nitrogen, and Hydrogen. E. MATHIAS, C. A. CROMMELIN, and H. KAMERLINGH ONNES (*Compt. rend.*, 1922, **174**, 1395—1397; cf. A., 1921, ii, 256).—A mathematical paper in which formulæ are given for the four gases by means of which it is possible to calculate the latent heat and hence the reduced heat of vaporisation, and also the difference between the specific heats of the saturated vapour and the saturated liquid. W. G.

The Vapour Pressure of Hydrogen and New Thermometric determinations in the Domain of Liquid Hydrogen. J. PALACIOS MARTINEZ and H. KAMERLINGH ONNES (*Arch. Néerland.*, 1922, [iii], **6**, 31—39).—Cath and Onnes (A., 1918, ii, 218) could not explain the disagreement of their results for the vapour pressure of hydrogen in the neighbourhood of its boiling point with the values found by Keesom and Onnes. The former authors determined the temperatures directly by means of a helium thermometer, whilst the latter employed a platinum resistance thermometer, the values obtained being reduced to the hydrogen and helium thermometers by Onnes and Holst. The present authors redetermined these measurements between the absolute temperatures 14·10° and 20·63°, particular attention being directed to the boiling point of hydrogen, an important point on the temperature scale. The boiling point was found to be 20·35°, a value agreeing with that found by Keesom and Onnes; for points immediately below the boiling point the results agreed with those of Cath and Onnes. All the results are given in a table. The calibration of platinum thermometers is also described. W. T.

The Distillation of Greatly Frothing Liquids. KLÄNHARDT (*Chem. Ztg.*, 1922, **46**, 493).—The difficulty in effecting the distillation of foaming liquids, such as aqueous solutions of soap or saponins, can be easily overcome by passing a current of compressed gas (such as carbon dioxide) over the surface of the liquid. A suitable apparatus for the introduction of the gas is a tube terminating in a ball of 2 cm. diameter in which three series of holes about

1 mm. in diameter are blown. Distillations with steam can also be smoothly effected in this manner. H. W.

Maintenance of the Adiabatic Condition in Calorimetry. FREDERICK BARRY (*J. Amer. Chem. Soc.*, 1922, 44, 899—937; cf. A., 1920, ii, 734).—An extensive series of measurements is described, which were undertaken to establish a standard of accuracy for the calorimetry of slow processes. It is shown that to attain the accuracy defined by the loss or gain of 0.5 gram-calorie in ten hours, the following precautions are necessary and sufficient. A closed calorimetric system must be used which allows no measurable evaporation into the insulating air gaps, and these gaps must be dry. The outer environment must be thermostatically controlled, whenever there are conducting parts which pass directly from the calorimeter to the surrounding air, with a precision characteristic of the system employed. This is about $\pm 0.2^\circ$ when encased mercury thermometers are used. It is always advisable to keep the environmental temperature close to the calorimetric within the characteristic range of negligible direct leakage; and it is necessary to correct for fluctuation beyond this range. For this control, a constant temperature chamber is essential. A convenient structure for this purpose is described. With the usual system of about 1000 cal. heat capacity the air gaps should not be less than 40 mm. wide and there is no advantage in larger gaps. With a 40 mm. gap, when the temperature fluctuates within $\pm 0.01^\circ$ and the environment within 0.2° of the calorimetric temperature, the system is practically adiabatic within the error defined above, the total leakage being little more than three times that of a perfect vacuum gap. The leakage rate in such systems is the same in both directions and for heads below 0.1° is proportional to the head. Corrections for imperfect adiabaticity due to bath fluctuation may be made on this basis, since transmission lag has no measurable effect on the leakage. A calorimeter based on the above-named conditions is described. With this calorimeter, the initial conditions are perfectly adiabatic; and the total calorimetric lag for the temperature changes characteristic of long-continued operations is negligible. It is permissible and advantageous to stir the closed system reciprocally. In the system described, this may be done without significant loss of heat. The heat of stirring is developed identically in different assemblages and at the same rate for any one speed, whether stirring is intermittent or continuous. It may vary at low speeds with the fourth and higher powers of the speed. The continuous stirring necessary to ensure mixture in chemically reacting systems is productive of negligible error; and similar stirring at low speeds throughout determinations is shown to be practicable, although in very protracted operations intermittent stirring is better. Strictly uniform speeds are necessary for long, continuous stirring, but not for intermittent stirring, nor for continuous stirring during mixture if the approximate rates of heat production under different speeds be known for the system used. The open calorimeter is unavailable in protracted calori-

metry, since, even with an initially dry gap, evaporation leakage is inconstant as the result of uncontrollable variability in prior conditions. This is due in great part to the condensation of water vapour in the air gap on the walls of the calorimeter, which occurs on clean as well as on dirty surfaces, and is probably due to the simple adsorption of moisture, an effect which can be neither eliminated nor controlled even under approximately isothermal conditions. With change of temperature, its effects are greater and still more irregular, and in gaps saturated with water are complicated in the extreme. The close system is best for all types of calorimetry; but in the adiabatic measurement of swift reactions the open calorimeter may be safely used, either with dry surfaces in an initially dry gap, or with invisibly wetted surfaces in a saturated gap which holds no water, provided always that correction for aggregate thermal disturbances due to evaporation leakage be made by interpolation from the results of observations taken before and after the reaction period. The initially dry gap, with which leakage is less and more nearly uniform, is preferable in such measurements; but correction for imperfect adiabaticity is safer with the saturated gap. Under all heads less than 1° , the total leakage from a closed calorimeter system in a dry gap shows slight divergence from Newton's law, since the convection increases very slowly with increase in gap width, and varies as powers of the thermal head less than two except when the gap is unusually wide. Similar leakage from the open system is irregular, but yields itself to a rough analysis which gives some indication of the character of the thermal disturbances and the magnitude of the uncertainties characteristic of the merodiabatic measurements which have yielded the bulk of the present thermochemical data. J. F. S.

Calorimetric Researches : (I) The Standardisation of a Calorimetric System ; Comparison of the Heats of Combustion of the Substances Used for Standardisation : Benzoic Acid and Naphthalene. P. E. VERKADE, J. COOPS, jun., and H. HARTMAN (*Rec. trav. chim.*, 1922, **41**, 241—277).—A discussion of previous work on this subject and its bearing on the authors' experiments. Standardisation of the apparatus was effected as the result of thirty-seven experiments in the combustion of benzoic acid and thirty-six in the combustion of naphthalene. A detailed description is given of the apparatus used, the method of carrying out the experiments and the corrections applied to the results. Adiabatic calorimetry was discarded in favour of the ordinary methods for various reasons; there is little difference in the results obtained by the two methods (cf. Dickinson, A., 1914, ii, 802, and Swientoslawski, A., 1921, ii, 679). The ratio between the values obtained for the two substances used agrees with that of Dickinson, the figure given by Swientoslawski (A., 1918, ii, 32) is criticised on both theoretical and practical grounds. In order to carry out an absolute standardisation of a calorimetric system, a standard substance should be chosen the heat of combustion of which is known with the greatest possible accuracy. The results obtained

by the authors show that benzoic acid should be preferred to naphthalene, as it gives more constant results, is easily obtained pure, and is practically non-hygroscopic. Naphthalene, on the other hand, does not undergo complete combustion in many experiments, and also loses weight by evaporation; figures are given in illustration of the latter objection. Other substances suggested for the purpose are not so easily obtained pure as is benzoic acid. The value finally recommended as the standard ratio of the heats of combustion of naphthalene and benzoic acid is 1.5203. H. J. E.

Heat of Combustion of Lactic Acid. OTTO MEYERHOF (*Biochem. Z.*, 1922, 129, 594—604).—The author has determined the heat of combustion of lactic acid (in dilute aqueous solution) from the heat of combustion of zinc lactate, the heat of dilution of lactic acid, the heat of neutralisation of zinc oxide and lactic acid, and the heat of solution of zinc lactate in water. The value found for 1 gram of lactic acid is 3615 cal. H. K.

Heat of Solution of Zinc in Hydrochloric Acid. THEODORE W. RICHARDS and THORBERGUR THORVALDSON (*J. Amer. Chem. Soc.*, 1922, 44, 1051—1060).—A new apparatus is described, suitable for dissolving substances in a comparatively small excess of liquid and providing a conveniently large heat capacity for the reception of the heat. With this apparatus the heat of the reaction $\text{Zn} + 2(\text{HCl}, 200\text{H}_2\text{O}) = \text{ZnCl}_2, 400\text{H}_2\text{O} + \text{H}_2$ was found to be 36.32 Cal. (or 151.8 kilojoules) if the hydrogen is dry, and 36.07 Cal. (or 150.8 kilojoules) if the hydrogen is moist at 20°. The total energy change, U , of this reaction is therefore 36.90 Cal. (or 154.2 kilojoules) at 20°. The heats of dilution of factors and products needed for the calculation of the above results were determined experimentally. The temperature-coefficient of the heat of solution of zinc in concentrated hydrochloric acid is shown to be negative and of considerable magnitude; even with diluted acid it probably amounts to -30 cal. per degree. J. F. S.

Heat of Solution of Cadmium in Hydrochloric Acid. THEODORE W. RICHARDS and SETSURO TAMARU (*J. Amer. Chem. Soc.*, 1922, 44, 1060—1066).—Using the apparatus previously described (cf. preceding abstract), the heat of the reaction $\text{Cd} + 2(\text{HCl}, 200\text{H}_2\text{O}) = \text{CdCl}_2, 400\text{H}_2\text{O} + (\text{H}_2)_p$ is found to be 17.23 Cal. (or 72.0 kilojoules) at 20° if the hydrogen is dry, and 16.98 Cal. (or 71.0 kilojoules) if the hydrogen is moist at 20° and 760 mm. The total energy change, U , of the reaction is therefore 17.81 Cal. (or 74.4 kilojoules) at 20°. The temperature-coefficient of the heat of solution of cadmium in concentrated hydrochloric acid is shown to be negative -71 cal. per degree, and probably with diluted acid it amounts to -30 cal. per degree. All the necessary factors used in the present calculation were determined experimentally. J. F. S.

Heat of Transformation of Austenite into Martensite and of Martensite into Pearlite. NOBUO YAMADA (*Sci. Rep. Tohoku Imp. Univ.*, 1922, 10, 453—470).—The heat of dissolution of carbon

in iron and also the heat content of austenite and pearlite are determined in a series of carbon steels ranging from 0.38—1.74% carbon. The heat of dissolution of carbon in iron is shown to be a linear function of the carbon content of the steel and amounts to 1130 cal. per gram of carbon. By combining the results obtained with those of Meuthen (cf. A., 1913, ii, 385) for the total heat of transformation, the heat of allotropic transformation of austenite to martensite is calculated. It is a linear function of the carbon content and has a maximum value of 5.6 cal. per gram for eutectic steel. For a carbon content of 0.38% carbon, the value is 2.4 cal. and for 1.74% carbon 4.8 cal. Martensite has a definite heat content corresponding with the quantity of dissolved carbon, a definite crystalline form stable at ordinary temperatures, characteristic physical properties as compared with austenite or pearlite, and must therefore be considered as an independent phase. It is shown that troostite, sorbite, and pearlite have the same specific heats within the limits of experimental error. It is confirmed that the transformation of austenite into pearlite involves the intermediate production of martensite.

J. B. F.

Some Remarkable Properties of Gases. FRIEDRICH WÄCHTER (*Z. anorg. Chem.*, 1922, **121**, 225—239).—A theoretical paper which derives from Naumann's formula the lowest value for the molecular weight of gases to enable them to remain in the atmosphere of a planet. Hydrogen can exist in our atmosphere, but not in the atmosphere of the moon. In the atmosphere of the sun, a gas with a molecular weight 1/1864 can remain. The author identifies these imponderable molecules with the light ether, which he thus regards as a very light gas susceptible to the Lorenz contraction. He concludes from the above that the apparent mass of electrons as found by Kaufmann and Bucherer is merely a proof of the Lorenz contraction in the ether; thus the idea of apparent mass in the interior of the atom is not justified. Instead of the idea of β -particles or atoms of electricity, the author suggests the atoms of the lightest ponderable gas of invariable mass, and on this hypothesis finds that the enormous velocities in the inner atom such as Bohr calculated for the Rutherford atom are unnecessary. These results are also applied to the radioactive atoms. An earlier paper dealt with the astronomical implications, such as the Michelson's experiment, the abnormal motion of mercury, and the displacement of spectral lines (*Astron. Z.*, 1921, **4**, 62). W. T.

Specific Characters of Hydrolytic Decomposition. S. T. J. TROMP (*Rec. trav. chim.*, 1922, **41**, 278—295; cf. Schoorl, A., 1920, i, 531).—A study of the specific volume occupied by the molecule of water which is fixed in various reactions. An examination of esters, cyanides, carbylamines, nitro-compounds, anhydrides, ethers, oximes, amides, and acetals shows that the volume of the water taken up varies little in one type of reaction, but considerably from one type to another. This method of investigation shows that neither cyanides nor carbylamines should be regarded as esters of hydrocyanic acid and also provides a means of dis-

tinguishing between nitrites and nitro-compounds, also between cyanides and carbylamines. The values obtained for anisole and phenetole show that these substances are esters of phenol rather than ethers, and the view of saccharose as an ether is corroborated. Comparison of the specific refraction of fixed water with that of water as such gives no such trustworthy data as the discrepancies are considerable and numerous.

H. J. E.

Thin Films of Binary Mixtures (Glycerides). (MILLE) PAULE COLLET (*J. Phys. Radium*, 1922, [vi], 3, 128—132).—Mixtures of various glycerides in benzene solution are applied to the surface of water and determinations made of the surface covered. No additive relationships were found for binary mixtures, the curves always showing either maxima or minima. These maxima and minima appear to correspond with changes of physical state and with a simple ratio of the number of molecules of the two constituents. On the assumptions that all the molecules are associated at the maximum and minimum value of the area covered and that the free molecules of the glycerides cannot exist side by side, it is possible to derive theoretical curves which are in good agreement with the experimental results.

W. E. G.

Examination of the Compression Equation of Liquids by means of the Data of Amagat and Bridgman. H. CARL (*Z. physikal. Chem.*, 1922, 101, 238—268).—The compression equation put forward by Wohl (this vol., ii, 117) which has the form $e^{(v_0-v)/A} = 1 + (p-p_0)/(k+p_0)$; $v_0\alpha = A$; $k = br - c$ has been tested by means of Amagat and Bridgman's isothermals for twelve liquids. In the expression, K is a pressure quantity for the internal pressure of liquids and α a constant, which expresses the influence of different initial resistances and the distances of molecules and atoms in the liquid. Every normal liquid has a constant α value over the whole temperature range 0—80°, and the K value in the limit is sufficient in the case of normal liquids for rising temperature in the above equation. Water exhibits here the known anomaly in its volume change up to pressures of 2500 kg./cm.² At 50°, it shows first a regular behaviour with increasing temperature. The constant α is constant over the whole range of temperature, but the K values cannot, as is to be expected from the foregoing, be brought into a linear equation, since from 50° downwards K decreases with increasing temperature. Similar anomalies are shown by ethyl alcohol for lower pressures, according to Amagat's data for 1—1000 atmos., α is not constant over the temperature range 0—180°, and K does not decrease so regularly with increasing temperature, as is shown to be the case for all normal liquids. The agreement between the v value calculated by the above equations and those from Amagat's isothermals is excellent. A mean divergence of 0—2 units is found in the last place of decimals, the error of experiment being 6 units. Larger divergences have been shown from a series of differences to be due to experimental errors. In the case of Bridgman's results for higher pressures, the agreement has a mean divergence of 0.5%. A reduced form of the equation, in

which the pressure values are expressed as fractions of p_k , has been obtained. By inserting $K/p_k = b\tau/p_k - c/p_k = 1/\beta(\tau - \gamma)$ the equation $(v_0 - v)/v_0 = \alpha \log_e[1 + \beta(\tau - \pi_0)/(\tau - \gamma - \pi_0/\beta)]$ is obtained, and on placing $p_0 = 0$ in this the simplified form $(v_0 - v)/v_0 = \alpha \log_e[1 + \beta\tau/(\tau - \gamma)]$ results. A table containing the values of α , β , and γ and the critical data of the substances examined is included in the paper.

J. F. S.

Negative Viscosity. ADOLPH I. RABINOVICH (*J. Amer. Chem. Soc.*, 1922, 44, 954—964).—The term negative viscosity is applied to the viscosity of solutions which have lower viscosities than the solvents. With the object of explaining negative viscosity, the author discusses a large number of factors which may determine the viscosity of solutions. The factors considered are, (a) viscosity of the solvent, (b) viscosity of the solute, (c) solvation, (d) formation of complex ions and molecules, (e) the electric fields of the ions, (f) depolymerisation of associated solvents, (g) electrostriction, and (h) depolymerisation of the solute.

J. F. S.

Significance of Surface Tension Phenomena for the Dairy Practice. OTTO RAHN (*Kolloid Z.*, 1922, 30, 341—346).—The author has studied the formation of the foam which accompanies the pouring of skim milk from the cream separator. It is shown that foaming is due to a reduction of the surface tension caused by the accumulation in the surface of an albuminous substance which passes into the walls of the foam. These walls are shown to contain a solid substance, and the drying of such material is shown to be similar to the drying of many albuminous substances inasmuch as it is irreversible. It is probable that this solid albuminoid is the main constituent of the skin which forms on the surface of milk which has been heated at 60° or above. On the basis of the foregoing, the formation of whipped cream is explained as follows: on whipping cream a network of foam is produced, which is stiffened by the solidified fat so that it does not fall like the foam but retains its form. A microscopic examination of whipped cream shows that it consists of a solid structure of albumin penetrated by layers of solid fat which has the same form as the foam. On warming whipped cream above the melting point of the fat, the structure collapses, but remains somewhat frothy because the albumin cannot melt. Butter formation is explained as follows: The cream contains much of the foam producing albumin; this surrounds the particles of fat. On churning, a large volume of air is entrapped by the cream, so that the surface is very much enlarged. The albumin passes into the walls of the foam and takes the fat with it, so that a foam rich in fat stands above a liquid poor in fat (buttermilk). The fat particles lie very close together in the foam walls and are compressed by the surface pressure to form conglomerates of fat particles. This causes the albumin in the walls to solidify and further churning breaks up the foam and mixes the fat conglomerates with the solid albumin to form lumps of butter. It is shown that butter may be formed at 34°, but not at 39°.

J. F. S.

Theory of Adsorption Processes. M. POLANYI (*Z. Elektrochem.*, 1922, **28**, 110—111).—A criticism of Eucken's paper (this vol., ii, 262) on the theory of adsorption processes. J. F. S.

The Adsorption Problem. Sorption of Vapours by Charcoal. BROR GUSTAVER (*Koll. Chem. Beihefte*, 1922, **15**, 185—338).—The sorption isotherms for water, alcohol, and acetic acid vapours have been determined at 20°, 25°, 30°, and 35° with animal charcoal as adsorbent. These show that sorption is not a single process. The isotherms, which over the first portion are straight lines (water) or concave to the pressure axis (alcohol and acetic acid), show a well-defined inflexion at the point *O*. At this point the curve rises steeply and becomes convex to the pressure axis (water), or turns, and forms a wedge-shaped peak pointed outwards with respect to the pressure axis (alcohol and acetic acid). In the experiments, the vapours were added to the charcoal in small portions and the sorption equilibria were reached in all cases from both sides. The first part of the curves before the point *O* is reached represents a reversible process, since the points obtained by both methods lie on the same curve, whilst the equilibria points above the point *O* lie on parallel curves. Equilibrium is set up much more rapidly from below than from above. The experimental results are explained in much the same way as the sorption by gels. The sorption occurs first through adsorption and then from the point *O* mainly through condensation of liquid in the pores of the charcoal. The hysteresis observed is explained by the varying curvature of the meniscus during sorption and desorption. This depends on the fact that during sorption the walls of the charcoal pores are only slowly wetted, which in its turn is probably due to the presence of air. When the equilibrium is established from the under side, desorption, a rapid evaporation of the vapour from the charcoal, occurs, whereby the meniscus in the capillaries take on a greater curvature than previously and the equilibrium is rapidly stable. Sorption takes place slowly, because the radius of curvature of the meniscus decreases uninterruptedly while the walls are moist. Thereby a fresh condensation of vapour takes place and the establishment of equilibrium is delayed. The taking up of each portion of vapour added occurs at first by means of adsorption, and the adsorption equilibrium is established in about thirty seconds. Only in the case of charcoal which had been exhausted immediately before the experiment could the adsorption equilibrium be recognised. For this process the formula $k=1/tp_{\infty} \cdot [\log p/(p-p_{\infty}) - \log p_0/(p_0-p_{\infty})]$ is representative. The structure of charcoal has been discussed, and the radii of the capillaries have been calculated by means of the Trouton and Andersson formula. The taking up of alcohol and acetic acid vapour at 20° occurs in such a way that only capillaries of the same diameter are active. Approximately 75% of the total quantity is taken up at very low pressures, that is by capillaries of very small radius of the order $r < 6.2 \times 10^{-8}$ cm., which indicates that charcoal is possessed of an exceedingly fine structure. The

course of the taking up of water differs from that of the two other liquids inasmuch as the calculated radii are partly much greater and that the maximum volume is smaller. These two circumstances are explained by the smaller power of water to wet the charcoal and by the slow penetration of liquid into the pores of the charcoal. The surface of the charcoal has been approximated from the experimentally determined volume of the empty space of the charcoal and the calculated radii of the capillaries in which condensation occurs. The free surface of the capillaries of radius $r > 6 \times 10^{-8}$ cm. is about 600 sq. metres per gram of charcoal and the total area of all capillary walls > 3000 sq. metres. Trouton's sorption rule is shown to be incorrect and in its place it is shown that the volume taken up, and not the mass, of a given liquid is independent of the temperature and only a function of the pressure.

J. F. S.

More General Theory of the Adsorption of Solutions. WOLFGANG OSTWALD and RAMON DE IZAGUIRRE (*Kolloid Z.*, 1922, 30, 279—306).—A theoretical paper in which a large number of types of adsorption curves and formulæ have been collected and criticised. The view is expressed that every adsorption is at first a process of "unmixing," whereby a most concentrated solution, the absorbed solution, forms on the surface of the adsorbent whilst the equilibrium solution remains behind. Solutions of substances are adsorbed and not the "dry" dissolved substances. In opposition to the previously held views, the author postulates that the adsorption of the solvent, along with the dissolved substance, is an integral and equally important process in every adsorption from solutions. The assumption is made that the adsorption of the dissolved substances, without reference to the simultaneously adsorbed solvent, increases asymptotically up to the highest concentration in accordance with Boedecker's logarithmic formula. It is shown that this formula is not in keeping with the difference in concentration before and after adsorption, except in the case of very dilute solutions. Since the actual amount of adsorbed substance continuously increases, the difference in concentration ($c_0 - c$) must show a maximum and at the highest concentrations approach the zero value. It is shown that the whole of the adsorption curves may be constructed on the basis of the three assumptions made above and the simultaneous adsorption of the solvent. A number of formulæ are developed which give a quantitative relationship between adsorption and concentration on the basis of the above assumptions. The most important of these are: (1) $N/m(c_0 - c) = kc^n(100 - c)$ for the case where the adsorption of the solvent may be neglected; (2) $N/m(c_0 - c) = kc^n(100 - c) - l(100 - c)^a \cdot c$ for cases where the solvent is adsorbed directly by the adsorbent; (3) $N/m(c_0 - c) = kc^n[100 - c(1 + s(100 - c)^b)]$ for cases in which the solvent of the adsorbed solution is brought into the surface layer by the solvation of the dissolved substance; (4) $N/m(c_0 - c) = kc^n[100 - c(1 + s(100 - c)^b)] - l(100 - c)^a \cdot c$ for cases in which the adsorption of the solvent takes place by both methods

provided for in equations 2 and 3. In these equations N is the weight of solution in grams at the commencement, m the weight of adsorbent, c_0 the initial percentage composition of the solution, c the equilibrium percentage concentration of the solution, k , l , s , n , α , β are constants. A graphic method for evaluation of the constants is described. The second equation has been tested by means of the measurements of Schmidt-Walter for the adsorption of acetic acid and water by charcoal (A., 1914, ii, 542) and Gustafson's measurements of the adsorption of phenol and alcohol by charcoal (A., 1915, ii, 824) and a remarkably good agreement between the calculated and experimental values found.

J. F. S.

Relationship of Dolezalek's Theory of Solutions to that of Planck. H. CASSEL (*Z. physikal. Chem.*, 1922, **101**, 235—237).—In earlier papers, Wagner (A., 1920, ii, 596) has shown that Dolezalek's theory of solutions is a special case of Planck's theory of concentrated liquid mixtures, which is of no special importance. The present author shows mathematically that Wagner's conclusions are based on an error.

J. F. S.

Solubility. IV. Principle of the Specific Interaction of Ions. J. N. BRÖNSTED (*J. Amer. Chem. Soc.*, 1922, **44**, 877—898; cf. A., 1920, ii, 536; this vol., ii, 199).—It is shown that the activity coefficient of an ion may be determined by two factors, one of which is due to the salting-out effect of the salt solution serving as solvent and the other to electrical interaction between this ion and the ions of the solvent. Ions are uniformly influenced by ions of their own sign; their activity coefficients depend, therefore, only on the action of ions of the opposite sign and the salting-out effect of the solvent. The salting-out effect of a salt solution can be represented as a product of the salting-out effects of the separate ions. A number of relationships are derived for the solubility of salts at constant concentration of solvent solution and methods for determining the ratios of salting-out coefficients, coefficients of interaction, and activity coefficients. Thermodynamic relationships between activity coefficients and osmotic coefficients in solutions of constant total concentration and a relationship between solubility and freezing point are derived. The above theoretical conclusions and relationships are experimentally verified by solubility determinations in solutions of highly complex salts. A method of determining liquid junction potentials, on the basis of the present considerations, is put forward. In the case $0.1M KNO_3 | 0.1M NaNO_3$, a small concentration of silver nitrate is added and the *E.M.F.* of the cell $Ag | KNO_3(0.1M) AgNO_3(0.001M) || AgNO_3(0.001M) NaNO_3(0.1M) | Ag$ measured. The *E.M.F.* is given by $\pi_1 = RT \log_e f_{Ag}(NaNO_3) / f_{Ag}(HNO_3) + \pi_d$, where π_d is the junction potential and f is the activity coefficient of the Ag in the solution named. Also $\pi_1 = RT \log_e r_{(Na/K)} + \pi_d$ where $r_{(Na/K)}$ is the ratio of the salting-out coefficients of the sodium and potassium ions. This ratio can be determined from solubility measurements and consequently π_d may be calculated.

J. F. S.

Hydration of Electrolytes. S. ASCHKENASI (*Z. Elektrochem.*, 1922, 28, 106—110).—A theoretical paper in which the deviations of osmotic quantities of moderately and very dilute solutions have been collected together with the inflexion points of the experimental series. The Born-Fajans's polarisation-hydration hypothesis is discussed, and it is shown that the forces assumed in this hypothesis would produce a greater effect on the vapour pressure than is observed experimentally. In Fajans's method of calculating the heat of hydration, the influence of the dielectric constant on the work of dissociation is not taken into account. The author presents a chemical mechanism for hydration in salt solutions on the assumption that the associated liquid molecules of water are modified in the process. On the basis of this view, the author attempts to overcome the objections which have been offered to a step-wise hydration. J. F. S.

Calculation of the Osmotic and Activity Functions in Solutions of Uni-univalent Salts. J. N. BRÖNSTED (*J. Amer. Chem. Soc.*, 1922, 44, 938—948).—A theoretical paper in which from the point of view that the individualities of salt solutions have an origin similar to that of the individual salting out effect in the case of non-electrolytes, equations are deduced to govern the osmotic and activity functions of uni-univalent salts in pure and mixed solutions. These equations have been tested by means of the most accurate experimental results, which include measurements of depression of the freezing point, solubility, and electromotive force. Ratios of activity coefficients have been calculated on the basis of these equations in conjunction with the principle of the specific interaction of ions. J. F. S.

The Mutual Solubility of Liquids at High Pressures. JEAN TIMMERMANS (*Arch. Néerland.*, 1922, [iii], 6, 147—151).—This work completes the previous researches of Kohnstamm and Timmermans (*A.*, 1913, ii, 481). The apparatus employed is described in the previous communication.

A.—System water-*sec.*-butyl alcohol. The curve obtained for this system under high pressures is analogous to that of methyl ethyl ketone. With increasing pressure, the lower critical solution temperature (normally below the freezing point, -8.45°) is raised and the higher critical temperature, 113.8° , is lowered. Under a pressure of 830 kilos. they meet at about 65° , so that at these high pressures the liquids are miscible in all proportions. The values of dt/dp for both critical values are given. As demanded by theory, dt/dp for the freezing point of water was found to be less than for alcoholic solutions. The alcohol employed had a boiling point $99.50^{\circ} \pm 0.01$. $d_4^4 = 0.82263 \pm 0.00002$.

B.—System water-*isobutyric* acid (b. p. 154.35° , f. p. -47.0° , $d_4^4 = 0.96819$). The three-phase system crystallises at -3.0° at ordinary pressures. The critical solution temperature decreases more rapidly than the temperature of crystallisation as the pressure is increased so that the three-phase region soon disappears below the curve of solidification. By extrapolation, this was found to occur

at -8.2° under a pressure of 625 kilos. dt/dp was found to have a fairly constant value of 0.054.

C.—System water-phenol. dt/dp was found to be very small, but increases with the pressure 0.0036 to 0.0055.

D.—System cyclohexane-methyl alcohol. dt/dp has a fairly large value, but decreases with increasing pressure 0.0314 to 0.0191.

W. T.

A New Method of using X-Rays in Crystal Analysis.

GEORGE L. CLARK and WILLIAM DUANE (*Proc. Nat. Acad. Sci.*, 1922, 8, 90—96).—Certain alterations in the procedure of X-ray analysis is described. X-Rays belonging to the continuous spectrum are used instead of the line spectrum usually employed. For this purpose, the ordinary X-ray tube with a tungsten target is very convenient. The method has the advantage of flexibility, for short penetrating X-rays may be utilised, and with these it is possible to investigate crystals of high absorbing power. The method is suitable for both powders and large crystals. The values obtained for the distances between the planes of atoms are independent of the constants of any other crystal. The new methods of measurement are applied to the crystal structure of potassium iodide. One of the "peaks" obtained corresponds with the characteristic line spectrum of iodine.

W. E. G.

Isomorphism and Ionic Structure. HANS GEORG GRIMM (*Z. Elektrochem.*, 1922, 28, 75—81).—A theoretical paper in which author discusses the connexion between ionic structure and isomorphism. It is shown that ions built on the argon type differ considerably in size from those built on the neon type, whilst those of the krypton type are only little larger than those of the argon structure, and those built on the xenon type differ from those of the krypton type by intermediate amounts. Thus the expression $r_A - r_{Ne} > r_X - r_{Kr} > r_{Kr} - r_A$ represents the differences of the ionic radii. The characteristic gradation, of the lattice distances of similarly constructed compounds, in which the variable ions have the same number of exterior electrons, is conditioned by the gradation of the atomic radii. This is shown by the fact that the lattice distances of binary compounds is approximately given by the linear equation $r = \alpha a + \beta k$, in which a and k are the radii of the anion and cation respectively and α and β numerical constants. The relationship between isomorphism and ionic structure is shown by means of a table. It is shown that miscibility between ions with the neon and helium structure is unknown, although aluminium hydroxide and boric acid show a slight tendency. In the case of ions of the neon and argon structure, no crystallochemical relationship exists between O" and S", but F', Cl', Na' and K' exhibit miscibility in the case of large molecules and at high temperature, that is, under conditions where the influence of the difference in the radii is overcome. Complete miscibility is found between the ions of the argon and krypton type on account of the small difference between the atomic radii. Ions of the

krypton and xenon configuration also show close crystallo-chemical relationships, but the somewhat greater difference between the radii is expressed in the limited miscibility of potassium bromide and iodide. The ions of the argon and xenon structures form the end members of the well-known isomorphous series, but here also the difference in the ionic radii is shown in the existence of a large gap in the mixture series of potassium chloride and iodide. In general, the isomorphic relationships are determined by the characteristic gradation of the ionic radii in the case of similarly constituted ions.

J. F. S.

Fundamental Principles of Colloidal Chemistry. I. Electrical Charge of the Particles and New Conceptions of Micellæ. RICHARD ZSIGMONDY (*Z. physikal. Chem.*, 1922, **101**, 292—322; cf. this vol., ii, 129).—A theoretical paper in which it is shown that the conception expressed by the term micella has been entirely adopted in the various branches of colloidal chemistry. After considering a number of widely differing colloidal systems it is shown that the term micella has been applied to a series of different structures, which the author proceeds to characterise more exactly. Whilst the micellæ of Nägeli are crystalline individual particles, the term is employed by others for particles of very different constitution. It is pointed out that in most modern colloidal research the electrical charge has become the important factor, whilst the study of the particles themselves has been neglected. It is known that colloidal particles can be either crystalline primary particles (colloidal gold, vanadium pentoxide), secondary particles partly filled with liquid (stannic acid, purple of Cassius), or aggregates of molecules and ions of little stability combined with condensation nuclei (soap solutions). Both the primary particles and the secondary particles, depending on the electric charge can exhibit a considerable stability, and then they behave as multivalent ions of substances of very large molecular weight.

J. F. S.

Energy Principle of the Coagulation of Colloids, particularly of the Thermo-kinetic Mechanism. ROBERT FRICKE (*Z. physikal. Chem.*, 1922, **101**, 185—192).—A theoretical paper in which the heat change of some ideal coagulation processes of hydrosols and gas-dispersoids has been calculated on the basis of the thermo-kinetic shadow theory (A., 1920, ii, 740; 1921, ii, 387). The small heat changes obtained were compared with those calculated from the osmotic work and the work of lattice separation. All values are found to be of small dimensions. A short discussion of the different interdependent heat effects accompanying coagulation is entered on, and it is shown that the electrostatic discharge work must furnish a large portion of the heat of coagulation. The coagulation of discharged colloids in the case of hydrosols is mainly brought about by intermolecular forces (lattice forces) in the sense applied by Zsigmondy, and to a lesser degree by thermokinetic shadow action. In the case of gas-disperse systems the latter cause assumes an importance which may not be neglected in considering the actual precipitation.

J. F. S.

Physico-chemical Investigation of Colloidal Sulphur. G. ROSSI (*Kolloid Z.*, 1922, 30, 228—230).—In a previous communication (A., 1912, ii, 1037) it was shown that the electrical conductivity of a solution of colloidal sulphur containing sulphuric acid and sodium sulphate is lower than that of a similar solution containing no colloidal sulphur. It is now shown that the freezing point of a solution of the same kind as the above is lower after the colloidal sulphur has been removed than when the colloid is present. Further, if a colloidal sulphur solution containing sulphuric acid and sodium sulphate is dialysed to remove the electrolytes and the same amount of electrolytes added to the dialysed solution, then the electrical conductivity has the same higher value of the solution from which the colloid has been removed. It is thus shown that the action of the colloidal sulphur extends to sulphuric acid and sodium sulphate even when these are present in quantities larger than those required to stabilise the sol, provided that the excess of these substances was present when the colloid was formed. Further quantities of these substances added after the formation of the colloid are not affected by the colloid. It is therefore assumed that crystalloids, in the form of sulphuric acid and sodium sulphate, undergo a change in their physical nature in the presence of colloids, which is shown by a reduction in the electrical conductivity and the osmotic pressure.
J. F. S.

Cell for the Observation of Colloidal Solutions for Use with Substage Ultra-Condensers. FRED FAIRBROTHER (*J. Amer. Chem. Soc.*, 1922, 44, 1071—1072).—A cell is described for use in quantitative work on colloids with substage ultra-condensers. The apparatus eliminates the necessity of readjusting the optical arrangement every time a fresh portion of the colloidal solution has to be examined, as is the case with Burton's method of using a hemacytometer slide ("Physical Properties of Colloidal Solutions," 1921, p. 124). The apparatus consists principally of a circular brass block which has a circular hole in the centre and is bored at two diametrically opposite points to receive two copper tubes 2 mm. diameter. The holes are made so that they pass from the edge of the brass ring to the base. The bottom of the block, which is ground perfectly flat, is carefully cemented to a cover-glass and two holes are bored through the latter to meet the two holes in the base. The arrangement has the advantage of a reinforced cover-glass, and as both surfaces of cover-glass and slide are flat it is not necessary to cement the two together. The apparatus is held in position on the slide while being fixed in position by plasticine. To the copper tubes are fixed a delivery funnel and clip and an exit tube.
J. F. S.

Protective Colloids. XII. Gelatin as a Protective Colloid.
I. Colloidal Silver. A. GUTBIER, J. HUBER, and A. ZWEIGLE (*Kolloid Z.*, 1922, 30, 306—313; cf. A., 1921, ii, 312).—The effect of chloroform, toluene, acetone-chloroform, and ethyl acetate in small quantities on solutions of gelatin has been investigated and it is shown that all four substances act as excellent preservatives

for gelatin solutions. In the case of chloroform (10 c.c. per litre of solution), the solutions are stable, and only after weeks does a small quantity of a white precipitate separate; with toluene, a slight turbidity appears in a 5% gelatin solution after two weeks, but with more dilute solutions no turbidity is observed. The addition of 1% acetone-chloroform increases the stability of gelatin solutions, whilst with 0.5% of ethyl acetate a turbidity appears in two days; this turbidity speedily redissolves and the solution then remains unchanged for forty-two days. The efficiency of gelatin as a protective colloid for colloidal silver has been investigated, and it is shown that colloidal silver obtained by reduction with hydrazine may be protected with a small concentration of gelatin and the stability increased, but an excess of gelatin exerts a disturbing influence on the stability. The most stable and at the same time the most concentrated silver sol is prepared by reducing 10 parts of 0.1*N*-silver nitrate in 30 parts of 0.42% gelatin with hydrazine and immediately dialysing for six hours. This sol is reddish-brown by transmitted light and olive by reflected light; after dialysing for ten days a preparation was obtained which contained 40.45% of silver and was stable for three-quarters of a year. When the colloidal silver was produced by means of sodium hyposulphite, sols were obtained which were much less stable than those reduced by hydrazine. The effect of electrolytes on protected silver sols was investigated and it is shown that chlorides (hydrogen, sodium, magnesium, and barium) effect a rapid precipitation of the colloid; *N*-sulphuric acid produces a precipitation which is not complete in five days, sodium thiosulphate brings about a slow sedimentation, whilst sodium hydroxide and sodium carbonate in *N*-solutions are without coagulating action.

J. F. S.

Dependence of the Base Equilibrium in Permutite on the Concentration of the Surrounding Solution. A. GÜNTHER-SCHULZE (*Z. Elektrochem.*, 1922, **28**, 85—89).—The author has investigated the equilibrium in solutions of two salts of varying concentrations when shaken with permutite containing the same kations as the solution, with the object of ascertaining the concentration of the solution which is in equilibrium with the permutite. The experiments were carried out at 22° with 50% silver and 50% ammonium, 33.33% copper and 66.67% ammonium, and 4% lanthanum and 96% ammonium permutites, and the solutions contained the respective salts as nitrates. The results show that in a certain measure the demands of the law of mass action are fulfilled, but until the ionic concentration can be definitely determined it cannot be proved that the equilibrium is always set up in keeping with this law.

J. F. S.

The System *iso*Butyl Alcohol-Water. A. MICHELS (*Arch. Néerland.*, 1922, [iii], **6**, 127—146).—The *isobutyl* alcohol was specially purified, b. p. = 107.47°, d_4^{20} = 0.8197. The higher critical solution temperature was found to be 132.8° and the pressure 332.5 cm. of mercury, the critical concentration being 37% alcohol. Crystal-

lisation of the mixture prevented the realisation of the lower critical temperature. dt/dp for the critical temperature was found to be less than 0.01. The solubility of the alcohol in water was found to have a minimum value at 58° with a composition corresponding with 7% alcohol. The influence of impurities on the temperature of separation into two layers was very marked; 1% of ethyl alcohol lowered it by 50° , other impurities raised the temperature. Measurements were also carried out of the vapour pressure of the system; these are given in tables for $t-x$ and $p-x$; the latter shows a maximum but the composition corresponding with it could not be definitely fixed; it lies in the three-phase region. Further experiments are being carried out. W. T.

Influence of the Time Factor on the Interpenetration of Solids by Chemical Reaction. H. WEISS and P. HENRY (*Compt. rend.*, 1922, 174, 1421—1423; cf. this vol., ii, 207).—The time necessary for a given increase in the zone of interpenetration may be divided into two parts, namely, that required for the chemical reaction once the molecules of the two metals, in this case silver and antimony or copper and antimony, are in contact, and the time necessary for the molecules to traverse the layer already formed. The experiments tend to show that the duration of the chemical reaction is negligible as compared with the time required for the molecules to traverse the layer formed. The experimental points at each temperature fall on two distinct curves which are closely connected with the intensity of the blows of the hammer used to establish contact between the two metals. W. G.

The Rate of Evolution of Carbon Dioxide from Solution in Presence of Colloids. ALEXANDER FINDLAY and OWEN RHYS HOWELL (*T.*, 1921, 121, 1046—1052).

The Influence of Solvents on the Velocity of Formation of Quaternary Ammonium Salts. JAMES ALEXANDER HAWKINS (*T.*, 1922, 121, 1170—1176).



The Velocity of Absorption of Carbon Dioxide by Alkaline Solutions. PAUL RIOU (*Compt. rend.*, 1922, 174, 1463—1466).—Velocity curves have been plotted for the absorption of carbon dioxide by solutions of potassium carbonate under different conditions of concentration and temperature and with or without the presence of potassium hydrogen carbonate or potassium chloride. With potassium carbonate alone the velocity of absorption at first increases with the concentration very rapidly to a maximum and then slowly diminishes. The presence of potassium hydrogen carbonate causes a very marked diminution in the velocity of absorption, but the effect of potassium chloride is less marked. In every case, rise in temperature is accompanied by an increase in the velocity of absorption. Other factors being equal, the velocity of absorption is practically proportional to the concentration of the carbon dioxide in the air mixture. W. G.

Acceleration of the Solubility of Metals in Acids by Reducible Compounds. H. J. PRINS (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, **23**, 1449—1454).—The velocity of solution of a metal in an acid is enhanced by the presence of a reducible substance (cf. A., 1918, ii, 13). Nitrobenzene and benzaldehyde accelerate the rate of solution of iron, zinc, lead, tin, copper, nickel, aluminium, and silver in a large number of acids. In the presence of nitrobenzene, the rate of solution is increased up to 1000 times. The increase in the attack on the metal is accompanied frequently by an increased evolution of hydrogen. Benzaldehyde is selective in its action. Benzophenone retards the rate of solution of the metal and the rate of generation of hydrogen from zinc and an acid. Co-action has been realised in a system of four components—a metal, a salt of hydrochloric acid, a feeble acid (acetic acid), and nitrobenzene.
W. E. G.

Landolt's Reaction. ANTON SKRABAL (*Z. Elektrochem.*, 1922, **28**, 224—244).—The Landolt reaction which takes place in solutions of sulphurous acid and excess of iodic acid has been investigated. This reaction takes place according to the equations (1) $\text{IO}_3' + 3\text{SO}_3'' = \text{I}' + 3\text{SO}_4''$, (2) $\text{IO}_3' + 6\text{H}' + 5\text{I}' = 3\text{I}_2 + 3\text{H}_2\text{O}$, (3) $3\text{I}_2 + 3\text{SO}_3'' + 3\text{H}_2\text{O} = 6\text{I}' + 6\text{H}' + 3\text{SO}_4''$. Of the three partial reactions, 1 and 2 take place slowly with respect to 3 and determine the time of the reaction. Because the rapid reaction 3 follows the slow reaction 2 the sum of the equations of these reactions give equation 1 and consequently until the change of colour occurs equation 1 represents the total reaction. If reaction 2 is made more rapid than 1 by adding iodide to the mixture, then the whole of the sulphite will be oxidised according to 2 and 3 and reaction 2 alone will determine the time of the reaction. According to the measurements of Dushman (A., 1921, ii, 315), the expression $-d[\text{IO}_3']/dt = K_2[\text{IO}_3'][\text{I}']^2[\text{H}']^2$ represents this reaction. At the point of change, the whole of the sulphite has been oxidised, and therefore the initial concentration of this substance is a measure of the time required by reaction 2 to occur. If all the variable concentrations are chosen large with respect to the sulphite concentration, then reaction 2 takes place with constant velocity and the time of the reaction must be proportional to $[\text{SO}_3'']/[\text{IO}_3'] \cdot [\text{I}']^2 \cdot [\text{H}']^2$. This simple relationship has been experimentally verified. In this form, the Landolt reaction is particularly suited as a lecture experiment for demonstration of reaction velocity. The time law $-d[\text{IO}_3']/dt = K_1[\text{IO}_3'] \cdot [\text{SO}_3''] \cdot [\text{H}']$ is deduced for both reactions 1 and 2 individually. The constants K_1 and K_2 have been evaluated and tested over a wide range of concentration and shown to be in keeping with Dushman's experimental results (*loc. cit.*). In connexion with the foregoing experimental investigation, chemical kinetics are discussed generally. It is shown that chemical reactions only follow a definite time law over a limited range of concentrations and velocities. They are subject to a change of time law, so the constant K_2 of the iodate-iodide reaction becomes markedly smaller with increasing velocity, as is shown by

the present experiments and those of Dushman. The temperature quotients, $Q_{10}=K_{t+10}/K_t$, holds only for a definite total reaction. In the case of the total reactions, $\text{IO}_3' + 6\text{H}' + 8\text{I}' = 3\text{I}_3' + 3\text{H}_2\text{O}$ and $\text{IO}_3' + 6\text{HSO}_4 + 8\text{I}' = 3\text{I}_3' + 6\text{SO}_4'' + 3\text{H}_2\text{O}$, the value Q_{10} is 1.3 for the first and 0.85 for the second. These two temperature quotients are connected through the time law of the total reaction with the heat change of reaction $\text{H}' + \text{SO}_4'' = \text{HSO}_4'$. Both total reactions have their own heat changes, and the relationship between this and Q_{10} rules the adiabatic course of the reaction. J. F. S.

Kinetics of Reactions in which Halogen Atoms pass from the Inner Sphere of Activity to the Outer. H. FREUNDLICH and R. BARTELS (*Z. physikal. Chem.*, 1922, 101, 177—184).—The velocity of reaction of the following reactions has been determined quantitatively in some cases and semi-quantitatively in others at 0° and 25° . (1) $[\text{Cr}(\text{NH}_3)_5\text{Br}]^{++} + \text{H}_2\text{O} = [\text{Cr}(\text{NH}_3)_5\text{H}_2\text{O}]^{+++} + \text{Br}'$; (2) $[\text{Cr}(\text{NH}_3)_5\text{I}]^{++} + \text{H}_2\text{O} = [\text{Cr}(\text{NH}_3)_5\text{H}_2\text{O}]^{+++} + \text{I}'$;



The velocity has been compared with the reactions previously studied (A., 1914, ii, 448). The whole of the reactions are shown to conform with an equation of the first order. The velocity constants are dependent on the nature of the halogen and also on the nuclear atom; they increase very strongly, up to 100 times, in the order $\text{Cl} < \text{Br} < \text{I}$, and in the chlorine series they increase 50 times in the order $\text{Co} < \text{Cr} < \text{C}$. As far as the temperature coefficients were determined, they were found to be large; the constant A of the Arrhenius equation was about 10,000. This corresponds with the rule propounded by Halban, that apparent as well as true unimolecular reactions are characterised by their marked dependence on temperature. J. F. S.

Course of Simultaneous Reactions. RUDOLF WEGSCHEIDER (*Z. Elektrochem.*, 1922, 28, 99—100).—In connexion with the statement of Martin and Fuchs (*ibid.*, 1921, 27, 155) that in the chlorination of methane the concentration of methane and its chlorinated product are dependent on the concentration of one of these substances and quite independent of the time and the chlorine concentration, it is shown that this is not a special case, but is generally true for simultaneous reactions, which with regard to one of the reacting substances, H , are of the same order (n^{th}). If the reaction equation for two such actions are: $A_1 + nH \rightleftharpoons B_1$ and $A_2 + nH \rightleftharpoons B_2$, where A and B are complex molecules which do not contain H , the equilibrium constants are $K_1 = [B_1]/[A_1][H]^n = k_1/k'_1$ and $K_2 = [B_2]/[A_2][H]^n = k_2/k'_2$. On subtracting the second equation from the first, the equation $A_1 + B_2 \rightleftharpoons A_2 + B_1$ is obtained which must represent a possible equation of reaction. If the equilibrium constant of this is K , then $K = K_1/K_2 = k_1 \cdot k'_2/k_2 \cdot k'_1 = n_{12}/n_{12}'$, where n_{12} is the relationship between the equilibrium constants of the two reactions in the direction left to right and

n_{12}' in the opposite direction. This shows that the concentrations of all molecular types of A and B at equilibrium are determined by equations which do not contain the absolute values of the velocities.

J. F. S.

Mode of Action of Platinum in Oxygen-Hydrogen Catalysis and the Application of Titanium Sulphuric Acid for the Control of the Course of the Change. K. A. HOFMANN (*Ber.*, 1922, 55, [B], 1265—1274; cf. A., 1920, ii, 240; this vol., ii, 276).—Sensible quantities of hydrogen peroxide are only produced in the surrounding acid during the catalysis of dissolved molecular oxygen and activated hydrogen at platinum surfaces when the reducing power of the hydrogen-platinum electrode is abnormally reduced, when the liquid surrounding the pole contains large amounts of dissolved oxygen, and when the pole is as free as possible from oxides of platinum. It is highly probable that the hydrogen-oxygen catalysis proceeds normally in two stages (1) $O_2 + 2H = H_2O_2$ and (2) $H_2O_2 + 2H = 2H_2O$. The velocity of the second action at an active platinum surface is, however, so great that hydrogen peroxide does not escape into the surrounding liquid. A specific reducing action towards a third substance of hydrogen peroxide which is liberated at the best only in minute traces is not to be expected in any circumstances. Further, the reduction potential of hydrogen peroxide is much less than the potential of hydrogen activated at a platinum surface.

In preparative or other work in which reductions are to be effected with activated hydrogen, the author recommends the use of the acid titanium sulphate solution for indicating the conditions favourable to maximum reduction or for controlling the course of the change, since the activity of the system is readily shown by the formation of the characteristic violet-brown solutions of tervalent titanium sulphate. In addition, it is to be expected that the reduced titanium will both accelerate and regulate the reducing action since reduced titanium solutions have been successfully utilised on many previous occasions as reducing agents. H. W.

A Graphical Study of the Stability Relations of Atom Nuclei. WILLIAM D. HARKINS and S. L. MADORSKY (*Physical Rev.*, 1922, 19, 135—156; cf. A., 1921, ii, 445, 582, 690).—The following properties are considered graphically: P , the number of positive electrons in the nucleus, which is taken to be numerically equal to the atomic weight; M , the net positive charge, which is equal to the atomic number; N , the number of negative electrons, which is equal to $(P - M)$; $[(N/P) - 0.5]$, the excess of the relative negativeness N/P over the minimum 0.5; n , the isotopic number, which is equal to $(P - 2M)$ and also to $(N - M)$. Except in the case of hydrogen and of the helium isotope $P=3$, N/P is never less than 0.5, and M/P is never greater than 0.5. For 85% of the atoms composing the earth's crust, both these ratios are 0.5 and $n=0$. As the atomic number increases, the relative negativeness necessary to stability increases above 0.5; that is, as α -particles are added, extra cementing electrons are required to overcome

the increasing mutual repulsion of the positive units of the nucleus. An α -ray transformation does not change n , but increases N/P , whilst a β -ray change decreases n by two units and also decreases N/P ; thus in each group of isotopes, those with larger values of N/P exhibit greater β -ray and less α -ray instability. Stability considerations require that the number of isotopes is on the whole smaller for the lighter atoms. Further, the number of isotopes is larger for even than for odd numbered elements, especially where M is greater than 29. For most atoms, M , n , and P are either all even or all odd, but N is usually even. The curve for the frequency of occurrence of atoms as a function of n shows periodic maxima four units apart, whilst as a function of M or N the periodic maxima are two units apart. It thus becomes possible to predict the existence of the more abundant isotopes of elements of which the mean atomic weights are accurately known.

A. A. E.

Spectra of X-Rays and the Theory of Atomic Structure. D. COSTER (*Phil. Mag.*, 1922, [vi], 43, 1070—1107).—The X-ray spectra of the heavier elements are arranged in a scheme which involves the existence of one K -level, three L -levels, five M -levels, and seven N -levels in the electronic structure of the atom. These levels are associated with the presence of various sub-groups or shells of electrons in the atom. The rules governing the transference of electrons between the various levels are discussed, and further experimental work is adduced in support of these conclusions. New measurements are made of the L -series of X-ray spectra of the elements from rubidium to barium. Fourteen lines are recorded and the wave-lengths offer strong arguments in support of Bohr's theory (this vol., ii, 363). Diagrams, based on these lines, are given of the various transitions which have been found to occur between any two levels. Niton, xenon, and krypton are chosen as typical elements. The lines which disappear with decreasing atomic number fall out at the places in the periodic table predicted by Bohr. Certain faint lines in the L -series which are connected with the M -levels cannot be classified. These and other single lines suggest the existence of new levels associated with only one line. They are attributed, in part, to a more complicated excitation of the atom, due to the removal of two electrons at the same time.

W. E. G.

Structure of Atoms and Molecules in the Light of the Dispersion Theory. JARL A. WASASTJERNA (*Z. physikal. Chem.*, 1922, 101, 193—217).—A theoretical paper in which it is shown that the atomic volume is the volume of a sphere constructed round the middle point of the nucleus of an atom and on the surface of which the outside electron system is situated. At a collision of two atoms the minimum distance between the nuclei is equal to the sum of the radii of the orbits of the outside electrons of both atoms. An electrolyte molecule can exist in two modifications, of which the one consists of uncharged and the other of charged atoms or atomic groups. When a molecule is present in

the latter form, it may split into ions if it is present in a medium of sufficiently high dielectric constant. An equilibrium exists between the two modifications, which lies strongly on the one side for weak electrolytes and strongly on the other side for strong electrolytes. Amphoteric electrolytes may exist in three modifications. Typical strong electrolytes consist almost entirely of polarised molecules. The above makes possible the calculation of the true ionic refractions, and thereby the Kossel hypothesis of the structure of the atom can be confirmed in some of its important points. From the results of the above-named calculations it follows that the outside sheath of the atoms of the alkali metals contains only a single electron, which in salt formation passes over to the acid radicle. In the case of the alkaline-earth metals, the outside sheath has two electrons, which in the same way pass over to the acid radicle. The next sheath of electrons of both the alkali and alkaline-earth atoms is identical with the outside system of the elements in the preceding horizontal row, which in consequence of the increased attractive force was nearer the centre of the nucleus. The structure of the ions undergoes no great change during the formation of polarised molecules, and the electron orbit of the ions can in most respects be regarded as unchanged. The above can be extended, in all probability, to non-polarised molecules and electro-neutral atoms, except in cases where double and treble linkings or rings in the sense adopted by Baeyer are present.

J. F. S.

Separation of Isotopes by Thermal and Pressure Diffusion.

ROBERT S. MULLIKEN (*J. Amer. Chem. Soc.*, 1922, **44**, 1033—1051).—The theory of the separation of isotopes by thermal diffusion and by centrifuging is discussed. Equations are developed giving the difference in atomic weight obtainable in any operation, similar to the equations for diffusion and evaporation processes obtained in a previous paper (this vol., ii, 295). For thermal diffusion, the difference in atomic weight between portions of an isotopic gas at temperatures T_1 and T_2 respectively is $\Delta_r M = K \times B \log_e T_1/T_2$ approximately, the atomic weight being greater at the colder end; B is the ordinary separation coefficient, previously defined, and K is an approximate constant for each element, having a value which is probably about 0.25 and depending on the behaviour of the molecule during impacts. The term KB may be named the thermal separation coefficient. The method of thermal diffusion is shown to be much less effective as a means of separating isotopes than ordinary diffusion or evaporation. A somewhat more advantageous modification of the method is described under the name evaporative thermal diffusion. For the centrifuging of a gas, the difference in atomic weight between the central and peripheral regions is $\Delta_r M = P(v^2 - v_0^2)$, where P , the centrifugal separation coefficient, is a characteristic constant for each element and v and v_0 are the velocities at the peripheral and central regions of the material under treatment. The values of P for several elements have been calculated and the following are recorded for 20°:

lithium, 1.16×10^{-12} ; boron, 1.85×10^{-12} ; neon, 7.38×10^{-12} ; magnesium, 8.97×10^{-12} ; chlorine, 14.5×10^{-12} ; nickel, 18.4×10^{-12} ; zinc, 53×10^{-12} ; bromine, 20.4×10^{-12} , and mercury, 47×10^{-12} . For ordinary air, the coefficient would be 62×10^{-12} . The value of P is unaffected by the state of combination of the element, even if the compound contains other isotopic elements. Thus the separation is equally great for all compounds of a given element. This is in contrast to the situation with all the other diffusion methods, for which the degree of separation of a given element in one operation is inversely proportional to the molecular weight of the compound. Further, the value of P for any element is independent of the atomic weight, whilst the ordinary separation coefficient B is inversely proportional to the latter. Hence, the theory is on this basis relatively increasingly more favourable to the centrifugal method as the atomic weight increases. $P = (M_2 - M_1)^2 x_1 x_2 / 2RT$ for a mixture of two isotopes and for a mixture of several isotopes it has the value $\sum_a \sum_b x_a x_b (M_a - M_b)^2 / 2RT$. P , unlike B , is inversely proportional to T , but depends on the atomic or molecular weight intervals $(M_a - M_b)$ and the molecular fractions, x , in the same way as B . Although for the heavy elements the theory predicts, for a peripheral velocity of 10^5 cm./sec., a separation many times that obtainable in a single diffusion or evaporation, it is shown that compression and condensation of the gas or vapour into the peripheral region make such large separations impracticable if carried out in the ordinary way. The pressure ratio between the two regions is given by $\log_e p/p_0 = M\Delta_r M / 2RT$. P , and so increases with atomic and molecular weight. A special method, named evaporative centrifuging, is proposed whereby gas condensed in the periphery of the centrifuge at high speed would be allowed to evaporate very slowly, the light fraction being drawn off very gradually at lower pressure from the centre of the apparatus. The process would be in effect precisely analogous to an evaporation in which the separation coefficient was increased from B to Pv^2 . This method, applicable at the ordinary temperature to hydrogen chloride, bromide, selenide, telluride, and silicide, might be expected, with peripheral velocities up to 10^5 , to yield a separation ten or fifteen times as great in a single operation as would ordinary diffusion or evaporation. The method may be the most rapid method of separating the isotopes of some of the elements of medium atomic weight, but for the lighter or heavier elements the method is less promising. The theory of the separation of isotopes by centrifuging liquids is discussed, and it is shown that the degree of separation for a given apparatus is identical for liquids, gases, and intermediate states of matter. An attempt has been made to test the theory in the case of mercury, but negative results were obtained which are attributable to a slight vibration of the centrifuge. This effect is likely to prove a limiting factor in any attempt to use the method of liquid centrifuging. A statement made in a previous paper in connexion with the diffusion of an isotopic gas through a thin layer of another gas enclosed in the pores of a membrane (*loc. cit.*) is corrected. This method

is now found to be always less favourable than molecular diffusion, instead of more favourable as previously stated. This is due to the effect of the increased mean free path of the heavier molecules which results from persistence of velocity. This factor was expressly neglected previously. Further, it is shown that mass motion, either positive or negative, should have practically no effect on the rate of separation, except indirectly as it may affect back pressure.

J. F. S.

The Hydrogen Molecule. III. ALBERT C. CREHORE (*Phil. Mag.*, 1922, [vi], 43, 1025—1038; cf. A., 1921, ii, 632; this vol., ii, 438).—A theoretical investigation into the stability of the author's model of the hydrogen molecule. On this view, the forces between the electrons and nuclei cause slight deformations of the shape of the electrons. Two of the electrons in the molecule are flattened and two are expanded in the direction of the minor axis, and these processes may be accompanied by energy changes which may be either positive or negative in sign. These energy changes, which are slight compared with the total energy of the electron, correspond with the heat of dissociation or heat of reaction. The change in mass which occurs on dissociation is so small that it must escape experimental detection.

W. E. G.

The Abnormally Long Free Paths of Electrons in Argon. J. S. TOWNSEND and V. A. BAILEY (*Phil. Mag.*, 1922, [vi], 43, 1127—1128).—A continuation of an investigation into the mean free path of the electron in various gases (cf. this vol., ii, 277). From experiments on mixtures of hydrogen and argon, it is deduced that the mean free path in argon at 1 mm. pressure is about 15 mm., or fifty times the mean free path calculated from the atomic radius of argon.

W. E. G.

Some Linear Relationships in the Periodic System. WILHELM BILTZ (*Z. Elektrochem.*, 1922, 28, 65—70).—A theoretical paper in which a number of linear relationships between the properties of the elements are described. The relationships described are in every case between the inactive gases and the other elements. In the case of atomic volume, it is shown that this quantity for a halogen is proportional to the value for the inactive gas which precedes it in the periodic system. The expression: at. vol. (halogen) = $a \times \text{at. vol. (inactive gas)} + b$, where a and b are constants and have in this case the values 0.8015 and zero respectively. Taking the atomic volumes of the inactive gases as known, the corresponding values for the halogen are given with an error of $\pm 0.2\%$. Similar straight line relationships are found for the atomic volumes of potassium, rubidium, and caesium; titanium, zirconium, and cerium; and silicon, germanium, and tin; but for no other cases. In the same way, the boiling points of the halogens are connected with those of the inactive gases by the equation $\text{b. p. (halogen)} = 2.753 \times \text{b. p. (inactive gas)} - 1.8$. Here the accuracy is the same as before. If hydrogen is included with the halogens, the relationship $\text{b. p. (halogen)} = 2.705 \times \text{b. p. (inactive gas)} + 8.5$ is

accurate to $\pm 2.5\%$. In the case of the critical temperature, the expression $T_c(\text{halogen}) = 2.648 T_c(\text{inactive gas}) + 18.1$ is true absolutely both for the halogens or hydrogen. The atomic refraction of the halogens and hydrogen are connected with the similar value of the inactive gases by the expression $R_D(\text{halogen}) = R_D(\text{inactive gas}) + 0.41$. In this case, the divergence between the calculated and experimental values nowhere exceeds the experimental error. For the elements of the sulphur group $R_D(\text{element}) = 1.713 R_D(\text{inactive gas}) + 1.15$ is true to $\pm 1.8\%$, whilst for hydrogen and the alkali metals $R_D(\text{alkali metal}) = 1.13 R_D(\text{inactive gas}) + 0.51$ with an error of $\pm 1.5\%$. The atomic radii of the halogens, elements of the sulphur group, and the alkali metals are connected to within $\pm 1.0\%$ with the corresponding values for the inactive gases by the expressions $r(\text{halogen}) = 0.78 r(\text{inactive gas}) + 0.26$, $r(\text{S element}) = 0.645 r(\text{inactive gas}) + 0.53$, and $r(\text{alkali metal}) = 1.18 r(\text{inactive gas}) - 0.23$. From a consideration of the foregoing, the author deduces periodic constants.

J. F. S.

The Determination of the State of Valency of Lower Compounds by means of Ammoniacal Silver Solutions. OSCAR OLSSON COLLENBERG (*Z. anorg. Chem.*, 1922, **121**, 281—297).—The author has investigated this method of determining valency (introduced by Philipp and Schwehel, *Ber.*, 1897, **12**, 2234) with respect to suitable conditions of experiment and accuracy of results. If the reaction takes place on the water-bath, the best results are obtained with concentrated ammonia and not too great an excess of silver nitrate in a comparatively small volume. The results obtained are low, but suitable for valency determinations, wholly unsuitable, however, for accurate work, such as determination of purity. If the reaction has to be carried out at a high temperature and under pressure, then a large excess of silver salt in concentrated ammonia is necessary; the temperature should be kept as low as possible and not maintained longer than necessary. The best results were obtained by using silver chloride or sulphate, in the presence of cyanides; silver nitrate should not be used. Ammonia generally contains reducing impurities (organic bases) and a blank experiment should be carried out.

W. T.

Observations on the Construction and Use of the Steele-Grant Microbalance. E. J. HARTUNG (*Phil. Mag.*, 1922, [vi], **43**, 1056—1064).—A description of the Steele-Grant microbalance used in the study of the action of light on the silver haloids (T., 1922, **121**, 682). A balance with a double knife-edge and giving a sensitivity of 1 in 10^6 was sufficiently accurate for the proposed work. For this type the simple double triangle type of beam is recommended; a single knife-edge balance constructed in this manner gave a sensitivity of 1 in 10^7 . Details are given of the grinding of the knife-edges and the drawing of the suspension fibres. Damage to the fibre suspension is avoided by the attachment of a double hook to the end of the quartz weights; this facilitates the removal of the objects from the balance. The author discusses the various sources of error in manipulation,

affecting the sensitivity of the balance, and producing apparent changes in weight.

W. E. G.

Extraction of Small Quantities of Liquids by the Soxhlet Apparatus. HEINRICH HANDORF [*Z. angew. Chem.*, 1922, **35**, 257—258 (D.R.-P., 807757 and 809512)].—A glass vessel is inserted in the Soxhlet apparatus. In cases where the liquid is to be extracted by a solvent lighter than itself, this vessel has above it a funnel, the stem of which extends to the bottom of the vessel and ends in a rose. The liquid to be extracted is placed in the vessel and is covered by a layer of the solvent. Condensed solvent drops into the funnel, emerges in drops from the rose, and rises through the liquid. In the case of a liquid being extracted by a solvent heavier than itself, a short funnel ending in a capillary rests on the top of the vessel, and a tube opening near the bottom of the vessel passes through its wall at a suitable height. The condensed solvent falls through the liquid and passes through the tube to the outside of the vessel. The apparatus is suited for the extraction of colloidal liquids. Quantitative extraction is attained.

H. M.

Rhythmic Deposition of Precipitated Vapours. ENOCH KARRER (*J. Amer. Chem. Soc.*, 1922, **44**, 951—954).—When two narrow cylinders containing respectively small quantities of aqueous solutions of ammonia and hydrochloric acid are placed under a large glass container and so placed that there is a lateral temperature gradient in the container, a series of rings of ammonium chloride forms on the inside of the cylinder which contains the hydrochloric acid. The number of rings varies with the conditions from 1 per cm. to a very large number. These rings mark the edges of disks, so that the vapour space above the hydrochloric acid may be regarded as divided into a number of compartments by disks of ammonium chloride. Within the cylinder each compartment marks the boundaries of a system of vertical and horizontal convection currents, the necessary elements for the formation of which are a lateral temperature gradient and a vertical density gradient.

J. F. S.

Inorganic Chemistry.

Some Properties of Hydrogen Desorbed from Platinum and Palladium. PAUL ANDERSON (*T.*, 1922, **121**, 1153—1161).

The Radius of Hydrogen Atoms in Crystals. G. AMINOFF (*Geol. Förr. Förrh.*, 1921, **43**, 389—396).—It is calculated that the radius of the hydrogen atom in magnesium hydroxide and manganese hydroxide, the elementary parallelepipeds of which are hexagonal, is, respectively, 1.15 and 1.01 Å.U., whilst in ice the radius is 1.12 Å.U. It is shown that the structure $R^{\text{II}}(\text{OH})_2$ may be regarded as ice, H_2OOH_2 , of which one H_2 group has been

replaced by a bivalent metal ion, R^{II} . This explains why the group $\begin{smallmatrix} HO \\ OH \end{smallmatrix}$ of ice has practically the same dimensions in these hydroxides. The relatively high value calculated for the radius of the hydrogen atom is considered to be in keeping with the relation of hydrogen to the alkali metals. CHEMICAL ABSTRACTS.

An Attempt to Separate the Isotopes of Chlorine. E. B. LUDLAM (*Proc. Camb. Phil. Soc.*, 1922, **21**, 45—51).—(Sir) J. J. Thomson has suggested (*Proc. Roy. Soc.*, 1921, [A], **99**, 87) a method of separation of isotopes based on the number of impacts per second of the molecules of a gas on a liquid surface. The number of impacts will be in inverse ratio to the square root of the mass. If on striking the surface the molecules are removed by solution or chemical action, the composition of the residual gas should steadily alter. Hydrochloric acid at a pressure of a few centimetres of mercury was passed over (a) a water surface, and (b) ammonia gas, and a small fraction allowed to remain uncombined. No change in the molecular weight was observed after either process. The failure to obtain a separation indicates that the determining factor in the recombination of ammonia with hydrochloric acid is not mere collision. W. E. G.

Fractional Diffusion of Iodine Vapour. EMIL KOHLWEILER (*Z. physikal. Chem.*, 1922, **101**, 218—234).—Iodine has been subjected to a careful fractional diffusion, and the combining weight of the original material as well as that of the various fractions has been determined by conversion of the iodine into iodide and precipitation as silver iodide. Seventeen determinations with ordinary pure iodine gave a mean value of 126.93, the extreme values being $126.92 + 0.073\%$ and $126.92 - 0.079\%$. Seven determinations with the first fraction of the diffusion gave the value 126.07, which differs by 0.67% from the accepted value for iodine, whilst fourteen determinations with the end fraction gave 127.18, with a divergence of 0.21% from the usual value. These results indicate that, in addition to the isotope of combining weight 127 indicated by Aston, there are at least two others, one lighter and the other heavier than Aston's isotope. It is shown that Aston's results and the present results are not necessarily opposed to one another, for the sensitiveness of Aston's measurement is of the order 5%, whilst that of the present work is much sharper.

J. F. S.

Revision of the Density of Oxygen Gas. E. MOLES and F. GONZÁLEZ (*Anal. Fís. Quím.*, 1922, **20**, 72—80; cf. A., 1921, ii, 546).—Measurements were made on oxygen prepared in different ways as described in the former paper. The arithmetical mean of forty-five determinations gave the value 1.4892 ± 0.00007 as the weight of a litre of oxygen at 0° and 760 mm. This value is identical with the mean of two hundred and thirteen previous determinations by other workers. The corresponding value calculated for latitude 45° is 1.42897. G. W. R.

Researches on Sulphuryl Chloride. II. A New Chlorinating Agent : Preparation of Polychloro-derivatives of Benzene. OSWALD SILBERRAD (T., 1922, 121, 1015—1022).

The "Dynamic" Allotropy of Tellurium. A. DAMIENS (*Compt. rend.*, 1922, 174, 1344—1346).—Density determinations made on tellurium carefully purified by distillation in hydrogen and then in a vacuum, the material being subsequently crystallised either by vaporisation or by slow cooling of the molten material, do not confirm the views of Cohen and Kröner (A., 1913, ii, 315) as to the dynamic allotropy of tellurium. Tellurium obtained by vaporisation has d 6.310, and is not changed by long heating at various temperatures. Specimens prepared in other ways show a slightly lower density, as, owing to the manner of preparation, they are slightly porous. Amorphous tellurium has d 5.85—5.87; on heating, it is transformed into crystalline tellurium with development of heat and its density rises. W. G.

The Reactivity of Ammonia. EDWARD CHARLES CYRIL BALLY and HERBERT MAXWELL DUNCAN (T., 1922, 121, 1008—1014).

Polymorphism of Arsenic. PORPHYRY NICOLAEVITCH LASCHTSCHENKO (T., 1922, 121, 972—978).

Flocculation of Colloidal Arsenic Sulphide. Influence of the Concentration of the Colloid, Shaking, and Temperature. A. BOUTARIC and M. VUILLAUME (*Compt. rend.*, 1922, 174, 1351—1353).—The influence of different factors varies with the nature of the electrolyte used to cause flocculation. With chlorides of potassium, barium, magnesium, or manganese, the velocity of flocculation increases with the concentration of the colloid, but with aluminium or cadmium chlorides the velocity diminishes as the concentration increases. In comparing once shaking to mix the colloidal solution and the electrolyte, and continuous stirring, the opacity curves for the two conditions superpose one another. With potassium chloride, flocculation of the continuously stirred solution only occurs after the opacity limit has been reached, but with barium or aluminium chlorides flocculation occurs before the opacity limit is reached. The velocity of flocculation varies inversely as the temperature where potassium, sodium, ammonium, or lithium chloride is the electrolyte, directly as the temperature with barium, calcium, strontium, magnesium, or cadmium chloride, and is independent of the temperature with aluminium chloride. W. G.

The Isotope Ratio of New Zealand Boron. ALEXANDER DONALD MONRO (T., 1922, 121, 986—988).

The Potassium Salt of Hexahydrodioxidyboron. RAMES CHANDRA RAY (T., 1922, 121, 1088—1094).

The System Potassium Sulphate-Aluminium Sulphate-Water at 25°. HUBERT THOMAS STANLEY BRITTON (T., 1922, 121, 982—986).

Use of Röntgen Rays in Determining the Structure of the Crystals of Lithium and of some of its Compounds with Light Elements. I. J. M. BLJVOET and A. KARSEN (Proc. K. Akad. Wetensch. Amsterdam, 1922, 23, 1365—1370).—The lithium atoms in lithium metal are arranged in a centred cubic lattice with a lattice parameter $a=3.50 \times 10^{-8}$ cm. The relative intensities of the lines are explained by either (i) a simple centred cubic lattice or (ii) a centred cubic lattice with electrons revolving in circles normal to the trigonal axes. There is no evidence for the presence of a lattice of stationary valence electrons. The atomic diameter is 3.04×10^{-8} , in agreement with the value given by Bragg (3.00×10^{-8}). W. E. G.

Crystal Structures of Alkali Haloids. EUGEN POSNJAK and RALPH W. G. WYCKOFF (*J. Washington Acad. Sci.*, 1922, 12, 248—251).—Further X-ray determinations (this vol., ii, 214) are given for lithium chloride, bromide, and iodide, and for sodium, potassium, rubidium, and caesium fluorides. The X-ray constants for the whole series of twenty salts RX are tabulated. The structures of caesium chloride, bromide, and iodide are of the body-centred type, whilst all the others are of the rock-salt type of the simple cubic lattice. L. J. S.

The Preparation of Ammonium Nitrate. (Mlle) WURMSER (*Gompt. rend.*, 1922, 174, 1466—1468).—Le Chatelier has previously given a square diagram showing the states of equilibrium between the solution of the four salts ammonium chloride, sodium nitrate, sodium chloride, and ammonium nitrate and the solid salts at 16° . A similar diagram showing the states of equilibrium at 100° is now given. The curves at 16° and 100° are essentially of the same shape, and from them it is possible to determine the best experimental conditions for obtaining the maximum yield of ammonium nitrate from a solution of the four salts. W. G.

The Properties of Ammonium Nitrate. II. Ammonium Nitrate and Water. IDA L. MILLICAN, ALFRED FRANCIS JOSEPH, and THOMAS MARTIN LOWRY (*T.*, 1922, 121, 959—963).

The Properties of Ammonium Nitrate. III. Ammonium Nitrate and Sodium Nitrate. REGINALD GEORGE EARLY and THOMAS MARTIN LOWRY (*T.*, 1922, 121, 963—969).

The Transformation of Ammonium Carbamate. C. MATIGNON and M. FRÉJACQUES (*Bull. Soc. chim.*, 1922, [iv], 31, 394—412).—A résumé of work already published (*A.*, 1920, ii, 250; 1921, ii, 33; this vol., ii, 272). W. G.

Crystal Structure of Ammonium Fluosilicate. RICHARD M. BOZORTH (*J. Amer. Chem. Soc.*, 1922, 44, 1066—1070).—The crystal structure of ammonium fluosilicate has been shown to be like that of ammonium chloroplatinate, ammonium chlorostannate, and potassium chlorostannate, namely, the structure is that of fluorspar in which each fluorine atom is replaced by an ammonium

group, and each calcium atom by a fluosilicate group with the six fluorine atoms equidistant from the silicon atom in the directions of the axes of the crystals (cf. this vol., i, 441). The length of the cube constituting the unit of structure containing four molecules was found to be 8.38 Å.U.; the corresponding lengths for the three complex salts mentioned above being respectively 9.84, 10.05, and 9.96 Å.U. The shortest distance between the centres of the atoms of fluorine and silicon is 1.72 Å.U., whilst the sum of the Bragg radii for these atoms is 1.84 Å.U. The corresponding distances between the platinum or tin and the chlorine atoms in the other three salts are 2.3, 2.46, and 2.44 Å.U., respectively.

J. F. S.

The Preparation of a Silver Amalgam of the Composition Hg_3Ag_2 by Precipitation from a Solution of Silver Nitrate in Pyridine. ROBERT MÜLLER and REINHOLD HÖNIG (*Z. anorg. Chem.*, 1922, 121, 344—346).—Metallic needles were formed by keeping a drop of mercury in contact with a solution of silver nitrate and cuprous iodide in anhydrous pyridine. With silver nitrate solution alone, crystals of mercurous nitrate were deposited and a liquid amalgam was formed. The *E.M.F.* of the cell $\text{Ag}|0.1\text{NAgNO}_3$ in pyridine| Hg is 0.038 volt, the silver being negative. For the formation of the metallic crystals, the addition of an approximately equal amount of cuprous iodide is necessary (1 : 1); the addition of a much smaller amount (1 : 10) causes the precipitation of finely divided black silver, too large amounts (5 : 1) change the drop of mercury into a granular powder. The concentration of the silver ion in pyridine is not increased by the addition of cuprous iodide, the concentration cell $\text{Ag}|0.1\text{NAgNO}_3$ in pyr. sat. with $\text{Cu}_2\text{I}_2|0.1\text{NAgNO}_3|\text{Ag}$ having an *E.M.F.* of 0.37 volt, the silver in the mixed solution being negative. The formation of a blue precipitate of cupric iodide indicates that the silver salt is reduced to metallic silver, whilst the cuprous salt is oxidised to the cupric salt. The cuprous-cupric potential is +0.18 volt, being greater than that of mercury. Cuprous iodide can be replaced by cerous chloride, the separation of crystals being, however, much slower. Ferrous bromide gave a negative result; in water, the ferrous ferric potential is +0.75 volt, but in pyridine it is evidently less than the mercury potential. The amalgam which crystallises in needles corresponds in composition with the formula Hg_3Ag_2 . The potential of the cell $\text{Hg}_3\text{Ag}_2|\text{AgNO}_3.\text{Cu}_2\text{I}_2|\text{Ag}$ equals 0.37—0.38 volt, the silver being negative against the amalgam.

W. T.

Attempts at a Synthetic Manufacture of Mother-of-Pearl by Production of Chemical Tracery. CLÉMENT and RIVIÈRE (*Compt. rend.*, 1922, 174, 1353—1356).—By precipitating calcium carbonate in the presence of colloidal protein material, such as gelatin in thin layers, a deposit has been obtained having a nacreous structure. The deposit becomes opaque on prolonged drying at 50°, just as mother-of-pearl or pearl itself “dies” when dried for a long period. These experiments support the accepted views as

to the physical and chemical composition of mother-of-pearl and fine pearl.

W. G.

Action of Various Analytical Reagents on Chemical Glassware. W. E. S. TURNER and T. E. WILSON (*J. Soc. Glass Technology*, 1922, 6, 17—20).—The tests were made on three types (two English, one French) of chemical glassware; in most cases the solution was kept in contact with the glass for three hours at 100°. Nitric acid (*d* 1.2), and 2*N*-ammonium sulphide solution had but little action on the glasses, but the latter were attacked to an appreciable extent by *N*/2- and *N*/4-sodium phosphate solutions. The French glass was attacked to a greater extent than were the English glasses by the sodium phosphate solutions, whilst there was but little difference between the three types as regards their resistance to alkali hydroxide solutions, showing that the corrosive action of the alkali salt solutions was not due entirely to their alkalinity.

W. P. S.

Solubility of Crystallised Strontium Hydroxide. D. SIDERSKY (*Bull. Assoc. Chim. Sucr.*, 1921, 39, 167—177).—Tables and graphs are given showing the solubility of crystallised strontium hydroxide in water at temperatures varying from 0° to 101.2° (b. p. of the solution); at 0°, 100 grams of the solution contain 0.90 gram of $\text{Sr}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$; at 15°, 1.46 grams, and at 101.2°, 49.75 grams. Other tables give the solubility of the hydroxide in sucrose solutions at different concentrations and temperatures. At 15°, 100 grams of 10% sucrose solution dissolve 3.79 grams of $\text{Sr}(\text{OH})_2 \cdot 8\text{H}_2\text{O}$, whilst 100 grams of a 25% solution dissolve 7.31 grams of the hydroxide.

W. P. S.

The Reaction of Magnesium Salts with Acid Potassium Pyrantimonate. MILIVOJ OBAJDIN (*Rad. Acad. Sci. Zagreb*, 1922, 226, 220—233).—The preparation of *magnesium dihydrogen orthantimonate* by addition of a magnesium salt to a solution of dipotassium dihydrogen pyrantimonate is described; according to the concentration, it is obtained either as hexagonal or monoclinic crystals or in the amorphous form. From a solution containing 1% of the pyrantimonate, both hexagonal and monoclinic crystals and a large quantity of the amorphous precipitate are deposited. From a solution containing 0.35% of the pyrantimonate, monoclinic crystals are alone deposited, whilst a solution containing 0.07% of the potassium pyrantimonate yields hexagonal crystals only. The composition of both hexagonal and monoclinic crystals corresponds with the formula $\text{Mg}(\text{H}_2\text{SbO}_4)_2 \cdot 10\text{H}_2\text{O}$. The crystals lose 10[?] H_2O if heated at 176°. When heated further, they lose 2 H_2O and change their colour successively from colourless to pink, orange-yellow, yellowish-green, greyish-green, greyish-blue, and violet, a fact which is probably due to the formation of various polymerides. Finally, the crystals spontaneously incandesce and turn white. The composition corresponds now with magnesium metantimonate, $\text{Mg}(\text{SbO}_3)_2$. The magnesium dihydrogen orthantimonate, *d* 2.57, is insoluble in alcohol; the solubility in water at 16° is 0.7075 gram in 1000 grams of solution.

S. S. M.

Phosphorescent Zinc Sulphide. A. A. GUNTZ (*Compt. rend.*, 1922, 174, 1356—1358).—The wurtzite and sphalerite forms of zinc sulphide prepared by heating the precipitated sulphide are both phosphorescent. They may be distinguished under the microscope by their crystalline structures, but show no difference in density. Phosphorescence only develops in the presence of a trace of a heavy metal. The phosphorescence of wurtzite is more persistent than that of blende, but in both cases the decrease of luminous intensity is extremely rapid. Under the influence of heat and shock, the two varieties are extremely thermoluminescent and triboluminescent. W. G.

The Substance between the Crystallites of Metallic Substances. G. TAMMANN (*Z. anorg. Chem.*, 1922, 121, 275—280).—A piece of metal formed from a regulus consists of crystallites surrounded by a film of impurities which form the residue when the metal is dissolved in a solvent. To study the structure of these thin layers of impurities it is necessary to employ a transparent solvent; the evolution of gas and convection currents should be avoided, as these destroy the structure. A sheet of cadmium was dissolved in ammonium nitrate, and the film was found to have a net-like structure, in the meshes of which were micro-crystals of the metal. When the cadmium was dissolved in hydrochloric acid, the net structure was destroyed by the evolution of gas and the foreign substance was left as suspended particles. The amount of impurity was very much decreased by distilling the cadmium in a vacuum. The author points out the desirability of methods to render these non-metallic impurities visible, to estimate them quantitatively, and to find their influence on the properties of the metal. W. T.

Monoclinic Double Selenates of the Cadmium Group. A. E. H. TUTTON (*Proc. Roy. Soc.*, 1922, [A], 101, 245—264).—As in the manganese group (this vol., ii, 505), only three salts, those of rubidium, caesium, and ammonium, could be obtained in the cadmium group of hexahydrated double selenates (cf. preceding abstract). The rubidium and caesium salts were only obtained with great difficulty at a low temperature. No optical investigation was possible of these two salts, and a few optical determinations on the ammonium salt were only just possible, as the crystals rapidly become opaque. The salts crystallise in the holohedral-prismatic class of the monoclinic system.

Rubidium cadmium selenate, $\text{Rb}_2\text{Cd}(\text{SeO}_4)_2 \cdot 6\text{H}_2\text{O}$, $a : b : c = 0.7402 : 1 : 0.5026$; $\beta = 105^\circ 7'$. The crystals are metastable down to 0° . Caesium cadmium selenate, $\text{Cs}_2\text{Cd}(\text{SeO}_4)_2 \cdot 6\text{H}_2\text{O}$, $a : b : c = 0.7319 : 1 : 0.5011$; $\beta = 106^\circ 22'$. This salt is less unstable than the rubidium salt. Ammonium cadmium selenate, $a : b : c = 0.7418 : 1 : 0.5026$; $\beta = 106^\circ 1'$; $d_4^{20} 2.450$.

The results of thirty-two years' work on the double sulphates and selenates of the series $\text{R}_2\text{M}\left(\begin{smallmatrix} \text{S} \\ \text{Se} \end{smallmatrix} \text{O}_4\right)_2 \cdot 6\text{H}_2\text{O}$ are discussed, in the course of which seventy-five salts, comprising nineteen distinct

groups, have been examined. In each group it is shown that there is a regular progression in the minutest details of the structural and physical properties of the crystals as the atomic number of the alkali metal increases in passing from potassium (19) to rubidium (37) and caesium (55). The ammonium salt is always practically isostructural with the rubidium salt of the same group. These results are in agreement with the recently developed theories of atomic and crystalline structure of Langmuir and W. L. Bragg.

E. H. R.

Tellurium Lead and Tellurium Antimony Alloys. MAX DREIFUSS (*Z. Elektrochem.*, 1922, 28, 100—101, 224).—When lead is added in small quantities of molten tellurium, a portion of the lead forms a slag with the tellurium and a small amount of tellurium passes into solution. Tellurium has a deoxidising action on lead, so that lead containing tellurium may be heated to redness without the free liquid surface losing its brightness. Alloys containing tellurium and lead may be prepared by mixing tellurium with molten antimony; here a continuous series of mixed crystals is formed. To the molten alloy small quantities of antimony-lead alloy are added and eventually pure lead. A stiff, pasty mass is obtained which on heating at 700° becomes liquid and does not readily lose its bright surface. In the same way, small quantities of tin and copper alloy with the antimony tellurium alloy. The addition of 2% tellurium to lead-antimony alloys has no marked effect on the hardness. The addition of 5—6% of tin to antimony-lead alloys produce much more desirable properties than does tellurium.

J. F. S.

The Structure of the Isomorphous Group, $\text{Pb}(\text{NO}_3)_2$, $\text{Ba}(\text{NO}_3)_2$, $\text{Sr}(\text{NO}_3)_2$, $\text{Ca}(\text{NO}_3)_2$. L. VEGARD (*Z. Physik*, 1922, 9, 395—410).—The space lattice of this isomorphous group of compounds has been determined by the application of the powder method. Little difficulty was experienced with the nitrates of lead, barium, and strontium, but the deliquescent calcium salt gave less satisfactory results. The positions of the maxima of reflection show that the crystals are not built up entirely of face-centred lattices. The metal atoms are arranged, however, in a face-centred lattice, for the intensity of the maxima of this lattice increases with increasing atomic weight of the metal. Three oxygen atoms and one nitrogen atom form a group at four corners of a cube. This group, which possesses a trigonal axis, is surrounded by four atoms of the metal.

The values for the atomic radii of nitrogen and oxygen atoms are appreciably higher than in those substances investigated by W. L. Bragg. The atomic radii vary with the nature of the chemical combination and the geometrical arrangement in space. The X-ray method provides a useful method for the determination of the density of solid substances. The presence of impurities scarcely affects the lattice constants or the atomic radii, from which the density may be calculated. Thus the density of the pure substance can be accurately determined from impure material. The values

for the densities of the compounds in the isomorphous group of nitrates, calculated from the atomic radii, are in good agreement with the experimental results. W. E. G.

Thallic-Thallous Sulphate. I. A. BENRATH and H. ESPENSCHIED (*Z. anorg. Chem.*, 1922, **121**, 361—362).—Many double salts of thallic and thallous sulphates are claimed by different authors, but the present authors could obtain only two corresponding in composition with Ti_2SO_4 , $\text{Ti}_2(\text{SO}_4)_3$ and $5\text{Ti}_2\text{SO}_4$, $\text{Ti}_2(\text{SO}_4)_3$; all the other so-called double salts are to be regarded as mixtures of these. The preparations and methods of analysis are given. The double salt, Ti_2SO_4 , $\text{Ti}_2(\text{SO}_4)_3$, has a transition temperature of 48° with a fairly large transition interval. The double salt interval for $5\text{Ti}_2\text{SO}_4$, $\text{Ti}_2(\text{SO}_4)_3$ has its upper limit between 23° and 30° . Between this transition temperature and 48° the whole system is in a transition state, so that at these temperatures mixtures of variable composition will be obtained. W. T.

Determination of the Complex Formation in Aqueous Solutions of Copper Salts by Means of Permutite. A. GÜNTHER-SCHULZE (*Z. Elektrochem.*, 1922, **28**, 89—99).—The complex formation in solutions of copper chloride, acetate, formate, sulphate, chlorate, nitrate, and bromide has been investigated by shaking known concentrations of the salts with pure potassium permutite. From measurements of the exchange of kations between the permutite and the solution, conclusions may be drawn as to the presence of complexes in the solution. It is shown, by this method, that the simple complex kation CuR^+ is present in practically all the copper salt solutions examined even at the greatest dilutions. The presence of the complex Cu_2R_3^+ has been proved only in the case of copper chloride and bromide, whilst it is probable that the other salts exhibit a higher complex formation, which, however, is not present in such great concentrations. The complexity of copper salt solutions increases at constant concentration with the strength of the acid, and is greater with the salts of halogen acids than with the salts of oxygen acids of equal strength. J. F. S.

Extraction and Purification of Scandium from Thorveitite from Madagascar. PIERRE URBAIN and G. URBAIN (*Compt. rend.*, 1922, **174**, 1310—1313).—The finely ground mineral is fused with sodium hydroxide and the product extracted with water. The insoluble residue is dissolved in the least possible quantity of sulphuric acid, and the rare earth metals are precipitated as fluorides by the addition of an excess of hydrofluoric acid. The fluorides are decomposed by sulphuric acid, and scandium hydrogen sulphate crystallises out on concentrating the solution. The sulphates are converted into nitrates by passage through the hydroxides, and the solution of nitrates is evaporated and heated on a water-bath for a long time to free it from excess of acid. To the concentrated solution an excess of powdered potassium sulphate is added until the solution is saturated with this salt. After three days, the double scandium potassium sulphate is collected and washed as

quickly as possible with a saturated solution of potassium sulphate. Any scandium remaining in the mother-liquors is precipitated as hydroxide and put through the same process. The double sulphate may be dissolved in aqueous ammonium carbonate, and from the solution, on warming, scandium hydroxy-carbonate is precipitated. The residue of the scandium in the mother-liquors may be recovered by conversion into its acetylacetonate, which is soluble in chloroform and sublimes when heated in a vacuum at 200° . W. G.

The Atomic Numbers of Neo-ytterbium, Lutecium, and Celtium. G. URBAIN (*Compt. rend.*, 1922, 174, 1349—1351).—An historical paper in which the author points out that Dauvillier's work (this vol., ii, 463) completes the work of Moseley, himself, and others, and definitely establishes celtium as a chemical element with the atomic number 72, and having an atomic weight between that of lutecium, 175, and that of tantalum, 181.5. W. G.

The Measurements of the Densities of the Haloids of Aluminium with Mercury as Pyknometer Liquid. WILHELM BILTZ and WALTER WEIN (*Z. anorg. Chem.*, 1922, 121, 257—265).—The author in previous communications (A., 1921, ii, 437; 1922, ii, 487) has discovered an abnormality in the linear relations in the atomic and molecular volumes of the haloids of aluminium. Former workers (Deville and Troost, *Ann. Chim. Phys.*, 1860, [iii], 58, 279) used "oil of naphtha" as the pyknometer liquid. Inaccuracies due to the solubility of the bromide and iodide of aluminium in this liquid led the author to replace it by mercury. Toluene was employed in the case of the chloride, on which it has no action. Improved methods of preparation of the haloids are given, and density determinations are described. The densities and molecular volumes were found to be: aluminium fluoride (d_4^{25} 3.10), 27.1; aluminium chloride (d_4^{25} 2.44), 54.7; aluminium bromide (d_4^{25} 3.01), 88.7; aluminium iodide (d_4^{25} 3.98), 102.5. These values, however, do not show the linear relationship exhibited by all other haloids. Other abnormalities are found in the physical constants of the haloids of aluminium, for example, the melting point of the bromide does not lie between those of the iodide and chloride. These are ascribed to polymerisation to double molecules, $[\text{Al}_2\text{F}_6]$, $[\text{Al}_2\text{Cl}_6]$, $[\text{Al}_2\text{Br}_6]$, and $[\text{Al}_2\text{I}_6]$. These double molecules correspond with cryolite, $[\text{AlF}_6]\text{Na}_3$, the three univalent sodium atoms being replaced by one trivalent aluminium atom thus: $[\text{AlF}_6]\text{Al}$, $[\text{AlCl}_6]\text{Al}$, $[\text{AlBr}_6]\text{Al}$ and $[\text{AlI}_6]\text{Al}$. An X-ray examination is necessary to test this explanation. W. T.

Monoclinic Double Selenates of the Manganese Group. A. E. H. TUTTON (*Proc. Roy. Soc.*, 1922, [A], 101, 225—245).—In the group of the series $\text{R}_2\text{M}(\text{SeO}_4)_2 \cdot 6\text{H}_2\text{O}$, in which M is manganese, only three salts could be obtained, namely, those in which R is rubidium, caesium, or ammonium. Potassium manganese selenate, like the corresponding sulphate, only exists in the tetra-hydrated form, even at 0° . The morphological relationships between the three salts are similar to those observed in every

other group of the series (cf. A., 1920, ii, 690). The salts all crystallise in the holohedral-prismatic class of the monoclinic system. Rubidium manganous selenate, $\text{Rb}_2\text{Mn}(\text{SeO}_4)_2 \cdot 6\text{H}_2\text{O}$, $a:b:c=0.7422:1:0.5008$; $\beta=105^\circ 9'$; $d_4^{20} 2.763$. Cæsium manganous selenate, $\text{Cs}_2\text{Mn}(\text{SeO}_4)_2 \cdot 6\text{H}_2\text{O}$, $a:b:c=0.7319:1:0.4957$; $\beta=106^\circ 22'$; $d_4^{20} 3.008$. Ammonium manganous selenate, $(\text{NH}_4)_2\text{Mn}(\text{SeO}_4)_2 \cdot 6\text{H}_2\text{O}$, $a:b:c=0.7427:1:0.4979$; $\beta=106^\circ 16'$; $d_4^{20} 2.158$. The ammonium salt, besides the cleavage parallel to $\{201\}$ common to the series, exhibits an even more perfect one parallel to $\{010\}$. The optical constants of the crystals of all the salts were measured.

E. H. R.

Equilibria in the Systems Iron-Carbon-Oxygen and Iron-Hydrogen-Oxygen, and the Free Energies of the Oxides of Iron. E. D. EASTMAN (*J. Amer. Chem. Soc.*, 1922, **44**, 975—998).—A theoretical paper in which the data obtained by a number of authors in connexion with the two systems named in the title have been correlated and criticised. A number of calculations have been made from the correlated data, and the following conclusions drawn. Ferric oxide and ferroso-ferric oxide at 1100° and above form a continuous series of solid solutions, although there are indications that at temperatures much below 1000° they may not be soluble in each other in all proportions. Ferroso-ferric oxide and ferrous oxide up to 1100° at least definitely do not form a continuous series of solutions, although they do dissolve in each other to a limited extent. Ferrous oxide appears to be soluble in iron to the extent of 20—25 mol. % (6—8% oxygen), although metallic iron is not appreciably soluble in the oxide. Ferrous oxide is unstable with respect to ferroso-ferric oxide and iron below about 565° , and ferroso-ferric oxide shows an inversion point in the same region. When carbon monoxide reacts with pure iron in a closed tube above 500° , carbon is not ordinarily deposited as a separate phase. At temperatures above 800° , a trivariant equilibrium with iron containing dissolved carbon or carbide as the solid phase may be established, or one in which the solid phase contains dissolved oxygen as well as carbon, and the gas is richer in carbon dioxide than in the former case. It is possible that bivariant equilibria corresponding with these, with a separate dissolved carbon or carbide phase, may also be established on repeated treatment, but this does not appear to be proved. With a stream of carbon monoxide acting on iron in the vicinity of 650° , separate phases consisting of carbon, ferrous oxide, and iron containing dissolved oxygen and carbon (or carbide) are probably formed. At higher temperatures, the oxide and separate carbon phases do not appear, and the amount of dissolved carbon decreases. Carbon dioxide acting on iron in a closed tube probably produces again a trivariant system. At temperatures in the neighbourhood of 600° and lower, carbon is present in the solid phase, decreasing in amount as the temperature is increased. The individual results, of the investigators of univariant and bivariant equilibria involving iron or ferrous oxide, ferrous oxide or ferroso-ferric oxide, carbon

monoxide, carbon dioxide, and carbon under conditions such that the presence of separate and definite solid phases in large amounts was assured, are in sufficient agreement to permit the use of equilibrium constants and free energies based on them. These constants are tabulated for the range 600—1000°. Bivariant equilibria involving iron or ferrous oxide, ferrous oxide or ferroso-ferric oxide, and hydrogen yield such different results that the data cannot be safely used.

J. F. S.

The Penetration of Tempering in Steel. GEORGES CHARPY and LOUIS GRENET (*Compt. rend.*, 1922, **174**, 1273—1276).—A square bar of the steel to be examined is used for the purpose. It is heated in the furnace at the required temperature until its temperature is uniform. The upper part of the bar is then wrapped round with asbestos, and on to the base is directed a regulated stream of cold water. When the cooling is complete, the hardness is determined on one face at different distances from the base. The bar may then, if necessary, be submitted to further thermal treatments and the other three faces used for determining the hardness. The results are in accord with those obtained on a fragment of metal cut out from a thick piece after its tempering. With a nickel-chrome steel, the hardness was found to vary from 387 at 3 mm. from the base to 180 at 90 mm. With an auto-tempering steel, after tempering at 800°, the hardness was practically uniform along the whole length of the bar.

The method may also be used for metals which are not homogeneous, and permits of a study of the influence of cementation on thick plates. In this way, it was found that, in certain cases, the hardness of the part situated behind the cemented region is less than that obtained with a bar of the same composition, tempered under the same conditions, but showing no cemented region.

W. G.

Centrifugal Method for Preparing Colloidal Ferric Hydroxide, Aluminium Hydroxide, and Silicic Acid. RICHARD BRADFELD (*J. Amer. Chem. Soc.*, 1922, **44**, 965—974).—Colloidal ferric hydroxide, aluminium hydroxide, and silicic acid may be prepared from the freshly formed precipitates of these substances by removing the excess of precipitating agent by a very thorough washing, which is achieved by means of a centrifuge with a rotation of 32,500 per minute. This method has many advantages over the older methods, chief of which are, (a) the removal of the electrolytes formed on precipitation is more complete; (b) the addition of a peptising agent and its consequent incomplete removal by either prolonged boiling or dialysis is unnecessary; (c) sols of a more uniform degree of dispersion can be prepared, since particles of similar size and of similar degrees of hydration are deposited in the same zone of the centrifuge bowl; (d) sols of any desired concentration from a semi-gel to the merest trace can be prepared by the addition of water to the more concentrated form, and all concentrations are very stable. It is shown that aluminium hydroxide

precipitated by ammonium hydroxide in the presence of an excess of sulphate ions is not irreversible. J. F. S.

The Reduction of Oxides by Hydrogen. E. BERGER (*Compt. rend.*, 1922, **174**, 1341—1343; cf. A., 1914, ii, 656; Sabatier and Espil, A., 1914, ii, 276).—From a study of the velocity curves for the reduction, by hydrogen, of specimens of nickel oxide prepared in different ways, the author finds that the discontinuity in the curves occurs at different stages in the reduction varying with the specimen. This discontinuity cannot, therefore, be explained on the basis of the formation of an intermediate oxide, and doubt is therefore cast on the existence of sub-oxides of nickel. W. G.

New Properties of the Green Chromium Sulphate. A. RECOURA (*Compt. rend.*, 1922, **174**, 1460—1463).—It has been shown by Colson (A., 1907, ii, 177, 267, 474) that the green chromium sulphate, immediately after its solution in water, exists in a condensed form, which is slowly depolymerised. In this condensed form, it is found to be capable of masking the sulphate ion in other sulphates to the extent of several hundred molecules for each molecule of the green sulphate present. The number of molecules which can be thus masked depends on the length of time during which the green sulphate has been in solution, the acidity, and the concentration of the solution. W. G.

X-Ray Examination of the Trioxides of Tungsten, Molybdenum, and their Hydrates. H. C. BURGER (*Z. anorg. Chem.*, 1922, **121**, 240—242).—The highest oxide of tungsten, WO_3 , can exist in various forms as large, bright yellow crystals, or long, thin, dark blue, or green needles. All these modifications gave identical angles of reflection which were, however, different from that of the hydrate. This proves that the hydrate is a definite chemical compound with its own characteristic crystal form. It corresponds in composition with $\text{WO}_3 \cdot \text{H}_2\text{O}$, that is, H_2WO_4 ; this, on heating, loses water and gives the above trioxide. A definite chemical compound was also indicated in the case of the other hydrate, $\text{WO}_3 \cdot 2\text{H}_2\text{O}$; that is, H_4WO_5 tungstic acid. In the case of molybdenum, the existence of H_2MoO_4 as a definite chemical compound was proved. This is a much more convenient method for the solution of such problems than that of measuring the vapour pressure as a function of the composition. The phases in the mixture can be found and the whole work carried out in a short time with a few mg. of the substance. W. T.

The Valency of Tungsten and Molybdenum in their Complex Octacyanides. OSCAR OLSSON COLLENBERG (*Z. anorg. Chem.*, 1922, **121**, 298—312).—The author reviews previous work on the subject and shows the sources of error. He determined the valency by means of ammoniacal silver chloride and found that the central atoms were in each case quadrivalent in agreement with their formulæ, $[\text{W}(\text{CN})_8]\text{K}_4$ and $[\text{Mo}(\text{CN})_8]\text{K}_4$. These results were confirmed by another method, using ferric sulphate and potassium

permanganate, this method, due to Friedheim and Hoffmann (A., 1902, ii, 265), being modified and so rendered applicable in the presence of cyanides.


W. T.

The Chlorides of Bivalent Molybdenum, Tungsten, and Tantalum. I. KURT LINDNER [with EMMA HALLER, HERBERT HELWIG, ANTONIE KÖHLER, and HEILMUTH FEIT] (*Ber.*, 1922, 55, [B], 1458—1465).—The formulæ $\text{HMo}_3\text{Cl}_7 \cdot 4\text{H}_2\text{O}$, $\text{HW}_3\text{Cl}_7 \cdot 4 \cdot 5\text{H}_2\text{O}$, and $\text{Ta}_3\text{Cl}_{14} \cdot 7\text{H}_2\text{O}$ have been ascribed by Rosenheim and Kohn (A., 1910, ii, 300), Hill (A., 1917, ii, 36), and Chapin (A., 1910, ii, 303), respectively, to the dichlorides of molybdenum, tungsten, and tantalum. The preparation of these chlorides has been subjected to exhaustive investigation. It is shown that they are uniformly constituted in accordance with the general formula, $\text{HX}_3\text{Cl}_7 \cdot 4\text{H}_2\text{O}$, in which the metal is bivalent. The methods used in their preparation are the treatment of the metals with gaseous chlorine under suitable conditions of temperature and pressure and the reduction of the higher chlorides by gases or other reducing agents such as metals.

At all temperatures up to 1200° , molybdenum is converted by chlorine into the pentachloride. The dichloride is obtained conveniently by passing carbonyl chloride over powdered molybdenum at $700\text{--}800^\circ$; a small quantity of the pentachloride sublimes, after which the metal gradually becomes coated with a heavy yellow layer of molybdenum dichloride, which ultimately protects the metal from further action. The dichloride is then extracted with concentrated hydrochloric acid and the residual metal is again treated with carbonyl chloride. The compound, $\text{HMo}_3\text{Cl}_7 \cdot 4\text{H}_2\text{O}$, separates almost quantitatively from the acid solutions in the form of long, yellow needles. At about 600° , molybdenum trichloride is the main product of the action of carbonyl chloride on molybdenum, whereas below this temperature the pentachloride is produced. Less favourable but useful results are obtained by the reduction of molybdenum pentachloride with finely divided aluminium in the presence of ignited quartz powder (to moderate the violence of the reaction). The product is subsequently extracted with hydrochloric acid. The yield is considerably diminished by the reduction of molybdenum pentachloride to other stages and even to metallic aluminium and the consequent difficulty of working up the residues.

Tungsten dichloride cannot be prepared from the metal and carbonyl chloride, which, over a wide interval of temperature, give carbon and the "red oxychloride," WOCl_4 , in addition to small quantities of tungsten hexachloride and probably of tungsten pentachloride. The reduction of tungsten hexachloride by sodium amalgam (Hill, *loc. cit.*) is uncontrollable, and even if successful only gives minimal quantities of the dichloride; the results with an alloy of sodium and potassium are even less promising. On the other hand, tungsten hexachloride is readily reduced to the dichloride by ignition in a stream of nitrogen with powdered aluminium and quartz; the product is extracted with hydrochloric acid and the

reddish-yellow extract is concentrated and saturated with hydrogen chloride, thus causing the separation of the compound



in slender, yellow needles, which are considerably less stable than the corresponding molybdenum compound and rapidly lose hydrogen chloride when exposed to air.

Tantalum is converted into its pentachloride by carbonyl chloride. Tantalum pentachloride is reduced by powdered metals to the dichloride and, since the reaction proceeds less violently than with the molybdenum and tungsten compounds, the addition of powdered quartz may be omitted. Aluminium, zinc, and lead are efficient. The residues are extracted exhaustively with hydrochloric acid. The isolation of the compound $\text{HTa}_3\text{Cl}_7 \cdot 4\text{H}_2\text{O}$, a blackish-green, unusually stable, crystalline powder, from the extracts is a matter of some difficulty, since it only separates from very concentrated solutions and its crystallisation is greatly impeded by the presence of dissolved aluminium chloride. It is preferable, therefore, to use zinc which can be removed as sulphide after addition of sodium acetate to the solution or lead, the dissolved traces of which can be precipitated directly with hydrogen sulphide. Molten tantalum pentachloride cannot be reduced electrolytically between carbon poles, since the substance is an insulator (cf. Biltz and Voigt, this vol., ii, 302). Electrolysis takes place after addition of alkali chlorides, and the dichloride is formed to some extent; the method has no preparative value. H: W.

Electrolytic Removal of Alkali from Salts. A. LOTTER-MOSER (*Kolloid Z.*, 1922, **30**, 346—348).—The author shows that by the electrolysis of normal sodium tungstate it is possible to prepare paratungstates and other complex tungstates by the removal of sodium hydroxide. In the case of sodium paratungstate, the preparation is carried out as follows. A solution of normal sodium tungstate ($\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$), 51 grams in 200 c.c. of water, is placed in a 250 c.c. porous pot which stands in a beaker containing exactly 750 c.c. of *N*/10-sodium hydroxide. An anode consisting of a coil of platinum wire is placed in the porous cell, and the electrolysis carried out with a current of 2 amperes until the sodium hydroxide has increased in concentration by the calculated amount. When this state is reached, the anode solution is allowed to crystallise; sodium paratungstate separates in large, vitreous crystals. In the same way, sodium ditungstate and metatungstate were also prepared (cf. Kröger, this vol., ii, 212). J. F. S.

The Hydrates of Uranium Tetroxide and Uranium Trioxide. GUSTAV F. HÜTTIG and EDITH VON SCHROEDER (*Z. anorg. Chem.*, 1922, **121**, 243—253).—Details are given of the preparation of the bright yellow, hygroscopic hydrate of uranium tetroxide, which corresponds with the formula $\text{UO}_4 \cdot 3\text{H}_2\text{O}$. This was heated in a vacuum, but the tetroxide, UO_4 , could not be obtained, the final product being the trioxide, UO_3 . The compound loses oxygen before all the water is expelled, and the authors consider the existence of the tetroxide as very doubtful. The changes in composition were

studied in a Hüttig tensi-eudiometer (A., 1921, ii, 195), and the following hydrates were indicated: $\text{UO}_4 \cdot 4.5\text{H}_2\text{O} = \text{UO}_3 \cdot \text{H}_2\text{O}_2 \cdot 3.5\text{H}_2\text{O}$; $\text{UO}_3 \cdot 3\text{H}_2\text{O} = \text{UO}_3 \cdot 0.5\text{H}_2\text{O}_2 \cdot 2.5\text{H}_2\text{O} = \text{H}_2\text{U}_2\text{O}_8 \cdot 5\text{H}_2\text{O}$; $\text{UO}_3 \cdot 5.2\text{H}_2\text{O} = \text{UO}_3 \cdot 0.5\text{H}_2\text{O}_2 \cdot 1.5\text{H}_2\text{O} = \text{H}_2\text{U}_2\text{O}_8 \cdot 3\text{H}_2\text{O}$ [?]; peruranic acid pentahydrate, $\text{UO}_3 \cdot 0.5\text{H}_2\text{O} = \text{H}_2\text{U}_2\text{O}_7$, pyrouanic acid. It is assumed that these are derivatives of the trioxide and that hydrogen peroxide is contained in the first, second, and third compounds.

In the case of uranium trioxide, indication was given of the following hydrates: $\text{UO}_3 \cdot 2\text{H}_2\text{O} = \text{H}_2\text{UO}_4 \cdot \text{H}_2\text{O}$, $\text{UO}_3 \cdot \text{H}_2\text{O} = \text{H}_2\text{UO}_4$, uranic acid.

W. T.

The Triboluminescence of Uranium Salts and Three New Organic Uranium Compounds. J. A. SIEMSEN (*Chem. Ztg.*, 1922, 46, 450; cf. A., 1919, ii, 346).—A specimen of uranium nitrate which had been kept in a stoppered bottle for forty-one years showed weak triboluminescence compared with other preparations, and its luminosity was very slight. A specimen of sodium uranate which was not triboluminescent was converted into uranium nitrate which showed fairly strong triboluminescence. A preliminary announcement is made of the discovery of three salts, obtained by the action of pyrazolone derivatives on uranium compounds, which possess powerful bactericidal properties. No further chemical particulars are given.

E. H. R.

The Behaviour of the Stannic Acids towards Hydrochloric Acid. GEORGE ERNEST COLLINS and JOHN KERFOOT WOOD (*T.*, 1922, 121, 1122—1133).

The Anodic Corrosion of Bismuth. Bismuth Compounds. E. B. R. PRIDEAUX and H. W. HEWIS (*J. Soc. Chem. Ind.*, 1922, 41, 167—171r).—As a possibly advantageous alternative to attack by nitric acid, the anodic corrosion of bismuth in an anolyte of sodium nitrate was investigated. The bismuth anodes were cast on copper wire, the cathode was a strip of nickel and in order to prevent reduction of the nitrate to ammonia it was placed in a catholyte of dilute alkali in a porous pot.

In the first series of experiments, 10% sodium nitrate solutions were used, with a current density of 12.5 amperes per sq. dm. and an average of 4.6 volts. The current efficiency calculated from the anodic loss was 100%, and 1.76 kilowatt-hours were consumed per kilo. of bismuth dissolved. The product was an oxynitrate containing on the average about 30% of bismuth oxide, and came down partly as a white deposit and partly as a crust on the anode. The amount of bismuth in the solution was negligible. In the second series of experiments, saturated solutions of sodium nitrate were used as the anolyte, and in this case the bismuth oxynitrate came down entirely as a loose precipitate, and one-half to three-quarters of the total bismuth remained in solution, from which it may be obtained as oxynitrate by dilution, or used directly for other bismuth preparations. The fully hydrated white bismuth hydroxide, $\text{Bi}(\text{OH})_3$, can be prepared from this anodic solution by allowing it to run into an equal volume of $N/2$ -sodium hydroxide

diluted about five times. The white precipitate is washed by decantation. An examination of xeroform (bismuth tribromophenoxide) prepared by the usual method of adding bismuth nitrate to a solution of tribromophenol in sodium hydroxide, showed that it contained an appreciable quantity of oxynitrate, and the alcoholic extract contained nitrobromophenols in addition to much tribromophenol. On ignition, moreover, it gave a lemon-yellow residue containing bromine, and it does not therefore yield pure bismuth oxide, as usually stated. Commercial xeroform appears therefore to be very impure, although doubtless quite suitable for the purpose intended. A purer preparation was obtained by using the anodic solution of bismuth nitrate in sodium nitrate. It forms a bulky, pink precipitate, which, although it has a definite bismuth content (15.46%), does not appear to correspond exactly with any of the simpler compounds which might be expected, for example, $(C_6H_2Br_3O)_3Bi$ requires 17.36%. It is possibly a hydrate, and the matter is being further investigated. G. F. M.

Double Haloids of Bismuth and Thallium. G. CANNERI and G. PERINA (*Gazzetta*, 1922, 52, i, 241—246; cf. this vol., ii, 378; Ephraim and Barteczko, A., 1909, ii, 236).—*Bismuth thalious bromide*, $2TlBr, BiBr_3$, prepared by dissolving bismuth carbonate in nitric acid and adding potassium bromide, crystallises in lustrous, lemon-yellow, hexagonal lamellæ, and in neutral aqueous solution undergoes rapid hydrolysis to thalious bromide, bismuth oxybromide, and bromine. *Bismuth thalious iodide*, $2TlI, BiI_3$, prepared by adding potassium iodide to an acid solution containing thallium and bismuth, forms red, regular hexagonal plates, exhibits physical and chemical properties similar to those of the preceding compound, and is soluble in about 20,000 parts of water.

The formation of this insoluble double iodide may be used as a means of detecting and estimating bismuth after other metals forming insoluble iodides have been removed. A solution containing 0.029 gram of bismuth per litre yields a voluminous red precipitate with a thalious salt and potassium iodide, whereas with hydrogen sulphide only a faint brown coloration is obtained. For the estimation, the solution of the bismuth salt is acidified with hydrochloric acid and then treated with a known volume, in excess, of standard potassium iodide solution. Standard thalious sulphate solution is next run in from a burette until precipitation of the red double iodide ceases and that of yellow thalious iodide is evident. The precipitate is collected on a Gooch crucible and washed thoroughly with absolute alcohol, the residual excess of potassium iodide in the solution being estimated by means of potassium iodate. If a represents the amount in grams of iodine added as iodide to the solution, b that combining with the thallium, and c that remaining in excess, calculated on the basis that 3 atoms of iodine correspond with 1 atom of bismuth, the weight of the latter in grams is given by: $208[a - (b + c)]/380.76$.

Addition to a solution of $K_3Bi(CNS)_6$ (cf. Vanino and Hauser, A., 1902, i, 14) of a proportion of thallium acetate corresponding

with one equivalent of the potassium results in the formation of an orange-yellow *precipitate* of the composition $K_2TlBi(CNS)_6$, whilst with double this proportion of thallium acetate, the *compound*, $KTl_2Bi(CNS)_6$, is obtained as a paler yellow precipitate. All attempts to prepare the trisubstituted salt, $Tl_3Bi(CNS)_6$, have proved unsuccessful, the molecule undergoing demolition with formation of thalious thiocyanate.

T. H. P.

The Reduction of Tantalum Pentachloride. OTTO RUFF and FRITZ THOMAS (*Ber.*, 1922, 55, [B], 1466—1473).—Tantalum pentachloride is reduced by metallic aluminium in the presence of aluminium chloride at 300° to a mixture of various chlorides of tantalum. A larger excess of aluminium remains unchanged even when, as is occasionally the case, metallic tantalum is formed. The temperature is limited by the resistance of the tubes. The dark green product of the reaction evolves aluminium chloride and tantalum pentachloride when heated, the latter being given off in such a manner that the composition of the residue can be regulated by the choice of temperature and pressure. A mixture corresponding approximately with tantalum tetrachloride is obtained at $250^\circ/2-3$ mm.; at $350-400^\circ$ the residue has the composition of a trichloride which is itself decomposed at about 500° , leaving a mixture of di- and tri-chlorides.

The mixtures of chlorides are moderately stable towards air. Treatment with water causes the formation of intensely green solutions of the trichloride, whilst the pentachloride is hydrolysed to hydrochloric and tantalic acids and the dichloride remains undissolved as a blackish-green residue. The green solutions become oxidised slowly on exposure to air, the action being checked by the presence of acids. The di- and tri-chlorides are soluble in cold, sufficiently dilute sodium hydroxide solutions without evolution of hydrogen or alteration in the valency of the element. The alkaline solutions are oxidised with unusual readiness by air and evolve hydrogen when they are warmed; a brown tantalous oxide is thereby produced as an intermediate step in the formation of tantalic acid. The production of this substance is also observed during rapid oxidations (with hydrogen peroxide or nitric acid) in acid solution. When oxidation is effected by hydrogen peroxide in alkaline solution, a brown solution of a pertantalate is obtained which evolves oxygen and becomes colourless when boiled.

The behaviour of aqueous solutions of titanium trichloride towards common reagents is described in detail.

H. W.

Gold Sulphides. A. GUTBIER and E. DÜRRWÄCHTER (*Z. anorg. Chem.*, 1922, 121, 266—274).—Various sulphides of gold are described in the literature. The authors could not obtain the sulphides with the formulæ Au_2S and Au_2S_2 . A rapid stream of hydrogen sulphide passed at the ordinary temperature into a 2% solution of chloroauric acid, $[AuCl_4]H$, in *N*-hydrochloric acid gave a black precipitate of auric sulphide, Au_2S_3 , mixed with small amounts of metallic gold. The relative amounts of sulphide and metal depend on temperature, concentration of the chloro-

auric acid, and the rate of passing in the hydrogen sulphide. At low temperatures -2° , with a low concentration of the chloro-acid and a very rapid stream of gas, the pure sulphide was obtained. The low concentration of the tetrachloro-acid is necessary, because it reacts with the sulphide formed. Auric sulphide reacts more rapidly with auric chloride than with the tetrachloro-auric acid and pure auric sulphide is not obtained as above if the gas is bubbled into a solution of auric chloride. W. T.

Some Complex Chlorides containing Gold. III. A New Cæsium Auric Chloride. HORACE L. WELLS (*Amer. J. Sci.*, 1922, [v], 3, 414—416).—This salt, $\text{Cs}_5\text{Au}_3\text{Cl}_{14}$, was obtained from a concentrated solution of cæsium chloride containing comparatively little auric chloride. It is best prepared in the presence of concentrated hydrochloric acid, because it is quickly decomposed by water, with the formation of a yellow precipitate of the usual double salt, CsAuCl_4 . The new complex salt was obtained as very minute deep red crystals. W. T.

Rubidium (Cæsium) Silver Gold Bromides. EUGEN SUSCHNIG (*Monatsh.*, 1921, 42, 399—404).—Triple bromides of rubidium (or cæsium), silver, and gold have been prepared corresponding with the triple chlorides described by Bayer (A., 1920, ii, 688) and discussed by Emich (*ibid.*, 689). The silver and gold are vicarious constituents, and their proportions in the compounds depend on their concentrations in the solutions from which crystallisation takes place. The compounds correspond with the formulæ $\text{Rb}_3\text{Ag}_x\text{Au}_{(2-x/3)}\text{Br}_9$ and $\text{Cs}_3\text{Ag}_x\text{Au}_{(2-x/3)}\text{Br}_9$. In the rubidium compound, the atomic proportion of silver varied from 0.77 to 1.59, and in the cæsium compound from 0.89 to 1.14. The rubidium salt forms almost black needles, deep violet in thin sections, whilst the cæsium salt is similar to Bayer's triple chloride. The salts can only be prepared in strong hydrobromic acid solution; they are decomposed by water. An analogous potassium salt could not be obtained, but microscopical tests showed that an ammonium compound is formed, and also compounds with organic bases such as guanidine, caffeine, pyridine, hexamethylenetetramine, and methyl- and ethyl-amine instead of rubidium or cæsium. The compounds are adapted to the microchemical identification of gold, silver, and rubidium or cæsium in absence of disturbing elements. Microchemical tests indicate that aluminium, chromium, zinc, nickel, cobalt, uranium, mercury, and manganese can replace silver in the triple chlorides and that ferric iron, calcium, strontium, and barium can replace the alkali metal. E. H. R.

Determination of the Werner Structure of Inorganic Compounds by means of X-Rays. PAUL SCHERRER and P. STOLL (*Z. anorg. Chem.*, 1922, 121, 319—320).—The space lattices of the following complex salts were determined: $[\text{PdBr}_6]\text{Rb}_2$, $[\text{Ni}(\text{NH}_3)_6]\text{Cl}_2$, and $[\text{PtCl}_6]\text{K}_2$. The six co-ordinated groups were placed at the six corners of a regular octahedron with the central atom at the centre. The salts resembled calcium fluoride with

the co-ordinated complex taking the place of the atom calcium. The platinum and chlorine were 1.6×10^{-8} cm. apart and the edge of the octahedron measured 9.7×10^{-8} cm. W. T.

Chloroiridates of Complex Metal Bases. A. BENRATH, W. BÜCHER, and H. ECKSTEIN (*Z. anorg. Chem.*, 1922, **121**, 347—360).—Jørgensen prepared a large number of salts of the amines of cobalt and chromium by combining the complex bases with different acid radicles. The complex acids of the noble metals formed beautifully coloured, crystalline salts. The present authors employed the complex hexachloroiridate as the acid radicle. The salts formed with cobalt as the central atom of the complex base were found to crystallise only with difficulty and to be more unstable than the corresponding platinum compounds. In the hexammine salts the most stable were those in which all the acid radicles were replaced by the complex iridium radicle; the others were very unstable. The hexammine and pentammine salts as well as the aquo-salts with four or five molecules of ammonia are insoluble, stable, yellow to brown in colour, decomposed only by long boiling with an alkali hydroxide, micro-crystalline and decomposed when heated alone at 150° . The tetrammine salts are more soluble, can be precipitated with alcohol, are crystalline and golden-yellow to bronze-red in colour. They decompose in hot water, and when heated alone at 105 — 120° . The preparation and properties of the following are described: Hexamminecobaltic hexachloroiridate, $[\text{Co}(\text{NH}_3)_6]_2[\text{IrCl}_6]_3$. Hexamminecobaltic nitrate hexachloroiridate, $[\text{Co}(\text{NH}_3)_6]\text{NO}_3[\text{IrCl}_6]$. Basic hexamminecobaltic hexachloroiridate, $[\text{Co}(\text{NH}_3)_6]\text{OH}[\text{IrCl}_6]$. Chloropentamminecobaltic hexachloroiridate, $[\text{Co}(\text{NH}_3)_5\text{Cl}][\text{IrCl}_6]$. Aquopentamminecobaltic hexachloroiridate, $[\text{Co}(\text{NH}_3)_5\text{H}_2\text{O}]_2[\text{IrCl}_6]_3$. *trans*-Dinitrotetramminecobaltic hexachloroiridate trihydrate, $[\text{Co}(\text{NH}_3)_4(\text{NO}_2)_2][\text{IrCl}_6] \cdot 3\text{H}_2\text{O}$.


cis-Dinitrotetramminecobaltic hexachloroiridate,
 $[\text{Co}(\text{NH}_3)_4(\text{NO}_2)_2][\text{IrCl}_6]$.

cis-Dichlorotetramminecobaltic hexachloroiridate,
 $[\text{Co}(\text{NH}_3)_4\text{Cl}_2]_2[\text{IrCl}_6]$.

Aquanitrotetramminecobaltic hexachloroiridate,
 $[\text{Co}(\text{NH}_3)_4(\text{H}_2\text{O})\text{NO}_2][\text{IrCl}_6]$.

The salts with chromium in the basic complex resemble the corresponding cobalt compounds. They are dirty yellow in colour. The hexammine and pentammine salts separated as very fine suspensions, the tetrammine salts crystallised much more readily. The following salts are described: Hexamminechromic hexachloroiridate, $[\text{Cr}(\text{NH}_3)_6]_2[\text{IrCl}_6]_3$. Hexamminechromic nitrate hexachloroiridate, $[\text{Cr}(\text{NH}_3)_6]\text{NO}_3[\text{IrCl}_6]$. Basic hexamminechromic hexachloroiridate, $[\text{Cr}(\text{NH}_3)_6]\text{OH}[\text{IrCl}_6]$. Hexamminechromic sulphate hexachloroiridate, $[\text{Cr}(\text{NH}_3)_6]_2(\text{SO}_4)_2[\text{IrCl}_6]$. Chloropentamminechromic hexachloroiridate, $[\text{Cr}(\text{NH}_3)_5\text{Cl}][\text{IrCl}_6]$. Hydroxo-aquo-tetramminechromic hexachloroiridate, $[\text{Cr}(\text{NH}_3)_4(\text{H}_2\text{O})\text{OH}][\text{IrCl}_6]$. Dithiocyanatotetramminechromic hexachloroiridate,
 $[\text{Cr}(\text{NH}_3)_4(\text{CNS})_2]_2[\text{IrCl}_6]$.

Dichloroaquatriamminechromic hexachloroiridiate,



The nitrate iron could not be separated from the chloroiridiate ion by means of nitron as the complex ion also forms an insoluble product, thus resembling the chloro-acids of the other noble metals. The compound, $(\text{C}_{20}\text{H}_{16}\text{N}_4)_2[\text{IrCl}_6]\text{H}_2$, separates as a dark red, crystalline precipitate. Ruthenium can be partly separated from the other noble metals by fractional crystallisation of the nitrous compounds, that of ruthenium being more soluble than the others.

W. T.

Mineralogical Chemistry.

Calcium Uranium Autunites. F. HENRICH (*Ber.*, 1922, 55, [B], 1212—1215; cf. A., 1918, ii, 96).—The detailed analysis is recorded of a calcium uranium autunite from Steinbruch Fuchsbau, near Leopoldsdorf, in the Fichtelgebirge, the analysis (I) of which agrees closely with the formula $\text{CaO} \cdot (\text{UO}_3)_2 \cdot \text{P}_2\text{O}_5 \cdot 10\text{H}_2\text{O}$. The ratio of radium to uranium is 2.8×10^{-7} , or 87% of the equilibrium value between the two elements. This appears to be the highest value yet recorded for a calcium uranium autunite. A Portuguese calcium uranium autunite from the Beira Alta Province (analysis II) had a ratio of radium to uranium which is only 38% of the theoretical value and was appreciably poorer in radium than the Bavarian specimen.

	UO ₃ .	CaO.	P ₂ O ₅ .	Insol.	Ignition.	Total.
I.	59.71	5.92	14.30	1.50	18.17	99.6
II.	60.72	6.04	14.55	0.53	18.27	99.93

H. W.

Ceruleofibrite, a New Mineral. EDW. F. HOLDEN (*Amer. Min.*, 1922, 7, 80—83).—As radiating tufts of bright-blue fibres and as blackish-blue, fibrous masses, this was found on specimens of cuprite from Bisbee, Arizona. The needles are orthorhombic ($a : b : c = 0.78 : 1 : 0.49$) with angles near to those of brochantite, d 3.54, H. 3. The optical characters are stated. The mineral is readily soluble in cold dilute hydrochloric acid. Analysis gives the formula $\text{CuCl}_2 \cdot \frac{1}{3}\text{Cu}_3\text{As}_2\text{O}_8 \cdot 6\text{Cu}(\text{OH})_2$, representing a very basic chloroarsenate.

CuO.	Cl.	As ₂ O ₅ .	H ₂ O.	Insol.	Total less O for Cl.
73.8	7.5	7.5	11.7	0.3	99.1

L. J. S.

Chemical Investigation of Japanese Minerals containing Rarer Elements. IV. Samarskite and an Unnamed Mineral from Ishikawa, Iwaki Province. YŪJI SHIBATA and KENJIRŌ KIMURA (*J. Chem. Soc. Japan*, 1922, 43, 301—312; cf. this vol., ii, 220).—Samarskite, d 5.50, hardness 5—6, from Ishikawa-machi, Iwaki Province, gave on analysis :

MgO.	CaO.	MnO.	FeO.	Al ₂ O ₃ .	Rare earths.	UO ₂ .	SnO ₂ .	TiO ₂ .	SiO ₂ .	Cb ₂ O ₅ .	Ta ₂ O ₅ .	H ₂ O.	Total.
0.25	1.25	0.84	9.64	0.61	17.34	16.87	0.73	0.10	0.37	41.11	10.00	0.85	99.96

From which is deduced the formula $2[M_2(\text{Cb}, \text{Ta})_2\text{O}_7]_n\text{R}_2(\text{Cb}, \text{Ta})_2\text{O}_8$, where M=Fe, Ca, Mg, Mn, UO, and R rare earth metals.

On spectroscopic examination, lines of lead, barium, strontium, tungsten, yttrium, erbium, dysprosium, gadolinium, samarium, scandium, and terbium were observed.

Imperfect black crystals, with resinous lustre of an unnamed mineral (d 6.2—6.4, hardness 5—6, streak, blackish-brown) from the same locality gave on analysis :

MgO.	CaO.	MnO.	FeO.	Al ₂ O ₃ .	Rare earths.	UO ₂ .	SnO ₂ .	SiO ₂ .	TiO ₂ .	Cb ₂ O ₅ .	Ta ₂ O ₅ .	H ₂ O.	Total.
1.07	0.86	0.40	11.78	0.87	8.40	21.88	1.20	0.30	0.21	36.80	15.00	0.89	99.66

On spectroscopic examination, lines of lead, barium, strontium, zinc, copper, bismuth, beryllium, yttrium, erbium, gadolinium, dysprosium, samarium, lanthanum, and scandium were observed.

K. K.

Composition of Thortveitite from Madagascar. CH. BOULANGER and G. URBAIN (*Compt. rend.*, 1922, 174, 1442—1443; cf. A., 1920, ii, 627; this vol., ii, 306).—The mean of five analyses is :

SiO ₂ .	Sc ₂ O ₃ .	ZrO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	Gl, Mn, Sn, Sb, Pb.	Total.
44.1	42.4	8.4	3.3	2.0	traces	100.2

The figures for scandia include about 0.5% of yttrium earths (yttrium, neoytterbium, and lutecium). The mineral is only slightly attacked by hydrofluoric acid or by fused bisulphates. Two or three fusions with sodium carbonate are necessary for its decomposition.

L. J. S.

Velardeñite from California. EARL V. SHANNON (*Proc. U.S. Nat. Mus.*, 1922, 60, Art. 22, 1—4).—A dull olive-green, coarsely-granular mineral with greasy to resinous lustre, from Tulare Co., California, is colourless under the microscope. It is optically uniaxial and negative, $\omega=1.657$, $\epsilon=1.653$. Analysis I corresponds with $14\text{CaO}, 2\text{MgO}, 5\text{Al}_2\text{O}_3, 9\text{SiO}_2$, which is interpreted as a mixture of five parts of velardeñite ($2\text{CaO}, \text{Al}_2\text{O}_3, \text{SiO}_2$) with two of åkermanite ($\text{MgO}, 2\text{CaO}, 2\text{SiO}_2$).


	SiO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	CaO.	MgO.	H ₂ O.	Total.
I.	27.88	25.52	1.59	0.43	40.86	4.18	0.34	100.80
II.	32.66	5.95	24.15	—	35.51	2.59	—	100.86

Mixed with it are grains of a yellowish-green garnet which (with some spinel) gave analysis II, indicating andradite.

L. J. S.

White Chlorites. EARL V. SHANNON and EDGAR T. WHERRY (*J. Washington Acad. Sci.*, 1922, 12, 239—241).—Colerainite (A., 1918, ii, 325) and sheridanite (A., 1912, ii, 1181) are classed as white chlorites. Analysis I is of a white chlorite from Brinton's quarry, Chester Co., Pennsylvania, II from Nottingham, Chester

Co., and III of a greenish-grey schistose rock from Miles City, Montana. I and III agree with the sheridanite formula



but II gives the ratios $9\text{MgO}, \text{Al}_2\text{O}_3, 6\text{SiO}_2, 7\text{H}_2\text{O}$. III agrees also with sheridanite in its optical constants, whilst II is different.

	SiO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	MnO.	CaO.	MgO.	H ₂ O-.	H ₂ O+.	Total.
I.	28.10	26.20	1.66	nil	trace	30.36	0.56	14.00	100.88	
II.	36.70	10.38	1.22	trace	trace	0.86	36.44	1.06	13.80	100.46
III.	27.78	24.30	1.43	0.35	nil	trace	32.71	0.06	13.01	99.64

L. J. S.

Crocidolite from Eastern Pennsylvania. EDGAR T. WHERRY and EARL V. SHANNON (*J. Washington Acad. Sci.*, 1922, 12, 242—244).—Blue pleochroic amphiboles ("glaucamphiboles") occur at many localities in eastern Pennsylvania. Analysis of material from Oley Line shows it to be a semimagnesian crocidolite with the formula $\text{H}_2\text{O}, \text{Na}_2\text{O}, \text{MgO}, \text{FeO}, \text{Fe}_2\text{O}_3, 6\text{SiO}_2$.

SiO ₂ .	TiO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	CaO.	MgO.	Na ₂ O.	K ₂ O.	H ₂ O-.	H ₂ O+.	Total.
51.62	2.27	0.92	18.36	10.93	0.48	5.92	5.62	0.66	1.04	2.57	100.39

Titanium is regarded as replacing aluminium and iron, rather than silicon, and to it the deep blue is ascribed. The low content of alkalis is accounted for by the replacement of sodium by hydrogen.

L. J. S.

Analytical Chemistry.

A New Method of Colorimetry. PAUL DOSNE (*Bull. Soc. Ind. Mulhouse*, 1922, 88, 73—77).—A beam of light (from a Nernst lamp or other suitable source) is directed upwards through a column of a solution of known strength of the coloured substance under investigation and the height of the column necessary for the complete absorption of the transmitted light, as observed through a spectroscope, is determined. Since under these conditions the colouring power of a substance is directly proportional to the height of the column and to the strength of the solution, the method allows various colouring matters to be compared when dissolved in water, alcohol, or other suitable solvents.

A. J. H.

Colorimetry. PAUL FLEURY (*Bull. Soc. Chim. Biol.*, 1922, 4, 223—232).—A discussion of the conditions under which colorimetric methods yield the most accurate results.

E. S.

Colorimetry. (MILLE) Y. GARREAU (*Bull. Soc. Chim. Biol.*, 1922, 4, 233—234).—The accuracy of the colorimetric method is indicated by the concordant results obtained during a large number of estimations of sugar by the picric acid method.

E. S.

Improved Form of Gas-analysis Apparatus. E. W. BLAIR and T. SHERLOCK WHEELER (*J. Soc. Chem. Ind.*, 1922, **41**, 187—188t).—A modification of the usual Bone-Wheeler apparatus is described, in which compressed air and reduced pressure are employed to alter the mercury levels; this saves much of the labour involved in raising and lowering the mercury reservoirs. The pressure of the air supply is about two atmospheres. In other respects, the method of working the apparatus is not altered.

W. P. S.

The Absorption Meter, an Apparatus for Gas Analysis. L. MOSER (*Z. anorg. Chem.*, 1922, **121**, 313—318).—The author describes a simplification of the apparatus he employed for the gravimetric estimation of hydrogen phosphide (A., 1922, ii, 393). The apparatus has been tested and gives good results in the estimation of sulphur dioxide, cyanogen, hydrogen cyanide, silicon fluoride, the hydrides of arsenic and antimony, ozone, and chlorine. It is unsuitable for carbon dioxide or any gases which requires an absorption liquid affected by the air (baryta water).

W. T.

Apparatus for Determining Hydrogen-ion Concentration. Application to the Detection of Mineral Acids in Vinegar. ANDRÉ KLING, A. LASSIEUR, and (MME) A. LASSIEUR (*Ann. Chim. analyt.*, 1922, **4**, 135—137).—A simple apparatus is described for determining hydrogen-ion concentration electrometrically. It is shown that the hydrogen-ion concentration of vinegars containing from 5—7% of acetic acid varies from p_H 2.54 to 2.83 and that the addition of 0.24% of sulphuric acid causes this value to become 1.48 to 2.02.

W. P. S.

The Clark Hydrogen-electrode Vessel and Soil Measurements. DANIEL J. HEALY and PERRY E. KANAKER (*Soil Sci.*, 1922, **13**, 323—328).—Satisfactory measurements of the hydrogen-ion concentration of fresh and air-dried soils can be made with the Clark hydrogen-electrode vessel. Air drying increases the values obtained over those of fresh samples.

A. G. P.

The Estimation of Chlorine in Organic Material. STEFAN VON BOGDÁNDY (*Z. physiol. Chem.*, 1922, **120**, 30—41; cf. A., 1913, ii, 426).—The author shows that if his method is carried out according to his instructions, high results are not obtained as is asserted by Gutmann and Schesinger (A., 1914, ii, 379).

S. S. Z.

Application of Amalgams in Volumetric Analyses. III. Estimation of Iodic, Bromic, and Chloric Acids. SUETARÔ KIKUCHI (*J. Chem. Soc. Japan*, 1922, **43**, 173—177).—Iodic acid reacts at 60° with tervalent titanium obtained by reduction of titanous acid as described previously (A., 1921, ii, 596), the excess of tervalent titanium being then titrated with iron alum. The reaction proceeds as follows: $IO_3' + 6Ti''' + 6H' = I' + 6Ti'''' + 3H_2O$. Bromic and chloric acids can be estimated by similar procedure.

K. K.

Sodium Peroxide. JAROSLAV MILBAUER (*Chem. Listy*, **16**, 84—86).—The estimation of the active oxygen of sodium peroxide by means of acid potassium permanganate solution is examined. The author finds that the best results are obtained by the addition of boric acid to the solution, as this fixes the hydrogen peroxide liberated by the acid. This method is very exact and rapid, and does not require the use of ice.

The effects produced by the presence of various metallic salts, and of spongy platinum and palladium are also investigated. It is found that if the salt is added to the water before the sodium peroxide, the results for the estimation of oxygen will be low; if the salt is added after the solution of the sodium peroxide in water, the results are high. This explains the failure of the use of permanganate solution alone, as the manganous sulphate formed acts as a catalyst, which must be neutralised by an anti-catalyst (in this case boric acid).

The iodometric method of Kingzett (T., 1880, **37**, 792) is also examined and found satisfactory.

The influence of catalysts on the method whereby the peroxide is analysed by estimation of the oxygen liberated by its decomposition in acid or alkaline solution is also studied. From the results obtained for this, the author formulates, as a method for the estimation of the active oxygen of sodium peroxide, its decomposition in alkaline solution, in the presence of cupric sulphate, and the measurement of the gaseous oxygen so liberated. The advantage of this method over that of Archbutt (A., 1895, ii, 185) is that the decomposition is complete in two minutes, and that the evolved oxygen is free from carbon dioxide.

By means of these new methods of estimation the author shows that the action of the atmosphere on sodium peroxide is principally due to the presence of moisture.

R. T.

Estimation of Sulphur Dioxide. G. BASIL BARHAM (*Chem. News*, 1922, **124**, 279—281).—Titration with permanganate solution under definite conditions is recommended for the estimation of sulphur dioxide. Dilute sulphuric acid solution is treated with a definite volume of standard permanganate solution, and the mixture then divided into two equal portions; the sulphur dioxide solution under examination is added to one portion and the permanganate solution is added until the coloration is again equal in intensity to that of the other portion, which has been reserved for comparison. To ensure the colorations being of the same tint, the acid solution should be reduced and oxidised once or twice (with sulphite and permanganate solutions, respectively) before it is used for the actual estimation.

W. P. S.

Examination of Foods for the Presence of Sulphites. A. CHASTON CHAPMAN (*Analyst*, 1922, **47**, 204—205).—When foods containing onions or mustard are acidified with phosphoric acid and distilled in a current of carbon dioxide, the distillate on oxidation with bromine will yield a distinct quantity of sulphuric acid; if the oxidation is made with hydrogen peroxide in place of bromine,

these foods will not yield any sulphuric acid unless they actually contain sulphites. W. P. S.

Factors Influencing the Estimation of Sulphates in Soil. C. T. HIRST and J. E. GREAVES (*Soil Sci.*, 1922, **13**, 231—249).—A comparison was made between the gravimetric method of estimating sulphates as barium sulphate and the volumetric chromate method. By the latter method an average of 97% of the sulphate in soil extracts could be recovered. Sodium and potassium chlorides had little effect on the results obtained; but in the presence of iron and aluminium salts the volumetric method tended to give low results. The presence of nitrates caused high figures by the gravimetric method and low ones by the volumetric method. A 1:5 water extract of the soils used, was found to be sufficient to remove all sulphates after forty minutes' shaking. Soils particularly rich in gypsum needed shaking with water for a longer period. Extracts should be clarified by a Pasteur-Chamberland filter.

Very small quantities of sulphates can be detected more easily by the chromate than by the gravimetric method. A. G. P.

The Analysis of Polythionates. ALBIN KURTENACKER and ALBERT FRITSCH (*Z. anorg. Chem.*, 1922, **121**, 335—343).—The authors have already found that tetrathionates react quantitatively with a cyanide to form thiosulphates (A., 1921, ii, 502). Trithionates were found by Raschig (A., 1920, ii, 756) to react quantitatively with cyanides in hot alkaline solutions to form sulphites. The latter reaction is incomplete in neutral solutions, whereas tetrathionates in alkaline solutions give sulphites as well as thiosulphates. This reaction is therefore unsuitable for the estimation of a mixture of these two polythionates. The estimation of a mixture of polythionates by means of a sulphite as recommended by Raschig was found to be unsatisfactory. The method of Riesenfeld and Feld (this vol., ii, 45) was tested. In the absence of other polythionates, the trithionate reacts quantitatively with copper sulphate, but the estimation is not applicable to mixtures. A tetrathionate boiled for half an hour with copper sulphate gave a precipitate of copper sulphide and sulphur in proportions corresponding with the decomposition of the tetrathionate as follows: $S_4O_6^{2-} = S_2O_6^{2-} + S$. The authors found, contrary to the results of Riesenfeld and Feld, that tetrathionates were quite stable; a solution showed no decomposition after keeping for fourteen days, the instability reported being due to impurities. W. T.

Pregl's Microanalysis. FRIEDRICH HOLTZ (*Ber.*, 1922, **55**, [B], 1496—1497).—The use of a bomb furnace in the estimation of halogen and sulphur is avoided by placing the tubes in brass tubes which are closed at one end and stoppered at the other and are heated in the vapour of boiling diphenylmethylamine, b. p. 295°.

The estimation of nitrogen in difficultly-combustible substances is effected in silica tubes which are filled in the following order: asbestos plug, oxidised copper wire (7 cm.), reduced copper wire

(11 cm.), asbestos plug, copper oxide powder mixed with the substance and a little potassium chlorate (4 cm.), oxidised copper wire (5 cm.). The tube is supported at either end on gutters lined with asbestos in such a manner that the portion of the tube containing the reduced copper is left exposed to the direct heat of the burner. This portion of the tube is doubly protected on its upper side. After displacement of the air by carbon dioxide, the reduced copper is raised to a white heat and after a second passage of carbon dioxide the heating of the remainder of the tube is effected very gradually in the direction towards the reduced copper. In this manner, the decomposition of the oxides of nitrogen is effected with certainty.

Substances may be conveniently dried in a hollow copper cube the walls of which are 1 cm. thick and 6.5 cm. long internally.

H. W.

A "Micro-Kjeldahl" Method of Estimating Nitrogen.

ARTHUR R. LING and WILLIAM JOHN PRICE (*J. Soc. Chem. Ind.*, 1922, 41, 149—151r).—A weighed portion of the substance containing 1.0—0.1 mg. of nitrogen is boiled until colourless with 8 c.c. of concentrated sulphuric acid, 1 gram of dry potassium sulphate, 0.02 gram of anhydrous copper sulphate, and 2 drops of 2.5% platinic chloride solution. After cooling and diluting with 15 c.c. of water, the liquid is distilled from a 300 c.c. flask with sufficient 40% sodium hydroxide solution to render it alkaline, and the ammoniacal distillate is collected in 50 c.c. of water in a graduated 250 c.c. flask. All the ammonia will have been expelled when 100 c.c. of liquid has distilled. The distillate is then "nesslerised," made up to 250 c.c., and the tint compared with a standard ammonium sulphate solution containing 1 mg. of nitrogen per c.c. The comparison will give the apparent nitrogen content which must be corrected by a factor (obtained from a curve given in the original paper) in order to arrive at the actual nitrogen content. This is due to the fact that intensity of colour and ammonia concentration are not directly proportional. G. F. M.

Kjeldahlisation of Mononitrophenols, Mononitrobenzoic Acids, and Mononitrocinnamic Acids. R. M. MARGOSCHES and ERWIN VOGEL (*Ber.*, 1922, 55, [B], 1380—1389).—A continuation of previous work (A., 1920, ii, 50), in which it was shown that the Kjeldahl-Gunning method gives accurate results only for the ortho-derivatives when applied to the mononitrophenols and mononitrobenzoic acids.

The Kjeldahlisation of mononitro-phenols and -benzoic acids by sulphuric acid without addition of potassium sulphate gives correct values for the ortho-, and low values for the meta- and para-compounds; it is remarkable that the latter are considerably higher than those given by the Kjeldahl-Gunning method.

Stepwise Kjeldahlisation according to Kjeldahl-Gunning shows that in the case of *o*-nitrophenol and *o*-nitrobenzoic acid, the major part of the nitrogen value is attained during the stage of the initial discoloration which occurs at the commencement of the heating

and the theoretical nitrogen value is almost attained at the period of most violent action. With *p*-nitrobenzoic acid, on the other hand, practically no nitrogen value is attained at the stage of initial darkening and the nitrogen is almost entirely present in such a form that the calculated nitrogen value is observed after addition of phenol; at the period of most violent action, the "converted" nitrogen increases somewhat, whereas that "convertible" by subsequent addition of phenol declines, ultimately becoming reduced to zero when the solution is colourless.

In the case of *p*-nitrobenzoic acid, the replacement of potassium sulphate in the Kjeldahl process by copper oxide or mercuric oxide leads to analytical results which are in harmony with the calculated values. The activity of copper oxide is not affected by the presence of potassium sulphate, whereas that of mercuric oxide is greatly depressed. The activity of sodium sulphate is inferior to that of potassium sulphate and approximately equal to that of lithium sulphate.

A further series of experiments is described in which the Kjeldahlisation of *p*- and *m*-nitrobenzoic acid and of *p*-nitrophenol is effected by the presence of potassium sulphate; the volatile products are collected and heated with concentrated sulphuric acid, phenol, and copper oxide. *p*-Nitrobenzoic acid and *p*-nitrophenol are found to behave in an entirely different manner; the former yields a volatile nitrogenous compound in which the nitrogen can be converted into ammonia by sulphuric acid, phenol, and copper oxide, whereas the latter undergoes a far more complex decomposition with the formation of products which do not yield ammonia under the described conditions.


In striking contrast to the nitrobenzoic acids, the three nitrocinnamic acids yield almost the whole of their nitrogen as ammonia when heated with sulphuric acid and potassium sulphate. Satisfactory results are also obtained with sulphuric acid alone or mixed with copper oxide.

H. W.

A Micro-method for the Estimation of Ammonia in Blood and in Organic Fluids. K. L. GAD-ANDRESEN (*J. Biol. Chem.*, 1922, 51, 367—372).—One c.c. of blood is used for the estimation. After addition of borate, it is evaporated to dryness by heating at 25° in a current of dry air. The ammonia thus removed is absorbed in sulphuric acid, decomposed by hypobromite, and the nitrogen measured in Krogh's microrespirometer as in the author's urea method (*A.*, 1920, ii, 646). The method is also applicable to secretions, but not to tissues.

E. S.

New Insoluble Organic Nitrates. JOSÉ GIRAL PEREIRA (*Anal. Fis. Quím.*, 1922, 20, 84—92).—In a study of organic compounds giving precipitates or turbidities with nitrates, the following compounds are mentioned as new precipitants: *p*-dianisylphenethylguanidine hydrochloride (acoine),



dicyanodiamide, $\text{NH}\cdot\text{C}(\text{NH}_2)\cdot\text{NH}\cdot\text{CO}\cdot\text{NH}_2$, produced by the hydrolysis of dicyanodiamidine, $\text{NH}\cdot\text{C}(\text{NH}_2)\cdot\text{NH}\cdot\text{CN}$; the hydrochloride

of hydrocupreine ethyl ether (optoquine), hydrastine hydrochloride. The common features in the constitution of nitrate precipitants is discussed. G. W. R.

So-called Di-9 : 10-monohydroxyphenanthrylamine suggested as a Reagent for Nitric Acid and Phenanthroxazine. B. FORESTI (*Gazzetta*, 1922, 52, i, 278—285).—The author has investigated Bamberger and Grob's phenanthroxazine (A., 1901, i, 280) and finds that the melting point $350\text{--}355^\circ$ given by these authors is that of a mixture of the compound with its decomposition products. The value obtained for the melting point depends on the initial temperature of the heating bath. With a bath at 422° (non-corrected), fusion occurs after the lapse of six seconds from the time of immersion, and since this is the length of time required for the tube and substance to assume the temperature of the bath, the melting point is taken as 422° .

It is found, further, that the substance obtained by Schmidt and Lumpp (A., 1910, i, 312; ii, 450) by the action of aqueous ammonia on dihydroxyphenanthrene, and described by these authors as di-9(10)-monohydroxyphenanthrylamine, consists of a mixture of phenanthroxazine with impure phenanthrazine and that the characteristic behaviour of the substance towards nitrates is due to the presence of the phenanthroxazine. In testing for nitrate, use is made of a solution of 0.01 gram of phenanthroxazine in 100 c.c. of concentrated sulphuric acid, and as dilution of this blue solution with water may give rise to a violet coloration similar to that obtained with nitrates, the substance to be tested must not be in aqueous solution. The dry residue left in a porcelain dish by a drop of the liquid to be examined is treated with a drop of the reagent; in presence of a nitrate, a reddish-violet coloration is formed, but if the amount of nitrate is very small, only a violet ring appears after some time. In this way 1×10^{-7} gram of nitric acid is detectable. This reaction is not disturbed by moderately high proportions of chlorate or chromate, and although slightly less sensitive than that of brucine with nitric acid, is not given by nitrous acid. The coloration is not destroyed by sulphur dioxide, but is changed to pale yellow by stannous chloride. T. H. P.

Estimation of the Nitro-group in Aromatic Organic Compounds. II. T. CALLAN and J. A. RUSSELL HENDERSON (*J. Soc. Chem. Ind.*, 1922, 41, 157—161T).—The low results obtained with certain compounds by the Knecht-Hibbert reduction method with titanous chloride for the estimation of nitro-groups is due in the case of readily chlorinated substances such as o-nitroanisole, α -nitronaphthalene, etc., to chlorination by the titanous chloride and the reducing action of the thereby liberated hydrogen, whereby a lesser amount of titanous salt is oxidised than would otherwise be the case. This source of error can be entirely obviated by using titanous sulphate solutions instead of the chloride, or may be very largely avoided even by using sulphuric acid instead of hydrochloric acid to acidify the titanous chloride solution. In general, the greater the excess of hydrochloric acid used the greater will be

the error due to chlorination. Another source of error, due to the volatility in steam of the simpler nitro-compounds, may be obviated by sulphonating, but much more simply by reducing under a ground-in reflux condenser about 9 inches long. By adopting these modifications and precautions, it is shown that, contrary to statements by English (A., 1920, ii, 776), nearly all types of nitro-compounds can be accurately estimated, including mononitro-hydrocarbons, chloronitro-compounds, and other negatively substituted nitrohydrocarbons. Recrystallised *p*-nitroaniline is recommended as the ultimate standard for the titanous solutions.

G. F. M.

Separation of Phosphoric Acid in Qualitative Analysis.

D. BALAREFF (*Z. anorg. Chem.*, 1922, **121**, 254—256).—The precipitates of the ammonium sulphide group (Groups III and IV taken together) are dissolved in hydrochloric acid (solution is thus free from cobalt and nickel) and the solution neutralised with ammonia. A large excess of lead nitrate is added and then sodium acetate until the solution is neutral to methyl-orange. The phosphoric acid is precipitated as lead phosphate, mixed with lead chloride and phosphates of ferric iron, chromium, and aluminium. The large excess of lead-ion, however, ensures that some ferric, chromium, and aluminium-ions remain in solution. The excess of lead is then removed by hydrochloric acid. The author claims that this method possesses advantages in speed and completeness of separation over the older methods.

W. T.

Effect of Adsorption by Stannic Sulphide on the Estimation of Phosphates. ŠUETARŌ KIKUCHI (*J. Chem. Soc. Japan*, 1922, **43**, 329—333).—When phosphate is determined by the usual method, it is found after the tin in solution is removed as sulphide, that there is a deficiency of phosphate. This is due to adsorption of the phosphate by stannic sulphide; stannous sulphide, on the contrary, does not adsorb phosphate; therefore the quadrivalent tin in the solution is reduced to the stannous state by zinc in the presence of carbon dioxide and removed as sulphide. Estimation of the phosphate in the filtrate then gives correct values. K. K.

The Citric Solubility of Mineral Phosphates. J. F. TOCHER (*J. Agric. Sci.*, 1922, **12**, 125—143).—The proportion of phosphoric acid extractable by citric acid from basic slags and mineral phosphates is closely dependent on the conditions of experiment. The three variables being the amount of citric acid used, the amount of mineral phosphate used, and the volume of liquid used, the author has investigated the influence on the amount of phosphoric acid extracted of varying one of these factors whilst keeping the other factors in a constant ratio. With constant weights of mineral phosphate and citric acid, the amount of phosphoric acid dissolved increases with the volume. With constant weight of mineral phosphate and constant volume, the amount of phosphoric acid dissolved increases with increase of the amount of citric acid used. With constant weight of citric acid and constant volume, the

amount of phosphoric acid extracted decreases with increase in the amount of mineral phosphate used. The quantitative relationships are connected with considerations of mass action and it is shown that the solubility curves obtained can be made to fit an empirical formula. It is held that citric solubility is influenced by so many factors that it is an untrustworthy test of the agricultural value of mineral phosphates and slags. The only standard tests to be accepted are total phosphate content, fineness of grinding, and freedom from injurious substances. G. W. R.

Invisible "Mirrors" of Arsenic, Antimony, and Bismuth. HERMANN SCHEUCHER (*Monatsh.*, 1921, 42, 411—420).—Bettendorff's test for arsenic, consisting in reducing an oxide of arsenic with stannous chloride and hydrochloric acid, when arsenic is thrown down as a voluminous, brown precipitate, has been developed into an extremely sensitive microchemical test. The reactants are introduced into a capillary tube which is then sealed to a point at one end and heated for a short time in a bath of boiling amyl alcohol (130°). The tube is then centrifuged and the arsenic appears as a brown precipitate in the point of the capillary. In this way, 0.01γ of arsenic ($\gamma=0.001$ mg.) can be detected at a dilution of 1 in 250,000. By contrast, the magnesium ammonium arsenate test is uncertain with 0.1γ of arsenic. The test is not interfered with by antimony, tin, lead, copper, or cadmium. The Marsh test, using Lockemann's method (A., 1905, ii, 353), is sensitive to 0.1γ , whilst with less arsenic the mirror becomes scarcely visible, and invisible with 0.04γ . If, however, the glass capillary containing the mirror, or supposed mirror, is treated with bromine vapour to oxidise the arsenic, and the Bettendorff test is then applied as above, as little as 0.01γ of arsenic can be detected.

The Marsh test applied to antimony is sensitive to 0.2γ , but the presence of antimony in invisible mirrors can be demonstrated by the luminescence test described by Paneth (A., 1919, ii, 68). Bismuth can be detected in a similar way when the hydride is prepared by Paneth's method. The mirror is dissolved in a drop of nitric acid, which is then transferred to a piece of ignited calcium carbonate held in a platinum loop and the flame test applied. For antimony, the limit of sensitiveness is 0.001γ . An invisible mirror can also be "developed" with a solution of ferrous sulphate in acetic acid containing silver nitrate. After a short time, silver is deposited in the place where the mirror should be. The reaction is extraordinarily sensitive, but, unfortunately, a glass tube which has merely been ignited with only a current of air passing through may give the reaction, although a quartz tube will not do so.

E. H. R.

Colorimetric Estimation of Arsenic by means of Quinine Molybdate. D. CHOUGHAK (*Ann. Chim. analyt.*, 1922, 4, 138—142).—A solution containing a trace of arsenic is oxidised with nitric acid, evaporated to dryness, and the residue treated with 0.5 c.c. of 17.5% nitric acid, 4.5 c.c. of water, and 20 c.c. of quinine molybdate reagent; the turbidity produced is compared with that

given by a known amount of arsenic under similar conditions. As little as 0.00002 mg. of arsenic gives a visible turbidity. The reagent is prepared by dissolving 0.5 gram of quinine hydrochloride in 10 c.c. of water, adding 5 c.c. of arsenic acid solution (containing 0.01 mg. of arsenic per c.c.), 10 c.c. of 17.5% nitric acid, and 1 c.c. of sodium molybdate solution (3.5 grams of sodium carbonate and 9.5 grams of molybdic acid per 100 c.c.); after a few minutes, the mixture is diluted with water to 120 c.c., and filtered through a filter-paper moistened previously with nitric acid and washed. If heavy metals are present in the solution under examination, the arsenic must be separated by distillation with hydrochloric acid, potassium bromide, and hydrazine sulphate, the distillate oxidised with nitric acid, evaporated to dryness, and the dry residue used for the estimation.

W. P. S.

Inadequacy of the "A.R." test for Alkalis in Calcium Carbonate. W. SINGLETON and H. WILLIAMS (*Analyst*, 1922, 47, 252—254).—Only about 50% of the total extractable sodium carbonate is washed out on boiling 5 grams of calcium carbonate with 50 c.c. of water for ten minutes. The alkalis present in 5 grams of "A.R." calcium carbonate were therefore determined by the Lawrence Smith method (this Journal, 1871, 442) and the amount of sodium carbonate obtained was more than ten times that given by the "A.R." method. The result was checked by decomposing 5 grams of the calcium carbonate with hydrochloric acid and precipitating the calcium twice with ammonium carbonate and ammonium oxalate. Substantially the same result was obtained. It therefore appears that only about 10% of the alkali carbonates present in the calcium carbonate are removed by one extraction with water.

H. C. R.

A Comparison of the Calcium Content of some Virgin and Cultivated Soils of Kentucky by an Improved Method for the Estimation of this Element. O. M. SHEDD (*Kentucky Agr. Exp. Sta. Bull.*, 236 (October 1921)).—In estimating the total calcium in the soils, 1 gram of soil (passing through a 100-mesh sieve) is fused with 5 grams of a mixture of sodium and potassium carbonates (10 : 13) for ten minutes. The fused mass is dissolved in hydrochloric acid, evaporated to dryness, and the silica dehydrated. The residue is extracted with dilute hydrochloric acid and the silica filtered off. Filtrate and washings should not exceed 100 c.c. The solution is made just alkaline to litmus with strong ammonia and then just acid with hydrochloric acid. The solution, which is not always perfectly clear at this stage, is heated to boiling and the calcium precipitated by adding 1—2 grams of solid ammonium oxalate. Further addition of hydrochloric acid may now be necessary to maintain the solution acid to litmus. After two to three minutes' boiling, the solution is placed on the steam-bath for a few hours and allowed to remain over-night. The precipitate is filtered, washed twice with water, and ignited to convert it into oxide or carbonate. This is redissolved in acid and manganese precipitated by the addition of bromine

and excess of ammonia. After heating until only a little ammonia remains, the solution is made faintly acid with oxalic acid and filtered. The calcium is then precipitated from the filtrate as before and finally weighed as oxide.

A. G. P.

Flame Reactions of Thallium. JACOB PAPISH (*Proc. Indiana Acad. Sci.*, 1918, 166—169).—Thallous chloride dissociates in a Bunsen flame to the free metal which colours the flame and may be precipitated on a cold surface as a brown mirror. In a hydrogen flame in air, thallous chloride produces five zones of luminescence. Metallic thallium will condense from the innermost cone on a cold surface but not from the outer cone where the luminescence is probably due to oxidation of the thallium.

CHEMICAL ABSTRACTS.

Tervalent Titanium. II. Estimation of Copper and Iron in the Presence of each other. WILLIAM M. THORNTON, jun. (*J. Amer. Chem. Soc.*, 1922, 44, 998—1001; cf. A., 1921, ii, 279).—A method is described for the estimation of iron and copper in the presence of one another by means of a standard solution of titanium trisulphate. The method is as follows. The mixture of cupric and ferric sulphates is dissolved in less than 100 c.c. of water, acidified with 10 c.c. of 1:1-sulphuric acid, and cooled to 15°. A quantity of 10% ammonium thiocyanate solution from 2.5 to 5 c.c., depending on the amount of copper anticipated, is added and the solution titrated with a standard solution of titanium trisulphate until the pink colour just vanishes. The titre gives the sum of the copper and iron present. The whole is then heated to incipient ebullition, to coagulate the cuprous thiocyanate, cooled, and filtered. The precipitate is thoroughly washed with cold water and the filtrate and washings are cooled to 15°. The solution has by this time probably become pink again, due to oxidation by air. If so, the colour is bleached by the careful addition of titanium trisulphate, and sufficient silver nitrate (2.5—5 c.c. of 25% solution) is added to precipitate the whole of the thiocyanate. The ferrous iron is then titrated with a standard solution of potassium permanganate. The method is shown to be accurate and trustworthy, but the precaution of previously testing the titanium solution for iron must be taken, and if such is found the permanganate value corrected for this amount.

J. F. S.

Quantitative Analysis of Aluminium Alloys, especially of Duralumin. E. MENDES DA COSTA-VET (*Chem. Weekblad*, 1922, 19, 249—251).—The alloy of aluminium introduced for airships by the Dürerer Metallwerke contains Cu 3.5—4.5%, Mn 0.5—1.0%, Mg 0.5%. The methods of Collit and Regan (A., 1918, ii, 175) and of Withey (A., 1916, ii, 538) for analysis of this alloy are unsatisfactory, since in the first the magnesium is almost completely carried down in the ammonia precipitation of iron, manganese, and aluminium, whilst in the second it is impossible to precipitate the magnesium from the tartaric acid solution by means of pyrophosphate.

It was found that magnesium can best be estimated after removal of copper by means of hydrogen sulphide, and of iron, aluminium, and part of the manganese by means of ammonium chloride and ammonia, the latter separation being performed twice, by precipitation with phosphate. Some of the manganese remains from the ammonia precipitation, and is weighed with the magnesium as pyrophosphate; this is estimated by redissolving the precipitate and titrating with permanganate.

The ammonia precipitation must be carefully carried out, and the liquid filtered immediately to avoid separation of magnesium at this stage.

S. I. L.

The Quantitative Estimation and Separation of Aluminium.

GERHART JANDER and ERWIN WENDEHORST (*Z. angew. Chem.*, 1922, 35, 244—247).—Precipitation of large amounts of aluminium hydroxide from solutions containing other metals such as magnesium and manganese leads to such high results that the direct estimation of aluminium in alloys consisting chiefly of aluminium is not possible by the usual methods. In such cases, the authors recommend heating the alloy in a current of well-dried hydrogen chloride at 200°, whereby all the aluminium sublimes as chloride together with small amounts of silicon, magnesium, and manganese chlorides, whilst iron, copper, and zinc remain in the boat. The sublimate is dissolved in dilute nitric acid and the solution evaporated repeatedly with further quantities of the same acid until all chlorine is expelled. The silica that separates is removed by filtration and the solution is evaporated to dryness. The residue is ignited, gently at first, then strongly, and weighed as aluminium oxide containing some manganese and magnesium oxides, which are determined separately, the former colorimetrically and the latter as pyrophosphate, and deducted. [Cf. *J. Soc. Chem. Ind.*, 1922, July.]

A. R. P.

A Proposed Estimation of Chromium and Nickel in Steel.

FRITZ SIMION (*Chem. Ztg.*, 1922, 46, 504).—The sample is dissolved in aqua regia, the solution neutralised and poured into an excess of a solution of sodium hydroxide containing hydrogen or sodium peroxide. After boiling to expel oxygen, the precipitate is collected on a Nutsche filter, well washed, then returned to the beaker and digested with ammonia and ammonium carbonate to extract the nickel. The chromate in the filtrate is estimated by any suitable method. [Cf. *J. Soc. Chem. Ind.*, 1922, July.]

A. R. P.

Applications of Amalgams in Volumetric Analyses. IV.

The Uses of Cadmium Amalgam. NAOTSUNA KANÔ (*J. Chem. Soc. Japan*, 1922, 43, 333—340).—In earlier papers (A., 1921, ii, 596, 714), the volumetric estimation of molybdenum, titanium, etc., after reduction by means of liquid zinc amalgam was described. To extend the method, the zinc is now replaced by cadmium. The method of preparation of the amalgam and the reducing operation are practically the same as in the former case. In the cases of molybdenum and iron, the cadmium amalgam acts

as well as the zinc amalgam, but in the cases of vanadium and titanium, the reduction is not complete. With uranium, it is necessary to carry out the reduction in an atmosphere of carbon dioxide.

K. K.

The Estimation of Uranium in Presence of Phosphoric Acid. A. SCHOEP and W. STEINKUHLER (*Bull. Soc. chim. Belg.*, 1922, **31**, 156—159).—The analysis of dewindtite (this vol., ii, 305) and of stasite (this vol., ii, 386), phosphates of uranium, and lead, necessitated more satisfactory methods than those hitherto used. The following procedure has given trustworthy results for both phosphate and uranium in a series of control experiments. Silica and elements which are precipitated by hydrogen sulphide in acid solution are first eliminated, the uranium is oxidised by means of nitric acid, and the phosphoric acid then precipitated with molybdate solution. Addition of ammonia transforms the uranium into ammonium uranate; from this the metal is precipitated as uranium sulphide on the addition of ammonium sulphide, the precipitate being washed until the washings are free from molybdenum. No solution and reprecipitation is necessary.

H. J. E.

Iodometric Studies. II. BOHDAN KÖHLER (*Chem. Listy*, 1921, **16**, 87—91; cf. A., 1921, ii, 410).—The best results in the iodometric titration of tin are obtained by dissolving the tin in 6*N*-hydrochloric acid, adding to this a known amount of iodine dissolved in potassium thiocyanate, and determining the excess of iodine with a standardised sodium thiosulphate solution.

Contrary to the observation of Zulkowski and Bruhns (*J. pr. Chem.*, 1859, **103**, 351), the velocity of the reaction $\text{Cr}_2\text{O}_7^{''} + 14\text{H}^+ + 6\text{I}^- \rightarrow 2\text{Cr}^{+++} + 7\text{H}_2\text{O} + 6\text{I}$ is quite considerable.

The titration of tin by means of potassium dichromate solution gives exact results if atmospheric oxygen is excluded from the solutions. Since it is, in practice, difficult to have air-free solutions of potassium dichromate, a correction of +0.068 mg. of tin per c.c. of aqueous aerated potassium dichromate solution is advocated. This correction is based on the assumption that the solubility of oxygen in decinormal potassium dichromate solution is little different to that in pure water.

R. T.

Reduction of Vanadic Acid Solutions with Mercury. LE ROY W. MCCAY and WILLIAM T. ANDERSON, jun. (*J. Amer. Chem. Soc.*, 1922, **44**, 1018—1021; cf. this vol., ii, 217).—When vanadic acid in the presence of sulphuric or hydrochloric acid is vigorously shaken with mercury the vanadium is reduced to the quadrivalent state according to the equation $2\text{HVO}_3 + 2\text{Hg} + 3\text{H}_2\text{SO}_4 = 2\text{VOSO}_4 + \text{Hg}_2\text{SO}_4 + 4\text{H}_2\text{O}$. If the solution before shaking is mixed with a little more sodium chloride than is equivalent to the mercurous sulphate formed, the whole of the mercury is precipitated as mercurous chloride and a pure blue solution left which may be quantitatively titrated with potassium permanganate when the reaction,

represented by the equation $10\text{VOSO}_4 + 2\text{KMnO}_4 + 12\text{H}_2\text{O} = 10\text{HVO}_3 + \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 7\text{H}_2\text{SO}_4$ takes place. A method of estimating vanadic acid by means of the above reactions is described. The solution of vanadic acid containing sulphuric acid and the requisite amount of sodium chloride (0.3 gram for each 50 c.c. of *N*/10-permanganate required) is diluted to 100 c.c., placed in a stoppered bottle, and vigorously shaken with 20 c.c. of mercury for five minutes. The liquid above the grey mercury mixture is decanted into a small beaker and poured through a suction filter. The residue in the bottle is washed four times by decantation, 20 c.c. of water being used for each washing, and the washings are severally decanted into the beaker and then through the filter. The total filtrate is diluted to 250 c.c., heated at 80–90°, and titrated with permanganate to the faintest pink. The whole process may be completed in thirty minutes; it gives results which are identical with those obtained by the sulphur dioxide method, and since uranic acid and arsenic acid are not reduced by mercury, the method may be employed in the presence of these acids.

J. F. S.

Detection of Antimony in Analysis. TH. SABALITSCHKA and H. SCHMIDT (*Ber. deut. Pharm. Ges.*, 1922, **32**, 132–135).—To obviate the necessity of using platinum foil, antimony may be detected in the solution of the sulphides of antimony and tin in concentrated hydrochloric acid as ordinarily obtained in the course of analysis, by diluting a portion of this solution with an equal volume of water, adding a piece of arsenic-free zinc to the cooled liquid in a test-tube, and allowing the issuing gases to impinge on a piece of filter-paper freshly moistened with a 10% solution of silver nitrate. In presence of antimony the paper will become quite black on the under side within two minutes, or immediately if considerable quantities are present. A slight brown coloration is to be disregarded, and it is important that the silver solution is not too concentrated, otherwise small amounts of hydrogen antimonide may cause only a brownish-yellow coloration instead of black, as is always obtained with dilute silver solutions.

G. F. M.

Detection of Benzene in Light Petroleum. P. SCHWARZ (*Chem. Ztg.*, 1922, **46**, 401).—If 5 c.c. of pure light petroleum are added to 2 c.c. of a mixture consisting of equal volumes of aniline and 95% alcohol, the aniline separates as a layer; should as little as 5% of benzene be present in the light petroleum, the aniline does not separate from the mixture and a clear solution is obtained. If desired, the light petroleum may be fractionated previously, and the test applied to the fraction having b. p. 80° to 110°.

W. P. S.

The Detection and Estimation of Acidity and Alkalinity in Cotton Fabrics. HUBERT FRANK COWARD and GLADYS MARY WIGLEY (*Trans. Text. Inst.*, 1922, **13**, 121–126).—The testing of cotton fabrics for traces of strong acids, such as might

give rise to "tendering," is complicated on the one hand by the fact that cotton exhibits a preferential attraction for the basic constituent of neutral salts and thus produces an acid reaction in their aqueous solutions, and, on the other, by the fact that cotton retains absorbed mineral acids so tenaciously that they can only be completely removed by repeated extractions with boiling water. It is now shown that the first factor is of far less moment in the titration of acid in the presence of the fabric than the second factor would be in a titration of the extract. Acidimetry of bleached cloth, correct to 0.01% of hydrochloric or sulphuric acid, and alkali-metry, correct to 0.02% of soda, calculated as sodium hydroxide, can be carried out by titrations with 0.02*N*-solutions, at the boil, in the presence of the fabric, using phenolphthalein as indicator. A fabric which contains up to 0.1% of sulphuric acid might, however, retain up to 0.05% after two extractions with boiling water.

The use of organic indicators, and a mixture of potassium iodide, iodate, and starch, for "spot" tests on the fabric is also described, and the sensitivity limits are indicated. Methyl-red is responsive to about 0.005% of strong acid or alkali. Difficulties arising from the presence of weak acids, soap, "finishing" compositions, and dyes are briefly discussed, but methods for testing raw cotton fabrics have still to be investigated.

J. C. W.

Estimation of Volatile Fatty Acids. G. WIEGNER and J. MAGASANIK (*Mitt. Geb. Lebensmitt. Hyg.*, 1919, 10, 156—174; from *Bied. Zentr.*, 1922, 51, 140—145).—In the distillation of aqueous solutions of volatile fatty acids the following relationship is established, $(\log y_1 - \log y_2)/(\log x_1 - \log x_2) = \text{constant}$, where y_1, y_2 , are the amounts of acid in terms of c.c. of standard alkali required for neutralisation of the residue at the beginning and end of a distillation interval and x_1, x_2 , the corresponding amounts of water. Values of the constant are given for the volatile fatty acids. The formula is applied to the estimation of volatile acids in silage. An example is given in which an aqueous extract containing acetic and butyric acids is made up to a known volume and distilled to half volume, the operation being repeated three times. From the titre of the distillates the acid value of each successive residue is obtained, whence by calculation the amounts of acetic and butyric acids respectively in the original extract are obtained. The method may be applied to the estimation of volatile acids in wines. Analytical figures are given for the amounts of acetic, butyric, and lactic acids in different kinds of silage.

G. W. R.

Analysis of Acetic Anhydride. A. RECLAIRE (*Perf. Essent. Oil Rec.*, 1922, 13, 148—149).—Ten grams of acetic anhydride are weighed out into a 250 c.c. flask, diluted with 75 c.c. of water free from carbon dioxide, and heated on a water-bath for fifteen minutes under a reflux condenser. The product is diluted to 500 c.c. with a further quantity of carbon dioxide-free water, and 50 c.c. are titrated with *N*/2-potassium hydroxide solution free from carbonate. The percentage of acetic anhydride present is then (c.c. *N*/2 KOH used $\times 17.144$)—570.45. The necessity for using

* water and alkali free from carbon dioxide is emphasised. The latter is readily obtained by mixing 1000 c.c. of ordinary potassium hydroxide solution with 50 c.c. of milk of lime, and after keeping one or more days the clear liquid is syphoned off and standardised.

G. F. M.

Detection and Estimation of Free Sulphuric Acid and Sulphoacetates in Cellulose Acetates. M. ENTAT and E. VULQUIN (*Ann. Chim. analyt.*, 1922, 4, 131—135).—To estimate free sulphuric acid, 10 grams of the cellulose acetate are digested for thirty minutes with 200 c.c. of water at 15°, the mixture is then filtered, and the filtrate is titrated with standardised barium hydroxide solution, the end-point being determined electrometrically. There is a sharp break in the titration curve when all the free sulphuric acid has been neutralised. None of the samples examined by the authors contained free sulphuric acid, and it is suggested that this acid is not present in normal cellulose acetates, the quantities found by other observers being actually due to the presence of alkali or alkaline-earth sulphates, acetic acid giving the acid reaction. Cellulose acetate prepared by methods involving the use of sulphuric acid as catalyst invariably contains sulphoacetic acid combined with the cellulose ester. The sulphoacetate is estimated by heating 10 grams of the cellulose acetate with 50 c.c. of water in an autoclave at 125° for five hours; the sulphuric acid resulting from this hydrolysis is then titrated electrometrically. The amount of sulphuric acid present as sulphoacetate in well-prepared cellulose acetate is always less than 0.03%. W. P. S.

Determination of the Iodine Value of Aliphatic and Aromatic Unsaturated Compounds. D. HOLDE, P. WERNER, IDA TACKE, and C. WILKE (*Chem. Umschau*, 1922, 29, 185—188).—The determination of the iodine value is a valuable criterion of the purity of higher unsaturated fatty acids and their derivatives when other methods, such as molecular weight and melting point determinations, fail, and has been used by Holde and Wilke in the case of erucic acid (this vol., i, 317). For this purpose, the Hanus reagent (10 grams of iodine monobromide in 500 c.c. of glacial acetic acid) is preferred, as it gives results in close agreement with those obtained with Hübl's reagent, and is more easily prepared, more stable, and more rapid in action, fifteen minutes being sufficient, with about 50% excess of the reagent, for the completion of the reaction. Accurate results are also obtained with aromatic unsaturated compounds, for example, saffrole, under similar conditions. Nevertheless, as is well known, anomalous results are given by the Hübl-Waller reagent, and by Wijs's reagent for cholesterol and phytosterol, the former giving abnormally low, and the latter abnormally high results. With the original Hübl solution, however, consistent, although slightly high results (73—77 compared with 65.7 theoretical) were obtained with cholesterol, but with phytosterol values varying from 41 to 76 were obtained, according to the duration of the reaction and the excess of Hübl solution employed. With both alcohols, Wijs's solution gave values of

about 135, that is, approximately double the theoretical. A similar difference in the results obtained with the Hübl-Waller and with Wijs's reagents has been observed with mineral lubricating oils and with naphthenic acids, which may indicate the presence of cholesterol derivatives in these products. G. F. M.

Estimation of the Various Acids of Wine. TH. VON FELLENBURG (*Z. Unters. Nahr. Genussm.*, 1922, 43, 217—255).—The author has developed a scheme of analysis by means of which the separate proportions of weak organic acids (tannic acids) and of tartaric, malic, succinic, and lactic acids may be estimated. [Cf. *J. Soc. Chem. Ind.*, 1922, 514A.] T. H. P.

A Colorimetric Method for the Estimation of Homogentisic Acid in Urine. A. P. BRIGGS (*J. Biol. Chem.*, 1922, 51, 453—454).—The method depends on the reduction of phosphomolybdic acid by homogentisic acid (cf. Bell and Doisy, A., 1920, ii, 769). The diluted urine is treated first with a solution containing 5% of ammonium molybdate in 5*N*-sulphuric acid and then with a 1% solution of potassium dihydrogen phosphate. The colour produced is compared with a quinol standard, experiments having shown that 1 mg. of quinol is equivalent to 0.79 mg. of homogentisic acid. Albumin and sulphides interfere with the reaction, but these may be removed, if present, by means of trichloroacetic acid and silver sulphate. E. S.

Estimation of the Iodine-Bromine Number without using Potassium Iodide. L. W. WINKLER (*Z. Unters. Nahr. Genussm.*, 1922, 43, 201—204).—The use of potassium iodide in this estimation can be avoided by treating the oil or fat with excess of bromine, adding a solution of arsenious oxide, and titrating the excess of the latter with potassium bromate solution. A blank estimation is carried out, using the same quantities of the reagents as for the actual estimation and allowing them to react for the same length of time. The results are almost identical with those obtained from using potassium iodide solution according to the original method (*ibid.*, 1916, 32, 358). H. C. R.

Use of Semi-microchemical and Microchemical Methods in Place of Macrochemical Methods in the Analysis of Foods. H. LÜHRIG (*Pharm. Zentr.-h.*, 1922, 63, 218—221, 227—232).—The Reichert-Meissl, Polenske, saponification, and iodine numbers of fats and oils may be estimated on quantities of substance varying from 0.02 to 0.2 gram. The procedure employed is similar to that used in the methods in their usual form and is described in detail; in the case of the Reichert-Meissl number, the distillation requires about five minutes, 10 c.c. of distillate are collected (when 0.1 gram of fat is taken), and the result obtained must be multiplied by 0.9 to obtain the true Reichert-Meissl number. W. P. S.

The Constants of Indian Beeswax. O. D. ROBERTS and H. T. ISLIP (*Analyst*, 1922, 47, 246—251).—Tables are given of the constants of twenty-three samples of honeycomb and wax

collected under the supervision of District Officers in Bengal, Eastern Bengal, and Assam. The low acid values obtained are remarkable, being in many cases lower than 6, the figure hitherto accepted as the lowest representing unadulterated Indian beeswax. Both Weinwurm's test and Salamon and Seaber's test are shown to be untrustworthy.

H. C. R.

The Estimation of Aldehydes and Ketones by means of Hydroxylamine. A. H. BENNETT and F. K. DONOVAN (*Analyst*, 1922, 47, 146).—The method originally proposed for the estimation of citral in lemon oil (A., 1909, ii, 192) can be applied to other aldehydes and ketones. With formaldehyde, acetone, and benzaldehyde, very close results are obtained. Carvone requires rather more carefully standardised conditions. The analyses of citronellal and citral are slightly less accurate. No satisfactory results could be obtained for camphor. The hydroxylamine hydrochloride used for this process should be recrystallised from water, and not from alcohol.

A. G. P.

The System Camphor-Alcohol-Water in Relation to the Titration of Camphor-spirit. K. SCHERINGA (*Pharm. Weekblad*, 1922, 59, 389—395).—The proportions of camphor, alcohol, and water which can remain in equilibrium in a single-phase system at various temperatures have been determined, and the results expressed graphically on the ordinary equilateral triangle presentation. Known solutions of alcohol and camphor were treated with water until a permanent separation occurred, and the increase in weight was then determined.

It is shown that many mixtures can be prepared to satisfy the requirements laid down in the Pharmacopœia, and that in analysing the preparation as recommended by titration with water the temperature must be kept between 20° and 25°, and also that the amount of camphor which can be added and held in solution at 15° should be specified.

S. I. L.

The Detection of Vernin. H. STEUDEL and R. FREISE (*Z. physiol. Chem.*, 1922, 120, 126—129).—Vernin when boiled with acetic anhydride containing a small particle of sodium acetate and then cooled yields lustrous, pale rose needles, which after twice recrystallising from absolute alcohol give colourless prismatic needles of triacetylvernin, m. p. 226°, $[\alpha]_D^{25} + 2.306^\circ$.

S. S. Z.

The "Cotton Wool Plug" Test for Indole. S. N. GORE (*Indian J. Med. Research*, 1921, 8, 505—507).—The under-surface of the plug is evenly moistened, first with a few drops of a solution of 1 gram of *p*-dimethylaminobenzaldehyde in 95 c.c. of absolute ethyl alcohol and 20 c.c. of hydrochloric acid, and then with a few drops of a solution of 1 gram of potassium persulphate in 100 c.c. of distilled water. The tube containing the plug, pushed down to about an inch above the surface of the broth culture, is placed upright in a gently boiling water-bath and heated for about fifteen minutes. Indole, when present to so small an extent as

0.0005 mg. per c.c. of the broth culture, volatilises and develops on the plug a rose colour which is viewed through the tube.

CHEMICAL ABSTRACTS.

The Detection and Estimation of Monoamino-acids. R. ENGELAND (*Z. physiol. Chem.*, 1922, 120, 130—140).—The author estimated proline by means of his betaine method in glutin and elastin. The former protein showed a content of 10.9% and the latter of 4.3% of proline. The same method was applied in the case of ascitic fluid. Six and a half litres of fluid yielded 0.06 gram. of the chloraurate of the betaine of leucine and 0.025 gram. of the chloraurate of the betaine of lysine by this method. S. S. Z.

Colorimetric Estimation of the Amino-acid Nitrogen in Normal Urine. OTTO FOLIN (*J. Biol. Chem.*, 1922, 51, 393—394).—Ammonia is removed by shaking with permutedite and the estimation is then carried out as for blood (cf. this vol., ii, 540). E. S.

The Manipulations Preparatory to the Estimation of Amino-acids in Blood. FERNAND PETITJEAN (*Bull. Soc. Chim. Biol.*, 1922, 4, 108—114).—For the preparation of blood for the estimation of its amino-acid content by Van Slyke's method, the author recommends, from the numerous methods which have been proposed, decomposition of urea by means of the soja bean, followed by precipitation of proteins by the addition of trichloroacetic acid. Details of the technique used are given in the original. E. S.

Estimation of Urea from the Carbon Dioxide Produced by Urease. ZOLTÁN ASZÓDI (*Biochem. Z.*, 1922, 128, 391—395; cf. A., 1920, ii, 519).—An improvement in Partos's apparatus is described and examples given of its use. H. K.

A Micro-urease Method for the Estimation of Urea in Blood, Secretions, and Tissues. K. L. GAD-ANDRESEN (*J. Biol. Chem.*, 1922, 51, 373—376).—The urea is first decomposed by means of urease and the ammonia then estimated by the method described for ammonia (this vol., ii, 523). A correction is made for the ammonia originally present. E. S.

The Estimation of Guanidine. A. H. DODD (*J. Soc. Chem. Ind.*, 1922, 41, 145—147r).—The presence of guanylcarbamide, which is likely when guanidine has been prepared from dicyanodiamide and dilute acids, interferes with the estimation of guanidine as picrate, as this substance also gives an insoluble picrate with picric acid or ammonium picrate. It is found, however, that guanylcarbamide picrate is much more soluble in sodium hydroxide solution than the guanidine compound, and by choosing suitable concentrations it is possible to precipitate the latter whilst the former is retained in solution. The guanidine solution, of less than 1% strength, is added to 50 c.c. of the precipitant consisting of a solution of 20 grams of picric acid in 100 c.c. of *N*/1-sodium hydroxide solution. The precipitation is carried out in hot solutions

(90°), and the mixture kept for at least six hours before filtering through a Gooch crucible. The correction which would have to be applied to the weight of picrate obtained on account of its slight solubility in water can only be determined for each individual case, but empirically it was found that under the conditions above mentioned washing the precipitate with only 50 c.c. of water left just about sufficient sodium picrate behind to compensate for the solubility of guanidine picrate. It was found incidentally that precipitation of guanylcarbamide by neutral sodium picrate was a distinct improvement on the use of picric acid. G. F. M.

The Method of Estimating Allylthiocarbimide in Mustard Flour. E. LUCE and A. DOUCET (*J. Pharm. Chim.*, 1922, 25, 458—464).—In estimating allylthiocarbimide by Dieterich's method, using ammoniacal silver nitrate, exact results may be obtained either volumetrically or gravimetrically, providing certain precautions are observed. In the case of mustard flour, the reaction is complete after six hours' contact even in the cold. Rise in temperature and extension of the time of maceration are both followed by a secondary reaction which destroys some of the essence formed at first, and hence a low figure is obtained. During the first hour of maceration, this secondary reaction is negligible whatever the temperature may be. Hence to obtain accurate results the duration of the maceration should be reduced to one hour and the essence left in contact with the ammoniacal silver nitrate in the cold for six hours or at 80—85° for one hour, using a flask with a reflux condenser. W. G.

Characteristics of Identity of a Water Distilled from the Cherry Laurel HENRI PECKER (*J. Pharm. Chim.*, 1922, [vii], 25, 424—429).—The distillate from the cherry laurel as officially prepared and containing 1 gram of hydrocyanic acid per litre should also contain a marked amount of benzaldehyde, usually above 3 grams per litre. Under such conditions, the amount of free hydrocyanic acid does not exceed 0.25 gram per litre. The distillate should give an almost immediate precipitate, in the cold, with Denner's phenylhydrazine reagent, and with a solution of ammonium molybdate in sulphuric acid should give a deep blue colour. W. G.

Potentiometric Titrations of and by means of Potassium Ferrocyanide. I. The Titration of Potassium Ferrocyanide by means of Potassium Permanganate. I. M. KOLTHOFF (*Rec. trav. chim.*, 1922, 41, 343—352; cf. Kelley and Bohn, A., 1920, ii, 134).—In the potentiometric titration of ferrocyanides by means of potassium permanganate, the former solution should contain sufficient acid to prevent the precipitation of potassium manganese ferrocyanide, $K_2MnFeCy_6$. The acid may be either hydrochloric or sulphuric, the change of potential denoting the end-point being smaller with the former, but constant potential is more rapidly attained than with sulphuric acid. The author's experiments fail to confirm the objection that an error is intro-

duced owing to the liberation of chlorine by the action of hydrochloric acid on the permanganate. The method is accurate with ferrocyanide solutions of very small concentration. It is advantageous to add a definite quantity of potassium ferrieyanide before titration, as, by this means, atmospheric oxidation of the ferrocyanide is avoided. The necessary stirring is effected by passing a current of carbon dioxide through the mixture. Details of experimental results with ferrocyanide solutions of different concentrations and in the presence of different amounts of both acids are given.

H. J. E.

The Estimation of Uric Acid in Tissue Extracts. H. STUDEL and K. SUZUKI (*Z. physiol. Chem.*, 1922, 119, 166—171).—The fresh spleen from cattle contains substances other than uric acid which reduce phosphotungstic acid to blue compounds. It is therefore pointed out that in order to obtain trustworthy results with Folin's method for the estimation of uric acid in tissues it is necessary to isolate the uric acid quantitatively previously to estimating it colorimetrically.

S. S. Z.

Methods for the Estimation of the Alkaloids. PAUL HERZIG (*Arch. Pharm.*, 1922, 259, 249—308).—A comprehensive and critical survey of all the known methods, gravimetric, volumetric, colorimetric, refractometric, and polarimetric, for the estimation of alkaloids. Direct weighing, although possessing many obvious advantages, has the great disadvantage that it is difficult without loss to isolate the alkaloid in a sufficiently pure condition. Of all the methods suggested for weighing as insoluble salts or double salts only precipitation by picric acid, phospho- or silico-tungstic acid, or picrolonic acid (dinitrophenylmethylpyrazolone) have any practical value, and of these only the latter is considered really useful for practical purposes, and even here comparative estimations made on nux vomica seeds and extracts showed that the results obtained were always 2.5—5% low. Of the volumetric methods titration with standard acid is the most generally employed, iodometric estimations are not to be recommended, but iodometric estimation with iodide-iodate of the excess of acid unneutralised by the alkaloid gives good results except in the case of feebly basic alkaloids the sulphates of which are much dissociated, for example, narcotine, pilocarpine, and purine derivatives, and is particularly useful if much colouring matter is present. Volumetric precipitation methods using Meyer's reagent, phosphomolybdic acid, potassium ferrocyanide, or picric acid are regarded as inexact except in special cases such as ferrocyanide for the estimation of strychnine in presence of brucine, and picric acid for the cinchona alkaloids. Quantitative colorimetric and optical methods all demand the isolation of the alkaloid in a colourless condition, and they are often cumbersome as well, and consequently are very little used.

G. F. M.

The Titration of the Cinchona Alkaloids and their Salts. N. SCHOORL (*Rec. trav. chim.*, 1922, 41, 228—235; *Pharm. Weekblad*, 1922, 59, 369—374).—The neutralisation curves of quinine

and hydrochloric acid have been determined, using methyl-orange, methyl-red, neutral-red and phenolphthalein as indicators, both in aqueous solution and in 50% alcohol. The addition of alcohol to the aqueous solution diminishes the dissociation constant of quinine, thus rendering the base less alkaline and increasing the hydrolysis of the salts; but as the dissociation constant of the indicators is diminished at the same time, the experimental results obtained in alcoholic solution may appear to contradict this statement. Methyl-orange and methyl-yellow are unsuitable owing to indefinite end-points. Titration of quinine as a monacid base can be carried out very accurately with the use of methyl-red. The monohydrochloride is alkaline to methyl-red and acid to neutral-red; this property may be made use of, not only for quinine, but also in the case of the other cinchona alkaloids. Cinchonine gives similar results. Experimental details of the titrations are given.

H. J. E.

[Detection of] Quinotoxine in Quinine Salts. DOMENICO GANASSINI (*Boll. Chim. Farm.*, 1922, **61**, 193—199).—The presence of quinotoxine in a salt of quinine may be detected as follows: The salt is dissolved in water, if necessary with addition of sulphuric acid, and the solution rendered alkaline by means of sodium hydroxide and extracted with ether or chloroform. After separation of the solution and evaporation of the solvent, part of the residue is carefully evaporated to dryness with a few drops of concentrated nitric acid. If quinotoxine is present, the residue thus obtained is deep yellow and is turned intense brownish-yellow by ammonia solution. The remainder of the residue from the solvent is dissolved in a little warm water acidified with acetic acid and the solution filtered if necessary. One portion is shaken with a little sodium nitrite, the nitrous acid liberated converting the quinotoxine into its nitroso- and nitroso-oximino-derivatives, which form a yellow precipitate; on quinine, nitrous acid has no action. Another portion is shaken with a drop of phenylhydrazine, the quinotoxine phenylhydrazone formed being gradually deposited as a precipitate which ultimately becomes deep orange-yellow; here also quinine remains unaffected. Bromophenylhydrazine gives a deep red coloration with quinotoxine. Solutions of quinine salts which turn yellow when sterilised probably contain quinotoxine and should not be employed for medicinal purposes. T. H. P.

Colorimetric Methods for the Separate Estimation of Tyrosine, Tryptophan, and Cystine in Proteins. OTTO FOLIN and JOSEPH M. LOONEY (*J. Biol. Chem.*, 1922, **51**, 421—434).—Certain defects in the method of Folin and Denis (*A.*, 1912, ii, 1012) for the estimation of tyrosine are remedied, and an extension is made to include the separate estimation of tryptophan and cystine. When tryptophan is to be estimated, the protein is hydrolysed by means of barium hydroxide to avoid loss due to humin formation. In the case of cystine, which is decomposed by boiling alkali hydroxides, acid-hydrolysis is used. For the estimation of tyrosine and tryptophan, the hydrolysate, which must contain between 3.5% and

7.5% of sulphuric acid, is filtered from barium sulphate, and tryptophan is precipitated from an aliquot part of the filtrate by the addition of Hopkins and Cole's mercuric sulphate reagent. After separation by centrifuging, tyrosine is estimated colorimetrically in the supernatant liquid by means of Folin and Denis's phenol reagent in the presence of sodium carbonate and sodium cyanide. The estimation of tryptophan is made similarly in the solution obtained by dissolving the mercuric sulphate precipitate in sodium cyanide. In the case of cystine, sodium carbonate is added to the hydrolysate and the cystine is reduced by means of sodium sulphite. The colour produced on addition of the uric acid reagent is then compared with a standard. Results of the application of the method to a number of proteins are given. E. S.

A System of Blood Analysis. Suppl. III. A New Colorimetric Method for the Estimation of the Amino-acid Nitrogen in Blood. OTTO FOLIN [with HSIEN WU] (*J. Biol. Chem.*, 1922, 51, 377—391; cf. A., 1919, ii, 308; A., 1920, ii, 337).—The method described is applicable to the tungstic acid filtrate and depends on the red coloration which is produced when amino-acids react with β -naphthaquinonesulphonic acid in alkaline solution. Five or 10 c.c. of the filtrate are used. One per cent. sodium carbonate and a 0.5% solution of the sodium salt of the sulphonic acid are added under specified conditions and the mixture is left for thirty hours in the dark. The colour is then intensified by the addition of an acetic acid-sodium acetate solution, after which the surplus quinone is decolorised by means of a 4% solution of sodium thiosulphate. A glycine solution is used as a standard. In the cases of histidine and tryptophan only one nitrogen atom reacts. Proline, tryptophan, arginine, and, to a lesser extent, alanine give unsatisfactory results. Ammonia also gives the coloration with the reagent, but this may be neglected in blood analysis. The blue colour which indole gives with β -naphthaquinonesulphonic acid is not produced under the conditions of alkalinity used in the estimation. E. S.

Use of the Conventional Carbon Factor in Estimating Soil Organic Matter. J. W. READ and R. H. RIDGELL (*Soil Sci.*, 1922, 13, 1—6).—In soil analysis it is usually assumed that 58% of carbon represents 100% of organic substance, this ratio being based on the results obtained by various observers. Estimations of the carbon content of thirty-seven surface soils from different parts of the United States showed that the percentage varied from 30.20 to 56.27, the average being 49.26, which is about 9% below the accepted value. The authors consider that the trustworthiness of any carbon factor is doubtful; if one is employed, more accurate results will be obtained if it is based on 50 to 52% of carbon.

W. P. S.

General and Physical Chemistry.

Optical Properties of certain Substances important in Microchemistry. LUIGI RAITERI (*Atti R. Accad. Lincei*, 1922, [v], **31**, i, 112—116).—The following refractive indices have been measured: sodium uranylacetate, 1.5082 ($\lambda=535$), 1.5044 (589), 1.5003 (671); sodium silicofluoride, 1.3112 (677), 1.3125 (589), 1.3132 (535), 1.3146 (458); potassium silicofluoride, 1.339; potassium platinichloride, 1.8103 (677), 1.8209 (606), 1.8353 (535), 1.856 (458).
T. H. P.

Aspects of the Neon Spectrum. LAURENCE ST. C. BROUGHALL (*Phil. Mag.*, 1922, [vi], **44**, 204—214; cf. this vol., ii, 213).—An attempt to explain the spectrum of neon by the application of the principles of Bohr and by the employment of the Langmuir hypothesis of the cubic atom. After making certain assumptions with regard to the axes of revolution of the outer electrons and taking the value $2c=1.30 \times 10^{-9}$ cm. for the radius of the neon atom (W. L. Bragg, A., 1920, ii, 537), it is shown that the radius of the orbit of the two inner electrons in the normal atom is $r=6.19 \times 10^{-9}$ cm. This figure does not apply to the atom in the gaseous state. From these two values of c and r the wave-lengths of the spectral lines have been calculated. The discussion of the ionised atom is very complex, the general mathematical treatment being impossible without further experimental evidence. Ionisation may occur by an electron penetrating the nucleus of the neon atom. The result of the alteration in the charge on the nucleus is to increase the number of lines in the part of the spectrum of greater wave-length.
W. E. G.

The Structure of the Line $\lambda=6708 \text{ \AA}$. of the Isotopes of Lithium. J. C. McLENNAN and D. S. AINSLIE (*Proc. Roy. Soc.*, 1922, [A], **101**, 342—348).—When strong arcs are maintained in the vapour of metallic lithium the wave-length $\lambda=6708 \text{ \AA}$. consisted of a close quartet with separations of 0.128 \AA ., 0.173 \AA ., and 0.165 \AA .. It is possible that these lines are two doublets of Li^6 and Li^7 . The doublet of Li^7 should be about sixteen times as intense as that of the Li^6 form. If a and c of the four components is ascribed to Li^6 and the components b and d to Li^7 , then the relative intensities would fit the relative amounts of isotopes present. On this basis, the doublet separation of Li^6 would be 0.301 \AA . and for Li^7 0.338 \AA ., instead of 0.55 \AA . calculated on the basis of the law known to hold for the other alkali elements. The isotope separation on this view would be 0.128 \AA . to 0.165 \AA .. If a and b and c and d are the two doublets, the average isotope separation would be 0.32 \AA .. Line $\lambda=6103.77 \text{ \AA}$. consisted of at least three well-defined components and $\lambda=4602.37 \text{ \AA}$., although difficult to resolve, had as a minimum two components.
W. E. G.

The Absorption Spectrum of Sulphur for X-Rays.

AXEL LINDH (*Compt. rend.*, 1922, 175, 25—27).—A study of the absorption spectra of sulphur in its different combinations shows that, as in the case of chlorine (A., 1921, ii, 344), the limits of absorption K for sulphur are displaced towards the shorter wave-lengths for the higher valencies. The differences between bivalent and quadrivalent and sexavalent sulphur respectively corresponds with 7 and 11 volts. The absorption spectra of the thiosulphates offer a special interest in that they show two different limits of absorption, one corresponding with bivalent sulphur and the other with sexavalent sulphur. W. G.

The Precise Measure of the Layers of Energy of the Barium Atom and the Appearance of the L -Spectrum of Ionisation.

A. DAUVILLIER (*Compt. rend.*, 1922, 174, 1543—1546).—Using the method previously described (A., 1921, ii, 421, 475 ; this vol., ii, 463), the author has studied the high frequency spectrum of barium. The photographs show two white rays, L_1 and L_2 , the positions of which coincide with those of the limits of absorption. These two rays are immediately followed by two weak black lines of slightly greater wave-length. From these rays the energy of the layers of low frequency has been measured, and for the layer P the value found is 7.0 ± 0.3 volts, although the ionisation potential of barium vapour has been found to be 5.2 volts. The doublet $\gamma_3 \gamma_7$ has been found for barium, γ_3 being much more intense than γ_7 , and each of these rays has a satellite of greater wave-length. Further, a number of weak rays have been found which are satellites of the principal rays, and certain lines of ionisation have been noted and measured. The wave-lengths of the L -rays for barium are tabulated and their intensities given, and amongst these are certain new lines which cannot at present be attributed with certainty to barium. W. G.

A New Absorption Phenomenon Observed in the Domain of X-Rays.

M. DE BROGLIE and A. DAUVILLIER (*Compt. rend.*, 1922, 174, 1546—1548).—The existence of absorption rays in the neighbourhood of the limits of high frequency has previously been recorded (A., 1920, ii, 655 ; 1921, ii, 475). A study of the L emission spectrum of barium, without the interposition of any screen, has shown such rays to be much more apparent and without any absorption band being visible (preceding abstract). This phenomenon is attributed to the state of intense ionisation in which the barium atoms occupying the focus of the anticathode are found. W. G.

Spectral Study of the Triboluminescence of Sucrose.

HENRI LONGCHAMON (*Compt. rend.*, 1922, 174, 1633—1634).—The spectrum of triboluminescence of sucrose shows almost all the constituents of the second positive band spectrum of nitrogen and no others. The triboluminescence of sugar is thus due to a

discharge in the air between two solid particles which have just been suddenly separated and are electrically charged. A triboluminescent effect may be obtained in a tube in which sugar has been finely powdered by shaking, the tube being evacuated by sudden compression or expansion of the residual air. W. G.

The Fluorescence of Mercury Vapour Excited by Röntgen Rays. J. STEPH. VAN DER LINGEN (*Z. Physik*, 1922, **10**, 38—40).—The fluorescence spectrum of mercury vapour at low densities, obtained with soft Röntgen rays, consists of two structureless bands and five emission lines. The bands reach from 4350 to 5500 and from 3100 to 3600 Å., and have their maxima at 4350 and at 3300 Å. The emission lines are 3140 Å., 3660 Å., 4050 Å., 4350 Å., and 5500 Å. The spectrum excited by Röntgen rays is in no way similar to that excited by ultra-violet light. between 1860 and 2550 Å., with the possible exception of bands in the visible spectrum. In no case was any trace found of the lines 2536 and 2346 Å. discovered by Steubing (*Physikal. Z.*, 1909, **10**, 797).

W. E. G.

The Characteristic Röntgen Radiation from Carbon and Boron. J. HOLTSMARK (*Physikal. Z.*, 1922, **23**, 252—255).—Measurements are made of the *K* radiation for boron and carbon, and these are compared with the previous values of Richardson and Bazzoni (this vol., ii, 14), Hughes (this vol., ii, 184), and Kurth (*Physical Rev.*, 1921, **18**, 461). The wave-lengths of the *K* radiation of carbon and boron are $\lambda=42.9$ and $\lambda=83.6$, respectively. The boron point is not in line with the *K_a* emission points for other elements. Theoretical reasons are advanced to account for this deviation.

W. E. G.

The Chemical Action of Penetrating Radium Rays.
XIV. The Action on Oxalic Acid, Potassium Tetroxalate, and Potassium Chlorate. ANTON KAILAN (*Monatsh.*, 1922, **43**, 1—12).—When a 0.01 *N*-solution of oxalic acid was exposed for 2136 hours to the action of the rays from about 100 mg. of radium, a considerable loss of acidity occurred and a smaller loss of reducing power when titrated with permanganate. The radiated solution contained also formic acid and an aldehyde. The quantitative determinations showed that the greater part of the oxalic acid which disappears decomposes according to the equation $C_2H_2O_4 = CO_2 + H \cdot CO_2H$. Some hydrogen peroxide is formed according to the equation $C_2H_2O_4 = 2CO + H_2O_2$, and this causes the apparent loss of oxalic acid, as shown by the permanganate titration, to be smaller than the loss estimated by acidity. From previous work on the equilibrium of hydrogen peroxide under the influence of radium rays (this vol., ii, 466) it is concluded that the decomposition of the oxalic acid is not conditioned by hydrogen peroxide formation. In presence of uranium nitrate, oxalic acid disappeared at the same rate under the influence of radium rays as in its absence,

but no formic acid was found in the radiated solution. Experiments showed that the decomposition of formic acid by the rays was not accelerated by uranium nitrate, and it must be concluded that no formic acid is produced from the oxalic acid when the uranium salt is present. On the other hand, more hydrogen peroxide is formed, as shown by the permanganate titre of the radiated solution. The ratio of number of molecules affected by the rays, m , to number of ion pairs, n , is in each case 0.3, whether calculated on oxalic acid or hydrogen peroxide.

Under the influence of rays from a quartz mercury lamp, in absence of uranium salt decomposition of the oxalic acid occurred without formation of formic acid, whilst in presence of uranium salt much formic acid appeared and the permanganate titre fell more rapidly than the acidity.

When potassium tetroxalate solution was exposed to radium rays, less formic acid and more hydrogen peroxide was produced than when oxalic acid was used. The lower hydrogen-ion concentration might have led to more rapid decomposition of the formic acid. The rays from the quartz mercury lamp resulted in hydrogen peroxide formation but no formic acid; in presence, however, of uranium nitrate 1 mol. of formic acid was found for each mol. of oxalic acid disappearing.

Potassium chlorate in 0.1*N*- to 0.01*N*-solution undergoes very slight reduction to chloride by prolonged action of radium rays at 8–14°. After sixty-four hours' exposure at 45–50° to the rays of a quartz mercury lamp there was only slight reduction, about equal to that produced by 1000 hours' exposure to radium rays.

E. H. R.

The Binding of Electrons by Atoms. J. W. NICHOLSON (*Phil. Mag.*, 1922, [vi], 44, 193–203).—A mathematical paper in which it is shown that the hyperbolic orbits of Epstein which have been used extensively in the interpretation of certain groups of X-rays rest on a mathematical error. A determinate and finite value of W cannot be obtained for an electron moving about any atomic nucleus, if the path involved takes the electron to infinity.

W. E. G.

New Differential Method for the Measurement of the Conductivity of Electrolytes. L. ROLLA and L. MAZZA (*Gazzetta*, 1922, 52, i, 421–428).—The apparatus used in this method consists of an induction coil, I , a resistance box, R , a telephone, T , and three coils, P_1 , P_2 , and S , wound on an iron core. P_1 and P_2 have equal resistances and coefficients of self-induction, and the three coils together form a differential transformer with two primary circuits and one secondary.

The electrolytic resistance, X , to be measured is connected, in series with the induction circuit, P_1 , to the terminals of the induction coil; the resistance box is similarly connected, in series with P_2 , to the same terminals so that the current at any moment flows

through P_1 and P_2 in opposite directions. S is connected directly to the telephone.

Consideration of the two circuits XP_1I and RP_2I shows that P_1 and P_2 produce at any instant two opposite magnetic fluxes, F_1 and F_2 , the resultant flux being $F_1 - F_2$. When $F_1 - F_2 = 0$, the *E.M.F.* induced in S is zero and the telephone is silent; this condition is evidently satisfied when X and R are equal. The arrangement of P_1 , P_2 , and S on a cylindrical iron core gives poor results, but if the core consists of two L-shaped pieces juxtaposed to form a rectangle and P_1 and P_2 be wound together on one long side and S , composed of three separate coils capable of being grouped variously, be wound on the opposite long side of the core, excellent results are obtained. The sensitiveness of the method may be greatly increased by the use of the audion. T. H. P.

The Electromotive Properties of Magnesium and the Potentiometric and Thermal Analysis of the System Magnesium—Mercury. R. PH. BECK (*Rec. trav. chim.*, 1922, 41, 353—399).—The normal potential of magnesium has not been ascertained with certainty, and as some magnesium amalgams in contact with solutions appear to yield as great a potential difference as the metal itself, the author has extended the work of Cambi and Speroni (A., 1915, ii, 453) on the thermal analysis of magnesium amalgams in an endeavour to complete the fusion diagram, to investigate the potential differences due to amalgam as compared with those due to the pure metal, and to ascertain whether the values obtained for the potential are in accord with the dilution law. The figures given by Cambi and Speroni (*loc. cit.*) for amalgams containing from 70 to 84 atomic % of mercury are confirmed; no measurements were made for higher percentages. The following compounds were found in the system: $MgHg_2$, transition temperature 170° , at which it is coexistent with $MgHg$; $MgHg$, m. p. 625° ; Mg_3Hg_2 , m. p. 562° ; Mg_2Hg , m. p. 580° ; and the thermal diagram is given and discussed in detail. The existence of $Mg_{10}Hg$, stated by Kremann and Muller, is disproved; it is suggested that their mixtures were not sufficiently heated to attain equilibrium. The vapour pressures of two amalgams were measured, but little value is claimed for the results owing to experimental difficulties; similar measurements made by Kremann and Muller are criticised for the same reason.

Determinations of the potential differences between amalgams varying from 0—60 atomic % of mercury in a solution of anhydrous magnesium chloride of 0.1982*N*-concentration in ethyl alcohol against a calomel electrode were made. The results obtained show considerable fluctuations with change of composition of the amalgam, but no conclusion is drawn that these correspond with new phases. It seems, however, that no mixture or compound of the two constituents is more negative than magnesium itself. The author points out that the usual method of preparing amalgams for such

experiments does not exclude the possibility of working with a metastable substance and that measurements made against aqueous solutions may give too low a value but never too high. Further, if the potential difference undergoes the normal change on dilution, this would tend to confirm the accuracy of the figure obtained. But in the case of magnesium, the potential becomes less positive with increasing dilution. In order to compare the potential of magnesium and of magnesium amalgams in contact with neutral acid and alkaline solutions, several series of experiments were carried out in which the measurement was made in open vessels, using a normal calomel electrode. The conclusions drawn are that the potential of magnesium may be measured in acid solution and that the figures obtained differ but little from those for magnesium amalgams, that in neutral solution atmospheric oxygen may cause the metal to be somewhat passive, and that in alkaline solution there is a considerable tendency towards passivity. In the case of magnesium in contact with solutions of magnesium sulphate of various concentrations, the results appear to contradict the hypothesis put forward by Kistiakowsky (A., 1910, ii, 258), that magnesium passing into solution from an electrode is primarily transformed into the hydroxide. In acid solutions of different concentrations, the potential becomes more negative with increasing dilution; the figure finally reached is $E_H = -1.854$. Experiments on cathode polarisation of magnesium showed that measurements of potential made after interruption of the polarisation are never more negative than the highest values obtained for the metal and from its amalgam with no current passing. Thus this last figure should be considered as very close to the equilibrium potential which the author suggests is most probably $E_H = 1.856-1.876$ volts.

H. J. E.

Absorption of Hydrogen by Elements in the Electric Discharge Tube. F. H. NEWMAN (*Phil. Mag.*, 1922, [vi], 44, 215—226).—A further study of the behaviour of hydrogen in the presence of various elements in a discharge tube (cf. A., 1921, ii, 295). Evidence is obtained that the "clean up" of hydrogen in a discharge tube is partly due to chemical changes. On the admission of hydrogen, activated by an electric discharge, to the clean surface of a sodium-potassium alloy a white, crystalline compound was produced which was probably composed of the hydrides of the two metals. The white layer slowly changed to a dark grey deposit. Similarly, the activated gas gave hydrogen sulphide with sulphur. These effects were not due to the ions of the discharge tube, as these were eliminated by charged platinum strips before the gas reached either the alloy or the sulphur. Measurements were made of the rates of disappearance of hydrogen in the presence of sodium-potassium alloy, and of sulphur and other elements. A considerable proportion of this absorbed hydrogen was recovered when the tube was heated at 300°. Nitrogen, on the other hand, cannot be again liberated, even when the tube is heated to the softening point. The

modifications of hydrogen and nitrogen which are absorbed in the discharge tube are probably triatomic.

The absorption is not entirely due to chemical action, for the law of constant proportions does not appear to be followed.

W. E. G.

Rapid Electrolysis without Rotating Electrodes. GRAHAM EDGAR and R. B. PURDUM (*J. Amer. Chem. Soc.*, 1922, **44**, 1267—1270).—Rapid electrolysis may be carried out without the use of rotating electrodes if the solution is stirred by means of a current of air. A piece of apparatus is described for such analyses. The cathode is a platinum gauze cylinder in the centre of which is a spiral platinum electrode. The electrodes are contained in a glass tube 31 mm. diam. to which are sealed three glass "air lifts," connected in a single tube at the bottom and entering symmetrically about half-way up the wide tube, the seals being made tangentially. When air is forced into the apparatus half filled with liquid, both vertical and rotational stirring is effected. Experiments are described which show that the results obtained with this apparatus are equally as good as those obtained with rotating electrodes.

J. F. S.

An Electrochemical Study of the Reversible Reduction of Organic Compounds. J. B. CONANT, H. M. KAHN, L. F. FIESER, and S. S. KURTZ, jun. (*J. Amer. Chem. Soc.*, 1922, **44**, 1382—1396).—The authors have developed the general equation $\pi = \pi'_0 + 0.0295 \log \{ [A]/[AH_2]_r - 0.0295 \log K_1 K_2 + 0.059 \log [H^+] + 0.0295 \log \{ K_1(K_2 + [H^+]/[H^+]^2) + 1 \}$ (in which A represents the oxidised and AH_2 the reduced form) to express the oxidation-reduction potentials of compounds of the type of quinone in both acid and alkaline solutions. The potentials of six anthraquinonesulphonic acids have been measured by a titration method, using titanous chloride and sodium hyposulphite, and the results obtained over a wide range of hydrogen-ion concentration have been found to be in accord with the above equation. The validity of the method is shown by determinations of the potentials of mixtures of the reduced and oxidised compounds.

W. G.

Thermo-regulator. E. B. STARKEY and N. E. GORDON (*J. Ind. Eng. Chem.*, 1922, **14**, 541).—A thermo-regulator is described which is capable of giving a steady temperature to 0.05°. The regulator consists of a U-tube, connected through a short vertical capillary with a toluene container. The U-tube and capillary are filled with mercury, the two ends of which are connected with wires which form one of the leads of the heating lamp. Expansion of the toluene causes the mercury to be pushed away from one wire and so break the circuit, whilst contraction remakes the circuit. Corrosion of the mercury is prevented by filling the space above the mercury with an inert gas.

J. F. S.

Entropy of the Elements and the Periodic System. W. HERZ (*Z. Elektrochem.*, 1922, 28, 258—259).—A theoretical paper in which the author shows that the entropy of the elements varies regularly in keeping with the arrangement of the elements in the periodic system. Using the entropy values calculated by Lewis and Gibson (A., 1918, ii, 29), it is shown that in any group of the periodic system the value of the entropy increases with increasing atomic weight in the case of the metallic elements, the elements of the fourth group and the inactive gases, whilst the electro-negative elements of the oxygen and halogen groups have decreasing entropies with increasing atomic weights. J. F. S.

Proposal for the Fixing of a Unit of Measurement in Thermochemistry. W. SWIENTOSLAWSKI (*Roczniki Chemji*, 1921, 1, 479—487; cf. A., 1921, ii, 679).—The following proposals were put forward at the International Chemical Conference at Brussels in 1921. The heat of combustion of 1 gram of benzoic acid, in cal. at 15°, should be taken as the unit, rather than that of cane-sugar, which is hygroscopic, or of naphthalene, which is liable to sublime. By using one substance only for the standard, many constant calorimetric corrections would not be needed, and a simpler formula could be used. Certain discrepancies in the values for the heat of combustion of benzoic acid, as obtained by various workers, are pointed out, and also in the ratios for the heats of combustion of benzoic acid, cane-sugar, and naphthalene. It is proposed that the heat of combustion of benzoic acid at constant volume be taken provisionally as 6321 cal., with the proviso that the determination of this value be repeated as soon as possible, with the greatest possible accuracy, using the method of adiabatic measurements.

R. T.

Temperatures of Combustion. J. BRONN (*Z. angew. Chem.*, 1922, 35, 328).—The temperature of combustion of a gas is calculated from the heat of combustion and the specific heat of the products of combustion. An accepted figure for the combustion of hydrogen in pure oxygen is 6670°, which would presume the specific heat of steam to be 0.485. It is shown that the specific heat at 4000° may be taken at 0.85, from which the temperature of combustion may be calculated to be 3900°. The temperatures for carbon monoxide (5000°), methane (4400°), and acetylene (6200°) are given. The temperatures of combustion of gases are greatly influenced by their velocities of flame propagation, except in so far as flameless combustion affects the result. The velocities of flame propagation of ethylene, acetylene, and benzene differ. H. M.

Thermochemical Researches on Oximes. I. Ketoximes not Exhibiting Stereoisomerism. ALICJA DORABIALSKA (*Roczniki Chemji*, 1921, 1, 424—447).—The heats of certain reactions of some ketoximes are measured. From the results so obtained, and from those obtained previously (W. Swientoslawski, A., 1919, 2, 336), various generalisations are made. The ketoximes used were

acetoxime, phenylmethylketoxime, and diphenylketoxime. The chief reactions studied are the formation of the hydrochloride of the oxime, its solution in 10% sodium hydroxide solution, and the formation of the sodium salt of the oxime. The results obtained, together with those calculated from these, are given in the following table, where q_{HCl} is the heat of formation of the hydrochloride, in ethereal solution. For aqueous solution this is expressed by q'_{HCl} . The heat of the solution of the hydrochloride in 10% sodium hydroxide solution is Q_{HCl} , in water, S_{HCl} . The heat of solution of the oxime in dilute sodium hydroxide solution is q , in water, S , in ether, S' . The degree of hydrolysis of sodium salts of these oximes is given as $\nu\%$. Results are expressed in calories per millimole.

	q_{HCl}	q'_{HCl}	q	$\nu\%$	Q_{HCl}	S	S_{HCl}	S'
Acetoxime .	10.93	1.19	5.70	39.6	17.16	-1.28	-1.95	-2.89
Phenylmethyl- ketoxime .	8.17	0.51	6.34	40.3	16.52	-2.98	-3.52	—
Diphenyl- ketoxime .	7.71	—	5.80	—	12.51	-4.06	-6.99	—

In the case of phenylmethylketoxime, the hydrochloride, prepared at different temperatures, and from different media, gave the same Q_{HCl} , thus showing that no stereoisomeric modification of this oxime is produced under the conditions in question.

It is concluded that the heat of formation of the hydrochlorides of the ketoximes differs little from that of the aldoximes, and diminishes as the acidity of the ketoxime increases. The heat of solution of these hydrochlorides diminishes with increasing acidity of the ketoxime, owing to a parallel increase in the negative heat of the solution, and decomposition of the former in water. Finally, the negative heat of solution of ketoximes in water appears to be greater than that of aldoximes.

R. T.

Pycnometry. R. SAAR (*Chem. Ztg.*, 1922, **46**, 433—435).—Tables and formulæ are given for calculating specific gravities from one temperature to another.

W. P. S.

An Experimental Comparison of the Viscous Properties of (a) Carbon Dioxide and Nitrous Oxide; (b) Nitrogen and Carbon Monoxide. C. J. SMITH (*Proc. Physical Soc.*, 1922, **34**, 155—164).—The viscous properties of these gases have been compared directly by observing the time required by a mercury pellet to force a certain volume of the different gases through a capillary tube. Measurements were carried out at different temperatures, absolute viscosities were obtained by comparison with air, and the mean area of collision deduced by using Chapman's formula.

The viscosities expressed in C.G.S. units $\times 10^{-4}$ are: carbon dioxide and nitrous oxide, 0.0° , 1.366; 15.0° , 1.441; 100.0° , 1.845. Carbon monoxide and nitrogen, 0° , 1.665; 15.0° , 1.737; 100.0° , 2.118. The mean collision areas in square centimetres are: carbon dioxide and nitrous oxide, 0.834×10^{-15} . Carbon monoxide and nitrogen, 0.767×10^{-15} .

The fact that carbon dioxide and nitrous oxide (or carbon monoxide and nitrogen) have molecular dimensions which are identical within experimental error is consistent with Langmuir's view that the outer electron arrangements of these molecules are the same; this is also supported by the fact that the viscosities are equal at the three temperatures chosen.

W. T.

Theory of Adsorption Processes. A. EUCKEN (*Z. Elektrochem.*, 1922, **28**, 257—258).—Polemical; an answer to Polanyi's criticism (this vol., ii, 479) of the author's paper on the theory of adsorption processes (this vol., ii, 262).

J. F. S.

Adsorption in Solution and at Interfaces of Sugars, Dextrin, Starch, Gum Arabic, and Egg-albumin, and the Mechanism of their Action as Emulsifying Agents. GEORGE L. CLARK and WILLIAM A. MANN (*J. Biol. Chem.*, 1922, **52**, 157—182).—Measurements were made of the surface tension and viscosity of solutions of the above substances for a large number of concentrations both with and without the addition of electrolytes. The interfacial tension between each solution and benzene was also measured, and the efficiency of the emulsifying agent with respect to both benzene and kerosene estimated. The factors of predominating importance in an emulsifying agent appear to be viscosity and ability to lower interfacial tension. In the case of sugar, viscosity is the more important, whilst with egg-albumin, which gave the best emulsions, film formation is of primary importance. Two types of agglomeration are distinguished, namely, that accompanied by an increase in viscosity, in which coalescence of the particles is accompanied by enclosure of water, and that producing a decrease in viscosity, in which there is a decrease both in surface and amount of bound water. Dextrin and starch, considered separately, provide examples of the former type. When these substances are considered in relation to one another, however, it is found that the viscosity curves run parallel, the curve for dextrin being slightly above that for starch, thus resembling the viscosity curves of sulphur sols of different degrees of dispersion (cf. Odén, A., 1912, ii, 1143). This is considered to be evidence that starch and dextrin differ merely in the degree of agglomeration (second type) of the particles, and thus to furnish experimental proof of the view put forward by Herzfeld and Klinger (A., 1920, i, 713).

E. S.

Carrying Down by Precipitates. PAUL DUTOIT and ED. GROBET (*J. Chim. physique*, 1922, **19**, 328—330).—The carrying down of soluble salts by precipitates is explained by the authors as due to the fact that the adsorption of these substances on the surface of the precipitate causes the soluble substance to attain locally a value which exceeds its solubility product and hence causes its precipitation. With the object of testing this hypothesis, the author has estimated calcium in the presence of magnesium

salts as oxalate, and barium in the presence of calcium both as sulphate and chromate. Two series of experiments were carried out, in the first stirring by hand with a glass rod was employed, and in the second series very rapid mechanical stirring was used. Very different results were obtained in the two series, but the values of the second series are in good agreement with the theoretical values, thus showing that soluble substances are not appreciably carried down by precipitates if the solution is rapidly stirred during the mixing.

J. F. S.

The Sorption of Neutral Soap by Wool, and its Bearing on Scouring and Milling Processes. BRITISH RESEARCH ASSOCIATION FOR THE WOOLLEN AND WORSTED INDUSTRIES (*Trans. Text. Inst.*, 1922, **13**, 127—142).—The sorption of soap from aqueous solution by carefully cleansed wool at 25° has been studied. Analyses of the solution were made after an arbitrarily fixed interval, without waiting for equilibrium to be established, and corrections were applied for the amount of water sorbed by the wool on the assumption that the material contains 15% of moisture under ordinary atmospheric conditions and 33% when immersed. The results indicate preferential sorption of the kation, and it is suggested that the fatty acid is removed from solution both by sorption and by precipitation on the fibre. With mixtures of soaps of oleic and palmitic acids, preferential sorption of oleic acid occurs, but the potassium and sodium oleates are almost identical in behaviour. The effect of the fatty acid is apparently to lower the sorption of alkali.

J. C. W.

Mordants. I. WILDER D. BANCROFT (*J. Physical Chem.* 1922, **26**, 447—470).—A historical review of the use of mordants in dyeing.

J. S. G. T.

The Dissociation of Barium Platinichloride. G. GIRE (*Compt. rend.*, 1922, **174**, 1700—1703).—The author has measured the dissociation pressures of barium platinichloride over the temperature range 428° to 665°, and has obtained the equation

$$\log P = -3446.316/T + 7.702 \log T - 17.41669.$$

The dissociation of barium platinichloride is an equilibrium of the type, $\text{sol.} + \text{gas} \rightleftharpoons \text{sol.} + Q$, and the value of Q is calculated to be 30.35 cal. for the temperature 948°, which corresponds with a pressure of 760 mm.

W. G.

A Rapid Dialyser. A. GUTBIER, J. HUBER, and W. SCHIEBER (*Ber.*, 1922, **55**, [B], 1518—1523).—The apparatus consists of a wooden disk supporting a framework composed of glass rods in the form of a hollow cylinder. The membrane is passed over the framework and secured in its natural folds to the wooden disk. A stirrer is placed within the vessel, which is mounted in such a manner that the frame and stirrer can be rotated in opposite directions (the apparatus is fully figured in the original). The

apparatus is placed in an inverted, tubulated bell-jar, through the tubulus of which water is admitted, which can be heated or cooled by coils placed in the jar; the level of the water and its discharge are governed by a lever arrangement. Comparative experiments show the dialyser to be much more rapid in its action than those of Graham and Zsigmondy-Heyer. In addition to rapidity of action, the following advantages are claimed for the apparatus: extensive exclusion of external air from the internal liquid, from which, however, samples may readily be taken; small liability to changes in volume, particularly to dilution of the internal liquid; great safety in action, since only a single piece of parchment is used; simple method of securing a continuous change of external water; the possibility of dialysis at raised or lowered temperatures without complicated arrangements. H. W.

Dolezalek's Theory of Solutions. K. HERRMANN (*Z. angew. Chem.*, 1922, **35**, 349—351, 353—355).—A theoretical paper in which liquid mixtures and solutions are differentiated and discussed generally and in connexion with the Dolezalek theory of solutions. J. F. S.

Measurement of Solubility by Floating Equilibrium. Solubility of Lead Acetate. MERLE L. DUNDON and W. E. HENDERSON (*J. Amer. Chem. Soc.*, 1922, **44**, 1196—1203).—A new method for the determination of solubility is described. The method consists in placing a float, which has been calibrated so that it will just sink in a solution of known composition of the substance under investigation, into a weighed quantity of the saturated solution and adding the solvent until the float just sinks. The quantity of solvent added is noted, and from this, the weight of the solution taken, and the calibration constant of the float the solubility is readily calculated. This method has been applied to the determination of the solubility of lead acetate in water at temperatures from 0° to 50°. The following values in grams of lead acetate per 100 grams of water are recorded: 0°, 19.7; 5°, 23.7; 10°, 29.3; 15°, 35.6; 20°, 44.3; 25°, 55.2; 30°, 69.7; 35°, 88.9; 40°, 116.0; 45°, 153.0; 50°, 221.0. The method is capable of considerable speed or great accuracy, and in ordinary circumstances of a considerable amount of both. Some variation is found in duplicate experiments at the higher temperatures, which is shown to be due to the hydrolysis of the acetate followed by subsequent evaporation of acetic acid. J. F. S.

Influence of Electrolytes on the Solubility of Non-electrolytes. ANDREW MCKEOWN (*J. Amer. Chem. Soc.*, 1922, **44**, 1203—1209).—The heats of solution of ether in various solutions of sodium chloride have been calculated from Thorne's solubility data at 15° and 25° (T., 1921, **119**, 262), by means of an expression which connects solubility, heat of solution, and temperature, of the form of the van't Hoff isochore. Statistical

treatment of the phenomena of solution and precipitation leads to an expression for solubility, $S = Ae^{Q/RT}$, which is in agreement with the isochore relation mentioned above. The salting-out effect of salt on ether has been examined so far as the effect depends on the activities of the ions of the salt. It is shown that the relative specific influences of the kation and anion are -21.7 and $+671$, that is, the anion is the predominating factor in the process.

J. F. S.

Amorphous Precipitates and Crystalline Sols. F. HABER (*Ber.*, 1922, 55, [B], 1717—1733).—The method of distinguishing between the crystalline and amorphous condition depends on the ability of crystalline substances to give Röntgen-ray interference fringes. The trustworthiness of the method is dependent on the conditions (i) that the lattice constants are not less than the half wave-length of the Röntgen rays employed, (ii) that they are not much greater, and (iii) that the material under investigation is uniform in its lattice properties.

Crystalline or amorphous, solid masses are obtained by super-saturation processes which may be considered to occur in two distinct phases, the formation of aggregates and the rearrangement of such aggregates, with loss of free energy, into ordered lattice formations. The former process can be studied conveniently in supercooled, molten masses of a uniform chemical material which may be regarded as unarranged molecular aggregates the further fate of which depends on their rate of arrangement. With regard to the process of arrangement, its energy is known (latent heat of liquefaction), and also that it is divisible into two parts, the formation of nuclei and their growth. Tammann's work on this subject has shown that the formation of nuclei frequently only occurs with considerable velocity at temperatures far below the melting point at which the nuclei can only grow very slowly; if, however, the mass is heated to just below its melting point, the nuclei grow rapidly and crystallisation occurs throughout the entire mass. Thermodynamical treatment of the subject shows that the minute aggregates are only stable when the condition is fulfilled that $AT/T = 2S/rqd$, in which AT/T is the fraction of the mass melting point T in absolute values by which cooling must occur if the solid phase, regarded as spheres of radius r , is to be in (labile) equilibrium with the molten mass. S is the specific free energy of the boundary surface, q the latent heat, and d the density of the solid phase.

The processes involved in the formation of precipitates and sols differ from those described above in that the rate of aggregation can be controlled by exceeding the limit of solubility to a greater or less extent. The connexion between the mass solubility H_M (solubility as generally accepted) and the solubility of the smallest aggregates, l , is given by the equation $2S'M/rd = RT \log l/L_M$, in which S' is the specific free surface energy of the particles, T the temperature of the experiment, d the density of the solid phase,

M the molecular weight, and r the radius of the particles regarded as spheres.

The considerations outlined above are applied to the formation of very sparingly soluble substances. If the rate of aggregation is high and as far as possible in excess of that of arrangement, amorphous precipitates are to be expected which gradually, particularly on warming, pass into the crystalline condition. If, however, the rate of aggregation is depressed by only slightly exceeding the solubility limit, the rate of arrangement may be sufficient to cause the orderly formation of crystals before the formation of visible particles has occurred. This, however, involves an alteration in the rate of aggregation due to electrical phenomena at the boundary of the molecules and liquid, the net result of which is that the growth of the aggregates is greatly impeded and sols are produced. These conclusions are illustrated by the cases of the precipitates and sols of aluminium or ferric hydroxides which, however, are to be regarded as the ideal case from which deviations are to be expected in two directions. If the rate of arrangement is greater, it is not to be expected that the precipitate will be obtained in the amorphous condition and be retained as such during the Röntgen exposure. The rate of arrangement may be expected to be at its maximum in binary, heteropolar compounds in which the bipolar character is most marked. (This is illustrated in the cases of the silver haloids and the sulphides of mercury, zinc, and cadmium). On the other hand, with molecules in which the bipolar character is not strongly accentuated, the tendency towards the formation of amorphous sols and precipitates must be more marked; this is illustrated experimentally by the cases of the hydroxides of zirconium and thorium.

H. W.

Molecular Arrangement and Liquid Crystal Formation.

D. VORLÄNDER (*Z. angew. Chem.*, 1922, **35**, 249—250).—A close relationship is shown to exist between the symmetry of the molecule and its capacity to form liquid crystals. This property is shown to the greatest extent by those molecules containing long chains of atoms and possessing a definite axis of symmetry. Thus para-substitution in the benzene ring is especially favourable to the formation of liquid crystals. The introduction of groups which destroy the symmetry of the molecule either reduces the range of stability or else completely destroys the power of formation of liquid crystals.

Benzidine derivatives $(\text{CHR}:\text{N} \langle \text{benzene ring} \rangle \text{N}:\text{CHR})$ give liquid crystals with a great range of stability, but the introduction of a CH_2 group between the two benzene nuclei causes the disappearance of this enantiotropic form. On account of the angle between the carbon valencies ($109^\circ 28'$) the molecule will no longer be symmetrical. Similar results are obtained by the introduction

of CO, CS, S, and O groups into the same position. The NH group reduces the range of stability practically to zero, although monotropy is sometimes observed in these compounds. Further lengthening of the chain with the formation of diphenyl ethane derivatives causes a reappearance of the property, although it is not so strongly developed as in the diphenyl derivatives. The diphenylpropanes do not yield liquid crystals, which, however, appear again with the derivatives of diphenylbutane. The author explains these phenomena by the departure of the molecule from the straight line arrangement.

W. E. G.

Electrical Precipitation of Colloids. CLAUDE HAINES HALL, jun. (*J. Amer. Chem. Soc.*, 1922, **44**, 1246—1249).—Finely divided suspensions of arsenious sulphide, mercuric sulphide, prussian blue, barium sulphate, aluminium, copper, lead, silver, and iron, in transformer oil of resistance 1.5×10^{16} ohms per c.c., have been subjected to alternating and direct currents of voltages from 10^4 to 2×10^5 with the object of effecting a precipitation. In no case was any precipitation brought about, even although the current was applied for three hours. A theoretical treatment of the subjects shows that the rate of precipitation would be 1/20000 of that for smoke for the same application of energy. J. F. S.

A Static Method of Study of Hydration. MARCEL GUICHARD (*Bull. Soc. chim.*, 1922, [iv], **31**, 552—554).—A simple form of apparatus is described in which it is possible to follow the equilibrium between an absorbent substance and water vapour in a vacuum and over a fairly considerable range of temperature. It consists of a glass tube to contain the absorbent substance and having sealed to it at right angles a graduated tube which contains the water. The whole is evacuated and sealed off. The two parts of the apparatus can be maintained at different temperatures as desired.

W. G.

The Geometric Representation of Saline Equilibria. HENRY LE CHATELIER (*Compt. rend.*, 1922, **174**, 1501; cf. A., 1894, ii, 223).—A reaffirmation of priority in the method of representing saline equilibria by square diagrams as opposed to the claim of Jänecke (this vol., ii, 427).

W. G.

Effect of an Electrolyte on Solutions of Pure Soap. Phase-rule Equilibria in the System Sodium Laurate-Sodium Chloride-Water. JAMES WILLIAM MCBAIN and ARTHUR JOHN BURNETT (*T.*, 1922, **121**, 1320—1333).

The System Silver Perchlorate-Water-Benzene. ARTHUR E. HILL (*J. Amer. Chem. Soc.*, 1922, **44**, 1163—1193; cf. A., 1921, ii, 261).—The ternary system silver perchlorate-water-benzene and the binary systems made up of pairs of the components have been investigated. It is shown that the system silver perchlorate-water has a eutectic point at -58.2° , at which temperature the saturated solution contains 73.9 % of the silver salt. This is

the lowest eutectic known for a true salt and water. The solubility of silver perchlorate rises to 88.8 % at 99°. The eutectic of silver perchlorate-benzene is at +5.12° and the solution contains 3.44 % of the silver salt. The solubility curve rises steeply with the temperature, reaching an almost perpendicular form between 50° and 145°; at the latter temperature the saturated solution contains 63.0 % of silver perchlorate. Attention is directed to the similarity between this solubility curve and those which are found where the system gives rise to two liquid phases, either in the stable or metastable region. The hydrate $\text{AgClO}_4 \cdot \text{H}_2\text{O}$ has been isolated and its transition temperature found to be 43.1°; the compound $\text{AgClO}_4 \cdot \text{C}_6\text{H}_6$ has also been isolated and its transition point found to be 145°. The ternary system, silver perchlorate-water-benzene is marked by the possession of six realisable invariant equilibria, and a very large number of univariant equilibria, which furnish at least one example of every type of equilibrium theoretically possible for a ternary system containing volatile components. The ternary eutectic representing three solid phases, solution, and vapour, lies at -58.4°, and the eutectic of highest temperature is at 42.1°. The course of the twenty-four univariant equilibria originating at these two quintuple points and the four intermediate quintuple points has been investigated. Isothermal diagrams have been constructed to show the various equilibria existing at temperatures between -58.4° and +145°. Between 5.24° and 23.4° three coexistent liquid phases can be prepared in stable equilibrium with each other and their vapour. The occurrence of two separate fields of unsaturated solution is explained as due to the intersection of the solubility curve of silver perchlorate with the bi-nodal curve representing the limited solubility of benzene and water. The occurrence of a third field is explained as due to the existence, between -2.7° and +30°, of an additional closed bi-nodal curve which does not extend to any one of the two component axes at any temperature. It appears that this is the first demonstrated case of a closed bi-nodal curve. J. F. S.

Use of Iron Pyrites in a Friedel-Crafts' Reaction. JOHN ARMSTRONG SMYTHE (T., 1922, 121, 1270--1279).

Combination in Detonating Gas in Presence of Colloidal Palladium Solution. C. SANDONNINI and A. QUAGLIA (*Gazzetta*, 1922, 52, i, 409-416; cf. this vol., ii, 557).—In mixtures containing only hydrogen and oxygen, the reaction in presence of colloidal palladium solution is proportional to the concentration of the detonating mixture. If, however, this concentration is the same in two cases, the velocity of reaction is somewhat the greater when the hydrogen is in excess than when the oxygen is in excess. Since it is also found that the reaction of combination in pure detonating gas may be accompanied by phenomena of reduction or hydrogenation, the conclusion is drawn that in this case the combination is due to the action of active hydrogen.

The hydrogenation of ethylene and the formation of water have velocities of the same order when the ratio between the amounts of hydrogen and oxygen is that normal to detonating gas. When ethylene and oxygen are in concentrations requiring the same quantity of hydrogen and the latter is present in sufficient amount for both reactions, the latter proceed with almost equal velocities; when, however, the amount of hydrogen is insufficient for either reaction, only the formation of water occurs to an appreciable extent.

T. H. P.

Limits for the Propagation of Flame in Vapour-Air Mixtures. I. Mixtures of Air and One Vapour at the Ordinary Temperature and Pressure. ALBERT GREVILLE WHITE (T., 1922, **121**, 1244—1270).

Limit of Inflammability of the Vapours of the System, Alcohol-Petrol, and of a Triple System with a Basis of Alcohol and Petrol. ROGER G. BOUSSU (*Compt. rend.*, 1922, **175**, 30—32).—A study of the variation of the lower limit of inflammability of the binary system petrol-alcohol and of the ternary system petrol-alcohol-ether, using the method of Le Chatelier and Boudouard (A., 1898, ii, 574). The results verify the formula $n/N + n'/N' = 1$, where N and N' are the limits of inflammability of each of the two vapours and n and n' the proportions in which they are present in the mixture under examination.

W. G.

A Study of the Rate of Saponification of Oils and Fats by Aqueous Alkali under Various Conditions. MABEL HARRIET NORRIS and JAMES WILLIAM MCBAIN (T., 1922, **121**, 1362—1375).

Behaviour of certain Metals as Catalysts. I. C. SANDONNINI (*Gazzetta*, 1922, **52**, i, 394—408).—The author has carried out a number of series of experiments with the object of ascertaining the behaviour of a catalyst, highly active in the case of two separate, analogous reactions, when it is introduced into a system in which both these reactions can occur. The catalysts employed were platinum black, reduced nickel, copper, and silver, and the systems were composed of hydrogen and oxygen, mixed with either ethylene, or phenanthrene, or nitrobenzene, or allyl alcohol, these being compounds able to undergo ready oxidation or hydrogenation.

In the system, $C_2H_4 : H_2 : \frac{1}{2}O_2$, the possible reactions are: (1) formation of water from the hydrogen and oxygen, (2) hydrogenation of the ethylene, (3) oxidation of the ethylene, and (4) decomposition of the ethylene. In presence of finely divided nickel, reaction (1) becomes explosive at the ordinary temperature, (2) becomes appreciable at 30° and has its optimum at 130—250°, whilst (3) is not markedly influenced, its products being essentially water and carbon dioxide. The results now obtained show that the velocity of reaction (1) is diminished enormously by the presence of ethylene, although at the ordinary temperature this reaction is the only one

taking place to an appreciable extent: when the temperature is raised, reaction (2) assumes an increased velocity, so that most of the ethylene is hydrogenated to ethane at 225°, and reaction (3) also becomes of measurable velocity. Various possible causes of the alternations in the velocities of reactions (1) and (2) are discussed. The retardation of reaction (1) by ethylene is analogous, up to a certain temperature, to the similar effect observed by Henry (*Phil. Mag.*, 1836, **65**, 329) for carbon monoxide, with which oxygen combines more rapidly than with hydrogen in presence of platinum. The presence of atomic or active hydrogen produced either, as Sabatier supposed, by way of a hypothetical hydride or in some other way, is insufficient to explain the author's results, which may, however, be determined in some degree by the specific adsorption of gases by metallic catalysts studied by Taylor and Burns (*A.*, 1921, ii, 630). The possible formation of an unstable compound of nickel and ethylene, analogous to the formation of nickel tetracarbonyl from nickel and carbon monoxide, is also discussed.

In presence of copper, the velocities of the different reactions are only slightly increased, the greatest velocity at all temperatures being that of the formation of water; the copper undergoes marked oxidation, which favours oxidation phenomena. None of the reactions is appreciably accelerated by the presence of silver. The formation of water under the influence of platinum is retarded by the presence of ethylene, but is the principal reaction taking place. In the system containing nitrobenzene, the reduction of the latter in presence of nickel is more rapid than the formation of water from its elements. The results of Bone and Wheeler (*T.*, 1904, **85**, 1637) are discussed.

T. H. P.

Catalytic De-hydroxidation of Formic Acid. ERICH MÜLLER [with K. SPONSEL] (*Z. Elektrochem.*, 1922, **28**, 307—310).—The catalytic conversion of formic acid into hydrogen and carbon dioxide, by means of rhodium, first studied by Deville and Debray in 1874, is due to the presence of impurities in the catalyst, the catalytic activity decreasing as the purity of the metal increases. Osmium exhibits considerable catalytic activity, and a detailed investigation has been made of the dependence of the activity on the mode of preparation of this catalyst, more especially on the temperature of reduction of the metal, the composition of the catalysed solution as regards the concentration of the acid and the presence or absence of sodium formate, and the mass of catalyst employed. Osmium prepared by reduction of the hydrated dioxide at 150° possesses very considerable activity, 0.3 gram effecting the evolution of 10 litres of gas in three hours. The activity is increased by any conditions favouring the existence of the catalyst in the colloidal state.

J. S. G. T.

The Use of the Oxides of Platinum for the Catalytic Reduction of Organic Compounds. I. V. VOORHEES and ROGER ADAMS (*J. Amer. Chem. Soc.*, 1922, **44**, 1397—1405).—When

chloroplatinic acid is fused with sodium nitrate a brown oxide of platinum is obtained which is an excellent catalyst for the reduction of various types of organic compounds. The speed of reduction with this catalyst is greater than when ordinary platinum black is used. The most satisfactory conditions for preparing this oxide in its most active form have yet to be worked out. W. G.

Effect of a Magnetic Field on Catalysis by Ions in the Presence of a Paramagnetic Salt. WILLIAM EDWARD GARNER and DOUGLAS NORMAN JACKMAN (T., 1922, **121**, 1298—1299).

Analysis of the Electronic Structure of the Elements.

A. DAUVILLIER (*J. Phys. Radium*, 1922, [vi] **3**, 154—180).—A further discussion of the theory of atomic structure (cf. A., 1921, ii, 475, and this vol., ii, 43). A new arrangement of the periodic table is proposed which conflicts with the chemist's view as to the positions of the elements of the manganese and iron groups in the periodic table. The spectra and atomic volumes of these elements are in accord with the view that they should be placed, together with oxygen, in the sixth group. They should possess only six electrons in the outer shell of the atom. This would remove certain anomalies present in the modern theory of the periodic table. An arrangement of the electrons in the shells is advanced which differs in some respects from that of Langmuir. The maximum number of the electrons in the successive shells of the atom are respectively 2, 8, 18, 18, 32, 8 instead of 2, 8, 8, 18, 18, 32 according to Langmuir. An apparatus is described by which the author has measured the *K*, *L*, and *M* series of the heavy elements. These results, which will be communicated in a future paper, throw light on the distribution of the electrons in the various shells of the atom. W. E. G.

Complex Compounds. A. MAGNUS (*Physikal. Z.*, 1922, **23**, 241—247).—A theoretical paper in which the theory of Kossel (A., 1916, ii, 243) is applied to the stability relations and the valency forces of complex compounds. On this theory, the most stable complex is one which is formed from its components with the greatest liberation of energy. The case of positively charged central atoms is considered. The stability of a complex is dependent on the number of positive charges (*n*) on the central atom and on the number of groups (*p*) held in combination. It is shown that the stability of complexes with different values of *n* and *p* may be derived from purely electrostatic considerations. Maximum values are found for the stability at values of *p* which are in agreement with experience. For complexes with a small charge on the central atom, high co-ordination numbers are not to be expected unless the groups attached are bipolar in nature. Ions of the form XO_6 , in consequence of the energy conditions, are rarely stable. The stability of hydrates and compounds with ammonia is shown to

be dependent on the bipolar nature of the water and ammonia molecules.

W. E. G.

The Electron Theory of Valency as Applied to Organic Compounds. JULIUS STIEGLITZ (*J. Amer. Chem. Soc.*, 1922, **44**, 1293—1313).—A theoretical paper in which the author discusses the relative merits of the theories of polar and non-polar valencies in organic compounds as applied to various types of compounds. In the molecular rearrangement of *s*-bistriphenylmethylhydrazine the theory of polar valencies gives an evident cause for the rearrangement, revealing a "fault" in the molecule, and expresses the intramolecular oxidation and reduction characteristic of the rearrangement. This theory is also applicable to the analogous molecular rearrangements of chloro- and bromo-amides, of hydroxylamine derivatives, and of peroxides. In the absorption reactions of carbon dioxide and of a large number of organic compounds containing unsaturated groups of the types C:O, C:S, C:N, polar valencies are recognised as the directing forces. The same applies to ethylene hydrocarbons.

The author develops the Crum Brown-Gibson rule of substitution in the benzene series from a polar electronic structure for benzene, which does not require the assumption of negative hydrogen, but is based on the application to benzene of results firmly established in the aliphatic series. The different behaviour of negative halogen, in compounds of the type RX, and of positive halogen, in compounds of the type RNHX, is taken as further evidence in support of polar valencies.

Evidence of the complete polar structure of acids, such as sulphuric, phosphoric, or organic acids is found in (a) Faraday's observation of the deposition of sulphur at the negative electrode in the electrolysis of concentrated sulphuric acid, (b) the work of Bird and Diggs (*A.*, 1914, ii, 614) on the use of yellow phosphorus as the equivalent of a metal for the production of an electric current, (c) the production of currents by the use of organic compounds as the source of escape of electrons in oxidation-reduction cells.

In the oxidation and reduction of organic compounds at the ordinary temperature, exposed valencies on double and triple bonds, in bivalent and tervalent carbon, etc., are the common seat of oxidation and reduction reactions. Although polarity in organic compounds is insisted on as giving an invaluable guide in following organic reactions, it need not be of the extreme character shown by common salts, but may well be of the character proposed by Bohr, Lewis, and Kossel, where the transfer of electrons from atom to atom is not so complete as in the case of common electrolytes.

W. G.

A Discussion of Triple Salts. HORACE L. WELLS (*Amer. J. Sci.*, 1922, [v], **4**, 27—30).—In some cases where analogous triple salts are known, these occur in extensive series, but there are a great many cases where analogy is lacking between salts of analogous

metals, and there appear to be no definite laws, based on the valency or other characters of the constituent salts, according to which they are formed. A number of examples of triple chlorides, thiocyanates, and nitrites are quoted. E. H. R.

New Small Autoclave for Hydrolysis Experiments.

R. EBERHARD GROSS (*Z. physiol. Chem.*, 1922, **120**, 185—188).—A description of a small autoclave which can be heated and cooled quickly. It is therefore suitable for hydrolysis experiments in which samples need to be withdrawn at different times for testing for various reactions. S. S. Z.

Inorganic Chemistry.

The Rectilinear Diameter of Hydrogen. E. MATHIAS, C. A. CROMMELIN, and H. KAMERLINGH ONNES (*Ann. Physique*, 1922, **17**, 463—474).—A résumé of work already published (*A.*, 1921, ii, 256; this vol., ii, 440). W. G.

The Rectilinear Diameter of Oxygen. E. MATHIAS and H. KAMERLINGH ONNES (*Ann. Physique*, 1922, **17**, 416—441).—A résumé of work already published (*A.*, 1910, ii, 771, 829; 1911, ii, 387). W. G.

Equilibrium between Sulphur and Iodine in Solution. M. AMADORI (*Gazzetta*, 1922, **52**, i, 387—394).—In view of the fact that the results of recent investigations indicate that the so-called compounds of sulphur with iodine described by various authors are merely mixtures of the two elements, the author has examined the solubility and cryoscopic relations of sulphur and iodine when present together. It is found that the separate solubilities of each of these elements in carbon disulphide, benzene, and bromoform are less than those which obtain in presence of the other element; in the first of these solvents both solubilities are increased by more than 100% by mixing the sulphur and iodine. The corresponding solubility curves of the mixtures in benzene and in carbon disulphide consist of two branches meeting in a point of double saturation.

Cryoscopic measurements of the mixtures in bromoform solution show that the depression of the freezing point of the solvent is somewhat less than that calculated from the molecular weights of the sulphur and iodine. No evidence is obtained of the existence of a solid compound of the two elements. T. H. P.

The Crystallisation of Amorphous Tellurium. A. DAMIENS (*Compt. rend.*, 1922, 174, 1548—1550).—From a study of the heats of reaction of different forms of tellurium with a mixture of bromine and bromine water, Berthelot and Fabre have found the relationship $\text{Te}_{\text{cryst}} = \text{Te}_{\text{amorph}} + 12.096 \text{ cal. (for 64 grams)}$. Previous work on the allotropy of tellurium did not confirm this equation (cf. this vol., ii, 498), and the author has accordingly repeated the work of Berthelot and Fabre, using, however, a mixture of bromine and concentrated hydrochloric acid diluted with its own volume of water. His results do not confirm those of Berthelot and Fabre, but establish the equation $\text{Te}_{\text{amorph.}} = \text{Te}_{\text{cryst}} + 2.63 \text{ cal.}$, which is in accord with his work on the allotropy of tellurium (*loc. cit.*).
W. G.

The Rectilinear Diameter of Nitrogen. E. MATHIAS, H. KAMERLINGH ONNES, and C. A. CROMMELIN (*Ann. Physique*, 1922, 17, 455—463). A more detailed account of work already published (*A.*, 1915, ii, 143).
W. G.

Normal Density of [Chemically Pure] Nitrogen. E. MOLES (*J. Chim. physique*, 1922, 19, 283—289).—The results of twenty-nine determinations of the density of pure nitrogen have been systematically discussed, and from calculations, the weight of the litre of nitrogen at 0°, 760 mm., and 45° latitude is found to be $L_N = 1.2507 \pm 0.0001 \text{ grams}$. The measurements used are those of Rayleigh and Ramsay, Leduc, Gray, and Moles, in which two entirely different methods were used, and the material was drawn from eight different sources. Under the conditions 0°, 760 mm., and $g = 980.665$, the rounded value is $L_N = 1.2507 \text{ grams}$. The actual value obtained from nitrogen obtained from sodium nitrite is $L_N = 1.2506 \text{ grams}$, a value which is almost identical with the above-cited mean. This result is remarkable, since it was obtained from volumetric determinations.
J. F. S.

The Synthesis of Ammonia by Collision with Slow-moving Electrons. E. BUCH ANDERSEN (*Z. Physik*, 1922, 10, 54—62).—A study of the action of a stream of electrons on the production of ammonia from mixtures of nitrogen and hydrogen at low pressures. A current was passed between a glowing tungsten wire and two platinum electrodes in an atmosphere of the two gases. The current was kept constant and the effect of variations in the composition of the gas mixture and in the applied *E.M.F.* were investigated. The maximum rate of formation of ammonia did not correspond with the stoichiometric mixture, but with mixtures containing a great excess of nitrogen. This result is in agreement with the view that the first step in the production of ammonia is the ionisation of the nitrogen molecule. Curves are given showing the velocity of formation of ammonia plotted against the applied electromotive force. These curves show a series of maxima at 22, 26, and 33 volts. No appreciable quantities of ammonia were produced below an applied electromotive force of

17.7 volts. This potential is identical with the ionisation potential of nitrogen (17.75 volts). It is, however, difficult to distinguish by direct measurement between the ionisation potentials of hydrogen and nitrogen.

W. E. G.

Oxidation of Nitric Oxide and the Recovery of Nitrogen Oxides from Mixtures with Air. E. BRINER, S. NIEWIAZSKI, and J. WISWALD (*J. Chim. physique*, 1922, **19**, 290—309; *Helv. Chim. Acta.*, 1922, **5**, 432).—The work which has been done on the commercial oxidation of nitric oxide and the recovery of oxides of nitrogen is reviewed and a number of points requiring further investigation are emphasised. Some of these points have been investigated by the authors. It is shown that the formation of nitrite by absorption of a gaseous system $\text{NO}-\text{NO}_2$ by alkali takes place according to the scheme proposed by Leblanc (*Z. Elektrochem.*, 1906, **12**, 541), based on the fact that nitrous anhydride exists in small concentrations, and this is in keeping with Raschig's experiments. Experiments on the oxidation of nitric oxide at low temperatures confirm the results of Lunge, Bodenstein, and Wourtsel, and show that the oxidation proceeds directly from nitric oxide to nitrogen tetroxide without the intermediate formation of nitrous anhydride. Comparative experiments made at ordinary temperatures, the temperature of carbon dioxide "snow" (-80°), and the temperature of liquid air (-190°) show that a low temperature is very favourable to the oxidation of oxides of nitrogen and to the recovery of the oxides. It follows from the experiments that cooling to a very low temperature will be the best means of effecting a complete recovery of the oxides of nitrogen from very dilute mixtures of these gases such as are obtained after passage through the electric arc, provided that the technical difficulties can be overcome, whilst without this low temperature very large oxidation chambers will be necessary.

J. F. S.

Economic Realisation of Oxidation Reactions in Factories where Nitric Acid is Synthesised. Applications. CAMILLE MATIGNON (*Bull. Soc. chim.*, 1922, [iv], **31**, 555—561).—The author proposes to use the dilute nitric acid such as is obtained at first in the fixation of atmospheric nitrogen or in the oxidation of ammonia, for chemical preparations in which it can be used as an oxidising agent, providing the reduction of the nitric acid does not proceed beyond the stage of nitric oxide. The oxides of nitrogen can be reconverted into nitric acid by atmospheric oxygen and again absorbed in the towers. In this way there is practically no loss of nitric acid and thus the expense of the oxidation process is very small. Suggested applications are the preparation of copper sulphate from copper by the combined action of sulphuric and nitric acids, and the preparation of oxalic acid from sugar or molasses by oxidation with nitric acid. This latter process was satisfactorily performed on a cargo of sugar damaged by sea water.

W. G.

The Accessory Elements of the Dephosphoration Slags.

A. DEMOLON (*Compt. rend.*, 1922, 174, 1703—1706).—Estimations have been made on a number of slags of the amounts of calcium soluble in solvents under definite conditions. The solvents used were: distilled water, 5% sugar solution, 2% phenol, a neutral solution of ammonium humate, cold solutions of ammonium chloride of different strengths, a saturated solution of carbon dioxide, and mineral acids. The amount of free calcium oxide in the slags was small, varying from 1% to 3%. In all the solvents the amount of calcium going into solution increased at first very rapidly with the time of shaking and then only very slowly over a long period of time. This is considered to be due to the presence of calcium silicates which react slowly with the solvents. The amounts of magnesium oxide present in the slags varied from 3% to 15% with an average of 8.90%. The amount of manganese found was fairly constant at about 4—5%, which was easily soluble in 2% citric acid.

W. G.

Magnetochemical Investigations of Constitutions in Mineral Chemistry. The Acids of Arsenic.

PAUL PASCAL (*Compt. rend.*, 1922, 174, 1698—1700).—The results of measurements of the magnetic susceptibilities of a large number of arsenic compounds indicate that combined arsenic has two atomic susceptibilities according to the degree of saturation of its compounds. The results furnish further evidence in support of the rule that the logarithm of the atomic susceptibility of an element is a linear function of the atomic weight in each natural family. It is verified exactly for the group phosphorus, quinquevalent arsenic, and quinquevalent antimony and for the group trivalent arsenic, antimony and bismuth.

W. G.

Reaction between Boron Nitride and Various Metallic Oxides with Production of Nitric Oxide.

U. SBORGI and A. G. NASINI (*Gazzetta*, 1922, 52, i, 369—387).—Estimations have been made of the yields of nitric oxide obtained when air is passed over mixtures of boron nitride with various metallic oxides heated at different temperatures. Four samples of the nitride, prepared and stored in different ways, were employed. The yield of nitric oxide obtained varies greatly with these different samples and also with the nature of the admixed oxide. With the oxides Fe_2O_3 , Ni_2O_3 , Co_2O_3 , MnO_3 , Mn_3O_4 and CuO , yields as high as 72% are given, and the residue remaining after the reaction gives an increased yield of nitric oxide with a fresh quantity of boron nitride; this result is repeated four or five times, the yield increasing each time until the residue becomes converted into a compact mass.

T. H. P.

Carbon and its Neighbours in the Periodic System.

ALFRED STOCK (*Z. angew. Chem.*, 1922, 35, 341—343).—A discussion of the regularities occurring between the compounds of

the elements boron, carbon, nitrogen, and silicon. The greatest resemblances are found between the hydrides and their derivatives containing oxygen, nitrogen, or chlorine. In these compounds, carbon unites the chemical properties of its neighbours. It resembles boron in its tendency to form long chains of atoms, nitrogen in forming volatile compounds, and silicon and boron in giving rise to high-boiling derivatives. The property of combining equally readily with either electro-positive or electro-negative elements is shared to some extent by nitrogen and silicon. W. E. G.

The Preparation of Carbon from Carbon Monoxide by means of a Catalyst. J. P. WIBAUT (*Rec. trav. chim.*, 1922, **41**, 400—401).—An attempt to prepare pure carbon which, from the method of preparation, could not contain hydrogen or hydrocarbons, was unsuccessful. Carbon monoxide was passed over iron oxide at a temperature of about 450° and a good yield of carbon obtained. The resulting gas contained from 80—90% of carbon dioxide. The catalyst, however, was found to be fairly evenly distributed throughout the carbon, and prolonged attempts to extract it yielded a product the ash content of which was above 6%. H. J. E.

The Rectilinear Diameter of Argon. E. MATHIAS, H. KAMERLINGH ONNES, and C. A. CROMMELIN (*Ann. Physique*, 1922, **17**, 442—455).—A résumé of work already published (*A.*, 1913, ii, 112, 478). W. G.

The Analysis by Positive Rays of the Heavier Constituents of the Atmosphere; of the Gases in a Vessel in which Radium Chloride had been stored for Thirteen Years, and of Gases given off by Deflagrated Metals. (Sir) J. J. THOMSON (*Proc. Roy. Soc.*, 1922, [A], **101**, 290—299).—The positive ray method of analysis is applied to the solution of several problems. The residues from a thousand tons of liquid air have been examined for the presence of molecules heavier than krypton and xenon. Two new constituents of the atmosphere were found with molecular weights 163 and 260. The lighter of these is the more abundant, but the quantity in the air is very small compared with xenon. There is evidence that the lighter molecule carries two positive charges. No place in the periodic table can be found for a new element with this atomic weight, and it is possible that the two new lines are due to diatomic molecules of xenon and krypton.

The gases from 70 mg. of radium, stored in an evacuated vessel by (Sir) J. Dewar in 1909, have been analysed by the positive ray method. Helium and hydrogen were both present, but no neon or H₂ was found. In addition, a faint line, for which $m/e = 5$, seems to indicate a compound of helium with one atom of hydrogen. The analysis was also applied to gases which had stood over radium and to gases lit by deflagrating wires. The presence of doubly, triply, and quadruply charged atoms of oxygen and nitrogen, and

of doubly and triply charged atoms of carbon, was detected. A compound $m/e=10$ is believed to exist, consisting of an atom of oxygen and four atoms of hydrogen. This invariably carries a double charge. The deflagration of fine wires of tungsten, gold, or copper by powerful electric currents did not yield any helium, although H_2 was detected. W. E. G.

Constitution of Metallic Substances. CHARLES A. KRAUS (*J. Amer. Chem. Soc.*, 1922, **44**, 1216—1239).—A theoretical paper in which it is pointed out that the constitution of substances cannot be derived from their properties in a condensed state with any considerable degree of accuracy. In order to determine the constitutions of metallic substances, it is necessary to study their properties at low concentrations. At low concentrations, the elementary metals are salt-like substances, and it is suggested that metallic compounds will exhibit salt-like properties in even a more decided manner. In solutions of metallic compounds in liquid ammonia, the more electro-negative element functions as anion, as follows from the fact that it is precipitated on the anode on electrolysis. All but the strongly electro-positive elements exhibit a negative as well as a positive valency. The normal anions of metallic elements form complex anions in the presence of the element in question. These complex metallic compounds are similar to the complex iodides and sulphides which have been studied in aqueous solution. The property of forming complex anions is one common to many metallic as well as non-metallic elements. It is shown that the complex telluride-ion carries two charges. In solution in ammonia, the complex anions $TeTe^-$ and Te_3Te^- exist. It is inferred that the anions of the metals of the fifth group carry three charges, and those of the fourth four charges. Since it has been shown that selenium, tellurium, bismuth, antimony, arsenic, tin, and lead form complex anions in ammonia solution, it may be inferred that other metals not soluble in ammonia form similar complex anions. Metallic compounds of this class are, therefore, virtually salts. The multiplicity of compounds derivable from a given pair of metallic elements is accounted for on the hypothesis that in their compounds the more electro-negative elements form complex anions, and this view brings metallic compounds into line with the present conceptions of atomic structure. The physical properties of metallic compounds are in keeping with the hypothesis that they possess a salt-like structure. It is pointed out that the energy effect accompanying the formation of metallic compounds is of the same order of magnitude as that accompanying the formation of salts, and as in the case of salts, the energy change is the greater the more electro-negative one element is with respect to the other. Electro-positive groups, such as the ammonium group, are virtually metals, although their stability in the free state is not sufficient to permit their isolation in most instances. The most stable groups possess sufficient stability to demonstrate their

metallic properties in the free state. Such groups resemble elements in their properties. The property of metallicity is not an atomic one, it may be imparted to non-metallic elements by combination with other non-metallic elements, and it is due to the presence of uncombined negative electrons. The electrons which impart metallic properties to an element are those to which the chemical reactions of this element with other elements are due. The reaction between strongly electro-positive and strongly electro-negative elements, or groups of elements, consists essentially in a combination of the negative electrons of the electro-positive constituent with the atoms of the electro-negative constituent. In metal ammonia complexes of the type $\text{Ca}(\text{NH}_3)_6$, which are metallic substances, the negative electrons are not primarily concerned. The experimental data on which the foregoing conclusions are based are to be published shortly. J. F. S.

The Possible Existence of Metallic Compounds in the State of Vapour. A. EUCKEN and O. NEUMANN (*Z. Elektrochem.*, 1922, 28, 322—324).—Sodium amalgam was distilled at pressures between 5 and 10 mm. of mercury, and the boiling point and corresponding composition of the vapour and liquid phases were determined. The results indicate that, contrary to the suggestion of von Wartenberg (A., 1915, ii, 226), no appreciable amount of a compound of sodium and mercury was formed in the state of vapour at about 500°. J. S. G. T.

Interaction of Sodium Chloride and Silica. FRANCIS HERBERT CLEWS and HUGH VERNON THOMPSON (T., 1922, 121, 1442—1448).

Chlorites of Sodium and other Metals. G. R. LEVI (*Gazzetta*, 1922, 52, i, 417—420; *Atti R. Accad. Lincei*, 1922, [v], 31, i, 212—216; cf. this vol., i, 527).—The following new chlorites are described. *Sodium chlorite*, $\text{NaClO}_2 \cdot 3\text{H}_2\text{O}$, prepared from barium chlorite and sodium sulphate, forms lustrous laminæ, and, when anhydrous, explodes on percussion. *Lithium chlorite*, similarly prepared, forms anhydrous, deliquescent crystals and explodes on percussion. *Calcium chlorite*, prepared from calcium peroxide and chlorine dioxide, explodes on percussion and decomposes completely in contact with a heated wire. *Strontium chlorite*, similarly obtained, explodes on percussion, and in contact with a hot wire decomposes with evolution of heat. *Thallous chlorite* explodes on percussion and decomposes immediately at 70—80°. T. H. P.

Oxidation and Reduction Reactions with Chlorites. GIORGIO RENATO LEVI (*Atti R. Accad. Lincei*, 1922, [v], 31, i, 370—373; cf. preceding abstract).—At 180—200°, sodium chlorite undergoes decomposition quantitatively in accordance with the equation $3\text{NaClO}_2 = 2\text{NaClO}_3 + \text{NaCl}$. The action of ozone on the salt in aqueous solution yields chlorine dioxide, $2\text{NaClO}_2 + \text{O}_3 + \text{H}_2\text{O} = 2\text{NaOH} + \text{O}_2 + 2\text{ClO}_2$; if the dioxide is not carried away

by the ozone, it reacts either with the sodium hydroxide, $2\text{NaOH} + 2\text{ClO}_2 = \text{NaClO}_2 + \text{NaClO}_3 + \text{H}_2\text{O}$, or with hydrogen peroxide, if this has been formed by the excess of ozone, $2\text{NaOH} + \text{H}_2\text{O}_2 + 2\text{ClO}_2 = 2\text{NaClO}_2 + 2\text{H}_2\text{O} + \text{O}_2$. In aqueous solution, sodium chlorite and nitrite react, with development of heat, forming the chloride and nitrate, $2\text{NaNO}_2 + \text{NaClO}_2 = \text{NaCl} + 2\text{NaNO}_3$. The reaction between a chlorite and a ferrocyanide is expressed by the equation $4\text{K}_4\text{Fe}(\text{CN})_6 + \text{NaClO}_2 + 2\text{H}_2\text{O} = \text{NaCl} + 4\text{K}_3\text{Fe}(\text{CN})_6 + 4\text{KOH}$, and may be rendered complete either by gradual neutralisation of the alkali hydroxide by means of dilute sulphuric acid or by addition of a salt such as magnesium sulphate, magnesium hydroxide being then deposited. With an iodide, sodium chlorite reacts with liberation of iodine, $\text{NaClO}_2 + 4\text{KI} + 2\text{H}_2\text{O} = 4\text{I} + 4\text{KOH} + \text{NaCl}$ (cf. Bray, A., 1906, ii, 222, 223, 278), and here too the reaction may be carried to completion by addition of a substance, such as boric acid, capable of neutralising the alkalinity; if the liquid is heated on a water-bath and in an apparatus which prevents sublimation of the iodine, the reaction may be used for the estimation of a chlorite in presence of a chlorate, which does not liberate iodine under these conditions. In neutral solution, chlorites do not act on alkali bromides, but in presence of sulphuric acid bromine is set free; the behaviour towards bromates is similar. From acid solutions of iodates and periodates no iodine is liberated, oxygenated compounds of chlorine, formed by decomposition of the chlorous acid, being obtained.

In neutral solution sodium chlorite reacts slowly with a sulphite, thus: $2\text{Na}_2\text{SO}_3 + \text{NaClO}_2 = 2\text{Na}_2\text{SO}_4 + \text{NaCl}$; with solutions of free sulphurous acid the corresponding reaction takes place immediately. The reaction, $\text{Na}_2\text{S}_2\text{O}_3 + 2\text{NaClO}_2 + \text{H}_2\text{O} = 2\text{NaCl} + 2\text{NaHSO}_4$, which occurs with development of heat, is complete only when either a slight excess of chlorite is used or the liquid is kept neutral by means of magnesium oxide; otherwise a little sulphurous acid is formed. With thiocyanates, chlorites react principally according to the equation $2\text{NaCNS} + 3\text{NaClO}_2 + 2\text{H}_2\text{O} = 2\text{NaHSO}_4 + 2\text{HCN} + 3\text{NaCl}$, slight excess of chlorite being required for the complete oxidation of the sulphur of the thiocyanate to sulphuric acid. Chlorites oxidise hydrogen sulphide to sulphuric acid, and sodium formate and oxalate to sodium hydrogen carbonate. T. H. P.

Borates. VI. The System $(\text{NH}_4)_2\text{O} - \text{B}_2\text{O}_3 - \text{H}_2\text{O}$ at 45° and at 90° . UMBERTO SBORGI and LELIO FERRI (*Atti R. Accad. Lincei*, 1922, [v], 31, i, 324—329; cf. A., 1921, ii, 580).—In this system at 45° are found the 1 : 5 : 8- and the 1 : 2 : 4-compounds, in addition to the 1 : 4 : 6-compound occurring at 60° ; the first two compounds are able to exist unchanged in presence of pure water, but the 1 : 4 : 6-compound is found only in solutions containing an excess of $(\text{NH}_4)_2\text{O}$ and having temperatures above that of the surrounding air. The same three compounds are formed also at 90° . T. H. P.

The Electrolytic Preparation of Sodium Perborate. KURT ARNDT and ERNST HANTGE (*Z. Elektrochem*, 1922, 28, 263—273).—The conditions under which sodium perborate may be prepared electrolytically have been investigated. It is shown that this salt is most efficiently prepared from a solution of borax and sodium carbonate containing 120 grams of anhydrous sodium carbonate and 30 grams of borax per litre of solution. A small quantity of sodium chromate (0.5 gram per litre) and one drop of turkey red oil added to the electrolyte increases the efficiency, since this reduces the cathodic reduction to a minimum. A large anodic current density is requisite and to obtain this an anode of bright platinum is employed, whilst a water-cooled length of tin tubing serves as cathode. The most suitable current density lies between 10 and 20 amp./dm.², and the most suitable temperature is 14—16°; above this temperature the yield of perborate is greatly reduced. During electrolysis, the bath loses carbon dioxide, and if this loss is not replaced the current yield becomes very poor owing to the large concentration of hydroxyl-ions. With an increasing concentration of the per-salt, the yield decreases during the electrolysis. The analysis of the hydrogen-oxygen gas mixture evolved during the electrolysis shows that a strong decrease in the anodic oxidation is responsible for the falling off of the yield. The merest trace of platinum salt in the electrolyte has a very bad effect on the yield, and the presence of iron in the sodium carbonate has a similar bad effect. Both substances decrease the anodic oxidation and increase the cathodic reduction. The addition of cyanide only partly eliminates the deleterious effect of the iron. The curve produced by plotting the current density against the anodic potential shows a sharp inflexion point at 1.47 volts. This potential is due to the concentration of a higher oxide of platinum, presumably PtO₃, which is formed as a solid solution in the anode material by the discharge of hydroxyl ions. It is likely that this higher oxide, in keeping with its potential and oxidising power, acts as an intermediate product in the formation of sodium perborate. J. F. S.

Structure of Crystals of Lithium and some of its Compounds with Light Elements. II. Lithium Hydride. J. M. BIJVOET and A. KARSSSEN (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 25, 26—30; cf. A. 1921, ii, 200).—Using the method of Debye and Scheerer, the authors have prepared and investigated the X-ray photographs of lithium hydride. The results show that lithium hydride crystallises in the regular system with four molecules in the elementary cell which has a side of length $a = 4.10 \times 10^{-8}$ cm. The density of lithium hydride is calculated to be 0.76 ± 0.01 . The structure is that of sodium chloride, with positive lithium ions and negative hydrogen ions. Systems of two-electron rings are round both the lithium and hydrogen nuclei with radii $0.05a$ and $0.6a$, respectively, and the planes of which are normal to non-intersecting trigonal axes. J. F. S.

Calcium-Ammonium. E. BOTOLFSEN (*Bull. Soc. chim.*, 1922, [iv], 31 561—567); cf. cf. Biltz and Hüttig, A., 1921, ii, 201).—Solid calcium-ammonium is formed within the temperature limits -15° and $+30^{\circ}$. The pasty state sometimes recorded in the literature is probably due to the presence of traces of sodium in the calcium used. The author confirms previous work on this compound (*loc. cit.*) in certain respects and finds that when it is heated in a vacuum at about 34° it decomposes explosively, the temperature depending slightly on the vacuum obtained. The products of this decomposition are the nitride and hydride of calcium and hydrogen. It is suggested that calcium is capable of forming a series of compounds with ammonia in much the same way as is iodine. W. G.

The Transformation of Gypsum into Ammonium Sulphate. C. MATIGNON and M. FRÉJACQUES (*Compt. rend.*, 1922, 175, 33—35).—The authors obtained a 96% yield in the interaction of an industrial sample of gypsum with ammonium carbonate (cf. Neumann, A., 1921, ii, 587). The curves giving the velocity of the reaction show the existence of two distinct phases. During the first phase the velocity is slower than in the second phase. The presence of ammonium sulphate in the original solution causes a diminution in the velocity of the reaction, but in all cases equilibrium is complete at the end of two and a half hours. W. G.

Magnesium-Cadmium Alloys. LÉON GUILLET (*Rev. Met.*, 1922, 19, 359—365).—The magnesium-cadmium alloys consist of a series of solid solutions and the compound CdMg, which is soluble in all proportions in either metal. Cadmium increases the hardness of magnesium, but the alloys are malleable up to a content of 55% Cd. With more than this they become brittle at ordinary temperatures, due to the formation of a new solid solution stable below 246° and consisting of the compound CdMg with excess of either magnesium or cadmium [cf. *J. Soc. Chem. Ind.*, 1922, 553A]. A. R. P.

The Quaternary System Potassium Sulphate-Magnesium Sulphate-Ammonium Sulphate-Water. ARNOLD WESTON (T., 1922, 121, 1223—1237).

Chemical and Electrochemical Behaviour of Salts of the Acids of Lead. G. GRUBE (*Z. Elektrochem.*, 1922, 28, 273—289).—Two new methods for the preparation of alkali salts of plumbic acid are described, the first of which consists in the anodic solution of a pure lead anode in a strong solution of alkali hydroxide by means of a direct current of high current density on which is superimposed an alternating current. In the second method, a solution of an alkali plumbite in concentrated alkali hydroxide is submitted to anodic oxidation at a platinum electrode. In both methods the alkali plumbates crystallise from the electrolyte in the pure condition during the electrolysis. The oxidation potentials of the processes

$\text{Pb}^{++} + 2 \oplus \longrightarrow \text{Pb}^{+++}$ and $\text{PbO}_2'' + 2\cdot\text{OH}' + 2 \oplus \longrightarrow \text{PbO}_3'' + \text{H}_2\text{O}$ have been systematically investigated in alkaline solution and also the potential of lead in solutions of alkali plumbite has been measured. It is shown that the normal potential of lead against potassium plumbite in 8.42 *N*-potassium hydroxide at 18° is $\epsilon_h - 0.613$ volt, whilst the normal potential of plumbite-plumbate in potassium hydroxide of the same concentration is $\epsilon_h + 0.208$ volt. From these values the normal potential of lead against alkali plumbate solution is calculated to be $\epsilon_h - 0.203$ volt. In all cases a decrease in the concentration of the alkali hydroxide displaces the potential toward more negative values. The nature of the plumbo-plumbates which separate from highly concentrated solutions of alkali plumbates and plumbites in alkali hydroxides has also been investigated, and it is shown that the composition of the deposit varies with the alkalinity of the solution in the sense that from solutions with a concentration of alkali hydroxide less than 4.9*N*, the compound $\text{Pb}_2\text{O}_3\cdot 3\text{H}_2\text{O}$ separates, whilst from solutions more concentrated than 6.9*N*, Pb_3O_4 separates. Hence it follows that Pb_3O_4 is to be regarded as the lead salt of ortho-plumbic acid, and that in very alkaline solutions of plumbates the anions of meta-plumbic acid, PbO_3'' and ortho-plumbic acid PbO_4'''' , are both present. To ascertain whether the alkali plumbates are salts of the acid $\text{H}_2[\text{Pb}(\text{OH})_6]$ or of H_2PbO_3 , the author has investigated the dehydration of sodium plumbate, $\text{Na}_2\text{PbO}_3\cdot 3\text{H}_2\text{O}$. It is shown that on heating this salt the water is all expelled without decomposition of the salt, from which the conclusion is drawn that the three molecules of water are to be regarded as water of crystallisation and that sodium plumbate is a salt of metaplumbic acid. J. F. S.

Effect of Grinding on the Apparent Density of Lead Oxides.

O. W. BROWN, S. V. COOK, and J. C. WARNER (*J. Physical Chem.*, 1922, **26**, 477—480).—The apparent density of heavy, crystalline lead oxide is diminished initially by grinding, the minimum value being attained by about ten hours' grinding in a pebble mill making 50 revolutions per minute. Thereafter, the apparent density increases, rapidly at first, to a constant value greater than the initial value. The initial decrease of apparent density is not shown by a light, amorphous lead oxide. J. S. G. T.

Thermal Analysis of the System $\text{Ti}_2\text{O}-\text{B}_2\text{O}_3$. G. CANNERI and R. MORELLI (*Atti R. Accad. Lincei*, 1922, [v], **31**, i, 109—111; cf. Buchtala, A., 1914, ii, 135).—The fusion diagram of this system exhibits three maxima corresponding with thallous metaborate, TiBO_2 , m. p. about 474°; thallous pyroborate, $\text{Ti}_4\text{B}_2\text{O}_5$, m. p. about 434°; and normal thallous borate, Ti_3BO_3 , m. p. 370° (decomp.). T. H. P.

Phenomena of Diffusion in Metals in the Solid State and Cementation of Non-ferrous Metals. II. Cementation of Copper by means of Chromo-manganese. G. SIROVICH and A. CARTOCETI (*Gazzetta*, 1922, **52**, i, 436—442).—By means of

the arrangement previously used (this vol., ii, 68) experiments have been made on the cementation of copper by chromo-manganese containing 30.4% and 63.8% of chromium and manganese respectively. At 900°, the cementation is so intense that, in a few moments a layer of copper-manganese containing more than 20% of manganese is formed; as this alloy melts below 900°, the succeeding experiments were carried out at 800°. Although the iron of the ferro-manganese and the chromium of the chromo-manganese do not migrate to the copper, these metals evidently influence the migration of the manganese; the chromium having a greater effect than the iron. The mechanism of the process of cementation is discussed briefly.

T. H. P.

Action of Sulphur on Cuprous Chloride. FREDERICK WILLIAM PINKARD and WILLIAM WARDLAW (T., 1922, **121**, 1300—1302.)

The Structural Formula of Copper Sulphide. W. GLUUD (*Ber.*, 1922, **55**, [B], 1760—1761; cf. this vol., ii, 446).—If hydrogen sulphide is passed into a 1.5% solution of copper sulphate in ammonia (10%) until only a faint blue colour remains, a variety of copper sulphide is precipitated which yields elementary sulphur on immediate oxidation by air, whereas after preservation for four or five hours it is oxidised to copper sulphate and thiosulphate.

It is suggested that the formulæ Cu_2S and $\text{Cu} \begin{smallmatrix} \text{Cu} \\ \text{Cu} \end{smallmatrix} \text{S}_2$ are to be assigned to the two modifications. This conception is in harmony with the known tendency of copper sulphide to pass into cuprous sulphide. The modification which yields free sulphur does not give potassium thiocyanate when treated with potassium cyanide; this reaction is shown strongly by the other variety. The difference is probably shown in the equations (i) $2\text{CuS} + 10\text{KCN} = \text{K}_6\text{Cu}_2\text{C}_8\text{N}_8 + 2\text{K}_2\text{S} + \text{C}_2\text{N}_2$ and $\text{C}_2\text{N}_2 + \text{H}_2\text{O} = \text{HCN} + \text{HCNO}$; (ii) $\text{Cu}_2\text{S}_2 + 8\text{KCN} = \text{K}_6\text{Cu}_2\text{C}_8\text{N}_8 + \text{K}_2\text{S}_2$ and $\text{K}_2\text{S}_2 + \text{KCN} = \text{K}_2\text{S} + \text{KCNS}$.

H. W.

The Dissociation Pressures of Hydrated Double Sulphates. I. Hydrated Cupric Alkali Sulphates. ROBERT MARTIN CAVEN and JOHN FERGUSON (T., 1922, **121**, 1406—1414).

The Alloys of Cerium. LÉON GUILLET (*Rev. Mét.*, 1922, **19**, 352—358).—The constitutional diagrams of alloys of cerium with iron, copper, aluminium, magnesium, bismuth, tin, and silicon are reproduced and discussed. Except in the case of iron in the iron-rich alloys, no solid solutions are formed with any of these metals, but a large number of definite compounds are known, some of which melt at a much higher temperature than either of the constituents; for example, Bi_2Ce_3 melts at 1,630° and CeAl_2 at 1,475°. There are two allotropic modifications of the compounds CeFe_2 and CeAl_4 . As cerium increases the brittleness of metals with which it alloys, it is not suitable for use as a deoxidiser.

A. R. P.

Reactions of Sodium Hydroxide with Salts of Aluminium.

ÉDOUARD GROBET (*J. Chim. physique*, 1922, 19, 331—335).—Making use of the thermal method of titration previously described (cf. this vol., ii, 578), it is shown that the addition of sodium hydroxide to dilute solutions of aluminium nitrate forms successively aluminium hydroxide, sodium metaluminate, and sodium ortho-aluminate. The addition of sodium hydroxide to dilute solutions of aluminium chloride, aluminium sulphate, and potash alum forms successively aluminium hydroxide, basic sodium aluminate, $\text{Al(ONa)}_3\text{Al(OH)}_3$, and sodium ortho-aluminate. Sodium hydroxide added to concentrated solutions of aluminium chloride, nitrate, or sulphate produces a basic salt of the type $\text{AlX}_3\text{Al(OH)}_3$, which is followed by the hydroxide, meta-aluminate and ortho-aluminate. Concentrated solutions of potash alum yield under the same treatment a basic salt, $\text{Al}_2(\text{SO}_4)_3\cdot 2\text{Al(OH)}_3$, aluminium hydroxide, a basic aluminate, $\text{Al(ONa)}_3\text{Al(OH)}_3$, and the ortho-aluminate.

J. F. S.

Crystal Structures of the Hexa-ammoniates of the Nickel Haloids. RALPH W. G. WYCKOFF (*J. Amer. Chem. Soc.*, 1922, 44, 1239—1245).—The crystal structure of the complex compounds of the type $\text{NiX}_2\cdot 6\text{NH}_3$, formed when ammonia is added to solutions of nickel chloride, bromide, and iodide respectively, has been examined by means of X-ray photographs. It is shown that the arrangement of the atoms in the crystals of these substances is such that the salts are strictly isomorphous with those of ammonium platinichloride (this vol., ii, 214). The dimensions of the unit cell and the values of the variable parameters defining the positions of the nitrogen atoms in the chloride and iodide have been estimated. The following values are recorded: $\text{NiCl}_2\cdot 6\text{NH}_3$, side of unit cell 10.09 Å.U.; $\text{NiBr}_2\cdot 6\text{NH}_3$, length of side of unit cell, 10.48 Å.U.; $\text{NiI}_2\cdot 6\text{NH}_3$, length of side of unit cell 11.01 Å.U. In each case there are four molecules in the unit cell. The nickel atoms occupy the position of platinum, and the halogen atoms those of nitrogen in the ammonium platinichloride crystal.

J. F. S.

Composition and Crystal Structure of Nickel Nitrate Hexammoniate. RALPH W. G. WYCKOFF (*J. Amer. Chem. Soc.*, 1922, 44, 1260—1266).—From analyses, X-ray spectrum measurements, and considerations of space group arrangements, it is shown that neither of the formulæ $\text{Ni(NO}_3)_2\cdot 4\text{NH}_3\cdot \text{H}_2\text{O}$, or $\text{Ni(NO}_3)_2\cdot 4\text{NH}_3\cdot 2\text{H}_2\text{O}$ is correct for the pale blue isotropic octahedra which are obtained when an excess of a concentrated solution of ammonia is added to an aqueous solution of nickel nitrate. The true formula is $\text{Ni(NO}_3)_2\cdot 6\text{NH}_3$. A study of the Laue photographs of these crystals by the generally applicable methods which use the theory of space groups indicates that they have a structure similar to that of the corresponding complex haloids (preceding abstract), in which the nitrate groups replace the halogen groups, and in which the ammonia groups are related to the nickel atoms in exactly the same

way in both compounds. The position of the atoms of the nitrate groups cannot be determined with great accuracy. Both the symmetry characteristics and the diffraction data permit the same kind of displacement of the nitrate nitrogen atoms as is experienced by the sulphur atoms in pyrites. The unit cube contains four molecules and the length of its side is 10.96 Å.U. J. F. S.

Solubility. VI. Solubility of Ammoniates. FRITZ EPHRAIM and PAUL MOSIMANN (*Ber.*, 1922, **55**, [B], 1608—1619; cf. A., 1921, i, 508).—In a previous communication it has been pointed out that the similarity between dissolved substance and solvent has an important influence on solubility and that salts can achieve this similarity with water as solvent by the formation of hydrates or solvates. An experimental examination of this hypothesis has now been attempted by investigating the solubility in water of salts in which the formation of hydrates is rendered difficult. This can be achieved by occupying the co-ordination positions which would eventually be filled by water with other neutral groups, for example, ammonia. A relationship might therefore be expected between the solubility of a series of ammoniates such as $[\text{Ni}(\text{NH}_3)_6]\text{X}_2$ and their ammonia tension. A difficulty, however, arises, since many of these salts are decomposed by water and their solubility has therefore been determined in aqueous ammonia solution, d^{20}_4 0.950, to which half its volume of alcohol (96%) has been added. With the hexamminenickelo-salts there is a parallelism between solubility and ammonia tension in the series iodide \rightarrow bromide \rightarrow chloride \rightarrow thiocyanate, whilst the perchlorate and formate occupy the expected position in the series; on the other hand, in the series tetrathionate \rightarrow thiosulphate \rightarrow sulphide \rightarrow chloride \rightarrow nitrate \rightarrow chlorate, the solubility increases with decreasing ammonia tension. In general, therefore, there is no parallel between ammonia tension and solubility, at any rate in the cases of the ammoniates of the salts containing oxygen. The experiments have been extended to the amines of copper and cadmium salts; in the oxygen-free salts, a relationship is observed between ammonia tension and solubility, but it is doubtful if this is more than accidental, since the solubility is so greatly affected by the nature of the medium. The amines of cobalt and chromium are more suitable, since they dissolve in water to a reasonable extent and are not decomposed thereby. The two series $[\text{Co}(\text{NH}_3)_6]\text{X}_3$ and $[\text{Cr}(\text{NH}_3)_6]\text{X}_3$ are so similar in their general properties that it would be expected to find the solubilities of the salts falling in the same order. This is actually the case to a great extent, but the parallelism is not complete. Particularly in the case of the cobalt compounds, the salts fall into two distinct groups, in which the solubilities are of widely differing orders of magnitude, the boundary lying between the perchlorate and oxalate. Possibly the two portions comprise the true salts and pseudo-salts respectively.

The following compounds have not been described previously :

hexamminecobaltic naphthalene- β -sulphonate $[\text{Co}(\text{NH}_3)_6][\text{C}_{10}\text{H}_7\text{SO}_3]_3$, pale yellow, microscopic needles; *hexamminecobaltic picrate*, $[\text{Co}(\text{NH}_3)_6][\text{O}\cdot\text{C}_6\text{H}_2(\text{NO}_2)_2]_3$, microscopic needles; *hexamminechromic dichromate*, $[\text{Cr}(\text{NH}_3)_6]_2[\text{Cr}_2\text{O}_7]_3$, lustrous, hexagonal needles; *hexamminechromic picrate*, small, matted needles; *hexamminechromic naphthalene- β -sulphonate*, pale yellow, microscopic needles; *hexamminechromic perchlorate*, pale yellow needles; *hexamminechromic chlorate*, slender, pale yellow needles. The hexamminechromi-salts resemble the alkaloids in giving *precipitates* with the following reagents: potassium mercuric iodide (slender, yellow needles), potassium bismuth iodide (microcrystalline, reddish-brown powder), sodium cobaltinitrite (very small, reddish-brown crystals). The solubilities of the following hexamminecobalti-salts (gram.-mol. per litre) in water at 9° have been estimated: chloride (0.20), bromide (0.040), iodide (0.015), nitrate (0.033), sulphate (0.015), perchlorate (0.013), chlorate (0.215), chromate (0.00055), dichromate (0.00029), oxalate (0.00069), phosphate (0.00055), naphthalene- β -sulphonate (0.00059), picrate (0.00042). The hexamminechromi-salts have the following solubilities at 17.5°: chloride (0.42), bromide (0.118), iodide (0.043), nitrate (0.072), dichromate (0.005), sulphate (0.046), phosphate (0.0032), oxalate (0.00084), picrate (0.00037), naphthalene- β -sulphonate (0.00069), chlorate (0.062), perchlorate (0.0199). H. W.

Hydrous Oxides. III. HARRY B. WEISER (*J. Physical Chem.*, 1922, 26, 401—434).—In continuation of previous work (A., 1920, ii, 760), the author has studied the properties of various hydrous chromic oxides prepared under different conditions. The results indicate that no definite hydrates of chromic oxide are formed by precipitating a chromic salt with alkali. The evidence that Guignet's green is a definite hydrate is inconclusive. Hydrous chromic oxide freshly precipitated in the cold is readily soluble in acids, but becomes insoluble on keeping or heating. Between the two extremes of solubility an indefinite number of hydrous oxides exists. By precipitating hydrous chromic oxides at temperatures from 0° to 225°, substances ranging in colour from greyish-blue to bright green have been prepared. Positively charged colloidal hydrous chromic oxide may be prepared by peptising the hydrous oxide with chromic chloride, and by hydrolysis of the chloride or nitrate, but not of the acetate. The negatively charged colloid is prepared by peptising the hydrous oxide with sodium or potassium hydroxides. The properties of the respective colloidal suspensions are discussed, more especially the precipitating and peptising action of alkali salts. J. S. G. T.

Precipitation from Uranyl Nitrate by means of Sodium Hydroxide. Radioactivity of the Precipitate. PIERRE JOLIBOIS and ROBERT BOSSUET (*Compt. rend.*, 1922, 174, 1625—1628).—When sodium hydroxide is added to a dilute aqueous solution of uranyl nitrate precipitation only commences when an equimolecular

amount of sodium hydroxide has been added. From this stage up to the addition of two molecules a precipitate is obtained, and if this is filtered off and the filtrate is boiled, a further precipitate is deposited. Either precipitate contains 1.5% or more of sodium hydroxide which is not removed by washing. If more than two molecules of alkali are added precipitation is complete and the precipitate contains still higher percentages of alkali. Radioactivity measurements on the different precipitates show that the intensity of radiation is proportional to the amount of radium present. The first precipitate which is formed contains a very important proportion of the uranium-X, the oxide of which is apparently less basic than that of uranium. W. G.

Solubility in the Solid State of Bismuth and Cadmium in Lead. CLARA DI CAPUA (*Atti R. Accad. Lincei*, 1922, [v], 31, i, 162—164).—By means of the method devised by Plato (A., 1906, ii, 521; 1907, ii, 239), the author has investigated the fusion diagrams of the systems Pb-Bi and Pb-Cd. The results obtained give for the solid solubility of lead in bismuth the value 4% and for that of bismuth the value 34%. For lead in cadmium or cadmium in lead the solid solubility is practically zero; in this system the specific thermal conductivity varies linearly with the composition. T. H. P.

Mineralogical Chemistry.

A New Occurrence of Ilsemannite. CHAS. W. COOK (*Amer. J. Sci.*, 1922, [v], 4, 50—52).—A large number of molybdenum deposits were examined, but ilsemannite was detected only at Ouray in Utah (A. 1917, ii, 491) and near Gilson in California. At the latter locality, molybdenite occurs with pyrites in aplite, and is often completely altered to yellow molybdenite. A brown tarnish suggests a first stage in the alteration to molybdenum dioxide; and a later stage to a mixture of dioxide and trioxide is suggested by the presence of a blue material which is insoluble in water. Ilsemannite, the blue soluble material, is perhaps a hydrated compound of dioxide and trioxide. L. J. S.

Chemical and Spectrochemical Study of Spanish Bismuth Minerals. S. PIÑA DE RÚBIAS and F. GILA ESTEBAN (*Anal. Fis. Quím.*, 1921, 19, 347—386; cf. Piña de Rúbias, A., 1921, ii, 267).—A description with chemical analyses and spectrographical data of bismuth minerals occurring in Spain. A series of minerals is described having a chemical composition varying between bismuthospherite, $(\text{BiO})_2\text{CO}_3$, and bismuthite, $\text{BiO}\cdot\text{CO}_3\cdot\text{Bi}(\text{OH})_2$.

Another class of minerals includes oruetite, Bi_8TeS_4 , and antimoniferous bismuthine. Oruetite and similar minerals are probably eutectic mixtures of bismuth, tetradymite, Bi_2Te_3 , and bismuthine, Bi_2S_3 . The carbonate minerals of the first series may be derived from the oruetite minerals. Native bismuth, scheelite, and molybdenite are also described.


G. W. R.]

Ardennite from Ceres, in the Ala Valley (Piedmont).

FERRUCCIO ZAMBONINI (*Atti R. Accad. Lincei*, 1922, [v], 31, i, 147—151).—This mineral occurs, intimately associated with piemontite and mixed with various other species, in aggregates of colophony-brown, vitreous fibres or rods, often twisted or bent. The crystals exhibit marked pleochroism and the other optical characters observed with the ardennite found in the Belgian Ardennes, and have the following percentage composition :

SiO_2 .	As_2O_5 .	P_2O_5 .	V_2O_5 .	Al_2O_3 .	Fe_2O_3 .	CuO .	CaO .	SrO .	BaO .
29.53	10.30	0.12	0.25	22.40	3.11	0.75	5.61	trace	trace
MnO .	NiO .	$[\text{MgO}]$.	Na_2O .	K_2O .	$\text{H}_2\text{O}-$.	$\text{H}_2\text{O}+$.			Total.
18.76	0.04	4.03	0.07	0.12	0.12	5.42			100.63

These results correspond well with the formula



The mineral is almost pure arsenioardennite, the proportion of vanadioardennite being very small.]

T. H. P.

The Melting of Potash Felspar. G. W. MOREY and N. L. BOWEN (*Amer. J. Sci*, 1922, 4, 1—21).—A pure synthetic orthoclase was prepared by crystallising glass of the composition KAISi_3O_8 in a bomb containing water vapour, and this material was used for determining the melting point of the pure substance. The temperature usually given for the melting point, determined on natural crystals, is 1200° . It was found that orthoclase has no true melting point. When kept for a week at 1200° , it forms a glass which, under the microscope, is seen to have a cross-lined structure. At higher temperatures, the structure becomes more distinct and develops into typical leucite crystals. The temperature range of incongruent melting, in which leucite crystals are in equilibrium with liquid, is $1170-1530^\circ$. Three natural potash feldspars showed the same behaviour, namely, microcline from N. Carolina, sanidine from Laacher See, and adularia from St. Gotthard, but in these the upper limit was lowered somewhat by the foreign matter present. It follows that orthoclase must be considered as a binary compound in the leucite-silica ($\text{KAISi}_2\text{O}_6-\text{SiO}_2$) system.* The general form of the diagram of this system is shown, but only a few points on it have yet been accurately determined. The incongruent melting of orthoclase is of particular importance for the theory of rock formation. It shows that leucite can form from a liquid containing an adequate amount of silica, or even excess, for orthoclase formation and that a mass

may have leucite as early crystals (phenocrysts) together with free silica as late crystals (ground-mass). The regular behaviour of leucite in breaking up into orthoclase and nephelite suggests that the early separation of leucite may afford a key to the origin of many nephelitic rocks as well as of leucite rocks. E. H. R.

Tridymite from the Euganean Hills, Italy. EDOARDO BILLOWS (*Mem. Accad. Lincei*, 1922, [v], **13**, 506—524).—A detailed crystallographic description of tridymite from the trachyte of Zovon in the Euganean Hills near Padua. The crystals are cloudy and now consist of an aggregate of quartz. L. J. S.

Analytical Chemistry.

Behaviour of Platinum on Strong Ignition. D. BALAREFF (*Chem. Ztg.*, 1922, **46**, 573).—On heating either new or carefully cleaned old platinum crucibles in a blast flame or over a powerful Teclu burner, a fairly constant loss in weight of 0.6 mg. per hour was noticed, whilst the side of the crucible nearest the blast was considerably brighter than that more remote. No experiments were made, however, to find an explanation of the loss.

A. R. P.

Heavy Liquids for the Separation of Minerals. ENRICO CLERICI (*Atti R. Accad. Lincei*, 1922, [v], **31**, i, 116—118; cf. A., 1911, ii, 257).—The following three liquids, useful for the mechanical separation of minerals of different densities, are colourless and may be diluted and recovered: Aqueous barium bromomercurate solution d^{11} 3.11, d^{18} 3.14. Saturated aqueous thallium formate solution, d^{10} 3.31, d^{20} 3.40, d^{50} 4.10. Aqueous solution of thallium formate and thallium malonate in equal proportions, d^{10} 4.00, d^{50} 4.70, $d^{\text{about } 100}$ above 5. Thallium formate, melting at 95° to a highly mobile liquid, may also be used, and a mixture of thallium formate and malonate in equal amounts melts below 95° and has d above 5. Fused thallium formate has d below 5, but it dissolves powdered thallium carbonate, giving a liquid with d above 5. T. H. P.

A New Physico-chemical Method of Volumetric Analysis applied to some Problems of Inorganic Chemistry. PAUL DUTOIT and ED. GROBET (*J. Chim. physique*, 1922, **19**, 324—327).—A method is described by which solutions of acids may be titrated

with bases using a thermometer graduated in $1/100^\circ$ as indicator. The solution to be titrated is placed in a small Dewar vessel, which stands in a somewhat larger Dewar vessel, a mechanical stirrer is placed in the solution, and the alkali is added at regular intervals in amounts which cause the temperature to increase by not more than 0.02° . The burette is surrounded by asbestos paper and the stopcock is operated by a long pair of pincers. The number of c.c. of alkali added are plotted as abscissæ and the temperature after each addition as ordinate, and the points joined. It is found that the end-point is marked by a decided change of direction in the curve. Not only is this point fixed by the curve, but all other points at which a change in the nature of the reaction occurs are fixed. Thus with sulphuric acid the points corresponding with the completion of the formation of the hydrogen sulphate and the normal sulphate are both accurately shown. In the titration of phosphoric acid with sodium hydroxide the points where the formation of NaH_2PO_4 , Na_2HPO_4 , and Na_3PO_4 , respectively, is complete are well marked. In the titration of normal sodium phosphate with nitric acid, the points where the formation of Na_2HPO_4 , NaH_2PO_4 , and H_3PO_4 is complete are clearly marked. The titration of the nitrates of zinc, lead, and magnesium by this method indicates the completion of the formation of definite basic salts, and in the case of the two first-named metals, of zincates and plumbites, respectively. Titration of salts of cobalt, copper, and nickel with ammonia in the same way indicates the formation of the various ammonia complexes. This method yields identical results with those obtained by electrometric and electroconductivity titrations and in addition it also indicates the formation of derivatives which these methods do not. J. F. S.

Simple Method of Electrometric Titration in Acidimetry and Alkalimetry. PAUL FRANCIS SHARP and F. H. MACDOUGALL (*J. Amer. Chem. Soc.*, 1922, **44**, 1193—1196).—The object of the work described is the preparation of a number of constant and reproducible electrodes which are electrometrically equivalent to hydrogen electrodes dipping in solutions of various known hydrogen-ion concentrations. Such electrodes are extremely useful in cases where it is necessary to titrate a solution of an acid to an end-point which shall have a definite hydrogen-ion concentration. In such a case, the half cell containing a hydrogen electrode and the solution being titrated is connected with the electrode which has the same potential difference as the hydrogen electrode will have when the titration is completed. A solution of alkali is then run into the acid until a galvanometer indicates that the total cell, comparison electrode|sat. KCl|titration liquid| H_2 Pt, has a zero *E.M.F.* The following are the comparison electrodes together with the potential against a normal calomel electrode and the equivalent hydrogen-ion concentration :

HgPb—	
(12—12.5%)	0.52 c.c. KI in 100 c.c. solution ; <i>E.M.F.</i> 0.5195 = 10^{4N-H}
"	2.90 c.c. KI „ 100 „ „ 0.5609 = $10^{4.7N-H}$
"	6.14 c.c. KI „ 100 „ „ 0.5786 = 10^{5N-H}
"	67.50 c.c. KI, in 100 „ „ 0.6378 = 10^{6N-H}
HgCd—	
(12—12.5%)	100 c.c. CdSO ₄ + 0.2 c.c. KI „ 0.6967 = 10^{7N-H} !
"	10 c.c. CdSO ₄ + 10 c.c. KI in 100 c.c. sol. <i>E.M.F.</i> 0.7560 = 10^{8N-H}
"	100 c.c. CdSO ₄ + 26.40 c.c. KI <i>E.M.F.</i> 0.8151 = 10^{9N-H}
"	2c.c. CdSO ₄ + 48.7c.c. KI in 100c.c. sol. <i>E.M.F.</i> 0.8743 = 10^{10N-H}

The number of c.c. of potassium iodide and cadmium sulphate refer, respectively, to 2.0*N*- and 0.5*N*-solutions, respectively (cf. Pinkhof A., 1920, ii, 121). J. F. S.

The Use of Potassium Ferrocyanide in Potentiometric Titrations. II. The Potentiometric Titration of Zinc. I. M. KOLTHOFF (*Rec. trav. chim.*, 1922, 41, 425—437).—The possibility of this titration has been established for some time (Koninck and Prost, A., 1896, ii, 675), but no worker has yet stated the degree of accuracy with which determinations may be made (cf. Bichowsky, A., 1917, ii, 219). The author recommends the use of a solution of potassium ferrocyanide, the concentration of which is 1/40 mol., to this is added 1 gram per litre of potassium ferriyanide. The standardisation of the zinc solution for the control experiments, which is described in detail, is a matter of difficulty; in previous work sufficient attention does not seem to have been directed to accuracy in this respect. The titration is best carried out at 70°, as at this temperature the reaction is quicker and the change in potential which denotes the end-point is greater. As the results obtained in neutral solution are too low, 1—2 c.c. of 4*N*-sulphuric acid should be added. Excess, however, should be avoided, otherwise the precipitate of potassium zinc ferrocyanide is partly dissolved and the change in potential at the end-point diminished. In such a solution 0.2 mg. of zinc may be determined within 1%. In presence of potassium sulphate at 70% and of ammonium sulphate at ordinary temperatures, results are obtained which are 0.5% too high; the latter salt at 70° has the reverse effect, which is also given by ammonium chloride. The titration may be carried out speedily and accurately in the inverse direction. H. J. E.

Estimation of Chlorine in Organic Compounds. J. KLIMONT (*Chem. Ztg.*, 1922, 46, 521—522).—Attempts were made to estimate chlorine in organic compounds by boiling the latter with alcoholic potassium hydroxide solution under a reflux apparatus and titrating the excess of alkali subsequently with standard acid. Acetylene dichloride, monochlorohydrin, monochloroacetone, dichloroacetone, α -chloronaphthalene, dichloronaphthalene, bornyl chloride, and pinene hydrochloride yielded either low or untrustworthy results; trichloroethane, acetylene tetrachloride, and α -dibromohydrin yielded high results, whilst monochloroacetic acid gave a correct result. W. P. S.

Estimation of Free Chlorine and Hypochlorous Acid in Concentrated Salt Solutions. M. C. TAYLOR and C. A. GAMMAL (*J. Ind. Eng. Chem.*, 1922, 14, 632—635).—The estimation of free chlorine and hypochlorous acid in concentrated salt solutions, which may contain either sodium hypochlorite or hydrochloric acid, is usually carried out by estimations of total "available" chlorine, and of free chlorine by an aeration process. The latter, however, is inaccurate owing to decomposition of hypochlorous acid or of hypochlorite during aeration. It is now shown that the rate of the removal of the free chlorine is a linear function of the amount present at any moment, whilst when all free chlorine has been removed the rate of removal varies as the square of the amount of hypochlorous acid present. The curves obtained by plotting the rate of removal of chlorine against the total amount removed as measured by absorption in potassium iodide solution and titration with thiosulphate will therefore be two distinct lines having a point of intersection, the abscissa of which is the free chlorine initially present as such. To obtain the amount of chlorine initially present as hypochlorous acid and hydrochloric acid, the increase in acidity after aeration is determined by means of potassium iodide-iodate solution in terms of thiosulphate solution, and this is equal to the actual loss of chlorine during aeration due to such reactions as $\text{HOCl} \rightarrow \text{HCl} + \text{O}$. G. F. M.

Estimation of the Chlorometric Degree of Bleaching Chlorides. J. ROYER (*Ann. Falsif.*, 1922, 15, 146—148).—For the estimation of available chlorine in bleaching solutions the method of Poncius is recommended as being rapid and exact. It consists in titrating the solution with standard potassium iodide in the presence of sodium hydrogen carbonate. The first reaction consists in the oxidation of the iodide to iodate, but as soon as the free chlorine has been used up the next drop of iodide solution reacts with the iodate with the liberation of free iodine, which is indicated by starch paste. W. G.

Electrometric Titrations with Silver Nitrate. Estimation of Chlorides, Bromides and Iodides, and of Iodides in the Presence of Chlorides and Bromides. I. M. KOLTHOFF (*Z. anal. Chem.*, 1922, 61, 229—240).—In most cases the solubility of the silver compound indicates whether the end-point of a titration with silver nitrate solution may be ascertained electrometrically. The method is trustworthy for chlorides, iodides, and bromides, even in very low concentration, and iodides may be titrated in ammoniacal solution in the presence of chlorides and bromides. Certain complex salts, such as ferrocyanides, pyrophosphates, etc., cannot be estimated by the method, although with ferricyanides and thiocyanates the results obtained are accurate. The method may also be used for the estimation of cyanides, chromates, oxalates, tartrates, succinates, and salicylates. W. P. S.

Estimation of Minute Amounts of Gaseous Oxygen and its Application to Respiratory Air. HOWARD M. SHEAFF (*J. Biol. Chem.*, 1922, **52**, 35—50).—The method is intended mainly for use in following the oxygen consumption of plant and animal tissue under different conditions. By means of the apparatus, which is described and illustrated in the original, it is stated that oxygen can be estimated in amounts as small as 1×10^{-7} gram. The method depends on the conversion of the oxygen, in the presence of nitric oxide and sodium hydroxide, into sodium nitrite, and the estimation of the latter colorimetrically by means of sulphanilic acid and α -naphthylamine. E. S.

The Oxygen-absorption and Concentration of Pyrogallol Solutions used in Gas Analysis. FRITZ HOFFMANN (*Z. angew. Chem.*, 1922, **35**, 325—328).—The absorptive powers of solutions of pyrogallol and potassium hydroxide of all proportions are tabulated in a Gibbs triangular diagram for the ternary system pyrogallol, potassium hydroxide, and water. A line of maximum absorptions was found to correspond with mixtures containing pyrogallol and potassium hydroxide in the proportions 3 to 2. The field to one side of the line, corresponding with a smaller proportion of the latter, falls sharply in power of absorption. The optimum proportion is given as pyrogallol 20 parts, potassium hydroxide 20 parts, and water 60 parts. H. M.

Rapid Estimation of Sulphur. I. LUIGI LOSANA (*Giorn. Chim. Ind. Appl.*, 1922, **4**, 204—206).—The method here described depends on the fact that, if a compound containing sulphur is heated with powdered iron in absence of air, the sulphur in the residual mass is liberated completely as hydrogen sulphide on subsequent treatment with hydrochloric acid. This gas is absorbed by zinc acetate solution and the sulphide thus formed estimated by titration with iodine solution. Tests made on flowers of sulphur and on copper, lead, barium, and cobalt sulphates gave excellent results. The method is applicable also to the estimation of sulphur in organic compounds, but the latter must previously be heated with copper oxide or lead chromate in a porcelain crucible, or, if the organic compound is volatile, in a narrow hard glass tube, in which it is covered with a deep layer of the oxidising material [cf. *J. Soc. Chem. Ind.*, 1922, 614A]. T. H. P.

The Kleemann Modification of the Kjeldhal Process. FRANTIŠEK SKUTIL (*Chem. Listy*, 1922, **16**, 173—177).—The various methods of estimating the nitrogen of foodstuffs and manures, using modifications of the Kjeldhal process, are reviewed, and the advantages of the Kleemann modification (*Z. angew. Chem.*, 1921, **34**, 625) pointed out. Results obtained by a modification of this method are found to be in satisfactory agreement with those obtained by the standard analytical method used in Czecho-Slovakia. This method differs from that of Kleemann in that, after the addition of mercury, 30% of hydrogen peroxide is added, and then the

concentrated sulphuric acid, slowly, and with cooling. After some time, the contents of the flask are heated until a deep coloration appears. Finally, anhydrous potassium sulphate is added, and the process completed. R. T.

The Estimation of the Total Non-protein Nitrogen of Serum. Comparative Study of Trichloroacetic and Metaphosphoric Acids as Protein Precipitants. PAUL CRISTOL (*Bull. Soc. Chim. Biol.*, 1922, 4, 267—271).—Higher values are obtained for total non-protein nitrogen when proteins are precipitated by means of metaphosphoric acid than when trichloroacetic acid is used for this purpose. This is due to the partial hydrolysis of the proteins by the former reagent. E. S.

The Estimation of Non-protein Nitrogen in Blood. ERIC PONDER. (*Biochem. J.*, 1922, 16, 368—369).—Blood (0.2 c.c.) is added to water (1 c.c.) and the pipette used is washed out twice, each time with a further 0.2 c.c. of water. The protein is precipitated with 0.2 c.c. of a 10% solution of sodium tungstate and 0.2 c.c. of 2/3*N*-sulphuric acid as in Folin's method for the preparation of blood filtrates. 0.5 C.c. of this filtrate is boiled very gently with 0.2 c.c. of a digestion mixture diluted 1 in 4. The digestion mixture consists of 50 c.c. of a 5% copper sulphate solution, 100 c.c. of 85% phosphoric acid, and 300 c.c. of pure sulphuric acid. After boiling for two minutes, water is added to make the volume 3.5 c.c., and the solution is directly nesslerised and compared with a standard solution of ammonium sulphate. W. O. K.

Detection and Estimation of Nitrate Nitrogen in Urine and Serum. O. NOLTE (*Z. anal. Chem.*, 1922, 61, 278—282).—The presence of nitrates may be detected by means of the diphenylamine reaction or by the less sensitive ferrous sulphate reaction; the diphenylamine reaction, however, is obtained with oxidising substances other than nitrates. For the estimation of nitrates in urine, the Schlösing-Grandeau gasometric method appears to be the most trustworthy, but the nitric oxide content of the volume of gas obtained must be estimated subsequently. W. P. S.

Analytical Determination of Oxides of Nitrogen in Gas Mixtures. CHARLES L. BURDICK (*J. Ind. Eng. Chem.*, 1922, 14, 308—310).—Mixtures of nitrogen peroxide and nitric oxide are absorbed in dilute alkali in accordance with the equations $3\text{NO}_2 + 2\text{NaOH} = 2\text{NaNO}_3 + \text{NO} + \text{H}_2\text{O}$ and $\text{NO} + \text{NO}_2 + 2\text{NaOH} = 2\text{NaNO}_2 + \text{H}_2\text{O}$. If previous to absorption there is a development of mist in consequence of cooling, the nitrogen peroxide present in the hot gas will be partly converted into nitric acid and nitric oxide. An absorption apparatus must therefore provide for the measurement of nitric oxide passing through the alkali unabsorbed. The absorption bulb in the apparatus described is a tube containing a number of glass bells with side perforations placed vertically above one another and with the inlet tube passing down through

their central axes. This is filled with a known volume of $N/10$ alkali hydroxide free from carbonate diluted as required. The gas passes through this into an aspirator, in which it is treated with hydrogen peroxide, absorbed in excess of alkali hydroxide, and estimated as usual. The solution in the absorption bulb is titrated with $N/10$ -sulphuric acid, using a few drops of a solution of methyl-red in $N/100$ -alkali hydroxide as indicator. To the neutralised solution a known excess of permanganate with 5 c.c. of sulphuric acid are added, the solution is left for a few minutes, a slight excess of ferrous sulphate added, and the excess titrated back. If there is no acid mist present and if A c.c. of alkali and P c.c. of permanganate are neutralised in the absorber, it will be seen that of this $(3A + P)/2$ c.c. of alkali hydroxide have been neutralised by nitrogen peroxide and the total alkali hydroxide— $(3A + P)/2$ c.c. by nitric oxide.

If nitric acid mist is present, it is absorbed as nitrate in the bulb and is measured by the excess of alkali hydroxide neutralised over the nitrite formed, whilst the nitrite formed contains twice the quantity of nitrogen present as nitrogen peroxide. If more than 5% of excess oxygen is present in the gas mixture, a correction for oxidation during the passage of the gas through the sampling tube should be applied.

C. I.

A Possible Source of Error in the Bell-Doisy Method for the Estimation of Phosphates in Blood Plasma. W. DENIS and L. VON MEYSENBUG (*J. Biol. Chem.*, 1922, **52**, 1—3).—The method of Bell and Doisy (A., 1920, ii, 769) for the estimation of phosphates in blood gives accurate results when applied to serum, but low values are obtained when plasma is used. Serum should therefore be used for this estimation; if, however, plasma is used, the accuracy may be increased by restricting the amount of anti-coagulant and increasing the quantities of molybdic acid and quinol.

E. S.

Microchemical Investigation of Arsenic. ARNALDO PIUTTI and ENRICO BOGGIO-LERA (*Mem. Accad. Lincei*, 1922, [v], **13**, 475—479).—In the detection of arsenic in very small proportions, good results are obtained by using a reagent composed of 1 c.c. of 3% ammonium molybdate solution, 10 c.c. of 3.4% ammonium nitrate solution, and 39 c.c. of 40% nitric acid solution. With liquids containing 0.02—0.004 mg. of arsenic per c.c., this reagent is used directly, but for lower concentrations of arsenic, the reagent is diluted five times with 40% nitric acid. A drop of the arsenic solution yields characteristic microscopic crystals when evaporated with a drop of the reagent. As little as 0.00005 mg. of arsenic is detectable in this way.

T. H. P.

Estimation of Arsenic Acid. L. ROSENTHALER (*Z. anal. Chem.*, 1922, **61**, 222—229).—Diffused light does not interfere in the iodometric estimation of arsenic acid, but atmospheric oxygen

reacts with the hydriodic acid formed, yielding further quantities of free iodine. This may be prevented by adding 5 grams of sodium hydrogen carbonate before the potassium iodide is introduced. The titration solution should contain at least 16% of hydrochloric acid or 33.3% of sulphuric acid. When sulphuric acid is used, the following procedure should be adopted. The iodate solution is treated with concentrated sulphuric acid in quantity sufficient to make the concentration of the latter 33.3%, the mixture is cooled, and 5 grams of sodium hydrogen carbonate are added in small quantities at a time; concentrated potassium iodide solution is now added, and the iodine is titrated with *N*/10-thiosulphate solution. A precipitate which forms on the addition of the iodide consists chiefly of arsenic tri-iodide, and should be dissolved by adding a small quantity of water before the titration is commenced.

W. P. S.

Iodometric Estimation of Arsenic and Antimony Sulphides. FERDINAND NIKOLAI (*Z. anal. Chem.*, 1922, 61, 257—272).—Arsenites and the corresponding antimony salts may be estimated volumetrically by adding their solutions to an excess of standard iodine solution containing dilute acetic acid and sodium acetate and then titrating the excess of iodine with thiosulphate solution. Arsenic trisulphide, or antimony trisulphide, may be titrated by dissolving it in sodium hydroxide solution and adding this solution to iodine solution containing acetic acid and sodium acetate, with subsequent titration of the excess of iodine. The alkaline solution of the sulphide is oxidised readily by atmospheric oxygen, but this may be prevented, or the rate of oxidation retarded to a considerable extent, by the addition of a small quantity of gelatin.

W. P. S.

Tests of an Iodine Pentoxide Indicator for Carbon Monoxide. S. H. KATZ and J. J. BLOOMFIELD (*J. Ind. Eng. Chem.*, 1922, 14, 304—306).—The "Hoolamite" carbon monoxide indicator consists of a glass tube containing granulated pumice impregnated with 1 part of iodine pentoxide to 5 parts of fuming sulphuric acid (60% SO_3). The gas is drawn through cotton-wool filters and a tube of activated charcoal into a rubber hand bulb and discharged through the tube containing the pumice. The colour produced on the latter is matched with a series of permanent colours in a sealed glass tube. A series of tests with a large number of observers showed that by this instrument a minimum concentration of 0.07% of carbon monoxide can be detected, and that a rough quantitative estimation of higher concentrations can be made. In the absence of the activated charcoal most hydrocarbons, hydrogen sulphide, and hydrogen chloride interfere; methane, chlorine, carbon dioxide, sulphur dioxide, and some other gases do not. The "Hoolamite" tube may be used for six to eight estimations, after which it deteriorates.

C. I.

The Direct Estimation of Small Quantities of Radium by the Penetrating Rays. B. SZILARD (*Compt. rend.*, 1922, 174, 1695—1698).—The electrometer previously described (*ibid.*, 1922, 174, 1618) has been modified so as to provide a transportable instrument working without either a high tension battery or a projection mirror. By means of it rapid measurements can be made of the penetrating rays. The sensitiveness of the instrument corresponds with 10^{-4} U.E.S.

W. G.

Ash-alkalinity (of Foodstuffs). B. PFYL. (*Z. Unters. Nahr. Genussm.*, 1922, 43, 313—339).—To utilise the titration values of the ash of foodstuffs as indications of their composition, it is necessary to choose the values, to be determined in such a way that they are definite and as independent as possible of chance circumstances, and to employ simple and unexceptionable methods for determining them. The chemical processes taking place during the incineration of foodstuffs have to be considered more closely than has been done in the past, and the necessity for avoiding loss of mineral acids and for the conversion of the whole of the phosphorus into tribasic phosphate must be kept in view. In many cases this can only be attained by the addition of a measured quantity of alkali before incineration. The most useful values to be determined are (a) the intrinsic alkalinity, (b) the methyl-orange alkalinity, and (c) the total phosphate of the ash. The first may be defined as the excess, expressed in milli-equivalents, of the kations Na^+ , K^+ , Ca^{++} , Mg^{++} , which remains after combination with the anions PO_4''' , SO_4'' , Cl' for O'' , and the weakly acid anions CO_3'' , SiO_3'' , BO_2 , MnO_3'' , MnO_4'' , AlO_2'' . If such excess does not exist, the difference is expressed as "intrinsic acidity." (b) is the excess, expressed in milli-equivalents, of the same four kations which remains after combination with the anions $\text{H}_2\text{PO}_4'$, SO_4'' , and Cl' for O'' , and the above enumerated weakly acid anions. The absence of such excess is indicated as methyl-orange acidity. (c) The total phosphates is expressed in milli-equivalents of PO_4''' . These three values can be obtained by titration against methyl-orange, followed by a titration against phenolphthalein. Full details of simple methods of carrying out the above determinations are given; these are the result of many years of experience, and are founded on sound scientific principles. Modifications necessary in special cases and precautions and corrections to be applied where special accuracy is needed are also indicated. The above three values are usually calculated to 100 grams of dry food-stuff or 1 litre of liquid, but occasionally to 1 gram of ash or 1 gram of ash soluble in hydrochloric acid. In cases of adulteration, one or other of these values is affected. Methods proposed by others for the determination and presentation of ash-alkalinity are criticised.

H. C. R.

Atkinson's Process for the Estimation of Potassium in the Presence of Sodium, Magnesium, Sulphates, and Phosphates. S. J. WATSON (*Analyst*, 1922, 47, 285—288).—The

process is considered unsatisfactory because the compensating errors necessary to overcome the loss due to the solution of some potassium perchlorate do not occur to an extent sufficient to balance this loss, and the quantity of methyl alcohol and perchloric acid used is too large to make the process economical. Keeping the contents of the beaker at the boiling point for an hour, whilst perhaps necessary to ensure complete solution of undesirable substances, must also favour the solution of potassium perchlorate itself.

H. C. R.

The Estimation of calcium in Blood. ARTHUR ROBERT LING and JOHN HERBERT BUSHILL (*Biochem. J.*, 1922, 16, 403—406).—Two to 5 c.c. of blood are incinerated in a platinum dish, treated with 1.5 c.c. of concentrated hydrochloric acid, and washed into a special centrifuge tube. After adjusting the reaction, the calcium is precipitated with oxalic acid and ammonium oxalate. The calcium oxalate is centrifuged and washed, sulphuric acid is added, and the oxalic acid titrated with permanganate.

W. O. K.

A Colorimetric Method for the Estimation of Small Amounts of Magnesium. F. S. HAMMETT and E. T. ADAMS (*J. Biol. Chem.*, 1922, 52, 211—215).—The method is designed for application to urine, blood, and tissue extracts and is a modification of Kramer and Tisdall's method (*A.*, 1921, ii, 595). The precipitate of ammonium magnesium phosphate obtained in the latter method is dissolved in 0.01*N*-hydrochloric acid and the phosphorus estimated colorimetrically by Bell and Doisy's method (*A.*, 1920, ii, 769). The value for magnesium is then calculated from this result.

E. S.

Volumetric Estimation of Lead Peroxide in Miniums. Practical Modification of Diehl's Method. A. BONIS (*Ann. Falsif.*, 1922, 15, 157—159).—The following modified procedure for Diehl's method of estimating lead peroxide in samples of minium is advocated. Half a gram of the sample is macerated with 2.5 c.c. of nitric acid (*d* 1.080) and the mixture is washed into a conical flask with 25 c.c. of a saturated solution of sodium acetate. To this is added 10 c.c. of a 12% solution of potassium iodide in saturated sodium acetate, and the whole is well shaken, and the iodine liberated is titrated with *N*/10-thiosulphate solution. If the sample of minium contains iron oxide, instead of titrating the free iodine an excess of standard thiosulphate is added and the whole is made up to 100 c.c. with the saturated sodium acetate solution. After filtering, an aliquot portion of the filtrate is titrated back with *N*/10-iodine solution.

W. G.

Rapid Electro-analysis. A. KLING and A. LASSIEUR (*Ann. Chim. Analyt.*, 1922, 4, 171—177).—For the rapid electrolytic deposition of copper or zinc a silver cathode is recommended,

together with an anode of platinum alloyed with iridium or rhodium. The silver cathode is readily cleaned without loss in weight by immersion in a cold solution of 10 grams of trichloroacetic acid in 50 c.c. of strong ammonia and 50 c.c. of water. Copper is deposited from a sulphate solution acidified with nitric acid and containing 5 grams of sodium metaphosphate if iron is present. Zinc is deposited from a solution containing sodium acetate and 5 c.c. of acetic acid per 100 c.c., together with 10 c.c. of a saturated solution of sodium fluoride if iron is present; large quantities of iron should, however, be separated. Lead may be deposited on the anode as peroxide from a solution containing 15 c.c. of nitric acid per 100 c.c., or as metal on the cathode by electrolysis at 70° with 5 amperes a solution containing less than 0.4 gram of lead and 3 c.c. of nitric acid (*d* 1.3), 2.5 grams of crystallised gallic acid, and 5 c.c. of 95% alcohol in 100 c.c. total bulk. Antimony may be electrolysed from a solution containing 80 c.c. of sodium sulphide solution (*d* 1.14), 60 c.c. of water, and 5 grams of potassium cyanide by a current of 5 amperes at 60–70°, or from a boiling solution containing 20 c.c. of hydrochloric acid and 4 grams of hydroxylamine hydrochloride in 130 c.c. of water. The latter solution, with the addition of 10 grams of ammonium oxalate, may be used for the deposition of tin. The above methods are adapted to the rapid analysis of brasses, bronzes, and white metals [cf. *J. Soc. Chem. Ind.*, 1922, 551A]. A. R. P.

Iodometric Estimation of Copper in the Presence of Iron.

A. WÖBER (*Z. angew. Chem.*, 1922, **35**, 336–337).—Solutions containing copper salts in the presence of ferrous iron always contain a small amount of cuprous ions. These cannot be oxidised by nitric acid or hydrogen peroxide in ammoniacal solution, as the small amount of nitrites formed interferes with the subsequent titration. The solution is therefore made ammoniacal and treated with a current of air at 70°, whereby all the copper dissolves as cupric salt and the iron is precipitated as ferric hydroxide, which is collected, re-dissolved and re-precipitated to recover the adsorbed copper. The combined filtrate is boiled to expel ammonia, acidified, treated with potassium iodide, and titrated as usual [cf. *J. Soc. Chem. Ind.*, 1922, 545A]. A. R. P.

Separation of Aluminium from Iron by means of o-Phenetidine. K. CHALUPNY and K. BREISCH (*Z. angew. Chem.*, 1922, **35**, 233–234).—The solution containing the iron and aluminium salts is slightly acidified with hydrochloric acid and treated with a current of hydrogen sulphide for ten minutes; carbon dioxide is then passed through the solution until all excess of hydrogen sulphide has been expelled. These operations are carried out in a flask closed with a cork and provided with inlet and outlet tubes. Ammonium carbonate is then added until a precipitate forms, the latter is dissolved by the addition of dilute hydrochloric acid, and an excess of 5% alcoholic phenetidine

solution is added. The flask is closed, the contents are heated at 80° , and the precipitate is then collected, washed with hot, dilute ammonium nitrate solution, ignited, and weighed as aluminum oxide. If copper and other metals are present with the iron and aluminium, the copper must be separated electrolytically, the iron and aluminium then precipitated as basic acetates, the latter dissolved in dilute hydrochloric acid, and the solution used for the separation of the iron and aluminium as described. It is essential that the iron should be present in the ferrous state; ferric salts yield a precipitate with phenetidine solution. W. P. S.

Gravimetric Analysis. XXV. Estimation of Manganese. L. W. WINKLER (*Z. angew. Chem.*, 1922, **35**, 234—235; cf. A., 1921, ii, 656).—The manganese is precipitated as manganese ammonium phosphate, the method being that described previously (*loc. cit.*) for the estimation of cadmium. Small corrections are applied to the weight of precipitate obtained; for instance, 1 mg. is deducted when the precipitate weighs 0.2 to 0.3 gram. The presence of potassium chloride does not interfere, but sodium chloride must not be present. W. P. S.

Oxidation of Manganese to Permanganate in Alkaline Solution. J. HESLINGA (*Chem. Weekblad*, 1922, **19**, 274).—Manganese in ores and alloys is readily detected by heating the solution with potassium hydroxide and excess of copper sulphate (which acts as a catalyst), and adding a few drops of a mixture of bromine water and potassium hydroxide; the violet coloration due to permanganate develops at once. 0.005 Mg. of manganese may be detected. The reaction is not suitable for quantitative estimation, since it is only complete where very small quantities of manganese are present; also the presence of iron affects the colour, probably by formation of potassium ferrate.

The catalytic action of the copper oxide is probably due to its tendency to form easily dissociated oxygen compounds, and its tendency to form compounds with manganese hydroxide. More copper than manganese must be present, otherwise the catalyst remains in this combination. S. I. L.

The Adsorption of Iron by Precipitates of Manganese Dioxide. MAX GELOSO (*Compt. rend.*, 1922, **174**, 1629—1631; cf. Nicolardot, Geloso and Réglade, A., 1920, ii, 334).—When manganese is precipitated as its dioxide by the addition of ammonium persulphate to an acid solution of manganese sulphate containing some iron salt, the amount of iron adsorbed by the precipitate varies directly with the amounts of iron and manganese present in the solution and inversely with the acidity. The presence of ammonium sulphate has no effect. The iron adsorbed is mostly in the form of hydroxide. W. G.

Estimation of Chromium in Steels. LUIGI LOSANA and ENRICO CAROZZI (*Giorn. Chim. Ind. Appl.*, 1922, **4**, 197—200).—Gravimetric methods for estimating chromium in steel give good

results but are too tedious for industrial use, and colorimetric methods are not to be recommended. Provided that attention is directed to various points, satisfactory results are obtained by Stead's method, in which the chromium is oxidised by means of nitric acid and by permanganate to chromic acid and this is treated with ferrous sulphate, the excess of the latter being titrated with permanganate. Modifications are suggested which render the ammonium persulphate method more rapid and exact [cf. *J. Soc. Chem. Ind.*, 1922, 594A].

T. H. P.

New Qualitative Test for Uranium. HAROLD D. BUELL (*J. Ind. Eng. Chem.*, 1922, 14, 593).—Uranium may be detected in slags or ores by preparing a nitric acid solution of the material, too great an excess of acid being avoided, and adding an excess of granulated zinc. When the reaction with the acid has subsided, a yellow deposit will appear on the zinc if uranium is present. Gold, platinum, thorium, lead, tungsten, titanium, chromium, mercury, and copper do not interfere with the test. Iron and vanadium only interfere if present in large quantities, and in that case the spent liquid is removed, and the zinc and the deposit are again treated with nitric acid. The deposit dissolves, but reappears when the acid is again exhausted, and vanadium and iron remain in solution. As regards the delicacy of the test, in a solution of pure uranyl nitrate it was possible to detect 0.88 mg. of uranium per c.c. The test is not applicable in presence of sulphuric or hydrochloric acids. The yellow deposit is apparently the hydrated trioxide, $\text{UO}_3 \cdot 2\text{H}_2\text{O}$.

G. F. M.

Colorimetric Estimation of Vanadium in Steel. A. KROFF (*Z. angew. Chem.*, 1922, 35, 366—367).—The steel is dissolved in a mixture of nine parts of sulphuric and one part of phosphoric acid to prevent separation of tungstic acid. The solution is oxidised first with nitric acid, then with ammonium persulphate to destroy carbonaceous matter, and the colour produced by hydrogen peroxide in this solution is matched with that produced by a known amount of vanadium pentoxide added to a solution containing approximately equal amounts of chromium, nickel, and hydrogen peroxide [cf. *J. Soc. Chem. Ind.*, 1922, 594A].

A. R. P.

Rapid Detection of Bismuth in Urine and in Saliva. DOMENICO GANASSINI (*Boll. Chim. Farm.*, 1922, 61, 321—326).—For the detection in urine of small proportions of bismuth, which is administered in the form of sodium or potassium bismuthotartarate (cf. Sazerac and Levaditi, this vol., i, 89) in cases of syphilis, good results are obtained by means of stannous chloride solution, prepared by dissolving 10 grams of the crystallised salt in 10 c.c. of concentrated hydrochloric acid and heating the liquid with a few pieces of tin on a water-bath until it becomes almost clear (cf. Vanino and Treubert, A., 1898, ii, 461). Ten c.c. of the urine are treated with 2 c.c. of this reagent and then with 5 c.c.

of 20% sodium hydroxide solution, and the whole is shaken. If the urine contains salts of bismuth, these undergo reduction to the metal, which is deposited with the precipitated calcium and magnesium phosphates and gradually colours these brown or black; in some cases, also, the supernatant liquid becomes brown owing to dissolution of part of the bismuth in the colloidal condition. Saliva may be tested similarly. That the dark precipitate actually contains bismuth may be shown by conversion of a little of it into the vivid red bismuth rubidium iodide, $\text{BiI}_3 \cdot 2\text{RbI} \cdot 2 \cdot 5\text{H}_2\text{O}$, and microscopic examination of this.

Nylander's test for dextrose (A., 1884, 1433) may be used inversely as a test for bismuth, but is not so sensitive as that given above.

T. H. P.

Estimation of Minute Quantities of Methane. ERNST MURMANN (*Oesterr. Chem.-Ztg.*, 1922, 25, 90).—The estimation of small quantities of methane, involving the absorption of the carbon dioxide derived therefrom in baryta solution, has hitherto required two or more absorption flasks to ensure the complete removal of carbon dioxide from the gases owing to the short time in which the bubbles are in contact with the liquid. One ordinary Erlenmeyer flask only is sufficient, however, if the gases are led into the baryta solution through a capillary tube and 0.5—1.0 c.c. of 1% gelatin solution is added to the liquid so as to make a foam on the surface which increases the duration of contact of the gases and the absorbent solution to at least thirty seconds, and renders absorption thereby complete. The capillary may conveniently be fixed into the flask through the straight arm of a T-piece fitted into a rubber stopper, and the connexion made gas-tight with a piece of rubber tube. The artifice may also be useful for other cases where complete absorption of a gas by a liquid is required.

G. F. M.

The Estimation of Benzene in Gases. E. BERL (*Z. angew. Chem.*, 1922, 35, 332).—The benzene is absorbed by carbon, which is then heated at 100—120° in a current of steam (in a salt-bath). A small amount of benzene remains behind, and therefore a test is made with carbon containing benzene to control the results. The burette filled with water in which the benzene is collected is provided with an overflow in such a manner that the benzene remains in the burette. This method is simpler and better than that by which the benzene is washed out of the gases with light petroleum.

H. M.

Some New Colour Reactions of Cholesterol. LOUIS KAHLBERG (*J. Biol. Chem.*, 1922, 52, 217—225).—Cholesterol dissolves in arsenic trichloride to a pink solution changing to cherry-red; isocholesterol yields a cobalt blue solution changing through violet, purple, and dark red to dark green; phytosterol, however, dissolves to a colourless solution. The colours are discharged by the addition of solvents such as benzene, toluene, and

chloroform. The colour reactions are also obtained with hot solutions of arsenious oxide in concentrated hydrochloric acid, but are not so permanent as with the anhydrous chloride. Coloured solutions, which are not, however, sufficiently characteristic to be used for distinguishing between the sterols, are also obtained with certain other acid chlorides. E. S.

Analytical Observations on the True Blood-sugar Value in Normal and Pathological Individuals. WILHELM STEPP (*Arch. exp. Path. Pharm.*, 1922, 90, 105—128).—A series of analyses of blood-sugar was carried out in the following way: the proteins were removed with phosphotungstic acid and the sugar estimated by a reduction method (that of Bertrand or Laquenne); a further amount of the filtrate was freed from phosphotungstic acid with lead acetate, the excess of lead removed, and the solution concentrated; the concentrated fluid was analysed for sugar by a reduction method, by fermentation, and polarimetrically.

The values obtained by the reduction methods on the untreated filtrate were 20% to 100% higher than those obtained on the filtrate after concentration; the difference was greatest in cases where there was nitrogen retention, and was proved to be not due to loss of sugar in the process of analysis.

The agreement between the results obtained by the three methods on the concentrated filtrate was, on the whole, good, the polarimetric and fermentation methods giving especially close agreement. It is suggested that nitrogenous reducing substances are carried down in the process of removing the excess of phosphotungstic acid, and that the lower values obtained after this treatment represent the true concentration of sugar in the blood.

C. R. H.

The Inversion of Sucrose in the Alkaline Copper Solution. E. CANALS (*Bull. Soc. chim.*, 1922, [iv], 31, 583—588).—Results indicate that sucrose is slowly hydrolysed during the estimation of reducing sugars in its presence by means of some form of Fehling's solution. The actual amount of sucrose hydrolysed increases with the concentration of the sucrose, with the time of heating, and with the temperature at which the liquid is kept. It is essential therefore to make a blank estimation, and it is advisable to heat by immersion in a water-bath. W. G.

Estimation of Humus by Oxidation with Chromic Acid. A. GEHRING (*Z. anal. Chem.*, 1922, 61, 273—278).—The soil is first treated with dilute sulphuric acid in a flask through which a current of air free from carbon dioxide is passed. When all carbon dioxide produced by the decomposition of carbonates has been expelled, potassium dichromate is added to the flask and the mixture is heated; the carbon dioxide and other gases resulting from the oxidation of the organic matter are passed through a combustion tube containing copper oxide and lead chromate, and the carbon dioxide is then absorbed in potash bulbs. W. P. S.

Oxidation with Mixtures of Sulphuric Acid and Chromates. L. J. SIMON (*Compt. rend.*, 1922, 174, 1706—1708).—For the wet combustion of organic compounds by means of chromic acid, the most satisfactory results are obtained by the use of concentrated sulphuric acid and silver chromate, using about 15 c.c. of the acid and 12 grams of the chromate for about 0.1 gram of substance. This oxidising mixture may be used for the combustion of acetates or acetyl derivatives, theoretical results being obtained.
W. G.

Analysis of Acetic Anhydride. C. EDWARD SAGE (*Perf. Essent. Oil Rec.*, 1922, 13, 172).—In the method described by Reclaire (this vol., ii, 532) for the estimation of acetic anhydride, the phenolphthalein used as indicator seems with certain samples to be peculiarly insensitive. The actual cause of this is obscure, but the peculiarity seems to run parallel with the development of the iodoform reaction when the sample in question is tested with sodium hydroxide and iodine. The unknown impurity giving this reaction may be traced back to the original acetic acid from which the anhydride was prepared, and it seems to be associated with faults in odour which sometimes develop in products prepared from the anhydride, for example, aspirin, or synthetic "acetyl" perfumes. The insensitiveness of phenolphthalein above mentioned may be overcome, and a sharp end-point obtained if the titration is made by adding an excess of $N/2$ -hydroxide, keeping for some time without heating until hydrolysis is complete, then adding a known quantity of $N/2$ -acid and finishing off the titration with more $N/2$ -hydroxide. By this procedure errors due to heating under reflux will also be eliminated.
G. F. M.

Estimation of Fatty Acids (and Cholesterol) in Small Amounts of Blood Plasma. W. R. BLOOR, K. F. PELKAN, and D. M. ALLEN (*J. Biol. Chem.*, 1922, 52, 191—205).—Bloor's method (*A.*, 1914, ii, 392) for the estimation of fatty acids in blood probably gives incorrect results owing to the different nephelometric values which the various fatty acids and cholesterol possess (cf. Csonka, *A.*, 1918, ii, 277). To overcome this two modifications are suggested: (1) the separation of cholesterol from the fatty acids and the separate estimation of the two fractions thus obtained (cf. *A.*, 1916, ii, 275), (2) the use of a standard, consisting of 60% oleic and 40% palmitic acid, which melts at approximately the same temperature as the mixture obtained from blood. Using these modifications, the results are accurate to within 5%.
E. S.

Estimation of Small Amounts of Lactic Acid. S. W. CLAUSEN (*J. Biol. Chem.*, 1922, 52, 263—280).—The method of Fürth and Charnass is slightly modified. Lactic acid is decomposed either by acid permanganate at 95° or by 50% sulphuric acid at 140°, the aldehyde produced being swept into excess of sodium hydrogen sulphite. The excess of the sulphite is removed by addition of iodine, the end-point being adjusted to a definite blue

to starch. Saturated sodium hydrogen carbonate is then added until the blue coloration is just discharged, and the sulphite of the aldehyde bisulphite compound titrated with standard iodine, the end-point being determined by a blank control. The errors which arise when the method is applied to blood or urine may be reduced, but not entirely eliminated, by first extracting the lactic acid from the protein-free filtrate by ether, using a special extraction apparatus. E. S.

Detection and Estimation of Oxalic Acid and its Use in Standardising Iodine and Silver Solutions. L. ROSENTHALER, (*Z. anal. Chem.*, 1922, 61, 219—222).—Iodine is liberated when potassium iodate solution is heated with the addition of a small quantity of oxalic acid; the reaction may be obtained with 3mg. of oxalic acid and permits of the detection of the latter in the presence of tartaric and malic acids which reduce the iodate much more slowly. Oxalic acid and oxalates may be estimated by heating their dilute sulphuric acid solution with an excess of standard iodate solution until all free iodine has been expelled, cooling the mixture, and titrating the excess of iodate with thiosulphate solution after the addition of potassium iodide. It is proposed to use sodium oxalate for the standardisation of iodine solutions and of silver nitrate solutions; in the latter case, a known quantity of the oxalate is treated with a slight excess of silver nitrate solution, the mixture diluted to a definite volume, filtered, and the excess of silver titrated in an aliquot portion of the filtrate with thio-cyanate solution. W. P. S.

Oxidation of Oxalic Acid in the Absence of other Acids. J. C. WITT. (*J. Physical Chem.*, 1922, 26, 435—446).—It is shown that oxalic acid may be titrated with potassium permanganate without the addition of any other acid, the oxalic acid functioning both as reducing agent and acid. The end-point is marked by the appearance of turbidity followed by a slight permanent precipitate and is affected by the temperature of titration, the concentration, and the presence of electrolytes. When a solution of an oxalate is titrated in the presence of sulphuric acid, the colloid precipitating power of the sulphate-ion is effective in the presence of a fairly high concentration of the hydrogen-ion, and hence the quantity of sulphuric acid required is much greater than the theoretical value. J. S. G. T.

Standard Method for the Estimation of Soap in Wool. BRITISH RESEARCH ASSOCIATION FOR THE WOOLLEN AND WORSTED INDUSTRIES (*Trans. Text. Inst.*, 1922, 13, 143—149).—Traces of soap left on a woollen fabric may cause serious disturbances in dyeing, and consequently may require estimation. It is shown that extraction at the boiling point with neutral absolute alcohol, in a glass apparatus with ground-in joints, satisfactorily removes the soaps and free fatty acids from the material. J. C. W.

Olive Oils and the Villavecchia Reaction. JEAN PRAX (*Ann. Falsif.*, 1922, **15**, 159—161).—It has been shown previously (*ibid.*, 1921, **14**, 270) that certain olive oils, particularly Tunisian, give a red coloration with Villavecchia's reagent, similar to that given by an olive oil adulterated with sesamé oil. This abnormal reaction is not obtained if such an oil is shaken with its own volume of 90% alcohol containing 10% of ammonia and the alcohol and ammonia then evaporated on a water-bath before applying the test. It is now shown that it is as a result of the action of the ammonia that these abnormal oils no longer give the red coloration with the Villavecchia reagent. W. G.

The Orcinol Reaction with Furfuraldehyde. Application to the Colorimetric Estimation of Small Quantities of Furfuraldehyde. PAUL FLEURY and GABRIEL POIROT (*Bull. Soc. Chim. Biol.*, 1922, **4**, 252—266).—For the estimation, 1 c.c. of the furfuraldehyde solution and 5 c.c. of a hydrochloric acid reagent (*d.* 1.19 and containing about 60 mg. of ferric chloride per litre) are added successively to 4 c.c. of an orcinol reagent (1 gram of orcinol in 1600 c.c. of glacial acetic acid). The mixture is then heated for one minute on a water-bath, left for half an hour at the ordinary temperature, and the blue coloration compared with that produced in a standard solution treated simultaneously in the same manner. The standard solution contains 0.1 mg. of furfuraldehyde per c.c.; that to be estimated should contain between 0.01 and 0.30 mg. E. S.

A Cause of Error in the Application of the Colour Test for Acetone. ANTONIO TROISE (*Ann. Chim. Analyt.*, 1922, **4**, 177—178).—On testing a sample of urine for acetone by means of ammonia and sodium nitroprusside in acetic acid (Lieben's reaction) the characteristic violet colour was obtained, although the distillate did not give the reaction. The cause of the colour was found to be the presence of istyline (1.8 dihydroxyanthraquinone) which had been taken medicinally. A. R. P.

Differentiation and Estimation of Saponins. L. KOFLER (*Z. Unters. Nahr. Genussm.*, 1922, **43**, 278—287).—The usual methods of investigation do not suffice to arrive at the identity of a saponin when it is not available in the pure state. The author makes use of the foaming power in addition to the hæmolytic action of saponins. The "foam number" is obtained by shaking up 10 c.c. of each of a series of solutions of different concentrations in test-tubes 16 mm. internal width for fifteen seconds and allowing them to remain for fifteen minutes. The foam number is given by the dilution in that tube in which the foam stands 1 cm. high. The hæmolytic index is determined in the usual way, the details of which are specified. If the hæmolytic index is divided by the foam number, a quotient is obtained which is independent of the state of purity of the saponin. This quotient has a characteristic value for each of the six different saponins investigated, varying

from zero in the case of glycyrrhizin to 10 in the case of Merck's digitonin. By this means saponins separated from artificial lemonades by Brunner's method (*ibid.*, 1902, 5, 1197; 1908, 16, 165; 1912, 23, 586; 1914, 27, 192) can be readily identified and quantitatively estimated by comparing the hæmolytic index and foam value obtained with those tabulated for the saponin in question. It would be possible to specify a maximum value for the "poison/foam" quotient which should not be exceeded by the saponins used in food-stuffs. The figure 1.0 or 0.5 is suggested.

H. C. R.

The Defecation of Blood for the Estimation of Carbamide.

AL. IONESCU (*Bul. Soc. Chim. Romania*, 1922, 4, 13—17).—The defecating agent recommended consists of a saturated aqueous solution of sodium chloride to which has been added 10% of glacial acetic acid. To the serum an equal volume of this reagent is added and the mixture is heated to boiling and filtered. The filtrate is made slightly alkaline with sodium hydroxide and the carbamide estimated in an aliquot portion by the hypobromite method.

W. G.

The Testing of Foodstuffs for Vitamins. J. C. DRUMMOND and A. F. WATSON (*Analyst*, 1922, 47, 235—246).—Details are given of the method of carrying out physiological tests on rats for the presence of the three vitamins in foodstuffs. Young healthy rats of not more than 50 grams body weight are fed on a ration of purified foodstuffs from which all traces of the vitamin to be tested for have been removed. These rations are specified for each of the three vitamins. When the rats have shown no further increment of weight for fourteen days, the substance to be tested is administered in a daily ration of known weight. The method has been used to show that samples of butter vary very considerably in the amount of vitamin they contain.

The amount of vitamin-B in milk depends entirely on the food of the cow. The monkey is the best animal for testing for vitamin-C, but the guinea pig is generally used. Testing is rendered more difficult by the fact that the guinea pig is entirely herbivorous. The potency of the food supplement added is judged from the daily dose necessary to prevent the onset of, or to cure established, scurvy. The antiscorbutic value of milk is dependent on the diet of the animals. Lemon juice possesses a very much higher antiscorbutic potency than lime juice.

H. C. R.

The Necessity of Checking the Quality of Sodium Tungstate used in the System of Blood Analysis. OTTO FOLIN (*J. Biol. Chem.*, 1922, 51, 419—420).—Specimens of sodium tungstate which are not alkaline to phenolphthalein contain complex tungstates. They may be made suitable for use in the author's system of blood analysis by addition of the requisite quantity of alkali.

E. S.

General and Physical Chemistry.

The Luminescence of Incandescent Solids. E. L. NICHOLS and H. L. HOWES (*Physical Rev.*, 1922, **19**, 300—318).—Certain oxides, when heated to temperatures lying within a definite and sometimes narrow range, emit radiation in a limited region of the spectrum far in excess of the radiation emitted in that region by a black body at the same temperature. The effect is ascribed to luminescence, and is particularly marked in the case of the blue radiation from columbium oxide; the blue glow is also shown by the oxides of glucinum, magnesium, calcium, aluminium, silicon, and zirconium. Intensity-temperature curves for the red, green, and blue radiations from the oxides of samarium, gadolinium, gallium, columbium, erbium, cerium, praseodymium, and neodymium show that all except cerium emit a blue or greenish-blue glow, and all but columbium and neodymium have one or two red, green, and blue outbursts above 1000°. Luminescence is essentially different from temperature radiation, and the effect for any oxide is sensitive to slight impurities. It is considered that luminescence is probably a phenomenon of instability associated with a chemical change such as oxidation, or with the physical disturbances which the oxides undergo as they are heated.

A. A. E.

Optics of Disperse Systems. V. I. LIFSCHITZ and GEORG BECK (*Kolloid Z.*, 1922, **31**, 13—15; cf. A., 1920, ii, 137).—A number of misprints and errors of calculation contained in a previous paper (*loc. cit.*) are corrected. Tables of the recalculated values are given and a repetition of the measurements of the refractive index of arsenic trisulphide sol has brought the authors' measurements into agreement with those of Wintgen. The discrepancy previously found was due to a film of the arsenic sulphide separating on the prism. It is also shown that the influence of the degree of dispersion on the refraction is very small in comparison with the influence of constitution.

J. F. S.

The Mass of the Particles that Emit the Spectrum of Carbon Monoxide. M. DUFFIEUX (*Compt. rend.*, 1922, **175**, 159—161).—On the supposition that the widening of the spectral lines emitted by a luminous gas is solely due to thermal vibration, the limit of the interference fringes which can be obtained with these lines is given by the equation $N = K\sqrt{M/T}$, where K is a constant, T the absolute temperature, and M the mass of the luminous particles on the atomic weight scale. This method was applied to an investigation of the origin of the spectra of nitrogen and carbon monoxide. The values found for N for the nitrogen lines, 3998 and 4200, were 270,000 and 274,000, respectively, whilst very concordant results averaging 313,000 were found for four of

the carbon monoxide bands, and 260,000 for the fifth band, 4123. Adopting on the one hand Lord Rayleigh's value for K , namely, 1.37, as an upper limit, and, alternatively, Fabry and Buisson's, 1.22, as the lower limit, M for nitrogen becomes 11.6 or 14.5, and for carbon monoxide 15.3 or 18.3 for the four bands, and 10.6 or 13.2 for the fifth band. The conclusion is drawn that the two bands of nitrogen are emitted by atomic nitrogen (14), and that the four bands 5610, 5198, 4833, and 4509 are due to atomic oxygen (16), whilst the fifth band is due to carbon atoms (12), the masses of all these atoms falling well within the upper and lower limiting values obtained for M .
G. F. M.

The Excitation of the Enhanced Spectra of Sodium and Potassium in a Low Voltage Arc. PAUL D. FOOTE, W. F. MEGGERS, and F. L. MOHLER (*Astrophys. J.*, 1922, 55, 145—161; cf. this vol., ii, 4).—It is shown that the quantum theory requires that the enhanced spectra of sodium and potassium should be excited at comparatively low voltages. The experimental results demonstrate that this is the case, and clearly show the three-stage development in the spectra of these elements, namely, (1) the single pair stage, 2.1 to 5.1 volts for sodium; 1.6 to 4.3 volts for potassium; (2) the arc spectrum stage, 5.1 to 30 volts for sodium; 4.3 to 20 volts for potassium; (3) the enhanced spectrum stage, above 30 volts for sodium; above 20 volts for potassium.

A. A. E.

The Difference between Series Spectra of Isotopes. P. EHRENFEST (*Nature*, 1922, 109, 745—746; cf. following abstract).—A criticism of the use of Bohr's formula for the change in frequency, ν , due to the motion of the nucleus, in the case of atoms in which several electrons move round the nucleus. The formula $\nu_2 : \nu_1 = M_2/M_2 + m : M_1/M_1 + m$, where M_1 , M_2 , and m are, respectively, the masses of the nuclei of the isotopes and of the electron, and ν_1 and ν_2 the frequencies of the corresponding lines, cannot be true in general in such cases, although it will probably be possible to derive a sufficiently approximate formula for the case of the p - and d -motions of lithium; this must be very difficult, however, for the case of the $1.5S$ -path.

A. A. E.

The Difference between Series Spectra of Isotopes. N. BOHR (*Nature*, 1922, 109, 746; cf. preceding abstract).—It is admitted that the effect of the mass of the nucleus on the spectrum of an atom containing more than one electron is a complex problem; not only may the mass effect disappear completely in such cases, but also, in the case of the motions actually met with in the emission of the series spectra, this effect may be different from that calculated for an atom with one electron. It is shown that the problem differs essentially from the problem of two bodies in celestial mechanics. The discrepancies, observed by Merton, in the wave-lengths of certain lines in the spectra of lead isotopes are considered with reference to the mass effect, and to the possibility of their being due to a slight difference in the field of force

surrounding the nucleus, arising from the difference in the internal nuclear structure of the lead isotopes. A. A. E.

The Difference between Series Spectra of Isotopes. J. W. NICHOLSON (*Nature*, 1922, 110, 37; cf. preceding abstracts).—The differences found by McLennan in the spectra of the lithium isotopes are three times as great as those calculated by Bohr's formula. This large separation cannot be explained by the quantum theory, and it is suggested that the new series may be a combination series or a spark series. In an investigation on some of the simpler possible orbits in a lithium atom with only two electrons, a specially simple class of orbits was found. The principal spark-line of lithium has a value very close to λ 6708, the red line shown in the ordinary spectrum. The results appear to indicate that when there are many electrons in an atom, a ratio roughly of order 1/10 exists between the orbital radii of the two outer consecutive electrons. An immediate consequence is that the Bohr formula would never be very far wrong in its use for a rough determination of the separation to be looked for in the spectra of isotopes.

A. A. E.

The Vacuum-spark Spectra of the Metals. EDNA CARTER (*Astrophys. J.*, 1922, 55, 162—164).—Vacuum-spark spectra of calcium, magnesium, cadmium, titanium, and iron were obtained, the iron spectrum being also photographed on a dyed film from λ 4000 to λ 6600 Å.; the new spectra show no striking new characteristics. The vacuum-spark spectrum of calcium and magnesium is practically identical with the spark spectrum in air; with cadmium, and particularly with titanium, the arc lines are more intense. In the case of iron, the vacuum-spark spectrum is more like the arc spectrum than the spark spectrum. In general, the vacuum-spark spectra resemble the luminescence spectra produced by cathode-ray bombardment in a high vacuum; the conditions of emission in the two cases are considered to be very similar.

A. A. E.

The Limits of Absorption L of the Elements Barium to Antimony. LINDSAY (*Compt. rend.*, 1922, 175, 150—151).—The limits of absorption, L , of the elements barium, caesium, iodine, tellurium, and antimony were measured by means of a vacuum spectrograph, using a prism of calcite, and a 0.1 mm. slit. The current used was between 10 and 50 milliamperes, at such a voltage that the second order did not appear in the spectral region studied, for example, 8000 volts for antimony. The screens consisted of a thin sheet of paper which had been dipped in a saturated solution of a suitable compound of the element. The following results, expressed in units of 10^{-11} cm., were obtained.

	L_1 .	L_2 .	L_3 .
Barium	2357.7	2199.5	2060.2
Cæsium	2467.8	2307.3	2160.5
Iodine	2712.4	2548.3	2381.9
Tellurium	2847.0	2683.7	2502.6
Antimony	2994.5	2831.0	2632.7

A comparison of the values of the Sommerfeld doublet (expressed as $1/\lambda$) for the limits of absorption (L_2-L_1) and the two lines $\beta_1-\alpha_2$ measured by Hjalmar shows a very satisfactory agreement.

G. F. M.

The Variations of the Spectrum of the Mercury Arc with the Conditions of Emission. ST. PROCOPIŮ (*Compt. rend.*, 1922, 175, 217—219).—If the arc is produced in a vacuum with a current of 14—15 volts and 1.9 amperes, the spectrum contains more lines towards the extremity of the ultra-violet than if a more powerful current (65 volts, 3.5 amperes) is used with a correspondingly increased vapour density. Under both conditions of current, the cathodic spectrum is richer in lines than the anodic, and with the higher voltage the cathodic lines become diffuse, sometimes so much so as to form a continuous spectrum. If the arc is struck in air, under water, or especially in coal gas, the lines of the two secondary series are weakened and widened compared with the lines of the vacuum arc, this being the more pronounced the higher the position of the line in the series. The line 2537 appears strongly inverted in the three media, and a number of lines are strengthened, notably 3984, 2857, 2848, and 2224. The author considers that the widening of the lines is related to the Stark effect and explainable by the assumption of an intermolecular electric field, whilst the weakening or disappearance of certain lines when the arc is struck in air, water, etc., may be explained by Bohr's theory.

G. F. M.

Fine Structure of Röntgen Spectra. ADOLF SMEKAL (*Sitzungsber. Akad. Wiss. Wien*, 1921, 130, 25—30; from *Chem. Zentr.*, 1922, i, 240; cf. A., 1921, ii, 292, 615).—The scheme for the *L* series can only be tested for tungsten since in this case alone are sufficient experimental data available. There are five *M* layers and at least five *N* layers.

G. W. R.

The Absorption Spectrum of Potassium Permanganate. A. HAGENBACH and R. PERCY (*Helv. Chim. Acta*, 1922, 5, 454—468).—Spectrophotometric measurements were made of the absorption of aqueous potassium permanganate solutions at twelve dilutions from 2^{-6} to 2^{-17} mol. per litre at sixty-one points in the visible spectrum between wave-lengths 438 and 701 $\mu\mu$. From the results, curves were plotted in which the percentage absorption was plotted against the wave-length at each dilution, and from the curves obtained the positions of the absorption bands were deduced. New figures are given for the wave-lengths of the eight known bands, differing slightly in some instances from those given by Formánek. The new figures are 4388, 4537, 4695, 4866, 5050, 5249, 5461, 5695. These bands are related in such a manner that there is a constant difference between the oscillation frequencies of successive members of the group. This law has not previously been observed to hold for an inorganic substance. The broad band which Formánek ascribed to the wave-length 6370 has been resolved into four bands at 5949, 6264, 6528, 6862. These then

form a continuous series with the other eight bands. The Beer-Lambert law was found to hold within the limits of 5%. E. H. R.

Structure of the Second Cyanogen Band. JUNZÔ ÔKUBO (*Sci. Rep. Tôhoku Imp. Univ.*, 1922, **11**, 55—86).—The second cyanogen band radiated by the ordinary carbon arc in air was examined under higher dispersion and resolving power than that used by previous investigators. The wave-length of all lines lying between the first head at λ 4216 Å. and λ 3900 Å. (about 1555 in all) were measured and calculated in international units. This band contains many doublet lines with close components, some of which have been previously described as singlets. Three remarkable series were traced, one triplet series beginning from the first head and two closed doublet series beginning from the second head. The intensity distribution in these series is described. The wave-length of the heads of cyanogen bands were measured and some simple relations are shown to exist between the differences in the ratios of the wave numbers of the heads; for example, the differences in the ratios for successive heads are generally in an arithmetical progression. T. H. B.

Absorption Spectra of Triphenylmethane Colouring Matters. EMILIO ADINOLFI (*Atti R. Accad. Lincei*, 1922, [v], **31**, i, 461—464).—The author has shown (*Rend. Accad. Sci. Fis. Mat. Napoli*, 1921, [iii], **27**) that triphenylmethane colouring matters, when dissolved in water, alcohol, ether, benzene, glycerol, toluene, or xylene, exhibit absorption curves of similar types with maxima which undergo gradual displacement as the molecular weight of the solvent increases. Solutions of 19 of these colouring matters in water and in methyl, ethyl, isobutyl, and amyl alcohols have now been examined spectroscopically by the diffusion method (A., 1920, ii, 720), the positions of the mid-points of the two absorption bands being tabulated. With the alcoholic solutions the velocity of diffusion is greater than with those in other solvents, this velocity diminishing as the molecular weight of the alcohol increases. The diffusion of some of the compounds examined is characterised by two distinct phases: the first consists in a rapid propagation of the coloration with a very low gradient of the concentration, the spectrum showing only one of the characteristic bands which assumes constant breadth; the original surface of separation remains and gives rise to a second, slower diffusion, as a result of which the absorption curve characteristic of this group of compounds becomes complete. That this phenomenon of double diffusion appears to be due to two different atomic groupings is supported by other evidence to be published later.

The gradual displacement of the maxima of the absorption curves as the molecular mass of the alcoholic solvent increases is observed with all the compounds examined with the exception of uranine and rhodamine-B. T. H. P.

The Calculation of the Colour of the Azo-dyes and Related Coloured Substances. JAMES MOIR (T., 1922, **121**, 1555—1562).

Absorption Spectra of Benzeneazophenol and its Derivatives. ALPHEUS W. SMITH and C. E. BOORD (*J. Amer. Chem. Soc.*, 1922, **44**, 1449—1455).—The absorption spectra of benzeneazophenol and each of the three isomeric tolueneazophenols, chlorobenzeneazophenols, and nitrobenzeneazophenols have been measured in alcohol and in alcohol containing sodium hydroxide. The measurements were made by the visual method and in all cases continuous absorption curves were obtained. It is shown that the substitution products absorb light more strongly in the visible region than does benzeneazophenol, and the alkali salts more strongly than the free dyes. The order of absorption for the isomerides in the dilutions examined is para > meta > ortho, except in the case of the nitro-derivatives, when it becomes para > ortho > meta. The deviation of the latter derivatives may be due to a difference in structure. The changes in absorption brought about by substitution are discussed briefly in the light of McClelland's mutual induction theory of absorption of light. J. F. S.

Disturbance of the Power of Fluorescing of Fluorescent Solutions by Light, and the Photochemical Equivalent Law. PETER PRINGSHEIM (*Z. Physik*, 1922, **10**, 176—184).—The change in the intensity of the fluorescence of fluorescein, eosin, and other substances in aqueous, alcohol, glycerol, and amyl alcohol solutions, when submitted to an intense beam of light, has been investigated. It is shown that the disturbance of the power of fluorescence, discovered by Perrin (*A.*, 1918, ii, 418; 1919, ii, 177) and investigated by Wood (*Proc. Roy. Soc.*, 1921, [A], **99**, 362; this vol., ii, 334) is due to the concentration of the acting light, a result also obtained by Wood. Of two solutions of equal concentration, that with the weaker fluorescence is changed more rapidly than the other, and of two solutions with the same intensity of fluorescence and with identical absorption and fluorescence spectra, the one may be changed more rapidly than the other and thereby a different product produced. The fluorescence spectrum of a newly-formed substance is displaced in the same sense as the absorption spectrum of this substance. The intensity of fluorescence is not markedly changed at low temperatures, but the chemical reaction is practically inhibited. For these reasons, the hypothesis of Perrin (*loc. cit.*), according to which fluorescence is an attendant phenomenon of chemical change, cannot be maintained. On the other hand, there is the second possibility, that either fluorescence or chemical change is of importance for the validity of the photochemical equivalent law. J. F. S.

The Influence of Constitution on the Rotatory Power of Optically Active Substances. XV. A New Constant for Calculating the Curve of Rotation-dispersion. H. RUPE (*Annalen*, 1922, **428**, 188—198).—In place of the customary magnitudes $\lambda\alpha$ and λ_0^2 , it is now proposed to use $\lambda\alpha$ and λ_0 , and, in particular, their product, $\lambda\alpha \cdot \lambda_0 = k$, which is termed the "Rotation-Dispersion Product," and is signified by the letters "*P.R.D.*" It is shown (by reference to camphor derivatives) that the *P.R.D.*'s

exhibit a fair degree of constancy for a series of substances of the same "optical type."

C. K. I.

Molecular Ælotropy in Liquids. C. V. RAMAN (*Nature*, 1922, 110, 11).—In experiments with many liquids on the molecular scattering of light, the scattered beam in a direction transverse to the primary rays shows a large admixture with unpolarised light. It is therefore suggested that the polarised and unpolarised parts of molecularly scattered light may arise in two distinct ways, the former being a mass effect, and the latter a molecular effect. Passing from the condition of vapour to that of liquid considerably increases the ratio of unpolarised to polarised light, as has been experimentally shown in the cases of ethyl ether, benzene, and chloroform; such increase would naturally be in proportion to the number of molecules per unit volume. Moreover, when a substance is in the state of vapour under small pressures, both the positions and orientations of its molecules are absolutely at random, and assuming the molecules to be ælotropic, the degree of imperfection of polarisation of the light scattered by it may be calculated; in the liquid state, however, the packing of the molecules is so close that their ordering in space is no longer at random, but the orientations may still be considered to be arbitrary. Viewed in this way, the same results are obtained.

A. A. E.

The Rotatory Power of Crystals and Molecular Rotatory Power. LOUIS LONGCHAMON (*Compt. rend.*, 1922, 175, 174—177).—A comparison of the crystalline rotatory power with the molecular rotatory power expressed as $[\alpha]d/100$, where $[\alpha]$ is the specific molecular rotation, and d the density of the crystal, showed with the substances examined, including camphor, sucrose, rubidium, and ammonium tartrates, and ammonium molybdomalate, no definite constant relationship. On the other hand, the rotatory dispersions, that is to say, the ratio of the rotations for two given spectral lines, exhibited practically constant values, in spite of great variations in the rotatory power in passing from the liquid to the crystalline state. In the case of camphor, remarkably constant values were obtained for the rotatory dispersion in solution in ethyl ether, and in hexane, and in the liquid, gaseous, and crystalline states. If it is admitted with Traube and Landolt that there is a structural rotatory power in crystals superimposed on a molecular rotatory power, the former must therefore present the same rotatory dispersion as the latter, which, a priori, does not seem reasonable. The only alternative to this is that there is no structural rotatory power, and the rotatory power of the crystal is therefore the molecular rotatory power in a determined direction, it being established that the rotation is variable with the direction of observation. The rotatory dispersion, however, is in all cases independent of the direction of observation.


G. F. M.

A Case of Total Anomalous Rotation-Dispersion. H. RUPE and H. SCHMID (*Helv. Chim. Acta*, 1922, 5, 437—443).—Tschugaev and Glinin (A., 1912, ii, 1020) consider that the cause of anomalous

rotation-dispersion is to be found in the influence of the optical superposition of several asymmetric carbon atoms on an absorption band in the ultra-violet. The authors dispute this hypothesis, and suggest that the phenomenon is due to the disturbing influence of unsaturated residues on the asymmetric carbon atom. In agreement with this view, they find that triphenylmethyl- α -aminocamphor exhibits total anomalous rotation-dispersion in benzene and in acetone solution, whilst the condensation product of α -aminocamphor with hydroxymethylenecamphor is normal.

α -Triphenylmethylaminocamphor is prepared by condensing triphenylchloromethane with α -aminocamphor, m. p. 153–154°, $[\alpha]_D -1.34^\circ$, $[\alpha]_D -1.68^\circ$, $[\alpha]_{H_K} -1.68$ ($\lambda=546.3$), $[\alpha]_F -2.13^\circ$ in benzene.

Methylenecamphor- α -aminocamphor,



forms white to pale yellow leaflets, m. p. 228–230° (decomp.); $[\alpha]_D +90.51^\circ$, $[\alpha]_D +121.8^\circ$, $[\alpha]_{H_K} +151.7^\circ$, $[\alpha]_F +211.4^\circ$ in benzene.
E. H. R.

Coagulation of Colloids by Sunlight. P. B. GANGULY and N. P. DHAR (*Kolloid Z.*, 1922, **31**, 16–19).—The action of sunlight on colloidal gold, silver, platinum, copper, sulphur, selenium, a number of colloidal sulphides, hydroxides, and complex colloidal solutions, both pure and in the presence of protecting colloids, has been investigated. In most cases, it is shown that sunlight increases the rate of coagulation. In the case of the metallic sulphides, the sensitiveness toward sunlight follows the order: arsenic, cadmium, mercury, and antimony sulphide, whilst in the presence of gelatin as protecting colloid the order is: nickel, platinum, zinc, gold, iron, cobalt, and silver sulphide. In the case of a manganese dioxide sol containing 0.3 gram of manganese dioxide per litre, sunlight has the same coagulating action as 2.5 c.c. of a 2*N*-solution of copper sulphate acting on 25 c.c. of the sol. Colloidal solutions of stannous, bismuth, lead, manganese, and ferrous sulphides, when acted on by sunlight in the presence of air, are oxidised with the formation of colloidal sulphur, which on further action of light is coagulated.
J. F. S.

Decomposition of Iodoform in Solutions by Means of Radiant Energy. E. H. BUTLER (*Chem. News*, 1922, **125**, 38–39).—Iodoform in benzene or carbon tetrachloride solutions is decomposed by light, but in alcoholic solution there is no decomposition.
J. F. S.

Thermotropy. JOSÉ RODRÍGUEZ MOURELO (*Anal. Fis. Quím.*, 1922, **20**, 139–145).—Some experiments are recorded on the effect of heat on the phototropy of certain sulphides. Phototropy is diminished by low temperatures and increases with rise of temperature up to a maximum, after which it diminishes. Phototropy may, under certain conditions, be induced by simple heating in the absence of light.
G. W. R.

Conductivity of the Latent Image. A. G. RABINOVICH (*J. Physical Chem.*, 1922, 26, 577—590).—With the object of furnishing evidence as to the nature of the undeveloped photographic image, the author has determined the electrical resistance of celluloid photographic films and silver bromide gelatin emulsions both before and after illumination. The results show that the resistance undergoes no change measurable by the most sensitive methods, and consequently this method throws no light on the nature of the latent image. The electrical conductivity of a dry film 5×10 mm. at $18-20^\circ$ is of the order 10^{-11} /ohm cm. and the temperature coefficient about 10% per degree, whilst the specific conductivity of a silver bromide gelatin emulsion at 22° is 8.0×10^{-4} ohm $^{-1}$ and the temperature coefficient 3% per degree.

J. F. S.

Absorption and Dispersion of Radiation. FRITZ WEIGERT (*Z. physikal. Chem.*, 1922, 44, 414—444).—A theoretical paper in which the absorption and dispersion of radiation are considered in the light of modern work. The experimental results of the effects of linearly polarised light on the different photo-sensitive systems and the adaptation of the system to the colour of the exciting radiation are mainly considered. It is shown that for these effects not a single molecule, but always a whole system of numerous unordered and closely-packed molecules of different types, is responsible. The actual process of adsorption, which is closely connected with the adaptation, can only occur through the simultaneous action of different molecules or molecular groupings. The combination of the individual parts of such a closely packed aggregate occurs through electrons in an internal photo-electric effect, in which an electron from a molecule is projected in the direction of the radiation with an energy given by the relationship $1/2mv^2 = h\nu$, and taken up by an electron receiver. The frequency and direction of the absorbed radiation is determined by the distance between the two molecules and their relative orientation; since almost all combinations are possible, the adaptation to the colour and direction of the radiation is practically unlimited. The absorption spectrum varies with the spatial arrangement and the relative proportion of the electron emitter and receiver, in the sense that the bands are displaced toward the red when the proportion of receivers is greater than that of emitters, and toward the blue when the proportion is the reverse. It is possible by spectrometric measurements to differentiate the two constituents. The inconstancy of the absorption of the chief coloured substances, the finely divided disperse systems, organic dyes, and inorganic complex compounds can be shown in the case of the photo-chlorides, cyanin collodion layers, and the blue copper-ammonia complexes respectively. In all these cases, Beer's law does not hold, and the reason for this is to be found in the method of action of the electron emitters and receivers, so that here a perfectly general phenomenon is to be found. The emitters expel their electrons with an energy $h\nu$ when they encounter a radiation of frequency ν in a perfectly

definite phase, which probably is the maximum of the magnetic displacement. The electron then probably takes a cometary path, and if, at the limit of its path, it meets an electron receiver it stays there, and the system now richer by one energy quantum takes from the radiation the amount of energy still failing. In the more likely case where the electron does not meet with a receiver, it returns to the emitting molecule and thereby emits an impulse wave of the frequency ν . This impulse is, however, emitted with a certain delay, which is equal to the time of flight of the electron. During the passage of radiation through a material system, a diminution of the light velocity is recorded which is greater the greater the frequency, for the electron has a longer path and the time of flight is longer. Here is to be found a meaning for normal dispersion. When the time of flight is exactly a half period of the radiation, the emitted impulse destroys the exciting radiation entirely by interference. If this corresponds with the characteristic frequency of the substance, then the radiation does not pass through the medium, and a zero velocity of light and an infinitely large index of refraction are observed. When the time of flight is only a little larger than the half period, then it appears as though the emitted wave commences before the exciting radiation disappears. Here there is an apparent increase in the velocity of light and abnormal dispersion. This repeats itself when the time of flight t_ν is equal to an odd multiple of the half period which is expressed by the formula $t_\nu = (n - 1/2)1/\nu$. The abnormal dispersion of spectrum lines is probably explained by this formula, when for n the natural series of numbers is inserted.

J. F. S.

Determination of the Velocity of the α -Rays of Polonium. (MLLE) IRÈNE CURIE (*Compt. rend.*, 1922, 175, 220—222).—The velocity of emission of the α -rays of polonium was determined by the magnetic deviation method, the distance between the impressions produced by the α -rays on a photographic plate in the presence and in the absence of the magnetic field being measured with great precision by means of the instrument used for spectrographic measurements, from which it was calculated that the velocity of the rays is 1593×10^9 cm. per second, and the ratio of the emission velocities of the α -rays of polonium compared with those of radium-*C* is accordingly 0.829. This is in close agreement with the value 0.826 given for the ratio of the cube roots of the penetrating power of the α -rays of these two elements.

G. F. M.

α -Particles as Detonators. G. H. HENDERSON (*Nature*, 1922, 109, 749).—If it is considered that when an α -particle passes through matter, the matter in its proximity is momentarily raised to a high temperature, the detonation of certain unstable substances would be expected to take place on exposure to the action of these particles. Air-dried nitrogen iodide is detonated in this way. The detonation is not caused by the first α -particle which happens to strike the substance, but appears to be a probability effect.

A. A. E.

Multiple Valency in the Ionisation by α -Rays. T. R. WILKINS (*Physical Rev.*, 1922, **19**, 210—220; cf. Millikan, Gottschalk, and Kelly, *ibid.*, 1920, **15**, 157; *Proc. Nat. Acad. Sci.*, 1919, **5**, 591).—Slight modifications were introduced into Millikan's oil-drop method for determination of the valency of individual ions, polonium being used instead of radium bromide as a source of α -rays. In air, not more than 3% of the positive ions could have been doubly charged; the number was probably considerably smaller. In helium, a determination was made of the variation of the number of "doubles" with the range; the maximum proportion was found to be about 10% for the range of maximum ionising power. The corresponding figure for hydrogen was 0.5%. The relative stopping power for α -rays of air and helium was found to be about 3.8, a result which is in agreement with the Bragg-Kleeman law. A. A. E.

Theory of Röntgen Spectra. II. ADOLF SMEKAL (*Sitzungsber. Akad. Wiss. Wien*, 1920, **129**, 635—660; from *Chem. Zentr.*, 1922, **i**, 240).—The error introduced by the use of the Loschmidt number in the standard value for the lattice constant of sodium chloride is considered. Attention is also directed to a theoretical L_{α} arrangement which takes account of the relativity correction. The hypothesis used in the proof of the electron ring theory may be reduced to Bohr's frequency hypothesis. The presence of three electrons in the K ring follows qualitatively from the assumption of a spacial arrangement of electrons in the sheath. The determination of the quantum conditions of the L and M electrons permits the number of L and M absorption edges to be obtained. G. W. R.

Röntgen Spectra and Chemical Valency. GREGOR WENTZEL (*Naturwissenschaften*, 1922, **10**, 464—468).—A discussion of the dependence of the K absorption limits of chlorine and phosphorus on the valency as observed by Lindh (*Compt. rend.*, 1921, **172**, 1175) and Bergengren (*A.*, 1920, **ii**, 654). It is probable that the properties of the K limits of titanium, vanadium, and chromium reported by Fricke (*A.*, 1921, **ii**, 6) are due in so far as this structure extends to the weaker part to the presence of ions of different valency. The structure of the $K\beta$ line of the light elements is explained similarly (cf. Hjalmar, *A.*, 1920, **ii**, 655; 1921, **ii**, 145, 292). CHEMICAL ABSTRACTS.

Radium Synthesis of Carbon Compounds from Air. F. HARRISON GLEW (*Nature*, 1922, **109**, 714).—Quartz or glass fibres, supported immediately above an uncovered radioactive surface, the whole being enclosed in a box in a dark room, become covered in a few days with a white, viscid liquid, tending to form droplets. Prolonged exposure causes increase in size of the droplets, coloration to dark brown, and a tendency to solidify with irregular contraction. The gaseous emanation of radium seems to be necessary, since deposits cannot be obtained by using α -, β -, or γ -rays, either separately or in combination. A chemical examination of the products has not yet been made. A. A. E.

An Attempt to Influence the Rate of Radioactive Disintegration by Use of Penetrating Radiation. G. HEVESY (*Nature*, 1922, **110**, 216; cf. Ellis, this vol., ii, 339).—It is assumed that the emission of γ -rays from the nucleus of the atom precedes the disintegration process. If, therefore, the nucleus could be induced to take up a γ -ray impulse supplied by an exterior source, it would mean a change in its stability, and so most probably a change in the rate of its disintegration. Experiments with uranium in radioactive equilibrium with uranium-*X*, and with radium-*D* in equilibrium with radium-*E*, failed to show a change in β -radiation greater than 0.1% and 0.2%, respectively. A. A. E.

Measurement of the Resonance, Radiation, and Ionisation Potentials of Several Gases and Vapours. PAUL EDWARD BOUCHER (*Physical Rev.*, 1922, **19**, 189—209; cf. Mohler and Foote, A., 1920, ii, 464).—A modification of Compton's variable-area electrode was employed. The resonance potentials were obtained from the difference in voltage between successive peaks of the resonance curves; the ionisation potentials from the Lenard current-voltage curves corrected as to zero by the resonance curves; the relative importance of radiation and ionisation at various voltages from the ratio of the Lenard currents obtained with the plate to the corresponding currents obtained with the gauze electrode having only one-fourth the area. The critical resonance potentials, potentials at which radiation is predominant, and ionisation potentials, in volts, accurate to about ± 0.1 volt except in the case of toluene, are as follows: hydrogen, 10.1, 10.1—13.6, 13.6, and 15.6; nitrogen, 8.4, 8.4—15.8, 15.8; oxygen, 8.0, none, 14.0; ethyl ether, 6.6, 8.1—10.1, 13.6; benzene, 6.0, none, 9.6; toluene, 6.2 ± 0.2 , none, 8.5 ± 0.5 ; xylene, 6.5, none, 10.0; chloroform, 6.5, none, 11.5. A. A. E.

The Effect of Oxygen and Hydrogen on the Emission of Electrons from Hot Platinum. LAURICE L. LOCKROW (*Physical Rev.*, 1922, **19**, 97—113).—By using a tube with a platinum cylinder surrounding a V-shaped filament, it was found that the effect of oxygen and air was slightly to decrease the emission, the observations of Richardson, Wilson, and others being thereby confirmed. The effect is, however, temporary and is not due to the formation of heavy ions, neither is there evidence of the formation of an oxide of platinum. Pure hydrogen probably has no effect on the emission from pure platinum, the observed effects, although considerable, being ascribed to impurities. Richardson's constants for pure platinum were found to be: $A=9900$; $b=52000$, whence the thermionic work function for platinum is 4.46 volts. These values are lower than those obtained by other methods, and are subject to errors inherent in the resistance method of measuring temperatures. A. A. E.

Behaviour of Free Electrons toward Gas Molecules. H. B. WAHLIN (*Physical Rev.*, 1922, **19**, 173—186).—To account for the appearance of abnormal negative mobilities at low pressures,

Wellisch (*Amer. J. Sci.*, 1915, [iv], **39**, 583; 1917, [iv], **44**, 1; A., 1917, ii, 352) assumes that in order to form an ion an electron must have more than a certain amount of energy, whilst Thomson (*Phil. Mag.*, 1915, [vi], **30**, 321) considers that it must have made, on the average, n impacts. The author finds that when the energy of the electrons is increased the number of ions is not increased; hence Thomson's theory is preferred. The Thomson constant n was determined for carbon monoxide, ammonia, ethylene, acetylene, ethane, ethyl chloride, and chlorine. The results, which depend on the values assumed for the mobility of the electron in the various gases, indicate a progressive decrease in the value of n in the above order from about 10^8 for carbon monoxide to 10^3 or less for chlorine. It is concluded that electronegative gases attach electrons more readily than do electropositive gases, and that the degree of saturation of hydrocarbons has but little effect. The mobility of negative ions in chlorine, ethane, and ethylene was found by Rutherford's method to be, respectively, 0.73, 1.30, and 0.91 cm./sec./volt/cm. at atmospheric pressure. A. A. E.

Piezo-electricity of Potassium Sodium Tartrate Crystals.

E. KILBURN SCOTT (*Trans. Faraday Soc.*, 1922, **17**, 748—752).—An account of the piezo-electric properties of Rochelle salt crystals, based on the recent work of A. McLean Nicolson. Suitable crystals are best prepared by the hurried cooling of a saturated solution of potassium sodium tartrate, in which a temperature gradient is established and seed crystals are immersed. Piezo-electric properties are best developed in crystals having one side slightly concave and the other convex, and in which the "hour-glass" marking is developed. An average-sized crystal weighs 100 grams. The piezo-electric effect is very pronounced when the crystal is dried in 90% alcohol for twenty-four hours, then in 100% alcohol for about four hours, and afterwards baked at 40° for several days. In the case of crystals showing the "hour-glass" structure, the piezo-electric effect is best developed by the application of a twisting couple about the principal axis of the crystal. Potentials as high as 500 volts can be developed, and the crystals are readily applicable to the reproduction and transmission of sounds.

J. S. G. T.

The Electrodeless Discharge in certain Vapours.

J. K. ROBERTSON (*Physical Rev.*, 1922, **19**, 470—477; cf. A., 1921, ii, 668).—The effect of the discharge, which was produced by electromagnetic induction, varied with the vapour pressure and with the *E.M.F.* Potassium at 250—300° gave a bright ring with a spectrum including seven or more members of each subordinate series. Sodium above 300° gave a bright yellow discharge showing the *D*-lines, four members of the diffuse series, and three members of the sharp series. Lithium up to 500° gave only a feeble discharge due to impurities. Kowalski's observations in the case of mercury were confirmed and extended. A white ring discharge (at 70—110°), at first dazzling, became fainter and was followed (at 110—115°) by a green glow of which the spectrum showed

a few lines superimposed on a continuous band extending from the violet to the yellow. The glow may be associated with poly-atomic molecules formed at the higher vapour pressures. Iodine at -5° to $+5^{\circ}$ gave a pale yellow ring showing a band spectrum, changing with increase in the spark gap to a green, pink-bordered ring with a line spectrum. The change may be due to dissociation of molecules into atoms. It is suggested that the method may be applied to the investigation of isotopy. A. A. E.

Construction of Platinum Film Electrodes and their Method of Use. A. EILERT (*Z. angew. Chem.*, 1922, 35, 445—446).—The method of preparing durable electrodes from burnt-in layers of platinum is described. A thin piece of platinum wire is sealed into the end of a glass tube and while the glass is still soft pushed inward, so that a deep depression is produced. This is allowed to cool and then coated with one of the numerous platinising liquids in such a way that both the outside walls of the tube and the walls of the depression are thickly coated. The tube is then heated, at first cautiously and then to the softening point. In this way, a layer of platinum is burnt into the glass and is in good contact with the platinum wire. A drop of mercury inside the tube makes contact with the rest of the apparatus. Measurements are given to show the durability and permanence of such electrodes. J. F. S.

Construction of Platinum Film Electrodes and their Method of Use. A. EILERT (*Z. angew. Chem.*, 1922, 35, 452—455).—Examples are given of the application of the film electrodes described in the preceding abstract. Only in the case of film electrodes used in the electro-analysis of metals do the films require periodical renewal. A simplified method of electrometric titration of solutions of salts of weak acids is described. J. S. G. T.

Sodium Amalgam Electrode for the Determination of Sodium Ion [Concentration]. BENJAMIN S. NEUHAUSEN (*J. Amer. Chem. Soc.*, 1922, 44, 1411—1416).—The trustworthiness of a sodium amalgam electrode for the measurement of the concentration of sodium ions has been investigated by measuring cells of the type $\text{NaHg}|\text{NaCl}(c)||\text{sat. KCl}||\text{sat. KCl, Hg}_2\text{Cl}_2|\text{Hg}$ at 25° . Further measurements have also been made with sodium chloride containing various amounts of potassium, ammonium, calcium, and zinc chlorides. The electrode was made up in a vessel similar to that employed by Lewis and Kraus (*A.*, 1910, ii, 1027), except that at the bottom of the vessel there was a tap for draining off the amalgam which had dropped from the electrode. An amalgam containing about 0.16% of sodium was used and measurements were made against the normal electrode. The following values were obtained for the concentrations of sodium chloride stated: 0.20*N*, 2.1483 volt; 0.10*N*, 2.1650 (2.1649); 0.02*N*, 2.2035 (2.2042); 0.01*N*, 2.2198 (2.2213); the values in brackets are the *E.M.F.* values calculated from the value obtained for the 0.20*N*-solution. Further readings were made in each case at five-minute intervals to see how rapidly

the electrode lost its sodium, and then a new surface was produced and a further reading made. The following figures show the type of change: 0.2*N*-sodium chloride, voltage at start, 2.1482; after five minute intervals, 2.1482, 2.1480, 2.1476, 2.1470, 2.1460; new surface, 2.1483. Long series of measurements are recorded for solutions containing the salts mentioned above, gelatin, serum, and proteins and no difficulties were observed in the work.

J. F. S.

Modification of the Clark Hydrogen Electrode Vessel to permit Accurate Temperature Control. GLENN E. CULLEN (*J. Biol. Chem.*, 1922, 52, 521—524).—The Clark electrode vessel (A., 1916, ii, 75) is provided with an additional opening by means of which a thermometer may be inserted in the solution. E. S.

Electrolysis of Aqueous Solutions of Alkali Nitrites with a Lead Anode, and an Electrometric Determination of the Constitution of the Complex Ion formed. F. H. JEFFERY (*Trans. Faraday Soc.*, 1922, 17, 709—718).—In continuation of previous work on the electrolysis of aqueous sodium nitrite solutions employing a silver or copper anode (A., 1920, ii, 662; 1921, ii, 374), the author has investigated the phenomena occurring when a pure lead anode is employed in such electrolysis. A cathode of smooth platinum was used. The strength of the anolyte solution varied from 6.9 grams to 27.6 grams of sodium nitrite per 100 grams of water. The temperature was maintained at 17—18°, and the anode area was 100 sq. cm. The current ranged from 0.099 to 0.77 ampere. Lead was in no case deposited on the cathode, but was dissolved at the anode, forming a complex ion which gave a bright orange colour to the anolyte. For small concentrations of lead in alkali nitrite solutions, probably the only plumbo-nitrite complex formed is $[\text{Pb}(\text{NO}_2)_4]^-$. Crystals of $\text{Pb}(\text{NO}_2)_2 \cdot \text{H}_2\text{O}$ are in equilibrium with solutions obtained from anolytes of certain concentrations. Solid solutions derived from aqueous solutions containing potassium-, nitrate-, and nitrite-ions and the plumbo-nitrite complex are orange-coloured, but more red than the orange of lead nitrite. It is suggested that the complex exists as a definite atomic group in such crystals. J. S. G. T.

Electrolysis of Sodium Silicate Solutions. JAMES FREDERICK SPENCER and KATHLEEN PROUD (*Kolloid Z.*, 1922, 31, 36—37).—Orthosilicic acid may be prepared by the electrolysis of a 50% solution of sodium silicate in a divided cell, using a heavy anode current density. The cathode employed was a platinum dish 7.5 cm. in diameter, and this contained a small porous pot which held the anode, a coil of platinum wire of 1.80 cm.² surface. A current of 9 amperes was used, and this fell rapidly until no current passed, due to the formation of an insulating layer of orthosilicic acid on the anode. On bending the wire, the deposit fell away and the current again passed. The product is glass-like in appearance, entirely insoluble in water, and stable in the air.

J. F. S.

The Electrochemistry of Non-aqueous Solutions. I. Measurement of Current-density and Potential Difference in the Electrolysis of Metallic Salts in Pyridine. ROBERT MÜLLER (*Monatsh.*, 1922, 43, 67—74).—A form of electrolytic cell suitable for these measurements is described, and the preparation of an appropriate standard electrode investigated. The element $\text{Ag}/0.1N\text{-AgNO}_3$ in pyridine has a potential (measured with reference to the calomel electrode) of +0.28 volt ($H=0$), and the element $\text{Ag}/0.1N\text{-AgNO}_3$ in pyridine/ NH_4NO_3 saturated solution in pyridine, has +0.26 volt ($H=0$). C. K. I.

The Electrochemistry of Non-aqueous Solutions. II. Decomposition Potential and Electrode Potentials in the Electrolysis of Pyridine Solutions of Silver Nitrate, and the Potential of Silver in these Solutions. ROBERT MÜLLER and ALOIS DUSCHEK (*Monatsh.*, 1922, 43, 75—80; cf. preceding abstract).—The decomposition potentials are: $0.1N\text{-AgNO}_3$, 2.15 volts; $N\text{-AgNO}_3$, 2.05 volts; saturated- AgNO_3 , 1.95 volts. The silver potentials (invariant with current density) are: $\text{Ag}/0.1N\text{-AgNO}_3$, +0.24 volt; $\text{Ag}/N\text{-AgNO}_3$, +0.33 volt; $\text{Ag}/\text{saturated AgNO}_3$, +0.33 volt referred to the hydrogen electrode. The anodic potential differences are functions of the current density for each of the above concentrations. C. K. I.

Variations in the Magnetic Susceptibility of Oxychromic Salts with Addition of Sulphuric Acid. B. CABRERA and S. PIÑA DE RUBIES (*Anal. Fis. Quím.*, 1922, 20, 175—181).—The effect of the addition of sulphuric acid to solutions of an oxychromic salt is twofold and results in a depression followed by an elevation of the magnetic constant. In the experiments recorded the variations of the magnetic constant of solutions of an oxychromic salt with time were followed for different values of the ratio H^+/Cr^{+++} . For high values of this ratio, the change is so rapid that only the elevation in the magnetic constant is noticeable. For low values of the ratio, the initial depression is clearly shown by the experimental curves. G. W. R.

A New Adjustable Thermostat for all Temperatures between 0° and 100°. SAMUEL JUDD LEWIS and FLORENCE MARY WOOD (*Trans. Faraday Soc.*, 1922, 17, 696—700).—A form of thermostat employing a toluene-mercury thermo-regulator of the type due to Barnes and to Gouy is described. The individual parts of the apparatus were independently mounted on a framework extending the length of the thermostat. The surface of the mercury used in the electrical relay employed was covered with glycerol. The dimensions of the thermostat were 24"×16"×14", and any desired temperature was attained from cold in an hour and a half. After a further half hour, the temperature did not vary by more than 0.01°. J. S. G. T.

Electric Heating and Controlling Apparatus for a small Thermostat. S. O. RAWLING (*J. Soc. Chem. Ind.*, 1922, 41, 250—251r).—An electric heating and controlling apparatus is

described by means of which the steady temperature of a small thermostat may be speedily fixed at any point between 25° and 40° without making troublesome readjustments of the apparatus. The current is supplied through two resistances, R_1 and R_2 , in series to the heating resistance, and a toluene or calcium chloride regulator is so arranged that the second resistance may be cut out or left in the circuit as the temperature demands. The resistance R_1 is of such a size that when the current passes through it and the heater alone too much heat is developed, but when the current passes through R_1 , R_2 , and the heater, insufficient heat is supplied to keep the temperature at the required point. The thermo-regulator consists of a long bulb from the neck of which a side arm slopes downward and connects with the middle of a U-tube. The U-tube itself is divided into two parts by means of a glass seal through which a platinum wire is fused. One arm of the U-tube is provided with a short length of capillary tubing of about 1 mm. bore. A side tube is fitted above the top of the capillary and a rubber stopper fitted into the widened end of the U-tube carries a platinum electrode, which projects about 1.5 cm. into the capillary tube. The upper end of the tube leading from the main bulb is fitted with a glass tap, K , of fairly wide bore and very well fitting. To fill the regulator mercury is poured into the U-tube so that its level is about half-way up the capillary tube; the toluene is then placed in the bulb; electrical terminals are provided by the platinum wire dipping into the capillary and by a copper wire dipping into the other arm of the U-tube. These electrodes are connected with a small relay which cuts out or inserts the resistance R_2 in the circuit. Adjustment is effected by bringing the bath to the required temperature with the tap, K , open. The tap is then closed and the electrode tip adjusted to the surface of the mercury in the capillary. This regulator works extremely well, and with a bath of 27 litres it is capable of keeping the temperature constant to 0.02° at 25° . Calculations are given to show how the values of R_1 and R_2 may be arrived at for any particular temperature.

J. F. S.

The Specific Heats of some Sulphides Used in Metallurgy, with Special Reference to High Temperatures. K. BORNE-MANN and O. HENGSTENBERG (*Mitt. Metallhütt. Inst. Tech. Hochschule Breslau*, 1920, 18 pp.).—By means of Oberhoffer's vacuum method improved by substituting a Nernst metal calorimeter for the Bunsen ice calorimeter, the specific heats of the quartz used as container and of several sulphides were determined. The figures for quartz (to 1400°) agree best with those of Magnus (A., 1913, ii, 103) but also satisfactorily with those of White (A., 1909, ii, 966) and of Wüst. Results at 100° intervals given in tabular and graphic form can be summarised as follows: galena, $0-600^{\circ}$, $0.0500-0.0540$; cuprous sulphide, $0-900^{\circ}$, $0.1432-0.1369$; iron sulphide, $0-1200^{\circ}$, $0.1664-0.2216$ (heat of fusion calculated as 60 cal. per gram); magnetic pyrites, $0-100^{\circ}$, 0.1531 ; pyrites, $0-100^{\circ}$, 0.1284 ; zinc blende, $0-900^{\circ}$, $0.1249-0.1351$; Emser blende,

0—900°, 0.1187—0.1311; Annam blende, 0—900°, 0.1131—0.1287. In zinc blendes containing much iron the relation between temperature and the average specific heats suggests a reciprocal solubility of ferrous sulphide and zinc sulphide at high temperatures. Between 720° and 760° the iron-zinc blendes showed two transition points not thus far known and these were confirmed by thermal analysis.

CHEMICAL ABSTRACTS.

The Energy of Gaseous Molecules. J. R. PARTINGTON (*Trans. Faraday Soc.*, 1922, 17, 734—741).—The experimental values of the molecular heats at constant volume of various monatomic, diatomic, and polyatomic gases are critically reviewed in the light of the equipartition of energy theory. It is suggested that exchange of energy in molecular collisions may occur in quanta, the energy being likewise distributed among the various degrees of freedom. The author discusses the translational energy of gaseous molecules, when the frequency in the energy quantum relation $\epsilon = h\nu$ is identified with the mean value of the collision frequency. It is suggested that although the translational energy is almost exactly equal to the equipartition value, at the ordinary temperature, the rotational energy may depend on the former through a whole-multiple relation existing between the respective frequencies. Vibrational frequencies are, as in the case of solid bodies, assumed to be independent of temperature. The values, deduced from experimental data, of the internal molecular heat of hydrogen at various temperatures, are compared with theoretical values calculated by means of the formulæ given respectively by Einstein, Nernst and Lindemann, Nernst, and an empirical formula of the author, in the form $\beta\nu' = 6541/\sqrt{T}$, where ν' is the rotational frequency, equal to 1000 ν , where ν is the translational frequency. The energy of the hydrogen and nitrogen molecules is considered from the point of view of Bohr's theory of atomic structure, as developed by Krüger (*Ann. Physik*, 1916, 50, 346; 51, 450). The theory indicates the molecular heat of nitrogen to be higher than that of hydrogen and less affected by temperature.

J. S. G. T.

Melting Interval of certain Undercooled Liquids. The Use of Liquid Air as a Refrigerant. JOHN BRIGHT FERGUSON (*J. Physical Chem.*, 1922, 26, 549—552).—It has been previously noted by McIntosh and Edson (*A.*, 1916, ii, 230) that solids formed by the sudden chilling of aqueous solutions of salts and acids do not show a melting interval extending from the eutectic point to the temperature at which such a liquid solution would be in equilibrium with a trace of pure ice, but rather melted exactly at this temperature. These authors prepared constant temperature baths by using such chilled materials. The present author has prepared and investigated a number of such chilled materials (glasses) from solutions of sodium chloride and hydrochloric acid. It is shown that the constant temperatures obtained on melting the glasses are dependent on the particular experimental conditions.

In the case of salt solution glasses, these temperatures approach the liquidus temperature, but are always below it. The analysis of the residual slushes left on partial melting indicate that the salt solution glasses may persist for some time at temperatures above the liquidus temperature. Since there does not appear to be an exact relation between the melting temperatures and the liquidus temperatures, the use of such glasses for constant temperature baths cannot be regarded as of general application. One explanation of the constant temperature obtained with these glasses is that this temperature is the point at which the heat absorbed by the melting glass exactly equals the heat of partial crystallisation of the excess component together with the heat lost through the walls of the container. The heat given out when unit weight of the excess component crystallises is probably very great when compared with the heat absorbed when unit weight of the glass melts. If the glass melts slowly, a constant temperature would result which would be near the liquidus temperature, but, on the other hand, if the glass melts quickly the constant temperature would be considerably below the liquidus temperature. When liquid air is used to "freeze" a solution, it is shown that the true melting curve may be obscured by the presence of the glass in the melting mass.

J. F. S.

Vapour Pressures and Heats of Vaporisation of Non-associated Liquids. F. SPENCER MORTIMER (*J. Amer. Chem. Soc.*, 1922, 44, 1429—1435).—A theoretical paper in which empirical methods for evaluating the constants in the vapour pressure and sublimation pressure equations: $\log P_v = C_v - S_v/T$ and $\log P_s = C_s - S_s/T$ are given and their significance is discussed. It is shown that S_v , the slope of the $\log P_v$ vs. $1/T$ curves is, for normal liquids, given by the expression $S_v = -68 + 4.877T_b + 0.0005T_b^2$. It is demonstrated from existing experimental data that the molecular heat of vaporisation is equal to $4.23S_v$, rather than $4.58S_v$, the value generally adopted. It is shown that the values of the molecular latent heats of vaporisation calculated from the equation $L_v = 4.23S_v$ are generally in better agreement with the experimental results than those calculated from the equations put forward by Trouton, Bingham, or Nernst. This advantage is especially noticeable when the equations are applied to substances boiling at high temperatures. Thus in the case of cadmium $T_b = 1057^\circ$, $L_{\text{obs.}} = 23480$, $L_{\text{calc.}} = 4.58S = 26300$, $L_{\text{calc.}} = 4.23S = 23920$; (Trouton) 22720; (Bingham) 30200; (Nernst) 22550. The thermodynamical equations connecting heats of vaporisation, heats of sublimation and heats of fusion and the relationships with vapour pressure and sublimation pressure are briefly reviewed and empirical methods for calculating each from the freezing and boiling points are put forward for normal liquids.

J. F. S.

The Heat of Formation of Aluminium Nitride. FR. FICHTER and ERNST JENNY (*Helv. Chim. Acta*, 1922, 5, 448—454).—The aluminium nitride used for the experiments was prepared from aluminium bronze and contained about 91% of AlN and 0.5%

of silicon, the remainder being alumina. The heat of combustion was measured in a bomb calorimeter, the nitride being mixed with a proportion of benzoic acid to facilitate combustion. The heat of combustion found for the reaction $2\text{AlN} + 3\text{O} = \text{Al}_2\text{O}_3 + \text{N}_2$ was 258.2 (254.8) Cal. Subtracting this from the heat of combustion of aluminium, $2\text{Al} + 3\text{O} = \text{Al}_2\text{O}_3 + 380.2$ Cal., the value for the heat of formation of aluminium nitride is $\text{Al} + \text{N} = \text{AlN} + 61.0$ (62.7) Cal. The figure in brackets is the corrected value, assuming that all the silicon present in the nitride was in the elementary form. The value found is considerably higher than that obtained by previous workers, 55—56 Cal. E. H. R.

The Graphitic Conception of Aromatic Carbon. A. L. VON STEIGER (*Ber.*, 1922, 55, [B], 1968—1979).—It has been shown previously that the heats of combustion of aromatic hydrocarbons can be calculated on the assumption that all aromatic C—C and C—H linkings are equivalent but differ from the corresponding aliphatic linkings. The value for the aromatic C—C linking which is thus deduced agrees very closely with that of the graphite main linking and thus important evidence of the graphitic conception of aromatic carbon, as postulated by Debye and Scherrer, is adduced. Attempts to apply similar methods to the calculation of molecular refractions have also been described (A., 1921, ii, 473), and it has been found that the results, based on linking constants, harmonise better with the observed values than do those calculated by Brühl's summation method. These results have been criticised adversely by von Auwers (this vol., ii, 98), who has pointed out that an incorrect value has been adopted for the molecular refraction of naphthalene; this is admitted, but it is pointed out that, although the deviation from strict additivity is thereby rendered more pronounced, the superiority of the newer method of calculation still remains.

The molecular refractions of the alkylbenzenes have been examined. With a compound such as ethylbenzene, the summation for the phenyl group is effected with aromatic, that of the ethyl radicle with aliphatic, constants; the character of the C—C linking joining the two halves of the molecule remains undecided, but is assumed provisionally to be of an aliphatic nature. The agreement between the calculated and observed values for the lower benzene homologues is very satisfactory; with the methyl compounds, depressions are observed which become more marked as the methyl groups are in closer proximity to one another and thus attain their maximum in compounds such as 1:2:3:4-tetramethylbenzene.

The communication concludes with a reply to the criticisms of Wibaut (this vol., ii, 239) and von Weinberg (A., 1921, ii, 668, 669). H. W.

The Heat of Combustion of Benzoic Acid, Cane-sugar, and Naphthalene. W. SWIENTOSLAWSKI and [MLLE] H. STARCZEWSKA (*Bull. Soc. chim.*, 1922, [iv], 33, 654—667).—A survey of previous work on the subject reveals discrepancies between the

results obtained by various workers (cf. Wrede, A., 1910, ii, 1038; Roth, A., 1910, ii, 584; Swientoslawski, A., 1915, ii, 315, and 1921 ii, 679; Dickinson, *Bull. Bur. Stand.*, 1915, **11**, 190; Richards, A., 1920, ii, 589; Jessy, this vol., ii, 1041; Henning, A., 1921, ii, 379; Verkade, this vol., ii, 474). Further measurements made by the authors are here compared with those of Dickinson and Verkade, and the opinions expressed by the latter (*loc. cit.*) as to the advisability of choosing benzoic acid as the standard substance are now supported by fresh experimental evidence. Reference is made to the consideration of this subject by the International Conference of Pure and Applied Chemistry, Lyons, 1922.

H. J. E.

Heat Developed on Mixing Sulphuric Acid, Nitric Acid, and Water. J. W. McDAVID (*J. Soc. Chem. Ind.*, 1922, **41**, 246—250r).—The heat developed on mixing various proportions of the following pairs of substances: 49.2% nitric acid and 96.16% sulphuric acid, 20.0% nitric acid and 96.16% sulphuric acid, 96.8% nitric acid and 60.0% sulphuric acid, 96.8% nitric acid and 99.5% sulphuric acid, and a mixed acid containing 49.8% sulphuric acid, 48.4% nitric acid, and 1.8% water with water, has been determined calorimetrically. From the results obtained together with Thomsen's values for the heat of dilution of sulphuric and nitric acids respectively ("Thermochemistry," Thomsen, Burke, p. 76), triangular diagrams have been constructed whereby the heat developed on making any mixture of the three substances may be obtained.

J. F. S.

Compressibility of Exhausted Flasks in the Determinations of the Densities of Gases. E. MOLES and R. MIRAVALLS (*Anal. Fis. Quím.*, 1922, **20**, 104—116).—The contraction of flasks used in the determinations of gas densities due to atmospheric pressure may be expressed by the empirical formula $x = 17.5 \cdot \frac{V}{W} \times 10^{-6}$, where x is the contraction per litre of volume, V the volume of the flask, and W its weight. Measurements made on six flasks by three methods showed good agreement of observed values with values calculated from the formula.

G. W. R.

The Compressibility at 0° and less than 1 Atmosphere and the Divergence from Avogadro's Law of Several Gases. PH. A. GUYE and T. BATUECAS (*Helv. Chim. Acta*, 1922, **5**, 532—543).—To determine accurately the molecular weight of a gas by physico-chemical methods, it is necessary to know accurately the weight of a normal litre, L_0 , of the gas and its divergence from Avogadro's law, $1 + \lambda$. A method is described for measuring with great precision the compressibility of a gas at 0°, from which $1 + \lambda$ can be calculated. The apparatus consisted of three cylindrical glass bulbs of approximately equal volume, accurately calibrated, in a vertical column, connected together by capillary tubing and at the top communicating with a barometer. The volume of the gas was measured in this apparatus at approximately 1, $\frac{1}{2}$, and $\frac{1}{3}$ atmosphere. Experiments were made with oxygen, hydrogen,

and carbon dioxide. The divergence from Avogadro's law was calculated from the formula $1+\lambda=(pv)_0/(pv)_1$ (cf. Guye, A., 1919, ii, 318).

The oxygen used was prepared by three different methods, from potassium permanganate, from a mixture of sodium and potassium chlorates, and electrolytically. The mean value obtained for $1+\lambda$ was 1.00085 ± 0.00002 , and the coefficient of compressibility per cm. is 11.3×10^{-6} . Taking $L_0=1.42891$, the value of the gas constant R is 22.414. Hydrogen was prepared by the action of concentrated potassium hydroxide on aluminium and also electrolytically. The mean value found for $1+\lambda$ was 0.99935 ± 0.00002 , which, taking $L_0=0.089858$, gives for the atomic weight of hydrogen 1.0077. The coefficient of compressibility is -8.6×10^{-6} . Carbon dioxide was prepared by heating pure sodium hydrogen carbonate and by the combustion of sucrose. The mean value of $1+\lambda$ was 1.00706 ± 0.00004 ; taking $L_0=1.97685$, this gives as the atomic weight of carbon 11.998, somewhat less than the value accepted by the International Committee, 12.005, but in good agreement with the probable correct value 12.00. The coefficient of compressibility is 92.2×10^{-6} . E. H. R.

The Compressibility at 0° and less than 1 Atmosphere and the Divergence from Avogadro's Law of Several Gases.

II. Ethylene. T. BATUECAS (*Helv. Chim. Acta*, 1922, 5, 544—546; cf. preceding abstract).—Ethylene was prepared by the action of sulphuric acid on ethyl alcohol and by the catalytic action of heated alumina on ethyl alcohol vapour. The mean value found for $1+\lambda$ was 1.00780 ± 0.00004 ; taking $L_0=1.26401$, the molecular weight is found to be 28.032, and the atomic weight of carbon 12.000. The coefficient of compressibility is 102×10^{-6} .

E. H. R.

Measurement of Surface Tension. ROBERT B. ELDER (*J. Physical Chem.*, 1922, 26, 558—562).—A theoretical paper in which the author discusses the measurement of surface tension in connexion with the work of Ferguson (A., 1914, ii, 768) and of Bhatnagar (A., 1921, ii, 169). It is shown that although the numerical values of the latter author are very close to those of Ferguson, the method used involves a number of errors which are partly compensating. The method and errors are discussed.

J. F. S.

Method for the Determination of the Surface of Adsorbing Powders. FRITZ PANETH and WALTER VORWERK (*Z. physikal. Chem.*, 1922, 101, 445—479).—The authors have put forward a rapid and trustworthy method for determining the surface of an adsorbing powder. The method is based on the following considerations. When a sparingly soluble powder is shaken with its saturated solution there will be an exchange of molecules between the surface of the powder and the solution; if it were possible to identify the molecules of the substance in solution, or at least a portion of them, then the partition of these between the solution

and surface could be determined, and at equilibrium the ratio of the identified molecules on the surface to the identified molecules in solution will be equal to the ratio of the total molecules on the surface to the total molecules in solution. This process can be actually carried out for substances which have isotopes; thus in the case of lead sulphate, thorium-*B* can be used and will constitute the identified molecules. The process can be followed by an electroscope and represented by the equation $\text{Isotope(ads)}/\text{Isotope(sol)} = \text{Element(surf)}/\text{Element(sol)}$. The ratio of the isotope adsorbed to the isotope in solution can be determined from the change in the β -ray activity of the solution, and the amount of lead in solution can be determined by analysis, hence, Element(surf) , that is, the weight of the surface layer, can be calculated. Using this method, the weight of the surface layer of one gram of various specimens of lead sulphate and lead chromate has been determined. The values obtained in grams of lead per gram of material are: lead sulphate 6.4×10^{-4} and lead chromate 69.2×10^{-4} . These values have been controlled by microscopic examination and found to be somewhat larger than the microscopic values, but of the same dimensions.

J. F. S.

Thickness of the Adsorbed Layer in the Adsorption of Dyes by Crystals. FRITZ PANETH and WALTER VORWERK (*Z. physikal. Chem.*, 1922, **101**, 480—488).—The adsorption of ponceau-2R by lead sulphate has been determined and from the amount adsorbed and the dimensions of the surface of the lead sulphate (cf. preceding abstract), it is shown that the adsorbed layer is one molecule thick.

J. F. S.

The Sorption of Carbonyl Chloride by Beechwood Charcoal. HUGH MILLS BUNBURY (*T.*, 1922, **121**, 1525—1528).

The Relation between Adsorption and Electrolytic Dissociation. M. A. RAKUSIN (*Biochem. Z.*, 1922, **130**, 282—285).—When a porous pot is immersed in solutions of sucrose or sodium chloride, the concentration of the solution increases owing to negative adsorption. This is correlated with the general adsorption of colloids and non-adsorption of electrolytes. H. K.

Chemical Adsorption. G. STADNIKOFF (*Kolloid Z.*, 1922, **31**, 19—32).—The author differentiates four types of adsorption: (1) solution adsorption, which is represented by Henry's law, $c_2 = kc_1$; (2) chemical adsorption, represented by the equation $c_2 = \sigma c_1 / (1 + \sigma_1 c_1)$; (3) combined adsorption, represented by the equation $c_2 = kc_1 + \sigma c_1 / (1 + \sigma_1 c_1)$, and (4) exchange adsorption, represented by $(mc_2 \cdot c'_1 v) / (mc'_2 \cdot c_1 v) = k$; or $c_2 c'_1 / c'_2 c_1 = k$. The author is of the opinion that chemical adsorption can only be definitely proved in the case of an adsorbent of definite composition possessing either marked acid or basic properties, adsorbing either base or acid from solution. With the object of ascertaining whether or no a chemical explanation for adsorption is possible, the adsorption of sulphuric, hydrochloric, nitric, and acetic acids and mixtures of two of these acids and also of iodine by aniline-black, prepared both

by the use of potassium dichromate and of potassium persulphate, has been investigated. In the case of sulphuric acid, the maximum adsorption is two molecules of sulphuric acid to one of aniline-black, and the adsorption is entirely reversible. The limiting value of the adsorption of hydrochloric acid is three molecules of hydrochloric acid to one molecule of aniline-black and in this case the formula for chemical adsorption is fully followed. The limiting adsorption of acetic acid is two molecules of the acid to one of the adsorbent, whilst in the case of the adsorption of iodine, twelve iodine atoms are adsorbed to each molecule of aniline-black, that is two iodine atoms to each quinonoid nitrogen atom of the tri-quinonoid aniline-black. The experimental results of other authors have been recalculated from the point of view of chemical adsorption (cf. Willstätter and Dorogi, A., 1909, i, 535, 975). J. F. S.

Dissociation of Hydrogen in a Tungsten Furnace and Low Voltage Arcs in Monatomic Gases. O. S. DUFFENBACK (*Science*, 1922, 55, 210—211).—A furnace consisting of a cylinder of thin sheet tungsten, and an axial tungsten filament, both heated electrically, was used to determine the voltages at which the arc was struck and broken. The results were in agreement with those required by Bohr's theory for the ionising (13·52 volts) and radiating (10·14 volts) potentials of the hydrogen atom, and the potential (16·4 volts) which, according to Bohr's theory, is approximately that necessary to accelerate an electron so that it will dissociate the molecule and ionise one of the atoms on impact. Calculated values for the percentage of monatomic hydrogen in equilibrium with diatomic hydrogen at temperatures from 1000° K. to 3000° K. appeared to be confirmed by the experimental results. A. A. E.

Formation and Dissociation of some Polyhalogen Compounds of Hydrogen in Aqueous Solution. PRIYADARANJAN RAY and PULIN VIHARA SARKAR (T., 1922, 121, 1449—1455).

Dissociation of Glucinum Sulphate. (MLLE) G. MARCHAL (*Compt. rend.*, 1922, 175, 270—272).—Glucinum sulphate decomposes under the influence of heat according to the equations $\text{GISO}_4 = \text{GIO} + \text{SO}_3$, and $2\text{SO}_3 \rightleftharpoons 2\text{SO}_2 + \text{O}_2$, and the total equilibrium pressure corresponding with temperatures from 590—830° was measured by means of a mercury manometer, the anhydrous sulphate being heated in an electric furnace in a previously exhausted porcelain tube, and the temperature measured with a le Chatelier platinum-rhodium couple. Utilising the known equation for the equilibrium of dissociation of sulphur trioxide, the partial pressures of sulphur trioxide, sulphur dioxide, and oxygen were calculated from the data obtained. The complete results are recorded in tabular form and include the following values for the partial pressures of sulphur trioxide, sulphur dioxide, and oxygen, respectively: at 700°, 3·30, 6·34, and 3·17 cm.; at 750°, 9·12, 18·24, and 9·12 cm., and at 785°, 17·23, 35·58, and 17·79 cm. Utilising the pressures at the above temperatures, the equilibrium equation was found to be $\log p_{\text{SO}_3} = -14907/T - 14·1 \log T + 57·97$, and the results obtained for other

temperatures from this equation agreed well with the pressures actually observed. The heat of dissociation of glucinum sulphate was calculated to be -53.5 cal. by comparison with that of copper sulphate and its decomposition pressures as determined by Plüddemann (*Diss.*, Berlin, 1907). G. F. M.

Thermostat Arrangement for the Determination of the Effect of Temperature on Diffusion. L. W. ÖHOLM (*Hyllnings-skrift tillägnad Ossian Aschan*, 1920, 9—17).—Experiments were carried out with potassium chloride and sugar solution (0.25N, 0.5N, and N) in a rectangular copper vessel with glass sides, in which the circulation was maintained by a propeller driven by an electric motor supported so as to prevent any shaking of the thermostat; the temperature was constant to $\pm 0.15^\circ$. In the equation $\alpha = A + B \log k$, where α is the temperature coefficient, A and B are constants, and k is the diffusion coefficient, α for potassium chloride between 13.5° and 29.5° was found to have an average value of 0.025, and for sugar between 13.7° and 28.7° the average value was 0.033. CHEMICAL ABSTRACTS.

The Determination of Molecular Weight by means of Osmotic Pressure. FOIX (*Bull. Soc. chim.*, 1922, [iv], 33, 653—654).—The author has measured osmotic pressures without using semi-permeable membranes in order to eliminate errors due to imperfections in the membrane. A solution is placed together with pure solvent in a closed system at constant temperature. If the weight of the solution remains constant, the distance between the free surfaces of the two liquids will be equal to the height of the corresponding osmotic column, since, for a perfect osmometer in a closed space, when equilibrium is reached the vapour should be in equilibrium both with the surface of the solution at the top of the column and with the pure solvent at its base. It is claimed that the method is successful, although it is difficult to maintain constant temperatures over the long period of time required to effect the measurement. H. J. E.

Solubility of Gases in Liquids. B. S. NEUHAUSEN (*J. Physical Chem.*, 1922, 26, 553—557).—The author has plotted the solubility data for ammonia, hydrogen chloride, sulphur dioxide, and carbon dioxide in methyl alcohol and ethyl alcohol and from the results shows that the formula $V = K(P\sigma/P_0)^{1/n}$ is applicable to the solubility of these gases in the solvents named, as was previously (this vol., ii, 264) shown for aqueous solutions. The values of the constants K and $1/n$ are given for all three solvents. J. F. S.

Melting Point, Latent Heat of Fusion, and Solubility. F. SPENCER MORTIMER (*J. Amer. Chem. Soc.*, 1922, 44, 1416—1429).—A theoretical paper in which a method of calculating the solubility of substances forming non-ideal solutions is described. It is shown that for those systems which do not form molecular complexes or solid solutions the ratio of the experimental to the ideal slope of the $\log Nvs. 1/T$ curves is a factor which may be determined by a graphical method. Three methods are described

whereby the ideal slope of the $\log Nvs. 1/T$ curves for any solute may be found. It is pointed out that the solubility or melting-point method of determining the latent heat of fusion of organic compounds gives very accurate results when interpreted in the light of the principles laid down in the paper. Some generalisations in connexion with the nature of binary systems obtained with various types of mixtures are put forward. The uses of these principles in determining the choice of a solvent for crystallisations and molecular-weight determinations are suggested. J. F. S.

Reactions in Fused Salt Media. II. Solvolysis. J. F. G. HICKS and WALLACE A. CRAIG (*J. Physical Chem.*, 1922, 26, 563—576; cf. this vol., ii, 147).—The authors have investigated the reactions which occur when lead chromate, lead oxide, silver chromate, and barium chromate, respectively, are dissolved in molten sodium chloride and in an equimolecular mixture of sodium and potassium nitrates at various temperatures up to about 850°. For comparative purposes the hydrolysis of lead chromate in water has been investigated both at ordinary pressure and under increased pressure. It is shown that the solvolysis of lead, silver, and barium chromates by a fused equimolecular mixture of sodium and potassium nitrates represents states of equilibrium analogous to hydrolysis. Of these reactions, the solvolysis of lead chromate is the most complete and that of silver chromate the least complete. The regularity with which the constants of solvolysis vary between consecutive observations follows the same order, and the marked irregularity in the case of silver chromate suggests the possibility of another factor entering into the reaction. The solvolysis of lead chromate by fused sodium chloride also represents a true state of equilibrium, whilst in the case of silver chromate the reaction goes to completion. In the solvolytic reactions studied, equilibrium is reached in about one hundred minutes, and the constants of solvolysis may be calculated with a fair degree of accuracy. The solvolysis of lead oxide by fused sodium chloride proceeds to completion at 850—870°, and that of silver chromate is also complete at the same temperature, but here the reaction occurs much more rapidly. The present work indicates that lead chromate and litharge combine by simple molecular addition after part of the former has undergone solvolysis in the nitrate flux. In other words, the red basic lead chromates are higher order compounds. The relatively small degree of solvolysis of lead chromate and the practically complete solvolysis of lead oxide by the same reagent under the same conditions explains the failure to obtain red basic lead chromates when a fusion containing lead chromate, lead oxide, and sodium chloride is rapidly cooled, and also the appearance of the red basic lead chromates when a similar fusion is slowly cooled. The results confirm the assumption of the dissociation $Pb_2CrO_5 \rightleftharpoons PbO + PbCrO_4$. This assumption was previously (*loc. cit.*) offered as an explanation of the failure to obtain red basic lead chromates when a fusion containing lead oxide, lead chromate, and sodium chloride was poured into water and agrees well with the evidence offered by

the thermal equilibrium diagram plotted for the system lead chromate-lead oxide. J. F. S.

Theory of Recrystallisation. H. ALTERTHUM (*Z. Elektrochem.*, 1922, 28, 347—356).—The present views of recrystallisation are considered and nucleus formation and growth are regarded as a consequence of a thermodynamic instability. It is shown that the energy changes which accompany the cold working of a metallic crystal may be divided into reversible and irreversible changes, and the different degrees of working as a continuous series of different modifications. On the basis of Boltzmann's theory of probability, a formula is developed which expresses the number of nuclei of a recrystallised metal as a function of the temperature and the degree of working, which is in excellent agreement with the experimental results of Czochralski (A., 1917, ii, 302). From the same point of view, a continuous series of modifications, which differ in the degree of dispersity, is assumed and a similar formula is derived for the number of nuclei in recrystallisation by absorption of other crystals. The proportionality factors are attributed, in part, to causes which are not thermodynamic, and the possible changes of these during the process of recrystallisation are discussed. J. F. S.

Loosening of Crystal Lattices. G. VON HEVESY (*Z. physikal. Chem.*, 1922, 101, 337—352).—The transport of material, that is, diffusion and electrolytic transport, in crystals is rendered possible by the gradual loosening of the crystal lattice by an increase of temperature. As a measure of the loosening tendency of a crystal, the dimensions of the change in electrical conductivity between solid and liquid at the melting point serve as the most suitable standard. This value is extraordinarily divergent for different substances, varying relatively from 1 for silver iodide to 20,000 for sodium nitrate. A close relationship exists between the loosening tendency and the electro-affinity of the ions which form the crystal. The greater the work necessary to convert the ions of a crystal into the neutral condition, the smaller is the tendency of the lattice to loosen. From this it follows that the loosening of the lattice is brought about by a disturbance of the normal condition of individual ions constituting the crystal. The increase in electrical conductivity which solid electrolytes undergo when illuminated is regarded as due to a loosening of the lattice which increases the mobility of the ions already present in the electrolyte, and the transport of electricity by electrons is regarded as a limiting case of this phenomenon. The relationship between the loosening and the overstepping of the Dulong and Petit value of the atomic heat at higher temperatures is discussed and the overstepping regarded as a preliminary condition of the melting process. J. F. S.

Röntgenographic Determination of Crystal Arrangement. M. POLÁNYI (*Naturwiss.*, 1922, 10, 411—416).—Röntgenographic methods must be employed to determine whether a crystal lattice is changed by elongation of a single crystal. By means of a fila-

ment diagram it was found that for a zinc crystal (1) the lattice changes its orientation with respect to the longitudinal axis when elongated; (2) in the section drawn out to a flat band (cf. Schiebold, *Z. Physik*, 1922, 9, 180; Carpenter and Elam, this vol., ii, 69; Gomperz, *Z. Physik*, 1922, 8, 184) the angle that the hexagonal axis of the crystal lattice makes with the longitudinal axis changes from 10° to 18° , and (3) the cylindrical filaments resulting from further elongation of the flat bands have the same orientation of their lattices as the flat bands. It is maintained that this same kind of shift in the orientation of the lattice occurs in a zinc wire as in the individual crystal.

CHEMICAL ABSTRACTS.

Graphical Determination of Hexagonal and Tetragonal Crystal Structures from X-Ray Data. ALBERT W. HULL and WHEELER P. DAVEY (*Physical Rev.*, 1921, 17, 549—570).—A graphical method of interpreting X-ray patterns of powders is described, for each type of lattice the logarithms of the theoretical spacings of the different planes being plotted as functions of the axial ratio. A number of graphs are reproduced for comparison with observed spacings. Three specific arrangements have been given for each system, namely, simple prism, centred, and face-centred arrangements in the cubic and tetragonal systems, and simple prism, close packed, and rhombohedral arrangements in the hexagonal system. Since (1) all possible arrangements of atoms in the cubic, tetragonal, or hexagonal systems may be obtained by the combination of one or more identical simple space lattices of cubes, right tetragonal prisms, and right 60° triangular prisms, respectively, and (2) the intermeshing of two or more identical lattices weakens or causes to disappear some of the lines due to a single lattice, but can add no new lines, it follows that the above, and all other possible arrangements, are obtained from the simple tetragonal and triangular prism lattices, respectively, by simply omitting part of the lines. The cubic forms may be found on either the tetragonal or hexagonal plots, whilst for the orthorhombic, monoclinic, and triclinic systems the relative crystal spacings must be expressed as a function of two, three, and five variables, respectively; a simple approximate method is, however, indicated.

Zinc is shown to be a hexagonal close-packed assemblage of prolate spheroids, with axial ratio 1.860 and the side of the unit triangle 2.760 Å. Cadmium has a similar structure, the corresponding measurements being 1.89 and 2.980. Indium has a structure very similar to that of aluminium (cubic close-packed), namely, a tetragonal close-packed arrangement of prolate spheroids, with axial ratio 1.06 and a unit square of side 4.58 Å. A. A. E.

X-Ray Crystal Analysis of Metals. ALBERT W. HULL (*Physical Rev.*, 1921, 17, 571—588; cf. *ibid.*, 1917, 10, 661; *Proc. Amer. Inst. Elec. Eng.*, 1919, 38, 1171; A., 1919, ii, 470; 1920, ii, 546; 1921, ii, 38, and preceding abstract).—The crystal structure of thirteen common metals has been determined. The lattices of chromium, molybdenum, and tantalum are body-centred cubes with sides 2.895, 3.143, and 3.272 Å., respectively; cobalt (α form),

nickel, rhodium, palladium, iridium, and platinum have face-centred cubic lattices with sides of cubes 3·554, 3·540, 3·820, 3·950, 3·805, and 3·930 Å., respectively; cobalt (β form), zinc, cadmium, and ruthenium have hexagonal lattices of the close-packed type with axial ratios 1·63, 1·86, 1·89, and 1·59, respectively, and with triangular sides 2·514, 2·670, 2·960, and 2·686 Å., respectively; indium has a face-centred tetragonal lattice with axial ratio 1·06 and side of elementary prism 4·58 Å. The structures of cadmium, zinc, and indium are close-packed arrangements of solid prolate spheroids, whilst that of ruthenium is a close-packed arrangement of oblate spheroids.

A. A. E.

Coagulation of Colloids. ARNE WESTGREN and JOSEF REITSTÖTTER (*J. Physical Chem.*, 1922, 26, 537—548).—A theoretical paper in which the authors summarise Smoluchowski's kinetic hypothesis of coagulation (A., 1917, ii, 297) and give an account of the work which has been published with the object of substantiating this hypothesis.

J. F. S.

Protecting Colloids. XII. Gelatin as Protecting Colloid.
II. Colloidal Selenium. A. GUTBIER and R. EMSLANDER (*Kolloid Z.*, 1922, 31, 33—36; cf. A., 1921, ii, 693; this vol., ii, 142, 283, 485).—The stabilising action of gelatin on colloidal solutions of selenium has been investigated. A very stable selenium sol may be prepared by dissolving 2—3 grams of selenium dioxide in a litre of 0·1% gelatin solution made up in chloroform water and slowly reducing at 18—25° with a solution of hydrazine hydrate (1 : 1000), taking care that the reduction is stopped before the last of the dioxide is reduced. The solution is then dialysed and preserved under a layer of chloroform. The solution is red in colour, quite clear and very stable at ordinary temperatures, but on heating a red, amorphous precipitate separates. Sols prepared as above have been preserved without change for eight years. The addition of hydrochloric acid or sodium hydroxide in small concentrations to the protected colloid causes the system to be more stable to freezing than the protected colloid without the addition of electrolyte. Sodium chloride causes the solutions to change to a lighter colour, but exerts no further action on the protected colloid. The colour change is attributed to the formation of a stable complex, salt-glutin-selenium.

J. F. S.

An Inhibition Period in the Separation of an Emulsion. T. C. NUGENT (*Trans. Faraday Soc.*, 1922, 17, 703—707).—The author has investigated the de-emulsifying action of sodium hydroxide solution on emulsions of benzene in water, containing 50% by volume of benzene and not less than 0·1% of gelatin as stabiliser. If sodium hydroxide is added immediately after production of the emulsion, de-emulsification commences at once. If the emulsion is left undisturbed for some time prior to adding the sodium hydroxide solution, de-emulsification is inhibited for a period following such addition. The inhibition period depends on the interval between the production of the emulsion and the addition of the sodium hydroxide

solution. It is suggested that the stability of the emulsion increases with its age owing to the gradual formation of protective layers of the stabilising agent about the benzene particles. J. S. G. T.

Precipitation of Metals by Hydrogen Sulphide. G. MCP. SMITH (*J. Amer. Chem. Soc.*, 1922, 44, 1500—1502).—A theoretical paper in which the author suggests that should it be thought best to regard the formation of sulphide precipitates from the ionic point of view alone, then the hydrogen sulphide (HS')-ion should be regarded as the active participating ion, and not the sulphide (S'')-ion. In support of this statement, it is shown that a 0.2*N*-solution of hydrochloric acid containing 0.2*N*-hydrogen sulphide contains in 5 c.c., 3×10^{20} non-ionised molecules of hydrogen sulphide, 15×10^{13} hydrogen sulphide ions, and only one sulphide ion. A further extreme case due to Knox (*A.*, 1908, ii, 830) is considered; according to this author, mercuric sulphide has a solubility product 2.8×10^{-54} , from which it follows that the maximum concentration of mercuric and sulphide ions is 1.7×10^{-27} in a saturated solution of mercuric sulphide, that is, one individual mercuric or sulphide ion in 1000 litres of solution, which implies that to maintain anything like equilibrium these ions must move with the velocity of light, a condition which is not fulfilled. J. F. S.

Force of Adhesion in Solutions. III. Partition of Substances between two Solvents. NIKOLAI SCHILOV, LIDIE LEPIN, MARIE JANTSCHAK, and MICHAEL DUBININ (*Z. physikal. Chem.*, 1922, 101, 353—402; cf. this vol., ii, 350).—A very large number of partition coefficients have been determined at temperatures ranging from 10° to 40°. These include: formic acid, acetic, propionic, succinic, benzoic, oxalic, tartaric, and nitric acids between ethyl ether and water; acetic, propionic, isobutyric, benzoic, mandelic, salicylic, trichloroacetic, and trichlorobutyric acids between benzene and water; iodine, pyridine, benzoic, and picric acids between toluene and water; benzoic and acetic acids between xylene and water; iodine, acetic, propionic, isobutyric, benzoic, and salicylic acids between chloroform and water; mandelic and benzoic acids between anisole and water; benzoic acid between phenetole and water; succinic acid between amyl ether and water; benzoic acid between light petroleum and water, and trichloroacetic and trichlorobutyric acids between amylene and water. It is shown that because of the mutual influence on solubility the use of the Henry-Dalton law, in connexion with the partition of substances between two liquid phases, is extremely limited, and the law therefore represents an ideal limiting condition which rarely occurs. The difference between dilute and concentrated solutions is not fundamental, but only quantitative. The partition coefficient is generally dependent on the total concentration of the solute, and the constancy of this quantity cannot alone be taken as a proof of the Henry-Dalton law, because mutual compensating volume changes in the liquids can occur, as has been actually shown in the present experiments. The limiting value of the partition coefficient at large concentrations of the solute is

not, in general, determined by the ratio of the individual solubilities in the two solvents, but by the ratio of the concentrations of the two phases at the triple point, solid-liquid-liquid. The partition data are readily presented graphically if the relative concentrations of the solute in the individual phases are plotted as a function of the total concentration. In this way, for each partition two coupled isotherms are obtained which in many cases have characteristic forms and serve to differentiate the various types. The partition coefficient can be expressed by the general approximation formula $K = C_1/C_2^n = C_1/C_2^{\log \beta_1 / \log \beta_2}$, in which β_1 and β_2 are defined by the change of concentration of the individual phases brought about by definite changes in the total concentration of the solute. The index n therefore becomes any suitable figure, and only in a few special cases has it a stoichiometric significance. In many of the cases examined now and previously the mean values of β_1 , β_2 , and n for the region of larger concentrations are approximately constant and may be followed up to the triple point. The partition of substances which are infinitely soluble in both phases is determined by the dissolving power of both solvents with respect to the solute. The coupled isotherms are convergent and meet at the critical mixture point of the two phases. This phenomenon can be investigated by partition and mixing experiments and furnishes information about the dependence of the dissolving power of water and organic solvents on constitution. Experiments on the kinetics of partition show definitely that with respect to the equilibrium concentrations partition is a time reaction of the first order. The slow diffusion process of the solute through the interface and through the concentrated layers which adhere to the interface is determinative of the velocity as in the case of heterogeneous systems of solid and liquid. The velocity constants obtained are to a degree proportional to the number of rotations per minutes of the reaction vessels and independent of the absolute amount of substance diffusing.

J. F. S.

System $\text{Na}_2\text{O}-\text{CO}_2-\text{NaCl}-\text{H}_2\text{O}$, considered as two Four Component Systems. F. A. FREETH (*Phil. Trans.*, 1922, [A], 223, 35—87).—The solubility relations in the systems $\text{Na}_2\text{CO}_3-\text{NaCl}-\text{H}_2\text{O}$; $\text{Na}_2\text{CO}_3-\text{NaOH}-\text{H}_2\text{O}$; $\text{Na}_2\text{CO}_3-\text{NaHCO}_3-\text{H}_2\text{O}$; $\text{NaOH}-\text{NaCl}-\text{H}_2\text{O}$; $\text{NaHCO}_3-\text{NaCl}-\text{H}_2\text{O}$; $\text{Na}_2\text{CO}_3-\text{NaOH}-\text{NaCl}-\text{H}_2\text{O}$ and $\text{Na}_2\text{CO}_3-\text{NaHCO}_3-\text{NaCl}-\text{H}_2\text{O}$ have been experimentally determined at 0° , 10° , 15° , 20° , 25° , 30° , 35° , 45° , and 60° . Numerous tables of the results and diagrams representing the relationships are included in the paper, from which the composition and quantity of the stable phases formed from any mixture of the components may be deduced.

J. F. S.

Liesegang's Phenomenon and Precipitate Formation. N. G. CHATTERJI and N. R. DHAR (*Kolloid Z.*, 1922, 31, 15—16).—In continuation of work previously published (this vol., ii, 205), it is shown that the peptisation of insoluble hydroxides of iron, cobalt, uranium, and thorium may also be brought about by

gelatin, agar, glue, gum arabic, and starch in the manner previously employed with glycerol. In the case of peptisation by glycerol, it is shown that this is governed by the concentration of the reacting substances. Thus glycerol does not peptise ferric hydroxide in the presence of concentrated solutions of ferric chloride. The authors describe experiments to show that the formation of Liesegang rings is closely connected with the above-named actions, and that here also the concentration of the reacting solutions is an important factor.

J. F. S.

Velocity of Chemical Change in Solid Substances. C. N. HINSHELWOOD and E. J. BOWEN (*Z. physikal. Chem.*, 1922, **101**, 504—505).—A discussion of Sieverts and Theberath's conclusions on the velocity of chemical reactions in solid systems (this vol., ii, 360), and a reference to the author's experimental work on the same subject (A., 1920, ii, 743; 1921, ii, 443; T., 1921, **119**, 721).

J. F. S.

Mechanism of the Reduction of Permanganate and its Physico-chemical Basis. II. Reaction of Permanganate and Formic Acid in Neutral Solution. JOSEF HOLLUTA and NIKOLAUS WEISER (*Z. physikal. Chem.*, 1922, **101**, 489—497; cf. this vol., ii, 448).—The reduction of potassium permanganate in neutral solution by formic acid has been examined at 17° and 26.75°. It is shown that the disturbances observed in the reductions carried out in acid solution disappear in neutral solutions; these disturbances are observed down to acid concentrations 0.001*N*, and are greatest at the start of the action. The reduction in acid solutions is represented by the equations $2(\text{MnO}_4' + \text{HCO}_2' + \text{H}_2\text{O} = \text{MnO}_4'' + \text{CO}_3'' + 3\text{H}')$ and $2\text{MnO}_4'' + \text{HCO}_2' + 7\text{H}^+ = 2\text{Mn}(\text{OH})_4 + \text{CO}_2$. The temperature coefficient of the reaction is 1.88, a perfectly normal value which is very close to the value observed for the action in dilute acid solutions (*loc. cit.*).

J. F. S.

Antioxidants and Antioxygenisers. A. SEYEWETZ and P. SISLEY (*Bull. Soc. chim.*, 1922, [iv], **33**, 672—676).—A review of the subject in which reference is made to the work of Bigelow (A., 1898, ii, 506), Titoff (A., 1904, ii, 113), Lumière and Seyewetz (A., 1905, ii, 379), Sisley (*Rev. gén. mat. col.*, 1911, 337), and Moureu and Dufraisse. The suggestion is made that the term antioxygeniser (*antioxygène*) applied by the last named is less suitable than antioxidant (*antioxydant*) suggested by Lumière and Seyewetz.

H. J. E.

Catalysis with Special Reference to Newer Theories of Chemical Action. I. The Radiation Theory of Chemical Action. (1) Radiation and Chemistry. JEAN PERRIN. (2) The Radiation Hypothesis of Chemical Reactivity and some of its Applications. W. C. McC. LEWIS. (3) A Theory of Chemical Reaction and Reactivity. E. C. C. BALY. (4) Is a True Unimolecular Action Possible? T. MARTIN LOWRY (*Trans. Faraday Soc.*, 1922, **17**, 546—572, 573—587, 588—595, 596—597).—(1) A review of the author's radio-chemical theory of reactions

("Les Atomes," 1913; A., 1919, ii, 177), which is in accord with the laws of Arrhenius and of Planck. Experimental verification of the theory is derived from a consideration of the phenomena of photochemistry, luminescence, and organic fluorescence. The theory is extended to the phenomena of radioactivity.

(2) A review of the author's application of the quantum radiation hypothesis to the investigation of the mode whereby energy necessary to produce chemical change in catalytic or non-catalytic processes is communicated to the reactant unit (cf. T., 1914, 105, 2330; 1915, 107, 233; 1916, 109, 796; 1918, 113, 471; 1920, 117, 1120).

(3) Chemical reaction between atoms is conceived as consisting in the joint loss of an equal amount of energy by the reacting atoms, whereby a stable molecule is produced. The atoms in combination retain their individuality as absorbers or radiators of energy. The least possible amount of energy that the molecule can lose is a multiple of the least common integral multiple of the frequency quanta of its component atoms. The theory is confirmed by the phenomena observed in the photochemical combination of hydrogen and chlorine, and the possibility of the existence of two samples of a gas, for example, ammonia in apparent thermal equilibrium but in different molecular phases.

(4) The author contends that none of the actions to which the radiation hypothesis has been applied involves only a single molecule. The application of the hypothesis to cases of radioactive disintegration and bombardment by α -particles, where single molecules are involved, is unnecessary. The foundation of the radiation hypothesis is consequently doubtful.

J. S. G. T.

Catalysis with Special Reference to Newer Theories of Chemical Action. II. Heterogeneous Reactions. (1) **Chemical Reactions on Surfaces.** (2) **The Mechanism of the Catalytic Action of Platinum in the Reactions $2\text{CO} + \text{O}_2 = 2\text{CO}_2$, and $2\text{H}_2 + \text{O}_2 = 2\text{H}_2\text{O}$.** IRVING LANGMUIR (*Trans. Faraday Soc.*, 1922, 17, 607—620, 621—654).—(1) The author reviews his theory of adsorption at solid surfaces and the *modus operandi* of chemical reactions occurring thereat. Films adsorbed at solid surfaces are exceedingly stable, are unimolecular in thickness, and are oriented with regard to the adsorbing surface. Adsorption occurs as the result of a time lag between condensation and evaporation occurring at the surface. The orientation of the molecules on the adsorbing surface is a vital factor in determining the activity of the surface towards reacting gases. Differences in the geometrical arrangement of the atoms in the surface are responsible for the activation of catalysts effected by the action that occurs on them. Reactions between separate phases of constant composition occur only at the boundaries of phases.

(2) A detailed account is given of a theoretical and experimental investigation of the conditions determining the combination of (a) carbon monoxide and oxygen, and (b) hydrogen and oxygen, occurring at a smooth platinum surface. In the case of the former

reaction, for the range of temperatures $500\text{--}700^\circ\text{K.}$, the platinum surface is nearly completely covered by a unimolecular carbon monoxide film and reaction occurs only when carbon monoxide molecules strike oxygen atoms which have become adsorbed in the spaces left vacant by the evaporation of carbon monoxide molecules. For the range of temperatures $750\text{--}1050^\circ\text{K.}$, the surface is nearly covered with oxygen when there is an excess of oxygen, and the reaction velocity is limited by the rate at which carbon monoxide molecules strike the surface. With an excess of carbon monoxide, the surface is largely bare, and the reaction velocity is limited by the rate at which oxygen molecules strike the surface. In the case of the reaction between hydrogen and oxygen, at low temperatures ($300\text{--}600^\circ\text{K.}$), the reaction velocity depends on the previous treatment of the platinum. With relatively inactive platinum, the results resemble those obtained with carbon monoxide and oxygen, the reaction velocity being roughly proportional to the pressure of oxygen and inversely to that of the hydrogen. At temperatures between 700°K. and 1900°K. , the results correspond closely with those obtained with carbon monoxide and oxygen. At these temperatures, the reaction is not sensitive to the previous treatment of the platinum. Adsorbed oxygen atoms are very reactive towards hydrogen, whilst under certain conditions adsorbed hydrogen atoms are relatively inactive towards oxygen molecules.

J. S. G. T.

Induced Reactions and Negative Catalysis. N. R. DHAR and N. N. MITTRA (*Trans. Faraday Soc.*, 1922, **17**, 676—680).—The phenomenon of induced reaction studied by Dhar (T., 1917, **111**, 690) is shown to be of general occurrence. Negative catalysis, due probably to the formation of intermediate compounds, occurs in oxidation reactions when the catalyst is readily oxidisable. It is concluded that one chemical change will promote or induce another chemical change of the same type.

J. S. G. T.

Adsorption and its Bearing on Catalysis. MARCEL GUICHARD (*Bull. Soc. chim.*, 1922, [iv], **33**, 647—653).—A study of the effect of variation of surface on catalytic activity. In the case of silica, a diagram is given showing the different amounts of water vapour adsorbed at various temperatures and constant pressure by the catalyst after preliminary heating to different temperatures; increase in the temperature of preliminary heating decreases activity in adsorption. Similar results are obtained in the case of finely divided nickel and also for various solids in respect of iodine vapour. The author concludes that the conditions which tend to decrease the amount of vapour adsorbed by a powdered solid are the elevation of equilibrium temperature and the lowering of equilibrium pressure, both being reversible; further, the temperature and duration of heating to which the solid has been subjected beforehand produce an effect which is irreversible. Reasons are given for the inference that the preliminary heating results in diminution of the surface available for adsorption.

H. J. E.

Catalysis by Platinum Black. G. VAVON and A. HUSSON (*Compt. rend.*, 1922, 175, 277—279).—A platinum black catalyst which has become poisoned and inactive in the hydrogenation of a particular substance is not necessarily inactive towards another substance, and experiments were undertaken to determine the point at which absorption of hydrogen ceased with various unsaturated substances when the catalyst was progressively poisoned by the gradual addition of small quantities of carbon disulphide. This limiting point was found to vary with the nature of the substance, with the nature of the solvent, with the quality of the platinum, and with the amount of catalyst used, but is independent of the concentration of the solution. In one instance, using 5.5 grams of substance, 50 c.c. of alcohol, and 0.2 gram of platinum, the amount of carbon disulphide required to inhibit hydrogenation was 1.1 mg. for cyclohexene, 0.8 mg. for nitrobenzene, 0.5 mg. for cinnamic acid, and 0.4 mg. for acetophenone, so that a catalyst which was inactive towards acetophenone would still hydrogenate nitrobenzene or cyclohexene. The author advances a physical explanation of the phenomenon, based on the adsorption of hydrogen by the metal.



G. F. M.

Rôle of Protective Colloids in Catalysis. II. THOMAS IREDALE (T., 1922, 121, 1536—1542).

Catalytic Addition of Hydrogen. Influence of Oxygen on the Catalyst. W. NORMANN (*Ber.*, 1922, 55, [B], 2193—2197).—Working under conditions which are considered to exclude the presence of oxygen and with hydrogen which is free from any trace of the latter, the author has been able to effect the hardening of cotton-seed oil in the presence of nickel (prepared by the reduction of the chloride or cyanide), nickel powder, or palladised kieselguhr. It appears therefore that Willstätter's assumption of the necessity for the presence of oxygen is invalid.

H. W.

Ion Activities in Homogeneous Catalysis. Formation of *p*-Chloroacetanilide from *N*-Chloroacetanilide. HERBERT S. HARNED and HARRY SELTZ (*J. Amer. Chem. Soc.*, 1922, 44, 1475—1484).—From theoretical considerations, it is deduced that in reactions in liquid systems catalysed by specific ions, it is the activities of such ions which determine the velocity of the reaction at any temperature when the catalysis depends on successive states of equilibria. The velocity constants of the conversion of *N*-chloroacetanilide into *p*-chloroacetanilide catalysed by hydrochloric acid have been determined at 17.65°, 25°, and 35° for concentrations of acid from 0.1*N* to 1.0*N*. This reaction according to Orton and Jones (*Brit. Assoc. Rep.*, 1910, 85) takes place thus:



reaction (1) is measurably slow, whilst reaction (2) is very rapid, so that the system approximates to a continuous series of successive equilibria and fulfils the conditions stated above. It is shown that the velocity constants at each temperature are proportional

to the product of the activities of the hydrogen and chlorine ions of the catalysing acid. The present work is the first case where homogeneous catalysis has been calculated with exactitude over a wide range of concentration and in concentrated solutions, without uncertainties arising in liquid junction potentials. The temperature coefficient has been considered and the critical increment roughly calculated and shown to vary considerably with rise of temperature. The relationship $K_T = k/\alpha_H \alpha_{Cl}$, which holds exactly for solutions of hydrochloric acid alone, fails when acid-salt mixtures are used as catalysts. J. F. S.

A Possible Reconciliation of the Atomic Models of Bohr and of Lewis and Langmuir. W. HUGHES (*Nature*, 1922, 110, 37—38).—If it is assumed that the electron shells are fixed and the nucleus rotates on an axis, a model is obtained which, when viewed with respect to the electron shells, is precisely the same as the Lewis-Langmuir model, with its very satisfactory representation of the mechanism of chemical combination; when viewed with respect to the whole atom, it possesses all the merits of Bohr's model, yielding an accurate explanation of the reaction of atoms and molecules with radiation. Further, the existence of isotopy can be predicted with its aid, for if the nucleus of a given atom possesses more than one stable axis of rotation with respect to itself, or to its surrounding shells of electrons, or to both, and these axes are associated with different amounts of energy, it is possible for the mass of the atom to be different for the different positions of the nuclear axis, since by the theory of relativity, energy possesses mass. A. A. E.

Atomic Structure. MAURICE L. HUGGINS (*Science*, 1922, 55, 459—460).—As an extension of Lewis's (A., 1916, ii, 310) theory of atomic structure, it is assumed that, although the number of electrons in each shell of the lighter atoms is the same as in the original theory, the fifth, sixth, seventh, and eighth electrons in the second and third shells pair with the first four, the distance between the electrons in each of these pairs, and also in each pair formed by bonding between atoms, being much less than the distance between pairs. The shells are therefore tetrahedra of pairs instead of cubes of single electrons. The electrons in each shell after the second tend to be placed opposite to the centres of the faces of the imaginary polyhedron formed by the electron groups in the underlying shell. When the nuclear charge becomes sufficiently great, the same forces which cause pairing of electrons in nitrogen result in the formation of triplets in the inner shells of the heavier atoms. As one after another of the outer electrons is drawn into an inner shell to form triplets, the remaining pairs are pushed further from the nucleus, with possible rearrangement of the kernel structure. The theory is applied in a number of instances. A. A. E.

The Quantum Mechanism in the Atom. E. T. WHITTAKER (*Proc. Roy. Soc. Edin.*, 1922, 42, 129—142).—Ewing's model of ferromagnetic induction (*Proc. Roy. Soc. Edin.*, 1922, 42, 30) is employed to investigate the mechanism within the atom which

compels all exchanges between kinetic energy and radiant energy to conform with the equation $U=h\nu$. In this model, a system of magnets is rigidly connected like the spokes of a wheel, having poles of the same name at the circumference and poles of a contrary sign at the centre of the circle. When an electron approaches such a model in the direction of the axis of the wheel, a rotation of the poles occurs which constitutes a magnetic current. The kinetic energy of the electron is expended in setting the magnetic structure of the atom in motion. When the velocity of the electron exceeds a critical value $2eM/\sqrt{Am}$, the magnetic structure is penetrated; M and A are quantities dependent on the atomic structure with which the electron collides. When its energy is less than this quantity, it is repelled without permanent loss of energy, the encounter being perfectly elastic. On penetrating the atom, the electron gives up $2e^2M^2/A$ of its energy and leaves the system with the remainder. This absorbed energy, which appears in the atom as a magnetic current, does not depend on the mass of the electron, but on its charge and on quantities dependent only on the atomic structure. This rotational energy is transformed into a radiant form by means of Hertzian oscillators. This view of the mechanism leads to the Planck equation $h\nu=U$. The emission of an electron already in the atom is a converse process and thus the equations may be applied to explain photo-electric phenomena. The connexion with the Bohr theory of series spectra is developed.

W. E. G.

The Quantum Mechanism in the Atom. (SIR) J. ALFRED EWING (*Proc. Roy. Soc. Edin.*, 1922, 42, 143—146).—A discussion of the mechanism suggested by Whittaker (see preceding abstract) for the conversion of the rotational energy of the wheel atom into radiation. There are two magnetic systems in his model of the atom: a central one forming the wheel, and another around it, which may be called the ring. An electron passing through the system produces relative angular displacement of the two parts of the atom, and magnetic forces are set up which tend to restore the whole to the original configuration. Thus oscillations are set up which expend their energy in the emitted radiation.

W. E. G.

The Quantum Mechanism in the Atom. R. A. HOUSTOUN (*Proc. Roy. Soc. Edin.*, 1922, 42, 221—222).—The Whittaker model is checked numerically. A calculation of the radius of the magnetic orbit gives values in satisfactory agreement with the dimensions of the hydrogen molecule. It is shown that the Planck constant comes out as the product of the unit electric charge and a unit magnetic quantity. It thus has the correct dimensions.

W. E. G.

The Importance of Whittaker's Atomic Model and of other Atomic Models. W. PEDDIE (*Proc. Roy. Soc. Edin.*, 1922, 42, 223—224).—A general discussion of the Whittaker model pointing out the inadequacy of the simply constituted atom of the electronic type.

W. E. G.

Sizes of Atoms in Diamond Type Crystals. ROBERT N. PEASE (*J. Amer. Chem. Soc.*, 1922, 44, 1497—1498).—On the basis of the results published in a recent paper (this vol., ii, 428), the author has recalculated the atomic radii of several elements of the rare gas type crystallising in the diamond type lattice. The following values are recorded: diamond 0.77 Å.U., silicon (S, Cl) 1.17 Å.U., germanium (Cu, Zn, Br) 1.22 Å.U., grey tin (Ag, I) 1.40 Å.U. The following interatomic distances are calculated: silicon carbide, 1.94 (1.90, 1.95); zinc sulphide, 2.39 (2.35, 2.35); cuprous chloride, 2.39 (2.38, 2.43); cuprous iodide, 2.62 (2.63, 2.78); silver iodide, 2.80 (2.83, 3.18); and cuprous bromide, 2.44 (2.52, 2.57). The values are given in Ångström units, and for comparison purposes the experimental value and the value from Bragg's atomic radii are included in brackets, the former value being placed first.

J. F. S.

The Properties of Elements and Salts as Related to the Dimensions of Atoms and Ions. GEORGE L. CLARK (*Science*, 1922, 55, 401—405).—It is shown that a considerable number of the properties of elements and salts are linearly related to the dimensions of the component atoms and ions. The relations are more general than those found by Biltz (A., 1921, ii, 487); thus it is found that if the atomic volumes of the five alkali elements are plotted on one axis (as the X -axis) of a three-dimensional rectangular co-ordinate system, the atomic volumes of the four halogens on the Z -axis, and the molecular volumes of the alkali haloids on the Y -axis, all the points (except some for caesium, the anomaly of which is attributed to difference in crystal-lattice form) lie approximately on a plane $z = bx + cy + d$ which passes nearly through the origin. When other properties are plotted on the Y -axis, in a very large number of cases the surface, whilst not a plane, is of a simple form ("doubly ruled surface"). The surfaces may commonly be expressed by the equation $z = axy + bx + cy + d$. By using this method, a number of possible new linear relationships may be predicted. Values already ascribed to the atomic and ionic radii of the halogens and alkali elements are considered, Henglein's procedure being regarded as questionable.

A. A. E.

Atomic Radii. I. MAURICE L. HUGGINS (*Physical Rev.*, 1922, 19, 346—353).—Bragg's (A., 1920, ii, 537) "atomic spheres" and "atomic radii" may, when considered from the point of view of Lewis's theory of atomic and molecular structure (A., 1916, ii, 310), be given a definite physical meaning. The "atomic sphere" of an atom is its valency shell, and its "atomic radius" is the distance from the atomic centre to a group of electrons (usually an electron-pair) in the valency shell. It is shown that Bragg's assumption, that the distance between two adjacent atomic centres is equal to the sum of their radii, is often not valid, since in many cases the group of electrons constituting the bond does not lie on the line joining the atomic centres; that is to say, the two "atomic spheres" overlap. Eight causes of variation of the atomic radius

of an element are considered, namely: (1) some of the valency pairs may not act as bonds, (2) a valency electron-pair may be forced away from its normal position in the valency shell, (3) widely varying pulls are exerted on electron-pairs by atoms of different elements, (4) an atom of the same element in different structures may pull an electron-pair away a different distance from the nucleus of the atom to which it is linked, (5) if an atom has a different number of electron-pairs in its valency shell in two crystals, their distances from the nucleus would not be expected to be the same, (6) valency electron-pairs are pulled out from the nucleus to a different extent, according to whether they are on the centre-lines or not, (7) many atoms are capable of more than one arrangement of electrons in that part of the atom inside the valency shell, (8) the number of electrons constituting a bond may be more than two.

A. A. E.

Structure of the Ions of the Rare Earths. H. G. GRIMM (*Z. physikal. Chem.*, 1922, **101**, 403—409).—A theoretical paper in which, on the basis of Bohr's arrangement of the electrons in xenon and niton, an attempt is made to draw conclusions on the structure of the ions of the rare earths, quinquivalent tantalum, sexavalent tungsten, and octavalent osmium ions which shall be in keeping with the chemical properties of these elements. It is shown that the outside sheath of the ions of the rare earths contains in all probability the same number of electrons, namely, eight or nine. The molecular volume and basicity series of the rare earths has been used to draw conclusions on the series of ionic radii (cf. A., 1921, ii, 127).

J. F. S.

Periodic System of the Atomic Ions. H. G. GRIMM (*Z. physikal. Chem.*, 1922, **44**, 410—413).—The author has constructed a table on the basis of the structure of the ions of the elements. This divides the elements into six groups, namely, those with no electrons in the outer sheath, those with two electrons in the outer sheath, those with eight and eighteen electrons, respectively, in the outside sheath. The last three groups are composed of elements which are built up of completed sheaths, that is, sheaths occupied by the same number of electrons as in sheaths of the inactive gases. The fifth group contains elements with an incomplete outside sheath, and the sixth group contains elements with an incomplete inner sheath but a constant outside sheath. The elements of the sixth group are those of the rare earths. In the table the anions lie to the left of the inactive gases and the kations to the right. In this arrangement several elements, such as Cl^- , Cl^{+++} , Se^{--} , and Se^{+++} , occur in two positions.

J. F. S.

Structure of some Gaseous Molecules of which Hydrogen is a Constituent. A. O. RANKINE (*Trans. Faraday Soc.*, 1922, **17**, 719—727).—From viscosity measurements the following values have been deduced for the respective magnitudes (expressed in $\text{cm}^2 \times 10^{-15}$) of the molecular "mean collision areas" of certain gaseous compounds containing hydrogen: HCl , 0.676; HBr ,

0.763; HI, 0.926; H_2S , 0.773; NH_3 , 0.640; PH_3 , 0.911; AsH_3 , 0.985; CH_4 , 0.772. The corresponding values for the inert gases have been recalculated and found to be as follows: Ne, 0.417; Ar, 0.648; Kr, 0.757; Xe, 0.915. Adopting the Lewis-Langmuir view of the mechanism of combination between hydrogen atoms and other elements, and assuming, on the basis of W. L. Bragg's X-ray measurements (A., 1920, ii, 537), that the atoms towards the end of each period of the periodic table are equal in size, it is shown to be probable that as the number of hydrogen atoms in the molecule increases, their nuclei become more remote from the nucleus of the central atom. This retreat of the hydrogen nuclei is due to their mutual repulsion, and eventually leads to the failure of the formation of molecules, otherwise possible, such as BH_5 and AlH_5 . The relative dimensions of CH_4 and Kr agree to within the limit of probable experimental error. It is to be anticipated, therefore, that NH_4 and Rb, which are correspondingly related to CH_4 and Kr, respectively, would occupy nearly equal spaces. This is confirmed from crystallographic data.

J. S. G. T.

Arabic Chemistry. E. J. HOLMYARD (*Nature*, 1922, 109, 778—779).—An historical note ascribing to Maslama al-Majriti the authorship of the section on chemistry in the "Letters" of the Brethren of Purity (10th century A.D.).

A. A. E.

[**Lecture Experiment.**] **Time Reaction.** MARTIN MEYER (*J. Amer. Chem. Soc.*, 1922, 44, 1498—1500).—A time reaction of the same type as that recently described by Forbes, Estill, and Walker (this vol., ii, 271) which is suitable for a lecture experiment is described. For example, 40 c.c. of *M*-sodium thiosulphate, 20 c.c. of 1.67*M*-potassium hydroxide containing 0.67*M* of potassium sodium tartrate and 44.46 grams of antimony trioxide per litre, are mixed and 10—15 c.c. of 4*N*-hydrochloric acid added. After an induction period of thirty to sixty seconds, a white turbidity appears which changes to the characteristic orange colour of antimony trisulphide. Variations of the quantities give different induction periods.

J. F. S.

Inorganic Chemistry.

New Revision of the Density of Oxygen Gas. E. MOLES and M. CRESPI (*Anal. Fis. Quím.*, 1922, 20, 190—192; cf. Moles and González, this vol., ii, 497).—Oxygen prepared from potassium permanganate contains traces of carbon dioxide and ozone and purification by phosphoric oxide is insufficient. Determinations on oxygen from potassium permanganate from which carbon dioxide and ozone had been removed by soda-lime and mercury, respectively,

gave the value (calculated for lat. 45°) 1.42895 for the density of the gas. This is in good agreement with the value previously given (*loc. cit.*).
G. W. R.

Ozone. E. H. RIESENFELD and G. M. SCHWAB (*Ber.*, 1922, 55, [B], 2088—2099).—The main difficulty in the preparation of pure ozone and the determination of its physical constants consists in its explosiveness. The recent advances in micro-analytical methods have enabled the authors to perform the necessary experiments with such small quantities of material that explosions are avoided or are not dangerous.

Dry oxygen is ozonised in the usual manner and the product condensed in small glass bulbs with long capillaries immersed in liquid air (the apparatus is figured and fully described in the original communication). Since the condensate is always richer in ozone than the supernatant gas, the latter is pumped away after each condensation and the vessel again filled with ozonised oxygen. When in this manner a dark blue condensate (solution of oxygen in ozone) has been obtained, the subsequent condensates consist of pale blue solutions of ozone in oxygen; when the exhaustion is renewed, the oxygen evaporates and the ozone passes into the first phase. The dark blue product still contains about 30% of oxygen, but its removal can be fairly readily effected by fractionation, since the boiling points of oxygen and ozone differ by about 70° . At the temperature of liquid air, the vapour pressure of liquid ozone is practically negligible so that the glass bulbs can be evacuated completely and the capillaries sealed off without danger. Analysis of the product is effected by breaking one of the bulbs containing a known weight of substance under potassium iodide solution containing boric acid, measurement of the volume of inactive oxygen, and estimation of that of active oxygen by titration with sodium thiosulphate. Within the limits of experimental error, the volume relationship of active to inactive oxygen is 1 : 2 and the purity of the ozone is controlled further by a micro-estimation of the molecular weight by Dumas's method. The following physical constants have been determined (the methods and requisite apparatus are fully described and figured in the original); m. p. -250° , b. p. -112.3° , critical temperature -5° , $d_{-182}^{25} 1.784$. Liquid ozone is not completely miscible with liquid oxygen at all temperatures, the critical temperature of solubility lying at -158° .

Very discordant results have been obtained during investigations of the thermal production of ozone from oxygen. According to the Nernst theorem, a minimal temperature of 4000° Abs. is necessary for the production of appreciable quantities of ozone, whereas Fischer has observed considerable formation of the gas at glowing Nernst filaments at a much lower temperature. The temperature attained in the explosion of pure ozone is calculated to be about 4000° Abs., and under these conditions a distinct odour of ozone is observed. On the other hand, no trace of ozone could be perceived in explosions of mixtures of ozone and oxygen

with which a temperature of more than 2000° Abs. was obtained. A purely thermal production of ozone is not involved, therefore, in Fischer's experiments.

The explosive decomposition of ozone is very sensitive to catalytic influences, but if these are excluded the gas is found to be unexpectedly stable. According to Warburg's calculation, the half life period of pure ozone at 16° is one hundred and sixty-seven hours, whereas the authors find that under certain conditions a period of several weeks elapses before decomposition has proceeded to this extent.

Violent changes of pressure cause the explosion of ozone in all states of aggregation. An explosion when the solid or liquid material is lightly touched or on sudden solidification of the liquid has not been observed.

The authors consider that their work brings conclusive evidence against the existence of oxozone. H. W.

The Oxidising and Reducing Properties of Sulphur Dioxide. I. Mercury Chlorides. LACHLAN MACQUARIE STEWART and WILLIAM WARDLAW (T., 1922, 121, 1481—1489).

The Physical Properties of Sulphur Trioxide. A. BERTHOUD (*Helv. Chim. Acta*, 1922, 5, 513—532).—Doubt is thrown by the author on the existence of sulphur trioxide in the so-called α - and β -forms. The fact that the two kinds can exist together indefinitely, with other observations of a similar character, indicates that the silky crystals, the so-called β -form, are a product of hydration. The compound, if compound it be, must be, however, of a unique type, containing something like a thousand or more mols. of sulphur trioxide to one of water. A number of physical properties of sulphur trioxide were determined with the greatest care. It has m. p. $16.85^{\circ} \pm 0.02$, agreeing closely with Lichty's result, 16.79° (A., 1912, ii, 1164) and b. p. $44.52^{\circ}/760$ mm. The vapour pressure curve, as deduced from determinations at 15 points between 24° and 47.8° , is represented by $\log p = -2314/T + 10.17$. The critical pressure and temperature were determined, using a modification of Pellaton's method (cf. A., 1916, ii, 245), and the critical temperature by direct observation of the disappearance and reappearance of the meniscus. The results are, $p_c = 83.8$ atm., $t_c = 218.3^{\circ}$. The density was determined at temperatures from 17° to 55° by a pycnometer method, and from 98° to 214° by a modification of Young's method (T., 1891, 59, 37), which gives the density of both liquid and vapour. The critical density, d_c , is 0.633. The surface tension was determined by the method of Ramsay and Shields, the results being, at 19° , 44.9° and 78.0° , respectively, $\gamma = 34.17$, 29.47 , and 22.63 . The molecular heat of vaporisation, calculated from the vapour tension curve, is 10,300 cal. The quotient L/T , Trouton's ratio, is surprisingly high, 32.5, the normal for liquids of similar b. p. being 21.0; the high value indicates association in the liquid form. The value of Eötvös's surface tension coefficient also indicates association of the liquid at lower temperatures, but at 78° the value is nearly

normal. The values of van der Waals's constants are : $a=0.01629$;
 $b=0.002684$. E. H. R.

Preparation of Selenium Dioxide. JULIUS MEYER (*Ber.*, 1922, 55, [B], 2082—2084).—Selenium in quantities of 60—75 grams is heated to its melting point in a porcelain boat placed in a wide hard glass tube; a very rapid current of oxygen which has been passed through fuming nitric acid is passed over it, causing the molten selenium to burn with a brilliant blue flame and to give a sublimate of selenium dioxide the purity of which increases with increasing rate of the supply of oxygen. The product, which contains small quantities of oxides of nitrogen, is purified by being sublimed in the same tube in a current of pure oxygen. To avoid loss of material, the tube is connected with a doubly tubulated vessel of two litres capacity in which the final traces of the dioxide are deposited. The combustion of 60—70 grams of selenium can be effected in about one and a quarter hours.

The oxidation of selenium appears to be greatly affected by catalytic influence, acidic substances causing acceleration whereas alkaline materials (including glass) cause retardation. H. W.

Active Hydrogen and Nitrogen. GERALD L. WENDT (*Nature*, 1922, 109, 749).—Newman's (this vol., ii, 279) failure to obtain a test for nitrides when sulphur, phosphorus, and iodine are treated with active nitrogen is not evidence of the absence of chemical reaction; in the cases of sulphur and phosphorus, the formation of sulphides and phosphides has been demonstrated by the author, whose experiments also show that phosphine and hydrogen sulphide are formed when phosphorus and sulphur are exposed to active hydrogen. A. A. E.

Active Hydrogen and Nitrogen. F. H. NEWMAN (*Nature*, 1922, 109, 749; cf. Wendt, preceding abstract).—The absorption of active nitrogen by sulphur, phosphorus, and iodine probably results in chemical combination, since the absorbed gas is not liberated on heating; in the case of active hydrogen, however, absorption occurs at temperatures above 0°. This absorption may be due in part to chemical action, but other processes, such as occlusion, have to be taken into account. A. A. E.

Reaction between Cathodic Hydrogen and Nitrogen at High Pressures. J. N. PRING and E. O. RANSOME (*Trans. Faraday Soc.*, 1922, 17, 689—694).—When hydrogen is liberated electrolytically at a cathode in contact with nitrogen, particularly at high pressures, the conditions would appear to be favourable to the synthesis of ammonia. A solution of sulphuric acid was electrolysed in the presence of nitrogen at pressures ranging from 1 to 500 atmospheres. At atmospheric pressure, the mean percentage yield of ammonia by direct union of the elements was 0.04%. At pressures from 60 to 104 atmospheres, the yield of ammonia was 0.09%. For pressures ranging from 300 to 500 atmospheres, no ammonia was synthesised. The results indicate

that no reaction occurs between nitrogen and cathodic hydrogen, the small yield of ammonia at the lower pressures being due to thermal action.

J. S. G. T.

Hydroxylamine. CARL KJELLIN (*Svensk Kem. Tidskr.*, 1921, **33**, 213—228).—In the reaction expressed by the equation: $\text{NH}_2\cdot\text{OH} + 2\text{Fe}(\text{OH})_2 + \text{H}_2\text{O} = \text{NH}_3 + 2\text{Fe}(\text{OH})_3$, the experimental results are in practically complete accord when the potassium hydroxide in the solutions does not exceed 1.75%. The reactions between hydroxylamine and stannous hydroxide and manganese hydroxide were also examined; the former yields ammonia in moderate agreement with theory, whilst in the latter case the amount of ammonia produced is negligible. The change undergone when hydroxylamine is subjected to heat in presence of aqueous potassium hydroxide appears to be expressed by the equation: $3\text{NH}_2\cdot\text{OH} = 3\text{H}_2\text{O} + \text{NH}_3 + \text{N}_2$. Compounds of the type $\text{CH}_2\text{R}\cdot\text{NH}\cdot\text{OH}$ ($\text{R} = \text{Me}$, Et , Pr) when hydrolysed in concentrated hydrochloric acid or potassium hydroxide, evolved up to 75% of the expected yield of ammonia. It is probable that *N*-isopropylhydroxylamine and bromine react in accordance with the scheme: $\text{CHMe}_2\cdot\text{NH}\cdot\text{OH} \rightarrow \text{CHMe}_2\cdot\text{NBr}\cdot\text{OH} \rightarrow \text{CHMe}_2\cdot\text{NO} \rightarrow \text{CMe}_2\cdot\text{NOH}$. *N*-Ethylhydroxylamine hydrochloride forms long, colourless, hygroscopic crystals, m. p. 37°.

CHEMICAL ABSTRACTS.

Crystal Structure of Phosphonium Iodide. ROSCOE G. DICKINSON (*J. Amer. Chem. Soc.*, 1922, **44**, 1489—1497).—The structure of the crystals of tetragonal phosphonium iodide has been investigated by means of X-rays, using chiefly Laue photographs. It is shown that the X-ray data are satisfactorily accounted for by a structure obtained by placing in a unit cell of the dimensions $6.34 \times 6.34 \times 4.62$ Å.U. phosphorus atoms at (000) and ($\frac{1}{2}\frac{1}{2}0$) and iodine atoms at ($0\frac{1}{2}u$) and ($\frac{1}{2}0\bar{u}$) where u has a value very close to 0.40, and it is shown that no simpler structure is capable of accounting for the data. The close relationship between this structure and the low temperature form of ammonium chloride is demonstrated.

J. F. S.

Hypophosphorous Acid. IV. Its Reaction with Cupric Chloride. ALEC DUNCAN MITCHELL (*T.*, 1922, **121**, 1624—1638).

The Volatilisation of Arsenic and Antimony by means of Methyl Alcohol. L. DUPARC and L. RAMADIER (*Helv. Chim. Acta*, 1922, **5**, 552—556).—By passing a current of air through a solution of arsenious or antimonious oxide in concentrated hydrochloric acid containing a suitable proportion of methyl alcohol, the whole of the arsenic or antimony can be carried over into an absorbing solution. In this way, 0.1 gram of arsenious oxide can be volatilised in one hour at 55° in 45 c.c. of methyl alcohol. In the case of antimonious oxide the vaporisation can be completely prevented by diluting the hydrochloric acid used with an equal volume of water. The method may be applicable for

separating arsenic and antimony from other metals and from each other.

E. H. R.

Revision of the Atomic Weight of Boron. Analysis of Boron Trichloride. O. HÖNIGSCHMID and L. BIRCKENBACH (*Anal. Fis. Quím.*, 1922, 20, 167—173; cf. A., 1921, ii, 646).—From determinations of the ratio $\text{BCl}_3 : 3\text{Ag}$ and $\text{BCl}_3 : 3\text{AgCl}$, following the methods used in the authors' revision of the atomic weight of bismuth (*loc. cit.*), the values of the atomic weight of boron obtained for three samples of specially purified boron trichloride were 10.840, 10.818, and 10.825, respectively. The first value is rejected and the value 10.82 is taken. It is pointed out that this value is in better accord with the theory of isotopes than the hitherto accepted value of 10.90.

G. W. R.

The Action of Diamond on Carbon Monoxide. FOIX (*Bull. Soc. chim.*, 1922, [iv], 33, 678—679).—An attempt to carry out the reaction $\text{CO}_2 + \text{C} \rightleftharpoons 2\text{CO}$ in the case of the diamond at temperatures above 1100° resulted in the deposition of amorphous carbon on the surface of diamond without the latter undergoing any change. The conclusion is drawn that the time of experiment, three hours, was insufficient for the attainment of equilibrium.

H. J. E.

The Preparation of Carbon Suboxide on a Larger Scale and the Properties of Pure Carbon Suboxide. ERWIN OTT and KARL SCHMIDT (*Ber.*, 1922, 55, [B], 2126—2130).—Diacetyl-tartaric anhydride is heated to its boiling point in a flask the neck of which is connected to a cylinder in which an electrically heated platinum wire is suspended. The upper portion of the cylinder is connected through a series of condensing vessels to a powerful pump which, in spite of the evolution of carbon monoxide, maintains a pressure of about 11 mm. (The apparatus is fully figured and described in the original.) The decomposition of one hundred and eighty-nine grams of anhydride can be effected in six to eight hours, the yield of carbon suboxide being 41% of that theoretically possible. From this point of view the process is superior to the improved method of Stock and Stoltzenberg (A., 1917, ii, 308), which depends on the action of phosphoric oxide on malonic acid, and has the further advantage of greater cheapness. In addition, the carbon suboxide obtained in this manner can be preserved unchanged (except for a slight darkening of colour which is probably due to incompleteness in the exclusion of moisture) for long periods, whereas that prepared with the help of phosphoric oxide rapidly becomes polymerised, probably owing to the presence of traces of phosphorous oxide in the latter (cf. Manley, T., 1922, 121, 331).

H. W.

The Crystal Structure of Quartz. MAURICE L. HUGGINS (*Physical Rev.*, 1922, 19, 363—368).—The suggested structure of quartz, obtained by means of the Lewis theory, is in accord with Bragg's conclusions. Each silicon atom is surrounded by four pairs of electrons at tetrahedron corners, which act as bonds

connecting it to four equidistant oxygen atoms. Each oxygen atom is also surrounded by four tetrahedrally oriented electron-pairs, two of which serve as bonds connecting the oxygen to silicon atoms. The crystal is thus not composed of silica units, but is a single molecule.

A. A. E.

Solubility of Helium in Water. HAMILTON P. CADY, HOWARD M. ELSEY, and EMILY V. BERGER (*J. Amer. Chem. Soc.*, 1922, **44**, 1456—1461).—The solubility of helium has been determined at a series of temperatures lying between 2° and 30°, using material which was spectroscopically pure. The following values of the absorption coefficient are recorded: 2°, 0.00938, 10°, 0.00895, 25°, 0.00861, and 30°, 0.00817. Plotting these values shows that there is no indication of the solubility passing through a minimum as has been repeatedly stated. The values are all smaller than those of Estreicher (A., 1900, ii, 205) and Antropoff (A., 1919, ii, 511), and larger than the single determination of Ramsay (T., 1895, 67, 697).

J. F. S.

The Preparation of Sodium Hydrogen Carbonate. ER. TOPODESCU (*Compt. rend.*, 1922, **175**, 268—270).—A study similar to that originally made at 15° (this vol., ii, 375) of the equilibrium of the four salts, sodium chloride, sodium hydrogen carbonate, ammonium chloride, and ammonium hydrogen carbonate with their saturated solutions has now been made at 35° and 50°. The geometric representation of the results obtained by means of a Le Chatelier diagram enables a calculation to be made of the theoretical yield, that is to say, the proportion of sodium chloride transformable into crystalline sodium hydrogen carbonate for a solution of any given initial composition.

G. F. M.

The Preparation of Ammonium Chloride at Low Temperatures. PAUL MONDAIN-MONVAL (*Compt. rend.*, 1922, **175**, 162—164).—The conditions governing the crystallisation of ammonium chloride at 0° from solutions containing in addition one or more of the salts sodium chloride, sodium carbonate, and ammonium carbonate, were studied in a similar manner to those previously described for 15°, and a Le Chatelier diagram is given showing the surfaces of saturation. The zone of crystallisation of sodium carbonate shows two distinct areas, one corresponding with the crystallisation of $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$, and the other with a hydrate of the formula $2\text{Na}_2\text{CO}_3 \cdot 5\text{H}_2\text{O}$, the existence of which, although disputed by Wegscheider (A., 1912, ii, 156), is now confirmed.

G. F. M.

A Preliminary Attempt to Transmute Lithium. RALPH W. G. WYCKOFF (*Science*, 1922, **55**, 130—131).—Inconclusive results were obtained when lithium or its salts was bombarded with a stream of electrons, in the hope of introducing one or two electrons into the nucleus, and detecting spectroscopically the formation of helium or hydrogen.

A. A. E.

Electrolytic Preparation of Calcium Amalgam. B. S. NEUHAUSEN (*J. Amer. Chem. Soc.*, 1922, **44**, 1445—1447).—The author discusses the methods which have been adopted previously for the preparation of calcium amalgam (Smith and Bennett, *A.*, 1909, ii, 663; 1910, ii, 500), and devises a more efficient method of preparing this material. The method consists in just covering the bottom of a crystallising dish, 25 cm. diam., with a layer of mercury which serves as cathode, and then filling the dish with a 1.75*N*-solution of calcium chloride. An anode of platinum foil (2×4 cm.) is placed in the solution parallel to the mercury surface at a distance of 4 cm., and a current of 3.5 amperes at 4.6 volts passed for thirty minutes. This arrangement prevents heating and foaming. After about thirty minutes, the generation of hydrogen sets in, but up to this point the solution remains neutral to litmus and there is no formation of a black powder which in other methods always forms and catalyses the decomposition of the amalgam. The amalgam is washed by allowing it to fall in a fine stream into two litres of distilled water. The product prepared in this way contains 0.069—0.075% of calcium, and is preserved in a bottle filled with carbon dioxide. J. F. S.

The Crystal Structures of Aragonite (CaCO₃) and Related Minerals. MAURICE L. HUGGINS (*Physical Rev.*, 1922, **19**, 354—362).—A proposed structure for aragonite is such that, as in calcite, each carbon atom is linked by double bonds to three oxygen atoms, each oxygen atom to two calcium atoms and one carbon atom, and each calcium atom to six oxygen atoms, at the corners of an irregular octahedron. The observed cleavage of aragonite is in accord with the principles that (a) cleavage tends to occur so as to leave the two new crystal surfaces electrically neutral, (b) where some bonds are weaker than others, cleavage will take place in such a way as to rupture the weaker bonds in preference to the stronger ones, (c) all bonds being equally strong, cleavage will occur between the planes connected by the fewest bonds per unit area. The structure of aragonite, and those of the isomorphous minerals strontianite, witherite, and cerussite, conform to the author's theory that the electron groups in an atomic shell tend to place themselves opposite to the faces of the imaginary polyhedron formed by the electron groups in the next underlying shell. A. A. E.

Dispersoid Chemistry of Gypsum. II. H. NEUGEBAUER (*Kolloid Z.*, 1922, **31**, 40—45).—The viscosity of suspensions of anhydrite, gypsum, burnt gypsum, dehydrated gypsum, anhydrite, and leucolith has been measured at various intervals of time after preparation, with the object of investigating the mechanism of the hydration of calcium sulphate. It is shown that in opposition to the statements of van't Hoff the preparations used in the present work could not be completely dehydrated at 115° or slightly higher. The water of crystallisation was never reduced to less than 4%. The common dihydrate lost its crystal water at 102° considerably more slowly and less completely than the other forms,

probably because of the size of the crystals. The viscosimetric investigation of the commercial "van't Hoff gypsum" gave very indefinite curves, whereas van't Hoff found extremely rapid combination with water and rapid hardening for this material. The formation of soluble anhydrite does not come into the question here. Gypsum prepared by slowly dehydrating precipitated calcium sulphate gave a normal combination curve, and the dihydrate gave curves analogous to those obtained with insufficiently dehydrated gypsum. A commercial product, anhydrite, prepared by the dehydration of natural anhydrite, exhibits, on dehydration, a behaviour analogous to that shown by "van't Hoff gypsum." Its viscosity curve is more pronounced than that of commercial gypsum obtained in the usual way, but less steep than that of precipitated gypsum. Towards the addition of alum it behaves in the same way as plaster of Paris. A further technical product, leucolith, showed no combination with water from its viscosity measurements. The behaviour in this case was analogous to that of estrich gypsum (flooring cement), the setting in these cases being due to the presence of free lime.

J. F. S.

The Constitution of Strontium-Lead Alloys. E. PIROWSKY (*Z. Metallk.*, 1922, **14**, 300—301).—Lead and strontium form one compound, Pb_3Sr , melting at 676° and containing 12.35% Sr, and a eutectic, containing very small amounts of strontium, which melts at the same temperature as lead. Strontium is completely insoluble in solid lead. [Cf. *J. Soc. Chem. Ind.*, 1922, Sept.]

A. R. P.

Barium Sulphuric Acid and Barium Selenic Acid. JULIUS MEYER and WALTER FRIEDRICH (*Z. physikal. Chem.*, 1922, **101**, 498—503).—The solubility of barium sulphate in 97.86% sulphuric acid has been determined at 25° and found to be 14.91 grams in 100 c.c. of acid. The saturated solution is shown to contain barium sulphuric acid, $\text{H}_2[\text{Ba}(\text{SO}_4)_2]$, and by electrolytic transport this compound can be accumulated in the anode compartment, where it crystallises. Dissolving barium selenate in selenic acid gives rise to a similar compound, barium selenic acid, $\text{H}_2[\text{Ba}(\text{SeO}_4)_2]$, which can similarly be crystallised by electrolytic transport. A concentrated solution of selenic acid saturated with barium selenate also deposits crystals of barium selenic acid.

J. F. S.

Physical Chemistry of the Oxides of Lead. IV. Red Lead and Lead Sesquioxide. SAMUEL GLASSSTONE (*T.*, 1922, **121**, 1456—1469).

Physical Chemistry of the Oxides of Lead. V. The Electromotive Behaviour of Lead Dioxide. SAMUEL GLASSSTONE (*T.*, 1922, **121**, 1469—1480).

Behaviour of Sulphides of Heavy Metals in Aqueous Solutions. O. WEIGEL (*Sitzungsber. Ges. Naturw. Marburg*, 1921, No. 2, 35—50; from *Chem. Zentr.*, 1922, i, 182—183; cf. *A.*, 1907, ii, 237).—The known solubilities of heavy metal sulphides

as experimentally determined are generally greatly in excess of the values calculated from their solubility products. The author's determinations of the solubilities of the sulphides of thallium, silver, and lead show that in each case the calculated solubility is less than that determined experimentally. From the effect of lead ions on the solubility of lead sulphide and the potential of a lead sulphide electrode, it is concluded that dissociation in aqueous solution takes place in successive stages and that most of the dissolved substance is in the form of complex ions. Removal of metallic ions by addition of sulphide ions depends on the formation of complex ions.

G. W. R.

The Blue Flame produced by Common Salt on a Coal Fire.

ARTHUR SMITHELLS (*Nature*, 1922, 109, 745; cf. Merton, *ibid.*, 1922, 109, 683).—A brief historical note on the recognition of the blue flame produced by common salt on a coal fire as being due to compounds of copper derived from pyrites in the coal.

A. A. E.

The Rate of Combination of Copper and Phosphorus at Various Temperatures. C. A. EDWARDS and A. J. MURPHY (*J. Inst. Metals*, Adv. Copy).—Penetration of copper rods by phosphorus vapour takes place at 640° at atmospheric pressure. The dissociation-temperature curve for alloys of copper and phosphorus containing up to 30.8% of the latter indicates the existence of the phosphides CuP and Cu_3P_2 .

CHEMICAL ABSTRACTS.

Corrosion of Copper by Salt Solutions. W. MÜLLER (*Z. Metallk.*, 1922, 14, 286—295).—Copper is rapidly corroded by sea water, sodium chloride, and magnesium chloride, with the formation of a green, basic chloride which readily detaches itself from the metal and therefore does not form a protective coating. Weak solutions of magnesium and calcium sulphates corrode copper very slowly, the metal becoming gradually covered with a white precipitate containing bluish-green flakes of basic sulphate. [Cf. *J. Soc. Chem. Ind.*, 1922, Sept.]

A. R. P.

Critical Constants of Mercury. S. WEBER (*K. Danske Videnskab. Selskab. Math.-fys. Medd.*, 1920, 3, No. 4; cf. A., 1921, ii, 699).—The critical pressure of mercury is 1036 atmospheres. If curves be plotted giving the density of liquid and vapour phases of mercury and hydrogen, the temperature and density being written in terms of the critical temperature and density, the two curves practically coincide. Vapour-pressure curves plotted in a similar manner also coincide, hence it is concluded that hydrogen at this low temperature is monatomic.

CHEMICAL ABSTRACTS.

Atomic Weight of Mercury from Different Sources. J. N. BRÖNSTED and G. HEVESY (*Nature*, 1922, 109, 780; cf. this vol., ii, 149).—Measurements of the density of mercury prepared by the same method from minerals of different origin exhibit no differences exceeding the possible experimental error, which corre-

sponds with 0.0004—0.0012 in the atomic weight. It is therefore concluded, with great probability, that the isotopic composition of mercury of terrestrial origin is the same. A. A. E.

Catalytic Decomposition of certain Oxides. GUY B. TAYLOR and G. A. HULETT (*J. Amer. Chem. Soc.*, 1922, **44**, 1443—1445).—A theoretical paper in which the authors criticise the conclusions of Kendall and Fuchs (this vol., ii, 147). It is pointed out that in the system $\text{HgO}:\text{Hg}:\text{O}_2$ it is inadmissible to speak of the equilibrium pressure of oxygen at a fixed temperature without specifying the partial pressure of mercury. Further, the grounds on which Kendall and Fuchs maintain that their results are in agreement with those of Lewis (A., 1906, ii, 284) in connexion with the dissociation of silver oxide are considered to require explanation. The conclusions with regard to the decomposition of barium dioxide are also criticised, and it is pointed out that the explanation of the discrepancy between the authors' results (A., 1913, ii, 932) and those of Kendall and Fuchs does not meet the case.

J. F. S.

Catalytic Decomposition of certain Oxides. JAMES KENDALL and FRANCIS J. FUCHS (*J. Amer. Chem. Soc.*, 1922, **44**, 1447—1448).—An answer to the criticisms of Taylor and Hulett (cf. preceding abstract). J. F. S.

Preparation of Pure Ceria-earth Compounds. R. H. MANSKE (*Can. Chem. Met.*, 1922, **6**, 83—84).—A study of the solubility curves shows that if a saturated solution of the sulphates of cerium (about 50%), lanthanum, praseodymium, and neodymium at 0° is rapidly heated to 41°, the sulphates of lanthanum and neodymium crystallise with cerium sulphate, but free from praseodymium. Separation from cerium is easy. If the saturated solution of the sulphates were heated to 65°, lanthanum would crystallise free from praseodymium and neodymium so long as cerium was present.

CHEMICAL ABSTRACTS.

The Solubility of Gases in Aluminium. J. CZOCHRALESKI (*Z. Metallk.*, 1922, **14**, 277—285).—Molten aluminium absorbs very little of the common gases below 900°, but with increasing temperature above this the amount absorbed and retained in the solid metal at ordinary temperatures increases approximately in the following order: nitrogen, carbon monoxide, air, oxygen, sulphur dioxide, carbon dioxide, illuminating gas, and hydrogen. In general, the gas is retained in the solid metal in numerous microscopic blow-holes, but very small amounts of nitrides, oxides, carbides, and sulphides are formed in atmospheres containing nitrogen, oxygen, carbon, or sulphur compounds. The technical significance of these facts is discussed with reference to the production and working of aluminium. [Cf. *J. Soc. Chem. Ind.*, 1922, Sept.] A. R. P.

Mordants. II. Alumina. WILDER D. BANCROFT (*J. Physical Chem.*, 1922, **26**, 501—536).—A general discussion on the

hydrolysis of aluminium salts and the adsorption of aluminium sulphate and alumina by wool, silk, and cotton. It is shown that all aluminium salts are hydrolysed to some extent in aqueous solution, and the amount of hydrolysis increases with increasing temperature. The actual hydrolysis is greater with salts of weak acids, but the apparent hydrolysis may be abnormally large in sulphate solutions owing to the coagulating effect of the sulphate ions on the colloidal alumina. Different fibres adsorb alumina to different degrees, wool having a much greater adsorbing power than cotton, and silk being probably slightly inferior to wool. Owing to this difference in specific adsorption, wool decomposes aluminium salt solutions which are distinctly acid, whilst cotton is effective only in more basic solutions. The colloidal alumina is taken up and held firmly. Coagulated alumina may be adsorbed to some extent, but it easily rubs off the material. It is probable that in all cases alumina is adsorbed and not a basic salt. The phenomena may be complicated by the fact that the alumina itself will adsorb some sulphuric acid, for example, and that the wool may, and probably does, adsorb some sulphuric acid also. Since alumina is adsorbed less strongly by cotton than by wool, it is also held less strongly by cotton than by wool. If to cotton is added some substance, such as tannin, which adsorbs alumina strongly, the cotton mordanted with tannin will be able to take alumina out of solutions of aluminium salts which are not decomposed by cotton alone, and alumina is held more strongly by mordanted cotton than by cotton alone. The increase in adsorbing power shown by mercerised cotton is due to structural differences in the cotton fibre. There is no evidence of the formation of any definite compound between alumina and either wool, silk, or cotton.

J. F. S.

Preparation of Metals by Goldschmidt's Aluminothermic Method. I. TÔTARÔ FUJIBAYASHI (*J. Chem. Ind. Japan*, 1922, 25, 499—511).—Pure manganese and chromium free from carbon have been prepared by Goldschmidt's method. For the former, an intimate mixture of trimanganic tetroxide (100 parts), manganese dioxide or sesquioxide (15—20 parts), and 90% of the calculated weight of powdered aluminium was used, the yield being 85—90% of the theoretical value. The product contained 95—97% of manganese, the remainder being mainly aluminium. For the preparation of chromium, an intimate mixture of chromium sesquioxide (100 parts), calcium chromate (10—15 parts), and 90% of the calculated weight of powdered aluminium was used, the yield being 85—92% of the theoretical; the product contained 95—97% of chromium and 3—5% of aluminium.

K. K.

Reducing Action of Ferrous Hydroxide. SUSUMU MIYAMOTO (*J. Chem. Soc. Japan*, 1922, 43, 397—438).—Nitrites or nitrates can be estimated as ammonia by reduction with ferrous hydroxide in alkaline solution. In the case of the nitrite, the sample (0.1—0.3 gram) is boiled with 15 grams of ferrous sulphate and 200 c.c. of saturated alkali hydroxide solution and the evolved

ammonia is passed into sulphuric acid of known concentration as in Kjeldahl's method. Nitrates are not so easily reduced as nitrites, a reflux condenser must be attached to the boiling-flask instead of a bulb, and the boiling continued for three to three and a half hours, a current of hydrogen, nitrogen, or air being introduced into the flask to carry over ammonia. Ferrous hydroxide also reduces nitrobenzene to aniline at the ordinary temperature.

The velocity of reaction between potassium nitrite and ferrous hydroxide in alkaline solution was measured at 25°, 35°, and 45°. When the initial concentration of the nitrite is very small compared with that of the alkali hydroxide, the reaction is linear. The velocity constant increases with the concentration of the alkali hydroxide, and is expressed by the formula: $k = e^{A-B/P(1+0.79 C_{\text{KOH}})}$, where $A = 20.609$ and $B = 8098.2$. The velocity constant is increased 2.35 times for each elevation of 10° in temperature. The presence of potassium sulphate in the system has no effect on the reaction velocity. Attempts to measure the velocity at the boiling point, 108°, were unsuccessful, the reaction occurring too quickly.

The velocity of reaction between potassium nitrate and ferrous hydroxide in alkaline solution was measured at 108° and 112.2°, the velocity constant being approximately 7.577×10^{-3} (calc. 7.726×10^{-3} ; $C_{\text{KOH}} : 5.5394$ mol. per litre) and 2.331×10^{-2} (calc. 2.326×10^{-2} ; $C_{\text{KOH}} : 6.9541$ mol. per litre respectively).

The electric potentials of the following cells were measured at 25°, using 1*N*-, 2*N*-, 3*N*-, 4*N*-, and 5*N*-sodium hydroxide solutions:

I. Pt. platinised $[\text{NaOH}, \text{Fe}(\text{OH})_2, \text{Fe}_3\text{O}_4, x\text{H}_2\text{O}] | \text{NaOH} | \text{HgO}, \text{NaOH} | \text{Hg}$; II. Pt. platinised $\text{H}_2 | \text{NaOH} | \text{HgO}, \text{NaOH} | \text{Hg}$. III. Pt. platinised $[\text{NaOH}, \text{Fe}(\text{OH})_2, \text{Fe}_3\text{O}_4, x\text{H}_2\text{O}] | \text{NaOH} | \text{NaOH} | \text{H}_2$, Pt. platinised. The potential of I is expressed by $E_1 = 0.8030 - 0.0015 C_{\text{NaOH}}$, that of II by $E_2 = -0.9270 - 0.00332 \log C_{\text{NaOH}}$, that of III by $E_3 = 0.1240 + 0.00332 \log C_{\text{NaOH}} - 0.0015 C_{\text{NaOH}}$. K. K.

A New Iron Salt. OTTO RÖHM (*Collegium*, 1921, No. 614, 282—284).—When a concentrated solution of ferrous sulphate is oxidised with chlorine and the excess of water allowed to evaporate, a crystalline substance having the composition $\text{FeSO}_4\text{Cl} \cdot 6\text{H}_2\text{O}$ is obtained. This compound, which is neither a mixture nor a double salt, and does not deliquesce in damp air, may also be prepared (a) by heating 1 mol. of ferric chloride, 1 mol. of ferric sulphate, and 18 mols. of water, and (b) by heating 1 mol. of ferric chloride, 1 mol. of sulphuric acid, and 6 mols. of water, and driving off the hydrogen chloride formed. CHEMICAL ABSTRACTS.

I. The Structure of Electrolytically Deposited Nickel. II. The Influence of Superposed Alternating Current on the Deposition and Solution Potential of Nickel. V. KOHLSCHÜTTER and H. SCHÖDL (*Helv. Chim. Acta*, 1922, 5, 490—512, 593—609).—I. Continuing the work of Kohlschütter and Vuilleumier (A., 1919, ii, 9) and Stäger (A., 1920, ii, 728) on the properties of electrolytically deposited nickel, experiments have been made with the object of determining whether there is any relation between

the micro-structure of the deposits and the contraction phenomenon which has been found to depend on the deposition potential. Comparative experiments were made, using direct current and alternatively direct current with a superposed alternating current, which considerably diminishes the observed contraction of the deposit. The previous conclusion was confirmed that, when deposits showing a high contraction are formed, deposition takes place at first in a thin deposit or skin of hydrogen. The same effect can be produced by saturating the electrolyte with carbon dioxide, although to a less extent. When there is free evolution of hydrogen at the cathode, the contraction is slight; the surface of the deposit is bright and metallic, but under the microscope it has a scaly appearance. When the evolution of hydrogen is small, the deposit shows a greater contraction and it has a brown appearance. The inner structures of the different deposits are practically identical when examined microscopically. The size of the particles is from 1.4 to 2.2 μ , the larger particles generally appearing when an alternating current is used. It is concluded therefore that the contraction is purely a surface phenomenon, due to the aggregation of particles, originally deposited in a highly disperse form, into particles of larger size.

Deposits obtained with direct current were brittle, those from sulphate solutions less so than those from chloride; those obtained with alternating current were flexible. There seemed, however, to be no relation between brittleness and the contraction phenomenon. The brittleness is probably due to the presence of hydrogen in the metal.

II. The superposition of the alternating current causes a lowering of the cathode potential more or less parallel with the lessening of the contraction effect in the deposit. A number of observations, however, indicate that the relation is not so close as might appear to be the case between the structure and contraction of the deposit on the one hand and the deposition potential on the other. Since, however, the contraction takes place almost instantaneously, it is not possible to observe the structure of the metal as it is originally deposited with a high cathode potential. There is no doubt that hydrogen is the important factor influencing both the polarisation and the structure of the deposit.

E. H. R.

New Hydrate of Uranyl Nitrate. FRANK E. E. GERMANN (*J. Amer. Chem. Soc.*, 1922, **44**, 1466—1469).—The density of solutions of uranyl nitrate has been determined at 24° for compositions up to 50% and a density curve plotted which exhibits a strong curvature towards the composition axis. Cooling and heating curves have been constructed, and both show the existence of an *icositetrahydrate* of uranyl nitrate, $\text{UO}_2(\text{NO}_3)_2 \cdot 24\text{H}_2\text{O}$. This compound forms spontaneously at -35° and exists only below -20° . The existence of five distinct fluorescence spectra due to uranyl nitrate as stated by Howes (*Physical Rev.*, 1915, **6**, 192) has been disproved. These are due to the hexahydrate, the icositetrahydrate, and various mixtures of the two depending on the concentration and rate of cooling.

J. F. S.

Complex Uranyl Hypophosphites. ARTHUR ROSENHEIM and GERT TREWENDT (*Ber.*, 1922, 55, [B], 1957—1960).—The analogy between sulphites and hypophosphites is illustrated further by the isolation of alkali uranyl hypophosphites which correspond in their composition with the alkali uranyl sulphites, $R_2[(UO_2)_2(SO_3)_3]$ and $R_2[(UO_2)_4(SO_3)_5] \cdot xH_2O$, described by Köhlschütter (*A.*, 1900, ii, 484).

Uranyl hypophosphite, $UO_2(H_2PO_2)_2$, microcrystalline, yellow prisms, is prepared by agitating solutions of one molecular proportion of uranyl nitrate and four molecular proportions of sodium hypophosphite (a microcrystalline, pale yellow *trihydrate* is also described); it is almost insoluble in water, but readily soluble in an excess of sodium hypophosphite or uranyl nitrate solution. It is transformed by a solution of sodium hypophosphite (4 molecular proportions) into the *salt*, $Na[(UO_2)_2(H_2PO_2)_5] \cdot 4.5H_2O$, slender, pale yellow needles (a *hexahydrate* is also described); the corresponding anhydrous *potassium salt*, yellow crusts, and *ammonium salt*, pale yellow leaflets, and the *dihydrated guanidinium salt*, aggregates of needles, were analysed. Uranyl hypophosphite is converted by six to eight molecular proportions of sodium hypophosphite into the *salt*, $Na[(UO_2)(H_2PO_2)_2] \cdot 3.5H_2O$, large, rectangular plates (the *pentahydrate* is also described). Attempts to prepare corresponding compounds with other alkali hypophosphites yielded salts of the series $R[(UO_2)_2(H_2PO_2)_5]$. The compounds from uranyl hypophosphite and ten or more molecular proportions of alkali hypophosphite could not be caused to crystallise. H. W.

The Isotopes of Tin. F. W. ASTON (*Nature*, 1922, 109, 813).—An immediate result of the application of a method for increasing the sensitivity of photographic plates towards positive rays has been the definite proof of the complex nature of tin (cf. *A.*, 1921, ii, 474). By using tin tetramethyl, eight lines corresponding approximately with atomic weights 120, 118, 116, 124, 119, 117, 122, 121 (in decreasing order of intensity) were definitely proved to be due to tin. The average atomic weight, in proportion to the intensities, agrees well with the accepted value. It is remarkable that the differences between the lines are integral to the highest accuracy, but the lines themselves compared with known lines give atomic weights always tending to be 2 or 3 parts per 1000 too light for the above whole numbers. It is strongly indicated that this divergence cannot be due to experimental error. The presence of the two faint components of xenon 128 and 130 previously suspected has now been confirmed. A. A. E.

Complex Mixed Antimony Iodobromides. A.-CH. VOURNAZOS (*Compt. rend.*, 1922, 175, 164—167).—An acid containing the complex anion $-SbI_3Br$ is obtained by the action of dry gaseous hydrogen bromide on antimony tri-iodide in a non-aqueous medium such as glacial acetic acid. It cannot be isolated in a solid state by the evaporation of the acetic acid solution, as, owing to dissociation, only a residue of the tri-iodide remains. The salts

of this acid are, however, obtained in crystalline form by triturating and finally warming equimolecular proportions of antimony triiodide and a bromide with a suitable non-aqueous medium, acetic acid, xylene, etc. The compounds form coloured crystals, which are rapidly decomposed by water or ethyl alcohol, giving antimonious oxide, hydriodic acid, and the metallic bromide. *Sodium antimoniodobromide*, SbBrI_3Na , prepared in xylene, forms small, orange-yellow crystals. *Potassium antimoniodobromide*, SbBrI_3K , is similar in colour, but the *ammonium* and *lithium* salts are reddish yellow. *Zinc antimoniodobromide*, $\text{ZnSbI}_3\text{Br}_2$, forms brown, tabular crystals, which are fairly stable and only slowly decomposed by water.

G. F. M.

Mineralogical Chemistry.

Haloës and Earth-history. A New Radioactive Element. J. JOLY (*Nature*, 1922, **109**, 517—518 and 578—579).—A description of colourless, spherical, halo-like forms, of average diameter 0.0104 mm., occurring in the Archæan black mica of Ytterby. It is suggested that they may be due to a radioactive element (for which the name "hibernium" is suggested) having an α -ray range in air of 1—1.5 cm.

A. A. E.

The Crystal Structures of Marcasite (FeS_2), Arsenopyrite (FeAsS), and Loellingite (FeAs_2). MAURICE L. HUGGINS (*Physical Rev.*, 1922, **19**, 369—373).—Arsenopyrite and loellingite are assumed to have the same general arrangement as marcasite, in which the sulphur atoms are considered to be in pairs, each sulphur atom being linked, by pairs of electrons, to one sulphur and four iron atoms, and each iron atom to six sulphur atoms.

A. A. E.

Minerals from near Oudjda, Morocco. J. BARTHOUX (*Compt. rend.*, 1922, **175**, 312—314).—An account is given of crystallised minerals (galena, vanadinite, pyromorphite, wulfenite, cerussite, dolomite, calcite, and aragonite) from a lead mine in dolomitic limestone at Gebel Mahser.

L. J. S.

Babingtonite from Japan. MANJIRÔ WATANABÉ (*Amer. J. Sci.*, 1922, [v], **4**, 159—164).—Babingtonite occurs with hedenbergite, garnet, chalcopyrite, magnetite, etc., in a metamorphic contact in the Yakuki mine, province Iwaki. A crystallographic and optical description is given of the material.

L. J. S.


Composition of Aerinite. J. ORCEL (*Compt. rend.*, 1922, **175**, 309—311).—Aerinite from Casserras, Huesca, prov. Aragon, consists of a mixture of a blue, transparent, strongly pleochroic mineral with pyroxene, quartz, and spinel. The first is easily separated

by reason of its ready solubility in acids. Analysis of hydrochloric acid solution gave :

SiO ₂ .	TiO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	MgO.	MnO.	CaO.	SrO.
39.26	0.49	19.80	7.13	1.35	3.52	0.03	9.08	0.20

P ₂ O ₅ .	V.	Alkalis.	H ₂ O (at 106°).	H ₂ O (at 400°).	H ₂ O (400° to 1100°).	Total.
0.07	trace	trace	5.43	11.03	2.92	100.31

corresponding approximately with



The mineral represents a new calcium-bearing type of the leptochlorites; this is supported by optical characters and its behaviour when heated. Spectroscopic analysis shows that the strontium and vanadium are present only in the hydrochloric acid extract of the mineral. The cause of the blue colour is discussed; it is regarded to be dependent on the molecular structure.

L. J. S.

Analytical Chemistry.

Nephelometry of Coloured Hydrosols. H. BECHHOLD and F. HEBLER (*Kolloid Z.*, 1922, **31**, 7—12).—In the nephelometric investigation of white turbidities it has been repeatedly shown that a strict proportionality exists between the concentration of the substance causing the turbidity and the amount of diffracted light, provided that no change occurs in the original substance. It is now shown that in the case of coloured sols and turbidities very marked deviations from this proportionality occur. Concentrated coloured sols such as colloidal indigotin and silver behave in comparison with dilute solutions as though they were less concentrated. This is shown to be due to the absorption of light by the coloured particles forming the turbidity. This action in the case of white particles is not great, but in that of coloured particles it is very strong. The disproportionality in the absorption is caused particularly by those wave-lengths which are absorbed by the turbid medium to a larger or smaller extent. This selective absorption must therefore be removed by the use of suitable light filters. The most efficient filters are shown to be isochromatic filters, that is thin layers composed of the substance under investigation. These filters may be employed either as solutions or as dry gelatin films between the source of light and the nephelometer. In this way it is possible to investigate nephelometrically coloured turbidities and coloured hydrosols in exactly the same way as colourless turbidities, and in these circumstances they show exactly the same proportionality between concentration and amount of diffracted light as white turbidities.

J. F. S.

Investigation, by Means of the Hydrogen Electrode, of the Chemical Reactions involved in Water Purification. R. E. GREENFIELD and A. M. BUSWELL (*J. Amer. Chem. Soc.*, 1922, **44**, 1435—1442).—Titration curves have been prepared, using the hydrogen electrode, for the neutralisation of the carbonates of sodium, magnesium, and calcium with a strong acid. It is shown that the shape and position of the curve are unaffected by the metallic ion, but that the inflection point occurs at a slightly higher Sørensen value in dilute solutions than in the more concentrated solutions. Precipitation curves of the precipitation of magnesium hydroxide have been plotted, and these show that the precipitation of magnesium is complete at a Sørensen value in the neighbourhood of $P_H=10.6$. Magnesium hydroxide does not form until the value is as high as $P_H=9.0$. Precipitation curves for the precipitation of calcium as carbonate, whilst not as regular as those obtained in the case of magnesium, tend to show that the reaction is complete, sufficient carbonate being present at $P_H=9.5$. Aluminium hydroxide is shown to start precipitating in solutions as acid as $P_H=4$, and to be completely precipitated at $P_H=6.5-7.5$. At values much higher than this, resolution commenced to take place, and this was complete at a value between $P_H=10$ and $P_H=11$. The precipitation reactions were not instantaneous, but took several hours to reach completion. This was the more noticeable the more dilute the solution. J. F. S.

Sodium Sulphide as a Substitute for Hydrogen Sulphide in Qualitative Analysis. GIORGIO VORTMANN (*Boll. Sci. tecn.* **3**, No. 5; *Giorn. Chim. Ind. Appl.*, 1921, **3**, 565).—A solution in hydrochloric acid is oxidised, treated with solid sodium carbonate in slight excess, warmed with sodium hydroxide solution (boiled to remove ammonia if necessary), and precipitated with excess of a 20% solution of sodium sulphide. The precipitate (i) may contain silver, copper, bismuth, cadmium, lead, iron, cobalt, nickel, manganese, and zinc as sulphides; uranium, chromium, and rare earths as hydroxides; barium, strontium, calcium, and magnesium as carbonates; whereas the filtrate (i) may contain the sulphides of mercury, nickel, arsenic, antimony, tin, gold, platinum, molybdenum, tungsten, and vanadium. On treatment of the precipitate (i) with dilute hydrochloric acid, iron, manganese, zinc, uranium, chromium, barium, strontium, calcium, magnesium (phosphate), lithium, and rare earths pass into solution (ii), whilst silver, copper, bismuth, cadmium, lead, cobalt, and nickel remain undissolved (ii). Solution (ii) is boiled to remove hydrogen sulphide, and bromine water is then added, followed by sodium carbonate in excess. The filtrate (iii) may contain chromium, uranium, and manganese, whilst the precipitate (iii) contains iron, manganese, zinc, barium, strontium, calcium, magnesium, and lithium. After redissolution of the precipitate (iii) in hydrochloric acid, iron is precipitated with sodium acetate, manganese with ammonium hydroxide and bromine water, barium, strontium, and calcium with ammonium carbonate, the zinc in the filtrate with sodium sulphide, and finally the magnesium

and lithium. The filtrate (i) is boiled with ammonium chloride to precipitate mercury, nickel, and aluminium, whilst antimony, tin, arsenic, molybdenum, tungsten, and vanadium remaining in solution can be separated by known methods. The alkali metals are tested for in another portion of the substance.

CHEMICAL ABSTRACTS.

Titration of Acids and Bases. J. L. LIZIUS and N. EVERS (*Analyst*, 1922, **47**, 331—341).—The theory of titration and the dependence of the hydrogen-ion concentration of the end-product on the nature of the salt formed is explained. A list of newer indicators with their colour changes and the P_H range over which they are applicable is given, and four mixed indicators are suggested. A table of common titrations with the hydrogen-ion concentrations at their end-points, suitable indicators, and the colours obtained at their end-points is also given. By titrating to a definite shade of colour instead of to the colour-change of the indicator, an increase in the accuracy of titrations results, and certain titrations are made possible which are impracticable by ordinary methods.

H. C. R.

Symmetrical Diphenylguanidine as a Standard in Acidimetry and Alkalimetry. C. A. CARLTON (*J. Amer. Chem. Soc.*, 1922, **44**, 1469—1474).—The suitability of symmetrical diphenylguanidine as a standard in acidimetry and alkalimetry has been investigated. It is shown that this substance can easily be obtained in a sufficiently high state of purity for the present purpose by three recrystallisations of the crude material from toluene. The pure compound is stable in air and soluble in alcohol. It may be titrated directly in cold solutions with either an alcoholic or an aqueous solution of hydrochloric acid, using either bromophenyl-blue or methyl-red as indicator. The results obtained with this substance are comparable with those obtained with sodium carbonate or silver chloride, and diphenylguanidine is more convenient in use. Diphenylguanidine is the only basic substance, proposed as a standard, which meets all the requirements of an ideal standard.

J. F. S.

Micro-incineration. A. SCHOELLER (*Ber.*, 1922, **55**, [B], 2191—2192).—The substance under investigation is spread evenly over a thin strip of glass (5—6 mm. wide) which is placed in a horizontal hard glass tube about 10 mm. in diameter and 12 cm. long. The tube is heated very gently until all volatile matter is expelled. The residue is allowed to cool and is subsequently heated rather more strongly in a current of moist oxygen; in most cases, the carbon disappears rapidly without at any time glowing. With difficultly combustible substances it is advisable to interrupt the heating again. Too powerful ignition is to be avoided. The minutest trace of ash is readily visible under the microscope. The method is particularly suitable for the incineration of sections of plant-tissue, since the original structure is better preserved than when they are heated over a free flame.

H. W.

The Estimation of Hydrogen and its Separation from Gaseous Paraffins by means of Palladious Chloride. J. A. MULLER and A. FOIX (*Bull. Soc. chim.*, 1922, [iv], 33, 713—717).—Hydrogen may be separated from gaseous paraffins and estimated by its reducing action on excess of palladious chloride. The precipitated palladium is dried and weighed, and hence the quantity of hydrogen is calculated. A correction should be made for the small quantity of the gas occluded in the metal; the authors state that 1 mg. of palladium represents 0.24 c.c. of hydrogen measured at 0° and 760 mm. when the reaction takes place under the prescribed conditions.
H. J. E.

Electrometric Titrations with Mercury Perchlorate. I. M. KOLTHOFF (*Z. anal. Chem.*, 1922, 61, 332—343).—Chlorides, bromides, or iodides may be titrated with mercury perchlorate solution, the end-point of the titration being determined with accuracy by electrometric means even in very dilute solutions, but the method cannot be applied to mixtures of these salts. The method is also trustworthy in the case of thiocyanates, cyanides, ferrocyanides, formates, acetates, monochloroacetates, lactates, benzoates, and salicylates, but it cannot be used for trichloroacetates owing to the ready hydrolysis of the mercury compound, or for the salts of polybasic aliphatic acids. The mercury perchlorate solution is prepared by saturating perchloric acid with mercuric oxide.
W. P. S.

Detection of Fluorine. B. FETKENHEUER (*Wiss. Veröffentl. Siemens-Konzern*, 1922, 1, [3], 177).—The substance is heated with sand and a few c.c. of sulphuric acid at 90° in a test-tube. On shaking the tube, the presence of fluorine is shown by the acid collecting into oily drops, which appear not to wet the surface of the glass. With 2 grams of substance, 0.01% of fluorine may be detected after heating for one minute. Insoluble fluorides such as aluminium fluoride must first be fused with sodium carbonate and sand; the fused mass is then heated as above with sulphuric acid.
A. R. P.

The Colour of Iodine Solutions at Low Temperature. JEAN PICCARD and E. HERRMANN (*Helv. Chim. Acta*, 1922, 5, 625—626).—The brown colour formed by iodine in solvents containing oxygen has been suggested as a qualitative test for oxygen (this vol., ii, 389), in absence of other elements or groups of an unsaturated character such as tervalent nitrogen. It has been observed that very dilute solutions of iodine in hydrocarbons such as light petroleum, which are violet at the ordinary temperature, become brown at lower temperatures, in the neighbourhood of -20°. When the iodine concentration is increased beyond a certain value, about 1 mg. in 10 c.c., however, the colour change does not take place. It is shown that the brown colour is due to an impurity in the commercial light petroleum, and that it does not appear with carefully purified specimens. The phenomenon can be reproduced by addition of a small quantity of alcohol to

the pure light petroleum, or to toluene, chloroform, and other solvents.

E. H. R.

Rapid Estimation of Sulphur. LUIGI LOSANA (*Giorn. Chim. Ind. Appl.*, 1922, 4, 297—299).—The method previously described (this vol., ii, 582) gives accurate results with pyrites, copper pyrites, and other mineral sulphides, ultramarine, adulterated white lead, cement, various rocks, sulphates of copper, sodium, etc., naphtha, anthracene, coke, ichthyol, "saccharin," ebonite, and caoutchouc.

T. H. P.

Determination of Sulphur in Vulcanised Rubber. J. W. W. DYER and AMY R. WATSON (*J. Soc. Chem. Ind.*, 1922, 41, 251—252).—Methods are described for the estimation of free and combined sulphur in vulcanised rubber. These are as follows. (a) Combined sulphur. The sample (0.5—3.0 grams) is added to 30—40 c.c. of nitric acid (d 1.42) in a suitable flask and the reaction started by gentle heating; subsequently the mixture is heated to the boiling point and boiling continued until the liquid is clear. Pure powdered potassium permanganate is now added, about 0.25—0.5 gram at a time, and the heating continued between additions. The addition of permanganate is continued until finally a small black precipitate remains; this requires generally about 2 grams of permanganate. The contents of the flask are poured into a dish and evaporated to dryness, taken up with 10 c.c. of concentrated hydrochloric acid, and again evaporated to dryness. The residue is treated with water, made up to 100 c.c., and made just acid to methyl-orange at the boiling point. The sulphur is then precipitated as barium sulphate. (b) Free sulphur. To 50—60 c.c. of acetone, 0.5—1 gram of material is added and the free sulphur extracted. To the extract 0.5—1 gram of powdered permanganate is added, the mixture shaken, and kept at the ordinary temperature for thirty minutes. If the purple colour disappears a little more permanganate is added. The acetone is then distilled off on the water-bath and the brown residue heated at 100—110° for a short time. About 3—5 c.c. of hydrochloric acid (d 1.16) is added for each gram of permanganate used, and the solution heated until colourless. It is then diluted a little and filtered, the filtrate is made up to 100 c.c., and ammonia cautiously added until the solution becomes yellow and slightly turbid. It is then made just acid and precipitated while boiling with 4—5 c.c. of $N/2$ -barium chloride solution. Both methods give results which are in good agreement with those obtained by the Carius method.


J. F. S.

Quantitative Micro-analysis of Mixtures with Special Reference to Organic Ultimate Analysis. A. BENEDETTI-PICHLER (*Z. anal. Chem.*, 1922, 61, 305—331).—The author describes the advantages of micro-chemical methods, points out the importance of obtaining representative samples in dealing with very small quantities of substances, and gives instances of the application of the methods and their limits of accuracy.

W. P. S.

Volumetric Estimation of Carbon and Hydrogen in Organic Compounds. JOSEF LINDNER (*Ber.*, 1922, 55, [B], 2025—2031).—The substance is burnt in the usual manner and the products of its combustion are passed over a substance which by its hydrolysis gives hydrogen chloride and thence into a known volume of standard barium hydroxide solution. The latter is titrated with *N*/10-hydrochloric acid in the presence of phenolphthalein; the carbon dioxide is subsequently removed after addition of an excess of hydrochloric acid and the resulting solution is again titrated with barium hydroxide. With *N*/10-solutions and burettes of 50 c.c. capacity, the analysis is conveniently performed with 20 mg. of material.

The most suitable substance for hydrolysis is chloronaphthylxy-chlorophosphine, which may contain chloronaphthyltetrachlorophosphine: it is decomposed by water in accordance with the equations:



It is prepared by passing chlorine into molten naphthyldichlorophosphine until three atomic proportions of the gas have been absorbed. It is not a well-defined individual. Hydrogen chloride is freely and quantitatively evolved when a current of moist air is passed through the molten material. It is, however, convenient to depress its melting point to some extent by a preliminary regulated treatment with moist air. The material prepared in this manner somewhat readily evolves considerable amounts of readily volatile acid products, which must be removed by protracted exposure to a current of dry air; the elimination of acid is thereby ultimately reduced to a small amount, for which a correction can be applied, but the presence of the acid has the drawback that the sharpness of the end-point in the ultimate titration is diminished.

Test analyses show that the method gives accurate results, which, however, are somewhat low for hydrogen, the source of error most probably lying in the uncertainty of the correction for the acidic products evolved spontaneously from the phosphorus compound. H. W.

Absorption of Carbon Monoxide by Acid Cuprous Chloride Solution in the Presence of a Reducing Agent. ALFRED KROPP (*Z. angew. Chem.*, 1922, 35, 451—452).—The rate of absorption of carbon monoxide by acid cuprous chloride solution is increased by the addition of stannous chloride to the solution. The stability of the solution is improved by the use of a slight excess of stannous chloride (cf. Krauskopf and Purdy, *A.*, 1920, ii, 267). J. S. G. T.

Comparison of Colorimetric and Electrometric Estimations of Hydrogen-ion Concentrations in Solutions containing Carbon Dioxide. GLENN E. CULLEN and A. BAIRD HASTINGS (*J. Biol. Chem.*, 1922, 52, 517—520).—Contrary to the

results obtained by Evans (A., 1921, ii, 271), the colorimetric and electrometric methods for the estimation of the hydrogen-ion concentration of solutions containing carbon dioxide yield identical results, provided precautions are taken to prevent the loss of carbon dioxide. E. S.

Estimation of Small Amounts of Potassium by the Lindo-Gladding Method. WILLIAM HAZEN (*J. Assoc. Off. Agric. Chem.*, 1922, 5, 456—460).—In the estimation of potassium by the Lindo-Gladding method, the use of 90% alcohol for washing gives better results than 80% alcohol, and when working with small amounts of potassium salts and high accuracy is desired, it is advisable to use the stronger alcohol for washing after the ammonium chloride treatment as well. The lower results obtained by using the weaker alcohol do not appear to be due to the sodium salts which may be present, as has been suggested, as equally low results are obtained in the absence of sodium salts. As, however, it takes a longer time to wash out the ammonium salts with 90% alcohol, it is preferable to use 80% alcohol for the second washing in ordinary fertiliser work, where the percentage error arising from this will not be very serious, as the samples generally contain relatively high amounts of potassium. G. F. M.

Identification of Cæsium and Rubidium. JOHN MISSENDEN (*Chem. News*, 1922, 124, 362).—The spectroscope affords the best means of distinguishing cæsium from rubidium; the former shows two bright blue lines (4557 and 4592) in the blue portion of the spectrum, whilst rubidium has two lines (4200 and 4237) in the violet and two other lines (7953 and 7810) in the red. The salts of the two metals resemble each other generally, but there is a considerable difference in the solubility of the two nitrates; 100 parts of water at 3.35° dissolve 11 parts of cæsium nitrate and 79.9 parts of rubidium nitrate. W. P. S.

Hydrotimetric Precision and Aqueous Preparation of the Standard Soap Solution. ED. JUSTIN-MUELLER (*J. Pharm. Chim.*, 1922, 26, 18—21).—Clark's method as modified by Boutron and Boudet is used, but a standard soap solution of only one-tenth the strength is recommended. It is prepared by dissolving 3.5 grams of white Marseilles soap in 200 c.c. of boiling water and then making the volume up to 900 c.c. This solution is diluted, if necessary, until 24 c.c. of it corresponds exactly with 40 c.c. of a 0.025% solution of calcium chloride or of a 0.059% solution of barium nitrate. W. G.

Estimation of Calcium in Natural Phosphates. R. MEURICE (*Ann. Chim. Analyt.*, 1922, [ii], 4, 198).—The phosphate is treated with mineral acid, the silica removed, and iron and aluminium removed as phosphates. The filtrate is neutralised with ammonia, treated with 20 c.c. of 20% hydrochloric acid, and 30 c.c. of 4% ammonium oxalate. The solution is warmed and 10% ammonium acetate solution added at a rate of about 30 drops per minute. A granular precipitate of calcium oxalate

is thus obtained, and is collected, washed, and ignited as usual. Test analyses carried out on known quantities of calcium in the presence of large excess of phosphate and magnesium sulphate gave very accurate results.

H. C. R.

Estimation of Magnesium and Alkalis in Minerals. L. H. BORGSTRÖM (*Hyllningsskrift tillägnad Ossian Aschan*, 1920, 118—123).—The precipitation of magnesium hydroxide is best effected with piperidine, a little of which should also be added to the washing water. Lawrence Smith's method for the separation of magnesium from alkalis in silicate analysis is the most satisfactory.

CHEMICAL ABSTRACTS.

Colorimetric Method for the Estimation of Small Amounts of Magnesium. A. P. BRIGGS (*J. Biol. Chem.*, 1922, 52, 349—355).—The method described is similar to that of Hammett and Adams (cf. this vol., ii, 587).

E. S.

Estimation of Magnesium in Blood, Plasma, and Serum. W. DENIS (*J. Biol. Chem.*, 1922, 52, 411—415).—The method is essentially the same as that of Briggs (preceding abstract) and of Hammett and Adams (this vol., ii, 587).

E. S.

Magnesium Compound of 8-Hydroxyquinoline. CARL TH. MÖRNER (*Pharm. Zentr.-h.*, 1922, 63, 399—402; cf. Griebel, A., 1921, ii, 606).—If an aqueous solution of 8-hydroxyquinoline (sulphate) is treated with a suspension of magnesium hydroxide, the latter goes into solution and a lustrous, sulphur-yellow precipitate of the magnesium compound of 8-hydroxyquinoline is formed. The precipitate is also given by "magnesia mixture." The precipitate is microcrystalline and consists of hexagonal platelets. The reaction takes place only in alkaline solution and is sufficiently sensitive to detect 1 part of magnesium in 25,000. The precipitate contains 6.25—6.29% of magnesium and approximates to the formula $(C_9H_6ON)_2Mg \cdot 4H_2O$. The compound is also formed by the action of 8-hydroxyquinoline on magnesium ammonium phosphate. The reaction thus permits the separation of magnesium from phosphate in ammoniacal solution. Its use as a dry antiseptic is suggested.

G. W. R.

Lead. I. Estimation of Minute Amounts of Lead in Biological Material. L. T. FAIRHALL (*J. Ind. Hyg.*, 1922, 4, 9—20).—The ash is dissolved in dilute hydrochloric acid, the solution neutralised to methyl-orange with sodium hydroxide, and then slightly acidified with hydrochloric acid. Lead is precipitated as sulphide, the washed precipitate dissolved in 2—5 c.c. of concentrated nitric acid, and boiled to expel hydrogen sulphide. After neutralising with sodium hydroxide and slightly acidifying with acetic acid, an excess of potassium chromate is added, and the solution boiled. The precipitate is dissolved in hydrochloric acid, and an excess of potassium iodide is added to the solution, which is titrated with 0.005*N*-sodium thiosulphate solution. For biological materials, the method appears to be accurate to $\pm 3\%$.

CHEMICAL ABSTRACTS.

Colorimetric Estimation of Manganese in Steels, Alloys, and Ores. J. HESLINGA (*Chem. Weekblad*, 1922, 19, 302—303).—The ordinary method of oxidising to permanganate in acid solution is unsatisfactory when iron is present, because of the yellow colour of iron salts. A simple and accurate estimation may be made by employing the compound H_2MnO_3 , which in alkaline solution gives a yellowish-brown to dark brown coloration.

One hundred to 200 mg. of material are dissolved in a suitable acid, and the solution is diluted and treated with excess of powdered zinc oxide, made up to 100—200 c.c., and filtered. The zinc oxide precipitates the trivalent metals and copper completely after a few minutes' shaking at the ordinary temperature. The filtered solution is poured into a burette, and allowed to fall into a solution containing hydrogen peroxide (3%) and excess of potassium hydroxide (10%) until a suitable colour is obtained; this is matched by means of a standard manganese solution. Several titrations are carried out in succession, adding more of the unknown solution, and matching the colour by means of the standard. Addition of only 0.02 mg. of manganese is easily detected.

Lead does not interfere if excess of potassium hydroxide is employed; if nickel and/or cobalt are present, potassium cyanide is added during the titration. Accurate results are readily obtained when the manganese content is not greater than 20%. S. I. L.

Estimation of Chromium in Metals. WILLI LOFFELBEIN (*Chem. Ztg.*, 1922, 46, 679).—Chromium is estimated in nickel bronzes and similar alloys by removing the heavy metals from a solution of the alloy with hydrogen sulphide, treating the oxidised filtrate with a slight excess of ammonia, collecting, igniting, and fusing the precipitate with sodium peroxide, acidifying the filtered solution of the melt with hydrochloric acid, adding potassium iodide, and titrating the liberated iodine with sodium thiosulphate solution. A. R. P.

Rapid Method for the Estimation of Chromium in Nickel-Chromium Steel. WILHELM HILD (*Chem. Ztg.*, 1922, 46, 702—703).—The chromium is oxidised to chromic acid in a sulphuric acid solution of the steel by means of potassium permanganate, the excess of which is destroyed with manganese sulphate. The chromic acid is then estimated by addition of an excess of standard ferrous sulphate solution and titration with permanganate or dichromate [cf. *J. Soc. Chem. Ind.*, 1922, 41, 671A.] A. R. P.

Estimation of Small Amounts of Molybdenum in Tungsten. DOROTHY HALL (*J. Amer. Chem. Soc.*, 1922, 44, 1462—1465).—Small amounts of molybdenum may be estimated in tungsten by converting the molybdenum into xanthate, extracting this with chloroform, and converting into sulphide and oxide. The method consists in dissolving 1 gram of the sample in a mixture of nitric and hydrofluoric acids and evaporating with sulphuric acid. The yellow oxides are dissolved in sodium hydroxide solution and trans-

ferred to a 200 c.c. separating funnel, acidified with sulphuric acid, and the solution diluted so that 1 c.c. contains about 0.1 mg. of molybdenum. The volume is then made up to 150 c.c., 0.5 gram of potassium xanthate and a few drops of 1 : 3-sulphuric acid are added and the mixture is shaken thoroughly. About 10 c.c. of chloroform are then added and the whole is shaken for several minutes. The highly-coloured chloroform layer settles at the bottom and is run into a second funnel for washing. The original solution is repeatedly treated with potassium xanthate, acid, and chloroform until the chloroform layer is colourless, each portion of chloroform solution being added to the first. The chloroform solution of molybdenum xanthate is washed with water several times, run into a 150 c.c. beaker, and evaporated to dryness. The residue is heated and converted into a mixture of oxide and sulphide. This is dissolved in nitric acid, evaporated with sulphuric acid, and precipitated as molybdenum sulphide by passing hydrogen sulphide into the hot molybdenum solution made alkaline with sodium hydroxide. The solution is then slowly acidified with dilute sulphuric acid and the precipitate filtered, washed with hot water, and carefully ignited to oxide. The results are easily reproducible and trustworthy, and the method is good for mixtures containing small amounts of molybdenum. J. F. S.

Rapid Analysis of Ferro-tungsten. LUIGI LOSANA and ENRICO CAROZZI (*Giorn. Chim. Ind. Appl.*, 1922, 4, 299—301).—Ferro-tungsten is readily dissolved in 50% nitric acid solution containing a little hydrofluoric acid, subsequent treatment with concentrated sulphuric acid resulting in the elimination of the hydrofluoric acid and precipitation of the tungsten as tungsten trioxide, which may be either weighed as such or estimated volumetrically (cf. *J. Soc. Chem. Ind.*, 1922, 41, 671A). T. H. P.

Electrometric Titration of Uranium with Potassium Permanganate and Potassium Dichromate. D. T. EWING and E. F. ELDRIDGE (*J. Amer. Chem. Soc.*, 1922, 44, 1484—1489).—When solutions of uranyl sulphate, reduced with zinc, are titrated electrometrically with potassium permanganate or when solutions of uranium chlorides similarly reduced are titrated with potassium dichromate, a change in the oxidation potential occurs when the tervalent uranium is all oxidised to the quadrivalent condition, and a second change occurs when the latter is all oxidised to the hexavalent condition. Thus the total amount of uranium may be calculated. A third change of potential is observed when iron is present, and this represents the point where all the ferrous iron is converted into ferric iron. The determination is carried out with apparatus of the usual type, except in the form of the titration vessel, since it is necessary to prevent the free access of air. The titration cell is a glass vessel fitted with a cover which has six holes bored in it. Through the holes pass a tube for leading in carbon dioxide, a platinum electrode, the connecting tube of a calomel cell, a stirrer fitted with a mercury seal, the tip of the burette, and the mouth of a Jones reductor. All the tubes passing

through the cover are fitted tightly. The method of procedure adopted is as follows: (a) *Estimation of uranium by permanganate.* Ten c.c. of a solution of uranium as sulphate (containing about 0.013 gram U_3O_8 per c.c.) are added to 40 c.c. of sulphuric acid solution A (2 c.c. conc. acid, 88 water) and warmed at 80–90°. Carbon dioxide is passed through the titration cell to remove the air, and the uranium solution slowly passed through the Jones reductor into the cell, the reductor being finally rinsed with 50 c.c. of sulphuric acid A. The final solution at this stage has a volume of 100 c.c., contains 2 c.c. of concentrated sulphuric acid, and is brown. The stirrer is set in motion and the permanganate slowly added. The *E.M.F.* rises slowly at first and then jumps to a steady value which represents the point at which all the trivalent uranium has been converted into the quadrivalent condition and the solution has become green. Continued addition of permanganate does not affect the *E.M.F.* noticeably until the second end-point is reached, and here the *E.M.F.* jumps rapidly. At this point, the whole of the quadrivalent uranium has been converted into the hexavalent condition. The volume of permanganate used between the first and second end-points is equal to the theoretical amount of oxidising agent required to oxidise uranium from the quadrivalent to the hexavalent condition. Hence from this quantity the total amount of uranium present, no matter what its state of oxidation provided that none of it is hexavalent, can be calculated. (b) *Estimation of uranium and iron with permanganate.* The process is carried out as above; the first and second end-points give, as before, the quantity of uranium. After the uranium has all been oxidised to the hexavalent condition, the oxidation of the ferrous iron commences and when this has been completed the third end-point is observed. Hence the volume of permanganate required to carry the *E.M.F.* from the second to the third end-point is equivalent to the amount of iron present. (c) *Estimation of uranium by dichromate.* The process is carried out exactly as in the first case, and two end-points are obtained which have the same significance as in the other cases, but the slope of the second end-point is not so clear as in the other cases. All methods yield excellent results.

J. F. S.

The Estimation of Small Quantities of Gold as Colloidal Gold by the Colorimetric Method. J. A. MULLER and A. FOIX (*Bull. Soc. chim.*, 1922, [iv], 33, 717–720).—Small quantities of gold may be rapidly and accurately estimated colorimetrically after separation from other metals, provided that the concentration of the gold is not greater than 1/40 mg. per c.c., and that the colloidal particles are of the same size in the two solutions to be compared. The latter condition is realised by precipitation in identical circumstances. The presence of vanadium leads to inaccurate results.

H. J. E.

Separation of Palladium and Platinum by means of Dimethylglyoxime. C. W. DAVIS (*U.S. Bur. Mines, Repts. Investigations*, 1922, No. 2351).—The weighed metals are dis-

solved in a little aqua regia, the solution is evaporated to dryness, the residue moistened with a drop of hydrochloric acid, diluted with cold water, and dimethylglyoxime added with constant stirring until no further precipitate is formed. After keeping for a few minutes, the voluminous precipitate is collected on a filter, well washed with cold water, and ignited, first in air and finally in hydrogen to obtain pure palladium. The method was tested with quantities of each metal ranging from 0.15 to 11 mg., and the greatest divergence in five analyses was 0.03 mg.

CHEMICAL ABSTRACTS.

A Micro-modification of Benedict's Method for the Estimation of Reducing Sugar in Urine. MILLARD SMITH (*J. Lab. Clin. Med.*, 1922, 7, 364—368).—The modification, like the original method, is applicable only to urines containing not less than 0.17% of sugar. Exactly 1.0 c.c. of Benedict's solution is run from a pipette into a special test-tube, 0.2—0.7 gram of anhydrous sodium carbonate is added, and a pebble or piece of quartz. Urine is slowly added to the mixture, maintained at the boiling point, from a Mohr pipette, graduated to read directly the percentage of sugar present.

CHEMICAL ABSTRACTS.

Critical Study of Bang's Method for the Estimation of Reducing Substances in Blood. HENRI LABBÉ, F. NEPVEUX, and MÉNÉLAS NOMIDIS (*J. Pharm. Chim.*, 1922, [vii], 26, 49—63).—Bang's micro-titrimetric method for the estimation of reducing sugars in blood, consisting in oxidising with iodic acid the cuprous oxide produced by reduction by about 100 mg. of the blood absorbed in blotting paper and clarified, and estimation of the excess of iodic acid by titration with thiosulphate in presence of an alkali iodide, gives satisfactory results provided the following points are attended to. All the reagents used must be absolutely pure. Notably the potassium chloride and uranium acetate used for preparing the clarifying solution should be specially purified, and the iodate is preferably prepared by oxidising potassium iodide with permanganate. Mercuric sulphate, or alcohol, or Patein's reagent was found to give inferior results as clarifying agents, as they all retain a quantity of sugar, and at the same time partly compensate for this by allowing a certain amount of reducing substances originating from the albumin to remain in solution. Bang's reagent exhibits these faults in a much less marked degree. The heating of the clarified blood with the iodate-copper solution should be continued for exactly four minutes, and heating by means of a current of steam generated under constant well-defined conditions in a 1500 c.c. flask containing a constant level of water is to be preferred to direct heating, as experiments showed that the method of heating had a considerable influence on the results. The errors should not on the average exceed 5% if all the above precautions are taken, and the estimation of a large number of samples to which known weights of invert-sugar had been added gave a mean of 97% of the added sugar actually found.

G. F. M.

Acetol. I. A New Test for Carbohydrates. OSKAR BAUDISCH and HARRY J. DEUEL (*J. Amer. Chem. Soc.*, 1922, **44**, 1585—1587).—The test depends on the production of acetol from carbohydrates. An aqueous solution of the carbohydrate (0.1 gram in 100 c.c.) is treated with solid sodium hydrogen carbonate (5 grams) and distilled nearly to dryness. With the carbohydrates which are more readily attacked, the solution in the distilling flask very quickly turns brown, and the distillate has a sweet odour. The residue in the flask after distillation has a strong typical caramel odour. The distillate is treated with *o*-aminobenzaldehyde (30 mg.) and enough potassium hydroxide solution to give a distinctly alkaline reaction; after addition of a piece of porous plate, it is evaporated over a free flame to about 1/3 volume, cooled, and acidified with hydrochloric acid. Then solid sodium hydrogen carbonate is added until the solution is alkaline to litmus. The presence of acetol is shown by a strong blue fluorescence, which can be seen in daylight, but more readily in light of short wave-lengths such as that yielded by the iron arc. To confirm the test, the 3-hydroxy-2-methylquinoline may be extracted by shaking the solution several times with alcohol-free ether; the ethereal solution is dried with sodium sulphate, and the solvent removed. The colourless, needle-like crystals of 3-hydroxy-2-methylquinoline which remain dissolve readily in a little alcohol, and the fluorescence shows up strongly after the addition of water to the alcoholic solution. The sensitiveness of the reaction has not been determined except for dextrose, with which quantities as small as 5 mg. give a positive result.

The test is shown by arabinose, xylose, ribose, lyxose, dextrose, fructose, mannose, galactose, glucosamine, lactose, sucrose, maltose, and dextrin, but not by acetoacetic acid, glycerol, β -hydroxybutyric acid, lactic acid, starch, or glycogen.

The formation of 3-hydroxy-2-methylquinoline when treated with *o*-aminobenzaldehyde and potassium hydroxide appears to be a specific reaction of acetol; it is not given by pure methylglyoxal, aldol, acetone, acetaldehyde, ethyl acetoacetate, crotonaldehyde, or ethyl alcohol. H. W.

Estimation of the Cellulose Content of Wood and other Raw Materials by the Action of Chlorine and Carbon Tetrachloride. E. HEUSER and H. CASSEUS (*Papierfabr., Fest-u. Auslandheft*, 1922, 80—93; cf. A., 1921, i, 708; Frank, D.R.-P. 323936).—1.2 to 1.5 Grams of material of medium fineness are shaken with 50 c.c. of a saturated solution of chlorine in carbon tetrachloride for one to two hours, and the fibre is collected on a filter and washed successively with carbon tetrachloride, alcohol, and water. The lignin chloride is then dissolved out with a 2% solution of sodium sulphite, the residue washed with water, dried, and weighed. The product, which contains 3—3.5% of pentosan and 0.04—0.05% of ash, has a copper number of 0.8—0.91 as compared with 3.14—3.23 by the older method. The yield of pure cellulose was 56%.

CHEMICAL ABSTRACTS.

Estimation of α -, β -, and γ -Cellulose. OFFERMANN (*Paper*, 1921, 28, 19—25).—The cellulose content of the pulp is estimated as follows: 10 grams of pulp are covered with 50 c.c. of mercerising liquid (17—18% sodium hydroxide) and allowed to stand for thirty minutes. The fibre mass is then rubbed with 50 c.c. of water, filtered on a Buchner funnel with a fine cotton filter, and carefully sucked dry. The residue is washed with cold water until the wash-water no longer has an alkaline reaction; this usually requires ten to twelve washings, 50 c.c. of water being used each time. Finally, the α -cellulose is washed with hot dilute acetic acid, then with hot water (six to eight times), dried, and weighed. β -Cellulose is estimated by adding concentrated acetic acid to the alkaline filtrate until a marked acid reaction is produced, when β -cellulose separates in a finely divided state, and the colour of the brown filtrate becomes considerably lighter. To assist coagulation, the precipitate is heated on the water-bath at 100° until it settles. It is filtered on a fine cloth filter, washed six to eight times with hot water, transferred to a porcelain dish, dried, and weighed. γ -Cellulose is estimated by difference.

CHEMICAL ABSTRACTS.

Determination of the Iodine Value of Aliphatic and Aromatic Unsaturated Compounds. D. HOLDE (*Chem. Umschau*, 1922, 29, 253—254; cf. this vol., ii, 533).—Attention is directed to the different behaviour of the Hanus reagent with simple aliphatic oils such as sesame oil and mineral oils containing unsaturated cyclic structures. In the first case, even with varying excess of iodine, constant values are obtained, whilst in the second case variable values far above the theoretical are obtained with more or less considerable excess of the reagent. This is due to the rupture of bridge structures, and in certain circumstances the behaviour of an oil with excess of the Hanus reagent can be utilised to determine whether cyclic unsaturated compounds are present.
G. F. M.

The Sulphuric Acid Test for Liver Oils. J. C. DRUMMOND and A. F. WATSON (*Analyst*, 1922, 47, 341—348).—The substance present in liver oils which is responsible for the well-known purple coloration with sulphuric acid was found in the following species: Man, horse, ox, pig, cat, monkey, rabbit, guinea pig, chicken, duck, pigeon, rat, mouse, frog, shark, cod, haddock, ling, coal fish, dog fish, sprat, and skate. The substance appears to be a normal constituent of the liver, and is not derived from the bile or from products of autolysis or putrefaction. Evidence is presented to show that it is probably derived from the food, although an examination of the stages in the food of the cod did not reveal its ultimate origin with certainty. The chemical nature of the substance has not been ascertained. It forms a low proportion of the unsaponifiable fraction, is not cholesterol, and probably not a member of the lipochrome pigments. It is thermo-stable in the absence of air or oxygen, but is rapidly destroyed by oxidation. The few properties of the substance which are known, as

well as the available data regarding its distribution in natural products, show certain resemblances to the unidentified dietary unit known as vitamin-A; and, without assuming the identity of the two factors, it is suggested that the association may be of some significance. The colour test cannot be regarded as specific for liver fats, although they usually give the most intense reactions. The body fat and fat from other organs of animals, especially if they have been fed on liver oils, may give the reaction.

H. C. R.

Chemical and Physiological Investigation of the Fats and Lipoids of the Blood. II. The Kumagawa-Suto Method of Estimation of Lipoids. P. LEMELAND (*Bull. Soc. Chim. biol.*, 1922, 4, 300—321; cf. A., 1921, i, 633).—In the Kumagawa-Suto method, errors occur which are due to oxidation of the unsaturated acids during the heating and drying processes, and to loss of volatile constituents of the total unsaponifiable matter. The author promises to describe a new technique for the estimation.

E. S.

Estimation of Small Quantities of Furfuraldehyde Colorimetrically. PAUL FLEURY and GABRIEL POIROT (*J. Pharm. Chim.*, 1922, [vii], 26, 87—96).—Small quantities of furfuraldehyde may be estimated colorimetrically with an accuracy of about $\pm 1\%$ by means of the blue coloration which is formed in acidic medium with orcinol in a suitable solvent, preferably acetic acid, and is intensified by the presence of traces of iron. The reagents required are hydrochloric acid (*d* 1.19) containing 20 mg. of iron as ferric chloride per litre; glacial acetic acid containing 1 gram of orcinol in 1600 c.c., and a standard solution of furfuraldehyde containing 1 gram in 10 litres of 1% acetic acid. One c.c. of the standard solution or of the solution to be estimated is mixed with 4 c.c. of the acetic acid-orcinol solution and 5 c.c. of the hydrochloric acid, and warmed for one minute in a water-bath. After keeping for thirty minutes, the maximum coloration is developed, and comparison is made with the standard in a Duboscq colorimeter. The reaction has a sensibility of about 4 mg. of furfuraldehyde per litre.

G. F. M.

New Analytical Applications of Nessler's Reagent. Characterisation of Ketones; Estimation of Aldehydes. J. BOUGAULT and R. GROS (*J. Pharm. Chim.*, 1922, 26, 5—11).—Certain ketones, such as acetone, methyl ethyl ketone, cyclohexanone, trimethylcyclohexenone, acetophenone, and *p*-methoxyacetophenone, give with Nessler's reagent in the cold a yellowish-white precipitate which dissolves in acid or on the addition of potassium cyanide with liberation of the ketone. This reaction may be used for the detection of acetone in urine, by suspending a small capsule, containing 10 c.c. of Nessler's reagent, over 200 c.c. of urine, previously acidified, the whole being covered by a bell-jar. The time required for the formation of a precipitate varies inversely as the concentration of the acetone in the urine.

Certain aldehydes react quantitatively with Nessler's reagent according to the equation $R \cdot CHO + H_2O + 2I = 2HI + R \cdot CO_2H$, the iodine coming from the mercuric iodide. To a known volume of the aldehyde solution, corresponding with 0.01—0.05 gram of the aldehyde, is added an excess (about 30 c.c.) of Nessler's solution, and 10 c.c. of sodium hydroxide solution. The mixture is well shaken, and after some time is neutralised with hydrochloric acid. A known volume of standard iodine solution is added, and then the excess of iodine is titrated back with standard thiosulphate solution. It is necessary to make a control estimation on the sodium hydroxide. Accurate results were obtained with formaldehyde, furfuraldehyde, benzaldehyde, and piperonaldehyde, but the method is not satisfactory for vanillin or acetaldehyde.

W. G.

The U.S.P. Test for Acetone in Alcohol. J. W. E. HARRISSON (*J. Amer. Pharm. Assoc.*, 1922, **11**, 16).—Some zinc salts and antimony compounds interfere with the test, which depends on the development of a violet colour in presence of sodium hydroxide and sodium nitroprusside. The best test consists in the addition of 1 c.c. of potassium hydroxide solution to 5 c.c. of the ethyl alcohol, followed by a solution of iodine in potassium iodide until coloration occurs; heat must not be used. A precipitate is obtained immediately if 0.25% of acetone is present.

CHEMICAL ABSTRACTS.

Analysis of Camphor Preparations. MAURICE FRANÇOIS and ÉMILE LUCE (*J. Pharm. Chim.*, 1922, [vii], **25**, 500—507).—Tincture of camphor (French Codex) contains 10 grams of camphor dissolved in 90 grams of 90% alcohol, and should have $\alpha +6.5^\circ$ in a 200 mm. tube. The optical rotation varies somewhat with the strength of the alcohol, but, provided that the alcoholic concentration is correct, a lower rotation than that mentioned will indicate a deficiency in the camphor content or that artificial (inactive) camphor is present. The optical rotation of camphor decreases slightly with increase of temperature and in alcoholic solution the value decreases considerably for a decrease in the alcoholic concentration. For instance, in absolute alcohol camphor has $\alpha_D +43.0^\circ$, and in 90% alcohol $\alpha +39.66^\circ$. To ascertain whether a low value for the optical rotation of a sample of tincture of camphor is due to deficiency in camphor or to the presence of artificial camphor, 50 c.c. of the sample are evaporated under a bell-jar over sulphuric acid, the crystalline residue is dried for a further forty-eight hours over sulphuric acid, and its optical rotation then determined in alcoholic solution; natural camphor has $\alpha_D +43^\circ$. The alcoholic strength of tincture of camphor may be ascertained from the specific gravity, allowance being made for the fact that each 1% of camphor present increases the specific gravity by 0.00116.

W. P. S.

Estimation of Cyanogen. HIDEKICHI YANAGISAWA (*J. Pharm. Soc. Japan*, 1922, 369—377).—The method is based on

the conversion of cyanogen into ammonia. A cyanogen compound is heated with dilute sulphuric acid in a sealed tube at 200° during four hours; after cooling, the product is transferred to a Kjeldahl flask, and after making alkaline with sodium hydroxide, the ammonia is distilled into $N/4$ -sulphuric acid; the excess is titrated with $N/4$ -ammonia as usual. K. K.

Comparison of Methods for the Estimation of Urea. J. CARRA (*Biochim. terapia sper.*, 1921, 8, 225).—A comparison of various methods for the estimation of urea shows that a gravimetric method based on the formation of a compound of 1 mol. of carbamide and 2 mols. of xanthidrol is extremely exact. The urease method is also excellent. CHEMICAL ABSTRACTS.

Influence of Sodium Fluoride on the Estimation of Urea by the Xanthidrol Method. M. POLONOVSKI and C. AUGUSTE (*Compt. rend. Soc. Biol.*, 1922, 86, 1027—1028; from *Physiol. Abstr.*, 1922, 7, 214).—The presence of sodium fluoride renders the results obtained by this method too high. This is probably due to the liberation by the acetic acid of hydrofluoric acid. The error may be avoided by the removal of the fluoride by calcium acetate. W. O. K.

Microchemical Detection of Tryptophan in the Plant. FRITZ KRETZ (*Biochem. Z.*, 1922, 130, 86—98).—Tryptophan can be detected in plant-tissues by Fürth's modification of Voisenet's reaction, if the tissues be preserved in a silica gel by soaking in sodium silicate and subsequent addition of concentrated hydrochloric acid. Tryptophan is found in moulds and richly in the embryonic tissues of higher plants and in all other parts containing protein. H. K.

Uric Acid. I. Examination of the Variables in the Folin and Wu Uric Acid Method. II. A Modification of the Folin and Wu Uric Acid Method. GEORGE W. PUCHER (*J. Biol. Chem.*, 1922, 52, 317—327, 329—334).—Using Folin and Wu's method (*A.*, 1919, ii, 308) for the estimation of uric acid in blood, only 75% of added uric acid was recovered. When, however, the solution was heated before filtering the coagulated proteins, a recovery of 93% was obtained. E. S.

Rapid Estimation of Uric Acid in Urine. E. VAILLANT (*Ann. Chim. Analyt.*, 1922, [ii], 4, 178—180).—The liquid is titrated with a standard iodine solution made by dissolving 3.78 grams of re-sublimed iodine and 15 grams of potassium iodide in a little water and diluting to 1 litre. Two c.c. of this solution oxidise 0.005 gram of uric acid (to alloxan and urea). The analysis is carried out by measuring into a small tube 2 c.c. of iodine solution and 1 c.c. of carbon disulphide and allowing the urine to fall a drop at a time from a burette, shaking the tube after each addition, until the carbon disulphide becomes first pale rose and then colourless. The uric acid content per litre of urine is then calculated by dividing the number of c.c. used into 5. Sugar and small amounts of albumin do not interfere; larger amounts of albumin should be

removed by heating to coagulate and then filtering. If the urine is alkaline, a few drops of acetic acid are added; if it contains crystals of uric acid, it is warmed until they dissolve and the analysis is conducted at 25–30°. A. R. P.

Direct Estimation of Uric Acid in Urine. STANLEY R. BENEDICT and ELIZABETH FRANKE (*J. Biol. Chem.*, 1922, **52**, 387–391).—The same reagents are employed as in the method for blood (this vol., ii, 405). To 10 c.c. of the diluted urine (containing 0.15 to 0.30 mg. of uric acid) are added 5 c.c. of the sodium cyanide solution and 1 c.c. of the arsenophosphotungstic acid reagent. After five minutes, the mixture is diluted to 50 c.c. and the colour compared with a standard containing 0.2 mg. of uric acid similarly diluted to 50 c.c. Owing to the larger amount of uric acid present in urine than in blood, it is unnecessary to heat the solution. Of the other constituents of urine, albumin appears to be the only one which interferes with the reaction. This may be removed, if present, by coagulation by heat in the presence of a drop of acetic acid. E. S.

Rapid Estimation of Quinine Salts. S. G. LIVERSEDGE and F. W. ANDREWS (*Pharm. J.*, 1922, **109**, 92–94).—Quinine in quinine salts is rapidly estimated by dissolving the alkaloidal salt in 20 c.c. of *N*/5-sulphuric acid, a known excess of *N*/2-sodium hydroxide is then added and the liberated quinine extracted with ether saturated with water, and subsequently the excess of alkali titrated with *N*/5-sulphuric acid, using phenolphthalein as indicator. The amount of acid originally combined with quinine is thus determined, whence the percentage of quinine or quinine salt is obtained by calculation. If boric acid is present, hæmatoxylin is used as indicator. G. F. M.

Estimation of Small Quantities of Atropine in Blood-serum. H. C. VAN DER HEYDE (*J. Lab. Clin. Med.*, 1922, **7**, 280–287).—Atropine was estimated in 1 c.c. of serum in maximum concentrations of 0.5%. Proteins are first precipitated with absolute alcohol, the precipitate being washed several times with absolute alcohol, and the alcoholic filtrates evaporated to dryness. The residue is treated with a small quantity of water and the solution filtered from fats on a microfilter. The dilution is then determined in which Mayer's reagent (potassium mercury iodide) will just give a precipitate, and compared with control solutions of known dilution. CHEMICAL ABSTRACTS.

Method for the Separation of Strychnine from Quinine. NORMAN EVERS (*Pharm. J.*, 1922, **109**, 90–91).—A practically quantitative separation of strychnine from quinine is achieved by taking advantage of the fact that chloroform extracts strychnine hydrochloride from strongly acid aqueous solutions, but does not remove quinine hydrochloride under similar conditions. The solution of the mixed alkaloids is acidified with hydrochloric acid to make it about 2*N*, and is extracted with five successive quantities of 50 c.c. each of chloroform. The chloroform extracts are united,

concentrated to 50 c.c., shaken with dilute aqueous ammonia, and evaporated to dryness. The residue is almost pure strychnine, and the quinine is extracted from the residual acid solution with chloroform after rendering alkaline in the usual way. G. F. M.

Detection of Thymine in the Presence of Sugar. HARRY J. DEUEL and OSKAR BAUDISCH (*J. Amer. Chem. Soc.*, 1922, **44**, 1581—1584).—A modification of the method, described recently by Johnson and Baudisch (this vol., ii, 238) for the detection of thymine, is given which makes it applicable in the presence of sugar. For this purpose, the thymine is precipitated as the mercury salt by the addition of saturated mercuric chloride solution (10 c.c.) and sufficient sodium hydroxide solution to make the mixture distinctly alkaline. The precipitate is separated by means of a centrifuge, and, after pouring off the supernatant liquid, it is thoroughly mixed with about 100 c.c. of distilled water and again centrifuged. It is then suspended in water (100 c.c.) and decomposed with hydrogen sulphide; the precipitated mercuric sulphide is filtered and excess of hydrogen sulphide is removed by boiling the filtrate. The remaining liquid is cooled and made up to 100 c.c. Oxidation of the thymine is effected by treating the solution with sodium hydrogen carbonate and ferrous sulphate in the presence of air and the test is completed as described previously. Thymine in quantities of 10—15 mg. may readily be detected in the daylight, whilst as small a quantity as 1 mg. may be detected by the use of the iron arc light. H. W.

Recognition of Proteins and Derivatives by Colour Reactions. M. A. RAKUSIN (*Biochem. Z.*, 1922, **130**, 268—281).—The author has examined a large number of animal and vegetable proteins, ferments, and toxins in respect of their behaviour towards eight colour reagents, namely, biuret, Millon's, xanthoprotein, Liebermann's, Adamkewitsch's, Molisch's, Pettenkofer's, and Ostromysslenski's, and the behaviour towards the sulphur reaction of Vohl and Rakusin. Each protein corresponds with a definite complex of reactions. H. K.

Estimation of Small Amounts of Albumin in Urine. G. GÉRARD (*Bull. sci. pharmacol.*, 1921, **28**, 466—467; from *Physiol. Abstr.*, 1922, **7**, 158—159).—Ten c.c. of urine are placed in each of two test-tubes. One is warmed and a few drops of trichloroacetic acid are added, giving the usual turbidity when albumin is present. To the other there is added drop by drop officinal spirit of peppermint. The tube and contents are shaken after each drop is added, and the turbidity of the two tubes is compared. When the opacity of both is alike, the test is completed. The author has found that 1 drop of the alcohol used is equivalent to 0.04 c.c. of albumin. No results are given. W. O. K.

The Differential Precipitation of the Proteins of Colostrum and a Method for the Estimation of the Proteins in Colostrum. PAUL E. HOWE (*J. Biol. Chem.*, 1922, **52**, 51—68).—When cow's colostrum is precipitated by addition of successive small quantities of sodium sulphate, a precipitation curve is

obtained which shows three "critical zones" at which increases in the concentration of sodium sulphate cause only small increases in the amount of protein precipitated. These zones occur at concentrations of 13.5–14.5, 17.4–18.4, and 21–22% of sodium sulphate. Up to concentrations of 13.5% euglobulin is precipitated, between 14.5 and 17.4% a mixture of caseinogen and ψ -globulin I is obtained, whilst the precipitate produced between 18.4 and 21% is termed ψ -globulin II. After removal of euglobulin, caseinogen may be precipitated separately by addition of either acetic acid or alum. Total precipitation, including albumin, is effected by means of trichloroacetic acid. By combining these precipitations with Kjeldahl estimations on the filtrate before and after precipitation, a method for the estimation of the proteins of colostrum and of milk is developed (cf. Woodman, A., 1921, i, 625).
E. S.

Estimation of Globulins in Blood-serum. R. R. HENLEY (*J. Biol. Chem.*, 1922, 52, 367–375).—Two methods are described. In the first, the globulins are precipitated by saturation with magnesium sulphate and the non-globulin nitrogen is estimated in the filtrate. The difference between this and the total nitrogen gives the globulin nitrogen. In the second method, the globulins are precipitated by half saturation with ammonium sulphate, filtered, dissolved in salt solution, and again precipitated. The precipitate is once more dissolved in salt solution and the globulins are finally precipitated by boiling, one or two drops of 10% acetic acid being added if necessary. The precipitate is then filtered on to a tared, hardened filter-paper, washed, dried at 100°, and weighed. The results obtained by both methods vary slightly according to the dilution employed, but are otherwise in agreement with those obtained by Cullen and Van Slyke's method (A., 1920, ii, 398).
E. S.

New Method for the Detection of Bile Pigments. AL. IONESCU and C. POP (*Bul. Soc. Chim. Romania*, 1922, 4, 20–26).—The following method is recommended as a quick and satisfactory means of detecting bile pigments in urine. To 10 c.c. of the urine are added 2 c.c. of a 0.5% solution of albumin and a few drops of 10% acetic acid. The mixture is boiled and filtered. The precipitate is washed with water and to it on the filter-paper is added one drop of a mixture of 4 c.c. of 6% potassium dichromate and 1 c.c. of 20% sulphuric acid. If the urine contains bile pigments the yellow precipitate turns a green or bluish-green colour. The depth of the colour depends on the amount of pigment present. The reaction is characteristic and very sensitive and is not given by other normal or abnormal urinary pigments.
W. G.

Estimation of Bilirubin in Human Serum. The Ehrlich-Pröscher Reaction. J. S. THANNHAUSER and E. ANDERSEN (*Deut. Arch. klin. Med.*, 1921, 137, 179–186; from *Chem. Zentr.*, 1922, ii, 425).—The colorimetric method of Hijmans van den Bergh, in which, after precipitation of protein from serum by means of ethyl alcohol, bilirubin is directly combined with diazobenzene-

sulphonic acid, is considered to be inexact owing to precipitation of a portion of the bilirubin by ethyl alcohol and also owing to the coloration obtained differing from that of the test solution. The latter difficulty can be avoided by the presence of excess of mineral acid. Directions are given for the estimation of bilirubin in different sera.

G. W. R.

Estimation of Trypsin. A Modification of Gross's Method. SOTARO KAI (*J. Biol. Chem.*, 1922, 52, 133—136).—The relative strengths of trypsin are estimated by measuring the time taken by equal amounts of each to digest 25 c.c. of a casein solution, the time of digestion being inversely proportional to the concentration of trypsin. The end-point of digestion is determined by adding a sodium hydroxide-acetic acid solution to test portions withdrawn from time to time, no precipitate being produced when digestion is complete. By arbitrarily choosing one preparation as standard, the concentrations of others may be expressed in terms of this.

E. S.

Acidosis. XVIII. Estimation of the Hydrogen Carbonate Concentration of the Blood and Plasma. DONALD D. VAN SLYKE (*J. Biol. Chem.*, 1922, 52, 495—499).—Two methods are described. In the first, total carbon dioxide is estimated by Van Slyke and Stadie's method (this vol., ii, 78) and P_H either electrometrically or colorimetrically by Cullen's method (following abstract). The hydrogen carbonate concentration is then calculated from the equation $[BHCO_3] = [CO_2]/(1 + 10^{P_K - P_H})$, which is derived from $[H'] = K' \cdot [H_2CO_3]/[BHCO_3]$, P_K being the negative logarithm of K' . The second is a titration method. One c.c. of plasma is treated with 5 c.c. of 0.01N-hydrochloric acid and shaken in a 100 c.c. round flask to remove carbon dioxide. It is then brought back to the original alkalinity by the addition of 0.01N-sodium hydroxide, phenol-red being used as indicator and 1 c.c. of the original plasma as standard. Sodium chloride solution (0.9%) is used for the preparation of the standard acid and alkali, and also for the dilution, where necessary, of the plasma. Special precautions must be observed to prevent loss of carbon dioxide from the standard.

E. S.

Acidosis. XIX. Colorimetric Estimation of the Hydrogen-ion Concentration of Blood Plasma. GLENN E. CULLEN (*J. Biol. Chem.*, 1922, 52, 501—515).—The hydrogen-ion concentration of diluted plasma or serum is estimated colorimetrically at 20° by comparison with Sørensen's phosphate standards, using phenol-red as indicator, and is then converted into that for undiluted plasma at 38° by use of an empirical factor. The latter has been determined by direct comparison of colorimetric with electrometric estimations, and has been found to be constant for plasma of the same species. It is different, however, for plasma of different species and also for plasma and serum of the same species. The success of the method depends on the use of a technique, which is described in the original, to prevent the loss of carbon dioxide during both the preparation of the plasma and the estimation.

E. S.

General and Physical Chemistry.

The Reflexion of X-Rays by Crystals. R. DARDORD (*J. Phys. Radium*, 1922, [vi], 3, 218—220).—When X-rays are reflected from a cylindrical crystalline surface, such as may be obtained by bending a sheet of mica, the analogy between the reflexion of X-rays and light becomes closer; the only difference is that the reflexion of the former is selective. An image of a rectilinear source of X-rays may be obtained from a cylindrical surface and received on a photographic plate. The theory of the formation of the image is discussed and this verified experimentally. W. E. G.

Errors arising in the Measurement of Unsymmetrical Spectrum Lines. T. R. MERTON and D. N. HARRISON (*Proc. Roy. Soc.*, 1922, [A], 101, 431—434).—One of the most serious sources of error in the measurement of the spectrum lines by juxtaposition of an arc spectrum is the drift of the spectrum across the plate due to either mechanical displacement or changes in the temperature of prism or grating. The methods frequently employed to reduce these errors are not satisfactory for unsymmetrical lines. It is shown that the position of the maximum of an unsymmetrical line in the case of a uniform drift lies at the intersection of the intensity distribution curves at the beginning and the end of the exposure, and hence in the comparison with symmetrical lines, errors amounting to a large fraction of the displacement may easily occur. For this reason, the spectra of gases at low pressures, in which the lines are known to be symmetrical, are to be preferred as standards of wave-length. W. E. G.

Atomic Hydrogen and the Balmer Series Spectrum. R. W. WOOD (*Phil. Mag.*, 1922, [vi], 44, 538—546).—From the investigation of the hydrogen spectrum developed by an electric discharge in long vacuum tubes containing hydrogen (A., 1921, ii, 665), it is concluded that the secondary spectrum is emitted by the hydrogen molecules, the Balmer series by the atoms. The latter series has been extended to the 20th line by abolishing the secondary spectrum and the faint continuous background of illumination and employing a sufficiently long exposure. It appears extremely probable that the part played by water vapour in enhancing the development of the Balmer series consists in supplying a "poison" (oxygen) to the catalysing wall of the discharge tube, thus permitting a high concentration of atomic hydrogen in the tube under the action of a heavy discharge. With a feeble discharge, the secondary spectrum predominates, as atomic hydrogen is not formed sufficiently fast to overtake the catalysing power of the tube wall. The whiteness of the discharge when the discharge tube is immersed in liquid air, observed by Merton, is due to the increased catalysing power of the wall at low temperatures. The catalysis theory is

likewise applicable to explain the characteristic spectrum developed when air or nitrogen is admitted to the discharge tube containing hydrogen, and the more complete development of the Balmer series in the solar corona and probably in nebulae. The wave-lengths of the components of the doublet H_γ were found to be 4340.494 and 4340.435 Å. J. S. G. T.

The Emission Spectrum of Monatomic Iodine Vapour. ST. LANDAU-ZIEMECKI (*Phil. Mag.*, 1922, [vi], **44**, 651—656).—The band spectrum of iodine vapour at low temperatures showed distinct maxima at $\lambda\lambda$ 422, 404, 395, 387, 380, 373, 368, 363, 340, 325, 323, 320, 319, 312, 310, 306, 304, 302, and 299 $\mu\mu$. The band spectrum became gradually fainter and the line spectrum developed as the temperature was raised. At 960—1000°, corresponding with almost complete dissociation of the iodine molecules, the band spectrum disappeared entirely and was replaced by the line spectrum, due to atomic iodine vapour, and consisting of wave-lengths 4868, 4769, 4680, 4324, 4132, 4100, 3384, 3281, 3081, 2879, 2593, 2583, and 2566 Å. Of these, λ 3281 Å. is certainly new. It is concluded that the band spectrum originates in the molecule, and the line spectrum in the atom. J. S. G. T.

The Spectrum of Helium in the Extreme Ultra-violet. THEODORE LYMAN (*Nature*, 1922, **110**, 278—279; cf. Fricke and Lyman, A., 1921, ii, 362).—When an improved apparatus is used, the line at 584.4 is of very great strength, and is accompanied by three new lines at 537.1, 522.3, and 515.7, of which the intensities decrease with the wave-length in a manner strongly suggesting a series relation. The spacing of these lines on the frequency scale is identical with that of the first four lines in the singlet principal series. The line at 584 is therefore regarded as the first member of a principal series, $oS-mP$. The wave-lengths of the new lines, however, are not in complete agreement with the accepted values of the resonance and ionisation potentials. A. A. E.

Some Spectrum Lines of Neutral Helium derived theoretically. LUDWIK SILBERSTEIN (*Nature*, 1922, **110**, 247—248).—Owing to the prohibitive nature of the general problem of three or more bodies, Bohr's quantum theory can account only for those spectrum lines which form a series of the simple Balmerian type, that is (apart from X-ray spectra of the higher atoms), only those of atomic hydrogen and ionised helium. An attempt is therefore made to obtain a special solution to the problem in the case of neutral helium, the spectrum of which, containing some 105 lines, has not been accounted for theoretically. Assumption that the nucleus and the two electrons are always collinear with each other, the latter describing two equal and oppositely situated ellipses round the former, is evidently erroneous, since the corresponding formula does not cover a single observed line of helium. On the other hand, assumption that the mutual perturbation of the two electrons is negligible yields a spectrum formula which covers about forty lines of the spectrum of neutral helium, and will

probably be found to cover fifty more lines. It therefore appears that the electrons within the atom do not repel each other even with a small fraction of the force usually attributed to them. Lithium appears to yield similar, although less striking results. It is concluded that the field of force of a bound electron is entirely engaged by the nucleus, at least in the case of helium and probably lithium, but possibly also in that of the higher atoms. A. A. E.

Spectrum Lines of Neutral Helium. W. M. HICKS (*Nature*, 1922, 110, 309).—The constitution of the secondary spectrum of hydrogen may be examined by a method similar to that used by Silberstein (preceding abstract) for helium. Practically the whole of this spectrum depends on the sequence of the Balmer series. If $f(m)$ denote the m th sequent, the wave-length of any line is of the form $\Sigma k_m f(m)$, where the k_m are positive or negative integers. The spectrum is a kind of linkage spectrum in which the usual links are replaced by the separations between the successive lines of the primary, namely, 5331.57, 2467.75, etc. In Silberstein's theory, as in all orbital theories, the two electrons are considered to be moving independently, and each to pass between two of its corresponding paths; if, however, the combined change of energy is radiated, these two events must be simultaneous. A. A. E.

Ionisation in the Solar Atmosphere. HENRY NORRIS RUSSELL (*Astrophys. J.*, 1922, 55, 354–359).—Barium is much more highly ionised in the sun than sodium, although their ionisation potentials are the same. The absorption of photospheric radiation by atoms in the solar atmosphere tends to increase the degree of ionisation, both directly, by shifting an electron into a position from which its removal is easier, and indirectly, when enhanced lines are absorbed, by getting the ionised atoms into states in which they are probably less likely to combine with electrons. The case of lithium, however, is anomalous. Barium lines and enhanced rubidium lines in the solar spectrum are tabulated. The ionisation potential of manganese (7.6 volts) agrees well with the value predicted from the behaviour of its lines in the solar spectrum. A. A. E.

The Spectrum of Glucinum and its Noteworthy Relation to the Spectrum of Aluminium. L. C. GLASER (*Ann. Physik*, 1922, [iv], 68, 73–88).—Employing the arc or spark between electrodes of glucinum, the author has investigated the spectrum of the metal in the visible and ultra-violet regions. The line spectrum observed confirms, with certain limitations, the previous measurements of Rowland and Tatrall. In the region $\lambda\lambda$ 5500–4900 Å., no lines, apart from those constituting part of the band spectrum, attributable with certainty to glucinum, were observed. The line spectrum observed was constituted of $\lambda\lambda$ 4607?, 4600?, 4572, 4486.7?, 4277.8?, 4263.6?, 4216.0, 4079.0, 3994.0, 3936.0, 3909.3, 3891.0. A more accurate measurement of the strongest line, λ 4572, gave the result 4572.73 ± 0.0098 Å., in terms of the normal international iron spectrum. Attention is directed to

certain analogous physical features characterising the production of the band spectra of glucinum and aluminium. The wave-lengths of the constituent bands in the spectra of both elements are tabulated, and it is remarked that there is considerable agreement between the respective first differences of wave-lengths of successive lines in corresponding bands in the two spectra. Deslandre's rule is true only as a first approximation. Attention is also directed to the approximate constancy of certain numerical values calculated from the wave numbers of the heads of the bands $\lambda\lambda$ 4708 and 4842 in the case of the glucinum and aluminium spectrum, respectively. It is concluded that the atomic structures of glucinum and aluminium, so far as the emission of their band spectra is concerned, are the same. J. S. G. T.

The Band Spectra of Cadmium. ERIK HULTHÉN and ERNST BENGTTSSON (*Compt. rend.*, 1922, **175**, 423—426).—Two systems of bands have been measured at $\lambda=4500$ and $\lambda=4300$ and their structures studied. The results show that the bands of the elements zinc, cadmium, and mercury can be embodied in the same expressions, and the constants of the series in the bands of cadmium are given. W. G.

A Continuous Spectrum from Mercury Vapour. C. D. CHILD (*Astrophys. J.*, 1922, **55**, 329—344; cf. A., 1921, ii, 3).—A continuous spectrum identical with the fluorescence spectrum is emitted in addition to the ordinary line spectrum when (a) the temperature of the vapour is above 120° and below $300\text{--}400^\circ$, depending on the pressure, (b) the pressure is above 1 mm., (c) the current density is low, (d) very little air or other contaminating gas is present. Ionisation of the vapour is not, however, necessary. It is suggested that the carriers of the spectrum are molecules consisting of two or more atoms, which emit the fluorescence spectrum as a result of excitation by ultra-violet radiation λ 2536 emitted by atoms struck by electrons with energies corresponding with at least 4.9 volts. The ionisation potential for mercury appears to vary with the temperature, reaching a minimum at about 140° . Above 120° , mercury vapour reacts with air to form a dark compound; the reaction is associated with the appearance of the continuous spectrum. A continuous spectrum of sodium, much fainter than the line spectrum, has also been observed.

A. A. E.

The Lanthanum Violet Bands and the Associated Lines. JUNZŌ ŌKUBO (*Sci. Rep. Tohoku Imp. Univ.*, 1922, **11**, 95—104).—The edges of the bands in the violet and green regions of the lanthanum spectrum consist of closed doublets, and the edges of these are determined accurately in order to seek a relation between the band and line spectra. The best conditions for observing the bands and diffused lines are obtained when the temperature of the arc is low, or by using lanthanum as the negative electrode with a moderate current; under these conditions, the sharp lines are comparatively faint. It was found that one pair of double

lines exists nearly midway between the corresponding heads of each series of bands, and another between the heads of successive series so that the bands are not independent of each other. By the introduction of the quantum theory, the conclusion is drawn than the lanthanum violet bands are emitted as a consequence of the molecular rotation when line emission takes place in the atom, thus accounting for the double structure of the bands.

W. E. G.

The Arc Spectra of Gallium and Indium. H. S. UHLER and J. W. TANCH (*Astrophys. J.*, 1922, 55, 291—301).—By the use of a concave grating with 590 lines per mm. and a radius of curvature of 640 cm., accurate determinations were made of the wave-lengths for air and the vacuum wave numbers of twenty-three gallium lines and thirty-four indium lines, λ 4511 to λ 2170. All the gallium lines and twenty-eight of the indium lines were found to belong to the two subordinate series, for each of which from five to eight terms were identified. Four indium lines, previously supposed to be single, were resolved. An attempt to obtain the absorption spectrum of indium vapour gave only the two strongest lines, λ 4511 and λ 4102.

A. A. E.

Optical Experiment Demonstrating the Bragg Method of Crystal Analysis. BILFRIED QUARDER (*Physikal. Z.*, 1922, 23, 350—352).—An optical analogue to the distribution of intensities in the various orders of X-ray spectra reflected from the 1, 1, 1, planes of sodium chloride or the 1, 0, 0 planes of zinc sulphide is afforded by the use of a photographic grating consisting of equally spaced, alternately thick and thin lines. By reducing the grating aperture, broadening of the lines occurs, a result analogous to the broadening of the lines in the X-ray spectrum, occurring on powdering the diffracting crystals, as observed by Debye and Scherrer (*A.*, 1917, ii, 437).

J. S. G. T.

Spectra of X-Rays and the Theory of Atomic Structure. IV and V. D. COSTER (*Phil. Mag.*, 1922, [vi], 44, 546—573).—In continuation of previous work (*A.*, 1922, ii, 491), the author has investigated the *L*-series of the characteristic X-ray spectra of the rare elements from lanthanum (atomic number 57) to lutecium (atomic number 71). In general, the lines for these elements are much sharper than those for elements of lower atomic number, and most of the stronger lines are accompanied by fairly intense satellites. Measurements have been made of a great number of lines which may be arranged in a simple diagram and are conveniently referred to as diagram lines, a number of non-diagram lines, and some absorption discontinuities. The measured frequencies of the diagram lines are in agreement with results to be anticipated from Bohr's theory of the successive development of the shells of electrons in the various atoms. Most of the non-diagram lines lie on the short wave-length side of an intense diagram line, and the difference of frequency between the two lines is approximately proportional to the atomic number of the element.

A theoretical discussion is given of the origin of the non-diagram lines and of the fine structure of the absorption discontinuities. In criticism of the views of Wentzel (this vol., ii, 249) in this connexion, experimental results have been obtained showing that the assumption of the successive ionisation of the inner shells of the atom is untenable, and that part of the non-diagram lines are emitted by atoms which have lost more than one electron at the same time by the impact of a single high-speed β -particle. An explanation is given of the origin of some non-diagram lines lying on the long wave-length side of diagram lines, and it is suggested that these lines should appear both in the emission and in the absorption spectra. J. S. G. T.

Analysis of the Electronic Structure of the Elements.

A. DAUVILLIER (*J. Phys. Radium*, 1922, [vi], 3, 221—251).—The author has investigated the *L*-series of lines in the characteristic X-ray spectra of uranium, gold, platinum, iridium, osmium, tungsten, and antimony, and the *K*-lines in the spectra of copper and molybdenum (cf. A., 1921, ii, 669; this vol., ii, 43, 101, 559). The numbers of observed lines have been at least doubled, and owing to the correction of certain errors, it is shown that Sommerfeld's theory is applicable to the calculation of the frequencies of these lines. The results have been applied to determine the energy levels of the atoms of most of the elements and afford a complete qualitative explanation of their absorption spectra; they have likewise been correlated with the ionisation potentials of certain gases and metallic vapours, by the application of the quantum hypothesis. The numbers of electrons in the various corpuscular layers of the respective elements have been calculated, and it is shown that the number n of such electrons is related to the quantum number τ characterising the layer by the equation $n=2\tau^2$. A theory is advanced to account for the complex structures of the *K* and *L* series of lines in the X-ray spectra of the light elements, and these spectra are, by the aid of the theory, correlated with the corresponding spectra of the elements in the visible region. The variation of atomic volume and chemical valency of the elements, as shown by the periodic table, is also at least partly explained by the theory developed. J. S. G. T.

The Absorption Spectrum of Potassium Vapour.

SNEHAMOY DATTA (*Proc. Roy. Soc.*, 1922, [A], 101, 539—547).—The principal series lines in the absorption spectrum of potassium vapour have been measured up to $m=42$ (compared with $m=24$ previously). Satisfactory agreement is shown between the measured frequencies and those calculated by the series formula $p_1(m)=35008\cdot480-109736\cdot1/(m+1\cdot29667-0\cdot06154/m)^2$, except for $m=3$ and $m=39$ to 42. The value for $m=3$ is abnormal, and the latter values receive a possible explanation along the lines of Sommerfeld's assumption of the occurrence of collisions between the electrons emitted from the atom and other electrons or atoms. The first seven members of the principal series have been resolved, compared with five previously. New absorption lines, first

appearing at 20 mm. pressure, and apparently not corresponding with known lines in the emission spectrum, were found at higher pressures. The combination lines $1s-2d$ and $1s-3d$ were found to be absorbed, the first as a pair, confirming the presence of a satellite to the lines of the diffuse series. The appearance of these absorption lines contradicts the selection principle. J. S. G. T.

Absorption of Potassium Vapour in the Associated Series.

A. L. NARAYANA and D. GUNNAIYA (*Nature*, 1922, **110**, 250).—Traces of absorption occur at about 1100° . In the diffuse series, distinct traces of absorption have been detected; the bands 5780, 5340, 5300, and 5100 apparently correspond with 5782, 5340, 5323, and 5100 of $(2p-m.d)$. A. A. E.

Absorption of Light by Sodium and Potassium Vapours.

GEORGE R. HARRISON (*Proc. Nat. Acad. Sci.*, 1922, **8**, 260—263).—Vapour density alone determines the development of the continuous absorption spectrum of sodium vapour. No visible effect is attributable to hydrogen present in pressures up to 400 mm. of mercury. Absorption begins at the last resolved line of the principal series. No definite break in the continuous absorption spectrum was observed, but the absorption decreases rapidly with decreasing wave-length, the region λ 2200 Å. being one of the most transparent regions studied. With unsaturated sodium vapour, the band spectrum is much weaker than the line and continuous spectra. It is suggested that the bands are due to loose molecular aggregates. Fluorescence, which was marked in saturated vapour, disappeared on superheating the vapour. The continuous absorption spectrum of potassium vapour resembled that of sodium. J. S. G. T.

The Absorption and Fluorescence Spectra of Benzene.

VICTOR HENRI (*J. Phys. Radium*, 1922, [vi], **3**, 181—211).—The ultra-violet spectrum of benzene has been studied over pressures ranging from 0.01 to 65 mm. At the low pressures the absorption bands are resolved into a series of narrow bands, about 0.15 Å. apart, which fuse very readily on passing a certain limited pressure. They are distributed in eight groups, of which only five are visible at 0.01 mm. More than 350 bands have been counted.

The application of Bohr's theory to the study of these band spectra permits of their classification into four series, each band being given by an equation of the form $1/\lambda = A - B + na - (pb + p^2b') + \alpha(m^2 - q^2)$, in which A , B , a , b , b' , and α are constants and n , p , m , and q whole numbers. The intensities of the heads of the bands are greatest for $p=0$ and diminish rapidly as p increases, which is in agreement with the theory. The coefficients deduced from the results of the ultra-violet absorption are available for the calculation of the infra-red absorption spectra of benzene. The terms A and B are due to electronic movements and do not enter into the frequency of the infra-red absorption bands, for which a simple formula is deduced, $1/\lambda = an - bp$; the agreement between the calculated and experimental values is very satisfactory. The moment of inertia of the benzene molecule ($I = 1.45 \times 10^{-39}$) is

derived from the value of the constant α and compared with that calculable from the crystal models of benzene derivatives deduced by Bragg.

The ultra-violet spectra of liquid benzene and its solutions correspond closely with the spectra of the vapour, except for a slight shift towards longer wave-lengths. On dilution, the specific effect of the solvent becomes more marked, the bands widen and move towards the red; this effect, which is analogous to the Stark effect, may be due to the action of an intense electric field of the solvent molecules. The displacement does not bear any relation to the dielectric constant of the solvent. The fluorescent spectrum of liquid benzene consists of six bands, which are excited by monochromatic light of wave-lengths lying within the benzene absorption bands *D*, *E*, *F*, *G*, and *H*; lines which correspond with the *A* and *B* absorption bands do not excite any fluorescence. The molecules absorbing the monochromatic light do not return to their original state on the emission of the fluorescent light, and the fluorescent bands lie towards the red with respect to the absorption bands. When the vapour of benzene is subjected to a high frequency discharge, a new absorption band appears ($\lambda = 2750.5 \text{ \AA.}$).

W. E. G.

The Ultra-violet Absorption Spectra of Dihydric Phenols.

F. W. KLINGSTEDT (*Compt. rend.*, 1922, 175, 365—367).—The quantitative study of the absorption spectra of dihydric phenols in hexane solution has given new results. Diagrams of the normal spectra of phenol, resorcinol, catechol, and quinol are given. These show that *m*- and *o*-derivatives yield results similar to those obtained with phenol, whilst the *p*-compound possesses a series of eight distinct bands in place of three exhibited by the other substances. This is a general result, as shown by the spectra of other benzene derivatives. The spectra are considerably modified by the solvents used, especially by alcohol; the authors consider that, as alcohol has been used for much work on absorption spectra, the observations should be repeated with hexane solutions.

H. J. E.

Phosphorescent Zinc Sulphide containing Copper.

B. GUDDEN and R. POHL (*Ann. Physik*, 1922, [iv], 68, 154—156).—Wilde has recently investigated the variation of the dielectric constant of phosphorescent zinc sulphide containing copper due to illumination, a phenomenon previously studied by the authors and by Molthan. Briefly, the results indicate that the increase in the value of the dielectric constant due to illumination is a maximum at about 140° . The effect is small at high temperatures. Temporary fluctuations of the effect at all temperatures, and for all exciting wave-lengths, is correlated with the luminosity of the phosphorescent substance. Regarding the phenomenon as originating in certain centres of excitation surrounding the relatively few copper atoms present, it follows that the effect must decrease with increasing temperature, and increase to a limiting value with decrease of temperature. The occurrence of the experimentally determined maximum at 140° is explained by assuming that the mobility of the electrons

within the excited centres diminishes as the temperature is reduced, an assumption which is in accord with other experimental results. The failure to detect the effect in other varieties of these phosphorescent substances is regarded as a matter of accuracy of measurement.
J. S. G. T.

Fluorescence, Photochemical Action, and Einstein's Law. FRITZ WEIGERT (*Z. Physik*, 1922, 10, 349—351).—A theoretical paper in which the author criticises Perrin's theory of fluorescence (cf. A., 1918, ii, 418; 1919, ii, 177). The part played by oxygen in the photochemical change of most fluorescent substances demonstrates the independence of the fluorescence and the sensitivity to light. For example, an oxygen-free solution of fluorescein is stable to light. The photochemical reaction velocity decreases with increasing partial pressure of oxygen, whereas the brightness of the fluorescence is unaffected. The experiments of Wood and Pringsheim are to be explained along these lines: Einstein's law cannot be applied to these photochemical processes, since under the most favourable conditions 200 energy quanta are necessary for the absorption of one molecule of oxygen by the fluorescent substance.
W. E. G.

Ionisation of Potassium Vapour by Light. R. C. WILLIAMSON (*Proc. Nat. Acad. Sci.*, 1922, 8, 255—260).—By employing a jet of hot potassium vapour, directed into a cool vacuum chamber and traversed by a pencil of light derived from a mercury arc in quartz, evidence has been obtained of the ionisation of potassium vapour by light. The limiting long wave-length (λ 2856 Å.) of the exciting radiation can be calculated from the ionising potential V by the quantum relation $Ve = h\nu$. Below this limit, the ionising power of the radiation increases continuously as the wave-length decreases.
J. S. G. T.

Rise of γ -Ray Activity of Radium Emanation. F. P. SLATER (*Phil. Mag.*, 1922, [vi], 44, 300—304; cf. this vol., ii, 13).—Curves are given showing the rise of γ -ray activity from pure radium emanation when enclosed by lead and aluminium of varying thicknesses. The fraction of the ionisation due to radium-*B* when in radioactive equilibrium with radium-*C* has been deduced by trial for thicknesses of lead up to 20 mm. The data have been utilised in deducing the absorption coefficients of the heterogeneous γ -radiation from radium-*B*.
W. E. G.

The Distribution of Radioactive Substances in Solutions. HILARY LACHS and MATHILDE WERTENSTEIN (*Physikal. Z.*, 1922, 23, 318—322).—In alkaline or neutral solutions of polonium, radium-*D*, radium-*E*, and radium, allowed to remain for a period of hours, the radioactivity exhibited at various depths increases as the depth increases. In acid solutions, the radioactivity is very approximately uniformly distributed throughout the solution. No conclusive results were obtained with solutions of radium-*A*, -*B*, and -*C*. The effects are attributed to the presence of minute particles, derived from the air or the vessel, in the solutions, the

radioactive material being adsorbed by these particles. The results are applied to explain the low values obtained by Paneth for the diffusion coefficient of polonium in neutral and alkaline solution, and of thorium-*B* in alkaline solution. J. S. G. T.

The Disintegration of Elements by α -Particles. (SIR) E. RUTHERFORD and J. CHADWICK (*Phil. Mag.*, 1922, [vi], 44, 417—432).—The nature of the particles ejected from various elements by bombardment with swiftly moving α -particles (cf. A., 1921, ii, 293), has been investigated by determining the deflexion of the particles in a magnetic field. The results show that the particles ejected from aluminium, phosphorus, and fluorine consist of positively charged hydrogen nuclei. The same is also probably true of boron and sodium. The ranges of the particles ejected, in the forward and backward directions, from nitrogen, aluminium, boron, fluorine, sodium, and phosphorus have been more accurately determined. In all cases, hydrogen particles were ejected in all directions, and the maximum range in the backward direction was less than in the forward direction. The maximum ranges, in cms. of air, of the ejected particles, in the forward and backward directions, respectively, were found to be: boron, 58 and 38; nitrogen, 40 and 18; fluorine, 65 and 48; sodium, 58 and 36; aluminium, 90 and 67; and phosphorus, 65 and 49. Only in the case of nitrogen was the maximum range of the ejected particles in the backward direction much less than that of free hydrogen particles. No particles of range greater than 30 cm. of air, in the forward direction, were ejected from either lithium or glucinum, and there was no evidence of the ejection of long range particles from magnesium, silicon, or chlorine. Of the series of elements from hydrogen to potassium hitherto examined, the active elements (those from which hydrogen nuclei are ejected) are odd-numbered in the order of atomic number in the regular sequence 5, 7, 9, 11, 13, 15. The atomic masses of the active elements are expressed by $4n+a$, where n is a whole number, and $a=3$ for all except nitrogen, for which $a=2$. With the exception of boron, which has two isotopes, the active elements are pure elements. The results indicate that neither lithium nor chlorine has any lightly bound satellites in its nuclear structure. The hydrogen satellite of nitrogen is relatively close to the main nucleus. The nuclei of even light elements are very complex systems, and it is difficult to advance any simple rule to account for the wide difference in structure of successive elements.

J. S. G. T.

Temperature Coefficient of Electromotive Force of Galvanic Cells and the Entropy of Reactions. ROSCOE H. GERKE (*J. Amer. Chem. Soc.*, 1922, 44, 1684—1704).—The *E.M.F.* of a number of electrolytic cells has been determined at a number of temperatures and the following values of the *E.M.F.* of the cell at 25° and the temperature coefficient are recorded: $\text{Ag}|\text{AgCl}|\text{HCl}(N)|\text{AgCl}|\text{Cl}_2$ (1 atm.), $E=1.1362$ volts, $\Delta E/\Delta T=-0.000595$ volts/degree; $\text{Hg}|\text{HgCl}|\text{HCl}(N)|\text{Cl}_2$ (1 atm.), $E=1.0904$ volts $\Delta E/\Delta T=-0.000945$; $\text{Ag}|\text{AgCl}|\text{KCl}(N)|\text{HgCl}|\text{Hg}$, $E=0.0455$,

$\Delta E/\Delta T = 0.000338$; $\text{Ag}|\text{AgCl}|\text{HCl}(N)|\text{HgCl}|\text{Hg}$, $E = 0.0455$, $\Delta E/\Delta T = 0.000337$; $\text{Pb}(\text{cryst.})|\text{Pb}(\text{OAc})_2(2N)|\text{Pb}(\text{Hg})$, $E = 0.0057$, $\Delta E/\Delta T = 0.000021$; $\text{Pb}(\text{stick})|\text{Pb}(\text{OAc})_2|\text{Pb}(\text{Hg})$, $E = 0.0057$, $\Delta E/\Delta T = 0.000015$; $\text{Pb}(\text{Hg})|\text{PbCl}_2|\text{HCl}(N) + \text{PbCl}_2|\text{AgCl}|\text{Ag}$, $E = 0.4843$, $\Delta E/\Delta T = 0.000202$; $\text{Pb}(\text{Hg})|\text{PbCl}_2|\text{HCl}(N) + \text{PbCl}_2|\text{HgCl}|\text{Hg}$, $E = 0.5299$, $\Delta E/\Delta T = 0.000129$; $\text{Ti}(\text{stick})|\text{Ti}(\text{OH})|\text{Ti}(\text{Hg})$, $E = 0.0027$, $\Delta E/\Delta T = 0.000020$; $\text{Ti}(\text{Hg})|\text{TiCl}|\text{NaCl}(N) + \text{HCl}(0.001N)|\text{TiCl} + \text{AgCl}|\text{Ag}$, $E = 0.7766$, $\Delta E/\Delta T = -0.0000794$; $\text{Pb}(\text{Hg})|\text{PbI}_2|\text{Pb}(\text{ClO}_4)_2(0.218M) + \text{HClO}_4|\text{PbI}_2|\text{I}_2$, $E = 0.8936$, $\Delta E/\Delta T = -0.000042$; $\text{Pb}(\text{Hg})|\text{PbI}_2|\text{KI}(N)|\text{PbI}_2 + \text{AgI}|\text{Ag}$, $E = 0.2078$, $\Delta E/\Delta T = -0.000189$; and $\text{Pb}(\text{Hg})|\text{PbI}_2|\text{KI}(0.1N)|\text{AgI}|\text{Ag}$, $E = 0.2135$, $\Delta E/\Delta T = -0.000173$. The free energy, ΔF_{298} , the entropy change, ΔS_{298} , and the change of heat content, ΔH_{298} , have been calculated from the *E.M.F.* measurements and the temperature coefficients of a number of reactions. The following are recorded: $\text{Ag} + \text{HgCl} = \text{AgCl} + \text{Hg}$, $\Delta F = -1050$ Cal., $\Delta S = 7.80$ Cal./degree, $\Delta H = 1275$ Cal.; $\text{Pb}(\text{cryst.}) = \text{Pb}(\text{Hg})$, $\Delta F = -263$, $\Delta S = 0.97$, $\Delta H = 26$; $\text{Pb}(\text{stick}) = \text{Pb}(\text{Hg})$, $\Delta F = -263$, $\Delta S = 0.69$, $\Delta H = -57$; $\text{Pb}(\text{solid}) = \text{Pb}(\text{Hg})$, $\Delta F = -263$, $\Delta S = 0.74$; $\Delta H = -42.4$; $\text{Pb}(\text{solid}) + 2\text{AgCl} = \text{PbCl}_2 + 2\text{Ag}$, $\Delta F = -22612$, $\Delta S = -8.58$, $\Delta H = -25170$; $\text{Pb}(\text{solid}) + 2\text{HgCl} = \text{PbCl}_2 + 2\text{Hg}$, $\Delta F = -24717$, $\Delta S = 6.70$, $\Delta H = -22720$; $\text{Ti}(\text{cryst.}) + \text{AgCl} = \text{TiCl} + \text{Ag}$, $\Delta F = -17975$, $\Delta S = -1.08$, $\Delta H = -18296$; $\text{Pb}(\text{solid}) + \text{I}_2 = \text{PbI}_2$, $\Delta F = -41501$, $\Delta S = -1.20$, $\Delta H = -41859$; $\text{Pb}(\text{solid}) + 2\text{AgI} = \text{PbI}_2 + 2\text{Ag}$, $\Delta F = -9852$, $\Delta S = -7.98$, $\Delta H = -12231$; $\text{Hg} + \frac{1}{2}\text{Cl}_2 = \text{HgCl}$, $\Delta S = -21.80$; and $\text{Ag} + \frac{1}{2}\text{Cl}_2 = \text{AgCl}$, $\Delta S = -13.73$. The third law of thermodynamics has been submitted to a most rigorous test and discrepancies found earlier by Lewis and Latimer (this vol., ii, 471) have been reduced tenfold. Assuming the validity of the third law, the entropies of chlorine and iodine have been calculated and the following values recorded: $\frac{1}{2}\text{Cl}_2$, $S_{298} = 27.1$; $\frac{1}{2}\text{I}_2$, $S_{298} = 13.5$.
J. F. S.

Electric Kataphoresis of Metallic Protein Compounds obtained by Treatment with Powdered Metals. A. BENE-DICENTI and S. REBELLO-ALVES (*Arch. intern. pharmacodynamie*, 1922, 26, 297—316).—In shaking a solution of egg-albumin or of blood-serum with powdered metals the metals become attached to the proteins and the properties of the latter change. They become coloured, their reactions are altered, they are not coagulated by heat, and resist putrefaction. The "fixation" of the metals is independent of the reaction of the protein solution and of its gas and salt content. The metals in this combination lose their ionic properties and are not dialysable. Metals vary in their power to combine with proteins: cobalt, copper, and iron combine to a greater extent than do aluminium and lead. Powdered oxides of the metals do not combine with proteins. By passage of an electric current through an alkaline solution of the metallic protein compound, the protein particles become negatively charged and move towards the positive electrode, carrying with them the fixed metal. In an acid solution the protein particles become negatively charged, move to the cathode and a part of the metal is deposited

on the electrode. At the chemically neutral (? isoelectric) point a protein coagulum forms, which gradually changes into a membrane upon which the fixed metal deposits. Boiling or heating a metallic protein solution to 50–60° changes profoundly its electrophoretic properties. It is suggested that the thermolability of sera, the fixation of metals by certain organs and their elimination, etc. depend on the formation of metastable protein compounds and on changes in the chemical equilibria of these compounds.

CHEMICAL ABSTRACTS.

Kataphoresis of Water in Organic Liquids. A. GYEMANT (*Z. physikal. Chem.*, 1922, **104**, 74–88).—The kataphoresis of drops of water and aqueous solutions of acids, bases, salts, and dyes in aniline, guaiacol, and benzonitrile has been measured by a microscopic process. In the case of guaiacol (less in aniline and still less in benzonitrile), it is found that water and aqueous solutions of inorganic electrolytes do not migrate at all or at best only very slowly, whilst solutions of salts which furnish organic ions possess a considerable kataphoric velocity. The sense of the direction is exactly that which would be expected from the kataphoresis of the same organic liquids in the corresponding aqueous solutions. In the case of organic kations, the water drops are negatively charged towards the organic medium and with organic anions positively charged. Some cases have been found which point to the assumption that only a portion of the total potential (ϵ), the so-called ζ -potential, is determinative of the kataphoresis. Thus, for example, it is found that drops of aqueous solutions of hydrochloric acid and aluminium chloride in benzonitrile are positively charged, although when benzonitrile emulsifies in the solutions it is positively charged; this phenomenon finds its explanation in the ζ -potential. The smaller velocity of migration of drops of aqueous solutions of electrolytes cannot be directly explained by the assumption that in these cases the ζ -potential in the organic liquid is much smaller. Probably it is to be explained by the inorganic ions being much less solvated in organic liquids than the organic ions. The Helmholtz formula for the velocity of kataphoresis cannot be directly applied to a water emulsion in organic liquids.

J. F. S.

Thermo-compression and Thermo-addition. MAURICE DELEUER (*Bull. Assoc. Chim. Sucr.*, 1922, **39**, 477–480).—A theoretical paper in which the conditions under which the equations, moist vapour+moist vapour=superheated vapour, and superheated vapour+superheated vapour=moist vapour, are true are discussed.

J. F. S.

Determination of Specific Heats at Low Temperatures. FRANZ SIMON (*Ann. Physik*, 1922, [iv], **68**, 241–280).—The accuracy of determination of specific heats of substances at low temperatures devised by Nernst and Schweser (*A.*, 1914, ii, 336), has been considerably increased by using a larger calorimeter designed so that the amount of liquid hydrogen employed is not

unnecessarily increased. Determinations have been made and the values tabulated of the respective specific heats of quartz glass, christobalite, "Lindemann" glass, mercury, ammonium chloride, dextrose, cuprous iodide, glycerol, and keton-resin ("Keton-harz," D.R.-P. 309224), at temperatures ranging from 15° Abs. to the ordinary temperature. The T^3 -law, expressing the value of the molecular heat divided by the number of atoms in the molecule, by the relation $c_p = 6.94 \times 10^{-6} T^3$, where T is the absolute temperature, is very closely obeyed by ammonium chloride, for temperatures between 20° and 32° Abs. Anomalous values of the specific heat of ammonium salts are due to a transformation occurring within a certain range of temperature—from about 140° Abs. to 243° Abs. in the case of ammonium chloride. The sudden drop in the value of the atomic heat of glycerol when the temperature is reduced just below 180° Abs. is correlated with the softening of the substance at this temperature. In the case of cuprous iodide, values of the free energy calculated by means of Nernst's theorem are in agreement with experimentally determined values. The values of the energy liberated by the combustion of dextrose at various temperatures are tabulated. The value of A , the free energy of the reaction in Nernst's relation, is 704,500 cal. at 290° and 706,700 at 310° T . It is shown that the value of the specific heat of complex bodies at different temperatures can be represented by the Nernst interpolation formula consisting of the sum of the Debye and Einstein functions. The number of the latter functions is small compared with the number of the former. The general characteristics of the variation with temperature of the atomic heats of complex bodies is discussed from the point of view of atomic vibrations, and it is shown why, in many cases, Berthelot's principle affords a close approximation to the energy relations involved.

J. S. G. T.

A New Physico-chemical Law. HAWKSWORTH COLLINS (*Chem. News*, 1922, **125**, 81—83, 97—98).—From tables of the atomic weight and specific gravity of the elements, and the specific gravity and heat of formation of a number of molecules, the relative volumes of the elements in combination and the heat of formation of the elements when combining to form molecules are deduced. The values calculated never differ more than 1% from the observed values. It is also shown that the heat of formation of each element is proportional to the product of the atomic weight and the change of volume. By change of volume is meant the difference between the volume of the element in the solid state at 15° and the volume of the element in combination with others at 15°.

J. F. S.

Heat of Crystallisation of Quartz. RAMES CHANDRA RAY (*Proc. Roy. Soc.*, 1922, [A], **101**, 509—516).—The heat of crystallisation of quartz has been determined by measuring the respective heats of solution of silver sand and silica glass in hydrofluoric acid. The following values were obtained for the specific heat of aqueous hydrofluoric acid solutions containing the respective percentages by weight of hydrofluoric acid: 10%, 0.908; 20%,

0.838; 30%, 0.756; 34.6%, 0.724. The mean values of the heats of solution in hydrofluoric acid per gram-molecule of silicon dioxide were found to be: silica glass, 37.24 Cal.; silver sand, 30.29 Cal. The heat of crystallisation of quartz, represented by the difference, is therefore 6.95 Cal. per gram-molecule. This value agrees with that calculated from Tschernobaeff and Wologdine's results, but is much higher than those obtained by Mulert (2.21) (A., 1912, ii, 626) and Wietzel (2.32), (A., 1921, ii, 504). The discrepancy is attributed to the effect of prolonged grinding in reducing part of the substance to the vitreous condition. The calculated value of the heat of crystallisation at 900° is 6.50 Cal. per gram-molecule. At 1300—1400°, the value is not much different from that at ordinary temperature. J. S. G. T.

Separation of Liquid Mixtures by Centrifuging. ROBERT S. MULLIKEN (*J. Amer. Chem. Soc.*, 1922, **44**, 1729—1730).—In an earlier paper (this vol., ii, 492) the author showed thermodynamically that the same separation of isotopes should be obtainable by centrifuging a liquid as a gaseous mixture, at a given speed. The same result was stated to be true for ideal solutions in general. The latter statement is now shown to be incorrect. The result depends not only on the ideality of the solutions, but also on equality of the vapour pressures and of the atomic or molecular volumes of the constituents. These conditions are fulfilled in few cases except that of a mixture of isotopes. J. F. S.

Viscosity and Molecular Dimensions of Sulphur Dioxide. C. J. SMITH (*Phil. Mag.*, 1922, [vi], **44**, 508—511).—By Rankine's method, the author has determined the viscosity of sulphur dioxide to be 1.253×10^{-4} C.G.S. units at 18.0°, and 1.630×10^{-4} at 100.0°. The value of Sutherland's constant calculated therefrom is 416, and the corresponding value of the viscosity at 0°, extrapolated by Sutherland's formula, is 1.168×10^{-4} C.G.S. units. The value of the mean collision area, calculated from the results by Chapman's formula, is 0.94×10^{-15} sq. cm. with a possible error of 2 or 3%. J. S. G. T.

Viscosity and Molecular Dimensions of Gaseous Carbon Oxysulphide. C. J. SMITH (*Phil. Mag.*, 1922, [vi], **44**, 289—292).—The viscosities of carbon oxysulphide have been determined at 15° and 100° by the method described by Rankine and Smith (cf. A., 1921, ii, 694, 696; this vol., ii, 549). The values obtained were $\eta_{15} = 1.200 \times 10^{-4}$ C.G.S. units, and $\eta_{100} = 1.554 \times 10^{-4}$ C.G.S. units; and by extrapolation from Sutherland's formula $\eta_0 = 1.135 \times 10^{-4}$ C.G.S. units. The mean area which the molecule presents in mutual collision is $\bar{A} = 1.06 \times 10^{-15}$ cm.². W. E. G.

Measurement of the Pressure of "Surface Fluids": Oleic Acid. A. MARCELIN (*Compt. rend.*, 1922, **175**, 346—348).—In order to measure the pressure exerted by thin layers of liquid resting on the surface of a second liquid, the author has

constructed a "surface manometer" which is sensitive and capable of rapid measurement. In the case studied, the addition of successive small quantities of oleic acid to the surface of water contained in a vessel results in a corresponding increase of pressure until a limiting value is reached, at which point a portion of the acid is floating as a drop of oil on the surface. This "surface saturation pressure" is comparable with the maximum tension of saturated vapour; it characterises the equilibrium between the "surface fluid"—a two-dimensional phase—and the same substance in a three-dimensional phase. At the equilibrium point, pressure on the surface liquid results in condensation into drops, and conversely. A further series of experiments was made in which the quantity of oleic acid was insufficient to cover the entire surface of the water. By surface compression, the pressure which is initially zero increases slowly at first, then more rapidly tending towards the saturation limit, and on allowing the surface liquid to expand, its pressure decreases in the inverse sense. If curves are drawn representing these pressures as functions of the surfaces, the second curve falls below the first; the two meet asymptotically, and the whole is of the same type as the curves of magnetic hysteresis. It is also shown that there is no limit to the extension of the surface liquid; this apparently corresponds with the limitless expansion of a gas.

H. J. E.

The Properties and Molecular Structure of Thin Films.

III. Expanded Films. N. K. ADAM (*Proc. Roy. Soc.*, 1922, [A], 101, 516—531).—In continuation of previous work, the author has investigated the effect of temperature on thin films of fatty acids spread on surfaces of distilled water, a phosphate solution, a dilute solution of hydrochloric acid, and of sodium hydroxide. At a certain temperature, dependent both on the nature of the substance in the film and on the composition of the solution, the films expand along the surface. The phenomena observed are discussed at length and the attempt is made to correlate the behaviour of such expanded films with that of a gas, the barriers on the surface over which the film is expanded representing the walls of the containing vessel in the case of the gas. The "gas" equation for the expanded films is of the Van der Waals's type, the b term, however, representing twice and not four times the area of the molecules, and the v term being replaced by the area of the film. It is shown that expanded films are one molecule thick, and that the molecules are probably arranged vertically, that is, they do not lie flat on the surface. It is shown that the steady increase in the expansion temperature with length of chain constituting the compound probably arises owing to the lateral attraction of chains lying side by side. The bearing of this result on the crystal structure of fatty acids and their esters is discussed. There appears to be no *a priori* reason for the view that lateral adhesion between the molecules prevents the films dissolving. Conclusions previously reached (this vol., i, 424) regarding the structure of *cis*- and *trans*-acids are now considered to be incorrect.

J. S. G. T.

Electric Endosmose. ALVIN STRICKLER and J. HOWARD MATHEWS (*J. Amer. Chem. Soc.*, 1922, 44, 1647—1662).—The rate and direction of the endosmotic flow of acetone, amyl alcohol, nitrobenzene, pyridine, benzaldehyde, *n*-butyl alcohol, furfuraldehyde, isopropyl alcohol, and *o*-nitrotoluene have been determined, using a modified form of Briggs's apparatus (A., 1917, ii, 236; 1918, ii, 214). The influence of the addition of acids, bases, and salts on the endosmosis has also been investigated. An adsorption orientation hypothesis has been suggested to explain the establishment of the original charge at the surface of contact of the membrane and solvent. This states that the molecules of the solvent, on adsorption, are orientated on the membrane surface producing a charge by their polarity. If dissociated molecules are present, they affect the membrane charge by specific adsorption of the ions produced. This does not eliminate adsorption, but suggests a more comprehensive view of the mechanism for the establishment of the original charge. The experiments show that the permeability of the membrane is an important factor in determining endosmotic flow, and that it is possible to use and duplicate membranes of filter-paper when using organic solvents. The rate of endosmosis is not quite a linear function of the voltage applied when this is varied over a wide range. Briggs's results (*loc. cit.*) as to temperature effects with water have been confirmed with non-aqueous solvents. The valency rule does not hold with the non-aqueous solvents investigated, and the order of adsorption of ions and their relative adsorption varies with the solvent employed. The concentration curve exhibits a maximum with non-aqueous solvents and the flow at low concentrations is in the same direction as that of the pure solvent. The addition of water to non-aqueous solutions changes the endosmosis in the direction it would manifest in pure water, the first additions showing the greatest effect. The direction of endosmosis is not determined by the dielectric constant ratio, although this is probably a measure of the magnitude of flow. The Hofmeister series is not consistent when considering endosmotic effects in non-aqueous solvents.

J. F. S.

Dependence of the Adsorption by Charcoal on the Quantity of Charcoal and the Nature of the Adsorption Isothermal. A. FODOR and B. SCHÖNFELD (*Kolloid Z.*, 1922, 31, 75—80).—The adsorption of acetic, lactic, and tartaric acids by various amounts of animal charcoal from solutions of various concentration has been determined. It is shown that with sufficiently dilute solutions of these substances and with a sufficient quantity of adsorbent the index $1/n$ of the adsorption equation approaches and reaches unity in the same way as was previously shown to be the case for amino-acids, polypeptides, and some carbohydrates (A., 1921, ii, 21). In sufficiently dilute solutions the value of n is very sensitive to changes in the amount of charcoal employed in the sense stated above. In less dilute solutions the value of n is practically independent of the amount of charcoal used. The

$(c_0 - c) - w$ curves, where w is the amount of adsorbent and $(c_0 - c)$ the quantity adsorbed, are parabolic and convex to the w axis and are steeper the greater the initial concentration, c_0 , and the smaller w is. The $c - w$ curves are also parabolic but concave to the w axis, the concavity being greater the larger c_0 and the smaller w . From this behaviour it is concluded that the expression $dc/dw = nc/w$ is correct, and interprets the adsorption equation $x = k'(c_0 - x)^{1/n}$ or $x^n = k(c_0 - x)$. In keeping with this interpretation of the index n , the rôle of this quantity in catalytic and fermentation processes is discussed.

J. F. S.

The Adsorption of Ions. JNANENDRA NATH MUKHERJEE (*Phil. Mag.*, 1922, [vi], 44, 321—346).—A critical résumé of the literature on the adsorption of ions in which the author applies his theory of the origin of the charge of a colloidal particle and its neutralisation by electrolytes (cf. this vol., ii, 198). The characteristic form of the adsorption isotherm, obtained when the ions are multivalent or complex, is explained on the basis of the greater chemical activity of the complex anions over that of the simple kations. It is assumed that the electrostatic attraction of the surface on the kation is less than the chemical affinity acting on the anion. With increasing concentration of electrolytes, the charge on the surface will decrease until a balance is reached between the chemical adsorption of the anion and the electrical adsorption of the kation. The second maximum is due to a balance between the electrical adsorption of the kation and anion. A theory of the action of acids and alkalis on the surface is developed for (1) an inert surface, and (2) a chemically active surface: preferential adsorption of one ion. The maximum charge due to acids gives a measure of the hydration of the surface. The conception of the chemical adsorption of the anion is applied to ionic equilibria on the surface of colloids and to the interchange of kations in the soil.

W. E. G.

The Connexion between Hydrolysis and Adsorption. W. MOELLER (*Z. Leder- u. Gerb.-Chem.*, 1922, 1, 160—165, 183—188; cf. *ibid.*, 1921, 1, 47; *Collegium*, 1920, 109, 152, 209, 267).—An investigation of the swelling and hydrolysis of gelatin in hydrochloric, acetic, lactic, and butyric acids yielded results similar to those with hide powder. Hydrolysis products with organic acids are adsorbed by the intact gelatin. With mineral acids hydrolysis, and with organic acids adsorption predominates.

CHEMICAL ABSTRACTS.

Free Energy of Gaseous Molecules with any [possible] Partition of Velocities. MAX PLANCK (*Sitzungsber. Preuss. Akad. Wiss. Berlin*, 1922, 63—70).—A theoretical paper in which the question is raised and discussed as to whether or no the free energy of a mixture of similar gaseous molecules is made up additively of the free energies of the individual groups of velocities, when the partition of velocities is anything possible and the internal energy is in accord with the temperature. It is shown that the

above question must be answered in the affirmative. It is also shown that in the diffusion of two molecular systems similar in all respects except that of velocity, the reduction of the free energy is independent of the difference in the velocities. Further, in connexion with the foregoing, it is shown how this case explains the so-called Gibbs paradox, despite the fact that the velocity is a continually changing quantity. J. F. S.

Diffusion and Intertraction. C. G. SCHONEBOOM (*Proc. Roy. Soc.*, 1922, [A], **101**, 531—539).—The phenomenon of “pseudo-podial” intertraction was originally observed by Wright, when blood-serum was run slowly on to an aqueous solution containing from 5 to 8% of common salt. The upper fluid is carried down into the lower, and the lower rises into the upper. The phenomenon may be observed with all inorganic and organic compounds soluble in water, provided the difference in specific gravity of the two solutions is not too great and the solutions do not react chemically. Surface tension alone is concerned in the production of the phenomenon, size of molecule and water of hydration exerting no influence. Distinction is drawn between the processes of “pseudopodial” intertraction and “simple” intertraction, a process intermediate between the former and the process of diffusion. J. S. G. T.

Preparation and Use of Collodion Osmometers. WILLIAM BROWN (*Ann. Bot.*, 1922, **36**, 433—439).—Details are given for the preparation and use of collodion thimbles for osmotic experiments. They are adapted for comparative rather than for absolute measurements. G. W. R.

Ionic Properties and Chemical Facts. IV. Lattice Energy and the Work of Ionisation of Inorganic Compounds. H. G. GRIMM (*Z. physikal. Chem.*, 1922, **102**, 113—140; cf. this vol., ii, 127, 483, 635).—A theoretical paper in which the author has collected and calculated the necessary data for comparing the regularities of the heat of formation of inorganic substances, and for showing the connexion between the heat of formation with the ionic properties, charge, radius, and structure. The lattice energy and work of ionisation have been calculated or approximated for about one hundred inorganic substances from the newer experimental data and where this is not available the differences of the lattice energy have been measured and the variation thus determined. J. F. S.

Ionic Properties and Chemical Facts. V. Connexion between Heat of Formation, Lattice Energy, and the Ionic Properties. H. G. GRIMM (*Z. physikal. Chem.*, 1922, **102**, 141—168; cf. preceding abstract).—The connexion between the heat of formation, lattice energy, ionic radius, ionic structure, and ionic charge has been investigated. It is shown that, in opposition to the heat of formation, the lattice energy of solid substances and the work of ionisation of gaseous substances, so far as these are known, are definitely related to the ionic properties, charge, structure, and radius. The lattice energy and work of ionisation

decrease with increasing radius both with anions and kations when the ions compared have the same number of exterior electrons. The characteristic gradation of the ionic radii is in general reproduced in the gradation of the lattice energy. The difference between the lattice energies of compounds of two outwardly similarly constructed kations or anions with a common anion or kation with the same exterior sheath, decreases with increasing radius of the common ion. Compounds which contain kations with eighteen exterior electrons have considerably greater lattice energies than compounds with the corresponding kation of the same period which have only eight electrons. The haloids of kations with eighteen or twenty exterior electrons show a much smaller dependence of the lattice energy on the halogen-ion radius than the haloids of kations with eight exterior electrons. In all cases investigated the series $U_{H^+} > U_{Li^+} > U_{Na^+}$ is true, where H^+ has 0, Li^+ has 2 and Na^+ has 8 exterior electrons. The empirical relationship $(U_{(SiX'_j)} - U_{(SiX''_j)})/4 > (U_{(AlX'_j)} - U_{(AlX''_j)})/3 > U_{(NaX')} - U_{(NaX'')}$ is shown to be true, where U represents the work of ionisation of the substance in the brackets and X' and X'' are halogens. A similar expression is also true for boron and lithium haloids. These relationships can be deduced when heteropolar linkings are assumed for all the substances concerned. The haloids of carbon and the halogen-substituted hydrocarbons do not follow a relationship corresponding with that followed by ions containing the helium sheath. This indicates, as also do preliminary and approximate calculations on the models of BH_3 and CH_4 , that the carbon compounds concerned do not possess a polar linking. Making use of Born's cycle, many of the known regularities of the heat of formation of solid substances and of dissolved substances as well as the exceptions to the regularities can be investigated. The gradation of the heat of formation by the variation of a single ion of the compound can be attributed to the variation of three thermochemical quantities which depend on the ionic properties in the simplest manner. The periodicity of the heat of formation of oxides and chlorides can, in the neighbourhood of the maxima and minima, be explained by similar analysis.

J. F. S.

Effect of Sucrose on the Activities of the Chloride- and Hydrogen-ions. J. W. CORRAN and W. C. McC. LEWIS (*J. Amer. Chem. Soc.*, 1922, 44, 1673—1684).—The effect of the addition of sucrose on the activity of the chloride-ion in 0.1*N*- and 0.5*N*-solutions of potassium chloride solutions has been investigated by means of the cell $Ag|AgCl|KCl(c_1)||KCl(c_2)+Sucrose|AgCl|Ag$. It has been found that the increase in activity could be accounted for by simply taking account of the decrease in the total water present in the solutions; that is, the potassium- and chloride-ions are soluble in the water of hydration of the sucrose. The effect of sucrose on the activity of the hydrogen-ion in 0.1*N*-hydrochloric acid has also been investigated, making use of the assumption of MacInnes (A., 1919, ii, 385) of the independent activity of the chloride-ion in solutions of chlorides of the same concentration.

It is shown by means of a comparison between the hydrogen-ion activities obtained in the present experiments, and the values obtained with a hydrogen electrode, that the hydrogen electrode measures the hydrogen-ion activity and not the geometric mean of the activities of the hydrogen-ion and the accompanying anion.

J. F. S.

Ionisation Constant of Hypochlorous Acid: Evidence for Amphoteric Ionisation. WILLIAM A. NOYES and THOMAS A. WILSON (*J. Amer. Chem. Soc.*, 1922, **44**, 1630—1637).—The electrical conductivity of dilute solutions of hypochlorous acid and mixtures of nitric acid and hypochlorous acid at 25° was measured, using a flowing electrolyte, with the object of testing the view of Stieglitz (A., 1902, ii, 66) that hypochlorous acid is an amphoteric electrolyte. The results show that the value k_a is 6.70×10^{-10} , the value of k_b for hypochlorous acid is too small to be determined by the change in the conductivity of 0.0001N-nitric acid when hypochlorous acid is added to it. When a current of air is passed through a solution of hypochlorous acid, chlorine monoxide and not the acid escapes. This indicates the presence of positive chlorine-ions in such a solution and confirms the amphoteric ionisation of the acid; $\text{HClO} \rightleftharpoons \text{H}^+ + \text{ClO}'$; $\text{HClO} \rightleftharpoons \text{Cl}' + \text{OH}'$. If the Lewis-Langmuir hypothesis is accepted, it requires an interpretation consistent with these facts.

J. F. S.

State of Aggregation and Liquid Crystals. O. LEHMANN (*Z. physikal. Chem.*, 1922, **102**, 91—101).—In a recently published work ("Die Aggregatzustände") Tammann has questioned the existence of liquid crystals because they do not fall into the scheme of states of aggregation. The author, therefore, has repeated the evidence on which the existence of the liquid crystal state of aggregation is based.

J. F. S.

Glass Suspensions produced by Rubbing Glass Walls with Glass Rods. ROBERT FRICKE (*Kolloid Z.*, 1922, **31**, 80—81).—Moderately concentrated glass suspensions may be prepared by rubbing the sides of a test-tube with a glass rod. These suspensions are alkaline to phenolphthalein and have a normality of the order 1/1000 which increases on keeping the suspension. They generally settle completely in twenty-four hours, but the settling is more rapid if electrolytes are added. The particles are negatively charged and wander to the anode on the application of an electric current. Examined in the ultramicroscope, the particles are seen to have diameters varying between 20 μ and 0.5 μ .

J. F. S.

A New Apparatus: The Nephelometer. I. NEWTON KUGELMASS (*Compt. rend.*, 1922, **175**, 343—345).—Changes in the degree of aggregation of a colloidal solution may be observed quantitatively by measuring variations in the degree of transparency. This is effected by allowing light of weak intensity from a constant source to fall on a sensitive thermopile after traversing a vessel containing the solution. The apparatus is

standardised by taking a series of readings after filling the vessel with distilled water. The author has used the apparatus in order to observe the coagulation of fibrinogen and the results obtained are expressed numerically. H. J. E.

Nephelometric Effect of Colloidal Systems of Different Size of Particles. H. BECHHOLD and F. HEBLER (*Kolloid Z.*, 1922, 31, 70—74).—The connexion between the turbidity and size of the suspended particles has been examined nephelometrically in the case of suspensions of barium sulphate in various mixtures of ethyl alcohol and glycerol. The degree of dispersion of the suspensions varied from $2.5\ \mu$ to $4\ \mu\mu$. The results show that for suspensions containing the same amount of barium sulphate but of varying degrees of dispersion, the turbidity increases from $2.5\ \mu$ downwards. The maximum turbidity is reached with particles of about $800\ \mu\mu$ for white light, that is, in the region of the extreme red. Further reduction of the size of the particles caused the turbidity to decrease strongly. It is in this region that Rayleigh's law holds. By the term turbidity is to be understood the effect, observed with the nephelometer, which is brought about by the diffraction of the light by the disperse phase. A method is described whereby the diameter of submicrons and amicrons may be determined if a comparison turbidity of a substance with particles of known diameter is available. J. F. S.

Turbidity Standard. H. BECHHOLD and F. HEBLER (*Kolloid Z.*, 1922, 31, 132—137).—The authors have prepared an easily reproducible and fairly stable standard turbidity and a sol suitable for use as a standard in work on turbidities and suspensions. They characterise as standard turbidity one which contains $1/1000$ mol. of barium sulphate; this is prepared by mixing $1/500$ mol. of hydroxylamine sulphate dissolved in glycerol with an equal volume of $1/500$ mol. barium chloride also dissolved in glycerol. The precipitate is chiefly made up of particles $2.5\ \mu$ in diameter. The standard sol is characterised as a turbidity which contains $1/200$ mol. of barium sulphate; this is prepared by mixing barium chloride and magnesium sulphate dissolved in a mixture of glycerol (6.5) and isobutyl alcohol (1.5). This is three times as turbid as the standard turbidity and can be made of the same turbidity by mixing with glycerol containing 15% of isobutyl alcohol. The standard sol consists of primary particles of not more than $90\ \mu\mu$ diameter. The standard sol is stable and constant for at least six months, whilst the standard turbidity is not trustworthy after eight days. J. F. S.

Theory of the Mechanical Synthesis of Colloids. F. SEKERA (*Kolloid Z.*, 1922, 31, 137—147).—The theory of the production of colloids in Plauson's colloid mill is briefly as follows. The mechanical grinding process bursts the active molecular compounds in the interior of the solid substance and produces particles of colloidal dimensions. From the surface of these particles the residual valencies tend to cause an aggregation of the small particles, so that secondary to the dispersion an aggregation process

is taking place. To retard or prevent the aggregation, the residual valencies are saturated by the addition of an electrolyte which produces salt complexes and charges the particles. J. F. S.

Dielectric Constants of Colloidal Solutions. J. ERRERA (*Kolloid Z.*, 1922, **31**, 59—64).—A method for measuring the dielectric constants of colloidal solutions, based on the Nernst condenser method is described and has been used to measure the dielectric constant of vanadium pentoxide sols. It is shown that a solution containing 14 per thousand has a dielectric constant of 400 in comparison with that of water of 81. This high value is not obtained with the freshly prepared sol, but appears only after long preservation. Thus a freshly prepared sol gave a value 74.7 which in seventeen days had risen to 94.8. The value given above was for a sol which had been kept for five years. The dielectric constant shows a characteristic dilution curve and a definite dependence on the temperature, current density, and the potential amplitude. The dielectric constant obtained is doubtless connected with the well-known double refraction of this sol.

J. F. S.

An Experimental Test of Smoluchowski's Theory of the Kinetics of the Process of Coagulation. JÑANENDRA NATH MUKHERJEE and B. CONSTANTINE PAPACONSTANTINOU (*Phil. Mag.*, 1922, [vi], **44**, 305—320).—According to the theory of Smoluchowski (cf. A., 1917, ii, 297), the successive stages in the coalescence of a colloidal solution are always the same and depend only on the time, and thus any property of the sol that varies continuously can be utilised to fix a definite stage in the coalescence. If his equations hold, all curves showing variations in the physical properties of the colloidal solution should be similar. Previous experimental work in which ultramicroscopical measurements have been made has yielded inconclusive data, possibly owing to difficulties inherent in the methods of measurement. The authors have tested the views of Smoluchowski by measurements of the changes in colour of gold sols on the addition of an electrolyte (cf. T., 1920, **117**, 1563). Since the absorption coefficient in the red region varies continuously with the coagulation, the attainment of a definite value of the absorption coefficient for a fixed wave length ($683\text{ }\mu\mu$) is taken as representing a definite stage in the coalescence. Experimental values of the absorption coefficients for the addition of several concentrations of potassium chloride, potassium nitrate, and barium chloride have been previously given (*loc. cit.*), and it is shown that the ratios of the times required to reach the same absorption coefficient are independent of the time or the stage of coalescence. The data thus supply the best evidence so far recorded in favour of the theory of Smoluchowski; T in his equations is a constant in the case of the gold sols within the limits of coagulation studied. The percentage of successful collisions between the colloidal particles increases rapidly with increase in concentration of electrolyte and varies markedly with changes in temperature.

W. E. G.

Colloidal Solutions of Carbon in Water. P. C. L. THORNE (*Kolloid Z.*, 1922, **31**, 119—132).—The author briefly summarises the literature on the formation of colloidal solutions of carbon and discusses the results obtained. An account is given of the preparation of colloidal solutions of carbon by the dispersion of charcoal anodes in the electrolysis of solutions of ammonia, sodium hydroxide, and dilute acids; similar experiments were carried out with graphite electrodes. A stable carbon sol may be prepared by the electrolysis of a solution of ammonia for six hours between two carbon electrodes with an anode current density of 0.6—1.2 amperes/cm.² and a voltage of 220. On filtering, the solution had a concentration 0.145*N* of ammonia and gave a residue of 0.034 gram per 100 c.c. This residue was insoluble in water, but redissolved in a dilute solution of ammonia; the excess of ammonia may be removed by boiling and the sol is then more stable than one which has been dialysed and still contains 0.039*N* ammonia. The ammonia-free sol is deep black in layers greater than 1 cm. thick and in thinner layers brown, and is made up of particles of about the same size as those of metallic sols. The influence of electrolytes on the sol has been investigated and the number of millimols. necessary to coagulate 1000 c.c. of the dialysed sol found to be: hydrochloric acid, 60; sodium hydroxide, 290; ammonium hydroxide, >9000; sodium chloride, 450; barium chloride, 0.5; and aluminium chloride, 0.3. These quantities are very similar to those required to coagulate a platinum sol. Similar experiments have been carried out with graphite sols and results of the same kind obtained. The most marked difference between the carbon sols and the graphite sols is that of colour, the former being brown and the latter black. An analysis of the dried sol shows that it is not a pure carbon sol which has been dealt with, but a mixture, the composition being C, 66.61%; H, 1.99%; O, 15.25%, insoluble and incombustible residue (probably silica), 16.15%.
J. F. S.

Electrical Properties of Silicic Acid Sols. OTTO LÖSENBECK (*Koll. Chem. Beihefte*, 1922, **16**, 27—46).—A number of experiments are described on the influence of silicic acid sols on the electrical conductivity of solutions of hydrochloric acid, and on the kataphoresis of the sol both pure and in the presence of hydrochloric acid of various concentrations. It is shown that the electrical conductivity of hydrochloric acid solutions of concentrations up to 0.01*N* is considerably reduced by the presence of silicic acid sol. This reduction does not take place instantaneously, but the amount of decrease increases with time according to a function of e , that is, the decrease is at first rapid and then continuously slower until it becomes asymptotic and a limiting value is approached. From the measurements of the migration velocity of the silicic acid particles under a known potential fall, the contact potential has been calculated by means of Smoluchowski's theory. It is shown that the particles of a pure silicic acid sol are negatively charged and that the charge is steadily reduced by the addition

of hydrochloric acid until the isoelectric point is reached. With further addition of acid the sign of the charge changes and the charge increases until the positive value is greater than the original negative value. The silicic acid particles are shown not to be uniform. The contact potential of the particles determined in different concentrations of the same preparation shows that this quantity is dependent on the concentration of the sol, in the sense that the isoelectric point of different sols lies at greater concentrations of hydrochloric acid the greater the concentration of the sol. The observed results are explained by the assumption that the silicic acid particles are saturated with hydrochloric acid, like a sponge, and that this is taken up slowly. The hydrogen-ions are expelled from these sponges with a definite tension, so that the contact potential between particle and solution may be approximately represented by the Nernst formula. J. F. S.

Colour of Photochlorides and Colloidal Silver. II. KARL SCHAUM and THEODOR MARX (*Kolloid Z.*, 1922, **31**, 64—70; cf. A., 1921, ii, 506).—In a previous paper (*loc. cit.*) the authors describe the presence of faintly illuminating red particles in silver sols of medium dispersion; these particles do not conform to Mie's theory. The authors now show that the red particles do not exist and the red colour observed was due to a contrast with the many green particles present. A method of determining the true colours of the colloidal particles in the presence of other particles is described. Previous attempts to reverse the order of colours obtained in photographic intensification by reduction failed, but it is now shown that a complete reversal may be obtained as follows. A blue silver sol is mixed with an equal volume of 10% gelatin solution and allowed to solidify, then a concentrated solution of ammonium hydroxide, potassium cyanide, or sodium thiosulphate is poured on to it and allowed to diffuse. After several days the following range of colours is observed, commencing at the top of the tube: colourless, yellow, orange, red, reddish-violet, bluish-violet, and blue. The action of electrolytes in definite amounts on silver sols is to produce definite colours. Thus in the case of magnesium sulphate and silver sols, 2 c.c. of the sol when treated with the following quantities of *N*/20-magnesium sulphate gives the colours indicated: 0 c.c., yellow; 0.10 c.c., orange; 0.20 c.c., red; 0.25 c.c., purple; 0.30 c.c., reddish-violet; 0.50 c.c., bluish-violet; and 1.50 c.c., blue. Similar changes were also obtained with ammonium chloride, sodium chloride, and potassium sulphate. J. F. S.

Behaviour of Carey Lea's Silver Sol towards Electrolytes and Hydrophilic Colloids. H. FREUNDLICH and E. LOENING (*Koll. Chem. Beihefte*, 1922, **16**, 1—26).—Carey Lea's silver sol behaves as an entirely hydrophobic sol, and since its particles are negatively charged the valency of the kations and the adsorbability are of most importance in its coagulation. The dependence of the coagulation value on the concentration of silver micellæ is bound up with the valency of the kations. The coagulum has a quite

different appearance when it is produced by univalent kations from that produced by bi- and ter-valent kations. Hydrophilic colloids such as saponin and gum arabic exert a protective action at all concentrations. This action increases strongly with the concentration of the hydrophilic colloid in the coagulation by tervalent kations, less strongly with bivalent kations, and weakly with univalent kations. This is explained by the fact that with the tervalent kations much smaller concentrations are necessary, so that the displacing action of the protective colloid is made more noticeable. Other hydrophilic colloids such as gelatin and caseinogen in smaller concentrations coagulate the silver sol, but above a definite concentration exert a protective action; this concentration increases with the concentration of the silver micellæ in the sol. The coagulation with small concentrations is, in the main, similar to the coagulation of acid gold sols by hydrophilic sols investigated by Gann (A., 1917, ii, 21). In neither case need complete coagulation occur, and an increase in the sensitiveness is brought about, that is, the metallic sol which contains small quantities of the hydrophilic colloid, is more sensitive toward electrolytes than the pure sol. Coagulation and increase in the sensitiveness by means of hydrophilic colloids are due to the fact that the sols of the hydrophilic colloids contain ions. In the case of the negative metallic sols, the colloid kations are operative and coagulate in the same manner as other kations. The silver sol is not discharged in solutions of colloids such as gelatin and caseinogen which are concentrated enough to exert a protective action, but here it is negatively charged. This is explained by the assumption that the amphoteric ions of gelatin lie with their positive ends toward the silver particles and the negative ends toward the interior of the solution. The above explanations are in keeping with the facts, that gelatin behaves in a protective but non-coagulating manner toward an alkaline gold sol, because even in alkaline solutions it contains too few colloidal kations; on the other hand, clupein sulphate, a protamine salt, which contains definite colloidal kations, coagulates an alkaline gold sol strongly. J. F. S.

Influence of Temperature on the Coagulation of Colloidal Gold. H. LACHS and STEPHANIE GOLDBERG (*Kolloid Z.*, 1922, **31**, 116—119).—The time required for the coagulation of gold sols prepared in various ways by a constant amount of sodium chloride has been determined at temperatures up to 70°, and in this way the influence of temperature on the rate of coagulation ascertained. The results show that the influence of temperature corresponds with that demanded by Smoluchowski's hypothesis, namely, that the coagulation time is inversely proportional to the absolute temperature. The sensitiveness, toward temperature changes, of the coagulation of colloidal gold is very much smaller than that of chemical reactions. J. F. S.

The Kinetics of the Coagulation of Proteins by Heat. HEINRICH LÜERS and MAX LANDAUER (*Z. angew. Chem.*, 1922, **35**, 469—471).—The velocity of coagulation of the plant albumin,

leucosin, was determined by the direct weighing of the coagulum produced at intervals during the reaction. The change of hydrogen-ion concentration during the reaction was very slight. It was concluded that the coagulation occurs in two stages; a chemical denaturation process followed by a colloidal phenomenon of coagulation, which is a flocculation of the denatured protein particles. The denaturation of the leucosin particles was shown to be a chemical reaction of the first order. The temperature coefficient of the denaturation in a medium of p_H 6.09 and at temperatures between 52.90° and 57.05° was 1.47 per 1°. A. G. P.

Is the Gelatin-Gum Arabic Coagulation a Chemical or a Colloid-chemical Process? F. W. TIEBACKX (*Kolloid Z.*, 1922, 31, 102—103).—A discussion of the coagulation of gelatin-gum arabic mixtures by acids, alkalis, and salts, from which it is shown that the coagulation is a mixed process which can be regarded as in part of a chemical nature and in part of a colloid-chemical nature. J. F. S.

Influence of Substituents on Various Chemical Reactions. SVEN BODFORSS (*Z. physikal. Chem.*, 1922, 102, 40—53).—It has been shown previously (A., 1918, i, 229) that a number of substituted aromatic aldehydes react with bromoacetophenone in the presence of sodium ethoxide according to the equation: $\text{Ph}\cdot\text{CHO} + \text{CO}\cdot\text{Ph}\cdot\text{CH}_2\text{Br} \rightarrow \text{CO}\cdot\text{Ph}\cdot\underset{\text{O}}{\text{CH}}\cdot\text{CH}\cdot\text{C}_6\text{H}_5$, whilst others do

not react in this way. The aldehydes which react all contain a negative substituent and those which do not may be brought to react by substituting such a negative group in them. It is now shown that the acids produced from the aldehydes which react are stronger than those produced from the aldehydes which do not react. The relationship between the reactivity of the aldehydes and the strength of the corresponding acid has been investigated. The strength of a number of substituted benzoic acids has been determined from conductivity measurements and the dissociation constant compared with the values for other acids found in the literature. The following newly determined values of K at 25° are recorded: 3-nitroanisic acid, 1.9×10^{-4} ; 3-bromoanisic acid, 7.2×10^{-5} ; cumic acid, 5.2×10^{-5} ; piperonylic acid, 4.5×10^{-5} . A further series of experiments has been carried out to ascertain the relationship between the maximum work of a given type of reaction and the nature of the substituting group. The reaction chosen is represented by the equation $\text{H} + \text{AgBz}(\text{solid}) = \text{HBz}(\text{solid}) + \text{Ag}$ and the maximal work of this reaction has been calculated from *E.M.F.* measurements of cells of the type $\text{Ag}|\text{AgBz}, \text{HBz}(\text{solid})||\text{HBz}(\text{solid})|\text{H}_2$ and $\text{Ag}|\text{AgBz}(\text{solid})||\text{HBz}(\text{solid})|\text{Quinhydrone}|\text{Pt.}$ at 25° (cf. Biilmann, A., 1921, ii, 372). The following values of the *E.M.F.* (π), the maximum work (A cal.), and the dissociation constant, respectively, are recorded: 5-nitrosalicylic acid, 0.066, 1520, 8.9×10^{-3} ; salicylic acid, 0.087, 2010, 1.0×10^{-3} ; *m*-nitrobenzoic acid, 0.109, 2510, 3.5×10^{-4} ; 3-nitroanisic acid, 0.135, 3110, 1.9×10^{-4} ; anisic acid, 0.144, 3320,

3.2×10^{-5} ; *m*-nitrocinnamic acid, 0.145, 3340, 8.3×10^{-5} ; *o*-chlorobenzoic acid, 0.147, 3390, 1.3×10^{-3} ; benzoic acid, 0.154, 3550, 6.7×10^{-5} ; *p*-nitrobenzoic acid, 0.160, 3690, 4.0×10^{-4} ; 3-bromoanisic acid, 0.163, 3760, 7.2×10^{-5} ; and piperonylic acid, 0.165, 3800, 4.2×10^{-5} . The above results show that generally the maximum work decreases with increasing dissociation constant. The cases where this relationship is not strict are probably to be explained either by the formation of hydrates by the silver salts or by the formation of acid silver salts.

J. F. S.

Equilibrium in Heterogeneous Systems. JA. K. SYRKIN (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1921, 4, 161—165).—Investigation of the reaction $\text{CaSO}_4 + \text{Na}_2\text{CO}_3 = \text{CaCO}_3 + \text{Na}_2\text{SO}_4$ with concentrations ranging from 0.01 to 0.002 gram-mol. per litre and thus corresponding with widely varying degrees of dissociation of the component salts, shows that the reaction does not conform to the law of equilibrium, the “constants” of the reaction exhibiting wide divergences. The reaction is reversible, the reversibility, that is, the proportion of non-reacting molecules, amounting to 58% at the ordinary temperature and at low concentrations. At 7°, scarcely any action occurs. The value of the reversibility at 38° is similar to that at 18°, probably owing to the solubility of calcium sulphate being at a maximum at 38°. The reversibility is a function of the concentration and increases with dilution, the more rapidly at low temperatures.

Thus, with reactions involving the formation of a new phase, the laws of equilibrium are incomparably more complex than with homogeneous reactions. At present, the dependence of the reversibility on the concentration is expressible only by an empirical relation of the form $Z = 1 + ac + bc^2$ T. H. P.

The Equilibria in Aqueous Solutions of the Alkali Metal Bisulphites. EDWARD CHARLES CYRIL BALY and ROBERT ARTHUR BAILEY (*T.*, 1922, 121, 1813—1821).

Equilibria and Reaction Velocities. W. F. BRANDSMA (*Chem. Weekblad*, 1922, 19, 318—322).—The treatment of reaction velocities based on molecular collisions of substances in the gaseous state or in dilute solution becomes difficult when the course of a reaction is not expressed by a simple equation, and experimental results do not agree with theory. The conditions leading to equilibrium may be regarded from two points of view, the first dealing with the forces involved, that is, the energy units, and the second with space factors, that is, with entropy units. The factors deduced from the first vary inversely with the absolute temperatures, and so, determine the temperature coefficients; the entropy factors are independent of temperature.

This division is of great importance in organic reactions, since the theoretical treatment can take due account of both energy and space factors.

S. I. L.

Theory of the Velocity of Chemical Reactions. J. N. BRÖNSTED (*Z. physikal. Chem.*, 1922, 102, 169—207).—A theoretical

paper in which the laws of the velocity of chemical reactions are considered. It is shown that chemical reactions between neutral molecules, or between neutral molecules and ions, are normal, that is, they follow very closely the law of mass action and are insensitive or only slightly sensitive to the addition of a neutral salt. Reactions between ions are abnormal, that is, the velocity constant is to a high degree dependent on the concentration of the reacting ions and on the concentration of an added neutral salt. For thermodynamical reasons, the concentrations or osmotic partial pressures in the ordinary kinetic reaction equations are to be replaced by the activities, and the equations are then true when the activities alone are changed. The reaction determinative of the velocity in a chemical change consists of the formation of an unstable critical complex in the sense put forward by Marcellin (A., 1915, ii, 328). The electrical charge of the complex is the sum of the charges of the reacting molecular species. The regularities and exceptions mentioned above are to be explained by the assumption that the velocity is inversely proportional to the activity coefficient of the critical complex, and that this is influenced by changes in the salt concentration in the same way as the ordinary ions. The data concerning the activity coefficients necessary for calculating the anomalies in the velocity are most easily obtained from solubility determinations of sparingly soluble salts in solvents containing dissimilar ions. The dependence of the activity coefficient on the concentration, obtained in this way, for the ions of various types makes it possible to calculate the extent of the neutral salt action, that is, the kinetic anomalies for the various types of reactions. On the basis of the foregoing, the following theoretical result is obtained. In reactions between ions of the same sign the salt action is positive, that is, the velocity is increased by an increasing salt concentration. In reactions where the reacting ions have opposite signs, the salt action is negative, that is, the reaction is retarded by increasing salt concentration. These theoretical results have been applied to a large number of chemical reactions in solution and for the greater number found to be confirmed, in some cases quantitatively. The kinetic and static anomalies may be removed by employing a concentrated solution of an indifferent salt as solvent.

J. F. S.

Velocity of Decomposition of Crystalline Substances.

A. SIEVERTS (*Z. physikal. Chem.*, 1922, **102**, 89—90).—An answer to Hinshelwood and Bowen (this vol., ii, 628; see also this vol., ii, 360; A., 1920, ii, 743; 1921, ii, 443).

J. F. S.

Mechanism of the Reduction of Permanganate and its Physico-chemical Foundation. III. Reaction between Manganate and Formic Acid. JOSEF HOLUTA (*Z. physikal. Chem.*, 1922, **102**, 32—39; cf. this vol., ii, 448, 628).—The velocity of the reaction between sodium manganate and sodium formate has been determined at temperatures varying between 27.4° and 22.6°, with constant concentrations of manganate and hydroxyl-ion and also with varying concentrations of all constituents. The results show

that the reaction as measured is of the second order, and is to be represented by the equation $\text{MnO}_4'' + \text{HCO}_2' + 2\text{H}_2\text{O} = \text{Mn}(\text{OH})_4 + \text{CO}_3'' + \text{OH}'$. Intermediate stages therefore do not need to be taken into account in the mechanism of this reaction. J. F. S.

Dependence of the Velocity of Alkaline Hydrolysis of Esters on the Constitution of the Alcohol. I. L. SMITH and HUGO OLSSON (*Z. physikal. Chem.*, 1922, **102**, 26—31).—The alkaline hydrolysis of propyl acetate and isopropyl acetate by 0.1145*N*-sodium hydroxide has been determined at 20°. The reaction constants are found to be 4.23 and 1.263, respectively, giving as the ratio between the propyl and isopropyl esters the value 3.4, a value which is very near the value (3.2) found by Menshutkin for the ratio of the rates of formation of the two esters.

J. F. S.

Catalysts and Chemical Equilibrium. J. F. DURAND (*Bull. Soc. chim.*, 1922, [iv], **31**, 759—762).—The conclusions drawn by Clarens (this vol., ii, 436) with regard to the effect of catalysts on the equilibrium of a system are criticised.

H. J. E.

Catalytic Formation of Water Vapour from Hydrogen and Oxygen in the Presence of Copper and Copper Oxide. ROBERT N. PEASE and HUGH S. TAYLOR (*J. Amer. Chem. Soc.*, 1922, **44**, 1637—1647; cf. this vol., ii, 148).—The catalytic combination of hydrogen and oxygen in the presence of metallic copper (produced by the reduction of copper oxide in hydrogen at 150—200°) has been investigated over the temperature range 100—200°, using gaseous mixtures containing up to 5% of oxygen. The results show that combination is practically complete at 200°, and is just measurable at 100°. In nearly all cases, copper oxide is formed simultaneously with water. After the oxygen is shut off and pure hydrogen alone passed over the catalyst, the oxide so formed is reduced at a greater rate than that of the formation of water while the oxygen is passing. Between 130° and 100°, the rate of oxidation of the copper and the rate of reduction of the oxide so formed, as well as the catalytic activity, suffer a marked reduction. At 130°, with oxygen equivalent to 10 milligrams of water passing in five minutes, the rate of formation of water passes through a maximum to a steady minimum value. The above results are satisfactorily explained by assuming that the combination takes place mainly as a result of the alternate oxidation and reduction of the catalyst, the reduction reaction being complicated by the known inhibitory influence of free oxygen on that reaction. J. F. S.

Catalysis of Hydrogen Peroxide by Finely Divided Platinum. The Influence of Inhibitors. EDWARD BRADFORD MAXTED (*T.*, 1922, **121**, 1760—1765).

Bohr and Langmuir Atoms. (SIR) OLIVER LODGE (*Nature* 1922, **110**, 341).—In view of the difficulty of explaining molecular combination in terms of electrical attraction between revolving electrons composing the peripheral parts of an atom (in accordance with

the demands of spectroscopic observations) rather than in terms of a static arrangement (more suited to the facts of chemical combination), it is considered that the idea of electrical attraction between the atoms as the major chemical force should give place to that of the interlacement of the magnetic fields which inevitably accompany rapidly revolving electric charges. A. A. E.

The Neuburger Nuclear Model. F. PABLO VALERAS (*Physikal. Z.*, 1922, 23, 304—305).—The relative merits of the Neuburger (cf. this vol., ii, 208, 365) and Meitner (cf. *ibid.*, ii, 15) models are discussed and an objection is raised to the latter model since it cannot be applied to the nuclei of boron¹⁰, boron¹¹, nitrogen, or fluorine. The Neuburger model is unsatisfactory, since it requires the existence of *isohelium* nuclei. The author proposes a new model in which the nuclear formulæ are given by $F_k = n\alpha + pH^+ + v\beta$ which takes account of all known atoms. W. E. G.

The Neuburger Nuclear Model. LISE MEITNER (*Physikal. Z.*, 1922, 23, 305).—A reply to Pablo Valeras (preceding abstract), pointing out that the new model is not in agreement with radioactive phenomena. W. E. G.

The Neuburger Nuclear Model. MAXIMILIAN CAMILLO NEUBURGER (*Physikal. Z.*, 1922, 23, 305—307).—A reply to F. Pablo Valeras (cf. preceding abstracts). The assumption of the existence of the *isohelium* nucleus is useful in explaining the structure of Gf^9 which cannot be expressed by the new model. A schematic representation is given of the structure and genesis of some of the lighter atoms. W. E. G.

The Stability of Atom Nuclei, the Separation of Isotopes, and the Whole Number Rule. WILLIAM D. HARKINS (*J. Franklin Inst.*, 1922, 194, 165—211; cf. A., 1915, ii, 543, 544, 814; 1920, ii, 479, 541).—The author has pointed out that of the first twenty-seven elements except hydrogen the atomic weights are very close to whole numbers on the basis of O=16, and those of which the atomic weights are multiples of 4 are much closer to whole numbers than the others. The simple explanation is that all the elements are intra-atomic compounds of hydrogen and that whilst in molecular compounds the latter has the atomic weight 1.0078, in these much closer combinations it has the atomic weight 1.000. Those atoms of which the atomic weights are divisible by 4 are regarded as being built up of α -particles which are known to be the nuclei of helium atoms. A system of notation denoting the electronic structure of the atoms has been introduced; for example, "Oxygen=16" is written $p_{16}e_{16}$ as a compound of 16 positive and 16 negative electrons. This is further elaborated to show the number of α -particles (α) present in the nucleus, and negative electrons in the nucleus are differentiated from those in the planetary system. The nuclei of elements of even atomic number can be represented as built up in this way of α -particles alone or with "cementing" negative electrons, usually two in

number. Those of odd number cannot be so represented but need the introduction usually of a "proton-electron" group, p_3e_2 .

From a consideration of the composition of meteorites, the earth's crust, and the radioactive series, the conclusion is drawn that the elements of even atomic number are much more stable than those which are odd. The atomic numbers of the five undiscovered elements are all odd. The stability is attributed to the great stability of the α -particle in the nucleus. The special relativity theory is used to explain the loss in weight of the four hydrogen atoms (4×1.0078) which form the helium atom (4.000), and the energy freed by this loss is shown to be 6.708×10^{11} cal. per gram-atom of helium formed. This theory is also applied to the calculation of weight losses in radioactive disintegrations, and it is shown that an experimental determination of the loss of mass of radiothorium over a period of three years would be a check on the validity of the theory. A. C.

Model of the Ionised Hydrogen Molecule. W. PAULI, jun. (*Ann. Physik*, 1922, [iv], **68**, 177—240).—A mathematical investigation of the conditions determining the stability of a model representing the motion of an electron under the influence of two positively charged nuclei, at rest and in equilibrium, a structure illustrating the ionised hydrogen molecule H_2^+ met with in positive rays. The analysis is based on an application of the quantum theory, and the extension of Bohr's correspondence principle to impacts between atoms or molecules and free electrons. Various orbits are investigated. Of quantised orbits symmetrical with regard to the medium plane bisecting at right angles the line joining the nuclei, that of minimum energy is characterised by three quantum numbers, 0, 1, 1, and fills up in a uniform manner the space contained within an ellipsoid of revolution. The possibility of the existence of a stable unsymmetrical quantised orbit remains for the present open. Consideration of the value of the ionisation potential of the hydrogen molecule indicates that the structure of the ionised hydrogen molecule is essentially meta-stable, and it is concluded that its real normal representation consists of an electron moving in a circular orbit in the medium plane between the nuclei, and characterised by conditions of stability and energy similar to those of the earlier Bohr model of the helium atom. Preliminary remarks are made concerning the application of the theory to the band emission spectrum of the ionised hydrogen molecule.

J. S. G. T.

The Mathematics of the Dicyclic Colour Theory, and a New Theory of the Structure of the Nitrogen Atom. JAMES MOIR (T., 1922, **121**, 1808—1813).

The Distribution of Electrons around the Nucleus in the Sodium and Chlorine Atoms. W. LAWRENCE BRAGG, R. W. JAMES, and C. H. BOSANQUET (*Phil. Mag.*, 1922, **44**, 433—449).—The distribution of electrons in the atoms of sodium and chlorine has been determined by measuring, by the method of X-ray analysis, the reflecting power per unit volume of the units constitut-

ing a crystal of sodium chloride. The experimental results indicate that the sodium atom comprises seven electrons arranged on a shell of radius 0.29 Å. and three electrons on a shell of radius 0.76 Å. The chlorine atom comprises ten electrons arranged on a shell of radius 0.25 Å., five electrons on a shell of radius 0.86 Å., and three electrons on a shell of radius 1.46 Å. Alternatively, the results are interpreted by continuous curves in which the electron density is plotted as ordinate against the corresponding distance of the electron from the centre of the atoms as abscissa. Considerable agreement is found between the experimentally determined values of the distribution factor F and those calculated for an atom of the type proposed by Bohr and comprising two electrons describing circular one-quantum orbits of radius 0.05 Å., four electrons describing two quantum circular orbits of radius 0.34 Å., and four electrons describing two-quantum elliptical orbits of semi-major axis 0.42 Å. The principal source of error in the conclusions arises owing to lack of information as to the part played by "extinction" in affecting the intensity of X-ray spectra. The results indicate that there cannot be, either in the sodium or chlorine atom, an outer shell of eight electrons. Eight electrons revolving in circular orbits of the same radius are equally inadmissible. J. S. G. T.

The Molecular Structure of Carbon Oxysulphide and Carbon Disulphide. A. O. RANKINE (*Phil. Mag.*, 1922, [vi], 44, 292—300).—The mean collision areas of molecules of carbon oxysulphide and carbon disulphide are calculated from certain assumptions based on the Lewis-Langmuir theory of valency (cf. A., 1921, ii, 192, 584). The electronic configuration in carbon oxysulphide is the same as that of an arrangement of argon and neon atoms in the sequence neon-neon-argon, and that of carbon disulphide corresponds with the sequence of inert atoms argon-neon-argon. The radii of the outer electron shells of atoms possessing the electronic configurations of neon and argon are 0.65 Å. and 1.03 Å., respectively, and the radii of the collision spheres of argon and neon atoms are 1.15 Å. and 1.44 Å., respectively. For the purpose of intermolecular encounters, the molecules consist of overlapping spheres the centres of which are separated by 1.30 Å. for neon-neon and 1.68 Å. for the argon-neon combination, and of which the radii are the collision radii of argon and neon atoms. The mean collision area of carbon oxysulphide should lie between the limits 1.00×10^{-15} cm.² and 1.09×10^{-15} cm.² and of carbon disulphide between the limits 1.23×10^{-15} cm.² and 1.37×10^{-15} cm.². In the former case, the agreement between the results of this examination and the value obtained by Smith (cf. this vol., ii, 686) is striking. In the latter case, comparison is not yet possible. W. E. G.

Valency Theories of Organic Chemistry. F. HENRICH (*Jahrb. Radioaktiv. Elektronik.*, 1922, 19, 1—38).—An historical account of the theory of valency leading up to a discussion of the modern views of Kossel, Hinsberg, J. Stark, Pauli, and Vorländer on the valency of organic compounds. W. E. G.

The Electron Theory of Valency as Applied to Organic Compounds. JULIUS STIEGLITZ (*J. Amer. Chem. Soc.*, 1922, 44, 1833—1834).—An addendum to a previous communication (this vol., ii, 560).

The rupture of the union of two carbon atoms by oxidation, involving the loss of two electrons by one of the atoms or jointly by both, takes place with particular ease when two neighbouring carbonyl groups are present in the molecule. In this case, there are no exposed valencies, but the structure is undoubtedly highly polar. Thus oxalic acid may be formulated,

$$\begin{array}{c} \text{H}^+-\text{O}-\overset{+}{\text{C}}-\overset{+}{\text{C}}-\text{O}-\text{H} \\ \quad \quad \quad \text{O}-\text{O}- \end{array}$$

and the high polarity of the field surrounding the carbon atoms evidently loosens the union between the two and exposes the electrons held by them somewhat in the same manner, but not to the same extent, as in the case of actual ionisation in a similar field. Similar considerations are applied to cyanogen and to dicarbonyl derivatives in general. H. W.

Lecture-room Demonstration of Atomic Models. LOUIS V. KING (*Phil. Mag.*, 1922, [vi], 44, 395—400).—Atomic models are described in which a number of steel spheres or soft iron rods are arranged in a strong alternating magnetic field. A simple form of the model consists of a coil of copper wire carrying alternating current over which is placed a large watch glass. Steel bearing-balls are used as magnetic elements. Since the spheres repel each other with a force varying as the inverse fourth power of the distance and the attraction to the centre varies as the distance, symmetrical stable groupings of the spheres are formed. When two coils are employed, mercury is used as a floating surface for the steel balls. The system of spheres may be set in oscillation by external magnets, giving a good illustration of the internal vibrations in the atom. The motion of molecules of a gas or the Brownian movement may be illustrated by employing short cylinders of iron or steel wire, which undergo violent movements if the field is suddenly applied. An experimental model illustrating the Rutherford atom may be constructed with two coils. Lengths of steel wire placed axially to the coils are used to make up the nucleus. Arrangements suitable for the demonstration of hydrogen or helium atoms are described. W. E. G.

Technical Sedimentation Analysis. I. FRIEDRICH-VINCENZ VON HAHN and DOROTHEA VON HAHN (*Kolloid Z.*, 1922, 31, 96—101).—The authors describe an apparatus, which is a slight modification of Ostwald and Hahn's two-limbed coagulation measurer, by means of which a rapid and accurate knowledge of the size of particles of commercial powders may be obtained. This apparatus has been employed to determine the relative size of the particles of a number of commercial preparations of soot and lamp-black. As a control of the results obtained with this apparatus, the older methods of differentiation have also been used, and it is shown that exactly similar results are obtained by all methods,

but that the new method is handier, more rapid, and more accurate (cf. this vol., ii, 205). J. F. S.

Gas Generator with Washing Apparatus. ROBERT MÜLLER, KOMANDIT-GESELLSCHAFT (D.R.-P. 346325; from *Chem. Zentr.*, 1922, ii, 978).—The apparatus consists of a conical flask fitted with a stopper which contains the essential features of the device. This consists of a tube, furnished with a funnel at the upper end and terminating in a trap at the lower end, through which the liquid used in generating the gas is introduced. The gas after generation passes through a washing apparatus of annular cross section fixed concentrically with the upper part of the tube.

G. W. R.

Inorganic Chemistry.

Solubility of Sulphur in certain Organic Liquids. R. DELAPLACE (*J. Pharm. Chim.*, 1922, [vii], 26, 139—140).—The solubility of sulphur in chloroform, carbon tetrachloride, benzene, toluene, and ether was determined at various temperatures ranging from 13° to 24°. The divergence of the results from those obtained by earlier workers is attributed in some cases to the slowness of saturation, twelve hours' continuous agitation being necessary to obtain saturated solutions in toluene and benzene, for example. From the results, which are given in tabular form, the following figures for the weights of sulphur dissolved in 100 grams of solvent, are abstracted: in chloroform at 15°, 0.874 gram; in carbon tetrachloride at 15.5°, 0.645 gram; in toluene at 20°, 1.857 grams; in benzene at 15°, 1.582 grams; in anhydrous ether at 13°, 0.188 gram.

G. F. M.

Autoreduction of Sulphurous Acid. GEORGE MACDONALD BENNETT (T., 1922, 121, 1794—1795).

Some Properties of Selenium Oxychloride. II. VICTOR LENHER (*J. Amer. Chem. Soc.*, 1922, 44, 1664—1667; cf. A., 1921, ii, 256).—Selenium oxychloride is extremely hygroscopic and the very greatest precautions must be taken in preparing the anhydrous material. The presence of small quantities of water increases the electrical conductivity enormously, and its reactivity towards carbonates is largely dependent on the amount of water present. The best qualitative test for the presence of water in selenium oxychloride makes use of cobalt carbonate. When cobalt carbonate which has been dried at 200° for three hours is sealed up with anhydrous selenium oxychloride, no action of any description occurs, but if the merest trace of water is present the oxychloride becomes blue and if the tube is sealed a pressure of carbon dioxide is set up. The dry carbonates of calcium, strontium, copper, nickel, cobalt, and ferrous iron are unattacked by anhydrous

selenium oxychloride, whilst the dry carbonates of barium, lithium, and magnesium slowly evolve carbon dioxide and that of zinc much more rapidly. The fused carbonates of sodium and potassium react readily with anhydrous selenium oxychloride with the development of a large amount of heat. Sulphur dioxide is without action on boiling anhydrous selenium oxychloride, but in the presence of water selenium is precipitated. Dry hydrogen sulphide in contact with selenium oxychloride produces a reddish-brown colour, after which hydrogen chloride is evolved and yellow selenium sulphide produced. At the same time, heat is developed which dissociates the sulphide into sulphur and red selenium. Liquid hydrogen disulphide reacts vigorously with selenium oxychloride in the cold, producing the same products as hydrogen sulphide. Carbon monoxide has no action on selenium oxychloride. Iodic acid, iodine pentoxide, and potassium iodate with selenium oxychloride give at first chlorine and then iodine chloride. Periodic acid with the oxychloride evolves chlorine and, when warmed, iodine chloride. Potassium bromate with selenium oxychloride in the cold gives chlorine followed rapidly by a mixture of bromine and chlorine. Selenium oxychloride warmed with potassium persulphate evolves chlorine, but in the presence of sulphuric acid the evolution occurs in the cold and with effervescence. Telluric and selenic acids evolve chlorine when gently warmed with the oxychloride. The metals tungsten, tantalum, and titanium are only slightly attacked by selenium oxychloride after long treatment at high temperatures. Phosphorus reacts with selenium oxychloride at low temperatures to form phosphoric oxide, selenium monochloride, and selenium tetrachloride. Selenium oxychloride dissolves readily in carbon tetrachloride, but after a time reaction takes place with the formation of selenium tetrachloride and carbonyl chloride. Gliadin, zein, glutenin, elastin, and albumin are readily attacked and dissolved by selenium oxychloride.

J. F. S.

Selenium Oxybromide. VICTOR LENHER (*J. Amer. Chem. Soc.*, 1922, **44**, 1668—1673).—Selenium oxybromide is best prepared by adding to pure sublimed selenium dioxide the calculated amount of fused selenium in a flask fitted with a rubber stopper through which passes a dropping funnel and a safety tube filled with fused calcium bromide. The flask is cooled to 0° and the amount of bromine necessary to convert the metal into tetrabromide cautiously added in small quantities. The mixture is then warmed until the whole of the oxide has dissolved to form the oxybromide. Selenium oxybromide is a reddish-yellow solid, m. p. 41.5—41.7°, b. p. 217°/740 mm. with considerable decomposition, d_{40}^{20} 3.38. The product decomposes so readily on heating, that it cannot be purified by distillation even under reduced pressure. It has an electrical conductivity at 40—50° of 6×10^{-5} ohms⁻¹. Water converts it slowly into selenious and hydrochloric acids. It dissolves in carbon disulphide, chloroform, benzene, toluene, and xylene, and the fused material is miscible in all proportions with these solvents. Carbon tetrachloride dissolves it readily, but the

fused material dissolves only to the extent of 6%, and on heating the solution for several days carbonyl chloride is evolved and selenium tetrabromide formed. Selenium oxybromide is a very active chemical reagent. Sulphur reacts readily with it in the cold with the evolution of sulphur dioxide; selenium dissolves in it to form the monobromide. Yellow phosphorus explodes with the solid, whilst red phosphorus takes fire with the solid and burns. Iodine dissolves in large quantities in selenium oxybromide, whilst chlorine displaces bromine. Silicon and carbon are not attacked. Sodium reacts explosively with the oxybromide, producing the bromide and selenium monobromide, whilst potassium reacts even more violently. Mercury, arsenic, antimony, tin, bismuth, iron, calcium, copper, lead, silver, molybdenum, thallium, gold, platinum, and zinc are all attacked, forming the metallic bromide and selenium monobromide. Zinc dust burns in the oxybromide; aluminium and magnesium are only slightly corroded after heating for a week at 100° in the oxybromide. Cadmium, chromium, nickel, cobalt, tungsten, and tantalum are not attacked when heated for ten days at 100°. The oxides of mercury, silver, calcium, and sodium react energetically with selenium oxybromide, those of arsenic, tin, and tellurium less energetically, and those of columbium, tantalum, vanadium, thorium, titanium, zirconium, and uranium are unattacked after heating for several days at 100°. With most carbonates, the anhydrous oxybromide reacts only sluggishly even on heating, but in the presence of moisture the reaction is vigorous. The commoner sulphides react with the oxybromide with the development of heat, forming metallic bromides and selenium monobromide. Potassium chlorate liberates bromine from the oxybromide, but potassium perchlorate, permanganate, dichromate, and chromium trioxide are without action on it. When dry air is bubbled through the oxybromide at 60°, bromine is set free, due to a primary dissociation into the dioxide and tetrabromide followed by a secondary dissociation of the tetrabromide into bromine and the monobromide. Sulphur dioxide and carbon monoxide are without action on the oxybromide, and hydrogen sulphide causes a slight decomposition.

J. F. S.

Production of Colourless Glass in Tank Furnaces with Particular Reference to the Use of Selenium. I. A. COUSEN and W. E. S. TURNER (*J. Soc. Glass Technology*, 1922, 6, 168—181).—Numerous small melts at 1380° and 1440° showed that not more than 0.09% of iron oxide in a glass could be effectively decolorised by selenium or sodium selenite. The greater corrosive action of the salt-cake on the pot and its action on the selenium at high temperatures rendered more decoloriser necessary for batches containing this ingredient than for those with soda ash alone. Experiments showed the distinct formation of sodium selenite from selenium when dropped on to fused sodium sulphate. In a batch containing only soda ash and no salt-cake, sodium selenite was much more effective than selenium in decolorising. Reheating the glasses only changed the tint of those which had been

over-decolorised, and then only when heated between 525° and 550°.

A. C.

Limits for the Propagation of Flame at Various Temperatures in Mixtures of Ammonia with Air and Oxygen. ALBERT GREVILLE WHITE (T., 1922, 121, 1688—1695).

Acetone in Commercial Ammonia. J. BOUGAULT and R. GROS (*J. Pharm. Chim.*, 1922, 26, 170—171).—In many cases in which ammonia is used as a reagent in the presence of iodine, the formation of small quantities of iodoform is observed. An explanation of this fact was put forward by Guérin (A., 1909, i, 126), but this is set aside by the authors, who show that iodoform results from the presence of acetone. The quantity of acetone was found to vary from 0.01—0.5 gram per litre in the different samples examined, and it has been found in so-called "pure ammonia for analysis" in addition to the commercial product. It is pointed out that the use of ammonia in testing for acetone in physiological products is liable to give misleading results.

H. J. E.

Viscous Properties and Molecular Dimensions of Silicane. A. O. RANKINE and C. J. SMITH (*Proc. Physical Soc.*, 1922, 34, 181—186).—The authors have determined experimentally the viscosity of gaseous silicon hydride at 15° and 100°, and on the assumption that Sutherland's law holds for this gas over the whole temperature range concerned the viscosity at 0° has been extrapolated. The data have been used to calculate the mean collision area presented by the silicon hydride molecule in the gaseous state. The experimental method employed was the same as that previously described (A., 1921, ii, 694). The following numerical data are recorded: viscosity in C.G.S. units $\times 10^{-4}$, 0°, 1.078; 15°, 1.134; 100°, 1.436; Sutherland's constant, (c), 229; mean collision area of the molecule, (A), 0.989×10^{-14} cm.². Using these data, an attempt is made to determine the structure of the silicon hydride molecule in relation to those of other gaseous hydrides the molecules of which have equal numbers of extra-nuclear electrons.

J. F. S.

Effect of Magnesia on the Resistance of Glass to Corroding Agents and a Comparison of the Durability of Lime and Magnesia Glasses. VIOLET DIMBLEBY, CONSTANCE M. M. MUIRHEAD, and W. E. S. TURNER (*J. Soc. Glass Technology*, 1922, 6, 101—107).—Magnesia glasses appear to possess a slight advantage over lime glasses as regards corrosion by water; they show no advantage in the case of hydrochloric acid, and are decidedly inferior as regards their resistance towards alkaline solutions.

W. P. S.

Crystal Structure of Potassium. L. W. McKEEHAN (*Proc. Nat. Acad. Sci.*, 1922, 8, 254—255).—Determination by the X-ray method of the crystalline structure of potassium cooled to -150° in a current of air shows that the atoms of the metal are arranged in a body-centred cubic lattice with $a = 5.20 \times 10^{-8}$ cm. The

corresponding value of the density is 0.92 gram/cm.³ compared with the commonly accepted value of 0.862 gram/cm.³. The difference is probably partly attributable to the imperfect crystallisation of samples previously examined. The observed crystalline structure does not persist as the temperature rises to about 20°.

J. S. G. T.

Symmetry and Crystal Structure of Sodium Hydrogen Acetate. RALPH W. G. WYCKOFF (*Amer. J. Sci.*, 1922, [v], 4, 193—198).—Cubes (d 1.402) having the composition $\text{NaH}(\text{C}_2\text{H}_3\text{O}_2)_2$ were obtained by the long digestion of fused sodium acetate, glacial acetic acid, and acetic anhydride. The X-ray analysis (reflexion spectra and Laue photographs) gives a unit cube of edge 15.98 Å.U. containing twenty-four chemical molecules. The space group is probably T_h^7 .

L. J. S.

Symmetry and Crystal Structure of Zinc Bromate Hexahydrate. RALPH W. G. WYCKOFF (*Amer. J. Sci.*, 1922, [v], 4, 188—192).—Good crystals having the composition $\text{Zn}(\text{BrO}_3)_2 \cdot 6\text{H}_2\text{O}$, mostly octahedral in habit, are formed by the slow cooling, or slow evaporation, of solutions. Reflexion X-ray photographs were obtained from the octahedral face, and Laue photographs through both octahedral and cube faces. These show an absence of planes of symmetry. From the various alternative arrangements, the space group T_h^6 is selected. The unit cube contains four chemical molecules and has length of edge 10.31 Å.U.

L. J. S.

Chemical Method of Separating the Isotopes of Lead. THOMAS DILLON, ROSALIND CLARKE, and VICTOR M. HINCHY (*Sci. Proc. Royal Dublin Soc.*, 1922, 17, 53—57).—The authors have attempted to effect a separation of the isotopes of lead by means of the Grignard reaction. Lead chloride prepared from some residues recovered from the manufacture of mesothorium was treated with lead tetraethyl and the reaction represented by the equation $2\text{PbCl}_2 + 4\text{MgRX} = \text{R}_4\text{Pb} + 2\text{MgCl}_2 + 2\text{MgX}_2 + \text{Pb}$ carried out. The lead tetraethyl and metallic lead remaining constitute two fractions which are used separately in a repetition of the process. The process was carried out twice and the atomic weight of the lead in the tetraethyl from one and the lead residue in the other process was then determined by the method employed by Soddy and Hyman (*T.*, 1914, 105, 1402). The relative values 207.1 and 207.4 were obtained, thus indicating that the above reaction furnishes a possible method of separating isotopes and that the chemical properties of isotopes although similar are not identical.

J. F. S.

The Mass-spectrum of Iron. F. W. ASTRON (*Nature*, 1922, 110, 312—313).—The mass spectrum of iron, examined as iron pentacarbonyl mixed with carbon dioxide, is characterised by a strong line at 55.94 ± 0.05 , and possibly a very faint line at 54.

A. A. E.

The Constitutional Diagram of the Iron-Carbon System based on Recent Investigations. KÔTARÔ HONDA (*Sci. Rep. Tohoku Imp. Univ.*, 1921, **11**, 119—129).—In a previous communication with Murakami (cf. A., 1921, ii, 699), it is shown that the graphitisation of iron always takes place after solidification of the fusion, the formation of graphite being due to the catalytic action of carbon monoxide on the solid cementite, just below the eutectic point, 1130° to 1100°. This being the case, the graphite lines on the usual double iron-carbon diagrams must be omitted. The shape of the solidus curve is frequently given incorrectly; according to the principles of thermodynamics, this curve, like the liquidus curve, should be bent upwards. In a discussion of the allotropic transformations of the various forms of iron, it is concluded that the α - β transformation is not a sudden change of phase, but a progressive change beginning from the lowest temperatures and ending at about 790°, the amount of change being dependent on the temperature. The magnetisation of iron is not accompanied by changes in the atomic configuration as in the case of allotropic change, but is a consequence of a gradual alteration in the energy of the atoms with rise in temperature. This transformation is represented by a dotted line passing through 790° for pure iron and 775° at 4% of carbon. Similarly, the cementite transformation is progressive and extends from the lowest temperature up to 215°. The break in the heating curve of cast iron observed by Ruer (A., 1921, ii, 198) is not evidence of the existence of a graphite eutectoid. W. E. G.

Crystal Structure of Iron and Steel. II. ARNE WESTGREN and GÖSTA PHRAGMÉN (*Z. physikal. Chem.*, 1922, **102**, 1—25; cf. this vol., ii, 152).—Röntgen photographs of iron wire at 800°, 1100°, and 1425°, obtained by the Debye and Scheerer method, show that iron within the so-called β and δ regions similarly to α -iron has a space-centred cubic lattice, whilst iron in the γ -iron region has a face-centred cubic lattice. The characteristic fact is therefore that the transition which occurs at 900° (A_3) is reversed at 1400° (A_4). The γ -iron lattice of austenite steel is extended by the dissolved carbon. It is found that a steel containing 1.98% of carbon possesses a larger iron lattice when it is quenched at 1100° than at 1000°, also that the α -iron lattice in martensite appears to be somewhat extended by the carbon. The homogeneous lattice region within the α -iron in martensite is shown to be extremely small. A steel containing 0.80% of carbon which has been quenched at 760° is shown to be practically amorphous. Debye photographs of cementite and spiegeleisen crystals are identical. By means of Laue photographs and an investigation of the interferences of a spiegeleisen crystal rotating in a Debye camera, it is shown that cementite belongs to the rhombic system and has an axial ratio 0.670 : 0.755 : 1. The dimensions of the elementary parallelepiped are 4.53, 5.11, and 6.77 Å.U., respectively. The unit crystal contains four molecules of Fe_3C , from which it follows that the specific gravity of cementite is 7.62. J. F. S.

The System Chromium Trioxide-Sulphur Trioxide-Water. LIONEL FELIX GILBERT, HAROLD BUCKLEY, and IRVINE MASSON (T., 1922, 121, 1934—1938).

Rôle of Chromates in Electrolysis for Chlorate. A. V. PAMFILOV (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1921, 4, 113—117).—Chromate plays a triple part in the electrolytic production of chlorate in an acid solution. (1) It acts as a "buffer" salt, regulating automatically the kinetic acidity of the liquid. (2) It forms a film on the cathode and thus prevents reduction of the hypochlorite formed as an intermediate product, and (3) this film protects the cathode from disintegration and hence allows the platinum electrodes to be replaced by those of metals more readily accessible, such as copper. These effects of chromate are specific and are not exhibited by any other compound. T. H. P.

The Adsorption of Uranium-X and its Isotope, Thorium, by Basic Ferric Acetate. ANDREW CHARLES BROWN (T., 1922, 121, 1736—1743).

Zirconium Ferrocyanide and Ferricyanide. F. P. VENABLE and E. O. MOEHLMANN (*J. Amer. Chem. Soc.*, 1922, 44, 1705—1707).—When solutions of potassium ferrocyanide are added to solutions of zirconyl salts, white precipitates are obtained which are unstable in the air and of varying composition. The precipitates are basic zirconium ferrocyanides and the composition depends on the extent to which the hydrolysis has proceeded. Thus, a freshly prepared solution of zirconyl chloride gives a substance of the formula $\text{ZrO}(\text{OH})_2 \cdot 2(\text{ZrO})_2\text{Fe}(\text{CN})_6$ and a solution which has been boiled gives a precipitate of the composition $9\text{ZrO}(\text{OH})_2 \cdot 2(\text{ZrO})_2\text{Fe}(\text{CN})_6$. Potassium ferricyanide gives no precipitate with freshly prepared solutions of zirconyl chloride, but on boiling the mixture the very basic compound $21\text{ZrO}(\text{OH})_2 \cdot 2(\text{ZrO})_3\text{Fe}(\text{CN})_6$

is precipitated.

J. F. S.

The Chemical Properties of Mesothorium. II. D. YOVANOVITCH (*Compt. rend.*, 1922, 175, 307—309).—A concentrated solution of the chlorides of barium and radioactive elements was treated with hydrochloric acid in the cold. The precipitated barium chloride carried down with it mesothorium-I, thorium-X, and radium, leaving mesothorium-II, radiothorium, and the active residue in solution. By this process, repeated at intervals, mesothorium-II was obtained in the acid solution, nearly pure and practically free from barium. The final purification of mesothorium-II was carried out by repeated co-precipitation with salts of iron or aluminium, and further by reprecipitation as hydroxide. To eliminate radiothorium a trace of thorium was added and later removed by means of hydrogen peroxide. Thorium-C and thorium-B were removed by precipitation with lead and bismuth, which in turn were removed as sulphides. Mesothorium-II in solution with elements of the cerium group was fractionally precipitated as oxalate hydroxide, and as the double

sulphate. Mesothorium-II in the presence of lanthanum and praseodymium appeared in greater quantity in the early fractions, but when mixed with neodymium and samarium was more concentrated in the last fractions. In the fractionation of the double nitrates of the rare earths and ammonium, mesothorium-II tends to accumulate with lanthanum rather than with samarium.

A. G. P.

Crystal Structure of Germanium. N. H. KOLKMEIJER (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 25, 125—126).—In continuation of previous work (A., 1921, ii, 200), the author has investigated the crystal structure of germanium, by the powder method. The structure, deduced from the observations, resembles that of the diamond, the lattice parameter a being equal to 5.61 \AA ., compared with the value 5.594 \AA . deduced from the value of the density and atomic weight, taking Avogadro's number equal to 6.062×10^{23} . As modifications of carbon, silicon, germanium, and tin resembling the diamond in structure are known, whilst the same is not true of titanium, zirconium, and thorium, it is concluded that carbon and silicon are more closely associated with the elements of group IVb than with those of group IVa.

J. S. G. T.

The Anode Oxidation of Gold. I. Anode Oxidation of Gold in Sulphuric Acid. FR. JIRSA and OT. BURYÁNEK (*Chem. Listy*, 1922, 16, 189—194).—A detailed examination is made of the oxidation of a gold anode in dilute sulphuric acid. The product obtained is shown to be auric hydroxide, which loses a molecule of water on keeping over phosphoric oxide, giving the compound $\text{AuO}(\text{OH})$, and this, on further dehydration, yields the oxide Au_2O_3 . It was observed that ozone is evolved during the passage of the current, and the amount produced per ampere-hour for various concentrations of acid is determined. The yields of hydroxide obtained with different currents and acid concentrations are also measured.

The solubilities of auric hydroxide, both freshly prepared and stale, and prepared both electrolytically and by hydrolysis, are determined for different concentrations of sulphuric acid, and from these results it is concluded that the product of solution is auric sulphate, and not the compound $\text{Au}(\text{OH})\text{SO}_4$.

R. T.

Mineralogical Chemistry.

The Composition and Calorific Power of Coal from the Deeper Strata of the Kenadza (Algerian) Mine. A. FOIX (*Bull. Soc. chim.*, 1922, [iv], 31, 813—816).—Coal from the deeper workings differs from that obtained nearer the surface by reason of its greater proportion of ash, nitrogen, and sulphur. The heat of combustion is about 8.255 calories per gram.

H. J. E.

Colloidal Phosphates. CARL ELSCHNER (*Kolloid Z.*, 1922, **31**, 94—96).—The author describes the colloidal formation of a clay found in the Fanning Islands which contains no silicates but is chiefly made up of calcium phosphates and organic materials. The mineral *naurite* is stated to be formed from a jelly-like mass containing tricalcium phosphate with uric acid and purine derivatives, which through the action of nitrifying bacteria has lost its organic matter and formed a substance of the composition $3(\text{Ca}_3\text{P}_2\text{O}_8), \text{Ca}(\text{OH})_2, x\text{H}_2\text{O}$. Another mineral, *meyersite*, found in the Hawaii Islands, is described. This is an aged gel of the composition AlPO_4 , 66.33%; FePO_4 , 2.52%; H_2O , 26.10%; and is to be distinguished from *variszite*, $\text{AlPO}_4 \cdot 2\text{H}_2\text{O}$. It is lighter in colour and is particularly characterised by its banded structure. The mineral is formed by the decomposition of lava by the constituents of guano.

J. F. S.

Olivine from Lithosiderites. P. CHIRVINSKII (*Bull. Soc. franç. Min.*, 1921, **44**, 170).—Olivine isolated from the pallasites of: I, Ahumada; II, Finmarken (*d* 3.366); III, Mt. Dyrring; and IV, Molong, gave:

	SiO_2	FeO	MnO	MgO	CaO	Total
I.	38.43	14.93	trace	46.27	0.40	100.03
II.	39.32	16.61	—	44.07	—	100.00
III.	40.52	13.13	—	47.30	trace	100.95
IV.	39.32	13.13	trace	47.70	trace	100.15

L. J. S.

Cyprine and Associated Minerals from Franklin, New Jersey. J. VOLNEY LEWIS and LAWSON H. BAUER (*Amer. J. Sci.*, 1922, [v], **4**, 249—251).—Sky-blue, granular to fibrous cyprine (a variety of idocrase), which was apparently homogeneous, was found on microscopical examination of thin sections to enclose about 33% of rounded grains of willemite. Analysis I gives, after deducting willemite, the figures under II agreeing approximately with the composition of idocrase. Associated with this mineral is pale pink, cleavable to coarsely fibrous rhodonite (var. *bustamite*), which gave analysis III; brown, granular garnet—the polyadelphite variety of andradite (analysis IV); also phlogopite, felspar, etc.

	SiO_2	Al_2O_3	Fe_2O_3	FeO	MnO	ZnO
I.	32.42	14.07	—	0.77	1.60	21.71
II.	35.14	20.86	—	—	—	—
III.	46.72	1.34	—	0.46	26.51	1.34
IV.	34.28	3.12	25.53	—	7.41	—

	CuO	CaO	MgO	H_2O	Total
I.	0.99	25.22	1.08	2.38	100.14
II.	1.47	37.40	1.60	3.53	100.00
III.	—	22.24	1.27	—	99.88
IV.	—	29.20	0.39	—	99.93

L. J. S.

Analytical Chemistry.

Refractometric and Interferometric Quantitative Analysis. J. BRÜKA (*Z. physiol. Chem.*, 1922, 121, 288—299).—The author has extended the observations of de Crinis (A., 1920, ii, 700) on the use of the refractometer for the analysis of solutions so as to make it more generally applicable. He has also described the use of the interferometer for more dilute solutions. The experiments have special reference to urine analysis. H. K.

Recovery of Solvents from Extracted Substances. A. B. PICHLER (*Chem. Ztg.*, 1922, 46, 698).—In the ordinary method of extracting substances in a Soxhlet apparatus a considerable quantity of the solvent is retained by the extraction thimble and its contents, particularly if the latter are of a bulky nature; this solvent may be recovered by transferring the thimble, at the end of the extraction, to a large flask and, when a suitable number of thimbles have thus been collected, submitting them to distillation. To recover solvent from a single thimble, the latter may be transferred to a tube which fits on to the lower end of the Soxhlet apparatus and the solvent then distilled into and collected in the Soxhlet apparatus.

W. P. S.

Apparatus for Measuring the Hydrogen-ion Concentration of a Solution. ANDRÉ KLING and A. LASSIEUR (*Ann. Chim. Analyt.*, 1922, [ii], 4, 232—238).—A further description of the electrometric method previously reported (this vol., ii, 234). G. F. M.

Colorimetric Estimation of Hydrogen Peroxide. M. L. ISAACS (*J. Amer. Chem. Soc.*, 1922, 44, 1662—1663).—A rapid method of estimating the concentration of hydrogen peroxide colorimetrically based on the method described by Denigès (A., 1890, 1185) and Crismer (A., 1892, 381) is worked out and fully described. About 30 c.c. of water, 10 c.c. of a 5% solution of citric acid, and 1 c.c. of the unknown hydrogen peroxide are placed in a 50 c.c. graduated flask and, after mixing, 1 c.c. of a 10% solution of ammonium molybdate is slowly added. Water is then added to make the volume 50 c.c. and the whole well shaken. The yellow colour develops at once and the solution is then compared with a standard potassium chromate solution (0.4 gram/litre) in a Dubosq colorimeter. When the standard is set at 20 the quantity of hydrogen peroxide is given by $x=0.05467/y$, where y is the colorimeter reading and x the number of grams of peroxide in the 50 c.c. of solution. The results obtained by this method compare extremely favourably with those obtained by direct titration with potassium permanganate. J. F. S.

The Error in Estimating Chloride by Bang's Micro-Method. RICHARD PRIGGE (*Biochem. Z.*, 1922, 130, 442—447).—The sources of error in this method have been examined

and full details of the technique required for the greatest exactitude are given. H. K.

Micro-analyses by Bang's Methods. I. LUDWIG PETSCHACHER (*Biochem. Z.*, 1922, **131**, 116—123).—The estimation of chlorides by microanalysis in drops of blood gives sufficiently accurate results, but the estimation of dextrose and residual nitrogen is not sufficiently accurate. H. K.

Colorimetric Estimation of Blood Chlorides. M. L. ISAACS (*J. Biol. Chem.*, 1922, **53**, 17—19).—The method is applicable to the tungstic acid blood filtrates and depends on the conversion of silver chromate into sodium chromate by the action of the blood chlorides. Excess of silver chromate (about 0.05 gram) is stirred into 10 c.c. of the blood filtrate previously neutralised by addition of a small quantity of magnesium carbonate. After centrifuging, the supernatant liquid (plus washings) is clarified by the addition of 1 c.c. of a 2% ammonium hydroxide solution, and diluted to 25 c.c. The colour is then compared with a potassium chromate solution which has been standardised by means of a solution of sodium chloride in the above manner. E. S.

Estimation of Chlorine in Tissues. O. M. PICO and J. MURTAGH (*Compt. rend. Soc. Biol.*, 1922, **86**, 405—406; from *Chem. Zentr.*, 1922, ii, 922).—The tissue is dissolved in concentrated sodium hydroxide solution and the proteins are precipitated by phosphotungstic acid. The chlorine is estimated in the filtrate by the method of Austin and Van Slyke. G. W. R.

Detection of Iodates in Potassium Iodide according to the French Codex, 1908. LACHARTRE (*J. Pharm. Chim.*, 1922, [vii], **26**, 134—138).—The test for iodates in potassium iodide described in the Codex lacks precision, and takes no account of the influence of such factors as the strength of the iodide solution, the amount of acid added, the duration of the reaction, etc. The following method is suggested. A 10% solution of the iodide in distilled water is prepared, and to 10 c.c., 3 c.c. of chloroform and 5 drops of glacial acetic acid are added. After shaking for thirty seconds, the chloroform, examined immediately after separation, should not be coloured. Performed in this manner, the limit of sensitiveness of the test is 0.08 part of iodate per 1000, which is considered to be all that is required for a reasonably pure product. The presence of copper in small quantities, for example, 0.0015 part per 1000, has no effect on the reaction with a perfectly pure iodide, but it appreciably accelerates the development of the iodine coloration when traces of iodate are present. The amount of copper in ordinary distilled water is, however, too small to have any effect on the reaction in either case. G. F. M.

Detection and Estimation of Fluorides. N. K. SMITT (*Bull. Bureau Biotech.*, 1922, 176—178).—Greeff's volumetric method for the estimation of fluorides (*A.*, 1913, ii, 975) may be simplified somewhat by the use of an alcoholic solution of ferric chloride. The presence of fluoride in neutral solution may be detected by

adding a little ammonium thiocyanate solution and then ferric chloride solution drop by drop; in the absence of interfering substances, rapid disappearance of the colour of the ferric thiocyanate indicates the presence of fluoride. Even insoluble fluorides, such as that of calcium, may be similarly detected if the substance is vigorously ground in a mortar with thiocyanate and ferric chloride solutions. A colorimetric method for estimating traces of fluorides, based on the above reaction, is also described. [Cf. *J. Soc. Chem. Ind.*, 1922, Oct.] T. H. P.

The Estimation of Oxygen in Organic Compounds. H. TER MEULEN (*Rec. trav. chim.*, 1922, **41**, 509—514; cf. Boswell, A., 1913, ii, 334).—The substance to be analysed is heated in a current of pure hydrogen and the resulting gases are passed first over metallic nickel in the form of thin wire and secondly over nickelised asbestos. The function of the former is that of transforming the decomposition products of the substance into compounds which are suitable for hydrogenation, which reaction occurs in the presence of the latter. With proper precautions, all the oxygen is transformed into water and weighed as such; but as a portion may remain as carbon dioxide, a second absorbent is required to obviate error due to incomplete transformation. Analyses carried out by this method have yielded satisfactory results, details of which are given. Halogens and sulphur act as poisons to the catalyst; the method is therefore not applicable to substances containing these elements. As calcium chloride is used to absorb the water formed in the reaction, nitrogenous substances may not be dealt with by this method on account of the ammonia which they yield on hydrogenation. H. J. E.

Estimation of Sulphides by Oxidation by means of Ferric Sulphate. P. P. BUDNIKOV and K. E. KRAUZE (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1921, **4**, 157—159).—The method described is based on the reactions $3\text{Na}_2\text{S} + \text{Fe}_2(\text{SO}_4)_3 = 3\text{Na}_2\text{SO}_4 + 2\text{FeS} + \text{S}$ and $2\text{FeS} + 2\text{Fe}_2(\text{SO}_4)_3 = 6\text{FeSO}_4 + 2\text{S}$, the ferrous sulphate thus formed being estimated by titration with permanganate solution. If the ferric sulphate is used in aqueous solution, it undergoes hydrolysis, whereas in presence of acid the sodium sulphide may be attacked with loss of hydrogen sulphide. These difficulties are avoided by adding sodium carbonate or hydroxide, and by acidifying, only after the oxidation, with phosphoric acid. Manganous sulphate is added prior to the titration. The method is applicable to calcium and zinc sulphides, but not to sulphides of copper, lead, silver, etc., which are insoluble. T. H. P.

Estimation of Non-protein Nitrogen by means of Ultrafiltration. M. RICHTER-QUITTNER and H. HOENLINGER (*Wiener Klin. Wochschr.*, 1921, **34**, 24).—Twenty c.c. of blood plasma or serum are diluted ten to twenty times with distilled water and filtered through a Buchner funnel and membrane (Zsigmondy and Jander, A., 1919, ii, 520), the non-protein nitrogen being estimated in the filtrate by Kjeldahl's or Pregl's micro-method.

The Triketohydrindene (Ninhydrin) Reaction as a Quantitative Colorimetric Method for the Estimation of Amino-acid Nitrogen. H. RIFFART (*Biochem. Z.*, 1922, **131**, 78—96).—The ninhydrin reaction is suitable for the estimation of amino-acid nitrogen if certain precautions are taken. All the amino-acids with free amino-groups examined, except histidine, give at P_H 6.976 the same standard colour if the solutions be equal in amino-acid nitrogen content, say, 10 mg. or less per litre. This result is obtained by taking 2 c.c. of such a solution, neutralising with $N/400$ alkali to P_H 6.976 by comparison with phosphate standards and methyl-red, adding 2 c.c. of phosphate solution of the same P_H and 1 c.c. of 1% ninhydrin solution, and heating for thirty minutes in the boiling-water-bath. The solutions are then diluted to 100 c.c. and estimated colorimetrically. Histidine gives the standard colour at P_H 6.24, thus involving a small error where mixtures of amino-acids contain this acid. Some amines and ammonium salts also give the colour reaction, but only in concentrations higher than 15 mg. per litre. H. K.

Estimation of Phosphorus in Minerals and in Coke Ash by a Colorimetric Method. GEORGES MISSON (*Bull. Soc. Chim. Belg.*, 1922, **31**, 222—225).—The phosphorus to be estimated is obtained in solution as phosphate, and it is essential that the solution should be free from chlorine. Potassium permanganate is added, followed by a solution of sodium peroxide in dilute nitric acid. After addition of ammonium vanadate, the solution is boiled and treatment with ammonium molybdate produces the colour which is compared with the standard tint. Three modifications of the method are given, (a) for minerals of small phosphorus content, 0.01—0.20%; (b) for minerals of greater phosphorus content, 0.20—2.50%; (c) for coke ash, together with details of the procedure in each case. A comparison of the results with those obtained by gravimetric methods shows that they are substantially in agreement, but the colorimetric method gives somewhat lower results for a phosphorus content of 1.7% and upwards.

H. J. E.

Modification of the Bell-Doisy Phosphate Method. A. P. BRIGGS (*J. Biol. Chem.*, 1922, **53**, 13—16; cf. A., 1920, ii, 769).—In the modification described, the coloration produced in acid solution, which is more stable than that in alkaline solution, is used for the estimation. The blood or plasma is diluted with 3 volumes of water and 1 volume of 20% trichloroacetic acid, shaken vigorously for a few seconds, and filtered after ten minutes. To 5 c.c. of the filtrate (corresponding with 1 c.c. of plasma), 2 c.c. of the molybdate solution, 1 c.c. of a 20% sodium sulphite solution, 1 c.c. of the quinol solution, and water to bring the volume to 10 c.c. are added. After thirty minutes, the coloration produced is compared with a standard. In the presence of the sodium sulphite reagent a blue coloration is obtained which is more intense than the green coloration produced in its absence. The method is applicable to urine. E. S.

Estimation of Lipoid Phosphoric Acid ("Lecithin") in Blood by Application of Bell and Doisy's Method for Phosphorus. F. S. RANGLES and ARTHUR KNUDSON (*J. Biol. Chem.*, 1922, 53, 53—59).—An alcohol-ether extract of blood or plasma is prepared according to Bloor's procedure (A., 1918, ii, 452). An aliquot part of this is evaporated, treated with sulphuric and nitric acids, and heated until the latter acid is completely driven off. Phosphoric acid is then estimated in the clear solution by Bell and Doisy's method (A., 1920, ii, 769). Disturbance of the final solutions produces bubbles which interfere with the reading. The solutions should therefore be left in the colorimeter cups for five minutes before making the comparison. E. S.

Conductometric Titration of Hydrofluosilicic Acid. NAOTSUNA KANÔ (*J. Chem. Soc. Japan*, 1922, 43, 556—560).—Using two adjustable parallel platinum plates as electrodes, the author has carried out the conductometric titration of hydrofluosilicic acid. When a mixture of 100 c.c. of water and 170—200 c.c. of 95% alcohol was used as solvent for the 0.138*N*-acid, the titration was conducted with sodium hydroxide solution as well as with potassium hydroxide solution without causing hydrolysis of the acid. K. K.

Critical Examination of Methods commonly used in Determining the Durability of Glass. W. E. S. TURNER (*J. Soc. Glass Technology*, 1922, 6, 30—45).—The methods investigated involved (1) the testing of the surface of glass vessels, (2) the use of plates, disks, or slabs of glass, (3) the treatment of the glass in the form of powder, and (4) the autoclave test. Whilst there are uncertainties in the results obtained by any of the methods, it is not difficult to distinguish qualitatively a good glass from a poor one. It is suggested that, in absence of a satisfactory general test, the tests should be made on plates of the glass or on the powder, preferably the latter. W. P. S.

New Apparatus for Estimation of Carbon Dioxide in Air. HENRIK LUNDEGÅRDH (*Biochem. Z.*, 1922, 131, 109—115).—The author has modified the Pettenkoffer method by replacing the flask by a zinc vessel in the form of a beaker with detachable cover, and fitted with three tubes to facilitate withdrawal and entry of solutions. A larger apparatus is also described on similar lines, suitable for use in the open. H. K.

Estimation of Gases in Metals. HAROLD LESTER SIMONS (*Chem. and Met. Eng.*, 1922, 27, 248—249).—A weighed quantity of the metal is introduced into a specially designed apparatus which is then exhausted by means of a Toeppler pump. Mercury is allowed to siphon over into the flask containing the metal and the whole is heated until the latter dissolves. The evolved gases are first examined spectroscopically in a Geissler tube, then transferred to the usual gas apparatus, and the quantity of each constituent estimated in the usual manner. [Cf. *J. Soc. Chem. Ind.*, 1922, 714A.] A. R. P.

Estimation of Small Quantities of Sodium in Aluminium and Alumina. R. GEITH (*Chem. Ztg.*, 1922, 46, 745).—The precipitation of aluminium hydroxide by even a very slight excess of ammonia is invariably not quite complete and the traces in solution are weighed with the sodium chloride. By electrolysis of a nearly neutral solution of aluminium chloride or sulphate containing iron, magnesium, calcium, and sodium salts as impurities, all the sodium may be deposited in a mercury cathode as amalgam whilst the bulk of the other metals are precipitated as hydroxides in the solution and the chlorine escapes as gas from the carbon anode. In a suitably arranged divided cell the mercury is made the cathode in the inner compartment and the anode in the outer, which contains dilute neutral sodium chloride solution and a nickel rod dipping into the mercury and forming a voltaic couple. The sodium then redissolves as hydroxide in the salt solution and its amount may be estimated by titration. The current required for the method is 0.15 ampere at 3.5 volts for two to three hours. [Cf. *J. Soc. Chem. Ind.*, 1922, 714A.] A. R. P.

A New Method of Qualitative Separation of the Alkaline-earth Metals. MICHEL POLONOVSKI (*Bull. Soc. chim.*, 1922, [iv], 31, 806—810).—The separation is based on the differences in solubility of the sulphites and thiosulphates of the three metals in ethyl alcohol-water mixtures. The carbonates as precipitated in the group are dissolved in acetic acid, and the solution is boiled and neutralised with ammonia. After addition of ethyl alcohol, the barium is precipitated by alcoholic sodium thiosulphate. A small proportion remains in solution; this is separated, after filtration, by addition of potassium sulphate, but not of sufficient concentration to precipitate any strontium. The strontium is then separated as sulphite by excess of alcoholic sodium sulphite or as sulphate by excess of potassium sulphate. The solution is then acidified with very dilute acetic acid and the calcium precipitated by ammonium oxalate. The concentrations of the reagents required to effect the separation are given, also the proportions in which they are to be used. H. J. E.

Approximate Volumetric Estimation of Barium. MICHEL POLONOVSKI (*Bull. Soc. chim.*, 1922, [iv], 31, 810—813; cf. previous abstract).—The separation of barium from strontium and calcium as thiosulphate may be utilised as a means of volumetric estimation by using excess of a standard sodium thiosulphate solution which contains ethyl alcohol. After allowing the mixture to remain for twenty-four to thirty-six hours to obtain as complete a precipitation of barium thiosulphate as possible, the solution is filtered and the excess of the reagent estimated. The quantity of barium present is thus found by difference. A correction is necessary by reason of the slight solubility of barium thiosulphate; this is applied by a formula given by the author, as the temperature at which the precipitation takes place and the percentage of alcohol present are factors which influence the result. H. J. E.

Use of Amalgams in Volumetric Analysis. V. Estimation of Iron, Titanium, Uranium, and Methylene-blue. SUETARÔ KIKUCHI (*J. Chem. Soc. Japan*, 1922, 43, 544—550; cf. A., 1921, ii, 596, 714; this vol., ii, 519, 529).—After the reduction of mixed solutions of two metals with liquid zinc- or cadmium-amalgam, their differential titration with two standard oxidising solutions was studied. A mixture of ferric oxide and titanium dioxide is titrated with potassium permanganate and iron alum, using ammonium thiocyanate as an indicator after reduction with zinc amalgam. They are quantitatively oxidised with iodine in the presence of sodium tartrate. Combining this reaction and that with iron alum, they can also be differentially titrated. A mixture of ferric oxide and uranium trioxide is titrated with potassium permanganate and iron alum, after reduction with cadmium amalgam in warm solution and in the presence of carbon dioxide. A mixture of titanium dioxide and uranium trioxide is titrated with potassium permanganate and iodine solutions, after reduction with cadmium amalgam in warm solution. Quadrivalent uranium is not oxidised by excess of iodine in the presence of sodium tartrate.

After a mixed solution has been oxidised with potassium permanganate, it is reduced with the amalgam and titrated differentially with iron alum or iodine. A definite quantity of titanous sulphate is reduced to tervalent titanium salt by the amalgam; methylene-blue is added at 60—70° until the solution is tinged with blue, and then it is titrated. The purity of Kahlbaum's methylene-blue B extra was 82.96%, and that of Merck's methylene-blue B.B., 85.07%.

K. K.

Use of Amalgams in Volumetric Analysis. VI. Electro-metric Titration of Iron, Molybdenum, Uranium, Vanadium, and Titanium. NAOTSUNA KANÔ (*J. Chem. Soc. Japan*, 1922, 43, 550—556; cf. preceding abstract).—Solutions of iron alum, ammonium molybdate, uranyl sulphate, vanadic acid, and titanium sulphate are reduced with zinc amalgam, separated from the amalgam, connected to the platinum and calomel electrodes, and their *E.M.F.* estimated, the standard solution of potassium permanganate meanwhile being gradually added. In these estimations, the volumes of the solution should be maintained as small as possible. In the case of titanium, a slight deficiency of the permanganate solution is first added, then excess of iron alum solution to oxidise tervalent titanium, and the solution then titrated with the permanganate solution. A mixture of iron and titanium can be electrometrically titrated with the permanganate solution, two break-points in the *E.M.F.* curve showing the end points.

K. K.

Estimation of Tin in Bearing Metals. JOSEF NÁGEL (*Chem. Ztg.*, 1922, 46, 698).—One gram of the metal turnings is dissolved by boiling in a flask with 100 c.c. of hydrochloric acid and 3 grams of potassium chlorate; after all free chlorine has been expelled, the solution is diluted with 30 c.c. of water, 7 grams of sodium chloride are added, and, after the addition of a strip of nickel having

an area of about 10 sq. cm., the solution is boiled for one hour. The stannic chloride is thus reduced to stannous chloride, whilst the antimony is deposited as flakes on the nickel plate. After the solution has been cooled in an atmosphere of carbon dioxide, the stannous chloride is titrated with standardised iodine solution.

W. P. S.

The Volumetric Estimation of Antimony and Tin in Red Brass. FRANZ JOHANN MÜCK (*Chem. Ztg.*, 1922, 46, 790).—The alloy is dissolved in nitric acid and the precipitated stannic and antimonic acids are collected, washed, and dissolved in hot sulphuric acid and potassium sulphate. The cooled solution is treated with 0.5 gram of tartaric acid, 180 c.c. of water, and 7 c.c. of hydrochloric acid and the antimony estimated by titration with permanganate. The solution is then boiled with a further 30 c.c. of water, 60 c.c. of hydrochloric acid, and 2–3 grams of metallic antimony, and the tin is estimated by titration with iodine [cf. *J. Soc. Chem. Ind.*, 1922, Oct.].

A. R. P.

Decomposition and Analysis of Monazite. P. WENGER and P. CHRISTIN (*Ann. Chim. Analyt.*, 1922, [ii], 4, 231–232).—The hitherto published methods for the attack of monazite are either long and delicate, or else leave large quantities of the rare earths undissolved. By the following method the whole of the mineral can readily be brought into solution. About 1 gram of the finely ground material is heated for five hours with 5 c.c. of sulphuric acid (d 1.84) at 230°. The resulting pasty mass is gradually poured into iced water, when all but a small residue passes into solution. This residue is collected on a filter and again heated with sulphuric acid, a little hydrofluoric acid being added to volatilise the silica. The residue in the crucible is then melted with six parts of sodium carbonate. On addition of water, the tantalum passes into solution, and any slight residue which still may remain will be zirconia, which is rendered soluble by fusion with potassium hydrogen sulphate. In the subsequent analysis the precipitation of the rare earths by oxalic acid should be conducted at 55° in a solution containing 0.15–0.75% of sulphuric acid and with an excess of 3% of oxalic acid. At this temperature the precipitation may be considered quantitative for yttrium, thorium, cerium, and didymium. Lanthanum oxalate, being the most soluble, is not completely precipitated, and 50 c.c. of the solution under the above conditions would retain 0.0011 gram of lanthanum in solution.

G. F. M.

The Use of Hanuš's Method for the Determination of the Iodine Figure of Mineral Oils. JAN HLADKÝ (*Chem. Listy*, 1922, 16, 207–211).—The method of Hanuš, whereby the iodine figure of mineral oils and bitumens is determined by means of a solution of iodine bromide in carbon tetrachloride, is examined. It is found that the reagent does not depreciate as rapidly as those used in the methods of Hübl and of Wijs (*E. Graefe, Petroleum*, 1905, 12, 631), and that the difference produced in the iodine

figure by increasing the time allowed for reaction is smaller than for the above methods. The concentration of the reagent, and the reaction temperature are not of great importance, whilst it is essential that the quantity of oil used is such as to leave a large excess of reagent after interaction. The presence of small quantities of water does not appear to affect the reaction. A further advantage of this method is that it permits the use of carbon tetrachloride, a universal solvent for the bitumens. The accuracy attainable is, for oils of a low iodine figure, from 1 to 2 units. R. T.

The Iodine Values of Aliphatic and Aromatic Unsaturated Compounds. D. HOLDE [with P. WERNER, IDA TACKE, and C. WILKE] (*Chem. Umschau*, 1922, 29, 185—188).—In the case of oleic acid anhydride and erucic acid, the iodine values found by the Hanuš method or by the Hübl-Waller method agree with the calculated values. Cholesterol, however, yields an iodine value of 73 to 77 by the Hübl method and phytosterol a value of 50 to 68, according to the time of contact (five to forty-eight hours); when estimated by the Wijs method, the iodine value of these two substances is identical, namely, 135. A similar difference in values obtained by the two methods has also been noticed in the case of certain mineral oils. W. P. S.

A New Method for the Detection of Methyl Alcohol. REINO HAMÄLÄINEN (*Acta Soc. Med. fennic. "Duodecim."*, 1921, 3, 1—6; *Ber. Ges. Physiol.*, 11, 262—263; from *Chem. Zentr.*, 1922, ii, 918).—Formaldehyde from the oxidation of methyl alcohol by potassium permanganate in ethyl alcohol-sulphuric acid solution is detected by addition of methyl violet-sulphurous acid decolorised by oxalic acid; a dark- to light-blue coloration is formed. G. W. R.

Estimation of the Alcohol Content of a Solution. A. LÉVÊQUE (*Bull. Sci. Pharmacol.*, 1921, 28, 549—554; from *Chem. Zentr.*, 1922, ii, 819).—The method of Rosset (A., 1921, ii, 598) is applied, using for miscibility temperature determinations, menthol, aniline sulphate, and phenol. G. W. R.

Estimation of Glycerol by the Dichromate Method. J. KELLNER (*Z. deut. Oel-Fett Ind.*, 1922, 42, 345; cf. this vol., ii, 399).—Complete oxidation of glycerol by dichromate is effected only when sulphuric acid, *d* 1.230, is employed. Sodium chloride, if present in the glycerol, should be removed by treatment with lead acetate and silver oxide, since chlorine is evolved even with sulphuric acid, *d* 1.1.

CHEMICAL ABSTRACTS.

Chemical Reaction of Phenol. ANGELO BELLUSSI (*Arch. Farm. speriment. Sci. aff.*, 1922, 33, 190—191).—When 2 c.c. of 2% phenol solution is gently heated on a water-bath with a small crystal of ammonium phosphate or arsenate and a few drops of 3% hydrogen peroxide solution, a pink or cherry-red coloration appears after a few seconds, the phenol evidently undergoing oxidation. The same coloration is obtained, only far more tardily, when the phenol solution is heated with hydrogen peroxide or

even alone, so that the action of the ammonium phosphate or arsenate is catalytic in nature. T. H. P.

Causse-Bonnan's Method for the Estimation of Dextrose. H. J. LEMKES and L. M. LANSBERG (*Pharm. Weekblad*, 1922, 59, 936—939).—It is confirmed that the results obtained vary considerably with the time occupied in titration. Attempts to get uniform results by substituting sodium for potassium compounds were unsuccessful. The white precipitate obtained at the end of the titration was found to contain Cu 23.05, Fe 11.2, K 21.55, (CN) 32.0, and water of crystallisation about 12.5%, but no empirical formula could be deduced. S. I. L.

Modification of Bang's Method for the Estimation of Sugar in the Blood. Z. ERNST and ST. WEISS (*Wiener klin. Wochschr.*, 1921, 34, 174—175).—The blood sample of 0.115 c.c. is aspirated by means of a calibrated capillary pipette which is attached to a 15 c.c. flask, and the pipette is rinsed into the flask by aspirating through it Bang's diluting salt solution. The stopper and pipette are removed, the flask is filled with the salt solution to the mark, the contents are mixed, filtered and 13 c.c. of the filtrate, corresponding with 0.1 c.c. of blood, are analysed by Bang's method (*A.*, 1918, ii, 278). Instead of the filtration, a centrifuge may be employed. CHEMICAL ABSTRACTS.

The Estimation of the Total Oxygen-combining Power of the Blood in the Van Slyke Apparatus. CHRISTEN LUNDGAARD and EGGERT MÖLLER (*J. Biol. Chem.*, 1922, 52, 377—378).—The blood is saturated with oxygen after, instead of before, introduction into the apparatus. The remainder of the estimation is then carried out by the modified method of Van Slyke and Stadie (this vol., ii, 78). E. S.

Qualitative and Quantitative Demonstration of Blood in Urine. A. JOHANNESSEN (*Ugesk. Læger*, 1921, 83, 1613; from *Physiol. Abstr.*, 1922, 7, 280).—The reagent recommended is prepared by adding 1 gram of phenolphthalein and 25 grams of potassium hydroxide to 100 c.c. of distilled water and then boiling with 10 grams of zinc dust until decolorised. After filtration, an equal volume of 96% alcohol is added. This solution keeps for six months. Immediately before use 1 c.c. of hydrogen peroxide is added to each 9 c.c. Equal volumes of the reagent and the suspected urine are mixed; if blood is present up to 1 in 20,000 or 1 in 30,000 a pinkish-red colour is formed. With blood in water, a positive reaction is obtained with dilutions as high as 1 in 500,000. Copper, even in traces, interferes with the test. E. S.

General and Physical Chemistry.

Influence of Pressure on the Refractive Index of some Organic Liquids. F. HIMSTEDT and IRMA WERTHEIMER (*Ann. Physik*, 1922, [iv], 67, 395—406).—The experimental arrangement of Röntgen and Zehnder was used, together with the Loewe-Zeiss interferometer. The refractive index and dn/dp were found for the following liquids, bromobenzene, chlorobenzene, nitrobenzene, chloroform, dichloroethylene, pinene, and hexane, and from these the compressibilities were calculated from the Newton, Beer, and Lorentz formulæ. The discussion of these values is postponed until the compressibilities of these liquids have been determined experimentally. W. E. G.

The Establishment of the International System of Normal Wave-lengths. FRIEDRICH MÜLLER (*Z. wiss. Photochem.*, 1922, 22, 1—20).—Wave-lengths of the carbon, iron, and nickel spectra were accurately measured and are tabulated. Methods of measurement and precautions taken are given. W. T.

Fine Structure of the Red Hydrogen Line. OTTO OLDENBURG (*Ann. Physik*, 1922, [iv], 67, 253—277).—Previous measurements of the Zeeman effect for H_α disagree among themselves and with the Sommerfeld quantum theory of fine structure. The divergences are shown to be mainly caused by errors arising in the photography of the spectrum from an echelon grating, and in the measurement of narrow diffuse bands. The true separation of H_α in the absence of a magnetic field, and at the temperature of liquid air, was obtained by resolution of the intensity curve into two Gauss curves. The value obtained was $0.140 \text{ \AA.} \pm 0.008$, whereas the measured distance between the two maxima was about 5% smaller. In weak fields (3500 gauss) the anomalous partial fusion of the p -components observed by Erochin was confirmed, but was shown to be due to a widening of the lines, without any appreciable alteration in the distance between the two components. At 18000 gauss, the two lines run together to form a single line which is narrower than the original doublet.

For weak fields, the Zeeman effect for the s -component is normal, but the widening of the lines at higher fields leads to the disappearance of the fine structure. Since the lines of H, He^+ , and Zn, which are most sensitive to the Stark effect, also become diffuse in a magnetic field, it is possible that an unexpected Stark effect was present. However, this is unlikely, since no appreciable change in potential along the capillary tube was observed under the influence of the magnetic field. Also the widening of the lines caused by the Stark effect disappears on the application of a magnetic field. The Zeeman effect for small magnetic fields is normal,

but the fusion of the fine structure components observed in the higher fields is in opposition to the Sommerfeld theory.

W. E. G.

The Balmer Series of Hydrogen. E. GEHRCKE and E. LAU (*Ann. Physik*, 1922, [iv], 67, 388—394).—Measurements of the fine structure of the Balmer series have been made at the temperature of liquid air, by means of a Lummer-Gehrcke interference spectroscope. The differences between the frequencies of the H_α , H_β , and H_γ doublets were, respectively, $\Delta\nu \cdot 10^{-9} = 8.69, 8.78, 8.74$. These values were independent of the time of exposure, so that no error due to the contraction of the gelatin of the photographic plate could have occurred. A considerable deviation occurs between these values and the theoretical fine structure constant, 10.95, of Sommerfeld. It is shown that this deviation from theory is not explained by the additional hypothesis of Sommerfeld, which states that each line of the doublet is composed of a number of sub-components. The experimental results are also in poor agreement with the theory of Silberstein. It is remarkable that the mean figure for $\Delta\nu$ is that derived by Lenz from the Abraham assumption of stationary electrons.

W. E. G.

Series and Other Regularities in the Spectrum of Manganese. MIGUEL A. CATALÁN (*Phil. Trans.*, 1922, 223A, 127—173).—The flame-arc, arc, and spark spectra of manganese have been photographed, using a Littrow spectrograph giving a linear dispersion of 16 Å. per mm. at λ 6300 and 5.5 Å. at λ 4700, a similar instrument with quartz prism giving a linear dispersion of 12 Å. per mm. at λ 4000, 4.5 Å. at λ 3000, and 2 Å. at λ 2300, and a concave grating ruled with 14438 lines per inch giving a normal dispersion of 5.5 Å. per mm. in the first order. It is shown that the flame, arc, and spark spectra of manganese, like those of other elements, are formed of two classes of lines, some belonging to the neutral atom and some to the ionised atom (Mn^+). In the spectrum of the neutral atom, triplet series and combination lines have been found. The series follows a Hicks formula with the Rydberg constant " N " and may be classified as follows: (a) A system of four triplets (sharp, diffuse, principal, and fundamental, the last two being rather doubtful), analogous to the ordinary triplet systems of series in the alkaline earths. (b) A system of three narrower triplet series (sharp, diffuse, and principal) which have strong analogies with the single line systems of series in the alkaline earths. (c) Another system of still narrower triplets (sharp, diffuse, and principal) which is parallel to the preceding system, the shift number being $C=10967.75$. The intercombination lines, $1S-1p_2$ and $1S-1p_3$ (λ 5394.677 and λ 5432.555), between the first two systems are found as two lines very prominent at low temperatures, and losing very much of their intensity at high temperatures. The ionisation and resonance potentials of manganese have been calculated from the term $1S$ and from the wave number of the line $1S+1p_2$. The values obtained are 7.4 volts and 2.3 volts, respectively. The lines of ionised manganese, enhanced lines, also

form triplets. The data do not permit of the series being traced, but there is evidence that they include a system of triplets (sharp, diffuse, and principal), and also another system of narrower triplets, corresponding with the single line systems in the alkaline earths, as in the case of the neutral atom. Each diffuse triplet, in neutral and ionised atom alike, is formed of nine lines, giving five d terms in contrast to the six lines giving three d terms which are found in the more familiar diffuse triplets of the alkaline earths. In the spectra of both the neutral and the ionised atom there are certain groups of prominent lines having the same character and showing similar variations with changes of temperature. The lines of each group are related by very exact numerical separations, and these lines are denoted by the term *multiplet*. It is pointed out that similar multiplets exist in the spectra of the neutral atoms of magnesium, calcium, barium, strontium, ionised aluminium, scandium, yttrium, and lanthanum, and neutral atoms of silicon and chromium, and that they may represent a form of regularity present in the spectra of many elements, and serve to indicate the most probable nature of the series in the spectra where the actual series have not been traced. The spectrum of manganese is discussed in connexion with the position of manganese in the periodic system. It seems most probable that the neutral atom of manganese has two electrons in the outermost ring, and that when this atom loses an electron, thus becoming ionised, another electron from the next ring comes out to take its place in the outermost ring, which again contains two electrons. This mechanism would account for the similarity of the neutral and ionised manganese atoms.

J. F. S.

Ultra-red Characteristic Frequencies of the Silicates.

CLEMENS SCHAEFER and MARTHA SCHUBERT (*Z. tech. Physik*, 1922, 3, 201—204; cf. this vol., ii, 179).—Since crystals containing CO_2 and similar groups show selective reflection which is characteristic of the group, and varies in a simple manner with the type of crystal, an analogous phenomenon was sought in the case of the silicates. The reflecting power of thirteen silicates was observed throughout the range 1 to 20μ , and twelve of these showed maxima near 10μ and 18μ . These maxima were complicated, varying greatly from one substance to another, and could not be correlated with the crystal structure in the simple manner noted in connexion with other groups such as CO_2 , ClO_2 , etc. One specimen, analcite, lacked the maximum of greater wave-length. Two specimens containing the group SiO_4 showed no particular similarity in their reflexion curves.

CHEMICAL ABSTRACTS.

The Regularities of Ultra-red Spectra and the Failure of the Classical Dispersion Theory. CLEMENS SCHAEFER (*Ann. Physik*, 1922, [iv], 67, 407—419).—A theoretical paper. In the absorption spectra of polyatomic gases, a simple relationship occurs between the various bands. In diatomic gases, the band with short wave-length is almost exactly the octave of the band with long wave-length. Similar octaves are present in the

absorption bands of solid substances. An explanation is put forward based on the assumption that the molecular vibrations are asymmetrical, which supports the quantum theory and is opposed to the classical theory of dispersion and absorption. W. E. G.

Absorption of Light by Sulphur at Various Temperatures. MITSU HARU FUKUDA (*Mem. Coll. Sci. Kyōto*, 1921, 4, 351—354).—The absorption spectrum of a layer of sulphur 0.3 mm. thick, prepared by heating the element slowly until melted, extends from the ultra-violet up to $\lambda 0.408 \mu$ at 0° , and extends a further 2μ towards the red end for every 10° rise of temperature up to 300° . No discontinuity, such as might be anticipated from the view of Kellas as to the constitution of the sulphur molecule (T., 1915, 113, 903), was observed in the absorption spectrum at about 160° . The absorption spectrum of plastic sulphur depends on the initial temperature to which the sulphur is heated during its preparation. The higher this temperature, the further does the absorption extend from the ultra-violet towards the red end. J. S. G. T.

Absorption of $\lambda 5460.97 \text{ \AA}$. by Luminous Mercury Vapour. J. C. McLENNAN, D. S. AINSLIE, and (MISS) F. M. CALE (*Proc. Roy. Soc.*, 1922, [A], 102, 33—45).—It has been shown that when the radiation constituting the green line of mercury is passed through moderately luminous mercury vapour the main component and the components No. +1, $\Delta\lambda = 0.0086 \text{ \AA}$. and No. -1, $\Delta\lambda = -0.0087 \text{ \AA}$. can be strongly absorbed. No marked absorption by luminous mercury vapour was observed in the case of the other nine components of the green line. Of the nine members constituting the magnetically resolved green line, it was found that the central undisplaced member was the only one that could be markedly absorbed by luminous vapour. It is also shown that absorption by luminous mercury vapour of the light constituting the green line in the mercury spectrum affords a means of clearly and easily demonstrating the existence of the components of the line with separations $\Delta\lambda = +0.0182 \text{ \AA}$. and $\Delta\lambda = -0.021 \text{ \AA}$., that is satellites Nos. +2 and -2. Some considerations are presented in support of the view that the components of the green line of mercury, for which $\Delta\lambda = +0.0182 \text{ \AA}$., $+0.0086 \text{ \AA}$., -0.0087 \AA ., -0.021 \AA ., and zero, may originate in atoms of the element having, respectively, the atomic weights 197, 198, 202, 204, and 200. J. F. S.

The Absorption of Mercury Vapour. J. FRANK and W. GROTRIAN (*Z. tech. Physik*, 1922, 3, 194—197).—To test the theory that the absorption bands of mercury vapour are due to diatomic molecules, the absorption of the band at 2540 \AA . was observed in superheated vapour. A sample saturated at $200^\circ/20 \text{ mm}$. was observed at temperatures up to 1200° . The strength of the band was found to decrease with increasing temperature. A second sample, saturated at $240^\circ/35 \text{ mm}$., showed greater absorption at low temperatures, but became nearly transparent at 1200° . This behaviour confirms the belief that the band is due to the molecule which is dissociated at high temperatures. From the rate of

variation of absorption with temperature, it is concluded that the work required for dissociation is of the order of one Cal. The number of molecules involved in the absorption is remarkably small, even at the lowest temperatures, where the partial pressure of molecules is less than 10^{-5} mm. CHEMICAL ABSTRACTS.

A Simple Method for the Measurement of Ultra-violet Absorption. CHR. WINTHER, BAGGESGAARD-RASMUSSEN, and E. SCHREINER (*Z. wiss. Photochem.*, 1922, 22, 33—46).—The photographic method of Merton (*T.*, 1913, 103, 124) for the quantitative measurement of absorption has been adopted for the ultra-violet. An improved fluorometer has been constructed for extinction measurements of certain wave-lengths and the accuracy of the instrument has been tested by thermo-electric controls. The quantitative photographic method was employed for the investigation of the nitrates of potassium and silver, the results differed from those obtained by qualitative methods. W. T.

The Emission of Light by Atoms. K. FÖRSTERLING (*Z. Physik*, 1922, 10, 387—392).—Previous experimental observations require that the light emitted by excited atoms is coupled with the atom in a continuous manner. This is contrary to the views of Sommerfeld, and in agreement with those of Mie. An external magnetic or electrical field does not appear to exert any effect on light free in the ether. Thus, if the emitted light is discontinuous, such fields should exert no effect on the radiated energy once it has left the electron. Those atoms in which the electrons have returned to an inner quantum orbit, after emitting the energy of excitation, should not show the Stark effect. The emitted light should be unchanged by the action of an electrical field and Stark, in his experiments, should have observed the unchanged line. For the *p*-components of H_α , H_γ , and H_δ and for the *s*-component of H_δ this was shown not to be the case. On this account, the assumption is made that the electron in its passage from the initial to the final orbit should give up its light continuously to the ether. If the assumption of the spherical wave, and the correspondence principle of Bohr, are still maintained, it would appear necessary to postulate some new energy form in the atom. W. E. G.

The Luminescence of Gas in the Electrodeless Discharge : The Induction Spectra of Cæsium and Rubidium. M. L. DUNOYER (*J. Phys. Radium*, 1922, 8, 261—292).—The vapours of cæsium and rubidium, when surrounded by a high frequency electromagnetic field, emit a brilliant luminescence at 100—300°. Rubidium vapour gives a feeble luminosity at the ordinary temperature, but no light can be detected from cæsium vapour below 90°. The colour of the luminescence of the saturated metallic vapour undergoes a sudden change at certain temperatures. It is uncertain to what extent this is due to an increase in density of the vapour. At the lowest temperatures, the induction spectra are very rich in lines, but the principal and secondary series are absent. At higher temperatures, all the cæsium and most of the

rubidium lines which occur in the spark spectra appear, and are present in great intensity. The secondary series are, however, completely absent, and only a few lines from the flame and arc spectra are to be found in the induction spectra of these metals. The ionised atoms are believed to be the source of the induction spectra, and not the emission of an electron by the Bohr mechanism. Such ions should resemble the atoms of the rare gases, and points of resemblance should occur between the induction spectra of the two groups of elements.

W. E. G.

The Luminescence of Decomposing Ozone. MAX TRAUTZ and WALTER SEIDEL (*Ann. Physik*, 1922, [iv], **67**, 527—572).—No chemiluminescence is observed when a carefully purified mixture of ozone and oxygen is heated to 400°. On the admission of coal gas to this mixture, light is emitted at the ordinary temperature, frequently with the formation of a mist. Purified air, carbon dioxide, and hydrogen do not give rise to this effect. Carbon monoxide when mixed with ozone produces chemiluminescence at temperatures above 150°, and a band of light is emitted between the wave-lengths 400 and 560 μ . This spectrum is independent of the thickness of the radiating gas.

The light emitted by mixtures of carbon monoxide and ozone is measured by means of an argon-filled rubidium cell. By a streaming method, it was shown that the amount of light is proportional to the product of the concentrations of the two gases; thus $\Delta\alpha = K_1[\text{CO}][\text{O}_3]/1 + K_2[\text{O}_3]/t$, where $\Delta\alpha$ is the electrometer deflexion, K_1 and K_2 are constants depending on the velocities of two simultaneous reactions, and t is the time of streaming for 50 c.c. of gas. These conclusions were confirmed by a study of the reaction in a stationary gas mixture. The velocity of emission of light obeys the second order law, and the deviations which were observed are probably due to irregularities in the temperature of the gas mixture. The velocity of emission increases with increase in temperature, but the total quantity of light emitted increases to a limiting value. The law of constant light summation propounded by Lenard for phosphorescent substances (*Heidelb. Akad.*, 1912, A, 5 Abh.) applies equally well to the chemiluminescence from the mixture of carbon monoxide and ozone, the quantity of light being independent of both the velocity and the temperature. The presence of indifferent gases such as nitrogen or carbon dioxide is without effect on the emission of light. The conclusion that the reaction is of the second order is supported by the observations of Angerer (A., 1921, ii, 257) on the decomposition of N_3 .

W. E. G.

Catalysis. XVI. Radiation as a Factor in Thermal and Photochemical Reactions. N. R. DHAR (*Z. anorg. Chem.*, 1922, **122**, 151—158).—The author has studied the velocity of certain chemical reactions in tropical sunshine and found that the light rays accelerated the reaction in all cases, more particularly so in the case of those reactions with a high temperature coefficient. The paper concludes with a brief review of the radiation hypothesis.

W. T.

The Effect of Colloids on Silver Bromide Emulsions. ROBERT SCHWARZ and HEINRICH STOCK (*Z. wiss. Photochem.*, 1922, **22**, 26—32).—The influence of catalysts on the photochemical decomposition of pure silver bromide has been investigated, the amount of decomposition being determined by the method previously described (A., 1921, ii, 614). Platinum sol delays the inversion point and solarisation. Silver and silicic acid sols act as positive catalysts, whilst aluminium hydroxide has a negative effect. Ferric hydroxide and stannic acid coagulated the emulsion. W. T.

The Photochemical Reduction of Titanium Salts. A. BENRATH and A. OBLADEN (*Z. wiss. Photochem.*, 1922, **22**, 65—71).—The sulphate and oxalate of titanium were not reduced by ethyl alcohol on long exposure to light, whereas the chloride reacted violently. For quantitative work, ethyl alcohol was an unsatisfactory reducing agent owing to the formation of hydrogen peroxide. Mandelic and lactic acids were found to be suitable, and measurements were carried out with the former, which was the more powerful reducing agent. The temperature coefficient of the reaction for 10° was found to be 2.21. The reaction obeyed the law of mass action, and was found to be bimolecular. Chlorides decreased the velocity, probably owing to the decrease of ionisation of the tetrachloride. W. T.

Elimination of Carbon Dioxide from Organic Compounds.
VI. The Action of the Light of the Electric Arc on Aqueous Oxalic Acid Solutions in Presence of Ferric, Chromic, and Mercuric Chloride, of Uranyl Acetate, and of Iodic Acid. HERMANN KUNZ-KRAUSE and PAUL MANICKE (*Ber. Deut. pharm. Ges.*, 1922, **32**, 209—216; cf. this vol., i, 6).—Many observations are recorded on the action of sunlight in decomposing aqueous solutions of oxalic acid, but none on the effect of artificial light. Experiments have been made on the decomposing effect of the light from an electric arc on oxalic acid in presence of a number of "catalysts." In presence of ferric chloride, 61.7% of the oxalic acid was decomposed into carbon dioxide in sixteen hours, the reaction being represented by the equation $\text{H}_2\text{C}_2\text{O}_4 + \text{Fe}_2\text{Cl}_6 = 2\text{FeCl}_2 + 2\text{HCl} + 2\text{CO}_2$. When chromic chloride was used, no decomposition occurred, but ammonium oxalate was oxidised in presence of mercuric chloride, the latter being reduced to mercurous chloride. The decomposition of oxalic acid was also promoted by uranyl acetate and by iodic acid, in the latter case with formation of free iodine. So far as qualitative experiments show, sunlight and the arc light are equivalent for the decomposition of oxalic acid in presence of the above substances. E. H. R.

Chemistry of Actinometers. A. BENRATH, E. HESS, and A. OBLADEN (*Z. wiss. Photochem.*, 1922, **22**, 47—64).—The mercuric chloride-ammonium oxalate actinometer was found to be trustworthy within narrow limits only. The ferric chloride and the oxalic acid-uranyl acetate actinometers were found to be in good agreement. Electrolytes retard the reduction of ferric chloride

and the decomposition of oxalic acid. The authors argue that the added electrolytes have no influence on the photochemical reaction, but decrease the dissociation of the iron salt and the oxalic acid, and they draw the conclusion that the ferric ion and the oxalate ion are photochemically active, whilst the undissociated molecules are inactive.

W. T.

The Photoelectric Effect of the Alkaline-earth Phosphors. KARL GÖGGELE (*Ann. Physik*, 1922, **67**, 301—319).—The relation between the photoelectric effect and the wave-length of the exciting light have been studied for CaBi_2 and SrBi_2 phosphors (cf. Lenard and Saeland, A., 1909, ii, 283). In order to maintain the phosphor in a condition of least excitation, it was illuminated simultaneously by the exciting radiation, and by either green or red light. On account of the low conductivity of the phosphor as compared with the metals, the photoelectric effect of these substances increases more slowly than the intensity of the exciting light. The normal photoelectric effect becomes appreciable at the violet end of the visible spectrum, and increases rapidly with decrease in wave-length. In addition, in the region of the wave-length of the phosphorescent light, selective effects are observed, which are especially large in the presence of green light. In the case of the CaBi_2 phosphor, the maxima for the photoelectric effect correspond exactly with the maxima of phosphorescence, but a shift of the photoelectric effect towards the red occurs with the SrBi_2 phosphor. The normal phosphors show a much greater photoelectric effect and phosphorescence than the decinormal phosphors. This is associated with the larger number of small phosphorescence centres in the normal phosphor. Phosphors when destroyed by pressure in the presence of light, become coloured, show practically no phosphorescence, and give the normal photoelectric effect. When destroyed in the dark, however, a much larger effect is obtained which decreases with time. It is suggested that this is caused by the destruction of the closed chain compounds forming the phosphorescence centres, into molecules containing free valency electrons. The magnitude of the photoelectric effect is proportional to the amount of the colouring of the phosphor.

The d - and u -emission of phosphorescent light is accompanied by the separation of electrons, whereas the m -emission, the fluorescence, is not accompanied by a photoelectric effect, and hence is not a consequence of electron emission. The electrons emitted from a phosphorescence centre do not necessarily return to the same centre or by the same path.

W. E. G.

β -Ray Spectra and their Connexion with γ -Radiation. LISE MEITNER (*Z. Physik*, 1922, **11**, 35—54).—By means of a more precise investigation, it is shown that the thorium- B line, which was previously and erroneously classified as a line belonging to the L_2 ring (this vol., ii, 416), in reality belongs to a second γ -radiation group, as was indicated by Ellis (A., 1921, ii, 422). Measurements of thorium- C and thorium- C' confirm those of von Baeyer, Hahn, and Meitner (A., 1911, ii, 567) as to the existence of a group of

β -rays of velocity about 95% that of light, which Ellis (*loc. cit.*) was unable to find, and, further, these measurements show that on using the exact energy of the K -disintegration of thallium some of these lines must be interpreted otherwise than was done by Ellis. Very exact experiments with radium- D which have been controlled by photometric registration show that the three L levels found by Ellis are not present in radium- D ; one only could be found and that is the L_2 level. It is concluded from the above-mentioned results and those of Ellis for radium- B that in the excitation of the characteristic radiation by γ -radiation proceeding from the nucleus of the atom itself practically only the L_2 level is affected, in contrast to the relationships in the excitation by Röntgen rays from outside the atom. An attempt is made to explain this result on the basis of the unsymmetrical atomic configuration postulated in Bohr's hypothesis. The criticism of the author's interpretation of the β -change put forward by Ellis is discussed, and it is shown that this interpretation is in keeping with the experimental facts.

J. F. S.

Wave-lengths of γ -Rays. LISE MEITNER (*Naturwiss.*, 1922, 10, 884; from *Chem. Zentr.*, 1922, i, 1352—1354; cf. this vol., ii, 416).—The lattice of crystals is too coarse for the determination of the wave-length of hard γ -rays. Primary β -rays from the disintegration of an atomic nucleus have in the simplest case uniform velocity, and energy corresponding with this velocity. The β -radiation may either pass out of the atom with unchanged velocity or else γ -rays may originate in the nucleus from the β -rays, having the same energy and the relation $E_\beta = h\nu_\gamma = hc/\lambda_\gamma$ (1) holds. The resultant γ -radiation detaches K -, L -, and M -electrons from the atom in which it originates and the energy of this secondary radiation is equal to the energy of the γ -radiation, less the energy used for disintegration or $h\nu_\gamma = E_\beta - E_k$ (2), where E_k is the energy used in detaching K electrons. E_β and E_k may be determined from the deviation of the β -rays in a magnetic field, and E_k from the K , L , and M absorption wave-lengths obtained by the lattice method. Applying equations (1) and (2), the author found the wave-length of the γ -rays from thorium- B to be 5.2×10^{-10} cm., those from radium 6.6×10^{-10} cm., and those from radium- D 2.9×10^{-9} cm. The shortest wave-length obtained by this method was that of the γ -rays from thorium- C'' , namely, 2.45×10^{-10} cm., approximately one-fifth of the wave-length of the shortest characteristic Röntgen rays. γ -Rays of even shorter wave-length may be expected to occur. The γ -rays originating from the nucleus may in some circumstances have a greater wave-length than the corresponding K -series. In the case of radium- D where the wave-length of the K series is 1.385×10^{-9} , the γ -rays cannot set free any K electrons.

G. W. R.

Excitation and Ionisation Potentials of Neon and Argon. G. HERTZ (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 25, 179—187).—The excitation potentials of argon and neon have been

determined by plotting the number of electrons which have a velocity below a definite small value (in the present experiments that below the velocity corresponding with 0.2 volt) against the applied tension. In this way, the excitation potentials are shown as sharp maxima in the curves. This method is unsuitable for measuring the ionisation potential, and a new method for measuring this quantity has been employed. The results show that, starting from the value 20.45 volts as the lowest excitation potential of helium, two pronounced excitation potentials are found at 17.35 and 19.15 volts for neon. The ionisation potential of neon is found to lie at 22.2 volts. In the case of argon, excitation potentials are found at 12.25 and 13.7 volts and the ionisation potential at 16.0 volts. It is shown in the case of neon that the values found are in harmony with the optical measurement of the spectrum, and, further, there is no room in the series spectrum for a resonance potential of 11.8 volts and ionisation potentials at 16.7 and 20 volts, as found by Horton and Davies (A., 1919, ii, 210), neither did the author find any indication of a resonance potential or ionisation potentials at the values stated. In the case of argon, a comparison with the series spectrum is not possible, but assuming that the argon spectrum is constituted similarly to the neon spectrum, then calculating from the observed excitation and ionisation potentials, the following mean values for the first groups of the terms are to be expected: $0.5s=130000\pm1000$, $1.5s\sim30400$; $2p\sim18600$, and higher terms <10500 .
J. F. S.

Chemical Reactions induced by the Corona Effect in Circuits traversed by Continuous Currents. C. MONTE-MARTINI (*Gazzetta*, 1922, 52, ii, 96—97).—The corona discharge is capable of effecting the following reactions: Formation of ozone from air; decomposition of carbon dioxide into carbon monoxide and oxygen; synthesis of ammonia from its elements; formation of carbon dioxide and hydrogen from carbon monoxide and water vapour; formation of hydrogen cyanide from acetylene and nitrogen; formation of formaldehyde from carbon dioxide and hydrogen; formation of benzoic acid from carbon dioxide and benzene vapour; decomposition of hydrogen sulphide into its elements and the inverse reaction; decomposition of hydrogen antimonide and its synthesis from its elements.
T. H. P.

Resistance Temperature Coefficient of Concentrated Solutions of Sodium in Liquid Ammonia. CHARLES A. KRAUS and WALTER W. LUCASSE (*J. Amer. Chem. Soc.*, 1922, 44, 1941—1949).—The resistance temperature coefficients of concentrated solutions of sodium in liquid ammonia have been measured from the saturation point up to $V=5.02$ and at temperatures from the boiling point of liquid ammonia down to -60° . Near the saturation point, the temperature coefficient of sodium in liquid ammonia has a value approximately 0.066%. With decreasing concentration, the temperature coefficient increases markedly, passing through a maximum value of approximately 3.6% at $V=1.1$ and then decreasing to a value of approximately 1.55 at dilutions above 4.0 litres

per atom. Taking into account the earlier work on the temperature coefficient of more dilute solutions, it is evident that the temperature coefficient passes through a minimum value. From the form of the curve at high concentrations, the temperature coefficient is seen to be approaching a negative value at high concentrations, although this value is not actually realisable, since the saturation point of the solution is reached. At the highest concentrations the solutions of the metals in liquid ammonia closely resemble metallic alloys. At lower temperatures, the temperature coefficient increases markedly at higher concentrations. As the concentration decreases, the influence of temperature on the coefficient decreases, and at a dilution $V=5.3$ the coefficient is relatively independent of temperature. From previous work on the influence of temperature on the temperature coefficient of more dilute solutions, it is clear that the influence changes sign, since at low concentrations the coefficient increases markedly with increasing temperature.

J. F. S.

Effect of Substitution on the Free Energy of Oxidation-reduction Reactions. I. Benzoquinone Derivatives. VICTOR K. LAMER and LILLIAN E. BAKER (*J. Amer. Chem. Soc.*, 1922, **44**, 1954—1964).—The oxidation reduction potentials of the substituted derivatives of quinone-quinol mixtures have been determined at 25°. The method consisted in adding the quinone to a measured volume of buffer solution, removing the oxygen by nitrogen, and then reducing a portion of the quinone to quinol with titanium chloride solution and measuring the potential of the mixture against a calomel electrode, using gold or platinum electrodes. Other measurements were made starting with the quinol and oxidising it with potassium dichromate. The concentration of hydrogen ions was also measured in each solution. The cells measured were therefore of the types: $\text{Hg}|\text{Hg}_2\text{Cl}_2, \text{KCl}(\text{sat})||\text{Q}+\text{QH}_2(yM)+\text{H}^+(xM)/\text{Pt}$; and $\text{Hg}|\text{Hg}_2\text{Cl}_2, \text{KCl}(\text{sat})||\text{H}^+(xM)|\text{H}_2(1\text{ atm.})/\text{Pt}$. The following values of the normal potentials and the reduction of free energy in the various cases are recorded: bromo-*p*-benzoquinone, 0.7151 volts, 138.0 kilojoules; chloro-*p*-benzoquinone, 0.7125, 137.5; *p*-benzoquinone, 0.6990, 134.9; toluquinone, 0.6454, 124.6; *p*-xyloquinone, 0.5900, 113.9; *p*-thymoquinone, 0.5875, 113.4; dimethoxy-*p*-benzoquinone, 0.5139, 99.2; naphthaquinone, 0.4698, 90.7. The effect produced by the substitution of the various groups is discussed in the light of the positive-negative rule of Michael. The validity of the method of work adopted has been established by (1) the constancy and reproducibility of the potentials under widely varying concentrations and following anodic or cathodic polarisation, (2) the use of different electrodes, (3) the agreement of the *E.M.F.* with those obtained by the method of pure mixtures, and (4) the approach of the equilibrium from both sides. The *E.M.F.* values are reproducible to 0.2 milli-volt.

J. F. S.

Stability of Phthalate Solutions as Standards in Hydrogen-ion Work. C. J. WOOD and P. P. MURDICK (*J. Amer. Chem. Soc.*, 1922, **44**, 2008—2009).—An answer to the criticisms of Oakes

and Salisbury (this vol., ii, 468), who have shown that phthalate solutions when measured against a hydrogen electrode a drift of the *E.M.F.* is observed (cf. also Merrill, this vol., i, 326). The authors have repeated their measurements with a 0.2*M.* solution of potassium hydrogen phthalate, which are shown to agree with those of Clark ("Determination of Hydrogen Ions," Baltimore, 1920). It is therefore contended that potassium hydrogen phthalate may be obtained in a sufficiently pure state to give constant potentials with the hydrogen electrode.

J. F. S.

An Electrolytic Current Intensification Effect, A New Electrolytic Displacement Effect, and the Connexion between Electrolysis and the Emission of Electrons in a Vacuum.

III. D. REICHINSTEIN and M. ROTHSCILD (*Z. physikal. Chem.*, 1922, 102, 209—234; cf. A., 1921, ii, 729, this vol., ii, 112).—A continuation of previous work. It is shown that the new displacement effect makes clear the mechanism of electrolytic detectors with a subsidiary source of current. It is shown experimentally that in such detectors the mechanism does not depend on a conversion into direct current, but that these detectors act as current intensifiers; the ratio of the direct current energy brought into existence by an alternating current to that of the alternating current itself may be greater than eighty-eight. An account is given of a new condenser commutator effect. A new electrolytic detector is described which does not make use of an external subsidiary current; this is characterised by the fact that in the absence of an alternating current it has an *E.M.F.*=0. An electrical current is capable of bringing about a reaction in an electrolyte without making any use of the migrations of the ions. When the new intensifier is connected with a source of alternating current, it intensifies the entering current one thousand times, consequently the intensifier and detector can be directly connected with the antennæ which collect high frequency undamped oscillations.

J. F. S.

The More Recent Conception of Electrolytes. IV. The Dissociation State of Acids of Medium Strength, especially Dichloroacetic Acid in Water and in Salt Solutions. ERLING SCHREINER (*Z. anorg. Chem.*, 1922, 122, 201—235).—Pure dichloroacetic acid was prepared and was found to be dimorphous with m. p. 12.15° and 13.25°, $d_{15}^{15}=1.5727$. The *E.M.F.* of this acid was measured in aqueous solutions as well as in 0.2 molar solutions of potassium chloride, sodium acetate, and magnesium sulphate. The quinhydrone electrode was used (Billmann, A., 1921, ii, 372). The conductivities of the acid and its sodium salts were also measured. Assuming the activity constant of the hydrogen ion to be 0.2, that of the dichloroacetate ion was found to be 0.37. Calculations based on the *E.M.F.* measurements and on the conductivity measurements were in complete agreement with the general law of mass action.

W. T.

Chromate Electrolysis with Diaphragms. A. LOTTERMOSER and KURT FALK (*Z. Elektrochem.*, 1922, 28, 366—376).—When

sodium chromate is electrolysed in a two or three compartment cell, the current yields of sodium dichromate and sodium hydroxide theoretically possible depend on the composition of the solutions originally present in the anode and cathode compartments. It was shown by Müller and Sauer (A., 1912, ii, 1037) that if the anolyte is a solution of sodium chromate and the catholyte one of sodium hydroxide, the yields are lower than they are when sodium chromate is used in both compartments. Their conclusions are now confirmed by extended observations on two compartment cells of the above types and on three compartment cells (a) with sodium hydroxide as the catholyte and sodium chromate in the other two compartments, and (b) with chromate in all three. The best yields of dichromate are obtained under the last conditions, as circumstances are then the least favourable for the migration of hydroxyl-ions into the anode compartment. The paper concludes with a mathematical discussion of the possible yield obtainable with these types of cell both with and without taking account of the influence of diffusion. [See *J. Soc. Chem. Ind.*, 1922, 857A.]

E. H. R.

Thermal Expansion of Concentrated Salt Solutions.

ROBERT MEYER (*Physikal. Ber.*, 3, 149; from *Chem. Zentr.*, 1922, i, 1214—1215).—The thermal expansions of aqueous solutions of sodium nitrate, potassium nitrate, zinc sulphate, and potassium carbonate, saturated at 20—70°, were measured at temperatures up to 90°. The coefficient of expansion of concentrated solutions increases with rise of temperature, but whilst in the case of water the rate of increase of the coefficient of expansion decreases smoothly with rise of temperature, the curves obtained for concentrated salt solutions are irregular and show points of inflexion the positions of which depend on the concentration of the solution. G. W. R.

Approximate Rule for the Specific Heats of Gases and Vapours.

RUDOLF MEWES (*Z. Sauerstoff Stickstoff Ind.*, 1922, 14, 26; from *Chem. Zentr.*, 1922, i, 1129—1135).—From the formulæ $cp_t = cp_0(1+\gamma)^t$ and $cp_t/cp_0 = (c'p_t/c'p_0)^n$ the specific heat of a substance can be calculated for all temperatures. Values are given of $\log(1+\gamma)$ and n , respectively, for a number of gases and vapours. The values of $\log(1+\gamma)$ are simple multiples of each other, whilst the values of n are simple integers. G. W. R.

The Specific Heats of Gases with Special Reference to Hydrogen.

CHRISTOPHER KELK INGOLD and EDITH HILDA USHERWOOD (T., 1922, 121, 2286—2291).

The Isotherms of Nitrogen, Oxygen, and Helium. L. HOLBORN and J. OTTO (*Z. Physik*, 1922, 10, 367—376).—The isothermals of oxygen and nitrogen at temperatures of 0°, 50°, and 100°, and at pressures up to 100 atm., and of helium from 50 to 100 atm. have been determined (cf. A., 1915, ii, 743; 1921, ii, 15). The coefficients are given for the equations $pv = A + Bp +$

Cp^2 , and $pv = (pv)_0(1 + B'/v + C'v^2)$; the pv values for helium may be expressed by a linear relationship.

	A.	B 10 ³ .	C 10 ⁶ .	B' 10 ³ .	C' 10 ⁶ .
Nitrogen...	0° : 1.00057	-0.58100	4.76327	-0.57070	4.58106
	50° : 1.18368	-0.02500	3.80816	-0.03281	4.79200
	100° : 1.36682	+0.36057	3.15102	+0.35016	4.98088
Oxygen ...	0° : 1.00130	-1.30143	0.46452	-1.27335	4.07229
	50° : 1.18456	-0.63543	0.27755	-0.62750	3.26394
	100° : 1.36782	-0.20357	0.22571	-0.20149	3.02964
Helium ...	0° : 0.99930	+0.69543	—	+0.69565	0.50594
	50° : 1.18223	+0.68887	—	0.68797	0.51689
	100° : 1.36518	+0.66804	—	0.67453	0.53306

Amagat's pv -values for 100 atm. are greater by 0.7% for nitrogen and 0.3% for oxygen, and the results of Kamerlingh Onnes and Hyndman for oxygen are, at the lower pressures, about 0.4% greater and, at the higher pressures, about the same amount smaller. The pv -values for air have been calculated for air from those for oxygen and nitrogen: the differences between these and the experimental values for air are within the error of the observations, except at 0° and 50° for the higher pressures. W. E. G.

Thermodynamic Treatment of the Possible Formation of Helium from Hydrogen. RICHARD C. TOLMAN (*J. Amer. Chem. Soc.*, 1922, 44, 1902—1908).—A theoretical paper in which it is shown that by combining the older ideas as to the possible formation of helium from hydrogen with the newer ideas as to the entropy of monatomic gases, it becomes possible to subject the reaction $4H = He$ to complete thermodynamical treatment. The equilibrium constant calculated from this treatment shows that hydrogen should combine practically completely to form helium at all temperatures below a million degrees and pressures above 10^{100} atmospheres. The conflict between this conclusion and the mutual presence of hydrogen and helium under solar, stellar, and terrestrial conditions has been discussed. Of the various possibilities presented for reconciling the thermodynamic conclusion as to equilibrium constant and the facts, the author regards the following as most hopeful; (1) the reaction to form helium from hydrogen is an extremely slow one even at 20000°, (2) hydrogen consists of a mixture of isotopes one of which has an atomic weight nearly unity, and (3) helium is decomposed in the sun and stars by the action of an extremely short wave-length radiation which is so penetrating as not to be in temperature equilibrium with the helium affected. J. F. S.

Correlation of Entropy and Probability. GEORGE A. LINHART (*J. Amer. Chem. Soc.*, 1922, 44, 1881—1886).—In a recent paper (this vol., ii, 257), the author has put forward two equations, the one $S = C\infty/K \cdot \log C\infty/(C\infty - C)$, connecting entropy and probability, and the other, $C = C\infty kT^K/(kT^K + 1)$, connecting specific heats and temperature. The validity of the former equation was substantiated for eighteen substances and the latter equation by a single example. In the present paper the second equation has been shown to hold for the specific heat of all substances which

have been measured by the thermo-electric methods. The results calculated for thirty-one substances agree with the experimental values in many cases better than the individual experiments agree with one another. Only in the case of ice and silver iodide is the agreement poor, and here it is shown that the specific heat data are quite erroneous. J. F. S.

The Relation of Melting Point to Boiling Point. R. LORENZ and W. HERZ (*Z. anorg. Chem.*, 1922, **122**, 51—60).—The ratio of the absolute melting point to the normal boiling point has been calculated for 412 elements and compounds (inorganic and organic).

No. of cases.	Ratio.	No. of cases.	Ratio.	No. of cases.	Ratio.
1	0.198	118 between	0.5—0.599	25 between	0.8—0.899
14 between	0.3—0.399	99	0.6—0.699	26	0.9—0.999
66	0.4—0.499	61	0.7—0.799	2 above	1.0

The mean of all the ratios (A) was found to be 0.62, or taking only those shown to occur more frequently (4, 5, 6, 7) the mean (B) was 0.5946. The following are the quotients for the different categories examined. Elements (A) 0.5583, (B) 0.4542. Inorganic compounds (A) 0.7183, (B) 0.6857. Organic compounds (A) 0.5839, (B) 0.5745. W. T.

Fused Salts. W. HERZ (*Z. Elektrochem.*, 1922, **28**, 390—391).—It has been shown that at the critical temperature the refractive index of non-associated liquids is a constant, 1.126 (A., 1921, ii, 529, this vol., ii, 97). From the measurements of Meyer and Heck of the molecular refractions of fused potassium and sodium nitrates (this vol., ii, 241), the critical refractive indices have been calculated and are, respectively, 1.136 and 1.138. It appears probable, therefore, that the requirements of the theory are fulfilled by fused salts. The critical densities, calculated from the law of corresponding states, are, respectively, 0.599 and 0.612. The boiling points of the alkali fluorides calculated by Lorenz and Herz from the surface tensions (this vol., ii, 261) are compared with the values found by von Wartenberg (this vol., ii, 146). The agreement is good except in the case of caesium fluoride. The critical pressures of the alkali fluorides, chlorides, bromides, and iodides are now calculated from the known relation between surface tension at the boiling point and critical pressure, $\gamma_s = 0.427p_k$. The values obtained are probably minimum values. They are used to calculate the constant of Cederberg's vapour pressure formula. When the differences between melting and boiling points of the alkali haloids are compared, regular variations are found as the atomic weight of the metal or halogen increases. E. H. R.

The Vapour Pressure of some Salts. III. H. VON WARTENBERG and O. BOSSE (*Z. Elektrochem.*, 1922, **28**, 384—386).—Continuing the work previously described on the vapour pressures of salts (this vol., ii, 156), the following new data have been obtained. Silver chloride, m. p. 449°, b. p. 1554°, $\log p = -44257/4.571T +$

5-298. Cuprous chloride, m. p. 430° , b. p. 1367° , $\log p = -19258/4.571T + 2.573$. Cuprous bromide, m. p. 483° , b. p. 1345° , $\log p = -19075/4.571T + 2.5794$. Cuprous iodide, m. p. 588° , b. p. 1293° , $\log p = -19257/4.571T + 2.690$. Lead chloride, m. p. 498° , b. p. 954° , $\log p = -28518/4.571T + 5.085$. Lead bromide, b. p. 916° , $\log p = -28180/4.571T + 5.183$. Lead fluoride, m. p. 818° , b. p. 1292° , $\log p = -39425/4.571T + 5.510$. Thallous chloride, m. p. 430° , b. p. 806° , $\log p = -25108/4.571T + 5.093$. Thallous bromide, m. p. 456° , b. p. 815° , $\log p = -25164/4.571T + 5.059$. Thallous iodide, m. p. 440° , b. p. 824° , $\log p = -25168/4.571T + 5.021$. Thallous fluoride, b. p. 298° , $\log p = -25152/4.571T + 9.639$. The thallous salts are abnormal in that the iodide has the highest boiling point and the fluoride the lowest. No vapour pressures of barium chloride, calcium chloride, calcium bromide, and boric anhydride could be measured at the highest attainable temperature, 1536 — 1546° . The opinion previously expressed that the fluorides form double vapour molecules is now withdrawn, an experimental error having been detected. The dissociation temperatures of the following acid fluorides were determined: lithium hydrogen fluoride, below 200° ; sodium hydrogen fluoride, about 270° ; potassium hydrogen fluoride, 310 — 440° ; caesium hydrogen fluoride, 500 — 600° ; rubidium hydrogen fluoride, 550 — 670° . E. H. R.

Calorimetric Determinations. P. E. VERKADE (*Chem. Weekblad*, 1922, 19, 389—393; cf. this vol., ii, 474).—The Third Conference of the International Union of Pure and Applied Chemistry, held at Lyon, June—July, 1922, recommended for the standardisation of calorimetric bombs the use of pure benzoic acid, prepared by the Bureau of Standards at Washington, and tested calorimetrically before issue; this special material is obtainable from the Bureau of the Institut International d'Etalons physico-chimiques at Brussels (The University, Rue des Sols). The other materials generally employed are reviewed, and the values given for the heats of combustion criticised. Pending exact determination of the mechanical equivalent of heat, which will permit of electrical measurements for standardisation, the Brussels benzoic acid should be used for all thermo-chemical determinations. The heat of combustion accepted by the Conference is 6324 cal. per gram weighed in air at 15° (6319, weighed in a vacuum). S. I. L.

Apparatus for the Rapid Determination of the Specific Gravity of Small Amounts of Liquid. ERICH WIEDBRAUK (*Z. anorg. Chem.*, 1922, 122, 167—170).—Water and the liquid in question are contained in two communicating U-tubes which also communicate with a manometer, the heights of the liquids which balance various pressures are measured, and hence the specific gravity is calculated. W. T.

Kinetic Theory of Adsorption. D. C. HENRY (*Phil. Mag.*, 1922, [vi], 44, 689—705).—A theoretical paper in which a kinetic theory of adsorption of gases has been developed by means of assumptions derived from Langmuir's conception of a unimole-

cular layer. Equations for the adsorption isotherm and isostere are deduced which are in keeping with experimental results. A method is suggested for applying these equations to adsorption from solution, and the results are compared with experimental data. A theoretical derivation of the well-known exponential formula is given which attributes a theoretical significance to the index $1/n$, which is in qualitative agreement with experimental data.
J. F. S.

Adsorption of Activated Sugar Charcoal. I. F. E. BARTELL and E. J. MILLER (*J. Amer. Chem. Soc.*, 1922, **44**, 1866—1880).—The adsorption of methylene-blue, a basic dye of the electrolyte type, by activated ash-free sugar charcoal has been investigated. The results show that this adsorption is partly, if not entirely, hydrolytic in character. Ash-free sugar charcoal will completely adsorb a certain maximum amount of methylene-blue from solution and still leave the solution neutral. During this process of adsorption a considerable quantity of hydrochloric acid is set free which is readily and completely adsorbed. Adsorption of quantities of methylene-blue greater than the above-mentioned maximum results in the liberation of hydrochloric acid which is not subsequently removed from solution by adsorption. The contradictory results obtained by previous investigators who have used blood charcoal as adsorbent have been reproduced with pure sugar charcoal and an explanation of these results is put forward.
J. F. S.

Adsorption and Catalysis in Fuller's Earth. ERIC KEIGHTLEY RIDEAL and WILLIAM THOMAS (T., 1922, **121**, 2119—2123).

Adsorption and Orientation of the Molecules of Dibasic Acids and their Etheral Salts in Liquid-Vapour Interfaces. H. H. KING and R. W. WAMPLER (*J. Amer. Chem. Soc.*, 1922, **44**, 1894—1902).—Surface tension measurements have been made at 20° by the drop-weight method of aqueous solutions of a series of concentrations of tartaric, malonic, succinic, fumaric, oxalic, malic, and maleic acids and the diethyl esters of the four first-named acids. The adsorption has been found for each solution by means of the Gibbs equation $u = -1/iRT \cdot \delta\gamma/\delta \log C$. A general explanation of the chemical theory of capillarity is given and this is shown to serve as an explanation of catalysis. Tables are drawn up giving the adsorption, surface tension, drop weight, and density of all the above-named solutions. It is shown that the adsorption of the dibasic acids is less than that of the monobasic acids; the degree of adsorption of the dibasic acids increases with increasing carbon content, but the increase is not regular for each carbon atom added to the chain. The presence of a hydroxyl group in the molecule decreases the amount adsorbed, and raises the surface tension, as shown by malic and *d*-tartaric acids. The esters are adsorbed more than the acids, due to the replacement of the polar carboxyl group by the group $-C(=O)-O-C_2H_5$. The molecules are oriented

in the surface with the polar groups in the liquid and the least active portions of the molecule forming the surface. J. F. S.

The Rate of Absorption of Carbon Dioxide by Ammoniacal Solutions. PAUL RIOU (*Compt. rend.*, 1922, 175, 472—474).—Investigation of the influence of temperature and of concentration on the rate of absorption of carbon dioxide by solutions of ammonium carbonate and hydrogen carbonate shows that at constant temperature it increases to a maximum with increase of concentration. In the presence of ammonium hydrogen carbonate, the absorption velocity is considerably less and a decrease takes place with higher concentration after the maximum has been reached. With increase of temperature, a maximum rate is also obtained; the presence of a larger proportion of hydrogen carbonate in the solution corresponds with a reduction in the temperature at which this maximum is reached. H. J. E.

Cataphoretic Charges of Collodion Particles and Anomalous Osmosis through Collodion Membranes Free from Protein. JACQUES LOEB (*J. Gen. Physiol.*, 1922, 5, 89—107).—If a collodion membrane separate a solution of a salt at a definite P_H from pure water, it is found that at $P_H 3$ the rate of diffusion of the water into the salt solution is normal, and is determined almost entirely by osmotic forces. If, however, the P_H be increased, "anomalous osmosis" is obtained. Measurements have been made of the potential difference between the salt solutions and pure water, and also of that between collodion particles and such salt solutions. It is found that the anomalous diffusion rates can be largely explained as due to the potential difference between the solutions on the two sides of the membrane, and that between the membrane and the liquid in its pores. The former potential difference appears to be due to the diffusion potential. W. O. K.

Interpretation of the Influence of Acid on the Osmotic Pressure of Protein Solutions. JACQUES LOEB (*J. Amer. Chem. Soc.*, 1922, 44, 1930—1941).—The effect of adding acid to casein solutions has been investigated with regard to the changes brought about in the osmotic pressure. The results show that the addition of a little acid to isoelectric casein solutions increases the osmotic pressure until a maximum is reached, after which the addition of more acid depresses the osmotic pressure. This is explained on the basis of the dispersion hypothesis by the assumption that the addition of a little acid increases the degree of dispersion and consequently the osmotic pressure of the solution, whilst the addition of more acid diminishes the degree of dispersion and consequently the osmotic pressure. The present observations show that when osmotic equilibrium is established between a solution of casein chloride enclosed in a collodion bag and an outside aqueous solution free from protein, the hydrogen-ion concentration is always greater in the outer solution than in the casein solution. There exists a membrane potential between the casein chloride solution enclosed in a collodion bag and the surrounding

aqueous solution free from protein with which the casein solution is in osmotic equilibrium, and this membrane potential can be measured with indifferent calomel electrodes and a Compton electrometer. When this is done at 24° , it is found that the number of millivolts of the observed membrane potential is equal to 59 (P_H inside $-P_H$ outside), the latter values being measured with the hydrogen electrode. This is the result to be expected if the inequality of the hydrogen-ion concentrations inside and outside the casein chloride solution at equilibrium is determined by Donnan's equation for membrane equilibria (A., 1911, ii, 848). The fact that the hydrogen-ion concentration inside and outside the casein chloride solution is not the same as that of the outside free from casein chloride which is in osmotic equilibrium shows that the observed osmotic pressure of the protein chloride solution cannot be entirely due to the protein, but must be partly due to the difference in the concentration of the crystalloid ions, hydrogen and chloride, in the inside and outside solution. It is therefore necessary to correct the observed osmotic pressure of a protein chloride solution for this difference of concentration of the hydrogen and chloride ions on the opposite sides of the membrane on the basis of the Donnan equilibrium. The method of evaluating this correction is demonstrated. When this evaluation is made, it is found that within the limit of accuracy of the observations and calculations the entire effect of the hydrogen-ion concentration on the osmotic pressure of the casein chloride solution is covered by the correction required and that there is little if anything left for the dispersion hypothesis to explain. This is in keeping with the conclusion previously drawn by the author, that the influence of electrolytes on the osmotic pressure of protein solutions is entirely, or almost entirely, the consequence of the difference in the concentration of crystalloid ions inside the protein solutions and the outside aqueous solutions at equilibrium, this difference being caused by the establishment of a membrane equilibrium.

J. F. S.

Selective Solvent Action by the Constituents of Aqueous Alcohol. ROBERT WRIGHT (T., 1922, 121, 2251—2256).

Variations in the Solubility of a Compound in Presence of Other Compounds. I. Water-Phenol-Diphenols. P. LEONE and E. ANGELESCU (*Gazzetta*, 1922, 52, ii, 61—74).—The critical solubility temperature for the system water-phenol is raised by addition of naphthalene, benzophenone, calcium chloride, benzoic acid, tartaric acid, etc., lowered by addition of succinic acid, sodium oleate, or mercuric cyanide, and first raised and then lowered by increasing proportions of acetone. The addition to this system of resorcinol, quinol, or catechol produces marked increase in the reciprocal solubility and a considerable lowering of the critical temperature. Thus, the compositions of the two layers formed at any temperature in a water-phenol mixture gradually approach one another on addition of a dihydroxybenzene and, for a certain concentration of the latter, become equal. The critical temperature of the system is 66.6° ; Rothmund (A., 1898,

ii, 503) found a rather higher temperature. The results obtained are given both graphically and in tabular form. T. H. P.

Variations in the Solubility of a Compound in Presence of Other Compounds. II. Water-Epichlorohydrin-Acetic Acid. P. LEONE and M. BENELLI (*Gazzetta*, 1922, 52, ii, 75—86; cf. preceding abstract).—The reciprocal solubilities of water and epichlorohydrin increase as the temperature is raised, but the critical solution temperature cannot be ascertained since chemical reaction occurs at temperatures above 80°. The authors have measured the reciprocal solubilities at different temperatures and also the depressions of the temperature of saturation by addition of varying proportions of acetic acid, which is miscible in all proportions with both water and epichlorohydrin. T. H. P.

Electronic Structures of Crystals. I. MAURICE L. HUGGINS (*J. Amer. Chem. Soc.*, 1922, 44, 1841—1850).—A theoretical paper in which the author shows how the Lewis hypothesis of atomic and molecular structures (A., 1916, ii, 310) may be applied to ascertain the relative position of electrons and electronic groups in crystals for which the atomic arrangements are known from X-ray investigation. It is shown that the valency electrons, in the crystals discussed, tend to form complete shells round each atomic kernel, each shell consisting of four or six electron pairs. Except for the elements of the first series of the periodic table, these pairs tend to be opposite to the faces of the kernel polyhedra. The atoms are held together by single bonds, each of which is a valency pair in the shell of two atoms and on or near the centre line between them, or by double bonds in each of which there are two electron pairs common to two atoms; the latter case obtains between the carbon and oxygen, and the nitrogen and oxygen in carbonates and nitrates, respectively. There are isolated pairs, that is pairs not acting as bonds in arsenic, antimony, bismuth, and quartz. From the point of view of the foregoing statements, the following crystals have been discussed with regard to their electronic structure: diamond, silicon, germanium, grey tin, zinc sulphide (cubic and hexagonal), cuprous chloride, bromide and iodide, silver iodide (cubic and hexagonal), zinc oxide, cadmium sulphide, carborundum, chalcopyrite, marcasite, hauerite, iron-nickel sulphide, arsenoferrite, loellingite, smaltite, safflorite, cloanthite, rammelsbergite, cobaltite, willyamite, gersdorffite, wolfchite, ullmanite, kallilite, arsenopyrite, glaucodotite, arsenic, antimony, bismuth, quartz, calcite, aragonite, carbonates of calcium, magnesium, iron (ferrous), manganese, zinc, strontium, barium, and lead, and several mixed carbonates of the above-named metals, and sodium and potassium nitrate. J. F. S.

The Effect of Scratching the Wall of a Vessel with a Glass Rod. L. DEDE (*Z. Elektrochem.*, 1922, 28, 364—365). ROBERT FRICKE (*ibid.*, 365—366).—The first author criticises Fricke's explanation (*ibid.*, 244) of the manner in which crystallisation of a solution or liquid is started when the wall of the containing vessel

is rubbed with a glass rod, namely, that detached particles of glass act as nuclei. The view is put forward that the phenomenon is electrostatic in character, and experiments with ammonium magnesium phosphate are described supporting this view. Fricke replies, criticising Dede's experiments and maintaining his opinion.

E. H. R.

Crystallisation and Proteins. ÉMILE CAVAZZANI (*Arch. internat. physiol.*, 1921, 18, 446—450; from *Chem. Zentr.*, 1922, i, 1214).—The crystallisation of ammonium chloride from saturated solutions is prevented by 0.05% of egg-albumin, or by 0.2—0.5% of gum arabic.

G. W. R.

An Electron Theory of Solids. (SIR) J. J. THOMSON (*J. Franklin Inst.*, 1922, 194, 281—289).—A theoretical paper giving a summary of the author's theory of the solid state (cf. this vol., ii, 252, 355). The potential energy of atoms in the crystals of the elements and the bulk moduli have been calculated, on the assumption that the attractive force between the atoms and nuclei varies as the inverse cube of the distance. The values for the uni-, bi-, and ter-valent elements are in absolute agreement with the experimental values of Richards. The excess of energy in the surface, the surface tension, has been deduced, and it is shown that this is proportional to $K^{\frac{1}{3}}$, where K is the bulk modulus, a relation due to Richards. The calculation of the potential energy of the atoms in chemical compounds is complicated by the unequal distributions of the electrons about the atoms, but the numbers obtained for the bulk moduli are a useful guide to the distribution of the electrons in the crystal.

W. E. G.

Hardness of Solid Substances and its Relationship to Chemical Constitution. A. REIS and L. ZIMMERMANN (*Z. physikal. Chem.*, 1922, 102, 298—358).—The method of determining hardness by Marten's sclerometric method has been investigated in connexion with a large number of substances of different types, including very soft substances which exhibit a scratch of 10μ in width when a load of only 1 gram is applied to the needle and shown to be trustworthy for this type of investigation. Experiments on the scratching of ninety crystals by one another yield results which on the whole are in agreement with those obtained by the foregoing method. The eroding action of a moving liquid on solid surfaces has been investigated and by using a stream of mercury from narrow nozzles it has been found possible to obtain trustworthy results on the resistance to erosion of solid substances. It is shown that the hardness and the resistance to erosion for the larger number of the substances examined (about one hundred and eighty) are parallel, but that the numerical value of the resistance to erosion, measured as time required for the removal of the same amount of material, increases much more than the hardness. The hardness depends on the particular crystal face employed, and on the direction of the scratch, whereas no such dependence could be observed in the erosion experiments. The relationship

between the hardness, crystal structure, and chemical constitution has been investigated for a large number of substances of all types. A general connexion is shown to exist on the basis of the classification of crystal lattices according to the strength and spacial partition of the forces of attraction between the atoms. In the case of those salts with lattices which are maintained by the electrostatic attractions of the ions, it is shown that generally the charges and distances between the ions are determinative of the hardness. In the case of compounds which have not the nature of salts, the hardness runs parallel with the strength of the residual affinities which the molecules exert outwardly. In keeping with the general chemical experience and with the tendency to association, it is shown from the hardness measurements that the residual affinity is smallest in the case of the hydrocarbons and increases most strongly under the influence of hydroxyl, carboxyl, and cyanogen groups. In the case of mixed crystals, it is shown that the mixture is harder than the mean of the components and in some cases harder than the harder component. The hardness of compressed pastilles has been compared with that of crystals of the same substances, and the two values are shown to be very nearly equal. A number of methods of producing large crystals have been investigated, and it is shown that the tendency of substances to form large crystals is to a certain extent a function of the chemical constitution and is often parallel with the hardness. J. F. S.

The Increase in the Size of Grain of Metallic Bodies (not previously Worked in the Cold) Formed from Powdered Material by Pressure or Sintering. F. SAUERWALD (*Z. anorg. Chem.*, 1922, 122, 277—294).—The application of pressure did not increase the size of the grain; nor does this occur at moderate temperatures, but at higher temperatures an increase in granular size was observed in the case of iron, nickel, cobalt, tungsten, and copper. This limiting temperature could be fixed within 100°, and an increase of temperature beyond this limit resulted in a rapid increase in the size of the grains. This limiting temperature is independent of the pressure applied in compressing the powder, and has a characteristic value for each metal. Increase in the size of grains was not observed in the cases of zinc and aluminium. These results are discussed in the light of Tammann's new theory of crystallisation (*A.*, 1921, ii, 172). W. T.

The Influence of Electrolytes on the Cataphoretic Charge of Colloidal Particles and the Stability of their Suspensions. I. Experiments with Collodion Particles. JACQUES LOEB (*J. Gen. Physiol.*, 1922, 5, 109—126).—The potential difference between collodion particles and the aqueous solution in which they are suspended has been determined by measuring the mobility of the particles under a given potential gradient. It is found that the influence of acid and of alkali and of most salts is to cause the negative charge on the collodion particle to increase to a maximum and then to decrease with increasing concentration of the solution.

A tervalent kation, however (for example, in the case of lanthanum chloride), may cause a steady decrease in the charge and even reverse it. The effects of hydrochloric acid, sodium hydroxide, lithium chloride, sodium chloride, and potassium chloride are very similar. If the P_H of the solution be varied, the variation of the charge with changing concentration of salt in the solution remains similar.

When the potential difference between the particles and the solution is reduced below a certain critical value, about 16 millivolts, the suspension of colloidal particles becomes unstable, and precipitation occurs.

W. O. K.

Elasticity of Purified Gelatin Jellies as a Function of the Hydrogen-ion Concentration. S. E. SHEPPARD, S. S. SWEET, and ANBER J. BENEDICT (*J. Amer. Chem. Soc.*, 1922, **44**, 1857—1866).—The rigidity and hydrogen-ion concentration of demineralised gelatin jellies, which are free from products of hydrolysis, has been determined for 4, 5, 7, and 10% jellies. The effect on the rigidity of the addition of 0.001% to 0.10% of alum to the above-named jellies has also been investigated. It is shown that except at very high ($P_H > 2$) and very low ($P_H < 11$) hydrogen-ion concentrations, the rigidity does not vary rapidly with changes of the hydrogen-ion concentration, the maximum being at $P_H = 7-9$. The addition of quantities of aluminium equivalent to 0.01—0.1% of alumina based on the dry gelatin produced considerable changes in the rigidity- P_H curve, and markedly increased the rigidity. These results are discussed in connexion with the theory of the viscosity and elasticity of gelatin. A misleading statement on Poisson's ratio for gelatin jellies in a previous paper (*A.*, 1921, ii, 311), and also in the literature, has been corrected.

J. F. S.

The Ultra-filtration of Soap Solutions: Sodium Oleate and Potassium Laurate. JAMES WILLIAM MCBAIN and WILLIAM JOB JENKINS (*T.*, 1922, **121**, 2325—2344).

The Reversibility of the Reaction between Nitrogen, Carbon, and Sodium Carbonate. CHRISTOPHER KELK INGOLD and DANIEL WILSON (*T.*, 1922, **121**, 2278—2286).

The Ammonium Carbonate-Carbamate Equilibrium. CARL FAURHOLT (*Z. anorg. Chem.*, 1922, **121**, 132—134).—A reply to Wegscheider's theoretical paper (this vol., ii, 376). The author shows that his results for the above equilibrium are in fair agreement with those obtained by Burrows and Lewis and by Fenton. The author also shows that the conclusions of Wegscheider (*A.*, 1916, ii, 617) are incorrect, because the latter found that carbamic acid was very weak and the ammonium salt completely hydrolysed in aqueous solution. The present author carried out colorimetric hydrogen-ion estimations and found that the ammonium salt is not hydrolysed to any appreciable extent.

W. T.


Equilibrium in the System Potassium Chlorate, Potassium Nitrate, and Water at 25°. SHIGERU TODA (*J. Chem. Soc. Japan*, 1922, **43**, 320—328; *Mem. Coll. Sci. Kyoto Imp. Univ.*, 1921, **4**, 377—382).—The equilibrium in the system potassium chlorate, potassium nitrate, and water at 25° has been studied. The solubility of potassium chlorate at 25° is 7.745 grams in 100 grams of solution or 8.395 grams in 100 grams of water, and that of potassium nitrate is 27.24 grams in 100 grams of solution or 37.44 grams in 100 grams of water. At 25°, potassium chlorate forms a solid solution with potassium nitrate taking up the latter to the extent of 14.37 molar %, whilst the nitrate takes up none of the chlorate. K. K.

Equilibrium in the System Copper Sulphate, Ammonium Sulphate, Potassium Sulphate, and Water at 25.0°. RYONOSUKE HAYAMI (*Mem. Coll. Sci. Kyoto*, 1921, **4**, 359—369; *J. Chem. Soc. Japan*, 1921, **42**, 421—441).—The equilibrium diagram for the system copper sulphate, potassium sulphate, ammonium sulphate, and water at 25° has been determined, and also the relation between the composition of the aqueous solutions and that of the solid solutions when the residue is either a mixture of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ and $\text{CuSO}_4 \cdot (\text{NH}_4, \text{K})_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$ or a mixture of $(\text{NH}_4, \text{K})_2\text{SO}_4$ and $\text{CuSO}_4 \cdot (\text{NH}_4, \text{K})_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$. J. S. G. T.

Dynamics of the Formation of Nitriles from Acid-anhydrides and Amides. II. Kinetic Study of the Reaction between Acetamide and Acetic Anhydride, using Phase Rule Methods. ROBERT KREMANN, ALFRED ZOFF, and VICTOR OSWALD (*Monatsh.*, 1922, **43**, 139—144).—The reaction: acetamide + acetic anhydride \rightleftharpoons acetonitrile + 2 acetic acid, is followed kinetically, and shown to reach an equilibrium corresponding with 83% conversion (from left to right in the sense of the above scheme) at 98°, and with 87% conversion at 78°.

The method consisted in taking the freezing points of mixtures of the components which had been heated at the required temperature for different lengths of time. The percentage compositions were then read from an experimentally determined curve of the freezing points of mixtures of the two systems involved in the reversible reaction. C. K. I.

Dynamics of the Formation of Nitriles from Acid-anhydrides and Amides. III. Reaction between Benzamide and Acetic Anhydride. ROBERT KREMANN, WILHELM RÖSLEB, and WILHELM PENKNER (*Monatsh.*, 1922, **43**, 145—159).—Two side reactions take place when equimolecular quantities of acetic anhydride and benzamide are heated together :



and in order to follow the process kinetically by the method previously employed (cf. preceding abstract) it was necessary

first to construct a ternary freezing-point diagram for the systems $[C_6H_5 \cdot CO \cdot NH_2 + (CH_3 \cdot CO)_2O]$, $[C_6H_5 \cdot CN + 2CH_3 \cdot CO_2H]$, and $[C_6H_5 \cdot CO \cdot NH \cdot CO \cdot CH_3 + CH_3 \cdot CO_2H]$. With the aid of this diagram the progress of the reactions is followed by a series of freezing-point determinations. The following values are deduced for the composition of the equilibrium mixture: benzamide 0.48, acetic anhydride 0.48, benzonitrile 0.14, acetylbenzoylimide 0.38, acetic acid 0.66 mol.

C. K. I.

Studies in Acidolysis. I. The Equilibrium between Acetic Acid, Trichloroacetic Acid, and their Ethyl Esters. II. The Equilibrium between Acetic Acid, Trichloroacetic Acid, and their Methyl Esters. J. J. SUDBOROUGH and D. D. KARVÉ (*J. Indian Inst. Sci.*, 1922, 5, 1—21).—As a preliminary to the study of the reaction between ethyl alcohol and a mixture of acetic and trichloroacetic acids, the equilibrium between the two acids and their ethyl esters was examined at 30°. The equilibrium constant of the reaction $CH_3 \cdot CO_2Et + CCl_3 \cdot CO_2H \rightleftharpoons CH_3 \cdot CO_2H + CCl_3 \cdot CO_2Et$ was found to be equal to unity, and the same result was obtained with the methyl esters. It follows that the relative affinities of the methoxy- and ethoxy-residues for acetyl and trichloroacetyl are practically identical, in spite of the difference between the dissociation constants of the two acids. In the absence of a catalyst the reaction $CH_3 \cdot CO_2Et + CCl_3 \cdot CO_2H \rightarrow CH_3 \cdot CO_2H + CCl_3 \cdot CO_2Et$ and the reverse reaction proceed very slowly at 30°, the time required for the attainment of equilibrium being 1200—1400 hours. The velocity constants in both directions are equal, the value of k/V being 0.00039, where $k/V = 1/2A_1t \log A_1/A_1 - 2x$. In this expression, V_1 = volume of 1000 grams of the mixture, A_1 = the number of gram-mols. per 1000 grams, and x the number of molecules which have reacted after t hours. With the methyl esters, the reaction is still slower, and $k/V = 0.000097$.

E. H. R.

The Ignition of Gases. II. Ignition by a Heated Surface. Mixtures of Methane and Air. WALTER MASON and RICHARD VERNON WHEELER (*T.*, 1922, 121, 2079—2091).

The Explosions of Varieties of Organic Dust. Experimental Investigation of the Simple Case of Sugar Dust. P. BEYERSDORFER (*Ber.*, 1922, 55, [B], 2568—2596).—The experiments consist in a thermal and an electrical investigation of the causes leading to the exploding of mixtures of very finely divided sucrose and air.

The explosion chamber consists of a glass vessel of about 600 c.c. capacity which is suspended symmetrically in a copper pot surrounded by an asbestos mantle and heated from below by a Franke burner. The vessel is provided with inlet and outlet tubes for gases and at its centre a cage of thin aluminium foil is suspended against which the sugar dust is blown through a horizontal side tube. The temperature is measured by a silver-constantan thermo-element. The sugar dust is passed through a sieve having

ten thousand meshes to the square centimetre, dried at $105^{\circ}/100$ mm., and preserved over concentrated sulphuric acid. The apparatus is heated to slightly above the desired temperature, and then allowed to cool to the desired extent, when the dust is blown against the foil. In this manner, the latter is the hottest part of the apparatus at the moment of the experiment.

The author defines the following terms. Ignition is a process which leads to visible combustion. Its cause may be of a dynamic, thermal, or an electrical nature. The ignition may lead to inflammation or explosion. Inflammation is a visible process of combustion which takes place throughout the material with small velocity and little change of pressure and has the power of causing further thermal ignition. Explosion is a visible process of combustion which is propagated through the material with high velocity and great change of pressure, and has the ability to ignite other combustible material by thermal action or by shock.

The author draws the following conclusions. The explosion of dusts can be caused by thermal or electrical processes. The temperature of ignition of sugar dust in air is $410 \pm 1^{\circ}$, and in oxygen 371° . The presence of small quantities of ozone lowers the temperature of ignition by 3° in air and by 4° in oxygen. The temperature of ignition is dependent on the oxygen content of the gas and on the nature and thermal conductivity of the relatively inert gases which accompany the oxygen, carbon dioxide having a noticeably greater extinguishing action than nitrogen. The dependence of the temperature of ignition of sugar dust on the oxygen content of mixtures of the latter gas with nitrogen is represented graphically. The curve approaches asymptotically to the temperature axis. At a point corresponding with the presence of about 9% of oxygen, the region of ignition becomes divided into an inflammation and an explosion region. Ultimately, the thermal explosion of sugar dust is a gaseous explosion. The process occurs in two phases: (i) the gasification of sugar dust such as takes place in the case of coal, and (ii) the oxidation of the "sugar gas" until ignition sets in. [In experiments on the temperature of ignition of sugar dust in air in which such ignition just does not occur, a rapid initial drop of 10 — 15° in temperature is invariably observed (due to gasification of the sugar) followed by a rise in temperature which generally exceeds the initial temperature ultimately (due to oxidation of the gas).] The upper limit of explosion is established at 13.5 kilos. of dust per cubic metre; the lower safety limit is calculated to be 17.5 grams per cubic metre.

Sugar dust becomes positively charged when projected through the air. If it is whirled through the air, tensions of multiples of ten thousand volts may be developed. Discharge of the loaded particles in a gas may occur silently, but they have no immediate connexion with the luminosity which accompanies the process and is readily visible in the dark. This luminosity is specific for each gas, and is identical with that observed for it in an alternating field.

Sugar dust may be caused to explode in an alternating field. The phases of the electrical explosion are: (1) the formation of

ozone and nitrogen pentoxide, (2) the reaction of these substances with the most minute sugar particles, and (3) explosion as a result of this reaction in the alternating field. Sugar dust forms an aerosol with air; if this suddenly approaches the isoelectric point, a flash occurs which ignites the dust cloud.

The conclusions elucidated for sugar dust are applicable to every explosive dust whether organic or inorganic.

Magnetic phenomena have no influence on the occurrence of dust explosions.


The violence of a dust explosion is attributed to the adsorption of oxygen by the dust.

Triboluminescence cannot be regarded as a cause of sugar dust explosions. H. W.

Michael's Addition and Separation Rule. L. SMITH (*Z. physikal. Chem.*, 1922, **102**, 359—363).—A theoretical paper in which on the basis of the author's investigations on the velocity of the reactions $\text{OH}\cdot\text{CH}_2\cdot\text{CH}(\text{OH})\cdot\text{CH}_2\text{Cl} + \text{OH}' \rightleftharpoons \text{OH}\cdot\text{CH}_2\cdot\text{CH}\cdot\text{CH}_2 + \text{Cl}' + \text{H}_2\text{O}$, and $\text{OH}\cdot\text{CH}_2\cdot\text{CHCl}\cdot\text{CH}_2\cdot\text{OH} + \text{OH}' \rightleftharpoons \text{OH}\cdot\text{CH}_2\cdot\text{CH}\cdot\text{CH}_2 + \text{Cl}' + \text{H}_2\text{O}$ (A., 1920, i, 658), an attempt is made to explain Michael's addition and separation rule. This rule may be stated as follows :

When two isomeric compounds may be formed from an unsaturated compound, a direct relationship exists between the relative quantities of the additive compounds produced and the velocities with which they may be reconverted into the original substance. J. F. S.


The Equilibrium Isomerism between Bornyl Chloride, *iso*Bornyl Chloride, and Camphene Hydrochloride. HANS MEERWEIN and KONRAD VAN EMSTER [with JACOB JOUSSEN] (*Ber.*, 1922, **55**, [B], 2500—2528; cf. A., 1920, i, 855).—In the molten condition and in solution, an equilibrium is established between bornyl chloride (I), *isobornyl* chloride (II), and camphene hydrochloride (III), in accordance with the scheme :



The equilibrium is dependent on the temperature and to a slight extent on the solvent. At the atmospheric temperature, it is greatly displaced in the direction of bornyl chloride. The transformation from right to left occurs exothermally; bornyl chloride is therefore the most stable, camphene hydrochloride the most labile isomeride. With increasing temperature, the equilibrium becomes more and more displaced in favour of *isobornyl* chloride and camphene hydrochloride. The interconversion of *isobornyl* chloride and camphene hydrochloride takes place readily at the atmospheric temperature, whereas the transformation of bornyl

chloride into *isobornyl* chloride occurs with greater difficulty and only at a more elevated temperature.

The velocity of the conversion of camphene hydrochloride into *isobornyl* chloride at 20° or 40° has been examined in the following solvents: cresol, sulphur dioxide, nitromethane, acetonitrile, nitrobenzene, benzonitrile, anisole, bromobenzene, ethyl bromide, chlorobenzene, benzene, light petroleum, and ether. The rate of change is enormously influenced by the solvent. In cresol, it occurs so rapidly that it is complete almost as soon as the camphene hydrochloride has passed into solution; ether, at the other end of the scale, acts practically as a stabilising agent towards camphene hydrochloride. The transformation is a unimolecular change. In general, the values of *k* exhibit a decreasing tendency during the course of an experiment, which is due to the partial dissociation of camphene hydrochloride in solution into camphene and hydrogen chloride and the catalytic acceleration of the transformation by the latter; with decreasing content of camphene hydrochloride, the concentration of the hydrogen chloride also diminishes and the rate of change becomes smaller. The results indicate that the conversion of camphene hydrochloride into *isobornyl* chloride is due to an actual, intramolecular atomic displacement and cannot be explained by elimination and re-addition of hydrogen chloride. An intimate connexion is obvious between the dielectric constant of the solvent and its influence on the rate of conversion of camphene hydrochloride into *isobornyl* chloride, the latter being greatest in strongly ionising solvents (nitromethane and acetonitrile) and smallest in non-ionising solvents (benzene, ether, light petroleum). From this it appears that the velocity of the change depends on the degree of the electrolytic dissociation of camphene hydrochloride and that the conversion only occurs subsequently to ionisation. The change, therefore, consists not in a wandering of the chlorine



scheme, whereby the more strongly ionised camphene hydrochloride passes into the less strongly ionised *isobornyl* chloride. A perfect parallelism between rate of change and dielectric constant of the solvent is not observed, exceptions being encountered in the cases of cresol, sulphur dioxide, and anisole, on the one hand, and of ether on the other. Precisely these solvents have an anomalous action towards triphenylmethyl chloride, and it seems probable that their effect can be explained in a uniform manner, and that phenol, sulphur dioxide, and (to a less extent) anisole form complex compounds with camphene hydrochloride in which the mobility of the chlorine atom is increased or the ionisation of camphene hydrochloride occurs to a greater degree. The influence of complex formation on the ionisation and therefore on the rate of transformation of camphene hydrochloride is further

illustrated by the catalytic acceleration of the change by the halogen acids, antimony pentachloride, stannic chloride, ferric chloride, mercuric chloride, and antimony trichloride (all of which yield additive compounds with triphenylmethyl chloride), whereas phosphorus trichloride and silicon tetrachloride, which show no tendency to complex formation, are inactive. In the presence of catalysts, the change is unimolecular, and the constancy of k is more satisfactory than in their absence, since the effect of the hydrogen chloride (see above) is of relatively less importance.

The influence of the solvent on the degree of ionisation of camphene hydrochloride is also observed in the alcoholysis of this substance. The conversion of camphene hydrochloride into camphene hydrate methyl ether by methyl alcohol has been examined kinetically in the presence of nitromethane, acetonitrile, nitrobenzene, benzonitrile, anisole, bromobenzene, chlorobenzene, benzene, ethyl bromide, pyridine, and ether and it is found that the effect of the solvents is approximately the same as that observed in the conversion of camphene hydrochloride into *isobornyl* chloride. The analogy between triphenylmethyl chloride and camphene hydrochloride is illustrated further by a study of the alcoholysis of the former by amyl alcohol in the presence of nitrobenzene, benzonitrile, bromobenzene, chlorobenzene, anisole, ethyl bromide, benzene, pyridine, and ether, and the complete parallelism thus established justifies the assumption of the ionisation of camphene hydrochloride which cannot be measured directly on account of the ease with which it is converted into *isobornyl* chloride in all solvents which are suitable for measurements of conductivity.

Unfortunately, it is not possible to measure the rate of the reverse transformation of *isobornyl* chloride into camphene hydrochloride, but it is shown to be subject to the same catalytic influences by observation on the rate of decomposition of *isobornyl* chloride into camphene or camphene hydrate and hydrogen chloride under differing conditions, this decomposition being consequent on the transformation of *isobornyl* chloride into camphene hydrochloride.

The mutual transformation of *isobornyl* and *bornyl* chlorides takes place with much greater difficulty and in non-ionising solvents in the absence of catalysts is extremely slow at 130°. In cresol, sulphur dioxide, or chlorobenzene in the presence of stannic chloride, the change occurs with considerable rapidity at 70°, but the conditions are unsuitable for physico-chemical examination.

The authors have previously described (*loc. cit.*) the isolation of "true" pinene hydrochloride. The compound had been obtained previously in 1914 by Aschan (cf. A., 1921, i, 795), whose work, however, was not generally accessible until recently. It has now been obtained in the crystalline condition (although not quite free from *bornyl* chloride) by the action of hydrogen chloride on a dilute solution of pinene in light petroleum at -60°. It is instantly decomposed by water and alcohol, and can be titrated in alcoholic solution in the same manner as hydrochloric acid. It resembles camphene hydrochloride in that its solutions can be greatly stabilised by the addition of ether.

The estimation of camphene hydrochloride, *isobornyl* chloride, and free hydrogen chloride in their solutions is effected in the following manner. Camphene hydrochloride is determined by adding an excess of *N*/5-absolute alcoholic sodium ethoxide to the solution together with methyl alcohol, and titrating the excess of alkoxide after the mixture has been preserved for an hour at 20° with *N*/10-alcoholic hydrogen chloride in the presence of phenolphthalein as indicator. *isoBornyl* chloride is estimated by boiling the mixture for an hour with *N*/5-alcoholic potassium hydroxide solution, with the addition of a little water. Free hydrogen chloride in the presence of camphene hydrochloride is estimated by taking advantage of the observation that the dissociation of the latter can be delayed to such an extent by the addition of ether that the free acid can be titrated with sodium ethoxide solution in the presence of iodoeosin as indicator.

Camphene hydrochloride is prepared by treating a solution of camphene in ether with hydrogen chloride and filtration of the product with exclusion of atmospheric moisture; the material, when most carefully prepared in this manner, contains about 9–10% of *isobornyl* chloride. A somewhat purer product (containing 93.3% of camphene hydrochloride) can be secured by using light petroleum as solvent, in which, however, the addition of the halogen acid occurs very slowly. *isoBornyl* chloride is obtained by saturating a solution of camphene in ethyl bromide with hydrogen chloride at 10–20°, heating the mixture uninterruptedly under a reflux condenser during six days at 55°, removing the solvent, and crystallising the residue from amyl alcohol. It has m. p. 161.5°, and contains 99.6% of pure *isobornyl* chloride; the presence of camphene hydrochloride cannot be detected. H. W.

The Velocity of Reaction in Mixed Solvents. III. The Influence of Temperature on the Velocity of Saponification of Esters. HAMILTON McCOMBIE, HAROLD ARCHIBALD SCARBOROUGH, and RICHARD HARDCASTLE SETTLE (T., 1922, 121, 2308–2318).

Rate of Hydrolysis of Wheat Gliadin. HUBERT BRADFORD VICKERY (*J. Biol. Chem.*, 1922, 53, 495–511).—Results are given of a large number of experiments in which measurements were made of the rate of hydrolysis of gliadin by acids of various concentrations and also by alkalis. Hydrolysis both of the amide nitrogen and of the peptide linkings proceeds more rapidly the greater the concentration of the acid. In the early stages, alkalis are more effective than acids. Owing to secondary decompositions, however, considerably larger quantities of ammonia are liberated by alkalis than by acids. Hydrolysis is more rapid with barium hydroxide than with sodium hydroxide of equivalent concentration. E. S.

Influence of the Catalyst on the Thermodynamic Quantities Regulating the Velocity of a Reaction. E. VAN THIEL (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 25, 199–209).—After a theo-

retical discussion of the velocity of chemical reactions in which the hypotheses of Trautz, Lewis, Scheffer, and Marcelin are considered, it is shown that the hypotheses of Trautz and Lewis demand that a reaction which is strongly catalysed shall have a smaller temperature coefficient than the same reaction uncatalysed or weakly catalysed, whilst Scheffer's hypothesis shows that a catalysed reaction can have a temperature coefficient and an energy increment which are greater than those of the uncatalysed reaction. An attempt has been made to decide between these two hypotheses by means of measurements of the rate of acetylation of diphenylamine at 0°, 20°, 30°, 40°, and 50° both without a catalyst and in the presence of *p*-bromobenzenesulphonic acid or *p*-toluenesulphonic acid as catalyst. The results of these experiments show that Scheffer's hypothesis is in keeping with the facts. J. F. S.

Dislocation Theory of Catalysis. J. BÖESEKEN (*Proc. K. Akad. Wetensch. Amsterdam*, 1922, 25, 210—215).—A theoretical paper in which the author discusses the mechanism of catalysis. He starts from the idea that ideal catalysts are those which are not temporarily fixed in intermediate compounds and so withdrawn from the reaction mass. Real catalysis is the interaction between the catalyst and the molecules which has nothing to do with the formation of a compound. This interaction is termed a *dislocation*, and may be regarded as consisting of a change in the paths of the electrons. The hypothesis is considered in the case of the photo-oxidation of alcohols in the presence of *p*-benzophenone and oxygen. In this case, the author has shown (A., 1921, ii, 500) that above a certain concentration of the ketone the velocity is independent of this concentration, and, further, it is proportional to the square of the intensity of the light and to the concentration of the alcohol, which is explained as follows: The ketone absorbs part of the light and is activated by it, the quantity of activated ketone is $I(1 - e^{-kcd})$, where k is the absorption coefficient, c the concentration of the ketone, and d the thickness of the layer. When k , c , and d are large, the quantity of activated ketone becomes proportional to I and independent of c . The whole process then is as follows: Under the influence of light a quantity of photo-ketone $kI(1 - e^{-kcd})$ is formed instantaneously, this amount approaches the value kI , so that the first stage of the process is represented by the equation (1) ketone + light = photo-ketone, and is followed by the reaction which regulates the process. (2) photo-ketone + alcohol = (photo-ketone), alcohol; the meeting of the two molecules activates two hydrogen atoms of the alcohol and is represented by (3) (photo-ketone), alcohol \rightarrow [(photo-ketone), active alcohol]. This process which probably takes place with the velocity of light, is the real catalysis, the dislocation. The alcohol molecules are thereby brought into the condition to react according to (4) $2[(\text{photo-ketone}) \text{ active alcohol}] + \text{O}_2 = 2 \text{ ketone} + 2 \text{ aldehyde} + 2 \text{H}_2\text{O}$, which also occurs with a great velocity. The photo-ketone may be replaced by palladium, and the same type of reaction will occur and is similarly explained. This explanation demands

intimate contact between the catalyst and the reacting molecules. It is shown that Scheffer's relation of the reaction constant, $\log k = (E_0 - E)/RT + B$ represents the thermodynamic aspect of the above hypothesis (cf. Thiel, preceding abstract). J. F. S.

The Influencing of the Catalysis of Hydrogen Peroxide with Platinum by Röntgen Rays. ANTON KAILAN (*Ber.*, 1922, 55, [B], 2492—2494).—Recalculation of the experimental results of Schwarz and Friedrich (this vol., ii, 436) show that the catalytic decomposition of concentrated hydrogen peroxide is a bi-molecular action and not unimolecular as stated by these authors.

In an addendum to the communication, the correctness of the criticism is recognised by Schwarz and Friedrich. H. W.

Catalysis. XIV. The Action of Nitric Acid on Metals and an Example of a Periodic Reaction. B. C. BANERJI and N. R. DHAR (*Z. anorg. Chem.*, 1922, 122, 73—80).—The rate of standard solution of copper, silver, lead, nickel, brass, mint silver, and of copper-nickel alloy, in nitric acid, was studied in the presence of different salts. Ferrous salts acted as catalysts; this catalytic effect is ascribed to the nitrous acid which is formed by the action of ferrous salts on nitric acid. Ferric salts were also found to act as positive catalysts; this is due to the formation of ferrous salts by the action of the metals copper, silver, etc., on the ferric salts. Oxidising agents, for example, potassium permanganate, potassium chlorate, are negative catalysts because they prevent the formation of nitrous acid. The dissolution of a 50% copper-nickel alloy in nitric acid takes place in well-defined periods of reaction, each period being followed by a period of no reaction. W. T.

Catalytic Action at Solid Surfaces. VIII. Action of Sodium Carbonate in Promoting the Hydrogenation of Phenol. E. F. ARMSTRONG and T. P. HILDITCH (*Proc. Roy. Soc.*, 1922, [A], 102, 21—27; cf. this vol., ii, 41).—The results of a number of experiments on the hydrogenation of phenol in the presence of nickel powder and sodium carbonate at 180° and under a pressure of hydrogen of 80 lb. per sq. inch are described. The object of the experiments was to ascertain the part played by a promoter in a catalytic change. The products of the reaction are cyclohexanol 95%, some cyclohexanone, with practically no unchanged phenol. The presence of sodium carbonate has an accelerating action which is dependent on the amount of carbonate present; if an excess is present the hydrogenation is retarded. The optimum amount of sodium carbonate bears no relationship to the amount of phenol present, but is closely connected with the amount of nickel catalyst employed, and in the present experiments is found to be at the point where the amount of sodium carbonate is about 25% of the weight of the nickel. The exact mechanism by which the sodium carbonate assists the hydrogenation has not been discovered, but it appears that the stimulation is not a true acceleration but the restoration of the nickel to its normal function by the suppression of a retardation, which is probably of the nature of a

poisoning of the catalyst by the phenol. It is found that the curvilinear nature of the time-absorption relationship in the absence of sodium carbonate becomes linear in the presence of this substance. This fact furnishes further evidence that the rôle of the promoter in this case is a protective one with respect to the catalyst, rendering the latter free to exert its normal functions.

J. F. S.

Catalytic Action at Solid Surfaces. IX. Action of Copper in Promoting the Activity of Nickel Catalyst. E. F. ARMSTRONG and T. P. HILDITCH (*Proc. Roy. Soc.*, 1922, [A], **102**, 27—32; cf. preceding abstract).—It is shown that a copper-nickel catalyst may be produced in hydrogen when suitably supported compounds are heated at 180°, whereas supported nickel compounds alone are not reduced by hydrogen below 300° and not rapidly below 350—400°. It is shown that for the reduction of the nickel at this low temperature an intimate mixture of the two compounds must be used, which must be more intimate than can be obtained by mechanical mixture. Co-precipitation is not always sufficient to obtain the desired degree of intimacy, for a comparison of the activity of the catalysts with the nature of the nickel cupricarbonates from which they are produced shows that both the nickel and copper must be in the same complex molecule to obtain any degree of catalytic activity. On the basis of the foregoing, the relatively low temperature of the reduction of the nickel is explained as follows: the reduction of copper carbonate or cupric oxide is strongly exothermic, and even although the exterior temperature is only 180°, it is evident that the temperature inside the molecule must momentarily, at least, be far above the outside temperature.

J. F. S.

Influence of *s*-Trinitrobenzene on Reactions in which Arylamines are involved. J. J. SUDBOROUGH and R. C. SHAH (*J. Indian Inst. Sci.*, 1922, **5**, 29—35).—To determine whether, on account of the readiness with which polynitro-aromatic compounds form additive compounds with most of the simple arylamines, such nitro-compounds would have a catalytic influence on reactions involving aromatic amines, a number of such reactions which are readily catalysed by iodine were carried out in presence of *s*-trinitrobenzene. The reactions studied were the methylation of aniline, formation of phenyl- α -naphthylamine from aniline and α -naphthylamine, condensation of aniline with benzophenone, the formation of thiodiphenylamine from diphenylamine and sulphur, and the acetylation of aniline. In no case had the *s*-trinitrobenzene any catalytic influence, whilst iodine was a powerful catalyst for all except the last of these reactions.

E. H. R.


Atoms and Electrons. ROBERT N. PEASE (*Nature*, 1922, **110**, 379—380).—In view of the facts (1) that the distances between the atomic centres in crystals of cerium and thorium are practically the same, the number of electrons being 58 and 90,

respectively, and (2) that in the atom of uranium (atomic volume 12·8) there are concentrated 32 more electrons than in the atom of neodymium (atomic volume 20·7). It is significant that elements of higher atomic number than uranium are not known to exist, and that most of those of immediately lower atomic number are unstable. In the case of elements of high atomic number, the electron-density must therefore be greater than in other cases, and be accompanied by a greater attractive force on the nucleus. It is suggested that periodical exertion by numbers of electrons of attractive forces on the positive nucleus in the same direction may result in the extraction of a positively charged constituent of the latter and its expulsion as an α -particle. The rate of decay would depend primarily on the stability of the nucleus, and the suggested mechanism would constitute the trigger action.

A. A. E.

A. A. E.

Periodic Structure of Atoms and Elements. H. NEWMAN ALLEN (*Nature*, 1922, **110**, 415).—In one of the models postulated by Crehore (A., 1921, ii, 632; this vol., ii, 438, 494), one of the particles marked 4 is also marked with a dot, evidently indicating that it has a charge $+e$ only. By calling these particles γ , it has been found possible to construct hypothetical models for a considerable number of atoms of the form indicated by the following examples representing nitrogen (I), fluorine (II or III), and uranium (IV), respectively:



The strokes represent electrons; in II, the α 's join to form a ring, in III, two γ 's join to the α to form a ring, and in IV, a ring structure is also obtained.

A. A. E.

The Rare Earths in the Periodic System. CARL RENZ (*Z. anorg. Chem.*, 1922, **122**, 135—145).—The author reviews the literature on the subject and discusses the difficulties. He suggests as a solution a modification of Soddy's helical representation of the periodic law. W. T.

W. T.

Determination of the Molecular Weight of Substances in Alcoholic Solution from the Elevation of the Flash Point.
ROBERT WRIGHT (T., 1922, 121, 2247—2250).

Polar and Non-polar Valency in Organic Compounds. W. E. GARNER (*Nature*, 1922, **110**, 543—544).—Two explanations may be given of the positive and negative relationships of groups in organic compounds: (a) there may be a partial transference of an electron between the group and the residue of the molecule, or (b) there may be a varying concentration of polar molecules in the typically non-polar substance. In the consideration of the second possibility, an equilibrium may be imagined to exist between the polar and non-polar substances which will be affected by the

temperature, nature of the solvent, etc. Thus, in those substances where the stability of the non-polar arrangement is very great, the occurrence of both forms, $\bar{A}\bar{X}^+$ and $\bar{A}\bar{X}^-$, will be possible, and in the presence of a suitable solvent these may give rise to the respective ions. A. A. E.

Melting Point Tubes as Reaction Vessels. AUGUST FUCHS (*Monatsh.*, 1922, **43**, 129—137).—Details are given for carrying out the commoner manipulations, using very small quantities of material (2—5 mg.). That satisfactory preparations and purifications can be effected by the methods indicated is proved by a record of examples actually worked out. C. K. I.

Perforation Apparatus for the Extraction of Liquids by Liquids. Non-miscible Liquids. FAYOLLE and CH. LORMAND (*Chim. et Ind.*, 1922, **3**, 273—274).—Two types of apparatus are described for the extraction of a liquid by another liquid with which it does not mix. One type is for use where the solvent is heavier than the liquid to be extracted, and the other where the solvent is the lighter of the two liquids. In either case the extraction is continuous. W. G.

A Shaking Machine for Large Quantities of Fluid. HARTWIG FRANZEN (*Z. physiol. Chem.*, 1922, **122**, 86—87).—An apparatus is described suitable for stirring large amounts of lead precipitates in aqueous suspension, while treating them with carbon dioxide or hydrogen sulphide. W. O. K.

Inorganic Chemistry.

Vapour Pressure of Hydrogen. Determinations in the Region of Liquid Hydrogen. H. KAMERLINGH ONNES and J. PALACIOS MARTINEZ (*Anal. Fis. Quím.*, 1922, **20**, 233—242).—The vapour pressures of liquid hydrogen at temperatures near its normal boiling point were measured using a helium thermometer. By interpolation, the boiling point of hydrogen at a pressure of 759·549 mm. of mercury is given as 20·35° K (Kelvin international scale). G. W. R.

Spontaneous Incandescence of Substances in Atomic Hydrogen Gas. R. W. WOOD (*Proc. Roy. Soc.*, 1922, [*A*], **102**, 1—9).—A number of experiments with very long hydrogen discharge tubes are described from which it is shown that some metals, oxides, and other substances are raised to incandescence when introduced into a stream of atomic hydrogen, the surface of the substance acting as a catalyst in bringing about the recombination of the atoms. Atomic hydrogen, practically free from molecular hydrogen, may be drawn by a pump from the central portion of a long discharge tube excited by a current of high potential. Fire

polished glass surfaces, such as the walls of a glass tube, have a comparatively feeble catalysing power whilst fractured surfaces cause the recombination of the atoms, and are strongly heated. The action of water vapour or oxygen in enhancing the Balmer spectrum, and suppressing the secondary spectrum of hydrogen, is probably due to its action on the walls of the tube, which, when dry, catalyse the atomic hydrogen as fast as it is formed by the current. The peculiar spectroscopic phenomena observed with long hydrogen tubes, described in an earlier paper (A., 1920, ii, 569) are explained. Methods are suggested for determining the physical and optical properties of atomic hydrogen gas. J. F. S.

Evidence of the Existence of Isotopes of Chlorine. MATAKICHI ISHINO (*Mem. Coll. Sci. Kyōtō*, 1921, 4, 311—315).—Employing the method of positive-ray analysis, using the crossed-deflexion method devised by Thomson, the author has obtained evidence of the existence of chlorine isotopes of respective atomic weights 34 and 36, and of positively charged chlorine atoms.

J. S. G. T.

The Decomposition of Chlorine into Atoms. MAX TRAUTZ and WALTER STÄCKEL (*Z. anorg. Chem.*, 1922, 122, 81—131).—The apparatus employed and methods of working are described in detail. The amount of chlorine decomposed was 1.50% at 1200°, 2.10% at 1240°, and 3.05% at 1280°. The limit of the absorption band for chlorine was found to be 390—420 $\mu\mu$. This corresponds with 67,500—73,000 cal. as the heat of decomposition according to Trautz's "approximate" equation $q_0 = Ah\nu$ (A., 1918, ii, 151). The heat of decomposition of chlorine calculated from the temperature coefficient was found to be $71,000 \pm 3,000$ cal. The same result was obtained by the use of the quantum theoretical constants and also by the use of Victor Meyer's vapour density measurements. With the exception of hydrogen, the heat of decomposition of the diatomic elements found experimentally agrees with the value obtained from the equation $d_0 = (5.78 \times 10^5 / \sqrt{M})$ cal., where M is the molecular weight.

W. T.

Vapour Pressure of Solid Chlorine and Bromine. F. A. HENGLEIN, G. VON ROSENBERG, and A. MUCHLINSKI (*Z. Physik*, 1922, 11, 1—11).—The vapour pressure of solid and liquid chlorine and solid bromine has been determined over a wide range of temperature. The following values are recorded for chlorine, 119.2° 0.0013; 126.0°, 0.0063; 133.0°, 0.028; 146.6°, 0.30; 161.1°, 2.75; 177.6°, 17.40; 194.5°, 64.40, and for bromine 177.6°, 0.0020; 210.0°, 0.26; 228.8°, 1.12; 227.0°, 1.71; and 241.1°, 6.44; the temperatures are expressed in absolute degrees and the pressures in millimetres of mercury. Vapour pressure formulæ have been deduced for both cases and have the form: $\log p = -1160/T + 7773$ for liquid chlorine; $\log p = -1530/T + 9950$ for solid chlorine; these formula are representative over the temperature ranges -95° to -78° and -154° to -112° , respectively, $\log p = -12150/T^{1.368} + 75030$ for solid bromine over the range -32° to

—96°. From the above data a number of constants have been calculated which include the following: Chlorine, m. p. 170·0° Abs.: vapour pressure at the melting point 8·9 mm., molecular heat of vaporisation at the melting point 5300 cal., heat of sublimation of solid chlorine at the melting point 6960 cal., molecular heat of fusion 1660 cal.; bromine: vapour pressure at the melting point 44·12 mm., molecular heat of sublimation at the melting point 9740 cal., specific heat of sublimation 60·91 cal., specific heat of vaporisation at the melting point 48·4 cal., molecular heat of fusion 12·5 cal. From the dissociation equilibrium of the chlorine molecule and the chemical constant as determined by Stern and Tetrode the vapour pressure of diatomic chlorine has been calculated and the chemical constant for diatomic chlorine in its normal condition has also been obtained.

J. F. S.

Physical Constants of Ozone. E. H. RIESENFELD and G. M. SCHWAB (*Z. Physik*, 1922, 11, 12—21; cf. this vol., ii, 637).—An account of the determination of a number of physical constants of pure ozone which was prepared as previously described by the authors (*loc. cit.*). The following values are put on record: m. p. —249·7°, b. p. —112·4°; critical temperature, —5°; specific gravity at —183°, $1·71 \pm 0·1$; change of density with temperature, $1/d = a + bT + cT^2$, where $a = 0·51193$, $b = 0·0004559$, $c = 0·000003929$; density at the boiling point, 1·46; coefficient of expansion, 0·0025; critical density, 0·537; critical pressure, 64·8 atmospheres. The authors show that neither in the gaseous nor in the liquid state is there any other molecular species present than that represented by the formula O_3 .

J. F. S.

Solubility of Sulphur Dioxide in Suspensions of Calcium and Magnesium Hydroxides. WM. THOMPSON SMITH and REGINALD B. PARKHURST (*J. Amer. Chem. Soc.*, 1922, 44, 1918—1927).—The solubility of sulphur dioxide has been determined in water, milk of lime, and milk of magnesia, at partial pressures of sulphur dioxide up to 760 mm., and temperatures from 5° to 60°. All proportions of calcium and magnesium hydroxides were used in suspensions of a total alkalinity of one equivalent per litre. It is shown that the concentration of sulphur dioxide as sulphurous acid is proportional to its partial pressure. With solutions of the same alkali concentration, the percentage salting-out effect increases with, but more than in proportion to, the temperature. With solutions of constant temperature, the percentage salting-out effect increases with the alkali concentration, but is less than proportional to it. Varying proportions of calcium and magnesium hydroxide have no effect on the equilibrium concentration of sulphur dioxide as sulphurous acid.

J. F. S.

The Physico-chemical Study of the Lead Chamber Process. MAX FORREER (*Bull. Soc. chim. Belg.*, 1922, 31, 254—293).—A detailed description of a form of apparatus in which sulphur dioxide, nitrogen peroxide, water, oxygen, and nitrogen may be brought into contact with each other under definite conditions

of pressure, temperature, and, in the case of liquids, surface of reaction. The proportions of these substances could be varied at will. It is shown that, in such a system, the formation of sulphuric acid only occurs in presence of a liquid phase, so that the system must be heterogeneous; the formation of the acid takes place in a shorter time and the yield is increased if a liquid, either sulphuric acid or water, is present at the outset. Further, the rate at which water vapour is supplied during the course of the reaction exerts a considerable effect both on the velocity of reaction and on the concentration of the product. Under certain conditions, the acid formed may disappear, which appears to indicate that the reactions assumed to take place in the chambers are at least partly reversible. For low concentrations of the gases, there is an optimum rate of intake for water vapour: this gives the best yield and, at the same time, the most concentrated acid. As the gas concentration increases, the optimum point shifts in the direction of diminution of water supply; a curve is given showing the speed of reaction plotted against concentration of gas. The relation between these two factors is much less complex when the composition of the liquid phase is constant. The author infers from his experimental results the existence of an intermediate substance of which water is a constituent; it is, however, decomposed in presence of an excess of water. He points out that, of the series of reactions which occur, the slowest is the "limiting reaction" in that it conditions the rate of formation of the product and shows that, in practice, the reaction velocity is strongly influenced by the rate of supply of nitrogen peroxide and water, but is practically unaffected by the supply of sulphur dioxide.

An attempt is made to calculate the order of the reaction, and, for this purpose, the heterogeneity of the system and the non-equivalence of the reactants are neglected. The values obtained vary from 1.26 to 3.67. However, when the concentration of the liquid phase does not vary, the values are in the neighbourhood of 2. In the case of temperature variations which have a similar effect to that of variations in water supply, it was not found possible to obtain a coefficient in agreement with the experimental data.

H. J. E.

Density of Atmospheric Nitrogen. A Small Anomaly in the Air of Madrid. M. PAYÁ and E. MOLES (*Anal. Fis. Quím.*, 1922, 20, 247—254).—The normal density of nitrogen obtained from the air of Madrid is given as 1.25681. The density of air in Madrid corresponds with a slightly higher proportion of oxygen than the normal for other places. The anomaly is attributed to the presence of higher proportion than normal of ozone and gases of the argon group.

G. W. R.

The Reactions between Gaseous Oxides of Nitrogen and Alkaline Solutions. A. SANFOURCHE (*Compt. rend.*, 1922, 175, 469—472).—The action of oxides of nitrogen on solutions of alkali hydroxide in which the calculated proportions of nitrate and nitrite are formed only occurs if the alkali is in excess at all points

in the solution. Otherwise, reaction with the water takes place with formation of nitric acid and nitric oxide, the latter being further oxidised if oxygen is present. Moreover, if all the oxides of nitrogen are not absorbed in their first passage through the solution, this reaction with water continues in the gaseous phase with water vapour. This is shown by the formation of a nitric acid mist and by its subsequent deposition as acid dew on the upper portion of the walls of the vessel in spite of the presence of excess of alkaline solution. It is pointed out that estimation of nitrous vapours by the method of absorption in alkali hydroxide is liable to lead to considerable error and for this reason it is preferable to use sulphuric acid as the absorbent. H. J. E.

The Solubility of Arsenic Trichloride in Concentrated Hydrochloric Acid at 100°. W. D. TREADWELL [with CH. MUSSLER] (*Helv. Chim. Acta*, 1922, 5, 818—821).—The solubility of arsenic trichloride in concentrated hydrochloric acid at 100° was determined by an indirect method which consisted in measuring the rate of volatilisation of the trichloride from the solution in a current of hydrogen chloride at 100°. From Henry's law, it follows that $\log(x_0 - x)/x_0 = -kV/V_0$, where x is the quantity of trichloride volatilised by the volume V of gas in a given time, x_0 is the quantity of trichloride originally present, V_0 the volume of the solution, and k the solubility coefficient. Experiments showed that the value of k remained constant for a considerable period of time, its mean value being 0.0233. From this it can be calculated that to volatilise 99% of the arsenic trichloride present in 100 c.c. of solution at 100° requires 19.8 litres of hydrogen chloride. From the known vapour pressure of arsenic trichloride, a litre of the vapour contains 2.34 grams. Hence the solubility in concentrated hydrochloric acid at 100° is $2.34/k = 100.3$ grams per litre. E. H. R.

Boric Acid Phosphors. RUDOLF TOMASCHEK (*Ann. Physik*, 1922, [iv], 67, 612—648; cf. Tiede and Wulff, this vol., ii, 245).—A number of boric acid phosphors have been prepared by the addition of phthalic anhydride, terephthalic acid, the sodium salt of fluorescein (uranin), naphthalic anhydride, and phenanthrene. The phosphorescence of these organic phosphors is in many ways simpler than that of the metal phosphors. In general, the bands of the former are all excited by the same wave-lengths of light, and owe their origin to the same phosphorescence centre. Thus the three mercury lines, 365.4, 313, and 253.6 $\mu\mu$, all appear to excite the four components of the band of the phenanthrene-phosphor. On the other hand, the different bands of the metal phosphors are excited independently. The relative intensities of the components of the bands of the boric acid substances may change with temperature; the uranin-phosphor emits a yellow light at -70° , but, when heated on metal foil, this radiation rapidly changes into a bluish-green phosphorescence. In opposition to the heavy metal phosphors, the duration and intensity of the emitted light are influenced only slightly by changes in temperature, being, in many

cases, the same at the temperature of liquid air as at the ordinary temperature. Neither can the phosphorescence be extinguished by exposure to red light. Phenanthrene-phosphor is especially sensitive to light, and it is shown that the decolorisation which occurs is not in any way the cause of the phosphorescence, as suggested by Perrin. The wave-lengths of the absorption bands of the transparent phosphor frequently coincide with those of the exciting radiation, and in the case of phenanthrene, correspondence is found to occur between the fluorescence bands and the absorption bands in the ultra-violet. The duration of the phosphorescence for organic boric acid phosphors is about one minute, and this comparatively short period points to the presence of very small centres; this is supported by the difficulty of fully exciting these centres. When the terephthalic acid phosphor is fully excited by an intense beam of light, the larger portion of the total phosphorescence takes place within the first second, so that the centres are mainly of very short duration. The centres do not all possess the same duration. The $1/L$ curves indicate that a bimolecular chemical reaction is not the cause of the phosphorescence, which can be explained satisfactorily on the assumption of a photo-electric process. Three kinds of emission take place, phosphorescence, fluorescence, and an emission in the ultra-violet. The conclusion is drawn that the fields of force are almost exclusively in the interior of the organic molecules.

W. E. G.

Borates. The System $(\text{NH}_4)_2\text{O}-\text{B}_2\text{O}_3-\text{H}_2\text{O}$. Temperature-Concentration Diagram. U. SBORGI and L. FERRI (*Mem. Accad. Lincei*, 1922, [v], 13, 570—591).—The results previously obtained (A., 1913, ii, 213, 318; 1915, ii, 686; this vol., ii, 568) are collected.

T. H. P.

The Modifications of Silicon. II. Silicon from Copper Silicide. WILHELM MANCHOT and HERBERT FUNK (*Z. anorg. Chem.*, 1922, 122, 22—26).—Specimens of silicon obtained from a copper silicide regulus have been studied; the results were similar to those obtained by the authors when employing aluminium and silver reguli (this vol., ii, 286). Conditions which favour the formation of crystals produce silicon which reacted but slowly with hydrofluoric acid. Crystalline silicon very soluble in this acid could not be obtained.

W. T.

Alkali Iodides. L. W. WINKLER (*Pharm. Zentr.-h.*, 1922, 63, 386—387).—Analyses of samples of commercial potassium iodide and sodium iodide showed that these substances have a high degree of purity. The respective quantities of iodide found corresponded with 98.6 to 99.6% of potassium iodide and with 97.0 to 99.9% of sodium iodide.

W. P. S.

Composition of the Liquid Phases in a Univariant System, Liquid-Liquid-Vapour, for Mixtures of Sodium and Ammonia. CHARLES A. KRAUS and WALTER W. LUCASSE (*J. Amer. Chem. Soc.*, 1922, 44, 1949—1953).—In an earlier paper (A., 1907, ii, 935), it has been shown that at low temperatures concentrated

solutions of sodium in liquid ammonia separate into two liquid phases which have an upper critical end-point at about -45° . The concentrations of these liquid phases have now been determined by electrical conductivity measurements and it has been found that the upper critical end-point lies at -41.6° and at a composition of 4.15 atom % of sodium. The phase relations in the system sodium-ammonia are briefly discussed. J. F. S.

Nature of the Complexes formed between Sodium and Tellurium in Liquid Ammonia. CHARLES A. KRAUS and C. Y. CHIU (*J. Amer. Chem. Soc.*, 1922, **44**, 1999—2008).—The reaction between sodium and tellurium in liquid ammonia has been investigated and the composition of the resulting solutions has been determined under various conditions. The initial compound formed is normal sodium telluride of the formula Na_2Te . In equilibrium with the normal telluride, the complex in solution has a composition corresponding with the formula Na_2Te_2 . In equilibrium with free tellurium the solution has a composition which varies as a function of the concentration. The maximum amount of tellurium present corresponds approximately with the formula Na_2Te_4 . J. F. S.

Crystal Structure of Silver Molybdate. RALPH W. G. WYCKOFF (*J. Amer. Chem. Soc.*, 1922, **44**, 1994—1998).—Making use of the data obtained from the reflection spectra and Laue photographs of silver molybdate, the crystal structure of this substance has been shown to be the same as that of the spinels and magnetite. The length of the side of the unit cube, which contains eight molecules, is 9.26 Å.U., and the parameter defining the position of the oxygen atoms is very close to $3/8$. The distance between molybdenum and adjacent oxygen atoms is 2.00 Å., between silver and the nearest oxygen atom 2.31 Å., and between the nearest molybdenum atom and silver 4.02 Å. J. F. S.

The Dissociation of the Chlorides of Bivalent Metals in Aqueous Solution. A. GÜNTHER-SCHULZE (*Z. Elektrochem.*, 1922, **28**, 387—389).—In a previous paper it was shown from a study of the basic exchange between copper salts in solution and permutite that the copper salt solution, even at high dilutions, contains complex kations (this vol., ii, 504). The investigation has now been extended to salts of barium, strontium, calcium, cadmium, manganese, cobalt, and nickel, using the chlorides of the metals and ammonium permutite. If a represents the number of equivalents of metal passing into the permutite in exchange for ammonia, and c the number of equivalents of chlorine likewise disappearing from the solution, then if the metallic chloride is present in solution only in the form of M^{++} and Cl^- ions, $c=0$ and $a/c=\infty$. If, however, the ions are MCl^+ and Cl^- , the MCl^+ ion passing into the permutite, $a/c=2$. If still more complex ions are formed, such as M_2Cl_3^+ , $a/c<2$. Applying this reasoning to the experimental results, it is found that at the highest concentrations of all the above salts, some MCl^+ ions are present and,

in the cases of magnesium, nickel, and copper, still more complex ions. The first stage of dissociation into MeCl' and Cl' preponderates in the case of those metals having the smallest ionic radius. At a dilution of 1.0*N*, a considerable proportion of MeCl' ions is found only in the case of those metals having the smallest ionic radius, namely, nickel and copper.

E. H. R.

Transference of the Acid Radicle in the Solid Phase. I. Carbonates of the Alkaline Earths and Magnesium. J. ARVID HEDVALL and J. HEUBERGER (*Z. anorg. Chem.*, 1922, **122**, 181—187).—It was found that if the oxides of the elements considered are arranged in order of increasing basicity MgO , CaO , SrO , BaO , then the oxide of the more basic will on heating with the carbonate of the less basic element be changed into the carbonate without the evolution of gas and at a temperature well below the temperature of dissociation of the carbonate taken. The temperature of the reaction decreases and the heat effect increases with increasing difference of basicity, for example, $\text{MgCO}_3 + \text{BaO} = \text{BaCO}_3 + \text{MgO} + 19 \text{ cal. } (t=230^\circ)$; $\text{SrCO}_3 + \text{BaO} = \text{BaCO}_3 + \text{SrO} + 6.14 \text{ cal. } (t=280^\circ)$.

W. T.

The Crystal Structure of Glucinum and Glucinum Oxide. L. W. MCKEEHAN (*Proc. Nat. Acad. Sci.*, 1922, **8**, 270—274).—The glucinum was obtained by electrolysis of sodium glucinum fluoride, $2\text{GlF}_2 \cdot \text{NaF}$, and contained a few per cent. of glucinum oxide. The diffracting centres in the metal and its oxide lie at the points of two symmetrically interpenetrating hexagonal space lattices. For glucinum oxide, the axial ratio, $c/a=1.63$, and $a=2.696 \times 10^{-8} \text{ cm.}$; for glucinum, $c/a=1.58$ and $a=2.283 \times 10^{-8} \text{ cm.}$ These values give computed densities within 1% of the best previous values. Glucinum more closely resembles magnesium, zinc, and cadmium, rather than calcium, strontium, and barium; the oxide is an analogue of zinc oxide, and the structure suggested by W. L. Bragg (*A.*, 1920, ii, 433) for the latter substance is also possible for glucinum oxide.

W. E. G.

Physical Chemistry of the Oxides of Lead. VI. The Anodic Behaviour of Lead and Lead Dioxide. SAMUEL GLASTONE (*T.*, 1922, **121**, 2091—2098).

The Isomerism of Metallic Oxides. I. Lead Monoxide. MALCOLM PERCIVAL APPLEBEY and ROBERT DOUGLAS REID (*T.*, 1922, **121**, 2129—2136).

The Hindrance of Precipitations with Hydrogen Sulphide by Neutral Chlorides. L. DEDE and P. BONIN (*Ber.*, 1922, **55**, [B], 2327—2331).—The action of hydrogen sulphide on a 1/1000-molar solution of lead chloride in water at 20° in the presence of hydrochloric acid and of neutral chlorides has been investigated. Precipitation of lead sulphide is completely inhibited by hydrochloric acid alone if present in a concentration of 1.4*N*. In the presence of increasing quantities of calcium, ammonium, or potass-

ium chlorides, decreasing concentrations of hydrochloric acid are sufficient to prevent precipitation entirely. In the presence of considerably smaller concentrations of hydrochloric acid, the precipitation of lead sulphide is incomplete. At higher temperatures, the effect of hydrochloric acid in the presence of neutral chlorides is still more marked. Similar results are obtained with cadmium salts, and apparently also with tin salts. The precipitation of lead sulphide is also hampered, although to a less degree, by the presence of perchloric acid. In this case, however, the addition of sodium perchlorate favours precipitation presumably by depressing the dissociation of the acid.

The precipitation of lead chromate in acetic acid solution is not quantitative in the presence of much chloride; in ammoniacal solution this is not the case.


H. W.

The Constitution of Basic Lead Salts. R. WEINLAND and RUDOLF STROH (*Ber.*, 1922, 55, [B], 2706—2718).—It has been assumed by Werner (A., 1908, ii, 42) that certain basic salts are to be regarded as aquo-salts in which the water molecules are replaced by metallic hydroxide, that is, as compounds containing a polynuclear kation; thus, atacamite, $\text{CuCl}_2 \cdot 3\text{Cu}(\text{OH})_2$, has been formulated $\left[\text{Cu} \left(\begin{smallmatrix} \text{OH} \\ \diagup \quad \diagdown \\ \text{OH} \end{smallmatrix} \right) \text{Cu} \right]_3 \text{Cl}_2$. The presence of such ions in several forms has been established by preparative substitution in the case of lead salts. Thus, in the basic perchlorates, $[\text{Pb}_2(\text{OH})_2](\text{ClO}_4)_2 \cdot 1.5\text{H}_2\text{O}$ (I) and $[\text{Pb}_3(\text{OH})_4](\text{ClO}_4)_2$ (II), the perchlorate residues could be replaced by other acid radicles. The stability of the metal-ol kations is not to be compared with that of the typical metal amines in aqueous solution, since the above basic perchlorates are to some extent decomposed into new basic salts when their solutions are diluted, so that, at any rate, a partial change has occurred in the complex. This instability is also shown by the fact that potassium ferricyanide and potassium cobalticyanide give salts with differently constituted lead kations from those obtained with alkali chlorides. The presence of such lead-ol complexes in basic lead acetates has also been established by the substitution method in the case of the basic lead acetate; one such ion is identical with that contained in the perchlorate (I).


Polynuclear lead kations are also present in the neutral lead acetate-perchlorates and lead acetate-nitrates (Weinland and Stroh, this vol., i, 981). They are distinguished from those contained in the basic salts only by containing acetato- instead of ol-bridges, thus: $[\text{Pb}_2(\text{OH})_2](\text{ClO}_4)_2 \cdot 1.5\text{H}_2\text{O}$ and $[\text{Pb}_2(\text{CH}_3\text{CO}_2)_2](\text{ClO}_4)_2 \cdot \text{H}_2\text{O}$. Application of these considerations to lead acetate itself causes the authors to regard it as the acetate of a polynuclear lead acetato-kation. In aqueous solution, undoubtedly more than one such kation is present, since the action of perchloric acid on lead acetate in neutral and acetic acid solution yields the perchlorates of four different lead acetato-kations. It has not yet been found possible to decide which kation is present in solid lead acetate, $\text{Pb}(\text{OAc})_2 \cdot 3\text{H}_2\text{O}$.

Basic lead perchlorate (I) is readily prepared from a solution of

lead monoxide in the calculated quantity of perchloric acid; its electrical conductivity indicates that three ions are present in its solutions. It is converted by double decomposition into basic lead chloride, $[\text{Pb}_3(\text{OH})_4]\text{Cl}_2$, a colourless powder which does not appear under the microscope to be crystalline; basic lead nitrate, $[\text{Pb}_3(\text{OH})_4](\text{NO}_3)_2$, *basic lead dithionate*, $[\text{Pb}_3(\text{OH})_4]\text{S}_2\text{O}_6$, lustrous leaflets and needles, and *basic lead nitroprusside*,




pale brownish-red cubes. With potassium ferricyanide and potassium cobalticyanide, it gives the analogously constituted *compounds*, $\text{Pb}_4(\text{OH})_2[\text{Fe}(\text{CN})_6](\text{ClO}_4)_3$, a brown powder, and




a colourless, heavy powder.

Basic lead perchlorate (II), tetrahedra and octahedra, is prepared by cooling a hot solution of perchloric acid which has been saturated with lead monoxide. It is converted by double decomposition into basic lead chloride, $[\text{Pb}_3(\text{OH})_4]\text{Cl}_2$, a yellowish-white powder; basic lead iodide, a yellow powder; basic lead nitrate, $[\text{Pb}_3(\text{OH})_4](\text{NO}_3)_2$, a heavy, white powder, and *basic lead dithionate*, $[\text{Pb}_3(\text{OH})_4]\text{S}_2\text{O}_6$, a colourless, crystalline precipitate. The following salts, also prepared by double decomposition, do not contain the original kation: *basic lead chromate*, $[\text{Pb}_2(\text{OH})_2]\text{CrO}_4$, a fine, red powder; *basic lead ferricyanide-perchlorate*, $\text{Pb}_3(\text{OH})_5[\text{Fe}(\text{CN})_6](\text{ClO}_4)_2$, a pale brown powder which explodes violently when heated; *basic lead ferricyanide*, $\text{Pb}_5(\text{OH})_7[\text{Fe}(\text{CN})_6]\cdot 3\cdot 5\text{H}_2\text{O}$, a yellow powder; *basic lead cobalticyanide-perchlorate*, $\text{Pb}_5(\text{OH})_5[\text{Co}(\text{CN})_6](\text{ClO}_4)_2\cdot 1\cdot 5\text{H}_2\text{O}$, a heavy, white powder; *basic lead nitroprusside-perchlorate*,



pale, brownish-red, crystalline aggregates.

Basic lead acetate is shown to contain the same kation as the perchlorate (I) by the fact that it gives by double decomposition the same basic chloride and dithionate. Concentrated solutions of the salt and potassium ferricyanide give the *salt*, $\text{Pb}_6(\text{OH})_7[\text{Fe}(\text{CN})_6](\text{CH}_3\cdot\text{CO}_2)_2\cdot\text{H}_2\text{O}$, brownish-red crystals, whereas dilute solutions yield a dark yellow, amorphous precipitate which fairly rapidly passes into the *compound*, $\text{Pb}_4(\text{OH})_5[\text{Fe}(\text{CN})_6]$, garnet-red prisms (the corresponding cobalt *compound*,



is prepared similarly). Treatment of the ferricyanide last mentioned with dilute acetic acid leads to the formation of the salt $\text{Pb}_3(\text{OAc})_5[\text{Fe}(\text{CN})_6]\cdot 2\text{H}_2\text{O}$, brownish-red needles. Solutions of lead acetate and sodium perchlorate give, under different conditions, the *salts*, $\text{Pb}(\text{OAc})_2\cdot\text{NaClO}_4$, rectangular prisms capped by pyramids at either end; $\text{Pb}_2(\text{OAc})_4\cdot\text{NaClO}_4\cdot 3\text{H}_2\text{O}$, colourless needles, and $\text{Pb}_4(\text{OH})_4(\text{ClO}_4)_4\cdot\text{NaClO}_4\cdot 3\text{H}_2\text{O}$, large, many-sided plates. H. W.

The Complex Cuprammoniates $\text{Cu}(\text{CN})_2\cdot\text{CuCN}\cdot 2\text{NH}_3$; $\text{Cu}(\text{CN})_2\cdot 4\text{CuCN}\cdot 4\text{NH}_3$; $\text{Cu}(\text{CNS})_2\cdot\text{Cu}(\text{CN})_2\cdot 5\text{Cu}\cdot 6\text{NH}_3$, and **the Analytical Methods for their Identification**. F. HALLE and K. HIRSCHKO (*Z. anorg. Chem.*, 1922, 122, 188—194).—The

above three new complex cuprammoniates have been prepared, and details are given of their analysis. W. T.

Critical Constants of Mercury. SOPHUS WEBER (*K. Danske Vid. Medd.*, 1920, 3, No. 4, 1—12; from *Chem. Zentr.*, 1922, iii, 27).—From theoretical considerations, the author derives the following critical constants for mercury: t_c 1450°; d_c 5.0; p_c 1042 atmospheres. The same value for p_c is given by extrapolation at 1450° with Knudsen's vapour pressure formula. The correlation of the Trouton constants with the critical data established by Cederberg is valid for the above values. The reduced vapour pressure curve for mercury approximately coincides with the curve for hydrogen. G. W. R.

Oxychlorides of Mercury. Equilibrium in the System Mercuric Chloride, Yellow Mercuric Oxide, and Water at 35°. SHIGERU TODA (*J. Chem. Soc. Japan*, 1922, 43, 312—320; *Mem. Coll. Sci. Kyōtō Imp. Univ.*, 1921, 4, 305—310).—The equilibrium in the system mercuric chloride, yellow mercuric oxide, and water at 35° has been studied. The solubility of mercuric chloride, 9.39 grams in 100 grams of water, and that of yellow mercuric oxide is 0.0014 gram in 100 grams of water. In the system investigated, two oxychlorides of mercury were found to exist: $\text{HgCl}_2 \cdot 2\text{HgO}$, purplish-scarlet needles, and $\text{HgCl}_2 \cdot 4\text{HgO}$, a brownish-yellow, amorphous substance. It is probable that the latter forms a solid solution with mercuric oxide. K. K.

The Rare-earth Magnesium Sulphide Phosphors. ERICH TIEDE and ARTHUR SCHLEEDE (*Ann. Physik*, 1922, [iv], 67, 573—580).—Phosphors containing cerium, lanthanum, didymium, praseodymium, neodymium, samarium, yttrium, gadolinium, erbium, ytterbium, scandium, thorium, and zirconium were prepared by the methods previously described (Tiede and Richter, this vol., ii, 215). They were all yellow in colour and, like the phosphors of the heavy metals, this colour was destroyed by pressure. The temperature relationships, and the spectrum of the phosphorescent light of the magnesium samarium phosphor were very similar to those of the calcium samarium phosphor of Hauer and Kowalski (A., 1914, ii, 320). Cerium, didymium, praseodymium, samarium, gadolinium, and scandium gave the best phosphors in the ordinary sense of the word, although the other metals gave strong phosphorescence in the cathode ray tube. The colour of the phosphorescence ranged from orange-yellow to pale green, and the duration of the emitted light was approximately twenty minutes. The spectra differed considerably from those of the phosphors of the heavy metals, being composed of intense sharp lines and bands. The exciting light, in all cases, lay in the ultra-violet. W. E. G.

The Separation of the Rare Earths by Basic Precipitation. IV. WILHELM PRANDTL and JOHANNA RAUCHENBERGER (*Z. anorg. Chem.*, 1922, 122, 311—314).—The influence of cadmium nitrate on the equilibrium between the nitrates of the rare earths and ammonia in ammonium nitrate solutions of various concen-

trations has been investigated. The conditions were the same as in previous experiments (cf. this vol., ii, 298). The presence of cadmium nitrate increased considerably the amount of rare earth kept in solution, especially lanthanum, this increase being much more marked than in the case of zinc nitrate. W. T.

Rare Earths. III. Atomic Weight of Lanthanum. B. S. HOPKINS and F. H. DRIGGS (*J. Amer. Chem. Soc.*, 1922, **44**, 1927—1929).—The atomic weight of lanthanum has been determined by a comparison of lanthanum chloride with silver. The material used was the less soluble portion of the magnesium rare-earth nitrates from which the whole of the cerium had been removed by James and Pratt's method (A., 1911, ii, 935). The sparingly soluble fractions containing lanthanum and praseodymium were fractionated further as double ammonium nitrates until the first twenty fractions failed to show the faintest trace of the praseodymium absorption spectrum. Fractions 7 and 8 of this series were precipitated with hot dilute oxalic acid solution, the precipitate filtered, washed, and ignited to oxide which was dissolved in nitric acid, and the solution treated with ammonia gas to precipitate the hydroxide. The alternate precipitation as oxalate and hydroxide was repeated four times, the last two taking place in conductivity water. The final oxide was suspended in conductivity water and treated with hydrogen chloride in a quartz flask. This material showed no spectroscopic lines of any of the related elements. The solution of lanthanum chloride was dried in a quartz flask in a current of air and hydrogen chloride until crystallisation commenced, the air current was shut off, and hydrogen chloride alone passed in until, at 85°, the first five molecules of water of crystallisation had come off, then the temperature was raised to 125°, when the final molecule of water commences to come off. As soon as this evolution was complete, the temperature was raised to 325°, and kept there for an hour, after which the salt was rapidly melted over a bunsen flame and then allowed to cool. The hydrogen chloride was displaced by dry air and the material weighed. The chloride was treated with silver solution and exact equivalence between the two substances determined nephelometrically. Using the data: density, $\text{LaCl}_3 = 3.947$, $\text{Ag} = 10.5$; atomic weights, $\text{Ag} = 107.88$, $\text{Cl} = 35.457$, it was found that 1.02450 grams LaCl_3 is equivalent to 1.35189 grams of silver, whence the mean atomic weight of lanthanum (10 expts.) = 138.89; the extreme values being 138.84 and 138.97. This value is very close to 138.91, the value found by Baxter, Tani, and Chapin (A., 1921, ii, 454), and seems to indicate that the value 139.0 adopted by the International Atomic Weight committee is slightly too high. J. F. S.

The Separation of the Rare Earths by Basic Precipitation. III. The Quantitative Separation of Cerium from the Other Earths. WILHELM PRANDTL and JOSEPH LÖSCH (*Z. anorg. Chem.*, 1922, **122**, 159—166).—The separation is based on the fact that ceric hydroxide is a weak base and its salts are highly hydrolysed. Cerous nitrate mixed with the nitrates of the other rare

earths was oxidised by trinitratocobalttriammine in acid solution with quantitative precipitation of the ceric hydroxide, $\text{Ce}(\text{NO}_3)_3 + [\text{Co}(\text{NH}_3)_3(\text{NO}_3)_3] + 4\text{H}_2\text{O} = \text{Ce}(\text{OH})_4 + \text{Co}(\text{NO}_3)_2 + 3\text{NH}_4\text{NO}_3 + \text{HNO}_3$. The formation of free acid prevents the precipitation of the other rare earths.

W. T.

Intermetallic Actions. The System Aluminium-Arsenic. QASIM ALI MANSURI (T., 1922, 121, 2272—2277).

The Constitution of Aluminates. JAROSLAV HEYROVSKY (*Chem. News*, 1922, 125, 198—199).—In order to investigate further the monobasicity of "aluminic acid" (cf. T., 1920, 117, 1013), cryoscopic and conductivity determinations were made on solutions of the alkaline-earth hydroxides when gradually saturated with aluminium hydroxide in the nascent condition, generated by the action of the solution on amalgamated aluminium. The regular fall in conductivity observed and the slight changes in the lowering of the freezing point show that only univalent aluminate ions exist. In contact with crystalline aluminium hydroxide, all the solutions are about 50% hydrolysed; with nascent aluminium hydroxide, the hydrolysis is only 1—1.5%. The process by which aluminium hydroxide dissolves in alkali is undoubtedly by complex-ion formation, $\text{Al}(\text{OH})_3 + \text{OH}' \rightarrow \text{Al}(\text{OH})_4'$, and not, as is often stated, by sending into solution one, two, or three of its hydrogen atoms as hydrions. The hydrolysis of aluminates is therefore, on this view, not due to scarcity of hydrions, but to the incompleteness of the complex formation.

G. F. M.

The Properties of Potassium Permanganate. G. FESTER and G. BRUDE (*Z. angew. Chem.*, 1922, 35, 527).—Crystals of pure potassium permanganate prepared in the dark and in air free from carbon dioxide and organic matter, are brown and have a bronze, metallic lustre. The finely powdered crystals, however, show a dark violet colour. When exposed to subdued daylight and ordinary air, the surfaces of the crystals develop a violet colour in a few hours and a steel blue lustre after two days. This change is confined to the surface layer, which acts as a protective layer to the remainder of the crystal. The change is particularly rapid in an atmosphere containing mineral acid. The crystals of potassium permanganate should be described as bipyramidal rhombs rather than as prisms. Potassium permanganate oxidises paper and most other organic substances, so that the statement that it is without action on litmus paper is untrue.

H. C. R.

Mechanism of the Reduction of Permanganate and its Physico-chemical Basis. IV. Reduction of Permanganate by Formate in Alkaline Solutions. JOSEF HOLLUTA (*Z. physikal. Chem.*, 1922, 102, 276—297; cf. this vol., ii, 448, 628, 700).—A continuation of previous work. It is shown that the reduction of permanganate by formates in alkaline solution takes place in two stages which are sharply separated from one another. In the first stage, the reduction of permanganate to manganate takes place, and this is reduced to an alkali manganite in the second

stage. The first stage takes place according to the equation $2\text{MnO}_4' + \text{HCO}_2' + 3\text{OH}' = 2\text{MnO}_4'' + \text{CO}_3'' + 2\text{H}_2\text{O}$, and it is shown that a disturbance occurs in the reaction at a point immediately before the completion of the reduction to manganate due to the commencement of the reduction of the manganate. The first stage is shown to be a second order reaction, inasmuch as one molecule of permanganate apparently oxidises one molecule of formate. To explain the behaviour of the manganate and the low order of the reaction it is assumed that it is not the permanganate itself which oxidises the formate, but atomic oxygen which is split off from the permanganate under the influence of the hydroxyl ions. It is further assumed that the hydroxyl ions, through the formation of an additive compound with the permanganate, act as oxygen carriers. It is shown that the above assumptions can be justified on the basis of the experiments and calculations of Sackur and Taegener (A., 1912, ii, 916). The temperature coefficient of this reaction is found to be 2.06. From the potential measurements of Sackur and Taegener (*loc. cit.*), the equilibrium constant of the reaction $2\text{MnO}_4'' + 2\text{H}_2\text{O} \rightleftharpoons 2\text{MnO}_2 + 4\text{OH}' + \text{O}_2$ is calculated to be 53.1 for 18–20°. It is shown that the presence of hydroxyl ions accelerates the reaction, the cause of which being in all probability that the hydroxyl ions affect the equilibrium pressure at which oxygen is separated from the permanganate hydroxyl ion complex.

J. F. S.

The Diffusion of Carbon in Metals and in Mixed Crystals of Iron. G. TAMMANN and K. SCHÖNERT (*Z. anorg. Chem.*, 1922, 122, 27–43).—Carbon was found to diffuse into iron at 750°, whereas no diffusion into the other metals took place at 980°. In mixed crystals of iron with tungsten, molybdenum, nickel, cobalt, and manganese containing a large amount of iron, the diffusion was found to increase with the addition of the second element until a maximum was reached; the diffusion then diminished. In the cases of tungsten and molybdenum, the maxima corresponded with 0.0306 molar concentration, and the diffusion increased by 0.166; with nickel, cobalt, and manganese as added metals, the maxima required a molar concentration of 0.114, and the increase in diffusion was 0.057. The diffusion coefficients found for different pieces of iron were found to vary; this was due to the foreign matter between the crystallites.

W. T.

Photochemical Examination of the Reaction between Ferric Salts and Iodides. NOBUJI SASAKI (*Mem. Coll. Sci. Kyōto Imp. Univ.*, 1922, 5, 315–327).—The reaction between ferric salts and iodides is retarded by exposure to sunlight or bright artificial light, whilst that between ferrous salts and free iodine is accelerated. These photochemical reactions are discussed, and are shown to follow a simple law.

W. P. S.

System $\text{Fe}_2\text{O}_3\text{--SO}_3\text{--H}_2\text{O}$. E. POSNJAK and H. E. MERWIN (*J. Amer. Chem. Soc.*, 1922, 44, 1965–1994).—The system ferric oxide-sulphur trioxide-water has been investigated over the

temperature range 50—200°, by heating various mixtures of the components at various temperatures and analysing the phases produced. The following crystalline phases have been obtained: Fe_2O_3 ; $\text{Fe}_2\text{O}_3 \cdot \text{H}_2\text{O}$; $3\text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 9\text{H}_2\text{O}$; $\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot \text{H}_2\text{O}$; $\text{Fe}_2\text{O}_3 \cdot 2\text{SO}_3 \cdot 5\text{H}_2\text{O}$; $2\text{Fe}_2\text{O}_3 \cdot 5\text{SO}_3 \cdot 17\text{H}_2\text{O}$; $\text{Fe}_2\text{O}_3 \cdot \text{SO}_3$ (2 forms); $\text{Fe}_2\text{O}_3 \cdot 3\text{SO}_3 \cdot 6\text{H}_2\text{O}$; $\text{Fe}_2\text{O}_3 \cdot 3\text{SO}_3 \cdot 7\text{H}_2\text{O}$; $\text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 3\text{H}_2\text{O}$, and $\text{Fe}_2\text{O}_3 \cdot 4\text{SO}_3 \cdot 9\text{H}_2\text{O}$. The conditions under which these substances are formed, the ranges of their stability, and some of their crystallographic and optical properties have been determined. J. F. S.

Some Poly-acids of the Elements of the Sulphur Group. JULIUS MEYER and VALENTIN STATECZNY (*Z. anorg. Chem.*, 1922, **122**, 1—21).—The poly-acids formed by elements of the sulphur group are formulated according to the co-ordination system with a co-ordination number equal to four. The authors have succeeded in preparing the following heteropoly-acids and some of their salts: chromosulphuric acid, $\text{H}_2[\text{CrO}_3(\text{SO}_4)]$, and its potassium, sodium, ammonium, barium, strontium, and calcium salts; chromoselenic acid, $\text{H}_2[\text{CrO}_3(\text{SeO}_4)]$, together with its potassium and barium salts. Methods of preparation are given in detail. The solubilities of chromic anhydride in sulphuric and selenic acids of various concentrations were determined and a method was given for the preparation of the pure anhydride. The existence of the sulphoselenic acids, $\text{H}_2[\text{SeO}_3(\text{SO}_4)]$ and $\text{H}_2[\text{SeO}_2(\text{SO}_4)_2]$, was indicated by thermal analysis. Sulphomolybdic acid, $\text{H}_2[\text{MoO}(\text{SO}_4)_2(\text{MoO}_4)]$, was also prepared, but the corresponding selenic acid could not be isolated. Indications were obtained of the existence of chromyl-sulphuryl chloride, CuSO_4Cl_2 . W. T.

Experimental Attempts to Decompose Tungsten at High Temperatures. GERALD L. WENDT and CLARENCE E. IRION (*J. Amer. Chem. Soc.*, 1922, **44**, 1887—1894).—Attempts have been made to decompose tungsten by rapidly discharging a condenser holding 30,000—45,000 volts through a tungsten wire 4 cm. long and 0.035 mm. diam., either in a vacuum or in carbon dioxide at 1 atm. pressure. In this way, momentary temperatures above 20,000° were produced in the wires which were entirely dissipated, no smoke, dust or solid residue being found after the explosion. Abundant quantities of gas were present after the explosions in a vacuum, and a spectroscopic examination of this showed the strong yellow helium line, and two red, one bright blue, and one pale violet line were also observed but have not been identified. The gas from the explosions in carbon dioxide was collected and measured, and it was found that about the same weight (mean 0.713 mg.) of tungsten yields in different experiments from 3.62 to 0.30 c.c. of gas unabsorbable by potassium hydroxide. It is pointed out that if the whole of the tungsten had been decomposed into helium the volume of gas would have been 4.0 c.c. J. F. S.

The Hydrates of Tungsten Trioxide. GUSTAV F. HÜTTIG and BRUNO KURRE (*Z. anorg. Chem.*, 1922, **122**, 44—50).—The

existence of hydrates of tungsten trioxide was investigated by the Hüttig tensi-eudiometer (A., 1921, ii, 195). Solutions of the pure white and yellow acids were employed, and the process of dehydration was followed. No compound was indicated in the case of the white acid, the temperature-composition curve indicating adsorption as the water content changes gradually. The dehydration of the yellow acid showed the existence of a monohydrate $\text{WO}_3 \cdot \text{H}_2\text{O}$. The authors found that at temperatures above 188° and below 76° the white acid is the more stable, whereas between 76° and 188° the yellow variety is the stable form. W. T.

Sodium Tungstates. I. EDGAR F. SMITH (*J. Amer. Chem. Soc.*, 1922, **44**, 2027—2036).—Tungstates in which the basic and acidic oxides are present in the ratio 4 : 10 constitute a very definite series of salts. The sodium compound, $4\text{Na}_2\text{O} \cdot 10\text{WO}_3 \cdot 23\text{H}_2\text{O}$, can be prepared by the method of Forcher and Gibbs, by passing carbon dioxide during several days through an aqueous solution of normal sodium tungstate or by the gradual addition of formic acid, until the action is distinctly acid, to a solution of normal sodium tungstate (100 grams) in water (100 c.c.). A mixture of the salts $4\text{Na}_2\text{O} \cdot 10\text{WO}_3 \cdot 23\text{H}_2\text{O}$ and $5\text{Na}_2\text{O} \cdot 12\text{WO}_3 \cdot 28\text{H}_2\text{O}$ is produced by the action of glacial acetic acid on a solution of sodium tungstate; the product, $9\text{Na}_2\text{O} \cdot 22\text{WO}_3 \cdot 51\text{H}_2\text{O}$, appears to be a mixture of the two salts described above. The salt $4\text{Na}_2\text{O} \cdot 10\text{WO}_3 \cdot 23\text{H}_2\text{O}$ forms monoclinic crystals which effloresce rapidly in dry air. It is soluble in water to the extent of 19 parts in 100 parts at atmospheric temperature; and has d 4.3. The following melting points are recorded for the various sodium tungstates: $\text{Na}_2\text{O} \cdot \text{WO}_3$, 665° ; $5\text{Na}_2\text{O} \cdot 12\text{WO}_3$, 705.8° ; $4\text{Na}_2\text{O} \cdot 10\text{WO}_3$, 680.8° ; $9\text{Na}_2\text{O} \cdot 22\text{WO}_3$, 683.3° ; $\text{Na}_2\text{O} \cdot 4\text{WO}_3$, 706° . Addition of an excess of the respective metallic chlorides to a boiling aqueous solution of the salt $4\text{Na}_2\text{O} \cdot 10\text{WO}_3 \cdot 23\text{H}_2\text{O}$ leads to the formation of the following compounds: *calcium* salt, $4\text{CaO} \cdot 10\text{WO}_3 \cdot 25\text{H}_2\text{O}$, a white, granular powder which does not melt when heated to intense redness; the *barium* salt, $4\text{BaO} \cdot 10\text{WO}_3 \cdot 22\text{H}_2\text{O}$, and the *strontium* salt, $4\text{SrO} \cdot 10\text{WO}_3 \cdot 26\text{H}_2\text{O}$, white, insoluble, granular powders; the *nickel* salt, $4\text{NiO} \cdot 10\text{WO}_3 \cdot 34\text{H}_2\text{O}$, a greenish-white powder; the *cobalt* salt, $4\text{CoO} \cdot 10\text{WO}_3 \cdot 35\text{H}_2\text{O}$, pink granules; the *manganese* salt, $4\text{MnO} \cdot 10\text{WO}_3 \cdot 30\text{H}_2\text{O}$, a colourless, granular powder. The action of a solution of the sodium salt, $4\text{Na}_2\text{O} \cdot 10\text{WO}_3 \cdot 23\text{H}_2\text{O}$, towards solutions of salts of many inorganic and organic bases has been examined qualitatively. The sodium salt loses the whole of its water of crystallisation when it is strongly ignited, but when dried at 100° until constant in weight it retains six molecular proportions. It therefore appears that it may be formulated definitely as an acid salt, $4(\text{Na}_2\text{O} \cdot \text{WO}_3) \cdot 6(\text{H}_2\text{O} \cdot \text{WO}_3) \cdot 17\text{H}_2\text{O}$, and similar considerations are extended to the tungstates of other series. Confirmation of its acidic character is found in the ready neutralisation of its solutions with *N*-sodium carbonate solution and also in conductivity experiments during its neutralisation with barium hydroxide solution. H. W.

The Binary Systems Na_2SiO_3 - Na_2WO_4 , K_2SiO_3 - K_2WO_4 , and Na_2WO_4 - K_2WO_4 . J. A. M. VAN LIEMPT (*Z. anorg. Chem.*, 1922, **122**, 175—180).—Klooster (A., 1911, ii, 111) examined the system Na_2WO_4 - Na_2SiO_3 and found no indications of a compound; the liquid salts were found to be immiscible. The present author has examined the viscosity and conductivity of aqueous solutions of the mixture. Curves representing the results showed no maxima or minima. The system K_2WO_4 - K_2SiO_3 gave similar results. The equilibrium curve for the system K_2WO_4 - Na_2WO_4 was obtained by the cooling method; a minimum freezing point was found for a mixture containing 80% of the sodium and 20% of the potassium salt. W. T.

Preparation and Reactions of Stannous Oxide and Stannous Hydroxides. FRANK WARD BURY and JAMES RIDDICK PARTINGTON (T., 1922, **121**, 1998—2004).

Reduction of Thorium Oxide by Metallic Tungsten. THE RESEARCH STAFF OF THE GENERAL ELECTRIC CO., LTD., LONDON (work conducted by COLIN JAMES SMITHELLS) (T., 1922, **121**, 2236—2238).

Germanium Hydride. JOHN H. MÜLLER and NICOL H. SMITH (*J. Amer. Chem. Soc.*, 1922, **44**, 1909—1918).—Germanium may be detected by a modification of Marsh's test in quantities as small as 0.00006 gram of the metal. The delicacy of the hydride relation is greatly increased by use of an alkaline generator as source of nascent hydrogen. Aluminium and dilute potassium hydroxide solution serve best for the formation of large quantities of germanium hydride, whilst sodium amalgam is best for small quantities. A modification of the usual form of Marsh apparatus is described which permits of the formation of a richer mixture of the hydride than has hitherto been prepared and still further increases the delicacy of the test. The apparatus is arranged as follows:—an electrolytic hydrogen generator, using 15% sodium hydroxide as electrolyte and having nickel electrodes, a washing tower containing a solution of a silver salt, a sulphuric acid drying tower, and a trap for exit of excess of hydrogen from the generator. The gas prepared and purified in the above train passes into the germanium hydride generator, which consists of a small wash-bottle the inner tube of which is cut off at a height 5 cm. above the bottom of the bottle; the generator is fitted with a funnel and stopcock through which the sodium amalgam and germanium solution may be added after the whole system has been filled with electrolytic hydrogen. To the generator is attached a small drying tube and to the end of this a constricted tube in which the germanium mirror may be formed. A rich mixture of germanium hydride and hydrogen is obtained by placing dry sodium amalgam in the generator and expelling the air by means of electrolytic hydrogen. The hydrogen stream is then interrupted and an aqueous solution of the germanium salt added and washed in with water. The germanium hydride is then allowed to form in the

closed apparatus, and when the action is over swept through the heated constricted tube by means of electrolytic hydrogen, when a mirror of germanium is immediately formed. The decomposition temperature of germanium hydride lies between 340° and 360° , and the deposition of the mirror is best obtained below a red heat. The reverse action, involving the loss of germanium in hydrogen, has been quantitatively investigated and shows that at $750\text{--}800^{\circ}$ 1.41855 grams of germanium loses 0.00647 gram when in a stream of hydrogen for thirty-eight hours. The dissimilarity of films of arsenic and germanium as obtained by the Marsh test is shown by a spectroscopic examination of the light transmitted by the films, and the thickness of the films has been determined. A study of the composition of the substance described by Voegelen (A., 1902, ii, 401) shows this substance to be a mixture which may contain little or much silver germanide Ag_4Ge along with metallic silver, depending on the method used in the preparation of the hydride. The precipitate contains much more germanide if the hydride is generated by the action of potassium hydroxide on aluminium and passed slowly through a solution of a silver salt, whilst if the formation of the hydride is brought about by means of sodium amalgam or the gas passed rapidly through the silver salt, the precipitate contains little germanide and the solution contains germanic acid. Evidence is put forward which indicates the presence of at least one hydride other than the tetrahydride, but the presence of this hydride has not been finally established. The richest mixture of hydrogen and germanium hydride examined contained 0.176% of the hydride. J. F. S.

Germanium Hydride. FRITZ PANETH and EDGAR SCHMIDT-HEBBEL (*Ber.*, 1922, 55, [B], 2615—2622).—Germanium hydride has been described previously by Voegelen, but its composition has not been settled definitely. The substance has now been re-examined, and since the authors had only a very limited amount of material at their disposal they have found it most convenient to prepare the gas by the action of zinc or, preferably, magnesium on a solution of germanium in the presence of 3—4*N*-sulphuric acid; the yield varies between 0.2 and 1.5%. The gas evolved from the solution is dried by passage over granulated calcium chloride and phosphoric oxide and subsequently led through a Landsiedl apparatus filled with potassium hydroxide solution (50%), a U-tube immersed in pentane cooled to -110° and finally through three U-tubes cooled in liquid air. The almost invisible white condensate in the last three tubes is analysed by the method used previously for antimony and tin hydrides (this vol., ii, 383). It is thus shown that the atomic ratio of hydrogen to germanium is exactly 4 : 1. The accuracy of the analyses excludes the possibility of the presence of more than the merest traces of possible hydrides, such as Ge_2H_6 , and the existence of these is the more improbable, since, under the conditions of the condensation, they must have become concentrated in the condensate; the pentane tube does not retain any germanium compounds, and about 15%

of germanium hydride passes uncondensed through the apparatus. It is shown that the molecular formula of germanium hydride is GeH_4 .

Germanium hydride is unusually stable, being unaffected by potassium hydroxide solution (50%) or phosphoric oxide, and requiring a relatively very high temperature to cause its certain decomposition. The colours of the germanium mirrors are extraordinarily varied and in this respect the metal is differentiated from all others which yield hydrides decomposable by heat. In doubtful cases, confirmation may be obtained by oxidising the mirror to colourless germanium dioxide by heating it in a current of oxygen and subsequently reducing it in hydrogen to copper-coloured germanium, which can then be dissolved in sodium hypochlorite. Arsenic exhibits greater solubility in the reagent, but its oxide would be volatilised completely under the experimental conditions.

H. W.

The System $\text{Bi}_2\text{O}_3\text{-PbO}$. L. BELLADEN (*Gazzetta*, 1922, 52, ii, 160—164).—The occurrence in nature of a series of minerals constituted of double sulphides of bismuth and lead which may mostly be regarded as derivatives of hypothetical ortho-, meta-, and pyro-sulphobismuthous acids renders possible the existence of similar compounds containing oxygen in place of the sulphur atoms. The author has investigated the fusion diagram of the system $\text{Bi}_2\text{O}_3\text{:PbO}$, the results obtained indicating the formation of the compounds: (1) $4\text{Bi}_2\text{O}_3\text{,PbO}$, m. p. 695° (decomp.); (2) $3\text{Bi}_2\text{O}_3\text{,2PbO}$, m. p. 686° , stable when fused; (3) $\text{Bi}_2\text{O}_3\text{,2PbO}$, m. p. 625° , stable on fusion. These indications are fully confirmed by micrographic examination.

Litharge has m. p. 870° and bismuth trioxide, m. p. 817° .

T. H. P.

Polonium Hydride. II. FRITZ PANETH and ADOLF JOHANNSEN (*Ber.*, 1922, 55, [B], 2622—2637).—Polonium hydride closely resembles bismuth hydride in the order of magnitude of its yields by all methods of preparation, its great instability during the condensation of minimal amounts, and its ready decomposability by chemical reagents.

The preparation of polonium hydride has been affected previously by the action of dilute acids on an electrolytic deposit of polonium on magnesium (A., 1919, ii, 76). Since it is difficult by this method to secure highly activated magnesium foil in uniformly good yield, a process has been evolved which depends on the deposition of polonium on magnesium powder by distillation. For this purpose, the polonium is first deposited electrolytically from a lead nitrate solution containing radium-*D* on the central portion of a thin piece of platinum foil. The latter is now suspended in a glass dish, the bottom and sides of which are covered with magnesium powder; the apparatus is brought into an evacuated desiccator and the foil is raised electrically to a bright yellow heat. The volatilised polonium is deposited mainly on the powder, which, after being carefully mixed, gives a preparation of uniform activity.

(A method for the recovery of polonium from its residual solutions and again precipitating it on magnesium is described in detail.) The yield of polonium hydride obtained by the solution of the activated powder in dilute acid is practically the same as from the activated foil, and lies intermediate between those of lead and bismuth hydrides. The formation of the hydride cannot be ascribed to the decomposition of a compound of magnesium and polonium, since the gas is also obtained by adding magnesium powder to a solution containing polonium. The method of "spark electrolysis" is also applicable to the production of polonium hydride, but the evolution of the gas by this method is too irregular for use for the present purposes.

Polonium hydride which has been condensed at the temperature of liquid air is very extensively decomposed during re-evaporation. The action appears to depend to some extent on the accompanying gases; thus, for example, it has not been found possible to re-volatilise polonium hydride which has been prepared by "spark electrolysis." The condensation and partial re-volatilisation of polonium hydride affords the most conclusive evidence of the gaseous nature of the compound.

Phosphoric oxide causes very extensive decomposition of the hydrides of polonium and thorium-C, whereas calcium chloride has a less strongly marked action; the latter, however, can only be used for the desiccation of polonium hydride when a very high initial activity can be used. Under the experimental conditions adopted, unboiled distilled water only allows 3% of polonium hydride to pass unchanged, whereas 50% of the gas passes through air-free, distilled water. *N/10*-Sodium hydroxide and *N/10*-silver nitrate solution cause very extensive decomposition of polonium hydride.

The rates of spontaneous decomposition of polonium and bismuth hydrides have been examined; the former is decomposed to the extent of 50% after four minutes, the latter after about twenty-five minutes. A more exact calculation is not warranted by the accuracy of the experiments, which suffer under the disadvantages that the gases cannot be dried. It is, however, established that moist polonium hydride is considerably more easily decomposed than bismuth hydride.

H. W.

The Recrystallisation of Pure Mechanically Unworked Gold, obtained by Solidification from the Fused Mass. W. FRAENKEL (*Z. anorg. Chem.*, 1922, **122**, 295—298).—Pure mechanically unworked gold showed no change in size or form of crystals on heating at 1000—1040°. The purity could not be guaranteed to be such that the crystallites might not be surrounded by films of impurity (cf. Tammann, A., 1921, ii, 172).

W. T.

Mineralogical Chemistry.

Colour of Fluorites. TOKUTARO SAKAO and MITSUIE HIROSE (*Mem. Coll. Sci. Kyōto*, 1921, 4, 349—350).—The colour of fluorite crystals has been attributed by Blount and Sequeira (T., 1919, 15, 705) to the presence of a small amount of organic matter. Employing the ultra-microscope, the authors were unable to detect the presence of such matter in a state of colloidal suspension in five varieties of coloured fluorite crystals. J. S. G. T.

Analytical Chemistry.

Qualitative Chemical Analysis. V. MACRI (*Boll. Chim. Farm.*, 1922, 61, 417—418).—By means of the following scheme, the use of sulphur compounds is avoided. The hydrochloric acid solution is placed, together with a rod of zinc, in a flask fitted with a gas delivery tube, the latter being heated so as to make a Marsh apparatus; the flask is eventually heated. The liquid is filtered, the precipitate containing silver, mercury, lead, bismuth, copper, cadmium, platinum, gold, tin, antimony, and arsenic. The filtrate is heated to boiling with zinc acetate, aluminium, iron, and chromium being precipitated. The filtrate from this is heated with hydrogen peroxide, which precipitates manganese. The filtrate from the latter is heated and treated with zinc oxide in presence of chlorine, nickel and cobalt being thus precipitated. Reference is made to various circumstances which prevent the subdivision into groups from being rigorously exact. [Cf. *J. Soc. Chem. Ind.*, 1922, 839A.] T. H. P.

Quantitative Microanalysis. MAURICE NICLOUX and GEORGES WELTER (*Chim. et Ind.*, 1922, 8, 269—272).—A summary of Pregl's methods of microanalysis for use with which a Longue's aperiodic microbalance is recommended. A micro-method for the estimation of urea in blood is outlined (cf. this vol., ii, 170). W. G.

Use of Resorcinol in Qualitative Inorganic Analysis. LAVOYE (*J. Pharm. Belg.*, 1921, 3, 889—890; from *Chem. Zentr.*, 1922, ii, 1154).—One c.c. of a 10% solution of resorcinol mixed with 2 c.c. of 10% ammonia solution gives characteristic colorations when added to 2 c.c. of different metallic salt solutions. The following colorations are obtained. Zinc salts, a yellowish-green colour, changing to deep blue; cadmium salts, a less intense blue colour; manganese salts, in the presence of ammonium salts, a bluish-green colour; nickel salts, bluish-green; copper salts, reddish-violet, changing to bluish-violet. With salts of the latter two metals the solutions should be so dilute that ammonia alone

gives no coloration. Platinum salts give a garnet-red colour; mercury salts give no colour, but the solution on evaporation gives crystalline needles. All the colorations mentioned change to red on the addition of acid. The reactions are sensitive with very small amounts of the metals in question. G. W. R.

Analysis of Minerals by the Röntgen Spectrograph. ASSAR HADDING (*Z. anorg. Chem.*, 1922, **122**, 195—200).—The suitability of this method is shown by the fact that monazite, which contains twelve oxides, only gives twelve lines. Quantities of substances are indicated roughly by the strength of the lines. Very small amounts of substances are shown by weak lines and the author suspects the presence of the unknown element of atomic number 61 in "fluocerite." W. T.

Use of Mixed Indicators. A. COHEN (*J. Amer. Chem. Soc.*, 1922, **44**, 1851—1857).—It is shown that differences of subjective colour are enhanced when two indicators are partially transformed at the same hydrogen-ion concentration. The colorimetric determination of hydrogen-ion concentration can, therefore, be rendered more precise by the use of suitable mixed indicators and in certain titrations sharper end-points can be obtained, particularly in coloured liquids. Thus with the indicator bromothymol-blue the value P_H 6.8 is indicated by a pure green colour, but in a yellow medium the colour is yellowish-green and difficult to observe; if, however, an indicator is added which is violet at this concentration, then the total effect of the mixed indicators will be a green end-point. This may be achieved by adding equal quantities of bromothymol-blue and bromocresol-purple. Several cases of the use of mixed indicators are described. J. F. S.

Modification of the Chlorometric Method of Hayem and Winter. J. MALGOYRE (*Bull. Soc. pharm. Bordeaux*, 1922, **60**, 59—64).—Gastric juice is treated with a reagent prepared by dissolving 2 grams of chromic acid in 100 c.c. of 95% ethyl alcohol to remove proteins, and then titrated directly. The free and total hydrochloric acid are estimated by the Topfer-Linossier method, and the combined acid found by difference. The total chlorine is estimated by neutralising 10 c.c. of gastric juice with a saturated solution of sodium carbonate, adding 10 c.c. of the chromic acid reagent, filtering, and titrating with 0.1*N*-silver nitrate. The fixed chlorine is equal to the difference between the total chlorine and total acidity. CHEMICAL ABSTRACTS.

Reductions with Cadmium and Lead in Volumetric Analysis. III. W. D. TREADWELL (*Helv. Chim. Acta*, 1922, **5**, 806—818).—[With P. HRISTIE, L. EGGER, and P. STURZENEGGER.] A warm solution of a chlorate in presence of sulphuric acid can be reduced quantitatively to chloride by passing it through a cadmium reduction tube (A., 1921, ii, 523), and the chloride can then be titrated by the Volhard process. Since perchlorates are not reduced under these conditions, chlorate can be determined in presence of perchlorate by this method. Perchlorate can be reduced to chloride

by boiling in sulphuric acid solution (about 10 c.c. of concentrated acid to 50 c.c. of 0.1N-chlorate solution) in presence of titanium sulphate for one-half to one hour with finely divided cadmium. The cadmium is only attacked to a slight extent by the acid, and a small amount of cadmium sulphide is formed which, however, does not interfere with the estimation of perchlorate. After cooling, the solution is diluted, the titanous sulphate present is carefully oxidised with permanganate and the chloride is titrated either by the Volhard process or electrometrically.

[With A. FREULER and A. WEBER.]—A reduction method for the estimation of columbium was described by Metzger and Taylor (A., 1909, ii, 702), depending on the stabilising effect of succinic acid on a solution of columbic acid in sulphuric acid. The solution was reduced with amalgamated zinc to the constant stage $\text{Cb}_2\text{O}_3 \cdot 107$. It has now been found that the succinic acid solution of columbic acid prepared in this manner is far from stable. In course of a few days it becomes opalescent and its reducibility gradually becomes less. The reducibility also becomes less when a solution of columbic acid in sulphuric acid is diluted, confirming the view that the reducibility depends on the dispersity of the colloidal acid. Consistent results were obtained when the reduction was carried out in a cadmium reduction tube in presence of ammonium molybdate or vanadate or of titanium sulphate, the subsequent titration being carried out with permanganate. The most stable solutions of columbic acid were obtained by dissolving the fluoride in concentrated hydrochloric acid, and consistent results were obtained when such solutions were reduced in a cadmium tube and titrated electrometrically with ferric chloride.

[With R. F. EDELMANN.]—A concentrated hydrochloric acid solution of a molybdate can be reduced quantitatively to the Mo^{III} stage by finely divided lead. Lead in a suitable form for a reduction tube was obtained by immersing a zinc rod in an acidified solution of lead acetate. The crystals of lead must be carefully washed with alcohol and ether, not pressed together. Stannic chloride was also reduced quantitatively to stannous chloride by means of lead. The oxidation titrations were carried out with permanganate.
E. H. R.

Chlorination of Mixed Silver Haloids in Gooch Crucibles.

M. G. MELLON and J. C. SIEGESMUND (*Proc. Indiana Acad. Sci.*, 1921, 197—199).—The crucible containing the silver haloids is placed inside a larger crucible which is covered with a watch glass with a hole in the centre. Chlorine is passed into the crucible through a tube extending to within 1 cm. of the bottom, the outer crucible being gently heated meanwhile.

CHEMICAL ABSTRACTS.

Electrometric Titrations with Lead Nitrate. I. M. KOLTHOFF (*Z. anal. Chem.*, 1922, 61, 369—377).—Electrometric titration of iodide solutions, of not too low concentration, with lead nitrate solution yields trustworthy results, even in the presence of chlorides and bromides; the method is applicable to the estim-

ation of ferrocyanides and also of sulphates provided that mineral acids, ferric salts, and aluminium salts are not present. The titration of pyrophosphates with lead nitrate solution gives unsatisfactory results, but the method may be used in the case of a few organic salts such as oxalates, tartrates, and succinates.

W. P. S.

A New Iodometric Method Based on the Formation and the Estimation of Cyanogen Iodide. RUDOLF LANG (*Z. anorg. Chem.*, 1922, **122**, 332—348).—A volumetric method for the estimation of iodide is given, based on the fact that an iodide in hydrochloric or sulphuric acid solution is oxidised by an iodate or permanganate in the presence of hydrocyanic acid to cyanogen iodide, the end-point being indicated by starch solution. This can be controlled by estimating the cyanogen iodide with thio-sulphate $\text{ICN} + 2\text{S}_2\text{O}_3^{''} + \text{H}^+ = \text{I}^+ + \text{HCN} + \text{S}_4\text{O}_6^{''}$. The method is found to be applicable in the presence of bromides and nitrates. Different stages of oxidation of iodine can also be estimated in mixtures.

W. T.

The Estimation of Sulphur in Vulcanised Rubber. J. W. W. DYER and AMY R. WATSON (*J. Soc. Chem. Ind.*, 1922, **41**, 332r).—With reference to the method described for the estimation of combined sulphur in vulcanised rubber (this vol., ii, 656), the authors acknowledge priority of publication to A. R. Pearson (*Analyst*, 1920, **45**, 405). They prefer to use the method for combined sulphur only, however, and not for total sulphur, unless the amount of free sulphur is small.

G. F. M.

Estimation of Sulphides by Oxidation with Ferric Salts. P. P. BUDNIKOFF and K. E. KRAUSE (*Z. anorg. Chem.*, 1922, **122**, 171—174).—The estimation is based on the reaction $3\text{Na}_2\text{S} + 3\text{Fe}_2(\text{SO}_4)_3 = 3\text{Na}_2\text{SO}_4 + 6\text{FeSO}_4 + 3\text{S}$. The amount of ferrous salt is found by permanganate. The acidity of ferrous sulphate is neutralised by the addition of alkali carbonate. The method is only applicable to sulphides which are soluble in water and acids.

W. T.

The Estimation of Sulphuric Acid as Barium Sulphate in the Presence of Aluminium. L. MOSER and P. KOHN (*Z. anorg. Chem.*, 1922, **122**, 299—310).—The results of the estimation of sulphuric acid in the presence of aluminium are too low, these low results being due to the presence of aluminium sulphate in the precipitate. The amount of this impurity increases up to a maximum with an increase in the concentration of the aluminium salt. Precipitated barium sulphate takes up aluminium sulphate from solution—a solid solution is thus indicated. The best results are obtained by working in hydrochloric acid solution when the maximum error is -0.15% . The errors observed in the presence of the aluminium-ion are only one-tenth those obtained in the presence of the ferric-ion.

W. T.

A New Accelerator for the Destruction of Organic Matter in the Kjeldahl Method for the Estimation of Nitrogen. M. SBOROWSKY and I. SBOROWSKY (*Ann. Chim. Analgt.*, 1922, 4, 266—267).—Mercurous iodide accelerates the destruction of organic matter in the digestion with sulphuric acid much more effectively than metallic mercury or other mercury salts. In two experiments carried out with 10 c.c. of sulphuric acid, with mercurous iodide 0.7 gram of sugar was destroyed in the time required for the destruction of only 0.1 gram with metallic mercury. A digestion was completed in fifty to sixty minutes which with metallic mercury required four to eight hours. H. C. R.

Estimation of Total Nitrogen in Fertilisers containing Nitrites and of Nitrite Nitrogen in the Presence of Nitrates. F. MACH and F. SINDLINGER (*Z. angew. Chem.*, 1922, 35, 473—474).—To estimate the total nitrite and nitrate nitrogen, 25 c.c. of an aqueous solution of the sample (this quantity of solution should contain not more than 0.05 gram of nitrogen) are added gradually to a boiling mixture of 30 c.c. of saturated potassium permanganate solution and 5 c.c. of dilute sulphuric acid (1 : 2); a further quantity of 10 c.c. of the acid is then added, the mixture is cooled, treated with 10 grams of ferrum redactum, and again boiled for about ten minutes. The resulting ammonia is estimated by distillation in the usual way. The nitrate nitrogen is then estimated in another portion of 25 c.c. of the solution by boiling this quantity for ten minutes with 30 c.c. of *N/3*-sulphuric acid and 10 c.c. of methyl alcohol (the whole of the nitrous acid is thus expelled in the form of its methyl ester), neutralising the cooled mixture, and reducing the remaining nitrate with ferrum redactum in sulphuric acid solution. The nitrite nitrogen is found by subtracting the nitrate nitrogen from the total nitrogen. If ammonium salts are present, the ammonia nitrogen must be estimated separately and an allowance made for its quantity. W. P. S.

Comparison of the Methods for the Estimation of Nitric Oxide. ALFONS KLEMENC and CORNELIE BUNZL (*Z. anorg. Chem.*, 1922, 122, 315—331).—Nitric oxide was prepared according to the method of Emich (A., 1892, ii, 939); this, however, contained 1% of impurity and was purified by fractionation, using liquid air. The estimation of the gas by passing it over heated copper was found to give accurate results. Estimations (1) with hydrogen in the Drehschmidt platinum capillary (Knorre and Arndt, A., 1899, ii, 806), (2) by the method of Baudisch and Klinger (*Ber.*, 1918, 45, 3231), and (3) by absorption with ferrous sulphate were found to give untrustworthy results. Good results were obtained by absorbing the gas by acidified bromate or *N/5*-potassium permanganate and estimating the excess. The latter method can also be used in the presence of carbon dioxide and other gases. W. T.

Detection of the Nitrous Anion. P. FALCIOLA (*Gazzetta*, 1922, 52, ii, 87—89).—Small proportions of nitrite may be detected : (1) By the transitory yellow coloration obtained by acidification

with sulphuric acid and gradual addition of sodium thiosulphate solution. (2) By the orange colour developed on addition, to a litre of the solution, of about 2 c.c. of sulphuric acid, excess of thiocyanate, and about 0.1 gram of ferrous sulphate (cf. Horst, A., 1921, ii, 461). (3) By the yellow coloration formed on addition of excess of aniline phosphate. [Cf. *J. Soc. Chem. Ind.*, 1922, 856A.]

T. H. P.

Separation of Phosphoric Acid in Qualitative Analysis.

N. TARUGI (*Boll. Chim. Farm.*, 1922, 61, 545—552).—The author's investigation of Gattermann and Schindhelm's proposal to eliminate phosphoric acid by means of stannic chloride (A., 1917, ii, 41) shows that, even under the best conditions, 15.4% of the total phosphoric acid passes into the filtrate from the stannic precipitate, and that the latter contains part of the iron, chromium, calcium, barium, etc., present. Further, when the proportion of phosphoric acid in the solution to be examined is unknown, it is impossible to add the exact quantity of ammonium phosphate to give a definite ratio between phosphate and stannic chloride.

T. H. P.

Effect of Iron on the Iodine Titration of Arsenite. F. L.

MELVILL (*J. S. African Chem. Inst.*, 1922, 5, 3—5).—The interference of ferric salts in the titration of arsenite solution by iodine solution may be prevented by the following procedure. The slightly acid arsenite solution, containing not more than 0.1% of iron (expressed as Fe_2O_3), is treated with about one-half the quantity of iodine solution required to oxidise all the arsenite, 10 c.c. of saturated sodium hydrogen carbonate solution are then added, and the titration is completed. In this way the whole of the iron is precipitated as ferric arsenate and no longer interferes with the titration.

W. P. S.

Estimation of Boric Acid. WILHELM STRECKER and ERNST

KANNAPPEL (*Z. anal. Chem.*, 1922, 61, 378—397).—The substance containing boric acid or its salts is mixed with 40% phosphoric acid and methyl alcohol and the mixture is distilled in a current of air saturated with methyl alcohol (the air is passed previously through a flask containing methyl alcohol) until all the methyl borate has been collected in a receiver containing sodium hydroxide solution. The distillate is then heated to expel methyl alcohol, treated with a slight excess of hydrochloric acid, using methyl-orange as indicator, boiled to expel carbon dioxide, and neutralised with sodium hydroxide solution. After the addition of mannitol in the proportion of 1 gram for each 10 c.c. of solution, the mixture is titrated with *N*/10-barium hydroxide solution, using α -naphtholphthalein solution as indicator.

W. P. S.

Sulphonated Derivatives of the Naturally Occurring Sulphidic Hydrocarbons. C. PÉPIN and G. REAUBOURG (*J. Pharm. Chim.*, 1922, [vii], 26, 258—261).—As the therapeutic

value of ichthyol is largely dependent on the amount of "sulphidic" sulphur it contains as distinguished from total sulphur which

comprises in addition sulphonic and sometimes sulphatic sulphur, it is desirable that more precise standards should be laid down to which commercial ichthyol preparations should conform. Ammonium or other sulphates should be almost completely absent, as indicated by the formation of nothing more than an opalescence on the addition of barium chloride to the filtrate after precipitation of an ichthyol solution by albumin and hydrochloric acid. Sulphidic sulphur can only be estimated by difference by Thal's method, which consists in an estimation of total sulphur, by Carius's method, for example, and subtracting from this the sulphur present as sulphate, if any, determined by weighing the above barium sulphate precipitate, and the sulphonic sulphur determined by an ammonia estimation, and calculating on the basis of one sulphonic group for each molecule of ammonia after making allowance for the ammonia combined as sulphate. As thus estimated, "sulphidic" sulphur should represent at least 15% of the weight of the dried ichthyol. The conservation of ethylenic linkings in the preparation of the ichthyol should be confirmed by the decolorisation of bromine water.

G. F. M.

Radioactive Indicators. FRITZ PANETH (*Z. angew. Chem.*, 1922, 35, 549—552).—Use has been made of radioactive isotopes of metals such as lead and bismuth for indicating the presence of minute and unweighable quantities of these elements or their compounds. Quantities of the radioactive isotopes of the order of a millionth of a milligram are easily detected by the electroscope. Once mixed with the inactive isotope, they cannot be separated by chemical means, so that the presence of the radioactive isotope indicates the presence of the inactive one also. Practical applications have been made in analytical chemistry in determining the solubility of very sparingly soluble substances and in the study of adsorption phenomena. In electrochemistry it has been established by this means that an interchange of the metallic portion of the molecule actually occurs on mixing solutions of two salts of the same metal and subsequently recovering them from solution, but that no such interchange occurs when one or both the substances are non-ionised. The actual deposition of an unweighable quantity of metal on the electrode before the minimum potential of electrolysis is reached has also been established. In colloid chemistry, the active surface of powders has been measured for the first time by the use of a radioactive indicator, and the distinction between crystalloid and colloid in the diffusion of the former through parchment membranes has been studied down to a dilution of 10^{-11} mol. per litre. In inorganic chemistry, the conditions for the preparation of bismuth and lead hydrides were studied by this means and the method was successfully applied to the measurement of the gas-tightness of rubber fabrics for gas-masks. The kinetic theory has received further experimental confirmation by the actual proof of the movement of the molecules in molten lead and measurements of the movement of lead ions in lead chloride even below its melting point.

H. C. R.

Estimation of Small Quantities of Calcium. PATRICK PLAYFAIR LAIDLAW and WILFRED WALTER PAYNE (*Biochem. J.*, 1922, 16, 494—498).—The method described is suitable for estimating calcium in amounts of the order of 0.1 mg., and gives results accurate to about 0.002 mg. The calcium is first precipitated as oxalate (in the case of blood serum, this may be done directly from the serum without previous ashing). After separation, the precipitate is dissolved in hydrochloric acid and the calcium reprecipitated in the form of calcium alizarinate by the addition of excess of alizarin in alcoholic solution, and, after warming, of a few drops of strong ammonia. When crystalline, the precipitate is collected in a Gooch crucible, washed with dilute ammonia, and decomposed with a solution of oxalic acid in 50% alcohol. The liberated alizarin is dissolved in 95% alcohol, made just alkaline with ammonia, and estimated colorimetrically by comparison with a standard solution of ammonium alizarinate. E. S.

The Separation of Zinc from Other Metals, especially Nickel. ALOIS LUDWIG (*Z. anorg. Chem.*, 1922, 122, 239—261).—Zinc can be separated from nickel by precipitation with hydrogen sulphide in the presence of ammonium tartrate. The latter salt forms stable complex compounds with nickel, which may perhaps explain its action; the latter might, however, be due to the replacement of the freed mineral acid by the weaker tartaric acid. Results are also given of attempts to effect the separation by precipitation of zinc from zinc ammonium phosphate in neutral solution in the presence of neutral tartrates and thiocyanates. These investigations are being continued. W. T.

The Influence of the Alkalis on the Titration of some Metals with Ferrocyanide. W. D. TREADWELL and D. CHERVET (*Helv. Chim. Acta*, 1922, 5, 633—639).—The electrometric titration of zinc with potassium ferrocyanide was described by Treadwell and Weiss (A., 1920, ii, 120). The sharpness of the end-point depends on the insolubility of the ferrocyanide of the heavy metal, but it is also dependent on the alkali metal used, since the precipitate is generally a double salt. When cadmium sulphate is titrated with potassium ferrocyanide in neutral or weakly acid solution, the end-point is indicated by a moderately sudden drop in potential, but only in a highly dilute solution has the precipitate the exact composition $\text{CdK}_2\text{Fe}(\text{CN})_6$. In the presence of a rubidium salt or a caesium salt, however, the end-point is much sharper, probably because the alkali metals with higher atomic volumes form less soluble double salts with cadmium ferrocyanide. When sodium ferrocyanide is used for the titration, the precipitate is the simple salt, $\text{Cd}_2\text{Fe}(\text{CN})_6$. When a zinc salt is titrated with potassium ferrocyanide the precipitate is $\text{Zn}_3\text{K}_2[\text{Fe}(\text{CN})_6]_2$, but with sodium ferrocyanide, $\text{Zn}_2\text{Fe}(\text{CN})_6$, is obtained. In presence of a potassium salt, however, sodium ferrocyanide gives the above zinc-potassium salt, whilst in presence of a caesium salt, potassium ferrocyanide gives $\text{ZnCs}_2\text{Fe}(\text{CN})_6$ with a very sharp end-point. Lead nitrate can be titrated with ferrocyanide in neutral solution,

but lead ferrocyanide is very sensitive to acid; consequently zinc can be titrated with ferrocyanide in hot acid solution in presence of lead. The precipitate obtained from lead nitrate and potassium ferrocyanide is $\text{Pb}_2\text{Fe}(\text{CN})_6$, but in presence of a cæsium salt the precipitate has the composition $\text{Pb}_2\text{X}_2[\text{Fe}(\text{CN})_6]_2$, where X is probably entirely cæsium. The ferrocyanides of the bivalent heavy metals appear to be less soluble the smaller the atomic volume of the heavy metal. E. H. R.

Estimation of Lead in Lead Amalgam. M. G. MELLON and H. F. REINHARD (*Proc. Indiana Acad. Sci.*, 1921, 189—195).—Existing methods are considered to be unsatisfactory for the estimation of small quantities of lead in the presence of large quantities of mercury. A weighed quantity of the amalgam is covered with 25 c.c. of a 10% solution of cupric nitrate and allowed to remain for fifteen to twenty-four hours; lead enters into solution, but mercury does not. After decantation through a filter, the amalgam is washed, and the filtrate treated with a few drops of acetic acid and sufficient potassium dichromate solution to precipitate the lead as lead chromate, which is washed in a Gooch crucible, dried at 120° , and weighed. CHEMICAL ABSTRACTS.

The Kastle-Meyer Reagent as a very Sensitive Reagent for Copper. OTTORINO CARLETTI (*Boll. Chim. farm.*, 1922, 61, 449).—A claim for priority (A., 1914, ii, 74) over Thomas and Carpentier (this vol., ii, 86). T. H. P.

New Method for the Estimation of Manganese. ST. MINOVICI and CONST. KOLLO (*Chim. et Ind.*, 1922, 8, 499—500).—Manganese may be estimated with accuracy, even when present in only small quantities, by precipitation as iodate by means of iodic acid according to the equation $\text{MnSO}_4 + 2\text{HIO}_3 = \text{Mn}(\text{IO}_3)_2 + \text{H}_2\text{SO}_4$. A solution of about twice the theoretical quantity of iodic acid is added to the manganese solution, and after warming for ten minutes on a water-bath twice the volume of alcohol is added. After keeping for a short time, the precipitate is collected on a filter or Gooch crucible and washed with 70% alcohol saturated with manganous iodate. The mother-liquor contains no traces of manganese, the iodate being insoluble in 70% alcohol in presence of iodic acid. In water at the ordinary temperature, its solubility is 0.195%, and in 70% alcohol 0.005%. The precipitate of manganous iodate is finally dried at 100° , and weighed. The absolute percentage error in the examples given varies from 0.16 to 0.63% when quantities of the order of 0.1—0.3 gram of the hydrated sulphate are taken for estimation. G. F. M.

A New Microchemical Method for the Identification of Tungsten. J. A. M. VAN LIEMPT (*Z. anorg. Chem.*, 1922, 122, 236—238).—A small quantity of sodium nitrite is melted on a platinum spoon and a little tungsten is thrown in; sodium tungstate is formed. This is dissolved in a little water and tungstic acid precipitated by the addition of hydrochloric acid (1:1). This is collected and then mixed with a few drops of concentrated

ammonia and placed on a slide. Ammonium paratungstate soon crystallises in four-cornered plates on the edges of the drop and as long needles in the centre. W. T.

¶ **Glacial Acetic Acid Method for Estimating Uranium in Carnotite.** WILFRED W. SCOTT (*J. Ind. Eng. Chem.*, 1922, 14, 531—532).—Half a gram of the ore is heated with 40 c.c. of dilute (1 : 1) nitric acid, the mixture is evaporated to dryness, the residue ignited, and then boiled for five minutes with a mixture of glacial acetic acid and nitric acid (100 : 5); the insoluble portion is separated by filtration and the filtrate is evaporated to dryness. This residue is ignited until it turns black and the extraction with the acetic acid-nitric acid mixture repeated; the filtrate obtained is evaporated, the residue dissolved in 10 c.c. of nitric acid and 40 c.c. of water, the solution neutralised partly with ammonia and ammonium carbonate is added in quantity sufficient to dissolve the precipitate of uranium carbonate which first forms. After the addition of an excess of 3 grams of ammonium carbonate and 5 c.c. of ammonia, the mixture is filtered, the filtrate is acidified with nitric acid, and heated to expel carbon dioxide. An excess of ammonia is then added and the boiling is continued until all the uranium has been precipitated, which is denoted by the yellow solution becoming colourless. The precipitate is collected, washed with 2% ammonium nitrate solution, ignited, and weighed as U_3O_8 . W. P. S.

Reductions with Cadmium in Volumetric Analysis. II. W. D. TREADWELL (*Helv. Chim. Acta*, 1922, 5, 732—743).—[With M. BLUMENTHAL.]—Uranyl salts can be titrated by first reducing to the uranous stage and titrating back with potassium permanganate (cf. A., 1921, ii, 523). When zinc is used for the reduction, the reaction tends to go beyond the uranous stage to the U''' stage and the same occurs, although to a less extent, when cadmium is used. Owing to the ease with which tervalent uranium is oxidised by air to the uranous stage, any over-reduction can be corrected by allowing the reduced solution to run from the reduction tube drop by drop into a vessel open to the air. Oxidation from the uranous to the uranyl stage by means of air depends on the degree of acidity of the solution. To prevent such oxidation, the reduction is carried out in a 3.0*N*-sulphuric acid solution. When the electrometric method of titration is used over-reduction can be ignored, as the titre is given by the distance between the two breaks in the potential curve. Titanium and uranium may be estimated together by reducing in very dilute solution and titrating electrometrically with permanganate in an atmosphere of carbon dioxide, the two breaks in the curve representing the completion of oxidation of titanous and uranous ions respectively.

[With M. STÄRKLE.]—The accuracy of the electrometric method for titrating titanium after reduction with cadmium (*loc. cit.*) was confirmed by comparison with a gravimetric method. It is shown that small quantities of titanium can be estimated accurately by this method in presence of large quantities of iron, using potassium dichromate for the titration.

[With M. Hooft.]—Careful experiments in which every precaution was taken to exclude air during the reduction and titration with permanganate show that vanadic acid is reduced quantitatively to the V'' stage by cadmium.

[With M. DREIFUSS and A. BOSSI.]—Indigotinsulphonic acid is reduced by finely divided cadmium in 0.2 to 0.5*N*-sulphuric acid solution to the leuco-stage, and can then be accurately titrated by means of ferric chloride by the electrometric method in an atmosphere of carbon dioxide. The course of the potential curve indicates that the oxidation proceeds in two stages, probably through a quinhydrone stage. Thioindigotin and methylene-blue can be estimated in the same way. [Cf. *J. Soc. Chem. Ind.*, 1922, Nov.]
E. H. R.

A Micro-method for the Estimation of Ethyl Alcohol in Blood. ERIK M. P. WIDMARK (*Biochem. Z.*, 1922, 131, 473—484).—A method is described suitable for estimating the alcohol in a drop of blood or in larger quantities. It depends on the absorptive power for alcohol of concentrated sulphuric acid containing a known amount of dichromate. The operations are conducted in a specially designed flask to avoid mixing and contamination of the standard dichromate by organic matter, and at a temperature of 50—60°. Absorption of the alcohol is complete within two hours. The excess of dichromate is estimated, after dilution with water, by potassium iodide and thiosulphate. Blank estimations are essential features.
H. K.

The Estimation of the Methyl Group in Methylated Thiolbenzenes. J. POLLAK and ANNA SPITZER (*Monatsh.*, 1922, 43, 113—120).—The estimation of the methylthiol group by a method analogous to Zeisel's process for estimating the methoxyl group is complicated by the fact that hydrogen sulphide and methylthiol are evolved along with methyl iodide during the decomposition by hydriodic acid. The process now described consists in decomposing the substance with hydriodic acid (*d* 1.7), and passing the volatile products through a paste of red phosphorus and weakly acidified cadmium sulphate solution, to remove hydrogen sulphide, and thence into alcoholic silver nitrate. The precipitate is ignited, and the mixture of silver and silver iodide completely converted into the iodide for weighing.
C. K. I.

Identification of Small Quantities of Terpeneol Hydrate in Complex Mixtures. G. DENIGÈS (*Bull. soc. pharm. Bordeaux*, 1922, 60, 49—54).—The sample is moistened with a drop of ethyl alcohol on a glass slide, terpeneol being deposited on the edge in tufts of prismatic needles; previous sublimation between two slides may be desirable. On addition of a drop of sulphuric acid, a yellow colour is produced, passing to orange on heating. Finally, on addition of a drop of a solution containing 0.5 gram of lævulose in 25 c.c. of ethyl alcohol and 75 c.c. of water, a carmine-red coloration is developed.
CHEMICAL ABSTRACTS.

Estimation of the so-called Total Geraniol Content of Citronella Oil. A. W. K. DE JONG and A. RECLAIRE (*Perf. Essent. Oil Rec.*, 1922, 13, 319—321).—Numerous samples of citronella oil were analysed by slightly varying methods in order to ascertain the reason for the different results often obtained by different operators. The maximum difference observed was 2.8%, the average difference 0.7%. The influence of acetic anhydride of higher percentage strength than usual (80%) was negligible, and sodium acetate dried over sulphuric acid did not give better results than the usual fused article. The method recommended is to heat 10 grams of the dry oil with 10 c.c. of 80% acetic anhydride and 2 grams of freshly fused sodium acetate for two hours. After cooling, 50 c.c. of water are added and the excess of anhydride is decomposed by warming and shaking. The acetylated oil is separated, washed with brine until neutral, and dried. About 1.5 grams are then saponified with *N*/2-alcoholic potassium hydroxide after neutralising any residual acidity. After two hours' boiling under reflux the excess of alkali is titrated back with *N*/2-sulphuric acid.
G. F. M.

Causse and Bonnan's Method for the Estimation of Dextrose. R. KRULL (*Pharm. Weekblad*, 1922, 59, 1035—1039).—The unsatisfactory end-point referred to by Lemkes and Lansberg (this vol., ii, 724) is due to the fact that towards the end of the titration, when most of the cuprous oxide has been removed from the solution, dextrose is being added to a strongly alkaline solution, to which, as is well known, it is very sensitive. The abnormal course of Fehling's reaction with urine may be considered in the light of Benedict's reaction. The iodometric estimation is accurate to 0.1%.
S. I. L.

Apparatus for Facilitating the Estimation of Dextrose by Titration with Permanganate by the Mohr-Bertrand Method. J. VAN DER HAAR (*Arch. Suikerind.*, 1922, 30, 213—216).—Precipitation is carried out in a 500 c.c. round flask, and the supernatant liquid siphoned slowly through a Soxhlet tube into an 800 c.c. bottle, the neck of which is of the same size as that of the round flask. The precipitate is washed with hot water and the washings are siphoned off. The Soxhlet tube is removed from the bottle and placed on the flask. A little ferrous ammonium sulphate solution is next run through the siphon into the Soxhlet tube and the remainder poured directly into the tube itself; the tube is rinsed, and the contents of the flask are titrated with permanganate after acidification with sulphuric acid.

CHEMICAL ABSTRACTS.

Estimation of Sucrose in Presence of Other Sugars by means of Alkaline-earth Hydroxides. A. BEHRE and A. DÜRING (*Z. Unters. Nahr. Genussm.*, 1922, 44, 65—70).—Material containing about 20 grams of sucrose is heated with 1.2 grams of freshly burnt lime in about 50 c.c. of water for one hour at 60—80°.

Other sugars are destroyed, and the sucrose can be determined polarimetrically.

A. G. P.

Flax and Kindred Fibres. I. Behaviour and Structure of Textile Fibres, and a Convenient Method of Distinguishing Flax from Hemp. C. R. NODDER (*Trans. Text. Inst.*, 1922, **13**, 161—171).—Flax and ramie fibres always twist in a clockwise direction when drying, whereas hemp and jute fibres twist in the reverse way, and cotton usually exhibits different twists in different parts of the same hair. An examination of about 100 fibres in a warm, dry room is sufficiently accurate for determining the proportion of flax and hemp in a mixture. The direction of twist is connected with the spiral, fibrillar structure of these "bast" fibres. This structure is best revealed by mounting the fibre in concentrated calcium chloride solution, tinged with iodine, and carefully compressing it by gentle pressure on the cover-slip. The fibre is easily made ten times its normal width and the spirals show as bright reddish-purple lines (the compressed parts) among brownish-yellow ones. The striations in the outer layer of flax and ramie cells form left-handed spirals, and those of hemp and jute right-handed spirals. It follows, therefore, that wetting of these fibres is accompanied by an untwisting of the fibrils which make up the outer layer of the cell-wall. The phenomena are discussed in the light of the physico-chemical properties of cellulose.

J. C. W.

Rapid Estimation of Acetic and Butyric Acids in Mixtures. L. NOTENBAART (*Natuurwetenschapp. Tijdschr.*, 1921, **3**, 131—134).—After previous experimental determination of the distribution of acetic acid and butyric acid between water and toluene, the total quantity of acid present is estimated by titration with 0.04*N*-sodium hydroxide. After extraction of most of the butyric acid and a little of the acetic acid by shaking with an equal volume of toluene, the aqueous solution is again titrated.

CHEMICAL ABSTRACTS.

Adaptation of the Pentabromoacetone Method to the Estimation of Citric Acid in Urine. WILLIAM B. MCCLURE (*J. Biol. Chem.*, 1922, **53**, 357—363).—The pentabromoacetone method for the estimation of citric acid (cf. Kunz, A., 1915, ii, 595) cannot be applied directly to urine owing to the formation of other precipitates. Satisfactory results may, however, be obtained if the urine is first made alkaline with sodium hydroxide and then treated, in the cold, with animal charcoal, and if the pentabromoacetone is finally separated from small quantities of impurities by volatilisation.

E. S.

Estimation of Meconic Acid in Opium. H. E. ANNETT and M. N. BOSE (*Analyst*, 1922, **47**, 387—391).—Five grams of opium are triturated with 50 c.c. of water and after keeping over-night 40—45 c.c. of the filtered liquid are treated with 6 c.c. of 50% calcium chloride solution and kept a further twenty-four hours.

29*

The precipitate, which is almost white, consists of calcium meconate and sulphate. It is collected, washed until the washings are colourless, and dissolved in 15 c.c. of 1.25*N*-hydrochloric acid. After keeping for twenty-four hours, pure meconic acid will have separated in white, crystalline scales. It is collected, washed twice with 0.5 c.c. of water, dried over sulphuric acid, and weighed as $C_7H_4O_7 \cdot 3H_2O$. To the weight found 0.0213 gram is added, to correct for the solubility of meconic acid in 15 c.c. of 1.25*N*-hydrochloric acid. The figure obtained is multiplied by the factor depending on the amount of the aliquot portion of the solution taken for analysis, and then by the factor 10/9, to correct for the amount of meconic acid unprecipitated by calcium chloride, in order to arrive at the weight of acid in 5 grams of opium. G. F. M.

Pregl's Solution. M. BACHSTETZ (*Ber. Deut. pharm. Ges.*, 1922, 32, 216—221).—Discrepancies between the analyses of Pregl's solution made by different observers indicate that its composition is variable in spite of assurances to the contrary given by the makers. A solution prepared by dissolving 0.0165 gram of sodium iodate and 0.65 gram of potassium iodide in 100 c.c. of water with the addition of 10 drops of *N*/10-hydrochloric acid gave very similar results to Pregl's solution when tested quantitatively for total, active, and free iodine. Its physiological and bacteriological properties were also similar to those of Pregl's solution. E. H. R.

The Sulphuric Acid Reaction for Liver Oil. H. D. RICHMOND and E. H. ENGLAND (*Analyst*, 1922, 47, 431).—Constant results were obtained in testing cod-liver oil by the dilution method (Drummond and Watson, this vol., ii, 665) by adding 1 c.c. of the cod-liver oil to 10 c.c. of liquid paraffin (B.P.), mixing, and transferring 10 drops of the mixture to a white porcelain basin, adding 1 drop of sulphuric acid (B.P.) and stirring with a glass rod. If a transient purple colour developed, liquid paraffin was added in successive quantities of 5 c.c. until no purple coloration was given, and the dilution at which a faint transient purple was seen was recorded. If no purple was seen, successive additions of 1 c.c., 1 c.c., 2 c.c., and 5 c.c. of cod-liver oil were made. H. C. R.

Properties and Preparation of Dutch East Indies Sandal Wood Oil. A. ROJDESTWENSKY (*Perf. Essent. Oil Rec.*, 1922, 13, 331—332).—The amount of oil obtainable from Dutch Indian sandal wood varies from 3.8—5.8%, and has the following characters: d_{20}^{20} 0.9745—0.9842, α_D^{20} -16.4 to -18.4 , santalol content 93.7—98%; n_D^{20} 1.5000—1.5013, soluble in 3—4 vols. of 70% alcohol. The quantity of oil obtainable from a sample may be ascertained by distilling not more than 0.5 kilo. of shavings with water after soaking for forty-eight hours. After ten to fifteen hours, the colour of the oil begins to darken, and in twenty hours the distillation is finished. The first and last runnings do not differ greatly in properties. G. F. M.

Importance of Dimethylhydroresorcinol for Detection of Volatile Aldehydes in Body Fluids. The Identification of Formaldehyde in Urine after Administration of Hexamethylenetetramine. WILHELM STEPP (*Biochem. Z.*, 1922, **130**, 578—581).—After administration of hexamethylenetetramine to a patient with kidney disease, formaldehyde was recognised in quantity in the urine by combination with dimethylhydroresorcinol.
H. K.

A Modified Schiff's Solution. E. WERTHEIM (*J. Amer. Chem. Soc.*, 1922, **44**, 1834—1835).—Rosaniline hydrochloride (0.005 gram) is dissolved in hot water (50—100 c.c.); the solution is filtered if necessary, diluted to 300 c.c. and cooled with running water. Sodium hyposulphite (6 grams) is added to the cooled solution. The salt dissolves completely in a few minutes, after which the solution is ready for use. If it is boiled for one minute (not longer) it first becomes red and subsequently practically colourless. It may now be heated in testing for aldehydes in order to hasten the test. Used in this manner, the solution gives the aldehyde test in about half the time required for the ordinary Schiff's test, although without heating it is usually a little slower than the ordinary Schiff's solution. Special precautions in preserving the solution are unnecessary.
H. W.

Comparative Estimations of Acetone in Urine. KURT KÄDING (*Biochem. Z.*, 1922, **130**, 448—458).—A comparison has been made of various methods, suitable for clinical purposes for the estimation of acetone in urine. Scharf's method (*Med. klin.*, 1921, **36**), like Schall's method, is a modification of Legal's test and is the most suitable.
H. K.

The Characterisation of the Colouring Matter of Saffron : Its Use in Investigations Relating to Laudanum Poisoning. MARCEL GUERBET (*J. Pharm. Chim.*, 1922, **26**, 218—220).—The coloration produced by the action of concentrated sulphuric acid on crocetin may be used for the detection of the colouring matter of saffron which yields crocetin on hydrolysis. This test may be used in the case of poisoning by drugs which are coloured with saffron.
H. J. E.

A Qualitative Test for Tannin. ETHEL ATKINSON and EDITH OLIVE HAZLETON (*Biochem. J.*, 1922, **16**, 516—517).—A piece of gold-beater's skin is pinned on a surface of paraffin wax, soaked in water, and then covered with an aqueous extract of the material to be tested. After about fifteen minutes it is washed, treated with a 1% solution of ferric chloride, and again washed. If the skin is stained, the presence of tannin is indicated.
E. S.

Estimation of Indican in Blood-serum. J. SNAPPER and W. J. VAN BOMMEL VAN VLOTEN (*Klin. Woch.*, 1922, **1**, 718—721; from *Chem. Zentr.*, 1922, ii, 1122).—In the estimation of indican

in serum, a stronger reaction is given after precipitation with trichloroacetic acid than after precipitation with alcohol. The following procedure is recommended. The serum is precipitated with an equal volume of 20% trichloroacetic acid. 2.5 C.c. of the filtrate are diluted to 10 c.c. with water and after addition of 1 c.c. of thymol in alcohol (5%) and 10 c.c. of Obermayer's reagent, left for twenty minutes. The liquid is then extracted with 2 c.c. of chloroform and the reading taken after thirty minutes. With a weak positive reaction a slightly increased indicanæmia is indicated, implying hypofunction of the kidneys (niereninsuffizienz). In acute nephritis, hyperindicanæmia implies a more marked intoxication than does retention of urea alone. The principal precautions to be observed in the estimation are given.

G. W. R.

Detection of Urea in Tissue by Means of Xanthydril. M. BONNET and J. HAUSHALTER (*Compt. rend. Soc. Biol.*, 1922, **86**, 395—397; from *Chem. Zentr.*, 1922, ii, 922).—For the microchemical detection of urea, the fixation of the tissue in a 10% solution of xanthydril in ethyl alcohol and pure acetic acid in the ratio 1 : 7 is recommended.

G. W. R.

A Reaction of Urea with *p*-Dimethylaminobenzaldehyde. H. K. BARRENSCHEEN and O. WELTMANN (*Biochem. Z.*, 1922, **131**, 591—595).—The yellowish-green coloration obtained in dilute urine on addition of Ehrlich's aldehyde reagent is due to urea. The reaction can be used for the recognition of residual nitrogen in serum freed from proteins by trichloroacetic acid when it exceeds 36—40 mg. %.

H. K.

Accurate Ureometer. ANTONIO SCIORTINO (*Arch. Farm. sperim. Sci. aff.*, 1922, **33**, 186—189).—The modified apparatus for the estimation of carbamide described allows of the use of varying quantities of the urine, blood, etc., and of arbitrary amounts of the sodium hypobromite. Equilibrium may be established between the external and internal pressures prior to the reaction and completion of the latter assured by shaking the reaction bottle under diminished pressure. The volume of the nitrogen formed is read in a burette graduated to 0.05 c.c.

T. H. P.

The Estimation of Urea in Urine by the Hypobromite Method with Complete Yield. (Mlle) M. JANET (*J. Pharm. Chim.*, 1922, **26**, 161—170).—Sodium hypobromite, under the usual experimental conditions, gives results which indicate only 90—92% of the actual urea present. The method described is said to be accurate within 1% when compared with results obtained by standard gravimetric methods. The solution containing urea is diluted, if necessary, so that the concentration is not much above 1% and a considerable excess of sodium hydroxide is added followed by the hypobromite. The presence of creatinine introduces a slight error; this is stated to be negligible. If ammoniacal com-

pounds are present, the whole of their combined nitrogen is liberated by the action of sodium hypobromite and sodium hydroxide so that the nitrogen due to urea is found by difference after estimating the ammonia.

H. J. E.

Electrometric Titration of Ferrocyanides. ERICH MÜLLER and HANS LAUTERBACH (*Z. anal. Chem.*, 1922, 61, 398—403).—The end-point of the titration of ferrocyanides in sulphuric acid solution with permanganate solution may be determined accurately electrometrically in cases where the titration has to be made under artificial light; the results obtained agree with those found where the end-point is denoted by the pink coloration of the slight excess of added permanganate.

W. P. S.

A Test for Pyridine. FRIEDRICH LEHNER (*Chem. Z.*, 1922, 46, 877).—If a few drops of aniline are added to a solution of pyridine containing water and a trace of cyanogen bromide, a red colour is immediately obtained and crystals of 2-anilinodihydropyridine phenyl bromide separate. One part of pyridine in 350,000 parts can be detected in this way.

H. C. R.

Detection of Small Quantities of Pyridine. A. GORIS and A. LARSONNEAU (*Bull. Sci. Pharmacol.*, 1921, 28, 497—498; from *Chem. Zentr.*, 1922, ii, 731).—By addition of aniline to a solution of pyridine in the presence of cyanogen bromide, 1-anilinodihydropyridinium phenyl bromide, a red substance, is formed. The coloration is noticeable even with one drop of pyridine in 10,000 c.c. of water. This reaction takes place even in the presence of pyrrole derivatives.

G. W. R.

A Reaction of Veronal and of the Hypnotics Derived from Barbituric Acid. RENÉ FABRE (*J. Pharm. Chim.*, 1922, [vii], 26, 241—249).—Veronal and other hypnotics of the barbituric acid series are readily condensed with xanthydrol to form crystalline dixanthyl derivatives by heating at 100° for about one minute an approximately 10% solution of the hypnotic in acetic acid with twice its weight of xanthydrol. After keeping for a few hours, the product is separated and washed with a little boiling alcohol, and is then sufficiently pure for melting-point determination. The m. p. of the principal dixanthylbarbituric acids are as follows: dixanthylveronal 245—246°, dixanthylphenylethylbarbituric acid 218—219°, and dixanthylallylbarbituric acid 242—243°. Similar derivatives are not formed by other hypnotics, and the reaction is therefore valuable for the characterisation and identification of the barbituric acids for both pharmaceutical and toxicological purposes, as it can easily be carried out with as little as 0.01 gram of the substance, or with the crude product obtained by the usual methods in toxicological investigations from the viscera, etc.

G. F. M.

The Estimation of Uric Acid. HENRY JACKSON, jun., and WALTER W. PALMER (*J. Biol. Chem.*, 1922, 53, 373).—A simplified

method for the preparation of the modified phosphotungstic acid reagent (this vol., ii, 328) is described. E. S.

Estimation of Uric Acid in Blood. L. BAUMAN and L. M. KEELER (*J. Lab. Clin. Med.*, 1922, 7, 551—552).—A modification of Folin and Wu's method in which calibrated Lovibond tintometer glasses replace standard uric acid solution. A red glass (0.4) is placed over the unknown solution and a blue glass (2.9) over the opposite prism in a Duboscq colorimeter; the blue compound is developed without the use of sodium sulphite. The glasses may also be used with the Bock-Benedict colorimeter if the red glass is placed over the immersion cylinder and the blue in front of and parallel to the standard cell. The method is of general application; the use of glasses shortens the time of procedure and removes uncertainty arising from the possible decomposition of the standard uric acid solution. Precipitation may be avoided by addition of three drops of a solution of gum acacia preserved with thymol.

CHEMICAL ABSTRACTS.

Estimation of Uric Acid and Urates in Blood. CH. O. GUILLAUMIN (*Compt. rend. Soc. biol.*, 1922, 86, 194—196; from *Chem. Zentr.*, 1922, iv, 111; cf. this vol., ii, 170—171).—Blood is freed from albuminous substances and a volume of the filtrate corresponding with 2 c.c. of plasma, clot, or total blood is treated with 1—2 c.c. of 0.9—1.0% sodium chloride solution and sufficient 40% sodium carbonate solution to give an alkaline reaction to litmus. Five c.c. of Folin's silver reagent are added, and, after mixing, the liquid is centrifuged. The clear liquid is removed and the precipitate treated with 2 c.c. of a solution of sodium chloride containing hydrochloric acid and 7—8 c.c. of water, and again centrifuged. The clear liquid obtained is treated with 0.5 c.c. of a solution of 2.5 grams of sodium cyanide and 5 grams of sodium sulphite in 100 c.c. of water, 1.5 c.c. of sodium carbonate solution, made up to 12.5 c.c., and, after addition of Folin and Denis's phosphotungstic acid reagent, compared with a standard uric acid solution. G. W. R.

Estimation and Constitution of a Fraction of the Uric Acid in Blood. CH. O. GUILLAUMIN (*Compt. rend. Soc. biol.*, 1922, 86, 258—260; from *Chem. Zentr.*, 1922, iv, 111; cf. preceding abstract).—Direct estimation of uric acid in serum freed from albuminous substances gives higher results than those obtained by the Folin and Denis phosphotungstate method. The difference corresponds with a portion of the uric acid of the blood which is associated with fractions of the original nuclein complex. For its estimation (by difference) the preliminary removal of albuminous substances is preferably effected by using metaphosphoric acid.

G. W. R.

The Volatilisation and Hydrolysis of Atropine in Toxicology. PAUL HARDY (*J. Pharm. Chim.*, 1922, 26, 220—226).—The author has investigated the error in estimations of atropine

due to losses by reason of volatility and hydrolysis, and finds that no loss occurs in extraction of the alkaloid, as it is not volatile in alcohol, ether, or chloroform vapour. In the case of water, atropine was detected in the distillate on boiling. Hydrolysis occurs readily in aqueous solution, increasing considerably with temperature; it is more rapid in alkaline than in neutral solution, but ammonia is not such an active hydrolytic agent as sodium hydroxide.

H. J. E.

The Estimation of Creatinine. K. PFIZENMAIER and S. GALANOS (*Z. Unters. Nahr. Genussm.*, 1922, 44, 29—41).—The method given in the *Schweizerisches Lebensmittelbuch* (3rd ed., p. 72) for the estimation of creatinine is rapid and simple, but can be successfully used only when light-coloured solutions are available for colorimetric comparison and a colorimeter of the Duboscq type is used. If the solutions are dark in colour or contain sugar, the method fails, as animal charcoal cannot be used because it adsorbs creatinine. The colorimetric comparison in cylinders as specified in the book was found to be unsatisfactory. The method of Sudendorf and Lahrmann (*Z. Unters. Nahr. Genussm.*, 1915, 29, 1) can always be used and is the only one available for use with dark-coloured solutions. If the authors' instructions are exactly followed and too great an excess of potassium permanganate—especially with very dark-coloured solutions—is avoided, accurate results are always obtained. The methods referred to are given in detail.

H. C. R.

An Apparatus for the Extraction of Theobromine and Caffeine by means of boiling Chloroform. O. P. A. H. SCHAAPE (*Pharm. Weekblad*, 1922, 59, 920—923).—The Soxhlet extractor is modified to allow of heating by means of a water-bath. Exact dimensions and procedure are laid down (cf. *J. Soc. Chem. Ind.*, 1922, 781A).

S. I. L.

Estimation of Yohimbine in Yohimba Bark. ARNOLD SCHOMER (*Pharm. Zentr.-h.*, 1922, 63, 385—386).—In applying a method described previously (A., 1921, ii, 360) to barks containing but little yohimbine, the results obtained are more trustworthy when the crystallisation of the yohimbine hydrochloride is made in the cold. The impure hydrochloride, after treatment with 5 c.c. of ether and evaporation of this solvent, is dissolved in 3 c.c. of absolute alcohol and the solution is kept at 0° for eighteen hours. The crystals which form are collected, washed with a small quantity of chloroform, then with ether, dried at 100°, and weighed.

W. P. S.

Identification of Traces of True Albumin in Urine. Separation of pseudoAlbumins. M. RENOUX (*J. pharm. Belg.*, 1922, 4, 381—382).—As a result of tests with a solution of egg-albumin, the author concludes that either neutral crystals of sodium sulphate or magnesium sulphate may be used for the defecation of urine

(after being made alkaline), preparatory to testing for traces of true albumin.

CHEMICAL ABSTRACTS.

The Estimation of Total Albumin in the Spinal Fluid. J. B. AYER and H. E. FOSTER (*Rept. pharm.*, 1922, **34**, 114—115).—The albumin is precipitated with thiosalicylic acid and compared with the precipitate obtained from blood serum of known albumin content. The method is very exact. The normal quantity of albumin in 100 c.c. of fluid varies between 16 and 40 mg. (average 25 mg.). It is usually increased in pathological conditions. The authors give the albumin content found in a number of cerebrospinal diseases.

CHEMICAL ABSTRACTS.

The Estimation of Albumose-Silver. J. HERZOG (*Pharm. Z.*, 1922, **67**, 802—803).—One gram of the silver-protein is dissolved in 10 c.c. of water and 10 c.c. of concentrated sulphuric acid added in a thin stream. Two grams of finely powdered potassium permanganate are added in small portions with vigorous shaking. After fifteen minutes, the solution is diluted with 50 c.c. of water and ferrous sulphate added in small portions to decolorise the solution. The silver is then titrated with *N*/10-ammonium thiocyanate solution, the oxidised iron acting as indicator.

H. K.

The Estimation of Fibrinogen. Protein Estimations in Salt Plasma. Question of the Utility of Serum for the Quantitative Investigation of Blood. G. LEENDERTZ and B. GROMELSKI (*Arch. exp. Path. Pharm.*, 1922, **94**, 114—123).—Indirect methods for the estimation of fibrinogen in blood are based on the fact that the fibrinogen content is equal to the difference between plasma- and serum-proteins. Since, however, serum formed by the spontaneous coagulation of blood contains water and chlorides emanating from the corpuscles, this relation only holds when the serum is obtained from separated plasma. Applications of the refractometric method to the indirect estimation of fibrinogen have not hitherto been successful when oxalate plasma has been used. Such failures have been due to two causes—the formation of a precipitate and the production of hypertonic plasma by the addition of sodium oxalate to blood. By using a 3.55% solution of sodium citrate, which is isotonic with blood and does not produce a precipitate, as anti-coagulant, the refractometric method may be readily applied. Two such methods, based on the above considerations, are described.

E. S.

Estimation of Hæmoglobin. E. MEULENGRACHT (*Fol. hæmatol. I. Archiv*, 1921, **27**, 1—9; from *Chem. Zentr.*, 1922, ii, 849).—Observations on the working details of hæmoglobin estimations by Sahli's and by Autenrieth's methods. It is held that the former is superior to the latter for clinical work.

G. W. R.

The Colorimetric Estimation of Hæmoglobin with Especial Reference to the Production of Stable Standards. EDWIN H. TERRILL (*J. Biol. Chem.*, 1922, 53, 179—191).—Stock solutions of acid hæmatin such as those of Cohen and Smith (A., 1919, ii, 532) and Robscheit (A., 1920, ii, 339) suffer from the disadvantages that the colour both changes and fades on keeping, whilst a turbidity is produced on dilution. Two modified methods for the preparation of acid hæmatin standards are therefore described. The first results in the formation of a concentrated stock solution which may be diluted, especially when the diluting fluid contains glycerol, without the production of a turbidity; although it fades slowly, it does not change qualitatively in colour. The second gives an acid hæmatin protein powder which is quite stable in the dry state and forms clear solutions in water and in 0.1*N*-hydrochloric acid. After standardisation, weighed quantities may be used for the preparation of standard solutions. A method is also described for the preparation from the powder of gelatin films of acid hæmatin which may be used with advantage in place of the glass plates of Newcomer (A., 1919, ii, 179).

In estimating hæmoglobin, the laking of the blood with water prior to acidification is recommended. Direct acidification produces a turbidity which leads to high results when a clear standard is used. Heat should not be employed to develop maximum colour as this again produces a turbidity. E. S.

Estimation of Bilirubin in Blood. PAUL HOLZER and HEINZ MEHNER (*Klin. Woch.*, 1922, 1, 66).—Meulengracht's (*J. Amer. Med. Assoc.*, 1920, 74, 68) method is worthless for the estimation of small amounts of bilirubin, and untrustworthy for the estimation of large amounts. To obtain accurate results it is necessary to enhance the bilirubin colour by diazotisation; Bergh's ("Der Gallenfarbstoff im Blute," 1918) method is the most trustworthy, whilst that of Haselhorst (A., 1921, ii, 472) gives inaccurate results for small concentrations of bilirubin.

CHEMICAL ABSTRACTS.

[Detection of Bile Pigments in] Gastric Juice. FRANZ UTZ (*Süddeutsch. Apoth.-Ztg.*, 1922, 62, 77—78; from *Chem. Zentr.*, 1922, ii, 921—922).—In examination of the contents of the stomach for bile pigments, the filtrate should not be used, as the pigments are retained by the filter. Directions are given for the use of the Gmelin and Plesch tests for bile pigments and for the detection of rennin. G. W. R.

Modification of the Iodine Test for Bile Pigments in Urine. ERNST SILBERSTERN (*Zentr. inn. Med.*, 1922, 43, 185—187; from *Chem. Zentr.*, 1922, ii, 850).—The ordinary reaction is modified, whereby the urine under examination is shaken with a solution of 0.5 gram of iodine in 36 grams of ether or 76.5 grams of chloroform (1 c.c. of this solution to 2—4 c.c. of the liquid to be tested). The excess of iodine remains in its original solvent whilst the

biliverdin colours the aqueous phase. In neutral solution, the reaction is more sensitive than the Gmelin test, although less sensitive in alkaline solution. The chloroform solution is preferable to the ethereal solution.

G. W. R.

Tests for Liver Function. LEPEHNE (*Münch. Med. Woch.*, 1922, 69, 342—344; from *Chem. Zentr.*, 1922, ii, 921).—Hay's flowers of sulphur test for bile acids in duodenal juice and urine is described in application to certain pathological conditions.

G. W. R.

Estimation of the Activity of Invertase. T. SWANN HARDING (*Sugar*, 1922, 24, 89—90; from *Chem. Zentr.*, 1922, ii, 1032).—A solution of sucrose, approximately 10%, is prepared so that on addition of 5 c.c. of water to 50 c.c. of the solution a reading of $+33.5^\circ V \pm 1^\circ$ is given in Schmidt and Haensch's polarimeter. The reading being accurately determined after acidifying with acetic acid, 5 c.c. of invertase solution are added and, after keeping at 30° for ten minutes, the solution is rendered alkaline by addition of sodium carbonate solution. The change in polarimeter reading is taken as a measure of the activity of the invertase solution. Ordinary invertase solutions change the polarimeter reading from $+33.5^\circ V$ to $+23-10^\circ V$, the most active invertases giving $10^\circ V$. Invertases which diminish the reading to $+18-20^\circ V$ invert a 10% sucrose solution within half an hour.

G. W. R.

Analysis of Blood. F. UTZ (*Pharm. Zentr.-h.*, 1922, 63, 425—430).—A procedure is given for a complete examination of blood, and the methods used are described in detail. The following estimations should be made, the figures in brackets being the values obtained for normal blood: *Blood*, sp. gr. (1.050 to 1.060) and total solids (21 to 22%). *Blood-serum*, sp. gr. (1.027 to 1.032), refractive index (n_D 1.343 to 1.350), freezing point (-56°), total solids (6 to 10%), total nitrogen (1.04 to 1.2%). *Blood-serum after removal of proteins by uranium acetate*, nitrogen (0.02 to 0.035%), urea (0.02 to 0.04%), uric acid (0.0025 to 0.0035%), creatinine (0.001%), reducing sugars (0.06 to 0.12%), sodium chloride (0.56 to 0.60%), and indican (0.045 mg. per 100 c.c.).

W. P. S.

General and Physical Chemistry.

The Refractive Indices of Phosphorescent Sulphides. MAURICE CURIE (*Compt. rend.*, 1922, 175, 617—619).—Direct determination of the refractive indices of phosphorescent sulphides gives, for yellow light, values between 2.10 and 2.15. An appreciable fraction of each substance gives figures ranging to a lower limit of 1.6, whilst for no portion did the refractive index exceed 2.20. According to the work of Lenard (*A.*, 1910, ii, 369) and Schmidt (*Ann. Physik*, 1922) the values should lie between 2.70 and 3.16, so that Lenard's interpretation of the constancy of the ratio λ/\sqrt{K} does not appear to be well founded. H. J. E.

Quantum Theory of Line Spectra. N. BOHR (*Danske Vid. Selsk. Skrifter, nat. mat. Afd.*, 1918, 4, [8], 1—100; from *Chem. Zentr.*, 1922, iii, 221—223).—Theoretical. The author develops his quantum theory of line spectra with special reference to the hydrogen spectrum, and to the Stark and Zeeman effects.

G. W. R.

The Intensities of the Lines in the Balmer Series of Hydrogen. MASAMICHI KIMURA and MITSUHARA FUKUDA (*Mem. Coll. Sci. Kyoto*, 1922, 5, 165—168).—The intensity distribution of the hydrogen lines in the Balmer series is greatly affected by the pressure of hydrogen. The addition of the vapours of hydriodic acid, iodine, sulphur, mercury, or sodium exerts an effect similar to that obtained on increased pressure. These gases suppress the higher members of the Balmer series and appear to exert a specific effect on the hydrogen atoms. The results on the action of iodine vapour are in disagreement with those of Holtzmark (*A.*, 1918, ii, 283).

W. E. G.

The Broadening of the Balmer Lines of Hydrogen with Pressure. E. O. HULBERT (*Astrophys. J.*, 1922, 55, 399—405).—Experimental evidence is adduced for abandoning, as a cause which interferes with the absolute homogeneity of spectrum lines, the consideration of disturbance depending on collision with other particles, and suggesting that the principal cause of the widening of the lines is the influence of an electric field on the radiating particle (cf. Merton, *Proc. Roy. Soc.*, 1915, 92, 322).

A. A. E.

Influence of Gases and Vapours on the Intensities of the Lines of the Secondary Spectrum of Hydrogen. MASAMICHI KIMURA and MITSUHARA FUKUDA (*Mem. Coll. Sci. Kyoto*, 1922, 5, 153—163).—The intensities of the lines in the hydrogen secondary spectrum are modified by pressure, and the addition of gases and vapours. Increase in pressure weakens those lines in the secondary spectrum on the long wave-length side of λ 6240, and in the region

between $\lambda 5690$ and H_{β} . Between these two regions, the lines undergo selective changes in intensity. The Fulcher group I lines are relatively intense at low pressures, whilst those belonging to group II are weak. The addition of helium, mercury, or bromine to the spectrum tube produces similar changes in the intensities of the lines to those obtained on increase of pressure, although helium exerts a specific effect on some of the hydrogen lines. The suggestion is made that the hydrogen molecule can be reformed either by the combination of a positively charged molecule and an electron, or by the association of two neutral atoms, and that the two groups of lines owe their existence to the occurrence of these two distinct processes. W. E. G.

The Continuous Spectrum of Hydrogen in the Schumann Region. E. P. LEWIS (*Physical Rev.*, 1920, **16**, 367—368; cf. *Science*, 1915, **41**, 947).—By the use of a fluorite vacuum spectrograph it has been found that the continuous spectrum of hydrogen extends with undiminished intensity well into the Schumann region. At $\lambda 1800$ it begins rather suddenly to diminish in intensity, and disappears at about $\lambda 1750$. Slightly below $\lambda 1700$, the line spectrum of hydrogen reappears, with no trace of a continuous background. Attention is directed to the fact that the continuous spectrum completely fills the region between the Balmer series and the Schumann line spectrum, with no lines whatever superimposed on it. The perfect uniformity and continuity of this spectrum make it the best background for the examination of absorption spectra (for example, of benzene) in the ultra-violet. Helium and neon give similar, although less intense, continuous spectra extending far into the ultra-violet. A. A. E.

Identification of Air Lines in Spark Spectra from $\lambda 5927$ to $\lambda 8683$. PAUL W. MERRILL, F. L. HOPPER, and CLYDE R. KEITH (*Astrophys. J.*, 1921, **54**, 76—77).—Data have been secured (cf. *ibid.*, 1920, **51**, 211) for the chemical identification of the air lines from $\lambda 5927$ to $\lambda 8683$ in spark spectra. The spark was operated alternately in oxygen and air, and the lines due to oxygen identified. The remaining air lines, except a few argon lines, are ascribed to nitrogen. A. A. E.

The Intensity Distribution in the β -Ray Spectra of Radium-*B* and -*C*. J. CHADWICK and C. D. ELLIS (*Proc. Camb. Phil. Soc.*, 1922, **21**, 274—280).—Two radically different views of β -ray disintegration have been put forward. That of Meitner (this vol., ii, 416) in which the entire β -ray emission of a radioactive substance consists of a series of homogeneous groups of electrons, and that of Ellis (this vol., ii, 466) according to which the disintegration electrons form a continuous spectrum. In the present paper, measurements are made of the intensity distribution in the β -ray emission of radium-*B* and -*C*, and it is shown that the continuous spectrum has a real existence, and is independent of the experimental arrangement. The five strong lines in the radium-*B* β -ray spectrum formed only one-fifth of the total emission,

the main feature of which was the continuous spectrum. The number of electrons received per second in the ionisation chamber, calculated from the observed ionisation currents, was in agreement with the number of atoms disintegrating per second at the source.

W. E. G.

The Spectrum of Radium Emanation. R. E. NYSWANDER, S. C. LIND, and R. B. MOORE (*Astrophys. J.*, 1921, **54**, 285—292).—Photographic and visual measurements were made of the spectrum of radium emanation, $\lambda\lambda$ 3982 to 7450. The relative intensities of the lines differed markedly, as a rule, from the previous observations of Watson and Royds and Rutherford; many strong lines of the one hundred and twenty previously measured were not found; of the forty-four lines obtained, nine were new. Changes of intensity with duration of the discharge were observed, some lines decreasing and some increasing in strength, whilst the colour of the discharge changed from a bright glow to violet. A modification is described of Duane's apparatus for the purification of radium emanation.

A. A. E.

The Spectrum of Neutral Helium. C. V. RAMAN (*Nature*, 1922, **110**, 700—701).—A criticism of Silberstein's attempt (this vol., ii, 674) to explain the spectrum of neutral helium on the assumption of the independence of the electrons. The view is expressed that the approximate agreements between the calculated and actual frequencies are merely fortuitous arithmetical coincidences, and a number of statements are made, based on a survey of the figures and a series of computations, in its support. The same considerations apply with even greater force in the case of lithium, when a choice of six numbers, compared with four in the case of helium, is permitted. Further, it is held that the Rydberg constant 109723 is appropriate only to the case of the ionised helium atom in which only one electron is coupled to the nucleus.

A. A. E.

The Structure of the Red Lithium Line. T. R. MERTON (*Nature*, 1922, **110**, 632).—A criticism of the views of McLennan and Ainslie (this vol., ii, 541; cf. Zeeman, A., 1913, ii, 812; *Proc. K. Akad. Wetensch. Amsterdam*, 1913, **15**, 1130; Kent, *Astrophys. J.*, 1914, **40**, 337; King, *ibid.*, 1916, **44**, 169; Takamine and Yamada, *Proc. Tokio Math. Phys. Soc.*, 1914, **7**, 339; Merton, *Proc. Roy. Soc.*, 1921, [A], **99**, 101) on the structure of the lithium line $\lambda=6708$. Under the appropriate conditions, the line appears to be a simple pair; if the four components were really two pairs due to the two isotopes, they should always appear together with an invariable intensity ratio of 1:16. This is not in accordance with the experimental results. Theoretical objections are also raised.

A. A. E.

The Exploded-wire Spectrum of Calcium. R. A. SAWYER and A. L. BECKER (*Physical Rev.*, 1921, **18**, 164).—By expending constant power on wires of varying mass of calcium and other elements, or varying power on wires of constant mass, spectra

have been obtained in which the relative intensities of the three calcium lines, *H*, *K*, and λ 4227, vary as in the spectra of stars of different classes. The calcium lines, particularly *H* and *K*, appear to be tremendously enhanced by the very high temperature of the explosion, since they appear more prominently in the spectra of exploded wires of other metals (in which the amount of calcium present must be very minute) than the spectra of the metals themselves.

A. A. E.

Excitation of Atoms to Light Emission by Electron Collisions. R. SEELIGER (*Z. Physik*, 1922, 11, 197—200).—According to the Bohr theory, the passage of an electron from an initial to a final orbit is determined by the two probabilities, *W* and *w*, where *W* is the probability that the electron shall be raised from the normal to the initial orbit, and *w* is the probability of the transition from the initial to the final state. The former (*W*) is affected by the conditions of excitation, and according to Franck, the latter (*w*) should be also dependent on the external conditions. To test this point, the effect of the velocity of the exciting electrons on the intensity of certain related mercury lines has been examined. Two pairs of lines, 4916 ($3\cdot5S-2P$) and 2857 ($3\cdot5S-2p_2$), and 2652 ($4d'-2p_2$) and 4339 ($4d'-2P$) were chosen, and the intensities of these measured in different parts of the glow-discharge. The form of the intensity curves were widely different for the individuals of both pairs, and the maxima occurred at different distances from the mercury electrode. Thus, in these cases, the probability of transition from one orbit to another is dependent on the external electric field.

W. E. G.

Mass-absorption Coefficients as a Function of Wavelength Above and Below the *K* X-Ray Limit of the Absorber. F. K. RICHTMYER (*Physical Rev.*, 1921, 17, 264—265).—An extension to silver and lead, and to longer wave-lengths, of the relation between X-ray wave-lengths and the mass-absorption coefficient μ/ρ (for aluminium, copper, and molybdenum, cf. this vol., ii, 105).

A. A. E.

The Evidence Regarding the so-called "*J*" Radiation in the Characteristic X-Ray Spectra of the Elements. F. K. RICHTMYER (*Physical Rev.*, 1921, 17, 433—434).—From an examination of the absorption of water, aluminium, copper, boron, silver, and molybdenum, the author concludes that, within 1%, there are no discontinuities in absorption suggestive of *J*-radiations.

A. A. E.

Absorption of X-Rays by Chromium, Manganese, and Iron. WILLIAM DUANE and HUGO FRICKE (*Physical Rev.*, 1921, 17, 529—531).—The *K* critical absorption wave-lengths ($\lambda \times 10^8$ cm.) of chromium, manganese, and iron are 2.0623, 1.8893, and 1.7377, respectively. The energy changes corresponding with the critical absorption appear to be the same for iron atoms in the bi- and ter-valent conditions.

A. A. E.

The Tungsten X-Ray Spectrum with a Mica Spectrometer. USABURO YOSHIDA and SHINSUKE TANAKA (*Mem. Coll. Sci. Kyoto*, 1922, 5, 173—178).—The arrangement is essentially that used previously (*ibid.*, 1921, 4, 343) and the grating constant was calculated from the wave-lengths of the prominent lines of the *L*-series of tungsten. With the mica spectrometer the X-ray spectrum of tungsten may be obtained up to the seventh and even to the tenth order. Eleven lines of unknown origin were detected on the photographs, but these might be due to reflection from the mica in an unknown manner. The wave-lengths of the known lines are in agreement with those given by Siegbahn within the limit of the experimental error. The existence of the line found by Overn at 1070 Å. has been confirmed. W. E. G.

Critical Potentials of the *L*-Series of Platinum. DAVID L. WEBSTER (*Physical Rev.*, 1920, 15, 238).—The *L*-series must be regarded as consisting of three sub-series, *L*₁ (containing the lines *l*, α_2 , α_1 , β_2 , β_5 , and β_6), *L*₂ (containing the lines γ , β_4 , β_1 , γ_1 , γ_2 , and possibly β_3), and *L*₃ (containing the lines γ_4 , γ_3 , and possibly β_3). A. A. E.

Temperature Shift in Near Ultra-red Bands. H. M. RANDALL, W. F. COLBY, and R. F. PATON (*Physical Rev.*, 1920, 15, 541—543; cf. Randall and Imes, A., 1920, ii, 570; Colby, A., 1920, ii, 655).—Curves are given showing the absorption of hydrogen chloride in the neighbourhood of 3.4 μ at 20°, 105°, and 250°. It appears that in spite of the outward shift of intensity to higher velocity values, no shift takes place in the fine structure. Further investigations at higher temperatures are in progress. A. A. E.

The Labile Nature of the Halogen Atom in Organic Compounds. VII. Absorption Spectra of the Halogen Derivatives of some Cyclic Compounds, and their Bearing on the Question of an Oxygen-Halogen Linking. HUGH GRAHAM and ALEXANDER KILLEN MACBETH (T., 1922, 121, 2601—2608).

Carbon Dioxide Absorption in the near Ultra-red. E. F. BARKER (*Astrophys. J.*, 1922, 55, 391—398).—New absorption curves have been obtained. The 2.7 μ region, previously considered to be a doublet, proves to be a pair of doublets, with centres at 2.694 μ and 2.767 μ (approx.). The 4.3 μ band appears as a single doublet with centre at 4.253 μ . The frequency difference between maxima is nearly the same for each of the three doublets, and equal to 4.5×10^{11} . Complete resolution of the band series was not effected, but there is evidently a complicated structure, with a "head" in each case on the side of shorter wave-lengths. If a linear configuration of the three atoms in the molecule of carbon dioxide is assumed, the moment of inertia of the molecule, computed from the doublet frequency difference, is 50×10^{-40} . This gives the distance of the central carbon atom from each oxygen atom as 0.97×10^{-8} cm. Since the linear molecule could have only three modes of vibration, it is suggested that the two doublets

at $2.7\ \mu$ may correspond with two successive changes in stationary state for the same vibration. A. A. E.

Absorption of Light by Inorganic Salts. E. F. GEORGE (*Diss., Ohio*, 1920).—The change in absorption produced by mixing sulphates in solution with other sulphates is small in comparison with corresponding changes in nitrates, and still smaller than in the case of chlorides. The influence of temperature on the absorption of solutions of mixed salts is greatest with chlorides. It therefore appears that Ostwald's view that the coloured ion alone is effective in the absorption of light is not well founded. When the ratio of the absorption of the mixture to the sum of the absorptions of the component parts is plotted as a function of the wavelength, the resulting curve generally shows a minimum near the middle of the spectrum. Many of the curves are shown to be exponential in form. CHEMICAL ABSTRACTS.

The Absorption Spectra of *o*-Cresolsulphonphthalein. W. R. ORNDORFF, R. C. GIBBS, M. SCOTT, and S. D. JACKSON (*Physical Rev.*, 1921, **17**, 437).—*o*-Cresolsulphonphthalein and other related compounds have in neutral aqueous solutions two absorption bands which are modified or replaced by new bands on the addition of acid or alkali. In the case of a dilute alkaline solution, the new type of absorption is not stable, but reverts to the two band absorption found in the corresponding neutral solution. It is considered that in neutral aqueous solutions the carbinol and hydrate forms of the phthalein are present and that on the addition of either acid or alkali a salt having a quinoid structure is formed. A. A. E.

Flame Excitation of Luminescence. E. L. NICHOLS and D. T. WILBER (*Physical Rev.*, 1921, **17**, 453—468).—The fluorescence developed by certain substances, notably lime, zirconia, magnesia, silica, alumina, zinc oxide, some phosphorescent sulphides, and a number of other compounds, by contact with a hydrogen flame, differs from photoluminescence and from mere temperature radiation. A definite temperature-range exists for each substance, and the effect does not appear to depend on the presence of traces of activating elements. Many substances which are strongly luminescent under the action of light do not respond to flame excitation. Excitation does not occur by heating in air or hydrogen or in a vacuum, or by contact with a jet of hydrogen while hot unless the hydrogen is ignited and conditions are favourable for free oxidation. The effect probably cannot be ascribed to the presence of H or H_2 molecules. The spectra obtained by flame excitation have been examined. A. A. E.

Chemoluminescence. I. A. A. GRINBERG (*J. Russ. Phys. Chem. Soc.*, 1920, **52**, 151—185).—A summary is given of previous work on this subject, particularly on bioluminescence (cf. Radziszewski, A., 1877, ii, 345; Lenard and Wolf, A., 1888, 1000; Trautz, A., 1905, ii, 662; Trautz and Schorigin, A., 1905, ii, 494; Dubois, A., 1913, i, 1021; Harvey, A., 1917, i, 365; ii, 436; Goss, A., 1917, ii, 436). The light effect obtained with the system, pyro-

gallol-hydrogen peroxide-potassium permanganate under various conditions has been investigated, the intensity of the light being measured photographically. At 16–17° the greatest intensity was obtained with a mixture of 4 c.c. of 0.005*M*-potassium permanganate, 2 c.c. of 0.01*M*-pyrogallol, and 2 c.c. of 3–12% hydrogen peroxide solutions. Sulphuric acid in the concentration 0.018–0.04% weakens the effect considerably, whilst 0.0046–0.026% of potassium hydroxide enhances it to a slight extent. T. H. P.

Mutarotation. I. C. N. REIBER (*Ber.*, 1922, 55, [B], 3132–3143).—The data with regard to the relationship of mutarotation to other physical changes in solution are very scanty and exhibit considerable discrepancies among themselves. The changes in refractive index and volume of a solution containing a substance undergoing mutarotation have now been studied.

The index of refraction of an aqueous solution of dextrose does not appear to vary when the ordinary Pulfrich refractometer is used, but the instrument is scarcely suitable for the purpose owing to the impossibility of maintaining the requisite degree of constancy in temperature. By means of a Hallwachs's prism, it can, however, be shown that the refractive index increases slightly with age in the case of dextrose solutions and that the change occurs with immeasurably greater rapidity when a little ammonia is added to the solution. The method, however, does not lend itself to the periodical observation of the action and attention has therefore been directed to the measurements of alteration of volume in the solution. The dilatometer employed resembles essentially a Sprengel pycnometer. One tube is firmly closed by a small rubber plate and the change in volume of the solution is indicated by the movement of the meniscus in the other limb and measured by means of a microscope. A movement of 0.005 mm. can be detected with certainty; this corresponds with an alteration in volume of less than 0.0000001. The capacity of the instrument is 66.7 c.c. The temperature is maintained constant to within 0.001°. It is thus shown that solutions of dextrose in water increase in volume during the process of mutarotation at a rate corresponding with that required for a unimolecular, irreversible reaction but subsequent observations prove that in reality a state of equilibrium is finally attained and indicate that water possibly takes some part in the change. The quantitative parallelism between mutarotation and change of volume makes it certain that the two phenomena have a common cause.

A solution of lactose the optical activity of which diminishes progressively shows a gradual expansion, whereas a contraction is observed with a solution of maltose with increase of specific rotation. Galactose solution suffers a primary expansion and a subsequent contraction. Solutions of sucrose and mannitol which do not exhibit mutarotation are constant in volume. Solutions of the optically inactive dulcitol do not undergo change in volume. Acetaldehyde, in 30 per cent. aqueous solution, exhibits a considerable expansion, whereas acetone under similar conditions gives

a marked and rapid contraction. Ethyl alcohol gives a slight expansion.

The applicability of the dilatometer just described is not restricted to reactions involving mutarotation, but since changes in volume are probably involved in all chemical reactions, it can be used for the investigation of processes taking place at the ordinary temperature. The accuracy of the procedure is at least as great as that of the polarimetric or refractometric method and it has the great advantage that it does not necessitate the use of transparent and colourless solution. It has been found very useful in the study of the inversion of sugar by invertin and the saccharification of starch by ptyalin; in each case a marked contraction takes place as is to be expected.

H. W.

Photochemistry of Chlorine Detonating Gas. FRITZ WEIGERT and KARL KELLERMANN (*Sitzungsber. Preuss. Akad. Wiss. Berlin*, 1922, 24, 315—320).—The early stages of the combination of hydrogen and chlorine have been investigated. It is shown that a characteristic streaky appearance is visible in the gas mixture during the time $1/100$ — $1/20$ second from the moment of excitation. A vanishingly small quantity of hydrogen chloride only is formed directly by the radiation, but this is followed by a large subsequent formation. From this result, the authors show that the primary reaction, contrary to many previous assertions, in all probability is in keeping with Einstein's photochemical equivalent law.

J. F. S.

Photochemical Oxidation of Hydriodic Acid. CHR. WINTHER (*K. Danske Vid. Medd. Math.-fys. Medd.*, 1920, 2, No. 2, 1—28; from *Chem. Zentr.*, 1922, iii, 592).—In the oxidation of hydriodic acid solutions in thin films, the velocity of reaction was measured at the beginning and also in the course of the reaction. The effect of shaking, oxygen concentration, and wave-length of light was also investigated. With vigorous shaking, the reaction shows an increase in velocity during an induction period and then remains constant. The sensitiveness of the oxidative process to light is proportional to the absorption of tri-iodide-ion which acts as an optical catalyst. This is an example of "optical autosensibilisation" where the catalyst is actually formed by the reaction which it catalyses. The oxidation of leuco-compounds to dyes, the oxidation of pyrogallol in alkaline solution, and the oxidation of sodium sulphide in aqueous solution are similarly catalysed.

G. W. R.

Photochemical Efficiency of Absorbed Radiation. CHR. WINTHER (*K. Danske Vid. Medd. Math.-fys. Medd.*, 1920, 2, No. 3, 1—35; from *Chem. Zentr.*, 1922, iii, 592—593).—The amount of energy absorbed in the photochemical oxidation of a certain amount of hydrogen iodide was found to be identical with the amount calculated as necessary for its activation. The absorbed energy is thus used exclusively for the activation of the hydrogen iodide molecule. The equivalence of absorbed energy and energy required

for activation is enunciated as a general principle in photochemical reactions. Certain consequences of this principle are discussed. Discrepancies are attributed to experimental errors or to the conversion of the absorbed radiation into radiation of shorter wave-length.

G. W. R.

The Decomposition of Hydrogen Molecules by Excited Mercury Atoms. G. CARIO and J. FRANCK (*Z. Physik*, 1922, **21**, 161—166).—On the collision of excited atoms with other atoms, the quantum energy may be given up without light emission (cf. Franck, this vol., ii, 464), and chemical change may take place. On exposing a mixture of hydrogen and mercury to the mercury line $2536\cdot7 \text{ \AA}$., the radiation was strongly absorbed, and the hydrogen dissociated. Powdered copper oxide was reduced, and the pressure of the hydrogen continuously decreased at a temperature of 45° . In the absence of hydrogen and mercury, no reduction occurred. The mercury is acting as a sensitiser. In the absence of copper oxide, the atomic hydrogen is adsorbed by the walls. At low pressures, the velocity of dissociation was very slow, and this behaviour indicates a short period of activity of the mercury atoms. The velocity of reaction increases up to 15 mm. pressure and then remains constant. Tungsten trioxide is also reduced, and the reduction is not in any way due to illumination by the light. Since tungsten oxide is not reduced by hydrogen below a red heat, it is unlikely that the phenomenon is caused by rapidly moving hydrogen molecules.

The hydrogen is dissociated by mercury atoms which are in the $2p_2$ condition. Calculations show that the upper limit for the heat of dissociation of hydrogen is 112 Cal.

W. E. G.

Photoelectric Effect of Alkali [Metal] Vapours and a New Determination of h . E. H. WILLIAMS and JAKOB KUNZ (*Physical Rev.*, 1920, **15**, 550).—The resonance potential of caesium vapour is 1.48 volts, whilst the ionisation potential is 3.9 volts. If the same amount of work is required for the breaking up of a neutral caesium atom by light as by moving electrons, then the long wave-length limit of the photoelectric effect of caesium vapour should be $l=318\cdot4 \mu\mu$, according to the equation $hn=eV$. If this is found to be the case, then (1) the long wave-length limit is quite different for caesium metal and for caesium vapour, and (2) a new photoelectric method for the determination of h is available. Experimentally, there is no photoelectric effect in caesium vapour down to $313 \mu\mu$, but the effect at $253 \mu\mu$ is very marked.

A. A. E.

The Photoelectric Long Wave-length Limit of Platinum and Silver. OTTO STUELMAN, jun. (*Physical Rev.*, 1920, **15**, 549—550; cf. *ibid.*, 1919, **13**, 109).—The photoelectric long wave-length limits of platinum and silver are $\lambda_0=284 \mu\mu$ and $\lambda_0=325 \mu\mu$, respectively.

A. A. E.

Soft X-Rays of Characteristic Type. E. H. KURTH (*Physical Rev.*, 1921, **17**, 528—529; **18**, 99—100).—The radiation curve of

aluminium shows breaks at 38 and 120 volts, corresponding with 326 Å.U. (possibly the *M*-series) and 103 Å.U. (possibly the *L*-series), respectively. For iron, 62.8 Å.U. and 48.4 Å.U. appear to represent the extremes of the *M*-series, whilst 16.3 Å.U. probably corresponds with the *L*-series. There was also some indication of a break in the curve at 50 volts, corresponding with 247 Å.U. Carbon yields a pronounced break at 43.6 Å.U., agreeing with the extrapolated value for the *K* series. The following approximate figures (Å.U.) are also recorded: *K*-series: carbon, 42.6; oxygen, 23.8. *L*-series: carbon, 375; oxygen, 248; iron, 16.3; copper, 12.3. *M*-series: iron, 54.3; copper, 41.6. *N*-series: iron, 247; copper, 119. A. A. E.

Absorption of X-Rays in Crystals. TYCHO E:SON AURÉN (*Medd. K. Vetenskapsakad. Nobel-Inst.*, 1922, 4, No. 10, 1—6).—It has been shown in a previous paper (A., 1921, ii, 367) that, in the case of amorphous substances, the molecular absorption coefficient can be calculated as a sum of the atomic absorption coefficients for X-rays of the constituent elements. It is now shown that the same holds true for a number of crystalline substances, including alum, calcite, and gypsum. The observed absorption coefficients agreed with considerable accuracy with those calculated from the atomic absorption coefficients derived from a study of amorphous substances, and the absorption showed no variation in different directions in the crystals. In the cases of rock salt and sylvine an abnormally low absorption was observed when the incident rays were parallel to the principal planes of the crystal. This effect, which is probably due to scattering, becomes less marked as the angle between the direction of the incident rays and the principal structural planes is increased. E. H. R.

The Durability of Radium Solutions. AUG. BECKER (*Z. anorg. Chem.*, 1922, 124, 143—152).—The permanency of the radium solution is defined as its constancy in emitting emanation. The author found that solutions containing from 0.02 to 0.0000005 mg. radium per gram solvent did not show any decay in eight years. The possible experimental error was 1%. W. T.

Electric Furnace Experiments involving Ionisation Phenomena. ARTHUR S. KING (*Astrophys. J.*, 1922, 55, 380—390; cf. Saha, A., 1920, ii, 457, 659; 1921, ii, 4, 162; *Proc. Roy. Soc.*, 1921, [A], 99, 135).—Results are obtained which verify certain predictions due to Saha. It is shown that the intensity of the enhanced lines of alkaline-earth metals is decreased by the presence of the more easily ionised potassium and caesium. The results with mixed gases are such as would be expected if it is assumed, with Saha, that flame and enhanced lines are associated with neutral and ionised atoms, respectively. A. A. E.

Thermal Ionisation of Gaseous Elements at High Temperatures. Confirmation of Saha's Theory. ARTHUR A. NOYES and H. A. WILSON (*Proc. Nat. Acad. Sci.*, 1922, 8, 303—307).—On the basis of Saha's hypothesis (A., 1920, ii, 659; 1921,

ii, 162), the authors show that the constant, K , of the reaction $M = M^+ + e^-$ can be calculated by means of the formula $\log k = -5048V/T + 2.5 \log T - 6.56$, where V is the ionisation potential and T the absolute temperature. From some earlier experiments of Wilson on the conductivity of salts in flames, it is possible to calculate approximately the value of K (A., 1916, ii, 72). These values have been calculated for the case of salts of the alkali metals and the value of $K \times 10^{12}$ obtained both by means of the above formula and Wilson's experiments. The values in brackets are obtained by the formula, caesium 3500 (8200), rubidium 780 (1600), potassium 430 (620), sodium 5.6 (6.2), and lithium 0.8 (1.4). The two sets of values are comparable and of the same order, and this is taken as a confirmation of Saha's hypothesis. From these ionisation constants, it is possible to calculate the degree of ionisation of an element into an electron and an ion. J. F. S.

Resonance and Ionisation Potentials of Helium. J. FRANCK (*Z. Physik.*, 1922, 11, 155—160).—A theoretical paper. The lines in the parhelium spectrum discovered by Lyman (this vol., i, 674) throw light on the relationship between the resonance and ionisation potentials, and the structure of the helium atom. The potentials calculated from the lines in this spectrum are in agreement with the experimental results of Franck and Knipping (A., 1920, ii, 72) and Horton and Davies (A., 1921, ii, 672), if 0.8 volt be subtracted from these values. It is not possible that this divergence is due to error of observation. Reasons are advanced to show that the first break in the curve corresponds with the change $0.5S-1.5s$, and the second with two successive changes, $0.5S-1.5S+0.5S-1.5s$. The difference is $0.5S-1.5S$, and not $0.5S-1.5s$ as has been assumed. This accounts for the difference of 0.8 volt.

The resonance and ionisation potentials of other gases, which are based on those of helium, must be reduced by this amount, and the following ionisation potentials are obtained after making this correction: H_2 , 16.4 ± 0.25 , 29.7 ± 0.4 ; N_2 , 17.05 ± 0.3 , 24.7 , 30.0 ; HI , 12.7 ; HBr , 13.3 ; HCl , 13.7 ; HCN , 14.8 . W. E. G.

Spectroscopic Investigation of the Ionisation of Argon by Electron Collisions. F. HORTON and ANN CATHERINE DAVIES (*Proc. Roy. Soc.*, 1922, [A], 102, 131—150).—The authors' previous determinations of the resonance radiation, 11.5 volts, and ionisation, 15.1 volts, of argon (cf. A., 1920, ii, 215) have been confirmed by both spectroscopic investigation and current-voltage curves. The apparatus used resembles that developed in their experiments with neon (cf. A., 1921, ii, 422). The brighter red lines of the argon spectrum appeared simultaneously, at a point a few volts higher than the ionisation voltage. The first visibility of the lines is accompanied by a sharp increase in the current which is due to the neutralisation of the space charge of the emitted electrons near the filament. On reducing the voltage the lines persist to an extent which depends on the electron current. At moderate gas pressures, the difference between the voltages of

appearance and disappearance of the lines sometimes amounted to 11 volts. On decreasing the voltage from about 25 volts, the lines in the "red spectrum," λ 4510.9, λ 4345.3, λ 4272.3, undergo a marked enhancement in the region of 19 to 16 volts. The intensity of the bright lines λ 7067.5 and λ 6965.8 decreases continuously in brightness as the voltage is lowered. The "blue spectrum" requires for its stimulation higher potential differences across the tube. At 19 volts, the lines λ 4430.4 and λ 4426.2 become visible, and with increasing voltage the intensity increases up to 24 volts and remains stationary until 34 volts, when the intensity again rapidly increases. The break at 19 volts corresponds with the removal of a second electron from an already ionised atom, and that at 34 volts with the double ionising voltage of the argon atom. The difference between the two values gives the normal ionising voltage. From a consideration of the minimum voltages at which the "blue spectrum" lines could be detected, it is shown that the Stark classification of the helium lines into "bivalent" and "tervalent" is inaccurate. The ionisation voltage is related to one of the *M* X-ray absorption limits of the elements in a manner analogous to that in which the helium ionisation voltage is related to the *K* absorption limits, and the neon ionisation voltage to the *L* absorption limits. W. E. G.

Luminous Discharge in Hydrogen and Mercury and a New Method of Measuring Ionisation Potentials. GEORGE E. GIBSON and W. ALBERT NOYES, jun. (*J. Amer. Chem. Soc.*, 1922, 44, 2091—2106).—The authors have examined the conditions of stability of the luminous discharge in hydrogen and mercury, and the variation of the potential, E_d , at which the luminous discharge vanishes, with changes of pressure. To explain the phenomena observed, they put forward the hypothesis that the potential gradient in the portion of the tube between the cathode and the place where ionisation first occurs diminishes as the pressure is decreased until at the minimum of E_d the potential drop in this region has fallen to a very low value, not more than a few tenths of a volt in the shorter tubes or more than about a volt in the longest tubes measured. At the higher pressures, ionisation occurs at regions intermediate in position between the anode and cathode, and as the pressure is diminished these regions recede from the cathode until at the minimum of E_d ionisation occurs only in the immediate neighbourhood of the anode. It is shown that the voltage at which the luminous discharge in hydrogen disappears in tubes furnished with heated tungsten cathodes is a function of the pressure for a given tube passing through a minimum value which is usually a simple multiple of the ionisation potential, 30.9 ± 0.4 volts. In mercury vapour the disappearance voltage of the luminous discharge is a simple integral multiple of the ionisation potential, 10.7 ± 0.4 volts. J. F. S.

Electrical Conductivity of Hydrochloric Acid and Potassium Chloride in Presence of Sucrose. ARTHUR JOSEPH KIERAN (*Trans. Faraday Soc.*, 1922, 18, 119—125).—Experimental deter-


minations have been made of the electrical conductivities at 25° of solutions of hydrochloric acid of concentrations ranging from 0.05*N* to 0.0005*N* in the presence of 5%, 10%, and 20% of sucrose. Similar determinations were made with potassium chloride solutions ranging from 0.01*N* to 0.0005*N* in the presence of 10% of sucrose. In the latter case, the equivalent conductivity increased normally with increasing dilution, to an asymptotic limit. In the case of hydrochloric acid solutions, however, the behaviour was abnormal, the equivalent conductivity passing through a maximum in the region $N/300$ — $N/500$, dependent on the concentration of the sucrose, and thereafter steadily decreasing with increasing dilution. This abnormality was traced to the presence of minute traces of an electrolytic impurity (probably an organic salt of calcium) in all samples of sucrose. Values of m in the relation $\Lambda = \Lambda' \eta^m$, in which Λ and Λ' are respectively the equivalent conductivities in media of unit viscosity and of viscosity η , have been determined as follows: hydrochloric acid and sucrose, 0.57; hydrochloric acid and dextrose, 0.68; potassium chloride and sucrose, dextrose, and glycerol, respectively, 0.73, 0.77, and 0.92. The results confirm the conclusion of Kraus (A., 1914, ii, 90) that m approaches the value unity as the molecules of the medium become smaller in size and the larger the size of the ions of the electrolyte.

J. S. G. T.

Conductivity of Sodium Iodide in Amyl Alcohol at very Low Concentrations. CHARLES A. KRAUS and J. EGBERT BISHOP (*J. Amer. Chem. Soc.*, 1922, 44, 2206—2212).—Exceedingly pure amyl alcohol may be prepared, as follows: ordinary amyl alcohol was submitted to an initial fractionation and the portion boiling above 130° dehydrated over quicklime, after which it was redistilled. The product was treated with a small amount of sodium and the resulting solution repeated fractionated. In this way, amyl alcohol with a specific conductivity of 2×10^{-8} was obtained and by effecting the fractionation under reduced pressure the specific conductivity became 1×10^{-8} . Solutions made up in this product were not stable, but if the alcohol was subjected to fractional condensation under reduced pressure and the distillate collected in the conductivity cell a product with specific conductivity 1.4×10^{-9} was obtained and solutions made up in this were very stable; solutions of sodium iodide in this specially prepared amyl alcohol did not vary more than 0.1% in fourteen hours. The electrical conductivity of solutions of sodium iodide in amyl alcohol was measured at 18° for concentrations ranging between $1.0 \times 10^{-5}N$ to $17 \times 10^{-5}N$. The mean variation of the results does not exceed 0.1%. In the more dilute solutions, the points conform to the requirements of the law of mass action within the limits of the experimental error. At the higher concentrations there is a small although constant deviation from the mass action relation. From an extrapolation of the curve to zero concentration the value 11.059 is obtained for Λ_0 and the value 2.933×10^{-4} for K .

J. F. S.

Variation of the Electrical Conductivity among the Ammoniacal Derivatives of Platinous Nitrite. L. A. TSCHUGAEV and N. A. VLADIMIROV (*J. Russ. Phys. Chem. Soc.*, 1920, **52**, 135—138).—The values of Λ_{1000} for tetra-amminoplatinous nitrite, $[\text{Pt}4\text{NH}_3](\text{NO}_2)_2$; nitrotriamminoplatinous nitrite,



cis- and *trans*-dinitrodiamminoplatinum, $[(\text{NO}_2)_2\text{Pt}(\text{NH}_3)_2]$; and potassium platinonitrite are found to be, respectively, 276.5, 96.5, 0.95, 2.42, and 302.4. The observed value of Λ_{2000} for nitrotriamminoplatinous platinonitrite, $[\text{NO}_2\text{Pt}3\text{NH}_3]_2\text{Pt}(\text{NO}_2)_4$, is 217, the value calculated by summation of the mobilities of the corresponding ions being 197.2 (cf. Tschugaev and Kiltinovitsch, T., 1916, **109**, 1286). T. H. P.

Passivity and Over-potential. ULICK R. EVANS (*Trans. Faraday Soc.*, 1922, **18**, 1—13).—It is suggested that the relative values of the interfacial tensions between the metal, corrosion product, and solution is the factor on which the protection against corrosion of the metal by the solution depends. The activation of passive metals by chlorides is related to the peptising action of metallic hydroxides by chlorides; the rendering passive by chromates is correlated with the flocculating action of these salts. The protective layer probably consists of a layer of oxygen atoms connecting the metal on the one side with solution on the other. Similarly, at a cathodically polarised electrode a layer of hydrogen nuclei connect the metal with the liquid, the hydrogen being probably in a state intermediate between the elementary and ionic forms. The theory is, in some respects, opposed to that of Friend (T., 1922, **119**, 932). J. S. G. T.

Chemical Passivity of Metals. J. GILLIS (*Natuurwetenschapp. Tijdschr.*, 1921, **3**, 66—68).—The phenomena of passivity may not be explained by the formation of an oxide membrane protecting the metal. The most accurate optical methods—which permit estimation of a layer of $0.8\ \mu\mu$ —fail to detect any sign of an oxide membrane. On the other hand, the metals which become passive by anodic polarisation are also those the ions of which do not readily discharge and precipitate on the cathode. There is, therefore, a co-existence of the anodic retardation (passivity) and of the cathodic retardation (slow metallic precipitation); the transformation $\text{M}^+ \rightleftharpoons \text{M} + \oplus$ is slow in both directions.

CHEMICAL ABSTRACTS.

Thermo-electric Force, the Entropy of Electrons and the Specific Heat of Metals at High Temperatures. WENDELL M. LATIMER (*J. Amer. Chem. Soc.*, 1922, **44**, 2136—2147).—A theoretical paper in which the thermo-electric force of a circuit of two metals has been expressed in terms of the entropy of the electrons in the two metals. This expression has the form

$$dE/dT = \int_0^T \sigma_2 \cdot dT/T - \int_0^T \sigma_1 \cdot dT/T$$

where σ_1 and σ_2 are the specific heats of the electrons of the two

metals. It has been postulated that the specific heat of a gram-molecule of electrons in a metal is given by the increase in the specific heat at constant volume of a gram-molecule of the metal above the equipartition value of $6/2R$. On this assumption, the absolute value of the thermo-electric force of the potassium sodium couple has been calculated at 25° from values of the specific heats of these metals. The changes in thermo-electric force of platinum, iron, cobalt, nickel, gold, copper, tungsten, molybdenum, calcium, aluminium, magnesium, and lead against silver have been calculated from specific heat data for temperature differences of 100° to 700° . Satisfactory agreement is obtained between the calculated and observed values. To make these calculations, the existing data on the thermo-electric force have been collected; new data for the thermo-electric force of lead, zinc, magnesium, aluminium, molybdenum, and tungsten against silver have been obtained, and these are summarised in a thermo-electric diagram. The entropy of the metal electrons at low temperatures has been discussed and the possibility of using thermo-electric data as a means of determining C_v for metals at higher temperatures pointed out. The C_v curves for cobalt, iron, calcium, nickel, palladium, aluminium, platinum, lead, magnesium, iridium, gold, copper, silver, molybdenum, and tungsten have been drawn in agreement with existing thermo-electric data. J. F. S.

Thermochemistry and Electromotive Force in Electric Elements. JARL A. WASASTJERNA (*Monograph, Helsingfors, 1918*).—A systematic, mathematical study of representative types of electrolytic cells, the *E.M.F.* being derived from the reaction energy and the affinity. The following results are recorded for the various cells, for the reaction energy, U , the affinity, A , and the electromotive force, E , respectively, the subscript figures representing absolute temperatures: (1) $\text{Cu}|\text{saturated solution CuI}|\text{I crystals}$; U_0 16296; A_{293} 15908; E_{293} 0.69. (2) $\text{Cd}|\text{saturated solution CdI}_2\text{—Hg}_2\text{I}_2\text{ paste}|\text{Hg}$; U_0 20856; A_{234} 20485; E 0.45. (3) $\text{Cu amalgam 10\%}|\text{saturated solution CuSO}_4\cdot 5\text{H}_2\text{O, —Hg}_2\text{SO}_4\text{ paste}|\text{Hg}$; U_{293} 25745; A_{293} 16560; E_{293} 0.353. (4) $\text{H}_2|0.1N\text{—HCl}|\text{Cl}_2$; E_{298} 1.38. (5) $\text{Pb}|\text{saturated solution PbCl}_2|\text{Cl}_2$; E_{291} 1.58. (6) $\text{H}_2|\text{equivalent weights of FeCl}_3\text{ and FeCl}_2\text{ in }0.1N\text{—HCl}|\text{Pt}$; A 3480 and 5220; E_{293} 0.42 and 0.46. (7) Daniell element with saturated solutions of $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$ and $\text{ZnSO}_4\cdot 7\text{H}_2\text{O}$; U_{291} 52045; A_{291} 49488 (subtract 220 cal. for osmotic work); E_{291} 1.69. (8) Daniell element with water as $100\text{H}_2\text{O}$; U_0 50350; A_{290} 50600; E 1.098. Cells delivering electrical energy, but without thermal changes, are considered, but no data are given. Values obtained for the Clark element are: U_0 67310; A_{288} 66218 (subtract 295 cal. for osmotic work); E_{288} 1.43.

CHEMICAL ABSTRACTS.

Complex Ion Formation in Hydrochloric Acid. RALPH M. HIXON (*Medd. K. Vetenskapsakad. Nobel-Inst., 1922, 4, No. 12, 1—8*).—If complex acids of the type H_2CaCl_4 exist in solutions of metallic chlorides in hydrochloric acid solution, a study of the

migration of the metallic ion when such solutions are electrolysed should reveal their presence. Solutions of magnesium, calcium, barium, and zinc chlorides at different concentrations in concentrated hydrochloric acid were electrolysed for six to eight hours at 12 volts, and samples of anolyte and catholyte were analysed. The solutions of magnesium and calcium chlorides both showed migration of the metal towards the anode at medium concentrations, but towards the cathode at higher and lower concentrations. The migration towards the anode is strong evidence for the existence of complex ions of the type $(\text{CaCl}_2)_x\text{Cl}_x^-$. The observed migration will depend on the relative concentrations of these complex ions and of the simple ions Ca^{++} , as well as on their migration velocities. At low concentrations, Ca^{++} will predominate. At concentrations of $[\text{CaCl}_2]$ approaching saturation, Ca^{++} may again predominate owing either to the decrease in the $[\text{HCl}]$ concentration or to the formation of a more complex acid $(\text{HCl})_x(\text{CaCl}_2)_y$ which would be less dissociated than the simpler acid formed at medium concentrations.

E. H. R.

The Hydration of the Lithium-ion. J. BABOROVSKY and J. VELISEK (*Chem. Listy*, 1922, 16, 250—256).—A new apparatus suitable for determining the transport numbers and hydration of ions, consists of two membranes of parchment paper separating the central portion of the solution from the anodic and cathodic portions and also of an arrangement whereby the anodic and cathodic vessels can be disconnected and weighed with their contents. The hydration of the lithium-ion was calculated from the change in weights and in concentrations of the solutions around the electrodes. The determination of the hydration of ions by this method is complicated by electro-osmosis. This the authors claim to have measured by replacing the two parchment paper membranes by a single membrane and estimating the differences in the concentration and specific gravity of the central portion after each experiment. The lithium-ion has one more molecule of water of hydration than the chlorine-ion. The transport number of lithium in a decinormal solution of its chloride is 0.3128 ± 0.0048 . The authors propose that the hydration of ions should be expressed in terms relative to that of the hydrogen-ion (the most mobile of all) which would be put equal to zero.

W. T.

Electrolysis with Drops of Mercury as the Electrode. J. HEYROVSKY (*Chem. Listy*, 1922, 16, 256—264).—To study the electrolysis of aqueous solutions of the alkali and alkaline-earth metals, the author employed drops of mercury as the electrode; the apparatus is figured. This enabled him to determine the exact course of the electrolysis. Electrolysis is produced when a certain *E.M.F.* is applied the value of which is characteristic of each salt and depends on the logarithm of the concentration. The rapid increase of the current is explained by a diffusion towards the interior of the drop of mercury of very dilute amalgams. This diffusion commences because the metal has not sufficient mercury for combination. The author calculated the potential at which

the kations deposit from their normal solutions on a mercury cathode: lithium, -2.023 ; sodium, -1.860 ; potassium, -1.883 ; rubidium, -1.796 ; caesium, -1.837 . The affinity of the metals for mercury is given by the difference between the characteristic potentials of polarised drops and the normal electrolytic potentials of the pure alkali metals. They were found to be lithium, 1.281 ; sodium, 1.138 ; potassium 1.325 ; rubidium, 1.409 ; caesium, (1.50) .
W. T.

Simultaneous Electrodeposition of Lead and Lead Dioxide.

M. G. MELLON and H. F. REINHARD (*Proc. Indiana Acad. Sci.*, 1921, 181—188).—The amounts of lead and lead dioxide deposited simultaneously on the cathode and anode respectively when a solution of lead nitrate is electrolysed under varying conditions are variable when free nitric acid is present, but for neutral solutions a fair constancy exists in the ratios for the amounts of each of the two deposits to that of the quantity of silver deposited in a coulometer in the circuit. The ratios of the anode and cathode deposits respectively to the lead equivalent of the silver were 1.0053 (alternatively, 0.990) and 0.9864 . The cause of the inaccuracy has not been traced.

CHEMICAL ABSTRACTS.

Electrolytic Migration of Sodium through Glass. M. PIRANI and E. LAX (*Z. tech. Physik*, 1922, 3, 232—235).—The film of water which persists inside an exhausted glass tube was removed by the migration of sodium ions through the glass. A tube shaped like a lamp bulb and containing a small tungsten wire was exhausted and immersed in molten sodium nitrate at 450° . The wire was heated to a high temperature and a potential of 200 volts was applied between the wire and the molten bath. Sodium ions migrated through the glass and a yellow luminescence was produced in the tube. The water film was displaced by this process and was attacked by the hot tungsten as evidenced by the formation of a film of blue tungsten trioxide on the tube.

CHEMICAL ABSTRACTS.

Method of Maintaining Small Objects at any Temperature between -180° and $+20^{\circ}$. P. P. CIOFFI and L. S. TAYLOR (*J. Opt. Soc. Amer.*, 1922, 6, 906—909).—The object is placed within a double-walled vacuum tube through which a stream of cold air, produced by the evaporation of liquid air contained in a Dewar vessel, is maintained. The rate of evaporation is controlled by means of an electric heater immersed in the liquid air, and determines the temperature to which the object is subjected. If desired, the object may be placed beyond the splayed-out end of the vacuum tube and an unbroken stream of cold air caused to flow across it by suction across the gap between the end of this tube and the wide mouth of a second tube connected to a vacuum pump. The latter tube should have a diameter about double that of the former and its length should be about twice that of the gap. The Dewar flask is preferably unsilvered, and is charged with liquid air through a funnel extending almost to the bottom and

provided with a plug of glass wool for filtering out ice and solid carbon dioxide. Temperature regulation may be maintained constant to within about 4° for many hours. J. S. G. T.

Heat of Evaporation. W. HERZ (*Z. anorg. Chem.*, 1922, **124**, 56—58).—Employing the values obtained by Young, the author finds that the quotients of heats of evaporation at equal fractions of the critical temperatures (on absolute scale) are fairly constant for the twenty-six organic compounds investigated. The quotient of latent heat of evaporation and absolute temperature is given for pentane at different temperatures, and remains fairly constant, 0.11—0.15, except in the neighbourhood of the critical temperature. The same was found to hold in the case of twenty-eight other organic compounds. Acetic acid is exceptional, this being probably due to association. W. T.

The Form of the Vapour Pressure Curve at High Temperature. I. The Curve for Lead. CHRISTOPHER KELK INGOLD (*T.*, 1922, **121**, 2419—2432).

High Temperature Investigations. XV. The Vapour Pressure of the Alkali Fluorides. OTTO RUFF, GERHARD SCHMIDT, and SUSANNE MUGDAN (*Z. anorg. Chem.*, 1922, **123**, 83—88).—The apparatus used was like that described by Ruff and Mugdan (*A.*, 1921, ii, 485) except that iridium crucibles were employed. The results are given for the fluorides of sodium, lithium, potassium, rubidium, and caesium. The latent heat of evaporation calculated by means of Liempt's equation (*A.*, 1920, ii, 588) was found to be, LiF, 52.2 Cal.; NaF, 51.1 Cal.; KF, 44.7 Cal.; RbF, 40.4 Cal.; CsF, 33.6 Cal. The critical temperatures calculated by the relationship $T_c = 1.55T_{760}$ are LiF, 3010° ; NaF, 3070° ; KF, 2750° ; RbF, 2600° ; CsF, 2370° .

W. T.

Heats of Combustion and Energy of Dissociation. WALTER HÜCKEL (*Ber.*, 1922, **55**, [B], 2839—2843).—A detailed criticism of the publications of von Weinberg on this subject (*A.*, 1920, ii, 668, 669).

In the final tables given by von Weinberg, only the previously known data for HCl, HBr, and HI are independent of hypothesis. The values for the C—C, C—H, C=C, C \equiv C, C—Cl, C—Br, and C—I linkings depend on the equality of Q_{CO} and Q_{CH} which has not been established. All the other data depend also on other unproved assumptions. The value of Weinberg's work lies in the fact that he has established for the first time a series of interesting relationships between the energy content of different organic compounds as illustrated by the scheme: (1) $Q_{(CO)} = Q_{C-O} + Q_{C-C}$, $Q_{(CO)} = Q_{C-O-C} + Q_{C-C}$, $Q_{C-O} = Q_{C-O-C}$ (2) $6Q_{NH} - Q_{N\equiv N} = 6Q_{CH} - 3Q_{CO}$ (3) $Q_{CO} - Q_{CH} = Q_{C-Cl} - Q_{H-Cl}$. H. W.

The Energy of the Atomic Linkings in Diamond and in Aliphatic Hydrocarbons. K. FAJANS (*Ber.*, 1922, **55**, [B], 2826—2838; cf. *A.*, 1920, ii, 354).—A reply to various criticisms

in many of which allowance has not been made for the admitted uncertainty of the author's data (cf. Hückel, A., 1920, i, 603; Polanyi, A., 1921, ii, 179; Swientoslawski, A., 1921, ii, 535; von Weinberg, A., 1920, ii, 668, 669; Thiel, A., 1920, ii, 667).

1. *Heat of sublimation of carbon and the absolute values of the linking energies.*—In the previous paper, the heat of sublimation of carbon was assumed for purposes of numerical illustration to be 287 Cal. Subsequently it has been shown by Kohn (A., 1921, ii, 302) that the heat of sublimation of carbon is 163.5 Cal. as a general mean of all measurements and that of diamond 168 Cal. The values for the most important linking energies have therefore been recalculated on a basis of 150 Cal. as the most probable value, on the supposition that carbon vapour is monatomic at 4200°, and that an equilibrium is established in Lummer's experiments.

2. *The energy of the double and triple linking.*—It is shown by a somewhat modified method that the heat of formation of a double or treble bond is greater than that of a single bond. Arguments are advanced in disproof of the criticism by Thiel (*loc. cit.*) that this conception is contrary to the known instability of substances with unsaturated linkings.

3. *Chemical forces in crystals.*—It has been asserted by Thiel (*loc. cit.*) that the chemical forces which are operative in the formation of a crystal are essentially those of subsidiary valency. This does not appear to be universally true, since the forces which hold the sodium chloride lattice together are identical with those which are operative in the vaporised salt molecule. The lattice is therefore held together by main valencies and the heat of sublimation is of the same order as that of the dissociation of the molecules into the free ions. On the other hand, the lattices of many organic compounds, for example solid hexane, are without doubt held together entirely by subsidiary valencies; this is shown by the fact that the molecular heat of sublimation of solid hexane is only 8 Cal., whereas the energy necessary for the disruption of the vaporised molecule into atoms is calculated to be 1511 Cal. The conception that the valency forces which unite the carbon atoms in the diamond lattice are identical with those uniting the carbon atoms of aliphatic compounds is not new; the advance made by Fajans consists in the exact quantitative demonstration of the correctness of the hypothesis by thermochemical methods.

4. The remainder of the communication is devoted to a question of priority (cf. von Weinberg, *loc. cit.*). H. W.

The Thermochemistry of Carbon Compounds. A. THIEL (*Ber.*, 1922, 55, [B], 2844—2845).—A reply to Fajans (preceding abstract). H. W.

Thermochemistry of Solutions. M. LEVANT-EZERSKY (*J. Russ. Phys. Chem. Soc.*, 1920, 52, 108—113).—Simple reasoning from fundamental principles leads to a number of equations which may be combined in the form: $q'_m = q_m - Bq_i = BQ_i - Q_m = \phi_i - \phi_m = BR_i - R_m$, where q'_m represents the heat of dilution of a solution containing $100m/(100-m)$ grams of solute per 100 grams

of water to the concentration $l\%$, and q_m^* and q_l^* have analogous significations; Q_l and Q_m are the heats of solution of $100l/(100-l)$ and $100m/(100-m)$ grams, respectively, of the solute in 100 grams of water; $\phi_m = Q_m$ and $\phi_l = m(100-l)Q_l/l(100-m)$, that is, $\phi_l - \phi_m$, corresponds with the difference between the heats of solution of one and the same quantity of the solute in such different amounts of water that solutions of $l\%$ and $m\%$ concentration, respectively, are obtained; R_l and R_m represent, respectively, the heats of reaction of $100l/(100-l)$ and $100m/(100-m)$ grams of the solute, liquefied at the temperature of the experiment, with 100 grams of water.

The above equation furnishes a convenient means of passing from the heats of solution of different amounts of a substance in one and the same amount of solvent to the heats of solution of one and the same quantity of solute in different quantities of solvent and from these to the heats of dilution.

T. H. P.

Effect of Variation in Weight of the Riders and Plummets of the Westphal Balance on the Accuracy of Specific Gravity Determinations. EDWARD A. TSCHUDY (*J. Amer. Chem. Soc.*, 1922, **44**, 2130—2135).—Accurate specific gravity determinations with the Westphal balance are possible only when the weight of water displaced by the plummet at 15° and the weights of the riders possess an exact multiple relationship amongst themselves. Variations in weight of the riders which disturb this relationship produce errors, often of considerable magnitude, in specific gravity determinations. Inaccurate plummet thermometers are also a source of error. The magnitude of ordinary variation in weight of riders and plummets has been ascertained, and equations have been derived for calculating the correction which must be applied to any indicated balance reading to obtain the true specific gravity of liquids lighter or heavier than water. Application of the corrections for any specific system of riders to indicated balance readings is facilitated by the construction and use of correction graphs.

J. F. S.

The Degree of Molecular Polymerisation of Substances under Critical Conditions. J. A. MULLER (*Compt. rend.*, 1922, **175**, 760—761).—A table of the calculated values of the degree of polymerisation of a number of elements and compounds under critical conditions is given. All the substances included, with the exception of helium, undergo polymerisation to a certain extent; those which, under normal conditions of pressure and temperature, approximate in their behaviour to perfect gases are least affected, the values of the mean degree of polymerisation ranging from 1.227 in the case of hydrogen to 1.285 in that of nitrogen. Liquid hydrocarbons and monohalogen derivatives of benzene give values from 1.39 to 1.45, whilst those for esters of saturated monobasic fatty acids vary from 1.45 to 1.48. Nitriles, water, and acetic acid yield results in the neighbourhood of 2.

H. J. E.

Corresponding States. W. HERZF (Z. anorg. Chem., 1922, 123, 132—136).—A theoretical paper in which the author investigates whether there is a constant relationship between the viscosity of liquids at the same fraction of the critical temperature. The viscosity at one-half the absolute critical temperature is divided by the viscosity at seven-twelfths the absolute critical temperature. In the case of 29 non-associating liquids, the ratio is 1.51—1.91. In the case of liquids which associate, the viscosity decreases more rapidly with increasing temperature and the ratio in the case of six liquids varies from 1.95 to 4.33. The author in the same way finds the ratio of the refractive indices at corresponding temperatures. The refractive index is calculated by equation $n = \sqrt{M + 2dR_L / M - dR_L}$. In the case of 29 liquids the approximate constancy of the ratio showed the validity of the law of corresponding states. W. T.

The Viscosity of Amphoteric Electrolytes in Solution. GUNNAR HEDESTRAND (Z. anorg. Chem., 1922, 124, 153—184).—Solutions of amino-acids are found to have a minimum viscosity at the isoelectric point. In the case of glycine and alanine, the viscosities of the anions is greater than that of the kations. Mixtures of glycine and alanine solutions have a viscosity which is the arithmetical mean of the corresponding unmixed solutions. The addition of neutral salts increases the viscosity of glycine and alanine solutions; it also displaces the minimum. The minimum viscosity of a gelatin solution is at the isoelectric point; the addition of salts increases the minimum value and at the same time shifts it to a lower hydrogen-ion concentration. On both sides of the minimum, the viscosity is decreased considerably by the salts. The author treated powdered gelatin with hydrochloric acid, and found that, on washing, a large amount of the gelatin went into solution; the same result was obtained by treating the powder with a mixture of hydrochloric acid and a salt. W. T.

Effect of Surface Tension. P. B. GANGULY and B. C. BANERJI (Z. anorg. Chem., 1922, 124, 140—142).—A stick of sodium hydroxide was held vertically in distilled water so that only one-half was immersed. In the absence of stirring, the part of the stick in contact with the surface dissolved the more rapidly. The same was observed in the case of sticks of potassium hydroxide, sodium nitrite, copper sulphate, and zinc chloride, and also of a zinc rod in dilute sulphuric acid. In the case of an iron rod in hydrochloric acid the reverse was observed and the part further from the surface dissolved the more rapidly. These effects are reduced by lowering the surface tension by a covering of oil, and are therefore assumed to be due to surface tension effects. W. T.

Negative Adsorption. II. Calculation of the Amount of Adsorbed Solvent. M. A. RAKUSIN and TATJANA GÖNKE (Biochem. Z., 1922, 132, 82—83).—The amount of adsorbed solvent

can be calculated by Gurvitch's equation for positive adsorption (A., 1914, ii, 435).
H. K.

The Sorption of Saturated Vapours by Charcoal. JOHN DRIVER and JAMES BRIERLEY FIRTH (T., 1922, 121, 2409—2414).

Salt Adsorption on Metal Surfaces. H. VON EULER and G. ZIMMERLUND (*Arkiv Kem. Min. Geol.*, 1922, 8, No. 14, 1—23).—Measurements have been made of the silver nitrate and potassium chloride adsorbed from aqueous solution by finely divided gold particles and by gold leaf. It has been found that the amount adsorbed increases with the amount of salt in solution, at first linearly and then tending to a maximum. At the maximum there is more adsorbed than would correspond with a unimolecular layer on the gold. Experiments are also described with mercury dropping through a calomel solution in which the amount of mercury in solution was determined by finding its effect as an enzyme poison.
W. O. K.

Mordanting of Wool with Potash Alum. W. W. PADDON (*J. Physical Chem.*, 1922, 26, 790—793).—Experiments are described from which it is concluded that, contrary to the opinion of Knecht, the mordanting of wool with potash alum does not lead to the formation of definite chemical compounds on the fibre, but is strictly an absorption phenomenon in which both alumina and sulphuric acid are involved.
J. S. G. T.

Mordants. III. Chrome. WILDER D. BANCROFT (*J. Physical Chem.*, 1922, 26, 736—772).—The author reviews critically a part of the literature, more especially the work of Liechti and Hummel (*J. Soc. Chem. Ind.*, 1893, 12, 244) concerned with the use of chromic oxide as a mordant in dyeing wool, cotton, and silk. The following are among the conclusions drawn: from dichromate solutions wool first adsorbs chromic acid and this is reduced to chromic oxide, which is the true mordant; within limits, increasing the acid concentration increases the chromic acid taken up; chromic acid oxidises organic compounds more readily in presence than in absence of wool; when wool is mordanted with chrome alum, a basic sulphate changing later to chromic oxide is first formed; silk adsorbs chromic oxide less strongly than wool does; cotton takes up scarcely any chromic oxide from chrome alum, but adsorbs it from an alkali solution; there is no evidence of the formation of any definite compound when wool is mordanted with chromic oxide.
J. S. G. T.

Tungsten Furnace for Experiments on Dissociation and Ionisation. K. T. COMPTON (*J. Opt. Soc. Amer.*, 1922, 6, 910—912).—Tungsten sheet is bent into the form of a cylindrical tube and clamped by end pieces each consisting of a steel ring fitting in a split rectangular steel block. The blocks are mounted on water-cooled brass tubes serving as leads for the heating current. Short glass tubes surround the leads for purposes of insulation. Loops of fine tungsten wire, drawn tightly round the cylindrical

tube, prevent bulging of the furnace. The central electrode consists of a straight length of "20 mil." tungsten wire welded to heavier molybdenum leads. A furnace 60 mm. overall length and about 5 mm. diameter was raised to a white heat by 100 amperes at 6 volts, and reached its melting point when a current of about 200 amperes was employed.

J. S. G. T.

The Thermal Decomposition of Chlorine. FR. A. HENGLEIN (*Z. anorg. Chem.*, 1922, **123**, 137—165).—A description of the apparatus suitable for a reactive gas at low pressure and high temperature is given in detail (cf. this vol., ii, 441). The apparatus was tested by measuring the dissociation of bromine. The dissociation of chlorine was investigated at a pressure of 1×10^{-5} atmospheres and between the temperatures of 700° and 900°. The following relationship was found to hold $\log K_{(\text{atmos.})} = -(5400/4.571T) + 1.75 \log T - 4.09 \times 10^{-4}T + 4.726 \times 10^{-8}T^2 + 0.93$ (cf. Bodenstein, A., 1916, ii, 554). The chemical constant of monatomic chlorine was calculated in three different ways: (1) by Stern's method (A., 1914, ii, 543), (2) by the reaction $\text{Ag} + \frac{1}{2}\text{Cl}_2 = \text{AgCl}$, the heat effect and electromotive force of which are accurately known, (3) by the reaction $\text{Hg} + \frac{1}{2}\text{Cl}_2 = \text{HgCl}$. The values found were respectively 0.68, 0.93, and 1.50. A linear relationship was found to exist between the heats of dissociation of chlorine, bromine, and iodine and their atomic radii and other spatial and thermal constants. The electroaffinity of chlorine was calculated to be 73 cal.

W. T.

Theory of Solvates. JA. K. SYRKIN (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, **6**, 267—280).—Solvates in general and hydrates in particular vary in composition and are stable only within narrow limiting conditions, so that they must be regarded as transitory systems of physico-chemical character. The structure of a hydrate may be represented as a molecule of solute surrounded by an envelope of water molecules. Such of the latter as are near to the nucleus are held by a force approximating to chemical force, whereas the more remote, semi-combined water molecules exist under the configurational influences of chemical and kinetic forces, the system being in a condition of stationary thermal equilibrium. Hence it is possible for hydrates of different composition to exist in one and the same solution. This equilibrium of hydrates possesses a macroscopic character. The gradual variation of the thermal effect with the temperature confirms these views and distinguishes the decomposition of a hydrate from the stoichiometric decomposition of an ordinary system. The existence of hydration is shown also by the change in the internal pressure, and the work of the internal forces may be expressed as the mechanical equivalent of chemical affinity (cf. Tammann, "Über die Beziehungen zwischen den inneren Kräften und Eigenschaften der Lösungen," 1907, 34).

T. H. P.

Solubility of Carbon Dioxide and Nitrous Oxide in certain Solvents. WILLIAM KUNERTH (*Physical Rev.*, 1922, **19**, 512—524).—In view of its bearing on the Lewis-Langmuir theory, a

comparison was made between the solubilities of carbon dioxide and of nitrous oxide between 18° and 36° in water, acetone, acetic acid, methyl alcohol, pyridine, ethyl alcohol, benzaldehyde, aniline, amyl acetate, ethylene bromide, isoamyl alcohol, and chloroform. In this series, the ratio of the solubility of carbon dioxide to that of nitrous oxide decreases regularly from 1.34 to 0.66. The ratio is nearly constant for each solvent, and the temperature coefficient, ds/dT , which is always negative, is in most cases nearly the same for the two gases. It is found that Raoult's law does not hold for the solubility of gases in liquids, and that there is little, if any, relation between solubility and the difference between internal or cohesion pressures of solvent and solute. Since, however, the ratio of the solubilities of carbon dioxide and nitrous oxide (the former of which is the more active chemically, and therefore has the stronger polarity) varies regularly with the dielectric constant of the solvent, and since this constant may be taken as an index of the polarity of the solvent, it is suggested that polarity may be an important factor in determining the relative solubility of gases in liquids. The following figures, for carbon dioxide and nitrous oxide, respectively, at 20°, are abstracted from a table giving the solubilities expressed in c.c. of the gas under existing barometric pressure per c.c. of solvent: water, 0.900, 0.675; acetone, 6.98, 6.03; acetic acid, 5.23, 4.85; pyridine, 3.85, 3.58; methyl alcohol, 3.57, 3.32; ethyl alcohol, 2.87, 2.99; benzaldehyde, 2.98, 3.15; aniline, 1.38, 1.48; amyl acetate, 4.65, 5.14; ethylene bromide, 2.27, 2.81; isoamyl alcohol, 1.91, 2.47; chloroform, 3.71, 5.60.

A. A. E.

Partition of a Substance between two Solvents and the Solution Field of Force. N. A. SCHILOV and L. K. LEPIN [with M. P. JANTSCHAK] (*Nachr. Phys.-Chem. Lomonossow-Ges.*, 1920, 1, 1—103; from *Chem. Zentr.*, 1922, iii, 541—542).—The partition of a dissolved substance between two liquid phases changes with varying concentrations and only remains constant in particular cases owing to the operation of compensating factors. The partition does not in general depend on the solubilities in the individual solvents; this relationship, in fact, only holds at the triple point of the three phases. The isotherms for the changes in concentration are different for each phase and are not linear. The relation $k=c_1^2/c_2$ is only a particular case of the relation $k=c_1^m/c_2=q \cdot c_1^n/c_2^n$, where n is an integer or fraction and given by $n=\log \beta_1/\log \beta_2$, β_1 and β_2 being factors expressing the changes in the solution field of force (Lösungskraftfeld) depending on concentration. The partition coefficient thus obtained is constant for many systems, excepting those where the two solvents have a considerable reciprocal solubility, and where chemical factors may play a part.

G. W. R.

Solubility of Liquids in Liquids. The Partition of the Lower Acids, particularly Formic, between Water and various Organic Solvents. NEIL E. GORDON and E. EMMET REID (*J. Physical Chem.*, 1922, 26, 773—789).—The partition

ratios at 25° of formic acid between water and the following respective solvents: cotton-seed oil, petroleum, benzene, toluene, xylene, carbon tetrachloride, carbon disulphide, and bromoform have been determined. In like manner, the respective partition ratios at 25° of acetic acid between water and either cotton-seed oil or petroleum, of propionic and butyric acids between water and cotton-seed oil have been determined. The results have been employed to determine the so-called solubilities of the acids in the solvents and vice versa.

J. S. G. T.

The Behaviour of Two Metals towards One Another when Dissolved in Mercury. G. TAMMANN and W. JANDER (*Z. anorg. Chem.*, 1922, **124**, 105—122).—The authors investigated the affinity of one metal for another when dissolved in mercury: (1) by measuring the freezing-point depression of the compound crystals and comparing this with that of the components, (2) by measuring the influence of the one metal on the *E.M.F.* concentration-curve of the other dissolved in a third more noble liquid metal. Thus the addition of small amounts of a more noble to a less noble metal has but a very small effect on the *E.M.F.* if the compound of the two metals is very highly dissociated, whereas, if a stable compound is formed, then a sudden change in the *E.M.F.* is caused when the concentrations of the metals correspond with the composition of this compound. The following intermetallic compounds were found to be very stable: AuZn , Mg_2Sn , CeSn_2 , and Ce_4Bi_3 , having very high affinity constants. AuZn $K=3.4 \times 10^7$. Mg_2Sn $K=1 \times 10^{10}$. CeSn_2 $K=1 \times 10^{40}$. Ce_4Bi_3 $K=3 \times 10^{120}$. Cf. water $K=4.6 \times 10^{78}$. Others were very highly dissociated in solution in mercury; for example, compounds of gold with cadmium and lead, of silver with zinc and cadmium, of copper with zinc and cadmium.

W. T.

Systems in which Metals Crystallise. JOHN L. HAUGHTON and G. WINIFRED FORD (*Trans. Faraday Soc.*, 1922, **18**, 112—118).—Of the twenty-nine elements the crystalline structures of which have been determined by X-ray analysis, five only are exceptions to the rule that all elements of the same periodic series crystallise in the same system. In nearly every case, alloys which form a homogeneous series of solid solutions right across the equilibrium diagram, crystallise in the same system. There appears to be no relationship between the changes which occur in the physical properties of metals at the melting point and the crystalline habit of the metals. A possible exception occurs in the case of the electrical resistivity of the metals in the odd series of group 5.

J. S. G. T.

Solutions, Suspensions, Colloids. N. P. PESKOV (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, **6**, 73—99).—The theoretical considerations developed by the author lead to the following conclusions. The modern doctrine of colloids deals with a conception of heterogeneity which is totally different from that assumed by the doctrine of phases, but is identical with the molecular-kinetic view of the universe; the latter is termed the generalised conception

of heterogeneity. The notion of dispersion current in the chemistry of colloids must be regarded as composed of two different conceptions, distinguished as qualitative and quantitative dispersion. The former of these is synonymous with the generalised conception of heterogeneity referred to above, and has no special bearing on systems represented by suspensions, colloids, and solutions. The latter, possessing a dimensional character, differs from the idea of a phase only in that the latter is characterised by the spacial magnitudes of its separate elements. Thus all systems to which the term quantitative is applied must be considered to be polyphasic.

Physical theories of colloids in general postulate that: (1) suspensions, colloids, and true solutions are systems, the qualitative differences between which may be regarded as functions of their degrees of dispersion, (2) colloids represent typical polyphase systems, and (3) all the characteristic properties of colloids are merely the results of the magnitudes of the molecules. The logical objection to such theories is that they accept as results of the doctrine of dispersion the hypothetical assumptions on which this doctrine is based.

The author considers that only careful investigation, from all points of view, of the process of coagulation and of the adsorption phenomena accompanying them will decide the question of the heterogeneity of colloids in their normal condition. T. H. P.

Electrical Colloid Synthesis. SVEN BODFORSS and PER FRÖLICH (*Koll. Chem. Beihefte*, 1922, **16**, 301—340).—Svedberg's method of synthesis of colloids has been investigated for the case where the induction coil current is replaced by a high tension symmetrical alternating current and the dependence of the amount of dispersion on the dimensions of the oscillation circuit determined. The dispersion of copper, tin, lead, zinc, antimony, aluminium, silver, magnesium, bismuth, and thallium by Svedberg's method and the present method has been compared, and it is shown that for the more easily dispersed metals the present method is inferior to the original Svedberg method, the yield being about one-half to two-thirds that obtained by the older method. In the case of the harder metals, the dispersion is about the same by both methods. Alloys belonging to the systems lead-tin and antimony-zinc have been dispersed by the present method and the sols produced examined. J. F. Š.

Winkelblech's Phenomenon or Pseudo-extraction and its Importance in Colloidal Chemistry. New Methods of Extracting Solids. K. CHARITSCHKOV (*J. Russ. Phys. Chem. Soc.*, 1920, **52**, 96—107).—Winkelblech (A., 1907, ii, 17) found that when certain pseudo-solutions are treated with various non-aqueous solvents insoluble in water, the solid disperse phase passes partly or completely into the liquid, non-aqueous phase, through-out which it becomes distributed in the form of a gel. This phenomenon is usually regarded as a special case of adsorption, but the results obtained on examining different combinations of disperse phases and non-aqueous solvents are completely incompatible

with this view. The effect seems to be due rather to mechanical seizure of the molecules of the solid phase by disintegrated liquid drops of lower specific gravity or to attraction of these molecules to the bottom by molecules of a heavier liquid.

From a pseudo-solution of gelatin part of the sol may be drawn to either the surface or the bottom of the liquid by treatment with light petroleum, chloroform, carbon disulphide, etc., and owing to the resemblance of the phenomenon to extraction the author terms it "pseudo-extraction."

For the velocity of transport of the molecules, Lewis's formula, $V = 2r^2S(S - S_1)/9\eta$, should be valid, S and S_1 being the specific gravities of the solid and liquid phases, r the radius of the molecules, and η the coefficient of viscosity, but greater exactitude would be attained by addition of the term, $2r^2S(S - d)/9\eta$, d denoting the specific gravity of air.

The fact that sols of silicic acid, Prussian blue, and almost all metals remain unchanged when treated with light petroleum, benzene, and similar liquids is explained on the assumption that, for certain values of S and S_1 the magnitude of V will be negative, the specifically lighter molecules of air and light petroleum being then not in a condition to draw upwards the heavier molecules of the sols of metals and various oxides.

The results of experiments on the pseudo-extraction of gelatin, agar-agar, and sodium oleate show that the quantity of the colloidal substance thus extracted is not proportional to the concentration; when the latter is very high, no pseudo-extraction takes place. For every disperse phase, the phenomenon appears to exhibit an optimum concentration and an optimum temperature. In no case, however, is it possible to remove by pseudo-extraction the whole amount of a sol occurring in the disperse phase. The excess above the normal proportion of potassium hydrogen tartrate or sodium sulphate present in supersaturated solution is withdrawn instantaneously from the latter by shaking with light petroleum. The same occurs when an aqueous alcoholic solution of sucrose is similarly treated, but in this instance the sucrose withdrawn partly redissolves in the aqueous alcoholic layer on prolonged keeping of the system after being shaken with the light petroleum. T. H. P.

General Method for Obtaining Gels of Inorganic Salts, and its Relation to Theories of the Colloidal State. K. CHARITSCHKOV (*J. Russ. Phys. Chem. Soc.*, 1920, 52, 91—96).—Treatment of a solution of sodium (potassium, etc.) naphthenate in a hydrocarbon containing excess of naphthenic acid with hydrogen chloride or carbon dioxide yields colloidal sodium chloride or carbonate. With heavy metal salts of naphthenic acid, which dissolve readily in benzene, toluene, or light petroleum, excess of naphthenic acid is not required; colloidal cupric and mercuric chlorides, etc., are obtainable in this way. The naphthenic acid may be replaced by oleic acid. The products formed show Brownian movement and the corresponding ultramicroscopic appearance. The sol first formed rapidly undergoes coagulation

but not throughout its entire mass, part remaining in a condition of pseudo-solution detectable ultramicroscopically.

Treatment of ferrous sulphate in the above way results in the formation of a white precipitate which, as oxidation proceeds, is gradually converted first into a black sol of triferric tetroxide and ultimately into a brown sol of ferric oxide. The action of hydrogen chloride on ferric naphthenate yields, not ferric, but ferrous chloride, the hydrogen chloride acting as a reducing agent under these conditions.

These and allied phenomena do not appear to be explicable by any of the existing theories of the colloidal state, or by von Weimarn's crystallisation theory, which is generally accepted. Neither this author's formula for expressing the course of separation of precipitates, nor that given by Nernst to indicate the velocity with which a precipitate forms round centres of crystallisation, takes into account the attraction or affinity between solvent and solute, which constitutes an important factor; in many cases, the whole process of crystallisation is the resultant of the force uniting the molecules and the affinity of the latter for the solvent.

The author regards gels and sols as the extreme and final stages of the interactions between solid and liquid phases, intermediate stages being represented by crystals of various degrees of symmetry.

T. H. P.

Sensitisation of Coagulation Processes: Colloids as Indicators of Photo-electric Effects. N. P. PESKOV (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, 6, 101—104).—Anthracene is found to exert a sensitising action on the coagulation of sols of arsenic trisulphide, platinum, and carbon, but it is not known if this action is exercised towards all sols or if it is limited to those carrying a definite charge. The action does not depend on chemical transformation of the anthracene or on its fluorescence, but lies in the electronic transpositions taking place in the system under the influence of insolation. Hence colloids constitute a new type of indicator for photo-electric effects. Reference is made to the various possible interpretations of the mechanism of the coagulation of colloids under the sensitising action of anthracene.

T. H. P.

Flocculation by Mixtures of Electrolytes. H. FREUNDLICH and P. SCHOLZ (*Koll. Chem. Beihefte*, 1922, 16, 267—284).—The authors have confirmed the results of Odén, to the effect that the coagulating properties of a salt are diminished or destroyed by the presence of a second salt, but in contradiction to Odén they have also found that the reduction is not due to a diminution of the kation action by the anion but rather to an antagonism between the kations. The antagonistic action is measured by mixing Odén's sulphur sol with a quantity of an electrolyte insufficient to cause flocculation and then determining the quantity of a second electrolyte necessary to effect this change. In the case of ion pairs with strong antagonism such as magnesium-lithium, or magnesium-hydrogen, the coagulation value of the magnesium

is a whole number multiple of the value obtained when the coagulation is effected by magnesium alone. The anions have also an influence in the diminution of the coagulation value in the order citrate">sulphate">chloride'. Definitely hydrophobic sols, such as Weimarn's sulphur sol and Donau's gold sol, do not exhibit the above-named phenomena. In these cases the coagulation values of sols containing electrolytes are always less than those of the pure sols; that is, the action of the two electrolytes is additive, as is to be anticipated from the coagulation hypothesis of hydrophobic sols. In these cases also, the coagulum cannot be peptised by electrolytes. The behaviour of Odén's sulphur sol is probably to be connected with its distinct hydrophilic character; that is, with the hydration of the micellæ. In the case of arsenic sulphide sol, a certain antagonism is observed for salts such as lithium chloride and magnesium chloride.

J. F. S.

Coagulation of Manganese Dioxide Sol. by Different Electrolytes. PHANI BHUSAN GANGULY and N. R. DHAR (*J. Physical Chem.*, 1922, 26, 701—714).—The coagulative powers of different electrolytes on manganese dioxide sol do not follow the Schulze-Hardy law, namely, the higher the valency of an ion the greater its precipitating action on a colloid. Changes of concentration of the sol alter the relative order of the electrolytes when arranged according to their coagulative powers. Coagulative power is not purely a function of valency, but is more a specific property of the ions, and is influenced, among other factors, by the time allowed for coagulation and the presence of anions.

J. S. G. T.

Rôle of the Hydrogen-ion Concentration in the Precipitation of Colloids. HERMANN V. TARTAR and ZALIA JENCKS GAILLEY (*J. Amer. Chem. Soc.*, 1922, 44, 2212—2218).—The effect of the hydrogen-ion concentration on the precipitation of mastic and gamboge sols by solutions of acids and salts has been investigated. It is shown that acids cause precipitation at the same hydrogen-ion concentration irrespective of the concentration of the colloid. The negative ion of the acid is without effect. Various potassium and ammonium salts precipitate the sols at the same concentration provided that the hydrogen-ion concentration is kept approximately constant. The precipitating values of the salts vary directly as the concentration of the colloid at the same hydrogen-ion concentration. The stabilising or peptising effect of the ion bearing a charge similar to that of the colloidal particle is shown to be very limited if it exists at all.

J. F. S.

Soaps and Proteins. II. Colloid Chemistry of Soap Manufacture, Analogies in the Colloid Chemistry of Soaps, Albumins, and Tissues. III. Colloid Chemical Behaviour of Fatty Acids and their Derivatives and the Analogous Behaviour of Neutral Proteins and their Derivatives. MARTIN H. FISCHER [with GEORGE D. McLAUGHLIN and MARIAN O. HOOKER] (*Koll. Chem. Beihefte*, 1922, 16, 99—179; cf. this vol., ii, 430).—A continuation of previous work. The principles of the hot and

cold processes in the soap industry are described. The significance of some of the constants of oils and fats, such as the density, saponification number, the iodine number, and the Reichert-Meissl number, is pointed out. The physical properties of soaps are discussed. An account of experiments on the systems, globulin-water, globulin-sodium hydroxide, globulin-acid, globulin-metallic chloride, gelatin-water, gelatin-alkali hydroxide, gelatin-acid, and gelatin-salt is given and peptisation and coagulation are discussed. The theory of poisoning by ammonium and heavy metal salts is discussed and it is shown that these substances combine with the protoplasm with the formation of compounds which are much less hydrated than the normal protoplasm compounds, and that this constitutes the poisoning. Reproductions of a large number of photographs, illustrating the experiments, are contained in an appendix to the paper. J. F. S.

Action of Anions, particularly the Hydroxyl-ion, on the Colloidal Condition of Night-blue. S. AKAMATSU (*Kolloid Z.*, 1922, 31, 209—215).—The effect of various anions on the surface tension of solutions of night-blue has been investigated to ascertain whether or no the position of the hydroxyl-ion in the activity series found by Traube (A., 1912, ii, 740) is correct. This series places the hydroxyl ion after the halogen ions instead of in the most active position as was expected. The surface tension of 0.2% solutions of night-blue containing concentrations varying from $N/10$ to $N/1000000$ of potassium iodide, bromide, chloride, nitrate, thiocyanate, chlorate, sulphate, hydroxide, and dihydrogen phosphate has been measured. It is shown that the addition of the various electrolytes causes the surface tension to rise to a maximum and then to fall to a value approaching the water value with decreasing concentration of the added electrolyte. The measurements showed that the hydroxyl ion is the most active of all anions and that the reason for the position given to it by Traube lies in the fact that he took as his concentration of hydroxyl ions the concentration of the hydroxide added. In the present work the concentration of the hydroxyl ion has been measured electrometrically. J. F. S.

The Equilibrium between some Organic Substances. H. HAMMERSTEN (*Arkiv Kem. Min. Geol.*, 1922, 8, No. 16, 1—14).—The equilibrium constant, K , for the reaction between phenylhydrazine and acetone or acetaldehyde has been investigated by the solubility method of Euler and Svanberg (A., 1921, i, 68), and also by the measurement of the refractive indices of the solutions. The various results are not very consistent, but in the case of acetone-phenylhydrazine the value of K appears to be about $7-9 \times 10^{-4}$, and in the case of acetaldehyde-phenylhydrazine about 10^{-3} . W. O. K.

The Solubility of Calcium Carbonate in Water in Equilibrium with a Gaseous Phase containing Carbon Dioxide. YUKICHI OSAKA (*Mem. Coll. Sci. Kyoto*, 1922, 5, 131—141).—The solubility of calcium carbonate in water has been calculated

from the data given by McCoy and Smith (A., 1911, ii, 379). It is assumed that carbon dioxide dissolved in water is a much stronger acid than acetic acid (cf. Strohecker, A., 1916, ii, 522, and Pusch, A., 1916, ii, 477, 557) and that the gas is only slightly combined with water. The apparent first dissociation constant and the second dissociation constant of water are taken as 3.50×10^{-7} and 4.91×10^{-11} , respectively; and the solubility product of calcium carbonate (calcite) at 25° was calculated to be 7.24×10^{-9} . The solubility of calcium carbonate in water, in the presence of carbon dioxide, is given by $C = [Ca^{++}] + 1 - \gamma/2\gamma[HCO_3']$, where γ is the degree of dissociation of calcium hydrogen carbonate, assuming that the solution is ionised to the same extent as that of calcium acetate of the same concentration. Formulæ are given for the relationship between $[Ca^{++}]$, and $[HCO_3']$, and the partial pressure of carbon dioxide.

W. E. G.

The Propagation of Explosion Waves in Gases contained in Tubes of Varying Cross-section. COLIN CAMPBELL (T., 1922, 121, 2483—2498).

Limits for the Propagation of Flame in Vapour-Air Mixtures. II. Mixtures of More than One Vapour and Air at the Ordinary Temperature and Pressure. ALBERT GREVILLE WHITE (T., 1922, 121, 2561—2577).

The Velocity of the Action of Oxygen, Hydrogen Sulphide, and the Halogens on Metals. G. TAMMANN and W. KÖSTER (*Z. anorg. Chem.*, 1922, 123, 196—224).—When iodine, chlorine, or air acts on a metal, a variety of surface colours is formed, and these colours give the rate of thickening of the film of compound formed (cf. A., 1920, ii, 248). The rate of chemical action is high but diffusion of the gas through the film of compound is slow. In the case of iodine, the rate is inversely proportional to the thickness of the film and is independent of the temperature. There is in the case of the action of iodine on silver a change in the velocity at 145° which is the transition temperature of silver iodide. In the case of the action of moist air containing chlorine and bromine on copper, silver, and lead there is a parabolic relationship between the time of action and the thickness of film. The same holds in the case of the oxidation of metals. At high temperatures oxidisable metals are almost immediately covered with a thin film of oxide which protects the metals.

Hydrogen sulphide reacts rapidly with copper and manganese, more slowly with silver, and still more slowly with lead. Iron, cobalt, nickel, chromium, antimony, bismuth, tin, aluminium, zinc, and cadmium show no change after several days' contact with the gas. The action of air on copper-zinc alloys was also investigated.

W. T.

The Formation of Oxides of Multivalent Metals from their Hydroxides. HANS VON EULER and ULF VON EULER (*Z. anorg. Chem.*, 1922, 124, 70—80).—The temperature-coefficient of the coagulation of albumin is very high, the rate being doubled

for an increase of temperature of 2° . The rate of dehydration of copper hydroxide was measured at different temperatures. The temperature-coefficient was very high, as in the coagulation of albumin, but was found to decrease with increasing temperature $40-45^{\circ}=3.0$ for 5° , $55-60^{\circ}=1.8$. The results were not in agreement with Arrhenius's formula (A., 1889, 1103), which led the authors to the conclusion that the dehydration consists of several reactions. The rate of dehydration was measured colorimetrically. The isoelectric points for copper hydroxide is 3×10^{-13} as obtained from the equation $I = \sqrt{K_s \cdot K_w / K_b}$. W. T.

Velocity of Hydrolysis of Ethyl Acetate. HERBERT S. HARNED and ROBERT PFANSTIEL (*J. Amer. Chem. Soc.*, 1922, **44**, 2193—2220).—The unimolecular velocity constants of the hydrolysis of ethyl acetate in the presence of hydrochloric acid of many concentrations have been accurately determined at 25° . A solution of the general equation for the velocity of hydrolysis has been obtained and the velocity constants have been calculated by the general equation. In four series of measurements, it has been found that the curve of the velocity constants divided by the molecular concentration of hydrochloric acid against $\log c_1$, where c_1 is the molecular concentration of hydrochloric acid, shows a minimum at $0.07-0.08N$ -acid. This is similar to the curve of the individual hydrogen-ion activity coefficient against $\log c_1$, which has a minimum at $0.15-0.18N$ -acid. It has been shown that the velocity constant divided by the product of the activities of the hydrogen ion and the water molecule is not a constant at different acid concentrations but has a maximum at $0.3N$ -hydrochloric acid. Some factors which may cause this deviation from constancy have been suggested. The kinetics of the hydrolysis of ethyl acetate are very complex, but the authors are of the opinion that the evidence obtained in the present work shows that the method employed is in general the correct method of attacking this problem. J. F. S.

Velocity of Inversion of Sucrose as a Function of the Thermodynamic Concentration of the Hydrogen-ion. HAROLD A. FALES and JACQUE C. MORRELL (*J. Amer. Chem. Soc.*, 1922, **44**, 2071—2091).—The velocity of hydrolysis of sucrose by solutions of sodium hydroxide, acetic acid, and hydrochloric acid of various concentrations has been determined at 35° and the thermodynamic hydrogen-ion concentration has been calculated from *E.M.F.* determinations of the reaction mixtures at the same temperature. It is shown that the addition of sucrose to solutions of hydrochloric acid causes a definite increase in the thermodynamic concentration of hydrogen-ion over the whole range of concentration of the acid. In the case of acetic acid, a maximum increase in hydrogen-ion concentration is observed which diminishes as the concentration of hydrogen-ion decreases, on the addition of sucrose. The addition of sucrose to sodium hydroxide appears to cause an increase in the thermodynamic concentration of

hydrogen-ion. The thermodynamic concentration of hydrogen-ion in the system remains appreciably constant during the course of the inversion process. The velocity of inversion of sucrose by hydrochloric acid is proportional to the thermodynamic concentration of hydrogen-ion only for the limited range of concentration of acid which lies between 0.01*N* and 0.001*N*. The bimolecular velocity constants, reduced to unit thermodynamic hydrogen-ion concentration, are constant with changing concentrations of sucrose only for each molecular concentration of hydrochloric acid between 0.01*N* and 0.3*N*. These constants vary considerably for different concentrations of acid. Evidence is adduced which shows that for acetic acid, the undissociated molecule is not a positive catalyst in the inversion process; or that there are other factors entering here which have never been considered. With strong acids, the inversion process does not appear to be strictly unimolecular, especially in the early periods of the reaction. Within the limits of the concentrations of the reactants used, no appreciable change in volume was noticed when the solutions of hydrochloric acid and sucrose were mixed.

J. F. S.

Generator Gas Equilibrium at High Pressures. KARL JELLINEK and ALOIS DIETHELM (*Z. anorg. Chem.*, 1922, **124**, 203—229).—The electric furnace employed was a modification of that used by Nernst (*Z. Elektrochem.*, 1907, **13**, 52); it allowed measurements to be carried out at 1300° and at 150 atmospheres. The reaction $C + CO_2 \rightleftharpoons 2CO$ was investigated between 800° and 1000° at pressures up to 50 atmospheres. Various forms of charcoal were employed, and the equilibrium, which was well defined, was practically the same in each case, and again the same in the case of graphite. The law of mass action was obeyed at all pressures. The integration constant of the reaction isochore of generator gas equilibrium was found to be 3.876; this agrees well with the value calculated by means of Nernst's heat theorem (3.80). The authors show that the maximum work of this reaction is equal to the heat effect if the reaction is carried out isothermally and reversibly and if the excess of oxygen has the same pressure as the carbon dioxide formed.

W. T.

Catalytic Activity of Copper. O. W. BROWN and C. O. HENKE (*J. Physical Chem.*, 1922, **26**, 715—727).—In continuation of previous work (this vol., i, 445), the authors have investigated the catalytic reduction of nitrobenzene to aniline by hydrogen in the presence of copper catalysts prepared by the reduction of the precipitated oxide. Sabatier's conclusion that a catalyst so prepared is more efficient than one prepared from an ignited nitrate was confirmed. By the use of the former catalyst, much lower rates of supply of hydrogen may be used without reduction in yield, and the reaction may be carried out at a much greater rate. The best temperature for carrying out the reduction was found to be about 260°. A small amount of iron in a copper catalyst prepared from an ignited oxide prevents the rapid decrease in activity of the

catalyst. Copper deposited on asbestos was found to be capable of effecting the reduction at a much greater rate per gram of copper than copper alone, but with high rates of supply of nitrobenzene, it quickly lost its activity. Copper on asbestos is a better catalyst than an equal volume of copper on pumice. J. S. G. T.

Selective Activation of Alumina for Decarboxylation or for Dehydration. HOMER ADKINS (*J. Amer. Chem. Soc.*, 1922, 44, 2175—2186).—Alumina has been preferentially activated for decarboxylation or for dehydration by modifying the method of preparation. This has been accomplished by obtaining aluminium hydroxide and hydrated alumina by precipitation from its salts or esters in water or xylene solution, and by preparing it by the action of water vapour on aluminium alkyl oxides, both in the pure state and when the latter are impregnated in pumice. The extent to which selective activation has been accomplished is indicated by the following figures, which give the relative "ethylene efficiency" of the variously prepared catalysts with ethyl acetate vapour at 465°: 34, 38, 41, 50, 107, and 150. The catalysts indicated were prepared as follows: by melting aluminium isopropoxide on pumice and exposing to moist air for eighteen hours (34); as last, but aluminium isobutoxide was used (38); by the action of moist air on aluminium methoxide; this was used as compressed tablets and contained traces of mercury (41); alumina from aluminium butoxide supported on pumice as above (50); tablets made from the precipitate produced by the action of hydrochloric acid on solutions of sodium aluminate (107); and tablets made from alumina obtained by the action of water on aluminium amalgam (150). An hypothesis has been advanced in which it is considered that the catalytic activity of alumina is conditioned by its molecular porosity, or the distances between the aluminium atoms and this, in part, is determined by the size, shape, and position of the radicles attached to the aluminium when the aluminium compound passes into the solid state. All the experimental evidence obtained with alumina catalysts is in complete harmony with this hypothesis. In terms of this hypothesis, it is held that decarboxylation is favoured by large pores in the alumina and that ethylene formation is favoured by small pores, large and small referring to units of molecular dimensions. The work of Milligan and Mead (this vol., ii, 447) on the dehydration and X-ray pattern of aluminium hydroxide is discussed in its relation to catalytically active alumina. The results obtained by Palmer (*A.*, 1921, ii, 542), in the use of electrolytic copper and copper by reduction for the hydrogenation of alcohol, are explained on the basis of the hypothesis noted above. Cf. Langmuir (this vol., ii, 629). J. F. S.

The Influencing of Catalysts and Specifically Active Catalysts. KARL W. ROSENMUND and F. ZETZSCHE (*Ber.*, 1922, 55, [B], 2774).—A reply to Abel (cf. Rosenmund and Zetzsche, *A.*, 1921, ii, 320, 392, 393; this vol., ii, 41; Abel, *A.*, 1921, ii, 542; this vol., ii, 26). H. W.

Kinetics of Inductive Processes : Schönbein's Reaction. A. V. PAMFILOV and N. N. PETIN (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, 6, 221—231).—The authors discuss previous work published on the influence of ferrous oxide on the liberation of iodine from an iodide by the action of hydrogen peroxide, and give the results of their own experiments on this reaction. It is found that not only the velocity of the initial stage of the reaction, but also the nature of its dependence on the concentration of the iron salt and on the corresponding acidity are completely different according as the iron is in the ferric or the ferrous condition. The liberation of iodine by ferric oxide or hydrogen peroxide separately is considerably accelerated when the two reactions occur together, and the velocities of the separate, and that of the combined, reactions increase with increase of the concentrations of all the components; the acidity of the medium is of comparatively little influence, but in general accelerates the reaction.

On the other hand, in neutral and faintly acid media the reaction is accelerated more by ferrous oxide than by ferric oxide in low concentration, but for definite conditions of concentration the reaction is retarded by increasing the concentration of the ferrous oxide either absolutely or in relation to the potassium iodide; this retardation is observed also in an acid medium. Increase in the acidity also retards the reaction considerably. This phenomenon disappears when the concentration of the potassium iodide increases and the solution is rendered strongly acid, especially as liberation of iodine by the acid becomes pronounced under these conditions.

Hence the diminution in the sensitiveness of Schönbein's reaction in a neutral medium caused by increase in the concentration of the catalyst (cf. Meissner, "Untersuchungen über den Sauerstoff," 1868, 78—80; Struve, *Z. anal. Chem.*, 1869, 8, 319) is not fortuitous but holds generally, both in acid and in neutral solutions. The results of replacement of the acid by the corresponding concentration of a sulphate show, as was found by Brode (A., 1901, ii, 443) for ferric-ions, that the influence of increased acidity is not to be referred to diminished concentration of the ferrous ions and that the hydrogen-ions must here play a part.

The mechanism of the action of ferrous oxide, typical of inductors in general, is probably based on its spontaneous oxidation, and the retarding influence exerted on the reaction by acidity is related to the great stability of ferrous-ions in an acid solution. T. H. P.

Union of Hydrogen with Acetylene Derivatives. XV. Velocities of Catalytic Reactions. J. S. ZALKIND (*J. Russ. Phys. Chem. Soc.*, 1920, 52, 199—211).—The author discusses the results obtained by various investigators on the course followed by reactions occurring under the influence of enzymes and other catalysts. The general conclusion drawn is that, even when catalysts so chemically simple as palladium and platinum are used, the course of the reaction is often quite complex, and that no essential difference exists between the actions of these simple catalysts and those of enzymes. In both cases, it is necessary to

assume the formation of intermediate compounds between the catalyst and the reacting compounds, although with regard to the means by which such intermediate compounds are formed—whether by chemical reaction or by simple adsorption—opinions differ. Adsorption phenomena may, indeed, exhibit complex character and a sharp distinction between adsorptive and chemical phenomena is scarcely possible.

T. H. P.

Speculations concerning the Positive Electron. [SIR] OLIVER LODGE (*Nature*, 1922, **110**, 696—697).—According to Larmor's theory, the positive and negative electrons can only differ in one being the mirror-image of the other. No positive electron in this sense has yet been discovered. Both positive and negative electrons are supposed to be extremely mobile, and the forces between them immense, so that they instantly bind themselves together into a compact and stable structure consisting of several thousands of each kind, the simplest and lightest of these composite structures being the hydrogen nucleus or proton. The proton for the first time allows a negative electron to revolve round it without being absorbed into its composition. On the mirror-image hypothesis, however, it becomes difficult to explain why only negative electrons should occur in the satellite systems; that is, why some atoms should not have nuclei with a surplus of negative charge, and be attended by positive electrons. The case of a number of primary electrons, both positive and negative, is considered. If, in the many possible ways of structural packing, something less than half the atomic nuclei formed were of the kind with positive satellites, whilst the other approximate half were of the negative satellite variety, the two classes would speedily combine with great violence and thereby form the tight-packed and stable nuclei of heavier atoms, until the complexity was too great to ensure stability. The combination of nuclei would cease only when they were protected by a screen of similar electrons, of which the sign might be positive or negative, but could not be both; that is, the resulting atoms could not be of opposite varieties. Thus the outlying satellites are all of one sign, either in every case or in so vast a preponderance of cases that no exceptions are as yet manifest. The free lifetime of the less plentiful variety would be too brief for ordinary detection, but it is suggested that by means of the shattering of nuclei and rapid methods of detection, the composite nature of the proton may possibly be demonstrated by the emission of something fractional of extreme instability. In this connexion, attention is directed to the fact that the atomic bombardment of aluminium yields particles of specially long range.

A. A. E.

Motions of Electrons in Argon and Hydrogen. J. S. TOWNSEND and V. A. BAILEY (*Phil. Mag.*, 1922, [vi], **44**, 1033—1052; cf. this vol., ii, 277, 494).—Further measurements have been made of the movement of electrons in argon and hydrogen, and in mixtures of the two gases, in a uniform electric field. The velocity of the electrons (W) in the direction of the electric force, their mean velocities of agitation in the gas (u), their mean free path (l), the

loss of energy on collision with a gas molecule (λ), and the factor (k) by which the energy of agitation of an electron exceeds the energy of a molecule of gas at 15° , are given for varying values of Z/p , where Z is the fall in potential in volts, and p is the gas pressure in millimetres. The loss in energy of an electron on collision with a molecule of argon is extremely small. When moving with a velocity of agitation of 12.6×10^7 cm. per sec., the fraction of energy lost by an electron on collision with a molecule is 1.6×10^{-5} in argon, 5×10^{-2} in nitrogen, and 4×10^{-2} in hydrogen, and the corresponding mean free paths are 0.147 cm. in argon, 0.029 cm. in nitrogen, and 0.035 cm. in hydrogen, the gases being at 1 mm. pressure.

In the case of argon, whereas W increases steadily with increase in Z/p , u increases to the constant value, 20.7×10^7 cm. per sec. Between the values of Z/p , 1.25—15, the velocity of agitation remains practically constant, whilst the loss of energy on collision steadily increases. This is due to a large loss in energy in collisions with velocities greater than the mean. The free path of an electron in argon at 1 mm. pressure has a maximum value of 1.6 cm. for the velocity 3.75×10^7 cm. per second, whereas the mean free paths of nitrogen and hydrogen show minima in the neighbourhood of 8×10^7 cm. sec. The large increases in the mean free paths of the electrons as the velocity diminishes are very striking.

The addition of argon to hydrogen, whilst reducing W , the velocity in the direction of the electric force, scarcely affects u , the mean velocity of agitation of the electrons. This is in agreement with the small loss in energy on collision with a molecule of argon. The mean free paths in argon, for velocities of agitation less than 11.5×10^7 cm. per second, were found by measurement of the mean free paths in mixtures of hydrogen and argon. W. E. G.

The Motion of Electrons in Carbon Dioxide. M. F. SKINKER (*Phil. Mag.*, 1922, [vi], 44, 994—999).—The apparatus used was similar to that of Townsend and Bailey (cf. this vol., ii, 277, 494). Curves are given comparing the speed of electrons in carbon dioxide, nitrogen, and hydrogen. Values for the mean free path of the electron, its speed in the direction of the electric force, its velocity of agitation, and the proportion of energy lost on collision with a molecule of gas, are tabulated against different values of Z/p (cf. Townsend and Bailey, preceding abstract). A sudden increase in the loss of energy which occurs on collision between an electron and a molecule of carbon dioxide is observed between the velocities of agitation 13×10^7 and 15×10^7 cm. per second. W. E. G.

The Constitution of the Chemical Elements. FRIEDRICH WÄCHTER (*Z. anorg. Chem.*, 1922, 123, 1—13).—A theoretical paper. The author assumes that the light ether is a ponderable gas, an atom of which has a mass $1/1800$ the mass of an atom of hydrogen. He regards the elements as different condensation forms of the light ether, the number of ether atoms in an atom of any element being 1800 times the atomic weight of the element. This is applied to the atomic heat of elements and to the spectral lines. W. T.

The Disintegration of Chemical Elements. FRIEDRICH WICHTER (*Z. anorg. Chem.*, 1922, 123, 14—30).—A theoretical paper. The author assumes that hydrogen and helium atoms are made up of light ether atoms, and that other elements are built up of hydrogen and helium atoms. He then endeavours to explain the disintegration of the radio elements. W. T.

Electronic Structures of Atoms. MAURICE L. HUGGINS (*J. Physical Chem.*, 1922, 26, 601—625).—A theoretical paper in which the Lewis hypothesis of valency and atomic structure is assumed to be correct for the lighter atoms. The evidence in favour of a tetrahedron of electron pairs, rather than a cube of single electrons, being the most stable arrangement for the valency shell of these atoms, is put forward. It is shown how the stability of such an arrangement may be the result of the deviation of the force law between two electrons from Coulomb's law at small distances. The chief requirement for this purpose is that as two electrons approach each other, the repulsion between them reaches a maximum at a distance of the order of magnitude of 1 Å., the force then decreasing, for a considerable proportion of the remaining distance at least. As a result of a law of this type, whenever two electrons, forming part of an atomic shell, are forced close enough together, they come still closer to each other, that is, the pair off, and similarly from three pairs aided by the attraction of an atomic nucleus and the repulsion of other electrons, triplets are formed. Based on this assumption and on the idea that the electrons in an atomic shell tend to place themselves opposite the centres of the faces of the imaginary polyhedron formed by the electron groups in the next underlying kernel shell, an arrangement of electrons is obtained for each element in each valency stage, with a few exceptions which have not yet been deduced. These arrangements conform with the known properties of the elements, and account for the periodic relationships in a simple and logical manner. J. F. S.

Bohr's Model of the Hydrogen Molecules and their Magnetic Susceptibility. KÔTARÔ HONDA (*Nature*, 1922, 110, 664—665).—Since, according to Langevin, hydrogen molecules should have a paramagnetic susceptibility, it has hitherto been assumed that Bohr's model does not explain the diamagnetic property of the gas. The actual diamagnetic susceptibility, as determined by Soné, is shown to lie between the two values calculated when the validity of Bohr's model is assumed in the case of $n+n'=1$ and $n+n'=2$, where n and n' are the azimuth and radial quantum numbers. A. A. E.

The Normal Helium Atom and the Quantum Theory. J. H. VAN VLECK (*Phil. Mag.*, 1922, [vi], 44, 842—869).—A study is made of the Kemble model of the helium atom. In this model the two electrons are arranged with axial symmetry. The energy of this atom has been computed, and the calculation has given a value of 20.7 volts for the ionisation potential, instead of 25.4

demanded by experiment, the discrepancy being slightly greater than for the Bohr model. It does not seem possible to devise a satisfactory symmetrical model of helium based on the conventional quantum theory of atomic structure. W. E. G.

The Neuburger Nuclear Model. MAXIMILIAN CAMILLO NEUBURGER (*Physikal. Z.*, 1922, 23, 389).—A reply to Meitner (cf. this vol., ii, 702). The author's model of the structure of the atomic nucleus is not dependent on the existence of *isohelium* in the free state. The nuclear formula requires, however, that groups, composed of three hydrogen nuclei and an electron, shall be capable of existence in the nucleus itself. W. E. G.

Negative Valency and Co-ordination Number. E. A. SCHILOV (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, 6, 281—290).—The most pronounced negative affinity is accompanied by non-metallic properties, so that, if it is assumed that negative affinity is the result of a tendency of the rings of valency electrons to attain saturation, it becomes necessary to make the supplementary assumption that only non-metallic rings of a special type exhibit such tendency. It is suggested that the origin of complex compounds may be represented on an analogous principle, co-ordinative affinity arising as a result of the tendency towards saturation of non-dissociating, non-metallic atomic skeletons.

T. H. P.

Quanta of the Ideal Monatomic Gas. K. SCHAPOSCHNIKOV (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, 6, 105—124).—A mathematical paper, in which the quantum theory is applied to gases at high and at low temperatures.

T. H. P.

Chemical Constants of some Diatomic Gases. J. R. PARTINGTON (*Phil. Mag.*, 1922, [vi], 44, 988—993).—The investigations of Sackur (cf. A., 1913, ii, 128), and Tetrode (*Ann. Physik*, 1912, [iv], 38, 434; 39, 225) on the chemical constants of monatomic gases have been extended to the case of a hypothetical diatomic molecule. In the first case considered, the gas molecule is assumed to consist of two identical atoms rigidly attached to each other at a fixed distance apart. The method of calculation is based on generalised statistical mechanics, and it is shown that the following equation holds over a certain range of temperatures, $C = 2.5 \log M + 2 \log K + 12.730$, where K = radius of gyration and M the molecular weight. This gives values for the chemical constants $C_{O_2} = 1.001$, $C_{N_2} = 0.904$, and $C_{H_2} = -2.255$. On account of the very small mass and diameter of the hydrogen molecule, hydrogen should behave in a manner intermediate between that of a monatomic and a diatomic gas. In the second case, that of two different atoms rigidly bound together, it is necessary to add $\log 2$ to the above equation, and $C_{CO} = 1.205$, and $C_{NO} = 1.263$. These results are in all cases of the right order. W. E. G.

A New Apparatus for Washing Gases. A. ERNST (*Chem. Listy*, 1922, 16, 274—276).—The tube leading the gas to be purified

into the washing bottle is surrounded by a long spiral, along which the gas has to make its way and is thus well washed. The tube leading in the gas and the spiral are easily removed from the vessel.

W. T.

Protection of Brass Weights. J. J. MANLEY (*Phil. Mag.*, 1922, [vi], 44, 948—950).—The method introduced by Faraday for protecting iron from rust has been employed for protecting brass weights. A satisfactory method of procedure is to cover the weight with a thin film of linseed oil, and heat it, appropriately supported in a covered "vitreosil" crucible or closed muffle, until the weight assumes a golden tint. The protecting film which is formed is tough and may be polished. The weight of 1 sq. cm. of normal film is 0.00005 gram.

W. E. G.

Lecture Experiments with Hydronitric [Hydrazoic] Acid and the Trinitrides [Azides]. A. W. BROWNE and A. B. HOEL (*J. Amer. Chem. Soc.*, 1922, 44, 2116—2119).—A series of lecture experiments with azoimide is described which includes the following. *Nitridation of hydriodic acid*: 5 c.c. of hydriodic acid (*d* 1.5) are placed in a 15 cm. test-tube and 1 c.c. of 1% hydrazoic acid is added when both iodine and nitrogen are liberated according to the equation $N_3H + 2HI = NH_3 + N_2 + 2I$. This experiment illustrates the similarity between hydrazoic acid and hydrogen peroxide. *Nitridation of hydrochloric acid*: 5 c.c. of hydrochloric acid are treated with 1 c.c. of 1% hydrazoic acid and raised to the boiling point for one minute; the contents of the tube are then poured into 2½ litres of water containing starch and potassium iodide, when the free chlorine liberates iodine. The reaction is represented by the equation $N_3H + 2HCl = NH_3 + Cl_2 + N_2$. *Formation of potassium manganate by the action of potassium azide on manganese dioxide*: 0.5 gram of finely powdered potassium azide and 0.5 gram of manganese dioxide are mixed and gently heated in a 100 c.c. porcelain crucible covered by a 200 c.c. crucible. The reaction is violent and proceeds with a hissing noise and the emission of numerous sparks. When the reaction has moderated, the larger crucible is removed and the heating continued for a short time. After cooling the mass, it is thrown into 2½ litres of water, when the green solution produced speedily changes to that of a permanganate solution. In this experiment it is advisable that the operator should not come too close to the crucible at the commencement of the experiment. *Detonation of silver azide by means of a fuse consisting of paper impregnated with potassium azide*: Silver azide, prepared by adding a dilute solution of silver nitrate cautiously to 25 c.c. of a 1% solution of hydrazoic acid, filtering and washing with water, is made into a thin disk 18 mm. diameter and 1 mm. thick, while still moist. This is then stuck on a piece of filter-paper prepared as follows: a strip 0.5—1.0 cm. wide and 0.25—0.50 metre long is painted on a filter-paper by means of a small brush dipped in a 5% aqueous solution of potassium azide and dried. The disk of silver azide is attached to one end of the strip and the whole thoroughly dried. The dried paper is then suspended and a glowing

splint brought in contact with the end of the strip removed from the silver azide; the combustion gradually progresses toward the silver azide, which finally detonates with a sharp report. J. F. S.

Inorganic Chemistry.

Diffusion of Hydrogen and Helium through Silica Glass and other Glasses. G. A. WILLIAMS and J. B. FERGUSON (*J. Amer. Chem. Soc.*, 1922, **44**, 2160—2167).—The rate at which hydrogen and helium, respectively, leak into evacuated tubes of silica glass, pyrex glass, and Jena glass has been determined at a series of temperatures and pressures. It is shown that silica glass is permeable to hydrogen at high temperatures. The permeability is proportional to the gas pressure and is an exponential function of the temperature. It becomes appreciable at 300°. Silica glass is similarly permeable to helium, and this is easily measurable at 182°. At 500°, the permeability for helium is about twenty-two times that for hydrogen. Neither pyrex glass nor Jena combustion glass was found to be permeable to hydrogen, but apparently hydrogen reacts with pyrex glass causing it to blacken. At 610°, pyrex glass is permeable to helium. The permeabilities of a number of samples of glass have been determined and the results compared with those of Wüstner (*Ann. Phys.*, 1915, [iv], **46**, 1095) and others. J. F. S.

The Solubility of Oxygen in Various Organic Solvents. FRANZ FISCHER and GEORG PFLEIDERER (*Z. anorg. Chem.*, 1922, **124**, 61—69).—The solubility was determined by pumping the gas from the saturated solution by means of a mercury-air pump and measuring the amount obtained. The results embodying the solubility of oxygen in sixteen organic solvents are tabulated; the specific gravities and vapour pressures are also given. W. T.

A New Form of Ozoniser. H. NĚMEČEK (*Chem. Listy*, 1922, **16**, 276).—Two stout copper wires serve as electrodes in a horizontal glass cylinder, the arrangement being such that the distance between the electrodes can be varied. The energy can be obtained by means of a Rhumkorff's coil or an electric machine. W. T.

Hydrophobic and Hydrophilic Sols of Sulphur. H. FREUNDLICH and P. SCHOLZ (*Koll. Chem. Beihefte*, 1922, **16**, 234—266).—The behaviour of the hydrophobic and several hydrophilic sols of sulphur has been compared. As a markedly hydrophile sol, that investigated by Odén, and prepared by the interaction of hydrogen sulphide and sulphur dioxide, and by the decomposition of thionic acids, has been considered, and as a hydrophobe sol, that prepared by Weimarn by pouring an alcoholic

solution of sulphur into water, is considered. The Weimarn sulphur sol has been shown to be completely hydrophobic; it is negatively charged and is coagulated by electrolytes according to the general laws concerning the coagulation of hydrophobic sols, namely, the strong influence of the absorbability and valency of the kations, the strong effect of the hydrogen-ion, and the slight effect of the hydroxyl-ion. The sols may only be prepared in relatively small concentrations, the coagulum is not peptised by washing with water, and the sols may be preserved for a day or two only. In keeping with Odén's results, the coagulation of Odén's sol is markedly different from that of Weimarn's sol, as is demanded by its greater hydrophilic behaviour. The difference is mainly shown in the following points: (i) alkali salts have a ten to twenty times weaker coagulating action than with Weimarn's sol; (ii) the lyotropic series of the kations is well defined; lithium-ions have a very weak coagulating action, whilst sodium-, potassium-, rubidium-, and caesium-ions are markedly and increasingly stronger; (iii) acids have a still weaker coagulating action than the alkali salts; (iv) alkalis convert the Odén sol into one which has similar properties to the hydrophobic Weimarn sol. The assumption that the micellæ of Weimarn's sol consist of λ -sulphur and those of Odén's sol of μ -sulphur is insufficient to explain these differences; for neither the behaviour towards acids nor that towards bases is in the least explained by the above assumption, whilst a very possible explanation is that Odén's sol contains pentathionic acid, whereas Weimarn's sol does not. The presence of pentathionic acid in Odén's sol is proved by the following data: (i) the acid can be detected in the filtrate from coagulated sulphur by the fact that on the addition of alkali more sulphur is deposited; (ii) prolonged action of alkali on coagulated sulphur or sulphur micellæ brings about the formation of thiosulphate; this action is brought about by ammonia without the sulphur itself being markedly attacked. It is probable that the pentathionic acid is adsorbed in the sulphur particles, since this substance is very difficult to remove by washing. The quantity of pentathionic acid bound to the sulphur is found to be 0.1—0.7 millimol. per gram of sulphur. The influence of pentathionic acid on the stability of Odén's sol is explained as follows. This acid has a constitution which is closely related to that of water and sulphur and therefore it facilitates the formation of micellæ, which, in addition to sulphur and pentathionic acid, contain large quantities of water. The greater water content of Odén's sol is manifested by its greater transparency when compared with Weimarn's sol of equal sulphur content. The micellæ are to be represented in Odén's sol by the

formula $\begin{array}{c} S\mu \\ S_5O_6H_2 \\ H_2O \end{array} S_5O_6''$, and in Weimarn's sols by $\begin{array}{c} S\lambda \\ S_2O \\ H_2O \end{array}'$. The

structure of the micellæ of Odén's sol explains the behaviour on coagulation as regards the following points: (i) alkalis decompose the pentathionic acid, whereby the loose combination

between the sulphur and water is also disturbed and converted into one which is similar to that of the Weimarn sol; (ii) acids have such a feeble coagulating power because they increase the stability of the pentathionic acid and produce more pentathionic acid from the pentathionates which may be present. The structure also is in keeping with the fact that hydrogen sulphide has an action on the sol similar to that of the alkalis; that is, it converts it into an hydrophobic sol which is much less stable because the pentathionic acid is decomposed by hydrogen sulphide with separation of sulphur. Further, sols of the same nature as Odén's sol are obtained by reactions which produce both sulphur and pentathionic acid; for example, the decomposition of sulphur monochloride by water: $5\text{S}_2\text{Cl}_2 + 6\text{H}_2\text{O} = 5\text{S} + \text{H}_2\text{S}_5\text{O}_6 + 10\text{HCl}$. The stability of Odén's sol, which is due to pentathionic acid, does not depend on the charge of the micellæ. Whether or no Odén's sol contains S_μ cannot be definitely answered, although the yellow colour of the sol makes it extremely likely that this variety of sulphur is present.

J. F. S.

The Existence of Sulphur Tetroxide. FRIEDRICH MEYER, GUSTAV BAILLEUL, and GERHARD HENKEL (*Ber.*, 1922, 55, [B], 2923—2929).—The experiments described owed their origin to a desire to examine Berthelot's sulphur heptoxide more fully, and as a considerable quantity of the product was desired, the original method was modified by submitting a current of sulphur dioxide and oxygen to the silent electric discharge instead of working with a fixed volume of the gases. Contrary to Berthelot's statements, the product which is formed does not appear to have an appreciable vapour tension even at 20° . The crystalline substance described by Berthelot is only formed when analysis shows that sulphur trioxide is present in excess. On the other hand, an amorphous product is obtained the analyses of which sometimes agree with the formula, S_2O_7 , but sometimes show more oxygen than is required by this expression. The latter substance has been prepared in a specially designed discharge tube (which is figured in the original), which is so constructed that it is sufficiently small and light to be weighed on an analytical balance, that it can be evacuated and kept vacuous, and has little dead space. The material, thus prepared in the tube from sulphur dioxide and oxygen, is weighed and analysed in situ. For the latter purpose, it is treated with an aqueous solution of potassium iodide with which, at a sufficiently low temperature, it reacts with unexpected slowness and with the liberation of a very small proportion of oxygen. The latter is measured together with the liberated iodine and sulphuric acid. The highest stage of oxidation yet reached corresponds with the formula, $\text{SO}_3 \cdot 2\text{SO}_4$. The authors, however, do not consider that this product represents the most highly oxidised form of sulphur which is possible, since in their work up to the present the imparted energy has not been sufficiently great in proportion to the volume of gas employed.

It appears possible that Berthelot's sulphur heptoxide is a

mixture of sulphur trioxide and sulphur tetroxide in equimolecular ratio.
H. W.

The Isotopes of Selenium and some other Elements. F. W. ASTON (*Nature*, 1922, 110, 664).—Very satisfactory mass-spectra have been obtained in the case of selenium by vaporising the element in the discharge tubes. Selenium is shown to consist of six isotopes, giving lines at 80, 78, 76, 82, 77, and 74, respectively, in decreasing order of intensity. The results agree with the atomic weight as determined by chemical methods, and measurement of the lines shows no detectable deviation from the whole-number rule. The mass spectra of cadmium and tellurium could not be obtained by this method, but the experiments have incidentally shown that chlorine has no isotope of mass 39, and that aluminium is a simple element of mass 27. Two new isotopes of xenon, 124 and 126, making nine in all, are found to exist in minute amounts. The first of these is isobaric with tin, and the selenium lines 78, 80, and 82 are isobares of krypton; it is noted that all isobares so far discovered have even atomic weights.
A. A. E.

Action of Compounds of Nitric Oxide and Hydroxylamine on Trisodium Arsenite. A. GUTMANN (*Ber.*, 1922, 55, [B], 3007—3012).—The oxidation of sodium arsenite to arsenate by sodium benzenediazoxide (which is not effected by the *iso*-diazoxide) has been attributed previously to the presence of a reactive, labile quadrivalent oxygen atom (A., 1912, i, 397; 1915, i, 127); an extension of the observations to a series of derivatives of nitric oxide and hydroxylamine indicates that sodium arsenite is a specific reagent for active oxygen.

Nitric oxide oxidises sodium arsenite in accordance with the equation: $2\text{NO} + \text{Na}_3\text{AsO}_3 = \text{N}_2\text{O} + \text{Na}_3\text{AsO}_4$; a similar change occurs when an excess of sodium arsenite is added to a solution of nitric oxide in freshly prepared, strongly alkaline potassium sulphite solution. Nitrosobenzene does not react with sodium arsenite at the atmospheric temperature but at the temperature of the boiling water-bath gives azoxybenzene and sodium arsenate.

The constitutions $\begin{array}{c} \text{N} \cdot \text{O} \\ || \quad || \\ \text{N} \cdot \text{O} \end{array}$ and $\begin{array}{c} \text{Ph} \cdot \text{N} \cdot \text{O} \\ | \quad || \\ \text{Ph} \cdot \text{N} \cdot \text{O} \end{array}$, are suggested for nitric oxide

and nitrosobenzene, respectively. Sodium nitroprusside does not appear to react with sodium arsenite. With hydroxylamine, the reactions $\text{NH}_2 \cdot \text{OH} + \text{Na}_3\text{AsO}_3 = \text{Na}_3\text{AsO}_4 + \text{NH}_3$ and $3\text{NH}_2 \cdot \text{OH} = \text{NH}_3 + \text{N}_2 + 3\text{H}_2\text{O}$ occur concurrently. β -Phenylhydroxylamine gives sodium arsenate, aniline, and azoxybenzene, $4\text{NHPH} \cdot \text{OH} + \text{Na}_3\text{AsO}_3 = \text{Na}_3\text{AsO}_4 + 2\text{NH}_2\text{Ph} + \text{O} < \begin{array}{c} \text{NPh} \\ | \\ \text{NPh} \end{array} + 2\text{H}_2\text{O}$, whereas the iso-

meric *p*-aminophenol is inactive. β -Ethylhydroxylamine, on the other hand, appears to give only arsenite and ethylamine. Dimethylaniline-*N*-oxide and sodium arsenite readily yield dimethylaniline and sodium arsenate. Quinine-*N*-oxide similarly gives quinine and sodium arsenate; the two oxides are therefore peroxidic in character.
H. W.

Properties and Structure of Pernitric Acid. IV. TRIFONOV (*Z. anorg. Chem.*, 1922, **124**, 123—135).—Pernitric acid is formed by the action of hydrogen peroxide on an acidified nitrite solution; low temperature and low concentration are not essential for its formation. Above 70°, it decomposes rapidly. The per-acid can be estimated by the addition of potassium bromide in dilute solution, and the liberated bromine estimated by means of a potassium iodide solution. The author finds that the reaction of formation is represented by the equation $2\text{HNO}_2 + 3\text{H}_2\text{O}_2 + (n-1)\text{H}_2\text{O} = \text{N}_2\text{O}_6 \cdot n\text{H}_2\text{O} + 3\text{H}_2\text{O}$, and not as given by Raschig (*Ber.*, 1907, **40**, 4585) or Schmidlin and Massini (*A.*, 1910, ii, 498). It thus receives the formula $\text{NO}_2 \cdot \text{O} \cdot \text{O} \cdot \text{NO}_2 \cdot n\text{H}_2\text{O}$. The reactions of pernitric acid with some organic compounds are given. W. T.

Phosphorus. W. MARCKWALD and K. HELMHOLTZ (*Z. anorg. Chem.*, 1922, **124**, 81—104).—The authors find the melting point of scarlet phosphorus to be $592.5 \pm 0.5^\circ$, and the critical temperature of the yellow modification to be 720.6° . Rapid cooling of phosphorus vapour deposits a mixture of the yellow and scarlet forms; slow cooling deposits the pure yellow modification. To explain this the authors assume that phosphorus vapour at 900° and 120 mm. pressure consists of P_4 molecules and 10% P_2 molecules. On cooling, the P_2 and P_4 molecules combine to form the complex molecules of scarlet phosphorus and this is to a certain extent deposited. Slow cooling, however, allows the scarlet phosphorus vapour to change into the vapour of the yellow form. The phenomenon observed on cooling liquid scarlet phosphorus is explained as follows. The liquid contains molecules of yellow (P_4) and scarlet (P_2) phosphorus in equilibrium: $n\text{P}_4 \rightleftharpoons m\text{P}_2$. Above 592.5° , most of the molecules are P_4 and on rapid cooling these solidify, giving the yellow variety. Slow cooling allows of a change of equilibrium to the right and therefore a deposit of the scarlet form on solidification. It is shown that the crystals isolated from a melt of the scarlet variety are identical with Hittorf's phosphorus. Black phosphorus changes into the scarlet form on being heated at 575° . W. T.

The Toxic Properties of Commercial Ferrosilicon. N. KURNAKOV and G. URASOV (*Z. anorg. Chem.*, 1922, **123**, 89—131).—Commercial ferrosilicon has been found to evolve poisonous and explosive gases. The alloys containing about 50% silicon, which is attacked by water, are very dangerous. The authors found that a solid phase of variable composition, *lebeaite*, makes its appearance in iron-silicon alloys with 33.4 to 100% by weight of silicon; *lebeaite* solidified at 1245° . *Lebeaite* can form ternary and quaternary solid solutions with phosphorus and aluminium. The ternary solid solutions (Fe, Si, P, and Fe, Si, Al) of *lebeaite* are stable towards water. The quaternary solution (Fe, Si, Al, P) is, however, attacked by water, phosphine being evolved. Ferrosilicon with less than 33.4% silicon in γ -iron can dissolve phosphorus and aluminium to the extent of 1.7% and 3%, respectively.

These ternary and quaternary solid solutions resist the action of water. Calcium phosphide will not dissolve in solid or liquid ferrosilicon.

W. T.

The Mechanism of the Formation of Silane from Magnesium Silicide. I. ROBERT SCHWARZ and ERICH KONRAD (*Ber.*, 1922, 55, [B], 3242—3252).—The decomposition of magnesium silicide by aqueous hydrochloric acid leads ultimately to the production of a solid, white substance, $\text{H}_2\text{Si}_2\text{O}_3$, which shows all the properties of dioxodisiloxan. The substance, however, cannot be a primary product of the decomposition of the silicide, but is to be regarded as the final result of a series of intermediate reactions which cannot be followed in purely aqueous solution. Attempts are therefore made to arrest the reaction at intermediate stages by replacing the water to a greater or less extent by alcohol. Under these conditions, monosilane is not evolved during the first period of the action, and the solid product contains magnesium, which cannot be removed by washing, but is free from chlorine. It is a colourless substance (the usual brown tint is due to admixed elementary silicon) which explodes on contact with air, yielding silicon and silicon dioxide, is decomposed by water, decolorises potassium permanganate and alcoholic bromine, and reduces silver nitrate. Analysis of it gives the ratio $\text{Si} : \text{Mg} : \text{H} = 1 : 2 : 2$. The primary hydrolysis of magnesium silicide may therefore be expressed by the equation: $\text{Mg}_2\text{Si} \cdot \text{Mg} + 2\text{H} \cdot \text{OH} = \text{H}_2\text{Si}(\text{Mg} \cdot \text{OH})_2$. In these circumstances, not more than 15% of the silicon actually present is found in the solid product, whilst also none escapes as monosilane. The remainder is found as a stable alcohol sol of silicic acid. A second reaction must therefore have occurred, possibly in accordance with the scheme: $\text{Mg}_2\text{Si} + 4\text{HCl} + 2\text{H}_2\text{O} = 4\text{H}_2 + 2\text{MgCl}_2 + \text{SiO}_2$, whereby account is rendered of the evolution of hydrogen. Approximately 10% of the magnesium silicide is decomposed in accordance with the first and about 90% in accordance with the second equation.

Many attempts have been made by varying the concentration of the alcohol and shortening the period of the action of the acid on the silicide to isolate other compounds intermediate between the primary product and the ultimate silicoformic anhydride. With increasing dilution of the alcohol, the production of silanes increases, but very little definite information is afforded by analysis of the solids. They can only be regarded as mixtures formed by further hydrolysis of the primary product, which they resemble in their ready inflammability on exposure to air. It appears probable that the initial product suffers further hydrolysis in two directions such as are suggested by the schemes: $\text{SiH}_2(\text{Mg} \cdot \text{OH})_2 + 2\text{H} \cdot \text{OH} = 2\text{Mg}(\text{OH})_2 + \text{SiH}_4$ and $\text{SiH}_2(\text{Mg} \cdot \text{OH})_2 + \text{H}_2\text{O} + 4\text{HCl} = 2\text{MgCl}_2 + 2\text{H}_2\text{O} + \text{SiH}_2\text{O} + 2\text{H}_2$.

Magnesium silicide is prepared by igniting an intimate mixture of finely-divided, ignited quartz and magnesium powder in an atmosphere of hydrogen and mechanical separation of the crystals of the silicide from the mixture. The crude product is freed from

admixed magnesium by the action of ethyl bromide and anhydrous ether. The purified material is a uniform, steel-blue, crystalline powder which contains traces of amorphous silicon and iron silicide. The apparatus used in studying the decomposition of the silicide is fully figured and described in the original. H. W.

Pressure of Saturated Carbonyl Chloride Vapour. N. I. NIKITIN (*J. Russ. Phys. Chem. Soc.*, 1920, 52, 235—249).—The vapour pressure of carefully purified carbonyl chloride has been measured at a number of temperatures ranging from 12.6° to -99° . The pressure has the value 889.2 mm. at 12.6° , and diminishes rapidly with fall of temperature to 540.4 mm. at -0.41° , and then continually more slowly; at -19.43° , the value is 232.2 mm. and at -94.2° a few tenths of a millimetre. T. H. P.

Interaction of Carbon Tetrachloride and Fuming Sulphuric Acid. A. I. LEFIN (*J. Russ. Phys. Chem. Soc.*, 1920, 52, 1—17).—The author has investigated the mechanism of the formation of carbonyl chloride and chlorosulphonic acid from carbon tetrachloride and fuming sulphuric acid (cf. Schützenberger, *Annalen*, 1870, 154, 375; Armstrong, *J. pr. Chem.*, 1870, [ii], 1, 246; Erdmann, A., 1893, i, 681). If the fuming acid contains 45% of dissolved sulphur trioxide, corresponding with the formula $\text{H}_2\text{S}_2\text{O}_7$, the reaction is expressed by the equation, $\text{CCl}_4 + \text{H}_2\text{SO}_4 \cdot \text{SO}_3 = \text{COCl}_2 + 2\text{SO}_3\text{HCl}$. If less than 45% of dissolved trioxide is present, so that the acid may be regarded as a solution of fuming in monohydrated acid, the fuming acid acts as above, the excess of monohydrate remaining together with the secondary product, chlorosulphonic acid. Finally, if the dissolved sulphur trioxide exceeds 45%, the acid may be considered as a mixture of pyrosulphuric acid with sulphur trioxide. In this case, part of the carbon tetrachloride reacts according to the equation $\text{CCl}_4 + \text{H}_2\text{SO}_4 \cdot \text{SO}_3 = \text{COCl}_2 + 2\text{SO}_3\text{HCl}$, and part according to $\text{CCl}_4 + 2\text{SO}_3 = \text{COCl}_2 + \text{S}_2\text{O}_5\text{Cl}_2$; both these reactions proceed quantitatively. T. H. P.

Reaction between Potassium Azide and Iodine in the Presence of Carbon Disulphide. A. W. BROWNE and A. B. HOEL (*J. Amer. Chem. Soc.*, 1922, 44, 2106—2116).—Potassium azide reacts with iodine in the presence of carbon disulphide, with the formation of potassium iodide and the liberation of nitrogen. The first stage in this reaction consists in the formation of potassium azidodithiocarbonate, $\text{KS} \cdot \text{CS} \cdot \text{N}_3$, by interaction of potassium azide and carbon disulphide, as expressed by the equation $\text{KN}_3 + \text{CS}_2 = \text{KS} \cdot \text{CS} \cdot \text{N}_3$. This reaction is irreversible, consequently there is no regeneration of carbon disulphide after it has once reacted. The second stage consists in the reaction between potassium azidodithiocarbonate with iodine, which results in the precipitation of azidocarbon disulphide $\text{S}_2(\text{CS} \cdot \text{N}_3)_2$ according to the equation $2\text{KS} \cdot \text{CS} \cdot \text{N}_3 + 2\text{I} = \text{S}_2(\text{CS} \cdot \text{N}_3)_2 + 2\text{KI}$. The halogenoid substance azidocarbon disulphide liberates triatomic nitrogen from potassium azide, with resultant evolution of ordinary nitrogen, as shown by the equations $2\text{KN}_3 + \text{S}_2(\text{CS} \cdot \text{N}_3)_2 = 2\text{KS} \cdot \text{CS} \cdot \text{N}_3 + 2\text{N}_3$; $2\text{N}_3 = 3\text{N}_2$.

Combination of these equations with the preceding one leads to the simple expression of the final result obtained when carbon disulphide, potassium azidodithiocarbonate, or azidocarbon disulphide, in relatively small amount, is brought into contact with a solution containing potassium azide and iodine. These results have been confirmed by iodometric and nitrometric determinations: $2\text{KN}_3 + 2\text{I} = 2\text{KI} + 3\text{N}_2$. J. F. S.

Potassium Azidodithiocarbonate. A. W. BROWNE and A. B. HOEL (*J. Amer. Chem. Soc.*, 1922, **44**, 2315—2320; cf. preceding abstract; Sommer, A., 1916, ii, 29).—Potassium azidodithiocarbonate may be prepared by the action of carbon disulphide on an aqueous solution of potassium azide at 40°. It is obtained in colourless, deliquescent crystals, which decompose quantitatively when gently heated, yielding potassium thiocyanate, sulphur, and nitrogen. When rapidly heated, the substance explodes, with the formation of potassium sulphide, carbon dioxide, and sulphur dioxide and trioxide in addition to the above. Solutions of the salt when treated with various oxidising agents or when electrolysed yielded azidocarbon disulphide (SCSN_3)₂. The authors confirm the structure, $\text{K}\cdot\text{S}\cdot\text{CS}\cdot\text{N}:\text{N}:\text{N}$, as assigned to it by Sommer (*loc. cit.*). W. G.

Melting and Freezing Point of Sodium Chloride. JOHN BRIGHT FERGUSON (*J. Physical Chem.*, 1922, **26**, 626—630).—The melting point and freezing point of sodium chloride have been determined by means of a platinum-platinum-rhodium thermometer. As the mean of a number of experiments, the value 803.1° is found for the freezing point and 803.4° for the melting point. The author recommends the value $803^\circ \pm 1^\circ$ as the best value for the freezing point. J. F. S.

The Properties of Ammonium Nitrate. IV. The Reciprocal Salt-pair, Ammonium Nitrate and Sodium Chloride. EDGAR PHILIP PERMAN (*T.*, 1922, **121**, 2473—2483).

Colloidal Calcium Hydroxide. M. VON GLASENAPP (*Kolloid Z.*, 1922, **31**, 195—196).—Two varieties of colloidal calcium hydroxide have been obtained from a marly dolomite of the composition CaO 22.75%, MgO 15.50%, Al_2O_3 7.68%, Fe_2O_3 1.87%, SiO_2 15.27%, CO_2 34.83%, H_2O 1.81%. The dolomite was burnt at 720—780° until the magnesium carbonate was entirely decomposed and the calcium carbonate half decomposed. After the burnt product had been kept for twelve hours in water, the two colloidal varieties of calcium hydroxide were microscopically visible. The primary product separates first from solution as small droplets which grow speedily, and pass into the secondary product which forms agglomerates of drops having the appearance of a cell network. Both colloids are optically isotropic but strongly refracting. In reflected light, the primary colloid is light blue and the secondary bright orange in colour. The secondary colloid does not maintain the colloidal state very long, but passes into crystalline calcium hydroxide. J. F. S.

Valency. XVI. The Ammonia Compounds of Calcium Haloids. GUSTAV F. HÜTTIG (*Z. anorg. Chem.*, 1922, 123, 31—42).—The methods of investigation were as previously communicated by the author (A., 1920, ii, 293 and 318). The following ammonia compounds were found to exist in the stable state:— $\text{CaCl}_2 \cdot 8\text{NH}_3$; $\text{CaCl}_2 \cdot 4\text{NH}_3$; $\text{CaCl}_2 \cdot 2\text{NH}_3$; $\text{CaCl}_2 \cdot \text{NH}_3$; $\text{CaBr}_2 \cdot 8\text{NH}_3$; $\text{CaBr}_2 \cdot 6\text{NH}_3$; $\text{CaBr}_2 \cdot 2\text{NH}_3$; $\text{CaBr}_2 \cdot \text{NH}_3$; $\text{CaI}_2 \cdot 8\text{NH}_3$; $\text{CaI}_2 \cdot 6\text{NH}_3$; $\text{CaI}_2 \cdot 2\text{NH}_3$, and $\text{CaI}_2 \cdot \text{NH}_3$.

These determinations were carried out at various temperatures ranging from -80° to 230° . The isothermal curves pressure-composition are given.

W. T.

The Solubilities of Calcite and Aragonite. HANS L. J. BÄCKSTRÖM (*Medd. K. Vetenskapsakad. Nobel-Inst.*, 1922, 4, No. 11, 1—11).—An account of work published more fully elsewhere (A., 1921, ii, 317).

E. H. R.

Setting and Velocity of Solution of Burnt Gypsum. P. P. BUDNIKOV and JA. K. SYRKIN (*Bull. Inst. Polyt. Ivanovo-Voznesensk.*, 1922, 6, 235—247).—The authors have investigated the velocities of dissolution of gypsum after the latter had been heated at various temperatures, the concentrations being determined at different stages of the dissolution by means of conductivity measurements. For the samples which have been heated at 115° and 125° , the conductivity and hence the solubility increases rapidly to a point corresponding with a solution considerably supersaturated with respect to $\text{CaSO}_4 \cdot \text{H}_2\text{O}$, and afterwards diminishes, at first rapidly and then slowly, but still indicates a certain degree of supersaturation after twenty-four hours. Supersaturated solutions are obtained also with gypsum heated at 400° , but not when a temperature of 500° is reached. In so far as the setting of gypsum is conditioned by crystallisation from supersaturated solutions, the temperature at which "dead-burnt" gypsum is formed may therefore be assumed to be about 450° .

Burnt gypsum may be regarded as consisting of two components, one soluble, which dissolves readily, and the other relatively insoluble. The solubility of samples which have been burnt at temperatures between 500° and 800° is rapid at first, this being explained by the presence of appreciable proportions of the soluble modification in dead-burnt gypsum.

The time occupied by dissolution does not furnish an exhaustive explanation of the disappearance of the ability of gypsum to set, and the authors suggest that the presence of the insoluble component in the interstices of the crystals prevents the latter from growing.

T. H. P.

The Silicates of Strontium and Barium. PENTTI ESKOLA (*Amer. J. Sci.*, 1922, [v], 4, 331—375).—The author has determined the equilibrium diagrams of the binary systems SrO-SiO_2 and BaO-SiO_2 , respectively, and also certain characteristics (crystal-line system and habit, refractive indices and dispersion, cleavage and density) of the compounds and eutectics formed in these

systems. The corresponding diagram and characteristics in the case of the system CaO-SiO_2 are given for purposes of comparison. In the system SrO-SiO_2 , compounds 2SrO, SiO_2 and SrO, SiO_2 were found, and these, together with SrO , occurred in one form only, although the temperature of formation of the silicates was varied from the melting point down to about 900° . SrO, SiO_2 was found to be closely isomorphous and optically very similar to $\alpha\text{CaO, SiO}_2$. Its crystals are apparently hemimorphic and belong either to the dihexagonal pyramidal or monoclinic domatic class, probably the latter. In the system BaO-SiO_2 , compounds 2BaO, SiO_2 , BaO, SiO_2 , 2BaO, 3SiO_2 , and BaO, 2SiO_2 were found. Of these, 2BaO, 3SiO_2 and BaO, 2SiO_2 were found to be isomorphous, of orthorhombic symmetry, forming a complete series of solid solutions. The melting-point diagram of this series is of Roozeboom's type I, without maximum or minimum, and the index of refraction varies continuously, but not linearly with the composition. Barium metasilicate, BaO, SiO_2 , is not isomorphous with the calcium and strontium meta-silicates. It is optically biaxial, probably of orthorhombic symmetry, and forms no solid solutions with $\alpha\text{CaO, SiO}_2$, but a double compound, 2CaO, BaO, 3SiO_2 is formed, which decomposes on heating into $\alpha\text{CaO, SiO}_2$ and liquid. Crystals of the double compound are uniaxial and negative, are probably hexagonal, and show good cleavages in their prismatic zone. Neither strontium nor barium metasilicate forms with magnesium metasilicate a double compound analogous to diopside. Felspars analogous to anorthite are formed by both strontium and barium. Strontium felspar and anorthite resemble one another very closely in optical properties, and the two appear to be completely miscible. Barium felspar is monoclinic and readily forms Carlsbad twins of the contact type.

J. S. G. T.

The System Ammonium Sulphate-Glucinum Sulphate-Water at 25° . HUBERT THOMAS STANLEY BRITTON (T., 1922, 121, 2612—2616).

Preparation and Properties of Magnesium Perchlorate and its Use as a Drying Agent. H. H. WILLARD and G. FREDERICK SMITH (*J. Amer. Chem. Soc.*, 1922, 44, 2255—2259).—The hexahydrate of magnesium perchlorate may be prepared by dissolving magnesia in perchloric acid, evaporating the solution until fumes of perchloric acid are evolved and crystallisation commences at the surface. The mass is then cooled to the ordinary temperature with the addition of sufficient water to keep the mass semi-fluid. The crystals are separated by centrifuging, redissolved in water, and crystallised. The hexahydrate, m. p. $145-147^\circ$, d_4^{25} 1.970, is now deliquescent. The trihydrate is prepared by keeping the hexahydrate over phosphoric oxide at $20-25^\circ$ for a month. The product has m. p. $145-147^\circ$, d_4^{25} 2.044. The anhydrous compound is obtained by heating either hydrate at 170° in a current of dry air for some time and then at 250° for a short time. Magnesium perchlorate decomposes at 400° without melting, forming a mixture of the oxide and chloride. When moist gas is

passed over anhydrous magnesium perchlorate at a rate not greater than 5 litres per hour, it is found to be as efficient a desiccating agent as phosphoric oxide. The trihydrate is as efficient as the anhydrous salt at 0° , but much less so at higher temperatures. J. F. S.

Crystal Structure of Cadmium Iodide. RICHARD M. BOZORTH (*J. Amer. Chem. Soc.*, 1922, 44, 2232—2236).—The crystal structure of cadmium iodide has been investigated by means of X-ray spectra and Laue photographs. It is shown that cadmium iodide is not hexagonal, as has generally been supposed, but trigonal. The unit parallelepiped contains one molecule, and the shortest distance between the centres of the cadmium and iodine atoms is 3.00 \AA.U. , which is exactly the sum of the radii of these atoms as found by Bragg (A., 1920, ii, 537). Because of the nature of the structure, the iodine atoms are also "in contact" on the Bragg hypothesis. The shortest distance between their centres is 4.21 \AA.U. , whilst Bragg's value of the diameter of this atom is 2.80 \AA.U. This separation indicates that the atoms which are presumably similarly charged may be expected to be farther apart than Bragg's hypothesis requires. The structure may be divided into electrically neutral layers by passing planes parallel to the base midway between the cadmium atoms, and this may well account for the excellent basal cleavage of the crystal. J. F. S.

The Ammoniates of Lead Haloids. Mixed Compounds and the Parent Salts. WILHELM BILTZ and WILHELM FISCHER (*Z. anorg. Chem.*, 1922, 124, 230—247).—The following ammoniates of the haloids of lead are described in literature: $\text{PbCl}_2 \cdot 1.5\text{NH}_3$; $\text{PbCl}_2 \cdot 2\text{NH}_3$; $\text{PbBr}_2 \cdot 2\text{NH}_3$; $\text{PbI}_2 \cdot \text{NH}_3$; $\text{PbI}_2 \cdot 2\text{NH}_3$; and $\text{PbI}_2 \cdot 4\text{NH}_3$. The authors were able to show also the existence of the following: $\text{PbCl}_2 \cdot \text{NH}_3$; $\text{PbCl}_2 \cdot 3.25\text{NH}_3$; $\text{PbBr}_2 \cdot \text{NH}_3$; $\text{PbBr}_2 \cdot 3\text{NH}_3$; $\text{PbBr}_2 \cdot 5.5\text{NH}_3$; $\text{PbI}_2 \cdot 0.5\text{NH}_3$; and $\text{PbI}_2 \cdot 5\text{NH}_3$. The method of investigation was the same as that employed in previous work (cf. Hüttig, A., 1920, ii, 293, 318; this vol., ii, 849). The authors regard the compound $\text{PbCl}_2 \cdot 3.25\text{NH}_3$ or $4\text{PbCl}_2 \cdot 13\text{NH}_3$ as a mixed compound of the salts $\text{PbCl}_2 \cdot 8\text{NH}_3$, $\text{PbCl}_2 \cdot 2\text{NH}_3$, and $2\text{PbCl}_2 \cdot 3\text{NH}_3$. Similarly, $2\text{PbBr}_2 \cdot 11\text{NH}_3$ is a compound of $\text{PbBr}_2 \cdot 8\text{NH}_3$ and $\text{PbBr}_2 \cdot 3\text{NH}_3$, and $2\text{PbI}_2 \cdot 10\text{NH}_3$ a compound of $\text{PbI}_2 \cdot 8\text{NH}_3$ and $\text{PbI}_2 \cdot 2\text{NH}_3$. These mixed compounds are not characterised by great stability or by instability; methods of detecting their formation are desirable. W. T.

Action of Ammonium Nitrate and of Aqueous Ammonia on Copper. Properties of Cupric Tetrammine Nitrite and Nitrate. HENRY BASSETT and REGINALD GRAHAM DURRANT (*T.*, 1922, 121, 2630—2640).

The Solubility of Copper Hydroxide in Sodium Hydroxide Solutions. G. SVENSEN MELBYE (*Medd. K. Vetenskapsakad. Nobel-Inst.*, 1922, 4, No. 8, 1—11).—The solubility of copper hydroxide in sodium hydroxide solutions between 2.65 and $6.6N$ increases at a linear rate with the concentration of sodium hydroxide according to the equation $[\text{Cu}(\text{OH})_2] = 0.045[\text{NaOH}] - 0.095$.

present authors carried out a systematic study of the ternary system iron-boron-carbon and their results are given in detail. Equilibrium sets in so slowly that the structure and properties of the iron-boron-carbon alloys depend to a large extent on the preliminary heating. The limit of the influence of boron on the structure was found to be 0.02%.

W. T.

The Constitution of Erdmann's Salt and its Derivatives.

E. H. RIESENFELD and R. KLEMENT (*Z. anorg. Chem.*, 1922, **124**, 1—21).—The authors claim that Erdmann's salt, $[(\text{NO}_2)_4\text{Co}(\text{NH}_3)_2]\text{R}$, has the two ammonia molecules in the *cis*-position. They base their conclusions on the fact that (1) by the action of ethylenediamine the trinitro-ethylenediamine-monammine derivative is obtained, $[(\text{NO}_2)_3\text{Co en NH}_3]\text{R}$; (2) the dioxalato-diammine derivative, $[(\text{C}_2\text{O}_4)_2\text{Co}(\text{NH}_3)_2]\text{R}$, prepared from Erdmann's salt, could not be resolved into optical antipodes; (3) the non-resolvability again of the monoxalato-derivative, $[\text{C}_2\text{O}_4(\text{NO}_2)_2\text{Co}(\text{NH}_3)_2]$; (4) the ease with which the dichloro-derivative was prepared, $[(\text{NO}_2)_2\text{Cl}_2\text{Co}(\text{NH}_3)_2]$.

The oxalate group could not be replaced by the malonate radicle on account of the solubility and reducibility of the malonate. Methods of preparing the above derivatives are given.

W. T.

Preparation of Colloidal Solutions of Nickel and Cobalt Hydroxides and some other Compounds of these Metals.

O. F. TOWER and MARTHA C. COOKE (*J. Physical Chem.*, 1922, **26**, 728—735).—Two methods are described for preparing colloidal suspensions of nickel hydroxide, one by dialysing a solution of nickel tartrate in the presence of an alkaline solution of potassium tartrate, and the other by treating a *N*/10-solution of nickel chloride with a little more than the equivalent quantity of a solution of potassium hydroxide, allowing the precipitated nickel hydroxide to settle, siphoning off the supernatant liquid, adding water, and repeating the process several times. Only very weak suspensions of cobaltous hydroxide could be obtained by these methods. Any solids obtained from the solutions referred to in a previous work on this subject (*A.*, 1900, i, 587) are composed of nickel tartrate with potassium tartrate adsorbed in it. Glycerol does not prevent the precipitation of nickel hydroxide from aqueous solutions of nickel salts. In alcoholic solution, interesting transformations occur in the gel on keeping, and finally a solution is obtained from which nickel hydroxide cannot be precipitated by dilution with water. Colloidal solutions of nickel sulphide, which are readily formed in the presence of an alkaline solution of a tartrate, immediately decompose with precipitation of the sulphide on dialysing out the tartrate.

J. S. G. T.

Hydrated Oxides. IV. HARRY B. WEISER (*J. Physical Chem.*, 1922, **26**, 654—686; cf. this vol., ii, 575).—A continuation of previous work in connexion with the oxides of tin. It is shown that the so-called stannic and metastannic acids are not acids, but are hydrated stannic oxides the composition and properties of which are

determined by the conditions of formation. Stannic oxide, freshly prepared by precipitation from a solution of a stannic salt at the ordinary temperature, is a loose, highly hydrated mass which is readily peptised by dilute acids and alkalis, forming a colloidal solution, and is dissolved by concentrated acids and alkalis, forming salts. The oxide is more compact, less hydrated, less soluble, and less easily peptised the higher the temperature of formation. Freshly-prepared stannic oxide undergoes these changes on keeping at the ordinary temperature. Experiments on the peptisation by nitric acid of oxides prepared under different conditions show that the different products are not mixtures of varying amounts of a definite α -oxide peptisable by nitric acid with a definite β -isomeride which is not peptisable by nitric acid. Hydrated stannic oxides prepared at different temperatures or aged for varying lengths of time are chemical individuals which differ from each other in the size of the particles and the structure of the mass, and hence in their adsorbability and mordanting action, their ease of peptisation, and their solubility. There are no definite hydrates of stannic oxide. Salts of the general formulæ SnR_4 and M_2SnO_3 are formed by the action of concentrated acids and alkalis on newly-formed hydrated stannic oxide. Numerous oxy-salts and complex stannates and metastannates have been prepared by the action of acids and alkalis on different hydrated stannic oxides. These are not definite compounds, but are adsorption products, the composition of which is determined by the age of the hydrated oxide and the concentration of acid or base with which they were treated. Colloidal solutions of hydrated stannic oxide can be prepared by peptisation of the freshly-formed oxide with dilute acids or alkalis. The colloid ages slowly on keeping and rapidly on heating. Any number of colloidal solutions are possible, each differing from the others in the size of the hydrated particles and hence in their reactivity, adsorbability, and stability under given conditions. Hydrated stannic oxide adsorbs colloidal gold, silver, and platinum, forming purple masses, the most common of which is the gold compound, purple of Cassius. All these purple substances are colloidal in character, the composition varying with the conditions of formation. When freshly prepared, they are readily peptised by dilute acids and alkalis, but when aged by drying they are peptised but slightly even by concentrated reagents. A mixture of the hydrated oxides of iron and tin in certain proportions is soluble in dilute ammonia. The reason for this is that hydrated stannic oxide is peptised by the hydroxyl-ion, whilst ferric oxide is not, but colloidal stannic oxide adsorbs ferric oxide and carries it into colloidal solution as long as tin is present in excess. At the same time, hydrated ferric oxide adsorbs stannic oxide and tends to take it out of colloidal solution, so that when the former is present in large excess none of the latter remains peptised. Stannic oxide does not precipitate in the usual way from a solution of tin in nitric acid containing a suitable amount of iron. The reason for this is that hydrated stannic oxide peptised by nitric acid coagulates spontaneously, since the aged oxide is neither peptised nor dis-

solved by this acid. Ferric nitrate peptises the oxide both when newly formed and when aged. Hence if freshly-prepared stannic oxide is peptised either by ferric nitrate or by a suitable mixture of ferric nitrate and nitric acid, coagulation does not take place on keeping or boiling on account of the stabilising action of the strongly adsorbed ferric ion; but if the concentration of ferric ion in the nitric acid solution is too low, complete or partial coagulation takes place on keeping or boiling. Stannic oxide jellies are prepared by coagulation of colloidal solutions of the oxide with suitable concentrations of electrolytes. The results of the present investigation support the author's general hypothesis of the formation of jellies.

J. F. S.

The Occurrence of Germanium in Cassiterite. ASSAR HADDING (*Z. anorg. Chem.*, 1922 **123**, 171—172).—The author examined cassiterite from Finbo by means of the X-ray spectrograph and found it to contain the following elements: tin, tantalum, iron, manganese, germanium, and zinc. The intensity of the lines shows that the amount of germanium was less than that of manganese and more than that of zinc. A similar analysis of cassiterite from Mamacka gave like results. No germanium was indicated in cassiterite from Finland and Bohemia. W. T.

Germanium Hydride. RUDOLF SCHENCK [with ALBERT IMKER] (*Rec. trav. chim.*, 1922, **41**, 569—575).—The preparation of an alloy of germanium and magnesium could not be effected by the action of magnesium on germanium oxide, as the reaction is too violent. The germanium was obtained as oxide from argyrodite; this was purified by conversion into germanium silicofluoride, thence into sulphide, and finally into oxide. Reduction to the metal was then effected by heating in a stream of hydrogen, and an alloy was prepared by heating one part of the powdered germanium so obtained with two parts of magnesium powder in an atmosphere of hydrogen. The hydride, prepared by the action of hydrochloric acid on the alloy in a stream of hydrogen, was obtained pure by cooling to the temperature of liquid air. Small quantities of germanium chloroform were produced simultaneously with the hydride and condensed in a freezing mixture in the preliminary stages of cooling. The uncondensed gases were passed through silver nitrate solution and again through a tube cooled in liquid air, but were not completely freed from the hydride, which could still be detected by its nauseating odour and by the deposition of metallic germanium on porcelain when the gas was burnt. The hydride condensed to a white solid which was purified by sublimation, m. p. -165° , b. p. $-126^{\circ}/757$ mm. Values for the vapour pressure from -164° to -125° are given. Three determinations of molecular weight by Regnault's method gave a mean value of 76.93, the theoretical value being 76.50. The gas readily decomposes into its elements on heating (cf. Paneth and Schmidt-Hebbel, this vol., ii, 776), and an analysis was carried out by passing it through a heated tube, the hydrogen formed being estimated in a eudiometer and the germanium weighed


as such. Three determinations gave 94.70% Ge and 5.29% H as mean values, the calculated percentages for GeH_4 being 94.73 and 5.27. The volume ratio of germanium hydride to the hydrogen obtained by its decomposition was found to be 1:2.05. The general reactions of the gas show that it is endothermic and analogous in properties to the hydrides of arsenic, antimony, tin, and bismuth (cf. Voegelen, A., 1902, ii, 401). H. J. E.

Sulphide Sols. III. Sol Preparation by Hydrolysis. F. V. VON HAHN (*Kolloid Z.*, 1922, **31**, 200—203; cf. A., 1921, ii, 577).—Fairly stable antimony pentasulphide sols may be prepared by the hydrolysis of the alkali salts of stibiothiosulphuric acid. The hydrolysis takes place extremely readily according to the equation $2\text{Na}_3\text{Sb}(\text{S}_2\text{O}_3)_3 = \text{Sb}_2\text{S}_5 + 3\text{Na}_2\text{SO}_4 + 3\text{SO}_2 + \text{S}$ (Szilágyi, A., 1920, ii, 207). The preparation is carried out as follows: a 0.5% solution of sodium antimony thiosulphate is kept until it commences to turn yellow and to show a definite Tyndall cone; it is then placed in a dialyser and washed with warm distilled water. After several hours' dialysis, the solution has taken on a deep red colour. The main point to be observed in the preparation is the rapid removal of the alkali sulphate. In the presence of sodium chloride the sol cannot be prepared, nor can a sol having a greater molecular concentration than 0.4 be obtained. Sols of one-tenth this concentration are stable for a few hours only. The sols prepared from the sodium salt are more stable than those prepared from either lithium or potassium antimony thiosulphate. Attempts to prepare arsenic pentasulphide sols by the hydrolysis of alkali arsenic thiosulphates failed entirely owing to the greater instability of these compounds. J. F. S.

Complex Platinum Compounds. III. Molecular Rearrangements observed with Complex Platinum Compounds. L. A. TSCHUGAEV and N. K. PSCHENICYN (*J. Russ. Phys. Chem. Soc.*, 1920, **52**, 47—62).—The depolymerisation of bicomplex salts of the type $[\text{Pt}4A]\text{PtX}_4$ (where A = a molecule of ammonia, amine, organic sulphide, etc.), resulting in the formation of the monomeric type $[\text{Pt}2A, \text{X}_2]$ and in the accompanying transference of one-half of the total A molecules from one platinum atom to another is frequently encountered among complex platinum compounds, but has not previously been observed with the simplest purely inorganic, in particular with the ammino-, complexes of platinum. This gap has now been filled by the authors, who show that tetra-aminoplatinous platinochloride (Magnus's green salt) is capable of undergoing depolymerisation in accordance with the scheme $[\text{Pt}4\text{NH}_3]\text{PtCl}_4 = [\text{Pt}2\text{NH}_3, \text{Cl}_2]$, the chloride of Reiset's Base II being thus formed. The reaction is effected by cautious heating over a naked flame to 270° , and never proceeds to completion, part of the green salt decomposing with separation of metallic platinum.

Experiments have been made also with certain heterometallic bicomplex compounds in which the platinum occurs only in the anion, $[\text{PtX}]_4$, whilst the kation is formed of another metal in com-

bination with a known number of A molecules giving the so-called neutral part of the complex : for example, $[M, nA]PtX_4$. If as the component M is taken an element such as silver, zinc, copper, mercury, etc., which, according to all the available data, holds molecules of ammonia, etc., less stably than does platinum, the transference of these molecules from the metal M to the platinum should take place with great readiness. With the aminiosilver platinichlorides, the transference of ammonia from the silver to the platinum atom occurs at comparatively low temperatures : At 60—70°, monoamminosilver platinichloride yields, together with silver chloride, also (a) tetra-amminoplatinous platinichloride (Magnus's green salt), (b) chlorotriamminoplatinous platinichloride (Cleve's salt), and (c) *cis*dichlorodiamminoplatinum (Peyrone's chloride)




whilst at a higher temperature (100°) the last two compounds disappear and are replaced by *trans*-dichlorodiamminoplatinum (chloride of Reiset's Base II); in either case, the loss of ammonia is negligible. At all temperatures between 100° and 200°, di-amminosilver platinichloride yields a new product, namely, tetra-amminoplatinous chloride (chloride of Reiset's Base I), the loss of free ammonia being considerable. At 100°, the only compound, besides silver chloride, accompanying the chloride of Reiset's Base I is Magnus's salt; at 150°, the chloride of Reiset's Base II also appears, whilst at 200° Magnus's salt disappears and the chlorides of Reiset's Bases I and II form the sole products. The Base II chloride, which appears in all cases at the highest temperatures, is evidently a secondary product formed from Magnus's salt.

When heated at 50—60°, diethylamminosilver platinichloride readily undergoes transformation into the analogues of the chloride of Reiset's Base I and Magnus's green salt, part of the ethylamine being liberated : $[Ag2NH_2Et]_2PtCl_4 = 2AgCl + [Pt4NH_2Et]Cl_4$ and $2[Ag2NH_2Et]_2PtCl_4 = 4AgCl + [Pt4NH_2Et]PtCl_4 + 4NH_2Et$. Under the same conditions, monoethylamminosilver platinichloride gives the analogues of Magnus's green salt and Peyrone's chloride, part of the original compound also decomposing with complete loss of the amine and formation of silver platinichloride.

Tetramminozinc platinichloride, $[Zn4NH_3]PtCl_4$, gives, at 111°, Peyrone's chloride, Magnus's green salt, and the chloride of Reiset's Base I.

T. H. P.

The Alkali-Ruthenium Double Sulphites. HEINRICH REMY (*Z. anorg. Chem.*, 1922, 124, 248—274).—The following definite double sulphites of ruthenium and the alkali metals were prepared,



There were also obtained substances which on analysis did not correspond with any simple formula, probably being mixtures of the above. The properties of the above double sulphites are given. The author discusses the difficulties of analysis. W. T.

Mineralogical Chemistry.

The Examination of Naturally Occurring Gases. II. F. HENRICH [with G. PRELL] (*Ber.*, 1922, 55, [B], 3021—3025).—The method for the collection and examination of naturally occurring gases has been simplified and extended (cf. A., 1920, ii, 767). The gases from a number of springs have been analysed, since such data appear to the author to form the most trustworthy basis for a judgment as to the atmospheric or terrestrial origin of the water. Re-examination of the gas from a cold spring near Leupoldsdorf, Bavaria, confirms the previous analysis (Henrich, *loc. cit.*), showing it to contain oxygen 13%; nitrogen and the rare gases being the only other constituents. The gas from a second spring situated about a hundred metres above the foregoing was also composed of the constituents of air, but with only 18.8% of oxygen. The gas from a third spring was found to contain oxygen (18.2%), nitrogen (81.8%), argon, and neon (1.1%). Since the ratio of nitrogen to rare gas in these gases is approximately the same as in air, it appears that the latter in some unexplained manner has got below the water in the spring. The low oxygen content of the gases in comparison with that of air is explained by the greater solubility of oxygen than of nitrogen in water, so that a gas richer in nitrogen is evolved. The extent to which the air is depleted of its oxygen appears to depend on the pressure of the water in the spring. H. W.

The Examination of Naturally Occurring Gases. III. F. HENRICH [with G. PRELL] (*Ber.*, 1922, 55, [B], 3026—3030).—Analyses are given of the gases from a number of pools on the Luisenburg near Wunsiedel and Alexandersbad (Fichtelgebirge) of which one may be quoted

H₂S (traces), CO₂ 3.1%, O₂ 1.7%, CH₄, 53.9%, N₂+rare gases 41.3%. Methane, carbon dioxide, and the small amounts of hydrogen sulphide appear to owe their origin to a bacterium which is not completely identical with *Bacillus methanigenes*. Nitrogen and the rare gases must be derived originally from the air. H. W.

Dopplerite. Studies in the Composition of Coal. FREDERICK VINCENT TIDESWELL and RICHARD VERNON WHEELER (*T.*, 1922, 121, 2345—2362).

Massive Troilite from Del Norte Co., California. ARTHUR S. EAKLE (*Amer. Min.*, 1922, 7, 77—80).—The bronze-tarnished

mineral (light greyish-brown when untarnished) resembles pyrrhotite, but differs from this in being non-magnetic and in being readily soluble in dilute sulphuric acid. Analysis of the soluble portion gave Fe 62.70, S 35.40% agreeing with the formula FeS ; d 4.67. The mineral is thus identical with the meteoric troilite. It occurs with chalcopyrite and magnetite in serpentine, and has probably resulted by the action of hydrogen sulphide solutions on the magnetite. Troilite and pyrrhotite are regarded as distinct species, and the suggestion that pyrrhotite is $\text{FeS} + \text{S}$ in solid solution (A., 1912, ii, 354) is not accepted. The magnetic character of pyrrhotite suggests the formula Fe_3S_4 for the end-member of the series $\text{Fe}_n\text{S}_{n+1}$. L. J. S.

Keeleyite, a new Lead Sulphantimonite from Bolivia, SAMUEL G. GORDON (*Proc. Acad. Nat. Sci. Philadelphia*, 1922, 74, 101—103).—This occurs as radiating aggregates of acicular (perhaps orthorhombic) crystals with pyrites and quartz in the San José mine at Oruro. It is dark grey with bright metallic lustre and gives a greyish-black streak; d 5.21, H 2. Analysis by J. E. WHITFIELD gave:

Pb.	Sb.	Cu.	Fe.	S.	Quartz.	Total.
25.80	43.46	2.25	2.77	24.54	(1.18)	100.00

Deducting 8.15% of admixed sulphides, this gives the formula $2\text{PbS}_3.3\text{Sb}_2\text{S}_3$, corresponding with rezbanyite ($2\text{PbS}_3.3\text{Bi}_2\text{S}_3$). The compositions of the known lead sulphantimonites are represented on a triangular diagram. L. J. S.

An Andorite-bearing Silver Ore from Nevada. EARL V. SHANNON (*Proc. U.S. Nat. Museum*, 1922, 60, art. 16, 1—5).—An ore specimen from Keyser mine, Nye Co., Nevada, shows rough prisms of andorite with pyrite and sphalerite in a gangue of rhodochrosite and quartz. Analysis I of the andorite gives, after deducting zinc and iron as sphalerite, the formula $\text{PbAgSb}_3\text{S}_8$. The crystals are usually altered and filled in a central cavity with a matted mass of fine fibres. Analysis II of this alteration product suggests the formula $\text{Pb}_3\text{AgSb}_3\text{S}_8$, and shows that there has been a removal of silver and antimony sulphides.

	Pb.	Ag.	Fe.	Zn.	Sb.	S.	Total.
I	23.35	12.09	1.55	3.56	37.64	22.63	100.82
II	45.14	7.78	2.72	1.56	23.22	[19.58]	100.00

The pale pinkish-buff rhodochrosite gave:

MnO.	FeO.	CaO.	MgO.	CO ₂ .	Total.	MnCO ₃ .
49.49	7.68	3.13	0.93	[38.87]	100.10	80.15

L. J. S.

Sodium Carbonate Minerals from Lake Magadi, Kenya Colony. P. WALTHER (*Amer. Min.*, 1922, 7, 86—88).—The extensive trona deposit of Lake Magadi consists of an aggregate

of pale-yellow, bladed crystals up to 9 cm. in length. Analysis of an average sample with *d* 2.14 gave :

Na_2CO_3 .	NaHCO_3 .	H_2O .	NaCl .	Fe_2O_3 , Al_2O_3 .	SiO_2 .	$\text{CaO}, \text{MgO}, \text{SO}_3$.	Total.
43.55	40.41	15.55	0.36	0.04	0.07	traces	99.98

agreeing with the formula $\text{Na}_2\text{CO}_3, \text{NaHCO}_3, 2\text{H}_2\text{O}$. The mineral has evidently been formed by the action of carbon dioxide and water on sodium silicates, since the rocks in the neighbourhood show extensive bands of flinty silica and carbon dioxide issues from numerous cracks at the surface. The salt crusts of Little Magadi, a dry salt pan 25 miles to the south, consist of harder, bladed (monoclinic ?) crystals which on analysis proved to be pure sodium hydrogen carbonate. This, owing to ready loss of carbon dioxide and conversion into trona, has not hitherto been recognised as a mineral. It here owes its existence to the presence of a large excess of carbon dioxide, which issues from cracks in the soil and rocks with a hissing sound. Optical examination by E. T. Wherry detected only trona, but the material had evidently been altered in the meantime. A new mineral name for sodium hydrogen carbonate is therefore avoided for the present. L. J. S.

Constitution of Thaumassite. EDW. F. HOLDEN (*Amer. Min.*, 1922, 7, 12—14).—Calculations by the Lorentz-Lorenz formula of the molecular refractivity of thaumassite, directly and from its components, point to the presence of 2OH (rather than 4OH) in the formula $\text{CaCO}_3, \text{CaSO}_4, \text{CaSiO}_3, 15\text{H}_2\text{O}$. L. J. S.

Minerals from Ljubija, Bosnia. RUDOLF KOEHLIN (*Tsch. Min. Mitt.*, 1921, 35, 1—12).—The iron-ore at Ljubija, near Prijedor, consists of crystalline-granular chalybite with impregnations of galena, pyrites, and chalcopyrite. Near the surface, the chalybite is altered to limonite, with which are various other secondary minerals. These, including the rarer species leadhillite, pyrochroite, and beudantite, are described. L. J. S.

A New Mineral which Contains the Rare Earths as its Main Component. F. HENRICH [with G. HILLER] (*Ber.*, 1922, 55, [B], 3013—3021).—*Weinschenkite* occurs in very small quantity in the Bavarian Oberpfalz in white, matted, globular deposits and in radiating needles on brown hematite from which it is readily detached. It appears to be mainly a hydrated phosphate of yttrium and erbium, $\text{PO}_4(\text{Er}, \text{Y}), 2\text{H}_2\text{O}$; but indications of the presence of other rare earths have been obtained. The small amount available was insufficient for a complete analysis.

Pseudowavellite, also occurring in minute amount in the same locality, resembles wavellite in that it contains aluminium as its main component, with calcium oxide (13%), barium oxide, and strontium oxide (about 1%) and rare earths (2—3%). Erbium and yttrium appear to be present in addition, possibly, to the cerium earths. H. W.

Supplementary Note on Ishikawaite, a New Mineral from Ishikawa, Iwaki Province. YŪJI SHIBATA and KENJIRO KIMURA (*J. Chem. Soc. Japan*, 1922, 43, 648—649).—The unnamed mineral from Ishikawa, of which an analysis was given (this vol., ii, 517), is now termed *ishikawaite*; the crystals are rhombic [$a : b : c = 0.9451 : 1 : 1.147$]. K. K.

Glaucosite from Lewes, Sussex : Constitution of Glaucosite. A. F. HALLIMOND [with analysis by E. G. RADLEY] (*Min. Mag.*, 1922, 19, 330—333).—A boring through the Gault at Iford Manor, near Lewes, struck the Greensand at a depth of 325 feet. The upper bed of this consists almost entirely of small grains of dark-green glaucosite. The grains consist of a felted mass of minute, birefringent flakes with mean n 1.62; d 2.70. Analysis gave :

SiO ₂ .	Al ₂ O ₃ .	Fe ₂ O ₃ .	FeO.	CaO.	MgO.
48.12	9.16	19.10	3.47	0.76	2.36
K ₂ O.	Na ₂ O.	H ₂ O at 105°.	H ₂ O >105°.	Total.	
7.08	0.22	4.78	5.28	100.33	

From this and a selection of earlier analyses, the formula is deduced as $R_2O_4(R_2O_3, RO), 10SiO_2, nH_2O$, in which $Al_2O_3 + Fe_2O_3$ and $FeO + MgO$ are regarded as mutually replaceable. In some cases, there is a considerable replacement of potassium by sodium, and the variety soda-glaucosite is suggested. L. J. S.

Structure and Composition of the Strathmore (Perthshire) Meteorite. W. F. P. McLINTOCK and F. R. ENNOS (*Min. Mag.*, 1922, 19, 323—329).—Four stones were observed to fall along a track six miles in length on December 3, 1917; the largest weighed 9932 grams and the total weight was 13,255 grams. The structure is that of the intermediate chondrite group; d 3.53. The bulk-analysis, deduced from analyses of the attracted and unattracted portions, is :

Fe.	Ni.	Co.	Fe.	S.	SiO ₂ .	TiO ₂ .	Al ₂ O ₃ .	Cr ₂ O ₃ .	Fe ₂ O ₃ .
7.31	1.26	0.05	4.00	2.29	40.32	0.14	2.57	0.44	0.35
FeO.	MnO.	CaO.	MgO.	K ₂ O.	Na ₂ O.	H ₂ O.	P ₂ O ₅ .	Cl.	Total.
12.99	0.29	1.84	24.96	0.11	0.90	0.17	0.23	0.03	100.25

From this is calculated the following mineral composition : feldspar 10.93, ilmenite 0.27, chromite 0.65, magnetite 0.51, chlorapatite 0.56, olivine 41.13, bronzite 30.91, troilite 6.29, nickel-iron 8.62, water 0.17%. L. J. S.

Analytical Chemistry.

A Microcolorimeter and Nephelometer. A. BAUDOUIN and H. BÉNARD (*Compt. rend. Soc. biol.*, 1920, 83, 602—603).—A description of a miniature Duboscq colorimeter, of which the cups have a capacity of 2 c.c., and the tube an adjustment of 20 mm. The apparatus can also be used as a nephelometer or ultraphotometer, and, by replacing the ocular with a small, direct-vision spectroscope, can be converted into a spectroscope comparator.

CHEMICAL ABSTRACTS.

Colorimeter. ED. MOREAU and A. BONIS (*Ann. Falsif.*, 1922, 15, 357—360).—The apparatus consists essentially of two graduated cylinders placed side by side in a suitable frame; one of these contains the test solution and the other the standard or comparison solution. The latter is admitted through a side tube at the bottom of the cylinder, this tube being connected with a tapped funnel or reservoir containing the solution.

W. P. S.

The Application of Coloured Glasses instead of Liquids in Colorimetric Researches. KLAS SONIÉN (*Arkiv Kem. Min. Geol.*, 1922, 8 No. 7, 1—10).—Combinations of coloured glasses are used as standards in colorimetry. Coloured glasses suitable for this purpose may be prepared by dyeing the gelatin films on photographic plates from which the silver has been removed by treatment with thiosulphate.

W. O. K.

A New Apparatus for Exact Gas Analysis. K. A. SCHALLER and W. BERNDT (*Chem. Ztg.*, 1922, 46, 972—973).—The apparatus consists of a manometer, a measuring tube immersed in a water-jacket with air agitation, and a detachable absorption vessel with an adjustable mercury reservoir. The passage of the mercury from the manometer into the measuring tube is prevented by placing between them a U-tube containing at its lowest point either a constriction or a throttle cock. All the operations of gas analysis can be carried out in this apparatus except explosions, for which the absorption vessel is replaced by a eudiometer. The procedure for estimating carbon dioxide, heavy hydrocarbons, oxygen, carbon monoxide, and the combustion of hydrogen, methane, and ethane is described.

H. C. R.

A Modified Methyl-orange Indicator. KENNETH CLAUDE DEVEREUX HICKMAN and REGINALD PATRICK LINSTAD (T., 1922, 121, 2502—2506).

Apparatus for Electrometric Titration. W. E. GARNER and C. A. WATERS (*J. Soc. Chem. Ind.*, 1922, 41, 337—338t).—The electrode vessel, which carries a small bulb, is attached to a rotating stirrer; two side tubes on the vessel are turned downwards and constricted at the ends. The bulb is convenient in the preparation of a calomel electrode. The side tubes are packed with asbestos, the

vessel is filled completely with electrolyte, and it is supported from the glass tube of the stirrer by means of a waxed cork. The metal used as an electrode is fused or waxed into a thin glass tube and attached by means of a small binding screw to the rotating spindle, electrical connexion being made through a small steel mercury cup to another binding screw. The outer electrode is of any convenient form; the turned-down side tubes on the electrode vessel make it possible to commence a titration with 5 c.c. of solution in the beaker.

W. P. S.

Quantitative Analysis by Measurement of the Degree of Supersaturation. ERNST FRITZ HÖPPLER (*Chem. Ztg.*, 1922, 46, 957—958).—The time elapsing between the addition of a reagent and the formation of a precipitate or coloration can be utilised as an approximate quantitative measure of traces of substances in solution. The method is particularly applicable to water analysis. Sulphuric acid and calcium can be estimated in this way between the dilutions of 2.3 and 100 mg. per litre. Nitrous acid can also be estimated by this means when not in greater concentration than 0.5 mg. per litre.

H. C. R.

The Rapid Estimation of Chlorine in Organic Compounds. E. VOROČEK (*Chem. Listy*, 1922, 16, 248—249).—The author combines the method of Marcussen and Dröscher (A., 1910, ii, 543) with his titration of chlorides against a mercuric salt in the presence of sodium nitroprusside as an indicator. The present author carries out the combustion in a funnel-shaped vessel of several litres capacity fitted with a stopcock. Since sufficient hydrogen is present in the molecule to convert all the chlorine into hydrogen chloride, the products of combustion are absorbed by water and titrated directly against mercuric nitrate. The estimation takes about an hour and a half.

W. T.

Potassium Dichromate as a Standard in Iodimetry and the Estimation of Chromates by the Iodide Method. WARREN C. VOSBURGH (*J. Amer. Chem. Soc.*, 1922, 44, 2120—2130).—It is shown that titration of iodine with thiosulphate in dilute acid solutions gives results which are 0.1—0.3% higher, and more concordant than the results of titration in a neutral solution. The presence of air affects the titration of iodine with thiosulphate when the acidity of the solution is equivalent to 0.3*N*-hydrochloric acid or greater, causing too much thiosulphate to be required. An apparent excess of oxidising action of dichromate is caused by the titration of the iodine in a solution of too high an acid concentration. This can be corrected by the exclusion of air, but more easily by dilution to such an hydrogen-ion concentration that the presence of air will not interfere. Chromates may be estimated to within 0.1% by the iodide method under the proper conditions. Potassium dichromate as a standard for thiosulphate solutions agrees with pure iodine to within 0.1%, but gives a slightly lower value. Errors in the iodine standardisation leading to such a difference appear to be about as probable as errors in the dichromate standardisation.

A standard solution of potassium permanganate is not trustworthy as a standard for thiosulphate solutions. J. F. S.

The Estimation of Sulphuric Acid as Barium Sulphate. Evidence of the Existence of a Complex Barium Sulphuric Acid. D. BALAREFF (*Z. anorg. Chem.*, 1922, **123**, 69—82).—The sources of error in the estimation of sulphuric acid as barium sulphate are: (1) occlusion of salts by the precipitate, for example, barium chloride; (2) occlusion of the mother-liquor; (3) the formation of the complex ion $[\text{Ba}_5(\text{SO}_4)_6]$, and this leads to the chief error. The author isolated the potassium salt of this complex ion $[\text{Ba}_5(\text{SO}_4)_6]\text{K}_2$. The most accurate results are obtained when barium sulphate is precipitated slowly from a dilute, boiling solution acidified with hydrochloric acid. Crystals in the precipitate have various forms; the prismatic form corresponds with the simple salt BaSO_4 . The chlorine content of barium sulphate is not caused by the salt $\text{SO}_4(\text{BaCl})_2$. W. T.

Application of Conductometric Titrations to Precipitation Analysis. V. Conductometric Titrations with Barium Salts. I. M. KOLTHOFF (*Z. anal. Chem.*, 1922, **61**, 433—448).—Sulphates may be accurately titrated conductometrically by means of barium chloride in solutions as dilute as 0.001*N*. The titration may be carried out more rapidly and exactly if alcohol is added to the solution than if an excess of barium sulphate is stirred in. The presence of mineral acids, aluminium, and calcium leads to low results, whereas the presence of iron salts generally causes the results to be too high. Nitrates in moderate amount do not interfere. The method is not applicable to the determination of sulphates in water owing to the presence of calcium.

Carbonates and chromates, as well as oxalates, tartrates, citrates, and malates in the presence of sufficient alcohol may all be titrated with barium chloride in a similar manner to sulphates, but phosphates, pyrophosphates, succinates, benzoates, and salicylates cannot be titrated by conductometric methods with barium chloride. A. R. P.

Some new Volumetric Methods. (Estimation of Sulphate, Lead, Acids, and Ammonia.) KARL JELLINEK and H. ENS (*Z. anorg. Chem.*, 1922, **124**, 185—202).—The volumetric methods of estimating sulphate are unsuitable on account of the filtrations involved. The authors have worked out the following method. Excess of barium nitrate is added to a sulphate solution, to this is added excess of potassium chromate and this excess estimated by running in a barium solution until the yellow colour disappears. Error by this method 0.3%. The use of silver salt as an indicator is not possible because the solubilities of silver and barium chromates are very near to one another. The following is given as a volumetric method of estimating lead; excess of chromate is added to a lead salt, a little silver nitrate is then added, and the excess of chromate is determined by titrating against a lead salt until the disappearance of the reddish-brown silver chromate. The latter method can also

be employed for the indirect estimation of sulphate. Titration of metals against arsenates and arsenites does not give positive results. The authors suggest the use of a copper solution as an indicator in acidimetry, the alkali being added until the precipitated hydroxide makes its appearance.

W. T.

Estimation of Selenium. LUIGI LOSANA (*Giorn. Chim. Ind. Appl.*, 1922, 4, 464—466).—The method used for estimating sulphur (this vol., ii, 582, 656) may be employed also for the estimation of selenium if a larger excess of iron is taken and the exclusion of air during the reduction is rendered as complete as possible. By suitable modification of the procedure, sulphur and selenium may be estimated simultaneously (cf. *J. Soc. Chem. Ind.*, 1922, Dec.).

T. H. P.

Rapid Method for Estimating Ammonia in Ammonium Salts. HERMANN BURKARDT (*Chem. Ztg.*, 1922, 46, 949).—Attention is directed to a rapid method for estimating ammonia in its salts, in which formaldehyde solution (containing no free acid) is added to the solution containing an ammonium salt, and the acid formed according to the equation $6\text{CH}_2\text{O} + 4\text{NH}_4\text{Cl} = 6\text{H}_2\text{O} + 4\text{HCl} + (\text{CH}_2)_6\text{N}_4$ is then directly titrated with a standard solution of sodium hydroxide, using phenolphthalein as an indicator.

A. J. H.

Titration of Nitrous Acid, and the Estimation of Nitrous and Arsenious Acids in the Presence of Each Other. ALFONS KLEMENC (*Z. anal. Chem.*, 1922, 61, 448—454).—The titration of acidified solutions of nitrites with permanganate in the air is attended with possible errors due to loss of nitrogen oxides and the slowness with which the reaction proceeds towards the end. These disadvantages may be overcome by adding the nitrite solution without acidifying to a solution containing an excess of potassium permanganate and sulphuric acid and enclosed in a glass-stoppered flask containing carbon dioxide under slightly reduced pressure. The solution is warmed at 40° to complete the reaction and the excess of permanganate is then titrated with standard oxalic acid.

[With F. POLLAK.]—The method is extended to the estimation of nitrous and arsenious acids in the same solution, one portion of which is titrated in the presence of sodium hydrogen carbonate with iodine for arsenious acid and another portion with permanganate as described above to obtain the sum of the two acids. Nitrous acid is then found by difference.

A. R. P.

The Use of Benzidine in the Detection of Phosphoric Acid. F. FEIGL (*Z. anal. Chem.*, 1922, 61, 454—457).—The yellow precipitate produced by ammonium molybdate in the usual test for phosphoric acid is often contaminated with arseno-molybdate and with molybdic acid, especially if an old solution of the reagent is used. To test this precipitate for the presence of phosphoric acid, it is collected on a close, ashless filter and washed somewhat, then moistened with a solution of benzidine hydrochloride containing acetic acid. The moist paper is held over an ammonia bottle, the vapour from which turns the precipitate blue in those parts con-

taining phosphoric acid. Even when no visible precipitate is produced in the test solution it should be poured through the paper and the paper tested as described. The blue colour is apparent when the original solution contains only 1 part of phosphorus as phosphoric acid in 460,000 parts of water. A. R. P.

The Volumetric Estimation of Phosphoric Acid and of Sodium Phosphate and Pyrophosphates. FRANK X. MOERK (*Amer. J. Pharm.*, 1922, **94**, 641—650).—Phosphoric acid and disodium hydrogen phosphate can be estimated volumetrically by titration with sodium hydroxide and hydrochloric acid, respectively, using a mixture of methyl-orange and indigo-carmin as indicator. To obtain results in consistent agreement with the silver phosphate precipitation method followed by titration of the liberated acid with alkali hydroxide until a permanent brown coloration of silver oxide is obtained, the dilution of the reacting solutions must be specified, and a definite quantity of sodium chloride must be added, the silver phosphate method being apparently unaffected by these factors. For the neutralisation method, using the indicator, the procedure is as follows. To 100 c.c. of water 0.2 c.c. each of 0.1% methyl-orange solution and 0.3% indigo-carmin solution are added, followed by dilute hydrochloric acid until the green colour changes without producing a violet colour. The solution thus prepared is divided into two equal parts, one being set aside and the other being added to the phosphoric acid or phosphate solution to be estimated, and the mixture titrated to match the tint of the reserved portion of the indicator solution with sodium hydroxide, or hydrochloric acid, as the case may be. The end-point corresponds in each case with the formation of NaH_2PO_4 . Under these conditions and using $N/2$ solutions, results in agreement with the silver phosphate method were obtained by titrating in presence of 7.5% of sodium chloride for the phosphoric acid, and 2.5% for the phosphate. Under similar conditions, sodium pyrophosphate can be titrated with acid as a diacid base. G. F. M.

Methyl-red in the Assay of Phosphoric Acid and Sodium Phosphate. FRANK X. MOERK and EDWARD J. HUGHES (*Amer. J. Pharm.*, 1922, **94**, 650—655).—Of the two methods described in the preceding abstract and the U.S.P. IX method for the volumetric estimation of phosphoric acid and phosphates, the latter process always gives low results. The mixed indicator method is the most rapid, but is influenced by the weight taken, by the amount of sodium chloride present, and by the strength of the volumetric solutions. The best results when working with unknown quantities, are obtained by the silver phosphate method, titrating back the liberated acid with alkali hydroxide and using methyl-red as indicator. The procedure is as follows. To 50 c.c. of standard silver nitrate solution one drop of methyl-red and a trace of alkali to produce a yellow colour are added, followed by 10 c.c. of the solution to be assayed. The liberated nitric acid (3 mols. per mol. of H_3PO_4 , or 1 mol. per mol. of Na_2HPO_4) is titrated with alkali hydroxide until the supernatant liquid is again yellow. During the neutralisation,

the pink colour may fade, in which case one or two drops more of methyl-red must be added.

G. F. M.

The Iodometric Micro-estimation of Phosphoric Acid, and of Phosphorus in Organic Compounds. O. SVANBERG, K. SJÖBERG, and G. ZIMMERLUND (*Arkiv Kem. Min. Geol.*, 1922, No. 10, 1—17).—An improvement on Neumann's method (A., 1903, ii, 243; 1905, ii, 68) for the micro-estimation of phosphorus is described, in which the phosphorus is precipitated as ammonium phosphomolybdate, and the nitrogen in the precipitate estimated by Bang's micro-Kjeldahl method, in which the ammonia is absorbed in acid, the excess of the latter being estimated by adding excess of potassium iodide and potassium iodate and titrating the liberated iodine with thiosulphate. If a correction is made for a systematic error, 0.05—1.00 mg. of phosphorus may be estimated to within 2—3%.

W. O. K.

Titration of Boric Acid in Presence of Phosphoric Acid. I. M. KOLTHOFF (*Chem. Weekblad*, 1922, 19, 449—450).—Addition of sodium citrate to solutions containing phosphoric and boric acids prevents interference by the latter in the titration of the phosphoric acid with sodium hydroxide. After the neutralisation, addition of mannitol allows of the titration of the boric acid by further addition of sodium hydroxide, addition of the alkali being continued until the pink colour of the phenolphthalein remains for at least three minutes, and is not destroyed by further addition of mannitol. Neither calcium nor magnesium interfere.

In boiling solutions containing boric acid, a reflux condenser is scarcely necessary, since after half an hour's boiling the loss is less than 1%.

S. I. L.

The Estimation of Boric Acid. W. W. DEERNS (*Chem. Weekblad*, 1922, 19, 480—481).—The author's method of estimating boric acid in presence of phosphoric acid by means of potassium iodide-iodate is simpler than the method proposed by Kolthoff (preceding abstract) of adding sodium citrate and titrating with alkali, and the interference of calcium compounds is not apparent. The citrate method is not new, having been proposed by Littmann (*Chem. Ztg.*, 1898, 22, 691) and Pfyl (A., 1914, ii, 290).

S. I. L.

Quantitative Estimation of Carbon and Hydrogen by means of the Sulpho-chromic Mixture. L. J. SIMON and A. J. A. GUILLAUMIN (*Compt. rend.*, 1922, 175, 525—527; cf. Guyot and Simon, A., 1920, i, 285, and ii, 332).—The method indicated is limited in scope, but is applicable to dibasic straight-chain acids, their methyl esters, and to all sugar derivatives which do not contain a methyl group directly linked to carbon, to aromatic acids, and to phenols which contain no alkyl substituents in the ring. The substance is completely oxidised by a known excess of reagent, the volume of carbon dioxide obtained is noted, and hydrogen estimated by determining volumetrically the excess of chromic acid. An alternative method depends on estimating the residual chromic

acid by addition of a known excess of a suitable organic substance and a second reading of the volume of carbon dioxide. A table is given of results obtained by the method as compared with standard figures for the same substances. The authors suggest that the method would be more generally applicable if silver chromate were used as the oxidising agent.

H. J. E.

The Function of Chromic Oxide in Oxidation by means of Sulpho-chromic Mixture. L. J. SIMON (*Compt. rend.*, 1922, 175, 768—770; cf. preceding abstract).—Chromic oxide formed in the reaction may play an important part in the oxidation. Control experiments in which chrome alum was added to the reagent showed that, in absence of oxidisable organic matter, the addition enables the liberation of oxygen to take place more readily and that the quantity of alum added influences the amount of oxygen evolved. If sufficient chromic oxide is added, sulphuric acid will completely decompose chromic anhydride on heating at 100°. Thus the addition of chromic oxide or its derivatives, when dealing with organic substances which are only partly oxidised by sulpho-chromic mixture, transforms what would otherwise be partial into complete oxidation, and in extreme cases gives rise to the evolution of free oxygen.

H. J. E.

The Tannic Acid Method for the Estimation of Carbon Monoxide in Blood. R. R. SAYERS and W. P. YANT (*U.S. Bur. Mines, Rep. Investigations*, 1922, No. 2356).—Standards are prepared containing 10—100% of carbon monoxide-hæmoglobin in 90—0% of oxyhæmoglobin as follows: 5 c.c. or more of blood are collected, 0.05 gram of potassium citrate or 0.02 gram of sodium fluoride being used for each 10 c.c. Half of the blood is saturated with carbon monoxide, and both parts are diluted to 10 vols. with water, and mixed in varying proportions. Then 0.1 c.c. of the blood to be tested is drawn from the finger into 1 c.c. of 0.05% potassium citrate or 0.03% sodium fluoride solution. To each of the blood samples contained in tubes of $\frac{5}{16}$ inch inside diameter is added 1 c.c. of 1% tannic acid in 1% pyrogallol solution. The tubes are inverted and comparison is made in eight to ten minutes. The standards, when properly sealed, may be kept for several weeks, but the reagent should be freshly prepared.

CHEMICAL ABSTRACTS.

Physical Method for the Estimation of Carbon Dioxide in the Respiratory Air. A. K. NOYONS (*Arch. Néerland. physiol.*, 1922, 7, 488—495).—A method is described for the estimation of carbon dioxide by the measurement of the heat conductivity of the mixture of gases.

W. O. K.

Estimation of Silica in Filtered Sea-water. ROGER C. WELLS (*J. Amer. Chem. Soc.*, 1922, 44, 2187—2193).—The author has investigated the estimation of silica in sea-water and finds that the most effective method is to add an aluminium salt if such is not already present and precipitate with a quantity of ammonia sufficient to produce a pink colour with rosolic acid; the silica carried down with the alumina is then recovered and estimated in

the usual way. It is shown that silica cannot be estimated by the loss on evaporation with hydrofluoric acid in the presence of calcium sulphate. An excess of at least two parts of alumina to one of silica is essential for the complete inclusion of silica in the alumina precipitate. An excess of ammonia favours the inclusion of silica in the ammonia precipitate, but this has a slightly solvent action on the aluminium hydroxide. A Sørensen value of 7—8, as shown by the pink colour with rosolic acid, is advisable in making the ammonia precipitation. A very small quantity of silica, roughly 0.3 mg., escapes precipitation and an equal quantity is generally found in the wash waters from the ammonia precipitate. In rock analysis, a single evaporation with hydrochloric acid is sufficient, provided silica is also determined in the ammonia precipitate. Eleven samples of water collected one mile south of Eastern Point Light, Gloucester, Mass., during eleven months gave contents of dissolved silica varying between 0.0029 and 0.0003 gram per litre. J. F. S.

The Estimation of Alkali Carbonates in Presence of Phenolphthalein. BONNIER (*Compt. rend.*, 1922, 175, 765—767).—Warder's method (A., 1881, 848) of estimating carbon in steel involves a titration of alkali hydroxide with sulphuric acid in presence of alkali carbonate. That of Winkler (Ireadwell, "Analytische Chemie," 9th ed., 2, 485) necessitates titration of the hydroxide in presence of barium chloride. Neither gives a sharp end-point. In the former case, the instability of the alkali hydrogen carbonate, on the formation of which the end-point depends, leads to errors and a study of the various factors which influence the result shows that the excess of sodium sulphate formed, the temperature, and the method of stirring have no effect. The initial colour of the indicator, the method of adding the acid, and the dilution (cf. Mestrezat, A., 1918, ii, 274) all affect the estimation to some extent. Winkler's method is satisfactory if carried out with normal solutions; with less concentrated solutions, however, the end-point tends to lose sharpness. H. J. E.

The Cobalt Nitrite Reaction for Potassium in Animal and Vegetable Cells. A. B. MACALLUM (*Arch. Néerland. physiol.*, 1922, 7, 304—308).—Sodium cobaltinitrite is a very sensitive reagent for the microchemical detection of potassium in animal and vegetable cells. The possibility of ammonia being precipitated by the cobaltinitrite may be avoided by the addition of formalin to the reagent, which converts the ammonia into hexamethylenetetra-amine. The only difficulty in the use of the reagent is that in animal tissue creatine may give a precipitate likely to be mistaken for potassium. W. O. K.

The Volumetric Estimation of Calcium. A. VÜRTHHEIM and G. H. C. VAN BERS (*Chem. Weekblad*, 1922, 19, 450—452).—Excess of ammonium oxalate is added to the solution, and after heating to boiling, calcium oxalate is precipitated by addition of ammonia; after filtration, the excess of ammonium oxalate is titrated with permanganate in the usual way. It is necessary to

use at least double the amount of oxalate theoretically required for the precipitation; on account of the rapid alteration of ammonium oxalate solutions, blank titrations must be carried out with each series of estimations.

The results agree very well with those obtained by the gravimetric estimation, and within very wide limits are independent of the temperature and degree of acidity of the solution to be titrated and of the strength of the permanganate solution employed.

S. I. L.

The Volumetric Estimation of Magnesium in Potassium Salts. A. VÜRTHEIM (*Chem. Weekblad*, 1922, 19, 461—462).—The method proposed by Precht (A., 1879, 1053), namely precipitation of magnesia by addition of carbonate-free alkali, and back titration of the excess after filtration, has been examined, and found to give satisfactory results for mixtures containing very large quantities of sodium, potassium, and calcium salts. Where calcium is present, oxalate is added to the alkali to ensure complete precipitation.

S. I. L.

Estimation of Lead in Lead Amalgam. M. G. MELLON (*J. Amer. Chem. Soc.*, 1922, 44, 2167—2174).—A résumé is given of the principles of the methods which have been used previously for the estimation of lead in lead amalgams. Data are presented which show the possibility of estimating quantities of lead of about 0.5 gram of lead in 30—50 grams of mercury with an accuracy of 0.05%. The method consists in the displacement of the lead by copper from an aqueous solution of copper nitrate, followed by the precipitation of the lead as chromate from the solution. It is shown that 0.4 gram of lead may be displaced from the amalgam in thirty minutes at the ordinary temperature if the mixture is stirred, but if allowed to remain stationary many hours are required for complete precipitation. The following procedure is recommended: the sample of amalgam is covered with water containing one drop of 10% acetic acid, 10 c.c. of a 2*N*-solution of copper nitrate are added and the mixture is stirred for thirty minutes and filtered to remove any metallic copper which has not dissolved in the mercury. One drop of 10% acetic acid is added to the filtrate, and the lead is precipitated as chromate by the method usually adopted.

J. F. S.

A Peculiar Catalytic Reaction for the Detection, and a Method for the Estimation, of the Smallest Traces of Copper [also a Lecture Experiment]. FRIEDRICH L. HAHN and G. LEIMBACH (*Ber.*, 1922, 55, [B], 3070—3074).—When a solution of a ferric salt is added to a solution of sodium thiosulphate a dark violet coloration is developed which gradually disappears as the ferric becomes reduced to ferrous salt and the sodium thiosulphate is converted into tetrathionate. The reaction is greatly accelerated by the presence of copper salts and its completion can be made more obvious by the addition of ammonium thiocyanate to the mixture; the latter substance has also the advantage of retarding the reaction.

The following solutions are required. Copper sulphate solution (3.9 grams of the hydrated salt per litre, of which 1 c.c. is diluted to 1 litre before use); ferric solution (5 grams of iron alum and 25 c.c. of 2*N*-hydrochloric acid per litre); 5*M*-ammonium thiocyanate solution; 1/15*M*-sodium thiosulphate solution. For the recognition of the minutest traces of copper, two litres of the iron solution should be mixed with 10—20 c.c. of thiocyanate; for the estimation of larger amounts more thiocyanate (up to 200 c.c.) may be used.

A neutral solution of the substance under investigation is placed in a beaker (500 c.c. capacity) and in a similar series of beakers known quantities of copper solution are placed. The solutions are made up to a fixed volume with distilled water, and to each are added 100 c.c. of the iron thiocyanate solution. Simultaneously (by means of a row of test-tubes fastened to a rod which can be rotated horizontally) 25 c.c. of thiosulphate solution are added to each beaker and the contents well mixed. The times required for decolorisation (which are conveniently 10—20 minutes) are then compared.

The velocity of the reaction is rather greatly dependent on small variations in the ratio of the concentrations, iron : thiocyanate : thiosulphate, on the degree of acidity and the temperature, so that it is scarcely possible at present to indicate definite intervals of time for definite quantities of copper. It is necessary to observe the progressive decolorisation in the solution under investigation and in solutions of known copper content simultaneously and to use a sufficient volume of solution (about 100 c.c.) to avoid the effect of local variations of temperature. The smallest amount of copper which can be identified with certainty under these conditions is 0.002 m.g.; further refinement is probable if a thermostat is used.

A similar effect on the course of the reaction has not been observed with any other substance yet investigated. Acceleration is caused by relatively large amounts of platinum, but the effect is so little marked that a confusion with copper is impossible. Aluminium, zinc, nickel, and arsenic in particular retard the action, especially in strongly acid solution. If such metals are to be investigated with regard to their copper content, it is essential that an equal weight of "foreign" metal should be added to the control solutions; in this manner, 0.001% of copper in nickel can be rapidly and certainly detected.

H. W.

Potentiometric Titration of Copper. E. ZINTL and H. WATTENBERG (*Ber.*, 1922, 55, [B], 3366—3370).—The method consists in the reduction of the cupric to cuprous salt in hydrochloric acid solution by means of titanium trichloride and subsequent re-oxidation with standard potassium bromate or dichromate solution in an atmosphere of carbon dioxide.

The apparatus consists of a beaker placed on an electrically heated hot plate and covered with a clock glass provided with four holes for the introduction of the nozzle of the burette, a stirrer, a T-tube which holds the indicator electrode (a stout smooth platinum

wire) and serves also for the introduction of carbon dioxide and the limb of the normal electrode (a calomel cell charged with solid potassium chloride and saturated potassium chloride solution). The *E.M.F.* of the titration cell is measured by compensation in the usual manner with the aid of a capillary electrometer. The copper solution is treated with a sufficient quantity of hydrochloric acid and reduced by a slight excess of titanium trichloride, whereby a colourless solution results if sufficient hydrochloric acid is present to prevent precipitation of cuprous chloride. The addition of standard bromate or dichromate solution causes the oxidation of the excess of tervalent titanium and subsequently that of the copper, the end-points of the two actions being marked by abrupt changes in the potential. Titration is effected at about 80°, at which temperature the potentials adjust themselves almost instantaneously; in cold solution, the bromate in particular reacts somewhat slowly with titanium trichloride. The concentration of the hydrochloric acid has not a very marked influence, and is suitably maintained at 4—8%. Considerable quantities of ammonium salts have no disturbing effect, but iron is titrated with the copper. Small quantities of nitric acid are reduced by the excess of titanium trichloride, after which they do not influence the change further. The use of stannous chloride in place of titanium chloride is not recommended, since the potentials, even in hot solution, only become constant slowly as long as an excess of stannous salt is present.

H. W.

Volumetric Estimation of Copper by means of Sodium Nitroprusside. GEORGES JORET (*Ann. Falsif.*, 1922, **15**, 354—356).—A solution of the copper salt, containing about 0.1 gram of copper, is neutralised with ammonia, then acidified slightly with nitric acid, treated with 40 c.c. of *N*/10-sodium nitroprusside solution (14.895 grams per litre), diluted to 200 c.c., and filtered; 100 c.c. of the filtrate are treated with 20 c.c. of *N*/10-silver nitrate solution, the mixture is filtered, and the excess of silver nitrate titrated with *N*/10-thiocyanate solution in an aliquot portion of the filtrate. The copper solution must not contain silver, nickel, cobalt, or halogens.

W. P. S.

Separation of Copper by means of Thiophenylhydantoic Acid. H. H. WILLARD and DOROTHY HALL (*J. Amer. Chem. Soc.*, 1922, **44**, 2253—2254).—In solutions slightly acidified with acetic acid, thiophenylhydantoic acid precipitates quantitatively copper, lead, mercury, cadmium, bismuth, and antimony, whilst arsenic, tin, and metals not precipitated by hydrogen sulphide are not precipitated under these conditions. When the copper precipitate is ignited to oxide, the product contains sulphate sufficient to cause an error of 0.1—0.6 mg. in the total weight. The copper compound is very unstable and quite unsuitable for direct weighing. To estimate copper by means of the precipitate with thiophenylhydantoic acid, it is best to fuse it with potassium pyrosulphate, dissolve in water, and estimate the copper by any of the well-known methods. The bismuth and antimony compounds

with thiophenylhydantoic acid are soluble in alcohol. To separate copper from the metals of the iron and zinc groups, the solution is neutralised with ammonia and 5 grams of glacial acetic acid are added. If metals precipitated by ammonia are present, 7 grams of citric acid are added before neutralising for every gram of such metal present. The solution is made up to 300—400 c.c., raised to the boiling point, and treated with 0.5 gram of thiophenylhydantoic acid in a little hot water, then dilute ammonia is added until a yellow precipitate begins to form. The solution is boiled, and if all the copper is precipitated the precipitate separates at the top, leaving a clear solution below. If this does not happen, a little more ammonia is added, but this may cause the precipitate to turn brown, due to the formation of sulphide. The precipitate is filtered hot, washed with hot water, and treated as stated above. The method is very good for small quantities of copper.

J. F. S.

Contradictions and Errors in Analytical Chemistry. I. The Precipitation of Aluminium by Thiosulphate and its Separation from Iron. II. The Ageing of Volumetric Thiosulphate Solutions. FRIEDRICH L. HAHN [with G. LEIMBACH (I) and H. WINDISCH (II)] (*Ber.*, 1922, 55, [B], 3161—3165).—I. The precipitation of aluminium by means of thiosulphate under the usual conditions (boiling the solution until sulphur dioxide is completely expelled and then adding ammonia) is only very incomplete; the apparent accuracy of the results is due to a balance of errors, the unprecipitated aluminium hydroxide being compensated by the alkali carried down by the precipitate. Precipitation is almost quantitative when the solution is boiled for only a short time and the remainder of the metal is precipitated by addition of a base. An almost complete separation from iron can be achieved if a very weak base, preferably phenylhydrazine, is used.

II. The addition of a very small amount of alkali enables thiosulphate solutions to be preserved without alteration in strength from the first day. The change which they otherwise suffer appears to be caused by the faintly acid reaction of distilled water; it certainly does not depend on the formation of sulphite or sulphide.

H. W.

The Separation of Ferric Oxide and Aluminium Oxide from Magnesium Oxide by the Nitrate Method. ANDRÉ CHARRIOU (*Compt. rend.*, 1922, 175, 693—695; cf. this vol., ii, 319).—In separating the oxides of iron and aluminium from that of magnesium by heating with ammonium nitrate, the retention of magnesium oxide by the other oxides increases with the magnesium content; aluminium oxide retains a relatively smaller amount than ferric oxide. By increasing the concentration of the ammonium nitrate solution from 4% to 10%, the magnesium oxide is almost completely removed, and if the mixture is treated three times in this way, no trace of magnesium remains.

H. J. E.

Separation and Estimation of Cobalt. I. Separation of Cobalt by means of Thiophenylhydantoic Acid. I. Separation of Cobalt from other Metals. H. H. WILLARD and DOROTHY HALL (*J. Amer. Chem. Soc.*, 1922, **44**, 2219—2226).—Cobalt may be quantitatively separated from arsenic, uranium, vanadium, titanium, tungsten, molybdenum, zinc, manganese, chromium, aluminium, magnesium, and calcium by precipitation with thiophenylhydantoic acid in slightly ammoniacal solution. In the presence of iron, the precipitate is rarely free from this impurity, and usually contains from one to five mg. regardless of the amount originally present. This does not, however, interfere with the volumetric estimation of cobalt. Nickel is always partly precipitated. The cobalt precipitate has not a definite composition, and is probably mixed with some cobalt sulphide, thus making it necessary to convert the cobalt into some other more definite form. The various separations of cobalt from other metals are effected as follows. *Separation from iron.* A solution containing 1 gram of iron and 25 mg. of cobalt in 300 c.c. is placed in an Erlenmeyer flask, treated with 8 grams of citric acid, and neutralised with ammonia (*d* 0.90) and 5 c.c. of excess ammonia are added. The solution is warmed at 35°, 0.7—1.0 gram of thiophenylhydantoic acid dissolved in 30 c.c. of water or alcohol added, and the mixture shaken vigorously for several minutes. The cobalt is precipitated and after boiling the precipitate may be easily filtered. *Separation from manganese.* This separation is effected as in the case of iron. *Separation from zinc.* The procedure is the same as for iron, except that an excess of 10 c.c. of ammonia is necessary and the solution must be boiled for several minutes to ensure complete precipitation of the cobalt. The other metals named are separated in the same way as zinc, except that 7 grams of citric acid are used. J. F. S.

Separation and Estimation of Cobalt. II. Gravimetric Estimation of Cobalt. H. H. WILLARD and DOROTHY HALL (*J. Amer. Chem. Soc.*, 1922, **44**, 2226—2231; cf. preceding abstract).—Various methods for the gravimetric estimation of cobalt have been examined, and it is shown that, although pure cobalt nitrate may be accurately ignited to the oxide Co_2O_3 and the latter reduced by hydrogen to the metal, the oxide obtained by the ignition of the thiophenylhydantoic acid precipitate (*vide supra*) contains a little sulphate. If the oxide obtained in this ignition is assumed, empirically, to be Co_2O_3 , the results for cobalt are fairly good. The ignition of cobalt sulphate at 550° is the most accurate method for the estimation of cobalt, but when this method is applied to the thiophenylhydantoic acid precipitate the results are usually slightly high and the sulphate is not completely soluble in water. The electrolytic estimation of cobalt is very satisfactory when the proper conditions and precautions are observed. Precipitation as cobalt ammonium phosphate followed by ignition to cobalt pyrophosphate is less accurate than the preceding methods. J. F. S.

Separation and Estimation of Cobalt. III. Volumetric Estimation of Cobalt. H. H. WILLARD and DOROTHY HALL (*J. Amer. Chem. Soc.*, 1922, **44**, 2237—2253; cf. preceding abstracts).—A number of methods have been examined for the volumetric estimation of cobalt. It is shown that in general volumetric methods are the most accurate and satisfactory for the estimation of cobalt. The most accurate method involves the oxidation of cobalt to cobaltic hydroxide in strongly alkaline solution by means of a perborate or hydrogen peroxide, followed by its volumetric reduction to a cobaltous salt. This reduction may be accomplished by the following methods: (a) By the addition of potassium iodide in acid solution, the iodine liberated being titrated with sodium thiosulphate; in this method iron must be absent. (b) By adding the cobaltic hydroxide to an acid ferrous sulphate solution, the excess of which is titrated with potassium permanganate; an empirical factor must be used in the calculation. (c) By the addition of a strongly acid solution of stannous chloride, the excess of which is titrated with iodine, iodate, bromate, or dichromate solution. The last, titrated electrometrically, is especially recommended. If iron is present, iodine will oxidise it quantitatively only in a neutral solution. (d) By adding titanous sulphate, the excess of which is titrated with permanganate. Since titanous hydroxide decomposes water with the liberation of hydrogen, the alkaline solution must be first almost completely neutralised by one of the several methods suggested. The presence of nickel does not interfere with this titration, but the oxidation is incomplete when more than 7 mg. of iron are present. In the presence of a large excess of potassium hydrogen carbonate, cobalt is oxidised by hydrogen peroxide to a green tervalent compound, which is reduced in the presence of pyrophosphate by ferrous sulphate, the excess of which is titrated with permanganate after acidifying with sulphuric acid. Iron and manganese in small amounts do not interfere in this method, but the principal source of error is in the removal of the excess of peroxide without decomposing the cobalt compound. J. F. S.

Gravimetric Estimation of Nickel as Nickel Dioxide. WILHELM VAUBEL (*Chem. Ztg.*, 1922, **46**, 978).—Nickelous hydroxide is precipitated from the solution with sodium hydroxide, collected, washed, and ignited. The ignited precipitate is again washed until free from alkali metals, dried, and dissolved in nitric acid. The excess of nitric acid is evaporated and the residue heated for thirty minutes in an air-bath at 280—330°. Pure black nickel dioxide results. The oxides of nickel at present known are NiO, Ni₃O₄, and NiO₂. The oxide Ni₂O₃ has not yet been obtained. H. C. R.

Electrometric Estimation of Nickel with Silver Nitrate. ERICH MÜLLER and HANS LAUTERBACH (*Z. anal. Chem.*, 1922, **61**, 457—464).—The nickel solution is treated with a slight excess of potassium cyanide solution, sufficient to decolorise the mixture, and connected with a normal calomel electrode, a galvanometer,

and a sliding resistance, the ends of which are connected with an accumulator, and one end and the sliding contact with a voltmeter. A silver wire in the solution acts as indicator electrode, and the solution is agitated by means of a motor-driven stirrer. The sliding resistance is adjusted until the voltmeter indicates 0.075 volt and silver nitrate solution is added slowly until the galvanometer indicates zero. The results are calculated as in the older chemical process.

A. R. P.

Electrometric Titration of Dichromate with Ferrous Sulphate. MARION EPPLEY and WARREN C. VOSBURGH (*J. Amer. Chem. Soc.*, 1922, **44**, 2148—2156).—The conditions under which dichromate may be estimated by electrometric titration with ferrous sulphate have been investigated. The practical limits of acidity for hydrochloric and sulphuric acids have been determined. It is shown that the amount of ferrous sulphate required to titrate a given amount of chromic acid varies with the concentration of the latter. Dissolved air has a negligible effect on the titration. Hydrofluoric acid in sufficient concentration reduces the amount of ferrous sulphate required by about 0.1%. When ferrous sulphate, standardised by means of a standard solution of potassium permanganate, is used to standardise a known amount of dichromate in 0.01*N*-solution, the value found for the latter is about four parts per thousand higher than the calculated value. When the dichromate concentration is 0.003*N*, the amount of ferrous sulphate required is about 1% higher than the calculated quantity. The titration of the ferrous ion with a potassium dichromate solution gives results which agree with those obtained by the reverse titration.

J. F. S.

The Analysis of High-percentage Tungsten Alloys. K. SEEL (*Z. angew. Chem.*, 1922, **35**, 643—644).—The serious loss of platinum involved in the customary fusions of tungsten alloys with sodium and potassium carbonates and nitrates may be avoided by fusing the finely divided alloy with sodium hydroxide and nitrate in silver crucibles. The fusion takes about 1 hour, and the crucibles are not seriously attacked. 0.5 Gram of the finely divided alloy is fused with 6 grams of sodium hydroxide and 3 grams of sodium nitrate. The fused mass is dissolved in water and filtered, the iron oxide remaining on the filter being dissolved in hydrochloric acid containing a little potassium chlorate, filtered, and the two filtrates united. The iron is estimated as ferric oxide as usual. The aluminium and the greater part of the silica are removed from the filtrate with ammonia, filtered off, and the tungsten in the filtrate precipitated with mercurous nitrate after acidifying, boiling off the carbon dioxide and concentrating to about 150 c.c. The tungstic acid obtained by ignition of the precipitate is strongly contaminated with alkali, which is removed by repeated extractions with 7% hydrochloric acid, and traces of silica are finally removed by treatment with hydrofluoric acid. In the case of alloys rich in iron, a residue containing iron and tungsten remains after the original fusion. This is fused with pyrosulphate and the tungsten

precipitated with nitric acid, estimated as tungstic acid and added to the figure obtained above. The iron in the filtrate is estimated in the usual way and added to the figure obtained above.

H. C. R.

Separation of Antimony from Arsenic and Tin. FRIEDRICH L. HAHN (*Z. anorg. Chem.*, 1922, **123**, 276).—Antimony can be separated quantitatively from arsenic and tin by oxidation to pyroantimonate (cf. A., 1916, ii, 266). This method is due to Hampe.

W. T.

The Estimation of Bismuth as Metal. ALBIN KURTENACKER and FELICITAS WERNER (*Z. anorg. Chem.*, 1922, **123**, 166—170).—The authors find that metallic bismuth can be accurately estimated by dissolving it in a ferric salt and titrating the ferrous salt produced by means of potassium permanganate, the reaction being $\text{Bi} + 3\text{Fe}^{+++} = \text{Bi}^{+++} + 3\text{Fe}^{++}$. Ferric chloride was found more suitable than the sulphate. The reduction of bismuth nitrate by formaldehyde, hypophosphorous acid, and alkali stannite gave results about 5% too low, this being due to the metal being mixed with some bismuthous oxide. Conditions for complete reduction to the metal could not be found.

W. T.

New Method for Estimating Volatile Substances in Air. E. FRITZMAN and K. MACJULEVITSCH (*J. Russ. Phys. Chem. Soc.*, 1920, **52**, 212—226).—To estimate the content of light petroleum vapour in air, the latter is first freed from dust, moisture, and carbon dioxide by passing it through tubes containing, in order, soda-lime, calcium chloride, phosphoric oxide, and cotton wool, the light petroleum being retained in two U-tubes packed with magnesium turnings and cooled in liquid air.

T. H. P.

Estimation of Toluene, Xylene, and Benzene. G. ZABOROWSKI (*Mat. grasses*, 1922, **14**, 6160—6161).—Commercial samples of the above hydrocarbons or their respective mixtures with "gasoline" may be examined, either volumetrically by sulphonation with a known quantity of sulphuric acid and titration of the excess of acid, or densimetrically. (a) About 3.5 grams of the sample and three times its weight (for xylene, four times) of (preferably 100%) sulphuric acid are warmed to 45° in a 50 c.c. glass-stoppered flask, shaken thoroughly for five minutes, allowed to cool, poured into 100 c.c. of water, boiled to remove light oils, and titrated with standard sodium hydroxide solution, using phenolphthalein. For toluene, the quantity of acid used is multiplied by 1.876; for xylene, by 2.163. In the case of benzene, sulphonation must be effected in a reflux apparatus. (b) The densimetric estimation consists in the use of the expression $T = -[100D(d_1 - d_2)] / (D - d_2)d_1$, where d_1 represents the d_{15} of the sample, d_2 the d_{15} of the un-sulphonated residue after sulphonation, and D the d_{15} of pure toluene (0.8706), xylene, or benzene.

CHEMICAL ABSTRACTS.

The Detection of Nitro-compounds. H. J. PRINS (*Perf. Essent. Oil Rec.*, 1922, **13**, 355).—The accelerating effect of nitro-

benzene on the reaction velocity of metals with weak acids is applied to the detection of this substance in benzaldehyde. The reaction is carried out by heating 2 c.c. of the oil and 6 c.c. of 80% acetic acid for one minute with a small piece of tin foil. In presence of 5% of nitrobenzene, the tin disappears immediately on boiling, with 1% the tin turns dark and disappears about ten minutes after the mixture has been heated, with 0.1% the tin turns black after about ten minutes, with 0.01% the tin becomes greyish-black after remaining over-night, whilst with pure benzaldehyde the metal retains its bright surface. In a similar way, the presence of 0.1% of artificial musk can be detected in perfumes. The reaction is not exclusively specific for nitro-compounds, as organic peroxides have the same property, but only oils rich in terpenes which contained peroxides owing to exposure to air show the reaction and then only feebly. Likewise worm-seed oil containing the peroxide caridol can scarcely give rise to confusion, as 5% of nitrobenzene in benzaldehyde reacts more rapidly than pure worm-seed oil. Peroxidised benzaldehyde does not show the nitrobenzene reaction at all.

G. F. M.

Estimation of Phenol in Mixtures of Tar Acids. W. H. HOFFERT (*J. Soc. Chem. Ind.*, 1922, 41, 334—337T).—The method proposed depends on the fact that the freezing point of phenol hydrate (16°) is lowered to the same extent by equal weights of each of the three cresols. In the case of a mixture of phenol and cresols, the freezing point may be determined directly, after the addition of water amounting to 10% of the weight of the mixture, if the quantity of phenol present exceeds 55%; if the proportion of phenol is less, a known quantity of pure phenol (m. p. 40.5°) must be added to bring the proportion above 55%. With a mixture of tar acids, it is first necessary to remove neutral hydrocarbons and pyridine bases by distilling with steam from a sodium hydroxide solution of the sample, or by extracting the same solution with ether or benzene; the recovered tar acids are then distilled. If the mixture distils completely below 203° , the estimation is made directly on the distillate, but otherwise, higher homologues must be separated by fractional distillation. A known weight of pure phenol is added to the distillate and the mixture is treated with 10% of its weight of water. Directions are given in detail regarding the procedure adopted for determining the freezing point and the amount of phenol corresponding with the freezing point obtained is found by reference to a graph. The method may be applied to the estimation of phenol in crude carboic acid.

W. P. S.

Iodometric Estimation of Sugars. FR. AUERBACH and E. BODLÄNDER (*Z. angew. Chem.*, 1922, 35, 631—632).—In the iodometric titration of the excess of Fehling's solution used in the estimation of a reducing sugar, the volume of $N/10$ -thiosulphate solution required is not strictly proportional to the amount of sugar present. A table is given showing the quantities of invert-sugar corresponding with different quantities of thiosulphate

solution. Replacement of a part of the potassium iodide used by potassium thiocyanate, as proposed by Bruhns, is not recommended.

W. P. S.

The Modified Lehmann Method for the Estimation of Dextrose. Adaptation to Small Quantities of Reducing Sugars. PAUL FLEURY and LOUIS BOUTOT (*Bull. Soc. Chim. Biol.*, 1922, 4, 361—374).—In the estimation of dextrose by Bruhns's method (A., 1920, ii, 773) the reagents should be added to the copper solution in the following order: iodide, acid, thiocyanate. The method has been used for the estimation of small quantities of reducing sugars in biological fluids. E. S.

Trustworthiness of the Benedict and Folin-Wu Blood-sugar Estimations. F. A. CSONKA and GRACE C. TAGGART (*J. Biol. Chem.*, 1922, 54, 1—3).—The different results obtained by the Folin-Wu (A., 1920, ii, 337) and the Benedict (A., 1918, ii, 247) methods are due to the presence in blood of a substance which reacts with picric acid but does not reduce copper solutions. The former method is thus the more trustworthy. E. S.

Estimation of Starch. I. Estimation of Starch in Barley and in Wheat. ARTHUR R. LING [with E. H. CALLOW and W. J. PRICE] (*J. Inst. Brewing*, 1922, 28, 838—853).—The method proposed depends on the hydrolysis of the starch by malt diastase as suggested by Brown and Morris (T., 1885, 47, 527). Since the amount of maltose yielded by a pure starch varies with the diastatic power of the malt, the authors have determined the relation between the percentage of maltose obtained from pure barley and wheat starches when these are hydrolysed under definite conditions, and the diastatic power of the malt employed. The results of these experiments are given in graphic form. To estimate starch in barley or wheat, 5 grams of the finely-powdered grain are extracted in a Soxhlet apparatus for 3—3½ hours with alcohol (d 0.920), the powder is then rinsed into a beaker with about 100 c.c. of water, and the mixture is boiled for ten minutes. The starch paste thus obtained is cooled to 57°, 10 c.c. of malt extract (prepared from a malt of known diastatic power) are added, the mixture is kept at 57° for one hour, then boiled, filtered, and the insoluble portion washed. The filtrate is diluted to 200 c.c. and 30 c.c. of this solution are diluted to 100 c.c. and titrated against 10 c.c. of Fehling's solution. The cupric reducing power of the malt extract is determined at the same time and under the same conditions, and the value obtained is used to correct the reducing power of the conversion product. W. P. S.

Identification of Inulin by a Mycological Method. ALDO CASTELLANI and FRANK E. TAYLOR (*Biochem. J.*, 1922, 16, 655—658).—*Monilia macedoniensis*, Castellani, ferments inulin with the production of gas. It also ferments dextrose, lævulose, galactose, and sucrose. The method consists of the utilisation of the above mould in conjunction with other fungi. S. S. Z.

Estimation of Volatile Acids in Wine. PHILIPPE MALVEZIN (*Ann. Falsif.*, 1922, 15, 360—362).—In reply to criticism, the author states that he sees no reason for altering the procedure described in his ether-extraction method for the estimation of volatile acids in wine. Tartaric and lactic acids are not extracted by the ether under the conditions given. W. P. S.

The Separation of Solid and Liquid Fatty Acids. W. MEIGEN and A. NEUBERGER (*Chem. Umschau*, 1922, 29, 337—342).—The methods of Varrentrapp, Bull and Fjellanger, Farnsteiner, David, Niegemann, and Facchini and Dorta were tested on known mixtures of solid and liquid fatty acids and by none was a true quantitative separation obtained. By the methods of Bull and Fjellanger, Facchini and Dorta, and David about one-half of the liquid acids was recovered in a nearly pure state. By recrystallising the solid fraction, part of this was also obtained nearly pure. By precipitating the aqueous solution of the potassium salts with excess of thallous sulphate, however, an almost quantitative separation of stearic, palmitic, and elaidic acids from oleic acid was attained, and no recrystallisation was necessary. A mixture of 0.502 gram of oleic acid with 0.505 gram of stearic acid gave 0.509 gram of solid acids of iodine value 0.3 and m. p. 68°, whilst a mixture of 0.200 gram of oleic acid and 0.502 gram of stearic acid gave 0.529 gram of solid acids of iodine value 0.9 and m. p. 67.5°. H. C. R.


The Neutralisation of Tartaric Acid by Potassium Hydroxide in Presence of Chlorides of Alkaline Earths. L. J. SIMON and L. ZIVY (*Compt. rend.*, 1922, 175, 620—622).—The fact that titration of alkali hydroxide against tartaric acid gives two end-points; that with helianthin indicating formation of the acid salt and that with phenolphthalein the normal salt, may be used in estimating mixtures of tartaric acid with other acids which give an end-point in presence of helianthin indicating the formation of normal salt. The presence of alkaline-earth chlorides in the solution introduces an error, as each molecule behaves towards potassium hydroxide as an equivalent quantity of hydrochloric acid. This effect is produced by any quantity and is quantitative in each case, the limiting value of the effect being the amount of tartaric acid present. Chlorides of other metals do not interfere with the titration in this manner. H. J. E.

Estimation of Salicylic Acid in Blood-serum and other Fluids of the Body. H. HERISSEY (*Compt. rend. Soc. biol.*, 1922, 87, 333—336).—A mixture of 10 c.c. of serum, 5 c.c. of water, 0.5 c.c. of a solution of 1 gram of sulphuric acid in 5 c.c. of water, and 40 c.c. of ether is shaken together for one minute in a stoppered glass bottle of 125 c.c. capacity. After an interval of a few minutes, 4.5 c.c. of sulphuric acid solution are again added and the contents well mixed by rotation of the flask, the supernatant ethereal layer


being decanted after five minutes. Of this solution, 30 c.c. are shaken in a funnel with 2—3 grams of anhydrous sodium sulphate; 3 c.c. of water containing one drop of dilute ferric chloride solution are then added, and the whole is vigorously shaken. In the presence of salicylic acid, a violet colour appears in the water layer on keeping. Should a positive reaction be given, the contents are thoroughly mixed and transferred to a small crystallising dish, the ether being allowed to evaporate spontaneously at room temperature. It is necessary to use a stirring-rod to prevent the formation of deposits on the sides of the dish. The solution is then filtered into a test-tube, together with a few drops of rinsing water, and its coloration compared with standards prepared from 10 c.c. of normal serum, 5 c.c. of water, and varying quantities of salicylic acid. A definite coloration is obtained with only 10 mg. of sodium salicylate per litre.

CHEMICAL ABSTRACTS.

Substituted Naphthalenesulphonic Acids. I. A Method for Identifying H-Acid and its Intermediates Obtained from Naphthalene-2 : 7-disulphonic Acid. D. F. J. LYNCH (*J. Ind. Eng. Chem.*, 1922, **14**, 964—965).—A method for the detection of 1 : 8-dinitronaphthalene-3 : 6-disulphonic, 1 : 8-diaminonaphthalene-3 : 6-disulphonic, 1-amino-8-naphthol-3 : 6-disulphonic (H-acid), and 1 : 8-dihydroxynaphthalene-3 : 6-disulphonic (chromotropic) acids in their mixtures by means of their reactions with solutions containing 2-cymidine sulphate, cobalt chloride, zinc sulphate, or the hydrochlorides of benzidine, α -naphthylamine, ψ -cumidine, toluidine, and *p*-nitrotoluidine, is developed. The tests are carried out with 3—5 c.c. of less than one-tenth gram-molecular solutions of the naphthalenesulphonic acids. The following substances were prepared. Benzidine salt of 1 : 8-dinitronaphthalene-3 : 6-disulphonic acid,



light yellow prisms, m. p. 275° (decomp.). Solubility 0.036 part at 20°. Cymidine 1 : 8-dinitronaphthalene-3 : 6-disulphonate, $\text{C}_{10}\text{H}_4(\text{NO}_2)_2(\text{SO}_3\text{H})_2\cdot(\text{C}_6\text{H}_3\text{MePr}^6\cdot\text{NH}_2)_2$, straw-coloured prisms, m. p. 265° (decomp.), solubility 0.059. Benzidine salt of chromotropic acid, $\text{C}_{10}\text{H}_4(\text{OH})_2(\text{SO}_3\text{H})_2\cdot\text{NH}_2\cdot\text{C}_6\text{H}_4\cdot\text{C}_6\text{H}_4\cdot\text{NH}_2$, brown prisms, m. p. 278° (decomp.), solubility 0.085. *p*-Nitrotoluidine 1 : 8-diaminonaphthalene-3 : 6-disulphonate,



greyish-tan needles, m. p. 270° (decomp.), solubility 0.186. Mono- α -naphthylamine salt of H-acid, $\text{NH}_2\cdot\text{C}_{10}\text{H}_4(\text{OH})(\text{SO}_3\text{H})_2\cdot\text{NH}_2\cdot\text{C}_{10}\text{H}_7$, light violet-grey prisms, m. p. 278° (decomp.), solubility 0.1006. Mono- ψ -cumidine salt of H-acid, $\text{C}_{10}\text{H}_2\text{O}_7\text{NS}_2\cdot\text{NH}_2\cdot\text{C}_6\text{H}_2\text{Me}_3$, grey prisms, m. p. 272° (decomp.), solubility 0.097. Mono-*p*-nitrotoluidine salt of H-acid, $\text{C}_{10}\text{H}_2\text{O}_7\text{NS}_2\cdot\text{NH}_2\cdot\text{C}_6\text{H}_3\text{Me}\cdot\text{NO}_2\cdot\text{H}_2\text{O}$, grey prisms, m. p. 265° (decomp.), solubility 0.106. Toluidine salt of H-acid, $\text{C}_{10}\text{H}_2\text{O}_7\text{NS}_2\cdot\text{NH}_2\cdot\text{C}_6\text{H}_3\text{Me}\cdot\text{C}_6\text{H}_3\text{Me}\cdot\text{NH}_2\cdot 3\text{H}_2\text{O}$, greyish-violet prisms, m. p. 260° (decomp.), solubility 0.102. Zinc 1 : 8-diamino-3 : 6-disulphonate, $\text{C}_{10}\text{H}_4(\text{NH}_2)_2\text{S}_2\text{O}_6\text{Zn}\cdot 3\text{H}_2\text{O}$, grey needles,

solubility 6.4; *cobalt* salt, $C_{10}H_4(NH_2)_2S_2O_6Co, 2H_2O$, brown needles, solubility 6 at 20°. The solubilities are parts per 100 parts of water, at room temperature unless otherwise specified.

A. J. H.

Estimation of Anthraquinone. O. A. NELSON and C. E. SENSEMAN (*J. Ind. Eng. Chem.*, 1922, 14, 956—957).—The following volumetric method has been devised for the estimation of anthraquinone in order to avoid the difficulties of filtration experienced in the method of Lewis (A., 1918, ii, 338). To about 0.5 gram of the sample of anthraquinone, intimately mixed with 3—4 grams of zinc dust, is added 100 c.c. of boiling 5% sodium hydroxide. After five minutes, the mixture is filtered through asbestos contained in a glass tube, about 24 cm. long and 3.5 cm. in diameter, and tapered at one end. This funnel is electrically heated at 90—95° and contains a mechanical stirrer. The red filtrate containing oxanthranol is received in a suction flask and is there directly titrated with a standard solution of potassium permanganate (3.8 grams per litre). Usually three to eight washings of the residue with hot sodium hydroxide solution are necessary in order completely to free it from unreduced anthraquinone, and the apparatus is arranged so that aerial oxidation of the filtrate is prevented. In the titration, the end-point is reached when the filtrate becomes colourless, except when phenanthraquinone is present when a bluish-green colour is obtained which changes to brownish-green on addition of excess of potassium permanganate. Accurate results are obtained by this method even when the anthraquinone contains phenanthraquinone, anthracene, phenanthrene, phthalic acid, phthalic anhydride, or other oxidation products of anthracene and phenanthrene.

A. J. H.

A Chemical Method of Assaying the Active Principles of Digitalis. ARTHUR KNUDSON and MELVIN DRESBACH (*J. Pharm. Expt. Ther.*, 1922, 20, 205—220).—The pharmacological activity of extracts of digitalis is found to be parallel with the colour developed on treating the digitalis preparation, decolorised with lead acetate, with an alkaline picrate solution. In order to evaluate the activity of the extract, comparison is made in the colorimeter with a standard ouabain solution treated in the same way.

W. O. K.

The Use of Sodium α -Naphthol-2-sulphonate for the Spectrophotometric Estimation of Aromatic Amino-compounds. W. E. MATHEWSON (*J. Assoc. Off. Agric. Chem.*, 1922, 6, 16—28).—The method consists essentially of coupling the naphtholsulphonate with the diazo-compound of the amine, and determining the concentration of the dye photometrically, in comparison with a standard solution of the pure dye. Sodium α -naphtholsulphonate is suited to the purpose because of the ease with which it is to be obtained pure, and to the rapidity and completeness of its reaction with diazo-compounds in dilute solution.

"Extinction coefficients" or "transmissive indices" (Priest, *J. Opt. Soc. Amer.*, 1920, 186) are determined by the König-Martens spectrophotometer, using the light from a mercury vapour lamp. Standard figures for aniline, *o*- and *p*-toluidines, α - and β -naphthylamines, benzidine, anthranilic acid, methyl anthranilate, and *p*-sulphanilic acid are given. The dye from methyl anthranilate is saponified in alkaline solution and direct determinations of its concentration are impossible. In this case the velocity coefficient of saponification is determined; and for purposes of estimation photometric readings are taken at intervals, and the original concentration is determined by calculation. The presence of small quantities of the nitroso-compound of α -naphtholsulphonate has a negligible effect on the transmissive index. Hydrazine may be used in certain cases (for example, β -naphthylamine) to remove excess of nitrous acid, but in others (for instance naphthionic acid) the results obtained are uncertain. Details of the method for estimating amino-impurities in dyes are given. The process may also be used for the estimation of nitrites. A. G. P.

Pernitric Acid as an Analytical Reagent. IV. TRIFONOV (*Z. anorg. Chem.*, 1922, 124, 136—139).—Pernitric acid reacts with aniline, yielding an intensely yellow product. The per-acid is formed by the action of hydrogen peroxide on an acidified solution of a nitrite. The above reaction with aniline can therefore be employed for the detection of hydrogen peroxide or a nitrite. Benzene is coloured by pernitric acid yellow, which changes into dark red on the addition of alkali. Toluene and xylene are also coloured yellow but fainter, which, however, does not change on the addition of alkali. Thus benzene in the presence of toluene, xylene, and the aliphatic hydrocarbons can be detected; the aliphatic members are not coloured. Similarly, benzene can be detected in methylated spirits. W. T.

The Analysis of β -Naphthylamine. H. R. LEE and D. O. JONES (*J. Ind. Eng. Chem.*, 1922, 14, 961—963).—Methods for the determination of β -naphthylamine, m. p. 110.1—110.2° (corr.), in the presence of its common impurities, β -naphthol, m. p. 121.2—121.3° (corr.), $\beta\beta$ -dinaphthylamine, m. p. 172.2° (corr.), and α -naphthylamine, m. p. 49.2—49.3° (corr.), are suggested, using specially purified samples of these substances. The direct nitrite titration method usually gives results which are too high, due partly to oxidation of the diazo- β -naphthylamine and partly, when β -naphthol is present, to formation of nitroso- β -naphthol. A more accurate method, based on the fact that β -naphthol-1-sulphonic acid does not absorb nitrous acid, consists of sulphonating 0.65 gram of the sample of β -naphthylamine with 20 c.c. of 25% of fuming sulphuric acid at 0—5° and finally at 15°, diluting the mixture with ice, boiling it to remove sulphur dioxide and afterwards adding 15 c.c. of concentrated hydrochloric acid and estimating the β -naphthylamine by titration with 0.1*N*-sodium nitrite

at 0—5°. For estimation of the β -naphthol present, the sample dissolved in dry benzene is saturated with hydrogen chloride, and after removal of β -naphthylamine hydrochloride by filtration, the β -naphthol in the filtrate is determined by titration with 0.05*N*-diazotised-*p*-nitroaniline solution. Impurities insoluble in acids are estimated by boiling 0.5 gram of the sample in 150 c.c. of 1.5% hydrochloric acid, filtering the solution through a Gooch crucible, washing the residue with hot water, and afterwards drying it to constant weight at 100°. Under these conditions, at least 10% of the β -naphthol passes into the filtrate, but only 0.10—0.20% of the $\beta\beta$ -dinaphthylamine is soluble, so that this may be extracted from the residue by means of solvents and weighed or analysed for nitrogen. Moisture is determined by drying 5 gram of the finely powdered sample over concentrated sulphuric acid in a vacuum. A modified Kjeldahl-Gunning method for the estimation of total nitrogen is described and the melting-point curves of the binary systems, β -naphthylamine- β -naphthol, β -naphthylamine- $\beta\beta$ -dinaphthylamine, and β -naphthylamine- α -naphthylamine are also given.

A. J. H.

Identification of Alanine by Crystallo-chemical Analysis.

IV. S. JAITSCHNIKOV (*J. Russ. Phys. Chem. Soc.*, 1920, **52**, 145—147).—Polarimetric identification of active alanine is sometimes impossible owing to the marked opalescence of the solution. In such case, use may be made of crystallographic measurements. The results of the author's measurements are compared with those of Groth and of Fischer (A., 1906, i, 145).

T. H. P.

Detection of Urea. Estimation of Proteolytic Ferments.

LUDWIG PINCUSSEN (*Biochem. Z.*, 1922, **132**, 242—244).—Urea is best detected in a liquid free from ammonia by addition of urease at P_H 7.0—7.2, keeping at 55° for fifteen minutes, then addition of two drops of saturated sodium carbonate solution and detection of the production of volatile ammonia by litmus paper. If the liquor contains ammonia originally, it is removed by shaking with powdered permutite, 0.2 gram per c.c.

A quantitative method for estimating proteolytic ferments is described which depends on the micro-Kjeldahl estimation of the non-protein nitrogen formed by the digestion. Unchanged protein is removed by colloidal iron.

H. K.

Detection of the Veronal Group. Diagnosis of Veronal Intoxication. HEINRICH HANDORF (*Z. ges. exp. Med.*, 1922, **28**, 56—80; from *Chem. Zentr.*, 1922, iv, 576).—A modification of the murexide test is described for the detection of barbituric acid derivatives in urine. The different members of the veronal group are distinguished by their behaviour when the reaction is carried out in the presence of ammonium chloride, sodium chloride, and barium chloride, respectively.

G. W. R.

Inconstancy of the Precipitation of Uric Acid from Urine in the Form of Cuprous Urate. J. KHOUBI (*Bull. Soc. Chim. Biol.*, 1922, 4, 375—378).—Denigès's method for the estimation of uric acid in urine gives low and irregular results owing to the incomplete precipitation of the copper salt. This may be remedied to some extent by first treating the urine (100 c.c.) with five drops of sulphuric acid and heating for one hour at 100°. E. S.

Use of the Newer Indicators in Titrations of Alkaloids. WILLIAM J. MCGILL (*J. Amer. Chem. Soc.*, 1922, 44, 2156—2160).—The Sørensen values of aqueous solutions of the hydrochlorides of quinine, cinchonine, morphine, atropine, and strychnine have been determined from *E.M.F.* determinations at various dilutions; measurements were also made in the presence of such quantities of alkali salts as would normally be present in the titration of an alkaloid hydrochloride by alkali. From the results, the most suitable indicators for the titration of such alkaloids are deduced and tested experimentally. It is shown that the indicators generally used in the titration of morphine, atropine, and quinine give results which are not nearly so accurate as should be obtained. Thus in the case of morphine the average error using methyl-red is 1.4%, whereas bromophenol-blue gives an average error of 0.5%, and in some cases it is as low as 0.1%. The most trustworthy indicator for quinine is bromocresol-purple, whilst this indicator is not superior to methyl-red in the titration of cinchona residues. Cocaine and methyl-red give low results with morphine, whilst bromophenol-blue gives a satisfactory value and end-point; the same is also true for atropine and mydriatic residues. Methyl-red can best be used in the titration of strychnine. J. F. S.

The Estimation of Carnosine in Muscle Extract. GEORGE HUNTER (*Biochem. J.*, 1922, 16, 640—654).—Further details are given for the estimation of carnosine in muscle by the application of Koessler and Hanke's diazo-method (this vol., ii, 328). Histidine can be estimated in muscle in the presence of carnosine by utilising Knoop's reaction, since the reaction is not characteristic of carnosine. This fact affords, further, the opportunity of estimating carnosine by a new method, namely, by hydrolysing it and determining the resulting histidine by Knoop's reaction. In ox-muscle, purines are responsible for about 3% of the diazo-colour value reckoned as carnosine, and 2% of the colour may be due to substances other than carnosine. The carnosine content of muscle varies both with the species of the animal and with the individual of the same species. S. S. Z.

Knoop's Test for Histidine. GEORGE HUNTER (*Biochem. J.*, 1922, 16, 637—639).—Different intensities of colour are produced in solutions of the same concentration of histidine by Knoop's test. This is due to the variation in the excess of bromine. The proportion of three atoms of bromine per molecule of histidine is found to give the maximum colour on heating. S. S. Z.

Precipitation of Proteins by Metaphosphoric Acid. Application to the Analysis of Blood, Pathological Liquids, and Cerebrospinal Fluid. A. GRIGAUT and P. ZIZINE (*Bull. Soc. Chim. Biol.*, 1922, 4, 388—406).—In the precipitation of proteins from serum, etc., by means of sodium metaphosphate and hydrochloric acid, the amount of acid employed, provided it is within the limits necessary for the complete precipitation of protein, has little influence on the amount of non-protein nitrogen which passes into the filtrate. Preliminary dilution of the serum, on the other hand, exerts considerable influence; the greater the dilution the smaller the amount of non-protein nitrogen which passes into the filtrate. For serum and pathological liquids the following method is recommended: The serum (10 c.c.) is diluted with water (6 c.c.). A 20% solution of sodium metaphosphate (2 c.c.) is then added, followed by 2*N*-hydrochloric acid (2 c.c.). In the case of certain liquids (for instance, whole blood, cerebrospinal fluid) modifications depending on the viscosity and content in protein must be made both in the dilution and the amount of reagents employed. Contrary to the statement of Cristol (this vol., ii, 583), metaphosphoric acid does not produce a partial hydrolysis of the proteins; the higher values obtained with this reagent are probably due to the presence of polypeptides in the filtrate. When trichloroacetic acid is employed these are retained by the proteins in the form of complexes owing to the high acidity of the solution. Metaphosphoric acid has been used for the removal of proteins in the estimation of urea, uric acid, non-protein nitrogen, and dextrose in blood and other fluids. E. S.

The Gradual Darkening of Hæmatin Solutions in Colorimetric Estimations, and its Prevention. H. C. GRAM (*Acta med. Scand.*, 56, 52—70; from *Chem. Zentr.*, 1922, iv, 576).—The darkening of hæmoglobin solutions is not attributed to change of oxyhæmoglobin to hæmatin. It is affected by strength of acid and to a less degree by temperature. It may be prevented, in the Autenrieth method, by addition of 2 c.c. of 3% hydrogen peroxide to 98 c.c. of the hydrochloric acid used. The wedge used for comparison must be specially graduated. The modification is not applicable to the Sahli method. G. W. R.

Estimation of Pepsin. MAKI TAKATA (*Tohoku J. Exptl. Med.*, 1921, 2, 127—130).—One gram of magenta-fibrin is added to gastric juice and the mixture incubated for thirty to forty-five minutes at 38°, then chilled, centrifuged, and the clear liquid compared colorimetrically with a magenta standard solution.

CHEMICAL ABSTRACTS.

Estimation of Total Protein, of Protein and Non-protein Nitrogen in Blood Plasma. H. BERRY and L. MOQUET (*Compt. rend. Soc. biol.*, 1922, 87, 329—331).—The authors found that in estimating the total proteins in blood by precipitation by heat and weighing the dry precipitate, attention must be directed to the hydrogen-ion concentration of the solution. The amount of precipitate and the quantity of nitrogen contained in it varies, the

maximum being obtained at $p_H=5.6$. Their own procedure is to neutralise 2 c.c. of plasma, diluted to 20 c.c. with water, either with 0.01*N*-hydrochloric acid or 0.1*N*-acetic acid, one drop of alizarin being used. This is then boiled for fifteen minutes, and the precipitate collected and dried. The nitrogen is estimated by the regular Kjeldahl method in a portion of the precipitate. The alternative procedure is to measure into a tall cylinder 1 c.c. of plasma, add to it 3 c.c. of acetone free from hydrogen sulphite and leave the precipitate to settle for twelve hours. This is then filtered, washed with acetone, boiling water, boiling alcohol, and finally with ethyl ether. The quantitative results are the same by both methods.

CHEMICAL ABSTRACTS.

Estimation of the Total Non-protein-nitrogen of Serum. Choice of a Suitable Albumin Precipitant. P. CRISTOL and M. SIMONNET (*J. Pharm. Chim.*, 1922, [vii], 26, 298—309).—Whilst trichloroacetic acid and tungstic acid are excellent precipitants for albumin in the estimation of the total non-protein-nitrogen of serum, and allow neither lipoid- nor protein-nitrogen to pass into the filtrate, metaphosphoric acid, on the contrary, is unsuitable as complete precipitation is not obtained and the results are always correspondingly high when estimation of total non-protein-nitrogen is in question, although the defecation with this precipitant is sufficiently good for carbohydrate or lactic acid estimations. Of all the precipitants tried, trichloroacetic acid is the best and most convenient, as the estimation of the non-protein-nitrogen can be carried out with as little as 2 c.c. of filtrate, whereas with tungstic acid 5 c.c. of filtrate are required. The evaporation and hydrolysis are more rapid and the estimation more exact. Further, any losses due to too violent ebullition are not to be feared. G. F. M.

Examination of the Duodenal Fluid obtained through a Tube. R. DAMADE (*Compt. rend. Soc. biol.*, 1922, 86, 947—948).—With a 2% solution of methyl-orange the alkalinity of the duodenal fluid was found to vary from 1.4 to 12.5 c.c. of *N*-hydrochloric acid for 10 c.c. Amylase is estimated by determining the amount, in milligrams, of dextrose found in 5 c.c. of 2% starch solution incubated at 37° for one hour with 1 c.c. of the duodenal fluid. This value generally varies from 15 to 35. For the estimation of lipase, 10 c.c. of a 10% solution of ethyl butyrate is treated with 1 c.c. of the duodenal fluid and neutralised to phenolphthalein with 0.1*N*-sodium carbonate. After incubating the mixture at 37° for one hour, the free acid formed is titrated. Usually 1.2—3.2 c.c. are required. For the estimation of trypsin, 1 c.c. of the fluid is added to 50 c.c. of 5% gelatin solution; this is then neutralised to phenolphthalein and kept at 37° for one hour, 20 c.c. of the digested mixture being titrated with 0.1*N*-sodium hydroxide solution, whilst another portion of 20 c.c. is titrated after the addition to it of 5 c.c. of formaldehyde solution and 5 c.c. of 90% ethyl alcohol. The tryptic activity is expressed by the sum of the volumes of sodium

hydroxide solution required; the value varies between 4.9 and 7.5, with an average of 5.35 c.c.

CHEMICAL ABSTRACTS.

Detection of Urobilin in Urine. GEORGES RODILLON (*J. Pharm. Chim.*, 1922, 26, 379—381).—An alcoholic solution of zinc acetate is prepared by mixing 15 grams of zinc oxide with 250 c.c. of 95% alcohol and 15 c.c. of glacial acetic acid, and, after frequently shaking during forty-eight hours, filtering the clear liquid. Equal volumes of this reagent and the urine are mixed in a test tube, 1 drop of a 1 in 10 dilution of tincture of iodine is added and a tenth of the total volume of chloroform. After shaking, the chloroform layer is allowed to separate, and if necessary is clarified by gentle warming. In presence of urobilin, an emerald-green fluorescence will be observed when a beam of light is directed on to the chloroform layer. When considerable quantities of urobilin are present, the chloroform solution will also show a rose-coloured tint.

G. F. M.

Mechanical Analysis of Humus Soils. GILBERT WOODING ROBINSON (*J. Agric. Sci.*, 1922, 12, 287—291).—The mechanical analysis of peaty soils by the usual methods is frequently of little value, owing to the cementing together of the finer soil particles by the organic colloids.

The removal of the latter by means of hydrogen peroxide causes a considerable increase in the clay fraction, which by microscopical examination is shown to consist almost entirely of mineral particles.

Ten grams of soil are heated for thirty minutes on a steam-bath with 50 c.c. of hydrogen peroxide (20 vols.). A further treatment with 25 c.c. of peroxide is usually sufficient to bring about a thorough dispersion of the soil particles.

A. G. P.

A New Method for the Mechanical Analysis of Soils and Other Suspensions. GILBERT WOODING ROBINSON (*J. Agric. Sci.*, 1922, 12, 306—321).—The existing method of soil analysis is considerably shortened by the means suggested. A suspension of soil (2%) in dilute sodium carbonate solution (about 0.025%) is placed in a cylinder and samples are withdrawn by means of a pipette at measured depths and after known periods of settling. The results are most conveniently represented by a graphical indication of summation percentages as a function of the logarithm of the settling velocity. By suitable choice of times and depths of sampling, figures corresponding with any of the systems of fractional sedimentation can be obtained. The diameter of the cylinder should not be less than 4 cm. Small errors in the depth of sampling have negligible effects on the final results.

The effect of a gel coating on the settling velocity of a particle is a reduction which is a simple function of the thickness of the gel. It is shown that below the first few centimetres of a settling column the change in concentration with depth is very small.

A. G. P.

INDEX OF AUTHORS' NAMES.

TRANSACTIONS AND ABSTRACTS.

1922.

(Marked T., and A., i and A., ii respectively.)

Anonymous.

grape oil from the Canadian vine
(*Vitis hederacea*), A., i, 97.

A.

Aarnio, B., absorption (by soil) of ammonium ions from solutions of ammonium salts and the effect of electrolytes thereon, A., i, 1227.

Abderhalden, Emil, are there protective enzymes against polysaccharides? A., i, 82.

alcoholic fermentation by means of yeast-cells under various conditions.

I. Influence of animal charcoal and other adsorbents on the course of the fermentation; formation of acetaldehyde, A., i, 92.

alcoholic fermentation by means of yeast-cells under various conditions. II., A., i, 92.

the isolation of amino-acids in blood, A., i, 287.

the rôle of vitamins in the chemistry of the cell, A., i, 607.

the composition of silk fibroin and its structure, A., i, 784.

Abderhalden, Emil, and *Andor Fodor*, functions of the yeast-cell; zymase and carboxylase action, A., i, 92.

Abderhalden, Emil, and *Ernst Wertheimer*, influence of substances obtained from yeast-cells and organs on the time course of the fission of substrates by polypeptidases, carbohydases, and esterases, A., i, 796.

Abderhalden, Emil, and *Bernhard Zorn*, the composition of the scales in psoriasis, A., i, 794.

Abel, Emil, the influencing of catalysts and specifically active catalysts, A., ii, 276.

Abelin, J., physiological action of proteinogenic amines. V., A., i, 610.

Aboulenc, Jean. See *Jean Baptiste Senderens*.

Abrahamson, Emanuel M. See *Marston Taylor Bogert*.

Abribat, Marcel, modifications of Kohlrausch's method for the measurement of the conductivity of electrolytes, A., ii, 344.

Acél, Desider, a micro-method for estimation of nitrogen, A., ii, 225.

Achard, Ch., A. Ribot, and *Léon Binet*, adrenaline hyperglycemia, A., i, 963.

Acree, Solomon Farley, R. R. Mellon, Pauline M. Avery, and *Edgar A. Slagle*, a stable single buffer solution, *p_H* 1 to *p_H* 12, A., ii, 157.

Acree, Solomon Farley. See also *R. R. Mellon*.

Acton, Hugh William, the cinchona alkaloids, A., i, 610.

[pharmacological action of] cinchona alkaloids, A., i, 899.

Adair, Gilbert Smithson. See *Walter Stiles*.

Adam, Matthew Atkinson, and *David Allston Legg*, production of butaldehyde and butyric acid therefrom, A., i, 222.

Adam, Neil Kensington, geometrical isomerism in unimolecular films, A., i, 424.

the properties and molecular structure of thin films. III. Expanded films, A., ii, 687.

Adam, William Gordon. See *William Lewcock*.

Adams, Chester E., and *Roger J. Williams*, laboratory preparation of acetaldehyde, A., i, 222.

Adams, E. T. See *Frederick S. Hammett*.

- Adams, Elliot Quincy**, rôle of hydrogen and hydroxyl-ion diffusion in nerve and muscle action, A., i, 1211.
- Adams, J. R.** See *Arthur W. Bull*.
- Adams, Roger**, the action of the Grignard reagent on thiocyanates, A., i, 531.
- Adams, Roger, M. F. Fogler, and C. W. Kreger**, the structure of disalicylaldehyde, A., i, 660.
- Adams, Roger, F. L. Roman, and W. N. Sperry**, the structure of the compounds produced from olefines and mercury salts; mercurated dihydrobenzofurans, A., i, 946.
- Adams, Roger.** See also *John R. Johnson, Oliver Kamm, Wilson D. Langley, Charles Shattuck Palmer, Armand J. Quick, and V. Voorhees*.
- Adickes, Franz.** See *Heinrich Wieland*.
- Adinolfi, Emilio**, absorption spectra of triphenylmethane colouring matters, A., ii, 601.
- Adkins, Homer**, selective activation of alumina for decarboxylation or for dehydration, A., ii, 834.
- Adrina, Homer, and A. C. Krause**, the action of alumina, titania, and thoria on ethyl and isopropyl acetates, A., i, 422.
- Adler, Erich.** See *Gustav Embden, and Solo Isaac*.
- Adler, Howard.** See *James Kendall*.
- Adler, Leo**, the function of the pancreas, A., i, 195.
- Adler, Oskar**, a reaction of wood and some observations on anethole, A., ii, 401.
- Adler, Oskar, and W. Wieschowski**, the physiological action of melanin acids, A., i, 498.
the formation of melanin from organic substances, A., i, 1170.
- Adolf, Mona, Wolfgang Pauli** [with *Franz Jandraschitsch*, physico-chemical analysis of aluminium oxy-salts and aluminium oxide sols, A., ii, 149.
- Adolph, Edward F., and Lawrence Joseph Henderson**, heat of reaction of oxygen with hæmoglobin, A., ii, 350.
- Agashe, Ganesh Sakharam**, obituary notice of, T., 745.
- Ahlberg, E.**, steric transformations with α -sulphonodialiphatic acids, A., i, 625.
- Ahlgren, Gunnar**, spontaneous reducing effect of muscle on methylene-blue; physiology of dehydrogenases, A., i, 792.
- Aiello, Giuseppe**, the distribution coefficients of diuretics and narcotics and the theory of narcosis, A., i, 301.
- Ainslie, D. S.** See *John Cunningham McLennan*.
- Airla, Yrjo.** See *Leonor Michaelis, and Peter Rona*.
- Akamatsu, S.**, action of anions, particularly the hydroxyl-ion, on the colloidal condition of night-blue, A., ii, 830.
- Åkerlöf, Gosta**, neutral salt action, at higher salt concentrations; the velocity of hydrolysis of ethyl acetate, and the hydrogen-ion activity of the catalyst, A., ii, 134.
- Aktien-Gesellschaft für Anilin-Fabrikation**, preparation of di- and poly-halogen substitution products of monohydric phenols, A., i, 1145.
- Albert, T. J.** See *Heinrich Wieland*.
- Albert, Talbot J., jun.**, tetra-substituted carbamides, A., i, 370.
- Albesco, G.** See (*Mme*) *Pauline Ramart-Lucas*.
- Aldrich, Martha.** See *Willey Denis*.
- Aldrich, Thomas Bailey, and Julia E. Blanner**, derivatives of trihalogen *tert*-butyl alcohols. IV. The benzoyl ester of tribromo-*tert*-butyl alcohol or brometone benzoyl ester, A., i, 910.
- Aldridge, (Miss) Joan, and Malcolm Percival Applebey**, the peroxidic compounds of copper, T., 238.
- Alessandri, Luigi**, singular formation of ketodinitrones and their behaviour, A., i, 558.
- Alken, Erika.** See *Fritz Mayer*.
- Allander, B.** See *Karl Gustav Denby*.
- Allen, D. M.** See *W. R. Bloor*.
- Allen, Eugene Thomas**, chemical aspects of volcanism with a collection of the analyses of volcanic gases, A., ii, 219.
- Allen, F. M., and Mary B. Wishart**, carbohydrate metabolism and diabetes. IV. Dextrose-nitrogen ratios in partially depancreatized dogs, A., i, 893.
- Allen, H. Newman**, periodic structure of atoms and elements, A., ii, 758.
- Alles, Richard.** See *Heinrich Wieland*.
- Allison, Vernon C., W. L. Parker, and G. W. Jones**, estimation of oxides of nitrogen [in air], A., ii, 313.
- Aloy, Jules, and E. Rodier**, action of light on uranium salts, A., ii, 337.
- Alsgaard, Peder Chr.**, electrolytic production of sodium perborate, A., ii, 375.
- Alterthum, Hans**, theory of recrystallisation, A., ii, 623.
- Altwegg, J., and D. Ebin**, 2:4-diketo-5-phenyl-5-ethyltetrahydro-oxazole, A., i, 53.
- Altwegg, J., and J. Landröven**, esters of ethylene halohydrins, A., i, 315.
ethyl esters of *p*- β -hydroxyethylamino-benzoic acid and of *p* bis- β -hydroxyethylaminobenzoic acid, A., i, 1022.

- Amadori, Mario**, equilibrium between sulphur and iodine in solution, A., ii, 561.
- Amantea, Giuseppe**, and **C. Kraskowsky**, crystallisation of hæmoglobin, A., i, 387.
- Amberger, Conrad**, and **Karl Bromig**, synthesis of fats (glycerides), A., i, 804.
- Aminoff, G.**, the radius of hydrogen atoms in crystals, A., ii, 496.
- Anagnostu, J. L.** See **Ludwig Pin-cussen**.
- Andersen, E.** See **Siegfried J. Thann-hauser**.
- Andersen, Erik Buch**, the synthesis of ammonia by collision with slow-moving electrons, A., ii, 562.
- Anderson, J. A.** See **Edwin Brown Fred**, and **W. H. Peterson**.
- Anderson, Paul**, some properties of hydrogen desorbed from platinum and palladium, T., 1153.
- Anderson, R. J.**, and **W. L. Kulp**, analysis and composition of maize pollen, A., i, 508.
- Anderson, V. G.** See **David Avery**.
- Anderson, William Theodore, jun.** See **LeRoy Wiley McCay**, and **Hugh Stott Taylor**.
- Ando, Kinji.** See **Yukichi Osaka**.
- Andouard, P.** See **A. Gouin**.
- André, Émile**, the actual state of the chemistry of the fats, A., i, 713.
the oils of grape seeds; the solid fatty acids; method of separation of stearic and palmitic acids, A., i, 908.
- André, Gustave**, the changes which oranges undergo on keeping, A., i, 209.
- Andrews, F. M.**, anthocyanin of *Beta vulgaris*, A., i, 96.
- Andrews, F. W.** See **Samuel G. Liver-sedge**.
- Andrews, James C.** See **James Kendall**.
- Andrews, T. M.** See **Alvin Sawyer Wheeler**.
- Angelescu, E.**, various oils of *Origanum vulgare* from different parts of Italy, A., i, 460.
- Angelescu, E.** See also **P. Leone**.
- Angeletti, A.** See **Michele Gua**.
- Angeli, Angelo**, diazo-compounds, A., i, 774.
cellulose nitrate, A., i, 922.
- Angeli, Angelo, Dino Bigiavi**, and **Gino Carrara**, azoxyphenols, A., i, 878.
- Angerer, E. von**, notes of spectrography, A., ii, 3.
- Annett, Harold Edward**, and **M. N. Bose**, estimation of meconic acid in opium, A., ii, 791.
- Anschütz, Richard**, and **Walter Claassen**, dialicylic or anhydrosalicylic acid and its transformation into xanthone-4-carboxylic acid, A., i, 456.
- Anschütz, Richard**, and **Siegfried Jaeger**, diglycollic acid or anhydro-glycollic acid, A., i, 428.
- Anselmi, S.** See **Arrigo Mazzucchelli**.
- Aoyama, Shinjiro.** See **Yoshiharu Murayama**.
- Apostolo, C.**, nitration of 5-iodo-2-nitro-phenetole, A., i, 335.
- Appelmans, R.**, effect of sugar on the production of indole, A., i, 794.
- Applebey, Malcolm Percival**, and **Robert Douglas Reid**, the isomerism of metallic oxides. I. Lead monoxide, T., 2129.
- Applebey, Malcolm Percival**, and **Sydney Herbert Wilkes**, the system ferric oxide-sulphuric acid-water, T., 337.
- Applebey, Malcolm Percival.** See also (*Miss*) **Joan Aldridge**.
- Appleyard, James Robert**, obituary notice of, T., 2898.
- Aptekmann, P.** See **J. V. Dubskey**.
- Arai, Kotoyoro**, choline as hormone of peristalsis, A., i, 970.
- Arai, Minoru**, the stimulatory action of amino-acid hydrochlorides on the pancreatic secretion, A., i, 297.
the bacterial degradation of *l*-leucine, A., i, 303.
- Araki, Chirô.** See **Heizaburô Kondô**.
- Arbuz, E.**, the phytin content of foods, A., i, 1102.
- Archibald, Ebenezer Henry**, and **William A. Gale**, the hydrolysis of platinum salts. II. Potassium platinibromide, T., 2849.
- Ariano, R.**, considerations on cooling and heating curves, A., ii, 470.
- Arinstein, Bernhard.** See **Carl Neuberg**.
- Ariola, V.**, toxicity of the metallo-albumins, A., i, 500.
toxicity of metallic powders, A., i, 500.
- Armenante, R.** See **Arrigo Mazzucchelli**.
- Armit, James Wilson**, and **Robert Robinson**, polynuclear heterocyclic aromatic types. I. Some indeno-quinoline derivatives, T., 827.
- Armstrong, Edward Frankland**, and **Thomas Percy Hilditch**, catalytic action at solid surfaces. VII. Influence of pressure on the rate of hydrogenation of liquids in the presence of nickel, A., ii, 41.
catalytic action at solid surfaces. VIII. Action of sodium carbonate in promoting the hydrogenation of phenol, A., ii, 756.

- Armstrong, Edward Frankland**, and **Thomas Percy Hilditch**, catalytic action at solid surfaces. IX. Action of copper in promoting the activity of nickel catalyst, A., ii, 757.
- Armstrong, Henry Edward**, enzyme action. XXIII. Homo- and heterolytic enzymes, A., i, 1078.
- Armstrong, Henry Edward**, and **Edward Victor Evans**, carbon monoxide in tobacco smoke, A., i, 1226.
- Arndt, Charles H.**, the growth of maize as affected by iron and aluminium salts, A., i, 1103.
- Arndt, Fritz** [with **E. Milde**, **F. Tschenscher**, (*Frl.*) **F. Bielich**, and **G. Eckert**], certain triazoles, A., i, 277.
- Arndt, Fritz**, **E. Milde**, and **F. Tschenscher** [with (*Frl.*) **F. Bielich**], ring closure with hydrazinedicarbonamides containing sulphur. II. Thiourazole, A., i, 375.
- Arndt, Kurt**, and **Ernest Hantge**, the electrolytic preparation of sodium perborate, A., ii, 569.
- Arnold, R.**, application of the biochemical method for characterising dextrose to the fruits of *Viburnum opulus* and to extracts of red *Cinchona* and of cola, A., i, 311.
- Arnold, W.**, the estimation of fatty acids based on their volatility in steam, A., ii, 324.
- Aron, Hans**. See **Richard Gralka**.
- Aron, Max**, the glycogen of the embryonic liver; the factor determining its formation, A., i, 702.
- Aronowaky, Alexander**. See **Hans Fringsheim**.
- Arrhenius, Olof**, clay as an ampholyte, A., i, 707.
adsorption of nutrients and plant growth in relation to hydrogen-ion concentration, A., i, 1097.
quantitative analysis by centrifuge, A., ii, 309.
- Arrhenius, Svante**, [theory of strong electrolytes], A., ii, 345.
- Asahina, Yasuhiko**, and **Michizô Asano**, spilanthol, the pungent principle of para cress. II., A., i, 505.
- Asahina, Yasuhiko**, and **Atsushi Fujita**, constitution of rutacarpine, A., i, 47.
anemonins, A., i, 946.
- Asahina, Yasuhiko**, and **Yoshitoyo Ishida**, oxidation of dihydroxystearic acid, A., i, 520.
- Asahina, Yasuhiko**, and **Yoshio Kondo**, 8-hydroxy-*o*-toluic acid, A., i, 657.
- Asahina, Yasuhiko**, and **Satoru Kuwada**, constitution of elsholtzic acid, A., i, 1047.
- Asahina, Yasuhiko**, and **Saburo Mithori**, oxidation of menthone by ferric chloride, A., i, 667.
- Asahina, Yasuhiko**, and **Toraji Shimidzu**, constituents of the seeds of *Pharbitis nil* chois. II., A., i, 506.
- Asahina, Yasuhiko**, and **Takao Tsukamoto**, hydration of caryophyllene, A., i, 845.
- Asakawa, Osamu**. See **Kozo Sakaguchi**.
- Asano, Michizô**, amides and anilides of some saturated fatty acids, A., i, 437.
- Asano, Michizô**. See also **Yasuhiko Asahina**.
- Aschan, Ossian**, colophenic acid, A., i, 221.
humoöcic acid, A., i, 714.
- Aschan, Ossian** [with **Nils Fontell**, and **P. E. Simola**], the resin acids of the Coniferae. V. The nitroso-chloride, nitrosite, and nitrosate of pinabietic acid and abietic acid [Levy]; constitution of abietic acid and abietine, A., i, 1152.
- Aschkenasi, Salo**, hydration of ions, A., ii, 253.
hydration of electrolytes, A., ii, 482.
- Asendorf, William F.** See **Frank Burnett Dains**.
- Asher, Leon**, and **Ernst Bernet**, physiology of the glands. LIII. Function of the spleen under normal and increased oxygen requirements, A., i, 491.
- Asher, Leon**, and **Francis H. Doubler**, physiology of the glands. XLIX. The respiratory interchange of the dog with extirpated spleen and diet deficient in iron, A., i, 286.
- Asher, Leon**, and **Walter Horrisberger**, the action of the thyroid gland hormone in phloridzin diabetes. XLVI., A., i, 293.
- Asher, Leon**, and **Chu Koda**, physiology of the glands. XLVIII. Experiments on the respiratory metabolism of the dog with extirpated spleen, A., i, 286.
- Asher, Leon**, and **Gengo Matsumo**, physiology of the glands. XLVII. The relations between the thymus, spleen, and bone-marrow, A., i, 298.
- Ashley, Julius Nicholson**. See **Archibald Edwin Goddard**.
- Askenasy, Paul**, and **Frithjof Grude**, action of nitrogen on mixtures of barium oxide and carbon at high temperatures, A., ii, 445.
- Aston, Francis William**, the isotopes of tin, A., ii, 650.
the mass-spectrum of iron, A., ii, 710.
the isotopes of selenium and some other elements A., ii, 844.

- Aston, Francis William, and R. H. Fowler**, some problems of the mass spectrograph, A., ii, 241.
- Astruc, A., E. Canals, and R. Bordier**, estimation of the alkaloids in extract of aconite, A., ii, 404.
- Astruc, A., and E. Renaud**, chloroform and pepsic digestion, A., i, 281.
- Astruc, A.** See also *F. Jadin*.
- Aszódi, Zoltán**, estimation of urea from the carbon dioxide produced by urease, A., ii, 536.
- Atack, Frederick William, and Charles William Soutar**, preparation of intermediates [3-chloro-2-aminoanthraquinone and 3-chloro-1-bromo-2-aminoanthraquinone] and a dyestuff of the anthraquinone series, A., i, 259.
- Atchley, Dana W.** See *Walter W. Palmer*.
- Aten, Adriaan Hendrik Willem**, electro-osmosis, A., ii, 114.
- Atkins, William Ringrose Gelston**, hydrogen-ion concentration of plant cells, A., i, 411.
relation of the hydrogen-ion concentration of the soil to plant distribution, A., i, 415, 509.
- Atkinson, C. P.**, apparatus for use in titrating intermediates with unstable diazo-solutions, A., ii, 237.
- Atkinson, Ethel, and Edith Olive Hasleton**, a qualitative test for tannin, A., ii, 793.
- Atkinson, H. V., David Rapport, and Graham Lusk** [with *James Evenden* and *G. F. Soderstrom*], animal calorimetry. XXII. The production of fat from protein, A., i, 966.
- Aub, Joseph C., and James H. Means**, the basal metabolism and the specific dynamic action of protein in liver disease, A., i, 193.
- Aubart.** See *Otto Diels*.
- Aubel, E.**, the attack of dextrose and levulose by the pyocyanic bacillus, A., i, 201.
- Aubry, Pierre**, detection of bismuth in urine, A., ii, 165.
- Auerbach, Friedrich, and E. Bodländer**, iodometric estimation of sugars, A., ii, 878.
- Auerbach, Rudolf**, substantive cotton dyeing, A., ii, 353.
- Augsted-Jensen, Halvard.** See *Wilhelm Steinkopf*.
- Auguste, C.** See *Michel Polonovski*.
- Aurén, Trycho E.-son**, absorption of X-rays in crystals, A., ii, 810.
- Ausenda, Camillo**, the carbamino-reaction of the blood-proteins and their alleged importance in the transport of carbon dioxide by the blood, A., i, 1209.
- Austin, Joshua Harold, Glenn E. Cullen, A. Baird Hastings, F. C. McLean, John P. Peters, and Donald D. van Slyke**, gas and electrolyte equilibria in blood. I. Technique for collection and analysis of blood, and for its saturation with gas mixtures of known composition, A., i, 1207.
- Austin, Joshua Harold.** See also *John P. Peters*, and *Donald D. van Slyke*.
- Austin, Percy Corlett.** See *Thomas Martin Lowry*.
- Auwers, Karl von**, [substitution processes], A., ii, 98.
the molecular refraction of aromatic hydrocarbons and "aromatic" carbon, A., ii, 98.
- Auwers, Karl von, E. Hilliger, and E. Wulf**, the capacity for migration of acyl residues in acyl-derivatives of the phenylhydrazones of hydroxyketones, A., i, 1192.
- Auwers, Karl von, and K. Hültenes**, 3-phenylindazole and 2-hydroxy-3-phenylindazole, A., i, 682.
- Auwers, Karl von, and Helene Jacobsen**, titrimetric and spectrometric analysis of keto-enol mixtures, A., ii, 168.
- Auwers, Karl von, and W. Jülicher**, transformation products of 1-methyl-1-trichloromethyl- $\Delta^{2,5}$ -cyclohexadiene-4-one, A., i, 841.
- Auwers, Karl von, and H. Kolligs**, molecular coefficients of refraction, A., ii, 174.
- Auwers, Karl von, and H. Lange**, halogenated indazoles and stereoisomerism of free indazoles, A., i, 684.
- Auwers, Karl von, and K. Saurwein**, 2-hydroxy-4:6-dimethylbenzaldehyde, hemimellitenol, and *iso*- ψ -cumenol, A., i, 1031.
- Auwers, Karl von, and H. Westermann**, the spectrochemistry of aliphatic dienes with conjugated double bonds, A., ii, 97.
- Auwers, Karl von, and Karl Ziegler**, hydrocarbons of the semibenzene group, A., i, 119.
carboxylic acids of the semibenzene group, A., i, 139.
chloro- and bromo-derivatives of alkylated cyclohexadienones, A., i, 144.
- Avasare, Mahadeo Dattatraya.** See *Kuverji Gosai Naik*.
- Avery, David, A. J. Hemingway, V. G. Anderson, and T. A. Read**, estimation of minute amounts of lead in water, with notes on certain causes of error, A., ii, 161.

- Avery, Pauline M.** See *Solomon Farley Acree*, and *R. R. Mellon*.
Ayer, J. B., and *H. E. Foster*, the estimation of total albumin in the spinal fluid, A., ii, 798.
Azadian, A., silicotungstic acid applied to the estimation of caffeine, A., ii, 237.

B.

- B.**, *H.*, characteristics and utilisation of beech nut oil, A., i, 96.
 nutmeg butter, A., i, 975.
Babcock, Harold D. See *Charles E. St. John*.
Baborovsky, Jiri, and *J. Velišek*, the hydration of the lithium-ion, A., ii, 816.
Bach, Alexis, and *Sophie Zubkova*, the ferment numbers of blood. I. Quantitative estimation of catalase, protease, peroxylase, and esterase in a drop of blood, A., i, 392.
Bach, Emerich. See *Peter Rona*.
Bachem, Carl, influence of lecithin on the excretion of veronal, A., i, 400.
 a new antiseptic with high iodine content (diethylene disulphide tetraiodide), A., i, 611.
Bachér, F. See *Richard Stoermer*.
Bachmann, Franz. See *Ernst Sieburg*.
Bacho, Ferruccio de, volumetric method for the estimation of hyposulphurous and sulphylic acids, A., ii, 311.
Bachster, Marcel, Pregl's solution, A., ii, 792.
Backer, Hilmar Johannes, and *J. V. Dubský*, the formation of salts of sulphocarboxylic acids. I. Cobaltous and cupric salts of sulphoacetic and α -sulphopropionic acids, A., i, 423.
Bacot, Arthur William, and *Arthur Harden*, vitamin requirements of *Drosophila*. I. Vitamins-B and -C, A., i, 493.
Baddiley, James. See *British Dyestuffs Corporation, Ltd.*
Badische Anilin- & Soda-Fabrik, production of alcohols, ketones, and the like [lithium formate, methyl alcohol, acetone, etc.], A., i, 218.
 introduction of arylamino-groups into aminoanthraquinones, A., i, 942.
 preparation of carbamide, A., i, 992.
Bäckström, Hans L. J., the solubilities of calcite and aragonite, A., ii, 849.
Bähr, Hans. See *Emil Knoevenagel*.
Bäjen, Wully. See *Erich Schmidt*.
Baer, Julius, localisation of the degradation of fats in the body, A., i, 490.
Baerts, Franz, the action of organo-magnesium compounds on nitriles, A., i, 817.
Baggesgaard-Rasmussen, H., and *Sven Werner*, alkylations. I. Alkylation of sodium sulphite, A., i, 104.
Baggesgaard-Rasmussen, H. See also *Chr. Winther*.
Bahlke, W. H. See *T. H. Rogers*.
Bailey, George C., and *Ralph S. Potter*, synthesis of indigotin from fumaric acid and aniline, A., i, 370.
Bailey, James R. See *Harry L. Lochte* and *De Witt Neighbors*.
Bailey, Kenneth C., direct synthesis of carbamide starting from carbon dioxide and ammonia, A., i, 815.
Bailey, Robert Arthur. See *Edward Charles Cyril Baly*.
Bailey, V. A. See *John S. Townsend*.
Bailleul, Gustav. See *Friedrich Meyer*.
Bailly, Octave, the action of epichlorohydrin on normal sodium phosphate in aqueous solution and the stability of a diglyceronmonophosphoric diester, A., i, 980.
Bain, Edgar C., and *James R. Withrow*, relative densities of alkali metal amalgams and mercury. II., A., ii, 145.
Baines, Harry, the mechanism of the bromination of phenol in aqueous solution, T., 2811.
Bak, Ada. See *I. M. Kolthoff*.
Baker, Herbert Brereton, change of properties of substances on drying, T., 568.
Baker, Julian Levett, and *Henry Francis Everard Hulton*, amylases of the cereal grains; the insoluble amylase of barley, T., 1929.
Baker, William E. See *Victor K. LaMer*.
Balaban, Isidore Elkanah, and *Frank Lee Pyman*, bromo-derivatives of glyoxaline, T., 947.
Balareff, D., the structure of pyrophosphoric acid. III., A., ii, 49.
 estimation of zinc as zinc pyrophosphate, A., ii, 228.
 suggestions for the construction of a periodic table on a wider basis, A., ii, 365.
 separation of phosphoric acid in qualitative analysis, A., ii, 525.
 behaviour of platinum on strong ignition, A., ii, 578.
 the estimation of sulphuric acid as barium sulphate; evidence of the existence of a complex barium sulphuric acid, A., ii, 864.
Baldwin, Wilmer. See *Alexander Lowy*.

- Bales, Sidney Hartley**, and **Stanley Arthur Nickelson**, hydrolysis of $\beta\beta$ -dichlorodiethyl sulphide; synthesis of divinyl sulphide and the preparation of a non-vesicant isomeride of $\beta\beta'$ -dichlorodiethyl sulphide, T., 2137.
- Ball, Nigel G.** See **Henry Horatio Dixon**.
- Ballauf, Fritz.** See **Hans Heinrich Schlubach**.
- Balls, A. K.** See **A. Elizabeth Hill**.
- Baltes, Max.** See **Wilhelm Strecker**.
- Baly, Edward Charles Cyril**, photosynthesis and the functions of pigments in the living plant, A., i, 307. catalysis with special reference to newer theories of chemical action. I. The radiation theory of chemical action. (3) A theory of chemical reaction and reactivity, A., ii, 628.
- Baly, Edward Charles Cyril**, and **Robert Arthur Bailey**, the equilibria in aqueous solutions of the alkali metal bisulphites, T., 1813.
- Baly, Edward Charles Cyril**, and **Herbert Maxwell Duncan**, the reactivity of ammonia, T., 1008.
- Baly, Edward Charles Cyril**, **Isidor Morris Heilbron**, and **Donald Pryre Hudson**, photocatalysis. II. The photosynthesis of nitrogen compounds from nitrates and carbon dioxide, T., 1078.
- Bancroft, Wilder Dwight**, mordants. I., A., ii, 551. mordants. II. Alumina, A., ii, 646. mordants. III. Chrome, A., ii, 822.
- Bandemar, Selma L.** See **Charles S. Robinson**.
- Banerji, B. C.**, and **Nitraton Dhar**, catalysis. XIV. The action of nitric acid on metals, and an example of a periodic reaction, A., ii, 756.
- Banerji, B. C.** See also **Phani Bhusan Ganguly**.
- Banus.** See **Garcia Banus**.
- Barbier, André.** See **Amé Pictet**.
- Barbour, Henry G.** See **Raymond L. Stehle**.
- Barcroft, Joseph, A. V. Bock**, **Archibald Vivian Hill**, **T. R. Parsons, W. Parsons**, and **R. Shoji**, the hydrogen-ion concentration and some related properties of normal human blood, A., i, 891.
- Barcroft, Joseph**, and **M. Nagahashi**, direct measurement of the partial pressure of oxygen in human blood, A., i, 890.
- Barendrecht, Hendrick Pieter**, the enzyme phosphatase-phosphatase, A., i, 67.
- Barham, G. Basil**, estimation of sulphur dioxide, A., ii, 520.
- Barker, E. F.**, carbon dioxide absorption in the near ultra-red, A., ii, 805.
- Barker, Thomas Vipond.** See **Gilbert Thomas Morgan**.
- Barlot, J.**, and (*Mlle*) **M. T. Brenet**, detection of fatty acids by the formation of their sodium uranyl salts, A., ii, 167.
- Barnett, Edward de Barry**, picolinoyl-aminoanthraquinones, A., i, 943.
- Barnet, Edward de Barry**, and **James Wilfred Cook**, pyridinium salts derived from some chloroacetylaminocompounds, T., 792. studies in the anthracene series. II., T., 1376.
- Barnett, Edward de Barry, James Wilfred Cook**, and **Herbert Henry Grainger**, studies in the anthracene series. III., T., 2059.
- Barnett, Edward de Barry**, and **L. J. Kay**, dinitrobenzil, A., i, 844.
- Barnett, W. Leigh**, syntheses with chloroacetyl chloride, A., i, 232.
- Barr, Guy**, and **Louis Leighton Bircumshaw**, viscosity of some cellulose acetate solutions, A., i, 232.
- Barratt, S.** See **Thomas Ralph Merton**.
- Barratt, Thomas**, and **J. W. Lewis**, the heat developed by the action of sodium hydroxide on cotton ("mercerisation"), A., i, 526.
- Barrenscheen, Hermann K.**, laevulosuria, A., i, 496.
- Barrenscheen, Hermann K.**, and **O. Weltmann**, a reaction of urea with *p*-dimethylaminobenzaldehyde, A., ii, 794.
- Barrenscheen, Hermann K.** See also **Hans Fischer**.
- Barrett, Gérald.** See **Charles Moureu**.
- Barrow, Fred, Evan Dalton Griffiths**, and **Edward Bloom**, *N*-oximino-ethers. II. *N*-Aryl ethers of 2:4- and 2:6-dinitrobenzaldoximes, T., 1713.
- Barry, Frederick**, maintenance of the adiabatic condition in calorimetry, A., ii, 478.
- Barsky, George.** See **Ralph H. McKee**.
- Barsky, Joseph.** See **Max Kahn**.
- Bart, Heinrich**, synthesis of aromatic arsinic acids, A., i, 1201. synthesis of aromatic arsinic acids by the interaction of isodiazo-compounds with the arsenite ion, A., i, 1202. two new syntheses of salvarsan base (*mm'*-diamino-*pp'*-dihydroxyarsenobenzene), A., i, 1202.
- Bartell, Floyd E.**, and **E. J. Miller**, adsorption of activated sugar charcoal. I., A., ii, 741.

- Bartell, Floyd E.**, and **L. B. Sims**, relation of anomalous osmosis to the swelling of colloidal material, *A.*, ii, 267.
- Bartels, R.**, and **Arnold Eucken**, equation of condition of nitrogen at small pressures and low temperatures, *A.*, ii, 117.
- Bartels, R.** See also **Herbert Freundlich**.
- Bartholomé, Walter.** See **Erich Schmidt**.
- Barthoux, J.**, minerals from near Oudjda, Morocco, *A.*, ii, 651.
- Bartlett, Edward P.** See **Theodore William Richards**.
- Bartlett, H. H.** See **W. Ellwood Cake**, and **Charles E. Sando**.
- Baru, R.** See **Benjamin Max Margosches**.
- Basch, E.** See **Robert Meyer-Bisch**.
- Bass, Lawrence W.** See **Oskar Baudisch**, and **Treat Baldwin Johnson**.
- Bassett, Henry**, and **Reginald Graham Durrant**, action of ammonium nitrate and of aqueous ammonia on copper; properties of cupric tetrammine nitrite and nitrate, *T.*, 2630.
- Batchelor, H. W.** See **P. L. Gainey**.
- Bate, Stanley Charles.** See **British Dyestuffs Corporation, Ltd.**
- Batta, G.**, and **H. Thyssen**, estimation of carbon in cast iron and steel by Corleis's apparatus, *A.*, ii, 395.
- Battegay, Martin**, and **J. Bernhardt**, the carbamides of anthraquinone, *A.*, i, 1041.
the urethanes of anthraquinone, *A.*, i, 1041.
- Battegay, Martin**, and **Ph. Brandt**, nitration of hydrocarbons in basic or neutral medium, *A.*, i, 1001.
- Battegay, Martin**, **Ph. Brandt**, and **J. Moritz**, action of light on 9-nitroanthracene, *A.*, i, 1001.
- Battegay, Martin**, and **J. Claudin**, the dibromoanthraquinones, *A.*, i, 43.
the dibenzoyldiaminoanthraquinones, *A.*, i, 43.
- Battegay, Martin**, and **G. Hugel**, 3:6-tetramethyldiaminocyanoselenopyronin, *A.*, i, 669.
- Battelli, Fred**, and (*Mlle*) **Lina Stern**, mechanism of action of oxidising and reducing ferments, *A.*, i, 1077.
- Batuecas, T.**, the compressibility at 0° and less than 1 atmosphere and the divergence from Avogadro's law of several gases. II. Ethylene, *A.*, ii, 618.
- Batuecas, T.** See also **Philippe Auguste Guye**.
- Bau, Arminius**, fermentation without yeast, *A.*, i, 807.
the oxalic acid content of early spring leaves, and some observations concerning this acid, *A.*, i, 369.
- Baudisch, Oskar**, mechanism of reduction of nitrates and nitrites in processes of assimilation, *A.*, i, 194.
- Baudisch, Oskar**, and **Lawrence W. Bass**, iron as photochemical catalyst. I. The decomposition of potassium ferrocyanide in daylight, *A.*, i, 993.
- Baudisch, Oskar**, and **Harry J. Denel**, acetol. I. A new test for carbohydrates, *A.*, ii, 664.
- Baudisch, Oskar**, and **Treat Baldwin Johnson**, new method for the detection of thymine, *A.*, ii, 238.
- Baudisch, Oskar.** See also **Harry J. Denel**, and **Treat Baldwin Johnson**.
- Baudouin, A.**, and **H. Bénard**, a microcolorimeter and nephelometer, *A.*, ii, 862.
- Baudrenghien, J.**, Δ^2 -butene- γ -ol [α -methylallyl alcohol], *A.*, i, 710.
- Bauer, Edouard.** See **Albin Haller**.
- Bauer, F. C.**, and **A. R. C. Haas**, the effect of lime, leaching, form of phosphate, and nitrogen salt on plant and soil acidity, and the relation of these to the feeding power of the plant, *A.*, i, 975.
- Bauer, K. Hugo**, and **R. Hardegg**, perilla oil, *A.*, i, 983.
- Bauer, K. Hugo**, and **K. Herberts**, china wood oil, *A.*, i, 806.
- Bauer, K. Hugo**, and **Fritz Werner**, the condensation of α -halogeno-ketones with aldehydes, *A.*, i, 1034.
- Bauer, Lawson H.** See **J. Volney Lewis**.
- Baughman, Walter F.**, and **George Samuel Jamieson**, the chemical composition of maize oil, *A.*, i, 414.
- Bauman, L.**, and **L. M. Keeler**, estimation of uric acid in blood, *A.*, ii, 796.
- Baumann, K.**, barium in the viscera, *A.*, i, 898.
- Baumgarten, Paul.** See **Helmut Scheibler**.
- Baur, Emil**, and **Haggenmacher**, photolysis of uranyl oxalate, *A.*, ii, 338.
- Baur, Emil**, and **Eugen Hersfeld**, fermentation without yeast, *A.*, i, 93.
rennet coagulation of milk as a stimulated process, *A.*, i, 284.
peptone fermentation, *A.*, i, 1097.
- Baur, Emil**, and **Adolf Rehmman**, photolysis of uranyl oxalate and uranyl acetate, *A.*, ii, 337.

- Baur, Fr.**, and **E. Oppenheimer**, theory of the retention and excretion of absorbed bromine salts, and the halogen content of the organism, *A.*, i, 968.
- Baxter, Gregory Paul**, [revision of the atomic weight of lanthanum], *A.*, ii, 298.
- Twenty-eighth Annual Report of the Committee on Atomic Weights; determinations published during 1921, *A.*, ii, 366.
- significance of the density of hydrogen bromide with reference to the atomic weight of bromine, *A.*, ii, 370.
- purity of atomic weight silver. II. Solid impurities, *A.*, ii, 377.
- Baxter, Gregory Paul**, and **Leon Woodman Parsons**, purity of atomic weight silver. I. Gases in pure silver and iodine, *A.*, ii, 376.
- Baxter, Gregory Paul**, and **A. F. Scott**, atomic weight of boron, *A.*, ii, 285.
- Bayer, Gustav**, the change of glycogen under the influence of light, *A.*, i, 323.
- Bayeux, Raoul**, subcutaneous absorption of oxygen in mountain climbing and aviation, *A.*, i, 79.
- Bazzoni, Charles Blizard**. See **Owen Willans Richardson**.
- Beaver, J. J.** See **James Kendall**.
- Bechhold, Heinrich**, and **F. Hebler**, nephelometry of coloured hydrosols, *A.*, ii, 652.
- nephelometric effect of colloidal systems of different size of particles, *A.*, ii, 693.
- turbidity standard, *A.*, ii, 693.
- Beck, Georg**. See **Israel Lifschitz**.
- Beck, R. Ph.**, the electromotive properties of magnesium and the potentiometric and thermal analysis of the system magnesium-mercury, *A.*, ii, 545.
- Bečka, J.**, refractometric and interferometric quantitative analysis, *A.*, ii, 715.
- Becker, Aug.**, the durability of radium solutions, *A.*, ii, 810.
- Becker, Alfred Gustav**. See **Edmund Speyer**.
- Becker, A. L.** See **R. A. Sawyer**.
- Becker, Ernst**, identification of oxycellulose by means of the barium compound, *A.*, ii, 94.
- Becker, Ernst**. See also **Carl Gustav Schwalbe**.
- Becker, Günther**. See **Edmund Speyer**.
- Becker, K.**, and **Willi Jancke**, Röntgen spectroscopic investigation of organic compounds. I. and II., *A.*, ii, 128.
- Becker, Oscar**, and **Jocelyn Field Thorpe**, the intermolecular condensation of methyl ethyl ketone in the presence of calcium oxide, *T.*, 1803.
- Becker, Paul**. See **Georges Chavanne**, and **Herman Decker**.
- Beckmann, Ernst**, a sodium press for the laboratory, *A.*, ii, 369.
- Beckmann, Ernst**, and **Erich Correns** [with **Otto Liesche**], migration of the methyl residue into the benzene nucleus; the transformation of methylaniline hydrochloride into toluidine hydrochloride, *A.*, i, 535.
- Hofmann's degradation of acid amides to amines, *A.*, i, 544.
- Beckmann, Ernst**, and **Otto Liesche**, extension of ebullioscopy and its application to binary mixtures, *A.*, ii, 117.
- Beckmann, Ernst**, **Otto Liesche**, and **Fritz Lehmann**, physico-chemical characterisation of lignin from winter rye straw, *A.*, i, 233.
- Bédos, Pierre**. See **Marcel Godchaot**.
- Behre, E.**, and **A. Düring**, estimation of sucrose in presence of other sugars by means of alkaline-earth hydroxides, *A.*, ii, 790.
- Behre, C. H., jun.**, native antimony from Kern County, California, *A.*, ii, 76.
- Behre, Jeanette Allen**, and **Stanley Rossiter Benedict**, creatine and creatinine metabolism. IV. The question of the occurrence of creatinine and creatine in blood, *A.*, i, 789.
- Behrendt, Hans**, the influence of the chemical composition and the physico-chemical structure on the function of frog muscles, *A.*, i, 197.
- Beisler, Walter H.**, and **Lauder William Jones**, 1-hydroxylaminoanthraquinone and some of its derivatives, *A.*, i, 1166.
- Bell, James**. See **Emil Alphonse Werner**.
- Bell, James Munsie**, and **Joseph L. McEwen**, nitrotoluenes. VIII. Binary systems of *m*-nitrotoluene with another nitrotoluene, *A.*, i, 726.
- Belladen, L.**, the system $\text{Bi}_2\text{O}_3\text{-PbO}$, *A.*, ii, 777.
- Bellis, B.** See **G. C. Supples**.
- Bellussi, Angelo**, chemical reaction of phenol, *A.*, ii, 728.
- Benade, Walter**. See **Gustav Heller**.
- Bénard, H.** See **A. Baudouin**.
- Benda, Ludwig**, arsenical acridine compound, *A.*, i, 888.
- Benedetti-Pichler, A.**, quantitative micro-analysis of mixtures with special reference to organic ultimate analysis, *A.*, ii, 656.

- Benedicenti, Alberico, and S. Rebello-Alves**, electric kataphoresis of metallic protein compounds obtained by treatment with powdered metals, A., ii, 688.
- Benedicks, Carl**, the Le Chatelier-Braun principle, A., ii, 358.
- Benedict, Anber J.** See *Samuel Edward Sheppard*.
- Benedict, Stanley Rossiter**, estimation of uric acid in blood, A., ii, 405.
- Benedict, Stanley Rossiter, and Elizabeth Franke**, direct estimation of uric acid in urine, A., ii, 669.
- Benedict, Stanley Rossiter.** See also *Jennette Allen Behre, Thomas P. Nash, jun., and R. C. Theis*.
- Benelli, M.** See *P. Leone*.
- Bengtsson, Ernst.** See *Erik Hulthén*.
- Bennett, Alexander Hutcheon, and F. K. Donovan**, the estimation of aldehydes and ketones by means of hydroxylamine, A., ii, 535.
- Bennett, George Macdonald**, autoreduction of sulphurous acid, T., 1794. monothioethylene glycol, T., 2139.
- Benoist, Eugène.** See *Albin Haller*.
- Benrath, Alfred, W. Bücher, and H. Eckstein**, chloroiridates of complex metal bases, A., ii, 515.
- Benrath, Alfred, and K. Drekopf**, electrical conductivity of salts and mixtures of salts, A., ii, 109.
- Benrath, Alfred, and H. Espenschied**, thallic-thalious sulphate. I., A., ii, 504.
- Benrath, Alfred, F. Hess, and A. Obladen**, chemistry of actinometers, A., ii, 731.
- Benrath, Alfred, and J. Oberbach**, behaviour of Fehling's solution in light, A., i, 108.
- Benrath, Alfred, and A. Obladen**, the photochemical reduction of titanium salts, A., ii, 731.
- Bentivoglio, (Miss) Marie.** See *John Read*.
- Benson, B.** See *Gabriel Bertrand*.
- Berényi, Ludwig**, the calculation of adsorption of vapours at different temperatures, A., ii, 426.
- Beretta, A.** See *G. Charrier*.
- Bergdahl.** See *Bernhard Neumann*.
- Bergell, Peter**, new diamine compounds, A., i, 720.
the nature of specific poisons, A., i, 970.
- Berger, Ernst**, the reduction of oxides by hydrogen, A., ii, 508.
- Berger, Ernst, and G. Crut**, the equilibrium in the reduction of nickel chloride by hydrogen, A., ii, 38.
- Berger, Emily V.** See *Hamilton Perkins Cady*.
- Berger, G.** See *Frans Maurits Jaeger*.
- Berger, W.** See *R. Doerr*.
- Bergh, A. A. Hijmans van den**, sulphohæmoglobinæmia, A., i, 793.
- Berglund, Hilding.** See *Otto Folin*.
- Bergman, A. G.**, reactions of combination with conjugated systems of double linkings. I. Bromination of isoprene, A., i, 1106.
reactions of combination with conjugated systems of double linkings. II. Combination of hydrogen bromide with di-isopropenyl [β -dimethyl- Δ - γ -butadiene], A., i, 1106.
- Bergman, Stig.** See *Hans von Euler*.
- Bergmann, Max**, degradation of *d*-saccharic acid to the dialdehyde of *l*-tartaric acid, A., i, 7.
- Bergmann, Max, and Arthur Miekeley**, the unsaturated reduction products of sugars and their transformations. IV. A glucoside-like derivative of a simple hydroxyketone, δ -acetyl-*n*-butyl alcohol; the constitution of fructose, A., i, 618.
- Bergmann, Max, Herbert Schotte, and Wolfgang Lechinsky**, unsaturated reduction products of the sugars and their transformations. III. 2-Deoxyglucose (glucodesose), A., i, 227.
- Bergmann, Max, Reinhold Ulpts, and Francisco Camacho**, aldehyde compounds of hydroxy-amines and the partial acylation of these amines, A., i, 1180.
- Bergstrom, F. W.**, vapour pressure of sulphur dioxide and ammonia, A., ii, 423.
- Berkeley, Cyril**, an organic constituent of the tube of *Mesochatopterus Taylori*, Potts, A., i, 400.
occurrence of manganese in the tube and tissues of *Mesochatopterus Taylori*, Potts, and in the tube of *Chatopterus variopedatus*, Renier, A., i, 493.
- Berkmann, Sophie.** See *N. Igarischew*.
- Berl, Ernst**, the estimation of benzene in gases, A., ii, 591.
- Berl, Ernst, and W. Schwebel**, separation of volatile substances from gases which are absorbed with difficulty. II. Use of cresols, A., i, 932.
additive products between cresols and alcohols, etc., A., i, 932.
- Berlingozzi, Sergio.** See *Mario Betti*.
- Bernard, Henri.** See *Pastureau*.
- Bernardi, Alessandro, and G. Rossi**, action of carbon disulphide on mercuric acetate, A., i, 421.

- Berndt, W.** See *K. A. Schaller*.
- Bernet, Ernst.** See *Leon Asher*.
- Bernhard, Hermann.** See *Jakob Meisenheimer*.
- Bernhardt, J.** See *Martin Battagay*.
- Bernoulli, August L., and Heinrich Jakubowicz,** velocity of decomposition of mono- and di-substituted malonic acids, A., ii, 40.
- Bernoulli, August L., and Fritz Schaaf,** benzaldehyde-copper and the heterogeneous rate of formation of this substance, A., i, 1029.
- Bernton, A.,** diphenylphosphoric acid [diphenyl hydrogen phosphate], a reagent for the amino-group, A., i, 1146.
- Berry, Arthur John,** studies on thallium compounds. I. Analytical, T., 394.
- Berry, Erwin H.** See *Arthur E. Paul*.
- Bers, G. H. C. van.** See *A. Vürthelm*.
- Berthelot, Albert,** antiseptic action of acraldehyde, A., i, 903.
- Berthoud, Alfred,** the physical properties of sulphur trioxide, A., ii, 638.
- Bertiaux, L.,** rapid analysis of white metal and similar alloys, A., ii, 396.
- Berton, A. L.** See *Gustave Vavon*.
- Bertrand, Gabriel, and B. Benson,** the importance of zinc in the feeding of animals; experiments on mice, A., i, 893.
- Bertrand, Gabriel, Freundler, and (Mlle) Y. Ménager,** the variations in the chemical composition of seawater and the evaluation of the saline content, A., ii, 441.
- Bertrand, Gabriel, and M. Mokragatz,** the presence of cobalt and nickel in arable soil, A., i, 975.
- the occurrence of cobalt and nickel in plants, A., i, 1099.
- Bertrand, Gabriel, and (Mme) M. Rosenblatt,** the distribution of manganese in the organism of higher plants, A., i, 95.
- the variation in the manganese content of leaves with age, A., i, 411.
- Bertrand, Gabriel, and R. Vladesco,** the zinc content of the organs of the rabbit and of some vertebrates, A., i, 493.
- Bethe, Albrecht,** influence of the hydrogen-ion concentration on the permeability of dead membranes, on adsorption by protein sols, and on the metabolism of cells and tissues, A., ii, 352.
- Bettag, Ludwig.** See *Heinrich Wieland*.
- Betti, Mario, and Sergio Berlingozzi,** isomerism of the isooxazolecaboxylic acids. VI., A., i, 52.
- Beutler, Hans,** atomic linkings in the carbon compounds. I. and II., A., ii, 259.
- method of formation of the atoms in carbon compounds, A., ii, 348.
- Beyer, P. H.,** nitration, chlorination, and bromination of *m*-hydroxybenzoic acid, A., i, 37.
- Beyersdorfer, P.,** the explosions of varieties of organic dust; experimental investigation of the simple case of sugar dust, A., ii, 749.
- Beyne, Edgar,** apparatus for the gasometric estimation of zinc in zinc powder, A., ii, 85.
- Beyschlag, C.** See *Georg Schroeter*.
- Bhatnagar, Shanti Swarupa,** reversal of phases in emulsions and precipitation of suspensoids by electrolytes, A., ii, 204.
- pure aniline and water emulsions, A., ii, 268.
- Bichowsky, F. Russell,** equilibrium in the reaction between sulphur dioxide and water, A., ii, 270.
- Biedermann, W.,** the action of pepsin and trypsin on diastase, A., i, 480.
- regeneration of diastase and its dependence on oxygen, A., i, 698.
- the diastatic action of albumoses and amino-acids, A., i, 1200.
- Biedermann, W., and Amin Rueha,** enzymes. VIII. Conditions of action of amylases, A., i, 65.
- Bieler, E. S.** See *James Chadwick*.
- Bielich, (Fr.) F.** See *Fritz Arndt*.
- Bierry, Henri, and L. Moquet,** estimation of total protein, of protein and non-protein nitrogen in blood plasma, A., ii, 886.
- Bierry, Henri.** See also *Alexandre Desgrez*.
- Bigelow, Lucius A.,** oxidation of side-chains with potassium permanganate. II., A., i, 998.
- Bigiavi, Dino,** glyoximes and peroxides, A., i, 352.
- Bigiavi, Dino.** See also *Angelo Angeli*.
- Billmann, Einar, and Hakon Lund,** the quinhydrone electrode, A., ii, 111.
- Bijvoet, J. M., and A. Karsen,** use of Rontgen rays in determining the structure of the crystals of lithium and of some of its compounds with light elements. I., A., ii, 499.
- structure of crystals of lithium and some of its compounds with light elements. II. Lithium hydride, A., ii, 569.
- Billon, (Mlle) Hélène,** the action of trimethylene chlorobromide on some aliphatic ketones, A., i, 717.

- Billows, Edoardo**, tridymite from the Euganean Hills, Italy, A., ii, 578.
- Biltz, Heinrich**, 7:9-dialkyldeoxyuric acids. I., A., i, 380.
- Biltz, Heinrich**, and **Hans Bülow**, 7:9-dialkyldeoxyuric acids. II. 7:9-Dimethyldeoxyuric acid, A., i, 381.
- 7:9-dialkyldeoxyuric acids. III. 7-Methyl-9-ethyldeoxyuric acid, A., i, 382.
- 7:9-dialkyldeoxyuric acids. V. 1:7:9-Trimethyldeoxyuric acid, A., i, 383.
- 7:9-dialkyldeoxyuric acids. VII. 7:9-Diethyldeoxyuric acid, A., i, 384.
- 9-methyl-8-iodo- Δ^7 -isoxanthine, A., i, 384.
- Biltz, Heinrich**, and **Dorothea Heidrich**, 7:9-dialkyldeoxyuric acids. IV. 9-Methyl-7-ethyldeoxyuric acid, A., i, 382.
- 7:9-dialkyldeoxyuric acids. VI. 1:3:7:9-Tetramethyldeoxyuric acid, A., i, 383.
- Biltz, Heinrich**, and **Hans Paetzold**, the two modifications of glycine; methylation with diazomethane, A., i, 528.
- Biltz, Heinrich, Karl Seydel**, and **Edith Hamburger-Glaser**, decompositions occurring in the acylation of 5:5-diphenyltetrahydro-4-glyoxalone, A., i, 871.
- Biltz, Wilhelm**, constitution of haloids, A., ii, 370.
- some linear relationships in the periodic system, A., ii, 494.
- Biltz, Wilhelm**, and **Wilhelm Fischer**, the ammoniates of lead haloids; mixed compounds and the par-nit salts, A., ii, 851.
- Biltz, Wilhelm**, and **Georg Hohorst**, heat of formation of compounds of metallic magnesium with metallic zinc, cadmium, aluminium, and calcium, A., ii, 350.
- Biltz, Wilhelm**, and **Gustav F. Hüttig**, ammoniates of magnesium haloids, A., ii, 59.
- Biltz, Wilhelm**, and **Wilhelm Stollenwerk**, ammoniates of cupro- and thallo-haloids, A., ii, 67.
- Biltz, Wilhelm**, and **Arthur Voigt**, are tantalum and columbium pentachlorides conductors of electricity in the fused state? A., ii, 302.
- Biltz, Wilhelm**, and **Walter Wein**, the measurements of the densities of the haloids of aluminium with mercury as pyknometer liquid, A., ii, 505.
- Binaghi, R.**, *Digitalis purpurea* growing wild on the uplands of Desulo (Sardinia), A., i, 412.
- action of polyhalogenated compounds of methane and ethane on magnesiyl [magnesium alkyl] compounds. II., A., i, 1002.
- Binaghi, R.** See also **Bernardo Oddo**.
- Binapf, Joseph**. See **Rudolf Pummerer**.
- Binda, Pietro**, experimental toxic hæmatoporphyria, A., i, 89.
- Binder, F. Otto H.**, constitution and heats of combustion, A., ii, 27.
- Binet, Léon**. See **Ch. Achard**.
- Binkert, A.** See **Hermann Staudinger**.
- Binswanger, Friedrich**, effect of carbon dioxide on sugar in blood, A., i, 962.
- Birch, Stanley Francis**, and **Jocelyn Field Thorpe**, the formation and stability of spiro-compounds. IX. The influence on stability of groups of high molecular weight, T., 1821.
- Birkenbach, Lothar**. See **Otto Hönigschmid**.
- Bircumshaw, Louis Leighton**, the surface tension of mixtures of alcohol and water at 25°, T., 887.
- Bircumshaw, Louis Leighton**. See also **Guy Barr**.
- Bischof, Friedrich**. See **Paul Horrmann**.
- Bishop, Edna R.**, **Esther B. Kittredge**, and **Joel H. Hildebrand**, titrations in ethyl alcohol as solvent, A., ii, 308.
- Bishop, Gerald**, and **Oscar Lisle Brady**, 2:4-dinitrobenzil, T., 2364.
- Bishop, John Egbert**. See **Charles August Kraus**.
- Bistrzycki, Augustin**, and **Bruno Brenken**, sulphur as the bridge atom in the middle ring of a derivative of anthracene, A., i, 268.
- Bistrzycki, Augustin**, and **Franz Kuba**, nuclear condensations of ethers of phenylmercaptan. I., A., i, 33.
- Bittner, Karl**. See **Rudolf Pummerer**.
- Bjerrum, Niels**, chromithiocyanates. I., II., III., and IV., A., i, 18, 19, 20, 21.
- Bjäre, Alfred**, and **John Svensson**, the effect of adrenaline on healthy individuals, A., i, 970.
- Black, E. M.**, **M. Hupper**, and **J. Rogers**, the effects of adrenal feeding on the iodine content of the thyroid gland, A., i, 966.
- Black, Ivan Armstrong**, **Edmund Langley Hirst**, and **Alexander Kullen Macbeth**, the labile nature of the halogen atom in organic compounds. VI. The action of titanous chloride and of ammonia on representative halogen compounds, T., 2527.

- Blagden, John William.** See **Howard & Sons, Ltd.**
- Blair, Ethelbert William, and Thomas Sherlock Wheeler,** the oxidation of hydrocarbons, with special reference to the production of formaldehyde, A., i, 917.
improved form of gas-analysis apparatus, A., ii, 519.
- Blair, Ethelbert William.** See also **Thomas Sherlock Wheeler.**
- Blair, J. S., and J. M. Braham,** mechanism of guanidine formation in fused mixtures of dicyanodiamide and ammonium salts, A., i, 1123.
- Blaise, Edmond Emile, and (Mlle) Montagne,** action of thionyl chloride on α -hydroxy-acids, A., i, 520, 715.
- Blanc, Louis Gaston.** See **Maurice François.**
- Blanner, Julia E.** See **Thomas Bailey Aldrich.**
- Blatherwick, Norman R.,** blood fat in diabetes, A., i, 89.
- Blatherwick, Norman R., and M. Louisa Long,** urinary acidity. I. Some effects of drinking large amounts of orange juice and sour milk, A., i, 967.
- Blatt, A. C.** See **George W. Raiziss.**
- Blechta, F.,** trimethylene dinitrate, A., i, 518.
- Bleibtreu, Max,** isolation of the coagulating enzyme from blood-serum, A., i, 1084.
- Bleyer, Benno, and R. Seidl,** ultra-microscopic investigation of casein, A., i, 386.
casein from cow's milk, A., i, 479.
- Bloch, Ernst.** See **Peter Rona.**
- Bloch, Ignaz, and Fritz Höhn,** symmetrical dichlorodimethyl sulphide, A., i, 218.
- Bloedorn, W. A., and J. E. Houghton,** the rôle of hexamethylenetetramine in the production of hæmaturia, A., i, 1091.
- Bloom, G. J. son, Carl Gustaf Santesson, and Hans von Euler,** physiological researches on vitamin-B and water-soluble biocatalysts, A., i, 1218.
- Blom, Axel Viggo,** new derivatives of diphenylamine, A., i, 27.
preparation of nitroanisole from nitrochlorobenzene, A., ii, 38.
- Blondeau, Joseph,** some dialkylated benzyl cyanides [phenylacetoneitriles] and the corresponding alcohols, amides, amines, and acids, A., i, 654.
- Bloom, Edward.** See **Fred Barrow.**
- Bloomfield, J. J.** See **Sidney H. Kats.**
- Bloor, W. R.,** lipæmia, A., i, 89.
nephelometric method for the estimation of phosphoric acid and its compounds in small quantities of blood, A., ii, 84.
- Bloor, W. R., K. F. Pelkan, and D. M. Allen,** estimation of fatty acids (and cholesterol) in small amounts of blood plasma, A., ii, 593.
- Bloor, W. R.** See also **Elsie Hill.**
- Blüthorn, K.,** calcium and phosphoric acid metabolism with large doses of calcium and sodium phosphate, A., i, 83.
- Blumenthal, M.** See **W. D. Treadwell.**
- Blunt, Katharine, Alta Nelson, and Harriet Curry Oleson,** basal metabolism of underweight children, A., i, 83.
- Boas, Friedrich,** the part played by lipoids in the metabolism of plant cells. I. and II., A., i, 94, 613.
action of saponins on yeast-cells, A., i, 502.
- Book, A. V.** See **Joseph Barcroft.**
- Bock, Johannes, and Poul Iversen,** phosphate excretion in the urine during water diuresis and purine diuresis, A., i, 1090.
- Bodansky, Meyer,** zinc and copper content of the human brain, A., i, 194.
- Bode, G., and K. Hemmel,** the manganese content of potatoes, A., i, 415.
- Bodenstein, Max, Friedrich Böhs, (Frl.) Lindner, and Ramstetter,** formation and decomposition of the higher nitrogen oxides, A., ii, 372.
- Bodfors, Sven,** influence of substituents on various chemical reactions, A., ii, 698.
- Bodfors, Sven, and Per Frølich,** electrical colloid synthesis, A., ii, 826.
- Bodfors, Sven.** See also **Oscar Olsson Collenberg.**
- Bodine, J. H.,** anaesthetics and carbon dioxide output. I. The effect of anaesthetics and other substances on the production of carbon dioxide by certain orthoptera, A., i, 889.
- Bodländer, E.** See **Friedrich Auerbach.**
- Boecker, Eduard,** the distribution of quinine alkaloids in the animal organism, A., i, 896.
- Boedecker, Fr., and H. Volk,** unsaturated bile acids. III. The relationships of apocholic acid, dioxycholenic acid [m. p. 260°], and cholic acid to deoxycholic acid, A., i, 1027.
- Böhs, Friedrich.** See **Max Bodenstein.**
- Bösesen, Jacob,** dislocation theory of catalysis, A., ii, 755.

- Böeseken, Jacob**, and *C. de Graaff*, the configuration of the β -phenylglyceric acids and phenylglycidic acid, A., i, 551.
- Böeseken, Jacob**, *Chr. van Loon*, *H. G. Derx*, and *P. Hermanns*, condition of motion of molecules in space, A., ii, 210.
- Bögel, J.** See *Fritz Verzár*.
- Böhm, Karl.** See *Ernst Späth*.
- Böhme, Otto.** See *Karl Freudenberg*.
- Boehringer & Söhne, C. F.**, preparation of nitrosulphonic acids of hydrogenated cinchona alkaloids, A., i, 46.
preparation of aminosulphonic acids of hydrogenated cinchona alkaloids, A., i, 46.
- Boenheim, Felix**, chlorine metabolism in pulmonary tuberculosis, A., i, 192.
- Bönniger, Max**, the dextrose content of the red blood-corpuscles of man and their behaviour in isotonic sugar solutions, A., i, 606.
- Boeree, Alford Reginald.** See *Dalziel Llewellyn Hammick*.
- Boessneck, Werner.** See *Gustav Heller*.
- Bogdándy, Stefan von**, the estimation of chlorine in organic material, A., ii, 519.
- Bogert, Marston Taylor**, and *Emanuel M. Abrahamson*, thiazoles. I. Derivatives of 2-phenylbenzthiazole; synthesis of an analogue of cinchophen (atophan), A., i, 578.
- Bogert, Marston Taylor**, and *Yu-Gwan Chen*, selenium organic compounds. I. Synthesis of 4-seleno-2-methylquinazolone, 2-phenylbenzoselenazole, and some derivatives of the latter, A., i, 1182.
- Bogert, Marston Taylor**, and *Martin Meyer*, thiazoles. II. 1-p-Tolybenzothiazole, dehydrothio-p-toluidine and some related compounds, A., i, 868.
- Boggio-Lera, Enrico.** See *Arnaldo Piutti*.
- Bogitch, B.**, the expansions of some refractory materials at high temperatures, A., ii, 115.
- Bogue, Robert Herman**, swelling and gelation of gelatin, A., i, 388.
the sol-gel equilibrium in protein systems, A., i, 782.
the structure of elastic gels, A., i, 782.
viscosity of gelatin sols, A., ii, 122.
- Bohn, Raymond T.** See *George Leslie Kelley*.
- Bohr, Niels**, atomic structure, A., ii, 277.
- Bohr, Niels**, the structure of the atom and the physical and chemical properties of the elements, A., ii, 363.
the difference between series spectra of isotopes, A., ii, 598.
quantum theory of line spectra, A., ii, 801.
- Boiry, F.**, the vulcanisation of caoutchouc in solution, A., i, 848.
- Bokorny, Thomas**, behaviour of diastase and other enzymes under unfavourable conditions; action of some nitrogenous compounds on germination, A., i, 93.
urea as a nutrient of yeasts and other plants, A., i, 1096.
urea and hippuric acid as nutrient materials for plants, A., i, 1222.
- Bolland, A.**, a microvolumenometer, A., ii, 221.
- Bollman, J. L.** See *William H. Welker*.
- Bollmann, Martin.** See *Otto Fischer*.
- Bolte, Fr.** See *Julius Tröger*.
- Bonacker, Iris.** See *Walther Borsche*.
- Bone, William Arthur**, *Archibald Ramsden Pearson*, *Eric Sinkinson*, and *W. E. Stockings*, researches on the chemistry of coal. II. The resinic constituents and coking propensities of coals, A., ii, 285.
- Bonin, P.** See *L. Dede*.
- Bonino, G. B.**, action of potassium ferrocyanide on silver haloids, A., ii, 78.
- Bonis, A.**, volumetric estimation of lead peroxide in miniums; practical modification of Diehl's method, A., ii, 587.
- Bonis, A.** See *Ed. Moreau*.
- Bonnet, Eugène**, action of soluble lead salts on plants, A., i, 412.
- Bonnet, M.**, and *J. Haushalter*, detection of urea in tissue by means of xanthydriol, A., ii, 794.
- Bonnier**, the estimation of alkali carbonates in presence of phenolphthalein, A., ii, 869.
- Booberg, Gunnar.** See *J. Arvid Hedvall*.
- Boomer, E. H.** See *Otto Maass*.
- Boord, Cecil E.**, and *F. F. Cope*, the action of selenium monochloride on propylene, butylene, and amylene, A., i, 421.
- Boord, Cecil E.** See also *Alpheus W. Smith*.
- Bordier, R.** See *A. Astruc*.
- Boresch, Karl**, water-soluble colouring matters of the Schizophyceæ, A., i, 210.
- Borgström, Johan Henrik Leonard**, estimation of magnesium and alkalis in minerals, A., ii, 659.

- Bornemann, G.**, preparation of bright metallic sodium and potassium, or their alloys, in nitrogen, A., ii, 444. the composition of Scheele's green, A., ii, 852.
- Bornemann, Karl**, and **O. Hengstenberg**, the specific heats of some sulphides used in metallurgy, with special reference to high temperatures, A., ii, 618.
- Bornstein, A.**, and **Kurt Holm**, respiratory metabolism in alimentary glycaemia. I., A., i, 890.
- Bornstein, A.**, and **Johannes Kerb**, distribution of chloride and water after poisoning with mercuric chloride, A., i, 404.
- Bornstein, A.**, and **Elisabeth Müller**, respiratory metabolism and toxic glycaemia, A., i, 392.
- Bornstein, A.**, and **Robert Vogel**, action of pilocarpine on the composition of the blood, A., i, 80.
- Bornstein, A.** See also **Robert Vogel**.
- Borsche, Walther**, and **Iris Bonacker**, fission of γ -pyrone by aniline and *N*-phenyl-4-pyridone, A., i, 50.
- Borsche, Walther**, and **F. Hallwax**, the constitution of the bile acids. IV. Keductodehydrocholic acid, A., i, 1158.
- the constitution of the bile acids. V. Transformations from the cholic to the lithocholic acid series, A., i, 1159.
- Borsche, Walther**, and **Robert Meyer**, isatin and its derivatives. III. 5:8'-Di-isatylmethane and its conversion into quinoline derivatives, A., i, 53.
- isatin and its derivatives. IV. The action of hydrazine on isatin and 1-methylisatin, A., i, 53.
- deoxyindigotin, A., i, 55.
- Borsche, Walther**, **O. Weickert**, and **Robert Meyer**, the bile acids. III. Biloidanic acid [Letsche's acid], A., i, 255.
- Bosanquet, C. H.** See **William Lawrence Bragg**.
- Bose, M. N.** See **Harold Edward Annett**.
- Bosman, Louis Pierre**, castelin, a new glucoside from *Castela Nicholsoni*, T., 969.
- Bosse, O.** See **H. von Wartenberg**.
- Boushard, Heinrich.** See **Elektrochemische Werke G. m. b. H.**
- Bossi, A.** See **W. D. Treadwell**.
- Bossuet, Robert.** See **Pierre Jolibois**.
- Borwell, Mailland Crease**, and **H. E. Corman**, preparation of anhydrous formic acid, A., i, 712.
- Bothe, W.** See **Hans Geiger**.
- Botolfesen, Erling**, calcium carbide, A., i, 1107.
- calcium-ammonium, A., ii, 570.
- Boucher, Paul Edward**, measurement of the resonance, radiation, and ionisation potentials of several gases and vapours, A., ii, 608.
- Boudin, Paul.** See **Albin Haller**.
- Bougault, J.**, and **R. Gros**, new analytical applications of Nessler's reagent; characterisation of ketones; estimation of aldehydes, A., ii, 666.
- acetone in commercial ammonia, A., ii, 709.
- Boulanger, Ch.**, and **Georges Urbain**, composition of thortveitite from Madagascar, A., ii, 517.
- Bourcet, Paul.** See **Joseph-Marie-Alphonse Chevalier**.
- Bourgeois, L.**, a simple process for obtaining crystallised gypsum, A., ii, 292.
- Bourguignon, Georges**, and **Conduché**, the introduction of the iodine ion by electrolysis in a human being, and its elimination by the urine, A., i, 704.
- Bourlet, Claude W.**, and **Walter Thomas**, the thermos flask in the chemical laboratory, A., ii, 139.
- Bourne, W.** See **Raymond L. Stehle**.
- Bournigault, A.** See **A. Robin**.
- Boussu, Roger G.**, limit of inflammability of the vapours of the system, alcohol-petrol, and of a triple system with a basis of alcohol and petrol, A., ii, 557.
- Boutaric, Augustin**, and **M. Vuillaume**, flocculation of colloidal arsenic sulphide; influence of the concentration of the colloid, shaking, and temperature, A., ii, 498.
- Boutin, A. M.** See **André Sanfourche**.
- Boutot, Louis.** See **Paul Fleury**.
- Bowen, Edmund John.** See **Harold Brewer Hartley**, and **Cyril Norman Hinshelwood**.
- Bowen, Norman L.** See **George W. Morey**.
- Boyd, David Runciman.** See **Paul Jacobsen**.
- Bozorth, Richard M.**, crystal structure of potassium cyanide, A., i, 441.
- crystal structure of ammonium fluosilicate, A., ii, 499.
- crystal structure of cadmium iodide, A., ii, 851.
- Braden, Maria.** See **Emil Heuser**.
- Bradfield, Richard**, centrifugal method for preparing colloidal ferric hydroxide, aluminium hydroxide, and silicic acid, A., ii, 507.

- Bradford, Samuel Clement**, an explanation of Liesegang's rings, A., ii, 358.
- Bradley, Harold Cornelius**, autolysis. VIII. The nature of the autolytic enzymes, A., i, 896.
- Bradshaw, Geoffrey Gordon**, and **Arthur George Perkin**, derivatives of 2-hydroxybenzanthrone. I., T., 911.
- Brady, Oscar Lisle**, the nitration of m-nitrotoluene, T., 328.
- Brady, Oscar Lisle, James Nelson Edmund Day**, and **William Joseph Woodgate Rolt**, the dinitrotoluidines, T., 526.
- Brady, Oscar Lisle**, and **Gerald Patrick McHugh**, the alkylhydrazones, T., 1648.
- Brady, Oscar Lisle**, and **Clifford Dane Thomas**, the isomerism of the oximes. X. Cinnam- and nitrocinnam-aldoximes, T., 2098.
- Brady, Oscar Lisle**. See also **Gerald Bishop**.
- Braecke, (Mlle) Marie**, application of Bourquelot's biochemical method to the investigation of sugars and glucosides in some of the Scrophulariaceae, A., i, 1225.
- Braecke, (Mlle) Marie**. See also **Marc Bridel**.
- Bragg, (Sir) William Henry**, structure of organic crystals, A., ii, 128.
the crystal structure of ice, A., ii, 440.
the significance of crystal structure, T., 2766.
- Bragg, William Lawrence, R. W. James**, and **C. H. Bosanquet**, scattering of Röntgen rays by the atoms of a crystal, A., ii, 184.
the distribution of electrons around the nucleus in the sodium and chlorine atoms, A., ii, 703.
- Braham, J. M.** See **J. S. Blair**.
- Brakeley, Elizabeth**. See **James Kendall**.
- Brand, Kurt**, and **H. Collischonn**, pyrogallol 1:3-dimethyl ether. I., A., i, 452.
- Brand, Kurt**, and **Karl Otto Müller**, the diphen succindene series. IV. 9:12-Dichloro- Δ^8 ,¹¹-diphen succindadiene and diphen succind-10-ene, A., i, 444.
- Brand, Kurt**, and **Joseph Steiner**, the catalytic reduction of aromatic nitro-compounds and a new method for the preparation of β -arylhydroxylamines. I., A., i, 536.
- Brandtsma, W. F.**, equilibria and reaction velocities, A., ii, 699.
- Brandt, Erich**, ionisation and excitation tension of nitrogen, A., ii, 186.
- Brandt, Ph.** See **Martin Battagay**.
- Brann, Martin**, the action of digitalis, calcium, and barium on strips of heart muscle (Löwe) and the autagonistic influence of cocaine, magnesium, and potassium, A., i, 1213.
- Brass, Kurt**, and **Erwin Ferber**, amino- and anilino-phenanthraquinones, A., i, 355.
- Brass, Kurt**, and **Ludwig Köhler**, dibenzothianthrenediquinone and dinaphthathiophendiquinone (conversion of the dithium ring into the thiophen ring), A., i, 1050.
- Braun, Julius von**, constitution of Grignard's compounds, A., i, 22.
- Braun, Julius von** [with **K. Moldánke, H. Dirlam**, and **H. Gruber**], benzopolymethylene compounds. IV. The two *ar*-aldehydes of tetrahydronaphthalene [tetralin], A., i, 748.
- Braun, Julius von, Otto Braunsdorf**, and **Kurt Rätz**, relationships between constitution and pharmacological action in the cases of the benzoic and tropic esters of hydroxyalkylamines, A., i, 759.
- Braun, Julius von**, and **Heinrich Gruber**, benzopolymethylene compounds. V. Synthesis of α -anthrapyridine [naphthazine] from tetralin [tetrahydronaphthalene], A., i, 762.
- Braun, Julius von, Erich Hahn**, and **Jon Seemann**, benzopolymethylene compounds. III. Dehydrogenation of tetrahydronaphthalene, hydrindene, and tetrahydroacenaphthene [tetraphthene] derivatives, A., i, 728.
- Braun, Julius von**, and **Wilhelm Kaiser**, the two isomeric phthalyl chlorides, A., i, 658.
- Braun, Julius von**, and **Georg Kirschbaum**, catalytic hydrogenations under pressure in the presence of nickel salts. I. Indene and acenaphthene, A., i, 727.
- Braun, Julius von**, and **Annemarie Nelken**, preparation of dihydroisomindole, A., i, 863.
- Braun, Julius von**, and **Josef Weismantel**, dealkylation of mixed secondary bases by phosphorus chloride, A., i, 1150.
- Braunholts, W.** See **Hermann Staudinger**.
- Braunholts, Walter Theodore Karl**, a comparison of three isomeric carbocyanines, T., 169.
reactions of the phosphazines, T., 300.
- Braunholts, Walter Theodore Karl**, and **William Hobson Mills**, the cyanine dyes. VI. Dyes containing a quinoline and a benzothiazole nucleus; the thioisocyanines, T., 2004.

- Braunkolts, Walter Theodore Karl.** See also *William Hobson Mills*.
- Brauniser, G.** See *Carlo Gastaldi*.
- Brauns, Dirk Hendrik,** crystalline chlorotetra-acetylmannose, A., i, 433.
- Brauns, Fritz.** See *Karl Freudenberg*.
- Braunsdorf, Karl.** See *Erich Schmidt*.
- Braunsdorf, Otto.** See *Julius von Braun*.
- Brasier, Sidney Albert.** See *Douglas Frank Twiss*.
- Bredig, Georg, and J. Michel,** chemical kinetics of perchloric acid and its salts, A., ii, 359.
- Bredt, Julius,** camphene-*epi-sec.*-carboxylic acids [1:1-dimethyl-6-methylene-(1:2:2)dicycloheptane-3-carboxylic acid], their preparation from bornylene-*epi*-carboxylic acid, and their transformation into *sec.*- β -hydroxycamphane-*epi*-carboxylic acid [2-hydroxy-3-methyl-*meso*-dimethyl-(1:2:2)dicycloheptanecarboxylic acid] and δ -hydroxycamphane-*epi*-carboxylic acid [4-hydroxy-3-methyl-*meso*-dimethyl-(1:2:2)-di-cycloheptanecarboxylic acid] (*o*- and *p*-borneolcarboxylic acids), A., i, 937.
- Breisch, K.** See *K. Chalupny*.
- Brenet, (Mlle) M. T.** See *J. Barlot*.
- Brenken, Bruno.** See *Augustin Bistrzycki*.
- Brenner, Anton,** the analysis of aluminium alloys, A., ii, 319.
- Brenner, Wladar,** antagonistic ion effects, A., i, 907.
- Breuer, Paul Karl.** See *Franz Feist*.
- Brewster, Ray Q.** See *Jean Piccard*.
- Bresina, Jindřich,** the solution of tetramethylammonium aluminate, A., i, 638.
- Bridel, Marc,** the presence of a glucoside in the stems and roots of *Sedum telephium*, L., A., i, 799.
- the presence of the glucoside of an essential oil in the leafy stems and roots of *Sedum telephium*, Linn., A., i, 1225.
- Bridel, Marc, and (Mlle) Marie Braecke,** the presence of sucrose and aucubin in the seeds of *Melampyrum arvense*, L., A., i, 209.
- the presence of melampyritol and aucubin in the foliated stems of *Melampyrum arvense*, L., A., i, 799.
- rhinanthin and aucubin; rhinanthin is impure aucubin, A., i, 1168.
- the presence of aucubin and sucrose in seeds of *Rhinanthus crista-galli*, L., A., i, 1225.
- Briegleb, Karl.** See *Emil Fromm*.
- Briellmann, P.** See *Hans Rupe*.
- Briggs, A. P.,** a colorimetric method for the estimation of homogentisic acid in urine, A., ii, 534.
- colorimetric method for the estimation of small amounts of magnesium, A., ii, 659.
- modification of the Bell-Doisy phosphate method, A., ii, 718.
- Briggs, A. P., and Philip Anderson Shaffer,** the excretion of acetone from the lungs, A., i, 190.
- Brigl, Percy,** the partial replacement of the acid groups in β -penta-acetylglucose, A., i, 225.
- carbohydrates. II. A new anhydride ($\alpha\beta$) of glucose, A., i, 1117.
- Brigl, Percy, and Edgar Fuchs,** lignoceric acid and its derivatives, A., i, 712.
- Brillouin, Marcel,** the Bohr atom; the Lagrange theorem applied to electronic orbits, A., ii, 438.
- Brimmer, Karl.** See *Hugo Weil*.
- Briner, Emil, S. Niewiazski, and J. Wiswald,** oxidation of nitric oxide and the recovery of nitrogen oxides from mixtures with air, A., ii, 563.
- Briner, Emil, and (Mlle) A. Trampler,** the mechanism of catalytic action in the hydrolysis of fats, A., ii, 208.
- Brinkley, Stuart R.,** equilibrium in the system ammonia-mercuric cyanide, A., i, 724.
- Brinkman, R.,** osmotic resistance and phosphatides of the blood; new quantitative methods, A., i, 1086.
- Brinkman, R., and (Frl.) H. Wastl,** the biochemistry of the phosphatides and sterols. IV. The importance of the proportion of cholesterol-lecithin of the erythrocyte surface for the stability of suspensions of blood-corpuscles and for natural hæmolysis, A., i, 289.
- Brinkman, R.** See also *S. van Creveld*.
- Brinton, Paul H. M.-P., and Charles James,** rates of hydrolysis of the rare earth carbonates and the serial order of the rare earth elements, A., ii, 39.
- concentration of the erbium earths, A., ii, 62.
- British Dyestuffs Corporation, Ltd., James Baddiley, Joseph Baron Payman, and Harry Wignall,** preparation of *o*-sulphonic acids of aromatic amines, A., i, 448.
- British Dyestuffs Corporation, Ltd., Arthur George Green, and Stanley Joseph Green,** preparation of phthalimide, A., i, 836.

- British Dyestuffs Corporation, Ltd.**, *Arthur George Green, Kenneth Herbert Saunders, and Stanley Charles Bate*, preparation of new triaryl-methane colouring matters, A., i, 1068.
- British Dyestuffs Corporation, Ltd.**, *Herbert Levinstein, and Georges Imbert*, preparation of phenylglycine compounds, A., i, 252.
- British Dyestuffs Corporation, Ltd.**, *Arthur George Perkin, and George Douglas Spencer*, preparation of benzanthrone derivatives, A., i, 941.
- British Dyestuffs Corporation, Ltd.** See also *Alexander Walker Fyfe, and David Segaller*.
- British Research Association for the Woollen and Worsted Industries**, the sorption of neutral soap by wool, and its bearing on scouring and milling processes, A., ii, 551.
- standard method for the estimation of soap in wool, A., ii, 594.
- Britton, Edgar C.** See *Moses Gomborg*.
- Britton, Hubert Thomas Stanley**, the system potassium sulphate-aluminium sulphate-water at 25°, T., 982.
- the system ammonium sulphate-glucinum sulphate-water at 25°, T., 2612.
- the separation of aluminium from glucinum. III., A., ii, 228.
- Brochet, André**, the preparation of cyclohexanol, A., i, 1144.
- Brodin, P., and P. Huchet**, a new antianaphylactic substance, sodium formaldehydesulphoxylate, A., i, 90.
- Brodin, P.** See also *A. Chauffard*.
- Brody, E.**, theoretical determination of the chemical constants of monatomic gases, A., ii, 191.
- Brooksmit, T. C. N.**, zinc borate, A., ii, 377.
- Brönsted, Johannes Nicolaus**, applicability of the gas laws to strong electrolytes. II., A., ii, 113.
- studies on chemical affinity. XIII. Reaction affinity in systems of solid salts, A., ii, 132.
- temperature dependence of the solubility, the activity and osmotic coefficients of salts, A., ii, 354.
- solubility. IV. Principle of the specific interaction of ions, A., ii, 481.
- calculation of the osmotic and activity functions in solutions of uni-univalent salts, A., ii, 482.
- theory of the velocity of chemical reactions, A., ii, 699.
- Brönsted, Johannes Nicolaus, and Georg von Hevesy**, separation of the isotopes of mercury, A., ii, 149.
- separation of the isotopes of chlorine, A., ii, 280.
- atomic weight of mercury from different sources, A., ii, 645.
- Brönsted, Johannes Nicolaus, and Agnes Petersen**, solubility. III. Solubility of metal ammonia salts in salt solutions, A., ii, 199.
- Brösamlen**, adrenaline hyperglycæmia, A., i, 1092.
- Brogie, Louis de**, the theory of absorption of X-rays by matter and the principle of correspondence, A., ii, 104.
- X-rays and thermodynamic equilibrium, A., ii, 249.
- Brogie, Maurice de**, the corpuscular spectra of the elements, A., ii, 330.
- Brogie, Maurice de, and A. Dauvillier**, a new absorption phenomenon observed in the domain of X-rays, A., ii, 542.
- Bromig, Karl.** See *Conrad Amberger*.
- Bronn, J.**, temperatures of combustion, A., ii, 548.
- Brooks, Matilda Moldenhauer**, the effect of hydrogen-ion concentration on the production of carbon dioxide by *Bacillus butyræus* and *Bacillus subtilis*, A., i, 201.
- penetration of kations into living cells, A., i, 308.
- Brose, W.** See *Carl Mannich*.
- Broughall, Laurence St. C.**, frequency of the electrons in the neon atom, A., ii, 213.
- aspects of the neon spectrum, A., ii, 541.
- Brown, Adrian**, obituary notice of, T., 2899.
- Brown, Andrew Charles**, the adsorption of uranium-X and its isotope thorium, by basic ferric acetate, T., 1736.
- Brown, Anson L.**, a new quantitative method for the estimation of iron in the blood, A., ii, 319.
- Brown, David Christie.** See *George Gerald Henderson*.
- Brown, Oliver W., S. V. Cook, and J. C. Warner**, effect of grinding on the apparent density of lead oxides, A., ii, 571.
- Brown, Oliver W., and C. O. Henke**, catalytic preparation of aniline. I. and II., A., i, 445, 535.
- catalytic activity of copper, A., ii, 833.
- Brown, Oliver W.** See also *C. O. Henke*.
- Brown, Ralph L.** See *Charles Mourou, and Julius Stieglitz*.

- Brown, William**, preparation and use of collodion osmometers, A., ii, 690.
- Brown, W. E. L., and Archibald Vivian Hill**, the physical chemistry of hæmoglobin in blood, A., i, 1199.
- Browne, (Miss) Agnes**. See *Douglas Norman Jackson*.
- Browne, Arthur Wesley, and A. B. Hoel**, lecture experiments with hydronitric [hydrazoic] acid and the trinitrides [azides], A., ii, 840.
reaction between potassium azide and iodine in the presence of carbon disulphide, A., ii, 847.
potassium azidodithiocarbonate, A., ii, 848.
- Browne, Frederick L., and Joseph Howard Mathews**, heat of coagulation of ferric oxide hydrosol with electrolytes, A., ii, 196.
- Browne, Harold H.** See *Clarke E. Davis*.
- Browning, Carl Hamilton, Julius Berend Cohen, R. Gaunt, and R. Gulbransen**, relationships between antiseptic action and chemical constitution, with special reference to compounds of the pyridine, quinine, acridine, and phenazine series, A., i, 612.
the antiseptic properties of cyanine dyes, A., i, 1097.
- Broy**. See *Bernhard Neumann*.
- Bruchhausen, F. von**. See *Johannes Gadamer*.
- Brude, G.** See *G. Fester*.
- Bruhat, Georges, and A. Delaygue**, determination of the higher point of inversion of the specific heat of the saturated vapour of benzene, A., ii, 348.
- Bruin, G. de**, cellulose nitrates, A., i, 13.
- Bruins, H. R.** See *Ernst Cohen*.
- Brühl, A.** See *Ludwig Moser*.
- Brun, Pierre**. See *Marcel Godehot*.
- Bruni, Giuseppe, and E. Romani**, mercaptothiazoles as accelerators in vulcanisation, A., i, 755.
- Brunkow, O. R., W. H. Peterson, and Edwin Brown Fred**, the influence of certain factors on the chemical composition of sauerkraut, A., i, 312.
- Bruylants, Pierre**, the action of organo-magnesium compounds on glutaronitrile, A., i, 531.
butenenitriles, A., i, 817, 924.
- Bucherer, Hans Theodor, and Rudolf Wahl**, 6-amino- α -naphthol-5-sulphonic acid (A-acid) and its derivatives, A., i, 247.
- Bucherer, Hans Theodor, and Rudolf Wahl**, the action of sulphites on aromatic amino- and hydroxy-compounds. IX. 6-Amino- α -naphthol-5-sulphonic acid (A-acid) and the sulphite reaction, A., i, 464.
- Bucherer, Hans Theodor, and Walther Zimmermann**, the action of sulphites on aromatic amino- and hydroxy-compounds. X. Action of phenylhydrazine-bisulphite mixture, particularly on aminonaphtholsulphonic acids, and azo-dyes, A., i, 465.
- Buchheim, Kurt**. See *Wilhelm Steinkopf*.
- Buchler, C. C.** See *Moses Gomborg*.
- Buchwaldt, Arnold**. See *Gustav Heller*.
- Buck, Johannes Sybrandt, and Isidor Morris Heilbron**, the reactivity of doubly-conjugated unsaturated ketones. III. Unsymmetrical hydroxy- and methoxy-derivatives, T., 1095.
benzopyrylium salts of distyryl ketones, I., T., 1198.
- Buckley, Harold**. See *Lionel Felix Gilbert*.
- Buckner, G. Davis, J. H. Martin, W. C. Pierce, and A. M. Peter**, calcium in egg-shell formation, A., i, 488.
- Buckner, Henry K.** See *George L. Clark*.
- Buddington, A. F.**, some natural and synthetic melilites, A., ii, 155.
- Budnikov, P. P., and K. E. Krause**, estimation of sulphides by oxidation with ferric salts, A., ii, 717, 782.
- Budnikov, P. P., and E. A. Schilov**, action of sulphur and sulphur compounds on terpenes, A., i, 944.
- Budnikov, P. P., and Ja. K. Syrkin**, setting and velocity of solution of burnt gypsum, A., ii, 849.
- Budnikov, P. P., and P. V. Zolotarev**, saccharification of cellulose, A., i, 922.
- Bücher, W.** See *Alfred Benrath*.
- Buell, Harold D.**, new qualitative test for uranium, A., ii, 590.
- Buell, Harold D., and C. R. McCrosky**, preparation and study of the rarer alkali bromates; rubidium bromate, A., ii, 146.
- Bülow, Hans**. See *Heinrich Bilts*.
- Bürger, Max**, the action of intravenous injections of hypertonic solutions of various sugars on the respiratory metabolism of the dog, A., i, 286.
- Bürklin, Elisabeth**. See *P. Karrer*.
- Bull, Arthur W., and J. R. Adams**, alizarin-iron lakes, A., i, 355.
- Bulli, Mario**. See *Luigi Bolla*.

- Ballis, D. E.** See *R. H. Robinson*.
- Balygina, Nad.** See *Nikolai Schilov*.
- Bump, Albert H.** See *James Bryant Conant*.
- Bunbury, Hugh Mills**, the sorption of carbonyl chloride by beechwood charcoal, *T.*, 1525.
- Bunzl, Cornélie.** See *Alfons Klemenc*.
- Burbidge, P. W.**, absorption of the *K* X-rays of silver in gases and gaseous mixtures, *A.*, ii, 184.
- Burdick, Charles Lator**, oxidation of nitric oxide and its catalysis, *A.*, ii, 272.
analytical determination of oxides of nitrogen in gas mixtures, *A.*, ii, 588.
- Burg, J. H. N. van der**, the preparation of acrylic acid and some of its derivatives, *A.*, i, 220.
- Burger, H. C.**, X-ray examination of the trioxides of tungsten, molybdenum, and their hydrates, *A.*, ii, 508.
- Burgess, Henry.** See *Thomas Martin Lowry*.
- Burkardt, Hermann**, rapid method for estimating ammonia in ammonium salts, *A.*, ii, 865.
- Burnett, Arthur John.** See *James William McBain*.
- Burns, Robert.** See *Forsyth James Wilson*.
- Burr, G. O.** See *Charles Shattuck Palmer*.
- Burrows, George Joseph**, and *Eustace Ebenezer Turner*, the preparation of certain ferrioxalates, *A.*, i, 916.
- Burton, E. F.**, and *E. D. MacInnes*, coagulation of colloidal solutions of arsenious sulphide by electrolytes, *A.*, ii, 130.
- Burton, Harold**, and *James Kenner*, the influence of nitro-groups on the reactivity of substituents in the benzene nucleus. V. Heteronuclear dinitro-derivatives, *T.*, 489.
the influence of nitro-groups on the reactivity of substituents in the benzene nucleus. VI. The elimination of halogen during the reduction of halogenated nitro-compounds, *T.*, 675.
- Bury, Charles R.**, Langmuir's theory of the arrangement of electrons in atoms and molecules, *A.*, ii, 43.
- Bury, Frank Ward**, and *James Riddick Partington*, preparation and reactions of stannous oxide and stannous hydroxides, *T.*, 1998.
- Buryánek, Ot.** See *Franz Jirsa*.
- Busch, Hedwig.** See *Emil Knoevenagel*.
- Bushill, John Herbert.** See *Arthur Robert Lang*.
- Busten, Harold W.**, and *Samuel Barnett Schryver*, method for the separation of amino-acids from the products of hydrolysis of proteins and other sources, *A.*, i, 182.
- Buswell, Arthur M.** See *R. E. Greenfield*.
- Butkewitsch, Wl.**, formation of oxalic acid and ammonia in cultures of *Aspergillus niger* on peptone, *A.*, i, 707.
utilisation of peptone as source of carbon by *Citromyces* species, *A.*, i, 707.
formation and accumulation of oxalic acid in *Citromyces* cultures on salts of organic acids, *A.*, i, 707.
ammonia as a product of protein transformation caused by mould fungi and the conditions of its formation, *A.*, i, 973.
formation of citric and oxalic acids in *Citromyces* cultures on sugar and a process for the estimation of these acids, *A.*, i, 973.
utilisation and formation of citric acid in cultures of *Citromyces glaber* on sugar, *A.*, i, 973.
- Butler, E. H.**, decomposition of iodoform in solutions by means of radiant energy, *A.*, ii, 604.
- Byk, Alfred**, conversion of energy in photochemical processes, *A.*, ii, 415.
- Bylinkin, J. S.** See *Alexei E. Tschitschibabin*.

C.

- Cabrera, B.**, magnetic properties and atomic structure, *A.*, ii, 469.
- Cabrera, B.**, and *Santiago Piña de Rubies*, variations in the magnetic susceptibility of oxychromic salts with addition of sulphuric acid, *A.*, ii, 612.
- Cady, Hamilton Perkins**, *Howard M. Elsey*, and *Emily V. Berger*, solubility of helium in water, *A.*, ii, 642.
- Cain, John R.**, and *J. Clyde Hostetter*, co-precipitation of vanadic acid with ammonium phosphomolybdate, *A.*, ii, 320.
- Cake, W. Ellwood**, the catalytic hydrogenation of dextrose, *A.*, i, 523.
- Cake, W. Ellwood**, and *H. H. Bartlett*, the carbohydrate content of the seed of *Asparagus officinalis*, *L.*, *A.*, i, 504.
- Cake, W. Ellwood.** See also *Hobart Hurd Willard*.
- Calcagni, Gennaro**, new hypothesis of the origin of natural fuels, *A.*, ii, 385.
- Caldwell, Mary L.** See *Henry Clapp Sherman*.

- Caldwell, *William E.*, and *William Gordon Lyle*, blood chemistry in normal and abnormal pregnancy, A., i, 489.
- Cale, *F. M.* See *John Cunningham McLennan*.
- Calingaert, *G.*, a homologue of ethylene sulphide; trimethylethylene [β -methyl- Δ -butylene] sulphide, A., i, 421.
- Callan, *Thomas*, and *James Alexander Russell Henderson*, estimation of the nitro-group in aromatic organic compounds. II., A., ii, 524.
- Callow, *E. H.* See *Arthur Robert Ling*.
- Callisen, *J.*, ethyl bromodiethylacetyl-allophanate, A., i, 1128.
- Camacho, *Francisco*. See *Max Bergmann*.
- Cameron, *Alexander Thomas*, and *M. S. Hollenberg*, the relative toxicity of the haloids and other anions, A., i, 499.
- Cameron, *Don H.* See *Harry N. Holmes*.
- Cambridge, *P. J.*, *J. A. C. Forsyth*, and *H. A. Howard*, normal sugar content of the blood, A., i, 81.
- Campbell, *Arthur Fred*, the speed of sulphonation of phenols. I. The effect of temperature and the methyl group, T., 847.
- Campbell, *Colin*, the propagation of explosion waves in gases contained in tubes of varying cross-section, T., 2483.
- Campbell, *H. L.* See *Victor K. LaMer*, and *Henry Clapp Sherman*.
- Campbell, *James Argyll*, and *Thomas Arthur Webster*, day and night urine during complete rest, laboratory routine, light muscular work, and oxygen administration, A., i, 197. effect of severe muscular work on the composition of the urine, A., i, 495.
- Canals, *E.*, effect of different kinds of solar radiation on the formation of essential oils in plants, A., i, 907. saccharase, A., i, 1075. the inversion of sucrose in the alkaline copper solution, A., ii, 592.
- Canals, *E.* See also *A. Astruc*.
- Cannan, *Robert Keith*. See *Jack Cecil Drummond*.
- Canneri, *G.*, chromates of thallium, A., ii, 297. thallium bismutho-, stibio-, and arseno-thiosulphates, A., ii, 378.
- Canneri, *G.*, and *R. Morelli*, thermal analysis of the system $Tl_2O-B_2O_3$, A., ii, 571.
- Canneri, *G.*, and *G. Perina*, double haloids of bismuth and thallium, A., ii, 512.
- Cant, *H. J.* See *James Riddick Partington*.
- Capps, *Julian H.*, estimation of metallic aluminium and aluminium oxide in commercial metal, A., ii, 319.
- Carlo, *G.*, and *J. Franck*, the decomposition of hydrogen molecules by excited mercury atoms, A., ii, 809.
- Carl, *H.*, examination of the compression equation of liquids by means of the data of Amagat and Bridgman, A., ii, 477.
- Carletti, *Ottorino*, the Kastle-Meyer reagent as a very sensitive reagent for copper, A., ii, 787.
- Carlton, *C. A.*, symmetrical diphenylguanidine as a standard in acidimetry and alkalimetry, A., ii, 654.
- Carmichael, (*Mies*) *Madel*, electro-synthesis of azelaic and thapsic acids, T., 2545.
- Carnot, *Paul*, and *Marc Tiffeneau*, a new hypnotic of the barbituric acid series; ethylbutylbarbituric acid, A., i, 900.
- Carothers, *W. A.* See *Charles Shattuck Palmer*.
- Carozzi, *Enrico*. See *Luigi Losana*.
- Carpenter, *Henry Cort Harold*, and *Constance F. Elam*, production of single crystals of aluminium and their tensile properties, A., ii, 69.
- Carpentier, *Georges*. See *Pierre Thomas*.
- Carr, *K. H.*, measuring soil toxicity, acidity, and basicity, A., ii, 172.
- Carr, *R. H.* See also *M. F. Showalter*.
- Carra, *J.*, comparison of methods for the estimation of urea, A., ii, 668.
- Carrara, *Gino*. See *Angelo Angeli*.
- Carré, *Marjory Harriotte*, changes which occur in the pectic constituents of stored fruit, A., i, 1222.
- Carré, *Marjory Harriotte*, and *Dorothy Haynes*, the estimation of pectin as calcium pectate and the application of this method to the estimation of the soluble pectin in apples, A., ii, 401.
- Carres, *C.*, clarification of urines by zinc ferrocyanide, A., ii, 233.
- Carriek, *L. L.*, solubilities and cooling curves of the mononitrophenols, A., i, 334.
- Carrière, *E.*, the aldehyde acids of the succinic series, A., i, 318.
- Carruthers, *Albert*, and *Edmund Langley Hirst*, methylation of xylose, T., 2299.
- Carter, *Edna*, the vacuum-spark spectra of the metals, A., ii, 599.
- Carter, *E. G.* See *Joseph E. Greaves*.
- Cartoceti, *A.* See *G. Sirovich*.

- Carus, M.**, improved method for the separation of iron and manganese, A., ii, 88.
- Casale, Luigi**, absorbent power of soils, and the absorption by plants of nutritive substances from the soil, A., i, 508.
- Case, Francis H.** See *Treat Baldwin Johnson*.
- Cashmore, Albert Eric, Hamilton McCombie, and Harold Archibald Scarborough**, the velocity of reaction in mixed solvents. II. The velocity of saponification of the ethyl esters of some monosubstituted benzoic acids, T., 243.
- Cassel, H.**, vapour pressures of binary mixtures, A., ii, 424.
relationship of Dolezalek's theory of solutions to that of Planck, A., ii, 481.
- Cassella & Co., Leopold**, preparation of a new acridine compound, A., i, 276.
preparation of compounds of pyridine-betaine with metallic salts, A., i, 860.
preparation of 1-amino-2-anthraquinone aldehyde, A., i, 942.
- Cassens, H.** See *Emil Heuser*.
- Castellani, Aldo, and Frank E. Taylor**, identification of inulin by a mycological method, A., ii, 879.
- Catalán, Miguel A.**, series and other regularities in the spectrum of manganese, A., ii, 726.
- Cate, J. Ten**, action of potassium, calcium and magnesium ions on the sympathetic nerve of the heart, A., i, 296.
- Cathcart, P. H.**, a simple hydrogen generator for use in making hydrogen-ion determinations, A., ii, 440.
- Catoire, M.** See *G. Malfitano*.
- Cavassani, Émile**, crystallisation and proteius, A., ii, 745.
- Caven, Robert Martin**, qualities of valency, A., ii, 279.
- Caven, Robert Martin, and John Ferguson**, the dissociation pressures of hydrated double sulphates. I. Hydrated cupric alkali sulphates, T., 1406.
- Chadwick, James, and E. S. Bieler**, collisions of α -particles with hydrogen nuclei, A., ii, 12.
- Chadwick, James, and C. D. Ellis**, the intensity distribution in the β -ray spectra of radium-B and -C, A., ii, 802.
- Chadwick, James.** See also (*Sir*) *Ernest Rutherford*.
- Challenger, Frederick, and Leslie Randal Ridgway**, organo-derivatives of bismuth. VI. The preparation and properties of tertiary aromatic bismuthines and their interaction with organic and inorganic halogen compounds, T., 104.
- Challenger, Frederick, and John Frederick Wilkinson**, organo-derivatives of bismuth. V. The stability of halogen, cyano- and thiocyno-derivatives of tertiary aromatic bismuthines, T., 91.
- Chalupny, K., and K. Breisch**, estimation of magnesium in technical nickel, A., ii, 227.
separation of aluminium from iron by means of *o*-phenetidine, A., ii, 588.
- Chandrasena, Juan Pedigo Charles, and Christopher Kelk Ingold**, the conditions underlying the formation of unsaturated and cyclic compounds from halogenated open-chain derivatives. IV. Products formed from halogen derivatives of muconic acid; the constitution of muconic acid, T., 1306.
effect of attached groups on the ease of formation of the cyclopentane ring, T., 1552.
- Chandrasena, Juan Pedigo Charles, Christopher Kelk Ingold, and Jocelyn Field Thorpe**, the chemistry of polycyclic structures in relation to their homocyclic unsaturated isomerides. III. Intra-annular tautomerism of α -campholytic acid, T., 1542.
- Chandratraya, V. L.** See *Howard James Winch*.
- Chantraine, H.**, the blood-sugar in narcosis and diseases of the nervous system, A., i, 192.
- Chanutin, Alfred**, animal calorimetry. XX. The influence of the ingestion of meat and of glycine and alanine on the carbon dioxide-combining power of blood plasma, A., i, 288.
- Chapas**, solubility of the isomeric toluic acids in the three xylenes, A., i, 339.
- Chapman, Alfred Chaston**, examination of foods for the presence of sulphites, A., ii, 520.
- Chapman, Arthur William**, imino-aryl ethers. I. *N*-Phenylbenzimidino-*m*-hydroxyphenyl ether and the synthesis of 2:4-dihydroxybenzophenone, T., 1676.
- Charitschkov, K.**, Winkelblech's phenomenon or pseudo-extraction and its importance in colloidal chemistry; new methods of extracting solids, A., ii, 826.

- Charitschkov, K.**, general method for obtaining gels of inorganic salts, and its relation to theories of the colloidal state, A., ii, 827.
- Charlton, Thomas J.** See *Howard W. Haggard*.
- Charpy, Georges**, and **Louis Grenet**, the penetration of tempering in steel, A., ii, 507.
- Charrier, G.**, *p*-nitrophenylstibinic acid, A., i, 1080.
- Charrier, G.** [with *C. Sala, S. Viola, G. Crippa, C. Cortassa, L. Demichelis, and M. Speirani*], aminoazo-, hydroxyazo-, and hydrazo-compounds, A., i, 769.
- Charrier, G.**, and **A. Beretta**, azimino-benzene (1:2:3-benzotriazol₃), A., i, 61.
- Charriou, A.**, the carrying down of calcium oxide by precipitates of ferric oxide, A., ii, 163.
the separation of the oxides of iron and aluminium from admixture with calcium oxide by the nitrate method, A., ii, 319.
the separation of ferric oxide and aluminium oxide from magnesium oxide by the nitrate method, A., ii, 873.
- Chattaway, Frederick Daniel**, and **Hinton John Harris**, the oxidation of sucrose by nitric acid, T., 2703.
- Chattaway, Frederick Daniel**, and **Henry Rowland Hill**, the interaction of diazonium salts and phenols, T., 2756.
- Chattaway, Frederick Daniel**, and **Deric William Parkes**, isomeric citraconyl hydrazides, T., 283.
- Chatterjee, Bibhu Charan.** See *Rasik Lal Datta*.
- Chatterjee, Kshetra Pada**, estimation of sulphate-ion as barium sulphate. I., A., ii, 390.
- Chatterji, Nitya Gopal**, and **Nilratan Dhar**, peptisation and precipitation, A., ii, 205.
Liesegang's phenomenon and precipitation formation, A., ii, 627.
- Chaudron, Georges**, and **G. Juge-Boirard**, the estimation of sulphur in iron pyrites, A., ii, 311.
- Chaudun, (Mlle) Andrée**, the inversion of sucrose by saccharase, A., i, 389.
- Chaudun, (Mlle) Andrée.** See also *H. Collin*.
- Chauffard, A.**, **P. Brodin**, and **A. Grigaut**, uric acid content of blood corpuscles, A., i, 1086.
- Chavanne, Georges**, and **Paul Becker**, dimethylcyclohexanes, A., i, 442.
- Chavanne, Georges**, and **B. Lejeune**, a new heptane; trimethylisopropylmethane [$\beta\beta$ -trimethylbutane], A., i, 417.
- Chavanne, Georges**, and **(Mlle) Hortense van Risseghem**, the viscosities of some hydrocarbons, A., ii, 352.
- Chazan, Sydney.** See *Gilbert Thomas Morgan*.
- Cheetham, H. C.** See *W. Lee Lewis*.
- Chemin, E.**, corrosive action of roots on marble, A., i, 94.
- Chemische Fabrik auf Aktien vorm. E. Schering**, preparation of a new diethylarbituric acid compound, A., i, 582.
preparation of camphene hydrochloride, A., i, 943.
preparation of 2-ar-tetrahydronaphthylquinoline-4-carboxylic acids, A., i, 952.
- Chemische Fabrik Flora**, preparation of sodium silver-thioglycollate, A., i, 425.
- Chemische Fabrik Griseheim-Elektron**, hydrogenation of unsaturated hydrocarbons, A., i, 977.
- Chemische Fabrik Worms Akt.-Ges.**, preparation of anthraquinone and its derivatives, A., i, 559.
- Chen, Yü-Gwan.** See *Marston Taylor Bogert*.
- Chéneveau, Charles**, an optical method for the determination of the reciprocal solubility of slightly miscible liquids, A., ii, 355, 427.
- Cherbulles, Émile**, and **Emanuel Feer**, formaldehyde derivatives of 2:5-diketopiperazine, A., i, 1065.
- Cherbulles, Émile**, and **K. N. Stavritsch**, new syntheses of pyrimidines, A., i, 581.
- Chernoff, Lewis H.** See *Carl Oscar Johns*.
- Chervet, D.** See *W. D. Treadwell*.
- Chesnut, Victor K.** See *Frederick Belding Power*.
- Chevalier, Joseph-Marie-Alphonse**, and **Paul Bourcet**, preparation of ethylene by reduction of acetylene, A., i, 801.
- Chevenard, P.**, expansion of chromium and of nickel-chromium alloys over a wide range of temperatures, A., ii, 153.
- Chiba, Chozo.** See *Riko Majima*.
- Chibnall, Albert Charles**, nitrogenous metabolism of the higher plants. II. The distribution of nitrogen in the leaves of the runner bean, A., i, 908.
nitrogenous metabolism of the higher plants. III. The effect of low-temperature drying on the distribution of nitrogen in the leaves of the runner bean, A., i, 1225.

- Chibnall, Albert Charles**, nitrogenous metabolism of the higher plants. IV. Distribution of nitrogen in the dead leaves of the runner bean, A., i, 1225.
- Chikashige, Masuri**, Honda's conception of the A¹ transformation and the quenching of steels, A., ii, 852.
- Child, C. D.**, a continuous spectrum from mercury vapour, A., ii, 676.
- Chiles, Howard M.**, and **William Albert Noyes**, optically active diazo-compounds. II., A., i, 924.
- Chirvinskii, P.**, olivine from lithosiderites, A., ii, 714.
- Chin, C. Y.** See **Charles August Kraus**.
- Chmelař, J.** See **Julius Stoklasa**.
- Chon, Sung-Sheng.** See **Hermann Staudel**.
- Chouchak, D.**, colorimetric estimation of arsenic by means of quinine molybdate, A., ii, 526.
- Choudhury, Kumud Nath.** See **Haridas Saha**.
- Chouke, K. S.** See **Edward A. Doisy**.
- Christiansen, J. A.**, a reaction between methyl alcohol and water and some related reactions, A., i, 3.
- Christiansen, Walter G.**, the relation between the mode of synthesis and toxicity of arsphenamine [salvarsan] and related compounds, A., i, 186. the sulphur content of arsphenamine (salvarsan) and its relation to the mode of synthesis and toxicity. I., II., and III., A., i, 601, 1202.
- Christie, George Hallatt**, and **James Kenner**, the molecular configurations of polynuclear aromatic compounds. I. The resolution of γ -6:6'-dinitro- and 4:6:4':6'-tetranitro-diphenic acids into optically active components, T., 614.
- Christin, P.** See **Paul Wenger**.
- Churchill, H. V.**, estimation of metallic aluminium and aluminium oxide in commercial metal, A., ii, 319.
- Chwolson, O.**, structure of the atomic nucleus, A., ii, 209.
- Ciacio, C.**, [estimation of] amino-nitrogen in the urine by the formol method, A., i, 88.
- Ciamicioian, Giacomo Luigi**, and **A. Galissi**, behaviour of certain organic compounds in plants. XIV., A., i, 503.
- Ciamicioian, Giacomo Luigi**, and **Ciro Ravenna**, biological signification of alkaloids in plants, A., i, 797.
- Cioffi, P. P.**, and **L. S. Taylor**, method of maintaining small objects at any temperature between -180° and $+20^{\circ}$, A., ii, 817.
- Ciusa, Riccardo**, "graphites" from pyrrole and from thiophen, A., i, 1057.
- Doebner's reaction.** IV., A., i, 1062.
- Ciusa, Riccardo**, and **M. Croes**, some constituents of lignites. II., A., ii, 385.
- Ciusa, Riccardo**, and **G. Bastelli**, certain salts with para-, ortho-, and meta-quinonoid structure. III. and IV., A., i, 1073.
- Ciusa, Riccardo**, and **Luigi Vecchiotti**, basic properties of the hydrazones. II., A., i, 474.
- Ciusa, Riccardo**, and **R. Vois**, fossil wax of Monte Falò, A., ii, 386.
- Claassen, Walter.** See **Richard Anschütz**.
- Clarens, J.**, catalysts and chemical equilibrium, A., ii, 436.
- Clark, A. H.** See **Lewis E. Warren**.
- Clark, A. J.**, mode of action of potassium on isolated organs, A., i, 399.
- Clark, A. W.**, and **R. F. Keeler**, a modified method for the estimation of phosphoric acid, A., ii, 84.
- Clark, Earl P.**, an improved method for preparing raffinose, A., i, 323.
- preparation of mannose, A., i, 433.
- Clark, George L.**, the properties of elements and salts as related to the dimensions of atoms and ions, A., ii, 634.
- Clark, George L.**, and **Henry K. Buckner**, properties of subsidiary valency groups. III. The preparation, properties, and molecular volume relationships of the hydrates and amines of cobalt fluoride, bromide, iodide, nitrate, carbonate, and citrate, A., ii, 300.
- Clark, George L.**, and **William Duane**, a new method of using X-rays in crystal analysis, A., ii, 483.
- Clark, George L.**, and **William A. Mann**, adsorption in solution and at interfaces of sugars, dextrin, starch, gum arabic, and egg-albumin, and the mechanism of their action as emulsifying agents, A., ii, 550.
- Clark, Guy W.**, micro-estimation of calcium in whole blood, plasma, and serum by direct precipitation, A., ii, 227.
- Clark, Guy W.** See also **Carl L. A. Schmidt**.
- Clark, Norman A.**, rate of formation and the yield of yeast in wort, A., i, 501.
- Clark, R. H.**, velocity of hydrolysis of sucrose, A., ii, 185.

- Clark, S. M. See *De Witt Neighbors*.
- Clark, Walter. See *Sidney Owen Rawling*.
- Clark, William Mansfield, instability of phthalate potentials, A., ii, 468.
- Clarke, Rosalind. See *Thomas Dillon*.
- Classen, W. See *A. Fischer*.
- Claude, Georges, the elimination of the heat of reaction in the synthesis of ammonia by high pressures, A., ii, 284.
- Claudin, J. See *Martin Battagay*.
- Clausen, S. W., estimation of small amounts of lactic acid, A., ii, 593.
- Clavera, José María, melting point of commercial terpin, A., i, 1042.
- Clawson, A. B. See *C. Dwight Marsh*.
- Clayson, Donald Herbert. Frank, Frederick Walter Norris, and Samuel Barnett Schryver, pectic substances of plants. II. Preliminary investigation of the chemistry of the cell-walls of plants, A., i, 206.
- Clayton, Herbert. See *William Davies*, and *Nevil Vincent Sidgwick*.
- Clément and Rivière, attempts at a synthetic manufacture of mother-of-pearl by production of chemical tracery, A., ii, 500.
- Clemo, George Roger, and William Henry Perkin, jun., introduction of the chloroethyl group into phenols, alcohols, and amino-compounds, T., 642.
- Clendinnen, Frederick William Jeffrey, mixed crystal formation in ternary systems containing water, ammonium chloride, and ferrous, cobaltous, or nickel chloride, T., 801.
- Clerici, Enrico, heavy liquids for the separation of minerals, A., ii, 578.
- Clews, Francis Herbert, and Hugh Vernon Thompson, the interaction of sodium chloride and silica, T., 1442.
- Clifford, Percy Herbert, and Robert George Fargher, "oxycellulose": its formation and reactions, A., i, 1121.
- Clifford, Winifred Mary, the distribution of carnosine in the animal kingdom, A., i, 398.
the effect of cold storage on the carnosine content of muscle, A., i, 791.
- Clogne, René, mercury or water ureometer for the estimation of urea in urine or in blood, A., ii, 237.
- Clogne, René. See also *Paul Couinaud*.
- Clover, Alphonso Morton, the autoxidation of ethyl ether, A., i, 619.
- Clatterbuck, Percival Walter, and Julius Berend Cohen, the aliphatic sulphonamides. I., T., 120.
- Cobet, Rudolf, and V. van der Reis, influence of arsenious acid on bacterial growth, A., i, 611.
- Cocconi, G. See *Efisio Mameli*.
- Cocks, L. V., and Arthur Henry Salway, a method for the estimation of trimethylene glycol in crude glycerol, A., ii, 232.
- Coffey, Samuel, the mechanism of the oxidation of drying oils as elucidated by a study of the true oxygen absorption. III. The action of driers, T., 17.
the reaction between sulphur monochloride and aniline, A., i, 132.
- Cohen, Abraham, xylenol-blue and its proposed use as a new and improved indicator in chemical and biochemical work, A., ii, 387.
use of mixed indicators, A., ii, 780.
- Cohen, Clara. See *Carl Neuberg*.
- Cohen, Ernst, and H. R. Bruins, use of the Zeiss water interferometer (Rayleigh-Lowe) for the analysis of non-aqueous solutions, A., ii, 77.
- Cohen, Ernst, and A. L. Th. Moersveld, determination of the specific heat of solid substances by the adiabatic electrical method, A., ii, 347.
- Cohen, Julius Berend. See *Carl Hamilton Browning*, *Percival Walter Clatterbuck*, *Victor Froelicher*, *John Richard Scott*, *Akira Shimomura*, and *Edward Johnson Wayne*.
- Cohn, Edwin Joseph, the physical chemistry of the proteins. I. The solubility of certain proteins at their isoelectric points, A., i, 882.
- Cohn, Robert, a solid water-soluble formaldehyde preparation, A., i, 918.
- Colby, Walter F., and Charles F. Meyer, the absorption spectrum of hydrogen chloride, A., ii, 5.
- Colby, Walter F. See also *H. M. Randall*.
- Cole, Harriet Isabelle. See *George Shannon Forbes*.
- Cole, Howard Irving, the dissociation of hexaphenylethane from the point of view of the octet theory of valence, A., ii, 438.
- Cole, R. M., thymol from nitrocymene, A., i, 31.
- Coleman, George H., and William Albert Noyes, chlorination and the formation of chloroamines by means of nitrogen trichloride, A., i, 133.
- Colin, H., and (Mlle) André Chaudan, the law of action of saccharase; velocity of hydrolysis and reaction of the medium, A., i, 389.

- Collidge, W. C.**, extraction and characterisation of alkaloids, A., ii, 827.
extraction of alkaloids from viscera, A., ii, 327.
- Collenberg, Oscar Olsson**, the determination of the state of valency of lower compounds by means of ammoniacal silver solutions, A., ii, 495.
the valency of tungsten and molybdenum in their complex octacyanides, A., ii, 508.
- Collenberg, Oscar Olsson**, and **Sven Bodfors**, kinetics of reactions in heterogeneous solutions; the reduction and oxidation actions of alternating currents, A., ii, 431.
- Collet, (Mlle) Paule**, thin films of binary mixtures (glycerides), A., ii, 477.
- Collie, John Norman**, and **(Miss) Amy Ada Beatrice Reilly**, diacetylacetone, T., 1984.
- Collins, George Ernest**, some physical properties of cotton cellulose and its modifications; a summary of existing data, A., i, 1120.
- Collins, George Ernest**, and **John Kerfoot Wood**, the amphoteric character of stannic hydroxide and its bearing on the isomerism of the stannic acids, T., 441.
the behaviour of the stannic acids towards hydrochloric acid, T., 1122.
the behaviour of the stannic acids towards solutions of alkaline hydroxides, T., 2760.
- Collins, Hawksworth**, relative volumes of the chemical elements, A., ii, 28
a new physico-chemical law, A., ii, 685.
- Collins, Sydney Hoare**, estimation of levulose (fructose) in straw, A., ii, 323.
- Collipp, J. B.**, the respiratory processes in *Mya arenaria* and other marine mollusca, A., i, 286.
- Collischonn, H.** See **Kurt Brand**.
- Comber, Norman M.**, the flocculation of soils. II., A., i, 212.
relation of the hydrogen-ion concentration of the soil to plant distribution, A., i, 416.
- Combes, Raoul**, the detection of the pseudo-bases of anthocyanidins in plant-tissues, A., i, 206.
the formation of anthocyanin pigments, A., i, 412.
- Combes, Raoul**, and **(Mlle) Denise Kohler**, what becomes of carbohydrates when the leaves of trees die? A., i, 1222.
- Comella, G.** See **E. Oliveri-Mandalà**.
- Compton, Arthur**, blood-enzymes. I. Occurrence of maltase in mammalian blood, A., i, 392.
blood enzymes. II. The influence of temperature on the action of the maltase of dog's serum, A., i, 1087.
- Compton, Karl T.**, tungsten furnace for experiments on dissociation and ionisation, A., ii, 822.
- Conant, James Bryant**, additive reactions of the carbonyl group involving the increase in valency of a single atom, A., i, 41.
- Conant, James Bryant, Albert H. Bump**, and **Harold S. Holt**, additive reactions of phosphorus haloids. III. The reaction with distyryl ketone and phenyl cinnamylidenemethyl ketone, A., i, 67.
- Conant, James Bryant, H. M. Kahn, L. F. Fieser**, and **S. S. Kurtz, jun.**, an electrochemical study of the reversible reduction of organic compounds, A., ii, 547.
- Conant, James Bryant, Alexander D. MacDonald**, and **A. McB. Kinney**, additive reactions of phosphorus haloids. IV. The action of the trichloride on saturated aldehydes and ketones, A., i, 186.
- Conant, James Bryant**, and **S. M. Pollack**, additive reactions of phosphorus haloids. II. The 1:4-addition of phosphenyl chloride, A., i, 67.
- Conant, James Bryant**. See also **Theodore William Richards**.
- Condelli, S.**, enantiomorphism of matter, Pasteur's theory, and life, A., i, 410.
- Conduché**. See **Georges Bourguignon**.
- Congdon, Leon A.**, and **Harry R. Ingersoll**, the influence of dextrose on the dialysis of sucrose through a parchment membrane; the possibility of the separation of dextrose from sucrose by dialysis, A., i, 322.
- Congdon, Leon A.**, and **Charles R. Stewart**, test for sucrose in the presence of dextrose, A., ii, 233.
- Conner, Samuel D.**, and **O. H. Sears**, aluminium salts and acids at varying hydrogen-ion concentrations, in relation to plant growth in water cultures, A., i, 613.
- Conover, Courtney**, and **Harry Drake Gibbs**, preparation of phthalic anhydride by the catalysis of the vapour phase reaction between naphthalene and atmospheric air, A., i, 454.
- Conrad, Eva**. See **Hermann Leuchs**.
- Consortium für Elektrochemische Industrie**, preparation of chlorinated acetyl chlorides, A., i, 315.

- Consortium für Elektrochemische Industrie**, preparation of crotonaldehyde, A., i, 1115.
- Constabel, Fr.**, the creatine content of the human heart muscle in various illnesses, A., i, 296.
- Conway, Ruth E.**, and **Florence V. Stephen**, the reaction of blood, A., i, 892.
- Cook, Charles W.**, a new occurrence of ilsemannite, A., ii, 576.
- Cook, James Wulfred.** See **Edward de Barry Barnett**.
- Cook, Leon W.**, a simpler method of determining acetyl values, A., ii, 824.
- Cook, S. V.** See **Oliver W. Brown**.
- Cooke, Martha C.** See **Olin Freeman Tower**.
- Coope, R.**, sugar of cerebrospinal fluid, A., i, 295.
- Cooper, Evelyn Ashley**, and **Hilda Walker**, nature of the reducing substance in human blood, A., i, 1084.
- Cooper, Evelyn Ashley.** See also **Gilbert Thomas Morgan**.
- Cooper, P. A.**, the X-ray structure of potassium cyanide, A., i, 1128.
- Coope, J., jun** See **P. E. Verkade**.
- Cope, F. F.** See **Cecil E. Boord**.
- Copisarow, Maurice**, theory of allotropy, A., ii, 137.
- Coppetti, Victor**, estimation of sulphurous acid, A., ii, 80.
- Corelli, R. M.**, preparation and constitution of a double potassium ammonium orthophosphate, A., ii, 287.
- Cori, Karl**, the action of intravenous injections of dextrose and gum arabic solution on diuresis, A., i, 1091.
- Corleis, Wilhelm.** See **Kurt Hess**.
- Corman, H. E.** See **Maitland Crease Boswell**.
- Cornog, Jacob**, simple method for the preparation of sodium hydroxide free from carbon dioxide, A., ii, 288.
- Corral, José Maria de**, influence of temperature on the reaction of the blood, A., i, 79.
- Corran, J. W.**, and **William Cudmore McCullagh Lewis**, effect of sucrose on the activities of the chloride- and hydrogen-ions, A., ii, 691.
- Correns, Erich.** See **Ernst Beckmann**.
- Cortassa, C.** See **G. Charrier**.
- Costantin, J.**, physiology of anthocyanin and chemistry of chlorophyll, A., i, 162.
- Costantino, A.**, the amino-acid content of plasma and corpuscles according to Bang, A., i, 81.
- Costa-Vet, E. Mendes da**, quantitative analysis of aluminium alloys, especially of duralumin, A., ii, 528.
- Coster, Dirk**, systematics of Rontgen spectra, A., ii, 180.
- the *L*-series of the X-ray spectrum, A., ii, 244.
- the *L*-series of the elements barium to rubidium, A., ii, 462.
- spectra of X-rays and the theory of atomic structure, A., ii, 491, 677.
- Costy, P.** See **A. Goris**.
- Couch, James F.** See **C. Dwight Marsh**.
- Couinaud, Paul**, and **René Clogne**, blood chemistry in puerperal infection, A., i, 401.
- Coulson, William.** See **Aquila Forster**.
- Cournot, Jean.** See **Léon Guillet**.
- Cousen, A.**, and **William Ernest Stephen Turner**, production of colourless glass in tank furnaces with particular reference to the use of selenium. I., A., ii, 708.
- Couture, John E.** See **Lemuel Charles Raiford**.
- Cow, Douglas V.**, and **Walter Ernest Dixon**, [physiological] action of dimethyl telluride dihaloids, A., i, 402.
- Coward, Hubert Frank**, and **Gladys Mary Wigley**, the detection and estimation of acidity and alkalinity in cotton fabrics, A., ii, 531.
- Coward, Katharine Hope.** See **Henry Lyster Jameson**.
- Cowlishaw, Geoffrey E.** See **G. F. Pickering**.
- Cox, Henry L.** See **Ben H. Nicolet**.
- Coysh, Reginald Henry.** See **Francis Ernest Francis**.
- Crabtree, Herbert Grace**, and **Robert Robinson**, a synthesis of isobrazilein and certain related anhydropyranol salts. II. Synthesis of isohæmatein, T., 1033.
- Craig, Wallace A.** See **John F. G. Hicks**.
- Crane, Marian M.**, effect of hydrogen-ion concentration on the toxicity of alkaloids for *Paramæcium*, A., i, 404.
- Cranston, John Arnold**, and **Robert Hutton**, the adsorption of radium-B and radium-C by ferric hydroxide, T., 2843.
- Craven, E. C.** See **William R. Ormandy**.
- Crawford, A.** See **Leonard A. Sayce**.
- Crawford, Archibald Barclay.** See **Forsyth James Wilson**.
- Crehore, Albert C.**, atoms and molecules. II., A., ii, 438.
- the hydrogen molecule. III., A., ii, 494.

- Creighton, Henry Jermain Maude**, a method for making methyl-violet, A., i, 478.
 electrolytic concentration of aqueous solutions of nitric acid. I., A., ii, 212.
 method of calculating fluidity, surface tension, and reaction (equilibrium) pressure, A., ii, 426.
- Cremer, C. J.** See *I. M. Kolthoff*.
- Crespi, M.** See *Enrique Moles*.
- Creutzfeldt, Walther Hildebrandt**, cathodic deposits from mixed solutions of two simple metallic salts, A., ii, 347.
- Creveld, S. van**, the distribution of chlorine in the blood, A., i, 287.
- Creveld, S. van**, and *R. Brinkman*, a direct demonstration of the impermeability of the corpuscles of man and of the rabbit for dextrose, A., i, 192.
- Creveld, S. van.** See also *J. de Haan*.
- Crippa, G.** See *G. Charrier*.
- Cristol, Paul**, zinc and cancer, A., i, 497.
 the estimation of the total non-protein nitrogen of serum; comparative study of trichloroacetic and metaphosphoric acids as protein precipitants, A., ii, 583.
- Cristol, Paul**, and *M. Simonnet*, estimation of the total non-protein nitrogen of serum; choice of a suitable albumin precipitant, A., ii, 887.
- Critchett, O. A.**, estimation of bismuth, A., ii, 90.
- Croce, M.** See *Riccardo Ciusa*.
- Crocker, Ernest C.**, application of the octet theory to single ring aromatic compounds, A., i, 927.
- Crommelin, Claude Auguste.** See *Émile Mathias*.
- Croxier, William John**, cell penetration by acids. V. The estimation of permeability changes, A., i, 897.
 cell penetration by acids. VI. The chloroacetic acids, A., i, 1090.
- Crut, G.** See *Ernst Berger*.
- Csányi, Wilhelm.** See *Richard Willstätter*.
- Csonka, Frank A.**, and *Grace C. Taggart*, trustworthiness of the Benedict and Folin-Wu blood-sugar estimations, A., ii, 879.
- Cullen, Glenn E.**, modification of the Clark hydrogen electrode vessel to permit accurate temperature control, A., ii, 611.
 acidosis. XIX. Colorimetric estimation of the hydrogen-ion concentration of blood plasma, A., ii, 672.
- Cullen, Glenn E.**, and *A. Baird Hastings*, comparison of colorimetric and electro-metric estimations of hydrogen-ion concentrations in solutions containing carbon dioxide, A., ii, 657.
- Cullen, Glenn E.** See also *Joshua Harold Austin*, *John P. Peters*, and *Donald D. van Slyke*.
- Cumming, William Murdoch**, the hydroferrocyanides and hydroferricyanides of the organic bases. I., T., 1287.
 apparatus for the estimation of methoxyl groups, A., ii, 232.
- Curie, (Mlle) Irène**, determination of the velocity of the α -rays of polonium, A., ii, 606.
- Curie, Maurice**, action of red and infra-red rays on the phosphorescent sulphides, A., ii, 246.
 the refractive indices of phosphorescent sulphides, A., ii, 801.
- Currey, Geoffrey Saunders**, the colouring matter of the scarlet pelargonium, T., 319.
 the colouring matter of red roses, A., i, 413.
- Curtis, W. E.**, structure of the band spectrum of helium, A., ii, 330.
- Curtius, Theodor**, and *Gustav Ehrhart*, decomposition of benzyl azide in indifferent media and in malonic ester, A., i, 775.
- Curtius, Theodor**, and *Friedrich Schmidt*, the action of sulphuryl azide on *p*-xylene, A., i, 776.
- Curtius, Theodor**, and *Wilhelm Sieber*, transformation of alkylated malonic acids into α -amino-acids. II. Syntheses of β -phenyl- α -alanine and of α -amino-*n*-butyric acid, A., i, 721.
- Cusmano, Guido**, catalytic reduction of nitrones, A., i, 148.
- Cutter, John Outram.** See *Thomas Martin Lowry*.
- Cuttica, Vittorio**, decomposition of potassium ferricyanide by the action of heat, A., i, 441.
 complex nitrites of nickel, A., ii, 448.
- Cuttica, Vittorio**, and *A. Paciello*, thalious nitrite in certain complex and double nitrites, A., ii, 377.
- Cwacha, J.** See *Julius Stoklassa*.
- Czechraleski, J.**, the solubility of gases in aluminium, A., ii, 646.

D.

Daeges, Karl, solubility limits of carbon in ternary steels. I. The system chromium-iron-carbon, A., ii, 70.

- Daevs, Karl**, solubility limits of carbon in ternary steels. II. The system tungsten-iron-carbon, A., ii, 70.
- Dafert, Otto von**, the effect of daylight on the content of active material in *Digitalis*, A., i, 97.
- Dalmer, J.** See *T. Merl*.
- Dalmer, Josef.** See *Roland Scholl*.
- Dains, Frank Burnett, Ruth Thompson, and William F. Asendorf**, the reactions of the formamides. X. The thioimidazolones [thiolglyoxalones], A., i, 1185.
- Dakin, Henry Drysdale**, synthesis of inactive para- and anti-hydroxy-aspartic acids (aminomalic acids) [aminohydroxysuccinic acids], A., i, 143.
resolution of hydroxyaspartic acids [aminohydroxysuccinic acids] into optically active forms, A., i, 430.
action of muscle tissue on fumaric, maleic, glutaconic, and malic acids, A., i, 792.
- Dale, Henry Hallett, and Harold Ward Dudley**, the pituitary active principles and histamine, A., i, 397.
physiological action of *N*-methyl-histamine and of tetrahydropyrido-3:4-iminazole ("iminazoleisopiperidin" of Frankel) [1:3:5-benztriazole], A., i, 403.
- Dalmer, O.** See *Carl Neuberg*.
- Damadé, R.**, examination of the duodenal fluid obtained through a tube, A., ii, 887.
- Damiens, A.**, the absorption of ethylene by sulphuric acid; preparation of ethyl alcohol, diethyl sulphate, and liquid hydrocarbons, A., i, 1105.
bromine normally present in animal tissues, A., ii, 79.
the "dynamic" allotropy of tellurium, A., ii, 498.
the crystallisation of amorphous tellurium, A., ii, 562.
- Damon, Samuel R.**, bacteria as a source of the water-soluble *B*-vitamin, A., i, 201.
- Danilof, Hildegard.** See *Gustav Reddellien*.
- D'Arbela, F.**, detection of lactic acid in organic liquids, A., ii, 460.
- Dardel, Jean Henri.** See *Friedrich Kehrmann*.
- Dardord, R.**, the reflection of *X*-rays by crystals, A., ii, 673.
- Darmois, Eugène**, two new ammonium molybdomalates, A., i, 220.
action of acids on ammonium molybdomalate, A., i, 522.
- Darwin, Charles Galton**, reflection of *X*-rays from imperfect crystals, A., ii, 416.
- Das, Radhakishen.** See *(Sir) Prafulla Chandra Ray*.
- Dash, Wilfred Marsden.** See *Nevil Vincent Sidgwick*.
- Datta, Rasik Lal, and Bibhu Charan Chatterjee**, halogenation. XXI. Some derivatives of carbamic esters; chlorine as a simultaneous oxidising and condensing agent, A., i, 815.
- Datta, Snehamoy**, the spectrum of glucinum fluoride, A., ii, 411.
the absorption spectrum of potassium vapour, A., ii, 678.
- Dandel, Victor**, spontaneous condensation of ethoxyacetone; formation of the corresponding aldol, α -diethoxy- β -methylpentan- β -ol- δ -one, A., i, 432.
- Dauvillier, A.**, analysis of the atomic structure, A., ii, 43.
the structure of the elements of mean atomic number, A., ii, 101.
the complexity of the *K*-series of the light elements and its theoretical interpretation, A., ii, 243.
the *L*-series of lutecium and ytterbium and the identification of celtium with the element of atomic number 72, A., ii, 463.
the precise measure of the layers of energy of the barium atom and the appearance of the *L*-spectrum of ionisation, A., ii, 542.
analysis of the electronic structure of the elements, A., ii, 559, 678.
- Dauvillier, A.** See also *Maurice de Broglie*.
- Davey, Wheeler P.** See *Albert W. Hall*.
- Davidhelser, L. Y., and Walter A. Patrick**, adsorption of ammonia by silica gel, A., ii, 262.
- Davidson, Arthur W.** See *James Kendall*.
- Davies, Ann Catherine.** See *Frank Horton*.
- Davies, Arthur Hugh, and Scottish Dyes, Ltd.**, production of hydroxy-derivatives of anthraquinone, A., i, 354.
- Davies, Arthur Hugh.** See also *John Thomas*.
- Davies, John Stanley Herbert.** See *Frederick Maurice Rowe*.
- Davies, William**, preparation of *p*-nitrophenylhydrazine and other aromatic hydrazines, T., 715.
the cumulative effect of the chlorine atom and the methyl and sulphonyl chloride groups on substitution in the benzene nucleus. III, T., 785.

- Davies, William**, an instance of the apparent effect of the entering group on the position of substitution in the benzene nucleus, T., 808.
- Davies, William**, and **Edgar Herbert Cuthbert Hickox**, the synthesis of *m*- α -benzopyrrole derivatives, T., 2840.
- Davies, William**, and **William Henry Perkin, jun.** [with **Herbert Clayton**], the chlorination and bromination of the toluic acids and the preparation of the phthalaldehydic acids, T., 2202.
- Davis, Clarke E.**, and **Earl T. Oakes**, physical characteristics of gelatin solutions, A., i, 597.
- Davis, Clarke E.**, **Earl T. Oakes**, and **Harold H. Browne**, viscosity of gelatin solutions, A., i, 63.
- Davis, C. W.**, separation of palladium and platinum by means of dimethylglyoxime, A., ii, 662.
- Davis, R. O. E.**, **L. B. Olmstead**, and **F. O. Lundstrum**, vapour pressures of ammonia-salt solutions, A., ii, 49.
vapour pressure of the system, lithium nitrate-ammonia, A., ii, 56.
- Davis, Tenney L.**, preparation of guanidine nitrate, A., i, 117.
the action of aqueous ammonia on dicyanodiamide, A., i, 118.
the action of sulphuric acid on nitroguanidine, A., i, 530.
the rôle of mercuric nitrate in the "catalysed" nitration of aromatic substances. II. Nitration of naphthalene, A., i, 818.
- Davison, C.**, and **C. H. Kunsman**, the scattering of electrons by nickel, A., ii, 251.
- Dawkins, Alfred Ernest**, heterogeneous equilibria; the ternary system sodium sulphate-sodium carbonate-water, T., 776.
- Day, James Nelson Edmund**. See **Oscar Lisle Brady**.
- Debucquet, L.**, composition of a rhinolith, A., i, 497.
- Decarrière, Eugène**, the rôle of gaseous impurities in the catalytic oxidation of ammonia; influence of hydrogen phosphide, A., ii, 284.
the rôle of gaseous impurities in the catalytic oxidation of ammonia gas, A., ii, 284.
- Decker, Herman**, natural system of carbon compounds. I. General statement of Mendeléev's law on the numerical relationships between primary, tertiary, and quaternary carbon atoms, A., i, 417.
- Decker, Herman**, natural system of carbon compounds. II. Empirical and rational allologous series and their graphical representation as a system, A., i, 513.
- Decker, Herman**, and **Paul Becker**, ring opening in the benzopyrrolum series, A., i, 358.
- Decker, Herman**. See also **Friedrich Kehrman**.
- Dede, L.**, the effect of scratching the wall of a vessel with a glass rod, A., ii, 744.
- Dede, L.**, and **P. Bonin**, the hindrance of precipitations with hydrogen sulphide by neutral chlorides, A., ii, 766.
- Deerns, W. W.**, the estimation of boric acid, A., ii, 867.
- Dehn, William Maurice**. See **George W. Fucher**.
- Deighton, Thomas**, some investigations on the electrical method of soil moisture estimation, A., i, 1227.
- Delange, Raymond**, the relationship of odour to molecular structure, A., i, 894.
- Delaplace, R.**, solubility of sulphur in certain organic liquids, A., ii, 706.
- Delauney, P.** See **Henri Hérissé**.
- Delague, A.** See **Georges Bruhat**.
- Delépine, Marcel**, the autoxidation of organic sulphur compounds, A., i, 621, 914.
complex iridium compounds, A., i, 859.
- Delépine, Marcel**, and **René Demars**, copper salts of aminosulphonic acids, A., i, 923.
- Delépine, Marcel**, and (*Mme*) **Pierre Jaffoux**, action of some acyclic halogenated derivatives on hexamethylenetetramine, A., i, 234.
- Deleuer, Maurice**, thermo-compression and thermo-addition, A., ii, 684.
- Delprat, G. D.**, and **G. H. Whipple**, liver function; benzoate administration and hippuric acid synthesis, A., i, 86.
- Demars, René**. See **Marcel Delépine**.
- Demichellis, L.** See **G. Charrier**.
- Demjanov, Nikolaus J.**, and **Marie Dojarenko**, methylenecyclobutane and dimethylcyclobutylmethylamine, A., i, 996.
the chlorohydrin and oxide of methylenecyclobutane and the transformation of the former into cyclopentanone, A., i, 1009.
vinylcyclopropane, certain derivatives of methylcyclopropylcarbinol and the isomerisation of the cyclopropane ring, A., i, 1014.

- Demjanov, Nikolaus J.**, and **Marie Dojar-enko**, preparation of cyclobutanone by the pyro-chemical decomposition of 1-hydroxycyclobutane-1-carboxylic acid, A., i, 1161.
- Demjanovski, S.**, the preparation of histidine from blood, A., i, 1052.
- Demolon, A.**, the sulphur-oxidising power of soils, A., i, 312.
the accessory elements of the de-phosphoration slags, A., ii, 564.
- Demoussy, Em.** See **Leon Maquenne**.
- Dempster, A. J.**, positive ray analysis of lithium and magnesium, A., ii, 417.
- Deneayer, M. E.**, the chemical constitution of felspars; analysis of two microclines, A., ii, 451.
- Denecke, G.** See **P. Morawitz**.
- Dengg, Rupert.** See **Alors Zinke**.
- Dengin, E. F.** See **Nicolai D. Zelinsky**.
- Denham, Henry George**, sub-salts of bismuth, A., ii, 218.
- Denham, Humphrey John**, the structure of the cotton hair and its botanical aspects, A., i, 615.
- Denigès, Georges**, identification of small quantities of terpeneol hydrate in complex mixtures, A., ii, 789.
- Denigès, Georges**, and **R. Tourrou**, micro-chemical reactions of "dulcin" [*p*-ethoxyphenylcarbamide], A., ii, 95.
- Denigès, Georges.** See also **C. Sauvageau**.
- Denis, Willey**, sulphates in blood, A., ii, 225.
estimation of magnesium in blood, plasma, and serum, A., ii, 659.
- Denis, Willey**, and **L. von Meysenbug**, a possible source of error in the Bell-Doisy method for the estimation of phosphates in blood plasma, A., ii, 584.
- Denis, Willey, Warren R. Sisson**, and **Martha Aldrich**, the effect produced on the composition of milk by the administration of certain inorganic and organic substances, A., i, 494.
- Denis, Willey.** See also **Warren R. Sisson**.
- Denison, Irving A.**, the nature of certain aluminium salts in the soil and their influence on ammonification and nitrification, A., i, 512.
- Dennett, John Horace.** See **John Albert Newton Friend**.
- Dennis, Louis Monroe**, and **F. E. Hance**, germanium. III. Germanium tetrabromide and germanium tetrachloride, A., ii, 302.
- Dennis, Louis Monroe**, and **Jacob Papish**, germanium. I. Extraction from germanium-bearing zinc oxide; non-occurrence in samarskite, A., ii, 150.
- Dernby, Karl Gustav**, extracellular bacterial proteases, A., i, 405.
- Dernby, Karl Gustav**, and **B. Allander**, influence of the hydrogen-ion concentration on the growth and formation of toxin of tetanus bacilli, A., i, 303.
- Dernikos, Diamandi.** See **Hans Pringsheim**.
- Dersin, Hans**, a new alkylamine and certain of its derivatives, A., i, 141.
- Derx, H. G.**, the configuration of ring systems in space, A., i, 651.
- Derx, H. G.** See also **Jacob Böseken**.
- Desgrez, Alexandre, Henri Bierry**, and **F. Rathery**, diabetes, β -hydroxybutyric acid, and *lævulose*, A., i, 1215.
- Deshapande, Shankar Shridhar**, and **Jocelyn Field Thorpe**, ring-chain tautomerism. II. The effect of the *gem*-diethyl group on the carbon tetrahedral angle, T., 1430.
- Dessemond, A.**, estimation of volatile matter in a coal, A., ii, 456.
- Detœuf, André**, monochlorocarbamide; preparation of chlorohydrins by its action on ethylenic hydrocarbons, A., i, 236, 327.
- Deuel, Harry J.**, and **Oskar Baudisch**, detection of thymine in the presence of sugar, A., ii, 670.
- Deuel, Harry J.** See also **Oskar Baudisch**.
- Dezani, Serafino**, genesis of thiosulphuric acid in animals, A., i, 968.
- Dhar, Nilratan**, some phenomena in the electromotive behaviour of certain metals, A., ii, 20.
the theory of reaction velocities, A., ii, 39.
the action of metals, such as copper and zinc, on an aqueous solution of ammonium nitrate, A., ii, 49.
new views on the constitution of chromic acid, A., ii, 382.
catalysis. XVI. Radiation as a factor in thermal and photochemical reactions, A., ii, 730.
- Dhar, Nilratan**, and **N. N. Mitra**, induced reactions and negative catalysis, A., ii, 630.
- Dhar, Nilratan.** See also **B. C. Banerji, Nitya Gopal Chatterji, Phani Bhusan Ganguly, N. N. Mitra, R. M. Purkayastha**, and **P. B. Sarkar**.
- Dhéré, Charles**, and **A. Schneider**, hæmocyanin. VI. and VII., A., i, 884.
- Di Capra, Clara**, solubility in the solid state of bismuth and cadmium in lead, A., ii, 576.

- Dickens, Frank, George Armand Robert Kon, and Jocelyn Field Thorpe**, the formation and stability of *spiro*-compounds. VIII. The Dieckmann-Komppa reaction, T., 1496.
- Dickinson, Roscoe G.**, crystal structures of complex cyanides of potassium with zinc, cadmium, and mercury, A., i, 532.
- crystal structures of potassium and ammonium stannichlorides, A., ii, 287.
- crystal structure of phosphonium iodide, A., ii, 640.
- Dickinson, Roscoe G., and Elbridge A. Goodhue**, crystal structures of sodium chlorate and sodium bromate, A., ii, 145.
- Dieckmann, Theodor, and E. Houdremont**, some compounds in the system $\text{CaO-P}_2\text{O}_5$, and their relation to basic slag, A., ii, 292.
- Dieckmann, Walter**, keto-enolic equilibria and Claisen's rule, A., i, 1020.
- Dieckmann, Walter, and Adolf Wittmann**, the relationship between the dimeric ketens and cyclobutane-1:3-dione and its derivatives, A., i, 1156.
- Diehl, Claus.** See *Emanuel Merck*.
- Diels, Otto.** The azo-ester reaction of amines and enols, A., i, 774.
- Diels, Otto** [with *Sörensen, Möhl, Aubart, Eckelmann, Kleinfeller, Harald Wackermann, and Fuldner*], the azo-ester reaction with amines and enols, A., i, 1194.
- Diels, Otto, and Harald Wackermann**, the structure of the compounds obtained by the oxidation of the additive products of β -naphthylamine and azo-esters, A., i, 1068.
- Dienes, Ludwig**, increase of nitrogen after fleshy and meal diets, A., i, 293.
- Dieter, Walter**, the capacity of yeast to degrade acid amides, A., i, 795.
- Dieterle, H.**, xanthosterol, A., i, 652.
- Diethelm, Alois.** See *Karl Jelinek*.
- Dietrich, C.** See *Georg Schroeter*.
- Dillon, Thomas, Rosalind Clarke, and Victor M. Hinchy**, chemical method of separating the isotopes of lead, A., ii, 710.
- Dilthey, Walther**, arylated pyridines. III. Quinodihydropyridines, A., i, 272.
- Dilthey, Walther** [with *H. Kaffer*], pyrylium compounds. XI. Pentaphenylpyrylium salts; the formulation of salts of dyes, A., i, 668.
- Dilthey, Walther** [with *H. Kaffer, Heinrich Meyer, and J. Nüsslein*], arylated pyridines and their relations to the corresponding pyrylium compounds. IV., A., i, 949.
- Dimbleby, Violet, Constance M. M. Muirhead, and William Ernest Stephen Turner**, effect of magnesia on the resistance of glass to corroding agents and a comparison of the durability of lime and magnesia glasses, A., ii, 709.
- Dimroth, Otto, and Theo Faust**, boric esters of hydroxyanthraquinones, A., i, 155.
- Dimroth, Otto, and Fritz Frister**, reduction of pyridine with zinc dust and acetic anhydride. II., A., i, 678.
- Dimroth, Otto, and Richard Heene**, reduction of pyridine with zinc dust and acetic anhydride, A., i, 48.
- Dimroth, Otto, and Valentin Hilleken**, anthradiquinones and anthraquinones, A., i, 158.
- Dimroth, Otto, Ernst Schultze, and Fritz Heinze**, the action of bromine on quinizarin and alizarin, A., i, 157.
- Dingemans, E.** See *J. V. Dubsky*.
- Dirken, M. N. J.**, relation between changes of temperature and consumption of oxygen by cold-blooded animals, A., i, 1207.
- Dirlam, H.** See *Julius von Braun*.
- Dische, Z.**, the distribution of chloride between corpuscles and plasma and the influence of carbon dioxide, A., i, 1086.
- Ditz, Hugo**, detection of manganese with benzidine and of cobalt by means of the thiocyanate reaction, A., ii, 229.
- Dixon, Henry Horatio, and Nigel G. Ball**, photosynthesis and the electronic theory. II., A., ii, 248.
- Dixon, Walter Ernest.** See *Douglas V. Cow*.
- Dobler, Friedrich**, kinetics of the reaction between ammonia and aromatic aldehydes, A., ii, 435.
- Dobrzanski, A. F.**, thermal analysis of the system *o*- and *p*-toluenesulphonamides, A., i, 1132.
- Doctor, E.** See *Ludwig Moser*.
- Dodd, Alfred Herbert**, the estimation of guanidine, A., ii, 536.
- Dodge, Francis D.**, vanillin glyceride, A., i, 748.
- Doerr, R., and W. Berger**, the oligodynamic effect of silver. IV., A., i, 1097.
- Dörries, W.** See *Adolf Windaus*.

- Doisy, Edward A., Emily P. Eaton,** and **K. S. Chouke**, buffer systems of blood-serum, A., i, 963.
- Dojarenko, Marie.** See *Nikolaus J. Domjanov.*
- Dolejšek, V.,** the lines K_{α} of the light elements, A., ii, 243.
N-series in X-ray spectra, A., ii, 411, 463.
- Dolenzek, Friedrich,** and **M. Schulze**, theory of binary mixtures. VII. Mixtures of ethyl ether and bromoform, A., ii, 118.
- Dombrowski, Stefan,** and **Stanislas Kozłowski**, the sterilisation of the intestine during fasting, A., i, 606.
- Dominik, W.,** the production of potassium hydrogen sulphate from ammonium hydrogen sulphate and potassium sulphate, A., ii, 145.
- Donat, Hans.** See *Wilhelm Steinkopf.*
- Donovan, F. K.** See *Alexander Hutcheon Bennett.*
- Dorabalska, Alicja,** thermochemical researches on oximes. II. The stereoisomeric ethyl esters of oximinacetoacetic acid, A., i, 725.
thermochemical researches on oximes. I. Ketoximes not exhibiting stereoisomerism, A., ii, 548.
- Doren, L. van, H. K. Parker,** and **Paul Lots**, use of the water interferometer as a pressure gauge, A., ii, 263.
- Doane, Paul,** a new method of colorimetry, A., ii, 518.
- Doubleday, (Miss) Ida,** boundary lubrication and chemical constitution; the optically active carbinols of the formula $C_2H_5 \cdot CH(OH) \cdot C_nH_{2n+1}$. T., 2875.
- Doubler, Francis H.** See *Leon Asher.*
- Doucet, A.** See *Émile Luce.*
- Doughty, Howard Waters,** and **Benjamin Freeman**, trihalogenmethyl reactions. IV. Reaction of trichloroacetic acid with copper, A., i, 427.
- Downs, Charles R.,** catalytic oxidation of naphthalene, A., i, 27.
preparation of maleic acid, A., i, 984.
preparation of anthraquinone, A., i, 1039.
- Downs, Charles R.** See also *John Maurice Weiss.*
- Dox, Arthur Wayland,** and **Lester Yoder**, *spiropyrimidines*. III. Condensation of cyclopropane-1:1-dicarboxylic ester with carbamides, A., i, 180.
pyrimidines from alkylmalonic esters and aromatic amidines, A., i, 374.
- Dox, Arthur Wayland,** and **Lester Yoder**, the reaction between *s*-dichlorodimethyl ether and ethyl malonate, A., i, 429.
benzylalkylbarbituric acids, A., i, 681.
some derivatives of *n*-butylmalonic acid, A., i, 808.
amide formation from esters of secondary alkylmalonic acids, A., i, 816.
- Doyon, M.,** use of frogs to demonstrate the anticoagulating action of nucleic acids, A., i, 82.
action of nucleic acid injected into the organism; immunisation by a single injection, A., i, 1087.
- Drakeley, Thomas James** [with *Frederick William Smith*], the ultimate composition of British coals, T., 221.
- Dreifuss, Max,** tellurium lead and tellurium antimony alloys, A., ii, 503.
- Dreifuss, Max.** See also *W. D. Treadwell.*
- Drekopf, K.** See *Alfred Benrath.*
- Dresbach, Melvin.** See *Arthur Knudson.*
- Dresel, K.,** and **F. H. Lewy**, sugar regulation in paralysis agitans, A., i, 969.
- Drew, Harry Dugald Keith.** See *Gilbert Thomas Morgan.*
- Dreyfus, Henry,** preparation of alkyl sulphates, A., i, 517.
- Driggs, F. H.** See *B. Smith Hopkins.*
- Driver, John,** and **James Brierley Firth**, the sorption of saturated vapours by charcoal, T., 2409.
- Druce, John Gerald Frederick,** isopropylstannonic acid and its derivatives, T., 1859.
the interaction of methyl iodide and potassium plumbite, A., i, 516.
the preparation and properties of organic stanno- and stanni-chlorides. IV. Some diamine stannichlorides, A., i, 639.
the preparation and properties of organic stanno- and stanni-chlorides. V. The salts of certain special bases, A., i, 1206.
- Drummond, Jack Cecil,** and **Robert Keith Cannan**, tethelin—the alleged growth-controlling substance of the anterior lobe of the pituitary gland, A., i, 491.
- Drummond, Jack Cecil,** and **Arthur Frederick Watson**, the testing of foodstuffs for vitamins, A., ii, 596.
the sulphuric acid test for liver oils, A., ii, 685.
- Drummond, Jack Cecil.** See also *Henry Lyster Jameson.*
- Duane, William,** calculation of the X-ray absorption frequencies of the chemical elements. I. and II., A., ii, 104.

- Duane, William**, and **Hugo Fricke**, absorption of X-rays by chromium, manganese, and iron, A., ii, 804.
- Duane, William**, and **R. A. Patterson**, X-ray spectra, A., ii, 463.
- Duane, William**. See also **George L. Clark**.
- Dubin, Harry E.** See **Casimir Funk**.
- Dubin, Michael**. See **Nikolai Schilov**.
- Duboe, (Mlle) T.**, the action of tribromoxyleneol on the tubercle bacillus, A., i, 972.
- Dubosc, André**, chemical analysis of caoutchouc articles, A., ii, 91.
- Duboux, Marcel**, calculation of the second dissociation constant of dibasic acids from the hydrogen-ion concentration, A., ii, 346.
- Dubrisay, René**, the action of boric acid on mannitol in alkaline solution, A., i, 1110.
the partial miscibility of liquid couples; applications to reactions between dissolved substances, A., ii, 428.
- Dubsky, J. V.** [with **P. Aptekmann**], neutralisation of the affinity of main and subsidiary valencies in compounds of a higher order. III., A., i, 104.
- Dubsky, J. V.**, and **E. Dingemane**, diketopiperazines. XI. 3:5-Diketol-1-benzylhexahydro-1:4-diazine, A., i, 56.
- Dubsky, J. V.**, and **E. Hoher**, diketopiperazines. XII. Attempted preparation of 1-benzoyl-3:5-diketohexahydro-1:4-diazine, A., i, 57.
- Dubsky, J. V.**, **E. Hoher**, and **E. Dingemane**, diketopiperazines. XIII., A., i, 57.
- Dubsky, J. V.** See also **Hilmar Johannes Backer**.
- Duchaček, Franz**, the diastatic value of malt, A., i, 974.
- Duchon, Frantisek**. See **Antonin Némec**.
- Duckham, (Miss) Rebecca**. See **William Howieson Gibson**.
- Duclaux, J.**, the mechanism of continuous luminous radiation, A., ii, 99.
- Ducloux, Enrique Herrero**, use of cesium chloride in microchemistry, A., ii, 77.
- Dudley, Harold Ward**. See **Henry Hallett Dale**.
- Düring, A.** See **A. Behre**.
- Dürrwächter, E.** See **Alexander Gutbier**.
- Dütsmann, Amadeus**. See **Fritz Straus**.
- Duff, James Cooper**, complex metallic amines. VII. Conductivities of diethylenediaminecobaltic bromides, T., 450.
- Duffenback, O. S.**, dissociation of hydrogen in a tungsten furnace and low voltage arcs in monatomic gases, A., ii, 620.
- Duffieux, M.**, the mass of the particles that emit the spectrum of carbon monoxide, A., ii, 597.
- Duffour, Alexis**, a new example of hemihedral forms not conforming to the sign of the optical activity, A., i, 829.
- Dufraisse, Charles**, ethylenic stereoisomerism, A., i, 534.
- Dufraisse, Charles**, and **Pierre Gérauld**, the supposed true dibenzoylmethane of Wislicenus; new experiments, A., i, 39.
the action of alcohols on α -bromobenzylidenacetophenone, A., i, 843.
- Dufraisse, Charles**. See also **Charles Mouren**.
- Dufton, Arthur Felix**, the separation of miscible liquids by distillation. II., T., 306.
- Duin, C. F. van**, steric hindrance of the sulpho-acid group, A., i, 139.
the addition of bromine to the trans-substituted cinnamic acids, A., i, 737.
- Dumesnil, Philippe**, the hydrolysis of the mono- and di-ethyl esters of diethylmalonic acid, A., i, 520.
the esterification of ethyl hydrogen diethylmalonate and of diethylmalonic acid, A., i, 622.
some derivatives of ethyl hydrogen diethylmalonate, A., i, 807.
- Dunbrook, Raymond F.** See **Alexander Lowy**.
- Duncan, Herbert Maxwell**. See **Edward Charles Cyril Baly**.
- Dundon, Merle L.**, and **William E. Henderson**, measurement of solubility by floating equilibrium; solubility of lead acetate, A., ii, 552.
- Dunham, Theodore, jun.** See **Theodore William Richards**.
- Dunn, Max S.**, and **Howard B. Lewis**, the action of nitrous acid on casein, A., i, 279.
the hydrolysis of casein and deaminised casein by proteolytic enzymes, A., i, 292.
- Dunn, Max S.**, and **Carl L. A. Schmidt**, the influence of position and of temperature on the reaction of aliphatic amino-nitrogen with nitrous acid, A., i, 1124.
- Dunoyer, Louis**, the luminescence of gas in the electrodeless discharge; the induction spectra of caesium and rubidium, A., ii, 729.

Dunstan, Albert Ernest, and Ferdinand Bernard Thole, the chemical nature of mineral lubricating oils, A., i, 213.

Dunstan, Arthur St. C., and Benjamin A. Wooten, arc-cathode spectra, A., ii, 99.

Duparc, Louis, and L. Ramadier, the volatilisation of arsenic and antimony by means of methyl alcohol, A., ii, 640.

Dupont, Georges, the composition of the essential oil of turpentine from Aleppo, A., i, 357.
the formulæ of dicyclic terpenes, A., i, 1042.

Dupont, Georges. See also *Maurice Vêres*.

Durand, J. F., catalysts and chemical equilibrium, A., ii, 701.

Durrans, Thomas Harold, the action of sulphuryl chloride on organic substances. I. Simple monosubstituted benzenes, T., 44.
absorption of ultra-violet light, A., ii, 7.

Durrant, Reginald Graham. See *Henry Bassett*.

Duschek, Alois. See *Robert Müller*.

Dutoit, Paul, and Édouard Grobet, carrying down by precipitates, A., ii, 550.

a new physico-chemical method of volumetric analysis applied to some problems of inorganic chemistry, A., ii, 578.

Dutt, Sikhibushan, dyes derived from phenanthraquinone. II. Naphthaflavindulines, T., 1951.

dyes derived from "saccharin"; the sulphamphthaleins, T., 2389.

Dutt, Sikhibushan, and Nirmal Kumar Sen, dyes derived from $\alpha\alpha'$ -dicyanodibenzyl diketone, T., 2663.

Dutt, Sikhibushan. See also *Anukul Chandra Sircar, and Edwin Roy Watson*.

Dux, Paul, and Artur Löw, osmotic behaviour of frog's muscle, deficient in water and poisoned by glycerol, and on the shrinkage of muscle protein, A., i, 398.

Duyzen, Franz. See *Erich Schmidt*.

Dvořák, K. See *V. Veselý*.

Dyer, J. W. W., and Amy R. Watson, determination of sulphur in vulcanised rubber, A., ii, 656, 782.

Dziwowski, Karol, and J. Suszko, the pyrogenic transformation of fluorene, A., i, 730.

E

Eakle, Arthur Starr, massive troilite from Del Norte Co., California, A., ii, 358.

Earl, John Campbell. See *James Colquhoun Irvine*.

Early, Reginald George, and Thomas Martin Lowry, the properties of ammonium nitrate. III. Ammonium nitrate and sodium nitrate, T., 963.

Eastman, Ermon Dwight, double and triple bonds, and electron structures in unsaturated molecules, A., ii, 367.

equilibria in the systems iron-carbon-oxygen and iron-hydrogen-oxygen, and the free energies of the oxides of iron, A., ii, 506.

Eaton, Elgar O., estimation of monobromocamphor, A., ii, 326.

Eaton, Emily P. See *Edward A. Doisy*.

Ebert, Ludwig, newer investigations on the anomaly of the strong electrolytes, A., ii, 113.

Ebin, D. See *J. Altwegg*.

Ebler, Erich, and A. J. van Rhyn, the radioactivity of mineral springs. II., A., ii, 16.

the adsorption of radioactive substances. II., A., ii, 16.

Eckart, Hanns, beef bone fat and neat's foot oil, A., i, 967.

Eckelmann, See Otto Diels.

Eckert, Alfred, and Gertrud Endler, highly condensed derivatives of xanthone, A., i, 941.

Eckert, Alfred, and Rudolf Endler, chlorination of quinol, A., i, 932.

Eckert, Alfred, Fritz Seidel, and Gertrud Endler, phenylcarbazole, A., i, 952.

Eckert, G. See *Fritz Arndt*.

Eckstein, H. See *Alfred Benrath*.

Edelmann, R. F. See *W. D. Treadwell*.

Eder, Alois, the estimation of cobalt in steel, A., ii, 458.

Eder, Robert, and C. Widmer, derivatives of β -methylantraquinone. I. Syntheses of chrysophanic acid [1:8-dihydroxy-3-methylantraquinone] and of 1:5-dihydroxy-3-methylantraquinone, A., i, 260.

Éderer, Stefan, the influence of the concentration of sugar on the synthesis of glycogen, A., i, 895.

Edgar, Graham, and R. B. Purdum, rapid electrolysis without rotating electrodes, A., ii, 547.

- Edgar, Graham**, and **W. O. Swan**, factors determining the hygroscopic properties of soluble substances. I. Vapour pressures of saturated solutions, A., ii, 949.
- Edgar, Graham**. See also **Arthur Becket Lamb**.
- Edge, Stephen Rathbone Holden**, benzobisthiazoles, T., 772.
- Edlbacher, Siegfried**, the free amino-groups of the proteins, A., i, 279. the proteic acids of urine. I. The hydroxyproteic acids, A., i, 692. the hydroxyproteic acids, A., i, 838.
- Edlbacher, Siegfried**, and **Berthold Fuchs**, the influence of β -naphthalene-sulphonyl chloride on the proteins, A., i, 279.
- Edwards, C. A.**, and **A. J. Murphy**, the rate of combination of copper and phosphorus at various temperatures, A., ii, 645.
- Effront, Jean**, the distinctive properties of amylases from different sources, A., i, 184. effect of filtration on amylases, A., i, 1076. characteristic properties of amylases of different origin, A., i, 1076.
- Ege, Rich**. See **Valdemar Henriques**.
- Egerer-Seham, Grete**, and **C. E. Nixon**, chemistry of blood and spinal fluid, A., i, 395.
- egger, L.** See **W. D. Treadwell**.
- Eggert, John**, sensitiveness of very sensitive explosives, A., ii, 138.
- Eggert, John**, and **W. Noddack**, verification of the photochemical equivalent law with photographic dry plates, A., ii, 9.
- Ehmer, Wilhelmine**. See **Alois Zinke**.
- Ehrenberg, Rudolf**, protein enzymes, A., i, 597.
- Ehrenberg, Rudolf**, and **Alfred Karsten**, urine iron, and renal function, A., i, 967.
- Ehrenfest, P.**, the difference between series spectra of isotopes, A., ii, 598.
- Ehrhart, Gustav**. See **Theodor Curtius**.
- Ehrlich, Josef**. See **Ludwig Moser**.
- Eichelberger, Marietta**, the carbohydrate content of navy beans, A., i, 799.
- Eichler, Fritz**. See **Erwin Ott**.
- Eichwald, Egon**, the action of the brush discharge on fatty acids and their glycerides, A., i, 982.
- Eiffaender, L.** See **A. Lüttringhaus**.
- Elliott, August**, construction of platinum film electrodes and their method of use, A., ii, 610.
- Elbeek, H.** See **Georg Schroeter**.
- Eirich, Lina**. See **H. Finger**.
- Eisenhardt, W.**, and **R. Schaefer**, variations in chloride-metabolism due to menstrual processes, A., i, 88.
- Eisenlohr, Fritz**, the molecular refraction of substances of higher melting point and the calculation of the corresponding refractive exponents to the temperature of comparison, 20°, A., ii, 1.
- Ekeley, John Bernard**, **E. C. Rogers**, and **Margaret Swisher**, the action of acetic anhydride on some benzyldeneanthranilic acids. V. A., i, 934.
- Ekkert, L.**, colour reactions of phenacetin and acetanilide, A., ii, 169.
- Elam, Constance F.** See **Henry Cort Harold Carpenter**.
- Elder, Robert B.**, measurement of surface tension, A., ii, 618.
- Eldridge, E. F.** See **Dwight T. Ewing**.
- Elektrochemische Werke G. m. b. H.**, **Heinrich Bosshard**, and **David Strauss**, preparation of water-soluble derivatives of aryl ethers of higher aliphatic alcohols, A., i, 913.
- Elias, Herbert**, and **Ubaldo Sammartino**, the part played by acid in carbohydrate metabolism. IV. The relation between acid and alkali and adrenaline-glycosuria, A., i, 86.
- Elias, Herbert**, and **E. A. Spiegel**, tetany, A., i, 401.
- Elias, Herbert**, and **St. Weiss**, the rôle of acid in carbohydrate metabolism. V. The action of acid and alkali on the carbohydrate metabolism of yeast-cells, A., i, 485. the action of the phosphate-ion on blood and urinary sugar, A., i, 1085.
- Eliasberg, Paul**. See **S. Kostytachev**.
- Eller, Wilhelm**, synthesis of the humic acids, A., i, 326.
- Eller, Wilhelm**, and **L. Klemm**, the action of sulphuryl chloride on aromatic amines, A., i, 448.
- Ellinger, Alexander**, and **S. M. Neuschloss**, comparison of the viscosity and velocity of ultra-filtration of serum, A., i, 484.
- Ellinger, Philipp**, the influence of potassium and of radioactivity on the oxidation velocity of the red corpuscles, A., i, 288. the pharmacology of cell respiration, A., i, 492.
- Elliott, Felix A.** See **Samuel Edward Sheppard**.
- Elliott, George Robert**, preparation and properties of the benzochloroamides, T., 202.

- Ellis, C. D., β -ray spectra and their meaning, A., ii, 389.
the interpretation of β -ray and γ -ray spectra, A., ii, 466.
- Ellis, C. D. See also James Chadwick.
- Ellsworth, H. V., and E. Poitevin, camesselite, a new borate mineral from British Columbia, A., ii, 804.
- Ellsworth, H. V. See also R. A. A. Johnston.
- Elschner, Carl, colloidal phosphates, A., ii, 714.
- Elsey, Howard M. See Hamilton Perkins Cady.
- Embsen, Gustav, and Erich Adler, physiological significance of the change in the condition of permeability in the limiting membrane of the muscle fibres, A., i, 196.
- Embsen, Gustav, and Heinz Lawaczek, formation of phosphoric acid in the contraction of frog's muscle, A., i, 492.
- Emerique, L. See L. Lapicque.
- Emery, William O., and Clarence D. Wright, distribution of certain drugs between immiscible solvents, A., ii, 205.
- Emmert, Bruno, and Paul Farr, action of iodine on *NN'*-dialkyltetrahydro-4:4'-dipyridyls, A., i, 179.
- Emmert, Bruno, and Otto Varenkamp, 1:1'-dialkyltetrahydro-4:4'-dipyridyls, A., i, 1064.
- Emmert, Bruno, and Otto Werb, 1:1'-dimethyltetrahydro-4:4'-dicollidyl [1:2:4:6:1':2':4':6'-octamethyltetrahydro-4:4'-dipyridyl], A., i, 680.
- Emelander, R. See Alexander Guthrie.
- Emster, Konrad van. See Hans Meerwein.
- Enderlin, Fr. See Karl W. Rosenmund.
- Endler, Gertrud. See Alfred Eckert.
- Endler, Rudolf. See Alfred Eckert.
- Endres, Gustav, the relation between the true reaction of the urine and the alveolar tension of carbon dioxide, A., i, 1214.
- Engeland, R., the free amino-groups of the proteins, A., i, 279.
the detection and estimation of mono-amino-acids, A., ii, 536.
- Engfeldt, N. O., the effect of Dakin's hypochlorite solution on certain organic substances, A., i, 812.
- England, E. H. See Henry Droop Richmond.
- Englis, Duane T., and Chuk Yee Tsang, the clarification of solutions containing reducing sugars by basic lead acetate; the effect of different de-leading agents, A., ii, 459.
- Engström, N. H., temperature correction in bomb calorimetry, A., ii, 27.
- Ennos, F. R. See W. F. P. McLintock.
- Ens, H. See Karl Jellinek.
- Entat, M., and E. Vulquin, detection and estimation of free sulphuric acid and sulphoacetates in cellulose acetates, A., ii, 533.
- Ephraim, Fritz, and Eduard Michel, metallic hydrides. II. Hydrides of the alkaline-earth metals and of lithium, A., ii, 58.
- Ephraim, Fritz, and Paul Mesimann, solubility. VI. Solubility of ammoniates, A., ii, 574.
- Eppinger, Hans, chemistry of amyloid degeneration, A., i, 497.
- Eppley, Marion, and Warren C. Vosburgh, electrometric titration of dichromate with ferrous sulphate, A., ii, 876.
- Erdenbrecher, A., sodium silicate, A., ii, 444.
- Erdeitein, F., and L. Fürth, the action of polished metals on toxius, A., i, 90.
- Erickson, E. Theodore, tschermigite (ammonium-alum) from Wyoming, A., ii, 304.
- Ericson, Goeta. See Hans von Euler.
- Ernst, A., a new apparatus for washing gases, A., ii, 839.
- Ernst, Z., and B. Szappanyos, formation of bilirubin in surviving spleen, A., i, 1089.
- Ernst, Z., and St. Weiss, modification of Bang's method for the estimation of sugar in the blood, A., ii, 724.
- Ernström, Efr., the temperature coefficients in the degradation of starch and the thermostability of malt diastase and ptyalin, A., i, 599.
- Errera, J., dielectric constants of colloidal solutions, A., ii, 694.
- Ertl, Arthur. See Roland Scholl.
- Ertl, K. See Ludwig Moser.
- Eschenhagen, Gerhard. See Carl Tubandt.
- Eskola, Pentti, the silicates of strontium and barium, A., ii, 849.
- Espenschied, H. See Alfred Benrath.
- Estermann, I. See M. Volmer.
- Estill, Howard Wilmot. See George Shannon Forbes.
- Etienne, G., and M. Vérain, partition of urea in blood, A., i, 963.
- Ettisch, M., Michael Polanyi, and K. Weissenberg, fibrous structure of hard drawn metal wires, A., ii, 201.
- Eucken, Arnold, theory of adsorption processes, A., ii, 262, 550.
moment of inertia and structure of the carbon dioxide molecule, A., ii, 849.

- Eucken, Arnold**, and **O. Neumann**, the possible existence of metallic compounds in the state of vapour, A., ii, 567.
- Eucken, Arnold**. See also **R. Bartels**.
- Euler, (Mme) Astrid Cleve von**, catechu and catechin, A., i, 45.
toxicity of methyl alcohol, A., i, 90.
chemical constituents of pine leaves. II., A., i, 100.
the lignin-like resins and tannins of spruce needles, A., i, 233.
- Euler, Hans von** [with **Arvid Hedelius** and **Zimmerlund**], ionic equilibria on metallic surfaces, A., ii, 251.
- Euler, Hans von**, and **Stig Bergman**, combination of iodine and starch, A., i, 921.
- Euler, Hans von**, and **Gösta Ericson**, degree of dispersion of saccharase, A., i, 886.
- Euler, Hans von**, and **Ulf von Euler**, the formation of oxides of multivalent metals from their hydroxides, A., ii, 831.
- Euler, Hans von**, and **A. Fahlander**, separation of halogens from organic compounds, A., ii, 360.
- Euler, Hans von**, and **K. Josephson**, *Saccharomyces marzianus* and top fermentation yeast *R*, A., i, 706.
a silver compound of saccharase, A., i, 1076.
- Euler, Hans von**, and **Signe Karlsson**, accelerators of fermentation, A., i, 972.
- Euler, Hans von**, and **Sture Landergren**, combination of iodine and starch. II., A., i, 921,
the inactivation of saccharase by iodine, A., i, 1076.
- Euler, Hans von**, and **Karl Myrbäck**, vitamin-B and co-enzymes. II., A., i, 293.
the thermolability of the sucrose-splitting enzyme of the human jejunum, A., i, 295.
compounds of iodine with constituents of starch, A., i, 527.
the conditions of acidity and thermolability of saccharase, A., i, 693.
inactivation of saccharase by small quantities of silver salts, A., i, 959.
the absorption of iodine by starch, A., i, 1120.
enzyme chemistry, A., i, 1201.
- Euler, Hans von**, and **Folke Nordlund**, the enzymic synthesis of fructose-diphosphate (hexosephosphate), A., i, 306.
- Euler, Hans von**, and **Olof Svanberg**, carboxylic esters as amphoteric electrolytes, A., i, 219.
the regeneration of inactivated saccharase by dialysis, A., i, 284.
the saccharase of the intestine, A., i, 296.
the phosphorus content of purified saccharase preparations, A., i, 1200.
- Euler, Hans von**, and **G. Zimmerlund**, salt adsorption on metal surfaces, A., ii, 822.
- Euler, Hans von**. See also **G. J:son Blohm**.
- Euler, Karl**. See **Stefan Goldschmidt**.
- Euler, Ulf von**. See **Hans von Euler**.
- Evans, Bernard Scott**, the estimation of small quantities of antimony in copper and brass, A., ii, 231.
- Evans, C. A. Lovatt**, acid production in shed blood, A., i, 890.
- Evans, Edward Victor**. See **Henry Edward Armstrong**.
- Evans, Richard Bromley**. See **Archibald Edwin Goddard**.
- Evans, Ulrick R.**, passivity and overpotential, A., ii, 814.
- Evans, William Lloyd**, and **Paul R. Hines**, the oxidation of potassium acetate to potassium oxalate, A., i, 803.
- Evans, William Lloyd**, and **Ora L. Hoover**, the oxidation of acetylcarbinol with potassium permanganate, A., i, 911.
- Evans, William Lloyd**, and **Cloyd D. Looker**, the influence of potassium hydroxide on the formation of vinyl alcohol from acetaldehyde, A., i, 102.
- Evans, William Lloyd**, and **Lily Bell Sefton**, the oxidation of isopropyl alcohol with potassium permanganate, A., i, 1108.
the oxidation of acetone with potassium permanganate, A., i, 1115.
- Eve, Arthur Stewart**, ionisation potential and the size of the atom, A., ii, 364.
- Evenden, James**. See **H. V. Atkinson** and **Graham Lusk**.
- Everest, Arthur Ernest**, and **Archibald John Hall**, the tinctorial properties of some anthocyanins and certain related pigments. II., A., i, 265.
- Evers, E. W.** See **George Leslie Kelley**.
- Evers, Fritz**. See **Carl Dietrich Harries**.
- Evers, Norman**, method for the separation of strychnine from quinine, A., ii, 669.
- Evers, Norman**. See also **John Leonard Lisius**.
- Ewart, Alfred James**, synthesis of sugars from formaldehyde, carbon dioxide, and water, A., i, 10.

- Ewbank, (Miss) Elinor Katharine.** See *Nevil Vincent Sidgwick.*
- Eweyk, C. van, and M. Tennenbaum,** secretin. I. and II., A., i, 397.
- Ewing, Dwight T., and E. F. Eldridge,** electrometric titration of uranium with potassium permanganate and potassium dichromate, A., ii, 661.
- Ewing, Dwight T.** See also *William Draper Harkins.*
- Ewing, (Sir) J. Alfred,** the quantum mechanism in the atom, A., ii, 633.
- F.**
- Fabre, René,** the constitution of resorcinol and some of its derivatives, A., i, 1147.
- a reaction of veronal and of the hypnotics derived from barbituric acid, A., ii, 795.
- Fabriques de Produits chimique de Thann et de Mulhouse,** preparation of borneol, A., i, 560.
- Fabris, E.,** solubility of potassium ferrocyanide in water; ice curve and cryohydric point, A., i, 329.
- Fabris, Stanislas,** cholesterol in cerebrospinal fluid, A., i, 969.
- Färber, Eduard.** See *Erik Häggglund.*
- Fahlander, A.** See *Hans von Euler.*
- Fabron, Wilhelm,** colophenic acids, A., i, 423.
- the analysis of partly hydrolysed fats. II., A., ii, 401.
- Faillebin,** mixed organometallic compounds of aluminium, A., i, 119.
- Fairbairn, (Miss) Ruth.** See *William Howieson Gibson.*
- Fairbrother, Fred,** cell for the observation of colloidal solutions for use with substage ultra-condensers, A., ii, 485.
- Fairbrother, Fred, and Enoch Swan,** the dissolution of gelatin, T., 1237.
- Fairbrother, Thomas H., and Arnold Renshaw,** the relation between chemical constitution and antiseptic action in the coal tar dyes, A., i, 612.
- Fairhall, L. T.,** lead. I. Estimation of minute amounts of lead in biological material, A., ii, 659.
- Fajans, Kasimir,** action of forces between separated atoms in diamond and aliphatic molecules, A., ii, 194.
- the energy of the atomic linkings in diamond and in aliphatic hydrocarbons, A., ii, 818.
- Falcicola, Pietro,** cinnamic acid in cryoscopy, A., ii, 421.
- reaction between thiosulphuric and nitrous ions, A., ii, 453.
- Falcicola, Pietro,** detection of the nitrous anion, A., ii, 783.
- Fales, Harold A., and Jacques C. Morrell,** new type of sodium lamp for polarimetry, A., ii, 43.
- velocity of inversion of sucrose as a function of the thermodynamic concentration of the hydrogen-ion, A., ii, 832.
- Fales, Helen L.,** analysis of camel's colostrum, A., i, 1090.
- Falk, Kurt.** See *Alfred Lottermoser.*
- Falkenheim, Susanne.** See *Adolf Sonn.*
- Falkner, (Miss) D. H.** See *(Miss) Mabel Harriet Norris.*
- Falta, Wilhelm, and M. Richter-Quittner,** the distribution of sugar between corpuscles and plasma, A., i, 696.
- Faltis, Franz, and Mathilde Krausz,** the course of the decomposition of apomorphine by reduction by Emde's method, A., i, 676.
- Faltis, Franz, and Felix Neumann,** alkaloids of the pareira root. II. isochondodendrine, A., i, 569.
- Faltis, Franz, and Carla Ruiz de Roxas,** some transformations of ethyl γ -bromo- α -dicarboxy-glutaconate, A., i, 624.
- Faragher, W. F., and Frederic Horace Garner,** the elimination of hydrogen chloride from chlorohydrocarbons, A., i, 3.
- Farbenfabriken vorm. F. Bayer & Co.,** preparation of derivatives of cholic acid, A., i, 554.
- preparation of complex iron compounds of the phosphoric esters of higher aliphatic polyhydroxyl compounds, A., i, 631.
- preparation of symmetrical aryl alkyl ethers, A., i, 934.
- preparation of bromodialkylacetylcarbamides, A., i, 992.
- preparation of mono-substituted sulphonamides, A., i, 999.
- Farbwerke vorm. Meister, Lucius, & Brünig,** preparation of anthraquinone derivatives [1:2-anthraquinonylisooxazoles], A., i, 178.
- preparation of esters of dihydroxy-diethyl sulphide, A., i, 420.
- preparation of new therapeutically active acridine derivatives, A., i, 468.
- preparation of dialkylaminoalkyl compounds, A., i, 529.
- preparation of β -aracyl- α -dialkyl-aminoethylbutyric esters, A., i, 639.
- preparation of ring ketones of the quinoline series, A., i, 867,

- Farbwerke vorm. Meister, Lucius, & Brüning**, preparation of aminoacetyl compounds of 4-amino-1-phenyl-2:3-dialkylpyrazolones, A., i, 954.
preparation of carbamide from cyanamide, A., i, 993.
preparation of aromatic selenium compounds, A., i, 1066.
- Fargher, Robert George**, and **William Herbert Gray**, chemotherapy of antimony; comparison of antimony tartrates with organic compounds of antimony, A., i, 404, 705.
- Fargher, Robert George**, and **John Charles Withers**, the chemical constituents of raw cotton, A., i, 615.
- Fargher, Robert George**. See also **Percy Herbert Clifford**.
- Farmer, Ernest Harold**, the addition of ethyl sodiocyanoacetate and ethyl sodiomalonate to ethyl muconate, T., 2015.
- Farmer, Ernest Harold**, **Christopher Kelk Ingold**, and **Jocelyn Field Thorpe**, the chemistry of polycyclic structures in relation to their homocyclic unsaturated isomerides. II. Intramolecular tautomerism, T., 128.
- Farmer, Robert Crosbie**, the velocity of decomposition of high explosives in a vacuum. III. Mercuric fulminate, T., 174.
- Fauré-Fremiet, E.**, constitution of the egg of *Sabellaria alveolata*, L., A., i, 85.
- Fauré-Fremiet, E.**, and (*Mlle*) **H. Garrault**, constitution of the ovarian egg of the carp (*Cyprinus carpio*), A., i, 700.
constitution of the egg of the trout (*Trutta fario*), A., i, 700.
- Fauré-Fremiet, E.**, and (*Mlle*) **du Vivier de Streel**, chemical constituents of the egg of the common frog (*Rana temporaria*) and their rôle in its embryonic development, A., i, 85.
- Faurholt, Carl**, the processes

$$\text{NH}_2 \cdot \text{CO} \cdot \text{ONH}_2 + \text{H}_2\text{O} \rightleftharpoons$$

$$(\text{NH}_4)_2\text{CO}_3 \text{ and } \text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons$$

$$\text{H}_2\text{CO}_3, \text{ A., ii, 272.}$$
the ammonium carbonate-carbamate equilibrium, A., ii, 747.
- Faust, E. S.**, camphoric acid derivatives, A., i, 748.
- Faust, Otto**, binary liquid mixtures, A., ii, 423.
- Faust, Theo**. See **Otto Dimroth**.
- Favreau, M.** See **A. Labat**.
- Fayolle**, and **Ch. Lormand**, perforation apparatus for the extraction of liquids by liquids; non-miscible liquids, A., ii, 759.
- Fazi, Remo de**, syntheses of new glucosides, A., i, 755.
- Fazi, Remo de**. See also **Romolo de Fazi**.
- Fazi, Romolo de**, and **Remo de Fazi**, action of ultra-violet rays on *Saccharomyces cerevisiae*, A., i, 1219.
- Feer, Emanuel**. See **Émile Cherbuliez**.
- Feigl, Friedrich**, the oxidising properties of carbon suspensions, A., ii, 51.
the use of benzidine in the detection of phosphoric acid, A., ii, 865.
- Feist, Franz**, isomerism in the glutamic acid series, A., i, 521.
action of bromine on methylal, A., i, 912.
- Feist, Franz**, and **Paul Karl Breuer**, the two β -methylglutaconic acids, A., i, 521.
 $\alpha\beta$ -dimethylglutaconic acid, A., i, 521.
- Feist, Franz**, **Paul Karl Breuer**, and **Bernh. Lubricht**, α -carboxy- β -phenyl- α -methylglutaconic esters and the isomerism of the phenylglutaconic acids, A., i, 553.
- Feist, Karl**, and **A. Futtermenger**, the optical activity of catechins, A., i, 565.
- Feit, Hellmuth**. See **Kurt Lindner**.
- Feld, G. W.** See **Ernst Hermann Riessenfeld**.
- Feldmeier, Gregor**. See **Karl Fleischer**.
- Feldsberg, Emil**. See **Sigmund Fränkel**.
- Felix, K.**, a basic protein derivative, A., i, 295.
histopeptone, A., i, 479.
the digestion of histone sulphate with pepsin and hydrochloric acid, A., i, 693.
a basic peptone-like substance in the thymus, A., i, 702.
- Fellenberg, Theodore von**, estimation of the various acids of wine, A., ii, 534.
- Fenger, Frederic**, chemical composition and physiological characters of brain cephalin, A., i, 315.
- Ferber, Erwin**. See **Kurt Brass**.
- Ferguson, Alfred L.**, and **Wesley G. France**, transport numbers of sulphuric acid by the concentration cell method, A., ii, 113.
influence of gelatin on the transport numbers of sulphuric acid, A., ii, 114.
- Ferguson, John**. See **Robert Martin Caven**.
- Ferguson, John Bright**, melting interval of certain undercooled liquids: the use of liquid air as a refrigerant, A., ii, 614.
melting and freezing point of sodium chloride, A., ii, 848.

- Ferguson, John Bright.** See also *G. A. Williams*.
- Ferjančić, S.** See *Maximilian Samec*.
- Fernández, Obdulio,** and *T. Garmendia*, Endo's reaction, biology of *Bacillus coli*, A., i, 405.
- Fernández, Obdulio,** and *A. Pizarroso*, the catalytic power of flour, A., ii, 94.
- Fernau, Albert,** and *Wolfgang Pauli*, action of penetrating radium radiation on inorganic and bio-colloids. III., A., ii, 202.
- Fernbach, Auguste,** and *Marcel Schoen*, the rôle of acetaldehyde in alcoholic fermentation, A., i, 203.
- Ferrara, Manfredi,** pharmacological action of magnesium sulphate and its application in strychnine poisoning, A., i, 500.
- Ferrer, Juan.** See *Heinrich Wieland*.
- Ferrer Tomás, Juan.** See *Antonio García Bandu*.
- Ferri, Lelio.** See *Umberto Sborgi*.
- Fester, G.,** and *G. Brude*, the properties of potassium permanganate, A., ii, 771.
- Fetkenheuer, B.,** detection of fluorine, A., ii, 655.
- Faulgen, R.** See *Wilhelm Stepp*.
- Fichter, Franz,** biochemical and electrochemical oxidation of organic compounds, A., ii, 23.
- Fichter, Franz,** and *Gustav Grisard*, the electrochemical oxidation of aromatic nitriles, A., i, 37.
- Fichter, Franz,** and *Wolfgang Jaeck*, the electrochemical oxidation of azobenzene, A., i, 62.
- Fichter, Franz,** and *Ernst Jenny*, the heat of formation of aluminium nitride, A., ii, 615.
- Fichter, Franz,** and *Hans Löwe*, the electrochemical oxidation of *o*-toluenesulphonamide, A., i, 242.
- Fichter, Franz,** and *Hugues de Montmollin*, electrochemical studies in the pyrazole group, A., i, 470.
- Fichter, Franz,** and *Emil Rothenberger*, electrochemical oxidation of dimethylaniline, A., i, 447.
- Fichter, Franz,** and *Richard Suter*, magnesium cyanide, A., i, 532.
- cathodic reduction of elementary nitrogen, A., ii, 372.
- Fielding, William R.,** behaviour of substances near the absolute zero, A., ii, 23.
- polymerisation at the critical temperature, A., ii, 421.
- Fierz, Hans Eduard,** constitution of carbenium dyes, A., i, 445.
- Fierz, Hans Eduard,** and *Richard Sellmann*, attempts to prepare perinaphthindigotin, and the behaviour of azo-dyes from naphthylglycines, A., i, 870.
- Fierz, Hans Eduard,** and *Richard Tobler*, 2:2'- $\beta\beta$ -naphthindigotin, A., i, 869.
- Fieser, L. F.** See *James Bryant Conant*.
- Fillon, Robert.** See *Gustav Hinard*.
- Finch, George Ingle,** and *Raymond Harold Kelsall Peto*, purification of phosphoric oxide, T., 692.
- Findlay, Alexander,** and *Owen Rhys Howell*, the rate of evolution of carbon dioxide from solution in presence of colloids, T., 1046.
- Findlay, Alexander,** and *Cyril Rosebourne*, decomposition and stabilisation of ammonium nitrate in presence of oxidisable material, A., ii, 291.
- Findlay, Alexander,** and *Vernon Harcourt Williams*, electrolytic reduction of dextrose, A., i, 523.
- Finger, H.,** and *Lina Elrich*, new synthesis of hydroxylated benzoylformic [phenylglyoxylic] acids, A., i, 254.
- Finks, A. J.** See *David Breese Jones*.
- Fioletova, Alexandra.** See *Nicolai Antonovich Pushin*.
- Fiori, Q.** See *G. de Sanctis*.
- Fioroni, W.** See *P. Karrer*.
- Firth, James Brierley.** See *John Driver*.
- Fischer, A.,** and *W. Classen*, the volumetric estimation of dithionates, A., ii, 453.
- Fischer, Artur.** See *Helmut Scheibler*.
- Fischer, E.** See *Fritz Mayer*.
- Fischer, Franz,** [lignite producer tar], A., i, 330.
- Fischer, Franz,** and *Georg Pfeleiderer*, the solubility of oxygen in various organic solvents, A., ii, 841.
- Fischer, Georg.** See *Rudolph Friedrich Weinland*.
- Fischer, Hans,** and *Hermann K. Barrenscheen*, the azo-dyes of bilirubin. I., A., i, 278.
- Fischer, Hans,** and *Marianne Hermann*, pyrroles and hydroxypyrroles, A., i, 1054.
- Fischer, Hans,** and *Maria Kean*, the iron salts of dipyrrolylphenylmethane dyes and triphenylpyrrolymethane. II. Diphenylpyrrolylcarbinol and its derivatives, A., i, 869.
- Fischer, Hans,** and *Viktoria Luckmann*, the iron salts of dipyrrolylphenylmethane dyes; triphenylpyrrolymethane. I., A., i, 276.

- Fischer, Hans, Karl Schneller, and Werner Zerweck**, [with *Max Schubert*], pyrroles. III. Ketones, ketonic acid esters, and ketonic acid nitriles of substituted pyrroles, A., i, 1055.
- Fischer, Hans, and Werner Zerweck**, pyrroles. I. Pyrrolealdehydes, A., i, 758.
- pyrroles. II. Nitration of substituted pyrroles, A., i, 758.
- Fischer, Heinrich**. See *Erich Schmidt*.
- Fischer, Martin Henry, and George D. McLaughlin**, theory of the Liesegang rings, A., ii, 206.
- Fischer, Martin Henry, George D. McLaughlin, and Marian O. Hooker**, soaps and proteins. I. Colloidal chemistry of soaps and the manufacture of soap, A., ii, 430.
- soaps and proteins. II. Colloid chemistry of soap; manufacture, analogies in the colloid chemistry of soaps, albumins, and tissues.
- III. Colloid chemical behaviour of fatty acids and their derivatives and the analogous behaviour of neutral proteins and their derivatives, A., ii, 829.
- Fischer, Otto**, [with *H. Kracker, Wolfgang Meier, and H. Schwappacher*], the constitution of the products arising from the condensation of *o*-aminoazo-compounds with aldehydes. I., A., i, 956.
- Fischer, Otto, and Martin Bollmann**, formation and properties of fluorescein, A., i, 936.
- Fischer, Richard**, pharmacology of carbon oxysulphide, A., i, 402.
- Fischer, Roger**, colloidal equilibrium of blood serum, A., i, 1209.
- Fischer, Wilhelm**. See *Wilhelm Biltz*.
- Fischler, M.**, device for the removal of fumes, A., ii, 388.
- Fisher, E. A.**, relation of the hydrogen-ion concentration of the soil to plant distribution, A., i, 510.
- Fiske, Cyrus H.**, the "alkaline tide" after meals. I., A., i, 88.
- inorganic phosphate and acid excretion in the post-absorptive period, A., i, 88.
- method for the estimation of total base in urine, A., ii, 408.
- Fitger, Peter**, stereoisomeric ethylthio-succinic acids, A., i, 107.
- certain products of the oxidation of inactive ethylthio-succinic acid, A., i, 108.
- Fitz, Reginald**, acetone substances in the blood in diabetes, A., i, 192.
- Flater, E.** See *Paul Pfeiffer*.
- Flecker, Oriel Joyce, and (Miss) Millicent Taylor**, studies of the constitution of soap solutions; sodium behenate and sodium monoate, T., 1101.
- Fleisch, Alfred**, an equilibrated sterilisable fluid of physiological hydrogen-ion concentration, A., i, 964.
- blood in impaired cell respiration; cause of avian beri-beri, A., i, 1215.
- Fleischer, Karl, and Gregor Feldmeier**, *ar*-aldehydes of tetrahydronaphthalene, A., i, 1160.
- Fleischer, Karl, and Ewald Retze**, synthesis of pyrene, A., i, 1138.
- Fleischer, Karl, and Karl Schranz**, 5-aminoacenaphthene, A., i, 1142.
- Fleischhauer, C.** See *Georg Schroeter*.
- Fleischmann, Walter**. See *Otto von Fürth*.
- Fleming, William D.**, vitamin content of rice by the yeast method; organic nitrogen as a possible factor in stimulation of yeast, A., i, 93.
- Fleury, Paul**, colorimetry, A., ii, 518.
- Fleury, Paul, and Louis Bontot**, the modified Lehmann method for the estimation of dextrose; adaptation to small quantities of reducing sugars, A., ii, 879.
- Fleury, Paul, and Gabriel Poirot**, the orcinol reaction with furfuraldehyde; application to the colorimetric estimation of small quantities of furfuraldehyde, A., ii, 595.
- estimation of small quantities of furfuraldehyde colorimetrically, A., ii, 666.
- Floros, Aristomenis**. See *Ludwig Pin-cussen*.
- Flütsch, Chr.** See *Karl W. Rosen-mund*.
- Fodor, Andor**, colloidal chemistry of the proteins. III., A., i, 691.
- Fodor, Andor, and B. Schönfeld**, dependence of the adsorption by charcoal on the quantity of charcoal and the nature of the adsorption isothermal, A., ii, 688.
- Fodor, Andor**. See also *Emil Abder-halden*.
- Föhrenbach, Erich**. See *Emil Fromm*.
- Földi, Zoltan**, alkylation of amines with sulphonic esters, A., i, 732.
- Foerster, Fritz**, nitrous anhydride, A., ii, 284.
- Försterling, Karl**, the emission of light by atoms, A., ii, 729.
- Fogg, H. C., and Charles James**, atomic weight of yttrium, A., ii, 297.
- Fogler, M. F.** See *Roger Adams*.

- Foix, A.**, the determination of molecular weight by means of osmotic pressure, A., ii, 621.
 the action of diamond on carbon monoxide, A., ii, 641.
 the composition and calorific power of coal from the deeper strata of the Kenadza (Algerian) mine, A., ii, 713.
- Foix, A.** See also *Joseph Auguste Muller*.
- Fokker, A. D.**, stationary electron vibrations without radiation resistance, A., ii, 106.
 electronic theory of the interior of the atom, A., ii, 137.
- Folin, Otto**, colorimetric estimation of the amino-acid nitrogen in normal urine, A., ii, 536.
 the necessity of checking the quality of sodium tungstate used in the system of blood analysis, A., ii, 596.
- Folin, Otto** [with *Hsien Wu*], a system of blood analysis. Suppl. III. A new colorimetric method for the estimation of the amino-acid nitrogen in blood, A., ii, 540.
- Folin, Otto**, and *Hilding Berglund*, the transportation, retention, and excretion of carbohydrates, A., i, 487.
 the retention and distribution of amino-acids with especial reference to the urea formation, A., i, 702.
 a colorimetric method for the estimation of sugars in normal human urine, A., ii, 400.
- Folin, Otto**, and *Joseph M. Looney*, colorimetric methods for the separate estimation of tyrosine, tryptophan, and cystine in proteins, A., ii, 539.
- Folkoff, Caspar**. See *A. J. Schaffer*.
- Fontell, Nils**. See *Ossian Aschan*.
- Fontés, G.**, and *L. Thivolle*, micro-estimation of lactose by means of potassium permanganate; application to milk, A., ii, 323.
- Foot, Paul D.**, *W. F. Meggers*, and *Fred L. Mohler*, excitation of the enhanced spectrum of magnesium in a low voltage arc, A., ii, 4.
 the excitation of the enhanced spectra of sodium and potassium in a low voltage arc, A., ii, 598.
- Foot, Paul D.**, and *Fred L. Mohler*, the significance of the half terms in spectral series formulae, A., ii, 410.
- Forbes, George Shannon**, and *Harriet Isabelle Cole*, solubility of silver chloride in dilute chloride solutions and the existence of complex argentichloride ions. II., A., ii, 291.
- Forbes, George Shannon**, *Howard Wilmot Estill*, and *Osman James Walker*, induction periods in reactions between thiosulphate and arsenite or arsenate; a useful clock reaction, A., ii, 271.
- Forbes, H. S.**, and *Louise Hompe*, the effect of carbon monoxide, illuminating gas, and benzene on blood coagulation time, A., i, 483.
- Ford, G. Winifred**. See *John L. Haughton*.
- Foresti, B.**, preparation and mechanism of formation of phenanthroxazine, A., i, 1062.
 so-called di-9:10-monohydroxyphenanthrylamine suggested as a reagent for nitric acid and phenanthroxazine, A., ii, 524.
- Forrer, Max**, the physico-chemical study of the lead chamber process, A., ii, 761.
- Forsman, J.**, effect of ether on anti-substances, A., i, 964.
- Forster, Aquila**, and *William Coulson*, nitro-derivatives of *m*-nitrodimethylaniline, T., 1988.
- Forster, Martin Onslow**, and *William Bristow Saville*, constitution of picrorocellin, a diketopiperazine derivative from *Roccella fuciformis*, T., 816.
 the triazo-group. XXII. Cinnamic acid chlorohydrin and its conversion into α -triazole- β -hydroxy- β -phenylpropionic acid, T., 2595.
- Forsyth, J. A. C.** See *P. J. Cambridge*.
- Fosse, Robert**, the synthesis of a nitrogenous principle of plants, hydrocyanic acid, by the oxidation of ammonia and carbohydrates, glycerol, or formaldehyde, A., i, 117.
- Fosse, Robert**, and *A. Hiculle*, synthesis of hydrocyanic acid by oxidation, in ammoniacal silver solution, of alcohols, phenols, and amines, A., i, 117.
 aptitude of formaldehyde to form hydrocyanic acid by oxidation in ammoniacal silver solutions, A., i, 523.
- Foster, A. L.** See *De Witt Neighbors*.
- Foster, Dorothy Lilian**, and *Dorothy Mary Moyle*, the interconversion of carbohydrate and lactic acid in muscle, A., i, 398.
- Foster, H. E.** See *J. B. Ayer*.
- Foulk, C. W.**, and *Samuel Morris*, comparative values of different specimens of iodine for use in chemical measurements, A., ii, 310.

- Fourneau, Ernest**, and **José Puyal**, the amino-alcohols; homologues of novocaine, A., i, 639.
- Fournier, Louis**, and **L. Guénot**, treatment of syphilis by bismuth, A., i, 301.
- Fournier, Louis**, **L. Guénot**, and **A. Schwartz**, first results of the treatment of syphilis by sodium *p*-hydroxy-m-aminophenylarsinate or "189," A., i, 300.
- Fowler, R. H.** See *Francis William Aston*.
- Fox, Francis William.** See *John Addyman Gardner*.
- Foxwell, G. E.**, thermal dissociation of ammonia with special reference to coke oven conditions, A., ii, 431.
- Fränkel, Sigmund**, and **Emil Feldsberg**, a new function of the tryptic ferment (anhydrase) and the preparation of *d*-tyrosine anhydride and *d*-tryptophan anhydride from the products of tryptic digestion, A., i, 184.
- Fränkel, Sigmund**, and **Oskar Gilbert**, lipoids. XVIII. The preparation of phosphosulphatides from brain, A., i, 294.
- Fränkel, Sigmund**, and **Josef Hager**, vitamins. II. Acceleration of fermentation by extracts of animal organs, A., i, 409.
- Fränkel, Sigmund**, and **Paul Jellinek**, products of prolonged tryptic digestion of casein, A., i, 965.
- Fränkel, Sigmund**, and **Artur Kász**, lipoids. XIX. A lecithin from human brain, A., i, 294.
- Fränkel, Sigmund**, and **Albert Scharf**, vitamins. III. Acceleration of fermentation by extracts of plants and the action of choline and aminoethyl alcohol on fermentation, A., i, 409.
- vitamins. IV. The adsorption of vitamins, A., i, 409.
- vitamins. V. Further experiments on the chemistry of vitamins, A., i, 409.
- Fraenkel, Walter**, the recrystallisation of pure mechanically unworked gold, obtained by solidification from the fused mass, A., ii, 778.
- Fraenkel, Waller.** See also *Richard Lorenz*.
- France, Wesley G.** See *Alfred L. Ferguson*.
- Francis, Francis Ernest** [with *John Olifford Pope*, and *Reginald Henry Coyah*], the composition of paraffin wax. I., T., 496.
- Francis, Francis Ernest**, *Cyril Mercer Watkins*, and *Reginald Wilfred Wallington*, the composition of paraffin wax. II. and III., T., 1529, 2294.
- Franck, J.**, the broadening of spectral lines, A., ii, 241.
- the theory of Klein and Rosseland applied to fluorescence, photochemical processes, and the electron emission from hot substances, A., ii, 464.
- resonance and ionisation potentials of helium, A., ii, 811.
- Franck, J.**, and *Walter Grotrian*, the absorption of mercury vapour, A., ii, 728.
- Franck, J.** See also *G. Cario*.
- François, Maurice**, estimation of mercury in the mercurial pills of the [French] codex, A., ii, 87.
- François, Maurice**, and *Louis Gaston Blanc*, preparation of the alkaloidal mercuri-iodides in crystalline form, A., i, 851.
- preparation of the alkaloidal bismuthic iodides in a crystalline form, A., i, 851.
- François, Maurice**, and *Ch. Lormand*, the photography of opaque crystals, A., ii, 127.
- stereoscopic photography of crystals, A., ii, 127.
- François, Maurice**, and *Émile Luce*, analysis of camphor preparations, A., ii, 667.
- Franke, Adolf**, and *Gerald Groeger*, action of α -bromoisobutaldehyde on sodiomalonic ester, A., i, 808.
- Franke, Elizabeth.** See *Stanley Rossiter Benedict*.
- Frankenthal, Kate.** See *Martin Jacoby*.
- Franklin, Edward Curtis**, ammonio-carbonic acids, A., i, 440.
- Franzen, Hartwig**, a shaking machine for large quantities of fluid, A., ii, 759.
- Franzen, Hartwig**, and *Fritz Helwert*, the chemical constituents of green plants. XX. The acids of the cherry (*Prunus avium*), A., i, 1102.
- Franzen, Hartwig**, and *Ernst Keyssner*, the chemical constituents of plants. XVII. The presence of lactic acid in the leaves of the blackberry (*Rubus fruticosus*), A., i, 310.
- Franzen, Hartwig**, and *Rudolf Ostertag*, the chemical constituents of green plants. XVIII. The acids from the berry of the mountain ash (*Pyrus aucuparia*) which are precipitated by lead acetate, A., i, 616.

- Fransen, Hartwig, and Rudolf Ostertag**, chemical constituents of green plants. XXI. The non-existence of Crassulaceae malic acid, A., i, 1223.
- Fransen, Hartwig, and Eugen Schumacher**, the chemical constituents of green plants. XIV. The acids in the currant (*Ribes rubrum*) which are precipitated by lead acetate, A., i, 310.
- Fransen, Hartwig, and Gustav Stäuble**, influence of substituents on reactions. VII. The preferential points of substitution in the naphthols and in α -naphthylamine, A., i, 450.
- Fransen, Hartwig, and Emmi Stern**, the chemical constituents of green plants. XV. The presence of lactic acid in the leaves of the raspberry (*Rubus idaeus*), A., i, 311.
- chemical constituents of green plants. XIX. Occurrence of lactic acid and succinic acid in the leaves of the raspberry (*Rubus idaeus*), A., i, 975.
- Fraps, George Stronach**, relation of the phosphoric acid of the soil to pot experiments, A., i, 616.
- Fraser, Ronald**, the constitution of benzene, T., 188.
- Frassinetti, Mario**. See *Luigi Rolla*.
- Frazer, Joseph Christie Whitney**. See *Paul Lotz*.
- Fred, Edwin Broun, W. H. Peterson, and J. J. Anderson**, characteristics of certain pentose-destroying bacteria, especially as concerns their action on arabinose and xylose, A., i, 201.
- Fred, Edwin Broun**. See also *O. R. Brunkow, and W. H. Peterson*.
- Fredenhagen, Karl**, theory of the electrolytic ionic condition and the calculation of the electrolytic solution constants, and of the related quantities from the chemical relationships, A., ii, 112.
- Freeman, Benjamin**. See *Howard Waters Doughty*.
- Freeth, F. A.**, system $\text{Na}_2\text{O}-\text{CO}_2-\text{NaCl}-\text{H}_2\text{O}$, considered as two four-component systems, A., ii, 627.
- Freise, R.** See *Hermann Stendel*.
- Fréjacques, M.** See *Camille Matignon*.
- Fresenius, Ludwig**. See *Otto Lemmermann*.
- Fresenius, Wilhelm, and Leo Grunhüt**, estimation of tannin and colouring matters in wines, A., ii, 96.
- detection of formic acid in wine, A., ii, 234.
- Freudenberg, Ernst, and Paul György**, fixation of lime by animal tissues. III.—VIII., A., i, 87, 291, 299, 608.
- Freudenberg, Karl**, catechin, A., i, 756.
- Freudenberg, Karl, Otto Böhme, and Ludwig Furrmann**, tannins and similar compounds. IX. Stereoisomeric catechins. II., A., i, 756.
- Freudenberg, Karl, and Fritz Brauns**, the configuration of the simple α -hydroxy-acids, A., i, 628.
- acetone sugars. I. Transformations of diacetonoglucose, A., i, 1117.
- Freudenberg, Karl, and Otto Ivers**, syntheses of mixed acylated halogen sugars, A., i, 523.
- Freudenberg, Karl, and Ludwig Orthner**, reduction of flavanone, A., i, 757.
- Freudenberg, Karl, and Wilhelm Seilasi**, tannins and similar substances. II. Chinese tannin, A., i, 1169.
- Freudenberg, Karl, and Olof Svanberg**, acetone sugars. II. Diacetonexylose, A., i, 1116.
- Freudenberg, Karl, and Erich Vollbrecht**, tannase, A., i, 285.
- tannins and similar compounds. X. The tannin of the native oak, A., i, 1046.
- Freuler, A.** See *W. D. Treadwell*.
- Freund, M.** See *Paul Jacobsen*.
- Freund, Waller**. See *Fritz Mayer*.
- Freundler**. See *Gabriel Bertrand*.
- Freundler, P., (Mlle) Y. Menager, and (Mlle) Y. Laurent**, iodine in the *Laminaria*, A., i, 98.
- the composition of the *Laminaria*, A., i, 98.
- Freundlich, Herbert**, concentration and potential fall at boundary surfaces, A., ii, 19.
- effective potential difference of electro-osmosis and allied phenomena, A., ii, 189.
- Freundlich, Herbert, and R. Bartels**, kinetics of reactions in which halogen atoms pass from the inner sphere of activity to the outer, A., ii, 489.
- Freundlich, Herbert, and A. Gyemant**, thermodynamic and electrokinetic potential difference at the surface of two liquids, A., ii, 342.
- Freundlich, Herbert, and E. Loening**, the protective and flocculating action of hydrophilic colloids and hydrophobic sols, A., ii, 356.
- behaviour of Carey Lea's silver sol towards electrolytes and hydrophilic colloids, A., ii, 696.
- Freundlich, Herbert, and P. Scholz**, flocculation by mixtures of electrolytes, A., ii, 828.
- hydrophobic and -hydrophilic sols of sulphur, A., ii, 841.
- Freundlich, Leo**. See *Siegfried Skraup*.

- Fricke, Hugo.** See *William Duane*.
- Fricke, Robert,** the analytical detection and differentiation of acetaldehyde, aldol, glyoxylic acid, and their presence in the urine of diabetic patients, A., i, 300.
- the presence of pyruvic acid in normal and diabetic urines, A., i, 495.
- use of the "silver method" in the estimation of acetaldehyde; its application in the estimation of other aldehydes; a convenient method of accumulation of aldehyde and other volatile substances from body fluids, A., ii, 326.
- the detection of aldol in the urine of diabetic patients, A., ii, 326.
- energy principle of the coagulation of colloids, particularly of the thermokinetic mechanism, A., ii, 484.
- glass suspensions produced by rubbing glass walls with glass rods, A., ii, 692, 744.
- Fricke, Robert.** See also *Wilhelm Stepp*.
- Fricke, Kurt.** See *Hermann Leuchs*.
- Frieber, Walther,** formation of indole, and indole reactions; behaviour of indole-negative bacteria, A., i, 901.
- Friedberg, Eduard,** quantitative measurement of the transient excretion of caffeine in man by a new biological method, A., i, 88.
- Friedebach, M.** See *H. P. Kaufmann*.
- Friedel, Georges, and L. Royer,** mixtures of anisotropic liquids and the identity of Grandjean's stratified liquids with liquids of the azoxyphenetole type, A., ii, 129.
- Friedemann, W. G.,** nitrogen distribution of proteins extracted by 0.2 per cent. sodium hydroxide solution from cotton-seed meal, the soja bean, and the coconut, A., i, 505.
- Friedländer, Paul,** the dyes from *Purpura aperta* and *Purpura lapillus*, A., i, 793.
- Friedländer, Paul, Walter Herzog, and G. von Voss,** indigoid dyes of the phenanthrene and indene series, A., i, 784.
- Friedländer, Paul, S. Karamessinis, and O. Schenk,** certain chloronaphthalene derivatives, A., i, 244.
- Friedländer, Paul, and K. Kunz,** 1:1'-diphenylindigotin, A., i, 765.
- Friedrich, Alfred.** See *Alois Zinke*.
- Friedrich, Walter.** See *Julius Meyer, and Robert Schwars*.
- Friend, Herman,** clinical method for the estimation of chlorides in blood, A., ii, 389.
- Friend, John Albert Newton,** lecture experiment; protective colloids, A., ii, 267.
- Friend, John Albert Newton, and John Horace Dennett,** the rate of solution of iron in dilute sulphuric acid both when stationary and under rotation, T., 41.
- Friend, John Albert Newton, and Reece Henry Vallance,** the influence of protective colloids on the corrosion of metals and on the velocity of chemical and physical change, T., 466.
- Fries, Karl,** 1:2-naphthaquinols, A., i, 42.
- Fries, Karl, and H. Hasenjäger,** the course of the reaction in the synthesis of oxindigo (2:2'-diketo- $\Delta^{1'}$ -dicoumaran), A., i, 45.
- Fries, Karl, and F. Kerkow,** linear benzonaphtha-*p*-thiazine [$\beta\beta$ -naphthaphenthiazine], A., i, 577.
- Fries, Karl, and H. Leue,** autoxidation of 2-acetyl- α -naphthol in alkaline solution, A., i, 462.
- Fries, Karl, and W. Lohmann,** the 2-methylnaphthalene series, A., i, 29.
- Frister, Fritz.** See *Otto Dimroth*.
- Fritsch, Albert.** See *Albin Kurt-nacker*.
- Fritschi, J.** See *Hermann Staudinger*.
- Fritzmann, E.,** preparation of hydrocyanic acid in large quantities in the laboratory, A., i, 1128.
- Fritzmann, E., and K. Macjulevitch,** new method for estimating volatile substances in air, A., ii, 877.
- Froboese, Kathe.** See *Victor Froboese*.
- Froboese, Victor, and Kathe Froboese,** estimation of aluminium in tungsten, A., ii, 397.
- Frölich, Per.** See *Sven Bodfors*.
- Froelicher, Victor, and Julius Berend Cohen,** the nitro- and amino-derivatives of *o*- and *p*-methoxybenzoic acids and of α - and β -methoxynaphthoic acids, T., 1652.
- Froeschl, Norbert.** See *Ernst Philipp*.
- Frog, F., and S. Schmidt-Nielsen,** fatty acids of butter, A., i, 494.
- Froidevaux, J.,** the estimation of ammoniacal nitrogen in nitrogenous organic substances, and particularly in proteins and their products of decomposition, A., ii, 454.
- Fromherz, Konrad,** the action of different groups of local anæsthetics, A., i, 705.
- Fromholdt and Nersessov,** pigment metabolism. I. and II., A., 387, 404.

- Fromm, Emil**, ring closure with hydrazinedicarboxamides containing sulphur; dithiourazole and iminothiourazole, A., i, 62.
- Fromm, Emil** [with *Hermann Wenzl*], syntheses from cyanamide; preparation of thiocarbamine cyanides, carbamine cyanides, and biurets, A., i, 437.
- Fromm, Emil**, and **Ernst Honold**, syntheses with cyanamide; cyanamido ethyl alcohol and guanidoethyl alcohol, A., i, 529.
- Fromm, Emil**, **Erich Kayser**, **Karl Briegleb**, and **Erich Föhrenbach**, disulphides with neighbouring single and multiple linkings; syntheses of triazoles and thiodiazoles, A., i, 377.
- Fromm, Emil**, and **Erich Siebert**, derivatives of dithioethylene and dithioacetylene and the additive capacity of multiple bonds, A., i, 542.
- Fuchs, August**, melting point tubes as reaction vessels, A., ii, 759.
- Fuchs, Berthold**. See **Siegfried Edlbacher**.
- Fuchs, Edgar**. See **Percy Brigl**.
- Fuchs, Friedrich**, anhydrides of *N*-carboxylic acid, A., i, 1152.
- Fuchs, Francis J.** See **James Kendall**.
- Fuchs, Karl**. See **Ernst Späth**.
- Fuchs, Walter**, [tautomerism of resorcinol], A., i, 336.
- Fuchs, Walker**, and **Heinrich Metzl**, the Wurtz-Fittig synthesis, A., i, 442.
- Fuchs, Walter**, and **Walter Stix**, tautomerism of phenols. V. 1:5- and 2:7-dihydroxynaphthalenes, A., i, 451.
- Füchtbauer, Chr.**, and **C. Joos**, intensity and broadening of spectrum lines, A., ii, 242.
- Fühner, Hermann**, the effective strengths of narcotics. I. Experiments on the isolated frog's heart, A., i, 199. the toxicology of arsine. II., A., i, 499.
- Fürth, Adolf**, strength of hypiodous acid, A., ii, 281.
- Fürth, L.** See **F. Erdstein**.
- Fürth, Otto von**, pressure of carbon dioxide or swelling of protein as the cause of muscle contraction, A., i, 398.
- Fürth, Otto von**, and **Walter Fleischmann**, the tyrosine content of proteins, A., i, 406.
- Fürth, Otto von**, and **Fritz Lieben**, colorimetric experiments on tryptophan. VI. The tryptophan content of some foods and the tryptophan requirement of men, A., i, 293.
- Fürth, Otto von**, and **Fritz Lieben**, decomposition of lactic acid by yeast and by blood-cells, A., i, 502. further experiments on the destruction of lactic acid by yeast, A., i, 1219.
- Fujibayashi, Tôtarô**, preparation of metals by Goldschmidt's aluminothermic method. I., A., ii, 647.
- Fujita, Atsushi**. See **Yasuhiko Asahina**.
- Fukuda, Mitsuharu**, absorption of light by sulphur at various temperatures, A., ii, 728.
- Fukuda, Mitsuharu**. See also **Masamichi Kimura**.
- Fukushima, Kanshi**, glycolytic action of leucocytes. I., A., i, 697.
- Fulda, H. L.** See **Paul Jacobsen**.
- Fuldner**. See **Otto Diels**.
- Fuller, Everett W.** See **Robert E. Wilson**.
- Funck, Albert**, the theory of blood-clotting, A., i, 290.
- Funck, Albert**. See also **B. Stuber**.
- Funk, Casimir**, and **Harry E. Dubin**, vitamin requirements of certain yeasts and bacteria, A., i, 203.
- Funk, Eberhard**, the influence of cobalt-ammines on the action of catalase and amylase, A., i, 481.
- Funk, Herbert**. See **Wilhelm Manchot**.
- Funke, G. L.**, influence of hydrogen-ion concentration on the action of the amylase of *Aspergillus niger*, A., i, 796.
- Futtermenger, A.** See **Karl Feist**.
- Fyfe, Alexander Walker**, and **British Dyestuffs Corporation, Ltd.**, preparation of 1-chloro-2-aminoanthraquinone, A., i, 260.

G.

- Gadals, L.**, estimation of sulphur in pyrites, A., ii, 79.
- Gadamer, Johannes**, and **F. von Bruchhausen**, the constitution of corydaline, A., i, 675.
- Gadamer, Johannes**, and **Carl John**, ecgonine, A., i, 167, 675.
- Gad-Andresen, Knud L.**, a micro-method for the estimation of ammonia in blood and in organic fluids, A., ii, 523. a micro-urease method for the estimation of urea in blood, secretions, and tissues, A., ii, 536.
- Galley, Zolia Jencks**. See **Hermann V. Tartar**.
- Gainey, P. L.**, and **H. W. Batchelor**, influence of hydrogen-ion [concentration] on the growth of *Azotobacter*, A., i, 1096.
- Galanos, S.** See **K. Pfizenmaier**.

- Galbraith, William Lyle.** See **William Lewcock.**
- Gale, William A.** See **Ebenezer Henry Archibald.**
- Galissi, A.** See **Giacomo Luigi Clamician.**
- Gallagher, Patrick Hugh,** phototropy and photoelectric effect, A., ii, 11.
- Gallagher, Patrick Hugh.** See also **Charles Moureu.**
- Gamble, C. J., and Margaret C. Herriek,** the utilisation of dextrose by the tubercle bacillus, A., i, 902.
- Gamble, James L.,** carbonic acid and bicarbonate in urine, A., i, 494.
- Gammal, C. A.** See **M. C. Taylor.**
- Gama, Alphonse, and O. Kaiser,** benzyl ester of 2-phenylquinoline-4-carboxylic acid, A., i, 51.
- Ganassini, Domenico,** [detection of] quinotoxine in quinine salts, A., ii, 539.
rapid detection of bismuth in urine and in saliva, A., ii, 590.
- Ganguly, Kanai Lal,** attempted synthesis of norpinic acid, A., i, 1153.
- Ganguly, Phani Bhusan, and B. C. Banerji,** effect of surface tension, A., ii, 821.
- Ganguly, Phani Bhusan, and Nilratan Dhar,** coagulation of colloids by sunlight, A., ii, 604.
coagulation of manganese dioxide sol by different electrolytes, A., ii, 829.
- García Bandás, Antonio, and Juan Ferrer Tomás,** derivatives of aminodiphenyl, A., i, 333.
- García Bandás, Antonio, and J. Pascual Vila,** organomagnesium derivatives, A., i, 734.
- Gardner, John Addyman, and Francis William Fox,** origin and destiny of cholesterol in the animal organism. XII. The excretion of sterols in man, A., i, 89.
origin and destiny of cholesterol in the animal organism. XIII. The autolysis of liver and spleen, A., i, 790.
- Gardner, John Addyman, George King, and Edwin Booth Powers,** the respiratory exchange in fresh-water fish. III. Gold fish, A., i, 1084.
- Garmendia, T.** See **Obdulio Fernández.**
- Garner, Frederic Horace.** See **W. F. Faragher.**
- Garner, V. C.** See **E. Weiss.**
- Garner, William Edward,** the Dushman equation for the velocity of a unimolecular reaction, A., ii, 358.
polar and non-polar valency in organic compounds, A., ii, 758.
- Garner, William Edward, and Douglas Norman Jackman,** note on the effect of a magnetic field on catalysis by ions in the presence of a paramagnetic salt, T., 1298.
- Garner, William Edward, and Kichimatsu Matsuno,** the explosion of acetylene and nitrogen. II., T., 1729.
- Garner, William Edward, and C. A. Waters,** apparatus for electrometric titration, A., ii, 862.
- Garrault, (Mlle) H.** See **E. Fauré-Fremiet.**
- Garreau, (Mlle) Y.,** colorimetry, A., ii, 518.
- Gastaldi, Carlo,** condensation of acetophenone. III., A., i, 367.
4:6-diphenyl-2-methylpyridine, A., i, 578.
2:4-diphenylpyridine, A., i, 679.
preparation of methylenecitric acid, A., i, 809.
- Gastaldi, Carlo, and G. Brauniser,** bisulphite compounds of oximino-ketones and glyoximes, A., i, 626.
- Gattefossé, Jean,** essential oils from Tonkin and Annam, A., i, 1167.
- Gaubert, Paul,** recrystallisation produced by annealing, A., ii, 35.
liquid crystals of calcium phosphate, A., ii, 445.
- Gault, Henri, and R. Guillemet,** the chlorination of *n*-butyl alcohol, A., i, 910.
- Gault, Henri, and T. Salomon,** the α -alkyl-levulic acids, A., i, 427.
themethylalkylpyridazinonecarboxylic esters, A., i, 873.
- Gault, Henri, and R. Weisk,** ethyl phenylpyruvate, A., i, 1024.
- Gaunt, R.** See **Carl Hamilton Brown-ing.**
- Gaviati, A., and T. Pavolini,** reduction reactions in the urine of patients treated with arsenobenzenes, A., i, 496.
- Gavron, Joseph L.** See **George W. Raisiss.**
- Gay, L.,** distillation and rectification, A., ii, 120.
- Geelmuyden, H. Chr.,** the rôle of ketonic compounds in intermediate exchanges and in the formation of sugar from fat, A., i, 607.
- Gehlhoff, Georg,** relationship between heat of solution and heat of fusion of organic substances, A., ii, 121.
- Gehroke, Ernst,** attempt [to find] a physical basis for atomic radiation, A., ii, 209.
- Gehroke, Ernst, and E. Lau,** the changeable fine structure of the Balmer series, A., ii, 8.

- Gehrke, Ernst**, and **E. Lau**, the Balmer series of hydrogen, A., ii, 726.
- Gehring, A.**, estimation of humus by oxidation with chromic acid, A., ii, 592.
- Gehrke, Max.** See **Burckhardt Helferich**.
- Geiger, E.**, and **Otto Loewi**, change of choline content of the frog's musculature through electric stimulation, A., i, 492.
- Geiger, Hans**, and **W. Bothe**, scattering of β -rays by thin metal sheets, A., ii, 13.
- Geiger, Hans**, and **A. Werner**, luminous path of α -rays in crystals, A., ii, 188.
- Geith, R.**, estimation of small quantities of sodium in aluminium and alumina, A., ii, 720.
- Gelissen, H. C. J. H.**, Golodetz's reaction (the benzoyl peroxide reaction), A., ii, 460.
- Geller, Hanns.** See **Walther Schrauth**, and **Georg Schroeter**.
- Geloso, Max**, the adsorption of iron by precipitates of manganese dioxide, A., ii, 589.
- Geloso, Max.** See also **Paul Nicolardot**.
- General Electric Co., Ltd.**, London, Research Staff, the disappearance of gas in the electric discharge. IV., A., ii, 417.
- General Electric Co., Ltd.**, London, Research Staff, [by **Colin James Smithells**], reduction of thorium oxide by metallic tungsten, T., 2236.
- Gennari, Marcella.** See **Ernesto Fuxeddu**.
- Gentle, Joseph Alfred Hector Roberts.** See **Nevil Vincent Sidgwick**.
- George, E. F.**, absorption of light by inorganic salts, A., ii, 806.
- Gérald, Pierre.** See **Charles Dufraisse**.
- Gérard, G.**, estimation of small amounts of albumin in urine, A., ii, 670.
- Gerard, R. W.**, intestinal intoxication. I. The presence and significance of histamine in an obstructed bowel, A., i, 790.
- Gericke, W. F.**, differences effected in the protein content of grain by applications of nitrogen made at different growing periods of the plant, A., i, 1226.
- Gerke, Roscoe H.**, temperature coefficient of electromotive force of galvanic cells and the entropy of reactions, A., ii, 682.
- Gerlach, Adolf.** See **William Küster**.
- Gerlach, Walter**, investigation of crystal space lattices by means of Röntgen rays; a simple Röntgen tube, A., ii, 36.
- Gerlach, Walter**, the K_{α} doublet, with new determinations of the lattice constants of some crystals, A., ii, 265.
- Gerlach, Walter**, and **Erich Koch**, a new method of absorption spectroscopy, A., ii, 380.
- Germann, Frank E. E.**, adsorption of radium by barium sulphate, A., ii, 16.
- new hydrate of uranyl nitrate, A., ii, 649.
- Germann, Frank E. E.**, and **Ralph N. Traxler**, adsorption of iodine by silver iodide, A., ii, 371.
- Germis, H. C.** See **Frans Maurits Jaeger**.
- Gersdorff, Charles E. F.** See **Carl Oscar Johns**, and **David Breese Jones**.
- Gersdorff, W. A.** See **M. C. Taylor**.
- Gesellschaft für Teerverwertung m.b. H.**, and **Rudolf Weissgerber**, preparation of thionaphthensulphonic acid, A., i, 1172.
- Gesellschaft für Teerverwertung m.b. H.**, **Rudolf Weissgerber**, and **Otto Kruber**, preparation of thionaphthencarboxylic acids, A., i, 567.
- Gessner, H.** See **Georg Wiegner**.
- Getman, Frederick Hutton**, spectrophotometric study of solutions of cupric chloride, A., ii, 411.
- absorption of light by solutions of cupric lithium chloride, A., ii, 412.
- Gettler, Alexander O.**, and **J. W. Jackson**, preparation of colloidal gold solution for testing spinal fluid, A., ii, 240.
- Ghosh, Jnanendra Chandra**, a general theory of solutions of electrolytes, A., ii, 125.
- Giaja, Jean**, decomposition of amygdalin from the point of view of conjugated fermentation, A., i, 185.
- Gibbs, Harry Drake.** See **Courtney Conover**.
- Gibbs, R. C.** See **William Ridgely Orndorff**.
- Gibson, Charles Stanley**, and **Patrick Playfair Laidlaw**, chloroform in the blood after death, A., i, 1218.
- Gibson, George Ernest**, and **William Albert Noyes, jun.**, luminous discharge in hydrogen and mercury and a new method of measuring ionisation potentials, A., ii, 812.
- Gibson, Robert B.**, and **C. P. Howard**, alcaptonuria and its metabolism, A., i, 401.
- Gibson, Robert B.**, and **Frances T. Martin**, creatine formation in a case of progressive pseudo-hypertrophic muscular dystrophy, A., i, 800.

- Gibson, William Howieson, (Miss) Rebecca Duckham, and (Miss) Ruth Fairbairn**, the products of nitration of toluene, T., 270.
- Gieckhorn, Jos.**, microchemistry of a new group of purple bacteria, A., i, 303.
- Gila Esteban, F.** See **Santiago Piña de Rubies**.
- Gilbert, Lionel Felix, Harold Buckley, and James Irvine Orme Masson**, the system chromium trioxide-sulphur trioxide-water, T., 1934.
- Gilbert, Oskar.** See **Sigmund Fränkel**.
- Gillallan, François A.**, catalysis. II. Dehydration and additive reactions of ethyl alcohol; the formation of acetal and mercaptans, A., i, 709.
- Gillespie, Louis J.**, colour standards for the colorimetric measurement of hydrogen-ion concentration, A., ii, 157.
- Gillespie, Louis J.** See also **Frederick G. Keyes**.
- Gillet, Alfred**, bivalent carbon, A., i, 213.
some compounds of bivalent carbon, A., i, 513.
- Gillis, J.**, chemical passivity of metals, A., ii, 814.
- Gillot, P.**, the changes and movements of the saccharine materials in *Mercuriale vivace* (*Mercurialis perennis*, L.) in the course of its annual growth, A., i, 1101.
- Gilmour, George van Barneveld**, the constitution and rotatory powers of mannitol and fructose complexes formed in solutions containing boric acid and sodium hydroxide, T., 1333.
- Gilmour, Hugh.** See **Gilbert Thomas Morgan**.
- Gilmour, Robert**, vapour pressure of acetaldehyde, A., i, 918.
- Gilta, G.**, the crystalline form of atoxyl [sodium *p*-aminophenylarsinate], A., i, 961.
- Giral Pereira, José**, new insoluble organic nitrates, A., ii, 523.
- Gire, G.**, the dissociation of barium platinichloride, A., ii, 551.
- Glaser, M.** See **Paul Karrer**.
- Gius, Michele**, oxidation of the trinitroxylenes, A., i, 534.
additive compounds of *s*-trinitroanisole with tertiary bases, A., i, 572.
a new red quinoline colouring matter, A., i, 681.
action of certain hydrazines and of pyridine on 5-bromo-1:2:4-trinitrobenzene, A., i, 691.
- Gius, Michele, and A. Angeletti**, action of certain primary bases on 5-bromo-1:2:4-trinitrobenzene, A., i, 649.
- Givaudan, L., & Co.** See **W. C. Sievers**.
- Glässner, Karl**, a new method for the estimation of pepsin, A., ii, 406.
- Glazoleva, Alexandra Alexandrovna.** See **Nicolai Antonovich Pushin**.
- Glasenapp, M. von**, colloidal calcium hydroxide, A., ii, 848.
- Glaser, L. C.**, the spectrum of glucinum and its noteworthy relation to the spectrum of aluminium, A., ii, 675.
- Glasson, J. L.**, beta rays and atomic number, A., ii, 183.
stopping power and atomic number, A., ii, 250.
- Glasstone, Samuel**, physical chemistry of the oxides of lead. III. Hydrated lead monoxide, T., 58.
physical chemistry of the oxides of lead. IV. Red lead and lead sesquioxide, T., 1456.
physical chemistry of the oxides of lead. V. The electromotive behaviour of lead dioxide, T., 1469.
physical chemistry of the oxides of lead. VI. The anodic behaviour of lead and lead dioxide, T., 2091.
- Glattfield, John W. E., and Frank V. Sander**, the C_6 -saccharinic acids. II. The preparation and resolution of *dl*- α -dihydroxybutyric acid; some derivatives of the optically active acids, A., i, 318.
- Glattfield, John W. E., and Edgar Wertheim**, the preparation of optically active hydrazines. II. The preparation of *dl*-*p*-sec.-butylphenylhydrazine; the resolution of *dl*-*p*-sec.-butylaniline, A., i, 385.
- Gleditsch, (Mlle) Ellen, and B. Samdahl**, the atomic weight of the chlorine in an ancient mineral, apatite, from Balme, A., ii, 281.
- Glew, F. Harrison**, radium synthesis of carbon compounds from air, A., ii, 607.
- Glover, C. C.** See **H. W. Valtielah**.
- Glund, Wilhelm**, copper sulphide, A., ii, 446.
the structural formula of copper sulphide, A., ii, 572.
- Godchot, Marcel, and Pierre Bédos**, cyclohexene oxide and *o*-methylcyclohexanol, A., i, 334.
- Godchot, Marcel, and Pierre Brun**, some derivatives of suberone, A., i, 350.
- Goddard, Archibald Edwin**, organo-derivatives of thallium. III. Some thalliumdialkyl salts and the preparation of thalliumdiaryl haloids, T., 36.

- Goddard, Archibald Edwin, Julius Nicholson Ashley, and Richard Bromley Evans**, organo-derivatives of tin and lead. I. Application of lead and tin tetraphenyls in the preparation of organo-metallic compounds, T., 978.
- Goddard, Archibald Edwin, and (Mrs.) Dorothy Goddard**, organo-derivatives of thallium. V. The preparation of thallium diaryl salts, T., 482.
- Goddard, Archibald Edwin, and James Bertram Ward**, metallic derivatives of nitrophenolic compounds. IV. Some complex nitrophenoxides of magnesium, silver, and lead, T., 262.
- Goddard, Archibald Edwin.** See also (Mrs.) **Dorothy Goddard**.
- Goddard, (Mrs.) Dorothy, and Archibald Edwin Goddard**, metallic derivatives of nitrophenolic compounds. III. nitrophenoxides of the alkali metals, T., 54.
organo-derivatives of thallium. IV. Action of thallium chlorides on the Grignard reagent and on organo-derivatives of tin, lead, and bismuth, T., 256.
- Goddard, (Mrs.) Dorothy.** See also **Archibald Edwin Goddard**.
- Goebel, Alfred.** See **Hans Lecher**.
- Goebel, H.**, the catalytic hydrogenation of methysticin, A., i, 657.
- Goebel, Walter F.** See **William Albert Noyes**.
- Göggel, Karl**, the photoelectric effect of the alkaline-earth phosphors, A., ii, 782.
- Göhring, Rudolf**, mechanism of the photochemical chlorine-hydrogen gas reaction and the question of the damping of the velocity [of reaction] of chlorine activated by light, A., ii, 9.
- Gönke, Tatjana.** See **Michael A. Rakusin**.
- Görbitts, Carl.** See **Heinrich Goldschmidt**.
- Goes, Eduard C.** See **Hans Heinrich Schlubach**.
- Götting, H.** See **H. P. Kaufmann**.
- Goiffon, R., and F. Nepveux**, organic acids in urine, A., i, 1215.
- Goldberg, Stephanie**, cataphoresis of colloidal carbon, A., ii, 429.
- Goldberg, Stephanie.** See also **Hilary Lachs**.
- Goldberger, Joseph, and W. F. Tanner**, amino-acid deficiency probably the primary etiological factor in pellagra, A., i, 1092.
- Goldschmidt, Heinrich, Carl Görbitts, Haakon Høngen, and Kristian Fahlø**, affinity dimensions of weak acids and bases in alcoholic solution and the alcoholysis of their salts, A., ii, 135.
- Goldschmidt, Heinrich, Einar Storm, and Odd Hassel**, reduction of nitro-compounds by stannous chloride. III., A., ii, 361.
- Goldschmidt, M.**, lipoids of the crystalline lens, A., i, 489.
- Goldschmidt, Stefan**, univalent oxygen. I., A., i, 1148.
- Goldschmidt, Stefan, and Karl Euler**, amine-oxidation. III. Bivalent nitrogen; diarylacylhydrazyls, A., i, 475.
- Goldschmidt, Stefan, and Konrad Benn**, amine-oxidation. IV. Bivalent nitrogen; α -diphenyl- β -trinitrophenylhydrazyl, A., i, 476.
amine-oxidation. V. Action of tetra-nitromethane on triphenylhydrazine, A., i, 478.
- Goldschmidt, Stefan, and Walter Schmidt**, univalent oxygen. II. Phenanthroxylys, A., i, 1149.
- Goldschmidt, Stefan, and Ludwig Strohmenger**, aromatic chloroamines. II., A., i, 1004.
- Goldschmidt, Stefan, and Bernard Wurschmitt**, amine-oxidation. VI. Radicles as intermediate stages in chemical reactions, A., i, 1139.
amine-oxidation. VII. The oxidation of aniline, A., i, 1139.
- Goldstein, Kurt.** See **Hans Fringsheim**.
- Gomberg, Moses, and Edgar C. Britton**, 2:2'-sulphonidotriphenylmethyl, A., i, 163.
- Gomberg, Moses, and C. C. Buchler**, benzyl ethers of carbohydrates, A., i, 112.
- Gomberg, Moses, and Wesley Minnis**, phenylthioxanthyl, A., i, 163.
- Gomberg, Moses, and F. W. Sullivan, jun.**, triphenylmethyl. XXX. Diphenyl- β -naphthylmethyl and the colour of free radicles, A., i, 929.
- González, F.** See **Enrique Moles**.
- Goode, Kenneth H.**, continuous reading electro-titration apparatus, A., ii, 307.
- Goodhue, Elbridge A.** See **Roscoe G. Dickenson**.
- Goodson, John Augustus**, the constituents of the flowering tops of *Artemisia afra*, Jacq., A., i, 1099.
- Goodyear Tire and Rubber Co.**, preparation of thiocarbamides, A., i, 236.
- Goos, Oskar.** See **Emil Knoevenagel**.
- Gordon, John.** See **James Walter McLeod**.
- Gordon, Neil E.**, origin of soil colloids, A., i, 1227.
- Gordon, Neil E., and Ebenezer Emmet Reid**, solubility of liquids in liquids; the partition of the lower acids, particularly formic, between water and various organic solvents, A., ii, 824.

Gordon, *Neil E.* and *E. B. Starkey*, influence of soil colloids on availability of salts, A., i, 1104.

Gordon, *Neil E.* See also *E. B. Starkey*.

Gordon, *Samuel G.*, kaeleyite, a new lead sulphantimonite from Bolivia, A., ii, 859.

Gore, *S. N.*, the "cotton wool plug" test for indole, A., ii, 535.

Goris, *A.*, and *P. Costy*, preparation and racemisation of hyoscyamine and its sulphate, A., i, 1174.

urea and urease in fungi, A., i, 1220.

Goris, *A.*, and *A. Larssonneau*, chemical composition of belladonna leaves, A., i, 1099.

detection of small quantities of pyridine, A., ii, 795.

Goris, *A.*, and *A. Liot*, the culture of *Bacillus pyocyaneus* on definite artificial media, A., i, 407.

Goris, *A.*, and *Ch. Vischniac*, essential oil of violet roots, A., i, 848.

Gorter, *K.*, the precursor of Indian-yellow, A., i, 932.

Gortner, *Ross Aiken*, and *Walter F. Hoffman*, an interesting colloid gel, A., ii, 130.

Gortner, *Ross Aiken*. See also *Walter F. Hoffman*.

Gottlieb-Billroth, *Hans*. See *Kurt Heinrich Meyer*.

Gottschaldt, *Else*. See *Otto Mumm*.

Gottschalk, *A.* See *Werner Lipschitz*.

Goudsmit, *S.*, doublets in the visible spectrum, A., ii, 462.

Gouin, *A.*, and *P. Andouard*, disappearance of nitrogen during digestion, A., i, 698.

Gould, *V. L.* See *Carl S. Marvel*.

Gowland, *William*, obituary notice of, T., 2907.

Goy, *S.*, and *E. Wende*, the wax of corpses, A., i, 969.

Graaff, *C. de*. See *Jacob Böeseken*.

Grab, *Max von*, pyruvic acid as an intermediary in the alcoholic fission of dextrose, A., i, 306.

Grabner, *Alfred*, so-called rapic acid, A., i, 519.

Gränacher, *Ch.*, the use of rhodanine in organic syntheses. I. Furylalanine, A., i, 849.

Gränacher, *Ch.*, *H. Reis*, and *E. Pool*, rhodanine. II., A., i, 576.

Gränacher, *Ch.*, and *P. Schauffelberger*, oxidation of aliphatic hydrocarbons with nitrogen peroxide. II., A., i, 513.

Gränacher, *Ch.* See also *Paul Karrer*.

Grafe, *Edward*, and *E. von Bedwitz*, the function of the thyroid gland in the regulation of temperature and in the metabolism of fever, A., i, 491.

Graham, *Hugh*, and *Alexander Killen Macbeth*, the labile nature of the halogen atom in organic compounds. III. The absorption spectra of bromomalonic derivatives and nitro-paraffins, and their bearing on the question of an oxygen-halogen linking, T., 1109.

the labile nature of the halogen atom in organic compounds. VII. Absorption spectra of the halogen derivatives of some cyclic compounds, and their bearing on the question of an oxygen-halogen linking, T., 2601.

Graham, *Hugh*. See also *Samuel Smiles*.

Graham, *J. J. T.*, and *C. M. Smith*, errors caused by nitrates and nitrites in the estimation of arsenic by the distillation method and a means for their prevention, A., ii, 314.

Grainger, *Herbert Henry*. See *Edward de Barry Barnett*.

Gralka, *Richard*, and *Hans Aron*, accessory food factors. II. Importance of water-soluble extractives, A., i, 395.

Gram, *H. C.*, fibrinogen content of human blood, A., i, 288.

estimation of the percentage of fibrin in blood and plasma, A., ii, 240.

the gradual darkening of hematin solutions in colorimetric estimations, and its prevention, A., ii, 886.

Gram, *H. C.* See also *A. Norgaard*.

Gramont, (*Comte*) *Arnaud de*, and *Gustave Adolphe Hemsalech*, the evolution of the spectrum of magnesium under the influence of increasing electric fields; applications to astrophysics, A., ii, 243.

Gramont, (*Comte*) *Arnaud de*. See also *Gustave Adolphe Hemsalech*.

Grandchamp, *L.* See *Philippe Malvezin*.

Grandmougin, *Eugene*, the homonuclear dibromoanthraquinones, A., i, 43.

the intermediate products in the synthesis of alizarin, A., i, 44.

octabromoindigotin, A., i, 53.

the halogenated indigotins, A., i, 180.

sulphobenzide [diphenylsulphone], A., i, 251.

some new derivatives of sulphobenzide [diphenylsulphone], A., i, 381.

the halogenated isatins, A., i, 368.

the acylated and alkylated leuco-indigotins, A., i, 470.

the quindolines, A., i, 584.

- Granger, F. S.**, and **John Maurice Nelson**, oxidation and reduction of quinol and quinone from the point of view of electromotive force measurements, A., i, 43.
- Grantham, G. E.**, the infra-red absorption spectra of alkali hydroxides, A., ii, 244.
- Graser, Johanna.** See **Richard Willstätter**.
- Grau, R.** See **Heinrich Ley**.
- Gray, H. Le B.** See **Gurney O. Gutekunst**.
- Gray, William Herbert.** See **Robert George Fargher**.
- Greaves, Joseph E., E. G. Carter,** and **Yeppa Lund**, influence of salts on asofication in soil, A., i, 973.
- Greaves, Joseph E.** See also **C. T. Hirst**.
- Grebe, Leonhard,** and **H. Konen**, band spectra of isotopes, A., ii, 4.
- Green, Arthur George.** See **British Dyestuffs Corporation, Ltd.**
- Green, Stanley Joseph.** See **British Dyestuffs Corporation, Ltd.**
- Greene, Herbert,** and **Robert Robinson**, the mechanism of the formation of benzoylbenzoin by treatment of benzoylmandelonitrile with an alcoholic solution of sodium ethoxide, T., 2182.
- Greenfield, R. E.,** and **Arthur M. Buswell**, investigation, by means of the hydrogen electrode, of the chemical reactions involved in water purification, A., ii, 653.
- Greer, Frank E.** See **Frederick W. Heyl**.
- Greeneke, Hellmuth.** See **Jakob Meisenheimer**.
- Gregory, John Walter**, ore deposits and their genesis in relation to geographical distribution, T., 750.
- Greinaacher, Heinrich**, luminescence of flame ions in the air spark, A., ii, 250.
- Greiner, Irens**, the estimation of small quantities of dextrose by Bertrand's process, A., ii, 400.
- Greinert, Wilhelm**, oxidation of some sugar acids, A., i, 1111.
- Grenet, Louis.** See **Georges Charpy**.
- Greulich.** See **G. Meyer**.
- Greune, H.** See **Theodor Zincke**.
- Griffin, Roger C.,** and **H. C. Parish**, the penetrability of filter-paper, A., ii, 309.
- Griffith, Robert Owen**, the hydrolysis of a salt formed from a weak acid and a weak base, A., ii, 420.
- Griffiths, Evan Dalton.** See **Fred Barrow**.
- Grigaut, A.**, estimation of uric acid in blood, A., ii, 405.
- Grigaut, A.,** and **P. Eizine**, precipitation of proteins by metaphosphoric acid; application to the analysis of blood, pathological liquids, and cerebrospinal fluid, A., ii, 886.
- Grigaut, A.** See also **A. Chauffard**.
- Griggs, Mary A.**, alkaline hydrolysis of casein, A., i, 182.
- Grignard, Victor,** and **A. C. Purdy**, $\alpha\beta$ -dichlorodiethyl ether, A., i, 802.
- Grim, F. V.** See **Walter A. Patriek**.
- Grimm, H.**, ionic properties and crystallochemical relationships. I. The properties of the ions which appear in crystals, A., ii, 127.
- Grimm, Hans Georg**, isomorphism and ionic structure, A., ii, 488.
structure of the ions of the rare earths. A., ii, 635.
periodic system of the atomic ions, A., ii, 635.
ionic properties and chemical facts.
IV. Lattice energy and the work of ionisation of inorganic compounds, A., ii, 690.
ionic properties and chemical facts.
V. Connexion between heat of formation, lattice energy, and the ionic properties, A., ii, 690.
- Grimme, Clemens**, oil from the seeds of *Jatropha curcas*, L., A., i, 98.
- Grimmel, Harry.** See **Adolf Windaus**.
- Grinberg, A. A.**, chemoluminescence. I., A., ii, 806.
- Grindley, Harry Sands.** See **T. S. Hamilton**.
- Grisard, Gustav.** See **Franz Fichter**.
- Grizskiewicz-Trochimowski, E.**, products of polymerisation of hydrocyanic acid, A., i, 723.
- Grobet, Edouard**, reactions of sodium hydroxide with salts of aluminium, A., ii, 573.
- Grobet, Edouard.** See also **Paul Dutoit**.
- Groeger, Gerald.** See **Adolf Franke**.
- Greenewege, J.**, the occurrence of emulsion in *Saccharomycetes* and the existence of a specific enzyme cellobiase, A., i, 903.
denitrification with formates; influence of the kation, A., i, 971.
- Groll, J. Temminck**, the influence of hydrogen-ion concentration on the action of pancreatic amylase, A., i, 600.
- Gromelski, B.** See **G. Leenderts**.
- Groningen, P. van.** See **Willem Reinders**.
- Gros, R.** See **J. Bougault**.
- Gross, Erwin G.,** and **Frank Pell Underhill**, metabolism of inorganic salts. I. The inorganic-ion balance of the blood in parathyroid tetany, A., i, 1210.

- Gross, Paul M.** See *James Kendall*.
- Gross, R. Eberhard**, the protamines, A., i, 784.
new small autoclave for hydrolysis experiments, A., ii, 561.
- Grottrian, Walter**, the *L* doublet of neon, A., ii, 179.
- Grottrian, Walter**. See also *J. Franek*.
- Grounds, Arthur**, the constitution of anthracite, A., ii, 385.
- Grube, Georg**, chemical and electrochemical behaviour of salts of the acids of lead, A., ii, 570.
- Gruber, Heinrich**. See *Julius von Braun*.
- Grude, Frithjof**. See *Paul Askenasy*.
- Grün, Adolf**, the constitution of glycerides from the point of view of the co-ordination theory, A., i, 420.
the co-ordination forms of glycerides, A., i, 621.
- Grün, Adolf**, and *Th. Wirth*, synthesis of Δ^8 -decanoic acid, A., i, 804.
 Δ^8 -decanoic acid, a previously unknown acid from butter, A., i, 806.
- Grün, Adolf**, and *Franz Wittka*, preparation and alkyl interchange of cellulose esters; cellulose stearate and laurate, A., i, 114.
- Grüneisen, E.**, and *E. Merkel*, velocity of sound in air and hydrogen at 0° and 1 atm., A., ii, 190.
- Grünhut, Leo**, and *J. Weber*, the action of amino-acids on sugars, A., i, 235.
- Grünhut, Leo**. See also *Wilhelm Frensenius*.
- Grünstein, Nathan**, preparation of aldol from acetaldehyde, A., i, 111.
preparation of butaldehyde and butyl alcohol from crotonaldehyde, A., i, 112.
- Gruyter, C. J. de**. See *Andreas Smits*.
- Grusowska, (Mme) Z.**, action of ferments on laminarin, A., i, 11.
- Gudden, B.**, and *Robert Pohl*, phosphorescent zinc sulphide containing copper, A., ii, 680.
- Guénot, L.** See *Louis Fournier*.
- Günther-Schulze, A.**, electromotive behaviour of aluminium, A., ii, 110.
dependence of the base equilibrium in permutite on the concentration of the surrounding solution, A., ii, 486.
determination of the complex formation in aqueous solutions of copper salts by means of permutite, A., ii, 504.
the dissociation of the chlorides of bivalent metals in aqueous solution, A., ii, 765.
- Günsberg, Ludwig**, theobromine excretion and theobromine diuresis, A., i, 703.
- Guerbet, Marcel**, the characterisation of the colouring matter of saffron; its use in investigations relating to laudanum poisoning, A., ii, 793.
- Guha, Praphulla Chandra**, constitution of the so-called dithiourazole of Martin Freund. I., A., i, 875.
constitution of the so-called dithiourazole of Martin Freund. II. New methods of synthesis, isomerism, and poly-derivatives, A., i, 876.
- Guichard, Marcel**, a static method of study of hydration, A., ii, 555.
adsorption and its bearing on catalysis, A., ii, 630.
- Guild, F. N.**, identity of flagstaffite with terpin hydrate, A., ii, 76.
- Guillaumin, A. J. A.** See *Louis Jacques Simon*.
- Guillaumin, Ch. O.**, estimation of uric acid in blood, A., ii, 170.
estimation of uric acid and urates in blood, A., ii, 796.
estimation and constitution of a fraction of the uric acid in blood, A., ii, 796.
- Guillemet, R.** See *Henri Gault*.
- Guillet, Léon**, the thermal treatment of certain complex aluminium alloys, A., ii, 69.
magnesium-cadmium alloys, A., ii, 570.
the alloys of cerium, A., ii, 572.
- Guillet, Léon**, and *Jean Cournot*, the variation of the mechanical properties of metals and alloys at low temperatures, A., ii, 261.
- Guinot, Henri**. See *André Job*.
- Guioth, Jean**. See *Marcel Sommelet*.
- Gulbransen, R.** See *Carl Hamilton Browning*.
- Gunkel, L.** See *Emil Heuser*.
- Gunnaiya, D.** See *A. L. Narayan*.
- Guntz, A. A.**, phosphorescent zinc sulphide, A., ii, 502.
- Gupta, Biraj Mohan**, and *Jocelyn Field Thorpe*, the formation of bromine derivatives of carbon compounds without the production of hydrogen bromide, T., 1896.
- Gustafson, A. F.**, the effect of drying soils on the water-soluble constituents, A., i, 703.
- Gustaver, Bror**, the adsorption problem; sorption of vapours by charcoal, A., ii, 479.
- Gutbier, Alexander**, and *K. Dürrwächter*, gold sulphides, A., ii, 513.

Guthier, *Alexander*, and *R. Emslander*, influence of freezing on colloidal selenium. III., A., ii, 283.
 protecting colloids. XII. Gelatin as protecting colloid. II. Colloidal selenium, A., ii, 625.
 Guthier, *Alexander*, *Fr. Heinrich*, and *J. Huber*, influence of freezing on colloidal selenium. II., A., ii, 142.
 Guthier, *Alexander*, and *J. Huber*, protective colloids. XI. Carrageen as a protective colloid. I. General colloid-chemical investigation of the extract of Irish moss, A., ii, 203.
 Guthier, *Alexander*, *J. Huber*, and *O. Kuppinger*, the analytical chemistry of colloidal disperse systems. I. Estimation of silver ion in the presence of colloidal silver, A., ii, 396.
 Guthier, *Alexander*, *J. Huber*, and *W. Schieber*, a rapid dialyser. A., ii, 551.
 Guthier, *Alexander*, *J. Huber*, and *A. Zweigle*, protective colloids. XII. Gelatin as a protective colloid. I. Colloidal silver, A., ii, 485.
 Guthier, *Alexander*, and *F. Krauss*, bromo-salts of ruthenium [ruthenium-bromides], A., i, 16.
 Guthier, *Alexander*, and *A. Mayer*, a simple dialyser, A., ii, 353.
 Guthier, *Alexander*, *E. Sauer*, and *F. Schelling*, action of alum on animal glue, A., i, 783.
 Guthier, *Alexander*, and *K. Staib*, estimation of zinc as sulphate, A., ii, 396.
 Guthier, *Alexander*, *A. Wolf*, and *A. Kiess*, protective colloids. XI. Carrageen as protective colloid. II. Colloidal silver, A., ii, 203.
 Gutekunst, *Gurney O.*, and *H. Le B. Gray*, the 6-alkyloxy-2-methylquinolines, A., i, 950.
 Guthrie, *Charles Claude*, a simplified form of apparatus for air analysis, A., ii, 78.
 a gas receiver of convenient and practical form for sampling expired air for analysis, A., ii, 158.
 Gutmann, *August*, action of compounds of nitric oxide and hydroxylamine on tertiary sodium arsenite, A., ii, 844.
 Guy, *Ruth A.*, limitations of the modified Lewis-Benedict method of blood sugar estimation, A., ii, 94.
 Guye, *Charles Eugene*, and *R. Rüdy*, new method of determining molecular diameters by the electromagnetic rotation of the discharge in gases, A., ii, 278.
 Guye, *Philippe Auguste*, obituary notice of, T., 2909.

Guye, *Philippe Auguste*, and *T. Batuecas*, the compressibility at 0° and less than 1 atmosphere and the divergence from Avogadro's law of several gases, A., ii, 617.
 Gyemant, *A.*, cataphoresis of water in organic liquids, A., ii, 684.
 Gyemant, *A.* See also *Herbert Freundlich*.
 György, *Paul*. See *Ernst Freudenberg*.

H.

Haan, *J. de*, the glycogen content of white blood corpuscles, A., i, 484.
 Haan, *J. de*, and *S. van Creveld*, the relation between blood-plasma and tissue fluids, especially the aqueous humour and the cerebrospinal fluid. I. The sugar content and the question of the combined sugar, A., i, 295.
 Haar, *Anne Wilhelm van der*, saponins. V. Hederin and its hederagenin, A., i, 160.
 oxydases, A., i, 284.
 saponins. VII., A., i, 565.
 saponins. VIII. The saponins from the leaves of *Aralia montana*, Bl. [galacturonoid-saponins and their magnesium and calcium salts], A., i, 1168.
 Haar, *Anne Wilhelm van der*, and *A. Tamburello*, saponins. VI. Hederagenin, A., i, 160.
 Haar, *J. van der*, apparatus for facilitating the estimation of dextrose by titration with permanganate by the Mohr-Bertrand method, A., ii, 790.
 Haas. See Röhm.
 Haas, *A. R. C.* See *F. C. Bauer*.
 Haas, *(Frl.) E.* See *Heinrich Wieland*.
 Haber, *Fritz*, amorphous precipitates and crystalline sols, A., ii, 553.
 Haber, *Fritz*, and *W. Zisch*, excitation of gas-spectra during chemical reactions, A., ii, 461.
 Hackl, *Oskar*, estimation of very small quantities of arsenic in silicate rocks, A., ii, 159.
 detection and estimation of small quantities of nickel and cobalt in silicate rocks, A., ii, 458.
 Hackmack, *D.* See *R. Siebeck*.
 Hadding, *Assar*, analysis of minerals by the Röntgen spectrograph, A., ii, 780.
 the occurrence of germanium in cassiterite, A., ii, 855.
 Haefelin, *Gerhard*. See *Paul Pfeiffer*.

- Haegermann, Gustav.** See *Richard Lorenz*.
- Hägglund, Erik, Nils Löfmann, and Eduard Färber,** cellulose acetate from wood cellulose, A., i, 323.
- Haehn, H.** See *F. Hayduck, and Georg Schroeter*.
- Hämäläinen, Reino,** a new method for the detection of methyl alcohol, A., ii, 723.
- Häuselmann, L.** See *H. Zschokke*.
- Haffner, F.** See *Alb. Jodlbauer*.
- Hagen, J.** See *Fritz Mayer*.
- Hagenbach, August, and R. Percy,** the absorption spectrum of potassium permanganate, A., ii, 600.
- Hager, Josef.** See *Stigmund Fränkel*.
- Haggard, Howard W., and Thomas J. Charlton,** the fate of sulphides in the blood, A., i, 287.
- Haggard, Howard W., and Vandell Henderson,** influence of hydrogen sulphide on respiration, A., i, 1206.
- Haggenmacher.** See *Emil Baur*.
- Hagman, Sidney M.** See *Hjalmar Johansson*.
- Hahn, Arnold,** "double-nitrogen," a diagnostic for endogenous protein-breakdown, especially for hidden supuration, A., i, 291.
- Hahn, Dorothea von.** See *Friedrich Vincenz von Hahn*.
- Hahn, Erich.** See *Julius von Braun*.
- Hahn, Friedrich L.,** still-head for the prevention of spray, A., ii, 368.
- Hahn, Friedrich L.,** still-head for the preparation of green manganous sulphide, A., ii, 880.
- Hahn, Friedrich L.,** separation of antimony from arsenic and tin, A., ii, 877.
- Hahn, Friedrich L. [with G. Leimbach, and H. Windisch],** contradictions and errors in analytical chemistry. I. The precipitation of aluminium by thiosulphate and its separation from iron. II. The ageing of volumetric thiosulphate solutions, A., ii, 873.
- Hahn, Friedrich L., and G. Leimbach,** a peculiar catalytic reaction for the detection and a method for the estimation of the smallest traces of copper [also a lecture experiment], A., ii, 870.
- Hahn, Friedrich Vincenz von,** kinetic and static coagulation measurements of suspensoids, A., ii, 37.
- Hahn, Friedrich Vincenz von,** sulphide sols. III. Sol preparation by hydrolysis, A., ii, 856.
- Hahn, Friedrich Vincenz von, and Dorothea von Hahn,** technical sedimentation analysis. I., A., ii, 705.
- Hahn, Friedrich Vincenz von.** See also *Wolfgang Ostwald*.
- Hahn, Otto,** the existence of the new radioactive element uranium-V described by Piccard and Stahel, A., ii, 340.
- Hahn, Otto, and Lise Meitner,** protactinium content of pitchblende residues and the branching relationship of the actinium series, A., ii, 185.
- Haller, Ekkehard,** the relation between bacteria, spores, and formaldehyde, A., i, 408.
- Haller, Ekkehard,** the bactericidal after-effect of formaldehyde solutions, A., i, 408.
- Hainsworth, William R., and Duncan A. MacInnes,** effect of hydrogen pressure on the electromotive force of a hydrogen-calomel cell. I., A., ii, 467.
- Hajós, K.** See *László Karczag*.
- Hakomori, Shin-ichirō,** catalytic analyses. I. Estimation of chloric acid, A., ii, 389.
- Halban, Heinrich von, and K. Siedentopf,** application of photo-electric cells to the measurement of the light absorption in solutions. II., A., ii, 332.
- Halberkann, Josef,** 5-methoxydioxindole and 5-methoxyisatin, A., i, 172.
- Halberkann, Josef,** derivatives of quinic acid, A., i, 174.
- Halberkann, Josef,** transformation of the diphenyl-, phenyl-*p'*-tolyl-, and di-*p'*-tolyl-amides of toluene-*p*-sulphonic acid, A., i, 1132.
- Haley, D. E., and J. F. Lyman,** castor bean lipase, its preparation and some of its properties, A., i, 390.
- Hall, Archibald John.** See *Arthur Ernest Everest*.
- Hall, Claude Haines, jun.,** electrical precipitation of colloids, A., ii, 555.
- Hall, Dorothy,** estimation of small amounts of molybdenum in tungsten, A., ii, 660.
- Hall, Dorothy.** See also *Hobart Hurd Willard*.
- Halla, F., and K. Hirschke,** the complex cuprammoniates
 $\text{Cu}(\text{CN})_2, \text{CuCN}, 2\text{NH}_3;$
 $\text{Cu}(\text{CN})_2, 4\text{CuCN}, 4\text{NH}_3;$
 $\text{Cu}(\text{ONS})_2, \text{Cu}(\text{CN})_2, 5\text{CuCN}, 6\text{NH}_3,$
 and the analytical methods for their identification, A., ii, 768.
- Haller, Albin, and Edouard Bauer,** syntheses by means of sodamide. IX. The preparation of $\beta\beta$ -dialkyl- α -hydri-
 dones or 2:2-dialkylindan-1-ones, A., i, 258.
- Haller, Albin, and Eugène Benoit,** syntheses by means of sodamide. XI. Substitution derivatives of benzyl-
 cyclopropane, A., i, 350.

- Haller, Albin**, and **Paul Bondia**, syntheses by means of sodamide. X. The *p*-aminobenzylidene- and *p*-aminobenzyl-camphors and some of their derivatives, A., i, 356.
- Haller, Albin**, and (*Mme*) **Pauline Ramart-Lucas**, new distinctive characteristics of the three propan- β -ol-camphorcarboxylolides melting respectively at 141°, 117–118°, and 89–90°, A., i, 460.
- Haller, Emma**. See **Kurt Lindner**.
- Hallimond, Arthur Francis**, [with **E. G. Radley**], glauconite from Lewes, Sussex; constitution of glauconite, A., ii, 861.
- Hallwass, F.** See **Walther Borsche**.
- Halphen, Hede**, standard dropping pipette, A., ii, 221.
- Hamburger, Hartog Jakob**, permeability of the glomerulus membrane for stereoisomeric sugars, with special reference to galactose, A., i, 490.
change of permeability with special reference to stereoisomeric sugars, A., i, 491.
estimation of the relative numbers of red blood-corpuscles of differing resistance (osmotic resistance curves) by means of sodium sulphate, A., i, 606.
the permeability of the glomerulus membrane for stereoisomeric sugars, A., i, 790.
- Hamburger, Robert**. See **Ernest Zerner**.
- Hamburger, Rudolf J.**, the importance of potassium and calcium ions for the production of artificial cedema and for the width of the blood vessels, A., i, 606.
- Hamburger, S.**, preparation of protocatechualdehyde, A., i, 556.
- Hamburger-Glaser, Edith**. See **Heinrich Biltz**.
- Hamer, (Miss) Frances Mary**. See **William Hobson Mills**.
- Hamilton, Cliff S.** See **W. Lee Lewis**.
- Hamilton, Leicester F.** See **Waldemar Lindgren**.
- Hamilton, T. S., W. B. Nevens**, and **Harry Sands Grindley**, estimation of the amino-acids of feeding stuffs, A., ii, 92.
- Hammarsten, Einar**, and **Erik Jorpes**, the "coupled" nucleic acid from the pancreas. II., A., i, 887.
- Hammarsten, H.**, the equilibrium between some organic substances, A., ii, 830.
- Hammarsten, Olof**, the action of chymosin and pepsin. VII. Further experiments on the purification of the enzyme of the stomach, A., i, 958.
the action of chymosin and pepsin. VIII. Relative sensitiveness to alkali of the stomach enzymes of the calf and pig, A., i, 958.
- Hammett, Frederick S.**, creatinine and creatine in muscle extracts. III. Concerning the presence of enzymes in muscle tissue which have creatine and creatinine as their substrates, A., i, 1090.
- Hammett, Frederick S.**, and **E. T. Adams**, a colorimetric method for the estimation of small amounts of magnesium, A., ii, 587.
- Hammett, Frederick S., Joseph E. Newrey**, and **John H. Müller**, erythropoietic action of germanium dioxide, A., i, 1086.
- Hamnick, Dalziel Llewellyn**, and **Alford Reginald Boeree**, preparation of atrioxymethylene and a new polymeride of formaldehyde, T., 2738.
- Hamnick, Dalziel Llewellyn**, and **George Hazlewood Locket**, preparation of sodium and potassium phthalimide, T., 2862.
- Hanak, A.**, estimation of sugar by titration with alkali of the cuprous oxide precipitated from Fehling's solution, A., ii, 166.
- Hance, F. E.** See **Louis Monroe Dennis**.
- Handorf, Heinrich**, extraction of small quantities of liquids by the Soxhlet apparatus, A., ii, 496.
detection of the veronal group; diagnosis of veronal intoxication, A., ii, 884.
- Handovsky, Hans**, sensitiveness of cells to poison as a function of their colloid-chemical conditions, A., i, 697.
- Hanke, Milton Th.**, and **Karl K. Koessler**, proteinogenous amines. XII. The production of histamine and other iminazoles from histidine by the action of micro-organisms, A., i, 406.
proteinogenous amines. XIII. The electronic interpretation of certain biochemical phenomena, A., i, 408.
proteinogenous amines. XIV. A microchemical colorimetric method for estimating tyrosine, tyramine, and other phenols, A., ii, 322.
proteinogenous amines. XV. A quantitative method for the separation and estimation of phenols including phenol, *o*-, *m*-, and *p*-cresols, and *p*-hydroxyphenylacetic, *p*-hydroxyphenylpropionic and *p*-hydroxyphenyl-lactic acids, tyrosine, and tyramine, A., ii, 322.

- Hann, Raymond M.** See *Edgar Theodore Wherry*.
- Hanner, A.** See *Balthasar Pfyl*.
- Hanselmayer, Franz.** See *Alois Zinke*.
- Hansgirt, Fritz**, perylene, A., i, 245.
- Hanson, A. W.**, method for the estimation of procaine [novocaine], A., ii, 405.
- Hantge, Ernst.** See *Kurt Arndt*.
- Hantzsch, Arthur [Rudolf]**, the alleged hexavalency of carbon in carbonium and dyestuff salts, A., i, 24.
the constitution of carbonium salts, A., i, 24, 820.
the coloured alkali salts of triphenylmethane, and triphenylmethyl as an amphoteric ion, A., i, 25.
the so-called halochromism of triphenylmethane derivatives, A., i, 26.
the co-ordination number of carbon, A., i, 26.
halochromism and "solvatochromism" of distyryl ketone and simpler ketones and of their ketonic chlorides, A., i, 556.
- Hantzsch, Arthur**, [with *Walther Meyer*], supposed cases of isomerism in the isatin series, A., i, 1177.
- Hanzawa, Toratoro.** See *Francis E. Rice*.
- Hara, H.** See *Frederick G. Keyes*.
- Haramaki, Katsumi**, the relative activities of the secretins of the digestive tract, A., i, 698.
- Hardegg, R.** See *K. Hugo Bauer*.
- Harden, Arthur**, and *Francis Robert Henley*, the function of phosphates in the oxidation of dextrose by hydrogen peroxide, A., i, 433.
- Harden, Arthur.** See also *Arthur William Bacot*.
- Harding, T. Swann**, the preparation of lævulose, A., i, 919.
use of invertase for sucrose estimation, A., ii, 167.
estimation of the activity of invertase, A., ii, 800.
- Hardy, Paul**, the relation between Vitali's reaction and the constitution of the alkaloids which give it, A., i, 948.
the Vitali reaction for cocaine, A., ii, 95.
the volatilisation and hydrolysis of atropine in toxicology, A., ii, 796.
- Harger, Rolla N.**, the oxidation of quinol in the presence of aliphatic amines, A., i, 539.
- Härl, Paul**, a red colouring matter produced by the action of *p*-dimethylaminobenzaldehyde on normal urine, A., i, 88.
- Harington, Charles Robert.** See *Jonathan Meakins*.
- Harker, George**, the temperature of the vapour arising from boiling saline solutions, A., ii, 28.
- Harkins, William Draper**, the separation of chlorine into isotopes, A., ii, 280.
the stability of atom nuclei, the separation of isotopes, and the whole number rule, A., ii, 702.
- Harkins, William Draper**, and *Dwight T. Ewing*, high pressure due to adsorption, and the density and volume relations of charcoal, A., ii, 123.
high pressure due to adsorption, A., ii, 197.
- Harkins, William Draper**, and *Anson Hayes*, separation of the element chlorine into isotopes (isotopic elements); the heavy fraction from the diffusion, A., ii, 140.
- Harkins, William Draper**, and *S. L. Madorsky*, a graphical study of the stability relations of atom nuclei, A., ii, 490.
- Harkins, William Draper**, and *Lathrop E. Roberts*, orientation of molecules in surfaces. VII. Vaporisation in steps as related to surface formation, A., ii, 422.
- Harkins, William Draper.** See also *Robert S. Mullikan*.
- Harle, H.**, viscosities of the hydrogen haloids, A., ii, 140.
- Harned, Herbert S.**, activity coefficients and colligative properties of electrolytes, A., ii, 255.
- Harned, Herbert S.**, and *Robert Pfannstiel*, velocity of hydrolysis of ethyl acetate, A., ii, 832.
- Harned, Herbert S.**, and *Harry Selts*, ion activities in homogeneous catalysis; formation of *p*-chloroacetanilide from *N*-chloroacetanilide, A., ii, 631.
- Harries, Carl Dietrich**, preparation of fatty acids, aldehydes, and ketones from mineral and tar oils, A., i, 514.
- Harries, Carl Dietrich**, and *Fritz Evers*, estimation of the molecular magnitude of caoutchouc by chemical methods, A., i, 357.
- Harries, Carl Dietrich**, and *W. Nagel*, aleuritic acid, A., i, 522.
- Harris, D. T.**, active hyperæmia, A., i, 696.
- Harris, Hinton John.** See *Frederick Daniel Chattaway*.
- Harris, John Edmund Guy**, and (*Sir*) *William Jackson Pope*, isoquinoline and the isoquinoline-reds, T., 1029.
- Harris, M.** See *Samuel Palkin*.

- Harrison, C. W.**, distillation method for the estimation of santalol in santal oil, A., ii, 400.
- Harrison, Douglas Creese.** See *Samuel Smiles*.
- Harrison, D. N.** See *Thomas Ralph Merton*.
- Harrison, Edward Frank**, memorial to, T., 2894, 2917.
- Harrison, George R.**, absorption of light by sodium and potassium vapours, A., ii, 679.
- Harrison, William**, electrical theory of adsorption, A., ii, 198.
- Harrisson, J. W. E.**, the U.S.P. test for acetone in alcohol, A., ii, 667.
- Hart, Harry B.** See *J. Newton Pearce*.
- Hart, Merrill C.**, and *Arthur D. Hirschfelder*, some derivatives of saligenin, A., i, 38.
- Hart, Merrill C.**, and *Wilbur B. Payne*, toxicity of neocarsphenamine [neosalvarsan], A., i, 704.
- Harter, L. L.**, and *J. L. Welmer*, pectinase produced by different species of *Rhizopus*, A., i, 507.
- Hartley, Harold Brewer, A. O. Ponder, Edmund John Bowen, and Thomas Ralph Merton**, attempt to separate the isotopes of chlorine, A., ii, 280.
- Hartley, Harold Brewer.** See also *Cyril Norman Hinshelwood*.
- Hartman, H.** See *P. E. Verkade*.
- Hartmann, Hellmuth.** See *Otto Ruff*.
- Hartmann, M.**, and *M. Seiberth*, dialkylamides of nicotinic acid, A., i, 679.
- Hartung, Ernst Johannes**, the action of light on silver bromide, T., 682. observations on the construction and use of the Steele-Grant microbalance, A., ii, 495.
- Harvey, E. Newton**, bioluminescence. XIV. The specificity of luciferin and luciferase, A., i, 299.
- Hasenbäumer, J.** See *Josef König*.
- Hasenjäger, H.** See *Karl Fries*.
- Hasenöhr, Rudolf**, and *Julius Zellner*, the chemistry of the higher fungi. XV. Chemical relations between the higher fungi and their substrate. II., A., i, 904.
- Hashimoto, Tokudji.** See *Takaoki Sasaki*.
- Hassé, H. R.** See *(Sir) J. B. Henderson*.
- Hassel, Odd.** See *Heinrich Goldschmidt*.
- Hasselblatt, Meinhard**, the influence of pressure on spontaneous crystallisation, A., ii, 35. crystallisation velocity under high pressure, A., ii, 35. the fusion diagram of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O} + \text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ at pressures of 1 to 3000 kilo./cm.², A., ii, 61.
- Hastings, A. Baird**, lactic acid in the blood of dogs in exercise, A., i, 81.
- Hastings, A. Baird**, and *Donald D. van Slyke*, determination of the three dissociation constants of citric acid, A., i, 985.
- Hastings, A. Baird.** See also *Joshua Harold Austin*, and *Glenn E. Cullen*.
- Hatcher, Robert A.** See *Soma Weiss*.
- Hattori, Kenzo**, colloidal structure of red blood corpuscles and hemolysis. III. Ultramicroscopic investigation of lipoids, A., i, 192.
- Hauenstein, J.**, blood and metabolism studies with radium emanations, A., i, 80.
- Haughton, John L.**, and *G. Winifred Ford*, systems in which metals crystallise, A., ii, 825.
- Haurowitz, Felix**, the gonads of *Rhizostoma cuvieri*, A., i, 1210.
- Hauser, Ernst.** See *Hermann Staudinger*.
- Haushalter, J.** See *M. Bonnet*.
- Hawkins, James Alexander**, the influence of solvents on the velocity of formation of quaternary ammonium salts, T., 1170.
- Haworth, Robert Downs**, and *Arthur Lapworth*, the direct acetalisation of aldehydes, T., 76.
- Haworth, Walter Norman**, and (*Miss*) *Grace Cumming Leitch*, the constitution of the disaccharides. VI. The biase of amygdalin, T., 1921.
- Hayami, Ryonosuke**, equilibrium in the system copper sulphate, ammonium sulphate, potassium sulphate, and water at 25°0, A., ii, 748.
- Hayashi, Toworu.** See *Seizaburo Okada*.
- Hayduck, F.**, and *A. Haehn*, zymase formation in yeast. I., A., i, 611.
- Hayes, Anson.** See *William Draper Harkins*.
- Haynes, Dorothy.** See *Marjory Harriott Carré*.
- Haywood, Percy Charles**, the reactivity of alkyl iodides with sodium benzyloxy and the effect of temperature on such reactions, T., 1904.
- Hazen, William**, estimation of small amounts of potassium by the Lindo-Gladding method, A., ii, 658.
- Hazelton, Edith Olive.** See *Ethel Atkinson*.
- Headden, William P.**, a tantalate and columbite from South Dakota, A., ii, 387.
- Healy, Daniel J.**, and *Perry E. Karraker*, the Clark hydrogen-electrode vessel and soil measurements, A., ii, 519.

- Heap, Joseph Greenwood, William Jacob Jones, and John Bamber Speakman**, the preparation of pyridine and of certain of its homologues in a state of purity, A., i, 171.
- Hebler, F.** See *Heinrich Bechhold*.
- Heck, Adolf.** See *G. Meyer*.
- Hedelius, Arvid.** See *Hans von Euler*.
- Hedestrand, Gunnar**, the viscosity of amphoteric electrolytes in solution, A., ii, 821.
- Hedin, Sven Gustav**, the proteolytic enzymes in albuminous urines, A., i, 609.
the proteolytic enzymes of the kidneys, A., i, 1212.
- Hedvall, J. Arvid**, examination by the X-ray spectrum of metallic oxides which are stable at red heat (prepared by different methods and having different properties, A., ii, 300.
the colour of ferric oxide, A., ii, 381.
- Hedvall, J. Arvid, and Gunnar Booberg**, the action of molten alkali chlorides on copper oxide, A., ii, 68.
- Hedvall, J. Arvid, and J. Heuberger**, transference of the acid radicle in the solid phase. I. Carbonates of the alkaline earths and magnesium, A., ii, 768.
- Heene, Richard.** See *Otto Dimroth*.
- Heerdt, Walter.** See *Hugo Weil*.
- Heidelberger, Michael**, preparation of crystalline oxyhæmoglobin, A., i, 962.
- Heidelberger, Michael, and Walter Abraham Jacobs**, syntheses in the cinchona series. IX. Certain quinine and benzoylcinchona salts, crystalline ethyldihydrocupreine (optochin) base, and other derivatives, A., i, 678.
syntheses in the cinchona series. X. Dihydrocinchonincol and the dihydroquinidinols, A., i, 673.
- Heidelberger, Michael.** See also *Walter Abraham Jacobs*.
- Heidrich, Dorothea.** See *Heinrich Biltz*.
- Heiduschka, Alfred, and E. Komm**, keratin, I., A., i, 967.
- Heiduschka, Alfred, and P. Roser**, the composition of beechnut oil (*Oleum fagi sylvaticæ*), A., i, 945.
- Helke, W.**, new detectors of high frequency vibrations, A., ii, 19.
simple methods for the determination of melting points and critical temperatures; the melting point of arsenic, A., ii, 25.
the solidification diagram of the zinc-arsenic alloys, A., ii, 60.
- Hell, Rudolf.** See *Fritz Mayer*.
- Hellbron, Isidor Morris.** See *Edward Charles Cyril Baly, and Johannes Sybrandt Buck*.
- Heilner, G.** See *Carl Mannich*.
- Heimann, Heinrich.** See *Erwin Ott*.
- Hein, Franz**, organo-chromium compounds. II. Abnormal salt formation of chromium pentaphenyl hydroxide; chromium tetraphenyl salts (elimination of a phenyl group), A., i, 76.
organo-chromium compounds. III. Chromium triphenyl hydroxide and its salts, A., i, 77.
- Heinrich, Fr.** See *Alexander Guthrie*.
- Heinze, Fritz.** See *Otto Dimroth*.
- Heinzelmann, Alfred**, rapid estimation of mercury in ores, A., ii, 182.
- Helbronner, André, and W. Rudolfs**, the attack of minerals by bacteria; oxidation of blende, A., i, 706.
- Helferich, Burckhardt**, emulsin, A., i, 390.
- Helferich, Burckhardt, and Max Gehrke**, γ -hydroxyaldehydes. IV. γ -hydroxyaldehydes with tertiary hydroxyl, A., i, 9.
- Helferich, Burckhardt, and Theodor Malkomes**, synthesis and properties of a δ -hydroxyaldehyde, A., i, 431.
- Helferich, Burckhardt, and Johann Adolf Speidel**, α -hydroxy-lactones, A., i, 6.
- Helferich, Burckhardt, and Rudolf Weidenhagen**, synthesis of substances resembling disaccharides from monohydroxyaldehydes, A., i, 1116.
- Heller, Gustav**, 8-hydroxy-2-phenylindazole, A., i, 1066.
- Heller, Gustav**, [with *Walter Benade and Otto Hochmuth*], new isomerism in the isatin series. V., A., i, 1058.
- Heller, Gustav, and Walter Benade**, the nature of isatoids, A., i, 582.
- Heller, Gustav, and Werner Boesneck**, isatogens, A., i, 369.
- Heller, Gustav, Hilde Lauth, and Arnold Buchwaldt**, reactivity of the nitrobenzaldehydes, A., i, 348.
cinnamoylformic acids, A., i, 1024.
- Heller, Gustav, and Paul Lindner**, tetranitroanthracrynone, A., i, 1040.
- Heller, Hans**, a modified reaction of tin, A., ii, 458.
- Heller, Ludwig**, the quantitative action of some factors in the clotting of blood, A., i, 291.
- Helmholtz, K.** See *Willy Marckwald*.
- Helmick, Homer H.**, estimation of thorium in monazite sand by an emanation method, A., ii, 164.
- Helwert, Fritz.** See *Harwig Franzen*.
- Helwig, Herbert.** See *Kurt Lindner*.

- Hemmd, K.** See *G. Bode*.
- Hemingway, A. J.** See *David Avery*.
- Hemmelmayer, Franz**, the dicarboxylic acid which is formed by heating 1 : 5-dihydroxynaphthalene with potassium hydrogen carbonate under pressure, A., i, 836.
- Hemmelmayer, Franz.** See also *Robert Kremann*.
- Hemsalech, Gustave Adolphe, and (Comte) Arnaud de Gramont**, occurrence of spark lines (enhanced lines) in the arc. I. Lead and tin, A., ii, 176.
the occurrence of spark lines (enhanced lines) in the arc. II. Magnesium, zinc, and cadmium, A., ii, 410.
- Hemsalech, Gustave Adolphe.** See also *(Comte) Arnaud de Gramont*.
- Henderson, George Gerald, John McGregor Robertson, and David Christie Brown**, the oxidation of sabinene with chromyl chloride, T., 2717.
- Henderson, G. H.**, α -particles as detonators, A., ii, 606.
- Henderson, James Alexander Russell.** See *Thomas Callan*.
- Henderson, (Sir) J. B., and H. R. Hassé**, thermodynamical theory of explosions. I and II., A., ii, 207.
- Henderson, Lawrence Joseph.** See *Edward F. Adolph*.
- Henderson, Thomas, and Alexander Kullen Macbeth**, the labile nature of the halogen atom in organic compounds. I. Titanium reductions of substituted nitroparaffins, T., 892.
- Henderson, William E.** See *Merle L. Dundon*.
- Henderson, Vandell.** See *Howard W. Haggard*.
- Henglein, Friedrich August**, molecular volumes, physical properties, and molecular models of the halogens, A., ii, 44.
a vapour pressure formula with a general integration constant, A., ii, 118.
regularities in the molecular volumes of inorganic compounds, A., ii, 280.
the thermal decomposition of chlorine, A., ii, 823.
- Henglein, Friedrich August, G. von Rosenberg, and A. Muchlinski**, vapour pressure of solid chlorine and bromine, A., ii, 760.
- Henglein, Friedrich August.** See also *H. von Wartenberg*.
- Hengstenberg, O.** See *Karl Bornemann*.
- Henke, C. O., and Oliver W. Brown**, catalytic preparation of azobenzene and aniline, A., i, 586, 1196.
- Henke, C. O.** See also *Oliver W. Brown*.
- Henkel, Gerhard.** See *Friedrich Meyer*.
- Henley, Francis Robert.** See *Arthur Harden*.
- Henley, R. R.**, estimation of globulins in blood-serum, A., ii, 671.
- Henning, B. H.**, the lipoids of the blood in tuberculosis, A., i, 963.
- Henning, Fritz**, numerical values of the gas constants, A., ii, 191.
- Henri, Victor**, absorption spectrum of benzene vapour and the fundamental magnitudes of the benzene molecule, A., ii, 381.
the absorption and fluorescence spectra of benzene, A., ii, 679.
- Henri, Victor, and Pierre Steiner**, absorption of ultra-violet rays by naphthalene, A., i, 928.
- Henrich, Ferdinand [August Karl]**, calcium uranium autunites, A., ii, 516.
valency theories of organic chemistry, A., ii, 704.
- Henrich, Ferdinand [with G. Hiller]**, a new mineral which contains the rare earths as its main component, A., ii, 860.
- Henrich, Ferdinand [with G. Frell]**, the examination of naturally occurring gases. II. and III., A., ii, 858.
- Henriques, Valdemar, and Rich. Ege**, the dextrose concentration in the arterial blood and in the venous blood from the muscles, A., i, 191.
- Henriques de Sousa.** See *Sousa*.
- Henry, D. C.**, kinetic theory of adsorption, A., ii, 740.
- Henry, P.** See *H. Weiss*.
- Henry, Thomas Anderson, and Thomas Marvel Sharp**, mercury compounds of hydroxybenzaldehydes, T., 1055.
- Henstock, Herbert**, the solubility of phenanthrene in various organic solvents, T., 2124.
- Hepburn, Joseph Samuel**, the enzymes of the abdominal adipose tissue of the common turkey, *Meleagris gallinavo*, A., i, 196.
- Hepburn, Joseph Samuel, and E. Q. St. John**, sugar content of the hen's egg, A., i, 294.
- Hepner, B. B.** See *Israel Lifschitz*.
- Hepworth, Harry**, some recent applications of magnesium in synthetic organic chemistry, A., i, 118.
- Herberts, K.** See *K. Hugo Bauer*.

- Herbst, Heinrich**, influence of water-content on the adsorption-capacity of an active charcoal, A., ii, 80.
- Hérissey, Henri**, biochemical synthesis of α -methyl-d-mannoside, A., i, 112.
- estimation of salicylic acid in blood-serum and other fluids of the body, A., ii, 880.
- Hérissey, Henri**, and **P. Delauney**, presence in several indigenous orchids of glucosides yielding coumarin on hydrolysis, A., i, 210.
- Hermann, Paul**, derivatives of dulcin, A., i, 1151.
- Hermanns, Leo**, the nature of Ehrlich's diazo-reaction. III., A., i, 1091.
- Hermans, P.** See **Jacob Böeseken**.
- Herold, Julius**. See **Wilhelm Steinkopf**.
- Herrera, A. L.**, diffusion in porous vessels, A., ii, 427.
- Herriek, Margaret C.** See **C. J. Gamble**.
- Herrmann, E.** See **Jean Piccard**.
- Herrmann, Erika**. See **Robert Schwarz**.
- Herrmann, K.**, Dolezalek's theory of solutions, A., ii, 552.
- Herrmann, Marianne**. See **Hans Fischer**.
- Herrmann, Walter**. See **William Küster**.
- Hertz, Gustav**, excitation and ionisation potentials of neon and argon, A., ii, 783.
- Herzheimer, Herbert**, effect of primary sodium phosphate on body power, A., i, 970.
- Herz, Walter [Georg]**, the properties of organic liquids, A., ii, 23.
- atomic and molecular volumes at the absolute zero. II., A., ii, 29.
- relationship of the internal friction of organic liquids to other properties, A., ii, 30.
- refraction of light at corresponding temperatures, A., ii, 97.
- coefficient of expansion of molten salts, A., ii, 116.
- the density of caesium at the absolute zero, A., ii, 289.
- validity of the laws of the internal friction of liquids, A., ii, 352.
- relationships between molecular refraction and other properties, A., ii, 409.
- entropy of the elements and the periodic system, A., ii, 548.
- fused salts, A., ii, 739.
- the heat of evaporation, A., ii, 818.
- corresponding states, A., ii, 821.
- Herz, Walter**, and **Paul Schuftan**, physico-chemical investigation of tetrahydronaphthalene and decahydronaphthalene, A., i, 647.
- Herz, Walter**. See also **Richard Lorenz**.
- Herzberg, Kurt**, detection of traces of lactose in urine by formation of form-aldehyde, A., ii, 167.
- Herzfeld, Eugen**. See **Emil Baur**.
- Herzfeld, Karl F.**, width of the absorption bands of the rare earths, A., ii, 6.
- theory of catalysis in homogeneous gas reactions, A., ii, 136.
- application of statistics to chemical equilibria, A., ii, 269.
- Herzig, Josef**, the action of diazomethane on the ureides and uric acid, A., i, 373.
- Herzig, Josef**, and **Hans Lieb**, deamino-proteins, A., i, 386.
- Herzig, Paul**, methods for the estimation of the alkaloids, A., ii, 538.
- Herzog, Johannes**, the estimation of albumose-silver, A., ii, 798.
- Herzog, Walter**, and **J. Kreidl**, relation between ability to form resins and chemical constitution. III. A new method for producing synthetic resins, A., i, 1168.
- a supposed method for the quantitative separation of "saccharin" from *p*-sulphaminobenzoic acid, A., ii, 237.
- Herzog, Walter**. See also **Paul Friedländer**.
- Heslinga, J.**, oxidation of manganese to permanganate in alkaline solution, A., ii, 589.
- colorimetric estimation of manganese in steels, alloys, and ores, A., ii, 660.
- Hess, E.** See **Alfred Benrath**.
- Hess, Kurt**, cellulose. V. A new degradation of cellulose; conversion of cellulose into a biose anhydride, A., i, 12.
- chemistry of the manufacture of artificial silk, A., i, 922.
- Hess, Kurt**, and **Wilhelm Corleis**, 1- α -N-methylpiperidylethan-1-one, A., i, 170.
- Hess, Kurt**, and **Ernst Messmer** [with (*Fr.*) **E. Jagla**], cellulose. VII. Cellulose copper compounds, A., i, 988.
- Hess, Kurt**, and **Ottmar Wahl**, scopoline. VI. The constitutions of scopolamine and scopoline; the Hofmann degradation of scopoline, A., i, 854.
- Hess, Kurt**, and **Wilhelm Weltzien**, triphenylethynylcarbinol and its analogues, A., i, 35.
- Hess, Kurt**, and **Walter Wittelsbach**, cellulose. VI. De-polymerisation of ethyl-cellulose, A., i, 116.

- Hess, W. R.**, the function of the vitamins in the chemistry of the cell, A., i, 399, 788.
- Hesse, Erich**, [physiological] effect of cyanamide, A., i, 1093.
- Hessler, John C.**, the preparation of phenylacetylene, A., i, 442.
- Hetényi, Géza**. See *Stefan Rusznyák*.
- Hettterschij, C. W. G.** See *J. Hudig*.
- Heubach, F.** See *Paul Jacobsen*.
- Heuberger, J.** See *J. Arvid Hedvall*.
- Heubner, Wolfgang**, and *Robert Meyer-Bisch*, the free and esterified sulphuric acid in normal and in pathological body-fluids, A., i, 291.
- Heuser, Emil**, pentosans, A., i, 921.
- Heuser, Emil** [with *Maria Braden*, and *E. Kürschner*], pentosans, A., i, 113.
- Heuser, Emil**, and *H. Casseus*, estimation of the cellulose content of wood and other raw materials by the action of chlorine and carbon tetrachloride, A., ii, 664.
- Heuser, Emil**, and *Wilhelm Ruppel*, methyl ethers of xylan, A., i, 810.
- Heuser, Emil**, and *Sigurd Samuelsen*, oxidation of lignin and lignosulphonic methyl ethers, A., i, 812.
- Heuser, Emil**, *R. Schmitt*, and *L. Gunkel*, methylation of lignin, A., i, 325.
- Heuser, Emil**, and *Fritz Stöckigt*, oxycellulose, A., i, 719.
- Hevesy, Georg von**, an attempt to influence the rate of radioactive disintegration by use of penetrating radiation, A., ii, 608.
- loosening of crystal lattices, A., ii, 623.
- Hevesy, Georg von**. See also *Johannes Nicolaus Brönsted*.
- Hewitt, H. W.** See *Edmund Brydges Rudhall Pridaux*.
- Hewitt, James Arthur**, and *David Henriques de Souza*, the metabolism of carbohydrates. II. On the possible occurrence of stereochemical changes in equilibrated solutions of reducing sugars introduced into the circulation, A., i, 395.
- Hewitt, James Arthur**, and *Dorothy Beatty Steabben*, fermentation of *i*-inositol, A., i, 406.
- Hewitt, James Arthur**. See also *John William Pickering*.
- Heyde, H. C. van der**, estimation of small quantities of atropine in blood-serum, A., ii, 669.
- Heyl, Frederick W.**, the phytosterols of ragweed pollen, A., i, 1224.
- Heyl, Frederick W.**, and *Frank E. Greer*, sodium hyposulphite, A., ii, 288.
- Heymann, Paul**, influence of minute concentrations of acid and alkali on the blood-vessels and other smooth muscle, A., i, 791.
- Heyrovsky, Jaroslav**, the constitution of aluminates, A., ii, 771.
- electrolysis with drops of mercury as the electrode, A., ii, 816.
- Hibbert, Eva**. See *Edmund Knecht*.
- Hickman, Kenneth Claude Devereux**, and *Reginald Patrick Linstead*, a modified methyl-orange indicator, T., 2502.
- Hickox, Edgar Herbert Cuthbert**. See *William Davies*.
- Hicks, John F. G.**, reactions in fused salt media. I. Basic lead chromates, A., ii, 147.
- Hicks, John F. G.**, and *Wallace A. Craig*, reactions in fused salt media. II. Solvolysis, A., ii, 622.
- Hicks, William Milchinson**, spectrum lines of neutral helium, A., ii, 675.
- Hieulle, A.** See *Robert Fosse*.
- Higginbotham (Miss) Lucy**, and *Arthur Lapworth*, the formation of substituted succinic acids from esters of α -unsaturated acids, T., 49.
- Higginbotham (Miss) Lucy**, and *Arthur Lapworth* [with *Charles Simpson*], formation of γ -alkylidene derivatives from ethylidenemalononic ester, T., 2823.
- Hijikata, Yoshizumi**, the cleavage products of the crystalline lens, A., i, 489.
- do the amino-acids occur in cow's milk? A., i, 494.
- influence of putrefaction products on cellular metabolism. II. The influence of phenylacetic and phenylpropionic acids on the distribution of nitrogen in the urine, A., i, 495.
- Hilcken, Valentin**. See *Otto Dimroth*.
- Hild, Wilhelm**, rapid method for the estimation of chromium in nickel-chromium steel, A., i, 660.
- Hildebrand, Joel H.**, and *Clarence A. Jenks*, solubility. VII. Solubility relations of rhombic sulphur, A., ii, 141.
- Hildebrand, Joel H.** See also *Edna R. Bishop*.
- Hilditch, Thomas Percy**. See *Edward Frankland Armstrong*.
- Hill, A. Elizabeth**, and *A. K. Balls*, a sulphonated naphthylarsinic acid, A., i, 1080.
- Hill, Arthur E.**, the system silver perchlorate-water-benzene, A., ii, 555.
- Hill, Arthur E.**, and *Thomas M. Smith*, hydrated oxalic acid as an oxidimetric standard, A., ii, 388.

- Hill, Arthur Joseph**, and **Erwin B. Kelsey**, thiocyanates and thiocarbimides. XV. The nature of the intramolecular rearrangement of thiocarbimidoacetanilides, A., i, 1141.
- Hill, Arthur Joseph**. See also *Treat Baldwin Johnson*.
- Hill, Archibald Vivian**, the combinations of hæmoglobin with oxygen and carbon monoxide, and the effects of acid and carbon dioxide, A., i, 193.
- the interactions of oxygen, acid, and carbon dioxide in blood, A., i, 696.
- Hill, Archibald Vivian**. See also *Joseph Basoreff*, and *W. E. L. Brown*.
- Hill, Elsie**, and *W. R. Bloor*, fat excretion, A., i, 968.
- Hill, Henry Rowland**. See *Frederick Daniel Chattaway*.
- Hiller, Alma**, and *Donald D. van Slyke*, protein precipitants, A., i, 1074.
- Hiller, Alma**. See also *Donald D. van Slyke*.
- Hillar, G.** See *Ferdinand Henrich*.
- Hilliger, E.** See *Karl von Auwers*.
- Hilton, O.** See *E. V. Lynn*.
- Himstedt, F.**, and *Irma Wertheimer*, influence of pressure on the refractive index of some organic liquids, A., ii, 725.
- Hinard, Gustav**, and *Robert Fillon*, the chemical composition of starfish, A., i, 87.
- Hinchey, Victor M.** See *Thomas Dillon*.
- Hines, Paul R.** See *William Lloyd Evans*.
- Hinkel, Leonard Eric**, and *William Dudley Williams*, action of the chlorides of phosphorus on chlorodimethyldihydroresorcinol, T., 2493.
- Hinshelwood, Cyril Norman**, and *Edmund John Bowen*, velocity of chemical change in solid substances, A., ii, 628.
- Hinshelwood, Cyril Norman**, and *Harold Hartley*, probability of spontaneous crystallisation of supercooled liquids, A., ii, 200.
- Hinshelwood, Cyril Norman**, *Harold Hartley*, and *B. Topley*, influence of temperature on two alternative modes of decomposition of formic acid, A., ii, 274.
- Hinshelwood, Cyril Norman**. See also *Harold Calvert Tingey*.
- Histikka, S. F.**, sulphite liquor lactones, A., i, 347.
- Hirabayashi, N.** See *Leonor Michaelis*.
- Hirose, Mituzie**. See *Tokutaro Sakao*.
- Hirose, W.** See *Georg Joachimoglu*.
- Hirsch, Julius**, the oxidative degradation of dextrose in the animal body, A., i, 87.
- carbolic acid. V. The binding of carbon to carbon biosynthetically in the aliphatic series, A., i, 978.
- Hirsch, Julius**. See also *Carl Neuberg*.
- Hirsch, Max**, detection of traces of osmium by means of potassium thiocyanate A., ii, 459.
- Hirsch, Paul**, physico-chemical studies on biological reactions. I., A., i, 785.
- Hirsch, Paul**, and *Rudolf Kunze*, physico-chemical studies on biological reactions. II. Spectro-chemical investigations on amino-acids and polypeptides, A., i, 781.
- Hirschfelder, Arthur D.** See *Merrill C. Hart*.
- Hirschko, K.** See *F. Halla*.
- Hirsch-Pogany, Margit**, is the heat-coagulation of protein a hydrolysis? A., i, 596.
- Hirst, C. T.**, and *Joseph E. Greaves*, factors influencing the estimation of sulphates in soil, A., ii, 521.
- Hirst, Edmund Langley**, and *Alexander Killen Macbeth*, the labile nature of the halogen atom in organic compounds. II. Action of hydrazine on nitrogen-halogen compounds and on bromomalonate esters, T., 904.
- the labile nature of the halogen atom in organic compounds. V. The action of hydrazine on the halogen derivatives of some esters and substituted cyclohexanes, T., 2169.
- Hirst, Edmund Langley**. See also *Ian Armstrong Black*, *Albert Carruthers*, and *James Colquhoun Irvine*.
- Hitchcock, David I.**, the colloidal behaviour of edestin, A., i, 698.
- the combination of gelatin with hydrochloric acid, A., i, 882.
- the colloidal behaviour of serum globulin, A., i, 1074.
- Hitchcock, David I.** See also *John Maurice Nelson*.
- Hixon, Ralph M.**, the effect of the reaction of a nutritive solution on germination and the first stages of plant growth, A., i, 1221.
- complex ion formation in hydrochloric acid, A., ii, 815.
- Hjalmar, Elis**, Röntgen spectra, A., ii, 180.
- Hladký, Jan**, the use of Hanus's method for the determination of the iodine figure of mineral oils, A., ii, 722.
- Hochmuth, Otto**. See *Gustav Heller*.
- Hodges, James H.** See *Theodore William Richards*.

- Höhn, Fritz.** See *Ignaz Bloch*.
- Hoel, A. B.** See *Arthur Wesley Browne*.
- Hölzl, Franz.** See *Robert Müller*.
- Hönig, Reinhold.** See *Robert Müller*.
- Hönigsberger, F.** See *Paul Jacobsen*.
- Hönigschmid, Otto** and **Lothar Brökenbach**, revision of the atomic weight of glucinum; analysis of glucinum chloride, A., ii, 214.
- revision of the atomic weight of boron; analysis of boron trichloride, A., ii, 641.
- Hoenlinger, H.** See *M. Richter-Quittner*.
- Höppler, E. F.**, quantitative analysis by measurement of the degree of supersaturation, A., ii, 863.
- Hoffert, W. H.**, estimation of phenol in mixtures of tar acids, A., ii, 878.
- Hoffman, Alfred**, the action of hydrogen phosphide on formaldehyde, A., i, 8.
- Hoffman, Waller Fred**, and **Ross Aiken Gortner**, sulphur in proteins. I. The effect of acid hydrolysis on cystine, A., i, 429.
- Hoffman, Walter Fred.** See also *Ross Aiken Gortner*.
- Hoffmann, Alex.** See *Géza Zemplén*.
- Hoffmann, Fritz**, the oxygen-absorption and concentration of pyrogallol solutions used in gas analysis, A., ii, 582.
- Hoffmann, G.**, experimental decision of the question of the radioactivity of all elements. II., A., ii, 184.
- Hoffmann-La Roche & Co., F.**, preparation of 3:7-dimethyl-1-allylxanthine, A., i, 585.
- preparation of allylarsinic acid, A., i, 818, 926.
- preparation of isopropylallylbarbituric acid, A., i, 872.
- Hofmann, Josef.** See *Hans Lecher*.
- Hofmann, Karl Andreas**, the oxygen-hydrogen catalysis by the platinum metals and the contact potentials in presence of aqueous electrolytes, A., ii, 276.
- mode of action of platinum in oxygen-hydrogen catalysis and the application of titanium sulphuric acid for the control of the course of the change, A., ii, 490.
- Hofmann, Karl Andreas**, and **Erich Will**, formation of acetylene and ammonia during incomplete combustion, A., i, 1107.
- Hogness, Thorfin K.**, surface tensions and densities of liquid mercury, cadmium, zinc, lead, tin, and bismuth, A., ii, 29.
- Hoher, E.** See *J. V. Dubsy*.
- Hohl, Heinr.** See *Robert Kremann*.
- Hohorst, Georg.** See *Wilhelm Biltz*.
- Holborn, Ludwig**, and **J. Otto**, the isotherms of nitrogen, oxygen, and helium, A., ii, 737.
- Holde, David**, determination of the iodine value of aliphatic and aromatic unsaturated compounds, A., ii, 665.
- Holde, David**, [with *P. Werner, Ida Tacke*, and *C. Wilke*], the iodine values of aliphatic and aromatic unsaturated compounds, A., ii, 723.
- Holde, David**, and **K. Schmidt**, brassidic anhydride, and anhydridisation by means of carbonyl chloride, A., i, 982.
- Holde, David**, and **Ida Tacke**, the electrical conductivity of anhydrides of the higher fatty acids, A., ii, 109.
- Holde, David, P. Werner, Ida Tacke**, and **C. Wilke**, determination of the iodine value of aliphatic and aromatic unsaturated compounds, A., ii, 533.
- Holde, David**, and **C. Wilke**, erucic acid and its anhydride. I, II., and III., A., i, 317, 519, 713.
- Holden, Edward F.**, ceruleofibrite, a new mineral, A., ii, 516.
- constitution of thaumasite, A., ii, 860.
- Holleman, Arnold Frederik**, monochlorotrinobenzenes, A., i, 997.
- Hollenberg, M. S.** See *Alexander Thomas Cameron*.
- Hollins, Cecil**, the mechanism of the Fischer indole synthesis, A., i, 863.
- Holluta, Josef**, mechanism of the reduction of permanganate and its physico-chemical basis. I. Reaction between [potassium] permanganate and formic acid in slightly acid solution, A., ii, 448.
- mechanism of the reduction of permanganate and its physico-chemical foundation. III. Reaction between manganate and formic acid, A., ii, 700.
- mechanism of the reduction of permanganate and its physico-chemical basis. IV. Reduction of permanganate by formate in alkaline solutions, A., ii, 771.
- Holluta, Josef**, and **Nikolaus Weiser**, mechanism of the reduction of permanganate and its physico-chemical basis. II. Reaction of permanganate and formic acid in neutral solution, A., ii, 628.
- Holm, Kurt.** See *A. Bornstein*.
- Holmberg, Bror**, stereochemical studies. VI. Stereoisomeric trithiocarbolactonic acids, A., i, 1113.
- Holmes, Edward O., jun.**, photochemical activity of the triphenylmethanesulphonic acids, A., ii, 465.

- Holmes, Edward O., jun., and Walter A. Patrick**, action of ultra-violet light on gels, A., ii, 338.
- Holmes, Harry N., and Don H. Cameron**, cellulose nitrate as an emulsifying agent, A., ii, 268.
- chromatic emulsions, A., ii, 269.
- Holmsen, J.** See **S. Schmidt-Nielsen**.
- Holmyard, Eric John**, Arabic chemistry, A., ii, 636.
- Holt, Harold S.** See **James Bryant Conant**.
- Holtmark, J.**, the characteristic Rontgen radiation from carbon and boron, A., ii, 543.
- Holtz, Friedrich**, Pregl's microanalysis, A., ii, 521.
- Holwerda, B. I.**, influence of lactic acid on lactic acid fermentation, A., i, 611.
- Holzer, Paul, and Heinz Mehner**, estimation of bilirubin in blood, A., ii, 799.
- Homolka, B.**, croconic acid and leuconic acid, A., i, 630.
- Hompe, Louise.** See **H. S. Forbes**.
- Honda, Kôtarô**, the constitutional diagram of the iron-carbon system based on recent investigations, A., ii, 711.
- Bohr's model of the hydrogen molecules and their magnetic susceptibility, A., ii, 838.
- Honold, Ernst.** See **Emil Fromm**.
- Hooft, M.** See **W. D. Treadwell**.
- Hooker, Marian O.** See **Martin Henry Fischer**.
- Hoop, L. de**, new observations on the constitution of the carbohydrates, A., i, 434.
- Hoover, Ora L.** See **William Lloyd Evans**.
- Hope, Edward**, ethyl α -cyano- β -methylglutaconate and its methyl homologues, T., 2216.
- Hope, Edward, and George Clifford Riley**, chlorination of benzoyl chloride. I, T., 2510.
- Hope, Edward, and Wilfrid Sheldon**, the addition of hydrogen cyanide to derivatives of glutacetic acid. I. The addition of hydrogen cyanide to ethyl α -cyano- β -methylglutaconate and its homologues, T., 2223.
- Hopkins, B. Smith, and F. H. Driggs**, rare earths. III. Atomic weight of lanthanum, A., ii, 770.
- Hopkins, B. Smith.** See also **C. C. Kloss, and L. F. Yntema**.
- Hopkins, Frederick Gowland.** See **E. J. Morgan**.
- Hopper, F. L.** See **Paul W. Merrill**.
- Hopper, Isaac Vance.** See **Forsyth James Wilson**.
- Horiuchi, Riki**, formation of triphenylpararosaniline hydrochloride from diphenylamine and chloral ammonias, A., i, 1067.
- Horlacher, E.** See **Paul Karrer**.
- Horn, Trude.** See **Heinrich Schroeder**.
- Horn, William John**, pyrimidines. XCI. Alkylation of 2-thiopyrimidines, A., i, 374.
- Hornemann, Curt**, the action of pilocarpine on the glycogen content of organs, A., i, 296.
- Horrisberger, Walter.** See **Leon Asher**.
- Horrnann, Paul, and Friedrich Bischof**, picrotoxin. XII. Picrotin ketone, $C_{14}H_{16}O_8$, A., i, 161.
- Horst, (Mlle) H. van der.** See **Jean Timmermans**.
- Horton, Frank, and Ann Catherine Davies**, spectroscopic investigation of the ionisation of argon by electron collisions, A., ii, 811.
- Hostetter, J. Clyde.** See **John B. Cain**.
- Houdremont, E.** See **Theodor Dieckmann**.
- Hengen, Haakon.** See **Heinrich Goldschmidt**.
- Houghton, J. E.** See **W. A. Bloedorn**.
- Houstoun, Robert Alexander**, the quantum mechanism in the atom, A., ii, 633.
- Howard, C. P.** See **Robert B. Gibson**.
- Howard, H. A.** See **P. J. Cammidge**.
- Howard, J. W.**, the enzyme hydrolysis of benzyl succinate, A., i, 960.
- Howard & Sons, Ltd., John William Blagden, and Maximilian Wierenstein**, preparation of amino-derivatives of hydrogenated cinchona alkaloids and their derivatives, A., i, 853.
- Howe, Paul E.**, an effect of the ingestion of colostrum on the composition of the blood of new-born calves, A., i, 80.
- relation between age and the concentrations of protein fractions in the blood of the calf and cow, A., i, 1084.
- the estimation of proteins in blood—a micro-method, A., ii, 171.
- the use of sodium sulphate as the globulin precipitant in the estimation of proteins in blood, A., ii, 172.
- the differential precipitation of the proteins of colostrum and a method for the estimation of the proteins in colostrum, A., ii, 670.
- Howell, Owen Rhys.** See **Alexander Findlay**.

- Hewes, Horace L.**, the spectra structure of the luminescence excited by the hydrogen flame, A., ii, 385.
- Hewes, Horace L.** See also **Edward L. Nichols**.
- Hoyt, Lester F.**, and **H. V. Pemberton**, estimation of glycerol in the presence of sugars, A., ii, 321.
- Hristie, P.** See **W. D. Treadwell**.
- Hubbard, Roger S.**, estimation of β -hydroxybutyric acid; estimation of acetone substances in the urine; estimation of acetone substances in the blood, A., ii, 234.
- Hubbard, Roger S.**, and **Samuel T. Nicholson, jun.**, acetonuria of diabetes, A., i, 969.
- Hubbard, Roger S.**, and **Floyd R. Wright**, blood acetone substances after the injection of small amounts of adrenaline chloride, A., i, 288.
- acetonuria produced by diets containing large amounts of fat, A., i, 496.
- Hubbuck, W.** See **A. Koenig**.
- Huber, J.**, estimation of manganese as sulphate, A., ii, 398.
- Huber, J.** See also **Alexander Guthrie**.
- Huber, Karl Jakob**, the excretion by the gastric mucous membrane and the salivary glands of alkaloids administered subcutaneously, A., i, 1214.
- Huber, L.** See **Paul Jacobsen**.
- Hubert, P.** See **Paul Ruggli**.
- Huehet, P.** See **P. Brodin**.
- Huebner, Julius**, and **F. Kaye**, effect of water and certain organic salts on celluloses, A., i, 435.
- Huebner, Julius**, and **J. N. Sinha**, action of iodine on celluloses, silk, and wool, A., i, 434.
- Hübner, Julius**, the graphical representation of the composition of chemical compounds, A., ii, 156.
- Hudig, J.**, and **C. W. G. Hetterschij**, Comber's reaction for acidity of soils, A., i, 1104.
- Hudson, Donald Pryce.** See **Edward Charles Cyril Baly**.
- Hückel, E.**, scattering of Röntgen rays by anisotropic liquids, A., ii, 14.
- Hückel, Walter**, the meaning of the atomic constants of heat of combustion and molecular refraction, A., ii, 195.
- heats of combustion and energy of dissociation, A., ii, 818.
- Hückel, Walter.** See also **Adolf Windaus**.
- Hültzen, K.** See **Karl von Auwers**.
- Huerre, R.**, the rôle played by the various elements of the wood of *Juniperus oxycedrus* in the formation of oil of cade, A., i, 505.
- Hüttig, Gustav F.**, valency. XVI. The ammonia compounds of calcium haloids, A., ii, 849.
- Hüttig, Gustav F.**, and **Bruno Kurre**, the hydrates of tungsten trioxide, A., ii, 773.
- Hüttig, Gustav F.**, and **Edith von Schroeder**, the hydrates of uranium tetroxide and uranium trioxide, A., ii, 510.
- Hüttig, Gustav F.** See also **Wilhelm Biltz**.
- Hugel, G.** See **Martin Battagay**.
- Huggins, Maurice L.**, conjugation and the structure of benzene, A., i, 928.
- the structure of benzene, A., i, 997.
- atomic structure, A., ii, 632.
- atomic radii. I., A., ii, 634.
- the crystal structure of quartz, A., ii, 641.
- the crystal structures of aragonite (CaCO_3) and related minerals, A., ii, 643.
- the crystal structures of marcasite (FeS_2), arsenopyrite (FeAsS), and loellingite (FeAs_2), A., ii, 651.
- electronic structures of crystals, A., ii, 744.
- electronic structures of atoms, A., ii, 838.
- Hughes, A. Ll.**, characteristic X-rays from boron and carbon, A., ii, 184.
- Hughes, Edward J.** See **Frank X. Moerk**.
- Hughes, W.**, a possible reconciliation of the atomic models of Bohr and of Lewis and Langmuir, A., ii, 632.
- Hughes, William Earl**, electro-deposition of lead from Mathers's perchlorate bath. I. Structure of the deposit, A., ii, 446.
- Hulbert, E. O.**, the broadening of the Balmer lines of hydrogen with pressure, A., ii, 801.
- Hulett, George Augustus.** See **Guy B. Taylor**.
- Hull, Albert W.**, X-ray crystal analysis of metals, A., ii, 624.
- Hull, Albert W.**, and **Wheeler P. Davey**, graphical determination of hexagonal and tetragonal crystal structures from X-ray data, A., ii, 624.
- Hulot, Pierre**, metallurgy of tellurium by the wet way, A., ii, 142.
- Hulthén, Erik**, and **Ernst Bengtsson**, the band spectra of cadmium, A., ii, 676.
- Hulton, Henry Francis Everard.** See **Julian Levitt Baker**.
- Hunt, Seth Bliss**, preparation of ketones from secondary alcohols, A., i, 810.

- Hunter, George**, estimation of carnosine in muscle extract, A., ii, 328, 885.
 Knoop's test for histidine, A., ii, 885.
Hupper, M. See *E. M. Black*.
Hurd, Charles D. See *Lauder William Jones*.
Hurst, Eric. See *John Bead*.
Hussey, Robert E. See *Brainerd Mears*.
Husson, A. See *Gustave Vavon*.
Hutton, Robert. See *John Arnold Cranston*.

I.

- Iatrides, D.** See *Ernst Winterstein*.
Ide, Toshio, the tryptophan content of important foodstuffs, A., i, 414.
Ikeda, Kikunae, and *Shintaro Kodama*, manufacture of chloro-acids, A., i, 219.
Imbert, Georges. See *British Dyestuffs Corporation, Ltd.*
Imker, Albert. See *Rudolf Schenck*.
Ingersoll, Harry R. See *Leon A. Congdon*.
Ingold, Christopher Kelk, the structure of the benzene nucleus. I. Intra-nuclear tautomerism, T., 1133.
 the structure of the benzene nucleus. II. Synthetic formation of the bridged modification of the nucleus, T., 1143.
 the form of the vapour pressure curve at high temperature. I. The curve for lead, T., 2419.
 the conditions underlying the formation of unsaturated and cyclic compounds from halogenated open-chain derivatives. V. Products derived from α -halogenated β -methylglutaric acids, T., 2676.
Ingold, Christopher Kelk, and *Lewis Charles Nickolls*, experiments on the synthesis of the polyacetic acids of methane. VII. *iso*-Butylene- $\alpha\gamma\gamma$ -tricarboxylic acid and methanetetracetic acid, T., 1638.
Ingold, Christopher Kelk, and *Edward Arthur Perren*, experiments on the synthesis of the polyacetic acids of methane. VI. Methanetricarboxylic acid and its unstable esters, T., 1414.
Ingold, Christopher Kelk, *Edward Arthur Perren*, and *Jocelyn Field Thorpe*, ring-chain tautomerism. III. The occurrence of tautomerism of the three-carbon (glutaconic) type between a homocyclic compound and its unsaturated open-chain isomeride, T., 1765.
Ingold, Christopher Kelk, and *Henry Alfred Piggott*, the mobility of symmetrical triad systems. I. The conditions relating to systems terminated by phenyl groups, T., 2381.
 the additive formation of four-membered rings. I. The synthesis and division of derivatives from 1:3-dimethindiazidine, T., 2793.
Ingold, Christopher Kelk, *Shinichi Sato*, and *Jocelyn Field Thorpe*, the influence of substituents on the formation and stability of heterocyclic compounds. I. Hydantoins, T., 1177.
Ingold, Christopher Kelk, and *(Miss) Edith Hilda Usherwood*, the specific heats of gases with special reference to hydrogen, T., 2286.
Ingold, Christopher Kelk, and *Daniel Wilson*, the reversibility of the reaction between nitrogen, carbon, and sodium carbonate, T., 2278.
Ingold, Christopher Kelk. See also *Juan Pedigo Charles Chandrasena*, and *Ernest Harold Farmer*.
Inichov, G. S., the chemical action of rennin, A., i, 960.
Inoue, Harushige, electrolytic reduction of some carboxylic acids, A., i, 36.
Ionescu, Al., chemistry of diabetic glycosuria, A., i, 198.
 barium chloride poisoning, A., i, 199.
 the defecation of blood for the estimation of carbamide, A., ii, 596.
Ionescu, Al., and *C. Pop*, new method for the detection of bile pigments, A., ii, 671.
Ionescu, Al. See also *Stefan Minovici*.
Iredale, Thomas, the rôle of protective colloids in catalysis. II., T., 1536.
Irion, Clarence E. See *Gerald L. Wendt*.
Irvine, James Colquhoun, and *John Campbell Earl*, mutarotation and pseudo-mutarotation of glucosamine and its derivatives, T., 2370.
 salicylidene derivatives of *d*-glucosamine, T., 2376.
Irvine, James Colquhoun, and *Edmund Langley Hirst*, 2:3:6-trimethyl glucose, T., 1213.
 the constitution of polysaccharides. V. The yield of glucose from cotton cellulose, T., 1585.
Irvine, James Colquhoun, and *Jocelyn Patterson*, the constitution of acetone derivatives of glucose and fructose, T., 2146.
 1:3:4:6-tetramethyl fructose, T., 2696.
Irvine, James Colquhoun, *(Miss) Elsie Stewart Steele*, and *(Miss) Mary Isobel Shannon*, the constitution of polysaccharides. IV. Inulin, T., 1060.

Irwin, Marian, sensory stimulation by alcohols and chlorohydrins, A., i, 899.

Isaac, Salo, and **Erich Adler**, the steric transformation of the hexoses through the agency of organs and cells (the so-called stereokinases), A., i, 297.

Isaacs, M. L., colorimetric estimation of hydrogen peroxide, A., ii, 715.

colorimetric estimation of blood chlorides, A., ii, 716.

Isgarischev, N., and **Sophie Berkmann**, overvoltage on electrodes and its connexion with the hydration of ions, A., ii, 253.

influence of colloids on overvoltage, A., ii, 254.

Ishida, Yoshitoyo. See **Yasuhiko Asahina**.

Ishihara, F., tetrodon poison and some of its chemical characteristics, A., i, 85.

Ishino, Matakichi, evidence of the existence of isotopes of chlorine, A., ii, 760.

Islip, Harold Thomas. See **Oswald Digby Roberts**.

Itagaki, Takeyoshi. See **Yoshiharu Murayama**.

Ivanitskaja, Agness. See **Nikolai Schilov**.

Ivers, Otto. See **Karl Freudenberg**.

Iversen, Poul. See **Johannes Book**.

Ives, Herbert E., proposed standard method of colorimetry, A., ii, 221.

Iwamoto, Yoshitora, properties and composition of tohaku oil, A., i, 98.

Iwanov, Nicolai N., the change undergone by nitrogenous substances in the final phases of yeast autolysis, A., i, 202.

the influence of fermentation products on the decomposition of proteins in yeast, A., i, 202.

protein decomposition in yeast during fermentation, A., i, 202.

Iyer, K. R. Krishna. See **Kishori Lal Mondgill**.

Isaguirre, Ramon de, surface tension of aqueous solutions of night-blue, A., ii, 262.

Isaguirre, Ramon de. See also **Wolfgang Ostwald**.

J.

Jablonskyński, K., and **F. J. Wiśniewski**, equilibrium law of electrolytes, A., ii, 190.

Jackman, Douglas Norman, and (*Miss*) **Agnes Browne**, the 25°-isotherms of the systems magnesium nitrate-sodium nitrate-water and magnesium sulphate-magnesium nitrate-water, T., 694.

Jackman, Douglas Norman. See also **William Edward Garner**.

Jackson, D. H. See **Francis P. Venable**.

Jackson, Henry, jun., and **Walter W. Palmer**, a modification of Folin's colorimetric method for the estimation of uric acid, A., ii, 328.

the estimation of uric acid, A., ii, 795.

Jackson, J. W. See **Alexander O. Gettler**.

Jackson, L. C., dielectric constants of some esters at low temperatures, A., ii, 252.

Jackson, Richard Fay, and **Clara G. Silsbee**, solubility of dextrose in water, A., i, 986.

Jackson, S. D. See **William Ridgely Orndorff**.

Jacobs, Walter Abraham, and **Michael Heidelberger**, aromatic arsenic compounds. IX. Diazoamino-compounds of p-aminophenylarsinic acid and its derivatives, A., i, 73.

aromatic arsenic compounds. X. Azodyes derived from p-aminophenylarsinic acid, A., i, 74.

syntheses in the cinchona series. VII. 5:8-Diaminodihydroquinine and 5:8-diamino-6-methoxyquinoline and their conversion into the corresponding aminohydroxy- and dihydroxy-bases, A., i, 671.

syntheses in the cinchona series. VIII. The hydrogenation of dihydrocinchonine, cinchonine, and dihydroquinine, A., i, 672.

Jacobs, Walter Abraham. See also **Michael Heidelberger**.

Jacobsen, Helene. See **Karl von Auwers**.

Jacobsen, Paul, summary of results on the isomeric changes of hydrazo-compounds, and considerations relating to the possibilities of their explanation, A., i, 596.

Jacobsen, Paul, [with **David Runckman Boyd**, **M. Freund**, **H. L. Fuld**, **F. Heubach**, **F. Hönigsberger**, **L. Huber**, **M. Jaenicke**, **H. Jost**, **G. Lockemann**, **A. Loeb**, **J. Pelzer**, **St. Pinkus**, **W. Sachs**, **P. Schmidt**, **H. Tigges**, and **K. Zaar**], the isomeric changes of hydrazo-compounds, A., i, 589.

Jacobsohn, Isadore M. See **Morris S. Kharasch**.

Jacobson, J., catalytic action of benzyl alcohol, A., i, 652.

- Jacoby, Martin**, artificial zymogens. II., A., i, 480.
formaldehyde as an intermediate step between the real assimilation and the formation of carbohydrate in the plant. II., A., i, 502.
- Jacoby, Martin**, and **Käle Frankenthal**, the importance of the amino-acids of hæmoglobin for the cultivation of the influenza bacilli, A., i, 302.
- Jacoby, Martin**, and **Tomihide Shimizu**, artificial zymogens. III., A., i, 481.
artificial zymogens. IV. Inactivation and reactivation of takadiastase, A., i, 481.
adsorption of ferments and zymogens. I. and II., A., i, 481.
- Jacquot, Raymond**. See **René Wurmser**.
- Jadin, F.**, and **A. Astruc**, relation between manganese content and proportion of ash in young and old leaves, A., i, 1098.
- Jaack, Wolfgang**. See **Franz Flechter**.
- Jaeger, Frans Maurits**, crystallography of derivatives of benzophenone, A., i, 353.
- Jaeger, Frans Maurits**, and **G. Berger**, the action of ultra-violet light on aqueous solutions of some organic acids and their salts, A., i, 316.
- Jaeger, Frans Maurits**, and **H. C. Germs**, binary systems of the sulphates, chromates, molybdates, and tungstates of lead, A., ii, 65.
- Jäger, Gustav**, theory of Brownian movement, A., ii, 30.
- Jaeger, Paul**. See **Wilhelm Steinkopf**.
- Jaeger, Siegfried**. See **Richard Anschütz**.
- Jäggi, A.** See **Hans Rupe**.
- Jänecke, Ernst**, Le Chatelier's "new" geometrical representation, A., ii, 427.
- Jaenicke, M.** See **Paul Jacobsen**.
- Jaffeux, (Mme) Pierre**. See **Marcel Delépine**.
- Jagla, (Fr.) E.** See **Kurt Hess**.
- Jahn, R.** See **Amé Pietet**.
- Jaitschnikov, Iv. S.**, hydrolysis of glycylglycine by hydrochloric acid, A., i, 1124.
identification of alanine by crystallochemical analysis, A., ii, 884.
- Jakób, W. F.**, researches on the photochemically sensitive compounds of molybdic and formic acids, A., i, 712.
- Jakubowicz, Heinrich**, the influence of temperature and constitution on the decomposition velocity of substituted malonic acid, A., ii, 361.
- Jakubowicz, Heinrich**. See also **August L. Bernoulli**.
- Jaloustre, Léon**. See **Laborde and Pierre Lemay**.
- James, Charles**. See **Paul H. M.-P. Brinton**, and **H. C. Fogg**.
- James, R. W.** See **William Lawrence Bragg**.
- Jameson, Henry Lyster, Jack Cecil Drummond**, and **Katharine Hope Coward**, synthesis of vitamin-A by a marine diatom (*Nitzschia closterium*, W. Sm.) growing in pure culture, A., i, 1099.
- Jamieson, George Samuel**. See **Walter F. Baughman**.
- Jancke, Willi**. See **K. Becker**.
- Jander, Gerhart**, chemical analysis with membrane filters. III. Their application to volumetric analysis, A., ii, 457.
- Jander, Gerhart**, and **Erwin Wendehorst**, the quantitative estimation and separation of aluminium, A., ii, 529.
- Jander, W.** See **Gustav Tammann**.
- Jandrasehitzsch, Franz**. See **Mona Adolf**.
- Janet, (Mlle) Marthe Paul**, the estimation of urea in urine by the hypobromite method with complete yield, A., ii, 794.
- Janet, (Mlle) Marthe Paul**. See also **W. Mestrezat**.
- Jansch, H.**, I. Estimation of methyl alcohol in remains for forensic purposes. II. Occurrence of methyl alcohol in the human body, A., ii, 232.
- Jánský, V.** See **Julius Stoklassa**.
- Jantschak, Marie**. See **Nikolai Schilov**.
- Jarisch, Adolf**, influence of temperature on hypotonic hæmolysis, A., i, 894.
soap and serum, A., i, 1087.
- Jassoy, H.** See **Adolf Sieglitz**.
- Jeffery, F. H.**, electrolysis of aqueous solutions of alkali nitrites with a lead anode, and an electrometric determination of the constitution of the complex ion formed, A., ii, 611.
- Jellinek, Karl**, and **Alois Diethelm**, generator gas equilibrium at high pressures, A., ii, 833.
- Jellinek, Karl**, and **H. Ens**, some new volumetric methods (estimation of sulphate, lead, acids, and ammonia), A., ii, 864.
- Jellinek, Paul**. See **Sigmund Fränkel**.
- Jemma, Giuseppe**, indican reactions for the detection of urine in stains, A., ii, 460.
- Jenze, Wilhelm**, the chemical and electrical behaviour of some series of alloys, A., ii, 64.
- Jenkins, William Job**. See **James William McBain**.
- Jenks, Clarence A.** See **Joel H. Hildebrand**.

- Jennings, J. M. See T. H. Rogers.
- Jenny, Ernst. See Franz Fischer.
- Jensen, H. See Adolf Windaus.
- Jeppsson, Carl Alex. See Erik Matteo Prochet Widmark.
- Jess, A., the content of the proteins of the lens in histidine, arginine, and lysine, A., i, 1199.
- Jirsa, Franz, the effect of telluric acid on the potential of a silver anode, A., ii, 418.
- Jirsa, Franz, and Ot. Buryánek, the anode oxidation of gold. I. Anode oxidation of gold in sulphuric acid, A., ii, 718.
- Joachimoglu, Georg, the action of some derivatives of chloroform with special reference to Traube's theory of the action of narcotics of the aliphatic series, A., i, 199.
- comparative experiments on the antiseptic action of some chloro-derivatives of methane, ethane, and ethylene, A., i, 304.
- the action of sublimate, phenol, and quinine on yeast, A., i, 903.
- the elective action of tellurium salts on bacteria of the colon-typhoid group, A., i, 1095.
- a lecture experiment for demonstrating the dependence of the antiseptic action of mercury compounds on the degree of ionisation, A., ii, 211.
- Joachimoglu, Georg, and W. Hirose, pharmacology of selenium and tellurium. III. The action of their acids on the organs of the circulation, A., i, 396.
- pharmacology of selenium and tellurium. II. Action of their acids on diphtheria bacilli, A., i, 406.
- Job, André, and Henri Guinot, preparation of dimethyl- and diphenylarsinecarboxylic acids, A., i, 75.
- Job, André, and René Reich, attempt at a systematic extension of the preparation of organometallic compounds; application to ferrous ethyl iodide, A., i, 645.
- Job, Paul, electrometric study of the hydrolysis of some complex cobalt-ammines under the action of barium hydroxide, A., ii, 301.
- the hydrolysis of the roseocobaltic salts, A., ii, 382.
- Jodlbauer, Alb., and F. Haffner, the relation between fluorescent substances which act in the dark and their photo-dynamic activity on cells, A., i, 94.
- Joël, Ernst, viscometric and stalagmometric measurements of urine, A., i, 198.
- Joffe, Jacob S. See Jacob Goodale Lipman, and Selman A. Waksman.
- Johannessen, A., qualitative and quantitative demonstration of blood in urine, A., ii, 724.
- Johannsen, Adolf. See Fritz Paneth.
- Johannsen, Otto. See Alois Zinke.
- Johansson, Hjalmar, and Sidney M. Hagman, β -halogen-substituted fatty acids and β -lactones, A., i, 425.
- John, Carl. See Johannes Gadamer.
- Johns, Carl Oscar, Lewis H. Chernoff, and Arno Vishover, saponin from *Agave lechuguilla*, Torrey, A., i, 797.
- Johns, Carl Oscar, and Charles E. F. Gersdorff, the proteins of the seed of the tomato, *Solanum esculentum*, A., i, 800.
- Johns, Carl Oscar. See also David Breese Jones.
- Johnson, Harlan W., relation of hydrogen-ion concentration in soils to their "lime requirement," A., i, 708.
- Johnson, John R., and Roger Adams, 4-carboxy-2-phenylquinoline-6-arsinic acid, A., i, 187.
- Johnson, Treat Baldwin, and Lawrence W. Bass, the spontaneous decomposition of imino-ethers, A., i, 736.
- Johnson, Treat Baldwin, and Oskar Baudisch, pyrimidines. XCII. New methods of identifying the pyrimidine, thymine, A., ii, 328.
- Johnson, Treat Baldwin, Arthur Joseph Hill, and Francis H. Case, the action of diazomethane on uracil, A., i, 471.
- Johnson, Treat Baldwin. See also Oskar Baudisch, and George W. Fucher.
- Johnston, R. A. A., and H. V. Ellsworth, the Annahme meteorite, A., ii, 306.
- Jolibois, Pierre, and Robert Bomuet, relationships between the different oxides of uranium, A., ii, 801.
- precipitation from uranyl nitrate by means of sodium hydroxide; radio-activity of the precipitate, A., ii, 575.
- Joly, John, haloes and earth-history; a new radioactive element, A., ii, 651.
- Jonas, K. G., natural and artificial humic acids, A., i, 326.
- Jones, A. J., the arsenic content of some of the marine algae, A., i, 905.
- Jones, David Breese, A. J. Finks, and Charles E. F. Gersdorff, proteins of the adzuki bean, *Phaseolus angularis*, A., i, 504.

- Jones, David Breese, Charles E. F. Gersdorff, Carl Oscar Johns, and A. J. Flaks**, proteins of the lima bean, *Phaseolus lunatus*, A., i, 1101.
- Jones, David Breese, and Carl Oscar Johns**, estimation of the monoamino-acids in the hydrolytic cleavage products of lactalbumin, A., i, 182.
- Jones, David Breese, and Henry C. Waterman**, the digestibility of proteins in vitro. III. The chemical nature of the nutritional deficiencies of arachin, A., i, 893.
- Jones, DeWitt O., and Henry R. Lee**, electrometric titration of azo-dyestuffs, A., ii, 239.
- Jones, DeWitt O.** See also **Henry R. Lee**.
- Jones, Edward William Taylor**, obituary notice of, T., 746.
- Jones, G. W., and W. L. Parker**, formation of nitrogen oxides in the slow combustion and explosion methods in gas analysis, A., ii, 223.
- Jones, G. W.** See also **Vernon C. Allison**.
- Jones, J. S., and J. C. Reeder**, the use of silica crucibles for the estimation of potassium in soils, A., ii, 85.
- Jones, Lauder William, and Charles D. Hurd**, rearrangements of some new hydroxamic acids related to heterocyclic acids and to diphenyl- and triphenyl-acetic acids, A., i, 248.
- Jones, Lauder William, and Alfred W. Scott**, new hydroxamic acids derived from cyclopropanecarboxylic acid, isobutyric acid, and dibenzylacetic acid; a comparative study of the Beckmann rearrangement of their derivatives, A., i, 453.
- Jones, Lauder William.** See also **Walter H. Beisler**.
- Jones, Martha R.**, calcium content of blood plasma and corpuscles in the new-born, A., i, 80.
- Jones, P. F.** See **Ellwood B. Spear**.
- Jones, S. Bayne.** See **A. J. Schaffer**.
- Jones, Walter**, the thermostable active agent of pig's pancreas, A., i, 479.
- Jones, William Jacob, and John Bamber Speakman**, some physical properties of aqueous solutions of certain pyridine bases, A., i, 171.
- Jones, William Jacob.** See also **Joseph Greenwood Heap**.
- Jonesco, Stan**, transformation, by oxidation, of the chromogens of some plants into a red pigment, A., i, 97.
- formation of anthocyanin in the flowers of *Cobaea scandens* at the expense of pre-existing glucosides, A., i, 97.
- Jonesco, Stan**, the distribution of anthocyanidins in the coloured organs of plants, A., i, 797.
- the transformation of a chromogen of yellow flowers of *Medicago falcata* under the action of an oxydase, A., i, 1224.
- Jong, Anne Willem Karel de**, the action of light on the cinnamic acids and the constitution of the truxillic acids, A., i, 339.
- biscoumaric acids, A., i, 1049.
- Jong, Anne Willem Karel de, and A. Reclaire**, estimation of the so-called total geraniol content of citronella oil, A., ii, 790.
- Jong, H. G. de.** See **Hugo Rudolph Kruyt**.
- Jonsson, Erik**, alkali silver thiosulphates and their ability to unite with ammonia, A., ii, 57.
- Joos, G.** See **Chr. Fuchtbauer**.
- Joret, Georges**, volumetric estimation of copper by means of sodium nitroprusside, A., ii, 872.
- Jorpes, Erik.** See **Einar Hammarsten**.
- Joseph, Alfred Francis, and Bernard Wyndham Whitfield**, Sudan essential oils, A., i, 754.
- Joseph, Alfred Francis.** See also (*Miss*) **Ida L. Millican**.
- Josephson, K.** See **Hans von Euler**.
- Jost, H.** See **Paul Jacobsen**.
- Joussen, Jacob.** See **Hans Meerwein**.
- Joyner, Reginald Arthur**, the viscosity of cellulose in cuprammonium hydroxide solution. I. The determination of the viscosity, T., 1511.
- the viscosity of cellulose. II. The lowering of the viscosity of cellulose by various reagents, T., 2395.
- Jüllcher, W.** See **Karl von Auwers**.
- Jüptner, Hunns von**, processes in gas generators and blast furnaces, A., ii, 374.
- Jürgens, J.** See **Johan Pieter Wibaut**.
- Juge-Bolard, G.** See **Georges Chaudron**.
- Jumelle, Henri**, barbasau nuts, A., i, 207.
- sakoa oil from Madagascar, A., i, 210.
- Jung, A.**, the influence of hydrogen-ion concentration on the solubility of uric acid, A., i, 1070.
- Jung, A., and H. Müller**, the biological difference of stereoisomerides, A., i, 486.
- Jung, J.**, estimation and distribution of chlorine in plants, A., i, 1098.
- Jung, R.** See **Fritz Mayer**.
- Jungkuns, R.** See **J. Pritzker**.
- Jurist, Alfred E.** See **Ben H. Nicolet**.

Justin-Mueller, Ed., comparative action of heat on cellulose, hydrocellulose, and oxycellulose and the characterisation of hydrocellulose by dry heat, A., i, 11.
 orcinol reaction of furfuraldehyde, A., ii, 95.
 hydrotimetric precision and aqueous preparation of the standard soap solution, A., ii, 658.

K.

Kaan, Maria. See *Hans Fischer*.
Kabeshima, I., toxin of the fish, *Plotosus anguillaris*, A., i, 608.
Käding, Kurt, comparative estimations of acetone in urine, A., ii, 793.
Kaehra, Hans. See *Hermann Leuchs*.
Käse, Artur. See *Sigmund Fränkel*.
Kaffer, H. See *Walther Dilthey*.
Kahho, Hugo, the effect of neutral salts on the heat coagulation of plant protoplasm, A., i, 94.
 action of neutral salts on plant plasma. II. and IV., A., i, 205, 308.
 the toxic action of heavy-metal salts on plant plasma. III., A., i, 311.
Kahlenberg, Louis, some new colour reactions of cholesterol, A., ii, 591.
Kahn, H. M. See *James Bryant Conant*.
Kahn, Max, and *Joseph Barsky*, acute yellow atrophy of the liver, A., i, 969.
Kai, Sotaro, estimation of trypsin; a modification of Gross's method, A., ii, 672.
Kailan, Anton, reactions in penetrating radium radiation and in ultra-violet radiation filtered by quartz glass. II. The hydrogen peroxide equilibrium set up in radium radiation, A., ii, 106.
 the chemical action of penetrating radium rays. XIII. The velocity of formation and equilibrium of hydrogen peroxide, A., ii, 466.
 the chemical action of penetrating radium rays. XIV. The action on oxalic acid, potassium tetroxalate, and potassium chlorate, A., ii, 543.
 the influencing of the catalysis of hydrogen peroxide with platinum by Röntgen rays, A., ii, 756.
Kallan, Anton, and *Emanuel Franz Neumann*, formation and hydrolysis of lactones. II., A., ii, 433.
Kaiser, O. See *Alphonse Gams*.
Kalser, Wilhelm. See *Julius von Braun*.
Kaiha, Sonkyô Kabushiki. See *Yoshitarô Suzuki*.

Kakiuchi, Samuro, physico-chemical properties of phospholipin. I. Precipitation of lecithin hydrosol by electrolytes, A., i, 711.
Kalb, Ludwig. See *Richard Willstätter*.
Kalle & Co., preparation of triazoles of the aromatic series (ψ -azimides), A., i, 61.
 preparation of α -chloronaphthalene derivatives, A., i, 819.
 preparation of naphthasultonesulphonyl chlorides, A., i, 819.
 preparation of 1-arylamino-4-hydroxynaphthalenes, A., i, 823.
 preparation of a derivative of 2-amino-5-hydroxynaphthalene-7-sulphonic acid, A., i, 824.
 preparation of diaminodinitrophenyl- and dinaphthacarbazoole-sulphonic acids, A., i, 867.
Kallmann, Hartmut, theory of strong electrolytes, A., ii, 125.
 carbon monoxide-oxygen cell with glass as electrolyte, A., ii, 467.
Kamm, Oliver, and *Roger Adams*, γ -di-*n*-butylaminopropyl *p*-aminobenzoate, A., i, 828.
Kamm, Oliver, and *Walter H. Newcomb*, $\gamma\gamma'$ -dihalogenodipropyl ethers, A., i, 105.
Kamm, Oliver, and *E. H. Volwiler*, γ -diallylaminopropyl *p*-aminobenzoate, A., i, 654.
Kamm, Oliver, and *John H. Waldo*, $\beta\beta'$ -dichlorodiethyl ether; the oxygen analogue of mustard gas, A., i, 105.
Kanal, T. See *Yoshitane Mori*.
Kanitz, Aristides, the significance of the second dissociation constant of uric acid in the equilibrium of monourate solutions, A., i, 277.
Kannappel, Ernst. See *Wilhelm Strecker*.
Kanô, Naotsuna, preservation of starch solution, A., i, 230.
 applications of amalgams in volumetric analyses. IV. The uses of cadmium amalgam, A., ii, 529.
 conductometric titration of hydrofluosillicic acid, A., ii, 719.
 use of amalgams in volumetric analysis. VI. Electrometric titration of iron, molybdenum, uranium, vanadium, and titanium, A., ii, 721.
Kapfhammer, Joseph, the formation of mercapturic acid during the ingestion of a protein minimum, A., i, 293.
Karamessinis, S. See *Paul Friedländer*.
Karozag, László, oxidation catalysis. I. and II., A., ii, 42, 137.

- Karcsag, László**, the importance of sequence in biology, A., i, 302.
- Karcsag, László**, and **K. Hajós**, the importance of sequence in biology. II., A., i, 302.
- Karlsson, Karl Gustav**, the maximum stability of esters of carboxylic acids, A., ii, 40.
- Karlsson, Signe**. See **Hans von Euler**.
- Karraker, Perry E.** See **Daniel J. Healy**.
- Karrer, Enoch**, rhythmic deposition of precipitated vapours, A., ii, 496.
- Karrer, Paul**, polysaccharides. XII. Glycogen, A., i, 11.
- hydroxycarbonyl compounds; 2:4:6:2'-tetrahydroxybenzophenone, A., i, 40.
- alkali-cellulose and the structure of cellulose, A., i, 231.
- a new degradation of cellulose; conversion of cellulose into a biose anhydride, A., i, 231.
- preparation of amino-alcohols, A., i, 991.
- Karrer, Paul** [with **W. Floroni**], polysaccharides. XVI., A., i, 1119.
- Karrer, Paul**, and **Elisabeth Bürklin**, polysaccharides. XIV. The amyloses, A., i, 435.
- Karrer, Paul, M. Gisler, E. Horlacher, F. Lochner, W. Mäder, and H. Thomann**, proteinogenous amino-alcohols and cholines. II., A., i, 813.
- Karrer, Paul, Ch. Gränacher, and A. Schlosser**, alkylation of the anhydrides of amino-acids, A., i, 235.
- Karrer, Paul, and E. Horlacher**, the decomposition of proteinogenous cholines into alcohols of the cinnamyl alcohol type, A., i, 825.
- Karrer, Paul, C. Nägeli, and Alexander P. Smirnov**, glucosides. X. The action of *dl*-acetobromoglucose on the silver salt of *dl*-mandelic acid, A., i, 253.
- Karrer, Paul, and J. Peyer**, methylated saccharic acid and methylated mucic acid, A., i, 809.
- Karrer, Paul, and J. O. Rosenberg**, sublimation experiments with carbohydrates, A., i, 812.
- Karrer, Paul, and Harry R. Salomon**, crystalline synthetic tannins. I., A., i, 266.
- Karrer, Paul, and Alexander P. Smirnov**, the constitution and configuration of the anhydro-sugars, A., i, 228.
- polysaccharides. XV. Constitution of diamylose and of the anhydro-sugar (cellosan) of cellulose, A., i, 435.
- Karrer, Paul, Max Staub, and A. Wälti**, polysaccharides. XIII. Inulin and the alkali hydroxide compounds of the anhydro-sugars, A., i, 229.
- Karszen, A.** See **J. M. Bijvoet**.
- Karsten, Alfred**. See **Rudolf Ehrenberg**.
- Karvė, D. D.** See **John Joseph Sudborough**.
- Karwat**. See **Bernhard Neumann**.
- Kashima, Kozo**. See **Shigeru Komatsu**.
- Katinssky, Hans von**. See **Hermann Lenchs**.
- Katsch, Gerhardt, and Géza Némethy**, alkapton chromogen, A., i, 198.
- Kattwinkel, R.**, the melting point of normal ammonium sulphate, A., ii, 445.
- Katz, Sidney H., and J. J. Bloomfield**, tests of an iodine pentoxide indicator for carbon monoxide, A., ii, 535.
- Kaufmann, Hugo**, constitution of carbonium dyes; halochromism, A., i, 820.
- nature of the ionogen linking, A., ii, 366.
- Kaufmann, Adolf**, synthesis of quinic acid, A., i, 464.
- Kaufmann, H. P.**, the exchange of halogen in unsaturated, aliphatic halogenated hydrocarbons. I., A., i, 213.
- bactericidal action of pyromucic acid, A., i, 304.
- keto-enolic tautomerism, A., i, 985.
- Kaufmann, H. P., and M. Friedebach**, a variety of wax from pine needles and certain abietic esters, A., i, 740.
- Kaufmann, H. P., and W. Kaufmann** [with **H. Götting**], substituted salicylic acids. I., A., i, 252.
- Kaufmann, H. P., and M. Schneider**, condensations of acetylene. I. Elucidation of the constitution of cuprene, A., i, 245.
- Kaufmann, H. P., and H. Zobel**, isomeric sulphonimides of naphtholic acid, a contribution to the theory of dulcogenic groups, A., i, 744.
- Kaufmann, H. P.** See also **Ludwig Knorr**.
- Kaufmann, W.** See **H. P. Kaufmann**.
- Kautsky, Hans, and H. Zecher**, the relationship between chemi- and photoluminescence in unsaturated silicon compounds, A., ii, 464.
- Kay, Herbert Davenport, and Henry Stanley Baper**, mode of oxidation of fatty acids with branched chains. II. The fate in the body of hydratropic, tropic, atrolactic, and atropic acids together with phenylacetaldehyde, A., i, 1093.

- Kay, L. J.** See *Edward de Barry Barnett*.
- Kaye, F.** See *Julius Huebner*.
- Kaysor, Erich.** See *Emil Fromm*.
- Keeler, L. M.** See *L. Bauman*.
- Keeler, R. F.** See *A. W. Clark*.
- Keeton, Robert W.**, excretion of ammonia following experimental administration of acids via the stomach and peripheral vein, A., i, 800.
- Kehrmann, Friedrich**, constitution of quinonoid organic onium salts, A., i, 259.
theory of carbonium compounds, A., i, 331.
constitution and colour. VIII., A., ii, 333.
- Kehrmann, Friedrich**, and *Jean Henri Dardel*, sulphonium bases derived from thiodiphenylamine and analogous substances, A., i, 1063.
- Kehrmann, Friedrich**, *Herman Decker*, and *Ch. Schmajewski*, oxonium salts from phenol ethers and phenols. II. Decomposition of oxonium salts from thymol ethyl ether and from anisole by water or substances of alkaline reaction, A., i, 32.
- Kehrmann, Friedrich**, *Herman Decker*, and *B. Solonina*, oxonium salts from phenol ethers and phenols. I. Nitrosophenol dyes. IV., A., i, 32.
- Kehrmann, Friedrich**, and *R. van der Laar*, oxonium salts of azo-substances, A., i, 384.
- Kehrmann, Friedrich**, and *St. Micevitz*, the history of the blue oxidation product from diphenylamine, A., i, 28.
- Kehrmann, Friedrich**, and *Gustave Roy*, the blue oxidation product of diphenylamine, A., i, 467.
- Kehrmann, Friedrich**, *Gustave Roy*, and *Marie Ramm*, the so-called peroxidation products of leuco-triphenylmethane dyes, A., i, 467.
- Kehrmann, Friedrich**, *Maurice Sandoz*, and *R. Monnier*, nitro-derivatives of quinol, A., i, 33.
- Kell, J.** See *Walter König*.
- Keith, Clyde R.** See *Paul W. Merrill*.
- Keller, Rudolf**, acidity and basicity, A., ii, 131.
- Kellermann, Karl.** See *Fritz Weigert*.
- Kelley, George Leslie**, and *E. W. Evers*, solid sodium hydroxide as an absorbent for carbon dioxide in steel analysis, A., ii, 160.
- Kelley, George Leslie**, and *J. A. Wiley*, estimation of chromium in ferrochromium by electrometric titration, A., ii, 164.
- Kelley, George Leslie, J. A. Wiley, Raymond T. Bohn**, and *W. C. Wright*, estimation of vanadium and chromium in ferrovanadium by electrometric titration, A., ii, 89.
- Kelley, Louise.** See *William Ridgely Oradoff*.
- Kellner, J.**, estimation of glycerol by the dichromate method, A., ii, 399, 723.
- Kellner, William**, obituary notice of, T., 2912.
- Kelly, Margaret W.** See *Arthur W. Thomas*.
- Kelsey, Erwin B.** See *Arthur Joseph Hill*.
- Kemp, Archie R.** See *Howard J. Lucas*.
- Kendall, F. E.** See *H. H. Mitchell*.
- Kendall, James**, application of ideal solution equations to dilute aqueous solutions, A., ii, 32.
abnormality of strong electrolytes and the ionisation theory of Ghosh, A., ii, 419.
- Kendall, James**, and *Howard Adler*, compound formation and solubility in systems of the type, formic acid-metal formate, A., i, 5.
- Kendall, James**, *Howard Adler*, and *Arthur W. Davidson*, prediction of solubility in polar solutions, A., ii, 34.
compound formation and conductivity in systems of the types formic acid-metal formate and sulphuric acid-metal sulphate, A., ii, 126.
- Kendall, James**, and *James C. Andrews*, solubilities of acids in aqueous solutions of other acids, A., ii, 34.
- Kendall, James**, and *J. J. Beaver*, compound formation in phenol-cresol mixtures, A., i, 136.
- Kendall, James**, and *Elizabeth Brakeley*, compound formation and viscosity in solutions of the types acid-ester, acid-ketone, and acid-acid, A., ii, 126.
- Kendall, James**, and *Francis J. Fuels*, catalytic influence of foreign oxides on the decomposition of silver oxide, mercuric oxide, and barium peroxide, A., ii, 147.
catalytic decomposition of certain oxides, A., ii, 646.
- Kendall, James**, and *Paul M. Gross*, factors affecting the stability of additive compounds in solution and their influence on ionisation equilibria. I., A., ii, 32.
compound formation and specific conductivity in solutions of the types, acid-ester, acid-ketone, and acid-acid, A., ii, 33.

- Kennaway, Ernest Laurence**, and **James McIntosh**, the action of whole blood on acids, A., i, 788.
- Kenner, James**, configurations of molecules of benzenoid substances, A., i, 533.
- Kenner, James**. See also **Harold Burton**, and **George Hallatt Christie**.
- Kenyon, Joseph**, a simplified method for the resolution of methyl-*n*-hexyl-carbinol, T., 2540.
- Kerb, Johannes**, and **Kurt Zeekendorf**, the course of alcoholic fermentation in presence of calcium carbonate, A., i, 305.
- Kerb, Johannes**. See also **A. Bornstein**.
- Kerecsy, Georg von**, and **Emil Wolf**, preparation of morphine allyl ether, A., i, 854.
preparation of aryl sulphonic esters of halogenated aliphatic alcohols, A., i, 1131.
- Kerkow, F.** See **Karl Fries**.
- Kermack, William Ogilvy**, **William Henry Perkin, jun.**, and **Robert Robinson**, harmine and harmaline. VI. The synthesis of *N*-methyltetrahydronorharmine and the constitution of harmaline and of the alkylated harmines, T., 1872.
- Kermack, William Ogilvy**, and **Robert Robinson**, an explanation of the property of induced polarity of atoms and an interpretation of the theory of partial valencies on an electronic basis, T., 427.
- Kern, E.** See **Walter Madelung**.
- Kesseler, H.**, **R. Röhm**, and **G. Lutz**, the estimation of nitrogen in nitric esters, A., ii, 392.
- Keyes, Frederick G.**, Joule-Thomson effect for air, A., ii, 24.
- Keyes, Frederick G.**, **Louis J. Gillespie**, and **Shinroku Mitsukuri**, a continuous flow calorimeter, and the determination of the heat of neutralisation of a solution of hydrochloric acid by one of sodium hydroxide, A., ii, 424.
- Keyes, Frederick G.**, and **H. Hara**, pressure of oxygen in equilibrium with silver oxide, A., ii, 353.
- Keyes, Harmon E.** See **Herman V. Tartar**.
- Keyser, Ernst**. See **Hartwig Franzen**.
- Kharasch, Morris S.**, aromatic mercuri-organic derivatives; the Hofmann rearrangement and the nature of valencies of mercury in mercuri-organic derivatives, A., i, 188.
an indirect method of preparation of organic mercuric derivatives and a method of linking carbon to carbon, A., i, 189.
- Kharasch, Morris S.**, and **Isadore M. Jacobssohn**, mercuri-organic derivatives; the mercurisation of aromatic amines and its relation to the theory of substitution, A., i, 189.
- Kharasch, Morris S.**, **Frederick W. M. Lommen**, and **Isadore M. Jacobssohn**, the nitroanilines, A., i, 608.
- Khoury, J.**, inconstancy of the precipitation of uric acid from urine in the form of cuprous urate, A., ii, 885.
- Kieckinger, Heinrich**, decomposition of citric acid of cow's milk by some bacteria, A., i, 1219.
- Kieran, Arthur Joseph**, electrical conductivity of hydrochloric acid and potassium chloride in presence of sucrose, A., ii, 812.
- Kiesel, Alexander**, yeast protein, A., i, 408.
the presence of ornithine in plants, A., i, 412.
the glutencasein of buckwheat, A., i, 412.
the enzymic degradation of arginine in plants. II., A., i, 413.
the action of arginase on agmatine and tetramethylenediguanidine; the specificity of enzymes, A., i, 413.
synthesis and properties of tetramethylenediguanidine, A., i, 531.
the constituents of the pollen grain of *Pinus sylvestris*, A., i, 799.
- Kiesel, Alexander**, and **Troitzki**, the distribution of urease in plants, A., i, 410.
- Kiess, A.** See **Alexander Gutbier**.
- Kiess, C. C.**, **B. Smith Hopkins**, and **Harry C. Kremers**, wave-lengths longer than 5500 Å. in the arc spectra of yttrium, lanthanum, and cerium, and the preparation of pure rare earth elements, A., ii, 244.
- Kikuchi, Mitzugi**, purine metabolism; fate of ingested purines, A., i, 698.
- Kikuchi, Sutarō**, application of amalgams in volumetric analyses. III. Estimation of iodic, bromic, and chloric acids, A., ii, 519.
effect of adsorption by stannic sulphide on the estimation of phosphates, A., ii, 525.
use of amalgams in volumetric analysis. V. Estimation of iron, titanium, uranium, and methylene-blue, A., ii, 721.
- Kiliani, Heinrich**, chemistry of the sugars. II. and IV., A., i, 223, 1111.
- Kiliani, Heinrich**, [with **Aug. Wingler**], chemistry of the sugars. III., A., i, 321.

- Killian, J. A.**, and **Carl P. Sherwin**, chemistry of normal and abnormal pregnancies, A., i, 489.
- Kilpatrick, Mary L.** See **Annie Louise Macleod**.
- Kimura, Kenjiro.** See **Yaji Shibata**.
- Kimura, Masamichi**, and **Mitsuharu Fukuda**, the intensities of the lines in the Balmer series of hydrogen, A., ii, 801.
- influence of gases and vapours on the intensities of the lines of the secondary spectrum of hydrogen, A., ii, 801.
- Kindermann, E.** See **Georg Schroeter**.
- Kindler, Karl.** See **Paul Rabe**.
- King, Anne Millicent**, the effect of high concentration of salt on the viscosity of a soap solution, A., i, 621.
- King, Arthur S.**, the electric furnace spectrum of scandium, A., ii, 100.
- electric furnace experiments involving ionisation phenomena, A., ii, 810.
- King, George.** See **John Addyman Gardner**.
- King, Harold**, the isolation of muscarine, the potent principle of *Amanita muscaria*, T., 1743.
- King, Harold**, and **Albert Donald Palmer**, the resolution of tropic acid and the stereochemical configuration of the cinchona alkaloids, T., 2577.
- King, H. H.**, and **E. W. Wampler**, adsorption and orientation of the molecules of dibasic acids and their ethereal salts in liquid-vapour interfaces, A., ii, 741.
- King, Herbert Joseph Seymour.** See **Gilbert Thomas Morgan**.
- King, Harold S.**, correlation of atomic structure and spectra, A., ii, 277.
- a proposed laboratory test of the theory of relativity, A., ii, 439.
- King, J. Fitch**, and **Walter A. Patrick**, measurement of dielectric constants, A., ii, 109.
- King, Jessie L.**, anticoagulating substances in the mucous membrane of the uterus, A., i, 701.
- King, Louis V.**, lecture-room demonstration of atomic models, A., ii, 705.
- Kinney, A. McB.** See **James Bryant Conant**.
- Kinose, Jiro.** See **Takaaki Sasaki**.
- Kinugasa, Yutaka**, and **Hisajiro Tatsuno**, detection of β -naphthol in foods, spices, and beverages, A., ii, 400.
- Kirby, W.**, determination of the melting- and boiling-points of anthracene, phenanthrene, and carbazole, A., i, 27.
- Kirchhof, Franz**, the action of concentrated sulphuric acid on natural and artificial caoutchoucs. II., A., i, 562.
- empirical and structural composition of natural and synthetic caoutchouc, A., i, 945.
- Kirschbaum, George.** See **Julius von Braun**.
- Kisch, Franz**, elimination of iron in urine, A., i, 898.
- Kishi, Niichiro.** See **Heisaburo Kondô**.
- Kiss, Julius**, physical chemistry of alexin-fixation reaction, A., i, 706.
- Kissling, A.** See **Rudolf Friedrich Weinland**.
- Kittredge, Esther B.** See **Edna R. Bishop**.
- Kjellin, Carl**, hydroxylamine, A., ii, 640.
- Klänhardt**, the distillation of greatly frothing liquids, A., ii, 472.
- Klason, Peter**, lignosulphonic acid, A., i, 14.
- the cellulose content of pine wood, A., i, 99.
- constitution of pine lignin. II., A., i, 324.
- lignin as it occurs in wood, A., i, 413.
- Kleeman, Richard Daniel**, an electrical doublet theory of the nature of the molecular forces of chemical and physical interaction, A., ii, 866.
- Klein, G.**, anthochlor, A., i, 1099.
- Klein, Paul.** See **Isidor Traube**.
- Klein, W.**, cellulose fermentation in the paunch of the ox and its importance for metabolic experiments, A., i, 82.
- Klein, W.**, and **Maria Steuber**, method for direct estimation of carbon dioxide and oxygen in the Berthelot bomb and its importance for the metabolic balance of herbivora, A., ii, 159.
- Kleiner, Israel Simon.** See **Isaac Neuwirth**.
- Kleinfeller.** See **Otto Diels**.
- Kleitman, Nathaniel.** See **William Salant**.
- Klemenc, Alfons**, behaviour of a non-attackable electrode in the process leading to the equilibrium $3\text{HNO}_2 \rightleftharpoons 2\text{NO} + \text{HNO}_3 + \text{H}_2\text{O}$, A., ii, 258.
- Klemenc, Alfons**, and **Cornelie Bunsel**, comparison of the methods for the estimation of nitric oxide, A., ii, 783.
- Klemenc, Alfons**, and **Friedrich Pollak**, decomposition of nitrous acid, A., ii, 432.
- titration of nitrous acid, and the estimation of nitrous and arsenious acids in the presence of each other, A., ii, 865.

- Klement, F.** See *David Reichinstein*.
- Klement, R.** See *Ernst Hermann Riesenfeld*.
- Klemm, L.** See *Wilhelm Eller*.
- Klemmer, Aloys**, high percentage hydrogen peroxide (perhydrol) for the estimation of the total sulphur in illuminating gas, A., ii, 224.
- Kleucker, Ernst**, condensations of *p*-nitrobenzyl chloride with cinnamaldehyde and furfuraldehydes, A., i, 784.
- Klimont, Isidor**, co-ordination forms of glycerides. I., A., i, 517.
apparatus for hydrogenation at ordinary pressures, A., ii, 369.
estimation of chlorine in organic compounds, A., ii, 580.
- Kling, André**, and **Arnold Lassieur**, separation and estimation of copper, lead, antimony, and tin; analysis of white metals, A., ii, 86.
the analysis of sour milk, A., ii, 402.
rapid electro-analysis, A., ii, 587.
apparatus for measuring the hydrogen-ion concentration of a solution, A., ii, 715.
- Kling, André**, **Arnold Lassieur**, and (*Mme*) **Arnold Lassieur**, apparatus for measuring the hydrogen-ion concentration of a solution; application to the detection of mineral acids in vinegar, A., ii, 234, 519.
- Klingstedt, F. W.**, ultra-violet absorption spectrum of phenol in different solvents, A., ii, 332.
the ultra-violet absorption spectra of dihydric phenols, A., ii, 680.
- Knauff-Lenz, E.**, intestinal saccharase, A., i, 485.
blood saccharase and the antigen properties of yeast saccharase, A., i, 694.
- Knaggs, (Miss) Isabel Ellie**, inorganic complex salts; crystallographic and optical study. I., T, 2069.
- Knapp, Arthur W.**, and **Raymond V. Wadsworth**, reactions between the higher fatty acids and salts of the lower fatty acids, A., i, 219.
- Knecht, Edmund**, and **Eva Hibbert**, preparation of α -pimaric acid of m. p. 212° , A., i, 1020.
- Knecht, Edmund**, and **F. P. Thompson**, behaviour of oxidised cellulose, A., i, 687.
- Knesabeek, Anna Marie von dem**, and **Fritz Ullmann**, the xanthone series, A., i, 559.
- Knies, W.** See *Fritz Mayer*.
- Knipping, Paul**, ionisation tension of the halogen hydrides, A., ii, 188.
- Knoevenagel, Emil**, keto-anils. IV. Reduction products of keto-anils, A., i, 1060.
- Knoevenagel, Emil**, and **Hans Bähr**, ketoanils. II. Constitution of the *N*-alkylketoanils and transformation of aliphatic ketoanils into derivatives of quinoline, A., i, 750.
- Knoevenagel, Emil**, and **Hedwig Busch**, an alkali-soluble modification of cellulose, A., i, 686.
- Knoevenagel, Emil**, and **Oskar Goos**, ketoanils. III. Fatty-aromatic keto-anils, A., i, 751.
- Knoll & Co.**, preparation of neutral soluble double compounds of the alkyl xanthines and their *N*-acyl derivatives, A., i, 464.
- Knoop, Franz**, reductions and oxidations and a coupled reaction in the intermediary metabolism of the animal body, A., i, 486.
- Knorr, Ludwig**, and **H. P. Kaufmann**, equilibrium in solution of the desmotic-isomeric diacetylsuccinic esters and its colorimetric estimation, A., i, 220.
- Knoth, G.** See *Hermann Ost*.
- Knowles, H. B.** See *Gustav Ernst Fred Lundell*.
- Knudson, Arthur**, and **Melvin Dresbach**, a chemical method of assaying the active principles of digitalis, A., ii, 882.
- Knudson, Arthur**. See also *F. S. Randles*.
- Kobayashi, Shōmei**, alligator and crocodile oils, A., i, 792.
a highly unsaturated hydrocarbon, and some higher alcohols in a commercial illipé fat, A., i, 1113.
- Kobayashi, Shōmei**. See also *Shōichi Nakatogawa*.
- Koch, Erich**. See *Waller Gerlach*.
- Koch, Peter Paul**, and **Fritz Schrader**, action of light on silver chloride, bromide, and iodide, A., ii, 182.
- Koda, Chu**. See *Leon Asher*.
- Kodama, Shintarō**, preparation of phenyl-acetaldehyde, A., i, 849.
- Kodama, Shintarō**. See also *Kikunae Ikeda*.
- Koeschlin, Rudolf**, minerals from Ljubija, Bosnia, A., ii, 860.
- Kögl, Fritz**. See *Heinrich Wieland*.
- Köhler, Antonie**. See *Kurt Lindner*.
- Köhler, Rohdan**, iodometric studies. II., A., ii, 530.
- Köhler, K.** See *Walter König*.
- Köhler, Ludwig**. See *Kurt Brass*.

- Keenig, A.**, and **W. Habbueh**, formation of hydrogen cyanide from nitrogen and hydrocarbons in the electric arc, A., i, 642.
- König, Fritz**, estimation of perchlorate by Rothmund's method, A., ii, 810.
- König, Josef, J. Hasenbäumer**, and **E. Kroger**, factors in the development of soil acidity, A., i, 510.
- König, Theodor**. See **Ernst Weitz**.
- König, Walter**, [with **H. Zorn**], the constitution of the pinacyanols, a contribution to the chemistry of the quinoxanines, A., i, 1188.
- König, Walter**, and **J. Keil**, 1:8-naphthasultam-4-sulphonic acid and certain of its derivatives, A., i, 822.
- König, Walter**, and **K. Köhler**, the mechanism of coupling reactions. II. 1:8-naphthasultam and its *N*-methyl derivative as azo-components, A., i, 879.
- Koessler, Karl K.** See **Milton Th. Hanke**.
- Köster, W.** See **Gustav Tammann**.
- Koetschau, Rudolf**, the ozonides of petroleum, A., i, 977.
- Kötz, Arthur**, and **H. Rathert**, simultaneous reduction and oxidation. III. Transformation of halogenaldehydes into aldehydes and acids through ketenes, A., i, 286.
- Köfler, Ludwig**, surface activity and toxic action of saponins, A., i, 610.
- differentiation and estimation of saponins, A., ii, 595.
- Kohl, C.** See **Fritz Mayer**.
- Köhler, (Mlle) Denise**. See **Raoul Combes**.
- Köhler, Elmer Peter**, the addition of malonic esters to benzoylphenylacetylene, A., i, 461.
- the action of bromine on certain β -ketonic esters, A., i, 552.
- Köhler, Elmer Peter**, and **L. I. Smith**, the reaction between alkalis and certain nitrocyclopropane derivatives, A., i, 457.
- Kohlshütter, Volkmar**, and **H. Schödl**, I. The structure of electrolytically deposited nickel. II. The influence of superposed alternating current on the deposition and solution potential of nickel, A., ii, 648.
- Kohlweller, Emil**, fractional diffusion of iodine vapour, A., ii, 497.
- Kohn, Moritz**, reducing actions of arsenious acid, A., ii, 143.
- behaviour of ammoniacal and alkaline copper solutions, A., ii, 149.
- Kohn, Moritz**, and **Avissim Mendelewitsch**, symmetrical dibromopivalic acid and 1-methylcyclopropane-1-carboxylic acid, A., i, 518.
- Kohn, P.** See **Ludwig Moser**.
- Koisumi, Shungo**, electrolytic oxidation of alcohols. I. *iso*Amyl alcohol, A., i, 979.
- Kolkmeijer, N. H.**, crystal structure of germanium, A., ii, 713.
- Kolligs, H.** See **Karl von Auwers**.
- Kollmann, Gustav**, synthesis of uric acid in the human organism, A., i, 298.
- Kollo, Constantin**. See **Stefan Minovici**.
- Kolmer, J. A.** See **J. F. Schamberger**.
- Kolthoff, I. M.**, the hydrolysis of anti-pyrene salicylate, A., i, 471.
- the normal potential of silver, A., ii, 20.
- estimation of small quantities of bromides and chlorides in iodides, A., ii, 79.
- the estimation of sulphates by means of a suspension of barium chromate, A., ii, 81.
- the sensitiveness of coloured indicators at temperatures above the ordinary, A., ii, 156.
- the separation of silver from mercurous salts, A., ii, 160.
- the identification of lævulose in presence of aldoses, A., ii, 166.
- electro-adsorption as a purely chemical process, A., ii, 197.
- the salt error of coloured indicators, A., ii, 222.
- the colorimetric estimation of hydrogen-ion concentration without buffer solutions, A., ii, 222.
- the titration of moderately strong acids or bases in the presence of very weak ones, A., ii, 228.
- potassium ferricyanide as a reagent in iodometry, A., ii, 224.
- the argentometric titration of phosphoric acid, A., ii, 314.
- the application of the iodine electrode in potentiometric titrations, A., ii, 385.
- the application of conductometric methods to precipitation analysis, A., ii, 452.
- qualitative reactions for arsenic, A., ii, 455.
- potentiometric titrations of, and by means of, potassium ferrocyanide. I. The titration of potassium ferrocyanide by means of potassium permanganate, A., ii, 587.
- the use of potassium ferrocyanide in potentiometric titrations. II. The potentiometric titration of zinc, A., ii, 580.

- Kolthoff, I. M.**, electrometric titrations with silver nitrate; estimation of chlorides, bromides and iodides, and of iodides in the presence of chlorides and bromides, A., ii, 581.
- electrometric titrations with mercury perchlorate, A., ii, 655.
- electrometric titrations with lead nitrate, A., ii, 781.
- application of conductometric titrations to precipitation analysis. V. Conductometric titrations with barium salts, A., ii, 864.
- titration of boric acid in presence of phosphoric acid, A., ii, 867.
- Kolthoff, I. M.**, and **Ada Bak**, the use of mercuric nitrate instead of silver nitrate in the estimation of the halogens, A., ii, 159.
- Kolthoff, I. M.**, and **C. J. Cremer**, the iodometric estimation of copper and arsenic present together, especially in Paris and Schweinfurth greens, A., ii, 86.
- Kolthoff, I. M.**, and **E. H. Vogelenzang**, the acidimetric estimation of dichromate, A., ii, 88.
- Komatsu, Shigeru**, and **Kozo Kashima**, constitution of polysaccharides. I. Xylan and its acetyl derivatives. I., A., i, 811.
- Komatsu, Shigeru**, and **Bunkichi Masumoto**, catalytic reduction of *d*-camphor. I., A., i, 752.
- Komm, E.** See **Alfred Heiduschka**.
- Komppa, Gustav**, and **R. H. Roschier**, the camphenilone group. III. The homogeneity of apobornylene; apocyclene, a new tricyclic hydrocarbon, A., i, 1167.
- Komuro, K.**, the minimum of odour perceptible in an absolutely inodorous space (camera inodorata), A., i, 84.
- Kon, George Armand Robert**, the formation and stability of *spiro*-compounds. VII. The application of the Dieckmann reaction to esters of the glutaric series, T., 513.
- Kon, George Armand Robert, Arnold Stevenson**, and **Jocelyn Field Thorpe**, ring-chain tautomerism. I. The occurrence and effect of keto-enol tautomerism between a ring compound and its open-chain isomeride, T., 650.
- Kon, George Armand Robert**, and **Jocelyn Field Thorpe**, a method for the preparation of $\alpha\beta\beta$ -trialkylated glutaric acids, T., 1795.
- Kon, George Armand Robert**. See also **Frank Dickens**.
- Kondô, Heisaburô, Niichirô Kishi**, and **Chârô Araki**, constitution of matrine. II., A., i, 269.
- Kondô, Heisaburô, Tomoichi Nakajima**, and **Gorô Murakawa**, 4-nitrosalicylic acid, A., i, 745.
- Kondô, Heisaburô**, and **Torizô Takahashi**, condensation of collidine with acetaldehyde, A., i, 1177.
- Kondo, Yoshio**. See **Yasuhiko Asahina**.
- Kondyrev, N. V.**, electrolysis of organo-magnesium compounds, A., i, 1128.
- Konek, Fritz von**, and **Nikolaus Szamák**, hydrindones. I. The hydrindone of the vanillin series, A., i, 458.
- Konen, H.** See **Leonhard Grebe**.
- Konishi, M.**, production of acetoacetic acid from urocanic acid in the surviving liver, A., i, 1211.
- Konishi, M.** See also **Yashiro Kotake**.
- Konrad, Erich**. See **Robert Schwarz**.
- Koopman, J.**, acetone in cerebrospinal fluid, A., i, 194.
- Kopatschek, Fritz**, action of uranyl acetate on some organic substances. I. Action of uranyl acetate on tartaric acid and its salts, A., i, 984.
- Kopetschni, Eduard**, and **Herta Wiesler**, a double decomposition catalysed by copper, A., i, 844.
- a new mode of formation of thiazole derivatives of the anthraquinone series, A., i, 868.
- Kornick, Erich**. See **Emanuel Merck**.
- Koser, S. A.**, trehalose fermentation in the differentiation of the paratyphoid-enteritidis group, A., i, 407.
- Kostytschev, S.**, photosynthesis. I. The ratio of carbon dioxide to oxygen in carbon assimilation, A., i, 307.
- photosynthesis. II. Does injury stimulate photosynthesis? A., i, 308.
- photosynthesis. III. Does carbon assimilation take place during summer night in subarctic regions? A., i, 308.
- photosynthesis. IV. Carbon dioxide assimilation by Leguminosæ, A., i, 613.
- Kostytschev, S.**, and **Paul Eliasberg**, the invertase of *Mucor racemosus*, A., i, 410.
- Kotake, Yashiro**, the deamination of amino-acids and the reversible transformations of the products so arising in the animal organism, A., i, 1218.
- Kotake, Yashiro**, and **M. Konishi**, the production of urocanic acid from histidine in the dog, A., i, 1217.

- Kotake, Yashiro, Y. Masai, and Yoshitane Mori**, the behaviour of phenylalanine in the animal organism, A., i, 1217.
the behaviour of amino-acids in vitally stained animals. I. and II., A., i, 1217.
- Kotake, Yashiro, Z. Matsuka, and M. Okagawa**, the deaminating of tyrosine in the animal organism, A., i, 1218.
- Kotake, Yashiro, and Yoshitane Mori**, the behaviour of phenyl-lactic acid in the animal organism. I., A., i, 1216.
the behaviour of phenylpyruvic acid in the animal organism, A., i, 1216.
- Kotake, Yashiro, and M. Okagawa**, influence of optical activity on cell permeability. I., A., i, 696.
the excretion of hydroxyphenyl-lactic acid after administration of tyrosine to rabbits, A., i, 1217.
- Koslowski, Antoine**, formation of the red pigment of *Beta vulgaris* by oxidation of the chromogens, A., i, 96.
- Koslowski, Stanislas**. See *Stefan Dombrowski*.
- Kracker, H.** See *Otto Fischer*.
- Kraemer, O.** See *A. Schönberg*.
- Kraft, K.**, barium compounds in the viscera, A., i, 301.
- Kraft, Edgar**. See *Wilhelm Schneider*.
- Kraisay, Anton**, re-testing the 100°-point of the saccharimeter. II. Preparation of chemically pure sucrose, A., ii, 233.
- Kramer, Benjamin, and Frederick F. Tisdall**, distribution of sodium, potassium, calcium, and magnesium between the corpuscles and serum of human blood, A., i, 1087.
- Kramer, S. P.**, behaviour of $\beta\beta$ -dichlorodiethyl sulphide, A., i, 914.
- Krase, Herbert J.**, simple formula for the calculation of the specific heats of solids, A., ii, 421.
- Kratzer, A.**, determination of the terms of the cyanogen bands, A., ii, 4.
regularities in systems of bands, A., ii, 409.
- Kraus, Charles August, Ghosh's theory** of electrolytic solutions, A., ii, 256.
equilibrium in mixtures of binary electrolytes A., ii, 269.
constitution of metallic substances, A., ii, 566.
- Kraus, Charles August, and John Egbert Bishop**, conductivity of solutions of ternary electrolytes in propyl alcohol, A., ii, 19.
- Kraus, Charles August, and John Egbert Bishop**, conductivity of sodium iodide in amyl alcohol at very low concentrations, A., ii, 813.
- Kraus, Charles August, and C. Y. Chiu**, nature of the complexes formed between sodium and tellurium in liquid ammonia, A., ii, 765.
- Kraus, Charles August, and Walter W. Lucasse**, conductivity of concentrated solutions of sodium and potassium in liquid ammonia, A., ii, 252.
resistance temperature coefficient of concentrated solutions of sodium in liquid ammonia, A., ii, 734.
composition of the liquid phases in a univariant system, liquid-liquid-vapour, for mixtures of sodium and ammonia, A., ii, 764.
- Kraus, Charles August, and Edward H. Zeltfuchs**, equilibrium in liquid mixtures of ammonia and xylene, A., i, 725.
- Kraus, Ernst Josef**, volumetric estimation of aluminium, A., ii, 87.
- Krause, A. C.** See *Homér Adkins*.
- Krause, Erich, and Rudolf Nitsche**, preparation of organic compounds of boron with the aid of boron fluoride. I. Boron alkyls and alkylboric acids, A., i, 22.
preparation of organic compounds of boron with the aid of boron fluoride. II. Boron triphenyl and phenylboric acid, A., i, 694.
- Krause, Erich, and Erich Pohland**, the fluorides of organo-metallic compounds. II. Lead alkyl and aryl fluorides, A., i, 644.
- Krause, Erich, and G. G. Reissaus**, lead triaryl, a parallel to triphenylmethyl. III. Lead triphenyl, tri-*p*-tolyl and tri-*o*-tolyl and dark red lead diaryls, A., i, 602.
- Krause, K. E.** See *P. P. Budnikov*.
- Krause, Leonhard**. See *Arthur Rosenheim*.
- Krauss, F.**, ruthenium tetroxide, A., ii, 75.
- Krauss, F.** See also *Alexander Galtier*.
- Krauss, Mathilde**. See *Franz Faltis*.
- Kreger, C. W.** See *Roger Adams*.
- Kreidl, J.** See *Walter Herzog*.
- Kreis, W.** See *Hermann Staudinger*.
- Kreitz, A. v.**, cobalt-tungsten alloys, A., ii, 381.
- Kremann, Robert, Franz Hemmelmayr, and Heinrich Reimer**, influence of substitution in the components on equilibria in binary solutions. XXXVI. The equilibria of dihydroxynaphthalenes with amines, A., i, 1010.

- Kremann, Robert, Heins Hohl, and Robert Müller**, influence of substitution in the components on equilibria in binary solutions. XXXV. The binary systems of triphenylcarbinol with pyrogallol, nitrophenols, polynitrobenzenes, and phenylenediamines, A., i, 138.
- Kremann, Robert, and Robert Müller**, influence of substitution in the components on equilibria in binary solutions. XXXIV. The binary systems of anthracene with nitro-derivatives of benzene, A., i, 131.
- Kremann, Robert, and Friedrich Odelga**, influence of substitution in the components on equilibria in binary solutions. XXXII. The binary systems of camphor with phenols, A., i, 159.
- Kremann, Robert, Friedrich Odelga, and Othmar Zawodsky**, influence of substitution in the components on equilibria in binary solutions. XXXI. The binary systems of triphenylmethane with amines and phenols, A., i, 131.
- Kremann, Robert, Wilhelm Bösl, and Wilhelm Fenkner**, dynamics of the formation of nitriles from acid anhydrides and amides. III. Reaction between benzamide and acetic anhydride, A., ii, 748.
- Kremann, Robert, and Hubert Strzelba**, influence of substitution in the components on equilibria in binary solutions. XXXIII. The binary systems of carbazole or acenaphthene with polynitro-derivatives of benzene and toluene, A., i, 176.
- Kremann, Robert, Alfred Zoff, and Victor Oswald**, dynamics of the formation of nitriles from acid anhydrides and amides. II. Kinetic study of the reaction between acetamide and acetic anhydride, using phase rule methods, A., ii, 748.
- Kremers, Harry C.** See *C. C. Kiess*.
- Kremers, Roland E.**, biogenesis of oil of peppermint, A., i, 357.
volatile oil of milfoil, A., i, 505.
the volatile oil of *Mentha aquatica*, Linné, and a note on the occurrence of pulegone, A., i, 848.
- Krestinski, V.**, magnesium compounds of the olefines. I., II., and III., A., i, 1128.
- Krets, Fritz**, microchemical detection of tryptophan in the plant, A., ii, 668.
- Kreutzer, A.** See *Hans Tropisch*.
- Křížka, P.** See *Julius Stoklasa*.
- Krieger, Wilhelm**. See *Fritz Mayer*.
- Krishna, Sri**, phenolcamphorein, T., 253.
- Krishna, Sri, and Frank George Pope**, the action of potassium iodide and iodate on some hydroxy-acids, T., 798.
- Kritschewsky, J. L.**, the action of salvarsan on the serum of animals and on blood-cells in vitro, A., i, 394.
- Kröger, E.** See *Josef König*.
- Kröger, M.**, preparation of silicic acid and tungsten hydroxide sols by means of Hildebrand cells, A., ii, 212.
influence of tungstic acid on the gelatinisation of silicic acid in concentrated hydrochloric acid solutions, A., ii, 213.
- Krönig, W.** See *Klothilde Meier*.
- Kropf, Alfred**, colorimetric estimation of vanadium in steel, A., ii, 590.
absorption of carbon monoxide by acid cuprous chloride solution in the presence of a reducing agent, A., ii, 657.
- Kruber, Otto**. See *Gesellschaft für Teerverwertung*.
- Krüger, R.** See *Leonor Michaelis*.
- Krull, H.** See *Alfred Wohl*.
- Krull, R.**, Causse and Bonnan's method for the estimation of dextrose, A., ii, 790.
- Kruyt, Hugo Rudolph, and H. G. de Jong**, capillary-electric phenomena in lyophilic sols, A., ii, 357.
- Kryz, Ferdinand**, colour reaction of sucrose, A., ii, 233.
- Krzaskowsky, C.** See *Giuseppe Amantea*.
- Kuba, Franz**. See *Augustin Bistrayeki*.
- Kudo, Yoshihiro**. See *Riko Majima*.
- Kühr, C. A. H. von Wolzogen**, occurrence of sulphate reduction in the deeper layers of the earth, A., i, 1228.
- Kürschner, E.** See *Emil Heuser*.
- Küster, William**, the prosthetic group of the blood pigments; hæmatin, A., i, 884.
bile pigments. XI. The preparation and purification of bilirubin from ox gall-stones, A., i, 885.
bile pigments. XII. The action of diazomethane on bilirubin and biliverdin, the oxidation of bilirubin in alkaline solution, and the action of hydrogen bromide-acetic acid on bilirubin, A., i, 885.
- Küster, William** [with *H. Maurer, W. Niemann, P. Schlaack, Schlayerbach, W. Weber, and Willig*], derivatives of pyrrole, A., i, 857.

Küster, William, and **Adolf Gerlach**, the prosthetic group of blood pigment; formoxyhæmin, A., i, 598.

Küster, William, and **Walter Herrmann**, bile pigments. XIII. Hexachlororubic acid, A., i, 886.

Kugelmann, I. Newton, a new apparatus; the nephelometer, A., ii, 692.

Kuhn, Alfred, swelling of gelatin in aqueous solutions of organic acids, A., i, 183.

Kuhn, Alfred. See also **Wolfgang Ostwald**.

Kuhn, Richard. See **Richard Willstätter**.

Kulp, W. L. See **R. J. Anderson**.

Kumagawa, H., the dismutation of various aldehydes by yeast, A., i, 805. the action of salts on the bleaching of methylene blue by various species of yeast, A., i, 306. decomposition of inositol and glycerol after the manner of true sugars by *Bacillus lactis aerogenes*, A., i, 972. production of the second and third forms of fermentation with *Saccharomyces sake*, *Zygosaccharomyces major*, and *Z. salvus*, A., i, 972.

Kunert, William, solubility of carbon dioxide and nitrous oxide in certain solvents, A., ii, 823.

Kunsman, C. H. See **C. Davisson**.

Kunz, Alphons. See **Géza Zemplén**.

Kunz, Jakob. See **E. H. Williams**.

Kunz, K. See **Paul Friedländer**.

Kunze, Rudolf. See **Paul Hirsch**.

Kuns-Krause, Hermann, occurrence of ellagic acid in *Rubus idæus*; the cause of the clouding of raspberry juice, A., i, 210.

Kuns-Krause, Hermann, and **Paul Manicke**, the action of mercurous formate on certain aliphatic halogen compounds, A., i, 6. elimination of carbon dioxide from organic compounds. VI. The action of the light of the electric arc on aqueous oxalic acid solutions in presence of ferric, chromic, and mercuric chloride, of uranyl acetate, and of iodic acid, A., ii, 731.

Kupfer, O. See **Hermann Staudinger**.

Kuppinger, O. See **Alexander Gutbier**.

Kurnakov, Nicolai S., and **G. Urasov**, the toxic properties of commercial ferrosilicon, A., ii, 845.

Kuroda, Chika. See **Rikô Majima**.

Kuroda, Sôzô. See **Jirô Takeda**.

Kurre, Bruno. See **Gustav F. Hüttig**.

Kurtenacker, Albin, and **Albert Fritsch**, the analysis of polythionates, A., ii, 521.

Kurtenacker, Albin, and **Josef Wagner**, the volumetric estimation of hydroxylamine and hydrazine, A., ii, 312.

Kurtenacker, Albin, and **Felicitas Werner**, the estimation of bismuth as metal, A., ii, 877.

Kurth, E. H., extension of the X-ray spectrum to the ultra-violet, A., ii, 410. soft X-rays of characteristic type, A., ii, 809.

Kurtz, S. S., jun. See **James Bryant Conant**.

Kuwada, Satoru See **Yasuhiko Asahina**.

Kyropoulos, S., metallographic investigations on the cathodic deposition of metals on aluminium and chromium, A., ii, 22.

L

L., purging nut-tree oil, A., i, 908.

Laar, Johannes Jacobus van, space-filling and the equation of condition, A., ii, 279.

Laar, R. van der. See **Friedrich Kehrman**.

La Barre, J. See **Edgard Zuns**.

Labat, A., and **M. Favreau**, chemical composition of amniotic fluid, A., i, 894.

Labbé, Henri, F. Nepveux, and **Ménelas Nomidis**, critical study of Bang's method for the estimation of reducing substances in blood, A., ii, 663.

Labes, Richard, increased velocity and intensity of action of groups of poisons or active drugs on bacteria and tadpoles produced by variation of the degree of acidity or alkalinity, A., i, 901. the beneficial action of charcoal suspensions and other substances with large surface development, such as colloidal silica, ferric phosphate, and agar-agar, on the formation of fermentation gases by *Bacillus coli* in protein-free nutrient media, A., i, 902.

Laborde, Léon Jaloustre, and **M. Leulier**, influence of radioactive substances on acetic fermentation, A., i, 1219.

Lacharte, detection of iodates in potassium iodide according to the French codex, 1908, A., ii, 716.

Lachman, Arthur, nitromalic acid, A., i, 107. dihydroxytartaric acid, A., i, 109. the rearrangement of benzil to benzilic acid, A., i, 459.

- Lachs, Hilary, and Stephanie Goldberg**, influence of temperature on the coagulation of colloidal gold, A., ii, 697.
- Lachs, Hilary, and Mathilde Wertenstein**, the distribution of radioactive substances in solutions, A., ii, 681.
- Ladenburg, Rudolf**, action of electrical fields on absorption lines (*D*-lines of sodium vapour), A., ii, 6.
effect of a strong electrical field on the absorption lines of sodium vapour, A., ii, 6.
- Ladenburg, Rudolf, and R. Minkowski**, chemical constants of sodium and potassium, A., ii, 191.
heat of vaporisation of sodium and the probability of the transition of the sodium atom from the resonance to the normal condition on the basis of optical measurements, A., ii, 194.
- Laer, Marc H. van**, action of hydrolytic enzymes. II., A., i, 64.
- Laer, Marc H. van, and R. Lombaers**, the formation of osazones, A., i, 113.
- Laidlaw, Patrick Playfair, and Wilfred Waller Payne**, estimation of small quantities of calcium, A., ii, 786.
- Laidlaw, Patrick Playfair**. See also *Charles Stanley Gibson*.
- Laing, (Miss) Mary Evelyn**. See *James William McBain*.
- Lal, Rattan**. See *Bawa Kartar Singh*.
- Lamb, Arthur Becket, Charles C. Scallione, and Graham Edgar**, preferential catalytic combustion of carbon monoxide in hydrogen, A., ii, 443.
- Lamb, Arthur Becket, and John P. Simmons**, heats of solution and of transformation of the acido- and aquo-cobalt pentammines, A., ii, 121.
- Lamb, Arthur Becket, and Victor Yngve**, strengths of cobaltamine bases and Werner's theory of bases, A., ii, 217.
- Lambourne, Herbert**, derivatives of methylstannonic acid; their bearing upon its constitution, T., 2533.
- Lambris, G.**, estimation of very small quantities of injurious acids in air, A., ii, 390.
- LaMer, Victor K.**, vitamins from the point of view of physical chemistry, A., i, 281.
- LaMer, Victor K., and Lillian E. Baker**, effect of substitution on the free energy of oxidation-reduction reactions. I. Benzoquinone derivatives, A., ii, 735.
- LaMer, Victor K., H. L. Campbell, and Henry Clapp Sherman**, the effect of temperature and the concentration of hydrogen ions on the rate of destruction of antiscorbutic vitamin (vitamin-C), A., ii, 275.
- LaMer, Victor K.** See also *Henry Clapp Sherman*.
- Landauer, Max**. See *Heinrich Lüers*.
- Landauer, Robert S.** See *Gerald L. Wendt*.
- Landau-Ziemecki, St.**, the emission spectrum of monatomic iodine vapour, A., ii, 674.
- Landergren, Sture**. See *Hans von Euler*.
- Landrien, Philippe**, a new class of active racemic substances, A., i, 808.
- Landrivon, J.** See *J. Altwegg*.
- Landsteiner, Karl**, heterogenetic antigen and haptens. XV., A., i, 195.
- Lang, Norbert**. See *Ernst Späth*.
- Lang, Rudolf**, a new iodometric method for the estimation of copper, A., ii, 318.
a new iodometric method based on the formation and the estimation of cyanogen iodide, A., ii, 782.
- Lange, Carl**, errors in the estimation of sugar by fermentation with yeast, A., ii, 93.
- Lange, H.** See *Karl von Auwers*.
- Lange, Hermann**, the influence of adrenaline on the permeability of the limiting membrane of muscle fibres, A., i, 791.
- Lange, Hermann, and Max Simon**, the liberation of phosphoric acid by the retina in the presence of light, A., i, 701.
- Langecker, Hedwig**, the practice of precipitation with lead, A., i, 315.
- Langhans, pyrofulmin**, a decomposition product of mercury fulminate, A., i, 328.
- Langley, Wilson D., and Roger Adams**, condensation of certain nitriles and various polyhydroxyphenols to form phenolic acids, A., i, 1153.
- Langmuir, Irving**, types of valency, A., ii, 137.
catalysis with special reference to newer theories of chemical action. II. Heterogeneous reactions; (1) chemical reactions on surfaces; (2) the mechanism of the catalytic action of platinum in the reactions $2CO + O_2 = 2CO_2$, and $2H_2 + O_2 = 2H_2O$, A., ii, 629.
- Langstein, Leo**, are the carbohydrate groups which appear in the acid hydrolysis of blood globulin a fission product of the protein molecule? A., i, 479.

- Langton H. M.**, saponification of oils and fats, A., i, 982.
- Lansberg, L. M.** See *H. J. Lemkes*.
- Lantz, K.** See *André Wahl*.
- Lapicque, L.**, and *L. Emerique*, variations in the chemical composition of Fucaceæ, A., i, 413.
- Lapkamp, K.** See *Rud. Seeliger*.
- Lapworth, Arthur**, a theoretical derivation of the principle of induced alternate polarities, T., 416.
- Lapworth, Arthur**, and *John Alexander McRae*, syntheses of alkylidenecyanoacetic acids and of substituted succinic acids. I. Acids containing aromatic residues, T., 1699.
- resorcinolphenylsuccinein, T., 2722.
- syntheses of alkylidenecyanoacetic acids and of substituted succinic acids. II. Preparation of acids containing saturated aliphatic residues and the constitution of the aliphatic alkylidenecyanoacetic esters, T., 2741.
- Lapworth, Arthur**, and *John Baldwin Shoesmith*, reciprocal induced polarity effects in cresols and their derivatives; properties of the isomeric methoxybenzyl bromides, T., 1391.
- Lapworth, Arthur**. See also *Robert Downs Haworth*, and (*Miss*) *Lucy Higginbotham*.
- Laquer, Fritz**, the degradation of carbohydrates in transversely striated muscles. I. and II., A., i, 298, 1089.
- a micro-extraction apparatus, A., ii, 325.
- Laqueur, Ernst**. See *J. Snapper*.
- Larson, Alfred T.**, and *Ernest C. White*, method of determining traces of oxygen in hydrogen, A., ii, 311.
- Larsonneau, A.** See *A. Goris*.
- Lasarev, P.**, investigations on the fundamental law of photochemistry, A., ii, 108.
- fundamental laws of photochemistry. II. Influence of cooling on the absorption of light by dyes, A., ii, 332.
- Laschtschenko, Porphyry Nicolaevitch**, polymorphism of arsenic, T., 972.
- Lasnitski, A.** See *Peter Rona*.
- Lassieur, Arnold**. See *André Kling*.
- Lassieur, (Mme) Arnold**. See *André Kling*.
- Lessmann, Max**. See *Hans Fringsheim*.
- Latiens, G.**, symmetrical di-iodoethyl-enes, A., i, 314.
- Latimer, Wendell M.**, distribution of thermal energy in the quadrivalent chlorides of carbon, silicon, titanium, and tin, A., ii, 256.
- Latimer, Wendell M.**, thermo-electric force, the entropy of electrons and the specific heat of metals at high temperatures, A., ii, 814.
- Latimer, Wendell M.** See also *Gilbert Newton Lewis*.
- Lau, E.** See *Ernst Gehrke*.
- Laurent, (Mlle) Y.** See *P. Freundler*.
- Lauterbach, Hans**. See *Erich Müller*.
- Lauth, Hilde**. See *Gustav Heller*.
- Lavers, Herbert**, tungsten, A., ii, 164.
- Lavoys**, use of resorcinol in qualitative inorganic analysis, A., ii, 779.
- Lawaczek, Heinz**. See *Gustav Embden*.
- La Wall, Charles H.** See *Heber Youngken*.
- Lawrance, Walter A.**, Friedel and Crafts' reaction; some substituted phthalic anhydrides with toluene and aluminium chloride, A., i, 340.
- Lawrance, Walter A.**, and *Harold G. Oddy*, Friedel and Crafts' reaction; diphenyl and ditolyl tetrahalogen phthalides, A., i, 455.
- Lax, E.** See *Marcello von Pirani*.
- Lax, Heinrich**, micro-estimation of total acetone in urine, A., ii, 326.
- Lax, Heinrich**. See also *Géza Petényi*.
- Leavenworth, Charles Samuel**. See *Thomas Burr Osborne*.
- Le Bas, Gervaise**, negative optical anomalies, A., ii, 9.
- relation between atomic volumes [of elements in combination] and optical refractivities, A., ii, 241.
- Lebeau, P.**, the oxides of uranium, A., ii, 302.
- Lebeau, P.**, and *M. Picon*, action of sodammonium on pyridine; preparation of the hydrate of tetrahydridipyridyl, A., i, 48.
- the action of sodammonium on hydrocarbons, A., i, 801.
- Le Brazide, Emilien**, some derivatives of anisylacetone (methoxyphenylpropanone) [*p*-methoxybenzyl methyl ketone], A., i, 456.
- Le Breton, (Mlle) Eliane**, presence and estimation in the total lipid ether-soluble phosphorus of phosphorus compounds other than phosphatides, A., ii, 168.
- Le Chatelier, Henri**, the manufacture of sodium carbonate by the ammonia process, A., ii, 375.
- the geometric representation of saline equilibria, A., ii, 555.
- Lecher, Hans**, and *Alfred Goebel*, disulphur dithiocyanate, A., i, 640.
- Lecher, Hans**, and *Josef Hofmann*, hydroxylamine. I. Simple method of preparation of free hydroxylamine, A., ii, 442.

- Lecher, Hans, and Kurt Simon**, the valency problem of sulphur. VII. *o*-Nitrophenyldithiochloride, A., i, 1013.
- Lecher, Hans, and Max Wittwer**, sulphur thiocyanate, A., i, 640.
alkylthiol thiocyanates. II., A., i, 641.
- Lechinaky, Wolfgang**. See **Max Bergmann**.
- Lecoq, Raoul**, the diastatic action of malt and its preparations (liquid, syrupy, and dried extracts), A., i, 312.
- Ledbury, Wilfrid**. See **Gilbert Thomas Morgan**.
- Lee, Henry, R.**, the estimation of *H*-acid, A., ii, 235.
- Lee, Henry R.**, and **De Witt O. Jones**, the analysis of β -naphthylamine, A., ii, 883.
- Lee, Henry R.** See also **De Witt O. Jones**.
- Lee, Olive P.**, and **Shiro Tashiro**, alkaligenesis. II. Ammonia production in muscle, A., i, 1089.
- Leerdertz, G.**, and **B. Gromelski**, the estimation of fibrinogen; protein estimations in salt plasma; question of the utility of serum for the quantitative investigation of blood, A., ii, 798.
- Legg, David Alliston**. See **Matthew Atkinson Adam**.
- Le Heux, J. W.**, choline as hormone for intestinal movement. III. Participation of choline in the action of various organic acids on the intestine. IV. Effect of choline on normal gastric movement, A., i, 85.
- Lehman, Edwin, P.**, inorganic blood phosphate, A., i, 191.
- Lehmann, Fritz**. See **Ernst Beekmann**.
- Lehmann, Jorgen**, the inhibition of blood coagulation by barium, strontium, and calcium chlorides, A., i, 892.
- Lehmann, Otto**, structure of [hanging] drops of mixed liquid crystals, A., ii, 201.
states of aggregation and liquid crystals, A., ii, 692.
- Lehner, Friedrich**, a test for pyridine, A., ii, 795.
- Lehrman, S.** See **W. H. Martin**.
- Leichtentritt, Bruno**, and **Margarete Zielskowsky**, the growth-promoting factor of lemon juice, A., i, 1094.
- Leimbach, G.** See **Friedrich L. Hahn**.
- Leitch, (Miss) Grace Cumming**. See **Walter Norman Haworth**.
- Lejeune, B.** See **Georges Chavanne**.
- Lemay, Pierre, and Léon Jaloustre**, the oxidising properties of certain radioactive elements, A., ii, 18.
- Lemay, Pierre, and Léon Jaloustre**, some oxidising properties of thorium-X, A., ii, 186.
- Lemeland, P.**, chemical and physiological investigation of the fats and lipoids of the blood. II. The Kumagawa-Suto method of estimation of lipoids, A., ii, 686.
- Lemkes, H. J.**, and **L. M. Lensberg**, Causse-Bonnan's method for the estimation of dextrose, A., ii, 724.
- Lemmermann, Otto, and Ludwig Fresenius**, soil acidity and its effect on germinating plants, A., i, 510.
- Lemmermann, Otto, and H. Wiessmann**, effect of silicic acid on crop production in the presence of insufficient amounts of phosphoric acid, A., i, 1103.
- Le Moal**. See **Warcollier**.
- Lenher, Victor**, some properties of selenium oxychloride. II., A., ii, 706.
selenium oxybromide, A., ii, 707.
- Lenher, Victor, Gilbert B. L. Smith, and George G. Town**, vapour pressure of selenium oxychloride, A., ii, 371.
- Lenher, Victor, Homer W. Stone, and Helen H. Skinner**, formation of potassium perchlorate from potassium chlorate, A., ii, 237.
- Lenher, Victor, and Martin Tosterud**, rapid analysis of potassium perchlorate, A., ii, 395.
- Leonard, Clifford Shattuck**, some compounds of piperidine with haloids, A., i, 362.
diethylrhodamine, A., i, 1182.
- Leonard, Helen A.** See **Burt E. Nelson**.
- Leone, P.**, and **E. Angelescu**, oil of *Satureja montana* of Italian origin, A., i, 357.
oil of *Thymus vulgaris* of Italian origin, A., i, 357.
Italian oil of *Thymus striatus*, A., i, 460.
variations in the solubility of a compound in presence of other compounds. I. Water-phenol-diphenols, A., ii, 743.
- Leone, P.**, and **M. Benelli**, variations in the solubility of a compound in presence of other compounds. II. Water-epichlorohydrin-acetic acid, A., ii, 744.
- Leopold, Ludwig**. See **Georg Sachs**.
- Lepape, Adolphe**. See **Charles Moureu**.
- Lepahne**, tests for liver function, A., ii, 800.
- Lepeschkin, W. W.**, the coagulation of proteins by heat, A., i, 1198.
- Lepin, A. L.**, interaction of carbon tetrachloride and fuming sulphuric acid, A., ii, 847.

- Lepin, Lidie.** See *Nikolai Schilov*.
- Lerolde, J.**, some properties of $\alpha\alpha$ -disubstituted esters, A., i, 215.
- Lespieau, Robert**, derivatives of acetylenic erythritol [hexinene- $\alpha\beta\epsilon\zeta$ -tetrol], $\text{HO}\cdot\text{CH}_2\cdot\text{CH}(\text{OH})\cdot\text{C}\equiv\text{C}\cdot\text{CH}(\text{OH})\cdot\text{CH}_2\cdot\text{OH}$, A., i, 108.
action of dichloroethyl ether, $\text{CH}_2\text{Cl}\cdot\text{CHCl}\cdot\text{OEt}$, on the mixed magnesium derivative of allyl bromide, A., i, 619.
- Lesser, Ernst J.**, the spacial separation of glycogen and diastase in the liver cells, A., i, 195.
- Leuchs, Hermann, Eva Conrad, and Hans von Katinusky**, spirans. IX. Preparation of bishydrocarbostryril-3:3-spiran and its by-products, A., i, 471.
spirans. X. Proof of the peculiar spiran asymmetry by the preparation of an optically active spiran, A., i, 873.
- Leuchs, Hermann, and Kurt Fricker**, strychnos alkaloids. XXXII. Transformations of the quinones from brucinesulphonic acid. I., A., i, 677.
- Leuchs, Hermann, and Hans Kaehrn**, strychnos alkaloids. XXXI. Violet and green colour reactions of cacotheline, A., i, 463.
- Leuchs, Hermann, Hans Mildbrand, and W. Robert Leuchs**, strychnos alkaloids. XXXIII. The degradation of cacotheline by bromine, A., i, 1052.
- Leuchs, Hermann, and Rudolf Nitschke**, strychnos alkaloids. XXXIV. The preparation of isostrychnine, A., i, 1175.
- Leuchs, Hermann, Fritz Osterburg, and Hans Kaehrn**, strychnos alkaloids. XXX. Reactions of cacotheline, A., i, 362.
- Leuchs, W. Robert.** See *Hermann Leuchs*.
- Leue, H.** See *Karl Fries*.
- Léulier, M.** See *Laborde*.
- Levaditi, C.** See *Robert Saxerac*.
- Levart-Ezersky, M.**, thermochemistry of solutions, A., ii, 819.
- Levene, Phœbus A.**, synthesis of 2-hexosamic acids and 2-hexosamines, A., i, 327.
ethyl benzylidenechitosamate and ethyl diazobenzylidengluconate (mannonate), A., i, 1028.
preparation and analysis of animal nucleic acid, A., i, 1075.
- Levene, Phœbus A., and Gustave Morris Meyer** [with *I. Weber*], phosphoric esters of some substituted glucoses and their rate of hydrolysis, A., i, 987.
- Levene, Phœbus A., and Gustave Morris Meyer** [with *I. Weber*], sulphuric esters of some substituted glucoses and their rates of hydrolysis, A., i, 987.
- Levene, Phœbus A., and Louis A. Mikeska**, possible asymmetry of aliphatic diazo-compounds. II., A., i, 818.
- Levene, Phœbus A., and Ida P. Rolf**, the unsaturated fatty acids of egg lecithin, A., i, 621.
- Levene, Phœbus A., and H. S. Simms**, the unsaturated fatty acids of liver lecithin, A., i, 424.
- Levene, Phœbus A., and F. A. Taylor**, synthesis of α -hydroxyisopentacosic acid and its bearing on the structure of cerebronic acid, A., i, 714.
- Lévêque, A.**, estimation of the alcohol content of a solution, A., ii, 723.
- Levi, Giorgio Renato**, chlorites of ammonium, tetramethylammonium, and certain amines, A., i, 527.
chlorites of sodium and other metals, A., ii, 567.
oxidation and reduction reactions with chlorites, A., ii, 567.
- Levi, Mario Giacomo**, formation of hydrocarbons from carbon monoxide and from formates, A., i, 801.
- Levine, M.**, bacteria fermenting lactose, and their significance in water analysis, A., i, 901.
- Levine, Victor E., Elmer Verner McCollum, and Nina Simmonds**, glacial acetic acid as a solvent for the antineuritic substance, water-soluble B, A., i, 975.
- Levinstein, Herbert.** See *British Dyestuffs Corporation, Ltd.*
- Lewcock, William, William Gordon Adam, Norman Edward Siderfin, and William Lyle Galbraith**, preparation of amino-phenols or aromatic amino-acids, A., i, 650.
- Lewin, L., and E. Stenger**, spectrographic detection of change in blood pigment due to certain organic and inorganic poisons, and the characteristic spectra of yolk of egg and urobilin, A., ii, 414.
- Lewis, Edgar**, the composition of the residue on distillation of crude glycerin, A., i, 419.
- Lewis, E. P.**, the continuous spectrum of hydrogen in the Schumann region, A., ii, 802.
- Lewis, Gilbert Newton, and Wendell M. Latimer**, revision of the entropies of the elements, A., ii, 471.

- Lewis, Howard B., and Daniel A. McGinty** [with *Lucie E. Root*], metabolism of sulphur. V. Cysteine as an intermediary product in the metabolism of cystine, A., i, 1088.
- Lewis, Howard B., and Lucie E. Root**, metabolism of sulphur. IV. The oxidation of cystine in the animal organism, A., i, 487.
- Lewis, Howard B.** See also *Max S. Dunn*.
- Lewis, J. Volney, and Lawson H. Bauer**, cyprine and associated minerals from Franklin, New Jersey, A., ii, 714.
- Lewis, J. W.** See *Thomas Barratt*.
- Lewis, Samuel Judd**, the ultra-violet absorption spectra and the optical rotation of the proteins of blood sera, A., ii, 245.
- the quantitative determination of the fluorescent powers (the spectro-fluorescometry) of cellulose, sugars, and other substances, A., ii, 334.
- Lewis, Samuel Judd, and Florence Mary Wood**, a new adjustable thermostat for all temperatures between 0° and 100°, A., ii, 612.
- Lewis, William Cudmore McCullagh**, radiation and chemical action, A. ii, 336.
- catalysis with special reference to newer theories of chemical action. I. The radiation theory of chemical action. (2) The radiation hypothesis of chemical reactivity and some of its applications, A., ii, 628.
- Lewis, William Cudmore McCullagh.** See also *J. W. Corran, Thomas Moran, and Henry Austin Taylor*.
- Lewis, W. Lee, and H. C. Cheetham**, arsenated benzophenone and its derivatives, A., i, 187.
- Lewis, W. Lee, and Cliff S. Hamilton**, 7-chloro-7:12-dihydro- γ -benzophenarsazine and some of its derivatives, A., i, 187.
- Lewy, F. H.** See *K. Dresel*.
- Lexow, Thor**, sea-wolf liver oil, A., i, 195.
- oil from the liver of *Acanthias vulgaris*, A., i, 490.
- Ley, Heinrich, and R. Grau**, complex tautomerism, A., i, 536.
- Ley, P.** See *Eberhard Rimbach*.
- Léys, Alexandre**, a rapid method for determining the acetyl value of oils and fats, A., ii, 167.
- Leyser, Felix.** See *Arthur Rosenheim*.
- Lieb, Hans.** See *Josef Herrig*.
- Lieben, Fritz**, the destruction of lactic acid by yeast-cells, A., i, 796.
- Lieben, Fritz**, behaviour of some amino-acids towards oxygenated yeast, A., i, 1219.
- Lieben, Fritz.** See also *Otto von Fürth*.
- Liebermann, Ludwig.** See *Carl Neuberg*.
- Liempt, J. A. M. van**, the green colour of tungsten trioxide, A., ii, 73.
- the equilibrium of tungsten and its oxides with hydrogen and water vapour; carbon monoxide and carbon dioxide and oxygen, A., ii, 301.
- the binary systems, $\text{Na}_2\text{SiO}_3\text{--Na}_2\text{WO}_4$, $\text{K}_2\text{SiO}_3\text{--K}_2\text{WO}_4$, and $\text{Na}_2\text{WO}_4\text{--K}_2\text{WO}_4$, A., ii, 775.
- a new microchemical method for the identification of tungsten, A., ii, 787.
- Liesche, Otto.** See *Ernst Beckmann*.
- Liesegang, Raphael Ed.**, a reversible turbidity phenomenon, A., ii, 369.
- Liévin, O.**, kinetic study of alkaline solutions of iodine, A., ii, 359.
- Lifschitz, Israel**, the formation of salts from aromatic nitro-compounds, A., i, 737.
- Lifschitz, Israel, and Georg Beck**, optics of disperse systems. V., A., ii, 597.
- Lifschitz, Israel, and B. B. Hepner**, colour isomerism and salt formation with iminoviolic acids. II., A., i, 767.
- Lifschütz, Isaac**, cholesterol dibromide, A., i, 251.
- the degradation products of cholesterol in the animal organs; (substances associated with cholesterol in the blood), A., i, 392.
- the chemical nature and the transformation of the fat of the blood, A., i, 392.
- metacholesterol and its by-products. III., A., i, 541.
- Light, Louis, and Frank Lee Pyman**, bromo-derivatives of 2-methylglucose, T., 2626.
- Limann, G.**, visible refraction and dispersion in aqueous solutions of salts, particularly coloured salts, A., ii, 173.
- Lind, Samuel Colville.** See *R. E. Nyswander*.
- Lindahl, Gunnar.** See *Erik Matteo Prochet Widmark*.
- Lindberg, Ernst**, activators of fermentation, A., i, 1219.
- Linderström-Lang, K.** See *Sören Peter Lauritz Sörensen*.
- Lindgren, Waldemar**, melanovanadite, a new mineral from Peru, A., ii, 155.

- Lindgren, Waldemar**, **Leicester F. Hamilton**, and **Charles Palache**, melano-vanadite, a new mineral from Peru, A., ii, 305.
- Lindh, Axel E.**, Röntgen absorption spectrum of chlorine, A., ii, 177.
the absorption spectrum of sulphur for X-rays, A., ii, 542.
- Lindh, Axel E.** See also **Arne Westgren**.
- Lindhard, J.**, colorimetric estimation of the concentration of hydrogen ions in very small quantities of blood by dialysis, A., ii, 240.
- Lindner, (Fr.).** See **Max Bodenstein**.
- Lindner, Egon.** See **Wacław Morawski**.
- Lindner, Josef**, the course of the quinaldine synthesis with 6-aminotetrahydroquinoline and 6-aminokairoline, A., i, 687.
volumetric estimation of carbon and hydrogen in organic compounds, A., ii, 657.
- Lindner, Kurt** [with **Emma Haller**, **Herbert Helwig**, **Antonie Köhler**, and **Hellmuth Feit**], the chlorides of bivalent molybdenum, tungsten, and tantalum. I., A., ii, 509.
- Lindner, Paul.** See **Gustav Heller**.
- Lindsay, the limits of absorption L** of the elements barium to antimony, A., ii, 599.
- Lineberry, R. A.** See **Francis P. Venable**.
- Ling, Arthur Robert** [with **E. H. Callow**, and **W. J. Price**], estimation of starch. I. Estimation of starch in barley and in wheat, A., ii, 879.
- Ling, Arthur Robert**, and **John Herbert Bushill**, the estimation of calcium in blood, A., ii, 587.
- Ling, Arthur Robert**, and **Dinshaw Rattongi Nanji**, crystalline glucose-ammonia and isoglucosamine, T., 1682.
the longevity of certain species of yeast, A., i, 92.
new method of preparing gluconic acid, A., i, 221.
the action of ammonia and of amino-compounds on reducing sugars. I. The action of ammonia on dextrose and levulose, A., i, 631.
the synthesis of glycine from formaldehyde, A., i, 1124.
- Ling, Arthur Robert**, and **William John Price**, a "micro-Kjeldahl" method of estimating nitrogen, A., ii, 522.
- Lingen, J. Steph. van der**, fluorescence of cadmium vapour, A., ii, 181.
- Lingen, J. Steph. van der**, the fluorescence of mercury vapour excited by Röntgen rays, A., ii, 548.
- Lingen, J. Steph. van der**, and **Robert Williams Wood**, the fluorescence of mercury vapour, A., ii, 245.
- Linhart, George Augustus**, relation between entropy and probability; integration of the entropy equation, A., ii, 257, 738.
- Linstead, Reginald Patrick.** See **Kenneth Claude Devereux Hickman**.
- Liot, A.** See **A. Goris**.
- Liotta, Domenico**, modification in the process of extraction of alkaloids, A., ii, 404.
- Lipman, Jacob Goodale**, **Selman A. Waksman**, and **Jacob S. Joffe**, the oxidation of sulphur by soil organisms, A., i, 303.
- Lipp, Peter**, comparison of α -campholanic acid with Mahla- and Tiemann's dihydrocampholenic acid and with isocampholic acid, A., i, 735.
- Lippmann, Edmund Oskar von**, the history of [ethyl] alcohol, A., i, 102.
botanical chemical observations, A., i, 311, 1223.
- Lipschitz, Werner**, and **A. Gottschalk**, reduction of the aromatic nitro-group as indicator of partial processes of respiration and of fermentation; a method for the comparative estimation of biological oxido-reduction. I. Experiments with respiring cells. II. Experiments with fermenting cells, A., i, 298.
- Listmann, F.** See **Fritz Mayer**.
- Liversedge, Samuel G.**, and **F. W. Andrews**, rapid estimation of quinine salts, A., ii, 669.
- Livingston, Marguerite B.** See **David I. Macht**.
- Lizius, John Leonard**, and **Norman Evers**, titration of acids and bases, A., ii, 654.
- Lloyd, Dorothy Jordan**, some properties of dialysed gelatin, A., i, 1075.
- Lloyd, Dorothy Jordan**, and **Charles Mayes**, the titration curve of gelatin, A., i, 280.
- Locher, F.** See **Paul Karrer**.
- Lochte, Harry L.**, **James K. Bailey**, and **William Albert Noyes**, symmetrical diisopropylhydrazine and its derivatives, A., i, 329.
- Lockemann, G.** See **Paul Jacobsen**.
- Lockemann, Georg**, rotary burner with fixed gas connexion, A., ii, 43.
- Locket, George Hazlewood.** See **Dalziel Llewellyn Hammick**.

- Lockrow, Laurice L.**, the effect of oxygen and hydrogen on the emission of electrons from hot platinum, A., ii, 608.
- Loequis, René, and Sung Wouseng**, the action of acetylene on the sodium derivatives of ketones and the preparation of dialkylethynyl-carbinols, A., i, 617.
the preparation of dialkylvinyl-carbinols, A., i, 710.
transformation of tertiary ethylenic alcohols (linalool type) into primary ethylenic alcohols (geraniol type), A., i, 710.
the preparation of aldehydes from tertiary alcohols, A., i, 810.
- Lodge, (Sir) Oliver**, Bohr and Langmuir atoms, A., ii, 701.
speculations concerning the positive electron, A., ii, 836.
- Loeb, A.** See **Paul Jacobsen**.
- Loeb, Jacques**, the significance of the isoelectric point for the preparation of ash-free gelatin, A., i, 387.
discrepancies between the observed and calculated potential difference of protein solutions near the isoelectric point, A., i, 692.
ionising influence of salts with tervalent and quadrivalent ions on crystalline egg-albumin at the isoelectric point, A., i, 881.
the mechanism by which tervalent and quadrivalent ions produce an electrical charge on isoelectric protein, A., i, 882.
the influence of aggregates on the membrane potentials and osmotic pressure of protein solutions, A., i, 883.
the origin of the potential differences responsible for anomalous osmosis, A., ii, 124.
electrical charges of colloidal particles and anomalous osmosis, A., ii, 354.
electrical charges of colloidal particles and anomalous osmosis. II. Influence of the radius of the ion, A., ii, 467.
cataphoretic charges of collodion particles and anomalous osmosis through collodion membranes free from protein, A., ii, 742.
interpretation of the influence of acid on the osmotic pressure of protein solutions, A., ii, 742.
the influence of electrolytes on the cataphoretic charge of colloidal particles and the stability of their suspensions. I. Experiments with collodion particles, A., ii, 746.
- Loeb, Jacques, and Robert F. Loeb**, the influence of electrolytes on the solution and precipitation of casein and gelatin, A., i, 182.
- Loeb, Leonard B., and Lloyd Schmiedekamp**, destruction of phosphorescent zinc sulphides by ultra-violet light, A., ii, 8.
- Loeb, L. Farmer**, caffeine concentration of the blood and urine of rabbits after parenteral administration, A., i, 705.
- Loeb, Robert F.** See **Jacques Loeb**, and **Walter W. Palmer**.
- Löffelbein, Willi**, estimation of chromium in metals, A., ii, 680.
- Löfmann, Nils.** See **Erik Hägglund**.
- Loening, E.** See **Herbert Freundlich**.
- Löpmann, Bernhard.** See **Erwin Ott**.
- Lösch, Joseph.** See **Wilhelm Prandtl**.
- Lösenbeck, Otto**, electrical properties of silicic acid sols, A., ii, 695.
- Lövgren, Sture**, urease, A., i, 185.
- Löw, Artur.** See **Paul Dux**.
- Löwe, Hans.** See **Franz Fichter**.
- Loewe, S.**, physical chemistry of lipoids; diffusion of methylene-blue through organic solvents, A., ii, 354.
- Loewi, Otto.** See **E. Geiger**.
- Lohmann, W.** See **Karl Fries**.
- Lohsner, Artur.** See **Jakob Meisenheimer**.
- Loisel, P.**, the existence of a new radioactive emanation in the springs of Bagnoles-de-l'Orne and its surroundings, A., ii, 15.
- Lombaers, R.** See **Marc H. van Laer**.
- Lombard, Maurice**, the action of nitrous acid on iodides in the presence of oxygen, A., ii, 313.
- Lombroso, Ugo**, action of hydrochloric acid on the fat exchange in the surviving liver, A., i, 702.
- Lommen, Frederick W. M.** See **Morris S. Kharasch**.
- Long, Esmond R.**, chemical problems in the bacteriology of tubercule bacillus, A., i, 1095.
- Long, M. Louisa.** See **Norman R. Blatherwick**.
- Longchambon, Henri**, spectral study of the triboluminescence of sucrose, A., ii, 542.
- Longchambon, Louis**, the rotatory power of crystals and molecular rotatory power, A., ii, 603.
- Locker, Cloyd D.** See **William Lloyd Evans**.
- Loomis, Albert G.**, system ammonia-magnesium-mercury; formation of magnesium hexammoniate, A., ii, 294.

- Loon, Chr. van.** See **Jacob Böese-ken.**
- Looney, Joseph M.** See **Otto Felin.**
- Lorens, Richard,** the theory of electrolytic ions. XXVI. How is the limiting value of the molecular conductivity of strong electrolytes determined? A., ii, 19.
- kation volumes in permutite, A., ii, 196.
- radii of the alkali metals and the alkali metal ions, A., ii, 211.
- Lorens, Richard, Walter Fraenkel, and M. Wormser,** electrochemical investigations of gold-copper mixed crystals, A., ii, 21.
- Lorens, Richard, and Gustav Haegermann,** the discovery of an equilibrium between cement and lime-water, A., ii, 59.
- Lorens, Richard, and Walter Herz,** surface tension of corresponding states, A., ii, 261.
- the relation of melting point to boiling point, A., ii, 739.
- Lormand, Ch.** See **Fayolle, and Maurice François.**
- Losana, Luigi,** colorimetric estimation of phosphorus, A., ii, 392.
- rapid estimation of sulphur, A., ii, 582, 656.
- estimation of selenium, A., ii, 865.
- Losana, Luigi, and Enrico Carossi,** estimation of chromium in steels, A., ii, 589.
- rapid analysis of ferro-tungsten, A., ii, 661.
- Losanitch, Sima M.,** decomposition of dithiocarbazates, T., 2542.
- Lottermoser, Alfred,** technical estimation of the colloidal part of tungsten powder, A., ii, 230.
- freezing of hydrosols, A., ii, 266.
- electrolytic removal of alkali from salts, A., ii, 510.
- Lottermoser, Alfred, and Kurt Falk,** chromate electrolysis with diaphragms, A., ii, 736.
- Lottermoser, Alfred, and Max Steude,** constitution of starch iodide, A., i, 10.
- Lots, Paul, and Joseph Christie Whitney Fraser,** osmotic pressures of concentrated solutions of sucrose as determined by the water interferometer, A., ii, 264.
- Lots, Paul.** See also **L. van Doren.**
- Lownes, Ashley Gordon,** abnormal crystallisation of lead azide by protective colloids, A., ii, 216.
- Lowry, Thomas Martin,** catalysis with special reference to newer theories of chemical action. I. The radiation theory of chemical action. (4) Is a true unimolecular action possible? A., ii, 628.
- Lowry, Thomas Martin, and Percy Corlett Austin,** optical rotatory dispersion. II. Tartaric acid and the tartrates, A., ii, 414.
- Lowry, Thomas Martin, and John Outram Cutter,** the rotatory dispersive power of organic compounds. X. The preparation and properties of pure ethyl tartrate, T., 532.
- Lowry, Thomas Martin, Victor Steele, and Henry Burgess,** new halogen derivatives of camphor. II. α' -Bromocamphor, T., 633.
- Lowry, Thomas Martin.** See also **Reginald George Early, and (Miss) Ida L. Millican.**
- Lowy, Alexander, and Wilmer Baldwin,** derivatives of 2:4:6-trinitrobenzaldehyde. II., A., i, 134.
- Lowy, Alexander, and Raymond F. Dunbrook,** compounds of tellurium tetrabromide with organic bases, A., i, 446.
- Lublin, Alfred,** Ambard's urea constant, A., i, 400.
- Lubowsky, K.,** emissivity of iron and copper, A., ii, 108.
- Lubricht, Bernh.** See **Franz Feist.**
- Lucas, Howard J., and Archie R. Kemp,** chromo-isomeric silver salts of pentabromophenol, and a theory of chromo-isomerism of solid compounds, A., i, 30.
- Lucasse, Walter W.** See **Charles August Kraus.**
- Luce, Emile, and A. Doucet,** the method of estimating allylthiocarbimide in mustard flour, A., ii, 537.
- Luce, Emile.** See also **Maurice François.**
- Luckmann, Viktoria.** See **Hans Fischer.**
- Ludlam, Ernest Bowman,** an attempt to separate the isotopes of chlorine, A., ii, 497.
- Ludwig, Alois,** the separation of zinc from other metals, especially nickel, A., ii, 786.
- Ludwig, Ernst.** See **Hermann Pauly.**
- Ludwig, Richard.** See **Ernst Welts.**
- Lübke, Alfred.** See **Erich Schmidt.**
- Lück, E. A.,** derivatives of *l*-menthol, A., i, 44.
- Lueck, Roger H.,** thermal decomposition of nitrogen pentoxide in solution, A., ii, 433.
- Lücke, Th.** See **Gerhard Carl Schmidt.**
- Lüdermann, Otto.** See **Erwin Ott.**

- Lüders, Hermann.** See *Adolf Windaus*.
Lüers, Heinrich, and *Max Landauer*, isoelectric point of the vegetable albumin leucosin, A., i, 881.
 the kinetics of the coagulation of proteins by heat, A., ii, 697.
Lührig, Heinrich, use of semi-microchemical and microchemical methods in place of macrochemical methods in the analysis of foods, A., ii, 534.
Lüscher, Ery, the nitrogen distribution in Bence Jones protein; a new colorimetric method for the estimation of tryptophan in protein, A., i, 1199.
Lüscher, G. See *Hermann Staudinger*.
Lüttichau, A., the action of heterogeneous proteins in the organism, A., i, 892.
Lüttringhaus, A., and *L. Eiffaender*, arylaminoanthraquinone derivatives, A., i, 355.
Luff, Bernard Dunstan Wilkinson, and *Benjamin Dawson Porritt*, estimation of available sulphur in golden sulphide of antimony, A., ii, 225.
Luff, G., the separation of zinc by means of ammonium phosphate, A., ii, 456.
Luger, Alfred, the action of metallic copper and silver on diastase; the so-called oligodynamic phenomena, A., i, 65.
 quinine hæmolytic, A., i, 81.
Lund, Hakon. See *Einar Billmann*.
Lund, Yappa. See *Joseph E. Greaves*.
Lundegårdh, Henrik, new apparatus for estimation of carbon dioxide in air, A., ii, 719.
Lundell, Gustav Ernst Fred, and *H. B. Knowles*, modified method for the estimation of iron and vanadium after reduction by hydrogen sulphide, A., ii, 88.
Lundin, Harry, the proteolytic enzymes of malt, A., i, 959.
Lundsgaard, Christen, and *Eggert Möller*, the estimation of the total oxygen-combining power of the blood in the van Slyke apparatus, A., ii, 724.
Lundstrum, F. O. See *R. O. E. Davis*.
Lunelund, Harald, absorption of light by solutions, A., ii, 8.
Lush, Ernest Joseph, preparation of formaldehyde or its polymerides from mixtures of carbon monoxide and hydrogen, A., i, 625.
Lusk, Graham [with *James Evenden*], animal calorimetry. XVIII. The behaviour of various intermediary metabolites on the heat production, A., i, 292.
Lusk, Graham. See also *H. V. Atkinson*.
Lutz, G. See *H. Kessler*.
Lutz, Oscar, sensitiveness and applicability of qualitative reactions. III. Strontium ions, A., ii, 227.
Lyle, William Gordon. See *William E. Caldwell*.
Lyman, J. F. See *D. E. Haley*.
Lyman, Theodore, the spectrum of helium in the extreme ultra-violet, A., ii, 674.
Lynch, D. F. J., substituted naphthalenesulphonic acids. I. A method for identifying H-acid and its intermediates obtained from naphthalene-2 : 7-disulphonic acid, A., ii, 881.
Lynn, E. V., and *O. Hilton*, the action of nitrosyl chloride on normal heptane, A., i, 417.

M.

- Maas, Otto,** and *E. H. Boomer*, vapour densities at low pressures and over an extended temperature range. I. Properties of ethylene oxide compared to oxygen compounds of similar molecular weight, A., i, 912.
Macallum, Archibald Byron, the cobalt nitrite reaction for potassium in animal and vegetable cells, A., ii, 869.
McAulay, A. L., recoil of hydrogen nuclei from swift α -particles, A., ii, 12.
Macaulay, Robert Milroy, the reaction between iodine and sulphurous acid, T., 552.
McBain, James William, and *Arthur John Burnett*, the effect of an electrolyte on solutions of pure soap, phase-rule equilibria in the system sodium laurate-sodium chloride-water, T., 1320.
McBain, James William, and *William Job Jenkins*, the ultra-filtration of soap solutions; sodium oleate and potassium laurate, T., 2325.
McBain, James William, (*Miss*) *Milliecent Taylor*, and (*Miss*) *Mary Evelyn Laing*, studies of the constitution of soap solutions; solutions of sodium palmitate, and the effect of excess of palmitic acid or sodium hydroxide, T., 621.
McBain, James William. See also (*Miss*) *Mabel Harriet Norris*.
Macbeth, Alexander Killen, the labile nature of the halogen atom in organic compounds. IV. The tautomeric hydrogen hypothesis, and the removal of the halogen atom from aromatic nitro-compounds, T., 1116.

- Macbeth, Alexander Killen**, and **John Pryde**, studies of the glucosides. I. The constitution of indican, T., 1660.
- Macbeth, Alexander Killen**, and **Robert Robinson**, cevadine. I., T., 1571.
- Macbeth, Alexander Killen**. See also **Ian Armstrong Black**, **Hugh Graham**, **Thomas Henderson**, and **Edmund Langley Hirst**.
- McCallie, S. W.**, the Pitts meteorite, A., ii, 306.
- McCallum, S. T.**, preparation of alcoholic potassium hydroxide volumetric solution, A., ii, 54.
- McCann, Wm. S.**, the protein requirement in tuberculosis, A., i, 497.
- McCay, LeRoy Wiley**, and **William T. Anderson, jun.**, reduction of solutions of ferric salts with mercury, A., ii, 217.
reduction of vanadic acid solutions with mercury, A., ii, 530.
- McClelland, Ernest Wilson**. See **Samuel Smiles**.
- McClure, William B.**, adaptation of the pentabromacetone method to the estimation of citric acid in urine, A., ii, 791.
- McCluskey, K. Lucille**, some new derivatives of 2-methylquinoline, A., i, 864.
- McCombie, Hamilton**, **Harold Archibald Scarborough**, and **Richard Hardcastle Settle**, the velocity of reaction in mixed solvents. III. The influence of temperature on the velocity of saponification of esters, T., 2308.
- McCombie, Hamilton**. See also **Albert Eric Cashmore**.
- McCollum, Elmer Verner**. See **Victor E. Levine**, and **C. R. Orton**.
- McCrosky, C. R.** See **Harold D. Buell**, and **Samuel Wilson**.
- McCrudden, Francis H.**, pharmacological and chemical study of the roes of the barbel and pike, A., i, 194.
- McDavid, James Wallace**, heat developed on mixing sulphuric acid, nitric acid, and water, A., ii, 617.
- MacDonald, Alexander D.** See **James Bryant Conant**.
- Macdonald, George William**, obituary notice of, T., 2913.
- MacDougall, D. T.**, the action of bases and salts on biocolloids and cell masses, A., i, 204.
- MacDougall, Frank Henry**. See **Paul Francis Sharp**.
- McEwen, Joseph L.** See **James Munsu. Bell**.
- McGill, William J.**, use of the newer indicators in titrations of alkaloids, A., ii, 885.
- McGinty, Daniel A.** See **Howard B. Lewis**.
- McGuigan, Hugh**, an explanation of Liesegang's rings, A., ii, 38.
- Mach, Felix**, and **F. Sindlinger**, estimation of total nitrogen in fertilisers containing nitrites and of nitrite nitrogen in the presence of nitrates, A., ii, 783.
- McHargue, James Spencer**, the rôle of manganese in plants, A., i, 906.
- Macheleidt**, volumetric estimation of potassium, A., ii, 227.
- Macht, David I.**, pharmacological examination of isopropyl alcohol, A., i, 1093.
- Macht, David I.**, and **Marguerite B. Livingston**, effect of cocaine on the growth of *Lupinus albus*, A., i, 798.
- McHugh, Gerald Patrick**. See **Oscar Lisle Brady**.
- Mellvaine, T. C.**, a buffer solution for colorimetric comparison, A., ii, 78.
- MacInnes, Duncan A.**, and **Eric B. Townsend**, an electro-volumetric method for lead, A., ii, 456.
- MacInnes, Duncan A.**, and **Yu Liang Yeh**, potentials at the junctions of univalent chloride solutions, A., ii, 252.
- MacInnes, Duncan A.** See also **William K. Hainsworth**.
- McIntosh, James**. See **Ernest Laurence Kennaway**.
- Macjulevitch, K.** See **E. Fritsmann**.
- Mackall, Colin M.** See **Ebenezer Emmet Beid**.
- McKee, Ralph H.**, and **George Barsky**, ligninsulphonic acids, A., i, 1121.
- McKeehan, Louis W.**, crystal structure of potassium, A., ii, 709.
the crystal structure of glucinum and glucinum oxide, A., ii, 766.
- McKelvy, Ernest C.**, and **Daniel H. Simpson**, equilibria in the systems, carbon disulphide-methyl alcohol and carbon disulphide-ethyl alcohol, A., ii, 271.
- McKenzie, Alexander**, and (**Miss**) **Isobel Agnes Smith**, catalytic racemisation of optically active acid amides, T., 1348.
- McKenzie, Alexander**, and (**Miss**) **Nellie Walker**, optical activation of racemic acid by *l*-malic acid, T., 349.
- Mackenzie, John Edwin**, the action of sodium methoxide and its homologues on benzophenone chloride and benzylidene chloride. III., T., 1695.
- McKeown, Andrew**, the potential of the iodine electrode and the activity of the iodide ion at 25°, A., ii, 417.

- McKeown, Andrew**, influence of electrolytes on the solubility of non-electrolytes, A., ii, 552.
- McLaughlin, George D.** See *Martin Henry Fischer*.
- McLean, F. C.** See *Joshua Harold Austin*.
- Maclean, (Mrs.) Ida Smedley**, nature of yeast fats, A., i, 304.
the conditions influencing the formation of fat by the yeast-cell, A., i, 795.
- McLean, Stuart**, thermal evolution of gases absorbed by charcoals and carbonised lignites, A., ii, 852.
- McLennan, John Cunningham, and D. S. Ainslie**, the structure of the line $\lambda = 6708$ Å. of the isotopes of lithium, A., ii, 541.
- McLennan, John Cunningham, D. S. Ainslie, and F. M. Cals**, absorption of $\lambda 5460.97$ Å. by luminous mercury vapour, A., ii, 728.
- McLennan, John Cunningham, and P. A. Petrie**, the spectra of helium, hydrogen, and carbon in the extreme ultra-violet, A., ii, 330.
- MacLeod, A. Garrard.** See *J. Lucien Morris*.
- MacLeod, Annie Louise, Marion C. Pfund, and Mary L. Kilpatrick**, the dinitro-derivatives of *p*-dichlorobenzene, A., i, 1131.
- McLeod, James Waller, and John Gordon**, production of hydrogen peroxide by bacteria, A., i, 1095.
- McLintock, W. F. P., and F. R. Ennos**, structure and composition of the Strathmore (Perthshire) meteorite, A., ii, 861.
- McMullen, T. C., Friedel and Crafts' reaction**; the preparation of 2-*p*-toluoylbenzoic acid, A., i, 140.
the Friedel-Crafts' reaction with phthalic anhydride, A., i, 1025.
- McRae, John Alexander.** See *Arthur Lapworth*.
- Maeri, V.**, qualitative chemical analysis, A., ii, 779.
- Madelung, Walter**, questions relating to the constitution and function of electrolytes, A., ii, 344.
- Madelung, Walter, and E. Kern**, dicyanamide, A., i, 438.
- Madinaveitia, Antonio**, hydroxybenzyl-dimethylamine, A., i, 133.
abiestic acid, A., i, 829.
- Madorsky, S. L.** See *William Draper Harkins*.
- Maeder, Horst.** See *Emanuel Merck, Richard Willstätter, and Otto Wolfes*.
- Mäder, W.** See *Paul Karrer*.
- Maestrini, Dario**, velocity of reaction of vegetable enzymes. I. Influence of the concentration of the enzymes on the velocity of action of the enzymes of germinated barley, A., i, 507.
velocity of reaction of vegetable enzymes. II. Effect of hydrogen ions and of salts on the velocity of action of the enzymes of germinated barley, A., i, 507.
velocity of reaction of vegetable enzymes. III. Effect of the quantity and volume of the substrate on the activity of the amylase of germinated barley, A., i, 508.
- Magasanik, J.** See *Georg Wiegner*.
- Magnus, A.**, complex compounds, A., ii, 559.
- Magnus, A., and E. Schmid**, a method for measuring vapour densities for the determination of atomic weights, A., ii, 260.
- Mahler, E. de**, a general method for the preparation of carbides of metalloids, and the existence of carbides of phosphorus and arsenic, A., i, 101.
a new process for the preparation of cadmium dimethyl, A., i, 240.
- Mailhe, Alphonse**, new preparation of cycloalkylamines, A., i, 332.
the catalytic decomposition of oleic acid, A., i, 423.
the catalytic decomposition of shark oil, A., i, 424.
catalytic transformation of vegetable and animal oils into petrol, A., i, 709.
catalytic decomposition of arachis oil, A., i, 712.
the catalytic decomposition of the lower acids, A., i, 803.
decomposition of aliphatic ketones, A., i, 985.
- Mains, Gerald H.**, the system, furfuraldehyde-water, A., i, 566.
- Majima, Rikô**, [with *Chozo Chiba and Yoshihiro Kudo*], the chief constituent of Japanese lac. IX. Chemical investigation of the different, naturally-occurring species of lac which are closely allied to Japanese lac, A., i, 263.
- Majima, Rikô**, [with *Yoshikide Tahara, Gilaro Takayama, Watanabe, and Okazaki*], the main constituent of Japanese lac. VIII. Position of the double bonds in the side chain of urushiol and demonstration that urushiol is not homogeneous, A., i, 262.

- Majima, Rikô**, and **Chika Kuroda**, the colouring matter of *Lithospermum erythrorhizon*, A., i, 946.
- Majima, Rikô**, **Kwanto Nagaoaka**, and **Keisuke Tamada**, the melting points of certain fatty-aromatic ketones, A., i, 257.
- Major, R. H.**, chronic acid nephritis, A., i, 610.
- Malfitano, G.**, and **M. Catoire**, amylo-cellulose considered as composed of silicic acid and amylose, A., i, 527.
- Malgoyre, J.**, modification of the chlorometric method of Hayem and Winter, A., ii, 780.
- Malkomes, Theodor**. See **Burckhardt Helfferich**.
- Malméjac, F.**, the synovial fluid, A., i, 703.
- Maltby, John Gwilliam**, optical rotations of the sugars. I. The aldohexoses and aldopentoses, T., 2608.
- Malvesin, Philippe**, sulphiformin (methanalsulphurous acid), A., i, 222.
the chemical composition and the bouquet of wines, A., i, 1220.
the estimation of tannin in wines, A., ii, 172.
estimation of volatile acids in wine, A., ii, 880.
- Malvesin, Philippe, Ch. Rivalland**, and **L. Grandchamp**, a new preparation of formaldehyde hyposulphite and an economical generator of hyposulphurous acid, A., i, 8.
- Mameli, Efisio**, cubebin. IV. Derivatives of cubebinolide, A., i, 847.
syntheses in the benzofuran group; 6-methylcoumaran-2-one and 3-methyl-6-isopropylcoumaran-3-one. I., A., i, 669.
mercuriation in the aromatic series. I. Phenolmercuri-acetates and -hydroxides and their derivatives, A., i, 695.
mercuriation in the aromatic series. III. Mercuriated derivatives of 6-iodothymol, A., i, 1082.
mercuriation in the aromatic series. IV. Dimercuriated derivatives of guaiacol, A., i, 1082.
- Mameli, Efisio**, and **G. Cocconi**, mercuriation in the aromatic series. V. Binary and ternary systems relating to mercuriation, A., i, 1083.
- Mameli, Efisio**, and **Anna Mameli-Mannessier**, mercuriation in the aromatic series. II. Thymolmercuri-acetates and their derivatives, A., i, 1080.
- Mameli-Mannessier, Anna**. See **Efisio Mameli**.
- Manchot, Wilhelm**, a modification of silicon soluble in hydrofluoric acid, A., ii, 144.
- Manchot, Wilhelm**, and **Herbert Funk**, modifications of silicon; solubility of silicon in hydrofluoric acid, A., ii, 286.
the modifications of silicon. II. Silicon from copper silicide, A., ii, 784.
- Manchot, Wilhelm**, and **Karl Ortner**, hydrates of selenium dioxide, A., ii, 283.
- Mancini, Mario A.**, modification in the Kossel-Neumann method for the estimation of phosphorus in organic substances, A., ii, 83.
- Manicke, Paul**. See **Hermann Kunz-Krause**.
- Manley, John Job**, the use as a drying agent of phosphoric oxide treated with ozone, T., 331.
protection of brass weights, A., ii, 840.
- Mann, Frederick George**, the interaction of aniline and acraldehyde, T., 2178.
- Mann, Frederick George**, and (*Sir*) **William Jackson Pope**, production and reactions of $\beta\beta'$ -dichlorodiethyl sulphide, T., 594.
the sulphilmines, a new class of organic compounds containing quadrivalent sulphur, T., 1052.
the β -chlorovinylarsines, T., 1754.
- Mann, William A.** See **George L. Clark**.
- Mannebach, O.**, a new apparatus for colorimetric estimations, A., ii, 158.
- Mannich, Carl** and **W. Brose**, the acetone [isopropylidene] compound of anhydroenneheptitol; the acetone compounds of polyhydroxy-alcohols, A., i, 1118.
- Mannich, Carl**, and **G. Heilner**, the synthesis of β -keto-bases from acetophenone, formaldehyde, and amine salts, A., i, 351.
transformations of a secondary ketobase, A., i, 371.
- Mannich, Carl**, and **C. A. Rojahn**, the colloidal nature of saccharated iron, A., i, 718.
- Manake, R. H.**, preparation of pure ceria-earth compounds, A., ii, 646.
- Mansky, (Frl.) S.**, the influence of sucrose on the greening of etiolated cotyledons at various stages of germination, A., i, 1222.
- Mansuri, Qasim Ali**, intermetallic actions; the system aluminium-arsenic, T., 2272.

- Maquenne, Léon**, the inversion of sucrose by alkaline copper solution, A., i, 920.
- Maquenne, Léon**, and **Ém. Demoussy**, vegetation in media poor in oxygen, A., i, 707.
- influence of calcium on the utilisation of the reserve material during the germination of seeds, A., i, 905.
- Marcel, Sigismund**. See **Hugo Weil**.
- Marcelin, A.**, measurement of the pressure of "surface fluids"; oleic acid, A., ii, 686.
- Marchal, (Mlle) G.**, dissociation of glucinum sulphate, A., ii, 620.
- Marchal, (Mlle) G.** See also **Camille Matignon**.
- Marchand, Robert**, preparation of terpincol, A., i, 262.
- Markwald, Willy**, and **K. Helmholtz**, phosphorus, A., ii, 845.
- Markwald, Willy**, and **F. Struwe**, guanidonium salts, A., i, 328.
- Marousson, Julius**, structure and formation of the humic acids and coal, A., i, 437.
- Marousson, Julius**, and **H. Smelkus**, the colouring constituents of montan wax, A., i, 807.
- Marden, John W.**, and **M. N. Rich**, zirconium, A., ii, 153.
- Margosches, Benjamin Max**, and **R. Baru**, chlorohydrocarbons and carbon chlorides. II. The knowledge of the saturation character of the di-, tri-, and tetra-chloroethylenes, A., ii, 235.
- Margosches, Benjamin Max**, and **Erwin Vogel**, kjeldahlisation of mononitrophenols, mononitrobenzoic acids, and mononitrocinnamic acids, A., ii, 522.
- Marinesco, G.**, the oxidising enzymes in the phenomena of life in its normal and pathological states, A., i, 790.
- the evolution of oxidative enzymes, A., i, 1211.
- Marinot, André**, estimation of water in fuels, A., ii, 223.
- estimation of sulphur in cast irons and steels, A., ii, 224.
- Mario, Tealdi**, red coloration of sodium hypochlorite solutions, A., ii, 457.
- Mark, Hermann**. See **Wilhelm Schlenk**.
- Markley, A. L.** See **F. H. Rhodes**.
- Marsh, C. Dwight**, **A. B. Clawson**, **James F. Couch**, and **Hadleigh Marsh**, western sneezeweed (*Helenium hoopesii*) as a poisonous plant, A., i, 413.
- Marsh, Hadleigh**. See **C. Dwight Marsh**.
- Marsh, J. K.**, and **Alfred H. alter Stewart**, a magnetic model of atomic constitution, A., ii, 277.
- Marshall, Eli Kennerly, jun.**, effect of loss of carbon dioxide on the hydrogen-ion concentration of urine, A., i, 494.
- Marshall, Eli Kennerly, jun.** See also **Benjamin S. Neuhausen**.
- Martin, Felix**, salts of codeine; the hydrobromide; preparation of solutions for injection; A., i, 948.
- Martin, Frances T.** See **Robert B. Gibson**.
- Martin, J. H.** See **G. Davis Buckner**.
- Martin, W. H.**, and **S. Lehrman**, scattering of light by dust-free liquids. II., A., ii, 335.
- Martinet, Jh.**, colouring matters of the isatin-yellow series, A., i, 278.
- Martinet, Jh.**, and **F. Vacher**, a naphthalenic di-isatin, A., i, 688.
- Martinez, J. Palacios**, and **Heike Kamerlingh Onnes**, the vapour pressure of hydrogen and new thermometric determinations in the domain of liquid hydrogen, A., ii, 472.
- Martinez, J. Palacios**. See also **Heike Kamerlingh Onnes**.
- Marvel, Carl S.**, and **V. L. Gould**, the preparation of mercury dialkyl compounds from the Grignard reagent, A., i, 329.
- Marx, Theodor**. See **Karl Schaum**.
- Masai, Y.** See **Yashiro Kotake**.
- Masing, G.**, the theory of resistance limits in mixed crystals, A., ii, 37.
- Mason, Walter**, and **Richard Vernon Wheeler**, the ignition of gases. II. Ignition by a heated surface; mixtures of methane and air, T., 2079.
- Massera, V.**, essential oil of juniper from Cyrenaica, A., i, 945.
- Massink, A.**, the minimum solubility of aluminium hydroxide in water, A., ii, 299.
- phenol-red as an indicator for acidity of media, A., ii, 307.
- Masson, James Irvine Orme**. See **Leonel Felix Gilbert**.
- Masters, (Miss) Helen**, reactions of cellulose with sodium chloride and other neutral salt solutions. I. Preliminary survey, T., 2026.
- Masumoto, Bunkichi**. See **Shigeru Komatsu**.
- Mathews, Joseph Howard**. See **Frederick L. Browne**, and **Alvin Strickler**.
- Mathewson, W. E.**, the use of sodium α -naphthol-2-sulphonate for the spectrophotometric estimation of aromatic amino-compounds, A., ii, 882.
- Mathias, Émile, Claude Auguste Crommelin**, and **Heike Kamerlingh Onnes**, rectilinear diameter of hydrogen, A., ii, 440, 561.

- Mathias, Émile, Claude Auguste Crommelin,** and **Heike Kamerlingh Onnes**, heat of vaporisation and the difference, $m'-m$, of the specific heats at the state of saturation for argon, oxygen, nitrogen, and hydrogen, A., ii, 472.
the rectilinear diameter of nitrogen, A., ii, 562.
the rectilinear diameter of argon, A., ii, 565.
- Mathias, Émile,** and **Heike Kamerlingh Onnes**, the rectilinear diameter of oxygen, A., ii, 561.
- Matignon, Camille**, economic realisation of oxidation reactions in factories where nitric acid is synthesised; applications, A., ii, 563.
- Matignon, Camille,** and **M. Fréjaques**, the synthetic preparation of carbamide from ammonia, A., i, 723.
the transformation of ammonia into carbamide, A., ii, 272.
conditions of formation and stability of ammonium carbamate, A., ii, 445.
the transformation of ammonium carbamate, A., ii, 499.
the transformation of gypsum into ammonium sulphate, A., ii, 570.
- Matignon, Camille,** and **(Mlle) G. Marchal**, the transformation of sodium formate into oxalate, A., i, 915.
- Matsumo, Gengo.** See **Leon Asher**.
- Matsumo, Kichimatsu.** See **William Edward Garner**.
- Matsuoka, Z.** See **Yashiro Kotake**.
- Matsuyama, Toshitane.** See **Kozo Sakaguchi**.
- Matsuyama, Yoshihiko.** See **Umetarō Suzuki**.
- Mattaar, Th. J. F.**, the replacement of halogen in 4-chloro-3-nitrobenzonitrile and in 4-bromo-3-nitrobenzonitrile. I and II., A., i, 251; ii, 275.
- Matthies, Max.** See **Fritz Paneth**.
- Mattson, Sante Emil**, relationship between precipitation, adsorption, and charge on the particles with particular reference to the hydroxyl ions, A., i, 800.
- Maurer, H.** See **William Küster**.
- Maurice, P.,** and **L. Servantie**, influence of the dextrose concentration and of the alkalinity on glycolysis in vitro, A., i, 121.
- Mauthner, Ferdinand**, the synthesis of aromatic ketones by means of mixed zinc organic compounds, A., i, 457.
new synthesis of isoferulic acid [3-hydroxy-4-methoxycinnamic acid], A., i, 936.
- Marted, Edward Bradford**, catalysis of hydrogen peroxide by finely divided platinum; the influence of inhibitors, T., 1760.
- Maxwell, T. B.**, some derivatives of fenchone, A., i, 753.
- May, Clarence E.,** 8-naphthol, A., i, 449.
- May, Clarence E.,** and **Harry P. Ross**, nesslerisation of ammonia solutions, A., ii, 312.
- Mayeda, Kenjiro.** See **Yoshiharu Murayama**.
- Mayeda, Minoru**, mannanase and levidulinase, A., i, 694.
- Mayen, Hans.** See **Emanuel Merck**.
- Mayer, A.** See **Alexander Gutbier**.
- Mayer, (Mlle) Anka.** See **Maximilian Samec**.
- Mayer, Fritz,** and **Erika Alken**, 2:6-dimethylnaphthalene, A., i, 999.
- Mayer, Fritz,** and **Walter Freund** [with **Kaspar Pfaff**, and **Hermann Wernicke**], the reaction of o-halogenated ketones with feebly basic amines and synthesis of derivatives of *ms*-phenylacridine, A., i, 865.
- Mayer, Fritz,** and **Rudolf Hell**, the constitution of pyrazoleanthrone-yellow, A., i, 877.
- Mayer, Fritz,** and **Wilhelm Krieger**, the diphenyleneoxide series, A., i, 746.
- Mayer, Fritz,** and **Heinrich Schonfelder**, the acenaphthene series. II., A., i, 1166.
- Mayer, Fritz,** and **Therese Schulte**, hydrogenation of 1:6-dimethylnaphthalene, A., i, 819.
- Mayer, Fritz,** and **Adolf Sieglitz**, investigations and ring closures in the methylnaphthalene series, A., i, 999.
- Mayer, Fritz,** and **Adolf Sieglitz** [with **E. Fischer**, **J. Hagen**, **R. Jung**, **W. Knies**, **C. Kohl**, **F. Listmann**, **W. Neugebauer**, and **Therese Schulte**], investigations and ring closures in the series of the methylnaphthalenes, A., i, 740.
- Mayer, Paul**, influence of mineral spring water on the carbohydrate interchange in yeast, A., i, 972.
- Mayes, Charles.** See **Dorothy Jordan Lloyd**.
- Massa, L.** See **Luigi Rolla**.
- Mazzocco, P.**, calcium in the blood of various species of animals, A., i, 788, 891.
- Mazzocco, P.,** and **R. Bustos Moron**, calcium content of blood serum in pregnancy and child-birth, A., i, 892.
- Mazzucchelli, Arrigo,** and **S. Anselmi**, density of aqueous solutions of ammonium perchlorate, A., ii, 876.

- Massuccelli, Arrigo**, and **R. Armenante**, hydrate of ethylene oxide, A., i, 620.
- Mead, W. J.** See **Lowell H. Milligan**.
- Meakins, Jonathan**, and **Charles Robert Harrington**, relation of histamine to intestinal intoxication. I. Presence of histamine in the human intestine, A., i, 396.
- Means, James H.** See **Joseph C. Aub**.
- Mears, Brainerd**, and **Robert E. Hussey**, use of perchloric acid as an aid to digestion in the Kjeldahl nitrogen estimation, A., ii, 159.
- Mecke, R.**, series regularities in the resonance spectrum of iodine, A., ii, 177.
- Meerwein, Hans**, and **Konrad van Emster** [with **Jacob Jousen**], the equilibrium isomerism between bornyl chloride, isobornyl chloride and camphene hydrochloride, A., ii, 751.
- Meerwein, Hans**, and **Jacob Jousen**, the preparation of bornylene, A., i, 1042.
- Mees, R. T. A.**, the detergent power of soap solutions, A., ii, 268.
- Meggers, W. F.** See **Paul D. Foote**.
- Mehner, Heinz.** See **Paul Holzer**.
- Meier, Klothilde**, and **W. Krönig**, blood-gas analysis. IX. Narcosis and charge on the colloids, A., i, 191.
- Meier, Klothilde.** See also **Felix Rosenthal**.
- Meier, Wolfgang.** See **Otto Fischer**.
- Meigen, W.**, and **A. Neuberger**, the separation of solid and liquid fatty acids, A., ii, 880.
- Meisenheimer, Jakob**, the Beckmann transformation, A., i, 152.
the nitrogenous constituents of yeast. II. The purine bases and the diamino-acids; results, A., i, 304.
optically active amine oxides. III., A., i, 813.
- Meisenheimer, Jakob**, and **Hermann Bernhard**, optically active amine oxides. III. (i). Methyl ethyl propylamine oxide, A., i, 813.
- Meisenheimer, Jakob**, **Hellmuth Greeske**, and **Amalie Willmersdorf**, behaviour of allyl- and benzyl-amino-oxides towards sodium hydroxide solution, A., i, 334.
- Meisenheimer, Jakob**, and **Artur Lohner**, optically active amine oxides. III. (ii). Methyl ethyl allylamine oxide, A., i, 813.
optically active amine oxides. III. (iii). Benzyl methyl ethylamine oxide, A., i, 822.
- Meisenheimer, Jakob**, and **Karl Weibesahn**, triaryl isooxazoles, A., i, 176.
- Meissner, R.**, acceleration of blood-clotting by euphylline, A., i, 193.
- Meltner, Lise**, radioactivity and atomic constitution, A., ii, 15.
the production of β -ray spectra by radioactive substances, A., ii, 416.
the relationship between β - and γ -rays, A., ii, 416.
the Neuburger nuclear model, A., ii, 702.
 β -ray spectra and their connexion with γ -radiation, A., ii, 732.
wave-lengths of γ -rays, A., ii, 733.
- Meltner, Lise.** See also **Otto Hahn**.
- Melamed, Dona.** See **Rudolf Pummerer**.
- Melander, Karl H. A.**, lignin. I. and II., A., i, 325.
estimation of methyl alcohol, A., ii, 320.
- Melbye, G. Svensen**, the solubility of copper hydroxide in sodium hydroxide solutions, A., ii, 851.
- Meldrum, William Buell.** See **Theodore William Richards**.
- Mellon, M. G.**, estimation of lead in lead amalgam, A., ii, 870.
- Mellon, M. G.**, and **H. F. Reinhard**, estimation of lead in lead amalgam, A., ii, 787.
simultaneous electro-deposition of lead and lead dioxide, A., ii, 817.
- Mellon, M. G.**, and **J. C. Siegesmund**, chlorination of mixed silver haloids in Gooch crucibles, A., ii, 781.
- Mellon, R. R.**, **Solomon Farley Acree**, **Pauline M. Avery**, and **Edgar A. Slagle**, the ionisation constants of glycerophosphoric acid and their use as buffers, especially in culture media, A., i, 405.
- Mellon, R. R.** See also **Solomon Farley Acree**.
- Melville, F. L.**, effect of iron on the iodine titration of arsenite, A., ii, 784.
- Menager, (Mlle) Y.** See **Gabriel Bertrand**, and **P. Freundler**.
- Menaul, Paul**, the hypobromite reaction on urea, A., ii, 403.
- Mende, H.**, the analysis of aluminium alloys, A., ii, 162.
- Mendelewitsch, Anissim.** See **Moritz Kohn**.
- Meneghetti, E.**, pharmacological action of colloidal arsenious sulphide, A., i, 301.
- Menzel.** See **Erich Müller**.
- Menzel, Heinrich**, dissociation and hydrolysis equilibria in solutions of salts of carbonic and boric acids, A., ii, 845.
- Menzel, W.** See **Julius Tröger**.
- Mezies, Robert Charles**, γ -methyl-fructoside, T., 2238.

- Mensies, Robert Charles**, the purification of methyl alcohol by means of sodium hypochlorite, T., 2787.
- Merck, Emanuel**, explosion of hydroargyrum oxycyanatum, A., i, 640.
preparation of hydrogenated 1-alkylpyridine-4-carboxylates, A., i, 949.
preparation of 1-alkylpyridine carboxylates, A., i, 950.
preparation of betaines of the pyridine series, A., i, 950.
- Merck, Emanuel, Claus Diehl, and Hans Mayer**, preparation of acetylsalicyl compounds of quinine and its derivatives, A., i, 46.
- Merck, Emanuel, and Otto Wolfes**, preparation of tropinonecarboxylic acid esters, A., i, 567.
- Merck, Emanuel, Otto Wolfes, and Erich Kornick**, preparation of carboxylic acids of the purine series, A., i, 1071.
- Merck, Emanuel, Otto Wolfes, and Horst Maeder**, preparation of tropinonecarboxylic acid esters, A., i, 568.
preparation of nortropinone and its derivatives, A., i, 1172.
preparation of tropinecarboxylic acid, A., i, 1173.
preparation of tropinouedicarboxylic esters, A., i, 1173.
- Merkel, E.** See *E. Grüneisen*.
- Merl, T., and J. Daimer**, a study of the catalase of flour, A., i, 184.
- Merrill, Alice R. Thompson**, cystine, A., i, 326.
- Merrill, David R., and Charles C. Scallione**, catalytic oxidation of carbon monoxide at ordinary temperatures, A., ii, 136.
- Merrill, George Perkins**, meteoric iron from Odessa, Ector Co., Texas, A., ii, 451.
meteoric irons from Alpine, Texas, and Signal Mountain, Lower California, A., ii, 452.
- Merrill, Henry Baldwin**, separation of molybdenum and tungsten by means of selenium oxychloride, A., ii, 229.
separation of columbium and tantalum by means of selenium oxychloride, A., ii, 230.
- Merrill, Paul W., F. L. Hopper, and Clyde R. Keith**, identification of air lines in spark spectra from λ 5927 to λ 8683, A., ii, 802.
- Merton, Thomas Ralph**, the structure of the red lithium line, A., ii, 808.
- Merton, Thomas Ralph, and S. Barratt**, spectrum of hydrogen, A., ii, 461.
- Merton, Thomas Ralph, and D. N. Harrison**, errors arising in the measurement of unsymmetrical spectrum lines, A., ii, 673.
- Merton, Thomas Ralph.** See also *Harold Brewer Hartley*.
- Merwin, Herbert Eugene.** See *Eugen Posnjak, and Henry Stephens Washington*.
- Messmer, Ernst.** See *Kurt Hess*.
- Messner, J.**, benzyl compounds [benzyl alcohol], A., i, 138.
- Mestrenat, W., and (Mlle) Marthe Paul Janet**, estimation of ammonia in urine by Schlesing's bell-glass method, A., ii, 453.
- Metzl, Heinrich.** See *Walter Fuchs*.
- Meulen, Henri ter**, a new method of estimating sulphur in organic compounds, A., ii, 311.
the estimation of oxygen in organic compounds, A., ii, 717.
- Meulengracht, E.**, estimation of hæmoglobin, A., ii, 798.
- Meurice, R.**, a rapid method for the estimation of ammoniacal nitrogen, A., ii, 225.
estimation of calcium in natural phosphates, A., ii, 658.
- Mewes, Rudolf**, approximate rule for the specific heats of gases and vapours, A., ii, 737.
- Meyer, Charles F.** See *Walter F. Colby*.
- Meyer, Friedrich, Gustav Bailleul, and Gerhard Henkel**, the existence of sulphur tetroxide, A., ii, 843.
- Meyer, G., and Grunlich**, spectroscopy of uranium and some rare earths, A., ii, 6.
- Meyer, G., and Adolf Heck**, molecular refraction of some molten salts and their degree of dissociation, A., ii, 241, 329.
- Meyer, Gustave Morris.** See *Phœbus A. Levens*.
- Meyer, Heinrich.** See *Walther Diltthey*.
- Meyer, Julius**, preparation of selenium dioxide, A., ii, 639.
- Meyer, Julius** [with *Leonhard Speich, and Hanns Moldenhauer*], complex selenates, A., ii, 70.
- Meyer, Julius, and Walter Friedrich**, barium sulphuric acid and barium selenic acid, A., ii, 644.
- Meyer, Julius, and Hanns Moldenhauer**, the preparation of telluric acid, A., ii, 49.
- Meyer, Julius, and Valentin Stacezny**, some poly-acids of the elements of the sulphur group, A., ii, 773.

- Meyer, Julius**, and **Walter Wagner**, organic derivatives of selenic acid, A., i, 620.
 nitrosylselenic acid, A., ii, 372.
- Meyer, Jules**. See **Leopold Ruzicka**.
- Meyer, Kurt Heinrich**, and **Hans Gottlieb-Billroth**, action of nitric acid on phenol ethers, A., i, 539.
- Meyer, Kurt Heinrich**, and **Ludwig Orthner**, synthesis of formamide from carbon monoxide and ammonia, A., i, 529.
- Meyer, Kurt Heinrich**, and **Kurt Schuster**, diphenylstyrylcarbinol and triphenylallene, A., i, 540.
 transformation of tertiary ethinylcarbinols into unsaturated ketones, A., i, 556.
- Meyer, Martin**, [lecture experiment], time reaction, A., ii, 636.
- Meyer, Martin**. See also **Marston Taylor Bogert**.
- Meyer, P.** See **Hermann Staudinger**.
- Meyer, Robert**, thermal expansion of concentrated salt solutions, A., ii, 737.
- Meyer, Robert**. See also **Walther Borsche**.
- Meyer, R. Ernest**. See **Paul Ruggli**.
- Meyer, Stefan**, and **Karl Preisbarm**, coloration and luminescence produced by the action of Becquerel rays, A., ii, 339.
- Meyer, Walther**. See **Arthur Hantzsch**.
- Meyer-Bisch, Robert**, and **E. Basch**, the fate of parenteral administered sulphur and its influence on metabolism, A., i, 83.
- Meyer-Bisch, Robert**. See also **Wolfgang Heubner**.
- Meyerhof, Otto**, energy exchanges in muscle. IV. Formation of lactic acid in cut muscle, A., i, 86.
 energy exchanges in muscle, A., i, 897.
 heat of combustion of lactic acid, A., ii, 475.
- Meysenbug, L. von**. See **Willey Denis**.
- Meiser, F.**, estimation of urea, A., ii, 170.
- Micewicz, St.** See **Friedrich Kehrman**.
- Michaelis, Leonor**, the determination of alkalinity in culture media, A., i, 405.
- Michaelis, Leonor**, and **Frjð Airila**, the electric charge of hæmoglobin, A., i, 63.
- Michaelis, Leonor**, and **N. Hirabayashi**, ionic synergism. II. Investigation of mastic sola, A., ii, 429.
- Michaelis, Leonor**, and **R. Krüger**, further elaboration of the indicator method without buffers, A., ii, 157.
- Michajlova, Olga**. See **Nikolai Schilow**.
- Michel, Eduard**. See **Fritz Ephraim**.
- Michel, J.** See **Georg Bredig**.
- Michels, A.**, the system isobutyl alcohol-water, A., ii, 486.
- Middleton, Edmund Burrus**. See **Frank C. Whitmore**.
- Miskeley, Arthur**. See **Max Bergmann**.
- Mikeska, Louis A.** See **Phæbus A. Levene**.
- Mildbrand, Hans**. See **Hermann Leuchs**.
- Milde, E.** See **Fritz Arndt**.
- Milbauer, Jaroslav**, sodium peroxide, A., ii, 520.
- Miller, E. J.** See **Floyd E. Bartell**, and **Charles S. Robinson**.
- Miller, Elizabeth W.**, effect of certain stimulating substances on the invertase activity of yeast, A., i, 202.
- Miller, G. von**. See **Richard Willstätter**.
- Miller, George E.** See **Ebenezer Emmet Reid**.
- Miller, Harry G.**, nitrogen compounds in lucerne hay, A., i, 414.
- Miller, J. E.** See **De Witt Neighbors**.
- Miller, M.** See **Edgar Wedekind**.
- Millican, (Miss) Ida L., Alfred Francis Joseph**, and **Thomas Martin Lowry**, the properties of ammonium nitrate. II. Ammonium nitrate and water, T., 959.
- Milligan, C. H.**, and **Ebenezer Emmet Reid**, the ethylation of benzene and naphthalene, A., i, 330.
 the transfer of hydrogen from an alcohol to an aldehyde, A., i, 337.
- Milligan, Lowell H.**, solubility of aluminium nitrate crystals in solutions of nitric acid of various strengths at various temperatures, A., ii, 380.
- Milligan, Lowell H.** [with **W. J. Mead**], mechanism of the dehydration of crystalline aluminium hydroxide and of the adsorption of water by the resulting alumina, A., ii, 447.
- Millikan, Robert Andrews**, extension of the ultra-violet spectrum and the progression with atomic number of the spectra of light elements, A., ii, 100.
- Mills, William Hobson**, the cyanine dyes. IV. Cyanine dyes of the benzothiazole series, T., 455.
- Mills, William Hobson**, and **Walter Theodore Karl Braunkholz**, the cyanine dyes. V. The virtual tautomerism of the thiocyanines, T., 1489.
- Mills, William Hobson**, and (Miss) **Frances Mary Hamer**, the quaternary salts of quinaldinic acid, T., 2008.

- Mills, William Hobson**, and (Sir) **William Jackson Pope**, 2-*p*-dimethylaminostyrylpyridine methiodide, a new photographic sensitiser, T., 946.
- Mills, William Hobson**, and **James Leonard Brierley Smith**, the reactivity of methyl groups in heterocyclic bases, T., 2724.
- Mills, William Hobson**. See also **Walter Karl Theodore Braunholtz**.
- Minaeff, W. J.**, and **Kurt Ripper**, anthracoumarin derivatives. I., A., 1, 162.
- Mingazzini, M.** See **Leopold Buzicka**.
- Minkowski, R.**, influence of the pressure of foreign gases on the *D*-lines in saturated sodium vapour, A., ii, 242.
- Minkowski, R.** See also **Rudolf Ladenburg**.
- Minnis, Wesley**. See **Moses Gomberg**.
- Minovici, Stefan**, and **Al. Ionescu**, new method for the volumetric estimation of copper, A., ii, 162.
- Minovici, Stefan**, and **Constantin Kollo**, new method for the estimation of manganese, A., ii, 787.
- Minton, Thomas Hooker**, and **Henry Stephen**, preparation of *o*-, *m*-, and *p*-nitrophenoxyacetic acids and various nitrotyloxyacetic acids and their derivatives, T., 1591.
- studies in the coumaranone series. II. The preparation of 4- and 6-chlorocoumaran-2-ones and their conversion into 2- and 4-chloroflavonols respectively, and some derivatives of *o*- and *p*-chlorophenoxycetic acids, T., 1598.
- Miolati, A.**, complex salt of mercuric acetate and sulphide, A., i, 982.
- Mirande, Marcel**, the influence of light on the formation of anthocyanin in the scales of lily bulbs (*Lilium candidum* and *L. martagon*), A., i, 1100.
- the relation between anthocyanin and oxydases, A., i, 1224.
- the relation between tissue-acidity and the presence of anthocyanin in the scales of lily bulbs exposed to light, A., i, 1224.
- Miravalles, R.** See **Enrique Moles**.
- Mislowitz, E.**, estimation of oxalic acid in urine, A., ii, 325.
- Missenden, John**, identification of caesium and rubidium, A., ii, 658.
- Misson, Georges**, detection and estimation of vanadium in steels, A., ii, 459.
- estimation of phosphorus in minerals and in coke ash by a colorimetric method, A., ii, 718.
- Mitchell, Alec Duncan**, studies on hypophosphorous acid. IV. Its reaction with cupric chloride, T., 1624.
- Mitchell, H. H.**, **W. B. Nevens**, and **F. E. Kendall**, the relation between the endogenous katabolism and the non-protein constituents of the tissues, A., i, 897.
- Mitchell, Philip H.**, **J. Walter Wilson**, and **Ralph E. Stanton**, the selective absorption of potassium by animal cells. II. The cause of potassium selection as indicated by the absorption of rubidium and caesium, A., i, 196.
- Mitscherlich, Alexander**, ignition point of detonating gas [hydrogen and oxygen, A., ii, 358.
- Mitsukuri, Shinroku**. See **Frederick G. Keyes**.
- Mitra, N. N.**, and **Niratan Dhar**, some induced reactions and their analogues in the animal body, A., i, 1210.
- Mitra, N. N.** See also **Niratan Dhar**.
- Mituhori, Saburo**. See **Yasuhiko Asahira**.
- Miura, Masataro**. See **Sylvester Solomon Zilva**.
- Miyadera, K.**, effect of radiothorium on metabolism, A., i, 966.
- Miyake, Koji**, and **S. Soma**, nitrification, A., i, 1096.
- Miyamoto, Susumu**, chemical reactions induced by the silent discharge. I. Ethylene and nitrogen. II. Benzene and carbon dioxide, A., i, 418.
- reducing action of ferrous hydroxide, A., i, 647.
- Miyashita, Chôji**. See **Akira Ogata**.
- Möhl**. See **Otto Diels**.
- Mochlmann, E. O.** See **Francis P. Venable**.
- Möller, Eggert**. See **Christen Lundsgaard**.
- Möller, Hans Peter**, rhythmic precipitation phenomena in cell membranes of plants, A., i, 94.
- Moeller, W.**, the connexion between hydrolysis and adsorption, A., ii, 689.
- Moerk, Frank X.**, the volumetric determination of phosphoric acid and of sodium phosphate and pyrophosphates, A., ii, 866.
- Moerk, Frank X.**, and **Edward J. Hughes**, methyl-red in the assay of phosphoric acid and sodium phosphate, A., ii, 866.
- Mörner, Carl Th.**, homogentisic acid. II. The behaviour of homogentisic acid when boiled with ferric chloride, A., i, 341.

- Mörner, Carl Th.**, homogentisic acid. III., A., i, 341.
magnesium compound of 8-hydroxy-quinoline, A., ii, 659.
- Moersveld, A. L. Th.** See **Ernst Cohen**.
- Moggi, Aldo.** See **Antonio Pieroni**.
- Mohler, Fred L.** See **Paul D. Foote**.
- Mohr, Ernst**, theory of the *cis-trans*-isomerism of decahydronaphthalene, A., i, 243.
two cycloheptane models free from strain, A., i, 441.
the accuracy of Dumas's method for the estimation of nitrogen in the cases of substances rich in nitrogen, A., ii, 82.
- Moir, James**, the calculation of the colour of the azo-dyes and related coloured substances, T., 1555.
the mathematics of the dicyclic colour theory, and a new theory of the structure of the nitrogen atom, T., 1808.
sensitive test for phenols, A., ii, 321.
colour and chemical constitution. XIII. Calculation of the colour of monocyclic dyes, A., ii, 333.
- Mokragants, M.** See **Gabriel Bertrand**.
- Moldasenke, K.** See **Julius von Braun**.
- Moldenhauer, Hanns.** See **Julius Meyer**.
- Moles, Enrique**, the atomic weight of carbon, A., ii, 51.
numerical revision of the data referring to the density of gaseous hydrogen bromide; atomic weight of bromine, A., ii, 140.
critical study of the modern value of the density of gaseous oxygen, A., ii, 141.
normal density of [chemically pure] nitrogen, A., ii, 562.
- Moles, Enrique**, and **M. Crespi**, thermal decomposition of potassium permanganate, A., ii, 374.
new revision of the density of oxygen gas, A., ii, 636.
- Moles, Enrique**, and **F. González**, revision of the density of oxygen gas, A., ii, 497.
- Moles, Enrique**, and **R. Miravalles**, compressibility of exhausted flasks in the determinations of the densities of gases, A., ii, 617.
- Moles, Enrique.** See **M. Payá**.
- Mollisch, Hans**, effect of transpiration on the disappearance of starch from leaves, A., i, 309.
the alleged development of hydrogen peroxide in carbon dioxide assimilation, A., i, 411.
- Molliard, Marin**, a new acid fermentation produced by *Aspergillus niger*, A., i, 611.
- Mollow, W.**, spleen and digestion, A., i, 397.
- Moloney, P. J.**, a quick-acting hydrogen electrode, A., ii, 253.
- Momferratos-Floros, Kate.** See **Ludwig Fincussen**.
- Monasch, E.**, the titration of zinc, A., ii, 160.
- Mond, Robert Ludwig**, and **Albert Edward Wallis**, researches on the metallic carbonyls, T., 29.
the action of nitric oxide on the metallic carbonyls, T., 32.
- Mondain-Monval, Paul**, preparation of ammonium chloride, A., ii, 444.
the preparation of ammonium chloride at low temperatures, A., ii, 642.
- Monnier, R.** See **Friedrich Kehrman**.
- Monro, Alexander Donald**, the isotope ratio of New Zealand boron, T., 986.
- Montagne, (Mlle).** See **Edmond Émile Blaise**.
- Montané, J.** See **Émile F. Terroine**.
- Montemartini, C.**, chemical reactions induced by the corona effect in circuits traversed by continuous currents, A., ii, 734.
- Montes, Zoila.** See **Augustus P. West**.
- Montmollin, Hugues de.** See **Franz Fichter**.
- Moore, B. E.**, excitation stages in open arc-light spectra. I. Sodium, potassium, calcium, strontium, barium, and magnesium. II. Silver, bismuth, cadmium, zinc, air, and copper, A., ii, 242.
- Moore, Forris Jewitt**, and **R. Thomas**, the constitution of the secondary product in the sulphonation of cinnamic acid, A., i, 454.
- Moore, R. B.** See **R. G. Nyswander**.
- Moquet, L.** See **Henri Bierry**.
- Moraczewski, Wacław**, and **Egon Lindner**, influence of intravenous sugar injections on the excretion of lactic acid, on the blood-sugar, and on the white blood-corpuscles, A., i, 402.
- Moran, R. C.**, dinitrodiphenylamine, A., i, 648.
- Moran, Thomas**, and **William Cudmore McCullagh Lewis**, studies in catalysis. XVI. The inversion of sucrose by hydrogen ion, T., 1613.
- Morawitz, P.**, and **G. Denecke**, swelling processes in subcutaneous tissues, A., i, 492.
- Moreau, Ed.**, and **A. Bonis**, colorimeter, A., ii, 862.

- Morelli, R.** See *G. Canneri*.
- Morey, George W., and Norman L. Bowen,** the melting of potash felspar, A., ii, 577.
- Morgan, Gilbert Thomas, and Sydney Chasan,** 6-amino-2-p-tolyl- $\alpha\beta$ -naphthatriazole, A., i, 181.
- Morgan, Gilbert Thomas, and Evelyn Ashley Cooper,** bactericidal action of the quinones and allied compounds, A., i, 204.
- Morgan, Gilbert Thomas, and Harry Dugald Keith Drew** [with *Evelyn Ashley Cooper*], researches on residual affinity and co-ordination. VIII. Interaction of tellurium tetrachloride and β -diketones, T., 922.
- Morgan, Gilbert Thomas, Harry Dugald Keith Drew, and Thomas Vipond Barker,** researches on residual affinity and co-ordination. XI. Interaction of selenium tetrachloride and β -diketones, T., 2432.
- Morgan, Gilbert Thomas, and Hugh Gilmour,** aminonaphthatriazoles as colour intermediates, A., i, 380.
- Morgan, Gilbert Thomas, and Herbert Joseph Seymour King,** researches on residual affinity and co-ordination. IX. Cobaltamine salts of the nitro-dyes, T., 1723.
- Morgan, Gilbert Thomas, and Wilfrid Ledbury,** researches on residual affinity and co-ordination. XIV. Interactions of metallic salts and dimethyldithiolethylene, T., 2882.
- Morgan, Gilbert Thomas, and John Ewart Moss,** researches on residual affinity and co-ordination. XII. Cobaltamine and ferric lakes of dinitrosoresorcinol, T., 2857.
- Morgan, Gilbert Thomas, and Hugh Norman Read,** the diazo-reaction in the carbazole series; carbazole-3-diazoimine and -3-diazonium salts, T., 2709.
- Morgan, Gilbert Thomas, and Harry Gordon Reeves,** azomethine derivatives of the 2- and 4-hydroxy- α -naphthaldehydes, T., 1.
- Morgan, Gilbert Thomas, and Horace Samuel Rooke,** β -naphthylmethylamine-6-sulphonic acid, A., i, 134.
- Morgan, Gilbert Thomas, and John David Main Smith,** researches on residual affinity and co-ordination. VII. Cobaltic lakes of the alizarin series, T., 160.
- researches on residual affinity and co-ordination. X. Salicylatotetraminocobaltic salts and the constitution of oxonium compounds, T., 1956.
- Morgan, Gilbert Thomas, and John David Main Smith,** researches on residual affinity and co-ordination. XIII. Cobaltamine and chromic lakes of the azosalicylic acids, T., 2866.
- Morgan, E. J., C. P. Stewart, and Frederick Gowland Hopkins,** the anaerobic and aerobic oxidation of xanthine and hypoxanthine by tissues and by milk, A., i, 1078.
- Morgen, A., G. Schöler, K. Windheuser, and Elsa Ohlmer,** replacement of protein by urea in rations, A., i, 293.
- Mori, Yoshitane,** the behaviour of phenyl-lactic acid in the animal organism. II., A., i, 1216.
- comparative researches on the production of acetoacetic acid from *d*- and *l*-phenyl-lactic acid and from *d*- and *l*-hydroxyphenyl-lactic acid in the surviving liver, A., i, 1217.
- Mori, Yoshitane, and T. Kanai,** the asymmetrical reduction of ketonic acids to the corresponding alcohols in organs, A., i, 1217.
- Mori, Yoshitane.** See also *Yashiro Kotake*.
- Morimoto, Yoshio,** urea content of cow's milk; estimation of urea, A., i, 703.
- Moritz, J.** See *Martin Battegay*.
- Moron, R. Bustos.** See *P. Mazzocco*.
- Morrell, Jacques C.** See *Harold A. Fales*.
- Morrell, Robert Selby,** the transformation of methyl α -elaeostearate into methyl β -elaeostearate, A., i, 932.
- Morris, J. Lucien, and A. Garrard Macleod,** the uric acid of human blood, A., i, 392.
- colorimetric estimation of uric acid; estimation of 0.03 to 0.5 mg. quantities by a new method, A., ii, 328.
- Morris, Samuel.** See *C. W. Foulk*.
- Mortimer, F. Spencer,** vapour pressures and heats of vaporisation of non-associated liquids, A., ii, 615.
- melting point, latent heat of fusion, and solubility, A., ii, 621.
- Moser, E.** See *Hugo Weil*.
- Moser, Eduard,** attainment of constant high temperatures, A., ii, 115.
- Moser, Ludwig,** the absorption meter, an apparatus for gas analysis, A., ii, 519.
- Moser, Ludwig, and A. Brukl,** gravimetric estimation of hydrogen phosphide and a new apparatus for gas analysis, A., ii, 393.
- Moser, Ludwig, and E. Doctor,** the preparation of hydrogen selenide from metallic selenides, A., ii, 46.

- Moser, Ludwig** and **Josef Ehrlich**, the separation of arsenic from tungsten, vanadium, and molybdenum by means of methyl alcohol in a current of air, A., ii, 814.
the theory of the distillation of arsenic and a new separation of arsenic from all metals in a current of air, A., ii, 815.
- Moser, Ludwig**, and **K. Ertl**, the preparation of hydrogen telluride from metal tellurides, A., ii, 48.
- Moser, Ludwig**, and **P. Kohn**, the estimation of sulphuric acid as barium sulphate in the presence of aluminium, A., ii, 782.
- Moser, Marie**. See **Siegfried Skrapup**.
- Mosimann, Paul**. See **Fritz Ephraim**.
- Moss, John Ewart**. See **Gilbert Thomas Morgan**.
- Moudgill, Kishori Lal**, 2:8-tetra-methyldiaminoacridine, T., 1506.
brominated isocyanines, T., 1509.
bromo-2-methylquinolines, A., i, 173
- Moudgill, Kishori Lal**, and **K. R. Krishna Iyer**, the essential oil from Inchi grass (*Cymbopogon celsus*, Stapf.), A., i, 945.
- Moudgill, Kishori Lal**, and **P. N. Vridhachalam**, essential oil of *Lantana camara*, A., i, 754.
- Mougue, G.**, the preparation of pure galactose, A., i, 630.
- Moureu, Charles**, and **Gerald Barrett**, the condensation products of ethyl β -chloropropionate and ethyl α -chloropropionate with magnesium ethyl bromide and some compounds which are derived from them, A., i, 4.
- Moureu, Charles** and **Ralph L. Brown**, the *o*- and *p*-nitrobenzyl bromides, A., i, 23.
- Moureu, Charles**, and **Charles Dufraisse**, auto-oxidation; the anti-oxygens, A., i, 250.
auto-oxidation; anti-oxygens, and various phenomena related thereto, II., A., i, 824.
- Moureu, Charles**, and **Patrick Hugh Gallagher**, derivatives of cinnamyl alcohol and phenylallyl alcohol; α -phenylglycerol, A., i, 84.
- Moureu, Charles**, and **Adolphe Lepape**, estimation of krypton and xenon in absolute values by spectrophotometry, A., ii, 394.
- Moyle, Dorothy Mary**. See **Dorothy Lilian Foster**.
- Muchlinski, A.** See **Friedrich August Henglein**.
- Mück, Johann**, unsaturated alcohols obtained from the fat of ovarialdermoid cysts, A., i, 1092.
- Mück, Franz Johann**, the volumetric estimation of antimony and tin in red brass, A., ii, 722.
- Mügge, Otto**, structure and simple displacement of iron, A., ii, 381.
- Müller, A.**, is the undecomposed hydrogen peroxide or the oxygen split from it the carrier of disinfecting action? A., i, 903.
- Müller, C.**, and **Otto Warburg**, energy exchange in carbon assimilation by green cells, A., i, 411.
- Müller, Elisabeth**. See **A. Bornstein**.
- Müller, Erich**, dehydroxidation of alcohols, A., i, 102.
internal or catalytic dehydroxidation of formaldehyde, A., i, 109.
alcoholic fermentation of formaldehyde by osmium, A., i, 110.
electrochemical oxidation of organic compounds, A., ii, 469.
- Müller, Erich** [with **Mensel** and **Schubert**], cathodic deposition of tellurium and selenium from their oxy-acids and their electroanalytical estimation, A., ii, 390.
- Müller, Erich** [with **K. Sponzel**], catalytic dehydroxidation of formic acid, A., ii, 558.
- Müller, Erich**, and **Hans Lauterbach**, the electrometric estimation of cyanides in the presence of haloids, A., ii, 403.
electrometric titration of ferrocyanides, A., ii, 795.
electrometric determination of nickel with silver nitrate, A., ii, 875.
- Müller, F.** See **Lothar Wöhler**.
- Müller, Fr.** See **Alfred Thiel**.
- Müller, Friedrich**, the establishment of the international system of normal wave-lengths, A., ii, 725.
- Müller, Hans**, the relation between fats and carbohydrates, A., i, 486.
the alcoholic fermentation of formaldehyde, A., i, 809.
- Müller, Hans**, and **Leo Müller**, the fermentation of glycerol in presence of sulphur, A., i, 904.
- Müller, Hans**. See also **A. Jung**, and **Hans Rupe**.
- Müller, John H.**, separation of germanium and arsenic, A., ii, 320.
- Müller, John H.**, and **Nicol H. Smith**, germanium hydride, A., ii, 775.
- Müller, John H.** See also **Frederick S. Hammett**.
- Müller, Karl Otto**. See **Kurt Brand**, and **Hans Pringsheim**.
- Müller, Leo**. See **Hans Müller**.

- Müller, Robert**, the electrochemistry of non-aqueous solutions. I. Measurement of current-density and potential difference in the electrolysis of metallic salts in pyridine, A., ii, 612.
- Müller, Robert**, and **Alois Duschek**, the electrochemistry of non-aqueous solutions. II. Decomposition potential and electrode potentials in the electrolysis of pyridine solutions of silver nitrate, and the potential of silver in these solutions, A., ii, 612.
- Müller, Robert**, and **Franz Hösl**, electromotive behaviour of aluminium, A., ii, 341.
- Müller, Robert**, and **Reinhold Hönig**, the preparation of a silver amalgam of the composition Hg_2Ag_2 by precipitation from a solution of silver nitrate in pyridine, A., ii, 500.
- Müller, Robert**. See also **Robert Kremann**.
- Müller, Robert**, **Komandit-Gesellschaft**, gas generator with washing apparatus, A., ii, 706.
- Müller, W.**, corrosion of copper by salt solutions, A., ii, 645.
- Mugdan, Susanne**. See **Otto Ruff**.
- Mugnet**, lead in the uranium minerals of Madagascar, A., ii, 216.
- Muirhead, Constance M. M.** See **Violet Dimbleby**.
- Mukai, G.**, the action of carbon dioxide on salt and water distribution in blood, A., i, 287.
- Mukerji, Dharendra Nath**, colouring matters from 1 : 2 : 4 : 5-tetrahydroxy-benzene and related substances, T., 545.
- vat dyes of the azo-series, T., 2879.
- Mukherjee, Jnanendra Nath**, the origin of the charge of a colloidal particle and its neutralisation by electrolytes, A., ii, 198.
- the adsorption of ions, A., ii, 689.
- Mukherjee, Jnanendra Nath**, and **Basil Constantine Papaconstantinou**, an experimental test of Smoluchowski's theory of the kinetics of the process of coagulation, A., ii, 694.
- Muller, Joseph Auguste**, the degree of molecular polymerisation of substances under critical conditions, A., ii, 820.
- Muller, Joseph Auguste**, and **A. Foix**, the estimation of hydrogen and its separation from gaseous paraffins by means of palladium chloride, A., ii, 655.
- the estimation of small quantities of gold as colloidal gold by the colorimetric method, A., ii, 662.
- Mulliken, Robert S.**, separation of isotopes by thermal and pressure diffusion, A., ii, 492.
- Mulliken, Robert S.**, separation of liquid mixtures by centrifuging, A., ii, 686.
- Mulliken, Robert S.**, and **William Draper Harkins**, separation of isotopes; theory of resolution of isotopic mixtures by diffusion and similar processes; experimental separation of mercury by evaporation in a vacuum, A., ii, 295.
- Mumm, Otto**, and **Else Gottschaldt**, the hydrogen esters of 2 : 6-dimethylcinchononic acid, A., i, 861.
- apophyllenic acid and *aa*'-dimethylapophyllenic acid, A., i, 862.
- Muntwyler, O.** See **Hermann Staudinger**.
- Murakawa, Gorô**. See **Heisaburô Kondô**.
- Murayama, Yoshiharu**, chemical constituents of a Chinese drug "Hsiung Ch'uang." I., A., i, 310.
- Murayama, Yoshiharu**, and **Shinjirô Aoyama**, constituents of the Japanese common earthworm. II., A., i, 898.
- Murayama, Yoshiharu**, and **Takeyoshi Itagaki**, essential oil of *Nepeta japonica*, Maxim. I., A., i, 44.
- Murayama, Yoshiharu**, and **Kenjirô Mayeda**, kawa-kawa resin, A., i, 265.
- Murdick, P. P.** See **C. J. Wood**.
- Muresanu, Augustin**, the distribution of chloride between corpuscles and plasma, A., i, 290.
- Murlin, John Raymond**, [imperfect digestion and] the amino-acid fractions and hippuric acid in the urine of pellagrins, A., i, 965.
- Murmann, Ernst**, laboratory notes [estimation of silica, phosphorus in iron, etc.], A., ii, 226.
- estimation of minute quantities of methane, A., ii, 591.
- Murphy, A. J.** See **C. A. Edwards**.
- Murray, Humphrey D.**, coagulation of colloids by electrolytes, A., ii, 37.
- Murschhauser, Hans**, which carbohydrates are excreted in the urine of sucklings when the sucrose in the food exceeds the assimilation limit? a method for the quantitative estimation of several carbohydrates simultaneously in the urine, A., i, 198.
- mutarotation of dextrose under the influence of sodium chloride, A., i, 322.
- the influence of sodium chloride on the mutarotation of dextrose in hydrochloric acid solution. I., A., i, 322.
- influence of sodium chloride on the mutarotation of dextrose in alkaline solution. I and II., A., i, 432.

Murschhauser, Hans, the law governing the constants of mutarotation of dextrose and the concentration of acid, A., i, 432.

the rotation of dextrose in solutions of trisodium phosphate; mutarotation as an analytical method, A., ii, 92.

Murtagh, J. See *O. M. Pico*.

Muschel, Anna, the darkening of carbohydrate containing nutrient media by *Bacillus mesentericus* var. *niger*, A., i, 1095.

Mussler, Ch. See *W. D. Treadwell*.

Muttele, C. F., detection of vegetable oils in animal fats; the phytosteryl acetate test, A., ii, 168.

detection of coconut oil in butter, A., ii, 236.

Myrbäck, Karl. See *Hans von Euler*.

Myers, Burton A., and *Marian C. Shevky*, the estimation of inorganic phosphorus in blood plasma by Bell and Doisy's method, A., ii, 455.

N.

Nägeli, C. See *Paul Karrer*.

Nagahashi, M. See *Joseph Barcroft*.

Nagai, Shôichirô, preparation of piperonaldehyde from isosafrole by the action of ozone, A., i, 839.

Nagai, Wally. See *Yûji Shibata*.

Nagai, W. N., [preparation of] dihydroxyphenylmethylaminoethanol hydrochloride, A., i, 652.

Nagaoka, H., and *Y. Sugiura*, the structure of the bismuth lines, A., ii, 101.

Nagaoka, Kwantô. See *Riko Majima*.

Nägel, Josef, estimation of tin in bearing metals, A., ii, 721.

Nägel, W. See *Carl Dietrich Harries*.

Naik, Kuverji Gosar, and *Mahadeo Dattatraya Avasare*, the formation and properties of dithioketones ($R_2C:S:S$) and dithio-ethers ($R_2S:S$). III., T., 2592.

Naiman, B. See *Alvin Sawyer Wheeler*.

Nakagawa, Tomouchi, the relation of salivary to gastric secretion, A., i, 789.

Nakajima, Tomouchi. See *Heisaburô Kondô*.

Nakatogawa, Shôichi, and *Shômei Kobayashi*, oil from the heads of sea animals of the family Delphinidae, A., i, 701.

Nanji, Dinshaw Rattonji. See *Arthur Robert Ling*.

Narayan, A. L., and *D. Gunnaiya*, absorption of potassium vapour in the associated series, A., ii, 679.

Narayan, A. L., and *G. Subrahmanyam*, a modified form of double slit spectrophotometer, A., ii, 329.

Nash, Thomas P., jun., the kidney factor in phloridzin diabetes, A., i, 497.

Nash, Thomas P., jun., and *Stanley Rossiter Benedict*, the ammonia content of the blood and its bearing on the mechanism of acid neutralisation in the animal organism, A., i, 191.

the ammonia content of blood, A., i, 483.

Nasini, A. G. See *Umberto Sborgi*.

Nathansohn, A., [biochemical and electrochemical oxidation of organic compounds], A., ii, 421.

Nauen, Ernst. See *Erwin Ott*.

Navias, Louis. See *Edward Wuyt Washburn*.

Naylor, Nellie M. See *Roemer Rex Renshaw*.

Neale, Sidney Maurice, influence of the solvent on ionisation and the accompanying heat effect, A., ii, 420.

Neave, A. S. See *Ellwood B. Spear*.

Neber, P. W., free *o*-aminophenylacetic acid, its esters and transformations, A., i, 545.

Negelein, Erwin. See *Otto Warburg*.

Neighbors, De Witt, A. L. Foster, S. M. Clark, J. E. Miller, and *James R. Bailey*, isopropyl-, menthyl-, and bornyl-semicarbazides; reduction of phenylhydrazones, A., i, 880.

Nelken, Annemarie. See *Julius von Braun*.

Nelson, Alta. See *Katharine Blunt*.

Nelson, Burt E., and *Helen A. Leonard*, identification of alkaloids under the microscope from the form of their picrate crystals, A., ii, 327.

Nelson, John Maurice, and *David I. Hitchcock*, the activity of adsorbed invertase, A., i, 184.

uniformity in invertase action, A., i, 388.

Nelson, John Maurice. See also *F. S. Granger*.

Nelson, O. A., and *C. E. Senseman*, vapour pressure determinations on naphthalene, anthracene, phenanthrene, and anthraquinone between their melting and boiling points, A., i, 245.

estimation of anthraquinone, A., ii, 882.

Němec, Antonín, and *František Dusehoň*, is it possible to determine the value of seeds by a biochemical method? A., i, 94.

- Némes, Antonin, and František Duchoň**, the occurrence and action of saccharo-phosphatase in the organism of the plant, A., i, 206.
an indicatory method for evaluating the vitality of seeds by a biochemical method, A., i, 411.
- Němeček, H.**, a new form of ozoniser, A., ii, 841.
- Német, Géza.** See *Gerhardt Katsch*.
- Nemours & Co., E. J. du Pont de.** See *Arthur Percival Tanberg*.
- Nepveux, F.** See *R. Goiffon, and Henri Labbé*.
- Nersesov.** See *Fromholdt*.
- Nets, A.** See *Robert Stollé*.
- Neuberg, Carl, and Bernhard Arinstein**, the nature of the butyric acid and butyl alcohol fermentation; fixation of acetaldehyde as a decomposition product; transformation of the aldol of pyruvic acid into butyric acid; production of higher fatty acids from sugar, A., i, 91.
- Neuberg, Carl, and Clara Cohen**, the formation of acetaldehyde and the realisation of the second form of fermentation with various fungi, A., i, 304.
- Neuberg, Carl, and O. Dalmer**, crystalline salts of some physiologically important sugar phosphates, A., i, 920.
- Neuberg, Carl, and Julius Hirsch**, classification of carboligase, A., i, 600.
- Neuberg, Carl, and Ludwig Liebermann**, monosulphates of dextrose and sucrose. III., A., i, 222.
carboligase. II., A., i, 305.
- Neuberg, Carl, and Heinz Ohle**, sulphur in agar, A., i, 323.
carboligase. III. and IV., A., i, 480, 540.
- Neuberg, Carl, Elsa Reinfurth, and Marta Sandberg**, stimulants of alcoholic sugar fission. VII., A., i, 306.
- Neuberg, Carl, and Marta Sandberg**, stimulants of alcoholic sugar fission. VIII. and IX., A., i, 408.
- Neuberger, A.** See *W. Meigen*.
- Neuburger, Maximilian Camillo**, isotope of the radio-elements and Meitner's nucleus model, A., ii, 107.
calculation of the branching relationships for dual α -disintegration and the Meitner nucleus model, A., ii, 107.
the Meitner nuclear model of the radio-elements as the basis of a relationship between the range and total number of nuclear particles of the α -radiators, A., ii, 183.
- Neuburger, Maximilian Camillo**, existence of isotopes of uranium and the Meitner nuclear model, A., ii, 185.
a new model of the nucleus, A., ii, 203.
isotope of the radio-elements, A., ii, 250.
existence of isotopes of the disintegration products of actinium and the Meitner nuclear model, A., ii, 340.
the atomic weight of isohelium, A., ii, 365.
the genesis of the elements, A., ii, 365.
the Neuburger nuclear model, A., ii, 702, 839.
- Neugebauer, H.**, dispersoid chemistry of gypsum. II., A., ii, 643.
- Neugebauer, W.** See *Fritz Mayer*.
- Neuhausen, Benjamin S.**, free and bound water in the blood, A., i, 696.
condition of electrolytes in the blood, A., i, 891.
sodium amalgam electrode for the determination of sodium ion [concentration], A., ii, 610.
solubility of gases in liquids, A., ii, 621.
electrolytic preparation of calcium amalgam, A., ii, 643.
- Neuhausen, Benjamin S., and Eli Kennerly Marshall, jun.**, electrochemical study of the condition of several electrolytes in the blood, A., i, 1085.
- Neuhausen, Benjamin S., and Walter A. Patrick**, organogels of silicic acid, A., ii, 144.
system, ammonia-water as a basis for a theory of the solution of gases in liquids, A., ii, 264.
- Neumann, Bernhard**, [with *Bergdahl, Broy, and Karwat*], the catalytic formation of hydrogen chloride from hydrogen and chlorine without explosion, A., ii, 44.
- Neumann, Emanuel Franz.** See *Anton Kailan*.
- Neumann, Felix.** See *Franz Faltis*.
- Neumann, Heinrich.** See *Roland Scholl*.
- Neumann, O.** See *Arnold Eucken*.
- Neuschloss, S. M.** See *Alexander Ellinger, and Otto Riesser*.
- Neuwirth, Isaac**, carbohydrate metabolism. III. A study of urinary sugar excretion in twenty-six individuals, A., i, 485.
- Neuwirth, Isaac, and Israel Simon Kleiner**, the blood-sugar content of capillary blood as compared with that of venous blood, A., i, 1208.
- Nevens, W. E.** See *T. S. Hamilton, and H. H. Mitchell*.

- Neville, Harvey A.** See *Hugh Stott Taylor*.
- Newbery, Edgar**, the overvoltage of the mercury cathode, T., 7.
- Newcomb, Walter H.** See *Oliver Kamm*.
- Newman, F. H.**, active modifications of hydrogen and nitrogen produced by α -rays, A., ii, 279.
absorption of hydrogen by elements in the electric discharge tube, A., ii, 546.
active hydrogen and nitrogen, A., ii, 639.
- Nicholas, Henry O.** See *Harry B. Weiser*.
- Nichols, Edward L.**, and *Horace L. Howes*, emission bands of erbium oxide, A., ii, 411.
the luminescence of incandescent solids, A., ii, 597.
- Nichols, Edward L.**, and *D. T. Wilber*, the luminescence of certain oxides sublimed in the electric arc, A., ii, 105.
flame excitation of luminescence, A., ii, 806.
- Nichols, H. J.**, the production of carbon dioxide by the typhoid bacillus and the mechanism of the Russell double sugar tube, A., i, 501.
- Nicholson, John William**, the binding of electrons by atoms, A., ii, 544.
the difference between series spectra of isotopes, A., ii, 599.
- Nicholson, Samuel T., jun.** See *Roger S. Hubbard*.
- Nickelson, Stanley Arthur.** See *Sidney Hartley Bales*.
- Nickolls, Lewis Charles.** See *Christopher Kelk Ingold*.
- Nieloux, Maurice**, and *Georges Welter*, does cyanic acid exist in the blood? A., i, 789.
quantitative gravimetric micro-analysis of urea; application to the estimation of urea in 1 c.c. of blood, A., ii, 170.
quantitative microanalysis, A., ii, 779.
- Nicolardot, Paul, Max Geloso**, and *Antoine Régla*, the estimation of manganese by Knorre's persulphate method, A., ii, 398.
- Nicolet, Ben H.**, C_{18} fatty acids. II. The relation of oleic and elaidic acids to their halogen additive products, A., i, 106.
the existence and reactions of positive halogen attached to carbon in aromatic compounds, A., i, 121.
- Nicolet, Ben H.**, and *Henry L. Cox*, C_{18} fatty acids. III. Four tetrahydroxy-stearic acids derived from linoleic acid, and their significance with regard to the linoleic acid of common oils, A., i, 320.
- Nicolet, Ben H.**, and *Alfred E. Jurist*, C_{18} fatty acids. IV. A rearrangement of the benzilic acid type in the aliphatic series, A., i, 622.
- Nicolet, Ben H.**, and *Joseph J. Pelc*, C_{18} fatty acids. V. Molecular rearrangements in some derivatives of unsaturated higher fatty acids, A., i, 643.
- Niemann, W.** See *William Küster*.
- Nierenstein, Maximilian**, catechutannins. I. Paullinia tannin, T., 23.
the constitution of catechin. IV., T., 604.
Waage's phytochemical synthesis of phloroglucinol from dextrose, A., i, 136.
gallotannin, A., i, 266.
- Nierenstein, Maximilian.** See also *Howard & Sons, Ltd.*
- Nieuwland, Julius A.** See *P. Benedict Oberdoerfer, Lawrence E. Rombaut*, and *Richard R. Vogt*.
- Niewiazski, S.** See *Emil Briner*.
- Niggli, Paul**, crystal structure and atomic constitution. I. and II., A., ii, 36.
- Nijk, D. R.**, comparative study of ring-substituted phenylphosphinic and phenylarsinic acids, A., i, 960.
- Nikitin, N. I.**, pressure of saturated carbonyl chloride vapour, A., ii, 847.
- Nikolai, Ferdinand**, iodometric estimation of arsenic and antimony sulphides, A., ii, 585.
- Nilsson, Harold**, hesperidine-like substances in the Umbelliferae, A., i, 211.
- Nishikawa, Hidejiro**, and *Robert Robinson*, the hydroxybenzoylphloroglucinols, T., 839.
- Nishizawa, Fushichi**, composition of bird-lime. II., A., i, 652.
- Nitsche, Rudolf.** See *Erich Krause*, and *Hermann Leuchs*.
- Nixon, C. E.** See *Grete Egerer-Scham*.
- Noddack, W.** See *John Eggert*.
- Nodder, Charles Reynolds**, flax and kindred fibres. I. Behaviour and structure of textile fibres, and a convenient method of distinguishing flax from hemp, A., ii, 791.
- Nolan, P. J.**, evidence for the existence of homogeneous groups of large ions, A., ii, 251.

- Nolte, Otto**, detection and estimation of nitrate nitrogen in urine and serum, A., ii, 583.
- Nomidis, Ménelas**. See *Henri Labbé*.
- Nonnenbruch, W.**, concentration of the blood. II. The action of diuretics of the purine group on the exchange of substances between the tissues and the blood, A., i, 79.
- Nordfeldt, E.**, importance of acidity for cyanohydrin synthesis and the non-existence of Rosenthaler's *syn-emulsin*, A., i, 66.
the asymmetric action of emulsin in the benzaldehydecyanohydrin synthesis, A., i, 1077.
- Nordlund, Folke**. See *Hans von Euler*.
- Nordström, A. M.**, phenylcamphenol, A., i, 944.
- Norgaard, A.**, influence of water supply on the content of reducing substances in the blood and urine, A., i, 892.
- Norgaard, A.** and *H. C. Gram*, relation between the chloride content of the blood and its volume per cent. of cells, A., i, 287.
- Norman, G. F.** See *Carl L. A. Schmidt*.
- Normand, G.** See *André Wahl*.
- Normann, W.**, catalytic addition of hydrogen; influence of oxygen on the catalyst, A., ii, 631.
- Norris, Frederick Walter**. See *Donald Herbert Frank Clayson*.
- Norris, (Miss) Mabel Harriet** [with (*Miss*) *D. H. Falkner*, and (*Miss*) *M. C. Price*], the constitution of soap solutions; hexadecanesulphonic (cetylsulphonic) acid and other sulphonates, T., 2161.
- Norris, (Miss) Mabel Harriet**, and *James William McBain*, a study of the rate of saponification of oils and fats by aqueous alkali under various conditions, T., 1862.
- Norrish, R. G. W.**, transition elements and the octet theory: a new arrangement of the rare earth elements in the periodic classification, A., ii, 211.
- Northrop, John H.**, the inactivation of trypsin. I. and II. The equilibrium between trypsin and the inhibiting substance formed by its action on proteins. III. Spontaneous inactivation, A., i, 282.
kinetics of trypsin digestion, A., i, 698.
- Notzitz, von**, basic exchange in soils, A., i, 511.
- Notenbaart, L.**, rapid estimation of acetic and butyric acids in mixtures, A., ii, 791.
- Nowak, Gisela**, and *Julius Zellner*, comparative plant chemistry. II. The berry fruit of some Caprifoliaceae, A., i, 615.
- Nowrey, Joseph E.** See *Frederick S. Hammett*.
- Noyes, Arthur Amos**, and *H. A. Wilson*, thermal ionisation of gaseous elements at high temperatures; confirmation of the Saha theory, A., ii, 810.
- Noyes, William Albert**, attempt to prepare nitro-nitrogen trichloride. II. Behaviour of mixtures of nitrogen and chlorine in a flaming arc, A., ii, 143.
- Noyes, William Albert**, and *Walter F. Goebel*, catalysis of the formation and hydrolysis of acetamide by acetic acid, A., i, 1127.
- Noyes, William Albert**, and *Thomas A. Wilson*, ionisation constant of hypochlorous acid; evidence for amphoteric ionisation, A., ii, 692.
- Noyes, William Albert**. See also *Howard M. Chiles*, *George H. Coleman*, and *Harry L. Lochte*.
- Noyes, William Albert, jun.** See *George Ernest Gibson*.
- Noyons, A. K.**, physical method for the estimation of carbon dioxide in the respiratory air, A., ii, 868.
- Nüsslein, J.** See *Walther Diltthey*.
- Nugent, T. C.**, an inhibition period in the separation of an emulsion, A., ii, 625.
- Nybergh, Bertil**, the mobility and valency demand of the ethyl group in the pinacolin transformation, A., i, 802.
the reverse pinacolin transformation, A., i, 918.
- Nyswander, R. E.**, *Samuel Colville Lind*, and *R. B. Moore*, the spectrum of radium emanation, A., ii, 803.
- Oakes, Earle T.**, and *Henry M. Salisbury*, use of phthalate solutions for hydrogen electrode standards, A., ii, 468.
- Oakes, Earle T.** See also *Clarke E. Davis*.
- Obajdin, Milivoj**, the reaction of magnesium salts with acid potassium pyrantimonate, A., ii, 501.
- Obaton, Fernand**. See *Marc Romieu*.
- Oberbach, J.** See *Alfred Benrath*.
- Oberdoerfer, P. Benedict**, and *Julius A. Nieuwland*, compounds of acetylene with silver phosphate and silver arsenate, A., i, 515.
- Obladen, A.** See *Alfred Benrath*.

- Oblata, J.**, low-voltage standard cells, A., ii, 848.
- O'Brien, H. R.**, and **W. L. Parker**, solubility of carbon monoxide in serum and plasma, A., i, 394.
- Ochi, Hideo**, semicarbazide hydrochloride, A., i, 723.
- Ochi, Shuichirō, Yoichi Onozawa**, and **The Tokyo Industrial Laboratory**, preparation of acetaldehyde from acetylene, A., i, 523.
- Ochs, Kurt**. See **Karl Ziegler**.
- Oddo, Bernardo**, and **R. Binaghi**, action of polyhalogenated compounds of methane and ethane on magnesiyl [magnesium alkyl] compounds, A., i, 313.
new synthesis of primary alcohols, and constitution of hydrogen peroxide, A., i, 314.
- Oddo, Bernardo**, and **G. Sanna**, new syntheses in the indole group. VIII., A., i, 371.
- Oddo, Giuseppe**, mesohydry. II., A., ii, 367.
mesohydry. III. Relation between the affinity of acids and the atomic ratio O:H in their functional groups, and the constitutional formulæ of the acids, A., ii, 367.
- Oddy, Harold G.** See **Walter A. Lawrance**.
- Odelga, Friedrich**. See **Robert Kramann**.
- Oehlert, H.** See **Richard Stoermer**.
- Ohlm, Lars William**, thermostat arrangement for the determination of the effect of temperature on diffusion, A., ii, 621.
- Oesterle, O. A.**, volatile substances from the bark of *Rhamnus frangula*, A., i, 100.
hyssopin, A., i, 849.
- Oesterlin, C.** See **Georg Schroeter**.
- Östling, Gustav Jim**, the Grignard reaction, A., i, 843.
- Offner, M.** See **Joaquin Enrique Zanetti**.
- Ogasawara, K.** See **E. Tokuko**.
- Ogata, Akira**, and **Chaji Miyashita**, homologues of menthol, A., i, 844.
- Ohle, Heinz**, carbohydrate-sulphates. IV., A., i, 986.
the constitution of vacciniin, A., i, 1018.
- Ohle, Heinz**. See also **Carl Neuberg**.
- Ohlmer, Elsa**. See **A. Morgen**.
- Ohlsson, Erik**, dependence of the action of succinodehydrogenase on hydrogen-ion concentration, A., i, 785.
- Ohma, S.**, classification of aromatic odours in sub-classes, A., i, 1090.
- Ohmann, O.**, preparation of aluminium carbide and of marsh gas, A., i, 2.
- Okada, Seizaburo**, and **Toworu Hayashi**, the amino-acid nitrogen content of the blood, A., i, 483.
- Okagawa, M.** See **Yashiro Kotake**.
- Okasaki**. See **Riko Majima**.
- Okubo, Junzō**, structure of the second cyanogen band, A., ii, 601.
the lanthanum violet bands and the associated lines, A., ii, 676.
- Okushima, Kwanichiro**, caffeine excretion in urine after tea and coffee drinking in man, A., i, 703.
- Oldenberg, Otto**, the structure of the Balmer series, A., ii, 410.
fine structure of the red hydrogen line, A., ii, 725.
- Oleson, Harriet Curry**. See **Katharine Blunt**.
- Oliveri-Mandalà, E.**, behaviour of azides of acids, A., i, 478.
action of hydrogen peroxide on nitriles and on amides: formation of hydroxamic acids, A., i, 544.
mechanism of reaction of aliphatic diazo-compounds, A., i, 994.
bases obtained in the decomposition of the azides of thiocarbamic acids, A., i, 1007.
addition of azoimide at contiguous double linkings. VIII. 5-Anilino-1-phenyltetrazole and the azide of dithiocarbamic acid, A., i, 1069.
- Oliveri-Mandalà, E.**, and **G. Comella**, normal chromium azide and the formation of complex salts, A., ii, 382.
- Olivier, E.**, the preparation of lead test-papers and notes on the titration of zinc with sodium sulphide, A., ii, 396.
- Olivier, Simon Cornelis Johannes**, the structural formula of caoutchouc according to Harries, A., i, 45.
the influence of some substituents in the benzene ring on the mobility of chlorine in the side-chain in its relation to the problem of substitution in the benzene ring, A., i, 646.
the chlorination of benzyl chloride, A., i, 726.
- Olmstead, L. B.** See **R. O. E. Davis**.
- Olsen, O.**, the action of kaolin on the terminal component and on the tributyrin hydrolytic capacity of guinea pig's serum, A., i, 292.
- Olsson, Hugo**. See **Lennart Smith**.
- Olsson, Urban**, inhibition phenomena in amylases, A., i, 390; ii, 401.

- Onnes, Heike Kamerlingh, and J. Palacios Martinez**, vapour pressure of hydrogen; determinations in the region of liquid hydrogen, A., ii, 759.
- Onnes, Heike Kamerlingh.** See also *J. Palacios Martinez, Émile Mathias, and Jean Timmermans.*
- One, Kashichi**, electrolytic reactions of naphthalene and its derivatives. III. Electrolytic oxidation of α -naphthylamine and *ar*-tetrahydro- α -naphthylamine, A., i, 1008.
- Onozawa, Yoichi.** See *Shuichirō Ochi.*
- Oparin, Alexander**, the green respiratory pigment and its importance in the oxidation of protein substances in the sprouting seeds of *Helianthus annuus*, A., i, 309.
- Opfermann**, estimation of α -, β -, and γ -cellulose, A., ii, 665.
- Oppenheimer, E.** See *Fr. Baur.*
- Oppenheimer, Gertrud.** See *Richard Willstätter.*
- Oreel, J.**, composition of aerinite, A., ii, 651.
- Orékhoﬀ, Alex., and Marc Tiffeneau**, the dehydration of benzylhydrobenzoin ($\alpha\beta$ -triphenylpropane- $\alpha\beta$ -diol); formation of benzyl diphenylmethyl ketone (semipinacolic transposition) and of diphenylindene (cyclisation), A., i, 458.
- Orlova, Lidie.** See *Nikolai Schilov.*
- Ormandy, William R., and E. C. Craven**, the system ethyl alcohol-water-aromatic hydrocarbons from 30° to -30°, A., i, 215.
- Orndorff, William Ridgely, R. C. Gibbs, M. Scott, and S. D. Jackson**, the absorption spectra of *o*-cresolsulphonphthalein, A., ii, 806.
- Orndorff, William Ridgely, and Louise Kelley**, *o*-4'-hydroxybenzoylbenzoic acid and some of its derivatives, A., i, 833.
- Orthner, Ludwig.** See *Karl Freudenberg, and Kurt Heinrich Meyer.*
- Orthner, Karl.** See *Wilhelm Manchot.*
- Orton, C. R., Elmer Verner McCollum, and Nina Simmonds**, the presence of the antineuritic substance, water-soluble B, in chlorophyll-free plants, A., i, 974.
- Osaka, Yukichi**, the solubility of calcium carbonate in water in equilibrium with a gaseous phase containing carbon dioxide, A., ii, 830.
- Osaka, Yukichi, and Kinji Ando**, the compounds of carbamide and benzoic acid, A., i, 1127.
- Osborne, Thomas Burr, Alfred John Wakeman, and Charles Samuel Leavenworth**, the proteins of the alfalfa plant [lucerne], A., i, 99.
- water-soluble constituents of the alfalfa plant [lucerne], A., i, 1104.
- Osswald, Philipp.** See *Wilhelm Windisch.*
- Ost, Hermann, and G. Knoth**, cellobiose, A., i, 526.
- Osterburg, Fritz.** See *Hermann Leuchs.*
- Osterhout, Winthrop John Vanleuren**, direct and indirect determinations of permeability, A., i, 308.
- Ostermeier, Hermann.** See *Hugo Weil.*
- Ostertag, Rudolf.** See *Hartwig Franzen.*
- Ostwald, Wa.**, homogeneity and dispersity, A., ii, 131.
- Ostwald, Wolfgang**, dispersoid analysis of non-aqueous systems, A., ii, 199.
- electro-adsorption as a purely chemical process, A., ii, 427.
- Ostwald, Wolfgang, and Friedrich Vincenz von Hahn**, a kinetic precipitation measurer, A., ii, 205.
- Ostwald, Wolfgang, and Ramon de Izaguirre**, more general theory of the adsorption of solutions, A., ii, 480.
- Ostwald, Wolfgang, and Alfred Kuhn**, acid taste, A., i, 84.
- role of acids in peptic digestion, A., i, 598.
- Oswald, Adolf**, the physiological action of metallic amines and allied compounds, A., i, 498.
- Oswald, Victor.** See *Robert Kremann.*
- Otsuka, Ichiro.** See *Takaoki Sasaki.*
- Ott, Erwin**, the constitution and tautomeric equilibrium of the two phthalic acid tetrachlorides, A., i, 834.
- preparation of cyanuric triazide, A., i, 1128.
- Ott, Erwin, and Fritz Eichler** [with *Otto Lüdermann, and Heinrich Heilmann*], natural and artificial pepper-substances. II. The chavicine of pepper-resin, the primarily active constituent of black pepper, A., i, 1026.
- Ott, Erwin, and Bernhard Löpmann**, simple cyano- and cyanuric compounds. III. Malononitrile and its halogenation, A., i, 643.
- Ott, Erwin, and Ernst Nauenz**, vanillin isomerides of the resorcylic series, A., i, 555.
- Ott, Erwin, and Karl Schmidt**, the preparation of carbon suboxide on a larger scale and the properties of pure carbon suboxide, A., ii, 641

- Ott, *Erwin*, and *Kurt Zimmermann*, natural and artificial pepper-substances and the relation between chemical constitution and peppery taste, A., i, 137.
- Ottenstein, *Berta*, estimation of urinary colloids by the gold number, A., i, 609.
- Otto, *J.* See *Ludwig Holborn*.
- Owen, *E. A.*, and *Winifred E. Page*, estimation of the radium content of radioactive luminous compounds, A., ii, 108.
- Oxley, *A. E.*, magnetism and atomic structure. II. The constitution of the hydrogen-palladium system and other similar systems, A., ii, 469.
- P.
- Paal, *Carl*, and *Hermann Steyer*, colloidal copper hydroxide, A., ii, 216.
variously coloured modifications of colloidal copper, A., ii, 294.
amalgams. III. Colloidal copper amalgam, A., ii, 446.
- Paassen, *P. van*, calcium in the blood, A., i, 1208.
- Paciello, *A.* See *Vittorio Cuttica*.
- Paddon, *W. W.*, dyeing of deaminated wool, A., i, 608.
mordanting of wool with potash alum, A., ii, 822.
- Padoa, *Maurizio*, absolute value of the energy of the linkings between the atoms of compounds, A., ii, 27.
specific heats. II. A., ii, 348.
- Pactold, *Hans*. See *Heinrich Biltz*.
- Page, *Winifred E.* See *E. A. Owen*.
- Pahle, *Kristian*. See *Haakon Goldschmidt*.
- Palache, *Charles*. See *Waldemar Lindgren*.
- Palfray, *L.*, neutral homocamphoric esters and their products of reduction, A., i, 548.
- Palkin, *Samuel*, and *M. Harris*, quinolines. I. Preparation of 6-ethoxy-2:4-dimethylquinoline, A., i, 951.
- Palladin, *Wladimir*, and *Helene Popov*, the genesis of amylase and maltase in plants, A., i, 614.
- Palmer, *Albert Donald*. See *Harold King*.
- Palmer, *Charles Shattuck*, and *Roger Adams* [with *W. A. Carothers*, *E. E. Parks*, *G. O. Burr*, and *J. S. Pierce*], the reactions of the arsines. II. Condensation of aromatic primary arsines with aldehydes, A., i, 785.
- Palmer, *Leroy S.*, influence of various antiseptics on the activity of lipase, A., i, 886.
the effects of impurities on the ionisation potentials measured in thermionic valves, A., ii, 108.
- Palmer, *William George*, the catalytic activity of copper, A., ii, 437.
- Palmer, *Walter W.*, *Dana W. Atchley*, and *Robert F. Loeb*, the regulation of osmotic pressure. II. The effect of albumin on the conductivity of a sodium chloride solution, A., i, 692.
- Palmer, *Walter W.* See also *Henry Jackson, jun.*
- Pamflov, *A. V.*, oxidation of *o*-toluenesulphonamide, A., i, 928.
rôle of chromates in electrolysis for chlorate, A., ii, 712.
- Pamflov, *A. V.*, and *N. N. Petin*, kinetics of inductive processes; Schönbein's reaction, A., ii, 835.
- Paneth, *Fritz*, peculiar catalytic action in chemical synthesis by glow discharge, A., ii, 363
radioactive indicators, A., ii, 785.
- Paneth, *Fritz*, and *Adolf Johannsen*, polonium hydride. II., A., ii, 777.
- Paneth, *Fritz*, *Adolf Johannsen*, and *Max Matthies*, the preparation of gaseous metallic hydrides from alloys and solutions, A., ii, 383.
- Paneth, *Fritz*, *Max Matthies*, and *Edgar Schmidt-Hebbel*, the preparation of gaseous metallic hydrides by the spark discharge, A., ii, 383.
- Paneth, *Fritz*, and *Edgar Schmidt-Hebbel*, germanium hydride, A., ii, 776.
- Paneth, *Fritz*, and *Walter Vorwerk*, method for the determination of the surface of adsorbing powders, A., ii, 618.
thickness of the adsorbed layer in the adsorption of dyes by crystals, A., ii, 619.
- Paolini, *Vincenzo*, isomeric amyrols, A., i, 754.
- Paolini, *Vincenzo*, and *S. Scelba*, polysilylides, A., i, 746.
- Papaconstantinou, *Basil Constantine*. See *Jnanendra Nath Mukherjee*.
- Papendieck, *A.* See *Otto Sohum*.
- Papish, *Jacob*, germanium. II. Identification of germanium by its visible arc spectrum, A., ii, 163.
flame reactions of thallium, A., ii, 528.
- Papish, *Jacob*. See also *Louis Monroe Dennis*.
- Paris, *G.*, the biochemistry of tobacco. II. Tobacco seeds, A., i, 211.
- Parish, *H. C.* See *Roger C. Griffin*.

- Parker, F. W.**, displacement method for obtaining soil solution, A., i, 511.
classification of soil moisture, A., i, 616.
- Parker, H. K.** See *L. van Doren*.
- Parker, William Bayley**, and *Gartha Thompson*, the variation of refractive index and density of benzene with temperature, T., 1341.
- Parker, W. L.** See *Vernon C Allison*, *G. W. Jones*, and *H. R. O'Brien*.
- Parkes, Deric William.** See *Frederick Daniel Chattaway*.
- Parkhurst, Reginald B.** See *William Thompson Smith*.
- Parks, E. E.** See *Charles Shattuck Palmer*.
- Parnas, Jakob K.**, and *Richard Wagner*, observations on sugar synthesis. I., A., i, 487.
estimation of small quantities of nitrogen by Kjeldahl's method, A., ii, 312.
- Parnas, Jakob K.** See also *Richard Wagner*.
- Parr, Paul.** See *Bruno Emmert*.
- Parsons, Llewellyn B.** See *James H. Walton*.
- Parsons, Leon Woodman.** See *Gregory Paul Baxter*.
- Parsons, T. R.** See *Joseph Barcroft*.
- Parsons, W.** See *Joseph Barcroft*.
- Partington, James Riddick**, the energy of gaseous molecules, A., ii, 614.
chemical constants of some diatomic gases, A., ii, 839.
- Partington, James Riddick**, and *H. J. Cant*, specific heats of ammonia, sulphur dioxide, and carbon dioxide, A., ii, 191.
- Partington, James Riddick.** See also *Frank Ward Bury*, and *(Mrs.) K. Stratton*.
- Partos, S.**, the hæmochrome of Herzfeld and Klinger, A., i, 597.
- Pascal, Paul**, preparation of ethyl alcohol, A., i, 8.
magneto-chemical investigation of the constitutions in mineral chemistry; the acids of phosphorus, A., ii, 285.
magneto-chemical investigations of constitutions in mineral chemistry; the acids of arsenic, A., ii, 564.
- Paschke, Fritz**, derivatives of straw lignin, A., i, 925.
- Pascual Vila, J.** See *Antonio Garcia Bandú*.
- Passerini, M.**, isonitriles. III. Reactions with the hydrates of halogenated aldehydes, A., i, 731.
- Passini, Fritz**, degradation of bile pigments by anaerobic putrefactive intestinal bacteria, A., i, 966.
- Pastureau, and Henri Bernard**, the chlorohydrin of mesityl oxide and its transformation into the chlorohydrin of tetramethylglycerol, A., i, 717.
- Paternò, Emanuele**, aniline arsenates, A., i, 731.
- Paton, R. F.** See *H. M. Randall*.
- Patrick, Walker A.**, and *F. V. Grim*, heat of wetting of silica gel, A., ii, 122.
- Patrick, Walter A.** See also *L. Y. Davidheiser*, *Edward O. Holmes, jun.*, *J. Fitch King*, and *Benjamin S. Neuhäusen*.
- Patterson, Jocelyn.** See *James Colquhoun Irvine*.
- Patterson, R. A.** See *William Duane*.
- Patterson, Thomas Stewart**, a property of ethyl tartrate, T., 1042.
- Patty, F. A.**, the production of hydrocyanic acid by *Bacillus pyocyaneus*, A., i, 407.
- Patzauer, A.**, reaction for blood with hydrogen peroxide in statu nascendi, A., ii, 172.
- Paul, Arthur E.**, and *Erwin H. Berry*, the Kjeldahl nitrogen method and its modifications, A., ii, 82.
- Pauli, Wolfgang**, mobility of protein ions, A., i, 478.
- Pauli, Wolfgang.** See also *Mona Adolf*, and *Albert Fernau*.
- Pauli, W., jun.**, model of the ionised hydrogen molecule, A., ii, 703.
- Pauly, Hermann**, the osculation of physical constants in homologous series, A., i, 1.
- Pauly, Hermann**, and *Ernst Ludwig*, glyoxalinedicarboxylic acid for the recognition and separation of organic bases, A., i, 953.
glucosamine as the basis of formation of heterocyclic compounds, A., i, 953.
- Pavlovič, R.** See *Peter Bona*.
- Pavolini, T.** See *A. Gaviati*.
- Payá, M.**, and *Enrique Moles*, density of atmospheric nitrogen; a small anomaly in the air of Madrid, A., ii, 762.
- Payman, Joseph Baron.** See *British Dyestuffs Corporation, Ltd.*
- Payman, William**, and *Richard Vernon Wheeler*, the combustion of complex gaseous mixtures, T., 363.
- Payne, Wilbur B.** See *Merrill C. Hart*.
- Payne, Wilfred Walter.** See *Patrick Playfair Laidlaw*.

- Peacock, David Henry.** See *David Segaller*.
- Pearce, J. Newton, and Harry B. Hart,** free energy of dilution and the activities of the ions of potassium bromide in aqueous solutions, A., ii, 254.
- Pearson, Archibald Ramsden.** See *William Arthur Bone*.
- Pease, Robert Norton,** sizes of atoms in crystals, A., ii, 428.
sizes of atoms in diamond type crystals, A., ii, 634.
atoms and electrons, A., ii, 757.
- Pease, Robert Norton, and Hugh Stott Taylor,** reduction of copper oxide by hydrogen, A., ii, 148.
catalytic formation of water vapour from hydrogen and oxygen in the presence of copper and copper oxide, A., ii, 701.
- Pecker, Henri,** characteristics of identity of a water distilled from the cherry laurel, A., ii, 537.
- Peddie, W.,** the importance of Whitaker's atomic model and of other atomic models, A., ii, 633.
- Pedley, Frank G.** See *Alfred T. Shohl*.
- Pedotti, Fausto,** the influence of lack of calcium in the diet on the respiratory basal metabolism, A., i, 286.
- Peiser, E.** See *Hermann Stendel*.
- Pekelharing, Cornelis Adrianus,** movement of pepsin in agar-agar gels with and without protein, A., i, 388.
- Pélabon, Henri (Joseph Léonard Ferdinand),** the constitution of selenium, A., ii, 141.
action of selenium on gold, A., ii, 303.
- Pelc, Joseph J.** See *Ben H. Nicolet*.
- Pelkan, K. F.,** physiology of the phenols. I. The estimation of phenols in the blood, A., ii, 399.
- Pelkan, K. F., and G. H. Whipple,** liver function. III. Phenol conjugation as influenced by liver injury and insufficiency, A., i, 490.
physiology of the phenols. II. Absorption, conjugation, and excretion, A., i, 499.
- Pelkan, K. F.** See also *W. R. Bloor*.
- Pellizzari, Guido,** action of cyanogen haloids on phenylhydrazine. VI. *o*-Phenylenesammelyl chloride, A., i, 585.
- Pelzer, J.** See *Paul Jacobsen*.
- Pemberton, H. V.** See *Lester F. Hoyt*.
- Penfold, Arthur Ramon,** a critical examination of the aromatic aldehydes occurring in certain eucalyptus oils, T., 266.
- Penfold, Arthur Ramon,** the essential oils of *Leptospermum flavescens* var. *grandiflorum* and *Leptospermum odoratum*, A., i, 44.
position of the double linking in piperitone. I. and II., A., i, 259, 1035.
essential oil of the leaves of *Doryphora sassafras*, A., i, 847.
- Penfold, Arthur Ramon.** See also *Henry George Smith*.
- Pénkava, J.** See *Julius Stoklasa*.
- Penkner, Wilhelm.** See *Robert Kremann*.
- Pennycuik, Stuart Wortley,** racemic acid in solution, A., i, 624.
- Pentimalli, F.,** protein intoxication. I. Introduction. II. Toxicity of albumin and its derivatives. Toxicity of peptones. IV. Toxicity of milk and its derivatives. V. Behaviour of blood pressure and of respiration. VI. Behaviour of body temperature. VII. Nystagmus. VIII. Morphological changes of the blood, A., i, 302.
- Pépin, C., and G. Réaumont,** sulphonated derivatives of the naturally occurring sulphidic hydrocarbons, A., ii, 784.
- Percy, R.** See *August Hagenbach*.
- Pereira.** See *Giral Pereira*.
- Perina, G.** See *G. Canneri*.
- Perkin, Arthur George, and George Douglas Spencer,** some reactions of benzanthrone, T., 474.
- Perkin, Arthur George, and Yoshisuke Uyeda,** occurrence of a crystalline tannin in the leaves of the *Acer ginnala*, T., 66.
- Perkin, Arthur George, and Thomas William Whattam,** some products of the reduction of 2-hydroxyanthraquinone, T., 289.
- Perkin, Arthur George.** See also *Geoffrey Gordon Bradshaw* and *British Dyestuffs Corporation, Ltd.*
- Perkin, William Henry, jun., and Alan Francis Titley,** conditions of formation of rings attached to the *o*-, *m*-, and *p*-positions of the benzene nucleus. I. The action of sodium on *o*-phenylenediacetic ester, T., 1562.
- Perkin, William Henry, jun.** See also *George Roger Clemons, William Davies, and William Ogilvie Kermack*.
- Perkins, Granville A.,** the expression of the octet theory of valence in structural formulae, A., ii, 138.
unsymmetrical addition to the double bond. I. A theory of the reaction mechanism of the direct union, A., ii, 438.

- Perkins, Granville A.**, the structure of chlorine dioxide and related compounds, A., ii, 441.
- Perling, A.**, the hydrolytic decomposition of the bismuth salts of phenol-carboxylic acids, A., i, 252.
- Perman, Edgar Philip**, the properties of ammonium nitrate. IV. The reciprocal salt-pair, ammonium nitrate and sodium chloride, T., 2473.
- Perquin, J. N. J.** See *Hein Israel Waterman*.
- Perren, Edward Arthur**. See *Christopher Kelk Ingold*.
- Perrin, Jean**, catalysis with special reference to newer theories of chemical action. I. The radiation theory of chemical action; (1) radiation and chemistry, A., ii, 628.
- Persch, Walter**. See *Hans Pringsheim*.
- Peski, A. J. van**, the mixed anhydrides of sulphuric acid and carboxylic acids. II. *n*-Butyrylsulphuric acid, A., i, 106.
- Peskov, N. P.**, solutions, suspensions, colloids, A., ii, 825.
sensitisation of coagulation processes; colloids as indicators of photo-electric effects, A., ii, 828.
- Petényi, Géza**, and *Heinrich Lax*, action of adrenaline on the blood-sugar, A., i, 403.
- Peter, A. M.** See *G. Davis Buckner*.
- Peters, John P., Glenn E. Cullen**, and *Joshua Harold Austin*, gas and electrolyte equilibria in blood. II The reversibility of the effects of changes in carbon dioxide and oxygen tensions on the carbon dioxide content of defibrinated horse blood, A., i, 1207.
- Peters, John P.** See also *Joshua Harold Austin*.
- Peters, Walter**, the addition of organic bases to metallic salts, A., i, 48.
- Petersen, Agnes**. See *Johannes Nicolaus Brønsted*.
- Peterson, W. H., Edwin Broun Fred.** and *J. A. Anderson*, fermentation of hexoses and related compounds by certain pentose-fermenting bacteria, A., i, 971.
- Peterson, W. H., Edwin Broun Fred.** and *E. G. Schmidt*, fermentation of pentoses by moulds, A., i, 1220.
- Peterson, W. H.** See also *O. R. Brunkow*, and *Edwin Broun Fred*.
- Petin, N. N.** See *A. V. Pamfilov*.
- Petitjean, Fernand**, the manipulations preparatory to the estimation of amino-acids in blood, A., ii, 536.
- Peto, Raymond Harold Kelsall**. See *George Ingle Finch*.
- Petrie, P. A.** See *John Cunningham McLennan*.
- Petry, Eugen**, conditions for the biological action of Röntgen rays. I, A., i, 204.
- Petschacher, Ludwig**, micro-analyses by Bang's methods. I, A., ii, 716.
- Peyer, J.** See *Paul Karrer*.
- Peytral, (Mlle) Églantine**, the mode of sudden pyrogenic decomposition of acetic acid at high temperature, A., i, 219.
the mode of pyrogenic decomposition of methyl acetate at high temperature, A., i, 219.
the mode of pyrogenic decomposition of acetone at high temperature, A., i, 222.
- Pfaff, Kaspar**. See *Fritz Mayer*.
- Pfannkuch, E.** See *Karl W. Rosenmund*.
- Pfanstiel, Robert**. See *Herbert S. Harned*.
- Pfleiderer, Georg**. See *Franz Fischer*.
- Pfeiffer, Paul**, betaines. I. Theory of betaines, A., i, 720.
- Pfeiffer, Paul** [with *E. Flater*], quinhydrones of the maleic anhydride series, A., i, 341.
- Pfeiffer, Paul**, and *Gerhard Haefelin*, betaines. II. Betaines of the cinnamic acid series, A., i, 738.
- Pfizenmaier, K.** and *S. Galanos*, the estimation of creatinine, A., ii, 797.
- Pfund, Marion C.** See *Annie Louise Macleod*.
- Pfyl, Balthasar**, ash-alkalinity (of food-stuffs), A., ii, 586.
- Pfyl, Balthasar, U. Reif**, and *A. Hanner*, detection of formaldehyde with phenols, A., ii, 94.
- Philipp, K.** See *Heinrich Kausch von Traubenberg*.
- Philippi, Ernst**, mellitic acid, pyromellitic acid, and their production from carbon by oxidation, A., i, 837.
- Philippi, Ernst**, and *Gertrud Rie*, mellitic acid, pyromellitic acid, and their production from carbon by oxidation. I. Oxidation of carbon by nitric acid, A., i, 837.
- Philippi, Ernst, Reinhard Seka**, and *Norbert Froeschl*, mellitic acid, pyromellitic acid, and their production from carbon by oxidation. III. Synthesis of pyromellitic acid from commercial xylene, A., i, 837.
- Philippi, Ernst, Reinhard Seka**, and *Lilly Robinson*, mellitic acid, pyromellitic acid, and their production from carbon by oxidation. IV. Synthesis of substituted pyromellitic acids, A., i, 837.

- Philippi, Ernst**, and **Richard Thelen**, mellitic acid, pyromellitic acid, and their production from carbon by oxidation. II. Oxidation of carbon by sulphuric acid, A., i, 887.
- Phillips, Max**, the preparation of 6:6'-di- α -hydroxyisopropylindigotin from *p*-cymene, A., i, 955.
- Phragmen, Gosta**. See **Arne Westgren**.
- Picard, A.**, and **E. Stahel**, a new radioactive substance, A., ii, 185.
- Picard, Jean**, the steric formula of the molecule of water, A., ii, 212.
preparation of catalytic copper, A., ii, 216.
detection of oxygen in organic compounds, A., ii, 389.
- Picard, Jean**, and **Ray Q. Brewster**, tetrapropylethane [5-dipropyloctane], A., i, 813.
the three aminotriphenylamines, A., i, 370.
- Picard, Jean**, and **E. Herrmann**, the colour of iodine solutions at low temperature, A., ii, 655.
- Piehler, A. B.**, recovery of solvents from extracted substances, A., ii, 715.
- Pickering, G. F.**, and **Geoffrey E. Cowlishaw**, the relation between the refractive index and the chemical characteristics of oils and fats (glycerides), A., ii, 325.
- Pickering, John William**, and **James Arthur Hewitt**, coagulation of the blood. I. Some physico-chemical aspects of coagulation, A., i, 393.
coagulation of the blood. II. Thrombin and anti-thrombins, A., i, 1208.
- Pickles, Alwyn**, reduction of ferric chloride, A., ii, 299.
- Pico, O. M.**, and **J. Murtagh**, estimation of chlorine in tissues, A., ii, 716.
- Picon, M.**, the action of sodammonium on hexamethylenetetramine, tetramethyldiaminomethane, and ethylideneethylamine, A., i, 1123.
- Picon, M.** See also **P. Lebeau**.
- Pictet, Amé**, and **André Barbier**, new synthesis of glycerol and α -glucoheptol, A., i, 4.
- Pictet, Amé**, and **R. Jahn**, a new depolymerisation product of starch, A., i, 987.
- Pictet, Amé**, and **J. H. Ross**, the polymerisation of levoglucosan, A., i, 525.
- Pictet, Amé**, and **Henry Vernet**, galactosan, A., i, 811.
- Pieraerts, J.**, copal oil, a new fat of the Belgian Congo, A., i, 908.
- Pierce, J. S.** See **Charles Shattuck Palmer**.
- Pierce, W. C.** See **G. Davis Buckner**.
- Pieroh, Kurt**. See **Heinrich Schulze**.
- Pieroni, Antonio**, iodo-derivatives of pyrrole, A., i, 763.
azoxyamides and diazo-compounds, A., i, 1071.
- Pieroni, Antonio**, and **Aldo Moggi**, constitution of certain polypyrroles, A., i, 766.
- Piettre, Maurice**, and **Antony Vila**, separation of the proteins of the serum, A., i, 63.
- Piggot, Charles Snowden**, manganese in the catalytic oxidation of ammonia, A., ii, 142.
- Piggot, Charles Snowden**. See also **T. H. Rogers**.
- Piggott, Henry Alfred**. See **Christopher Kell Ingold**.
- Pighini, Giacomo**, chemical and biochemical investigations of the nervous system under normal and pathological conditions. IX., A., i, 295.
- Pilling, Norman B.**, vapour pressure of metallic calcium, A., ii, 291.
- Piña de Rubies, Santiago**, and **F. Gila Esteban**, chemical and spectrochemical study of Spanish bismuth minerals, A., ii, 576.
- Piña de Rubies, Santiago**. See also **B. Cabrera**.
- Pincass, Heinrich**, attempts to prepare pure calcium cyanide, A., i, 532.
- Pincussen, Ludwig**, the fermentative properties of blood. V. Appearance of ferments in blood after various operations, A., i, 393.
behaviour of oxalic acid in the animal body, A., i, 403.
the influence of illumination on the metabolism of carbohydrates, A., i, 1088.
detection of urea; estimation of proteolytic ferments, A., ii, 884.
- Pincussen, Ludwig**, and **J. L. Anagnostu**, the influence of radiations on the hydrolysis of fats, A., i, 485.
- Pincussen, Ludwig**, and **Aristomenis Floros**, analysis of blood and urine. I., A., ii, 408.
- Pincussen, Ludwig**, and **Kate Momferatos-Floros**, influence of radiation on nucleic acid metabolism, A., i, 395.
analysis of blood and urine, A., ii, 408.
- Pinkard, Frederic William**, and **William Wardlaw**, the action of sulphur on cuprous chloride, T., 1300.
- Pinkard, Frederic William**. See also **William Wardlaw**.
- Pinkus, S.** See **Paul Jacobsen**.

- Pirak, J.**, asymmetric synthesis, A., i, 810.
- Pirani, Marcello von**, and **E. Lax**, electrolytic migration of sodium through glass, A., ii, 817.
- Pittarelli, Emilio**, source of error in tests for acetone, A., ii, 95.
- Piutti, Arnaldo**, and **Enrico Boggio-Lera**, microchemical investigation of arsenic, A., ii, 584.
- Piwowsky, E.**, the constitution of strontium-lead alloys, A., ii, 644.
- Pizarroso, A.** See **Obdulio Fernández**.
- Plagge, H.**, comparative experiments on the inhibitive action of some chlorine derivatives of methane, ethane, and ethylene on fermentation, A., i, 93.
- Planck, Max**, absolute entropy and chemical constants, A., ii, 191.
free energy of gaseous molecules with any [possible] partition of velocities, A., ii, 689.
- Plank, R.**, convertibility of chemical energy and the conception of affinity, A., ii, 357.
- Plantefol, L.**, the toxicity of different nitrophenols towards *Aspergillus niger*, A., i, 204.
- Platon, B.** See **Lennart Smith**.
- Plauson, Herman**, and **Georg von Tischenko**, preparation of hydrocarbons, A., i, 818.
- Plimmer, Robert Henry Aders**, and **John Lewis Rosedale**, distribution of enzymes in the alimentary canal of the chicken, A., i, 485.
- Plotnikow, Joh.**, photochemical studies. XIII. Photopolymerisation of vinyl chloride and the problem of caoutchouc, A., i, 419.
photochemical studies. XIV. Fundamental photochemical laws, A., ii, 248.
- Pohl, Julius**, the fate of methyl and isopropyl alcohols in the organism, A., i, 498.
- Pohl, Robert**. See **B. Gudden**.
- Pohland, Erich**. See **Erich Krause**.
- Poirot, Gabriel**. See **Paul Fleury**.
- Poittevin, E.** See **H. V. Ellsworth**.
- Polányi, Michael**, theory of adsorption processes, A., ii, 479.
Röntgenographic determination of crystal arrangement, A., ii, 623.
- Polányi, Michael**. See also **M. Ettisch**.
- Pollack, S. M.** See **James Bryant Conant**.
- Pollak, Friedrich**. See **Alfons Klemenc**.
- Pollak, J.**, and **Anna Spitzer**, the estimation of the methyl group in methylated thiolbenzenes, A., ii, 789.
- Pollak, Leo**, influence of amino-acids and fatty acids on the regulation of the blood sugar, A., i, 483.
- Polonovski, Michel**, a new method of qualitative separation of the alkaline-earth metals, A., ii, 720.
approximate volumetric estimation of barium, A., ii, 720.
- Polonovski, Michel**, and **C. Auguste**, influence of sodium fluoride on the estimation of urea by the xanthidrol method, A., ii, 668.
- Polonovski, Michel**. See also **C. Vallée**.
- Polushkin, E. P.**, alloys of iron and uranium, A., ii, 152.
- Pommereau, Hervé de**, the reduction of ethyl benzoate and some other benzene derivatives by sodium and absolute alcohol, A., i, 338.
the reduction of some aromatic compounds by means of sodium and absolute alcohol, A., i, 825.
the reduction of ethyl α naphthylacetate and the α -naphthylethanols by sodium and absolute alcohol, A., i, 828.
- Pondal, Martiniano Leguigamón**, fluorine in Spanish grapes, A., i, 1100.
- Ponder, A. O.** See **Harold Brewer Hartley**.
- Ponder, Eric**, hæmolytic action of sodium glycocholate, A., i, 292.
the estimation of non-protein nitrogen in blood, A., ii, 583.
- Ponndorf, Wolfgang**, preparation of coumarins, A., i, 565.
- Ponsio, Giacomo**, dioximes. I. and III., A., i, 17, 1037.
cobalt compounds of α -oximino-ketones, A., i, 661.
- Ponsio, Giacomo**, and **G. Ruggeri**, dioximes. II., A., i, 627.
- Pool, E.** See **Ch. Gränacher**.
- Pop, C.** See **Al. Ionescu**.
- Pope, Frank George**. See **Sri Krishna**.
- Pope, John Clifford**. See **Francis Ernest Francis**.
- Pope, (Sir) William Jackson**, and **James Leonard Brierley Smith**, the chlorinated dialkyl sulphides, T., 1166.
- Pope, (Sir) William Jackson**. See also **John Edmund Guy Harris**, **Frederick George Mann**, and **William Hobson Mills**.
- Popov, Helene**. See **Wladimir Palladin**.
- Popper, Egon**. See **Heinrich Wieland**.
- Porcher, Ch.**, and **A. Tapernoux**, the appearance of digestive enzymes during foetal life, A., i, 1088.
- Porcher, M.**, the iodohydrin derived from allylbenzene and its transformations, A., i, 539.

- Porcher, M.** See also *Marc Tiffeneau*.
- Porges, Otto**, a case of unusual acetone-uria, A., i, 496.
- Porritt, Benjamin Dawson**. See *Bernard Dunstan Wilkinson Luff*.
- Poznjak, Eugen**, and *Herbert Eugene Merwin*, system $\text{Fe}_2\text{O}_3\text{-SO}_3\text{-H}_2\text{O}$, A., ii, 772.
- Poznjak, Eugen**, and *Ralph W. G. Wyckoff*, crystal structures of alkali haloids, A., ii, 499.
- Poznjak, Eugen**. See also *Ralph W. G. Wyckoff*.
- Potter, Ralph S.** See *George C. Bailey*.
- Poucholle, A.**, tempering, A., ii, 299.
- Pound, James Robert**, properties of mixtures of ethyl ether, sulphuric acid, and water, T., 941.
- Povarnin, G.**, and *P. Tichomirov*, molecular compounds of diketopiperazine and phenols, A., i, 1185.
- Power, Frederick Belding**, and *Victor K. Chesnut*, the odorous constituents of apples, A., i, 96.
the occurrence of methyl anthranilate in grape juice, A., i, 97.
the odorous constituents of peaches, A., i, 99.
- Powers, Edwin Booth**, physiology of the respiration of fishes in relation to the hydrogen-ion concentration of the medium, A., i, 286.
- Powers, Edwin Booth**. See also *John Addyman Gardner*.
- Prandtl, Wilhelm**, density and molecular volume of the oxides of lanthanum, praseodymium, neodymium, samarium, and europium, A., ii, 379.
- Prandtl, Wilhelm**, and *Joseph Lösch*, the separation of the rare earths by basic precipitation. III. The quantitative separation of cerium from the other earths, A., ii, 770.
- Prandtl, Wilhelm**, and *Johanna Rauchenberger*, the separation of the rare earths by basic precipitation, A., ii, 298, 769.
- Pratt, David Doug**, and *Robert Robinson*, a synthesis of pyrylium salts of anthocyanidin type, T., 1577.
- Fraunhofer, P. H.**, some electro-osmotic experiments with de Haen's membrane filters, A., ii, 114.
- Prax, J.**, olive oils and the Villavecchia reaction, A., ii, 595.
- Prell, Ernst**. See *Rudolf Pummerer*.
- Prell, G.** See *Ferdinand Henrich*.
- Preli, M.** See *F. Traetta-Mosca*.
- Price, (Miss) M. C.** See *(Miss) Mabel Harriet Norris*.
- Price, William John**. See *Arthur Robert Ling*.
- Prideaux, Edmund Brydges Rudhall**, and *H. W. Hewis*, the anodic corrosion of bismuth; bismuth compounds, A., ii, 511.
- Prigge, C.** See *Georg Schroeter*.
- Prigge, Richard**, the error in estimating chloride by Bang's micro-method, A., ii, 715.
- Pring, John Norman**, and *E. O. Ransome*, reaction between cathodic hydrogen and nitrogen at high pressures, A., ii, 639.
- Pringsheim, Hans**, and *Alexander Aronowsky*, inulin. III., A., i, 635.
- Pringsheim, Hans**, and *Diamandi Dernikos*, chemistry of starch. VI. polyamyloses, A., i, 632.
- Pringsheim, Hans**, and *Kurt Goldstein*, chemistry of starch. VII. Relationship of the α - and β -polyamyloses to the content and integument substance of the starch granule, A., i, 633.
- Pringsheim, Hans**, and *Max Lassmann*, inulin. II. Inulin and glycogen, A., i, 634.
- Pringsheim, Hans**, and *Karl Otto Müller*, the physiology of the "polyamyloses." I., A., i, 410.
- Pringsheim, Hans**, and *Walter Persch*, chemistry of starch. IV. The methylation of polyamyloses, A., i, 113.
chemistry of starch. V. Methyl and acetyl products of the "polyamyloses," A., i, 632.
- Pringsheim, Hans**, and *Karl Schmalz*, tetraäxovoglucosan and tetraglucosan, A., i, 1118.
- Pringsheim, Peter**, divergence from Stokes's law during the excitation of the fluorescence of iodine vapour, A., ii, 178.
difference between the absorption spectrum and the complete fluorescence spectrum of iodine vapour indicated by Lenz's theory, A., ii, 178.
disturbance of the power of fluorescing of fluorescent solutions by light, and the photochemical equivalent law, A., ii, 602.
- Prins, Ada**, the conditions for the maximum precipitation of an amphoteric electrolyte, A., ii, 77.
- Prins, H. J.**, acceleration of the solubility of metals in acids by reducible compounds, A., ii, 488.
the detection of nitro-compounds, A., ii, 877.
- Pritsker, J.**, and *R. Jungkuns*, hazelnut oil and the estimation of arachidic acid, A., i, 208.

- Proscopiŭ, Stéfan**, electromotive force produced by the relative displacement of an electrode and an electrolyte, A., ii, 112.
the variations of the spectrum of the mercury arc with the conditions of emission, A., ii, 600.
- Proskouriakoff, A.** See *George W. Raisis*.
- Proude, Kathleen.** See *James Frederick Spencer*.
- Prud'homme, Maurice**, some relationships between the critical temperature and the melting and boiling points, A., ii, 349.
the corresponding states; the halogenated derivatives of benzene, A., ii, 426.
- Pryde, John.** See *Alexander Kull-n Macbeth*.
- Przibram, Karl.** See *Stefan Meyer*.
- Pscheneicyn, N. K.** See *Leo Alexandrovitch Tschugaev*.
- Pucher, George W.**, uric acid. I. Examination of the variables in the Folin and Wu uric acid method. II. A modification of the Folin and Wu uric acid method, A., ii, 668.
- Pucher, George**, and *William Maurice Dehn*, solubilities in mixtures of two solvents, A., ii, 126.
- Pucher, George W.**, and *Treat Baldwin Johnson*, the utilisation of ethyl- γ -diethoxyacetate for the synthesis of derivatives of glyoxaline; an attempt to synthesise histamine by a new method, A., i, 549.
- Pummerer, Rudolf**, and *Joseph Binapfl*, the behaviour of azo-compounds and their salts with aromatic hydrocarbons and aluminium chloride, A., i, 24.
- Pummerer, Rudolf**, *Josef Binapfl*, *Karl Bittner*, and *Karl Schuegraf*, the reaction between azobenzene hydrochloride and aromatic hydrocarbons. II., A., i, 1196.
- Pummerer, Rudolf**, *Dona Melamed*, and *Hans Puttfarcken*, oxidation of phenols. VII. Dehydrogenation of *p*-cresol, A., i, 1161.
- Pummerer, Rudolf**, and *Ernst Prell*, the addition of benzene to *p*-benzoquinone, A., i, 1164.
- Purdum, R. B.** See *Graham Edgar*.
- Purdy, A. C.** See *Victor Grignard*.
- Purdy, Helen A.**, and *L. E. Walbum*, the action of various metallic salts on hemolysis, A., i, 1087.
- Purgotti, Attilio**, detection of magnesium in presence of manganese and phosphoric acid, A., ii, 85.
- Purkayostha, R. M.**, and *Niratan Dhar*, catalysis. XIII. Temperature coefficient of catalysed and non-catalysed reactions, A., ii, 362.
- Purmann, Ludwig.** See *Karl Freudenberg*.
- Pushin, Nicolai Antonovich**, and *Alexandra Fioletova*, the equilibrium in the system *m*-dinitrobenzene-urethane, T., 1822.
- Pushin, Nicolai Antonovich**, and *Alexandra Alexandrovna Glagoleva*, the equilibrium in systems composed of water and alcohols; methyl alcohol, pinacone, glycerol, and erythritol, T., 2813.
- Putochin, N. J.**, the synthesis of pyrrolidine; reduction of pyrrole by catalytic hydrogenation, A., i, 1176.
the action of formaldehyde on pyrrolidine and piperidine, A., i, 1176.
- Puttfarcken, Hans.** See *Rudolf Pummerer*.
- Puxeddu, Ernesto**, degrees of hydrolysis of alkali salts of hydroxyazo-compounds, A., i, 589.
- Puxeddu, Ernesto**, and *Marcella Gennari*, facts and theories in the constitution of the hydroxyazo-compounds, A., i, 587.
- Puxeddu, Ernesto**, and *F. L. Vodret*, velocity of reaction in the photochemical dissociation of ferric chloride, A., ii, 415.
- Puyal, José**, migration of a double linking, A., i, 617.
- Puyal, José.** See also *Ernest Fourneau*.
- Pyman, Frank Lee**, orientation of the 1:4- and 1:5-dimethylglyoxalines; mode of fission of 5-aminoglyoxalines, T., 2616.
- Pyman, Frank Lee**, and *Laurence Barnett Timmis*, some arylazoglyoxalines, A., i, 1197.
- Pyman, Frank Lee.** See also *Isidore Elkanah Balaban*, and *Louis Light*.
- Quaglia, A.** See *Carlo Sandonnini*.
- Quagliarello, G.**, chemical and physical properties of muscle and muscle extracts. VI. Muscle extract of octopus, A., i, 608.
the absorption spectra of methæmoglobin, and the alleged transformation of methæmoglobin into oxy-hæmoglobin by the action of alkali, A., i, 883.

Quagliariello, G., effect of acids and alkalis on some chemical and physical properties of hæmoglobin, A., i, 1074.

Quarder, Bilfried, optical experiment demonstrating the Bragg method of crystal analysis, A., ii, 677.

Quick Armand J., and **Roger Adams**, aliphatic arsenic acids, and aliphatic-aromatic arsenious acids, A., i, 600.

Quisumbing, Francisco A., and **Arthur W. Thomas**, conditions affecting the quantitative estimation of reducing sugars by Fehling's solution; elimination of certain errors involved in current methods, A., ii, 92.

R.

Rabe, Paul, cinchona alkaloids. XXIII. Nomenclature, and isomerism phenomena, A., i, 360.

Rabe, Paul, Karl Kindler, and **Otto Wagner**, cinchona alkaloids. XXIV. Synthesis of vinyl-free quinatoxins and quinaketones, A., i, 361.

Rabinowitch, Adolph I., anomalous dissociation in aqueous solutions, A., ii, 186.

electrolytic dissociation of salts in concentrated solutions, and in the fused and solid states, A., ii, 187.

reversal of the molecular conductivity curve and abnormal dissociation, A., ii, 188.

negative viscosity, A., ii, 478.

conductivity of the latent image, A., ii, 605.

Rabinovitch, I. M., biochemistry of methyl alcohol poisoning, A., i, 963.

Racke, Fritz. See **Richard Willstätter**.

Radde, Erich, derivatives of amino-aldehydes, A., i, 1154.

Radley, E. G. See **Arthur Francis Hallimond**.

Rath, Kurt. See **Julius von Braun**.

Ragoss, A. See **Erich Tiede**.

Rahn, Otto, significance of surface tension phenomena for the dairy practice, A., ii, 478.

Rai, Raghunath. See **Bawa Kartar Singh**.

Raiford, Lemuel Charles, the nitration of halogenated phenols, A., i, 335.

Raiford, Lemuel Charles, and **John R. Couture**, the migration of acyl from nitrogen to oxygen, A., i, 931.

Raiteri, Luigi, optical properties of certain substances important in micro-chemistry, A., ii, 541.

Raith, Emmy. See **Anton Skrabal**.

Raiziss, George W., and **A. O. Blatt**, condensation products of arspenamine [salvarsan] with aldehydes, A., i, 1079.

Raiziss, George W., and **Joseph L. Gavron**, the colloidal properties of arspenamine [salvarsan] and allied products, A., i, 1202.

Raiziss, George W., and **A. Proskouria-koff**, organic nitro-compounds containing mercury, A., i, 604.

Raiziss, George W. See also **J. F. Schamberg**.

Rakshit, Jitendra Nath, morphine, codeine, and narcotine in Indian opium, A., ii, 96.

Rakuzin, Michael A., animal hides as amphoteric and colloidal protein; theory of dyeing, tanning, disinfection, and preservation of leather and the physiological action of tanning material, A., i, 702.

a new fractionation method for proteins and their derivatives, A., i, 957.

adsorption of proteins, ferments, toxins, and sera by aluminium hydroxide, A., i, 1199.

the relation between adsorption and electrolytic dissociation, A., ii, 619.

recognition of proteins and derivatives by colour reactions, A., ii, 670.

Rakuzin, Michael A., and **Tatjana Gönke**, negative adsorption. II. Calculation of the amount of adsorbed solvent, A., ii, 821.

Ramadier, L. See **Louis Duparc**.

Raman, C. V., anisotropy of molecules, A., ii, 102.

molecular structure of amorphous solids, A., ii, 201.

molecular anisotropy in liquids, A., ii, 603.

the spectrum of neutral helium, A., ii, 803.

Ramann, Emil, and **H. Sallinger**, reactions in heterogeneous systems; the systems $K_2CO_3 | BaSO_4$, $K_2SO_4 | BaCO_3$, $K_2CO_3 | CaC_2O_4$, $K_2C_2O_4 | CaCO_3$, and $K_2CrO_4 | AgIO_3-KIO_3 | AgCrO_4$, A., ii, 131.

Ramart-Lucas, (Mme) Pauline, the molecular transposition accompanying the dehydration of $\alpha\alpha$ -diphenyl- $\beta\beta$ -dimethylpropan- α -ol, A., i, 34.

Ramart-Lucas, (Mme) Pauline, and **G. Albesco**, two $\alpha\alpha\beta\beta$ -substituted propiophenones and their products of decomposition by sodamide, A., i, 662.

Ramart-Lucas, (Mme) Pauline. See also **Albin Haller**.

- Ramm, Marie.** See *Friedrich Kehr-mann*.
- Ramstetter.** See *Max Bodenstein*.
- Randall, H. M., Walter F. Colby, and R. F. Paton,** temperature shift in near ultra-red bands, A., ii, 805.
- Randles, F. S., and Arthur Knudson,** estimation of lipid phosphoric acid ("lecithin") in blood by application of Bell and Doisy's method for phosphorus, A., ii, 719.
- Rankine, Alexander Oliver,** structure of some gaseous molecules of which hydrogen is a constituent, A., ii, 635.
the molecular structure of carbon oxysulphide and carbon disulphide, A., ii, 704.
- Rankine, Alexander Oliver, and Clarence Joseph Smith,** viscous properties and molecular dimensions of silicane, A., ii, 709.
- Ransome, E. O.** See *John Norman Fring*.
- Raper, Henry Stanley.** See *Herbert Davenport Kay*.
- Rapport, David.** See *H. V. Atkinson*.
- Rast, Karl,** a micro-method for the determination of molecular weight in a melting-point apparatus, A., ii, 421.
- Rastelli, G.,** derivatives of isoeugenol, A., i, 1010.
- Rastelli, G.** See also *Riccardo Ciusa*.
- Rathert, H.** See *Arthur Köts*.
- Rathery, F.** See *Alexandre Desgrez*.
- Rathsam, G.** See *Hermann Staudinger*.
- Rathsburg, Hans,** mercury fulminate, A., i, 116.
reductions with titanium trichloride, A., i, 242.
- Rau, Madyar Gopal, and John Lionel Simonsen,** oils and fats from the seeds of Indian forest trees, A., i, 1100.
- Rau, Madyar Gopal.** See also *John Lionel Simonsen*.
- Rauch, Hans,** estimation of the carbon dioxide content of air, with special reference to the "aeronom," A., ii, 316.
- Rauchenberger, Johanna.** See *Wilhelm Prandtl*.
- Ravenna, Ciro,** constitution of the dipeptides of aspartic acid, A., i, 180.
- Ravenna, Ciro.** See also *Giacomo Luigi Ciamician*.
- Rawling, Sidney Owen,** electric heating and controlling apparatus for a small thermostat, A., ii, 612.
- Rawling, Sidney Owen, and Walter Clark,** the isoelectric condition of gelatin, T., 2880.
- Rây, (Sir) Prafulla Chandra,** triethylene tri- and tetra-sulphides. II., T., 1279.
- Rây, (Sir) Prafulla Chandra, and Radhakishen Das,** chloropicrin as a reagent for the diagnosis of mercaptans and potential mercaptans, T. 328.
- Rây, Priyadarajan, and Pulin Vihari Sarkar,** formation and dissociation of some polyhalogen compounds of hydrogen in aqueous solution, T., 1449.
- Ray, Rames Chandra,** the potassium salt of hexahydrodioxidyboron, T., 1088.
heat of crystallisation of quartz, A., ii, 685.
- Raymond, E.,** the fatty acids of colza oil, A., i, 798.
- Rayner, Archibald,** composition of the residue on distillation of crude glycerol, A., i, 711.
- Read, Hugh Norman.** See *Gilbert Thomas Morgan*.
- Read, John, and Eric Hurst,** the conversion of allyl alcohol to glyceryl chloro- and bromo-hydrins, T., 989.
the action of bromine water on indene, T., 2550.
- Read, John, and Henry George Smith,** piperitone. II. Benzylidene-*dl*-piperitone, T., 574.
piperitone. IV. The interaction of *dl*-piperitone and semicarbazide, and the isolation of pure *dl*-piperitone, T., 1863.
- Read, John, Henry George Smith, and (Miss) Marie Bentivoglio,** piperitone. III. The oximes of *dl*-piperitone, T., 582.
- Read, J. W.,** practical significance of the organic carbon; nitrogen ratio in soils, A., i, 416.
- Read, J. W., and R. H. Ridgell,** use of the conventional carbon factor in estimating soil organic matter, A., ii, 540.
- Read, T. A.** See *David Avery*.
- Read, William T.,** hydantoin; synthesis of the soporific 4-phenyl-4-ethylhydantoin [nirvanol], A., i, 954.
- Reaebourg, G.** See *C. Pépin*.
- Rebello-Alves, S.** See *Alberico Benedicenti*.
- Rebmann, Adolf.** See *Emil Baur*.
- Rechenberg, Carl von,** intersecting vapour pressure curves and deductions therefrom, A., ii, 120.
vaporisation of solutions of liquid pairs possessing intersecting vapour pressure curves, A., ii, 120.

- Rechenberg, Carl von**, and *W. von Rechenberg*, formation of additive products between cresols on the one hand and ethyl ether, ethyl alcohol, acetone, benzene, etc., on the other, A., i, 932.
- Rechenberg, W. von.** See *Carl von Rechenberg*.
- Reclaire, A.**, analysis of acetic anhydride, A., ii, 532.
- Reclaire, A.** See also *Anne Willem Karel de Jong*.
- Recoura, A.**, new properties of the green chromium sulphate, A., ii, 508.
- Reddellien, Gustav**, the mutual replacement of amine residues by anils, A., i, 145.
- Reddellien, Gustav**, and *Hildegard Danilof* fission of anils, A., i, 147.
- Redfern, Gladys M.**, absorption of ions by the roots of living plants. I. Absorption of the ions of calcium chloride by pea and maize, A., i, 614.
- Redwitz, E. von.** See *Eduard Grafe*.
- Reeder, J. C.** See *J. S. Jones*.
- Reedy, J. H.**, silver bromate, A., ii, 56. precipitation of arsenic sulphide from arsenates, A., ii, 225.
- Reeves, Harry Gordon.** See *Gilbert Thomas Morgan*.
- Regenbogen, (Mlle) A.** See *Nicolaas Schoorl*.
- Réglade, Antoine.** See *Paul Nicolardot*.
- Reglin, Werner.** See *Arthur Rosenheim*.
- Reich, René.** See *André Job*.
- Reichel, Heinrich**, water and ionic distribution in the organism A., i, 485.
- Reichenstein, David**, and *F. Klement*, an electrolytic current intensification effect, a new electrolytic displacement effect, and the connexion between electrolysis and the emission of electrons in a vacuum. II., A., ii, 112.
- Reichenstein, David**, and *M. Rothschild*, an electrolytic current intensification effect, a new electrolytic displacement effect, and the connexion between electrolysis and the emission of electrons in a vacuum. III., A., ii, 736.
- Reid, Ebenezer Emmet, Colin M. Mackall**, and *George E. Miller*, derivatives of anthraquinone, aliphatic thioethers, dithioethers, and thioether sulphonic acids, A., i, 154.
- Reid, Ebenezer Emmet.** See also *Neil E. Gordon, C. H. Milligan, and Raphael Rosen*.
- Reid, Robert Douglas.** See *Malcolm Percival Applebey*.
- Reif, G.** See *Balthasar Pfyl*.
- Reihlen, Hans**, the auxiliary valency of the hydroxyl group. I., A., i, 1146.
- Reihlen, Hans**, and *Adolf Sapper*, the auxiliary valency of the hydroxyl group; the complex salts of substituted pyrocatechol, A., i, 1147.
- Relly, (Miss) Amy Ada Beatrice.** See *John Norman Collie*.
- Reindel, Fritz.** See *Heinrich Wieland*.
- Reinders, Willem**, and *P. van Groningen*, equilibria in the system Fe-C-O; the equilibrium Fe₃-martensite-ferrous oxide-gas, A., ii, 153.
- Reinfurth, Elsa.** See *Carl Neuberg*.
- Reinhard, H. F.** See *M. G. Mellon*.
- Reinicke, Dora.** See *Peter Bona*.
- Reinger, Erich**, sativic acid, A., i, 623.
- Reis, A.**, and *L. Zimmermann*, hardness of solid substances and its relationship to chemical constitution, A., ii, 745.
- Reis, H.** See *Ch. Gränacher*.
- Reis, V. van der.** See *Rudolf Cobet*.
- Reissau, G. G.** See *Erich Krause*.
- Reissert, Arnold**, a new class of vat dyes containing sulphur and nitrogen, A., i, 583.
- Reist, Svend Hubert**, estimation of sugar in small quantities of blood by the Pavy-Sahli method, A., ii, 323.
- Reitstötter, Josef.** See *Arne Westgren*.
- Remy, Heinrich**, the alkali-ruthenium double sulphites, A., ii, 857.
- Renand, E.** See *A. Astruc*.
- Rendall, Arthur Geoffrey**, and (*Miss*) *Martha Annie Whiteley*, the oxime of mesoxamide (isonitrosomalonalonamide) and some allied compounds. IV. The ethers of isonitrosomalonalonilide, isonitrosomalondimethylamide, and isonitrosomalondibenzylamide, T., 2110.
- Rengade, Etienne**, the redissolution of a salt during the isothermal evaporation of a solution, A., ii, 31.
- Renn, Konrad.** See *Stefan Goldschmidt*.
- Renoux, M.**, identification of traces of true albumin in urine; separation of ψ -albumins, A., ii, 797.
- Renshaw, Arnold.** See *Thomas H. Fairbrother*.
- Renshaw, Roemer Rex**, and *Nellie M. Naylor*, dyes containing the furan ring, A., i, 566.
- Renz, Carl**, light reactions of the oxides of titanium, cerium, and the earth acids, A., ii, 61.
- photochemistry of thalious chloride.** II., A., ii, 66.
- the rare earths in the periodic system,** A., ii, 758.
- Report of the Council, T., 722.**
- Retze, Ewald.** See *Karl Fleischer*.

- Reuss, A.**, estimation of nitrates in drinking water by Mayrhofer's method, A., ii, 454.
- Reverdin, Frederic**, and **H.-P. André Boethlisberger**, nitro-derivatives of *p*-phenetidine, A., i, 537.
- Reychler, Albert**, atomic weights and frequencies, A., ii, 279.
- Rhind, Donald**, and **Francis Edward Smith**, tannase, A., ii, 407.
- Rhode, Heinrich**, solubility, capillary activity, and hemolytic activity of terpene derivatives, A., i, 964.
hemolysis by morphine and homologues, A., i, 1088.
- Rhodes, F. H.**, and **A. L. Markley**, freezing-point diagram of the system phenol-water, A., i, 135.
- Rhyn, A. J. van**. See **Erich Ebler**.
- Ribaud, G.**, spectrum of bromine vapour; Zeeman effect, A., ii, 3.
- Ribot, A.** See **Ch. Achard**.
- Rice, Francis E.**, and **Toratoro Hanzawa**, estimation of peroxydase in milk, A., ii, 407.
- Rich, M. N.** See **John W. Marden**.
- Richards, Theodore William**, magnitude of atoms, A., ii, 42.
- Richards, Theodore William, Edward P. Bartlett**, and **James H. Hodges**, compressibility of benzene, liquid and solid, A., ii, 28.
- Richards, Theodore William**, and **James Bryant Conant**, electrochemical behaviour of liquid sodium amalgams, A., ii, 340.
- Richards, Theodore William**, and **Theodore Dunham, jun.**, effect of changing the hydrogen-ion concentration on the potential of the zinc electrode, A., ii, 418.
- Richards, Theodore William**, and **William Buell Meldrum**, existence of tetra-hydrated sodium sulphate in mixed crystals with sodium chromate, A., ii, 54.
- Richards, Theodore William**, and **Allan W. Rowe**, heats of neutralisation of potassium, sodium, and lithium hydroxides with hydrochloric, hydrobromic, hydriodic, and nitric acids at various dilutions, A., ii, 425.
- Richards, Theodore William**, and **Charles P. Smyth**, solid thallium amalgams and the electrode potential of pure thallium, A., ii, 341.
- Richards, Theodore William**, and **Setsuro Tamaru**, heat of solution of cadmium in hydrochloric acid, A., ii, 475.
- Richards, Theodore William**, and **Thorbergur Thorvaldson**, heat of solution of zinc in hydrochloric acid, A., ii, 475.
- Richardson, Owen Willans**, and **Charles Blizard Bassoni**, excitation of soft characteristic X-rays, A., ii, 14.
- Richard, A.**, the comparative blood-pressure raising power of racemic and lævo-adrenaline, A., i, 891.
the action of the digestive juices on β -benzyl-d-glucoside, A., i, 1094.
the toxicity of β -benzylglucoside obtained by biochemical synthesis, A., i, 1094.
- Richmond, Henry Droop**, and **E. H. England**, the sulphuric acid reaction for liver oil, A., ii, 792.
- Richter, Friedrich**. See **Erich Tiede**.
- Richter, Rudolf**. See **Alots Zinke**.
- Richter-Quittner, M.**, the importance of imbibition for some biochemical problems, A., ii, 204.
chemical blood analysis. III. The importance of the ultra-filtration method for the analysis of blood, A., ii, 240.
blood analysis. IV. Ashing methods, A., ii, 407.
- Richter-Quittner, M.**, and **H. Hoenlinger**, estimation of non-protein nitrogen by means of ultra-filtration, A., ii, 717.
- Richter-Quittner, M.** See also **Wilhelm Falta**.
- Richtmyer, F. K.**, the laws of absorption of X-rays, A., ii, 105.
mass-absorption coefficients as a function of wave-length above and below the *K* X-ray limit of the absorber, A., ii, 804.
the evidence regarding the so-called "J" radiation in the characteristic X-ray spectra of the elements, A., ii, 804.
- Rideal, Eric Keightley**, the hydrogenation of ethylene in contact with nickel, T., 309.
- Rideal, Eric Keightley**, and **William Thomas**, the autoracemisation of potassium chromioxalate, T., 196.
adsorption and catalysis in Fuller's earth, T., 2119.
- Rideal, Eric Keightley**. See also **Charles George Lewis Wolf**.
- Ridgell, R. H.** See **J. W. Read**.
- Ridgway, Leslie Randal**. See **Frederick Challenger**.
- Rie, Gertrud**. See **Ernst Philipp**.
- Riebensahm, E.** See **Georg Schroeter**.
- Riedel, J. D. Akt.-Ges.**, preparation of derivatives of cholic acid, A., i, 553.
preparation of bile acids, A., i, 554.
preparation of ethers of *p*-hydroxyphenylcarbamide, A., i, 579.

- Riedel, J. D., Akt.-Ges.,** preparation of hexamethylenetetramine derivatives, A., i, 991.
 preparation of additive products of hexamethylenetetramine with esters of monohalogen fatty acids, A., i, 991.
 preparation of an unsaturated bile acid, A., i, 1160.
- Riemer, Heinrich.** See **Robert Krennman.**
- Riesenfeld, Ernst Hermann, and G. W. Feld,** polythionic acids and polythionates, A., ii, 45.
- Riesenfeld, Ernst Hermann, and R. Klement,** the constitution of Erdmann's salt and its derivatives, A., ii, 853.
- Riesenfeld, Ernst Hermann, and G. M. Schwab,** ozone, A., ii, 637.
 physical constants of ozone, A., ii, 761.
- Riesser, Otto,** the physiology of creatine, A., i, 791.
- Riesser, Otto, and S. M. Neuschloz,** mechanism of the contraction of striated muscle produced by poisons. V. The action of specific muscle poisons on lifeless colloids, A., i, 1212.
- Riffart, Hans,** the triketohydrindene (ninhydrin) reaction as a quantitative colorimetric method for the estimation of amino-acid nitrogen, A., ii, 718.
- Riffert, Elli.** See **Karl Schaum.**
- Riggert, K.** See **M. Volmer.**
- Riiber, Claus Nissen,** mutarotation. I., A., ii, 807.
- Riley, George Clifford.** See **Edward Hope.**
- Billiet, Auguste,** 6-aminoveratraldehyde and its derivatives, A., i, 839.
- Rimbach, Eberhard, and P. Ley,** acidifying action of hydroxy-organic compounds on boric acid and molybdic acid, A., ii, 342.
- Rindl, M.,** mafurra tallow, a product of the nuts of *Trichilia emetica*, A., i, 1102.
- Ringer, Wilhelm Eduard,** the influence of reaction on the action of trypsin. I., A., i, 282.
- Rinne, Friedrich,** space formulae, A., ii, 868.
- Rion, Paul,** the velocity of absorption of carbon dioxide by alkaline solutions, A., ii, 433, 487.
 the rate of absorption of carbon dioxide by ammoniacal solutions, A., ii, 742.
- Ripert, Jean,** the biology of the alkaloids of belladonna, A., i, 96.
- Ripper, Kurt.** See **W. J. Minaceff.**
- Risseghem, (Mlle) Hortense van,** synthesis of trimethylethylmethane [$\beta\beta$ -dimethylbutane], A., i, 813.
 γ -methyl- Δ^2 -pentene, A., i, 909.
- Risseghem, (Mlle) Hortense van.** See also **Georges Chavanne.**
- Rivalland, Ch.** See **Philippe Malvesin.**
- Rivett, Albert Cherbury David,** the quaternary system, ammonium chloride—sodium sulphate—ammonium sulphate—sodium chloride—water, T., 379.
- Rivière.** See **Clément.**
- Roaf, Herbert Eldon,** urochrome as a derivative of chlorophyll, A., i, 401.
- Robert, E.** See **Richard Stoermer.**
- Roberts, Lathrop E.** See **William Draper Harkins.**
- Roberts, Oswald Digby, and Harold Thomas Islip,** the constants of Indian beeswax, A., ii, 534.
- Robertson, John K.,** the electrodeless discharge in certain vapours, A., ii, 609.
- Robertson, John McGregor.** See **George Gerald Henderson.**
- Robin, A., and A. Sournigault,** cancerous anemia, A., i, 497.
- Robin, Paul,** action of nitrogen iodide and of cyanogen iodide on benzamidine, A., i, 36.
- Robinson, Charles S.,** hydrogen-ion concentration of human faeces, A., i, 899.
 the composition and preparation of a neutral solution of ammonium citrate, A., ii, 84.
- Robinson, Charles S., and Selma L. Bandemar,** the analysis of solutions of ammonium citrate, A., ii, 460.
- Robinson, Charles S., O. B. Winter, and E. J. Miller,** availability of organic nitrogenous compounds, A., i, 212.
- Robinson, Guy C.,** the acetone and butyl alcohol fermentation of various carbohydrates, A., i, 971.
- Robinson, Gilbert Wooding,** mechanical analysis of humus soils, A., ii, 888.
 a new method for the mechanical analysis of soils and other suspensions, A., ii, 888.
- Robinson, Lilly.** See **Ernst Philipp.**
- Robinson, Robert,** the atomic vibrations in the molecules of benzenoid substances, A., i, 593.
- Robinson, Robert.** See also **James Wilson Armit, Herbert Grace Crabtree, Herbert Greene, William Ogilvy Kermack, Alexander Killen Macbeth, Hidejiro Nishikawa, David Diyo Pratt, and James Scott.**

- Robinson, R. H.**, and **D. E. Bullis**, acid soils. III. The influence of calcium carbonate, calcium oxide, and calcium sulphate on the soluble soil nutrients of acid soils, A., i, 976.
- Robinson, William O.**, absorption of water by soil colloids, A., i, 1228.
- Robison, Robert**, the value of gelatin in relation to the nitrogen requirements of man, A., i, 488.
distribution of the nitrogenous constituents of the urine on low nitrogen diets, A., i, 495.
the estimation of total sulphur in urine, A., ii, 389.
- Robles, B. de Vries**, the blood calcium content in normal children and in tetany. I., A., i, 192.
- Rockefeller Institute for Medical Research**, preparation of aromatic aminoarsinic acids, A., i, 961.
preparation of aromatic arseno-compounds, A., i, 962.
- Rodier, E.** See **Jules Aloy**.
- Rodillon, Georges**, a source of error in testing urine for dextrose with *o*-nitrophenylpropionic acid, A., ii, 166.
detection of urobilin in urine, A., ii, 888.
- Rodriguez Mourello, José**, thermotropy, A., ii, 604.
- Röckemann, W.**, [state of tetrahydronaphthalene in the organism], A., i, 499.
- Röder, Hans.** See **Ernst Späth**.
- Röhm and Haas**, preparation of ethylene cyanohydrin [β -hydroxypropionitrile], A., i, 531.
- Röhm, Otto**, a new iron salt, A., ii, 648.
- Röhm, R.** See **H. Kessler**.
- Rørdam, H. N. K.**, influence of substitution in the ortho-, meta-, and para-positions on the absolute affinity of benzoic acid. I., A., i, 338.
- Rösler, Wilhelm.** See **Robert Kremann**.
- Roethlisberger, H.-P. André.** See **Frederic Reverdin**.
- Rogers, E. C.** See **John Bernard Ekeley**.
- Rogers, J.** See **E. M. Black**.
- Rogers, T. H.**, **Charles Snowden Piggot**, **W. H. Bahlke**, and **J. M. Jennings**, catalytic oxidation of carbon monoxide, A., ii, 136.
- Rogozinski, F.**, biochemistry of phosphorus, A., i, 1226.
- Rojahn, C. A.**, poly-ethers of trimethylene glycol, A., i, 103.
preparation of β -chloro- and β -bromopropionic acids from trimethylene glycol, A., i, 105.
- Rojahn, C. A.**, some ketones of the pyrazole series, A., i, 373.
3-chloro-1-alkylpyrazoles and 1-alkylpyrazol-3-one, A., i, 1183.
- Rojahn, C. A.** See also **Carl Mannich**.
- Rojdestwensky, A.**, properties and preparation of Dutch East Indies sandal wood oil, A., ii, 792.
- Rolf, Ida P.** See **Phæbus A. Levene**.
- Rolla, Luigi** [with **Mario Frassinetti**, and **Mario Bulli**], corrosion of certain aluminium alloys, A., ii, 379.
- Rolla, Luigi** and **L. Mazza**, new differential method for the measurement of the conductivity of electrolytes, A., ii, 544.
- Rolt, William Joseph Woodgate.** See **Oscar Lisle Brady**.
- Roman, F. L.** See **Roger Adams**.
- Romani, E.**, derivatives of benzthiazole, A., i, 466.
- Romani, E.** See also **Giuseppe Bruni**.
- Rombaut, Lawrence E.**, and **Julius A. Nieuwland**, catalytic synthesis of hexamethylenetetramine, A., i, 990.
- Romburgh, G. van**, nitro-derivatives of alkylated benzidines, A., i, 275.
- Romieu, Marc**, and **Fernand Obaton**, comparative spectroscopic study of the green pigment of the *Chetoptera* and of the chlorophyll of the *Ulva*, A., i, 793.
- Romijn, G.**, aluminium for the arsenic reaction, A., ii, 455.
- Rona, Elisabeth**, the ionium content of radium residues, A., i, 250.
- Rona, Peter, Yrjö Airila**, and **A. Lasnitski**, toxicity; the combined action of quinine and of narcotics on invertase and on the action of arsenic compounds on maltase, and on α -methylglucosidase, A., i, 959.
- Rona, Peter**, and **Emerck Bach**, toxicity. III. The action of *m*- and *p*-nitrophenols on invertase, A., i, 65.
- Rona, Peter**, and **Ernst Bloch**, toxicity. I. The action of quinine on invertase, A., i, 65.
the fixation of quinine by red blood-corpuses and the distribution of quinine in blood, A., i, 290.
fixation of quinine by corpuscles and its action on cell respiration, A., i, 484.
- Rona, Peter**, and **R. Pavlovic**, action of quinine and atoxyl on liver lipase, A., i, 887.
- Rona, Peter**, and **Dora Reinicke**, toxicity. II. The action of quinine on serum-lipase, A., i, 67.

- Rondoni, Pietro**, origin of melanin from pyrrole. II. Action of organ extracts on pyrrole; the sepia of the cuttlefish, A., i, 64.
- Rondou, Alb.**, the reaction of organo-magnesium compounds on nitriles; action of magnesium methyl bromide on phenylacetonitrile, A., i, 934.
- Rooke, Horace Samuel**. See *Gilbert Thomas Morgan*.
- Root, Lucie E.** See *Howard B. Lewis*.
- Roschier, R. H.** See *Gustav Komppa*.
- Rose, William Cumming**, the influence of food ingestion on endogenous purine metabolism. I. and II., A., i, 194.
- Rosebourne, Cyril**. See *Alexander Findlay*.
- Rosedale, John Lewis**, the amino-acids of flesh; the diamino-acid content of rabbit, chicken, ox, horse, sheep, and pig muscle, A., i, 492.
- Rosedale, John Lewis**. See also *Robert Henry Aders Plimmer*.
- Rosemann, R.**, the metabolism of calcium, A., i, 1210.
- Rosen, Raphael**, and *Ebenezer Emmet Reid*, sesqui-mustard gas or bis- β -chloroethyl ether of ethylene dithioglycol, A., i, 420.
- Rosenberg, G. von**. See *Friedrich August Henglein*.
- Rosenberg, H.**, fatigue of alkali metal cells in the neighbourhood of the discharge potential and the influence of this on the results of photometric measurements, A., ii, 189.
- Rosenberg, Max**, blood sugar. I. A critical survey of the methods of estimation of blood sugar and of the "threshold" concept, A., i, 482.
blood sugar. II. Alimentary hyperglycæmia under normal and pathological conditions, A., i, 789.
- Rosenberg, J. O.** See *Paul Karrer*.
- Rosenblatt, (Mme) M.** See *Gabriel Bertrand*.
- Rosenheim, Arthur**, and *Leonhard Krause*, selenious acid and heteropolyselenites, A., ii, 47.
- Rosenheim, Arthur**, and *Felix Leyser*, iso- and heteropoly-acids. XVII. Polyborates in aqueous solution, A., ii, 50.
- Rosenheim, Arthur**, and *Werner Reglin*, highly basic lithium salts of weak inorganic acids, A., ii, 289.
- Rosenheim, Arthur**, and *Gert Trewendt*, complex uranyl hypophosphites, A., i, 650.
- Rosenmann, Max**, fibrinolysis. II., A., 596.
- Rosenmund, Karl W.**, certain derivatives of arylated cinchonic acids, A., i, 51.
reduction of acid chlorides to aldehydes by means of nickel catalysts, A., i, 939.
- Rosenmund, Karl W.**, and *E. Pfannkuch*, gallaldehyde and its derivatives, A., i, 1030.
- Rosenmund, Karl W.**, and *Fritz Zetsche*, the influencing of catalysts and specifically active catalysts, A., ii, 41, 834.
- Rosenmund, Karl W.**, *Fritz Zetsche*, and *Fr. Enderlin*, preparation of aldehydes from acid chlorides. V. (Dialdehydes. II.) Synthesis of decanedial, A., i, 431.
- Rosenmund, Karl W.**, *Fritz Zetsche*, and *Chr. Flüttsch*, the preparation of aldehydes from acid chlorides. IV. Dialdehydes. I., A., i, 39.
- Rosenthal, Felix**, and *Klothilde Meier*, the types of reaction of the bile pigments and the quantitative relation of bilirubin to cholesterol in the blood during different forms of jaundice, A., i, 198.
- Rosenthaler, Leopold**, the hydrocyanic acid question. VII. *Cornus sanguinea*, A., i, 412.
asymmetric syntheses by means of enzyme action. IV., A., i, 480.
 σ -emulsin (oxynitrilase), δ -emulsin (oxynitrilase), and carboligase, A., i, 600.
the hydrocyanic acid question. VIII. Plants containing hydrocyanic acid and saponin. IX. Hydrocyanic acid content of cherry-laurel leaves infected by fungi. X. Influence of lesions on the hydrocyanic acid content of cherry-laurel leaves, A., i, 614.
iodic acid as a microchemical reagent for the detection of organic bases, A., ii, 327.
estimation of arsenic acid, A., ii, 584.
detection and estimation of oxalic acid and its use in standardising iodine and silver solutions, A., ii, 594.
- Roser, P.** See *Alfred Heiduschka*.
- Ross, Albert**. See *Wilhelm Schneider*.
- Ross, Harry P.** See *Clarence E. May*.
- Ross, J. H.** See *Amé Pictet*.
- Rossi, G.**, colloidal trimercuracetanilide acetate, A., i, 605.
physico-chemical investigation of colloidal sulphur, A., ii, 485.
- Rossi, G.** See also *Alessandro Bernardi*.
- Rost, E.**, zinc in the human and animal organism, A., i, 87.

- Rothenberger, Emil** See *Franz Fichter*.
- Rother, Julius**, the problem of nucleic metabolism. II. The influence of human faeces on yeast-nucleic acid, A., i, 292.
- Rothschild, M.** See *David Reichinstein*.
- Roure-Bertrand Fils**, essential oils, A., i, 845, 846, 847.
two Indo-Chinese oils, A., i, 846.
- Rousaud and Thiéry**, relation between the viscosity of blood and the ratio of uric acid in serum to that in whole blood, A., i, 394.
relation between viscosity and the ratio of cholesterol in serum to that in whole blood, A., i, 394.
- Rowe, Allan W.** See *Theodore William Richards*.
- Rowe, Frederick Maurice, and John Stanley Herbert Davies**, studies in the dihydronaphthalene series. III. The oxidation and bromination of 5:8-dihydro- α naphthylamine, T., 1000.
- Roxas, Carla Ruiz de.** See *Franz Faltis*.
- Roy, Gustave.** See *Friedrich Kehrman*.
- Royer, J.**, estimation of the chlorometric degree of bleaching chlorides, A., ii, 581.
- Royer, L.**, the inversion of the rotatory power in anisotropic liquids, A., ii, 415.
- Royer, L.** See also *Georges Friedel*.
- Rubricius, H.**, estimation of nickel in steels, A., ii, 163.
- Rudnick, Paul**, preparation of chloroplatinic acid by means of hydrogen peroxide, A., ii, 303.
- Rudolph, W.** See *André Helbronner*.
- Rüdy, R.** See *Charles Eugène Guye*.
- Rueha, Amin.** See *W. Biedermann*.
- Ruff, Otto**, reactions on metallic surfaces, A., ii, 363.
- Ruff, Otto, and Hellmuth Hartmann**, the absorption of nitrogen by calcium and its alloys, A., ii, 377.
- Ruff, Otto, Gerhart Schmidt, and Susanne Mugdan**, high temperature investigations. XV. The vapour pressure of the alkali fluorides, A., ii, 818.
- Ruff, Otto, and Fritz Thomas**, the reduction of tantalum pentachloride, A., ii, 513.
- Ruggeri, G.** See *Giacomo Ponsio*.
- Ruggli, Paul, and R. Ernest Meyer** [with *P. Hubert*], stilbene-2:2'-dicarboxylic acid, A., i, 343.
- Ruhemann, Siegfried**, lignite producer tar, A., i, 22.
- Runge, C.**, a new band spectrum of oxygen, A., ii, 329.
- Runnström, J.**, the action of some electrolytes and non-electrolytes on the velocity of sedimentation of the red blood-corpuscles of the horse, A., i, 289.
- Rupe, Hans**, trimethylcamphorylmethylammonium bromide, A., i, 666.
the influence of constitution on the rotatory power of optically active substances. XV. A new constant for calculating the curve of rotation-dispersion, A., ii, 602.
- Rupe, Hans, and P. Briellmann**, isocampholic acid, A., i, 1017.
- Rupe, Hans, and A. Jäggi**, the influence of constitution on the rotatory power of optically active substances. XIV. Ketonic derivatives of 1:2:2:3-tetramethylcyclopentane and 1:2:2-trimethylcyclopentane-3-carboxylic acid, A., i, 840.
- Rupe, Hans, and Hans Müller**, the products of the reduction of hydroxymethyleneacetophenone [phenyl β -hydroxyvinyl ketone] and of α -hydroxymethylene- α -benzylacetone [β -phenyl- α -hydroxy-methylene-ethyl methyl ketone], A., i, 40.
- Rupe, Hans, and H. Schmid**, reduction products of hydroxymethylenecamphor. V. Coupling of hydroxylamine with methylenecamphor, A., i, 1041.
a case of total anomalous rotation-dispersion, A., ii, 603.
- Rupe, Hans, and R. Wittwer**, condensation products of phenylhydroxylamine with hydroxymethylene compounds and carbinols. II. Hydroxymethylenedecybenzoin and phenylhydroxylamine, A., i, 448.
condensation products of phenylhydroxylamine with hydroxymethylene compounds and carbinols. III. Diphenylbromomethane and phenylhydroxylamine, A., 449.
- Ruppel, Wilhelm.** See *Emil Heuser*.
- Russell, Henry Norris**, ionisation in the solar atmosphere, A., ii, 675.
- Russell-Wells, Barbara**, carrageen (*Chondrus crispus*). III. The constitution of the cell wall, A., i, 1223.
- Rusznayák, Stejan, and Géza Hetényi**, physico-chemical investigations on body-fluids. IV. The state of sugar in serum, A., i, 291.
- Rutherford, (Sir) Ernest**, artificial disintegration of the elements, T., 400.
- Rutherford, (Sir) Ernest, and James Chadwick**, the disintegration of elements by α -particles, A., ii, 682.

- Rusicka, Leopold, and Jules Meyer**, higher terpene compounds. II. Abietic acid, A., i, 547.
higher terpene compounds. V. Conversion of abietic acid into methylretene, A., i, 829.
- Rusicka, Leopold, Jules Meyer, and M. Mingazzini**, higher terpene compounds. III. The naphthalene hydrocarbons cadalene and eudalene; two aromatic fundamental compounds of the sesquiterpene series, A., i, 560.
- Rusicka, Leopold, and M. Mingazzini**, higher terpene compounds. VI. The two methylisopropyl naphthalenes from cadalene, A., i, 1001.
- Rusicka, Leopold, and C. F. Seidel**, higher terpene compounds. IV. Synthesis of cadalene, A., i, 562.
the 4-piperidone ring, A., i, 1057.
- Ryschkewitsch, Eugen**, behaviour of carbon at high temperatures, A., ii, 443.
- S.
- Saar, R.**, pycnometry, A., ii, 549.
- Sabalitschka, Th.**, aniline glucoside (glucose anilide), A., i, 247.
- Sabalitschka, Th., and H. Schmidt**, detection of antimony in analysis, A., ii, 531.
- Sachs, Georg, and Ludwig Leopold**, chlorination of *p*-iododimethylaniline, A., i, 821.
- Sachs, W.** See *Paul Jacobsen*.
- Sage, C. Edward**, analysis of acetic anhydride, A., ii, 593.
- Saha, Haridas, and Kumud Nath Choudhury**, capsularin, a glucoside from jute leaf, T., 1044.
- Saha, Megh Nad**, atomic radius and ionisation potential, A., ii, 278.
- Saillard, E.**, composition of wild beet-roots, A., i, 415.
- St. John, Charles E., and Harold D. Babcock**, wave-length of lines in the iron arc from grating and interferometer measurements, λ 3870-8750, A., ii, 4.
- St. John, E. Q.** See *Joseph Samuel Hepburn*.
- Sakaguchi, Kozo, Osamu Asakawa, and Toshitane Matsuyama**, determination of the carbohydrate assimilative power of man, A., i, 699.
- Sakao, Tokutaro, and Mitsuie Hirose**, colour of fluorites, A ii, 779.
- Sakellarios, Euklid**, the action of bromine on nitrophenol-sulphonic and -sulphocarboxylic acids, A., i, 1144.
- Sala, C.** See *G. Charrier*.
- Salant, William, and Nathaniel Kleitman**, the [physiological] action of mercury, A., i, 794.
pharmacological studies on acetone, A., i, 794.
toxicity of scatole, A., i, 794.
- Salisbury, E.J.**, soil reaction and succession in relation to plant covering, A., i, 1104.
- Salisbury, Henry M.** See *Earle T. Oakes*.
- Salkind.** See *Zalkind*.
- Salkowski, Ernst [Leopold]**, yeast gum and saccharase, A., i, 304.
xylan, A., i, 323.
the chemical nature of toxins and antitoxins, A., i, 1216.
estimation of oxalic acid in urine, A., ii, 92.
the estimation of the purine bases in urine, A., ii, 405.
- Salles, Edouard.** See *A. Zimmern*.
- Sallinger, H.** See *Emil Ramann*.
- Sallmann, Richard.** See *Hans Eduard Fierz*.
- Salmon, Cyril Sebastian**, the effect of electrolytes on the constitution of soap solutions, as deduced from electromotive force, T., 711.
- Salomon, Harry R.** See *Paul Karrer*.
- Salomon, T.** See *Henri Gault*.
- Salway, Arthur Henry, and Percy Noel Williams**, the catalytic oxidation of saturated paraffin hydrocarbons and fatty acids, T., 1343.
- Salway, Arthur Henry.** See also *L. V. Cocks*.
- Samdahl, B.** See (*Mlle*) *Ellen Gleditsch*.
- Samec, Maximilian, and S. Ferjantšič**, plant colloids. XII. Action of formaldehyde on cellulose, A., i, 115.
- Samec, Maximilian, and (Mlle) Anku Mayer**, plant colloids. XIII. Synthetic amylo-phosphoric acids, A., i, 921.
- Samec, Maximilian, and V. Šeajević**, the composition of agar, A., i, 231.
- Sammartino, Ubaldo**, the chemistry of the lungs. I., A., i, 296.
the importance of the medium in the study of catalase, A., i, 391.
probable occurrence of proteinogenic amines in the thyroid, A., i, 966.
chemistry of the lungs; a new phosphosulphatide, A., i, 1089.
a new constituent of the thyroid, A., i, 1212.
- Sammartino, Ubaldo.** See also *Herbert Elias*.
- Samson, G.**, changes in the blood after oral administration of sodium chloride, A., i, 82.

- Samuelson, Sigurd.** See *Emil Henser.*
- Samuelsson, E.,** α -1-naphthylethylamine, A., i, 828.
- Sanctis, G. de, and Q. Fiori,** presence of aromatic hydroxy-acids in urine, A., i, 609.
- Sandberg, Marta,** the course of alcoholic fermentation in the presence of urea, A., i, 502.
- Sandberg, Marta.** See also *Carl Neuberg.*
- Sander, Frank V.** See *John W. E. Glattfield.*
- Sando, Charles E., and H. H. Bartlett,** the organic acids of *Pyrus coronaria*, L., *Rhus glabra*, L., and *Acer saccharum*, Marsh, A., i, 100.
- Sandonnini, Carlo,** behaviour of certain metals as catalysts. I., A., ii, 557.
- Sandonnini, Carlo, and A. Quaglia,** combination in detonating gas in presence of colloidal palladium solution, A., ii, 556.
- Sandos, Maurice.** See *Friedrich Kehrman.*
- Sands, Lila.** See *Fred W. Upson.*
- Sanfoureh, André,** the analysis of liquid nitrogen peroxide, A., ii, 454.
the reactions between gaseous oxides of nitrogen and alkaline solutions, A., ii, 762.
- Sanfoureh, André, and A. M. Boutin,** densities and refractive indices at 15° of mixtures of water, alcohol, and ether, A., i, 709.
- Sanna, G.** See *Bernardo Oddo.*
- Sano, Minoru,** phosphatides of fish sperm, A., i, 699.
optical properties of sphingomyelin, A., i, 701.
- Santesson, Carl Gustaf,** the action of poisons on enzymic processes. VII. Metal catalysis and catalase action. VIII. The volumetric method for the estimation of catalase, A., i, 1077.
- Santesson, Carl Gustaf.** See *G. J. von Blohm.*
- Sapper, Adolf.** See *Hans Reihlen.*
- Sarin, E.,** chemistry of the formation and ripening of honey, A., i, 197.
influence of organic acids on the formation and ripening of sugar-honey, A., i, 197.
- Sarkar, P. B., and Nitratan Dhar,** estimation of manganese by permanganate and investigation of some manganese, A., ii, 398.
- Sarkar, Pulin Vihari.** See *Priyadarajan Rây.*
- Sasaki, Nobuji,** the velocity of formation of barium peroxide, A., ii, 273.
- Sasaki, Nobuji,** photochemical examination of the reaction between ferric salts and iodides, A., ii, 772.
- Sasaki, Takaoki, and Tokudji Hashimoto,** benzylation and benzylation of 2:5-diketopiperazine, A., i, 56.
- Sasaki, Takaoki, and Jiro Kinose,** the degradation of *dl*- α -naphthylalanine by *Bacillus proteus*, A., i, 303.
- Sasaki, Takaoki, and Ichiro Otsuka,** the degradation of *l*-tryptophan by *Bacillus proteus*, A., i, 302.
- Saslavsky, Joh.** See *Nikolai Schilov.*
- Sato, Shinichi.** See *Christopher Kelk Ingold.*
- Sauer, E.** See *Alexander Gutbier.*
- Sauerwald, F.,** behaviour of carbon at high temperatures, A., ii, 443.
the increase in the size of grain of metallic bodies (not previously worked in the cold) formed from powdered material by pressure or sintering, A., ii, 746.
- Saunders, Horace Leonard,** the decomposition of ammonium nitrate by heat, T., 698.
- Saunders, Kenneth Herbert,** esters of the hydroxyalkylarylamines. I. Acid sulphuric esters of the simple monohydroxyethylarylamines, T., 2667.
- Saunders, Kenneth Herbert.** See also *British Dyestuffs Corporation, Ltd.*
- Saurwein, K.** See *Karl von Auwers.*
- Sauvageau, C., and Georges Denigès,** the efflorescences of *Rhodynenia palmata*; presence of a xylan in the Floridean algae, A., i, 507.
- Saville, William Bristow.** See *Martin Onslow Forster.*
- Sawyer, R. A., and A. L. Becker,** the production of enhanced line spectra, A., ii, 242.
the exploded-wire spectrum of calcium, A., ii, 803.
- Sayce, Leonard A., and A. Crawford,** estimation of carbon dioxide in mineral carbonates, A., ii, 316.
- Sayers, R. K., and W. P. Yant,** the tannic acid method for the estimation of carbon monoxide in blood, A., ii, 868.
- Sazerac, Robert, and C. Levaditi,** action of certain bismuth derivatives on syphilis, A., i, 89.
- Sborgi, Umberto, and Lelio Ferri,** borates. VI. The system $(\text{NH}_4)_2\text{O}-\text{B}_2\text{O}_3-\text{H}_2\text{O}$ at 45° and at 90°, A., ii, 568.
borates; the system $(\text{NH}_4)_2\text{O}-\text{B}_2\text{O}_3-\text{H}_2\text{O}$; temperature-concentration diagram, A., ii, 764.

- Sborgi, Umberto**, and **A. G. Nasini**, reaction between boron nitride and various metallic oxides with production of nitric oxide, *A.*, ii, 564.
- Sborowsky, I.** See **M. Sborowsky**.
- Sborowsky, M.**, and **I. Sborowsky**, a new accelerator for the destruction of organic matter in the Kjeldahl method for the estimation of nitrogen, *A.*, ii, 783.
- Scagliarini, Gino**, and **G. Torelli**, tervalent copper, *A.*, ii, 68.
catalytic action of copper in the oxidation of ammonia by means of persulphate, *A.*, ii, 81.
- Scalione, Charles C.** See **Arthur Becket Lamb**, and **David R. Merrill**.
- Scarborough, Harold Archibald.** See **Albert Eric Cashmore**, and **Hamilton McCombie**.
- Scatchard, George**, speed of reaction in concentrated solutions and the mechanism of the inversion of sucrose, *A.*, i, 230.
hydration of sucrose in water solution as calculated from vapour pressure measurements, *A.*, i, 230.
- Seelba, S.** See **Vincenzo Paolini**.
- Schaaß, Fritz.** See **August L. Bernoulli**.
- Schaap, O. P. A. H.**, an apparatus for the extraction of theobromine and caffeine by means of boiling chloroform, *A.*, ii, 797.
- Schaarschmidt, Alfred**, a new class of coloured reduction products of 1-benzoylanthraquinones or of 2:3-phthaloylbenzophenones, *A.*, i, 355.
- Schaarschmidt, Alfred**, **Maximilian Veidt**, and **Franz Schlosser**, compounds containing zinc derived from additive products of oxides of nitrogen and olefines, *A.*, i, 645.
- Schade, Heinrich**, physicochemistry of colloidal and supersaturated solutions of uric acid, *A.*, i, 1192.
- Schaefer, Clemens**, the regularities of ultra-red spectra and the failure of the classical dispersion theory, *A.*, ii, 727.
- Schaefer, Clemens**, and **Martha Schubert**, ultra-red reflection power of silica, *A.*, ii, 179.
characteristic ultra-red frequencies of selenates and chromates, *A.*, ii, 179.
characteristic ultra-red frequencies of chlorates, bromates, and iodates, *A.*, ii, 180.
ultra-red characteristic frequencies of the silicates, *A.*, ii, 727.
- Schaefer, R.** See **W. Eisenhardt**.
- Schaeppli, Hans**, permeability of cells and tissues. VIII. The question of the distribution of hormones and drugs in the blood, *A.*, i, 290.
- Schaffer, A. J.**, **Caspar Folkoff**, and **S. Bayne Jones**, the presence of nucleic acid in bacteria, *A.*, i, 1095.
- Schaller, K. A.**, and **W. Berndt**, a new apparatus for exact gas analysis, *A.*, ii, 862.
- Schaller, Waldemar Theodore**, sincosite, a new mineral, *A.*, ii, 450.
- Schamberg, J. F.**, **George W. Raiziss**, and **J. A. Kolmer**, sodium 2-nitro-4-hydroxymercuriphenoxide, *A.*, i, 391.
- Schames, Léon**, general improvement of the equation of condition; special equation of condition for hydrogen, *A.*, ii, 24.
- Schaposchnikov, K.**, quanta of the ideal monatomic gas, *A.*, ii, 839.
- Scharf, Albert.** See **Sigmund Fränkel**.
- Schanfelberger, P.** See **Ch. Gränacher**.
- Schaum, Karl** [with **Elli Riffert**], change of state of aggregation and polymorphism, *A.*, ii, 264.
- Schaum, Karl**, and **Theodor Marx**, colour of photochlorides and colloidal silver. II., *A.*, ii, 696.
- Scheibe, Gunter**, and **Gustav Schmidt**, di- and tri-quinolylmethanes united by the pyridine nuclei. III. Symmetrical di-2-quinolyl ketone, *A.*, i, 1190.
- Scheibler, Helmut**, and **Paul Baumgarten**, synthesis of *N*-alkylidene-amino-acids and their conversion into *N*-alkylamino-acids by hydrogenation, *A.*, i, 655.
- Scheibler, Helmut**, and **Artur Fischer**, synthetic experiments with ethenylcarbinols. I. Conversion of γ -methylbutinen- γ -ol into γ -methylbutenene, γ -methylbutanolone, and its condensation products, *A.*, i, 1108.
- Scheibler, Helmut**, and **Heinrich Ziegner**, metallic compounds of the enolic forms of carbonyl compounds and their application to syntheses. II. Synthesis of vinylideneglycol diethyl ether [keten-acetal]; explanation of the course of the reaction in the acetoacetic ester synthesis, *A.*, i, 426.
- Schellbach, Hans**, arsenical poisoning and the distribution of arsenic over the human body, *A.*, i, 900.
- Schellenberg, R.** See **Alfred Wohl**.
- Schelling, F.** See **Alexander Gutbier**.
- Schempp, Erick**, the behaviour of some cyclic compounds in the human and animal organisms, *A.*, i, 403.
- Schenck, A.** See **Aladar Skita**.
- Schenck, Martin**, the nitrogenous constituents of yeast, *A.*, i, 305.

- Schenck, Paul**, the influence of the thyroid gland on metabolism, with special reference to heat-regulation, A., i, 491.
the physiological action of β -amino-4-ethylglyoxaline (histamine), A., i, 498.
- Schenck, Rudolf** [with *Albert Imker*], germanium hydride, A., ii, 855.
- Schenk, O.** See *Paul Friedländer*.
- Schenk, Paul**, the influence of the thyroid on creatine-creatinine metabolism, A., i, 1212.
- Schenker, Robert**, the lipase of *Aspergillus niger* (van Tiegh), A., i, 203.
- Scherer, A.**, lung stones, A., i, 986.
- Scheringa, Klaas**, the system camphor-alcohol-water in relation to the titration of camphor-spirit, A., ii, 535.
- Scherrer, Paul**, and *P. Stoll*, determination of the Werner structure of inorganic compounds by means of X-rays, A., ii, 514.
- Schetselig, Jakob**, thorveitite, a silicate of scandium, A., ii, 306.
- Scheucher, Hermann**, invisible "mirrors" of arsenic, antimony, and bismuth, A., ii, 526.
- Scheuing, Georg**. See *Heinrich Wieland*.
- Schieber, W.** See *Alexander Gutbier*.
- Schiffner, Anton**. See *Anton Skrabal*.
- Schilov, E. A.**, action of silver sulphate in sulphuric acid solution on certain halogen derivatives, A., i, 913.
negative valency and co-ordination number, A., ii, 839.
- Schilov, E. A.** See also *P. P. Budnikov*.
- Schilov, Nikolai, Nad. Bulygina, Nik. Zerevtinov, Lidie Lepin, Olga Michajlova, Mich. Dubinin, Serg. Wosnesenski, Agness Ivanitakaja, Lidie Orlova, and Joh. Saslavsky**, adhesion forces in solutions. II. Adhesion series, A., ii, 350.
- Schilov, Nikolai**, and *Lidie Lepin* [with *Marie Jantschak*], partitioning of a substance between two solvents and the solution field of force, A., ii, 824.
- Schilov, Nikolai, Lidie Lepin, Marie Jantschak**, and *Michael Dubinin*, force of adhesion in solutions. III. Partition of substances between two solvents, A., ii, 626.
- Schillt, W.** See *Hermann Staudinger*.
- Schimmel & Co.**, estimation of chlorine in benzaldehyde, A., ii, 78.
- Schiets, A. B.**, electrolysis of aqueous solution of cerous salts, A., ii, 22.
- Schlack, P.** See *William Küster*.
- Schlatter, Gottfried**, lactic acid fermentation of dextrose by peptone, A., i, 1096.
- Schlayerbach**. See *William Küster*.
- Schleede, Arthur**. See *Erich Tiede*.
- Schlenk, Wilhelm, and Hermann Mark**, free pentaphenylethyl; the nature of the carbon linking, A., i, 1002.
analogues of pentaphenylethyl, A., i, 1003.
- Schlicht, A.**, bladder calculus of silicic acid, A., i, 496.
- Schlichting, Otto**. See *Heinrich Wieland*.
- Schliewiensky, H.**, reduction of acid chlorides to aldehydes by means of nickel catalysts, A., i, 938.
- Schlosser, A.** See *P. Karrer*.
- Schlosser, Franz**. See *Alfred Schaarschmidt*.
- Schlubach, Hans Heinrich, and Fritz Ballauf**, ammonium radicles. II. Tetraethylammonium. II. A., i, 15.
ammonium radicles. III. Ammonium, A., ii, 55.
- Schlubach, Hans Heinrich, and Edward C. Goes**, organo-metallic compounds. III. The mechanism of the Wurtz-Fittig synthesis, A., i, 1204.
- Schlundt, Herman**. See *Allen G. Shennstone*.
- Schmajewski, Ch.** See *Friedrich Kehrman*.
- Schmalz, Karl**. See *Hans Pringsheim*.
- Schmid, E.** See *A. Magnus*.
- Schmid, H.** See *Huns Rupe*.
- Schmidt, Carl L. A.**, preparation of cystine, A., i, 992.
- Schmidt, Carl L. A.**, and *Guy W. Clark*, fate of certain sulphur compounds when fed to the dog, A., i, 968.
- Schmidt, Carl L. A.**, and *G. F. Norman*, eosin hemolysis, A., i, 892.
- Schmidt, Carl L. A.** See also *Max S. Dunn*.
- Schmidt, Erich, Walter Barthelomé, and Alfred Lübke**, bromotrimethane. I, A., i, 826.
- Schmidt, Erich, and Karl Braunsdorf**, the natural proteins. I. Behaviour of chlorine dioxide towards organic substances, A., i, 781.
- Schmidt, Erich, and Franz Dausen**, in-crustive substances of plants. II., A., i, 206.
- Schmidt, Erich, and Heinrich Fischer**, preparation of *N*-nitroso-derivatives of secondary amines, A., i, 822.
- Schmidt, Erich, Richard Schumacher, Willy Bäjén, and Adalbert Wagner**, tetranitromethane. V. Tetranitromethane as nitrating agent. II., A., i, 738.

- Schmidt, Erich**, and **Rudolf Wilkendorf**, nitro- and amino-glycols. II., A., i, 314.
- Schmidt, E. G.** See **W. H. Peterson**.
- Schmidt, Friedrich**, hexamethyleneimine and its behaviour on oxidation, A., i, 761.
action of sulphuryl azide on benzene, A., i, 777.
- Schmidt, Friedrich**. See also **Theodor Curtius**.
- Schmidt, Gerhard**. See **Otto Ruff**.
- Schmidt, Gerhard Carl**, vapour pressure of binary mixtures, A., ii, 119.
ionisation brought about by quinine salts, A., ii, 182.
binary liquid mixtures, A., ii, 472.
- Schmidt, Gerhard Carl**, and **Th. Lücke**, diffusion of hydrogen through iron and platinum, A., ii, 198.
- Schmidt, Gustav**. See **Günter Scheibe**.
- Schmidt, H.** See **Th. Sabalitschka**.
- Schmidt, Hans**, aromatic compounds of antimony. V. Differing behaviour of lithium hydroxide from that of sodium or potassium hydroxide in the hydration of polymeric arylstibinic acids, A., i, 482.
aromatic antimony compounds, A., i, 1203.
- Schmidt, Harry**, relationship between the internal friction and the chemical constitution of gases, A., ii, 278.
- Schmidt, Julius**, and **Otto Spoun**, the phenanthrene series. XXXIII. Preparation of derivatives of phenanthraquinone and phenanthrene from 2-nitrophenanthraquinone, A., i, 665.
- Schmidt, John H.**, action of arsenious chloride on aniline, A., i, 285.
- Schmidt, K.** See **David Holde**.
- Schmidt, Karl**. See **Erwin Ott**.
- Schmidt, P.** See **Paul Jacobsen**.
- Schmidt, Walter**. See **Stefan Goldschmidt**.
- Schmidt-Hebbel, Edgar**. See **Fritz Paneth**.
- Schmidt-Nielsen, S.**, and **J. Holmsen**, composition of the urine of whales, A., i, 704.
- Schmidt-Nielsen, S.** See also **F. Frog**.
- Schmiedeskamp, Lloyd**. See **Leonard B. Loeb**.
- Schmitt, R.** See **Emil Heuser**.
- Schnabel, Alfred**, the distribution of cinchona alkaloids in the organism, A., i, 290.
a biological method for the estimation of substances which injure the cell and the embryo. II., A., i, 304.
- Schneider, A.** See **Charles Dhéré**.
- Schneider, M.** See **H. P. Kaufmann**.
- Schneider, Wilhelm**, and **Edgar Kraft**, sulphoacetic acid as condensing agent. IV. isoAcetovanillone, A., i, 749.
- Schneider, Wilhelm**, and **Albert Ross**, pyranhydrones. II. The constitution of diarylmethylpyrylium compounds, A., i, 1171.
- Schneller, Karl**. See **Hans Fischer**.
- Schödl, H.** See **Volkmar Kohlschütter**.
- Schöler, G.** See **A. Morgen**.
- Schoeller, A.**, micro-Kipp apparatus for the preparation of air-free carbon dioxide for use in the micro-estimation of nitrogen by Pregl's method, A., ii, 159.
micro-elementary analysis by Pregl's method, A., ii, 160.
simplified construction of the metal parts of the apparatus used in Pregl's micro-analytical method, A., ii, 160.
micro-incineration, A., ii, 654.
- Schoeller, Walter**, and **Walther Schrauth**, preparation of esters of complex mercuridicarboxylic acids and their products of hydrolysis, A., i, 605.
- Schöller, Walter**. See also **Fritz Weigert**.
- Schoen, Marcel**. See **Auguste Fernbach**.
- Schoen, Rudolf**, the non-haemoglobin nitrogen content of corpuscles a contribution to the nitrogen-metabolism of tissues, A., i, 484.
- Schönberg, A.**, o-quinones and 1:2-diketones. IV. Synthesis of acenaphthene, A., i, 27.
- Schönberg, A.**, and **O. Kraemer**, o-quinones and 1:2-diketones. V. Benzils of the peroxide type [ψ -benzils], A., i, 663.
- Schönert, K.** See **Gustav Tammann**.
- Schönfeld, B.** See **Andor Fodor**.
- Schönfelder, Heinrich**. See **Fritz Mayer**.
- Schoep, Alfred**, curite, a new radioactive mineral, A., ii, 77.
kasolite, a new radioactive mineral, A., ii, 154.
absence of cobalt in cornetite, A., ii, 219.
dewindtite, a new radioactive mineral, A., ii, 305.
stasite, a new mineral dimorphous with dewindtite, A., ii, 386.
becquerelite, a new radioactive mineral, A., ii, 450.
soddite, a new radioactive mineral, A., ii, 451.
- Schoep, Alfred**, and **W. Steinkuhler**, the estimation of uranium in presence of phosphoric acid, A., ii, 530.
- Schoepfle, C. S.**, phenyl-di- α -naphthylcarbinol and phenyl-di- α -naphthylmethyl, A., i, 337.

- Scholl, Roland** [with *Christian Seer, Richard Weitzenböck, and Arthur Ertl*], preparations in the naphthalene series, A., i, 650.
- Scholl, Roland, and Heinrich Neumann**, elimination of hydrogen from aromatic nuclei, and union of the latter by means of aluminium chloride. IV. Ring closure with doubly benzoylated naphthalenes, A., i, 261.
- Scholl, Roland, and G. Schwarzer**, elimination of hydrogen from aromatic nuclei and union of the latter by means of aluminium chloride. V. Experiments with benzil, stilbene, and phenanthrene, A., i, 331.
- Scholl, Roland, and Christian Seer**, elimination of hydrogen from aromatic nuclei and union of the latter by means of aluminium chloride. VI. Experiments with phenol ethers and with diphenylmethane, A., i, 336.
- Scholl, Roland, and Christian Seer** [with *Josef Dalmier*], elimination of hydrogen from aromatic nuclei and union of the latter by means of aluminium chloride, A., i, 258.
- Scholz, P.** See *Herbert Freundlich*.
- Schomer, Arnold**, estimation of yohimbine in yohimba bark, A., ii, 797.
- Schonebaum, C. W.**, the action of ozone on pure solutions of dextrose, levulose, and sucrose, A., i, 223.
the action of ozone on pure solutions of lactose, A., i, 717.
the action of hydrogen peroxide on pure solutions of dextrose, levulose, sucrose, lactose, and maltose, A., i, 920.
the action of ozone on pure solutions of maltose, A., i, 921.
- Schoneboom, C. O.**, diffusion and interaction, A., ii, 690.
- Schoofs, M.**, essential oils from *Myrica gale*, L., A., i, 1043.
- Schoorl, Nicolaas**, graphic deductions with regard to the dissociation of weak electrolytes, A., ii, 31.
the titration of the cinchona alkaloids and their salts, A., ii, 538.
- Schoorl, Nicolaas, and (Mlle) A. Regenbogen**, the system water-ethyl alcohol-chloroform; miscibility of the three components in different proportions and some practical applications, A., i, 215.
the system water-ethyl alcohol-carbon disulphide; miscibility of the three components in different proportions and some practical applications, A., i, 419.
- Schoorl, Nicolaas, and F. N. B. de Weerd**, the system, acetanilide-water, A., i, 247.
- Schoutissen, H. A. J.**, the action of nitrous acid on phenols, A., i, 135.
the velocity of reaction of diazotisation in its bearing on the problem of substitution in the benzene ring. II. The character of the diazonium group, A., i, 181.
- Schotte, Herbert.** See *Max Bergmann*.
- Schrader, Fritz.** See *Peter Paul Koch*.
- Schrader, Hans**, the spontaneous oxidation of lignin, natural humus material and coal and the influence of alkali thereon, A., i, 637.
- Schranz, Karl.** See *Karl Fleischer*.
- Schrauth, Walther, and Hanns Geller**, the preparation of α -amino- β -hydroxyacids from olefine-carboxylic acids, A., i, 1125.
- Schrauth, Walther.** See also *Walter Schoeller*.
- Schreinemakers, Frans Antoon Hubert**, non-, uni-, and bi-variant equilibria. XXI., A., ii, 430.
- Schreiner, Erling**, new conceptions of electrolytes. III. The hydration of the hydrogen ion, A., ii, 468.
new conception of electrolytes. IV. The dissociation state of acids of medium strength, especially dichloroacetic acid in water and in salt solutions, A., ii, 736.
- Schreiner, Erling.** See also *Chr. Winther*.
- Schroeder, E. von.** See *Gustav F. Hüttig*.
- Schroeder, Heinrich, and Trude Horn**, the relative distribution of carbohydrates in foliage in its dependence on water content, A., i, 906.
- Schroeter, Georg** [with *E. Kindermann, C. Dietrich, C. Beyschlag, C. Fleischhauer, E. Riebensahm, and C. Oesterlin*], hydrogenated naphthalenes and their transformations. II. Nitro- and amino-derivatives of tetrahydronaphthalene, A., i, 123.
- Schroeter, Georg** [with *Svanos, H. Einbeck, Hanns Geller, and E. Riebensahm*], hydrogenated naphthalenes and their transformations. III. Tetrahydronaphthalenesulphonic acids, tetrahydronaphthols and their derivatives, A., i, 126.
- Schroeter, Georg** [with *G. Vossen, F. Stahl, H. Haehn, and C. Prigge*], hydrogenated naphthalenes and their transformations. I., A., i, 122.

- Schroeter, Georg, and Tetralin**
G. m. b. H., preparation of α -keto-substituted hydrogenated naphthalenes, A., i, 1036.
preparation of symmetrical octahydro-anthracenes, A., i, 1136.
preparation of symmetrical octahydro-phenanthrenes, A., i, 1137.
- Schryver, Samuel Barnett.** See *Harold W. Buxton, and Donald Herbert Frank Clayton.*
- Schubardt, Wolfgang.** See *Hermann Strauss.*
- Schubert, Martha.** See *Clemens Schaefer.*
- Schubert, Max.** See *Hans Fischer.*
- Schuegraf, Karl.** See *Rudolf Pummerer.*
- Schufftan, Paul.** See *Walter Herz.*
- Schulenburg, Wilhelm.** See *Heinrich Wieland.*
- Schuler, J., NN'-di(p-allyloxyphenyl)-acetamide,** A., i, 179.
- Schulte, Therese.** See *Fritz Meyer.*
- Schultze, Ernst.** See *Otto Dimroth.*
- Schulz, Fr. N.,** the colouring matter and wax of the blood louse (*Schizoneura lanigera*), A., i, 494.
- Schulz, H.** See *H. von Wartenberg.*
- Schulze, Alfred,** the deviations from the gas laws of carbon disulphide, A., ii, 53.
binary liquid mixtures, A., ii, 424.
- Schulze, Heinrich, and Kurt Pieroh,** betulin, A., i, 1045.
- Schulze, M.** See *Friedrich Dolezalek.*
- Schumacher, Eugen.** See *Hartung Franzen.*
- Schumacher, Richard.** See *Erich Schmidt.*
- Schumm, Otto,** hæmatoporphyrin in the urine in cases of lead poisoning, A., i, 609.
the detection of small quantities of lead in urine, A., ii, 317.
- Schumm, Otto, and A. Papendieck,** the presence and detection of tyrosine in urine, A., i, 899.
- Schuster, Kurt.** See *Kurt Heinrich Meyer.*
- Schwab, G. M.** See *Ernst Hermann Riesenfeld.*
- Schwalbe, Carl Gustav, and Ernst Becker,** the elimination of furfuraldehyde from oxycelluloses; the solubility in alkali and the reduction capacity of oxycelluloses, A., i, 232.
- Schwappacher, H.** See *Otto Fischer.*
- Schwartz, A.** See *Louis Fournier.*
- Schwartz, Benjamin,** hæmatotoxins from parasitic worms, A., i, 493.
- Schwarz, P.,** detection of benzene in light petroleum, A., ii, 531.
- Schwarz, Robert, and Walter Friedrich,** the influencing of the catalysis of hydrogen peroxide with platinum by Röntgen rays, A., ii, 436.
- Schwarz, Robert, and Erika Herrmann,** metachromism of toluidine-blue, A., i, 930.
- Schwarz, Robert, and Erich Konrad,** the mechanism of the formation of silane from magnesium silicide. I., A., ii, 846.
- Schwarz, Robert, and Heinrich Stöck,** the effect of colloids on silver bromide emulsions, A., ii, 731.
- Schwarzenberg, K.** See *Julius Tröger.*
- Schwarzer, G.** See *Roland Scholl.*
- Schwebel, W.** See *Ernst Berl.*
- Schweiger, K.,** the solubility of cellulose acetate in the salts of the alkali- and alkaline-earth metals, A., i, 324.
- Schwen, Gustav.** See *Wilhelm Steinkopf.*
- Schwenk, Erwin,** condensation reactions of formic acid, A., i, 153.
preparation of hydroxyanthraquinones from nitroanthraquinones, A., i, 153.
- Seilasi, Wilhelm.** See *Karl Freudenberg.*
- Sciortino, Antonio,** accurate ureometer, A., ii, 794.
- Scofield, C. S.,** effect of alum on silicate colloids, A., i, 212.
- Scott, A. F.** See *Gregory Paul Baxter.*
- Scott, Alfred W.** See *Lauder William Jones.*
- Scott, E. Kilburn,** piezo-electricity of potassium sodium tartrate crystals, A., ii, 609.
- Scott, James, and Robert Robinson,** note on 2:3- and 2:5-dinitro-*p* toluidines, T., 844.
- Scott, John Richard, and Julius Berend Cohen,** the condensation of aromatic *o*-aminosulphonic acids with isocyanic acid, T., 2034.
- Scott, M.** See *William Ridgely Orndorff.*
- Scott, Wilfred W.,** glacial acetic acid method for estimating uranium in chertotite, A., ii, 788.
- Scottish Dyes, Ltd.** See *Arthur Hugh Davies, and John Thomas.*
- Sears, O. H.** See *Samuel D. Conner.*
- Sebor, J.** See *Julius Stoklass.*
- Seefried, Hermann.** See *Heinrich Wieland.*
- Seel, K.,** the analysis of high-percentage tungsten alloys, A., ii, 876.
- Seeliger, Rud.,** excitation of atoms to emission of light by electron collisions. IV. Behaviour of individual members of a series, A., ii, 17.

- Seeliger, Rud.**, excitation of atoms to emission of light by electron collisions, A., ii, 804.
- Seeliger, Rud.**, and **K. Lapkamp**, absorption of gases by crystals, A., ii, 80.
- Seeman, H.**, linear polarisation of the blue light from the focus of the Lillienfeld Röntgen tube, A., ii, 8.
a precision Röntgen spectrograph, A., ii, 15.
- Seemann, Jon.** See **Julius von Braun**.
- Seer, Christian.** See **Roland Scholl**.
- Sefton, Lily Bell.** See **William Lloyd Evans**.
- Segaller, David, David Henry Peacock, and British Dyestuffs Corporation, Ltd.**, manufacture of hydroxy- and sulphohydroxy-derivatives of anthraquinone, A., i, 560.
- Seiberth, M.** See **M. Hartmann**.
- Seidel, C. F.** See **Leopold Ruzicka**.
- Seidel, Fritz.** See **Alfred Eckert**.
- Seidel, Walter.** See **Max Trautz**.
- Seidell, Atherton**, isolation of vitamin, A., i, 887.
further experiments on the isolation of the antineuritic vitamin, A., i, 1078.
- Seidl, R.** See **Benno Bleyer**.
- Seka, Reinhard.** See **Ernst Philippi**.
- Sekera, F.**, theory of the mechanical synthesis of colloids, A., ii, 693.
- Seltz, Harry.** See **Herbert S. Harned**.
- Sen, Nirmal Kumar.** See **Sikhibhushan Dutt**.
- Senderens, Jean Baptiste, and Jean Aboulenc**, catalytic hydrogenation of polyphenols by the wet way, A., i, 136.
catalytic preparation of cyclohexanetriols, A., i, 337.
- Sen-Gupta, Hemendra Kumar, and Stanley Horwood Tucker**, the oxidation of α -dinaphthanthens, T., 557.
- Senseman, C. E.** See **O. A. Nelson**.
- Servantie, L.** See **P. Mauriac**.
- Settle, Richard Hardcastle.** See **Hamilton McCombie**.
- Seydel, Karl.** See **Heinrich Bilts**.
- Seyewetz, Alphonse, and P. Sisley**, antioxidants and antioxygenisers, A., ii, 628.
- Seyewetz, Alphonse, and Vignat**, action of sodium sulphite on nitrobenzene, A., i, 241.
- Shaffer, Philip Anderson**, antiketogenesis. III. Calculation of the ketogenic balance from the respiratory quotients, A., i, 83.
- Shaffer, Philip Anderson.** See also **A. P. Briggs**.
- Shaffer, Sherman**, substances dissolved in rain and snow, A., i, 512.
- Shah, R. C.** See **John Joseph Sudborough**.
- Shannon, Earl V.**, cristobalite in basalt from Washington, A., ii, 450.
velardeñite from California, A., ii, 517.
an andorite-bearing silver ore from Nevada, A., ii, 859.
- Shannon, Earl V., and Edgar Theodore Wherry**, white chlorites, A., ii, 517.
- Shannon, Earl V.** See also **Edgar Theodore Wherry**.
- Shannon, (Miss) Mary Isobel.** See **James Colquhoun Irvine**.
- Sharp, Paul Francis, and Frank Henry MacDougall**, simple method of electro-metric titration in acidimetry and alkalimetry, A., ii, 579.
- Sharp, Thomas Marvel.** See **Thomas Anderson Henry**.
- Shaver, W. W.**, the absorption spectrum of liquid and gaseous oxygen, A., ii, 331.
- Shaw, J. A.**, method for the estimation of free and combined carbon dioxide, A., ii, 226.
- Sheaff, Howard M.**, estimation of minute amounts of gaseous oxygen and its application to respiratory air, A., ii, 582.
- Shedd, Oliver March**, a comparison of the calcium content of some virgin and cultivated soils of Kentucky by an improved method for the estimation of this element, A., ii, 527.
- Sheehy, Edmond John**, origin of milk fat, and its relation to the metabolism of phosphorus, A., i, 400.
- Sheldon, Wilfrid.** See **Edward Hope**.
- Shenefeld, S. Lantz, Frank C. Villbrandt, and James E. Withrow**, sulphate-free sulphites for standard sulphur dioxide solutions, A., ii, 45.
- Shenstone, Allen G.**, an attempt to detect induced radioactivity resulting from α -ray bombardment, A., ii, 415.
- Shenstone, Allen G., and Herman Schlundt**, a determination of the number of α -particles per second emitted by thorium-C of known γ -ray activity, A., ii, 465.
- Sheppard, Samuel Edward, and Felix A. Elliott**, the drying and swelling of gelatin, A., i, 387.
- Sheppard, Samuel Edward, S. S. Sweet, and Amber J. Benedict**, elasticity of purified gelatin jellies as a function of the hydrogen-ion concentration, A., ii, 747.
- Sherman, Henry Clapp**, chemical investigation of amylases and related enzymes, A., i, 66.

- Sherman, Henry Clapp, and Mary L. Caldwell**, the influence of arginine, histidine, tryptophan, and cystine on the hydrolysis of starch by purified pancreatic amylase, A., i, 283.
- Sherman, Henry Clapp, Victor K. La Mer, and H. L. Campbell**, effect of temperature and of the concentration of hydrogen ions on the rate of destruction of antiscorbutic vitamin (vitamin-C), A., i, 207.
estimation of the antiscorbutic vitamin (vitamin-C), A., ii, 407.
- Sherman, Henry Clapp, and Florence Walker**, the influence of certain amino-acids on the enzymic hydrolyses of starch, A., i, 283.
- Sherman, Henry Clapp, and Marguerite Wayman**, effect of certain antiseptics on the activity of amylases, A., i, 282.
- Sherman, Henry Clapp.** See also *Victor K. LaMer*.
- Sherwin, Carl P.** See *J. A. Killian, and George J. Shipley*.
- Shewky, Marian C.** See *Burton A. Myers*.
- Shibata, Yûji, and Kenjiro Kimura**, chemical investigations of Japanese minerals containing rarer elements. II. Analysis of columbite and monazite of Ishikawa, Iwaki province, A., ii, 220.
chemical investigation of Japanese minerals containing rarer elements. IV. Samarskite and an unnamed mineral from Ishikawa, Iwaki province, A., ii, 516.
supplementary note on ishikawaite, a new mineral from Ishikawa, Iwaki province, A., ii, 861.
- Shibata, Yûji, and Willy Nagai**, the absorption of colouring matters of the flavone group. II. Absorption spectra of phenyl styryl ketone and its derivatives, A., ii, 413.
- Shibata, Yûji, and Taku Uemura**, Japanese minerals containing rare elements. III. Analyses of beryl of Naegi, Mino province, A., ii, 305.
- Shimizu, Toraji.** See *Yasuhiko Asahina*.
- Shimizu, Tomihide**, the fate of some polysaccharides in the digestive tract of mammals, A., i, 82.
hydrolysis of some polysaccharides (inulin, lichenin, and hemicellulose) in the digestive tract of mammals, A., i, 82.
influence of some polysaccharides (inulin, lichenin, and hemicellulose) on protein exchange, A., i, 83.
- Shimizu, Tomihide**, the chemical composition of brain, A., i, 85.
the behaviour of pyrrole in the animal body, A., i, 91.
behaviour of phrenosin in the animal body, A., i, 91.
- Shimizu, Tomihide.** See also *Martin Jacoby*.
- Shimo, Kôtarô**, constituents of *Phellodendron amurense*, A., i, 99.
- Shimomura, Akira, and Julius Berend Cohen**, the interchange of alcohol radicles in esters. I. and II., T., 883, 2051.
- Shipley, George J., and Carl P. Sherwin**, synthesis of amino-acids in animal organisms. I. Synthesis of glycine and glutamine in the human organism, A., i, 492.
fate of some of the phenylacetylated amino-acids in the animal organism, A., i, 1093.
- Shlager, M.** See *Ellwood B. Spear*.
- Shoesmith, John Baldwin.** See *Arthur Lapworth*.
- Shohl, Alfred T.**, effect of hydrogen-ion concentration on the estimation of calcium, A., ii, 395.
- Shohl, Alfred T., and Frank G. Pedley**, a rapid and accurate method for estimating calcium in urine, A., ii, 395.
- Shoji, R.** See *Joseph Barcroft*.
- Short, Wallace Frank, and John Charles Smith**, the condensation of phenols with the hydrochlorides of cyanamides and carbodi-imides, and its relation to the Hoesch reaction, T., 1803.
- Showalter, M. F., and R. H. Carr**, characteristic proteins in high- and low-protein maize, A., i, 1103.
- Siderfin, Norman Edward.** See *William Lewcock*.
- Sidersky, D.**, solubility of crystallised strontium hydroxide, A., ii, 501.
- Sidgwick, Nevil Vincent, and Herbert Clayton**, the solubility of the aldehydobenzoic acids, T., 2263.
- Sidgwick, Nevil Vincent, and Wilfred Marsden Dash**, the solubility and volatility of the nitrobenzaldehydes, T., 2586.
- Sidgwick, Nevil Vincent, and (Miss) Elinor Katharine Ewbank**, the solubility of the alkali salts of benzoic and the hydroxybenzoic acids in water, T., 1844.
- Sidgwick, Nevil Vincent, and Joseph Alfred Hector Roberts Gentle**, the solubilities of the alkali formates and acetates in water, T., 1837.

- Sidgwick, Nevil Vincent, and Thomas Weston Johns Taylor**, the solubility and volatility of 3:5-dinitrophenol, T., 1853.
- Sidgwick, Nevil Vincent, and Sydney Leonard Turner**, the solubility of the chlorophenols, T., 2256.
- Siebeck, R.** [with **D. Hackmack**], exchange of chlorine between the red blood corpuscle and the surrounding solution. I. The influence of narcotics on the exchange of chlorine, A., i, 1208.
- Siebeneck, H.**, the influence of the elements of the oxygen group on paraffin wax, A., i, 417.
- Sieber, Wilhelm.** See **Theodor Curtius**.
- Siebert, Erich.** See **Emil Fromm**.
- Sieburg, Ernst, and Franz Bachmann**, the influence of treatment with alkali or bromine on the physiological activity and foaming capacity of some saponaceous substances, A., i, 404.
- Siedentopf, H.** See **Heinrich von Halban**.
- Siegbahn, Manne**, new measurements of precision in the X-ray spectrum, A., ii, 104.
- Siegesmund, J. C.** See **M. G. Mellon**.
- Siegfried, K.**, the hydrocyanic acid question, A., i, 614.
- Sieglitz, Adolf** [with **Hans Spitzer**], fission of $\alpha\alpha$ -diphenylethyl- β -urethane, A., i, 828.
- Sieglitz, Adolf, and H. Jassoy**, the fluorene series. VII. Dibenzofulvene, A., i, 820.
- Sieglitz, Adolf.** See also **Fritz Mayer**.
- Sieke, Fritz**, formation of phenol by bacteria, A., i, 902.
- Siemssen, J. A.**, the triboluminescence of uranium salts and three new organic uranium compounds, A., ii, 511.
- Sievers, W. C., and L. Givaudan & Co.**, preparation of vanillin from acetyl-eugenol, A., i, 38.
- Sieverts, Adolf**, velocity of decomposition of crystalline substances, A., ii, 700.
- Sieverts, Adolf, and H. Theberath**, velocity of decomposition of silver permanganate, A., ii, 360.
- Silberrad, Oswald**, researches on sulphuryl chloride. II. A new chlorinating agent; preparation of polychloro-derivatives of benzene, T., 1016.
- the formation of hexachloroethane from chloropierin, A., i, 2.
- Silberstein, Adolf**, estimation of hydrogen-ion concentration in urine with indicators, A., ii, 452.
- Silberstein, Ludwik**, some spectrum lines of neutral helium derived theoretically, A., ii, 674.
- Silberstern, Ernst**, modification of the iodine test for bile pigments in urine, A., ii, 799.
- Silabee, Clara G.** See **Richard Fay Jackson**.
- Simion, Fritz**, a proposed estimation of chromium and nickel in steel, A., ii, 529.
- Simola, P. G.** See **Ossian Aschan**.
- Simmonds, Nina.** See **Victor E. Levine, and C. R. Orton**.
- Simmons, John P.** See **Arthur Becket Lamb**.
- Simms, H. S.** See **Phæbus A. Levens**.
- Simon, Franz**, determination of specific heats at low temperatures, A., ii, 684.
- Simon, Italo**, behaviour of 5-iodoguaiacol in the human organism, A., i, 970.
- Simon, Kurt.** See **Hans Lecher**.
- Simon, Louis Jacques**, the chromic oxidation of the homologues of acetic acid, A., i, 803.
- direct oxidation of esters of hydroxyacids by oxygen or air, A., i, 984.
- oxidation with mixtures of sulphuric acid and chromates, A., ii, 593.
- the function of chromic oxide in oxidation by means of sulpho-chromic mixture, A., ii, 868.
- Simon, Louis Jacques, and A. J. A. Guillaumin**, quantitative determination of carbon and hydrogen by means of the sulpho-chromic mixture, A., ii, 867.
- Simon, Louis Jacques, and L. Zivy**, the neutralisation of tartaric acid by potassium hydroxide in presence of chlorides of alkaline earths, A., ii, 880.
- Simon, Max**, the influence of asphyxiation on the permeability of the limiting membrane of the muscle fibres, A., i, 196.
- Simor, Max.** See also **Hermann Lange**.
- Simonnet, H.** See **R. Tiger**.
- Simonnet, M.** See **Paul Cristol**.
- Simons, Harold Lester**, estimation of gases in metals, A., ii, 719.
- Simonsen, John Lionel**, the constitution of the terpene present in the essential oil from *Andropogon Swarancusa*, Jones, T., 2292.
- the essential oil from the leaves of *Abies pindrow*, Spach, A., i, 753.
- Simonsen, John Lionel, and Madyar Gopal Rau**, the essential oil from *Blumea Malcomii*, T., 876.
- Simpson, Charles.** See (*Miss*) **Lucy Higginbotham**.

- Simpson, Daniel H.** See *Ernest C. McKelvy*.
- Simpson, Stephen G.**, effect of the presence of filter-paper on permanganate-oxalate titrations, A., ii, 234.
- Sims, L. B.** See *Floyd E. Bartell*.
- Sindlinger, F.** See *Felix Mach*.
- Singh, Bawa Kartar**, recent advances in stereochemistry, A., ii, 103.
- Singh, Bawa Kartar, Raghunath Rai, and Rattan Lal** studies in optically active dyes. I. Camphoreins, T., 1421.
- Singh, Gopal**, phototropy of inorganic salts; cuprous chloride and bromide, T., 782.
- Singleton, W., and H. Williams**, inadequacy of the "A.R." test for alkalis in calcium carbonate, A., ii, 527.
- Sinha, J. N.** See *Julius Huebner*.
- Sinkinson, Eric.** See *William Arthur Bone*.
- Sircar, Anukul Chandra, and Sukh-bhushan Dutt**, dyes derived from camphoric anhydride, T., 1283.
dyes derived from phenanthraquinone. I. Phenanthranaphthazines, T., 1944.
- Sirovich, G., and A. Cartoceti**, phenomena of diffusion in metals in the solid state and cementation of non-ferrous metals. I. Cementation of copper by means of ferro-manganese, A., ii, 68.
phenomena of diffusion in metals in the solid state and cementation of non-ferrous metals. II. Cementation of copper by means of chromo-manganese, A., ii, 571.
- Sisley, P.** See *Alphonse Seyewetz*.
- Sisson, Warren R., and Willey Denis**, inorganic constituents of milk. I. Chlorides in human milk, A., i, 87.
- Sisson, Warren R.** See also *Willey Denis*.
- Sjöberg, K.** See *Olof Svanberg*.
- Skaupy, Franz**, molecular condition of metals dissolved in mercury, A., ii, 297.
- Skinker, M. F.**, the motion of electrons in carbon dioxide, A., ii, 837.
- Skinner, Helen H.** See *Victor Lenher*.
- Skita, Aladar**, determination of the configuration of the stereoisomeric hexamethylenes, A., i, 534.
the mechanism of catalytic hydrogenation, A., ii, 207.
- Skita, Aladar, and A. Schenck**, stereoisomerism of cyclic hydrocarbons, A., i, 240.
- Skrabal, Anton**, Landolt's reaction, A., ii, 488.
- Skrabal, Anton, and Emmy Raith**, the alkaline hydrolysis of the esters of the two ethylenedicarboxylic acids, A., ii, 434.
- Skrabal, Anton, and Anton Schifferer**, velocity of hydrolysis of acetals, A., ii, 134.
- Skraup, Siegfried, and Leo Freundlich**, halochromic phenomena with carbinols, A., i, 539.
- Skraup, Siegfried, and Marie Moser**, benzoxazole derivatives, A., i, 574.
- Skutil, František**, the Kjelmann modification of the Kjeldhal process, A., ii, 582.
- Slagle, Edgar A.** See *Solomon Farley Acree, and R. R. Mellon*.
- Slansky, Pawel**, the drying of fatty oils, A., i, 1113.
- Slater, F. P.**, excitation of γ -radiation by α -particles from radium emanation, A., ii, 13.
rise of γ -ray activity of radium emanation, A., ii, 681.
- Sluiter, E.**, periodicity of enzymes; the lipase of the stomach, A., i, 887.
- Slyke, Donald D. van**, the measurement of buffer values and the relationship of buffer value to the dissociation constant of the buffer and the concentration and reaction of the buffer solution, A., i, 893.
apparatus for estimation of the gases in blood and other solutions, A., ii, 158.
acidosis. XVIII. Estimation of the hydrogen carbonate concentration of the blood and plasma, A., ii, 672.
- Slyke, Donald D. van, Joshua Harold Austin, and Glenn E. Cullen**, effect of ether anaesthesia on the acid-base balance of the blood, A., i, 1025.
- Slyke, Donald D. van, and Alma Hiller**, an unidentified base among the hydrolytic products of gelatin, A., i, 63.
- Slyke, Donald D. van, and William C. Stadie**, estimation of the gases of the blood, A., ii, 78.
- Slyke, Donald D. van.** See also *Joshua Harold Austin, A. Baird Hastings, and Alma Hiller*.
- Smekal, Adolf**, fine structure of Röntgen spectra, A., ii, 15, 600.
systematics of Röntgen spectra, A., ii, 181.
theory of Röntgen spectra. II., A., ii, 607.
- Smelkus, H.** See *Julius Marcusson*.
- Smiles, Samuel, and Hugh Graham**, derivatives of diphenylthiolbenzene, T., 2506.

- Smiles, Samuel**, and **Douglas Creese Harrison**, *p*-dithiobenzoic acid, T., 2022.
- Smiles, Samuel**, and **Ernest Wilson McClelland**, the interaction of aromatic disulphides and sulphuric acid, T., 86.
- Smirnov, Alexander P.**, 1-phenyl-4-pyridone, A., i, 464.
- Smirnov, Alexander P.** See also **Paul Karrer**.
- Smith, Alpheus W.**, and **Cecil E. Boord**, absorption spectra of benzeneazophenol and its derivatives, A., ii, 602.
- Smith, Clarence Joseph**, an experimental comparison of the viscous properties of (a) carbon dioxide and nitrous oxide; (b) nitrogen and carbon monoxide, A., ii, 549.
- viscosity and molecular dimensions of sulphur dioxide, A., ii, 686.
- viscosity and molecular dimensions of gaseous carbon oxysulphide, A., ii, 686.
- Smith, Clarence Joseph**. See also **Alexander Oliver Rankine**.
- Smith, C. M.** See **J. J. T. Graham**.
- Smith, Edgar Fahs**, sodium tungstates. I., A., ii, 774.
- Smith, Edwin S.** See **John Stewart**.
- Smith, Francis Edward**. See **Donald Rhind**.
- Smith, Frederick William**. See **Thomas James Drakeley**.
- Smith, Gilbert B. L.** See **Victor Lenher**.
- Smith, G. Frederick**. See **Hobart Hurd Willard**.
- Smith, George McPhail**, additive compounds of gold haloids with benzyl sulphide, A., i, 933.
- precipitation of metals by hydrogen sulphide, A., ii, 626.
- Smith, Henry George**, and **Arthur Ramon Penfold**, preparation of thymol, menthane, and menthol from eucalyptus oils, A., i, 81.
- Smith, Henry George**. See also **John Read**.
- Smith, Homer W.**, nature of secondary valency. III. Partition coefficients in the system water-ethyl ether, A., ii, 270.
- nature of secondary valency. IV. Partition coefficients in the system glycerol-acetone, A., ii, 270.
- nature of secondary valency. V. Partition coefficients in systems containing water as one component, with special reference to the absolute values of the series constants, A., ii, 430.
- nature of secondary valency. VI., A., ii, 431.
- Smith, (Miss) Isobel Agnes**. See **Alexander McKennie**.
- Smith, John Charles**. See **Wallace Frank Short**.
- Smith, John David Main**, Friend's theory of valency, A., ii, 279.
- Smith, John David Main**. See also **Gilbert Thomas Morgan**.
- Smith, James Leonard Brierley**. See **William Hobson Mills**, and (Sir) **William Jackson Pope**.
- Smith, Lennart**, preparation of β -chlorohydrins. A., i, 3.
- acetic acid esters of multivalent alcohols. I. Is there a connexion between the velocity of saponification and the method of preparation of the glycerol acetins? A., i, 915.
- titration of tin with ferric chloride, A., ii, 398.
- Michael's addition and separation rule, A., ii, 751.
- Smith, Lennart**, and **Hugo Olsson**, dependence of the velocity of alkaline hydrolysis of esters on the constitution of the alcohol. I., A., ii, 701.
- Smith, Lennart**, and **B. Platon**, kinetic determinations of the constitutions of hydroxy- and amino-chloropropanes, A., i, 1121.
- Smith, L. I.**, the action of halogens on ethyl acetoacetate, A., i, 318.
- Smith, L. I.** See also **Elmer Peter Kohler**.
- Smith, Millard**, a micro-modification of Benedict's method for the estimation of reducing sugar in urine, A., ii, 663.
- Smith, Maurice E.**, Friedel and Crafts' reaction; the carbomethoxybenzoyl chlorides with aromatic hydrocarbons and aluminium chloride, A., i, 140.
- Smith, Nicol H.** See **John H. Müller**.
- Smith, Thomas M.** See **Arthur E. Hill**.
- Smith, William Thompson**, and **Reginald E. Parkhurst**, solubility of sulphur dioxide in suspensions of calcium and magnesium hydroxides, A., ii, 761.
- Smithells, Arthur**, the blue flame produced by common salt on a coal fire, A., ii, 645.
- Smithells, Colin James**. See **General Electric Co., Ltd.**
- Smithey, Ira W.** See **Alwin Sawyer Wheeler**.
- Smits, Andreas**, electromotive behaviour of aluminium, A., ii, 20.
- application of the theory of allotropy to electromotive equilibria. IV., A., ii, 110.
- influence of intensive drying on internal change, A., ii, 358.

- Smits, Andreas**, and *C. J. de Gruyter*, electromotive behaviour of aluminium. III., A., ii, 21.
- Smitt, Nicholas Karrachy**, a rapid method for the estimation of acet-aldehyde, A., ii, 402.
- detection and estimation of fluorides, A., ii, 716.
- Smolenski, Eugenie**, and *Kazimir Smolenski*, preparation of amines from alcohols and ammonia, A., i, 234.
- Smolenski, Kazimir**. See *Eugenie Smolenski*.
- Smorodincev, A.**, reductases. I. Some conditions of the activity of starch reductase. I., A., i, 1201.
- the materials extracted from muscles. XXI. The organic bases of the flesh of swine. I., A., i, 1213.
- Smyth, Charles P.** See *Theodore William Richards*.
- Smythe, John Armstrong**, use of iron pyrites in a Friedel-Crafts' reaction, T., 1270.
- decomposition of benzyl disulphoxide, T., 1400.
- Smythe, William R.**, the spectrum of fluorine, A., ii, 99.
- Snapper, J.**, and *Ernst Laqueur*, estimation of hippuric acid in urine, A., ii, 92.
- Snapper, J.**, and *W. J. van Bommel van Vloten*, estimation of indican in blood-serum, A., ii, 793.
- Société Chimique des Usines du Rhône**, preparation of hydroxyaldehydes and their derivatives, A., i, 257.
- preparation of *n*-butyl *p*-aminobenzoate, A., i, 827.
- preparation of acetic anhydride, acet-aldehyde, or acetic acid from ethylidene diacetate, A., i, 915.
- preparation of a basic aluminium salicylate, A., i, 1151.
- Society of Chemical Industry in Basle**, preparation of *p*-methylaminophenol, A., i, 31.
- preparation of soluble derivatives of camphoric acid, A., i, 254.
- preparation of *o*-hydroxyazo-colouring matters, A., i, 385.
- preparation of intermediate products and colouring matters, A., i, 579.
- preparation of benzyl *p*-aminobenzoate, A., i, 828.
- preparation of a derivative of pyridine-3-carboxylic acid (nicotinic acid), A., i, 860.
- preparation of an unsaturated ether of *p*-dihydroxydiphenylacetamidine, A., i, 953.
- preparation of dialkylamides of nicotinic acid, A., i, 1058.
- Society of Chemical Industry in Basle**, preparation of β -thionaphthosatin, A., i, 1179.
- manufacture of amino-alcohols of the quinoline series, A., i, 1179.
- Soderstrom, G. F.** See *H. V. Atkinson*.
- Sörensen**. See *Otto Diels*.
- Sörensen, M.** See *Soren Peter Lauritz Sörensen*.
- Sörensen, Soren Peter Lauritz, M. Sörensen**, and *K. Linderström-Lang*, the "salt error" of the quinhydrone electrode, A., ii, 111.
- Solonina, B.** See *Friedrich Kehrmann*.
- Soma, S.** See *Kuji Miyake*.
- Sommelet, Marcel**, and *Jean Guioth*, hydrogenation by formic acid of some quaternary salts of hexamethylene-tetramine, A., i, 333.
- Somogyi, R.**, swelling of fibrin by acids, A., i, 182.
- action of acids on fermentation by yeast, A., i, 201.
- Somogyi, R.** See also *Isidor Traube*.
- Sondén, Klas**, the application of coloured glasses instead of liquids in colorimetric researches, A., ii, 862.
- Sonn, Adolf**, and *Susanne Falkenheim*, synthesis of fisetole, A., i, 1163.
- Soutar, Charles William**. See *Frederick William Attack*.
- Souza, David Henriques de**. See *James Arthur Hewitt*.
- Spacu, G.**, complex magnesium salts, A., i, 859.
- Späth, Ernst**, the anhalonium [cactus] alkaloids. II. Constitution of pello-tine, anhalonidine, and anhalamine, A., i, 163.
- constitution of kynmenic acid, A., i, 173.
- the anhalonium [cactus] alkaloids. III. The constitution of anhaline, A., i, 567.
- Späth, Ernst**, and *Karl Böhm*, constitution of the alkaloids of the calumba root, A., i, 1174.
- Späth, Ernst**, and *Karl Fuchs*, the active constituents of the true coto-bark; the synthesis of cotoin, A., i, 558.
- Späth, Ernst**, and *Norbert Lang*, conversion of berberine into paluatine, A., i, 166.
- the constitution of corydaline, A., i, 168.
- the synthesis of landanine, A., i, 568.
- Späth, Ernst**, and *Hans Röder*, anhalonium [cactus] alkaloids. IV. Synthesis of anhalamine, A., i, 852.
- Späth, Ernst**, and *Erich Tichelnitz*, the constitution of ricinine, A., i, 571.
- Späth, Wilhelm**, selenium [cells], A., ii, 189.

- Speakman, John Bamber.** See *John Greenwood Heap*, and *William Jacob Jones*.
- Spear, Ellwood B., P. F. Jones, A. S. Neave, and M. Schlager,** action of ultra-violet light on colloidal platinum, A., ii, 11.
- Speckan, Carl,** changes in the sweetness of dulcin (*p*-phenetolecarbamide) caused by chemical alterations of the molecule and the sweetening power of derivatives of *p*-hydroxyphenylcarbamide, A., i, 579.
- Speich, Leonhard.** See *Julius Meyer*.
- Speldel, Johann Adolf.** See *Burckhardt Hölferich*.
- Speirani, M.** See *H. Charrier*.
- Spencer, George Douglas.** See *British Dyestuffs Corporation, Ltd.*, and *Arthur George Perkin*.
- Spencer, James Frederick, and Kathleen Froud,** electrolysis of sodium silicate solutions, A., ii, 611.
- Sperry, W. N.** See *Roger Adams*.
- Speyer, Edmund, and Alfred Gustav Becker,** action of hydrogen peroxide on cinchona alkaloids, A., i, 674.
- Speyer, Edmund, and Gunther Becker,** morphine, A., i, 675.
- Speyer, Edmund, and Hermann Wieters,** hydrogenation of *isocodeine* and *ψ-codeine*, A., i, 47.
- codeineoxidesulphonic acids and their derivatives*, A., i, 169.
- Speyer-Haus, Georg,** preparation of derivatives of 3:3'-diamino-4:4'-dihydroxyarsenobenzene, A., i, 481.
- Spiegel, E. A.** See *Herbert Elias*.
- Spiegel, Leopold,** enzymic fat synthesis, A., i, 694.
- Spiller, John,** obituary notice of, T., 748.
- Spiro, Karl,** the calcium-potassium action, A., i, 85.
- lyotrope series and β-oxidation*, A., i, 489.
- decarboxylation*, A., i, 1213.
- Spiro, Karl, and A. Stoll,** the active constituent of ergot, A., i, 47.
- Spitzer, Anna.** See *J. Pollak*.
- Spitzer, Hans.** See *Adolf Sieglitz*.
- Sponsel, K.** See *Erich Müller*.
- Spoun, Otto.** See *Julius Schmidt*.
- Spreckelsen, Otto.** See *Ernst Thielepape*.
- Springer, Friedrich,** comparative plant chemistry. III. *Campanula rotundifolia*, L., A., i, 797.
- Sproemer, William C., and Guy B. Taylor,** vapour pressures of aqueous solutions of nitric acid, A., ii, 143.
- Srajevid, V.** See *Maximilian Samec*.
- Staden, A. von.** See *Adolf Windaus*.
- Stadie, William C.,** mechanical shaker and other devices for use with the van Slyke blood gas apparatus, A., ii, 78.
- Stadie, William C.** See also *Donald D. van Slyke*.
- Stadnikoff, George,** chemical adsorption, A., ii, 619.
- Stackel, Walter.** See *Max Trautz*.
- Stackling, Charles,** the radioactivity of the oxides of uranium, A., ii, 106.
- Stärkle, M.** See *W. D. Treadwell*.
- Stäuble, Gustav.** See *Hartwig Franzen*.
- Stahel, E.** See *A. Piccard*.
- Stahl, F.** See *Georg Schroeter*.
- Stalb, K.** See *Alexander Gutbier*.
- Stanek, Vladimir,** polarisation of normal sucrose solution, A., ii, 167.
- Stansell, Lionel William,** obituary notice of, T., 2916.
- Stanton, Ralph E.** See *Philip H. Mitchell*.
- Starcewska, (Mlle) H.** See *Wojciech Swientoslawski*.
- Stark, Johannes,** the structure of carbon chains, A., i, 2.
- Starkey, E. B., and Neil E. Gordon,** thermo-regulator, A., ii, 547.
- Starkey, E. B.** See also *Neil E. Gordon*.
- Starlinger, Wilhelm,** the flocculating capacity of human blood-plasma, A., i, 288.
- agglutination and velocity of sedimentation of corpuscles. II.*, A., i, 289.
- Starr, Henry E.,** human mixed saliva.
- I. Determination of the hydrogen-ion concentration of human mixed saliva.
- II. Variations in the hydrogen-ion concentration, A., i, 1209.
- Stasiak, A.,** blood sugar, A., i, 1208.
- Stateczny, Valentin.** See *Julius Meyer*.
- Staub, H.,** facilitation of intermediary sugar metabolism, A., i, 83.
- Staub, Max.** See *Paul Karrer*.
- Staudinger, Hermann,** ketens.
- XXXVIII. Aliphatic diazo-compounds. XXV. Aliphatic diazo-compounds and ketens, A., i, 238.
- ketens. XXXIX. Aliphatic diazo-compounds. XXVI. Behaviour of ring systems, A., i, 240.
- Staudinger, Hermann, and A. Binkert,** alkali salts of benzil and the benzilic acid transformation, A., i, 1016.
- Staudinger, Hermann, and W. Braunscholtz,** organic compounds of phosphorus. V. Action of carbonylene derivatives on phosphazines, A., i, 70.

- Staudinger, Hermann, and J. Fritsch,** isoprene and caoutchouc. V. The hydrogenation of caoutchouc and its constitution, A., i, 1043.
- Staudinger, Hermann, and Ernst Hauser,** ketens. XXXVII. Ketimine derivatives, A., i, 28.
new organic compounds of phosphorus. IV. Phosphineimines, A., i, 68.
- Staudinger, Hermann, W. Kreis, and W. Schilt,** the addition of hydrogen haloids to isoprene, A., i, 978.
- Staudinger, Hermann, and G. Lüscher,** aliphatic diazo-compounds. XXIV. Organic phosphorus compounds. VI. Preparation and reactions of phosphazines, A., i, 237.
- Staudinger, Hermann, and P. Meyer,** ketens. XLI. Methylenecarbonic acid derivatives [ketenacetals, etc.], A., i, 1015.
- Staudinger, Hermann, O. Muntwyler, and O. Kupfer,** isoprene dibromide, A., i, 979.
- Staudinger, Hermann, and G. Rathsam,** ketens. XL. Ketenacetals, A., i, 1014.
- Stavritsch, K. N. See Émile Cherbuliez.**
- Steabben, Dorothy Beatty. See James Arthur Hewitt.**
- Stearn, Allen E.,** ionic equilibria of strong electrolytes, A., ii, 420.
- Steel, Thos.,** the occurrence of calcium oxalate in the gidgee wattle (*Acacia cambagei*, Baker), A., i, 310.
- Steele, (Miss) Elsie Stewart. See James Colquhoun Irvine.**
- Steele, Lawrence L.,** abietic acid and certain metal abietates, A., i, 739.
- Steele, Victor. See Thomas Martin Lowry.**
- Stehle, Raymond L.,** the gasometric estimation of urea, A., ii, 403.
- Stehle, Raymond L., W. Bourne, and Henry G. Barbour,** effects of ether anaesthesia alone or preceded by morphine on the alkali metabolism of the dog, A., i, 1085.
- Steibelt, Werner. See Richard Willstätter.**
- Steiger, A. L. von,** the graphitic conception of aromatic carbon, A., ii, 616.
- Steinberger, Franz Karl. See Hans Stobbe.**
- Steiner, Joseph. See Kurt Brand.**
- Steiner, Pierre. See Victor Henri.**
- Steinkopf, Wilhelm, and Halvard Augsted-Jensen,** thiophen series. XIV. The condensation of diazoacetic ester with thiophen, A., i, 851.
- Steinkopf, Wilhelm, and Kurt Buchheim,** the action of pyrosulphuryl chloride on toluene, A., i, 122.
- Steinkopf, Wilhelm, Hans Donat, and Paul Jaeger,** organic compounds of arsenic. VIII. Action of cyanogen bromide on tertiary arsines, A., i, 994.
- Steinkopf, Wilhelm, and Julius Herold,** the thiophen series. XIII. The action of acetylene on pyrites, A., i, 850.
- Steinkopf, Wilhelm, and Gustav Schwen,** organic compounds of arsenic. VI. Fission of alkylated or arylated arsine hydroxybromides, A., i, 71.
organic compounds of arsenic. V. Action of cyanogen bromide on phenylated tertiary arsines, A., i, 72.
organic compounds of arsenic. VII. Additive compounds of iodoform and salts of organic bases of trivalent elements, A., i, 118.
- Steinkuhler, W. See Alfred Schoep.**
- Stenger, E. See L. Lewin.**
- Stephen, Florence V. See Ruth E. Conway.**
- Stephen, Henry. See Thomas Hooker Minton.**
- Stephens, H. N.,** the Friedel and Crafts' reaction, bromophthalic anhydrides, benzene, and aluminium chloride, A., i, 141.
- Stephenson, Marjory, and Margaret Dampier Whetham,** fat metabolism of the Timothy grass bacillus, A., i, 500.
- Stephenson, R. E.,** nitrification in acid soils, A., i, 416.
- Stepp, Wilhelm,** detection of acetaldehyde in urine, A., ii, 403.
analytical observations on the true blood-sugar value in normal and pathological individuals, A., ii, 592.
importance of dimethylhydroresorcinol for detection of volatile aldehydes in body fluids; the identification of formaldehyde in urine after administration of hexamethylenetetramine, A., ii, 793.
- Stepp, Wilhelm, and R. Feulgen,** the identification of the aldehyde-like substance in the urine of diabetics as acetaldehyde, A., i, 300.
acetaldehyde as a constituent of normal urine, A., i, 495.
- Stepp, Wilhelm, and Robert Fricke,** a simple and exact method for the direct estimation of acetaldehyde in the presence of acetone, A., ii, 236.
- Stern, Emmi. See Hartwig Fransen.**
- Stern, (Mlle) Lina. See Fred. Battelli.**
- Steuber, Maria. See W. Klein.**
- Steupe, Max. See Alfred Lottermoser.**

- Stendel, Hermann**, histochemistry of spermatozoa. IV. Chemical composition of the spermatozoa of the shad (*Clupea alosa*), A., i, 294.
the nucleic acids of the spleen of cattle, A., i, 297.
- Stendel, Hermann**, and *Sung-Sheng Chou*, the estimation of purine bases in urine, A., ii, 239.
- Stendel, Hermann**, and *R. Freise*, the origin of creatine and creatinine, A., i, 793.
the detection of vernin, A., ii, 535.
- Stendel, Hermann**, and *E. Feiser*, yeast-nucleic acid. II. and III., A., i, 279, 782.
nucleic acid-protein compounds, A., i, 1200.
- Stendel, Hermann**, and *K. Suzuki*, the estimation of uric acid in tissue extracts, A., ii, 538.
- Stevens, F. A.**, and *Randolph West*, peptase, lipase, and invertase of hæmolytic streptococcus, A., i, 903.
- Stevenson, Arnold**, and *Jocelyn Field Thorpe*, the formation of derivatives of tetrahydronaphthalene from γ -phenyl fatty acids. II., T., 1717.
- Stevenson, Arnold**. See also *George Armand Robert Kon*.
- Stevenson, L. D.**, comparative study of the sugar content of the spinal fluid in diseases of the nervous system, A., i, 295.
- Stewart, Alfred Walter**. See *J. K. Marsh*.
- Stewart, C. P.** See *E. J. Morgan*.
- Stewart, Charles R.** See *Leon A. Congdon*.
- Stewart, (Miss) Jessie**, aromatic sulphonyl chlorides, T., 2555.
- Stewart, John**, some relations of arsenic to plant growth. I., A., i, 1221.
- Stewart, John**, and *Edwin S. Smith*, some relations of arsenic to plant growth. II., A., i, 1222.
- Stewart, Lachlan Macquarie**, and *William Wardlaw*, the oxidising and reducing properties of sulphur dioxide. I. Mercury chlorides, T., 1481.
- Steyer, Hermann**. See *Carl Paal*.
- Stieglitz, Julius**, the electron theory of valency as applied to organic compounds, A., ii, 560, 705.
- Stieglitz, Julius**, and *Ralph L. Brown*, the molecular rearrangement of *s*-bistriphenylmethylhydrazine, A., i, 778.
- Stiles, Walter**, penetration of electrolytes into gels. IV. Diffusion of sulphates, A., ii, 125.
- Stiles, Walter**, and *Gilbert Smithson Adair*, penetration of electrolytes into gels. III. Influence of the concentration of the gel on the coefficient of diffusion of sodium chloride, A., ii, 125.
- Stix, Walter**. See *Walter Fuchs*.
- Stobbe, Hans**, and *Franz Karl Steinberger*, photo-reactions of the *trans*- and *cis*-cinnamic acids, A., i, 1018.
- Stock, Alfred**, carbon and its neighbours in the periodic system, A., ii, 564.
- Stock, Heinrich**. See *Robert Schwarz*.
- Stockings, W. E.** See *William Arthur Bone*.
- Stoddard, R. B.** See *W. G. Ungerer*.
- Stöckigt, Fritz**. See *Emil Heuser*.
- Stoermer, Richard**, and *F. Bachér*, configuration of the truxinic and truxillic acids. VI., A., i, 830.
- Stoermer, Richard**, and *H. Oehlert*, stereoisomeric derivatives of stilbene, A., i, 647.
- Stoermer, Richard**, and *E. Robert*, behaviour of crotonic acid in ultra-violet light. II., A., i, 519.
- Stoklasa, Julius**, influence of selenium and radium on the germination of seeds, A., i, 618.
influence of selenium on plant evolution in the presence or absence of radioactivity, A., i, 614.
- Stoklasa, Julius** [with *J. Chmelař, V. Janský, P. Křička, J. Pěnkava*, and *J. Zelenka*], action of selenium on the metabolism of plants in presence of the radioactivity of air and of soil, A., i, 974.
- Stoklasa, Julius** [with *J. Šebor, F. Týmich*, and *J. Cwacha*], the resorption of aluminium ions by the roots of plants, A., i, 502.
- Stoll, A.** See *Karl Spiro*.
- Stoll, P.** See *Paul Scherrer*.
- Stollé, Robert**, addition of hydrazonic acid to derivatives of carbodi-imide, A., i, 689.
preparation of *N*-substituted 3-dihalognoxindoles, A., i, 762.
- Stollé, Robert**, and *A. Netz*, the reaction of benzylidenebenzhydrazide chloride and dibenzhydrazide chloride with sodium azide, A., i, 690.
- Stollenwerk, Wilhelm**. See *Wilhelm Biltz*.
- Stone, Hosmer W.** See *Victor Lenher*.
- Storm, Einar**. See *Heinrich Goldschmidt*.
- Stosius, Karl**, the condensation of methyl β -naphthol-3-carboxylate with methyl *n*-opianate, A., i, 746.

- Stott, Verney**, pipettes, A., ii, 388.
- Stratton, (Mrs) K.**, and **James Riddick Partington**, latent heats of fusion. I. Benzophenone, phenol, and sulphur, A., ii, 258.
- Straus, Fritz**, and **Anadaus Dützmann**, dibenzylideneacetone [distyryl ketone] and triphenylmethane. X. Ionogenically linked halogen atoms, A., i, 148.
- Strauss, David**. See **Elektrochemische Werke G. m. b. H.**
- Strauss, Hermann**, and **Wolfgang Schubardt**, cholesterol content of blood-serum, A., i, 1209.
- Strecker, Wilhelm**, and **Max Baltes**, action of ozone on aliphatic and aromatic substitution products of ammonia, A., i, 14.
- Strecker, Wilhelm**, and **Ernst Kannappel**, estimation of boric acid, A., ii, 784.
- Streel, (Mlle) du Vivier de**. See **E. Faure-Fremiet**.
- Strickler, Alvin**, and **Joseph Howard Mathews**, electric endosmose, A., ii, 688.
- Strisower, Rudolph**, total nitrogen and residual nitrogen content of oedematous fluids, A., i, 964.
- Stroh, Rudolf**. See **Rudolf Friedrich Weinland**.
- Strohmenger, Ludwig**. See **Stefan Goldschmidt**.
- Struwe, F.** See **Willy Marckwald**.
- Strzelba, Hubert**. See **Robert Kremann**.
- Stuber, B.**, and **A. Funck**, blood clotting. IV., A., i, 393.
- Stuckert, Guillermo V.**, considerations on the solubility of the phosphatides, A., i, 1089.
- Stübel, H.**, microchemical recognition of urea in the kidney by means of xanthyrol, A., i, 397.
- Stuhlman, Otto, jun.**, the photoelectric long wave-length limit of platinum and silver, A., ii, 809.
- Stumper, Robert**, new observations on the venom of ants, A., i, 400.
- Sturzenegger, P.** See **W. D. Treadwell**.
- Subrahmanyam, G.** See **A. L. Narayan**.
- Sudborough, John Joseph**, and **D. D. Karvé**, studies in acidolysis. I. The equilibrium between acetic acid, trichloroacetic acid, and their ethyl esters. II. The equilibrium between acetic acid, trichloroacetic acid, and their methyl esters, A., ii, 749.
- Sudborough, John Joseph**, and **R. C. Shah**, influence of s.-trinitrobenzene on reactions in which arylamines are involved, A., ii, 757.
- Sugden, Samuel**, the determination of surface tension from the maximum pressure in bubbles, T., 858.
- Sugiura, Y.** See **H. Nagaoka**.
- Sullivan, F. W., jun.** See **Moses Gomborg**.
- Sundberg, Thure**, sensitiveness of certain tests for hydrocyanic acid, A., ii, 404.
- Supplee, G. C.**, and **B. Bellis**, citric acid content of milk and milk products, A., i, 197.
- Suschnig, Eugen**, rubidium (caesium) silver gold bromides, A., ii, 514.
- Suszko, J.** See **Karol Dziewoński**.
- Suter, Richard**. See **Franz Fichter**.
- Suydam, J. R., jun.** See **Joaquin Enrique Zanetti**.
- Suzuki, K.** See **Hermann Stendel**.
- Suzuki, Umetarō**, and **Yoshihiko Matsuyama**, succinic acid, A., i, 716.
- Suzuki, Yoshitarō**, and **Sankyō Kabushiki Kaisha**, preparation of aromatic esters in the presence of an organic base, A., i, 543.
- Svanberg, Olof, K. Sjöberg**, and **G. Zimmerlund**, the iodometric micro-estimation of phosphoric acid, and of phosphorus in organic compounds, A., ii, 867.
- Svanberg, Olof**. See also **Hans von Euler**, and **Karl Freudenberg**.
- Svanoe**. See **Georg Schroeter**.
- Svensson, John**. See **Alfred Bjure**.
- Swan, Enoch**. See **Fred Fairbrother**.
- Swan, W. O.** See **Graham Edgar**.
- Swarts, Frédéric**, some aliphatic fluorides, A., i, 101.
- Sweet, S. S.** See **Samuel Edward Sheppard**.
- Swientoslawski, Wojciech**, Eisenlohr's refractometric constants, A., ii, 173.
- physical properties of vapour-liquid systems. I. and II., A., ii, 192.
- divergences of the value of the mechanical equivalent of heat, A., ii, 194.
- new data in the thermochemistry of organic substances, A., ii, 195.
- proposal for the fixing of a unit of measurement in thermochemistry, A., ii, 548.
- Swientoslawski, Wojciech**, and **(Mlle) H. Starczewska**, the heat of combustion of benzoic acid, cane-sugar, and naphthalene, A., ii, 616.
- Swisher, Margaret**. See **John Bernard Ekeley**.
- Swoboda, Frederick K.**, nitrogen nutrition of yeast, A., i, 795.

- Syniewski, Wiktor**, oxidation of amylo-dextrin, A., i, 1120.
Syrkin, Ja. K., equilibrium in heterogeneous systems, A., ii, 699.
 theory of solvates, A., ii, 823.
Syrkin, Ja. K. See also *P. P. Budnikov*.
Szamá, Nikolaus. See *Fritz von Konek*.
Szányi, W. See *Fritz Versár*.
Szappanyos, B. See *Z. Ernst*.
Szillard, Béla, the direct estimation of small quantities of radium by the penetrating rays, A., ii, 586.

T.

- Tacke, Ida.** See *David Holdé*.
Tadokoro, Tetsutaro, ultra-violet spectroscopical studies on the antagonistic action of salts in organic colloidal solution, A., ii, 204.
 spectrochemical reaction of methyl-furfuraldehyde and hydroxymethyl-furfuraldehyde phloroglucides, A., ii, 236.
Tafel, Alan, thermal expansion of gelatin gels, T., 1971.
Taggart, Grace C. See *Frank A. Csonka*.
Tahara, Yoshihide. See *Riko Majima*.
Talstra, Sophia A., animal calorimetry. XIX. The influence of acids on the carbon dioxide-combining power of the blood plasma, A., i, 288.
Takahashi, Katsumi, fatty matters in the sea urchin, A., i, 609.
 nutritive value of fats and lipoids. I, A., i, 699.
 fatty matter in "herring roe," A., i, 700.
Takahashi, Torizô. See *Heisaburô Kondô*.
Takahashi, Yutaka, the origin of band spectra, A., ii, 3.
Takashima, Norikazu. See *Hidekichi Yanagisawa*.
Takata, Maki, gastric juice. III. Juice from the empty stomach, A., i, 698.
 estimation of pepsin, A., ii, 886.
Takayama, Gitaro. See *Riko Majima*.
Takeda, Jirô, and *Sajirô Kuroda*, new method for the preparation of alkalines. II, A., i, 272.
Takei, Takeo, the analysis of a volume curve of blood-corpuscles in hypertonic solutions, which renders possible the simultaneous differentiation of osmotic and colloido-chemical changes in volume, A., i, 289.
Tamada, Kensuke. See *Riko Majima*.
Tamaru, Setsuro. See *Theodore William Richards*.
Tamburello, A. See *Anne Wilhelm van der Haar*.
Tammann, Gustav, the chemical behaviour of crystallised binary compounds with one component nobler than hydrogen, A., ii, 63.
 the reaction limit of chemical agents on copper-gold alloys and their galvanic tension, A., ii, 75.
 electro-chemical behaviour of metallic conducting compounds, A., ii, 255.
 the substance between the crystallites of metallic substances, A., ii, 502.
 the rate of formation of some nitrides, A., ii, 852.
Tammann, Gustav, and *W. Jander*, the behaviour of two metals towards one another when dissolved in mercury, A., ii, 825.
Tammann, Gustav, and *W. Köster*, the velocity of the action of oxygen, hydrogen sulphide, and the halogens on metals, A., ii, 831.
Tammann, Gustav, and *K. Schönert*, the diffusion of carbon in metals and in mixed crystals of iron, A., ii, 772.
Tammann, Gustav, and *E. Vaders*, the electrochemical behaviour of alloys of manganese with copper, nickel, cobalt, and iron, A., ii, 380.
Tammann, Gustav. See also *Rudolf Vogel*.
Tanaka, Shunsuke. See *Usaburo Yoshida*.
Tanberg, Arthur Percival, and *Herbert Winkel (E. J. du Pont de Nemours & Co.)*, preparation of symmetrical tetra-substituted carbamides, A., i, 1009.
Tanch, J. W. See *H. S. Uhler*.
Tanner, W. F. See *Joseph Goldberger*.
Tanret, Georges, the chemical composition of the ergot of diss and of the ergot of oats, A., i, 504.
Tapernoux, A. See *Ch. Porcher*.
Tartar, Herman V., and *Zalia Jencks Gailey*, rôle of the hydrogen-ion concentration in the precipitation of colloids, A., ii, 829.
Tartar, Herman V., and *Harmon E. Keyes*, electrical conductivity of zinc sulphate solutions in the presence of sulphuric acid, A., ii, 186.
 measurement of overvoltage, A., ii, 343.
Tarugi, N., toxicology of arsenic, A., ii, 84.
 separation of phosphoric acid in qualitative analysis, A., ii, 784.

- Tashiro, Shiro.** See *Olric P. Lee*.
- Tatsuno, Hisajirō.** See *Yutaka Kinugasa*.
- Taylor, F. A.** See *Phœbus A. Levene*.
- Taylor, Frank E.** See *Aldo Castellani*.
- Taylor, Guy B., and George Augustus Hulett,** catalytic decomposition of certain oxides, A., ii, 646.
- Taylor, Guy B.** See also *William C. Sproesser*.
- Taylor, Henry Austin, and William Cudmore McCullagh Lewis,** studies in catalysis. XV. Absorption spectra of triethylsulphonium bromide in various solvents, in the short infra-red region, T., 665.
- Taylor, Hugh Stott, and William Theodore Anderson, jun.,** heat of formation of silver iodide, A., ii, 121.
- Taylor, Hugh Stott, and Harvey A. Neville,** catalysis in the interaction of carbon with steam and with carbon dioxide, A., ii, 143.
- Taylor, Hugh Stott.** See also *Robert N. Reese*.
- Taylor, John,** the interaction of aldehydes or ketones and thiocarbamides in the presence of acids, T., 2269.
- Taylor, L. S.** See *P. P. Cioffi*.
- Taylor, (Miss) Millicent.** See *Orsel Joyce Flecker, and James William McBain*.
- Taylor, M. C., and C. A. Gammal,** estimation of free chlorine and hypochlorous acid in concentrated salt solutions, A., ii, 581.
- Taylor, M. C., W. A. Gersdorff, and E. J. Tovrea,** electrolytic reduction of chromic chloride to the bivalent state, A., ii, 382.
- Taylor, Thomas Weston Johns,** radiation and chemical action, A., ii, 336.
- Taylor, Thomas Weston Johns.** See also *Nevil Vincent Sidgwick*.
- Taylor, William,** the non-protein nitrogen in goat's milk, A., i, 1213.
- Telecsky, J.** See *Ernst Winterstein*.
- Tennenbaum, M.** See *C. van Eweyk*.
- Terrill, Edwin H.,** the colorimetric estimation of hæmoglobin with especial reference to the production of stable standards, A., ii, 799.
- Terroine, Émile F., and René Wurmser,** the energy yield in the growth of *Aspergillus niger*, A., i, 706.
- Terroine, Émile F., René Wurmser, and J. Montané,** the influence of the constitution of nutritive media on the composition of *Aspergillus niger*, A., i, 1220.
- Tervaert, D. G. Cohen,** some new observations on the relation between the true sugar content of urine and the sugar content of blood, A., i, 1215.
- new methods of blood sugar estimation; estimation of the true sugar content of urine, A., ii, 166.
- Teschendorf, Werner,** the action of organic kations on the vascular system and its modification by inorganic ions, A., i, 90.
- physiology and pharmacology of the leech, *Hirudo medicinalis*, A., i, 970.
- Tetralin, G. m. b. H.,** preparation of ar-tetrahydronaphthylthiolacetic acids, A., i, 340.
- preparation of hydrogenated anthraquinones, A., i, 1039.
- Tetralin, G. m. b. H.** See also *Georg Schroeter*.
- Tezner, Otto, and Max Turolt,** the action of potassium and calcium ions on the surviving human stomach, A., i, 396.
- Thannhauser, Siegfried J., and E. Andersen,** estimation of bilirubin in human serum; the Ehrlich-Proschner reaction, A., ii, 671.
- Theberath, H.** See *Adolf Sieverts*.
- Theis, R. C., and Stanley Rossiter Benedict,** distribution of uric acid in the blood, A., i, 82.
- Thelen, Richard.** See *Ernst Philipp*.
- Thiel, Alfred,** mechanism of the colour change of some phthaleins, A., i, 455.
- disglomeration and formation of the autogenous lead tree, A., i, 63.
- the state of carbonic acid in aqueous solution, A., ii, 374.
- the thermochemistry of carbon compounds, A., ii, 819.
- Thiel, Alfred, and Fr. Müller,** chloro- and bromo-derivatives of phenolphthalein, A., i, 659.
- Thiel, E. van,** influence of the catalyst on the thermodynamic quantities regulating the velocity of a reaction, A., ii, 754.
- Thielepape, Ernst,** extension of the Kishner-Wolff method of reduction. I., A., i, 271.
- the quinoline series. I. Synthesis of 4-substituted quinolines and of quinoline-4-carboxylic acids, A., i, 271.
- Thielepape, Ernst, and Otto Spreckelsen,** extension of the Kishner-Wolff method of reduction, II., A., i, 1191.
- Thieme, Hans,** tryptaphavin [3:6-diamino-1-methylacridinium chloride], A., i, 59.

- Thierfelder, Hans**, constitution of glutamine, A., i, 326.
- Thiery**, application of Folin and Denis's phosphotungstic reaction to the estimation of uric acid in urine, A., ii, 238.
- Thiery**. See also **Rouzaud**.
- Thivolle, L.** See **G. Fontès**.
- Thole, Ferdinand Bernard**. See **Albert Ernest Dunstan**.
- Tholin, Th.**, the thermostability of the co-enzyme and its separation from vitamin-B from yeast, A., i, 305.
- Thomann, H.** See **Paul Karrer**.
- Thomas, Arthur W.**, and **Margaret W. Kelly**, the isoelectric point of collagen, A., i, 387.
- Thomas, Arthur W.** See also **Francisco A. Quisumbing**.
- Thomas, Clifford Dane**. See **Oscar Lisle Brady**.
- Thomas, F.** See **Douglas Frank Twiss**.
- Thomas, Fritz**. See **Otto Ruff**.
- Thomas, John, Arthur Hugh Davies**, and **Scottish Dyes, Ltd.**, production of dyestuff intermediates [aminoanthraquinones], A., i, 260.
- Thomas, K.**, decomposition of proteins of organs, A., i, 790.
- Thomas, Pierre**, and **Georges Carpentier**, a very sensitive reagent for copper; the Kastle-Meyer reagent, A., ii, 86.
- Thomas, Richard**, vapour pressures of dilute alcohol solutions, A., ii, 321.
- Thomas, Ruth**. See **Forris Jewell Moore**.
- Thomas, V.**, mixed organometallic compound of aluminium, A., i, 330.
- Thomas, Walter**. See **Claude W. Bourlet**.
- Thomas, William**. See **Eric Keightley Rideal**.
- Thompson, F. P.** See **Edmund Knecht**.
- Thompson, Gartha**. See **William Bayley Parker**.
- Thompson, Hugh Vernon**. See **Francis Herbert Clews**.
- Thompson, Ruth**. See **Frank Burnett Dains**.
- Thompson, T. J.** See **Fred W. Upson**.
- Thomson, G. P.**, anode rays of glucinum, A., ii, 106.
- Thomson, (Sir) Joseph John**, application of the electron theory of chemistry to solids, A., ii, 355.
the analysis by positive rays of the heavier constituents of the atmosphere; of the gases in a vessel in which radium chloride had been stored for thirteen years, and of gases given off by deflagrated metals, A., ii, 565.
an electron theory of solids, A., ii, 745.
- Thorne, Percy Cyril Lesley**, colloidal solutions of carbon in water, A., ii, 695.
- Thornton, William M., jun.**, tervalent titanium. II. Estimation of copper and iron in the presence of each other, A., ii, 528.
- Thorpe, Jocelyn Field**. See **Oscar Becker**.
- Stanley Francis Birch, Juan Pedigo Charles Chandrasena, Shankar Shridhar Deshapande, Frank Dickens, Ernest Harold Farmer, Biraj Mohan Gupta, Christopher Kolk Ingold, George Armand Robert Kon, and Arnold Stevenson**.
- Thorvaldson, Thorbergur**. See **Theodore William Richards**.
- Thunberg, Torsten**, is there a genetic relation between the oxygen inhaled and the oxygen of the exhaled carbon dioxide? A., i, 889.
- Thyssen, H.** See **G. Batta**.
- Tian, A.**, slow hydrolysis of salts, A., ii, 362.
- Tichomirov, P.** See **G. Fovarnin**.
- Tideswell, Frederick Vincent**, and **Richard Vernon Wheeler**, on dopplertite; studies in the composition of coal, T., 2345.
- Tiebackx, F. W.**, the colloids arabic acid and arabic acid plus gelatin, A., i, 638.
is the gelatin-gum arabic coagulation a chemical or a colloid-chemical process? A., ii, 698.
- Tiede, Erich**, and **Friedrich Richter**, inorganic luminescence phenomena. IV. Preparation of pure magnesium sulphide and its phosphorescence. II. Phosphorescent magnesium sulphides, A., ii, 215.
preparation of phosphorescent magnesium sulphide, A., ii, 247.
- Tiede, Erich**, and **Arthur Schleeds**, the rare-earth magnesium sulphide phosphors, A., ii, 769.
- Tiede, Erich**, and **Peter Wulff**, hydrates of boron trioxide as a constituent of systems capable of strong phosphorescence and containing organic compounds, A., ii, 245.
- Tiede, Erich, Peter Wulff**, and **A. Ragoss**, hydrated boric acid as the basis of systems capable of [showing] a high phosphorescence, A., ii, 8.
- Tiemann, Fr.**, new theory of the constitution of hydroxides, particularly those of the basic metallic oxides, A., ii, 53.
- Tiffeneau, Marc**, and **M. Porcher**, semipinacolic transposition in the benzylcyclohexene series; migration of the benzene radicle, A., i, 537.

- Tiffeneau, Marc.** See also *Paul Carnot*, and *Alex. Orékhoff*.
- Tiger, R., and H. Simonnet**, variations in the amylolytic activity of the pancreas and liver in avian polyneuritis, A., i, 195.
- Tigges, H.** See *Paul Jacobsen*.
- Tilley, C. E.**, density, refractivity, and composition relations of some natural glasses, A., ii, 220.
- Timmermans, Jean**, theory of concentrated solutions; separation of solids from organic mixtures, A., ii, 25.
the mutual solubility of liquids at high pressures, A., ii, 482.
- Timmermans, Jean, (Mlle) H. van der Horst**, and *Heike Kamerlingh Onnes*, the freezing points of pure organic liquids as thermometric constants at temperatures below 0°, A., ii, 258.
- Timmis, Laurence Barnett.** See *Frank Lee Pyman*.
- Tingey, Harold Calvert**, and *Cyril Norman Hinshelwood*, the catalytic decomposition of formic acid on surfaces of platinum and silver, T., 1668.
- Tingle, Alfred**, alleged adsorption of alumina from aluminium sulphate solutions by cellulose, A., i, 434.
- Tischenko, Georg von.** See *Hermann Plauson*.
- Tisdall, Frederick F.**, influence of the sodium-ion in the production of tetany, A., i, 1216.
a rapid colorimetric method for the estimation of the inorganic phosphorus in small amounts of serum, A., ii, 392.
- Tisdall, Frederick F.** See also *Benjamin Kramer*.
- Titley, Alan Francis.** See *William Henry Perkin, jun.*
- Titov, V. S.**, partition of chlorine between water and a gaseous phase, A., ii, 269.
- Tobler, Richard.** See *Hans Eduard Fierz*.
- Tocher, J. F.**, the citric solubility of mineral phosphates, A., ii, 525.
- Tochinal, Yoshihiko**, the food relations of *Eusarium lini*, A., i, 207.
- Toda, Shigeru**, equilibrium in the system potassium chlorate, potassium nitrate, and water at 25°, A., ii, 748.
oxychlorides of mercury; equilibrium in the system mercuric chloride, yellow mercuric oxide, and water at 35°, A., ii, 769.
- Tokuoka, E., and K. Ogasawara**, quantity of combined carbonic acid in cerebrospinal fluid, A., i, 895.
- Tokyo Industrial Laboratory.** See *Shuichirô Ochi*.
- Tolman, Richard C.**, entropy of electron gas, A., ii, 18.
thermal ionisation of metallic vapours, A., ii, 18.
relation between statistical mechanics and thermodynamics, A., ii, 257.
thermodynamic treatment of the possible formation of helium from hydrogen, A., ii, 738.
- Tomaschek, Rudolf**, boric acid phosphors, A., ii, 763.
- Tomíček, Oldřich**, some derivatives of indolinone, A., i, 679.
- Tomita, Masaji**, the behaviour of inactive malic acid in the organisms of the dog and rabbit, A., i, 801.
the decomposition of *D*-galactose according to the second mode of fermentation, A., i, 307.
phosphatases. I. and II., A., i, 960.
the influence of thyroxin on alcoholic fermentation, A., i, 972.
- Tomula, E. S.**, antimonite acid and the use of sodium antimonate in analysis, A., ii, 74.
- Topley, B.** See *Cyril Norman Hinshelwood*.
- Toporescu, E.**, the preparation of sodium hydrogen carbonate, A., ii, 375, 642.
- Torelli, G.** See *Gino Scagliarini*.
- Tosterud, Martin.** See *Victor Lenher*.
- Tourrou, R.** See *Georges Denigès*.
- Tovrea, E. J.** See *M. C. Taylor*.
- Tower, Olin Freeman**, and *Martha C. Cooke*, preparation of colloidal solutions of nickel and cobalt hydroxides and some other compounds of these metals, A., ii, 853.
- Town, George G.** See *Victor Lenher*.
- Townsend, Eric B.** See *Duncan A. MacInnes*.
- Townsend, John S.**, and *V. A. Bailey*, motion of electrons in gases, A., ii, 43.
motion of electrons in atoms, A., ii, 277.
the abnormally long free paths of electrons in argon, A., ii, 494.
motions of electrons in argon and hydrogen, A., ii, 836.
- Toyama, Yoshiyuki**, unsaponifiable constituents (higher alcohols) of shark and rayfish liver oils. II., A., i, 895.
derivatives of behenic and erucic acids, A., i, 1111.
composition of the fatty acids of raje oil, A., i, 1113.
- Toyama, Yoshiyuki.** See also *Mitsumaru Tsujimoto*.

- Traetta-Mosca, F.**, and **M. Preti**, action of *Aspergillus glaucus* on glycerol, A., i, 91.
- Trampler, (Mlle) A.** See **Emil Briner**.
- Traube, Isidor, P_H** again, A., i, 204.
a new viscostalagmometer for the estimation of surface tension and viscosity of liquids of very different limpidity, A., ii, 122.
- Traube, Isidor**, and **Paul Klein**, the theory of narcosis, A., i, 198.
submicrons visible in the Tyndall cone to the naked eye, A., ii, 201.
- Traube, Isidor**, and **R. Somogyi**, theory of disinfection, A., i, 204.
- Traube, Wilhelm**, alkaline copper hydroxide solutions and copper oxide-ammine-cellulose solutions. I. and II., A., i, 115, 718.
- Trautenberg, Heinrich Rausch von**, and **K. Philipp**, passage of α -rays through materials, A., ii, 12.
- Traun, H. Otto, Forschungslaboratorium**, preparation of diolefines and polymerisation products thereof, A., i, 514.
preparation of diolefines and derivatives thereof, A., i, 515.
preparation of vinyl compounds and polymerisation products thereof, A., i, 516.
preparation of vinyl haloids, A., i, 517.
preparation of vinyl sulphuric acid and homologues thereof, A., i, 517.
preparation of formaldehyde and methyl alcohol, A., i, 522.
preparation of hexamethylenetetramine and formaldehyde, A., i, 528.
- Traun, Max.** See **Hugo Weil**.
- Trautz, Max**, and **Walter Seidel**, the luminescence of decomposing orone, A., ii, 730.
- Trautz, Max**, and **Walter Stäckel**, the decomposition of chlorine into atoms, A., ii, 760.
- Trautz, Max**, and **Karl Winkler**, preparation of propylene in a pure condition, A., i, 909.
preparation of cyclopropane in a pure condition, A., i, 926.
problems of organic chemistry. I. The velocity of ring fission in gases; isomerisation of cyclopropane, A., i, 926.
- Traxler, Ralph N.** See **Frank E. E. Germann**.
- Treadwell, W. D.**, formulæ for the solubility of certain salts in aqueous ethyl alcohol and water, A., ii, 31.
- Treadwell, W. D.** [with **M. Blumenthal**, **A. Bossi**, **Max Dreifuss**, **M. Hoofst**, and **M. Stärkle**], reductions with cadmium in volumetric analysis. II., A., ii, 788.
- Treadwell, W. D.** [with **R. F. Edelmann**, **A. Freuler**, **P. Hristie**, **L. Egger**, **P. Sturzenegger**, and **A. Weber**], reductions with cadmium and lead in volumetric analysis. III., A., ii, 780.
- Treadwell, W. D.** [with **Ch. Mussler**], the solubility of arsenic trichloride in concentrated hydrochloric acid at 100°, A., ii, 763.
- Treadwell, W. D.**, and **D. Chervet**, the influence of the alkalis on the titration of some metals with ferrocyanide, A., ii, 786.
- Trewendt, Gert.** See **Arthur Rosenheim**.
- Trifonov, Iv.**, properties and structure of pernitric acid, A., ii, 845.
pernitric acid as an analytical reagent, A., ii, 883.
- Trimble, H. M.**, solubility of potassium permanganate in solutions of potassium sulphate and sodium sulphate, A., ii, 374.
- Tröger, Julius**, and **Fr. Bolte**, synthesis of α -benzopyrone derivatives and the rupture of the pyrone ring in these compounds, A., i, 267.
- Tröger, Julius**, and **W. Menzel**, synthesis of 8-arylsulphonylquinolines containing a side chain in the 2-position, A., i, 269.
- Tröger, Julius**, and **K. Schwarzenberg**, a new base from the residues of the hydrolytic products of cocaine, isomeric with tropine and ψ -tropine, A., i, 167.
- Troise, Antonio**, a cause of error in the application of the colour test for acetone, A., ii, 595.
- Troitzki.** See **Alexander Kiesel**.
- Tromp, S. T. J.**, specific characters of hydrolytic decomposition, A., ii, 476.
- Tropsch, Hans**, and **A. Kreutzer**, the acids of montan wax, A., i, 317, 804.
- Trümpler, G.**, electromotive behaviour of metallic compounds with electron conductivity, A., ii, 110.
- Tsang, Chuk Yec.** See **Duane T. Englis**.
- Tschelnitz, Erich.** See **Ernst Späth**.
- Tschenschner, F.** See **Fritz Arndt**.
- Tschirsch, E.**, [estimation of] vanillin, A., ii, 403.
- Tschitschibabin, Alexei E.**, and **J. S. Bylinkin**, the products of the benzylation of 2-aminopyridine, A., i, 573.

- Tschudy, Edward A.**, effect of variation in weight of the riders and plummets of the Westphal balance upon the accuracy of specific gravity determinations, A., ii, 820.
- Tschugaev, Leo Alexandrovitch**, and **N. K. Pscheniyn**, complex platinum compounds. III. Molecular rearrangements observed with complex platinum compounds, A., ii, 856.
- Tschugaev, Leo Alexandrovitch**, and **N. A. Vladimirov**, variation of the electrical conductivity among the ammoniacal derivatives of platinous nitrite, A., ii, 814.
- Tsuji moto, Mitsumaru**, clupanodonic acid, A., i, 917.
- Tsuji moto, Mitsumaru**, and **Yoshiyuki Toyama**, higher alcohols in the unsaponifiable matter from shark and ray-fish liver oils, A., i, 297.
- Tsukamoto, Takeo**. See **Yasukiko Asahina**.
- Tsukiye, Sogen**, preparation and properties of vitamin-B, A., i, 974.
- Tubandt, Carl**, and **Gerhard Eschenhagen**, behaviour of the photo-haloids in a direct electric current, A., ii, 346.
- Tucker, Stanley Horwood**. See **Hemendra Kumar Sen-Gupta**.
- Turban, Karl**, comparative investigations on the blood sugar content of the arterial and venous vascular systems, A., i, 482.
- Turina, Dozo**, action of selenium, sulphur, and tellurium salts on plants, A., i, 707.
- Turner, Eustace Ebenezer**, the action of cupric chloride on organometallic derivatives of magnesium, A., i, 23.
- Turner, Eustace Ebenezer**, and **F. H. H. Wilson**, decomposition of methyl oxalate by acetic acid, A., i, 916.
- Turner, Eustace Ebenezer**. See also **George Joseph Burrows**.
- Turner, Sydney Leonard**. See **Nevil Vincent Sidgwick**.
- Turner, William Ernest Stephen**, critical examination of methods commonly used in determining the durability of glass, A., ii, 719.
- Turner, William Ernest Stephen**, and **T. E. Wilson**, action of various analytical reagents on chemical glassware, A., ii, 501.
- Turner, William Ernest Stephen**. See also **A. Cousen**, and **Violet Dimbleby**.
- Turott, Max**. See **Otto Texner**.
- Tussenbroek, M. J. van**. See **Hein Israel Waterman**.
- Tutton, Alfred Edwin Howard**, monoclinic double selenates of the cadmium group, A., ii, 502.
- monoclinic double selenates of the manganese group, A., ii, 505.
- Twiss, Douglas Frank, Sidney Albert Brazier**, and **F. Thomas**, the dithiocarbamate accelerators of vulcanisation, A., i, 460.
- Týmich, F.** See **Julius Stoklasa**.

U.

- Uemura, Taku**. See **Yaji Shibata**.
- Ueno, Seichi**, promoters of the hydrogenation of oils. I., A., i, 983.
- Uhler, H. S.**, and **J. W. Tanch**, the arc spectra of gallium and indium, A., ii, 677.
- Ullmann, Alfred**, tyramine (*p*-hydroxy-phenylethylamine) as the active principle of the drug *Semina cardui* Mariæ, A., i, 616.
- Ullmann, Fritz**. See **Anne Marie von dem Knesebeck**.
- Ullpts, Reinhold**. See **Max Bergmann**.
- Ultée, A. J.**, identity of xanthosterol with lupeol, A., i, 826.
- stearic acid in the latex of *Ficus fulva*, Reinw., A., i, 1100.
- Underhill, Frank Pell**. See **Erwin G. Gross**.
- Ungerer, Ernst**, determination of the size of particles; attempts to explain the formation of layers in clay turbidities and their use in soil analysis, A., ii, 96.
- Ungerer, W. G.**, and **R. B. Stoddard**, odour value analysis, A., i, 894.
- Upson, Fred W.**, and **Lila Sands**, the decomposition of amines in the vapour stage, A., i, 1121.
- Upson, Fred W.**, and **T. J. Thompson**, the preparation and properties of several phenylalkylsuccinic acids, A., i, 343.
- Urasov, G.** See **Nicolai S. Kurnakov**.
- Urbain, Georges**, the atomic numbers of neo-ytterbium, lutecium, and celtium, A., ii, 505.
- Urbain, Georges**. See also **Ch. Boulanger**, and **Pierre Urbain**.
- Urbain, Pierre**, and **Georges Urbain**, extraction and purification of scandium from thorveitite from Madagascar, A., ii, 504.
- Urbasch, Stefan**, the volumetric and gravimetric estimation of zinc, A., ii, 317.

- Usherwood, (Miss) Edith Hilda**, tautomerism of dyads. I. Detection of tautomeric equilibria in hydrocyanic acid, T., 1604.
- Usherwood, (Miss) Edith Hilda**. See also *Christopher Kelk Ingold*.
- Utz, Franz**, [detection of bile pigments in] gastric juice, A., ii, 799.
analysis of blood, A., ii, 800.
- Uyeda, Yoshisuke**. See *Arthur George Perkin*.
- Uytenbogaart, jun.**, electrically heated apparatus for the determination of melting points, A., ii, 471.
- V.
- Vacher, F.** See *Jh. Martinet*.
- Vaders, E.** See *Gustav Tammann*.
- Vaillant, E.**, rapid estimation of uric acid in urine, A., ii, 668.
- Vaillant, Pierre**, variations in the conductivity of solid electrolytes, A., ii, 418.
- Valeras, F. Pablo**, the Neuburger nuclear model, A., ii, 702.
- Vallance, Reece Henry**, the solubility of potassium ferrocyanide, A., i, 724.
- Vallance, Reece Henry**. See also *John Albert Newton Friend*.
- Vallée, C.**, and *M. Polonowski*, micro-chemical estimation of nitrogen, A., ii, 312.
micro-estimation of albumin, A., ii, 406.
- Vallot, J.**, the measurement of the influence of heat and light on the activity of reduction of animal tissues, and applications to heliotherapy, A., i, 87.
- Valteich, H. W.**, and *C. C. Glover*, estimation of pepsin, A., ii, 406.
- Vanderveelde, Albert Jacques Joseph**, combined action of enzymes, A., i, 958.
- Vangindertaelen, A.** See *Henri Wuyts*.
- Varenkamp, Otto**. See *Bruno Emmert*.
- Vaubel, Wilhelm**, gravimetric estimation of nickel as nickel dioxide, A., ii, 875.
- Vavilov, S. I.**, dependence of the intensity of the fluorescence of dyes on the wave-length of the exciting light, A., ii, 181.
- Vavon, Gustave**, and *A. L. Berton*, the borneol obtained from the magnesium compound of pinene hydrochloride, A., i, 943.
- Vavon, Gustave**, and *A. Husson*, catalysis by platinum black, A., ii, 631.
- Vecchiotti, Luigi**, action of mercuric acetate on certain aminoazo-compounds, A., i, 478.
- Vecchiotti, Luigi**. See also *Riccardo Ciusa*.
- Vegard, L.**, the structure of the isomorphous group, $\text{Pb}(\text{NO}_3)_2$, $\text{Ba}(\text{NO}_3)_2$, $\text{Sr}(\text{NO}_3)_2$, $\text{Ca}(\text{NO}_3)_2$, A., ii, 503.
- Veidt, Maximilian**. See *Alfred Schaaerschmidt*.
- Velisek, J.** See *Jiri Baborovsky*.
- Venable, Francis P.**, and *D. H. Jackson*, chlorination by mixed carbon monoxide and chlorine, A., ii, 73.
- Venable, Francis P.**, and *R. A. Linberry*, zirconyl citrate, A., i, 917.
- Venable, Francis P.**, and *E. O. Moehlmann*, zirconium ferrocyanide and ferricyanide, A., ii, 712.
- Venkateswaran, Ramavenkatasubba**, the molecular scattering of light in *n*-pentane, T., 2655.
- Vérain, M.** See *G. Etienne*.
- Vereinigte Chemische Werke, Akt.-Ges.**, preparation of glycerol from sugar, A., i, 980.
- Vereinigte Chininfabriken Zimmer & Co.**, preparation of *O*-alkyl derivatives of hydrocupreine, A., i, 948.
- Vergelot, Charles**, application of Bourgelot's biochemical method to some members of the Caryophyllaceae and Papilionaceae, A., i, 207.
- Verkade, P. E.**, glutaric acid. II., A., i, 520.
calorimetric determinations, A., ii, 740.
- Verkade, P. E.**, *J. Coops, jun.*, and *H. Hartman*, calorimetric researches: (1) The standardisation of a calorimetric system; comparison of the heats of combustion of the substances used for standardisation: benzoic acid and naphthalene, A., ii, 474.
- Vermast, P. G. F.**, theory of disinfection in the light of the Meyer-Overton lipid theory, A., i, 406.
- Vermeylen, G.** See *André Wahl*.
- Vernadsky, Wladimir J.**, the decomposition of kaolin by organisms, A., i, 1096.
- Vernet, Henry**. See *Amé Pictet*.
- Verzár, Fritz, J. Bögel**, and *W. Szányi*, tension and extensibility of muscle during contraction by acids or chemical means, A., i, 1212.
- Verzár, Fritz**, and *W. Szányi*, the replaceability of potassium by uranium in cross-striped muscle, A., i, 1213.
- Vesely, V.**, and *K. Dvořák*, the replacement of diazo-groups by nitro-groups, A., i, 690.
- Vesterberg, K. Albert**, chemistry of terpenes, phytosterols, and resins. I. Extraction of amyryl, A., i, 825.

- Vesterberg, *K. Albert*, and *S. Westerland*, chemistry of terpenes, phyto-sterols, and resins. II. Separation of α - and β -amyrin; preparation of α -amyrilene, A., i, 825.
- Vésses, *Maurice*, and *Georges Dupont*, the composition of the essential oil of turpentine, A., i, 1043.
- Viale, *G.*, the presence of amino-acids in milk, A., i, 898.
- Vickery, *Hubert Bradford*, rate of hydrolysis of wheat gliadin, A., ii, 754.
- Viehöver, *Arno*. See *Carl Oscar Johns*.
- Vignat. See *Alphonse Seyewetz*.
- Vila, *Antony*. See *Maurice Piettre*.
- Vila, *M.*, the separation of the globulins of horse's serum, A., i, 1209.
- Vilbrandt, *Frank C.* See *S. Lantz Shenefield*.
- Villedieu, *G.*, and (*Mme*) *G. Villedieu*, copper sprays, A., i, 415.
- Villedieu, (*Mme*) *G.* See *G. Villedieu*.
- Viola, *S.* See *G. Charrier*.
- Violle, *P. L.*, periodicity in chloride excretion during dropsical nephritis, A., i, 963.
- Vischniac, *Ch.* See *A. Gorla*.
- Visco, *Sabate*, alcohol-soluble protein of the caryopsis of *Sorghum vulgare*. I. Extraction and identification, A., i, 211.
- Vladesco, *R.* See *Gabriel Bertrand*.
- Vladimirov, *N. A.* See *Leo Alexandrovitch Tschugaev*.
- Vleck, *J. H. van*, the normal helium atom and the quantum theory, A., ii, 838.
- Vliet, *Elmer B.* See *E. H. Volwiler*.
- Vloten, *W. J. van Bommel van*. See *J. Snapper*.
- Vodret, *F. L.* See *Ernesto Puxeddu*.
- Vogel, and *E. Weber*, effect of nitrogenous fertilisers on the alkaloid content of lupines, A., i, 798.
- Vogel, *Erwin*. See *Benjamin Max Margosches*.
- Vogel, *Hans*, investigations on potassium paralysis, A., i, 196.
- Vogel, *Heinz*. See *Robert Wintgen*.
- Vogel, *Robert*, and *A. Bornstein*, paradoxical behaviour of the sugar metabolism on simultaneous administration of pilocarpine and adrenaline ("dissimulatory reversal"), A., i, 395.
- Vogel, *Robert*. See also *A. Bornstein*.
- Vogel, *Rudolf*, and *Gustav Tammann*, the ternary system iron-boron-carbon, A., ii, 852.
- Vogelenzang, *E. H.* See *I. M. Kolthoff*.
- Vogt, *Richard R.*, and *Julius A. Nieuwland*, the rôle of mercury salts in the catalytic transformation of acetylene into acetaldehyde and a new commercial process for the manufacture of paracetaldehyde, A., i, 110.
- Voigt, *Arthur*. See *Wilhelm Bilts*.
- Voigt, *Waller*, German silver, A., ii, 295.
- Vois, *R.* See *Riccardo Ciusa*.
- Voit, *Kurt*, formaldehyde in the urine after administration of urotropin, A., i, 1218.
- Volk, *H.* See *Fr. Boedecker*.
- Vollbrecht, *Erich*. See *Karl Freudenberg*.
- Volmer, *M.*, and *I. Estermann*, vaporisation coefficients of solid and liquid mercury, A., ii, 193.
- coefficients of vaporisation and their relationship to Ostwald's step rule, A., ii, 193.
- Volmer, *M.*, and *K. Riggert*, dependence of velocity of reaction on the concentration in photochemical processes, A., ii, 336.
- Volwiler, *E. H.*, and *Elmer B. Vliet*, preparation and hydrolysis of benzyl esters, A., ii, 41.
- Volwiler, *E. H.* See also *Oliver Kamm*.
- Voorhees, *V.*, and *Roger Adams*, the use of the oxides of platinum for the catalytic reduction of organic compounds. I., A., ii, 558.
- Vorländer, *Daniel*, molecular arrangement and liquid crystal formation, A., ii, 554.
- Voroshcov, *N. N.*, synthesis of substantive azo-dyes derived from naphthalene, A., i, 956.
- chemistry of naphthalene and its derivatives; chemical peculiarities of the naphthalene nucleus, A., i, 1135.
- Vorschütz, *Joseph*, continuous current and permeability (in muscle). II. Effect of alkaloidal salts and other organic electrolytes, A., i, 791.
- Vortmann, *Giorgio*, sodium sulphide as a substitute for hydrogen sulphide in qualitative analysis, A., ii, 653.
- Vorwerk, *Waller*. See *Fritz Paneth*.
- Vosburgh, *Warren C.*, some errors in the study of invertase action, A., i, 61.
- potassium dichromate as a standard in iodimetry and the estimation of chromates by the iodide method, A., ii, 863.
- Vosburgh, *Warren C.* See also *Marion Eppley*.
- Voss, *G. von*. See *Paul Friedländer*.

- Vossen, G.** See *George Schroeter*.
Votocsek, Emil, the rapid estimation of chlorine in organic compounds, A., ii, 863.
Vournasos, Alexander Ch., complex mixed antimony iodobromides, A., ii, 650.
Vridhachalam, P. N. See *Kishori Lal Moudgill*.
Vürtheim, A., the volumetric estimation of magnesium in potassium salts, A., ii, 870.
Vürtheim, A., and *G. H. C. van Bers*, the volumetric estimation of calcein, A., ii, 869.
Vuillaume, M. See *Augustin Boutaric*.
Volquin, E. See *M. Entat*.

W.

- Wachtel, Curt**, detection and estimation of morphine and other alkaloids in animal excreta and organs, A., ii, 171.
Wackermann, Harald. See *Otto Diels*.
Wadsworth, Raymond V. See *Arthur W. Knapp*.
Wächter, Friedrich, some remarkable properties of gases, A., ii, 476.
 the constitution of the chemical elements, A., ii, 837.
 the disintegration of chemical elements, A., ii, 838.
Wälti, A. See *Paul Karrer*.
Waentig, Percy, preparation and properties of cellulose solutions, A., i, 988.
Wagner, Adalbert. See *Erich Schmidt*.
Wagner, Josef. See *Albin Kurtenacker*.
Wagner, Mario Basto, general methods for the determination of the true molecular constitution of pure substances and their mixtures, A., ii, 116.
 thermodynamics of mixtures. IX., A., ii, 116.
 theory of equations of state. II., A., ii, 117.
Wagner, Otto. See *Paul Rabe*.
Wagner, Richard, and *Jakob K. Parnas*, observations on sugar synthesis. II. An abnormal disturbance of carbohydrate exchange and its relation to diabetes mellitus, A., i, 965.
Wagner, Richard. See also *Jakob K. Parnas*.
Wagner, Walter. See *Julius Meyer*.
Wahl, André, and *R. Lantz*, the 2-hydroxy-1-arylnaphthylamines, A., i, 823.
Wahl, André, G. Normand, and *G. Vermeylen*, the monochlorotoluenes, A., i, 442.
Wahl, Carl, rupture of bridge linkings, A., i, 653.
Wahl, Ottmar. See *Kurt Hess*.
Wahl, Rudolf. See *Hans Theodor Bucherer*.
Wahlberg, H. E., Swedish pines and spruces, A., i, 1101.
Wahlin, H. B., behaviour of free electrons toward gas molecules, A., ii, 608.
Wakeman, Alfred John. See *Thomas Burr Osborne*.
Waksman, Selman A., micro-organisms concerned in the oxidation of sulphur in the soil. III. Media used for the isolation of sulphur bacteria from the soil, A., i, 706.
Waksman, Selman A., and *Jacob S. Joffe*, chemistry of the oxidation of sulphur to sulphuric acid by micro-organisms, and transformation of insoluble phosphates into soluble forms. A., i, 416.
Waksman, Selman A. See also *Jacob Goodale Lipman*.
Walbum, L. E., effect of manganous chloride on the formation of diphtheria toxin, A., i, 795.
 effect of different metallic salts on the formation of staphylococci, A., i, 795.
 formation of bacterial toxins. II. Diphtheria toxin, A., i, 902.
Walbum, L. E. See also *Helen A. Purdy*.
Walden, Paul, the λ -V curves of non-aqueous solutions of binary electrolytes, A., ii, 345.
Waldo, John H. See *Oliver Kamm*.
Waldschmidt-Leitz, Ernst. See *Richard Willstätter*.
Wales, H., naphthalenesulphonic acids. IV. Solubilities of some amine salts of naphthalenesulphonic acids, A., i, 727.
Walker, Florence. See *Henry Clapp Sherman*.
Walker, Hilda. See *Evelyn Ashley Cooper*.
Walker, (Sir) James, the rôle of the physicist in the development of chemical theory. T., 735.
Walker, (Miss) Nellie. See *Alexander McKenzie*.
Walker, Osman James. See *George Shannon Forbes*.
Wallington, Reginald Wilfred. See *Francis Ernest Francis*.
Wallis, Albert Edward. See *Robert Ludwig Mond*.
Walter, Heinrich, the chemical constitution of protoplasm, A., i, 308.

- Walters, F. M., jun.**, wave-length measurements in arc spectra photographed in the yellow, red, and infra-red, A., ii, 100.
- Walther, P.**, sodium carbonate minerals from Lake Magadi, Kenya Colony, A., ii, 859.
- Walton, James H., and Llewellyn B. Parsons**, preparation and properties of the persulphides of hydrogen, A., ii, 281.
- Walton, James H., and Clarence R. Wise**, equilibrium in the system, lithium chloride-quinoline, A., i, 370.
- Wampler, R. W.** See *H. H. King*.
- Wang, Chi Che, I.** The composition of Chinese edible birds' nests and the nature of their proteins. II. The isolation and the nature of the amino-sugar of Chinese edible birds' nests, A., i, 299.
- Warburg, Erik Johan**, carbonic acid compounds and hydrogen-ion activities in blood and salt solutions, A., i, 788.
- Warburg, Otto**, physical chemistry of cell respiration, A., i, 190.
surface reactions in living cells, A., i, 787.
- Warburg, Otto.** See also *C. Müller*.
- Warburg, Otto, and Erwin Negelein**, energy changes accompanying the assimilation of carbon dioxide, A., i, 1097.
- Wareoillier and Le Moal**, progressive disappearance of free sulphurous acid in a preserved apple juice, A., i, 415.
- Ward, Charles Frederick**, the bromination of acids in the α -position, T., 1161.
- Ward, James Bertram.** See *Archibald Edwin Goddard*.
- Warden, C. C.**, nature of alcoholic fermentation, A., i, 307.
- Wardlaw, William, and Frederic William Pinkard**, the oxidising properties of sulphur dioxide. III. Copper chlorides, T., 210.
- Wardlaw, William.** See also *Frederic William Pinkard, and Lachlan Macquarie Stewart*.
- Wark, Ian William**, a rapid iodometric estimation of copper and iron in mixtures of their salts, T., 358.
- Warner, J. C.** See *Oliver W. Brown*.
- Warren, Lewis E.**, estimation of digitoxin in digitan, A., ii, 403.
- Warren, Lewis E., and A. H. Clark**, the Bliss method for the separation of strychnine from quinine, A., ii, 171.
- Wartenberg, H. von, and O. Bosse**, the vapour pressure of some salts. III., A., ii, 739.
- Wartenberg, H. von, and Friedrich August Henglein**, the dissociation of chlorine, A., ii, 441.
- Wartenberg, H. von, and H. Schulz**, vapour pressure of some salts. II., A., ii, 146.
- Wasastjerna, Jarl A.**, refraction of light and atomic structure; refraction equivalents of ions, A., ii, 1.
optical properties of solutions; a theory of the structure of the molecules of electrolytes, A., ii, 2.
structure of atoms and molecules in the light of the dispersion theory, A., ii, 491.
thermochemistry and electromotive force in electric elements, A., ii, 815.
- Washburn, Edward Wight, and Louis Navias**, the relation of chalcedony to other forms of silica, A., ii, 304.
- Washington, Henry Stephens, and Herbert Eugene Merwin**, augite from Hawaii, A., ii, 220.
aphthitalite from Kilauea, A., ii, 386.
- Wasicky, Richard, *Capsella bursa pastoris***, Moench, A., i, 907.
- Wassermann, Paul.** See *Hugo Weil*.
- Wastl, (Frl.) H.** See *R. Brinkman*.
- Watanabe.** See *Riko Majima*.
- Watanabé, Manjirô**, babingtonite from Japan, A., ii, 651.
- Waters, C. A.** See *William Edward Garner*.
- Waterman, Henry C.** See *David Breeze Jones*.
- Waterman, Hein Israel, and J. N. J. Perquin**, evaluation of the degree of unsaturation of mineral oils in the Bergius process, A., ii, 90.
the estimation of aromatic hydrocarbons in mineral oil fractions, A., ii, 399.
- Waterman, Hein Israel, and M. J. van Tussenbroek**, the formation of formic acid during the decomposition of dextrose in alkaline solution, A., i, 433.
- Watkins, Cyril Mercer.** See *Francis Ernest Francis*.
- Watson, Arthur Frederick.** See *Jack Cecil Drummond*.
- Watson, Amy R.** See *J. W. W. Dyer*.
- Watson, Edwin Roy, and Sikhishushan Dutt**, attempts to prepare red sulphide dyes. I., T., 1939.
attempts to prepare red sulphide dyes. II. Mercaptan derivatives of azo-dyes, T., 2414.
- Watson, S. J., Atkinson's process** for the estimation of potassium in the presence of sodium, magnesium, sulphates, and phosphates, A., ii, 586.

- Wattenberg, H.** See *E. Zintl*.
- Wayman, Marguerite.** See *Henry Clapp Sherman*.
- Wayne, Edward Johnson, and Julius Berend Cohen,** the aldehydosalicyclic acids and their derivatives, T., 1022.
- Weber, A.** See *W. D. Treadwell*.
- Weber, E.** See *Vogel*.
- Weber, I.** See *Phabus A. Levene*.
- Weber, J.** See *Leo Grünhut*.
- Weber, Sophus,** critical constants of mercury, A., ii, 646, 769.
- Weber, W.** See *William Küster*.
- Webster, David L.,** critical potentials of the L-series of platinum, A., ii, 805.
- Webster, Thomas Arthur.** See *James Argyll Campbell*.
- Wecker, Albert.** See *Heinrich Wieland*.
- Wedekind, Edgar, and Cl. Weinand,** condensation products from acid haloids. X. Action of triethylamine on ketopinyl chloride, A., i, 548.
- Wedekind, Edgar, and Cl. Weinand** [with *M. Miller*], condensation products from acid haloids. IX. Ketonium compounds, A., i, 234.
- Weerd, F. N. B. de.** See *Nicolaas Schoorl*.
- Wegscheider, Rudolf,** transformations during the fission of racemic substances, A., i, 441.
equation of condition, A., ii, 192.
the ammonium carbonate-carbamate equilibrium, A., ii, 376.
influence of ethyl alcohol on the colour change of phenolphthalein, A., ii, 387.
course of simultaneous reactions, A., ii, 489.
- Wehmer, Carl,** action of coal gas on plants, A., i, 211.
- Weibezahn, Karl.** See *Jakob Meisenheimer*.
- Weichmann, Richard,** an absorption band spectrum for water in the region of wave-lengths of several decimetres, A., ii, 5, 331.
- Welck, R.** See *Henri Gault*.
- Welckert, O.** See *Walther Borsche*.
- Weidenhagen, Rudolf.** See *Burchhardt Helferich*.
- Weidert, F.,** absorption spectra of glasses of various composition containing didymium, A., ii, 413.
- Weigel, O.,** behaviour of sulphides of heavy metals in aqueous solutions, A., ii, 644.
- Weigert, Fritz,** physical chemistry of colour vision, A., i, 607.
photochemistry of the retina, A., ii, 11.
- Weigert, Fritz,** absorption and dispersion of radiation, A., ii, 605.
fluorescence, photochemical action, and Einstein's law, A., ii, 681.
- Weigert, Fritz, and Karl Kellermann,** photochemistry of chlorine detonating gas, A., ii, 808.
- Weigert, Fritz, and Walter Schöller,** photochemistry of silver compounds, A., ii, 10.
- Weigt, Hans,** electrical moments of carbon monoxide and dioxide molecules, A., ii, 109.
- Weil, Hugo, and Karl Brimmer,** reduction of substituted salicylic acids, A., i, 349.
- Weil, Hugo, and Walter Heerd, the** reduction of naphtholcarboxylic acids to aldehydes, A., i, 256.
- Weil, Hugo, and E. Moser,** the action of sodium hydrogen sulphite on nitro-compounds of the benzene series, A., i, 443.
- Weil, Hugo, and Hermann Ostermeier,** reduction of naphthalene- and naphthol-carboxylic acids, A., i, 139.
- Weil, Hugo, Max Traun, and Sigismund Marcel,** the reduction of substituted salicylic acids, A., i, 1023.
- Weil, Hugo, and Paul Wassermann,** the action of sodium hydrogen sulphite on the nitroanilines, A., i, 1006.
- Weil, K.** See *Adolf Windaus*.
- Weimer, J. L.** See *L. L. Harter*.
- Wein, Walter.** See *Wilhelm Biltz*.
- Weinand, Cl.** See *Edgar Wedekind*.
- Weinberg, Abraham Albert,** nephelometry; a nephelometer with constant standard, A., ii, 309.
- Weinland, Rudolf Friedrich, and Georg Fischer,** mangani-acetates and -benzoates, A., i, 421.
- Weinland, Rudolf Friedrich, and A. Kissling,** the constitution of some ferri-chloride (-thiocyanatenitrate) compounds with pyridine or quinoline, A., i, 363.
- Weinland, Rudolf Friedrich, and Rudolf Stroh,** complex compounds of lead acetate, A., i, 981.
the constitution of basic lead salts, A., ii, 767.
- Weiser, Harry B.,** adsorption by precipitates. V. Adsorption during the precipitation of colloids by mixtures of electrolytes, A., ii, 262.
hydrous oxides. III. and IV., A., ii, 575, 853.
- Weiser, Harry B., and Henry O. Nicholas,** influence of the concentration of colloids on their precipitation by electrolytes, A., ii, 266.

- Weiser, Nikolaus.** See *Josef Hollnatz*.
- Weismantel, Josef.** See *Julius von Braunn*.
- Weiss, E., and V. C. Garner,** chronic nephritis with an unusual degree of nitrogen retention, A., i, 704.
- Weiss, H., and P. Henry,** influence of temperature on the velocity of interpenetration of solids, A., ii, 207.
influence of the time factor on the interpenetration of solids by chemical reaction, A., ii, 487.
- Weiss, John Maurice, and Charles R. Downs,** catalytic oxidation of fluorene, A., i, 27.
the formation of malic acid, A., i, 622.
- Weiss, Pierre,** magnetism and the constitution of atoms, A., ii, 28.
- Weiss, R.,** estimation of the calcium content of blood, A., ii, 227.
- Weiss, Soma, and Robert A. Hatcher,** strychnine, A., i, 900.
- Weiss, St.** See *Herbert Elias, and Z. Ernst*.
- Weissenberg, K.** See *M. Ettisch*.
- Weissenberger, Georg, and A. Zoder,** analysis of chlorosulphonic acid, A., ii, 390.
- Weissgerber, Rudolf.** See *Gesellschaft für Teerverwertung, m. b. H.*
- Weitz, Ernst,** dissociation of the so-called 1:1'-dibenzyltetrahydro-4:4'-dipyridyl, A., i, 470.
- Weitz, Ernst, and Theodor König,** free ammonium radicles. IV. Further investigations on 1:1'-dibenzyl-4:4'-dipyridinium and its homologues and on the so-called 1:1'disubstituted tetrahydro-4:4'-dipyridyls, A., i, 1186.
- Weitz, Ernst, and Richard Ludwig,** free ammonium radicles. III. Existence of *N*-benzylpyridinium, A., i, 365.
- Weitzenböck, Richard.** See *Roland Scholl*.
- Welker, William H., and J. L. Bollman,** effect of subcutaneous injections of solutions of potassium cyanide on the catalase content of the blood, A., i, 192.
- Wells, Horace Lemuel,** complex chlorides containing gold. I. Pollard's ammonium silver auric chloride, A., ii, 449.
complex chlorides containing gold. II. Cæsium triple salts, A., ii, 449.
a chromophore grouping of atoms in inorganic triple salts and a general theory for the cause of the colours of substances, A., ii, 464.
some complex chlorides containing gold. III. A new cæsium auric chloride, A., ii, 514.
discussion of triple salts, A., ii, 560.
- Wells, P. V.,** simple theory of the nephelometer, A., ii, 308.
- Wells, Roger Clark,** estimation of silica in filtered sea water, A., ii, 868.
- Welter, Georges,** micro-analysis of organic substances, A., ii, 399.
- Welter, Georges.** See also *Maurice Nicloux*.
- Weltmann, Oskar,** the iodine number of urine, A., i, 88.
- Weltmann, Oskar.** See also *Hermann K. Barrenscheen*.
- Weltzien, Wilhelm.** See *Kurt Hess*.
- Welmüller, Ferdinand,** the diastatic action of cow's milk towards various starches, A., i, 400.
- Wende, E.** See *S. Goy*.
- Wendehorst, Erwin.** See *Gerhart Jander*.
- Wendt, Gerald L.,** active hydrogen and nitrogen, A., ii, 639.
- Wendt, Gerald L., and Clarence E. Irion,** experimental attempts to decompose tungsten at high temperatures, A. ii, 773.
- Wendt, Gerald L., and Robert S. Landauer,** triatomic hydrogen. II., A., ii, 369.
- Wenger, Paul, and P. Christin,** decomposition and analysis of monazite, A., ii, 722.
- Wentzel, Gregor,** spark lines in the Röntgen spectrum, A., ii, 249.
Röntgen spectra and chemical valency, A., ii, 607.
- Wenzl, Hermann.** See *Emil Fromm*.
- Werb, Otto.** See *Bruno Emmert*.
- Wernecke, Hermann.** See *Fritz Mayer*.
- Werner, A.** See *Hans Geiger*.
- Werner, Emil Alphonse,** the constitution of carbamides. XIV. The decomposition of urea by sodium hypobromite in alkaline solution, and an improved procedure for the estimation of urea by this means, T., 2318.
- Werner, Emil Alphonse, and James Bell,** the preparation of methylguanidine, and of $\beta\beta$ -dimethylguanidine by the interaction of dicyanodiamide, and methylammonium and dimethylammonium chlorides respectively, T., 1790.
- Werner, Felicitas.** See *Albin Kurtenacker*.
- Werner, Fritz.** See *K. Hugo Bauer*.
- Werner, P.** See *David Holde*.
- Werner, Sven.** See *H. Baggesgaard-Rasmussen*.
- Wertenstein, Mathilde.** See *Hilary Lachs*.
- Wertheim, Edgar,** a modified Schiff's solution, A., ii, 793.

- Wertheim, Edgar.** See also *John W. E. Glattfield.*
- Wertheimer, Ernst.** See *Emil Abderhalden.*
- Wertheimer, Irma.** See *F. Himstedt.*
- West, Augustus P., and Zoila Montes,** lumbago oil (candle-nut oil), A., i, 95.
- West, Randolph.** See *F. A. Stevens.*
- West, Ralph Winton,** quantitative reduction by hydriodic acid of halogenated malonyl derivatives. II. The *s.*-tetra-substituted amides of bromo- and chloro-malonic acid, T., 2196.
- Wester, D. H.,** the manganese content of (Dutch) seeds, A., i, 93.
the chemical constituents of some *Loianthaceae*, A., i, 208.
the manganese content of flowers, A., i, 309.
I. Culture experiments with soja beans. II. Occurrence of urease in parts of the plant other than the seeds, A., i, 311.
influence of different chemicals, of kations and anions, and of mixtures of electrolytes, on the ureolitic power of urease, A., i, 391.
- Westerlind, S.** See *K. Albert Vesterberg.*
- Westermann, H.** See *Karl von Auwers.*
- Westgren, Arne, and Axel E. Lindh,** crystalline structure of iron and steel, A., ii, 152.
- Westgren, Arne, and Gösta Phragmén,** crystal structure of iron and steel. II., A., ii, 711.
- Westgren, Arne, and Josef Reitzstötter,** coagulation of colloids, A., ii, 625.
- Weston, Arnold,** the quaternary system potassium sulphate-magnesium sulphate-ammonium sulphate-water, T., 1223.
- Wetterfors, Paul,** experimental proof of some theories of natural rotatory power of optically active solutions, A., ii, 247.
- Wetzel, Johannes,** water pump, A., ii, 139.
- Whatham, Thomas William.** See *Arthur George Perkin.*
- Wheeler, Alvin Sawyer, and T. M. Andrews,** hydroxynaphthaquinone. IV. New derivatives of 2:3:8-tribromo-5-hydroxy-1:4-naphthaquinone, A., i, 351.
- Wheeler, Alvin Sawyer, and B. Naiman,** hydroxynaphthaquinones. V. Derivatives of 2-bromo-5-hydroxy-1:4-naphthaquinone (monobromojuglone), A., i, 1165.
- Wheeler, Alvin Sawyer, and Ira W. Smithey,** *p*-cymene. III. The bromination of 2-amino-*p*-cymene, A., i, 332.
- Wheeler, Richard Vernon.** See *Walter Mason, William Payman, and Frederick Vincent Tidswell.*
- Wheeler, Thomas Sherlock, and Ethelbert William Blair,** the action of ozone on hydrocarbons with special reference to the production of formaldehyde. I. The action of ozone on methane, A., i, 1105.
a receiver for fractionation in a current of gas or under reduced pressure, A., ii, 258.
- Wheeler, Thomas Sherlock.** See also *Ethelbert William Blair.*
- Whelan, Julia,** oil of *Bystropogon canus*, A., i, 847.
- Wherry, Edgar Theodore,** optical-crystallographic properties of calcium oxalate monohydrate, A., ii, 450.
- Wherry, Edgar Theodore, and Raymond M. Hann,** crystallographic-optical properties of calcium fumarate and maleate, A., i, 716.
- Wherry, Edgar Theodore, and Earl V. Shannon,** crocidolite from Eastern Pennsylvania, A., ii, 518.
- Wherry, Edgar Theodore.** See also *Earl V. Shannon.*
- Whetham, Margaret Dampier.** See *Marjory Stephenson.*
- Whipple, G. H.** See *G. D. Delprat, and K. F. Polkan.*
- White, Albert Greville,** limits for the propagation of flame in vapour-air mixtures. I. Mixtures of air and one vapour at the ordinary temperature and pressure, T., 1244.
limits for the propagation of flame at various temperatures in mixtures of ammonia with air and oxygen, T., 1688.
limits for the propagation of flame in vapour-air mixtures. II. Mixtures of more than one vapour and air at the ordinary temperature and pressure, T., 2561.
- White, Charles Powell,** copper in tumours and normal tissues, A., i, 399.
application of the methods of correlation to the study of the urine, A., i, 1214.
- White, Ernest C.** See *Alfred T. Larson.*
- Whiteley, (Miss) Martha Annie.** See *Arthur Geoffrey Rendall.*
- Whitfield, Bernard Wyndham.** See *Alfred Francis Joseph.*
- Whitmores, Frank C., and Edmund Burrus Middleton,** organic mercury compounds prepared from *o*-chloromercuri-*p*-nitrobenzoyl chloride, A., i, 888.
- Whitney, Milton,** origin of soil colloids, A., i, 708.

- Whittaker, E. T.**, the quantum mechanism in the atom, A., ii, 632.
- Wibaut, Johan Pieter**, the energy of the atomic linkings in saturated and unsaturated hydrocarbons, A., i, 909.
behaviour of amorphous carbon and sulphur at high temperatures; carbon sulphides, A., ii, 52, 373.
comparison of the linking of carbon atoms in graphite and in aromatic hydrocarbons, A., ii, 259.
the preparation of carbon from carbon monoxide by means of a catalyst, A., ii, 565.
- Wibaut, Johan Pieter**, and **J. Jürgens**, action of mercuric acetate on nitrobenzene, A., i, 694.
- Widmark, Erik Matteo Prochet**, succinodihydrogenase, A., i, 600.
a micro-method for the estimation of ethyl alcohol in blood, A., ii, 789.
- Widmark, Erik Matteo Prochet**, and **Carl Alex. Jeppsson**, a defined organic catalyst with a hydrogen-ion optimum, A., ii, 437.
- Widmark, Erik Matteo Prochet**, and **Gunnar Lindahl**, chemical conditions for the maintenance of normal cell structure. IV. Difference in nitrogen and water content of muscle treated with sodium and calcium chloride, A., i, 607.
- Widmer, C.** See **Robert Eder**.
- Wieschmann, Ernst**, permeability of the red corpuscles of human blood for anions, A., i, 80.
permeability of the red blood corpuscles, A., i, 289.
- Wiechowski, W.** See **Oskar Adler**.
- Wiedbrauk, Erich**, apparatus for the rapid determination of the specific gravity of small amounts of liquid, A., ii, 740.
- Wiegner, Georg**, and **J. Magasanik**, estimation of volatile fatty acids, A., ii, 532.
- Wiegner, Georg, J. Magasanik**, and **H. Gossner**, hysteresis phenomena and coagulation of sols with ultramicros of rod-like structure, in particular vanadium pentoxide compared with fibrin sols, A., ii, 356.
- Wieland, Heinrich**, toad venom, A., i, 199.
- Wieland, Heinrich**, and **Franz Adickes**, bile acids. XIII. The formula of isodeoxybilanic acid; a contribution to orientation, A., i, 838.
- Wieland, Heinrich**, and **Richard Alles**, the poisonous substance of toads, A., i, 784.
- Wieland, Heinrich**, and **Ludwig Bettag**, the Friedel-Crafts' reaction, A., i, 1033.
- Wieland, Heinrich**, and **Fritz Kügl**, organic radicles with quadrivalent nitrogen. III., A., i, 726.
- Wieland, Heinrich, Egon Popper**, and **Hermann Seefried**, the occurrence of free radicles in chemical reactions; the radicles of the basic triphenylmethane dyes, A., i, 772.
- Wieland, Heinrich, Fritz Reindel**, and **Juan Ferrer**, diphenylene-ethylene, A., i, 1137.
- Wieland, Heinrich**, and **Georg Scheuing**, rosaniline-sulphurous acid and its colour reactions with aldehydes, A., i, 58.
- Wieland, Heinrich**, and **Otto Schlichting**, bile acids. XI. The oxidation of cholic acid, A., i, 554.
bile acids. XII. Cidoidanic acid, A., i, 838.
- Wieland, Heinrich**, and **Wilhelm Schulenburg**, bile salts. X. The further degradation of deoxycholic acid, A., i, 346.
- Wieland, Heinrich**, and **Albert Wecker** [with **T. J. Albert**, and (*Frl.*) **E. Haas**], hydrazines. XXV. A new class of (di)tertiary aromatic hydrazines, A., i, 779.
- Wieland, Hermann**, the mode of action of narcotic gases; nitrous oxide and acetylene, A., i, 497.
- Wiesler, Herta.** See **Eduard Kopetschni**.
- Wiesmann, H.** See **Otto Lemmermann**.
- Wieters, Hermann.** See **Edmund Speyer**.
- Wigley, Gladys Mary.** See **Hubert Frank Coward**.
- Wignall, Harry.** See **British Dyestuffs Corporation, Ltd.**
- Wilber, D. T.** See **Eduard L. Nichols**.
- Wilbrand, Eberhart**, excretion of sweat and the composition of the blood, A., i, 79.
- Wilder, Russell M.**, and **Malcolm D. Winter**, the threshold of ketogenesis, A., i, 893.
- Wiley, J. A.** See **George Leslie Kelley**.
- Wilke, C.** See **David Holde**.
- Wilke, E.**, aqueous carbonic acid solutions, A., ii, 52.
- Wilke, K.**, anthraquinone derivatives, A., i, 943.
- Wilkendorf, Rudolf.** See **Erich Schmidt**.
- Wilkes, Sidney Herbert.** See **Malcolm Percival Applebey**.
- Wilkins, T. R.**, multiple valency in the ionisation by α -rays, A., ii, 607.
- Wilkinson, John Frederick.** See **Frederick Challenger**.
- Will, Erich.** See **Karl Andreas Hofmann**.
- Willaman, J. J.**, preparation of inulin, with special reference to artichoke tubers as a source, A., i, 434.

- Willard, Hobart Hurd, and W. Ellwood Cake**, volumetric estimation of sulphide by oxidation to sulphate, A., ii, 80.
- Willard, Hobart Hurd, and Dorothy Hall**, separation of copper by means of thiophenylhydantoic acid, A., ii, 872.
- separation and estimation of cobalt. I. Separation of cobalt by means of thiophenylhydantoic acid. I. Separation of cobalt from other metals, A., ii, 874.
- separation and estimation of cobalt. II. Gravimetric estimation of cobalt, A., ii, 874.
- separation and estimation of cobalt. III. Volumetric estimation of cobalt, A., ii, 875.
- Willard, Hobart Hurd, and G. Frederick Smith**, preparation and properties of magnesium perchlorate and its use as a drying agent, A., ii, 850.
- Williams, Arthur G.**, estimation of phenanthrene, A., ii, 90.
- Williams, E. H., and Jakob Kunz**, photoelectric effect of alkali [metal] vapours and a new determination of h , A., ii, 809.
- Williams, E. J.**, a simple receiver for fractional distillation under diminished pressure, A., ii, 26.
- Williams, G. A., and John Bright Ferguson**, diffusion of hydrogen and helium through silica glass and other glasses, A., ii, 841.
- Williams, H.** See *W. Singleton*.
- Williams, J. Lisle**, the total non-protein nitrogen constituents of the blood in chronic nephritis with hypertension, A., i, 401.
- Williams, Percy Noel.** See *Arthur Henry Salway*.
- Williams, Roger J.** See *Chester E. Adams*.
- Williams, R. R.**, vitamins from the point of view of structural chemistry, A., i, 280.
- Williams, Vernon Harcourt.** See *Alexander Findlay*.
- Williams, William Dudley.** See *Leonard Eric Hinkel*.
- Williamson, E. C.**, ionisation of potassium vapour by light, A., ii, 681.
- Willig.** See *William Kuster*.
- Willmann, Karl**, the natural iron hydroxides, A., ii, 76.
- Willmersdorf, Amalie.** See *Jakob Meisenheimer*.
- Willstätter, Richard, and Wilhelm Csahny**, emulsin, A., i, 390.
- Willstätter, Richard, Johanna Graser, and Richard Kuhn**, invertase. III., A., i, 1200.
- Willstätter, Richard, and Ludwig Kalb** [with *G. von Miller*], the reduction of lignin and of carbohydrates with hydrogen iodide and phosphorus, A., i, 989.
- Willstätter, Richard, and Richard Kuhn**, remarks on the elution of saccharase and maltase from adsorbed substances, A., i, 283.
- the specific nature of saccharase and raffinase, A., i, 284.
- Willstätter, Richard, and Gertrud Oppenheimer**, the lactase content and the fermenting power of lactose-fermenting yeasts, A., i, 203.
- emulsin. II., A., i, 959.
- Willstätter, Richard, and Fritz Racke**, invertase. II., A., i, 598.
- Willstätter, Richard, and Werner Stalbelt**, maltase. III. The non-identity of maltase and α -glucosidase, A., i, 232.
- maltase. IV. The fermenting activity of yeasts poor in maltase, A., i, 306.
- Willstätter, Richard, and Ernst Waldschmidt-Leitz**, alkalimetric estimation of amino-acids and peptides, A., ii, 169.
- Willstätter, Richard, Otto Wolfes, and Horst Maeder**, tropinonecarboxylic acid esters, A., i, 938.
- Wilson, Daniel.** See *Christopher Kelk Ingold*.
- Wilson, F. H. II.** See *Eustace Ebenezer Turner*.
- Wilson, Forsyth James, and Robert Burns**, reactions of thiosemicarbazones. I. Action of halogen compounds, T., 870.
- Wilson, Forsyth James, Isaac Vance Hopper, and Archibald Barclay Crawford**, the action of amines on semicarbazones. I. Preparation of an optically active semicarbazide, T., 866.
- Wilson, H. A.** See *Arthur Amos Noyes*.
- Wilson, J. Waller.** See *Philip H. Mitchell*.
- Wilson, Robert E., and Everett W. Fuller**, reaction of carbonyl chloride with benzene and *m*-xylene in the presence of aluminium chloride, A., i, 827.
- Wilson, Samuel, and C. R. McCroskey**, polymorphic transformations of antimony trisulphide, A., ii, 154.
- Wilson, S. J.**, estimation of sodium in serum without the use of platinum dishes, A., ii, 395.
- Wilson, Thomas A.** See *William Albert Noyes*.

- Wilson, T. E.** See *William Ernest Stephen Turner*.
- Wilson, William H.** scorpion venom, A., i, 969.
- Wiltshire, Marion O. P.**, basal metabolism in menstruation, A., i, 395.
- Winch, Howard James**, and **V. L. Chandratreya**, the volumetric estimation of titanium dioxide in bauxite, A., ii, 459.
- Windaus, Adolf** [with *Harry Grimmel*, *H. Lüders*, and *A. von Staden*], cholesterol, A., i, 541.
- Windaus, Adolf**, *W. Dörries*, and *H. Jensen*, the behaviour of certain bisacylaminoethylene derivatives prepared from glyoxalines, A., i, 60.
- Windaus, Adolf**, and *Walter Hückel*, the formation of cyclic compounds from hydroaromatic dicarboxylic acids, A., i, 558.
- Windaus, Adolf**, and *Hermann Lüders*, the action of benzoyl peroxide on cholesterol, A., i, 453.
- Windaus, Adolf**, and *K. Weil*, digitonin and its derivatives, A., i, 848.
- Windheuser, K.** See *A. Morgen*.
- Windisch, H.** See *Friedrich L. Hahn*.
- Windisch, Wilhelm**, and *Philipp Osswald*, ionisation and surface activity of aqueous solutions of aliphatic acids, A., ii, 123.
- Wingårdh, K. A.**, absorption of Röntgen rays, A., ii, 249.
- Wingler, Aug.** See *Heinrich Kiliani*.
- Winkel, Herbert.** See *Arthur Percival Tanberg*.
- Winkler, Karl.** See *Max Trautz*.
- Winkler, Ludwig Wilhelm**, estimation of the iodine-bromine number without using potassium iodide, A., ii, 534. gravimetric analysis. XXV. Estimation of manganese, A., ii, 539. alkali iodides, A., ii, 764.
- Winter, Malcolm D.** See *Russell M. Wilder*.
- Winter, O. B.** See *Charles S. Robinson*.
- Winterstein, Ernst**, and *D. Iatrides*, taxine, the alkaloid from the yew, *Taxus baccata*. I, A., i, 572.
- Winterstein, Ernst**, and *J. Teleczky*, constituents of saffron. I. Picrocrocin, A., i, 563.
- Wintgen, Robert**, and *Heinz Vogel*, equilibrium, gelatin-hydrochloric acid. II., A., i, 280.
- Winther, Chr.**, theory of induced reactions, A., ii, 336. photochemical oxidation of hydriodic acid, A., ii, 808. photochemical efficiency of absorbed radiation, A., ii, 808.
- Winther, Chr., H. Baggesgaard-Rasmussen**, and *Erling Schreiner*, a simple method for the measurement of ultra-violet absorption, A., ii, 729.
- Wirth, Th.** See *Adolf Grün*.
- Wise, Clarence R.** See *James H. Walton*.
- Wisbart, Mary B.** See *F. M. Allen*.
- Wismer, K. L.**, pressure volume relation of superheated liquids, A., ii, 425.
- Wiśniewski, F. J.** See *K. Jabłoński*.
- Wiswald, J.** See *Emil Briner*.
- Withers, John Charles.** See *Robert George Fargher*.
- Withrow, James R.** See *Edgar C. Bain*, and *S. Lantz Shenefield*.
- Witt, J. C.**, oxidation of oxalic acid in the absence of other acids, A., ii, 594.
- Wittelsbach, Waller.** See *Kurt Hess*.
- Wittka, Franz.** See *Adolf Grün*.
- Wittmann, Adolf.** See *Walter Dieckmann*.
- Wittwer, Max.** See *Hans Lecher*.
- Wittwer, R.** See *Hans Rupe*.
- Witzemann, Edgar**, catalytic effect of ammonia on the oxidation of butyric acid with hydrogen peroxide, A., i, 6. transition from crystalloid to colloid properties within homologous series, A., ii, 423.
- Wöber, A.**, iodometric estimation of copper in the presence of iron, A., ii, 583.
- Wöhler, Lothar**, and *F. Müller*, calcium silicides, A., ii, 293.
- Wohl, Alfred**, preparation of acetaldehyde and acetic acid, A., i, 430. equation of condition. II. Principal equation of condition. III. Principal equation of condition and the equation of condition for individual substances. IV. Compressibility equations of liquids, A., ii, 117.
- Wohl, Alfred**, and *H. Krull*, saccharification of cellulose, A., i, 232.
- Wohl, Alfred**, and *R. Schellenberg*, conversion of active glyceraldehyde into active glyceric acid, A., i, 626.
- Wojnicz-Sianozencki, Z.**, kinetics of open saturated chains of carbon atoms in relationship to the Baeyer strain theory, A., i, 330.
- Wolf, A.** See *Alexander Guthrie*.
- Wolf, Charles George Lewis**, the mechanism of the reversal in reaction of a medium which takes place during growth of *Bacillus diphtheriae*, A., i, 1218.
- Wolf, Charles George Lewis**, and *Eric Keightley Rideal*, the properties of dibenzoylcytine, A., i, 1114.
- Wolf, Emil.** See *Georg von Kereszty*.

- Wolf, Elizabeth P.**, inflammation. I. Influence of chemicals on the chemotaxis of leucocytes in vitro, A., i, 401.
- Wolfe, Otto**, and **Horst Maeder**, tropinonecarboxylic acid esters, A., i, 938.
- Wolfe, Otto**. See also **Emanuel Merck**, and **Richard Willstätter**.
- Wolff, H.**, estimation of alcohols by acetylation, A., ii, 165.
- Wolff, (Mlle) N.**, furfurylidene-1-methylcyclohexan-2-one and some of its derivatives and the mono- and difurfurylidene-cyclohexanones, A., i, 668.
- Wolfenstein, Richard**, preparation of hydrogenated 1-alkylpyridine-3-carboxylates, A., i, 365, 949.
- preparation of N-alkyl haloids of methyl pyridine-3-carboxylate [methyl nicotinate], A., i, 365.
- preparation of quaternary ammonium salts of pyridine-3-carboxylic acid alkyl esters, A., i, 861.
- preparation of alkyl salts of hydrogenated 1-alkylpyridine-3-carboxylates, A., i, 950.
- Wolfram, H.**, rapid extraction funnel, A., ii, 221.
- Woltjer, H. R.**, the Zeeman effect, A., ii, 102.
- Wood, C. J.**, and **P. P. Murdick**, stability of phthalate solutions as standards in hydrogen-ion work, A., ii, 735.
- Wood, Florence Mary**. See **Samuel Judd Lewis**.
- Wood, John Kerfoot**. See **George Ernest Collins**.
- Wood, Robert Williams**, fluorescence and photochemistry, A., ii, 334.
- atomic hydrogen and the Balmer series spectrum, A., ii, 673.
- spontaneous incandescence of substances in atomic hydrogen gas, A., ii, 759.
- Wood, Robert Williams**. See also **J. Steph. van der Lingen**.
- Woog, Paul**, relation between the molecular properties and the capacity for fixation of iodine of certain hydrocarbons, A., i, 101.
- the velocity of extension of thin layers of oils on the surface of a sheet of water, A., ii, 197.
- Wooten, Benjamin A.** See **Arthur St. C. Dunstan**.
- Woring, Pierre**, degradation of fatty acids in the animal organism, A., i, 90.
- permeability of the intestine to sucrose, A., i, 1214.
- Wormser, M.** See **Richard Lorenz**.
- Worrall, David E.**, the action of hydroxylamine and of hydrazine on the aryl monothioamides of ethyl acetylmalonate, A., i, 874.
- Wosnessenski, Serg.** See **Nikolai Schilov**.
- Wouseng, Sung**. See **René Looquin**.
- Wrangell, M. von**, phosphorus nutrition of plants, A., i, 1098.
- Wrede, Fritz**, the synthesis of disaccharides containing sulphur and selenium by combining two dextrose residues in the C5-position; some new derivatives of ζ -bromoglucose, A., i, 226.
- thioglucose, A., i, 525.
- microchemical estimation of carbon and hydrogen, A., ii, 316.
- Wright, Clarence D.** See **William O. Emery**.
- Wright, Floyd R.** See **Roger S. Hubbard**.
- Wright, Robert**, determination of the molecular weight of substances in alcoholic solution from the elevation of the flash-point, T., 2247.
- selective solvent action by the constituents of aqueous alcohol, T., 2251.
- Wright, W. C.** See **George Leslie Kelley**.
- Wrzesniewski, A. N.**, examination of the pentose in a new case of pentosuria, A., i, 1215.
- Wu, Hsien**, separate analyses of the corpuscles and the plasma, A., i, 483.
- a new colorimetric method for the estimation of plasma proteins, A., ii, 406.
- Wu, Hsien**. See also **Otto Folin**.
- Wülfig, Johann A.**, preparation of water-soluble compounds of diethylbarbituric acid and its homologues, A., i, 1066.
- Wulff, E.** See **Karl von Auwers**.
- Wulff, Peter**. See **Erich Tiede**.
- Wunnerlich, H.** See **Robert Schwarz**.
- Wurmser, (Mlle)**, the preparation of ammonium nitrate, A., ii, 499.
- Wurmser, René**, and **Raymond Jaquot**, the relation between the colloidal state and the physiological functions of protoplasm, A., i, 1221.
- Wurmser, René**. See also **Émile F. Terroine**.
- Wurzschnitt, Bernard**. See **Stefan Goldschmidt**.
- Wuyts, Henri**, and **A. Vangindertaelen**, the quadrivalence of tin in its mercaptides, A., i, 250.
- Wyckoff, Ralph W. G.**, crystal structures of the alkali haloids. I., A., ii, 214.

- Wyckoff, Ralph W. G.**, crystallographic and atomic symmetries of ammonium chloride, A., ii, 290.
 crystal structure of silver oxide, A., ii, 291.
 crystal structures of the hexammoniates of the nickel haloids, A., ii, 573.
 composition and crystal structure of nickel nitrate hexammoniate, A., ii, 573.
 a preliminary attempt to transmute lithium, A., ii, 642.
 symmetry and crystal structure of sodium hydrogen acetate, A., ii, 710.
 symmetry and crystal structure of zinc bromate hexahydrate, A., ii, 710.
 crystal structure of silver molybdate, A., ii, 765.
Wyckoff, Ralph W. G., and **Eugen Posnjak**, crystal structure of ammonium platinichloride, A., ii, 214.
 crystal structures of the cuprous haloids, A., ii, 295.
Wyckoff, Ralph W. G. See also **Eugen Posnjak**.

Y.

- Yabuta, Teihiro**, a new organic acid (kojic acid) formed by *Aspergillus oryzae*, A., i, 939.
Yamada, Nobuo, heat of transformation of austenite into martensite and of martensite into pearlite, A., ii, 475.
Yamaguchi, Yohei, determination of high temperatures by effusion of gases, A., ii, 470.
Yamazaki, Eiichi, the successive stages in the hydrolysis of triacetin, A., i, 422.
Yanagisawa, Hidekichi, estimation of cyanogen, A., ii, 667.
Yanagisawa, Hidekichi, and **Norikazu Takashima**, Japanese bird-lime. II., A., i, 652.
Yant, W. P. See **R. R. Sayers**.
Yeh, Yu Liang. See **Duncan A. MacInnes**.
Yngve, Victor. See **Arthur Becket Lamb**.
Yntema, L. F., and **B. Smith Hopkins**, observations on the rare earths. XI. The arc spectrum of yttrium, A., ii, 462.
Yoder, Lester. See **Arthur Wayland Dox**.
Yoshida, Usaburo, and **Shinsuke Tanaka**, the tungsten X-ray spectrum with a mica spectrometer, A., ii, 805.
- Young, Elrid Gordon**, the optical rotatory power of crystalline ovalbumin and serum-albumin, A., i, 279.
 the coagulation of protein by sunlight, A., i, 386.
Youngken, Heber W., and **Charles H. La Wall**, sand spur, *Cenchrus tribuloides*, L., A., i, 1100.
Yovanovitch, D., the chemical properties of mesothorium. II., A., ii, 712.

Z.

- Zaar, H.** See **Paul Jacobsen**.
Zaborowski, G., estimation of toluene, xylene, and benzene, A., ii, 877.
Zalkind, J. S., union of hydrogen with acetylene derivatives. XIV. Hydrogenation of phenylacetylene, A., i, 1134.
 union of hydrogen with acetylene derivatives. XV. Velocities of catalytic reactions, A., ii, 835.
Zambonini, Ferruccio, ardenite from Ceres, in the Ala valley (Piedmont), A., ii, 577.
Zanetti, Joaquin Enrique, J. R. Snyder, jun., and **M. Offner**, the formation of butadiene from ethylene, A., i, 977.
Zawidzki, Jan, equation for the velocity of reaction of hydrogen peroxide and potassium permanganate, A., ii, 207.
Zawodsky, Othmar. See **Robert Kreman**.
Zeckendorf, Kurt. See **Johannes Kerb**.
Zehnder, Ludwig, conclusions from the spherical form of the simplest atom, A., ii, 208.
Zeitfuchs, Eduard H. See **Charles August Kraus**.
Zelenka, J. See **Julius Stoklasa**.
Zelinsky, Nicolai E., and **E. F. Dengin**, synthesis of aminohydroxy-acids and the amino-acid from cyclopropyl methyl ketone [acetyltrimethylene], A., i, 1126.
Zellner, Julius, comparative plant chemistry. I. *Lythrum salicaria*, L., A., i, 799.
 plant chemistry. IV. *Juncus effusus*, L., A., i, 1098.
Zellner, Julius. See also **Rudolf Hasenöhrl**, and **Gisela Nowak**.
Zemplén, Géza, and **Alex. Hoffmann**, salicin thiocyanate and disalicin sulphide, A., i, 563.
Zemplén, Géza, and **Alphons Kunz**, new derivatives of salicin containing nitrogen and polynuclear hydroxybenzylamines, A., i, 564.

- Zerovtsev, Nik.** See *Nikolai Schilov*.
- Zerner, Ernest,** and *Robert Hamburger*, the action of silver compounds on yeast, A., i, 307.
- Zerweck, Werner.** See *Hans Fischer*.
- Zetzsche, Fritz.** See *Karl W. Rosenmund*.
- Zeynek, R.**, the preparation of chloro- and bromo-tyrosine and the analogous tyramines, A., i, 254.
- Ziegler, Karl,** and *Kurt Ochs*, halochromic phenomena with diarylstyrylcarbinols, A., i, 151.
poly-arylated vinyl carbinols and their derivatives. II. Diarylstyrylcarbinols and the products of their transformations, A., i, 1047.
- Ziegler, Karl.** See also *Karl von Auwers*.
- Ziegner, Heinrich.** See *Helmuth Scheibler*.
- Zielaskowski, Margarete.** See *Bruno Leichtentritt*.
- Zijp, C. van**, benzidine hydrochloride as a reagent for wood cells, A., ii, 94.
- Zilva, Sylvester Solomon**, conditions of inactivation of the accessory food factors, A., i, 488.
- Zilva, Sylvester Solomon,** and *Masataro Miura*, quantitative estimation of the fat-soluble factor, A., i, 194.
- Zimmerlund, G.** See *Hans von Euler*, and *Olof Svanberg*.
- Zimmermann, Kurt.** See *Erwin Ott*.
- Zimmermann, L.** See *A. Reis*.
- Zimmermann, Walther.** See *Hans Theodor Bucherer*.
- Zimmern, A.**, and *Edouard Salles*, spectrographic study of the de-intensifying of barium platinocyanide in the Villard effect, A., ii, 104.
- Zinke, Theodor,** and *H. Greune*, sulphamidophthalic acid and sulphimido-phthalic acid, A., i, 550.
- Zinke, Alois**, preparation of dihydroxy-*perylene*, A., i, 132.
preparation of *perylene*, A., i, 132.
- Zinke, Alois,** and *Rupert Dengg*, synthesis of 1:12-dihydroxyperylene and *perylene*, A., i, 1013.
- Zinke, Alois, Alfred Friedrich, Otto Johannsen,** and *Rudolf Richter*, resin constituents. VIII. The amyryns from elemi resin. II. α -Amyrin, A., i, 667.
- Zinke, Alois, Franz Hanselmayer,** and *Wilhelmine Ehmer*, resin constituents. IX. The decomposition of *d*-siarésinolic acid and lubanyl benzoate, A., i, 668.
- Zintl, E.**, and *H. Wattenberg*, potentiometric titration of copper, A., ii, 871.
- Zisch, W.** See *Fritz Haber*.
- Zivy, L.** See *Louis Jacques Simon*.
- Zizine, P.** See *A. Grigaut*.
- Zobel, H.** See *H. P. Kaufmann*.
- Zocher, H.**, sols with non-spherical particles, A., ii, 102.
- Zocher, H.** See also *Hans Kautsky*.
- Zoder, A.** See *Georg Weissenberger*.
- Zoff, Alfred.** See *Robert Kremann*.
- Zollinger-Jenny, Ernst**, improvements in the process of converting organic acids into esters, A., i, 914.
- Zolotarev, P. V.** See *P. P. Budnikov*.
- Zondek, S. G.**, potassium and radioactivity, A., i, 296.
the ionic equilibrium of cells; the physiology of sodium, A., i, 296.
the rôle of inorganic salts in the functions of the cell, A., i, 399.
- Zorn, Bernhard.** See *Emil Abderhalden*.
- Zorn, H.** See *Walter König*.
- Zschokke, H.**, and *L. Häuselmann*, estimation of free acid in acid aluminium sulphate solutions, A., ii, 397.
- Zsigmondy, Richard**, some fundamental conceptions of colloidal chemistry, A., ii, 129.
fundamental principles of colloidal chemistry. I. Electrical charge of the particles and new conceptions of micellæ, A., ii, 484.
- Zubkova, Sophie.** See *Alexis Bach*.
- Zuckmayer, Fritz**, preparation of hydrogenated 2-phenylquinoline-4-carboxylic acid, its homologues, and their salts, A., i, 574.
preparation of substituted derivatives of hydrogenated 2-phenylquinoline-4-carboxylic acid, and their salts, A., i, 952.
- Zunz, Edgard**, chemical composition of the thyroid gland, A., i, 397.
- Zunz, Edgard,** and *J. La Barre*, the action of the phosphatides on the coagulation of blood, A., i, 697.
- Zwaardemaker, Hendrik**, odour and chemical structure, A., i, 607.
potassium-calcium equilibrium in animal systems, A., i, 1213.
- Zweigle, A.** See *Alexander Gutbier*.
- Zwikker, J. J. Lijnst**, starch, A., i, 10.
the constitution of polysaccharides A., i, 230, 434.
space structural formulæ of chemical substances in general, and of some alkaloids in particular, A., i, 567.

INDEX OF SUBJECTS.

TRANSACTIONS AND ABSTRACTS.

1922.

(Marked T., and A., i and A., ii respectively.)

A.

- Abies pindrow*, oil from the leaves of (SIMONSEN), A., i, 753.
- Abietic acid**, preparation of, and its salts (FREELE), A., i, 739.
constitution of (MADINAVEITIA), A., i, 829; (ASCHAN, FONTELL, and SIMOLA), A., i, 1152.
and its isomerides and derivatives (RUZICKA and MEYER), A., i, 547.
crystallography and optical activity of (DUFFOUR), A., i, 829.
cetyl and myricyl esters (KAUFMANN and FRIEDEBACH), A., i, 740.
- Abietine**, constitution of (ASCHAN, FONTELL, and SIMOLA), A., i, 1152.
- Abietinol** (RUZICKA and MEYER), A., i, 830.
- Absorption spectra**. See under Spectra.
- Acacia cambagei* (gidgee wattle), calcium oxalate in (STEEL), A., i, 810.
- Acanthias vulgaris*, oil from the liver of (LEXOW), A., i, 490.
- Acenaphthene**, synthesis of (SCHONBERG), A., i, 27.
catalytic hydrogenation of (v. BRAUN and KIRSCHBAUM), A., i, 727.
equilibria of, with nitrobenzenes and nitrotoluenes (KREMAN and STRZELBA), A., i, 176.
compounds of, with chloranil and dichlorophthalic anhydride (PFEIFFER and FLATER), A., i, 342.
- Acenaphthene**, 5-amino-, derivatives of (FLEISCHER and SCHRANZ), A., i, 1142.
- Acenaphthene series** (MAYER and SCHONFELDER), A., i, 1166.
- Acenaphthenequinone phenylhydrazones** (CHARRIER and SPEIKANI), A., i, 772.
- Acenaphthenesulphonic acids**, 5-amino-, and their derivatives (FLEISCHER and SCHRANZ), A., i, 1142.
- Acenaphthenone**, 5-bromo-, and its oxime (MAYER and STEGLITZ), A., i, 742.
- Acenaphthenone-7-aldehyde**, and its derivatives (FRIEDLANDER, HERZOG, and v. VOSM), A., i, 764.
- 5-Acenaphthylglycine** (FLEISCHER and SCHRANZ), A., i, 1144.
- Acenaphthyl-5-sulphamic acid**, and its sodium salt (FLEISCHER and SCHRANZ), A., i, 1142.
- Acer ginnala* (Korean maple), crystalline tannin in the leaves of (PERKIN and UYEDA), T., 66.
- Acer saccharum*, organic acids of (SANDO and BARTLETT), A., i, 100.
- Aceritol**, and its tetra-acetyl derivative (PERKIN and UYEDA), T., 70.
- Acertannin**, and its acetyl derivative (PERKIN and UYEDA), T., 68.
- Acetaldehyde**, formation of, in fermentation by micro-organisms (NEUBERG and COHEN), A., i, 304.
preparation of (ADAMS and WILLIAMS), A., i, 222; (WOHL), A., i, 430.
preparation of, from acetylene (VOGT and NIEUWLAND), A., i, 110;
(OCHI, ONOZAWA, and TOKYO INDUSTRIAL LABORATORY), A., i, 523.
preparation of, from ethylidene diacetate (SOCIÉTÉ CHIMIQUE DES USINES DU RHÔNE), A., i, 915.
vapour pressure of (GILMOUR), A., i, 918.
equilibrium of phenylhydrazine and (HAMMARSTEN), A., ii, 830.
preparation of aldol from (GRÜNSTEIN), A., i, 111.
condensation of collidine with (KONDÔ and TAKAHASHI), A., i, 1177.
formation of vinyl alcohol from, in presence of alkali (EVANS and LOOKER), A., i, 102.
rôle of, in alcoholic fermentation (FERNBACH and SCHOEN), A., i, 203.

- Acetaldehyde** in normal urine (STEPP and FEULGEN), A., i, 495.
 detection of, in urine (STEPP), A., ii, 403.
 detection of, in urine in diabetes (FRICKE), A., i, 300.
 estimation of (SMITT), A., ii, 402.
 estimation of, by the "silver" method (FRICKE), A., ii, 326.
 estimation of, in presence of acetone (STEPP and FRICKE), A., ii, 236.
- Acetals**, preparation of (HAWORTH and LAPWORTH), T., 76.
 velocity of hydrolysis of (SKRADAL and SCHIFFER), A., ii, 134.
- Acetamide**, catalysis by acetic acid of the formation and hydrolysis of (NOYES and GOEBEL), A., i, 1127.
 kinetics of the reaction of acetic anhydride with (KREMAN, ZOFF, and OSWALD), A., ii, 748.
- Acetamide**, tribromo-, and bromocyanide (GUPTA and THORPE), T., 1900.
- Acetanilide**, (*antifebrin*), equilibrium of water and (SCHOORL and DE WEERD), A., i, 247.
 detection of, colorimetrically (EKKERT), A., ii, 169.
- Acetanilide**, *p*-chloro-, formation of, from *N*-chloroacetanilide (HARNED and SELTZ), A., ii, 631.
- Acetanilide- α -pyridinium** salts, and *m*- and *p*-nitro- (BARNETT and COOK), T., 794.
- Acetanilide-*p*-sulphonyl chloride**, preparation of (STEWART), T., 2558.
- Acetic acid**, preparation of (WOHL), A., i, 430.
 preparation of, and its anhydride, from ethylidene diacetate (SOCIÉTÉ CHIMIQUE DES USINES DU RHÔNE), A., i, 915.
 and *trichloro*-, viscosity and compound formation in mixtures of, with esters and with ketones (KENDALL and BRAKELEY), A., ii, 126.
 equilibria of, with their ethyl and methyl esters (SUDBOROUGH and KARVÉ), A., ii, 749.
 equilibrium of epichlorohydrin, water and (LEONE and BENELLI), A., ii, 744.
 decomposition of methyl oxalate by (TURNER and WILSON), A., i, 916.
 and its methyl ester, pyrogenic decomposition of (PEYTRAL), A., i, 219.
- Acetic acid**, *N*-acetatomercuri-*m*-nitrophenylammonium salt (KHARASCH, LOMMEN, and JACOBSON), A., i, 604.
- Acetic acid**, alkali salts, solubility of, in water (SIDEWICK and GENTLE), T., 1837.
 basic ferric salt, adsorption of radioactive substances by (BROWN), T., 1736.
 guaiacol-4:6-dimercuri-salt (MAMELI), A., i, 1083.
 lead salt (DUNDON and HENDERSON), A., ii, 552.
 complex compounds of (WEINLAND and STROH), A., i, 981.
 basic lead salt, precipitation of (LANGCKER), A., i, 315.
 mercuric salt, action of, on aminoazo-compounds (VECCHIOTTI), A., i, 478.
 action of carbon disulphide on (BERNARDI and ROSSI), A., i, 421.
 complex compound of mercuric sulphide and (MIOLATI), A., i, 982.
 action of nitrobenzene with (WIBAUT and JÜRGENS), A., i, 694.
 potassium salt, oxidation of, to potassium oxalate (EVANS and HINES), A., i, 803.
 sodium hydrogen salt, crystal structure of (WYCKOFF), A., ii, 710.
 thymolmercuric salts (MAMELI and MAMELI-MANNESSIER), A., i, 1080.
 uranyl salt, photolysis of (BAUR and REBMANN), A., ii, 337.
 action of, on organic compounds (KOPATSCHEK), A., i, 984.
- Acetic acid**, esters, dielectric constants of (JACKSON), A., ii, 252.
 esters of multivalent alcohols (SMITH), A., i, 915.
 bromoaminotolyl esters (RAIFORD and COUTURE), A., i, 931.
 and chloro-, bromoethyl and chloroethyl esters, preparation of (ALTWEGG and LANDRIVON), A., i, 315.
 cellulose ester, preparation of, from wood cellulose (HÄGGLUND, LÖFMANN, and FARBER), A., i, 323.
 solubility of, in alkali and alkaline-earth salts (SCHWEIGER), A., i, 324.
 detection and estimation of sulphuric acid and sulphacetates (ENTAT and VULQUIN), A., ii, 533.
- ethyl ester, physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.
 as an amphoteric electrolyte (v. EULER and SVANBERG), A., i, 219.
 velocity of hydrolysis of (ÅKERLÖF), A., ii, 134; (HARNED and PFANSTIEL), A., ii, 832.

- Acetic acid**, ethyl and methyl esters, effect of hydrogen-ion concentration on the stability of (KARLSSON), A., ii, 40.
- ethyl and isopropyl esters, action of alumina, titania, and thoria on (ADKINS and KRAUSE), A., i, 422.
- ethylidene ester, preparation of acetic acid and its anhydride and acetaldehyde from (SOCIÉTÉ CHIMIQUE DES USINES DU RHÔNE), A., i, 915.
- glycogen ester (PRINGSHEIM and LASSMANN), A., i, 634.
- β -naphthyl and phenyl ester pyridinium chlorides (BARNETT and COOK), T., 797.
- 2:5-dinitro-4-aminophenyl ester (REVERDIN and ROETHLISSBERGER), A., i, 538.
- Acetic acid**, detection of, and its homologues, by their sodium uranyl salts (BARLOT and BRENET), A., ii, 167.
- estimation of, in mixtures with butyric acid (NOTENBAART), A., ii, 791.
- Acetic acid**, bromo-, thymyl ester, compound of hexamethylenetetramine and (RIEDEL), A., i, 991.
- bromo-, and chloro-, sodium salts, separation of halogens from (v. EULER and FARLANDER), A., ii, 360.
- chloro-derivatives, penetration of cells by (CROZIER), A., i, 1090.
- chloro-, *p*-chloroacetylaminophenyl ester (BARNETT and COOK), T., 795.
- dichloro-, dissociation of (SCHREINER), A., ii, 736.
- trichloro-, action of, with copper (DOUGHY and FREEMAN), A., i, 427.
- lead diphenyl salt (GODDARD, ASHLEY, and EVANS), T., 981.
- cyno-, ethyl ester, sodium derivative, condensation of, with ethyl muconate (FARMER), T., 2015.
- Acetic anhydride**, kinetics of the reaction of acetamide with (KREMANN, ZOFF, and OSWALD), A., ii, 748.
- kinetics of the reaction of benzamide with (KREMANN, RÜSLER, and PENKNER), A., ii, 748.
- action of, on benzyldeneanthranilic acids (EKELEY, ROGERS, and SWISHER), A., i, 934.
- analysis of (RECLAIRE), A., ii, 532; (SAGE), A., ii, 593.
- Acetic-*p*- α -hydroxyisopropylbenzoic acid**, *o*-amino- (PHILLIPS), A., i, 955.
- Acetoacetanilide**, dithio-, and its dinitro-derivative (NAIK and AVASARE), T., 2594.
- Acetoacetic acid**, production of, from phenyl-lactic acids in the liver (MORI), A., i, 1217.
- production of, from urocanic acid in the liver (KONISHI), A., i, 1211.
- catalytic decomposition of (WIDMARK and JEFFSSON), A., ii, 437.
- ethyl ester, synthesis of (SCHEIBLER and ZIEGNER), A., i, 426.
- action of halogens on (SMITH), A., i, 318.
- compound of methyl azodicarboxylate and (DIELS), A., i, 774.
- Acetoaceto-*o*- and -*p*-toluidides**, and dithio-, and dinitro-derivative of the latter (NAIK and AVASARE), T., 2594.
- Aceto-*p*-anisidide**, amino-, and its derivatives (HILL and KELSEY), A., i, 1141.
- Acetobromoglucoses** (KARRER, NAGELL, and SMIRNOV), A., i, 253.
- Aceto-*p*-chloroanilide**, amino-, and its derivatives (HILL and KELSEY), A., i, 1141.
- Acetol**. See Acetylcarbinol.
- Aceto- α - and - β -naphthalide-*o*-pyridinium salts** (BARNETT and COOK), T., 795.
- Acetone** in commercial ammonia (BOUGAULT and GROS), A., ii, 709.
- partition of compounds between glycerol and (SMITH), A., ii, 270.
- equilibrium of phenylhydrazine and (HAMMARSTEN), A., ii, 830.
- oxidation of, with potassium permanganate (EVANS and SEFTON), A., i, 1115.
- pyrogenic decomposition of (PEYTRAL), A., i, 222.
- compounds of cresols and (BERL and SCHWIBEL; C. and W. v. RECHENBERG), A., i, 932.
- sugar derivatives of (IRVINE and PATTERSON), T., 2146; (FREUDENBERG and SVANBERG), A., i, 1116; (FREUDENBERG and BRAUNS), A., i, 1117.
- excretion of (BRIGGS and SHAFFER), A., i, 190.
- pharmacological action of (SALANT and KLEITMAN), A., i, 794.
- picrate, compounds of thiocarbamides and (TAYLOR), T., 2271.
- detection of, in alcohol (HARRISSON), A., ii, 667.
- detection of, in presence of caoutchouc (PITTARRELLI), A., ii, 95.
- detection of, in urine (TROISE), A., ii, 595.
- estimation of, in urine (LAX), A., ii, 326; (KADING), A., ii, 793.

- Acetone**, chloro-oximino-, compound of sodium hydrogen sulphite with (GAS-TALDI and BRAUNIZER), A., i, 626.
- Acetone substances** in blood in diabetes (FITZ), A., i, 192.
estimation of, in blood and urine (HUBBARD), A., ii, 234.
- Acetone- δ -benzylsemicarbazone** (WILSON, HOPPER, and CRAWFORD), T., 867.
- Acetone- S -benzylthiosemicarbazone** (WILSON and BURNS), T., 873.
- Acetonedicarboxylic acid**, and its ethyl ester (INGOLD and NICKOLLS), T., 1642.
- $\alpha\beta$ -Acetoneglucose**, $\alpha\beta$ -isopropulidene-glucose, phosphoric and sulphuric esters of (LEVENE, MEYER, and WEBER), A., i, 987.
- Acetone- α -naphthyl**, and its salts (KNOEVENAGEL), A., i, 1061.
- Acetoneoxalic acid**. See Acetylpyruvic acid.
- Acetone- δ - α -phenylethylsemicarbazones** (WILSON, HOPPER, and CRAWFORD), T., 869.
- Acetone- α -tetrahydro- α -naphthyl**, and its hydrochloride (KNOEVENAGEL), A., i, 1061.
- Acetonethiosemicarbazone**, silver and sodium derivatives, and their derivatives (WILSON and BURNS), T., 873.
- Acetonitrotetrahydronaphthalides** (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 124.
- Acetonitrotetrahydronaphthalides** (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 125.
- Acetonuria** (HUBBARD and WRIGHT; PORGES), A., i, 496.
in diabetes (HUBBARD and NICHOLSON), A., i, 969.
- Acetylmalonic acid**, diethyl ester, and its phenylhydrazone (GAULT and SALOMON), A., i, 428.
- Acetophenone**, condensation of (GAS-TALDI), A., i, 367.
benzoylphenylhydrazone (v. AUWERS, HILLIGER, and WULF), A., i, 1193.
picrate, compound of thiocarbamide and (TAYLOR), T., 2272.
- Acetophenone**, o -chloro- (FARBWERKE vorm. MEISTER, LUCIUS, and BRÜNING), A., i, 469.
- Acetophenoneanil**, and its salts (KNOEVENAGEL and GOOS), A., i, 751.
- Acetophenone-2-nitro-4-cyanophenylhydrazene** (MATTAAR), A., i, 252.
- Aceto- m -toluidide**, amino-, and its derivatives (HILL and KELSEY), A., i, 1141.
- Aceto- p -toluidide**, 2:6 dichloro-3-nitro- (DAVIES), T., 813.
2:3- and 2:5-dinitro- (SCOTT and ROBINSON), T., 845.
- Aceto- m -toluidides**, nitro- (BRADY, DAY, and ROLT), T., 529, 532.
- Aceto- o - and - p -toluidide- ω -pyridinium salts** (BARNETT and COOK), T., 794.
- isoAcetovanillone**, and its derivatives (SCHNEIDER and KRAFT), A., i, 749.
- α -Acetoxyacetic acid**, $\alpha\beta$ -trichloro-, iminoether hydrochloride (KÖTZ and RATHERT), A., i, 237.
- Acetoxyacetyl chloride**, chloro- (BLAISE and MONTAGNE), A., i, 520.
- 3-Acetoxy-2-acetyl-5-methylpyrrole-4-carboxylic acid**, ethyl ester (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- α -Acetoxyacrylamide**, $\beta\beta$ -dichloro- (KÖTZ and RATHERT), A., i, 236.
- Acetoxyanthraquinone**, 1-mono- and 1:5-di-hydroxy-, and the diboroacetate of the latter (DIMROTH and FAUST), A., i, 157.
- 7-Acetoxy-3-benzenesulphonylconmarin** (TROGER and BOLIE), A., i, 268.
- o -Acetoxybenzoic acid** (*acetylsalicylic acid*; *aspirin*), ω -pyridinium chloride (BARNETT and COOK), T., 797.
- α -Acetoxybenzoyl ethylhydrocupreine hydrochloride** (MERCK, DIEHL, and MAYEN), A., i, 47.
- o -Acetoxybenzoylquinine hydrochloride** (MERCK, DIEHL, and MAYEN), A., i, 46.
- Acetoxycamphane-*epi*-carboxylic acids**. See Acetylborneolcarboxylic acids.
- i -Acetoxydecoic acid**, and its amyl and methyl esters (GRÜN and WIRTH), A., i, 805.
- 3-Acetoxy-2:4-dimethoxyacetophenone**, and its phenylhydrazone (BRAND and COLLISCHONN), A., i, 452.
- 4'-Acetoxy-3':4-dimethylazobenzene** (JACOBSEN), A., i, 592.
- 2-Acetoxy-4:6-dimethylbenzonitrile** (v. AUWERS and SAURWEIN), A., i, 1032.
- 4'-Acetoxy-3':4-dimethylhydrazobenzene**, isomerisation of (JACOBSEN), A., i, 592.
- 4'-Acetoxyhydrazobenzene**, 4-bromo-, isomerisation of (JACOBSEN), A., i, 593.
- α -Acetoxymercuri- γ - o -anisylpropan- β -ol** (ADAMS, ROMAN, and SPERRY), A., i, 947.

- p*-Acetoxymereuridimethylaniline, *o*- and *m*-nitro- (KHARASCH and JACOBSON), A., i, 189.
- o*-Acetoxymereuriethylaniline, *p*-nitro- (KHARASCH and JACOBSON), A., i, 189.
- Acetoxymercuri-*m*-hydroxybenzaldehyde (HENRY and SHARP), T., 1059.
- o*-Acetoxymercurimethylaniline, *p*-nitro- (KHARASCH and JACOBSON), A., i, 189.
- 1-Acetoxymercurimethyl-1:2-dihydrobenzofuran (ADAMS, ROMAN, and SPERRY), A., i, 947.
- Acetoxymercurinitroacetanilides (KHARASCH, LOMMEN, and JACOBSON), A., i, 604.
- Acetoxymercurinitroanilines (KHARASCH, LOMMEN, and JACOBSON), A., i, 603.
- p*-Acetoxymercuri-*o*-nitrodiacetanilide (KHARASCH, LOMMEN, and JACOBSON), A., i, 604.
- 4-Acetoxymercuri-2-nitroresorcinol (RAIZISS and PROSKOURIAKOFF), A., i, 604.
- 3-Acetoxy-4-methoxycinnamic acid, α -amino-, benzoyl derivative, lactone (SPATH and LANG), A., i, 568.
- 2-Acetoxy-5-methoxyccumarone (SONN and FALKENHEIM), A., i, 1163.
- 2-Acetoxy-3-methoxyphenyl-*n*-octaldehyde (MAJIMA and TAKAYAMA), A., i, 263.
- β -Acetoxy- α -methyleinnamic acid, methyl ester (DIECKMANN and WITTMANN), A., i, 1157.
- 4-Acetoxy-4-methylhydrazobenzene, isomerisation of (JACOBSEN), A., i, 592.
- 3-Acetoxy-5-methylpyrrole-4-carboxylic acid, ethyl ester (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERDACH, and WILLIG), A., i, 858.
- 2-Acetoxy-naphthalene, 3:6-dibromo- (FRANZEN and STAUBLE), A., i, 450.
- 5-Acetoxy-1:4-naphthaquinone, 2:3-dibromo-8-hydroxy-, and 8-chloro-2:3-dibromo- (WHEELER and ANDREWS), A., i, 354.
- β -4-Acetoxyphenylpropionolactone, β -2-hydroxy- (LANGLEY and ADAMS), A., i, 1153.
- α -Acetoxypropionitrile, $\beta\beta$ -dichloro- and $\alpha\beta\beta$ -trichloro- (KÜTZ and RATHERT), A., i, 236.
- p*-Acetoxystyrene, ω -nitro- (ROSENMUND and PFANNKUCH), A., i, 1030.
- 2-Acetoxytoluene, bromoamino-derivatives, acetyl derivatives of (RAIFORD), A., i, 335.
- 7-Acetoxy-3-*p*-toluenesulphonylcoumarin (TRÖGER and BOLTE), A., i, 268.
- Acetyl chloride, preparation of chloro-derivatives of (CONSORTIUM FÜR ELEKTROCHEMISCHE INDUSTRIE), A., i, 315.
- chlorosulphonyl derivative (BLAISE and MONTAGNE), A., i, 520.
- Acetyl chloride, chloro-, syntheses with (BARNETT), A., i, 232.
- Acetylacetones, metallic (MORGAN, DREW and BARKER), T., 2433.
- 4 Acetyl-3-2'-acetoxy-4'-methoxyphenyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 4-Acetyl-3- β -acetoxy-naphthyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 4-Acetyl-3-*o*-acetoxy-*p*-tolylidihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 936.
- Acetylaminocanthraquinone, 2-chloro-, and its pyridinium salts (BARNETT and COOK), T., 796.
- Acetylaminocazobenzene, *p*-chloro-, and its pyridinium chloride (BARNETT and COOK), T., 795.
- Acetylaminonaphthalene-1-azobenzene, 4-chloro-, and its pyridinium chloride (BARNETT and COOK), T., 795.
- p*-Acetylbenzeneazo- β -naphthylamine, and its derivatives (CHARRIER and CHIPPA), A., i, 771.
- 9 *p*-Acetylbenzeneazo-10-phenanthrol (CHARRIER and DEMICHELIS), A., i, 772.
- N*-Acetyl-*N*-benzylglycine (SCHEIBLER and BAUMGARTEN), A., i, 655.
- N*-Acetyl-*N*-benzylidenebetaine (SCHEIBLER and BAUMGARTEN), A., i, 656.
- Acetylbenzylideneresorcinol (FABRE), A., i, 1148.
- Acetylborneolcarboxylic acids (BREDT), A., i, 937.
- δ -Acetyl-*n*-butyl alcohol, and its semi-acetal (BERGMANN and MIEKLEY), A., i, 618.
- Acetylbutylenes (BAUDRENGHIEN), A., i, 710.
- Acetylcarbinol (*acetol*), production of, from carbohydrates (BAUDISCH and DRUEL), A., ii, 664.
- oxidation of, with potassium permanganate (EVANS and HOOVER), A., i, 911.
- 4-Acetyl-3-*o*-carboxyphenyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.
- Acetylinnamantialdoximes (BRADY and THOMAS), T., 2107.

- 4-Acetyl-3-*p*-diethylaminophenyldihydro-2:4-benzoxazine-1-one** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 4-Acetyl-3-*mp*-dimethoxyphenyldihydro-2:4-benzoxazine-1-one** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 5-Acetyl-2:4-dimethylpyrrole-3-azobenzenesulphonic acid** (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILBIG), A., i, 858.
- 5-Acetyl-2:4-dimethylpyrrole-3-azonaphthalenesulphonic acid** (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILBIG), A., i, 858.
- 4-Acetyl-2:5-dimethylpyrrole-3-carboxylic acid, 4-chloro-, ethyl ester** (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 5-Acetyl-2:4-dimethylpyrrole-3-carboxylic acid, 5-chloro-, ethyl ester, and its phenylhydrazones** (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 5-cyano-** (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 3-Acetyl-2:4-dimethylpyrrole-5-glyoxylic acid, and its ethyl ester** (FISCHER, SCHNELLER, and ZERWECK), A., i, 1057.
- Acetyldiphenylamine, trichloro-** (STOLÉ), A., i, 762.
- 4-Acetyldiphenylamine, 5-bromo-2:4-dinitro-** (GIUA and ANGELETTI), A., i, 650.
- Acetyldiphenylamine- ω -pyridinium salts** (BARNETT and COOK), T., 796.
- Acetylene, formation of, in combustion of air flames** (HOFMANN and WILL), A., i, 1107.
- preparation of, acetaldehyde from** (VOGT and NIEUWLAND), A., ii, 110; (OCHI, ONOZAWA, and TOKYO INDUSTRIAL LABORATORY), A., i, 523.
- explosion of nitrogen and** (GARNER and MATSUO), T., 1729.
- condensations of** (KAUFMANN and SCHNEIDER), A., i, 245.
- hydrogenation of derivatives of** (ZALKIND), A., i, 1134; ii, 835.
- reduction of** (CHEVALIER and BOURCET), A., i, 801.
- action of, on pyrites** (STEINKOFF and HEROLD), A., i, 850.
- compounds of, with silver arsenate and phosphate** (OBERDOERFER and NIEUWLAND), A., i, 515.
- action of, on sodium derivatives of ketones** (LOCQUIN and WOUSENG), A., i, 617.
- Acetylene, narcotic action of** (WIELAND), A., i, 497.
- 4-Acetyl-3-*p*-ethoxyphenyldihydro-2:4-benzoxazine-1-one** (EKELEY, ROGERS, and SWISHER), A.; i, 935.
- 6-Acetyl-4-ethyl-*m*-xylene** (PHILIPPI, SEKA, and FROESCHL), A., i, 887.
- Acetylisoegenol, dibromo- and diiodo-** (RASTELLI), A., i, 1010.
- Acetylglucosylsalicylic acid** (H. P. and W. KAUFMANN, and GÖTTING), A., i, 253.
- Acetylhexoyldioxime, nickel compound** (PONZIO), A., i, 18.
- N*-Acetylhydrazobenzene, *N*-nitroso-** (GOLDSCHMIDT and EULER), A., i, 475.
- N*-Acetylhydrazo-*p*-toluene, and its compound with triphenylmethyl** (GOLDSCHMIDT and EULER), A., i, 476.
- β -Acetyllindene and its phenylhydrazones** (RUPP and MÜLLER), A., i, 41.
- 1-Acetylisatin-acetylhydrazones** (BORSCHE and MEYER), A., i, 54.
- Acetylmalonic acid, ethyl ester, arylthioamides of, action of hydrazine and of hydroxylamine with** (WORRALL), A., i, 874.
- menthyl ester** (SHIMOMURA and COHEN), T., 2054.
- 4-Acetyl-3-*o*-methoxyphenyldihydro-2:4-benzoxazine-1-one** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 4-Acetyl-3-*p*-methoxy-*n*-tolylidihydro-2:4-benzoxazine-1-one** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 5-Acetylmethylaminococnaphthene** (FLEISCHER and SCHRANZ), A., i, 1144.
- Acetylmethylfuran, and its derivatives** (PONZIO and RUGGERI), A., i, 630.
- Acetylmethylglyoxime, preparation of, and its diacetyl derivative** (PONZIO and RUGGERI), A., i, 629.
- 3-Acetyl-2-methyl-5- α - γ -tetrahydroxybutylpyrrole** (PAULY and LUDWIG), A., i, 954.
- N*-Acetyl-7-methylisatin** (HELLER, BENADE and HOCHMUTH), A., i, 1060.
- 2-Acetyl-5-methylpyrrole-4-carboxylic acid, 3-hydroxy-, ethyl ester and its semicarbazone** (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILBIG), A., i, 858.
- N*-Acetyl-*O*-methyl-*L*-tyrosine** (KARRER, GISLER, HORLACHER, LOCHER, MÄDER, and THOMANN), A., i, 814.
- 2-Acetyl- α -naphthol, autoxidation of, in alkaline solution** (FRIES and LEUE), A., i, 462.
- Acetyl- β -naphthylmethylamine-6-sulphonyl chloride** (MORGAN and ROOKE), A., i, 134.

4-Acetyl-3-3'- and 3-5'-nitro-2'-acetoxyphenyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.

Acetyl-2-nitro-3-anilino-*p*-phenetidine (REVERDIN and ROETHLISBERGER), A., i, 538.

Acetyl-*p*-nitrobenzaldehyd methyldihydro-azone (BRADY and McHUGH), T., 1652.

Acetyl-2:4-dinitrohydrobenzoin (BISHOP and BRADY), T., 2367.

4-Acetyl-3-nitro-3':4'-methylenedioxyphenyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.

4-Acetyl-3-2':4'-dinitrophenyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.

Acetyl-*n*-nonoyldioxime, nickel compound (PONZIO), A., i, 18.

Acetyloxyhaemin (KUSTER and GERLACH), A., i, 597.

Acetylpalmitoyldioxime, nickel compound (PONZIO), A., i, 18.

1-Acetylphenylcarbinol, production of, during fermentation, and its derivatives (NEUBERG and OHLE), A., i, 540.

4-Acetyl-3-*p*-phenylenebisdihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 935.

Acetyl-*m*-phenylenediamine, mono- and di-thio- (EDGE), T., 774.

2-*p*-Acetylphenyl- α - β -naphthatriazole, and its phenylhydrazones (CHARNIER and CRIPPA), A., i, 771.

Acetylpyruvic acid (*acetonoxalic acid*), ethyl ester, action of *p*-anisidine with (THIELEPAPE), A., i, 271.

monthyl ester (SHIMOMURA and COHEN), T., 2053.

Acetylquinizarin (DIMROTH and HILCKEN), A., i, 158.

λ -Acetylricinoleoacetylhydroxamic acid (NICOLLET and PELC), A., i, 644.

Acetylsalicylic acid. See *o*-Acetoxybenzoic acid.

Acetylstearyldioxime, nickel compound (PONZIO), A., i, 18.

Acetylsuccinic acid, γ -bromo- ethylester, preparation of (HIRST and MACBETH), T., 2175.

5-Acetyltetrahydroacenaphthene, and its derivatives (V. BRAUN and KIRSCHBAUM), A., i, 728.

1-Acetyl-2-thiohydantoin-5-propionamide (THIERFELDER), A., i, 326.

4-Acetyl-2:4:5-trimethoxyphenyldihydro-2:4-benzoxazine-1-one (EKELEY, ROGERS, and SWISHER), A., i, 938.

1-Acetyl-1:2:2-trimethylcyclopentane-3-carboxylic acid, and its methyl ester, and their derivatives (RUPE and JÄGGI), A., i, 841.

3-Acetyl-2:4:5-trimethylpyrrole, 3-chloro-, and 3-cyano- (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.

***N*-Acetyl-*l*-tyrosine** (KARRER, GISLER, HORLACHER, LOCHER, MÄDER, and THOMANN), A., i, 814.

Acetylvanillin (SIEVERS and GIVAUDAN & Co.), A., i, 38.

***Achillea millefolium* (milfoil)**, constituents of oil from (KREMER), A., i, 505.

Acid, $C_6H_5O_6$, from hydrolysis of the cyanide from methyl hydrogen diglycolate (ANSCHUTZ and JAEGER), A., i, 429.

$C_5H_8O_3$, and its calcium salt, from oxidation of *isopropylisocrotyl*-carbinol (KRESTINSKI), A., i, 1130.

$C_9H_8O_4N_2$, from hydrolysis of methyl isatogenate oxime (HELLER and BORSNECK), A., i, 369.

$C_{14}H_{11}N_6$, and its salts and esters, from dibenzhydrazide chloride and sodium azide (STOLLÉ and NETZ), A., i, 690.

$C_{18}H_{34}O_4$, from dehydration of trihydroxystearic acid (REINGER), A., i, 624.

$C_{18}H_{34}O_5$, from dehydration of sativic acid (REINGER), A., i, 624.

$C_{15}H_{22}O_6N_4$, and its derivatives, from cacotheline and bromine (LEUCHS, MILDBRAND, and LEUCHS), A., i, 1052.

$C_{20}H_{30}O_3$, and its phenylhydrazones, from oxidation of caoutchouc (KIRCHHOF), A., i, 563.

$C_{20}H_{32}O_3$, and its derivatives, from reduction of abietic acid (RUZICKA and MEYER), A., i, 547.

$C_{20}H_{32}O_3$, from reduction of abietic acid (RUZICKA and MEYER), A., i, 548.

$C_{21}H_{30}O_5$, from oxidation of *d*-siarasinolic acid (ZINKE, HANSELMAYER, and EHMER), A., i, 668.

$C_{31}H_{52}O_8$, obtained by reduction of pyroprosolanellie acid (WIELAND and SCHULENBURG), A., i, 346.

$C_{22}H_{44}O_2$, from the oxidation of paraffin (GRANACHER and SCHAUFELBERGER), A., i, 514.

***A*-Acid**. See α -Naphthol-5-sulphonic acid, 6-amino-.

***H*-Acid**. See α -Naphthol-3:6-disulphonic acid, 8-amino-.

Acids, relation of the affinity of, to their constitutional formulæ (ODDO), A., ii, 367.

conductivity of mixtures of, with acids, esters, and ketones (KENDALL and GROSS), A., ii, 33.

- Acids**, solubility of, in aqueous solutions of other acids (KENDALL and ANDREWS), A., ii, 34.
 taste of, and their swelling power (OSTWALD and KUHN), A., i, 84.
 device for removal of fumes of, during oxidation (FISCHLER), A., ii, 388.
 from sugars, oxidation of (GREINERT), A., i, 1111.
 estimation of, volumetrically (LIZIUS and EVERS), A., ii, 654.
 aliphatic, ionisation and surface activity of (WINDISCH and OSSWALD), A., ii, 123.
 partition of, between water and organic solvents (GORDON and REID), A., ii, 824.
 oxidation of, with chromic acid (SIMON), A., i, 803.
 catalytic decomposition of (MAILHE), A., i, 803.
 lower, reactions between salts of, and higher fatty acids (KNAPP and WADSWORTH), A., i, 219.
 saturated, amides and anilides of (ASANO), A., i, 437.
*di*basic, second dissociation constants of (DUBOUX), A., ii, 346.
 and their esters, adsorption and orientation of the molecules of, in solutions (KING and WAMPLER), A., ii, 741.
 carboxylic, electrolytic reduction of (INOUE), A., i, 36.
N-carboxylic, anhydrides of (FUCHS), A., i, 1152.
 fatty, preparation of, from mineral and tar oils (HARRIES), A., i, 514.
 and their glycerides, action of the brush discharge on (EICHWALD), A., i, 982.
 properties and molecular structure of thin films of (ADAM), A., i, 424; ii, 687.
 catalytic oxidation of (SALWAY and WILLIAMS), T., 1843.
 with branched chains, oxidation of (KAY and RAPER), A., i, 1093.
 with eighteen carbon atoms (NICOLET), A., i, 106; (NICOLET and COX), A., i, 320; (NICOLET and JURIST), A., i, 622; (NICOLET and PELC), A., i, 643.
 degradation of, in the animal organism (WORINGER), A., i, 90.
 estimation of, by distillation in steam (ARNOLD), A., ii, 324.
 estimation of, in blood (BLOOR, PELKAN, and ALLEN), A., ii, 598.
- Acids**, fatty, higher, formation of, by sugar fermentation (NEUBERG and ARINSTEIN), A., i, 91.
 electrical conductivity of anhydrides of (HOLDE and TACKE), A., ii, 109.
 reactions between salts of lower aliphatic acids and (KNAPP and WADSWORTH), A., i, 219.
 solid and liquid, separation of (MEIGEN and NEUBERGER), A., ii, 880.
 volatile, estimation of (WIEGNER and MAGASANIK), A., ii, 532.
 mineral, detection of, in vinegar (KLING and LASSIEUR), A., ii, 234; (KLING, LASSIEUR, and LASSIEUR), A., ii, 519.
 organic, and their salts, action of ultra-violet light on aqueous solutions of (JAEGER and BERGER), A., i, 316.
 bromination of, in the α -position (WARD), T., 1161.
 esterification of (ZOLLINGER-JENNY), A., i, 914.
 phenolic, formation of (LANGLEY and ADAMS), A., i, 1153.
 strong, estimation of, volumetrically, in presence of weak acids (KOLTHOFF), A., ii, 223.
 $\alpha\beta$ -unsaturated, esters of, formation of substituted succinic acids from (HIGGINBOTHAM and LAPWORTH), T., 49.
 weak, alcoholysis of the salts of (GOLDSCHMIDT, GORBITZ, HOUGEN, and PAHLE), A., ii, 135.
- Acid amides**. See Amides, acid.
- Acid chlorides**, reduction of (ROSENMUND, ZEIZSCHE, and ENDERLIN), A., i, 431; (SCHLIEWIENSKY), A., i, 938; (ROSENMUND), A., i, 939.
 action of zinc organic compounds on (MAUTHNER), A., i, 457.
- Acid haloids**, condensation products of (WEDEKIND and WEINAND), A., i, 234, 548.
- Acidity** (KELLER), A., ii, 131.
- Acidolysis** (SUDBOROUGH and KARVÉ), A., ii, 749.
- Acidosis** (VAN SLYKE; CULLEN), A., ii, 672.
- Aconite**, estimation of alkaloids in extract of (ASTRUC, CANALS, and BORDIER), A., ii, 404.
- Acraldehyde** (*acrolein*), interaction of aniline and (MANN), T., 2178.
- antiseptic action of (BERTHELOT), A., i, 903.

- Aspartic acid**, amino-, chloro-, chloroamino-, chloronitro-, and nitroamino- derivatives (FARBWERKE VORM. MEISTER, LUCIUS, and BRETNING), A., i, 468.
- 2,7-diamino-**, and its diacetyl derivative, chloroacetates and chloroacetanilide (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.
- diaminodithiol-** (WATSON and DUTT), T., 1943.
- Asordine-9-carboxylic acid**, derivatives of, and 2-chloro- (FARBWERKE VORM. MEISTER, LUCIUS, and BRUNING), A., i, 469.
- Acrylic acid**, and its derivatives, preparation of (VAN DER BURG), A., i, 220.
- Actinium**, disintegration series of (HAHN and MEITNER), A., ii, 185; (NEUBURGER), A., ii, 340.
- Actinomometers**, chemistry of (BENRATH, HESS, and OBLADEN), A., ii, 731.
- Acy groups**, migration of, from nitrogen to oxygen (RAIFORD and COUTURE), A., i, 931.
- Additive compounds** in solution and their influence on ionisation (KENDALL and GROSS), A., ii, 32.
- Address**, presidential (WALKER), T., 735.
- Adhesion** in solutions (SCHILOV and others), A., ii, 350.
- Adipenic acid** (CARRIERE), A., i, 319.
- Adipic acid**, $\alpha\beta$ -dibromo- $\gamma\delta$ -dihydroxy-, ethyl ester (CHANDRASENA and INGOLD), T., 1316.
- trihydroxy-**, salts of (KILIANI), A., i, 225.
- Adrenaline** (*suprarenine*; *epinephrine*), effect of, on blood pressure (RICHAUD), A., i, 891.
- effect of, in health (BJURE and SVENSSON), A., i, 970.
- Adsorption** (SCHILOV and others), A., ii, 350.
- theory of (EUCKEN), A., ii, 262, 550; (POLÁNYI), A., ii, 479.
- electrical theory of (MUKHERJEE; HARRISON), A., ii, 198.
- kinetic theory of (HENRY), A., ii, 740.
- relation between electrolytic dissociation and (RAKUZIN), A., ii, 619.
- effect of, on catalysis (GUICHARD), A., ii, 630.
- and hydrolysis (MOELLER), A., ii, 689.
- chemical (STADNIKOFF), A., ii, 619.
- negative (RAKUZIN and GÖNKE), A., ii, 821.
- Adsorption** by charcoal (HERBST), A., ii, 30; (FODOR and SCHÖNFELD), A., ii, 688.
- by sugar charcoal (BARTELL and MILLER), A., ii, 741.
- by fuller's earth (RIDEAL and THOMAS), T., 2119.
- of gases, calculation of (BERÉNYI), A., ii, 426.
- by charcoal (GUSTAVAR), A., ii, 479.
- of saturated vapours by charcoal (DRIVER and FIRTH), T., 2409.
- of ions (KOLTHOFF), A., ii, 197; (MUKHERJEE), A., ii, 689.
- of liquids by charcoal (HARRIS and EWING), A., ii, 123, 197.
- by precipitates (WEISER), A., ii, 262.
- of radioactive substances (EBLER and VAN RHYN), A., ii, 16.
- at solid surfaces (LANGMUIR), A., ii, 629.
- of solutions (OSTWALD and DE IZAGUIRRE), A., ii, 480.
- of dibasic acids and their esters (KING and WAMPLER), A., ii, 741.
- Aerinite**, analyses of (ORCEL), A., ii, 651.
- "**Aeronom**," use of the (RAUCH), A., ii, 316.
- Affinity**, conception of (PLANK), A., ii, 357.
- chemical, studies on (BRÖNSTED), A., ii, 132.
- residual, and co-ordination (MORGAN and SMITH), T., 160, 1956, 2866; (MORGAN and DREW), T., 922; (MORGAN and KING), T., 1723; (MORGAN, DREW, and BARKER), T., 2432; (MORGAN and MOSS), T., 2857; (MORGAN and LEDBURY), T., 2882.
- Agar**, composition of (SAMEO and SRAJEVIĆ), A., i, 231.
- swelling of (MACDOUGALL), A., i, 204.
- sulphur compounds in (NEUBERG and OHLE), A., i, 323.
- Agave techuagilla*, saponin from (JOHNS, CHERNOFF, and VIEHOVER), A., i, 797.
- Agmatine**, action of arginase in plants on (KIESEL), A., i, 413.
- Air**. See Atmospheric air.
- Alanine**, detection of, by crystallographic measurements (JAFTSCHNIKOV), A., ii, 864.
- Alanine-choline**, and its salts (KARRER, GISLER, HORLACHER, LOCHER MADER, and THOMANN), A., i, 814.
- dl*-Alanine-*N*-methyramidine, salts of (PYMAN), T., 2623.

- Albumin**, adsorption in solutions of (CLARK and MANN), A., ii, 550.
 action of sulphosalicylic acid on the swelling of (OSTWALD and KUHN), A., i, 598.
 effect of, on the conductivity of sodium chloride solutions (PALMER, ATCHLEY, and LOEB), A., i, 692.
 precipitation of (CRISTOL and SIMONNET), A., ii, 887.
 egg, effect of salts on the heat coagulation of (LOEB), A., i, 881.
 toxicity of (PENTIMALLI), A., i, 302.
 serum, crystalline, rotation of (YOUNG), A., i, 279.
Ovalbumin, crystalline, rotation of (YOUNG), A., i, 279.
- Albumin**, detection of, in urine (RENOUX), A., ii, 797.
 estimation of, microchemically (VALLEE and POLONOWSKI), A., ii, 406.
 estimation of, in cerebrospinal fluid (AYER and FOSTER), A., ii, 798.
 estimation of, in urine (GÉRARD), A., ii, 670.
- Albumoses**, diastatic action of (BIEDERMANN), A., i, 1200.
- Albumose-silver**, estimation of (HERZOG), A., ii, 798.
- Alcaptonuria**, metabolism in (GIBSON and HOWARD), A., i, 401.
- Alcohol**, $C_{20}H_{40}O$, and its derivatives, from saponification of bird-lime (NISHIZAWA), A., i, 652.
- Alcohols**, preparation of (BADISCHE ANILIN- & SODA-FABRIK), A., i, 218.
 catalytic preparation of, from aldehydes (MILLIGAN and REID), A., i, 337.
 electrolytic oxidation of (KOIZUMI), A., i, 979.
 dehydroxidation of (MULLER), A., i, 102.
 action of ammonia on (E. and K. SMOLENSKI), A., i, 234.
 introduction of the chloroethyl group into (CLEMO and PERKIN), T., 642.
 sensory stimulation by (IRWIN), A., i, 899.
 estimation of, by acetylation (WOLFF), A., ii, 166.
 aliphatic halogenated, preparation of aryl sulphonic esters of (v. KERESZLY and WOLF), A., i, 1181.
 ethylenic tertiary, conversion of, into primary ethylenic alcohols (LOCQUIN and WOUSENG), A., i, 710.
 primary, synthesis of (ODDO and BINAGHI), A., i, 314.
- Alcohols**, secondary, conversion of, into ketones (HUNT), A., i, 810.
 tertiary, conversion of, into aldehydes (LOCQUIN and WOUSENG), A., i, 810.
- Alcoholimetry** by means of miscibility temperatures (LÉVÊQUE), A., ii, 723.
- Alcoholysis**, determination of the degree of (GOLDSCHMIDT, GÖRBITZ, HOUGEN, and PAHLE), A., ii, 135.
- Aldehydes**, preparation of (LOCQUIN and WOUSENG), A., i, 810.
 from acid chlorides (ROSEN-MUND, ZETZSCHE, and FLÜTSCH), A., i, 39; (ROSEN-MUND, ZETZSCHE, and ENDERLIN), A., i, 431; (SCHLIEWIENSKY), A., i, 938; (ROSEN-MUND), A., i, 939.
 from mineral and tar oils (HARRIES), A., i, 514.
 catalytic preparation of alcohols from (MILLIGAN and REID), A., i, 337.
 condensation of primary aromatic arsines with (PALMER and ADAMS), A., i, 785.
 condensation of o-aminoazo-compounds with (FISCHER), A., i, 956.
 condensation of a-halogenated ketones with (BAUER and WERNER), A., i, 1034.
 conversion of, into acetals (HAWORTH and LAPWORTH), T., 76.
 action of phosphorus trichloride on (CONANT, MACDONALD, and KINNEY), A., i, 186.
 compounds of salvarsan with (RAIZISS and BLATT), A., i, 1079.
 interaction of thiocarbamides and (TAYLOR), T., 2267.
 action of yeast on (KUMAGAWA), A., i, 305.
 aromatic, in eucalyptus oils (PEN-FOLD), T., 266.
 kinetics of the reaction of ammonia with (DOBLER), A., ii, 435.
 halogenated, conversion of, into aldehydes and acids (KOTZ and RATHER), A., i, 236.
 estimation of (BOUGAULT and GROS), A., ii, 666.
 estimation of, with hydroxylamine (BENNETT and DONOVAN), A., ii, 535.
- Aldehyde-amines**, phototropy of (GALLAGHER), A., ii, 11.
- 9-Aldehydoacridine**, 2-chloro-, and its derivatives (FARBWERKE VORM. MEISTER, LUCIUS, and BRÜNING), A., i, 470.
- Aldehydobenzoic acids**, solubility of (SIDOWICK and TURNER), T., 2256.
- Aldehydosulicylic acids**, and their derivatives, and bromo- and nitro- (WAYNE and COHEN), T., 1022.

- Aldol** (*β -hydroxybutaldehyde*), preparation of (GRUNSTEIN), A., i, 111.
detection of, in urine in diabetes (FRICKE), A., i, 330; ii, 326.
- Aleuritic acid**, and its methyl ester and acetyl derivative (HARRIES and NAGEL), A., i, 522.
- Alexin**, physical chemistry of fixation of (KISS), A., i, 706.
- Alfalfa**. See Lucerne.
- Algæ**, marine, arsenic content of (JONES), A., i, 905.
- Alizarin** (1:2-dihydroxyanthraquinone), preparation of (DAVIES and SCOTTISH DYERS, LTD.), A., i, 355.
synthesis of (GRANDMOUGIN), A., i, 44.
constitution of the iron lakes of (BULL and ADAMS), A., i, 355.
boracetate, and its acetyl derivative (DIMROTH and FAUST), A., i, 157.
- Alizarin**, 3-mono- and 3:4-di-bromo- (DIMROTH, SCHULTZE, and HEINZE), A., i, 158.
- Alizarin colouring matters**, cobalt derivatives (MORGAN and SMITH), T., 160.
- Alizandipyridinium** bromides, and their derivatives (BARNETT and COOK), T., 1386.
- Alizarinpyridinium** bromide, bromo-, and its phenobetaine (BARNETT and COOK), T., 1385.
- Alizarinpyridiniumnitrolbetaines** (BARNETT and COOK), T., 1388.
- Alizarinquinone** ethoxybromide, methoxybromide, and 3-mono-, and 3:4-di-bromo- (DIMROTH, SCHULTZE, and HEINZE), A., i, 158.
- Alkali** bromates, preparation and properties of (BUELL and McCROSKY), A., ii, 146.
carbonates, estimation of, in presence of phenolphthalein (BONNIER), A., ii, 869.
chlorides, action of heated copper oxide with (HEDVALL and BOBERG), A., ii, 68.
fluorides, vapour pressure of (RUFF, SCHMIDT, and MUGDAN), A., ii, 818.
haloids, crystalline structure of (WYCKOFF), A., ii, 214; (POSNJAK and WYCKOFF), A., ii, 499.
hydroxides, infra-red absorption spectra of (GRANTHAM), A., ii, 244.
reactions between nitrogen oxides and (SANFOURCHE), A., ii, 762.
compounds of anhydro-sugars and (KARER, STAUB, and WALTJ), A., i, 229.
- Alkali iodides**, purity of (WINKLER), A., ii, 762.
metals, atomic and ionic radii of (LORENZ), A., ii, 211.
alloys of mercury with (BAIN and WITHROW), A., ii, 145.
nitrophenoxides (D. and A. E. GODDARD), T., 54.
hydrogen sulphites, equilibria of, in aqueous solutions (BALY and BAILEY), T., 1813.
ruthenium sulphites (REMY), A., ii, 857.
silver thiosulphites, compounds of ammonia with (JONSSON), A., ii, 57.
- Alkaligenesis** (LEE and TASHIRO), A., i, 1089.
- Alkaline earth carbonates**, formation of, from the oxides (HEDVALL and HEUBERGER), A., ii, 766.
hydrides (EPHRAIM and MICHEL), A., ii, 58.
metals, qualitative separation of (POLONOVSKI), A., ii, 720.
phosphors, photoelectric effect of (GÖGGL), A., ii, 732.
- Alkaloids**, space formule for (ZWICKER), A., i, 567.
constitution of, which give Vitali's reaction (HARDY), A., i, 948.
extraction of, and their characterisation (COLLEDGE), A., ii, 327.
in plants, biological significance of (CIAMICIAN and RAVENNA), A., i, 797.
bismuthi-iodides and mercuri-iodides of (FRANÇOIS and BLANC), A., i, 851.
salts of, with aminohydroxysuccinic acids (DAKIN), A., i, 430.
excretion of, in the saliva and in the stomach (HUBER), A., i, 1214.
toxicity of, effect of hydrogen-ion concentration on (CRANE), A., i, 404.
anhalonium. See Cactus.
belladonna. See Belladonna.
cactus. See Cactus.
calumba root (SPÄTH and BÖHM), A., i, 1174.
cinchona. See Cinchona.
pareira root (FALTIS and NEUMANN), A., i, 569.
quinine. See Quinine.
detection of, by means of their pierate crystals (NELSON and LEONARD), A., ii, 327.
detection of, with iodic acid (ROSENTHALER), A., ii, 327.
estimation of (HERZIG), A., ii, 538.
estimation of, volumetrically, indicators for (MCGILL), A., ii, 885.

- Alkyl haloids**, action of silver sulphate in sulphuric acid solution on (SCHILOV), A., i, 913.
 iodides, reactivity of, with sodium benzyloxide (HAYWOOD), T., 1904.
 sulphates, preparation of (DREYFUS), A., i, 517.
- Alkylamines**, preparation of (TAKEDA and KURODA), A., i, 272.
- Alkylamines**, hydroxy-, benzoic and tropic esters of, and their pharmacological action (v. BRAUN, BRAUNSDORF, and RATH), A., i, 759.
- cycloAlkylamines**, preparation of (MAILHE), A., i, 332.
- Alkylarylamines**, hydroxy-, esters of (SAUNDERS), T., 2667.
- Alkylation** (BAGGESGAARD-RASMUSSEN and WERNER), A., i, 104.
- Alkylhydrazones** (BRADY and McHUGH), T., 1648.
- N-Alkylideneamino-acids**, synthesis of, and their hydrogenation (SCHEIBLER and BAUMGARTEN), A., i, 655.
- Alkylidenecyanoacetic acids**, syntheses of (LAPWORTH and McRAE), T., 1699, 2741.
- Alligator oil**, constituents of (KOBAYASHI), A., i, 792.
- Allotropy**, theory of (COPISAROW), A., ii, 137.
 and the electromotive behaviour of metals (SMITS), A., ii, 110.
- Alloxan**, action of diazomethane on (HERZIG), A., i, 373.
- Alloys**, formation of (MANSURI), T., 2272.
 mechanical properties of (GUILLET and COURNOT), A., ii, 261.
 metallic, chemical and electrical behaviour of (TAMMANN), A., ii, 63;
 (JENGE), A., ii, 64.
- Allyl alcohol**, conversion of, into glyceryl bromo- and chloro-hydrins (READ and HURST), T., 989.
- Allylamino-oxides**, action of sodium hydroxide with (MEISENHEIMER, GREEKE, and WILLMERSDORF), A., i, 334.
- Allylarsinic acid**, and its salts (HOFFMANN, LaRoche & Co), A., i, 818, 926.
- 5-Allylimino-2-thio-3-phenyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole** (GUHA), A., i, 877.
- 5-Allylimino-2-thio-3-tolyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole** (GUHA), A., i, 877.
- 8-Allyloxybutylonitrile** (BRUYLANTS), A., i, 924.
- 6-Allyloxy-2-methylquinoline**, and its ethiodide (GUTEKUNST and GRAY), A., i, 951.
- 6-Allyloxy-2-p-nitrobenzylthiol-4-methylpyrimidine** (HORN), A., i, 875.
- Allylthiocarbimide**, estimation of, in mustard flour (LUCE and DOUGET), A., ii, 537.
- 2-Allylthiol-3-phenyl-5-anilinomethylene-4-glyoxalone** (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- Allyl-p-toluidine** (FOLDI), A., i, 732.
- Altitudes**, high, effect of, on absorption of oxygen (BAYEUX), A., i, 79.
- Alum**, action of, on animal glue (GUTBIER, SAUER, and SCHELLING), A., i, 783.
- Alumina**. See Aluminium oxide.
- Aluminates**. See under Aluminium.
- Aluminioxalic acid**, potassium salt, crystallography and optical properties of (KNAGGS), T., 2072.
- Aluminium**, spectrum of, in relation to the spectrum of glucinum (GLASER), A., ii, 675.
 electromotive behaviour of (SMITS), A., ii, 20; (SMITS and DE GRUYTER), A., ii, 21; (GUNTHER-SCHULZE), A., ii, 110.
 and its amalgams, electromotive properties of (MULLER and HOLZL), A., ii, 341.
 cathodic deposition of (KYROPOULOS), A., ii, 22.
 solubility of gases of (CZOCZRALSKI), A., ii, 646.
 formation of large crystals of (CARPENTER and ELAM), A., ii, 69.
- Aluminium alloys**, corrosion of (ROLLA), A., ii, 379.
 complex, thermal treatment of (GUILLET), A., ii, 69.
 analysis of (MENDE), A., ii, 162;
 (BRENNER), A., ii, 319; (COSTA-VET), A., ii, 528.
 with arsenic (MANSURI), T., 2272.
 with magnesium, heat of formation of (BILTZ and HOBORST), A., ii, 350.
- Aluminium compounds**, effect of, on growth of plants (CONNER and SEARS), A., i, 613.
- Aluminium salts**, reactions of sodium hydroxide with (GROBET), A., ii, 573.
 in soils (DENISON), A., i, 512.
 absorption of, by the roots of plants (STOKLASA, ŠEBOR, TÝMICH, and CWACHA), A., i, 502.
- Aluminium carbide**, preparation of (OHMANN), A., i, 2.
 chloride, union of aromatic nuclei by means of (SCHOLL and SCHWARZER), A., i, 331; (SCHOLL and SEER), A., i, 258, 336.
 condensations with (PUMMERER and BINAPFL), A., i, 24.

Aluminium haloids, densities of (BILTZ and WEIN), A., ii, 505.
 hydroxide, colloidal, preparation of (BRADFIELD), A., ii, 507.
 crystalline, dehydration of (MILLIGAN and MEAD), A., ii, 447.
 adsorption of proteins, enzymes, toxins, and sera by (RAKUZIN), A., i, 1199.
 solubility of, in water (MASSINK), A., ii, 299.
 nitrate, solubility of, in solutions of nitric acid (MILLIGAN), A., ii, 380.
 nitride, heat of formation of (FIOHTER and JENNY), A., ii, 615.
 oxide (*alumina*), adsorption of, by cellulose (TINGLE), A., i, 434.
 activation of, for decarboxylation or for dehydration (ADKINS), A., ii, 334.
 catalytic action of, on acetates (ADKINS and KRAUSE), A., i, 422.
 use of, as a mordant (BANCROFT), A., ii, 646.
 separation of, from calcium oxide (CHARRIOU), A., ii, 819.
 separation of ferric oxide and, from magnesium oxide (CHARRIOU), A., ii, 873.
 oxychloride, composition of sols of (ADOLF, PAULI, and JANDRASCHITSCH), A., ii, 149.
 selenate (MEYER and SPEICH), A., ii, 71.
 sulphate, equilibrium of potassium sulphate, water and (BRITTON), T., 982.
 estimation of free acid in acid solutions of (ZSCHOKKE and HAUSELMANN), A., ii, 397.
 chlorosulphoxide and sulphur chloride, preparation of (SILBERRAD), T., 1017.
Aluminates, constitution of (HEYROVSKY), A., ii, 771.
Aluminium organic compounds (FAILLEBIN), A., i, 119.
 with methylene iodide (THOMAS), A., i, 330.
Aluminium estimation and separation:—
 estimation of, volumetrically (KRAUS), A., ii, 87.
 estimation of, and its oxide, in the commercial metal (CHURCHILL; CAPPS), A., ii, 319.
 estimation of, in tungsten (V. and K. FROBOESE), A., ii, 397.
 estimation of sodium in, and in its oxide (GEITH), A., ii, 720.
 estimation and separation of (JANDER and WENDEHORST), A., ii, 529.

Aluminium estimation and separation:—
 estimation of, and its separation from iron (HAHN and LEIMBACH), A., ii, 873.
 separation of glucinum and (BRITTON), A., ii, 228.
 separation of iron and (CHALUPNY and BREISCH), A., ii, 588.
Amalgams. See Mercury alloys.
Amanita muscaria, isolation of muscarine from (KING), T., 1743.
Ambrosia artemisiifolia. See Ragweed.
 Ambrosterol, and its acetate (HEYL), A., i, 1224.
Amides, formation of, from alkylmalonic esters (DOX and YODER), A., i, 816.
 action of hydrogen peroxide on (OLIVERI-MANDALÀ), A., i, 544.
 acid, Hofmann's conversion of, to amines (BECKMANN, CORRENS, and LIESCHE), A., i, 544.
 acid optically active, catalytic racemisation of (MCKENZIE and SMITH), T., 1348.
Amidines, aromatic, condensation of alkyl malonic esters with (DOX and YODER), A., i, 374.
 Amine, $C_{20}H_{35}O_3N_4$, and its salts, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
Amines, preparation of, from alcohols and ammonia (E. and K. SMOLENSKI), A., i, 234.
 Hofmann's conversion of acid amides to (BECKMANN, CORRENS, and LIESCHE), A., i, 544.
 equilibria of triphenylmethane with (KREMANN, ODELGA, and ZAWODSKY), A., i, 131.
 oxidation of (GOLDSCHMIDT and EULER), A., i, 475; (GOLDSCHMIDT and RENN), A., i, 476, 478; (GOLDSCHMIDT and WURZSCHMIDT), A., i, 1139.
 decomposition of the vapour of (UPSON and SANDS), A., i, 1121.
 reactions of, with azodicarboxylic esters (DIELS), A., i, 774, 1194.
 action of o-halogenated ketones with (MAYER and FREUND), A., i, 865.
 action of ozone on (STRECKER and BALTES), A., i, 14.
 action of, on semicarbazones (WILSON, HOPPER and CRAWFORD), T., 866.
 alkylation of, with sulphonic esters (FÖLDI), A., i, 732.
 compound of tellurium tetrabromide with (LOWY and DUNBROOK), A., i, 446.
 hydroferriacyanides and hydroferrocyanides of (CUMMING), T., 1287.

Amines, aromatic, mercuric compounds of (KHARASCH and JACOBSON), A., i, 189.

action of sulphuryl chloride on (ELLER and KLEMM), A., i, 448.

preparation of *o*-sulphonic acids of (BRITISH DYESTUFFS CORPORATION, LTD., BADDILEY, PAYMAN, and WIGNALL), A., i, 448.

influence of *s*-trinitrobenzene on reactions involving (SUDBOROUGH and SHAH), A., ii, 757.

proteinogenous (HANKE and KOESSLER), A., i, 406, 408; ii, 322.

occurrence of, in the thyroid (SAM-MARTINO), A., i, 966.

physiological action of (ABELIN), A., i, 810.

secondary, dealkylation of, by phosphorus chloride (v. BRAUN and WEISMANTEL), A., i, 1150.

N-nitroso-derivatives (SCHMIDT and FISCHER), A., i, 822.

Amine oxides, optically active (MEISENHEIMER; MEISENHEIMER and BERNHARD), A., i, 813; (MEISENHEIMER and LOHSENER), A., i, 813, 822.

Amino-acids, synthesis of, in the animal organism (SHIPLE and SHERWIN), A., i, 492.

from hydrolysis of proteins (BURTON and SCHRYVER; JONES and JOHNS), A., i, 182.

refraction of (HIRSCH and KUNZE), A., i, 781.

diastatic action of (BIEDERMANN), A., i, 1200.

alkylation of the anhydrides of (KARRER, GRÄNACHER, and SCHLOSSER), A., i, 235.

action of, on sugars (GRÜNHUT and WEBER), A., i, 235.

action of micro-organisms on (HANKE and KOESSLER), A., i, 406.

retention and distribution of, in reference to urea formation (FOLIN and BERGLUND), A., i, 702.

distribution of, in blood (COSTANTINO), A., i, 81.

content of, in blood (OKADA and HAYASHI), A., i, 483.

in muscle (ROSEDALE), A., i, 492.

behaviour of, in vitally stained animals (KOTAKE, MASAI, and MORI), A., i, 1217.

deamination of, in the animal organism (KOTAKE), A., i, 1218.

degradation of, in the animal organism (WÖRINGER), A., i, 90.

phenylacetyl derivatives, fate of, in the organism (SHIPLE and SHERWIN), A., i, 1093.

Amino-acids, aliphatic, rate of reaction of nitrous acid with (DUNN and SCHMIDT), A., i, 1124.

aromatic, preparation of (LEWCOCK, ADAM, SIBERFIN, and GALBRAITH), A., i, 650.

detection and estimation of (ENGLAND), A., ii, 536.

estimation of, by alkalimetry (WILLSTÄTTER and WALDSCHMIDT-LEITZ), A., ii, 169.

estimation of, by the ninhydrin reaction (RIFFART), A., ii, 718.

estimation of, in blood (PETITJEAN), A., ii, 536.

estimation of, in feeding stuffs (HAMILTON, NEVENS, and GRINDLEY), A., ii, 92.

α -Amino-acids, conversion of alkylated malonic acids into (CURTIUS and SIEBER), A., i, 721.

Amino-alcohols (FOURNEAU and PUTAL), A., i, 639.

preparation of (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 813; (KARRER), A., i, 991.

Amino-aldehydes, preparation of (RADDE), A., i, 1164.

Amino-compounds, introduction of the chloroethyl group into (CLEMO and PERKIN), T., 642.

diphenylphosphoric acid as a reagent for (BERNTON), A., i, 1146.

aromatic, action of sulphates on (BUCHERER and WAHL), A., i, 464; (BUCHERER and ZIMMERMANN), A., i, 465.

estimation of, spectrophotometrically (MATHEWSON), A., ii, 882.

Aminohydroxy-acids, synthesis of (ZELINSKY and DENGIN), A., i, 1126.

α -Amino- β -hydroxy-acids, preparation of, from olefinocarboxylic acids (SCHRAUTH and GELLER), A., i, 1125.

Ammunes, metallic, physiological action of (OSWALD), A., i, 498.

complex (DUFF), T., 450.

Ammonia, formation of, during combustion of air flames (HOFMANN and WILL), A., i, 1137.

synthesis of (ANDERSEN), A., ii, 562; (PRING and RANSOME), A., ii, 639.

removal of the heat of reaction in the synthesis of (CLAUDE), A., ii, 284.

thermal dissociation of (FOXWELL), A., ii, 431.

specific heat of (PARTINGTON and CANT), A., ii, 191.

vapour pressure of (BERGSTROM), A., ii, 423.

- Ammonia**, vapour pressure curves of, with salt solutions (DAVIS, OLMSTEAD, and LUNDSTRUM), A., ii, 49.
 vapour pressure of liquid mixtures of *m*-xylene and (KRAUS and ZEITFUCHS), A., i, 725.
 solubility of, in water (NEUHAUSEN and PATRICK), A., ii, 264.
 solubility of, in lithium nitrate (DAVIS, OLMSTEAD, and LUNDSTRUM), A., ii, 56.
 adsorption of, by silica gels (DAVID-HEISER and PATRICK), A., ii, 262.
 kinetics of the reaction of aromatic aldehydes with (DOBLER), A., ii, 435.
 equilibrium of, with magnesium amalgam (LOOMIS), A., ii, 294.
 liquid, equilibrium of mixtures of sodium and (KRAUS and LUCASSE), A., ii, 764.
 equilibrium of sodium and tellurium in (KRAUS and CHIU), A., ii, 765.
 ignition of mixtures of, with air and oxygen (WHITE), T., 1688.
 reactivity of (BALY and DUNCAN), T., 1008.
 effect of gaseous impurities on the catalytic oxidation of (DECARRIERE), A., ii, 284.
 use of manganese dioxide as catalyst in oxidation of (PIGGOT), A., ii, 142.
 oxidation of, by persulphates in presence of copper (SCAGLIARINI and TORELLI), A., ii, 81.
 compounds of calcium haloids with (HÜTTIG), A., ii, 849.
 conversion of, into carbamide (MATIGNON and FRÉJACQUES), A., i, 723; ii, 272.
 action of, on copper (BASSETT and DURRANT), T., 2630.
 compounds of cuprous and thalious haloids with (BILTZ and STOLLENWERK), A., ii, 67.
 compounds of lead haloids with (BILTZ and FISCHER), A., ii, 851.
 compounds of magnesium haloids with (BILTZ and HÜTTIG), A., ii, 59.
 compounds of mercuric cyanide with (BRINKLEY), A., i, 724.
 compounds of alkali silver thiosulphates with (JONESON), A., ii, 57.
 action of, on reducing sugars (LING and NANJ), A., i, 631.
 in blood (NASH and BENEDICT), A., i, 191; (NASH), A., i, 483.
 excretion of, in urine, following administration of acids (KEETON), A., i, 300.
- Ammonia**, nesslerisation of solutions of (MAY and BOSS), A., ii, 312.
 commercial, acetone in (BOUGAULT and GROS), A., ii, 709.
 estimation of, in its salts (BURKARDT), A., ii, 865.
 estimation of, in blood (GAD-ANDRESEN), A., ii, 528.
 estimation of, in nitrogenous organic substances (FROIDEVAUX), A., ii, 454.
 estimation of, in urine (MESTREZAT and JANET), A., ii, 453.
- Ammoniates** (BILTZ and STOLLENWERK), A., ii, 67.
 solubility of (EPHRAIM and MOSIMANN), A., ii, 574.
- Ammonium** (SCHLUBACH and BALLAUF), A., ii, 55.
 calcium (BOTOLFSSEN), A., ii, 570.
- Ammonium salts**, absorption of, by soils (AARNIG), A., i, 1227.
 estimation of nitrogen in (MEURICE), A., ii, 225.
- Ammonium borate**, equilibrium of the formation of (SBOGI and FERRI), A., ii, 568, 764.
 antimoniodobromide (VOURNAZOS), A., ii, 651.
 carbonate, equilibrium of, with ammonium carbamate (FAURHOLT), A., ii, 272, 747; (WEGSCHEIDER), A., ii, 376.
 perchlorate, density of aqueous solutions of (MAZZUCHELLI and ANSELMI), A., ii, 376.
- chloride**, preparation of (MONDAIN-MONVAL), A., ii, 444, 642.
 crystallography and symmetry of (WYCKOFF), A., ii, 290.
 equilibrium in the systems of, with water and ferrous, cobaltous or nickel chloride (CLENDDINEN), T., 801.
- chloride and sulphate**, equilibrium of, with sodium chloride and sulphate and water (RIVETT), T., 379.
- platinichloride**, crystalline structure of (WYCKOFF and POSNJAK), A., ii, 214.
- silver anic chloride** (WELLS), A., ii, 449.
- stannichloride**, crystal structure of (DICKINSON), A., ii, 287.
- chlorite** (LEVI), A., i, 527.
- diamminotetraamminocobaltiate** (KNAGGS), T., 2078.
- fluosilicate**, crystal structure of (BOZORTH), A., ii, 499.
- nitrate**, properties of (MILLICAN, JOSEPH, and LOWRY), T., 959; (EARLY and LOWRY), T., 963; (PERMAN), T., 2473; (WURMSER), A., ii, 499.

- Ammonium nitrate**, stability of (FINDLAY and RONEBOURNE), A., ii, 291.
 decomposition of, by heat (SAUNDERS), T., 698.
 action of, on copper (BASSETT and DURRANT), T., 2630.
 nitrite, action of, on copper (DHAR), A., ii, 49.
 cadmium nickel nitrite (CUTTICA), A., ii, 449.
 potassium phosphates (CORELLI), A., ii, 287.
 sulphate, preparation of, from gypsum and ammonium carbonate (MATIGNON and FRÉJACQUES), A., ii, 570.
 melting point of (KATTWINKEL), A., ii, 445.
 equilibrium of copper sulphate, potassium sulphate, water, and (HAYAMI), A., ii, 748.
 equilibrium of glucinum sulphate, water, and (BRITTON), T., 2612.
 equilibrium of magnesium and potassium sulphates, water, and (WESTON), T., 1228.
- Ammonium organic compounds**, quaternary, influence of solvents on the velocity of formation of (HAWKINS), T., 1170.
- Ammonium radicle**, the free (WEITZ and LUDWIG), A., i, 365; (WEITZ and KÖNIG), A., i, 1186; (SCHLUBACH and BALLAUF), A., ii, 55.
- Ammonia-carbonic acids** (FRANKLIN), A., i, 440.
- Amniotic fluid**, composition of (LABAT and FAVREAU), A., i, 894.
- Amorphous condition**, distinction between the crystalline and the (HABER), A., ii, 553.
- Ampelopsis hederacea**, formation of anthocyanin in (JONESCO), A., i, 97.
- Amygdalin**, decomposition of, by emulsin (GIAJA), A., i, 185.
 biase from (HAWORTH and LEITCH), T., 1921.
- Amyl alcohol**, pure, preparation of, and conductivity of sodium iodide in it (KRAUS and BISHOP), A., ii, 813.
- isoAmyl alcohol**, electrolytic oxidation of (KOIZUMI), A., i, 979.
 preparation of β -methyl- Δ^{α} -butylene from (PUYAL), A., i, 617.
- n- and iso-Amyl fluorides** (SWARTS), A., i, 101.
- 4-isoAmylamino-5-carboxybenzenesulphonic acid**, and its salts (JACOBS and HEIDELBERGER), A., i, 75.
- isoAmylammonium** ruthenibromides (GUTHRIE and KRAUSE), A., i, 16.
- Amylase**, genesis of, in plants (PALADIN and POPOV), A., i, 614.
 chemical investigation of (SHERMAN), A., i, 66.
 from different sources, characteristics of (EFFRONT), A., i, 184, 1076.
 effect of filtration on (EFFRONT), A., i, 1076.
 conditions of activity of (BIEDERMANN and RUEHA), A., i, 65.
 inhibition of the action of (OLSSON), A., i, 390.
 effect of antiseptics on the action of (SHERMAN and WAYMAN), A., i, 282.
 effect of cobaltamines on (FUNK), A., i, 481.
 of cereals (BAKER and HULTON), T., 1929.
 pancreatic, effect of hydrogen-ion concentration on the action of (GROLL), A., i, 600.
 hydrolysis of starch by, in presence of amino-acids (SHERMAN and CALDWELL), A., i, 283.
- isoAmylboric acid** (KRAUSE and NITSCHKE), A., i, 22.
- Amylene**, action of selenium chloride on (BOORD and COPE), A., i, 421.
- sec.-isoAmyl methyl ketone**, chloro- (WIELAND and BETTAG), A., i, 1034.
- Amylocellulose**, constituents of (MALFITANO and CATOIRE), A., i, 527.
- Amylodextrin**, oxidation of (SYNIEWSKI), A., i, 1120.
- Amylphosphoric acid**, synthetic (SAMEO and MAYER), A., i, 921.
- Amyloses** (KARRER and BÜRKLIN), A., i, 435; (KARRER and FIORONI), A., i, 1119.
- 6-isoAmyloxy-2-methylquinoline**, and its ethiodide (GUTEKUNST and GRAY), A., i, 951.
- isoAmylpyridinium salts** (EMMERT and PARR), A., i, 179.
- α -**Amyranone**, and bromo-, and their oximes (ZINKE, FRIEDRICH, JOHANNSEN, and RICHTER), A., i, 667.
- α -**Amyrene**, and its dibromide (ZINKE, FRIEDRICH, JOHANNSEN, and RICHTER), A., i, 667.
- α -**Amyrenol benzoate**, and bromo- (ZINKE, FRIEDRICH, JOHANNSEN, and RICHTER), A., i, 667.
- α -**Amyrilene**, preparation of (VESTERBERG and WESTERLIND), A., i, 825.
- α -**Amyrin** from elemi resin (ZINKE, FRIEDRICH, JOHANNSEN, and RICHTER), A., i, 667.
- Amyrins**, extraction and separation of (VESTERBERG; VESTERBERG and WESTERLIND), A., i, 825.

- Amyrole**, isomeric, and their esters (PAOLINI), A., i, 754.
- Anemia** *in. carcioma* (ROBIN and BOURNIGAUULT), A., ii, 497.
- Anesthesia**, ether, effect of, on the acid-base balance of blood (VAN SLYKE, AUSTIN, and CULLEN; STEHLER, BOURNE, and BARBOUR), A., i, 1085.
- Anesthetics**, effect of, on the output of carbon dioxide (BODINE), A., i, 889.
- local, action of (FROMHERZ), A., i, 705.
- Analysis**, contradictions and errors in (HAHN, LEIMBACH, and WINDISCH), A., ii, 873.
- with the interferometer (BEČKA), A., ii, 715.
- use of membrane filters in (JANDER), A., ii, 457.
- catalytic (HAKOMORI), A., ii, 389.
- colorimetric, buffer solution for (MCILVAINE), A., ii, 78.
- dispersoid, of non-aqueous solutions (OSTWALD), A., ii, 199.
- electrolytic (KLING and LASSIEUR), A., ii, 587.
- electrolytic precipitation (KOLTHOFF), A., ii, 452.
- electro-volumetric, apparatus for (GOODE), A., ii, 307; (GARNER and WATERS), A., ii, 862.
- electrodes for (SHARP and MACDOUGALL), A., ii, 579.
- in alcoholic solution (BISHOP, KITTREDGE, and HILDEBRAND), A., ii, 308.
- with barium salts (KOLTHOFF), A., ii, 864.
- gravimetric (WINKLER), A., ii, 589.
- microchemical (BENEDETTI-PICHLER), A., ii, 656.
- Pregl's (SCHOELLER), A., ii, 160; (HOLTZ), A., ii, 521; (NICLOUX and WELTER), A., ii, 779.
- of organic compounds (WELTER), A., ii, 399.
- physico-chemical volumetric (DUTOIT and GROBER), A., ii, 578.
- qualitative (MACRI), A., ii, 779.
- separation of phosphoric acid in (BALAREFF), A., ii, 525.
- inorganic, use of resorcinol in (LAVOYE), A., ii, 779.
- quantitative, by centrifuge (ARRHENIUS), A., ii, 309.
- by measurement of the degree of supersaturation (HÖPPLER), A., ii, 863.
- refractometric (BEČKA), A., ii, 715.
- sedimentation. See Sedimentation.
- volumetric (JELLINEK and ENS), A., ii, 864.
- Analysis**, volumetric, use of amalgams in (KIKUCHI), A., ii, 519, 721; (KANÓ), A., ii, 529, 721.
- Andropogon Swarancusa*, terpene from the essential oil of (SIMONSEN), T., 2292.
- Anemone**, white, calcium succinate on leaves of (V. LIPPMANN), A., i, 811.
- Anemonins** (ASAHINA and FUJITA), A., i, 946.
- Anethole**, colour reaction of (ADLER), A., ii, 401.
- o*-Anethole, β -nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.
- β -Anetholealkamine, and bromo-, and their salts (TAKEDA and KURODA), A., i, 273.
- Anethole-2-imino-oxazolidine, bromo-, derivatives of (TAKEDA and KURODA), A., i, 273.
- Angelica root oil (ROURE-BERTRAND FILS), A., i, 847.
- Anhalamine, constitution of (SPÄTH), A., i, 163.
- synthesis of (SPÄTH and RÖDER), A., i, 853.
- Anhaline, constitution of (SPÄTH), A., i, 567.
- Anhalonidine, constitution of (SPÄTH), A., i, 163.
- Anhalonium alkaloids. See Cactus alkaloids.
- 4:5-Anhydro-3-acetoxymereuri-4-hydroxybenzaldehyde (HENRY and SHARP), T., 1058.
- Anhydrobiase, and its hexa-acetyl derivative (HESS), A., i, 13.
- Anhydro- α -dimethylaconitic acid (FARMER, INGOLD, and THORPE), T., 149.
- Anhydroecgonine, constitution of (GADAMER and JOHN), A., i, 167.
- ethyl ether, and its salts (GADAMER and JOHN), A., i, 675.
- Anhydroenneheptitol, diacetone derivative of (MANNICH and BROSE), A., i, 1118.
- Anhydroglucose, constitution of (KARRER and SMIRNOV), A., i, 228.
- Anhydroglycollic acid. See Diglycollic acid.
- Anhydrohematein. See Isohematein.
- Anhydro- β -*d*-hydroxydinaphthylbutane. See 7-Methyl-7-ethyl- α -dinaphthaxanthene.
- Anhydro- β -1:1-*d*-hydroxydinaphthylpentane. See 7-Methyl-7-*n*-propyl- α -dinaphthaxanthene.
- Anhydro- γ -1:1-*d*-hydroxydinaphthylpentane. See 7:7-Diethyl- α -dinaphthaxanthene.

- Anhydro- β -1:1-dihydroxydinaphthylpropane.** See 7:7-Dimethyl- α -dinaphthaxanthen.
- Anhydroindoxyl- α -anthranilide** (HELLER and BERNADE), A., i, 582.
- Anhydro-6:7-methylenedioxy-2:3-indeno-(3:1)-quinoline methoxyhydrate** (ARMIT and ROBINSON), T., 834.
- Anhydro-6:7-methylenedioxy-2:3-(3-methylindeno)(2:1)-quinoline methoxyhydrate** (ARMIT and ROBINSON), T., 834.
- Anhydropterococellin** (FORSTER and SAVILLE), T., 821.
- Anhydrosalicylic acid.** See Disalicylic acid.
- Anhydro-sugars, constitution of** (KARER and SMIRNOV), A., i, 228.
- Aniline, catalytic preparation of** (BROWN and HENKE), A., i, 445, 535, 586; (HENKE and BROWN), A., i, 1196. emulsions of water and, coagulation of (BHATNAGAR), A., ii, 268. oxidation of (GOLDSCHMIDT and WURZSCHMITT), A., i, 1139. interaction of acetaldehyde and (MANN), T., 2178. action of, with arsenious chloride (SCHMIDT), A., i, 285. action of, with sulphur monochloride (COFFEY), A., i, 132. and *p*-bromo-, compounds of tellurium tetrabromide with (LOWE and DUNBROOK), A., i, 446. arsenates (SCHMIDT), A., i, 285; (PATERNO), A., i, 731. hydrochloride, action of sodium formylacetone with (THIELEPAPE), A., i, 271. hydroferrocyanide (CUMMING), T., 1298.
- ψ -Aniline, and its picrate** (SCHMIDT), A., i, 778.
- Anilines, di- ω -chloropentachloro- and di- ω -chloronitro-** (GOLDSCHMIDT and STROHMENGER), A., i, 1004. nitro-, tautomerism of, and their mercurisation (KHARASCH, LOMMEN, and JACOBSON), A., i, 603. action of sodium hydrogen sulphite on (WEIL and WASSERMANN), A., i, 1006.
- Aniline- d -glucoside, and its tetra-acetyl derivative** (SABALITSCHKA), A., i, 247.
- Aniline- α -sulphonic acid, *p*-bromo-, *p*-iodo-, and *p*-nitro-, salts of** (SCOTT and COHEN), T., 2038.
- Anilinesulphenitrile, salts of** (OLIVERI-MANDALA), A., i, 1008.
- 5-Ahilitaceneaphthene, 2':4'-di- and 2':4':6'-tri-nitro-** (FLEISCHER and SCHRANZ), A., i, 1144.
- 6-Anilinoanthracenecumarin** (MINARFF and RIPPER), A., i, 168.
- Anilinoanthraquinones, amino-** (BADISCHE ANILIN- & SODA-FABRIK), A., i, 942.
- 4-Anilinobenzonitrile, 3-nitro-4-*p*-amino-** (MATTAAR), A., i, 252.
- o*-Anilinobenzoylbenzene, 8-chloro-1-nitro- and 2':4'-dinitro-** (MAYER and FREUND), A., i, 866.
- 5-Anilino-3 benzylthiol-1:2:4-triazole, and its dibenzyl derivative** (FROMM, KAYSER, BRIGLEB, and FÖHRENBACK), A., i, 378.
- 2-Anilino-5:5-dimethylthiazoline** (DER-SIN), A., i, 142.
- 2-Anilino-diphenyleneglycollic acid, 2-*op*-di- and 2-*oo*'*p* tri-nitro-, and their lead salts** (BRASS and FERBER), A., i, 356.
- Anilinoethylitaconic acid, ethyl ester** (CARRIÈRE), A., i, 319.
- 1-Anilino-4-hydroxynaphthalene, and 1-*p*-mono- and 1-*op*-di-chloro-** (KALLE & Co.), A., i, 824.
- 1-Anilino-4-hydroxynaphthalene-2-carboxylic acid** (KALLE & Co.), A., i, 824.
- Anilinohydroxysuccinic acid, anil and dianilide of** (DAKIN), A., i, 143.
- Anilinoitaconic acid, ethyl ester** (CARRIÈRE), A., i, 318.
- 3-Anilino-5-ketoxoxazole, and its salts and 3-*p*-bromo-** (WORRELL), A., i, 874.
- 3-Anilino-5-ketopyrazole, and its salts and derivatives, and 3-*p*-bromo-** (WORRELL), A., i, 874.
- 2-Anilino-3-mercapto-1:4-naphthaquinone** (FRIES and KERKOW), A., i, 577.
- o*-Anilinomethylacetophenones, *o*-2':4'-dinitro-** (MAYER and FREUND), A., i, 866.
- o*-Anilinomethylbenzophenones, *o*-2':4'-dinitro-** (MAYER and FREUND), A., i, 866, 867.
- 5-Anilinomethylene-4-glyoxalone, 2-thiol-** (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- 7-Anilino-2-methylxanthone** (v. DEM KNESEBECK and ULLMANN), A., i, 360.
- 3-Anilino-naphthalavinduline nitrate** (DUTT), T., 1958.
- 2-Anilino-1:4-naphthaquinone, 3-chloro-2-*p*-amino-, and its 2-*p*-acetyl derivative** (FRIES and KERKOW), A., i, 578.
- 8-Anilino-1:4-naphthaquinone, 2:3-di- and 2:3:8-tri-bromo-5-hydroxy-** (WHEELER and ANDREWS), A., i, 354.
- 1-Anilino- α -naphthiminazole, and its derivatives** (FISCHER and SCHWAPFACHER), A., i, 957.

- 6-Anilino- α -naphthol**, benzoyl derivative (BUCHNER and WAHL), A., i, 464.
- 3-Anilino-4-oximino-5-ketoxoxazole** (WONKALL), A., i, 874.
- 5-Anilinophenanthranaphthazine**, 4-nitro- (SIRCAR and DUTT), T., i, 949.
- 2-Anilinophenanthraquinone**, and 2-*op-di*- and 2-*op*-*p*-tri-nitro- (BRASS and FERRER), A., i, 356.
- β -Anilino-*p*-phenetolecarbamide** (SPECK-AN), A., i, 580.
- 6-Anilino-1-phenyl-2:3-dibenzylquin-oxalium** chloride, *aa*-*di*cyno- (DUTT and SEN), T., 2667.
- 7-Anilino-*N*-phenyl-2-methyl-pyridaz-oxenanthone** (V. DEM KNESECK and ULLMANN), A., i, 360.
- 5-Anilino-1-phenyltetrazole**, and its derivatives (OLIVERI-MANDALÀ), A., i, 1069.
- o*-Anilinophenyl-*p*-tolylsulphone**, and *p*-amino-, and *p*-nitroso- (HALBER-KANN), A., i, 1132.
- α -Anilinopropion-amide** and -anilide (DUBSKY, HOHER, and DINGEMANSE), A., i, 58.
- 4:6-Anilino-1:3:5:7-tetramethoxy-anthraquinone**, 2:6-*di*nitro- (HELLER and LINDNER), A., i, 1040.
- 5-Aniline-1:2:4-triazole**, 3-amino-, 3-hydroxy-, and 3-thio-, and their derivatives (FROMM, KAYSER, BRIG-LEB, and FÖHRENBACH), A., i, 378.
- Anilinoourazole** (ARNDT, MILDE, and TSCHENSCHER), A., i, 376.
- 1-Aniloethoxymethylamino-5-phenyl-1:2:3:4-tetrazole** (STOLLÉ and NETZ), A., i, 690.
- 1-Anilomethoxymethylamino-5-phenyl-1:2:3:4-tetrazole** (STOLLÉ and NETZ), A., i, 690.
- Anilothiourazole**, and its derivatives (ARNDT, MILDE, TSCHENSCHER, BIELICH, and ECKERT), A., i, 277.
- Anils** (*Schiff's bases*), hydrolysis of (REDELLEN and DANILOF), A., i, 147.
- Animals**, zinc in the feeding of (BER-TRAND and BENZON), A., i, 893.
- cold-blooded, effect of temperature on the oxygen consumption of (DIR-KFN), A., i, 1207.
- marine, of the Delphinidae, oil from (NAKATOGAWA and KOBAYASHI), A., i, 701.
- Animal heat**, effect of intermediary metabolites on the production of (LUSK and EVENDEN), A., i, 292.
- tissues, effect of heat and light on rate of reduction by (VALLOT), A., i, 87.
- Animal tissues**, fixation of calcium by (FREUDENBERG and GYÖRGY), A., i, 87, 291, 299, 608.
- zinc in (ROST), A., i, 87.
- detection and estimation of mor-phine in (WACHTEL), A., ii, 171.
- estimation of bromine and iodine in (DAMIENS), A., ii, 79.
- o*-Anisidine** hydroferrocyanide (CUM-MING), T., 1295.
- p*-Anisidine**, action of ethyl acetoneoxa-late with (THIELEPAPE), A., i, 271.
- Anisole** (*phenyl methyl ether*), ultra-violet absorption spectrum of (DURRANS), A., ii, 7.
- Anisole**, nitro-, equilibrium of the form-ation of, from nitrochlorobenzene (BLOM), A., ii, 38.
- s*-trinitro-, additive compound of pyri-dine and (GITA), A., i, 572.
- Anisotropy** of molecules (RAMAN), A., ii, 102.
- Anisylacetone**. See *p*-Methoxybenzyl methyl ketone.
- 1-*p*-Anisyl-4-benzylidenehydantoin**, and 2-thio- (HILL and KELSEY), A., i, 1141.
- o*-Anisylcarbamine cyanide** (FROMM and WENZL), A., i, 438.
- p*-Anisylidimethylcarbinol** (SKRAUP and FREUNDLICH), A., i, 540.
- 1-*p*-Anisylhydantoin**, and 2-thio- (HILL and KELSEY), A., i, 1141.
- 1-Anisylidenamino-5-anisyl-1:3:3:4-tetrazole** (STOLLÉ and NETZ), A., i, 690.
- 4-*p*-Anisyl-5-methyloxasolidone**, and its acetyl derivative (TAKEDA and KU-RODA), A., i, 273.
- 4-*p*-Anisyl-5-methyloxazoline** mer-captan, and its acetyl derivative (TAKEDA and KURODA), A., i, 273.
- α -Anisyl- β methylpropane** (LE BRAZI-DEC), A., i, 457.
- β -Anisyl- α -methylpropionic acid**, α -hydr-oxy-, and its amide and nitrile (LE BRAZIDE), A., i, 456.
- α -*o*-Anisyl-naphthylamine**, β -hydroxy- (WAHL and LANTZ), A., i, 823.
- p*-Anisylisocitronacetonitrile**, potassium and sodium salts (MEISENHEIMER and WEIBEZAHN), A., i, 177.
- p*-Anisylnitromethane** (MEISENHEIMER and WEIBEZAHN), A., i, 177.
- o*-Anisylsulphamic acid**, sodium salt (WEIL and MOSER), A., i, 444.
- Annam brier oil** (ROURE-BERTRAND FILS), A., i, 846.
- Annual General Meeting**, T., 722.
- Anode**, lead, use of, in electrolysis of solutions of alkali nitrites (JEFFERY), A., ii, 611.

- Anode**, silver, effect of telluric acid on the potential of (JIRSA), A., ii, 418.
- Anta**, toxicity of the venom of (STUMPER), A., i, 400.
- Antagonism** of ions (BRENNAR), A., i, 907.
- Anthochlor** (KLEIN), A., i, 1099.
- Anthocyanidins**, distribution of, in the coloured organs of plants (JONESCO), A., i, 797.
detection of the ψ -bases of, in plants (COMBES), A., i, 206.
- Anthocyanins** in plants (ANDREWS; KOZLOWSKI), A., i, 96; (JONESCO), A., i, 97; (COMBES), A., i, 412.
development of, in lily bulbs (MIRANDE), A., i, 1224.
tinctorial properties of (EVEREST and HALL), A., i, 265.
physiology of (COSTANTIN), A., i, 162.
- Anthracene**, melting and boiling points of (KIRBY), A., i, 27.
vapour pressure of (NELSON and SENSEMAN), A., i, 245.
velocity of dissociation of, in hexane solution (VOLMER and RIGGERT), A., ii, 336.
equilibria of nitrobenzenes with (KREMAN and MÜLLER), A., i, 131.
- Anthracene**, dibromo-, dichloro-, and dinitro-, preparation of (BARNETT, COOK, and GRAINGER), T., 2065.
9-nitro-, action of light on (BATTEGAY, BRANDT, and MORITZ), A., i, 1001.
- Anthracene series**, studies in the (BARNETT and COOK), T., 1876; (BARNETT, COOK, and GRAINGER), T., 2059.
- Anthrachryson** (1:3:5:7-tetrahydroxy-anthraquinone), triamino- and tetra-nitro-derivatives (HELLER and LINDER), A., i, 1040.
- Anthracite**, constitution of (GROUNDS), A., ii, 385.
- Anthracenmarin**, 6-chloro- (MINAEFF and RIPPER), A., i, 162.
- Anthraquinones**, tri- and tetra-hydroxy- (DIMROTH and HILCKEN), A., i, 159.
- Anthranilic acid**, cobalt salt (MORGAN and MOSS), T., 2866.
methyl ester, in grape juice (POWER and CHESNUT), A., i, 97.
- Anthranilic acid**, tetrabromo-, glycine and formalde of (GRANDMOUGIN), A., i, 55.
- Anthranil-10-pyridinium salts**, 9-hydroxy- and 9-nitro- (BARNETT, COOK, and GRAINGER), T., 2064.
- Anthraquinone**, preparation of (DOWNS), A., i, 1039.
- Anthraquinone**, and its derivatives, preparation of (CHEMISCHE FABRIK WORMS), A., i, 559.
vapour pressure of (NELSON and SENSEMAN), A., i, 245.
benzoyl derivatives, coloured reduction products of (SCHAARSCHMIDT), A., i, 355.
hydroxy- and sulphohydroxy-derivatives, manufacture of (SEGALLER, PEACOCK, and BRITISH DYESTUFFS CORPORATION, LTD.), A., i, 560.
carbamides and urethanes of (BATTEGAY and BERNHARDT), A., i, 1041.
mono- and di-alkyl thioethers and sulphones (REID, MACKALL, and MILLER), A., i, 154.
estimation of (NELSON and SENSEMAN), A., ii, 882.
- Anthraquinone**, diamino-derivatives, and their dibenzoyl derivatives (BATTEGAY and CLAUDIN), A., i, 43.
- dibromo-derivatives** (BATTEGAY and CLAUDIN; GRANDMOUGIN), A., i, 43.
- 1-chloro-2-amino-**, preparation of (FYFE and BRITISH DYESTUFFS CORPORATION, LTD.), A., i, 260.
action of dimethylaniline with, in presence of copper (KOPETSOHN and WIESLER), A., i, 844.
- 3-chloro-2-amino- and 3-chloro-1-bromo-2-amino-** (ATAK and SOUTAR), A., i, 259.
- 2-hydroxy-**, reduction of (PERKIN and WHATTAM), T., 289.
- 1:2-dihydroxy-**. See Alizarin.
- 1:5-dihydroxy-**. See Anthrarufin.
- 1:3:5:7-tetrahydroxy-**. See Anthrachryson.
- 1-hydroxylamino-**, potassium salt, 1-nitro-, and 1-nitroso- (BREISLER and JONES), A., i, 1166.
- Anthraquinones**, hydrogenated, preparation of (TETRALIN G. M. B. H.), A., i, 1039.
- Anthraquinones**, amino-, preparation of (THOMAS, DAVIES and SCOTTISH DYES, LTD.), A., i, 260.
preparation of aryl derivatives of (LÜTTINGHAUS and EIFFLAENDER), A., i, 355; (BADISCHE ANILIN- & SODA-FABRIK), A., i, 942.
- hydroxy-**, preparation of (SCHWENK), A., i, 153; (DAVIES and SCOTTISH DYES, LTD.), A., i, 354.
boric esters of (DIMROTH and FAUST), A., i, 155.
- Anthraquinone-2-aldehyde**, 1-amino- (CASSELLA & Co.), A., i, 942.

- Anthraquinone-1-carboxylic acid**, 6-chloro- (SCHOLL, SEER, and DAIMER), A., i, 258.
- Anthraquinonedihydro-5-phenylphenazine** (BADISCHE ANILIN- & SODA-FABRIK), A., i, 942.
- Anthraquinone-2:1:6:5-di-*ms*-phenylacridine** (MAYER and FREUND), A., i, 866.
- 1:2-Anthraquinonemethylisooxazole** (WILKE), A., i, 943.
- 1:2-Anthraquinoneisooxazole**, and 5:6:7:8-tetrachloro-, and 5-nitro- (WILKE), A., i, 943.
- Anthraquinone-2:1-*ms*-phenylacridine**, and its derivatives (MAYER and FREUND), A., i, 866.
- Anthraquinone-3:9-*ms*-phenylacridine**, 1-nitro- (MAYER and FREUND), A., i, 866.
- Anthraquinonesulphonic acid**, *mono*- and *di*-alkyl thioethers and sulphones (REID, MACKALL, and MILLER), A., i, 154.
- o*-Anthraquinonyl-1'-aminobenzoylbenzene** (MAYER and FREUND), A., i, 866.
- o*-Anthraquinonyl-1'-amino-5-methylacetophenone**, and *o*-4'-hydroxy- (MAYER and FREUND), A., i, 866.
- $\alpha\beta$ -Anthraquinonylisooxazoles**, and 7:8:9:10-tetrachloro-, and 7-nitro- (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 178.
- Anthraquino-1-thiazole**, 2-amino-, and its acetyl derivative, and 3-chloro-2-amino- (KOPETSCHEK and WIESLER), A., i, 868.
- Anthrarufin** (1:5-dihydroxyanthraquinone) diboroacetate (DIMROTH and FAUST), A., i, 157.
- Anthrax spores**, effect of formaldehyde on (HAILER), A., i, 408.
- Anti-bodies**, effect of ether on (FORSSMAN), A., i, 964.
- Antigens**, heterogenic (LANDSTEINER), A., i, 195.
- Antiketogenesis** (SCHAFER), A., i, 83.
- Antimonic acid**. See under Antimony.
- Antimony**, native, from California (BEHR), A., ii, 76.
- absorption spectrum of (LINDSAY), A., ii, 599.
- L* series spectrum of (DAUVILLIER), A., ii, 101.
- volatilisation of, by means of methyl alcohol (DUPARC and RAMADIER), A., ii, 640.
- Antimony alloys** with tellurium (DREIFUSS), A., ii, 503.
- Antimony compounds**, chemotherapy of (FARGHER and GRAY), A., i, 404.
- Antimony iodobromides**, complex mixed (VOURNAZOS), A., ii, 660.
- Antimony** :—
- Antimonic acid**, and its salts, constitution of (TOMULA), A., ii, 74.
- trisulphide**, polymorphism of (WILSON and McCOSKEY), A., ii, 154.
- estimation of, iodometrically (NIKOLAI), A., ii, 585.
- estimation of available sulphur in (LUFF and PORRITT), A., ii, 225.
- pentasulphide**, preparation of stable sols of (v. HAHN), A., ii, 856.
- Antimony organic compounds**, aromatic (SCHMIDT), A., i, 482, 1203.
- Antimonytartaric acid**, salts of, and their chemotherapy (FARGHER and GRAY), A., i, 705.
- Antimony detection, estimation, and separation** :—
- detection of (SABALITSCHKA and SCHMIDT), A., ii, 531.
- detection of, by mirrors (SCHEUCHER), A., ii, 526.
- estimation of, in antimonates (TOMULA), A., ii, 74.
- estimation of, in copper and brass (EVANS), A., ii, 231.
- estimation of, volumetrically, in red brass (MÜCK), A., ii, 722.
- estimation of, and its separation from copper, lead and tin (KLING and LASSIEUR), A., ii, 86.
- separation of, from arsenic and tin (HAHN), A., ii, 877.
- Antioxidants** (SEYEWETZ and SISLEY), A., ii, 623.
- Anti-oxygens** (MOUREU and DUFRASSE), A., i, 250; (SEYEWETZ and SISLEY), A., ii, 623.
- Antipyrine** (1-phenyl-2:3-dimethyl-5-pyrazolone), salicylate, solubility of (KOLTHOFF), A., i, 471.
- Antiseptics**, action of, in relation to chemical constitution (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.
- effect of, on amylases (SHERMAN and WAYMAN), A., i, 282.
- Antitoxins**, chemical nature of (SALKOWSKI), A., i, 1216.
- Aphthalite** from Kilauea (WASHINGTON and MERWIN), A., ii, 386.
- isoApiole**, β -nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.
- Apophyllenic acid**, constitution of (MUMM and GORTSCHALDT), A., i, 862.
- Apples**, odorous constituents of (POWER and CHESNUT), A., i, 96.
- estimation of pectin in (CARR and HAYNES), A., ii, 401.

- Apples**, disappearance of free sulphurous acid in preserved juice of (WARCOLLIER and LE MOAL), A., i, 415.
- Aquopentamminecobaltiselenate**. See under Cobalt.
- Aquopentamminocobaltic 2:4-dinitro- α -naphthoxide** (MORGAN and KING), T., 1727.
- Aquopentamminocobaltic dipicrylamine** (MORGAN and KING), T., 1728.
- Aquopentapyridinemagnesium iodide** (SPACU), A., i, 859.
- Aquotetramminecobaltichloride selenate**. See under Cobalt.
- Arabic acid**, and its mixtures with gelatin (TIRBAKX), A., i, 633.
- Arabinose**, action of lactic bacteria on (FRED, PETERSON, and ANDERSON), A., i, 201.
- Arachidic acid**, estimation of (PRITZKER and JUNGKUNZ), A., i, 208.
- Arachin**, digestibility of (JONES and WATERMAN), A., i, 893.
- Arachis oil**, catalytic decomposition of (MAILHE), A., i, 712.
- β -Aracyl- α -dialkylaminoethylbutyric acids**, esters, preparation of (FABERWERKE VONM. MEISTER, LUCIUS, & BRÜNING), A., i, 639.
- Aragonite**, crystal structure of (HUGGINS), A., ii, 643.
equilibrium of calcium with, in aqueous solution (BÄCKSTRÖM), A., ii, 849.
- Aralia montana***, saponins from the leaves of (VAN DER HAAR), A., i, 1168.
- Araligenin**, and its derivatives (VAN DER HAAR), A., i, 1169.
- Ardennite**, from Ceres in Piedmont (ZAMBONINI), A., ii, 577.
- Arginase** in plants (KIESSEL), A., i, 413.
- Arginine**, degradation of, in plants (KIESSEL), A., i, 413.
- Argon**, free paths of electrons in (TOWNSEND and BAILEY), A., ii, 494, 836.
ionisation of, by electron collision (HORTON and DAVIES), A., ii, 811.
excitation and ionisation potentials of (HERTZ), A., ii, 733.
specific heat and heat of vaporisation of (MATHIAS, CROMMELIN, and ONNES), A., ii, 472.
rectilinear diameter of (MATHIAS, ONNES, and CROMMELIN), A., ii, 565.
- Aromadendral**, true nature of (PENFOLD), T., 266.
- Aromadendrin**, constitution of (NISHIKAWA and ROBINSON), T., 839.
- Aromatic compounds**, application of the octet theory to (CROCKER), A., i, 927.
- Aromatic compounds**, elimination of hydrogen from (SCHOLL and SEER), A., i, 258, 336; (SCHOLL and NEUMANN), A., i, 261; (SCHOLL and SCHWARZER), A., i, 331.
catalytic nitration of, by means of mercuric nitrate (DAVIS), A., i, 818.
reduction of (DE POMMEREAU), A., i, 825.
polynuclear, molecular configuration of (CHRISTIE and KENNER), T., 614.
heterocyclic (ARMIT and ROBINSON), T., 827.
- Arsenates**. See under Arsenic.
- Arsenic**, polymorphism of (LASCH-TSOHENKO), T., 972.
melting point of (HEIKE), A., ii, 25.
volatilisation of, by means of methyl alcohol (DUPARC and RAMADIER), A., ii, 640.
toxicology of (TARUGI), A., ii, 84.
- Arsenic alloys** with aluminium (MAN-SURI), T., 2272.
with zinc (HEIKE), A., ii, 60.
- Arsenic compounds**, magnetic susceptibilities of (PASCAL), A., ii, 564.
in marine algae (JONES), A., i, 905.
effect of, on plant growth (STEWART), A., i, 1221; (STEWART and SMITH), A., i, 1222.
poisoning by. See Poisoning.
- Arsenic carbide** (DE MAHLER), A., i, 101.
trichloride, solubility of, in concentrated hydrochloric acid (TREADWELL and MUSSLER), A., ii, 768.
trihydride (arsine), toxicity of (FÜHNER), A., i, 499.
- Arsenic acid**, estimation of (ROSENTHALER), A., ii, 584.
- Arsenates**, periodicity of reactions between thiosulphates and (FORBES, ESTILL, and WALKER), A., ii, 271.
- Arsenious acid**, reducing action of (KOHN), A., ii, 148.
estimation of, in presence of nitrous acid (KLEMENC), A., ii, 865.
- Arsenites**, periodicity of reactions between thiosulphates and (FORBES, ESTILL, and WALKER), A., ii, 271.
effect of iron salts on the iodometric estimation of (MELVILL), A., ii, 784.
- Arsenic trisulphide**, coagulation of sols of, by electrolytes (BURTON and MACINNES), A., ii, 180.
colloidal, flocculation of (BOUTANIC and VUILLAUME), A., ii, 498.
pharmacological action of (MENEGHETTI), A., i, 301.

- Arsenic trisulphide**, precipitation of, from arsenates (REEDY), A., ii, 225.
estimation of, iodometrically (NIKOLAI), A., ii, 585.
- Arsenic organic compounds** (MANN and POPE), T., 1754; (STEINKOPF and SCHWEN), A., i, 71, 72, 118; (JACOBS and HEIDELBERGER), A., i, 73, 74; (LEWIS and CHEETHAM; JOHNSON and ADAMS; LEWIS and HAMILTON) A., i, 187; (RAIZISS and BLATT), A., i, 1079; (STEINKOPF, DONAT, and JAEGER), A., i, 994; (BART), A., i, 1201, 1202.
with aniline (SCHMIDT), A., i, 285; (PATERNO), A., i, 731.
- Arsenious acids** (QUICK and ADAMS), A., i, 600.
- Arsenic detection, estimation, and separation**:—
detection of (KOLTHOFF; ROMIJN), A., ii, 455.
detection of, microchemically (PIUTTI and BOGGIO-LENA), A., ii, 584.
detection of, by mirrors (SCHEUCHER), A., ii, 526.
estimation of, colorimetrically (CHOUCHAK), A., ii, 526.
estimation of, iodometrically, in presence of copper (KOLTHOFF and CREMER), A., ii, 84.
estimation of, in presence of nitrates and nitrites (GRAHAM and SMITH), A., ii, 314.
estimation of, in silicate rocks (HACKL), A., ii, 159.
estimation and separation of (MOSER and EHRLICH), A., ii, 315.
separation of germanium and (MÜLLER), A., ii, 320.
separation of, from molybdenum, tungsten, and vanadium (MOSER and EHRLICH), A., ii, 314.
- Arsenious acid**. See under Arsenic.
- Arsenobenzene**, 3:3'-diamino-4:4-dihydroxy-, synthesis of (BART), A., i, 1202.
preparation of derivatives of (SPEYER-HAUS), A., i, 481.
dihydrochloride. See Salvarsan.
- Arseno-compounds**, aromatic, preparation of (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 962.
- Arsenopyrite**, crystal structure of (HUGGINS), A., ii, 651.
- Arsine**. See Arsenic trihydride.
- Arsine hydroxybromides**, alkylated and arylated, decomposition of (STEINKOPF and SCHWEN), A., i, 71.
- Arsines**, primary aromatic, condensation of, with aldehydes (PALMER and ADAMS), A., i, 785.
- Arsines**, tertiary, action of cyanogen bromide on (STEINKOPF, DONAT, and JAEGER), A., i, 994.
tertiary phenylated, action of cyanogen bromide on (STEINKOPF and SCHWEN), A., i, 72.
- Arsinetri-1-piperidinium chloride**, (LEONARD), A., i, 363.
- Arsinic acids** (QUICK and ADAMS), A., i, 600.
aromatic, synthesis of (BART), A., i, 1201, 1202.
- Arsinic acids**, amino-, aromatic, preparation of (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- Araphenamine**. See Salvarsan.
- Artemisia afra**, constituents of (GOODSON), A., i, 1099.
- Aryl alkyl ethers**, preparation of (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 934.
preparation of water-soluble derivatives of (ELEKTROCHEMISCHER WERKE, BOSSHARD, and STRAUSS), A., i, 913.
- β -Arylhydroxylamines**, preparation of (BRAND and STEINER), A., i, 536.
- Asarone**, β -nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.
- Asparagus officinalis**, carbohydrates in the seeds of (CAKE and BARTLETT), A., i, 504.
- Aspartic acid**, dipeptides of (RAVENNA), A., i, 180.
- Aspartic acids**, hydroxy-. See Suocinic acids aminohydroxy-.
- Aspergillus glaucus**, action of, on glycerol (TRAETTA-MOSCA and PRETI), A., i, 91.
- Aspergillus niger** (*Sterigmatocystis nigr*), acid fermentation produced by (MOLLIARD), A., i, 611.
effect of the nutritive media on the composition of (TERROINE, WURMSER, and MONTANÉ), A., i, 1220.
energy yield in the growth of (TERROINE and WURMSER), A., i, 706.
effect of hydrogen-ion concentration on the amylase of (FUNK), A., i, 796.
lipase from (SCHENKER), A., i, 208.
toxicity of nitrophenols towards (PLANTEFOL), A., i, 204.
formation of oxalic acid and ammonia in growth of, on peptone (BUTKEWITSCH), A., i, 707.
- Aspergillus oryzae**, formation of kojic acid by (YABUTA), A., i, 939.
- Assimilation** of nitrates and nitrites (BAUDISCH), A., i, 194.
plant. See Plants.

- Asymmetric compounds**, synthesis of, (PIRAK), A., i, 810.
 synthesis of, by means of enzyme action (ROSENTHALER), A., i, 480.
- Atmospheric air**, identification of the lines of, in spark spectra (MERRILL, HOPPER, and KEITH), A., ii, 802.
 positive ray analysis of (THOMSON), A., ii, 565.
 Joule-Thomson coefficients for (KEYES), A., ii, 24.
 velocity of sound in (GRÜNEISEN and MERKEL), A., ii, 190.
 propagation of flame in mixtures of gases with (PAYMAN and WHEELER), T., 363; (WHITE), T., 1244, 1688, 2561.
 ignition of mixtures of methane and (MASON and WHEELER), T., 2079.
 liquid, use of, as a refrigerant (FERGUSON), A., ii, 614.
 apparatus for analysis of (GUTHRIE), A., ii, 78.
 estimation of injurious acids in (LAMBEIS), A., ii, 390.
 estimation of carbon dioxide in (RAUCH), A., ii, 316; (NOYONS), A., ii, 868.
 apparatus for estimation of carbon dioxide in (LUNDEGÅRDH), A., ii, 719.
 estimation of nitrogen oxides in (ALLISON, PARKER, and JONES), A., ii, 313.
 estimation of petroleum vapour in (FRITZMANN and MACJULEVITSCH), A., ii, 877.
- Atoms**, structure of (DAUVILLIER; BURY), A., ii, 43; (ZEHNDER), A., ii, 208; (NEUBURGER), A., ii, 203, 365; (CHWOLSON), A., ii, 209; (BOHR), A., ii, 277, 363; (MARSH), A., ii, 277; (CREHORE), A., ii, 438; (HUGGINS), A., ii, 632; (LODGE), A., ii, 701.
 and refraction (WASASTJERNA), A., ii, 2, 491.
 in relation to spectra (KING), A., ii, 277.
 and Röntgen spectra (COSTER), A., ii, 491, 677; (DAUVILLIER), A., ii, 678.
 relation between magnetism and (WEISS), A., ii, 23; (CABRERA; OXLEY), A., ii, 469.
 electronic structure of (FOKKER), A., ii, 137; (DAUVILLIER), A., ii, 559; (PEASE), A., ii, 737; (HUGGINS), A., ii, 838.
 periodic structure of (ALLEN), A., ii, 758.
- Atoms**, models of (HUGHES), A., ii, 632; (VAN VLECK), A., ii, 838; (NEUBURGER), A., ii, 839.
 for lecture demonstration (KING), A., ii, 705.
 Bohr's model of (BRILLOUIN), A., ii, 438.
 Neuburger model of (VALERAN; MEITNER; NEUBURGER), A., ii, 702.
 Whittaker model of (PEDDIE), A., ii, 633.
 constitution of, and structure of crystals (NIGGLI), A., ii, 36.
 emission of light by (FÖRSTERLING), A., ii, 729; (SEELIGER), A., ii, 804.
 radiations from (GEHRCKE), A., ii, 210.
 thermodynamic equilibrium of electrons, radiation and (DE BROGLIE), A., ii, 249.
 dimensions of (RICHARDS), A., ii, 42.
 in diamond type crystals (PEASE), A., ii, 634.
 and their ionisation potentials (EVE), A., ii, 364.
 in relation to the properties of elements and compounds (CLARK), A., ii, 634.
 radii of (SAHA), A., ii, 278; (HUGGINS), A., ii, 634.
 stability of the nuclei of (HARKINS and MADORSKY), A., ii, 490; (HARKINS), A., ii, 702.
 induced polarity of (LAPWORTH), T., 416; (KERMACK and ROBINSON), T., 427.
 binding of electrons by (NICHOLSON), A., ii, 544.
 motion of electrons in (TOWNSEND and BAILEY), A., ii, 277.
 disintegration and transformation of (MEITNER), A., ii, 15.
 collisions between molecules and (FRANCK), A., ii, 464.
 electrical doublet theory of the attraction between (KLEEMAN), A., ii, 366.
 quantum mechanism within (WHITTAKER), A., ii, 632; (EWING; HOUSTOUN), A., ii, 633.
 energy of the linkings between (PADOA), A., ii, 27.
 stopping power and atomic number of (GLASSON), A., ii, 250.
- Atomic number**, scattering of β -rays and (GLASSON), A., ii, 183.
 and stopping power (GLASSON), A., ii, 250.
- Atomic weight** of boron (BAXTER and SCOTT), A., ii, 285; (HÖNIGSCHMID and BIRCKENBACH), A., ii, 641.
 of bromine (MOLES), A., ii, 140; (BAXTER), A., ii, 870.

- Atomic weight of carbon** (MOLES, A., ii, 51.
of chlorine (GLEDITSCH and SAMDAHL), A., ii, 281.
of glucinum (HÖNIGSCHMID and BROCKENBACH), A., ii, 214.
of isohelium (NEUBURGER), A., ii, 365.
of lanthanum (BAXTER), A., ii, 298 ; (HOPKINS and DRIGGS), A., ii, 770.
of mercury (BRÖNSTED and v. HEVESY), A., ii, 645.
of yttrium (FOGG and JAMES), A., ii, 297.
- Atomic weights, report of the Committee on** (BAXTER), A., ii, 366.
calculation of (REYCHLER), A., ii, 279.
determination of, from gas densities (BAXTER), A., ii, 370.
determination of, by vapour density measurements (MAGNUS and SCHMID), A., ii, 260.
purity of silver for use in determination of (BAXTER and WOODMAN), A., ii, 376 ; (BAXTER), A., ii, 377.
" **Atophan.**" See 2-Phenylquinoline-4-carboxylic acid.
- Atoxyl** (*sodium p-aminophenylarsinate*), crystallography of (GILTA), A., i, 961.
Atrolactic acid, fate of, in the body (KAY and RAPER), A., i, 1093.
l-Atrolactinamide, preparation and rotation of (MCKENZIE and SMITH), T., 1356.
- Atropic acid**, fate of, in the body (KAY and RAPER), A., i, 1093.
Atropine, estimation of (HARDY), A., ii, 796.
estimation of, in blood-serum (VAN DER HEYDE), A., ii, 669.
- Aucubin** from seeds of *Melampyrum arvense* (BRIDEL and BRAECKE), A., i, 209.
comparison of rhinanthin and (BRIDEL and BRAECKE), A., i, 1168.
- Augite** from Hawaii (WASHINGTON and MERWIN), A., ii, 220.
- Austenite**, heat of transformation of, into martensite (YAMADA), A., ii, 475.
- Autoclave** (GROSS), A., ii, 561.
Autolysis (BRADLEY), A., i, 896.
- Autoxidation** (MOUREU and DUFRAISSE), A., i, 250, 824.
- Autunite**, calcium uranium, analyses of (HENRICH), A., ii, 516.
- Azelaic acid**, preparation of (ASAHINA and ISHIDA), A., i, 520.
electro-synthesis of (CARMICHAEL), T., 2545.
- Azens** (STAUDINGER), A., i, 238.
- Azides**, formation of (OLIVERI-MANDALÀ), A., i, 473.
lecture experiments with (BROWNE and HOEL), A., ii, 840.
- Azidoanisylideneanilhydrazide** (STOLLÉ and NETZ), A., i, 690.
- Azidodibenzhydrazide** (STOLLÉ and NETZ), A., i, 690.
- Asiminobenzene**. See 1:2:3-Benztriazole.
- α 8-Asiminonaphthalenecarboxylic acid**, methyl ester (DIELS and WACKERMANN), A., i, 1069.
- p-Azoanisole**, oxonium salts of (KEHRMANN and VAN DER LAAR), A., i, 384.
- Azobenzene**, catalytic preparation of (HENKE and BROWN), A., i, 586
1196.
electrochemical oxidation of (FICHTER and JAECK), A., i, 62.
hydrochloride, reaction of aromatic hydrocarbons with (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1196.
- Azobenzene**, amino-, action of mercuric acetate on (VECCHIOTTI), A., i, 478.
decachloro- (GOLDSCHMIDT and STROHMENGER), A., i, 1006.
- Azobenzoyl-p-aminophenyl ether** (FICHTER and JAECK), A., i, 62.
- Azobenzoyl-p-nitrophenyl ether** (FICHTER and JAECK), A., i, 62.
- Azo-colouring matters** (JACOBS and HEIDELBERGER), A., i, 74.
calculation of the colour of (MOIR), T., 1555.
from bilirubin (FISCHER and BARRENSCHEEN), A., i, 278.
from naphthalene (VOROSHCHEV), A., i, 956.
action of phenylhydrazine-bisulphite mixture on (BUCHERER and ZIMMERMANN), A., i, 465.
vat (MUKERJI), T., 2879.
estimation of, electrometrically (JONES and LEE), A., ii, 239.
apparatus for estimation of intermediates for manufacture of (ATKINSON), A., ii, 237.
- Azo-colouring matters, o-hydroxy-**, preparation of (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 385.
- Azo-compounds**, action of, with aromatic hydrocarbons and aluminium chloride (PUMMERER and BINAPFL), A., i, 24 ; (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1196.
coupling of, with 1:8-naphthasultam and its derivatives (KÖNIG and KÖHLER), A., i, 879.

- Azo-compounds**, *o*-amino-, condensation of aldehydes with (FISCHER), A., i, 956.
 amino-, and hydroxy- (CHARRIER), A., i, 769.
 hydroxy-, constitution of (PUXEDDU and GENNARI), A., i, 587.
 hydrolysis of alkali salts of (PUXEDDU), A., i, 589.
- Azodicarboxylic acid**, esters, reactions of amines and enols with (DIELS), A., i, 774.
 ethyl and methyl esters, additive compounds of β -naphthylamine with, and their oxidation products (DIELS and WACKERMANN), A., i, 1068.
- Azoimide** (*hydrazoic acid*), lecture experiments with (BROWNE and HOEL), A., ii, 840.
 addition of, at contiguous double linkings (OLIVERI-MANDALA), A., i, 1069.
- Azomethine compounds** (MORGAN and REEVES), T., 1.
- 1:1'-Azonaphthalene-4:4'-disulphonic acid** (KALLE & Co.), A., i, 867.
- 1:1-Azo-ar-tetrahydronaphthalene** (SCHROETER, KINDERMANN, DIETRICH, BEYSLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 123.
- Azotobacter**, effect of hydrogen-ion concentration on the growth of (GAINIEY and BATCHELOR), A., i, 1096.
- Asoxyamides** (PIERONI), A., i, 1071.
- Asoxybenzene**, 3-nitro-4:4'-dihydroxy- (ANGELI, BIGIAYI, and CARRARA), A., i, 879.
- Asoxyphenol**, 3-nitro-4-hydroxy- (ANGELI, BIGIAYI, and CARRARA), A., i, 879.
- Asoxyphenols** (ANGELI, BIGIAYI, and CARRARA), A., i, 878.
- o*-Asoxyphenylacetic acid**, and its esters (NEBER), A., i, 546.
- 1:1-Asoxy-ar-tetrahydronaphthalene** (SCHROETER, KINDERMANN, DIETRICH, BEYSLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 123.
- B.**
- Babingtonite** from Japan (WATANABE), A., ii, 651.
- Bacilli**, diphtheria, effect of manganous chloride on formation of (WALBUM), A., i, 795.
 action of the oxy-acids of selenium and tellurium on (JOACHIMOGLU and HIROSE), A., i, 406.
 influenza, growth of (JACOBY and FRANKENTHAL), A., i, 302.
- Bacilli**, paratyphoid-enteritidis, fermentation of tetralose by (KOSER), A., i, 407.
 pyrocyanic, decomposition of sugars by (AUBEL), A., i, 201.
 tetanus, growth of (DERNEY and ALLANDER), A., i, 303.
 Timothy grass, fat metabolism of (STEPHENSON and WHETHAM), A., i, 500.
 tubercle, bacteriology of (LONG), A., i, 1095.
 action of tribromoxyleneol on (DUBOC), A., i, 972.
 utilisation of dextrose by (GAMBLE and HERRICK), A., i, 902.
 typhoid, production of carbon dioxide by (NICHOLS), A., i, 501.
- Bacillus butyricus***, production of carbon dioxide by (BROOKS), A., i, 201.
- Bacillus coli***, effect of colloidal substances on the growth of (LABES), A., i, 902.
 experiments on disinfection with (VERMAST), A., i, 406.
 Endo's reaction with (FERNÁNDEZ and GARMENDIA), A., i, 405.
- Bacillus diphtheriae***, reversal of reaction in media by the growth of (WOLF), A., i, 1218.
- Bacillus lactis aerogenes***, decomposition of inositol and glycerol by (KUMAGAWA), A., i, 972.
- Bacillus mesentericus niger***, darkening of carbohydrates by growth of (MUSCHL), A., i, 1095.
- Bacillus proteus***, action of, on *di*-naphthylalanine (SASAKI and KINOSE), A., i, 303.
 action of, on *l*-tryptophan (SASAKI and OTSUKA), A., i, 302.
- Bacillus pyocyaneus***, culture of, and the production of hydrocyanic acid (GORIS and LIOT; PATTY), A., i, 407.
- Bacillus subtilis***, production of carbon dioxide by (BROOKS), A., i, 201.
- Bacteria**, formation of hydrogen peroxide by (MCLEOD and GORDON), A., i, 1095.
 formation of indole by (FRIEBER), A., i, 901.
 formation of phenol by (SIEKE), A., i, 902.
 formation of toxins from (WALBUM), A., i, 902.
 action of, on racemic acids (CONDELLI), A., i, 410.
 importance of sequence in the action of antiseptics on (KARCZAG), A., i, 302.
 influence of arsenious acid on the growth of (COBET and VAN DER REIS), A., i, 611.

- Bacteria**, decomposition of citric acid in milk by (KICKINGER), A., i, 1219.
 effect of formaldehyde on (HAILER), A., i, 408.
 action of, on minerals (HELBRONNER and RUDOLFS), A., i, 706.
 nucleic acid in (SCHAEFFER, FOLKOFF, and JONES), A., i, 1095.
 influence of the reaction of the solution on the action of poisons or drugs on (LABES), A., i, 901.
 proteases in (DERNBY), A., i, 405.
 destruction of, by quinones (MORGAN and COOPER), A., i, 204.
 oxidation of sulphur by, in soils (WAKSMAN), A., i, 706.
 colon-typhoid, action of tellurium salts on (JOACHIMOGLU), A., i, 1095.
 intestinal, action of, on bile pigments (PASSINI), A., i, 966.
 lactose-fermenting (LEVINE), A., i, 901.
 pentose-fermenting (FRED, PETERSON, and ANDERSON), A., i, 201, 971.
 purple, microchemistry of (GICKLHORN), A., i, 303.
 soil, oxidation of sulphur compounds by (LIPMAN, WAKSMAN, and JOFFE), A., i, 303.
 estimation of substances which inhibit the reducing action of (SCHNABEL), A., i, 304.
- Balance**, micro-, Steele-Grant (HARTUNG), A., ii, 495.
 Westphal, accuracy of the (TSCHUDY), A., ii, 820.
- Barbel**, constituents and toxicity of the roes of the (MCCRUDDEN), A., i, 194.
- Barbituric acid**, action of diazomethane on (HERZIG), A., i, 873.
 detection of, and its derivatives (FABRE), A., ii, 795.
- Barbituric acids**, substituted, as hypnotics (CARNOT and TIFFENEAU), A., i, 900.
- Barium**, absorption spectrum of (LINDSAY), A., ii, 599.
L-spectrum of (DAUVILLIER; DE BROGLIE and DAUVILLIER), A., ii, 542.
- Barium salts**, conductometric titrations with (KOLTHOFF), A., ii, 864.
 poisoning by. See Poisoning.
- Barium carbonate**, action of nitrogen on carbon and, at high temperatures (ASKENASY and GRUDE), A., ii, 445.
 platinichloride, dissociation of (GIRE), A., ii, 551.
 chromate, solvolysis of (HICKS and CRAIG), A., ii, 622.
 platinocyanide, de-intensifying of, in the Villard effect (ZIMMERN and SALLES), A., ii, 104.
- Barium nitrate**, crystal structure of (VEGARD), A., ii, 503.
 thallous nitrite (CUTTICA and PACIELLO), A., ii, 378.
 peroxide, velocity of formation of (SASAKI), A., ii, 278.
 catalytic influence of oxides on the decomposition of (KENDALL and FUCHS), A., ii, 147.
 silicate (ESKOLA), A., ii, 849.
 sulphate, adsorption of radium by (GERMANN), A., ii, 16.
 equilibrium of potassium carbonate with (RAMANN and SALLINGER), A., ii, 131.
 tungstate (SMITH), A., ii, 774.
- Barium selenic acid** (MEYER and FRIEDRICH), A., ii, 644.
- Barium sulphuric acid** (MEYER and FRIEDRICH), A., ii, 644; (BALAREFF), A., ii, 864.
- Barium estimation**:—
 estimation of, volumetrically (POLONOVSKI), A., ii, 720.
- Barley**, amylase of (BAKER and HULTON), T., 1929.
 germinated, velocity of action of enzymes of (MAESTRINI), A., i, 507, 508.
 estimation of starch in (LING, CALLOW, and PRICE), A., ii, 879.
- Base**, C_8H_7N , and its salts, from *p*-xylene and sulphuryl azide (CURTIUS and SCHMIDT), A., i, 777.
 C_8H_7ON , and its salts and derivatives (TROGER and SCHWARZENBERG), A., i, 167.
 $C_{10}H_9ON$, and its salts, from *l*-tartaraldehydephenylhydrazones and hydrogen chloride (BERGMANN), A., i, 8.
 $C_{11}H_{17}O_3N$, and its salts and derivatives, from hydrolysis of *N*- γ -benzoxypropyldihydroisoindole (V. (BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
 $C_{13}H_{15}ONS$, and its salts, from thiodiphenylamine and sodium hydroxide (KEHRMANN and DARDEL), A., i, 1063.
 $C_{18}H_{20}ON_2$, and its salts, from aniline and acraldehyde (MANN), T., 2179.
 $C_{20}H_{24}ON_2$, from *p*-toluidine and acraldehyde (MANN), T., 2182.
 $C_{22}H_{26}N_2$, from acetoneanil and methyl iodide (KNOEVENAGEL and BÄHR), A., ii, 750.
 $C_{26}H_{28}O_3N_4$, and its salts, from oxidation of base $C_{18}H_{20}ON_2$ (MANN), T., 2180.
- Bases**, Werner's theory of (LAMBS and YNGVE), A., ii, 217.

- Bases**, heterocyclic, reactivity of methyl groups in (MILLS and SMITH), T., 2724.
- organic, additive compounds of metallic salts with (PETERS), A., i, 48.
- detection of, with iodic acid (ROSENTHALER), A., ii, 327.
- detection and separation of, by means of glyoxalin-dicarboxylic acid (PAULY and LUDWIG), A., i, 953.
- strong, estimation of, volumetrically, in presence of weak bases (KOLTHOFF), A., ii, 223.
- weak, alcoholysis of the salts of (GOLD-SCHMIDT, GÖRBITZ, HOUGEN, and PAHL), A., ii, 135.
- estimation of, volumetrically (LIZIUS and EVERS), A., ii, 654.
- Basilicum oil**, constants of (GATTE-FOSSE), A., i, 1167.
- Basicity** (KELLER), A., ii, 131.
- Bauxite**, estimation of titanium dioxide in (WINCH and CHANDRATREYA), A., ii, 459.
- Bean**, adzuki. See *Phaseolus angularis*.
lima. See *Phaseolus lunatus*.
navy, carbohydrates in (EICHELBERGER), A., i, 799.
pichurim. See Pichurim bean.
runner, nitrogen distribution in leaves of (CHIBNALL), A., i, 908, 1225.
soja. See Soja beans.
- Beckmann transformation** (MEISENHEIMER), A., i, 152.
- Becquerelite**, radioactive (SCHOEF), A., ii, 450.
- Beechnut oil**, constituents of (HEIDUSCHKA and ROSER), A., i, 945.
characteristics and utilisation of (B), A., i, 96.
- Bees-wax**. See under Wax.
- Beetroot** (*Beta vulgaris*), anthocyanin from (ANDREWS; KOZLOWSKI), A., i, 96.
wild, composition of (SAILLARD), A., i, 415.
- Behenic acid**, anilide and esters of (TOYAMA), A., i, 1111.
sodium salt, properties of solutions of (FLECKER and TAYLOR), T., 1101.
docosyl and phenyl esters (BRIGL and FUCHS), A., i, 713.
- Belladonna**, chemical constituents of the leaves of (GORIS and LARSONNEAU), A., i, 1099.
alkaloids from (RIFERT), A., i, 96.
- Benzaldehyde**, solubility of copper in dilute solutions of (BERNOULLI and SCHAAF), A., i, 1029.
reduction of, by sodium in alcohol (DE POMMEREAU), A., i, 339.
- Benzaldehyde** and *p*-nitro-, alkylhydrazones of (BRADY and McHUGH), T., 1650.
picrate, compound of thiocarbamide and (TAYLOR), T., 2269.
estimation of chlorine in (SCHIMMEL & Co.), A., ii, 78.
- Benzaldehyde**, 2:4-dihydroxy-. See *β*-Resorcyraldehyde.
- iodo-*m*-hydroxy- (HENRY and SHARP), T., 1059.
nitro-derivatives, solubility and volatility of (SIDGWICK and DASH), T., 2586.
2:4:6-trinitro-, derivatives of (LOWY and BALDWIN), A., i, 134.
- Benzaldehydes**, hydroxy-, mercury derivatives (HENRY and SHARP), T., 1055.
nitro-, reactivity of (HELLER, LAUTH, and BUCHWALDT), A., i, 348.
- Benzaldehyde-S-benzylthiosemicarbazone** (WILSON and BURNS), T., 874.
- Benzaldehyde cyanohydrin**, action of emulsin in synthesis of (NORDEFELDT), A., i, 1077.
- Benzaldehyde-2-nitro-4-cyanophenylhydrazones** (MATTAR), A., i, 252.
- Benzaldehyde-*p*-nitrophenylmethylhydrazones** (CIUSA and RASTELLI), A., i, 1073.
- Benzaldehyde phenylhydrazones**, additive compounds of, with aromatic polynitro-compounds (CIUSA and VECCHIOTTI), A., i, 474.
- Benzaldioximes**, 2:4- and 2:6-dinitro-, *N*-arylethers of (BARROW, GRIFFITHS, and BLOOM), T., 1714.
- Benzamide**, kinetics of the reaction of acetic anhydride with (KREMANN, RÖSLER, and PENKNER), A., ii, 748.
equilibria of, with dihydroxynaphthalenes (KREMANN, HEMMELMAYR, and RIEMER), A., i, 1012.
- Benzamidino**, action of cyanogen and nitrogen iodides on (ROBIN), A., i, 36.
- α -Benzamido- β -methylpropane, β -chloro-** (DERSIN), A., i, 142.
- Benzanilide**, 2:4-dinitro- (BARROW, GRIFFITHS, and BLOOM), T., 1714.
- Benzanthrone**, amino-, and hydroxy- (BRITISH DYE STUFFS CORPORATION, LTD., PERKIN and SPENCER), A., i, 941.
2-amino-, and its derivatives, and 2-hydroxy-, preparation of (PERKIN and SPENCER), T., 479.
2-hydroxy-, derivatives of (BRADSHAW and PERKIN), T., 911.
- 1:9-Benzanthrone**, 6- and 7-chloro- (SCHOLL, SEER, and DAIMER), A., i, 258.

Benzanthronecarboxylic acid, hydroxy-, and its salts and derivatives (BRADSHAW and PERKIN), T., 913.

m-a-Benzbispyrrole, 2:6-dihydroxy-, and its sodium salt (DAVIES and HICKOX), T., 2652.

Benzbisthiazoles (EDGE), T., 772.

meso-Benzdixanthylene (ECKERT and ENDLER), A., i, 941.

Benzene, constitution of (FRASER), T., 188.

structure of (CROCKER), A., i, 927; (HUGGINS), A., i, 928, 997.

nucleus, structure of the (INGOLD), T., 1133, 1143.

formation of rings attached to the (PERKIN and TITLEY), T., 1562.

existence and reactions of positive halogen attached to the (NICOLET), A., i, 121.

migration of methyl into the (BUECKMANN, CORRENS, and LIESCHE), A., i, 535.

substitution in the (DAVIES), T., 806.

effect of the mobility of chlorine in the side-chain on substitution in the (OLIVIER), A., i, 646.

influence of the chlorine atom and the methyl and sulphonyl chloride groups on substitution in the (DAVIES), T., 785.

influence of nitro-groups on the reactivity of substituents in the (BURTON and KENNER), T., 489, 675.

compounds, structure of molecules of (ROBINSON; KENNER), A., i, 533.

variation of the refractive index and density of, with temperature (PARKER and THOMPSON), T., 1341.

absorption and fluorescence spectra of (HENRI), A., ii, 679.

ultra-violet absorption spectrum of the vapour of (HENRI), A., ii, 331.

specific heat of the saturated vapour of (BRUHAT and DELAYGUE), A., ii, 348.

compressibility of (RICHARDS, BARTLETT, and HODGES), A., ii, 28.

equilibrium of ethyl alcohol, water and (ORMANDY and CRAVEN), A., i, 215.

equilibrium in the system, silver perchlorate, water and (HILL), A., ii, 555.

action of benzyl chloride and, in presence of iron pyrites (SMYTHE), T., 1276.

Benzene, reaction of carbon dioxide with, under the electric discharge (MIYAMOTO), A., i, 418.

action of carbonyl chloride on, in presence of aluminium chloride (WILSON and FULLER), A., i, 827.

compounds of cresols with (BERL and SCHWEBEL; C. and W. v. RECHENBERG), A., i, 932.

ethylation of (MILLIGAN and REID), A., i, 330.

action of sulphuryl azide on (SCHMIDT), A., i, 777.

chlorination of, with sulphuryl chloride (SILBERRAD), T., 1015.

derivatives, action of sulphuryl chloride on (DURRANS), T., 44.

detection of, in light petroleum (SCHWARZ), A., ii, 531.

estimation of (ZABOROWSKI), A., ii, 877.

estimation of, in gases (BERL), A., ii, 591.

Benzene, *tribromo*-derivatives, action of sodium with (FUCHS and METZL), A., i, 442.

5-bromo-1:2:4-*trinitro*-, action of amines on (GIUA and ANGELETTI), A., i, 649.

action of hydrazines and of pyridine on (GIUA), A., i, 691.

chloro-, physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.

p-dichloro-, nitration of (MACLEOD, PFUND, and KILPATRICK), A., i, 1131.

chloro-*trinitro*-derivatives (HOLLEMAN), A., i, 997.

halogen derivatives, corresponding states of (PRUD'HOMME), A., ii, 426.

1:2:4:5-*tetrahydroxy*-, colouring matters from (MUKERJI), T., 545.

nitro-derivatives, equilibrium of anthracene with (KREMANN and MÜLLER), A., i, 131.

equilibrium of, with carbazole and acenaphthene (KREMANN and STRZELBA), A., i, 176.

equilibria of triphenylcarbinol with (KREMANN, HOHL, and MÜLLER), A., i, 138.

nitro-, action of mercuric acetate with (WIBAUT and JÜRGENS), A., i, 694.

catalytic reduction of (BROWN and HENKE), A., i, 445, 535; (HENKE and BROWN), A., i, 586, 1196.

reduction of, by sodium in alcohol (DE POMMEREAU), A., i, 339.

action of sodium sulphate on (SEYEWETZ and VIGNAT), A., i, 241.

- Benzene, *m*-dinitro**, equilibrium in the system, urethane and (PUSHIN and FIOLETOVA), T., 2822.
- nitrochloro-**, equilibrium of the formation of nitroanisole from (BLOM), A., ii, 38.
- 5-Benzene(4'-arsinic acid)azo-4-amino-1,2-bisphenoxyacetic acid**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-3-amino-6-methoxyphenoxyacetic acid**, and its hydrochloride (JACOBS and HEIDELBERGER), A., i, 75.
- 6-Benzene(4'-arsinic acid)azoamino-methylphenoxyacetic acids**, and their salts (JACOBS and HEIDELBERGER), A., i, 75.
- Benzene(4-arsinic acid)azo- α -aminonaphthoxyacetic acids**, and their sodium salts (JACOBS and HEIDELBERGER), A., i, 75.
- 6-Benzene(4'-arsinic acid)azo-3-aminophenoxyacetic acid**, and its salts (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-2-ethoxyphenylglycine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 5-Benzene(4'-arsinic acid)azo-2-hydroxyphenoxyacetic acid**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-2-methoxyphenylaminomethanesulphonic acid**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-2-methoxyphenylglycine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-2-methylphenylglycine**, and its hydrochloride (JACOBS and HEIDELBERGER), A., i, 75.
- α -Benzene(4-arsinic acid)azo- α -naphthylglycine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-phenylaminomethanesulphonic acid**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 4-Benzene(4'-arsinic acid)azo-phenylglycine**, and its hydrochloride (JACOBS and HEIDELBERGER), A., i, 75.
- 1-Benzeneazo-3-*p*-aminophenylamino-6-sulphonic acid** (BUCHERER and ZIMMERMANN), A., i, 466.
- Benzeneazo-6-amino-2-phenylbenzthiazole, *p*-nitro-**; and its acetyl derivative (BOGERT and ABRAHAMSON), A., i, 576.
- 7-Benzeneazo-6-amino-3-*p*-tolyl α -naphthatriazole, 7-*p*-nitro-** (MORGAN and CHAZAN), A., i, 181.
- 4-Benzeneazo-3-anilino-5-ketoisoxazole** (WORRALL), A., i, 874.
- 6-Benzeneazohemimellitenol** (v. AUWERS and SAURWEIN), A., i, 1033.
- 6-Benzeneazo- α -hexahydrodeoxyinchonine** (JACOBS and HEIDELBERGER), A., i, 672.
- 5-Benzeneazo-2-hydroxy-4:6-dimethylbenzaldehyde**, and its sodium salt (v. AUWERS and SAURWEIN), A., i, 1032.
- Benzeneazo-4-hydroxy-2-phenyl mercaptan, 3-nitro-4-thiol-** (WATSON and DUTT), T., 2417.
- Benzeneazohydroxyquinol** (MUKERJI), T., 551.
- 4-Benzeneazo-1-methylaminonaphthalene-8-sulphonic acid, 4-*cp*-dinitro-** (KONIG and KOHLER), A., i, 880.
- 4-Benzeneazo-*N*-methyl-1:8-naphthasultam, 4-*p*-mono- and 4-*cp*-di-nitro-** (KONIG and KOHLER), A., i, 880.
- 4-Benzeneazo-1:8-naphthasultam, 4-*p*'-mono- and 4-*cp*-di-nitro-** (KONIG and KOHLER), A., i, 880.
- 2-Benzeneazo-1:8-naphthasultam-4-sulphonic acid, 2-*p*'-nitro-** and its potassium salt (KONIG and KEIL), A., i, 823.
- Benzeneazo- β -naphthol, *p*-amino-**, and its derivatives (CHARRIER and CORTASSA), A., i, 771.
- 2-Benzeneazo- α -naphthol-3:6-disulphonic acid, 7:8-diamino-** (BUCHERER and ZIMMERMANN), A., i, 466.
- Benzeneazo- β -naphthylamine, *p*-amino-**, acetyl derivative (CHARRIER and VIOLA), A., i, 771.
- Benzeneazo- β -naphthylglycine**, and *p*-nitro-, ethyl ester (FIEBZ and SALLMANN), A., i, 870.
- 4-Benzeneazo-1-naphthyl mercaptan** (WATSON and DUTT), T., 1940.
- Benzeneazo- β -naphthylmethylamine-6-sulphonic acid**, sodium salt (MORGAN and ROOKE), A., i, 134.
- 4-Benzeneazo-1-naphthylxanthic acid**, ethyl ester (WATSON and DUTT), T., 1940.
- Benzeneazo-4-nitro- α -naphthol, *p*-nitro-** (MUKERJI), T., 2880.
- Benzeneazo-4-oxybenzene-2':3- and 3:3'-dicarboxylic acids**, cobaltammine salts of (MORGAN and SMITH), T., 2872.
- Benzeneazoxycarboxylic acid**, potassium salt (PIERONI), A., i, 1073.
- Benzene-1-azo-2-oxy-3-naphthoic acid, *m*-nitro-**, and its cobaltammine salt (MORGAN and SMITH), T., 2873.

- 9-Benzeneazo-10-phenanthrol, *p*-bromo- and *o*-, *m*-, and *p*-nitro- (CHARRIER and DEMICHELIS), A., i, 772.
- Benzeneazophenol, and its derivatives, absorption spectra of (SMITH and BOORD), A., ii, 602.
- 2-Benzeneazophenol, 4'-bromo-, 4'-chloro-, and 4'-iodo- (CHATTAWAY and HILL), T., 2757.
- 3-Benzeneazophenylarsinic acid, 4-hydroxy- (BART), A., i, 1202.
- 1-*p*-Benzeneazophenylpiperidine, 4-hydroxy-, and its derivatives (BORSCHKE and BONACKER), A., i, 51.
- 5-Benzeneazosalicylaldehyde-4'-sulphonic acid, and its derivatives (WEIL, TRAUN, and MARCEL), A., i, 1024.
- Benzeneazosalicylic acid, *m*-nitro-, chromic, and cobaltammine salts of (MORGAN and SMITH), T., 2870.
- Benzeneazo-*ar*-tetrahydro-8-naphthols (SCHROETER, SVANOE, EINBROCK, GELLER, and RIEBENSAMM), A., i, 128.
- Benzeneazo-*p*-tolueneazophenols, *p*-bromo- (CHATTAWAY and HILL), T., 2760.
- Benzenedialdehydes, hydroxy-, and their derivatives (WEIL and BRIMMER), A., i, 350.
- 2-Benzenediazamino-3-bromo-*p*-cymene, and its hydrochloride and 2-*p*-nitro- (WHEELER and SMITHEY), A., i, 333.
- Benzenediasonium chloride, compound of bismuth trichloride and (CHALLENGER and WILKINSON), T., 102.
- Benzenedi-*m*-thiobenzoic acid, and its sodium salt (SMILES and GRAHAM), T., 2508.
- Benzenedi-*p*-thiodimethylaniline (SMILES and GRAHAM), T., 2510.
- Benzenehydrazo-4-nitro-*a*-naphthol, *p*-nitro- (MUKERJI), T., 2881.
- Benzenehydrazo-*o*-nitrophenol, *p*-nitro- (MUKERJI), T., 2881.
- Benzenesulphamphthalein, 1:2:4:5-tetrahydroxy-, and its acetyl derivative (DUTT), T., 2392.
- Benzenesulphonic acid, *N*-allylbrucinium, *N*-allylpapaverinium, and 1-allylpyridinium salts (FÖLDI), A., i, 732.
- Benzenesulphonic acid, bromoethyl, chloroethyl and dichloropropyl esters (v. KERESZTY and WOLF), A., i, 1131.
- Benzenesulphonpropyl-*n*-butylamide (v. BRAUN and WEISMANTEL), A., i, 1151.
- Benzenesulphonyl chloride, 2:5-dichloro- (STEWART), T., 2557.
- 5-Benzenesulphenylaminocacenaphthene (FLEISCHER and SCHRANZ), A., i, 1144.
- 3-Benzenesulphonylceumarin, 7-hydroxy- and 7-hydroxy-3-*p*-chloro- (TROGER and BOLTE), A., i, 268.
- Benzenesulphonyldiallylamine (FÖLDI), A., i, 733.
- 3-Benzenesulphonyl-2-methylquinoline, and 3-*p*-chloro-, and their salts and derivatives (TROGER and MENZEL), A., i, 270.
- 3-Benzenesulphonylquinophthalene, and 3-*p*-chloro- (TROGER and MENZEL), A., i, 270.
- Benzhydroxamic acid, thienoyl ester, and its salts (JONES and HURD), A., i, 250.
- Benzidine, compound of tellurium tetrabromide with (LOWY and DUNBROOK), A., i, 446.
- hydrochloride, action of, with lignin (VAN ZIJF), A., ii, 94.
- hydroferrocyanide (CUMMING), T., 1296.
- Benzil (*dibenzoyl*), action of aluminium chloride with (SCHOLL and SCHWARZER), A., i, 331.
- conversion of, into benzilic acid (LACHMAN), A., i, 459.
- alkali salts of (STAUDINGER and BINKERT), A., i, 1016.
- Benzil, 4:4'-*di*hydroxy-, and its dibenzoate (SCHÖNBERG and KRAEMER), A., i, 664.
- nitro-, preparation of, and its phenylhydrazone (BARNETT and KAY), A., i, 844.
- 2:4-*d*initro-, and its derivatives (BISHOP and BRADY), T., 2864.
- ψ-Benzils (SCHÖNBERG and KRAEMER), A., i, 663.
- Benzilic acid, conversion of benzil into (LACHMAN), A., i, 459.
- α-Benzilmonoxime, cobaltic salt (PONZIO), A., i, 662.
- Benziloximes, isomerism of (MEISENHEIMER), A., i, 152.
- Benzochloroamides, preparation and properties of (ELLIOTT), T., 202.
- 1:2:3:4-Benzodiquinone-1:3-dioxime (*dinitrosoresorcinol*), cobaltammine and ferric salts of (MORGAN and MOSS), T., 2857.
- Benzothylpropylamide (v. BRAUN and WEISMANTEL), A., i, 1151.
- Benzofuran series, syntheses in the (MAMELI), A., i, 669.
- Benzoic acid, and *p*-nitro-, chloroethyl esters, preparation of (ALTWEGG and LANDRIVON), A., i, 315.
- effect of substitution on the absolute affinity of (RØRDAM), A., i, 338.
- heat of combustion of (VERKADE, COOPS, and HARTMAN), A., ii, 474 ; (SWIENTOSLAWSKI and STARCZEWSKA), A., ii, 616.

- Benzoic acid**, compounds of carbamide and (OSAKA and ANDO), A., i, 1127.
- Benzoic acid**, and hydroxy-derivatives, alkali salts, solubility of, in water (SIDGWICK and EWBANK), T., 1844.
- Benzoic acid**, *o*-aminophenyl ester, hydrochloride, and its derivatives (BERGMANN, ULPTS, and CAMACHO), A., i, 1181.
- α*-amyrenol and bromo-*α*-amyrenol esters (ZINKE, FRIEDRICH, JOHANNSEN, and RICHTER), A., i, 667.
- benzylidene-*p*-aminophenylester (REDELLEN and DANILOF), A., i, 148.
- cinnamyl ester (RUPE and MULLER), A., i, 40.
- ethyl ester, reduction of, by sodium in alcohol (DE POMMERAU), A., i, 338.
- lubanyl ester, constitution of (ZINKE, HANSELMAYER, and EHMER), A., i, 668.
- and nitro-, tribromo- and trichloro-*tert*.-butyl esters (ALDRICH and BLANNER), A., i, 910.
- p*-chlorosulphonyl derivative (SMILES and HARRISON), T., 2023.
- Benzoic acid**, *p*-amino-, *p*-chloroacetyl derivative (QUICK and ADAMS), A., i, 601.
- and *p*-nitro-, esters of, and their derivatives (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- benzyl ester (SOCIETY FOR CHEMICAL INDUSTRY IN BASLE), A., i, 828.
- n*-butyl ester, and its hydrochloride (SOCIÉTÉ CHIMIQUE DES USINES DU RHÔNE), A., i, 827.
- γ*-diallylaminopropyl ester, preparation of (KAMM and VOLWILER), A., i, 654.
- γ*-dibutylaminopropyl ester (KAMM and ADAMS), A., i, 828.
- chlorodihydroxy-derivatives (FABRE), A., i, 1148.
- m*-hydroxy-, bromo-, chloro- and hydroxy-derivatives of (BEYER), A., i, 37.
- iodo-*m*-hydroxy- (HENRY and SHARP), T., 1059.
- p*-nitro-, lead diphenyl salt (GODDARD, ASHLEY, and EVANS), T., 982.
- 3:5-*d*-nitro-, compound of diethylaniline and (LEY and GRAU), A., i, 586.
- 2:4:6-*tri*-nitro-, mercuric salt (KHARASCH), A., i, 190.
- p*-dithio-. See *p*-Disulphidobenzoic acid.
- p*-thiol- (SMILES and HARRISON), T., 2024.
- Benzoic acids**, substituted, ethyl esters, velocity of saponification of (CASHMORE, MCCOMBIE, and SCARBOROUGH), T., 243.
- nitro-, estimation of nitrogen in, by the Kjeldahl method (MARGOSCHES and VOGEL), A., ii, 522.
- 4:5-Benzoin-1-one (MAYER and SIEGLITZ), A., i, 743.
- 5:6-Benzoin-1-one, 4-bromo-, and its oxime (MAYER and SIEGLITZ), A., i, 744.
- Benzonaphtha-*p*-thiazine. See *ββ*-Naphthaphenthiazine.
- Benzonitrile, 4-bromo-3-nitro-, and 4-chloro-3-nitro-, replacement of halogen in (MATTAAR), A., ii, 275.
- 2:5-dihydroxy- (FICHTER and GRISARD), A., i, 37.
- γ*-Benzophenarsazinic acid, and its sodium salt (LEWIS and HAMILTON), A., i, 183.
- Benzophenone**, latent heat of fusion of (STRATTON and PARTINGTON), A., ii, 258.
- chloride, action of sodium methoxide and its homologues on (MACKENZIE), T., 1695.
- diphenylhydrazone (CIUSA and VECCHIOTTI), A., i, 475.
- Benzophenone**, substituted derivatives of, crystallography of (JAEGER), A., i, 353.
- 2:4:6:2'-*tetra*hydroxy-, and its ketimide hydrochloride (KARRER), A., i, 40.
- 2:4:6:3'- and 2:4:6:4'-*tetra*hydroxy- (NISHIKAWA and ROBINSON), T., 842.
- Benzophenoneanil** methiodide (KNOEVENAGEL and BÄHR), A., i, 750.
- Benzophenoneanil-*p*'-carboxylic acid**, salts of (REDDELLEN), A., i, 146.
- ethyl and methyl esters (REDDELLEN and DANILOF), A., i, 147.
- Benzophenone-*p*-anilinoanil** (REDDELLEN), A., i, 146.
- Benzophenone-*p*-arsenious acid** and *p*-arsenious oxide (LEWIS and CHEETHAM), A., i, 187.
- Benzophenone-*p*-arsinic acid**, and nitro- (LEWIS and CHEETHAM), A., i, 187.
- Benzophenonebenzoylhydrazone** (REDDELLEN), A., i, 146.
- Benzophenonebenzoylimide** (REDDELLEN and DANILOF), A., i, 148.
- Benzophenone-*p*-hydroxyanil**, acetate and benzoate of (REDDELLEN and DANILOF), A., i, 148.
- Benzophenonemethylimide** (REDDELLEN), A., i, 146.
- Benzophenone-*β*-naphthyl** (REDDELLEN), A., i, 146.
- Benzophenone-*p*-tolyl** methiodide (KNOEVENAGEL and BÄHR), A., i, 750.

- Benzopolymethylene compounds** (v. BRAUN, HAHN, and SEEMANN), A., i, 728; (v. BRAUN, MOLDAENKE, DIRLAM, and GRUBER), A., i, 748; (v. BRAUN and GRUBER), A., i, 762.
- Benzopropyl-*n*-butylamide** (v. BRAUN and WEISMANTEL), A., i, 1151.
- α -Benzopyrone**, synthesis of derivatives of (TRÖGER and BOLTE), A., i, 267.
- Benzopyrylium compounds**, ring opening in (DECKER and BECKER), A., i, 353.
- p*-Benzoquinone**, electromotive equilibria of quinol with (GEANGER and NELSON), A., i, 43.
addition of benzene to (PUMMERER and PRELL), A., i, 1164.
derivatives, oxidation-reduction potentials of (LAMER and BAKER), A., ii, 785.
diphenylhydrazone, and its leuco-derivative (GOLDSCHMIDT and RENN), A., i, 476.
- Benzoselenamide**, *o*-amino- (BOGERT and CHEN), A., i, 1182.
- Benzotrichloride-*o*-carboxylic acid**, anilide and methyl ester (OTT), A., i, 836.
- Benzoxazole-2-aldehydeanil and -2-carboxyanilide** (SKRAUP and MOSER), A., i, 575.
- Benzoxazole-2-carboxyl chloride** (SKRAUP and MOSER), A., i, 575.
- Benzoxazole-2-carboxylic acid** (SKRAUP), and MOSER), A., i, 575.
- Benzoxazyl diphenylcarbinol** (SKRAUP and MOSER), A., i, 575.
- Benzoxazyl phenyl ketone**, and its oxime (SKRAUP and MOSER), A., i, 575.
- 5-Benzoxo-1-benzoyl-3-methylpyrazole**, and 4-bromo- (ROJAHN), A., i, 373.
- β -Benzoxo- α -diethylaminoethylbutyric acid**, ethyl ester, and its hydrochloride (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 640.
- β -Benzoxoethylconiline hydrochloride** (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- N*- β -Benzoxoethylidihydroisoindole hydrochloride** (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- N*- β - and - γ -Benzoxoethyltetrahydroisoquinoline hydrochlorides** (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 761.
- 2-Benzoxynaphthalene**, 3:6-*di*bromo- (FRANZEN and STÄUBLE), A., i, 450.
- N*- γ -Benzoxopropylconiline hydrochloride** (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- N*- β -Benzoxopropylidihydroisoindole**, and its derivatives (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- N*- γ -Benzoxopropylpiperidine hydrochloride** (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- Benzoyl chloride**, chlorination of (HOPE and RILEY), T., 2510.
peroxide, action of cholesterol on (WINDAUS and LÜDERS), A., i, 453.
Golodetz's reaction for (GELISSEN), A., ii, 460.
- Benzoylacetone**, 3-chloro-, and its copper derivative (MORGAN, DREW, and BARKER), T., 2456.
- Benzoylacetones**, selenium (MORGAN, DREW, and BARKER), T., 2458.
- Benzoylacyldiazomethane**, interaction of triphenylphosphine with (STAUDINGER and LÜSCHER), A., i, 238.
- Benzoyl acetyl ketone**, hydrazone of (STAUDINGER and LÜSCHER), A., i, 238.
- 1-Benzoyl-1-allylcyclopropane** (HALLER and BENOIST), A., i, 350.
- 5-Benzoylamino- γ -ketovaleic acid**, ethyl ester, and its semicarbazone (WINDAUS, DÖRRIES, and JENSEN), A., i, 61.
- α -Benzoylamino- β -[6-methoxyindyl(3)]-acrylic acid** (KERMACK, PERKIN, and ROBINSON), T., 1883.
- N*-Benzoylanthalamine**, *m*-nitro-, and its methyl ether (SPATH), A., i, 166.
- 1-Benzoyl-5-anilinomethylene-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- o*-Benzoylbenzoic acid**, *o*-3':5'-*di*bromo-4'-hydroxy-, *o*-3'chloro-4'-hydroxy-, and *o*-3':5'-*di*chloro-4'-hydroxy- (THIEL and MÜLLER), A., i, 660.
- o*-3':5'-*di*bromo-4'-hydroxy-, and *o*-4'-hydroxy-, and its salts and derivatives** (ORNDORFF and KELLEY), A., i, 833.
- Benzoylbenzoic acid**, mechanism of the formation of (GREENE and ROBINSON), T., 2182.
- Benzoyl- β -benzoxoethylcarbamide** (FROMM and HONOLD), A., i, 530.
- Benzoyl-3-benzoyloxy-4-methoxymandelonitrile** (GREENE and ROBINSON), T., 2195.
- γ -Benzoyl- β -benzylcrotonic acid**, γ -*p*-chloro- (KÖHLER and SMITH), A., i, 458.
- 1-Benzoylbenzylcyclopropane** (HALLER and BENOIST), A., i, 350.
- 2-Benzoyl-4-bromobenzoic acid** (STEPHENS), A., i, 141.

- α -Benzoyl- β -butoxy- β -phenylethanes**, α -bromo- (DUFRAISSE and GÉRALD), A., i, 848.
- α -Benzoyl- β -butoxy- β -phenylethylenes** (DUFRAISSE and GÉRALD), A., i, 843.
- 1-Benzoyl-2-*p*-chlorophenylcyclopropane**, 3-nitro- (KÖHLER and SMITH), A., i, 458.
- Benzoylcinchonidine** dihydrochloride (HEIDELBERGER and JACOBS), A., i, 673.
- Benzoylcinnamantialdoxime** (BRADY and THOMAS), T., 2104.
- Benzoyldiasoacetic acid**, methyl ester, interaction of triphenylphosphine with (STAUDINGER and LÜSCHER), A., i, 237.
- 4-Benzoyl-3:5-dibenzylthiol-1:2:4-triazole** (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 379.
- Benzoyldihydrocinchonidine** hydrochloride (HEIDELBERGER and JACOBS), A., i, 673.
- Benzoyldihydroquinine** hydrochloride (HEIDELBERGER and JACOBS), A., i, 673.
- 1-Benzoyl-3:5-diketo-1:4-hexahydrodiazine**, attempts to prepare (DUBSKY and HOHER), A., i, 57.
- N*-Benzoyl-5:7-dimethylisatoic acid** (HELLER, BENADE, and HOCHMUTH), A., i, 1060.
- 1-Benzoyl-3:4-dimethylpyrazole**, 5-chloro- (ROJAHN), A., i, 1184.
- 5-Benzoyl-2:4-dimethylpyrrole-3-carboxylic acid**, ethyl ester (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 4-Benzoyl-1:3-diphenylpyrazole**, 5-chloro- (ROJAHN), A., i, 373.
- α -Benzoyl- β -ethoxy- β -phenylethane**, α -bromo- (DUFRAISSE and GÉRALD), A., i, 39.
- 1-Benzoyl- β -*p*-ethoxyphenyl-4-glyoxal-*one***, 2-thiol- (DAINS, THOMPSON and ASENDORF), A., i, 1186.
- Benzoyliso Eugenol**, dibromo- and diiodo- (RASTELLI), A., i, 1010.
- Benzoyl-*p*-hydroxyphenylbenzoylcarbamide** (SPECKAN), A., i, 580.
- Benzoyl-*p*-hydroxyphenylcarbamide** (SPECKAN), A., i, 580.
- Benzoyliminodiacetic acid**, and its salts and methyl ester (DUBSKY and HOHER), A., i, 57.
- Benzoylkynurenic acid**, methyl ester (SPATH), A., i, 173.
- Benzoylmandelonitrile**, action of alcoholic sodium ethoxide on (GREENE and ROBINSON), T., 2182.
- α -Benzoyl- β -methoxy- β -phenylethane**, α -bromo- (DUFRAISSE and GÉRALD), A., i, 843.
- α -Benzoyl- β -methoxy- β -phenylethylene** (DUFRAISSE and GÉRALD), A., i, 848.
- α -Benzoyl- α -methylacetoacetic acid**, methyl ester (DIRCKMANN and WITTMANN), A., i, 1157.
- Benzoylmethylfurozan** (PONZIO), A., i, 1038.
- α - and β -Benzoylmethylglyoximes**, and their salts and derivatives (PONZIO), A., i, 1038.
- Benzoylmethylnaphthalenes** (MAYER and NIEGLITZ), A., i, 740, 743.
- 1-Benzoyl-1-methylcyclopropane**, and its derivatives (HALLER and BENOIST), A., i, 350.
- Benzoyl- β -naphthylmethylamine-6-sulphonyl chloride** (MORGAN and ROOKE), A., i, 135.
- Benzoyl-2-nitro-3-methylamino-*p*-phenetidine** (REVERDIN and ROETHLISBERGER), A., i, 538.
- Benzoyl-2:3-*di*- and -2:3:5-*tri*-nitro-*p*-phenetidines** (REVERDIN and ROETHLISBERGER), A., i, 538.
- Benzoyloxydiethylaminobutane**, amino- and nitro-, hydrochlorides (FOURNEAU and PUYAL), A., i, 639.
- Benzoyloxydiethylaminopropane**, amino- and nitro-, hydrochlorides (FOURNEAU and PUYAL), A., i, 639.
- Benzoyloxydimethylaminopropane**, nitro-, hydrochloride (FOURNEAU and PUYAL), A., i, 639.
- Benzoylphenylacetylene**, addition of malonic esters to (KÖHLER), A., i, 461.
- 1-Benzoyl-3-phenyl-5-anilinomethylene-4-glyoxal-*one***, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- Benzoylphenylbenzylcarbinol**, and its derivatives (GARCÍA BANÚS and PASCUAL VILA), A., i, 734.
- γ -Benzoyl- β -phenylbutyrolactone- α -carboxylic acid**, methyl ester (KÖHLER), A., i, 552.
- Benzoylphenylcarbamide** (FROMM and HUNOLD), A., i, 530.
- γ -Benzoyl- β -phenylcrotonic acid**, and its methyl ester (KÖHLER), A., i, 461.
- γ -Benzoyl- β -phenylethylmalonic acid**, methyl ester, and its bromo-derivatives (KÖHLER), A., i, 552.
- 1-Benzoyl-3-phenyl-4-glyoxal-*one***, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- γ -Benzoyl- β -phenyl- α -methylbutyrolactone- α -carboxylic acid**, methyl ester (KÖHLER), A., i, 552.
- 1-Benzoyl-2-phenylcyclopropane**, 3-nitro-1-*p*-chloro- (KÖHLER and SMITH), A., i, 458.

- Benzoylphloroglucinols**, hydroxy- (NISHIKAWA and ROBINSON), T., 839.
- Benzoylpipecerin** (GREENE and ROBINSON), T., 2187.
- α -Benzoyl- β -propoxy- β -phenylethane**, α -bromo- (DUFRAISSE and GÉRALD), A., i, 843.
- α -Benzoyl- β -propoxy- β -phenylethylenes** (DUFRAISSE and GÉRALD), A., i, 843.
- Benzoylpropylsemicarbazide** (NEIGHBOR, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.
- Benzoyl-5-quinoline**, *o*-amino- (GRANDMOUGIN), A., i, 584.
- N*- α -Benzoylstyryl-*p*-aminophenol** (RUPE and WITTEBERG), A., i, 449.
- β -Benzoyl- α -styrylethylphosphinic acid** (CONANT, BUMP, and HOLT), A., i, 68.
- 1-Benzoyl-3-*m*-tolyl-5-anilinomethylene-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- 1-Benzoyl-3-*m*-tolyl-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- 4-Benzoyl-1-*p*-tolyl-3-methylpyrazole**, 5-chloro- (ROJAHN), A., i, 373.
- Benzoyltriacetylchromoglucose** (FREUDENBERG and IVKRS), A., i, 524.
- Benzthiazole**, derivatives of (ROMANI), A., i, 466.
- ethiodide (MILLS), T., 460.
- Benzthiazole series**, cyanine dyes of the (MILLS), T., 455.
- Benzthiazolyl-1-*p*-benzoic acid** (BOGERT and MEYER), A., i, 868.
- 1:2:3-Benzthiazole**, preparation of, and its benzoyl derivative (CHARRIER and BERETTA), A., i, 61.
- 1:3:5-Benzthiazole**, physiological action of (DALE and DUDLEY), A., i, 403.
- Benzyl acetyl sulphide** and sulphoxide, and their derivatives (WAHL), A., i, 653.
- Benzyl alcohol**, instability of, and of its esters (MESSNER), A., i, 138.
- velocity of hydrolysis of esters of (VOLWILER and VLEET), A., ii, 41.
- catalytic action of (JACOBSON), A., i, 652.
- Benzyl azide**, decomposition of (CURTIUS and EHRHART), A., i, 775.
- bromides, *o*- and *p*-nitro- (MOUREU and BROWN), A., i, 23.
- chloride, chlorination of (OLIVIER), A., i, 726.
- action of benzene and, in presence of iron pyrites (SMYTHE), T., 1276.
- p*-nitro-, condensation of, with cinnamaldehyde and furfuraldehyde (KLEUCKER), A., i, 734.
- Benzyl chloroethyl ether** (CLEMO and PERKIN), T., 649.
- cyanide. See Phenylacetoneitrile.
- methyl ether, *p*-nitro- (KLEUCKER), A., i, 734.
- phenacyl sulphide and sulphoxide, and their derivatives (WAHL), A., i, 653.
- sulphide, compounds of gold haloids with (SMITH), A., i, 933.
- disulphoxide, decomposition of (SMYTHE), T., 1400.
- Benzylacetonylsulphone**, and its derivatives (WAHL), A., i, 654.
- Benzylamine hydroferrocyanide** (CUMMING), T., 1294.
- Benzylamine**, *m*- and *p*-nitro-, preparation of (INGOLD and PIGGOTT), T., 2384.
- Benzylaminoacetic acid**, cyano-, ethyl ester, and its derivatives (SCHEIBLER and BAUMGARTEN), A., i, 656.
- Benzylaminotacetic acid**, ethyl ester (CARRIÈRE), A., i, 318.
- Benzylaminomethylmalonic acid**, and its ethylester (CURTIUS and EHRHART), A., i, 776.
- Benzylamino-oxides**, action of sodium hydroxide with (MEISENHEIMER, GRESKE, and WILLMERSDORF), A., i, 334.
- α -Benzylaminopropionic acid**, hydrochloride (CURTIUS and EHRHART), A., i, 776.
- 1-Benzylaminopyrrolidine-5-one-3-carboxylic acid**, ethyl ester (CARRIÈRE), A., i, 318.
- 5-Benzyl-5-*iso*amylbarbituric acid** (DOX and YODER), A., i, 682.
- Benzylisoamylmalonic acid**, ethyl ester (DOX and YODER), A., i, 682.
- 2-Benzylbenzoxazole**, derivatives of (SKRAUP and MOSER), A., i, 575.
- Benzylbenzylideneamine**, polymeric (CURTIUS and EHRHART), A., i, 775.
- γ -Benzylbutane- $\alpha\beta$ -dicarboxylic acid**, $\alpha\beta$ -dicyno- γ -hydroxy-, and γ -hydroxy-, lactone (BIRCH and THORPE), T., 1833.
- γ -Benzylbutane- $\alpha\alpha\beta$ -tricarboxylic acid**, γ -hydroxy-, lactone (BIRCH and THORPE), T., 1833.
- 5-Benzyl-5-*n*- and -*iso*-butylbarbituric acids** (DOX and YODER), A., i, 682.
- Benzyl-*n*- and -*iso*-butylmalonic acids**, ethyl esters (DOX and YODER), A., i, 681.
- Benzylcamphor**, *p*-amino-, and its acetyl derivative, *p*-bromo-, *p*-chloro-, *p*-cyano-, *p*-hydroxy- (HALLER and OUDIN), A., i, 356.
- Benzylcetrin** (GOMBERG and BUCHLER), A., i, 112.

- N*-Benzylidihydroquinine hydrochloride (HEIDELBERGER and JACOBS), A., i, 678.
- Benzylidimethylamine, *o*-hydroxy- (MADINAVEITIA), A., i, 133.
- Benzyl diphenylmethyl ketone, formation of (ORÉKHOFF and TIFFENEAU), A., i, 458.
- 5-Benzyl-5-ethylbarbituric acid (DOX and YODER), A., i, 682.
- α -Benzyl- α -ethylbutyric acid, and its amide and acid chloride (HALLER and BAUER), A., i, 259.
- Benzylethylmalonic acid, ethyl ester (DOX and YODER), A., i, 681.
- Benzyl α -furyl ketone, *p*-nitro-, and its oxime (KLEUCKER), A., i, 735.
- β -Benzylglucoside, toxicity of (RICHAUD), A., i, 1094.
- β -Benzyl-*d*-glucoside, action of digestive juices on (RICHAUD), A., i, 1094.
- β -Benzylglutaric acid, and $\alpha\alpha'$ -dicyano-, diamide of (STEVENSON and THORPE), T., 1720.
- N*-Benzylglycine hydrochloride (SCHEIBLER and BAUMGARTEN), A., i, 656.
- Benzylcyclohexane series, semipinacolic transposition in (TIFFENEAU and PORCHER), A., i, 537.
- 1-Benzylcyclohexan-1-ol, 2-iodo- (TIFFENEAU and PORCHER), A., i, 537.
- 1-Benzylcyclohexan-2-one, and its semicarbazone (TIFFENEAU and PORCHER), A., i, 537.
- Benzylhydrobenzoin. See $\alpha\beta\gamma$ -Triphenylpropane- $\alpha\beta$ -diol.
- β -Benzylhydrocarboxystyryl- β -carboxylic acid. β -*o*-nitro-, ethyl ester (LEUCHS, V. KATINSKY, and CONRAD), A., i, 478.
- Benzylidene chloride, action of sodium methoxide and its homologues on (MACKENZIE), T., 1695.
- Benzylideneacenaphthene, 2-hydroxy-, glucoside of (BRUNI and ROMANI), A., i, 756.
- α -Benzylideneacetoacetic acid, α -*p*-nitro-, ethyl ester (HELLEN, LAUTH, and BUCHWALDT), A., i, 348.
- Benzylideneacetoneglucosamic acid (LEVENE), A., i, 1028.
- Benzylideneacetoneglucose (LEVENE, MEYER, and WEBER), A., i, 987.
- Benzylideneacetophenone. See Phenyl styryl ketone.
- 5-Benzylideneaminoacenaphthene, and chloro-, hydroxy-, and nitro-derivatives (FLEISCHER and SCHREANZ), A., i, 1148.
- Benzylideneaminoacetic acid, metallic salts (SCHREIBLER and BAUMGARTEN), A., i, 656.
- Benzylidene-*p*-aminobenzoic acid, ethyl ester (REDELLEN and DANILOF), A., i, 148.
- 1-*p*-Benzylideneaminophenyl-5-methylbenzthiazole (BOGERT and MEYER), A., i, 869.
- Benzylidene- $\alpha\beta$ -anhydrogluconic acid, ethyl ester and amide (LEVENE), A., i, 1029.
- 4-Benzylidene-2-anilino-5-ketoisooxazole, and its derivatives (WORRALL), A., i, 874.
- Benzylidene-*o*- and -*p*-anisidines, 2:4:6-trinitro- (LOWY and BALDWIN), A., i, 134.
- Benzylideneanthranilic acid, 2:4-dinitro- (EKELEY, ROGERS, and SWISHER), A., i, 935.
- Benzylideneanthranilic acids, action of acetic anhydride on (EKELEY, ROGERS, and SWISHER), A., i, 934.
- Benzylidenebenzhydrazide chloride, action of sodium azide with (STOLLÉ and NETZ), A., i, 690.
- Benzylidenebenzylamine, isomeric *mono*- and *di*-nitro-derivatives of (INGOLD and PIGGOTT), T., 2385.
- Benzylidenebis-*p*-nitrobenzaldehyde, *p*-nitro-, methylhydrazone (BRADY and McHUGH), T., 1651.
- Benzylidene-*p*-bromoaniline, *p*-hydroxy-, and *m*-nitro- (INGOLD and PIGGOTT), T., 2798.
- Benzylidene-camphor, *p*-amino-, and its salts, and *p*-chloro-, and *p*-cyano- (HALLER and BOUDIN), A., i, 356.
- Benzylidenechitosamic acid. See Benzylidene-glucosamic acid.
- Benzylidene-*o*- and -*p*-chloroanilines, 2:4:6-trinitro- (LOWY and BALDWIN), A., i, 134.
- Benzylidene-2:4-dichloroaniline, 2:4:6-trinitro- (LOWY and BALDWIN), A., i, 134.
- 1-Benzylidene-coumaran-2-ones, 4- and 6-chloro-, and their dibromides (MINTON and STEPHEN), T., 1602.
- Benzylidene-deoxygluconic acid, ethyl ester (LEVENE), A., i, 1029.
- Benzylidene-diacetoneglucose (FREUDENBERG and BRAUNS), A., i, 1117.
- Benzylidene-dicarbamic acid, ethyl ester (DATTA and CHATTERJEE), A., i, 818.
- Benzylidene-fluorene, 2-hydroxy-, glucoside of (BRUNI and ROMANI), A., i, 755.
- 9-Benzylidene-fluorene-2-carboxylic acid, 2:7-dibromo-, and its ethyl ester (SIEGLITZ and JASSOY), A., i, 821.
- Benzylidene-gluconic acid, β -bromo-, ethyl ester (LEVENE), A., i, 1029.

- Benzylidenegluconic acid**, α -chloro-, ethyl ester and amide (LEVENE), A., i, 1029.
- Benzylideneglucosamic acid**, and its ethyl ester (LEVENE), A., i, 1028.
- Benzylideneglycine** (SCHEIBLER and BAUMGARTEN), A., i, 656.
- 1-Benzylideneimino-5-phenyl-1:2:3:4-tetrazole**, and its iodochloride (STOLLÉ and NETZ), A., i, 690.
- Benzylideneindene**, 2-hydroxy-, glucoside of (DE FAZI), A., i, 755.
- Benzylideneisatinazine**, and *p*-nitro- (BORSCHÉ and MEYER), A., i, 54.
- 3-Benzylidenesulvulic acid**, and its ethyl ester, and their semicarbazones (KLEUCKER), A., i, 785.
- Benzylidenemannonic acid**. See **Benzylidenegluconic acid**.
- Benzylidenemethyl α -hydroxyisopropyl ketone**. See **α -Phenyl- δ -methyl- Δ -penten- δ -ol- γ -one**.
- Benzylidene-1-methylisatinazine**, and *p*-nitro- (BORSCHÉ and MEYER), A., i, 55.
- 2-Benzylidene-5-methylpyrrole-4-carboxylic acid**, 3-hydroxy-, ethyl ester (FISCHER and HERRMANN), A., i, 1054.
- Benzylidene- β -naphthylamine**, dimeride of (CIUSA), A., i, 1062.
- Benzylidene-*o*-nitroaniline**, and *m*- and *p*-nitro- (INGOLD and PIGGOTT), T., 2799.
- Benzylidene-*m*-nitroaniline**, *p*-hydroxy- (INGOLD and PIGGOTT), T., 2799.
- Benzylidene-*p*-nitroaniline**, *m*-nitro- (INGOLD and PIGGOTT), T., 2802.
- Benzylidenenitrobenzylamines**, *m*- and *p*-nitro- (INGOLD and PIGGOTT), T., 2801.
- Benzylidene-*o*- and -*p*-phenetidines**, 2:4:6-trinitro- (LOWY and BALDWIN), A., i, 134.
- Benzylidenephthalide-*o*-carboxylic acid**, methyl ester, and its dibromide (RUGGLI and MEYER), A., i, 344.
- Benzylidene-*dl*-piperitone** (READ and SMITH), T., 574.
- Benzylidene-*p*-toluidine**, *p*-nitro- (BOGERT and MEYER), A., i, 868.
- 4-Benzylidene-1:2:3-triethylresorcinol** (FABRE), A., i, 1148.
- Benzyliminodiacetic acid**, and its salts and derivatives (DUBSKY and DINGEMANN), A., i, 56.
- Benzyliminodiacetamide**. See **3:5-Diketo-1-benzyl-1:4-hexahydrodiazine**.
- Benzyliminodiacetonitrile**, and its hydrochloride (DUBSKY and DINGEMANN), A., i, 56.
- 3-Benzyl-2-indolinone**, and its 1-acetyl derivative, and 5:7-*di*bromo- (TOMIČEK), A., i, 679.
- Benzylmalonacidic acid** (CURTIUS and SIEBER), A., i, 722.
- Benzylmalonhydrazidic acid**, and its potassium salt and benzylidene derivative (CURTIUS and SIEBER), A., i, 722.
- Benzylmalonic acid**, *o*-nitro-, methyl ester (LEUCHS, v. KATINSKY, and CONRAD), A., i, 473.
- Benzyl-*p*-menthenol**, and its phenylurethane (READ and SMITH), T., 581.
- Benzylmethylamine**, *o*-hydroxy-, salts of (ZEMPLÉN and KUNZ), A., i, 564.
- Benzylmethylaniline**, and its oxide, picrates of (MEISENHEIMER and WILLMERSDORF), A., i, 334.
- 5-Benzyl 5-methylbarbituric acid** (DOX and YODER), A., i, 682.
- 2-Benzyl-2-methyl-2:3-dihydrofuran**, and 4(?)bromo- (HELPERICH and GEHRKE), A., i, 9.
- Benzylmethylethylamine**, and its oxide and their picrates (MEISENHEIMER and LOHSNER), A., i, 822.
- Benzylmethylethylhydroxylammonium salts** (MEISENHEIMER and LOHSNER), A., i, 822.
- 8-Benzylmethylheptenol** (HELPERICH and GEHRKE), A., i, 9.
- 4-Benzyl-1-methylcyclohexan-4-ol**, and 3-iodo- (TIFFENEAU and PORCHER), A., i, 537.
- 4-Benzyl-1-methylcyclo- Δ^3 -hexene** (TIFFENEAU and PORCHER), A., i, 537.
- 3-Benzylmethyl-2-indolinones** (TOMIČEK), A., i, 679.
- Benzyl methyl ketone**, *p*-hydroxy-, and its semicarbazone (LE BRAZIDEC), A., i, 456.
- 1-Benzyl-1-methylcyclopropane-2-carboxylic acid**, 2:3-dicyano-, and its amide (BIRCH and THORPE), T., 1830.
- 1-Benzyl-1-methylcyclopropane-2:3-dicarboxylic acid**, *cis*- and *trans*-forms, and their derivatives and 2:3-dicyanamide of (BIRCH and THORPE), T., 1830.
- N*-Benzylnor tropinone**, and its dibenzylidene derivative (MERCK, WOLFES, and MAEDER), A., i, 1173.
- Benzyl oxide**, sodium, reactivity of alkyl iodides with (HAYWOOD), T., 1904.
- 7-Benzyl-7:12-dihydro- γ -benzophenarazaine** (LEWIS and HAMILTON), A., i, 188.
- 5-Benzyl-3:4-dimethoxybenzaldehyde** (SPÄTH and RÖDER), A., i, 853.
- 5-Benzyl-3:4-dimethoxybenzoic acid** (SPÄTH and RÖDER), A., i, 853.
- β -5-Benzyl-3:4-dimethoxyphenylethylamine**, and its picrate (SPÄTH and RÖDER), A., i, 853.

- 5-Benzoyloxy-3,4-dimethoxystyrene**, *o*-nitro- (SPÄTH and RÖDER), A., i, 853.
- β-Benzoyloxytetraethylammonium iodide** (OLEMO and PERKIN), T., 649.
- 8-Benzoyltriethylamine** (CLEMO and PERKIN), T., 649.
- Benzylphenacylsulphone**, and its derivatives (WAHL), A., i, 654.
- 1-Benzylcyclopropane-1-carboxylic acid**, and its amide (HALLER and BENOIST), A., i, 350.
- 5-Benzyl-5-*n*-propylbarbituric acid** (DOX and YODER), A., i, 682.
- Benzylisopropylmalonic acid**, ethyl ester (DOX and YODER), A., i, 681.
- Benzylpyrasoleanthrone-yellow**, and *p*-chloro- and *o*-nitro- (MAYER and HEIL), A., i, 878.
- 1-Benzylpyridinium**, existence of (WEITZ and LUDWIG), A., i, 365.
hydrogen carbonate (WEITZ and KÖNIG), A., i, 1187.
- 3-Benzylsemicarbazide**, and its hydrochloride and benzylidene derivative (WILSON, HOPPER, and CRAWFORD), T., 868.
- Benzyl styryl ketone**, *p*-nitro-, and its oxime (KLEUCKER), A., i, 735.
- 2-Benzylthiol-1-*p*-anisyl-4-benzylidenehydantoin** (HILL and KELSEY), A., i, 1141.
- Benzylthiol-ψ-carbazinocarbothioxylic acid**, benzyl ester, and its benzoyl derivative (FROMM, KAYSER, BRIEGLER, and FÖHRENBACH), A., i, 880.
- 2-Benzylthiol-1-*p*-chlorophenyl-4-benzylidenehydantoin** (HILL and KELSEY), A., i, 1141.
- 2-Benzylthiol-1:4-dimethyldihydro-6-pyrimidone**, 2-*p*-nitro- (HORN), A., i, 375.
- 2-Benzylthiol-3-*p*-ethoxyphenyl-5-anilinomethylene-4-glyoxalone** (DAINE, THOMPSON, and ASENDORF), A., i, 1186.
- 2-Benzylthiol-4-methyldihydro-6-pyrimidone**, 2-*p*-nitro- (HORN), A., i, 375.
- 2-Benzylthiol-8-phenyl-5-anilinomethylene-4-glyoxalone** (DAINE, THOMPSON, and ASENDORF), A., i, 1186.
- 2-Benzylthiol-3-tolyl-5-anilinomethylene-4-glyoxalones** (DAINE, THOMPSON, and ASENDORF), A., i, 1186.
- 2-Benzylthiol-1-*m*-tolyl-4-benzylidenehydantoin** (HILL and KELSEY), A., i, 1142.
- S-Benzylthiosemicarbazide hydrochloride and sulphate** (WILSON and BURNS), T., 874.
- Benzyltrimethylammonium chloride**, *o*-hydroxy- (ZEMPLEN and KUNZ), A., i, 565.
- Benzylvinyl bromide** (PORCHER), A., i, 539.
- Berberine**, conversion of, into palmatine (SPÄTH and LANG), A., i, 166.
- Beri-beri**, avian, decrease in cell oxidation as the cause of (FLEISCH), A., i, 1215.
- Beryl**, Japanese (SHIBATA and URMURA), A., ii, 305.
- Beryllium**. See Glucinum.
- Beta vulgaris*. See Beetroot.
- Betaines** (PFEIFFER), A., i, 720; (PFEIFFER and HAEFELIN), A., i, 738.
- Betulin**, and its silver salt and hydrogen phthalate (SCHULZE and PIEROH), A., i, 1045.
- allo*Betulin, and its salts and esters (SCHULZE and PIEROH), A., i, 1045.
- apob*lloBetulin (SCHULZE and PIEROH), A., i, 1046.
- allo*Betulone, and its derivatives (SCHULZE and PIEROH), A., i, 1046.
- Bile acid**, unsaturated, preparation of an (RIEDEL), A., i, 1160.
- Bile acids** (BORSCHKE, WEICKEBT, and MEYER), A., i, 255; (WIELAND and SCHLICHTING), A., i, 554, 838; (RIEDEL), A., i, 554; (WIELAND and ADICKES), A., i, 838.
constitution of (BORSCHKE and HALLWASS), A., i, 1158, 1159.
unsaturated (BOEDECKER and VOLK), A., i, 1027.
detection of (LEPHENE), A., ii, 800.
- Bile pigments** (KÜSTER), A., i, 885; (KÜSTER and HERRMANN), A., i, 886.
action of intestinal bacteria on (PASINI), A., i, 966.
reactions of (ROSENTHAL and MEIER), A., i, 198.
detection of (IONESCU and POP), A., ii, 671.
detection of, in gastric juice (UTZ), A., ii, 799.
detection of, in urine (SILBERSTERN), A., ii, 799.
- Bile salts** (WIELAND and SCHULENBURG), A., i, 346.
- Billic acid**, diemicarbazone of (BORSCHKE and HALLWASS), A., i, 1159.
- Bilirubin**, preparation and reactions of (KÜSTER), A., i, 885.
constitution of (FROMHOLDT and NERSESSOV), A., i, 387.
formation of, in the spleen (ERNST and SZAFFANYOS), A., i, 1089.
azo-dyes from (FISCHER and BAREN-SCHEN), A., i, 278.
effect of injection of (FROMHOLDT and NERSESSOV), A., i, 404.

- Bilirubin**, estimation of, in blood (HOLZER and MEHNER), A., ii, 799.
estimation of, in serum (THANNHAUSER and ANDERSEN), A., ii, 671.
- Billverdin**, reactions of (KUSTER), A., i, 885.
- Biloidanic acid** (BORSCHKE, WICKERT, and MEYER), A., i, 255; (WIELAND and SCHLICHTING), A., i, 554.
- Binary systems**, influence of substitution on equilibria in (KREMANN and MÜLLER; KREMANN, ODELGA, and ZAWODSKY), A., i, 181; (KREMANN, HOHL, and MÜLLER), A., i, 188; (KREMANN and ODELGA), A., i, 159; (KREMANN and STRZELBA), A., i, 176; (KREMANN, HEMMELMAYR, and RIEMER), A., i, 1010.
- Biocolloids**, action of bases and salts on (MACDOUGALL), A., i, 204.
- Biological reactions**, physico-chemical studies on (HIRSCH and KUNZE), A., i, 781; (HIRSCH), A., i, 785.
- Biology**, importance of sequence in (KARCZAG; KARCZAG and HAJÓS), A., i, 302.
- Bioluminescence** (HARVEY), A., i, 299.
- Bird-lime**, Japanese (YANAGISAWA and TAKASHIMA; NISHIZAWA), A., i, 652.
- Birds' nests**, edible Chinese, constituents of (WANG), A., i, 299.
- Bis-5-acetyl-2:4-dimethylpyrrole 3'-disulphide** (FISCHER and HERRMANN), A., i, 1055.
- 4:4''-Bisazo-benzoyl ether**, 4'-hydroxy- and its acetyl derivative (FICHTER and JAECK), A., i, 62.
- 2:4-Bisbenzenesazophenol**, *mono-* and *di-*bromo-, -chloro-, and -iodo-derivatives (CHATTAWAY and HILL), T., 2758.
- Bis-1-benzyl-2:4-dimethylpyridinium** (WEITZ and KÖNIG), A., i, 1148.
- Bis-1-benzyl-2:4:6-trimethylpyridinium** (WEITZ and KÖNIG), A., i, 1188.
- Bis(3-carbethoxy-2:4- and -2:5-dimethylpyrrol)furylmethanes** (KÜSTER, WEBER, MAURER, NIRMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- Bis-3-carbethoxy-2:4-dimethylpyrrol-methane**. See Methenylbis-2:4-dimethylpyrrole-3-carboxylic acid, ethyl ester.
- p-Bis(3-carbethoxy-2:5-dimethylpyrrol)-methylbenzaldehyde**, and its hydrate (FISCHER and KAAH), A., i, 869.
- Bis(3-carbethoxy-2:4-dimethylpyrrol)-m-nitrophenylmethane** (KÜSTER, WEBER, MAURER, NIRMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- Bis- $\beta\beta'$ -dichlorodivinylchlorarsine-bis- $\beta\beta'$ -dichlorodivinylhydroxyarsine** platinichloride (MANN and POPE), T., 1759.
- Bis- β -chlorovinylbis- $\beta\beta'\beta''$ -trichlorotrivinylarsine**, platinum compound (MANN and POPE), T., 1758.
- Biscoumaric acids** (DE JONG), A., ii, 1049.
- Bisdiazobenzoylhistidine** (HERMANNS), A., i, 1092.
- Bis-4:4'-dibensthiazole**, 1:1'-dithiol- (ROMANI), A., i, 467.
- s*-Bisdibenzylmethylcarbamide** (JONES and SCOTT), A., i, 454.
- Bisdiethylaminoethylidiketocyclohexanedicarboxylic acid**, ethyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 525.
- Bisdihydrocarbestyryl-3:3'-spiran-6:6'-disulphonic acids**, and their barium salts (LEUCHS, CONRAD, and V. KATINSKY), A., i, 874.
- $\alpha\zeta$ -Bisdimethylamino- $\gamma\delta$ -diphenylhexane- $\gamma\delta$ -diols** (MANNICH and HEILNER), A., i, 351.
- Bis-*p*-dimethylaminotriphenylmethyl peroxide** (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Bisdiphenyldibiphenylenehydrazine** (WIELAND, WECKER, and ALBERT), A., i, 779.
- Bis-5-ethyl-5-methyltetrahydro-2-furyl ether** (HELFERICH and WEIDENHAGEN), A., i, 1115.
- Bisglucosyl ζ -selenides and ζ -sulphide**, and their derivatives (WEDE), A., i, 226.
- Bishydrocarbestyryl-3:3'-spiran**, and *p-mono-* and *pp'-di*hydroxy-, and their derivatives (LEUCHS, V. KATINSKY, and CONRAD), A., i, 471.
- Bishydrocoumarin-3:3'-spiran** (LEUCHS, V. KATINSKY, and CONRAD), A., i, 473.
- 3:3'-Bis-*o*- α -dihydroxybenzylamino-*di*-hydroxyarsenobenzene**, and its dihydrochloride (RAIZISS and BLATT), A., i, 1079.
- p-Bis- β -hydroxyethylaminebenzoic acid**, ethylester (ALTWEGG and LANDRIVON), A., i, 1023.
- 3:3'-Bis- α -hydroxy-*p*-methoxybenzylamino-4:4'-*di*hydroxyarsenobenzene** (RAIZISS and BLATT), A., i, 1079.
- 3:3'-Bis-*p*- α -*di*hydroxy-*m*-methoxybenzylamino-4:4'-*di*hydroxyarsenobenzene** (RAIZISS and BLATT), A., i, 1079.
- 3:3'-Bishydroxymethylamino-4:4'-*di*-hydroxyarsenobenzene dihydrochloride** (RAIZISS and BLATT), A., i, 1079.

Bishydroxymethyleneacetonedianilide.

See Dianilinovinyl ketone.

3:3'-Bis- α -hydroxy-*m*-nitrobenzylamino-

4:4'-dihydroxyarsenobenzene (RAIZISS and BLATT), A., i, 1079.

3:3'-Bis- α -hydroxy- γ -phenylallylamino-

4:4'-dihydroxyarsenobenzene (RAIZISS and BLATT), A., i, 1080.

Bis-1:3'-indil (*bis-1:3-di-indoyl*) (ODDO and SANNA), A., i, 371.**3:2'-Bis-*p*-methoxybenzylideneamino-**

1:1'-dinaphthyl (JACOBSEN), A., i, 594.

Bis-2-methyl-1-naphthoylhydrazine

(MAYER and SIEGLITZ), A., i, 742.

Bis-5-methyltetrahydro-2-furyl ether

(HELFERICH and WEIDENHAGEN), A., i, 1115.

Bis-6-methyltetrahydro-2- α -pyryl ether

(HELFERICH and WEIDENHAGEN), A., i, 1115.

Bismuth, spectrum of (NAGAOKA and SUGIURA), A., ii, 101.

L-series spectrum of (DUANE and PATTERSON), A., ii, 463.

N-series spectrum of (DOLEJSEK), A., ii, 463.

anodic corrosion of (PRIDEAUX and HEWIS), A., ii, 511.

liquid, density and surface tension of (HOGNESS), A., ii, 29.

solubility of lead in (DI CAPUA), A., ii, 576.

Bismuth compounds, preparation of (PRIDEAUX and HEWIS), A., ii, 511.**Bismuth salts, treatment of syphilis with** (SAZERAC and LEVADITI), A., i, 89;

(FOURNIER and GUENOT), A., i, 301.

Bismuth thallous bromide and iodide (CANNONI and PERINA), A., ii, 512.

trichloride, compounds of diazonium chlorides with (CHALLENGER and WILKINSON), T., 102.

hydrides (PANETH, JOHANNSEN, and MATTHIES; PANETH, MATTHIES, and SCHMIDT-HEBBEL), A., ii, 383.

subiodide and *suboxyiodide* (DENHAM), A., ii, 218.

oxide, fusion diagram of mixtures of lead oxide and (BELLADEN), A., ii, 777.

Bismuth organic compounds (CHALLENGER and WILKINSON), T., 91; (CHALLENGER and RIDGWAY), T., 104.

action of thallic chloride on (D. and A. E. GODDARD), T., 260.

dimethoxide and dimethyl (DENHAM), A., ii, 218.

Bismuthi iodides of alkaloids (FRANÇOIS and BLANC), A., i, 851.

Bismuth detection and estimation:—

detection of, by mirrors (SCHEUCHER), A., ii, 526.

detection of, in urine (AUBRY), A., ii, 165.

detection of, in urine and in saliva (GANASSINI), A., ii, 590.

detection and estimation of (CANNONI and PERINA), A., ii, 512.

estimation of (CRITCHETT), A., ii, 90.

metallic, estimation of (KURTENACKER and WERNER), A., ii, 877.

Bismuth minerals, Spanish, chemical and spectrographic study of (PIÑA DE RÚBIES and GILA ESTEBAN), A., ii, 576.**Bismuthines, tertiary aromatic** (CHALLENGER and WILKINSON), T., 91; (CHALLENGER and RIDGWAY), T., 104.***aa'*-Bisphenacylmethylmethylamine, and its hydrochloride** (MANNICH and HEILNER), A., i, 351.**Bis- γ -phenoxy-*n*-butyrylamine** (PUTOCHIN), A., i, 1176.***s*-Bistriphenylmethylhydrazine, molecular rearrangement of** (STIEGLITZ and BROWN), A., i, 778.**Bisulphites.** See Sulphites under Sulphur.**Bis*isovalerylamino*ethane** (WINDAUS, DÖRRIES, and JENSEN), A., i, 60.**Bis*isovalerylamino*ethylene, and its dibromide** (WINDAUS, DÖRRIES, and JENSEN), A., i, 60.**Biuret, preparation of derivatives of** (FROMM and WENZL), A., i, 437.

compound of copper ethylenediamine hydroxide with (TRAUBE), A., i, 719.

Blackberry. See *Rubus fruticosus*.**Bladder, silica calculus in the** (SCHLICHT), A., i, 496.**Bleaching chlorides, estimation of the available chlorine in** (ROYER), A., ii, 581.**Blende, oxidation of** (HELBRONNER and RUDOLFS), A., i, 706.**Blood, effect of poisons on the absorption spectrum of** (LEWIN and STENGER), A., ii, 414.

effect of radium emanation on (HAUENSTEIN), A., i, 80.

condition of electrolytes in (NEUHAUSEN), A., i, 891; (NEUHAUSEN and MARSHALL), A., i, 1085.

viscosity of, in relation to the distribution of cholesterol and of uric acid between plasma and corpuscles (ROUZAUD and THIÉRY), A., i, 394.

Blood, osmotic resistance and phosphatides of (BRINKMAN), A., i, 1086.
 increased supply of, during muscular activity (HARRIS), A., i, 696.
 effect of profuse perspiration on (WILBRAND), A., i, 79.
 chemistry of (EGERER-SEHAM and NIXON), A., i, 395.
 in pregnancy (KILLIAN and SHERWIN; CALDWELL and LYLE), A., i, 489.
 acetone substances in, after injection of adrenaline (HUBBARD and WRIGHT), A., i, 288.
 in diabetes (FITZ), A., i, 192.
 neutralisation of acids by (KENNAWAY and MCINTOSH), A., i, 788.
 effect of ether anaesthesia on the acid-base balance of (VAN SLYKE, AUSTIN, and CULLEN; STEHLE, BOURNE, and BARBOUR), A., i, 1085.
 amino-acids in (COSTANTINO), A., i, 181; (ABDERHALDEN), A., i, 287; (OKADA and HAYASHI), A., i, 483.
 ammonia in (NASH and BENEDICT), A., i, 191; (NASH), A., i, 483.
 calcium in (MAZZOCCHI), A., i, 788, 891; (VAN PAASSEN), A., i, 1208.
 of the new-born (JONES), A., i, 80.
 distribution of calcium, magnesium, potassium and sodium in (KRAMER and TISDALL), A., i, 1087.
 carbon dioxide equilibrium and hydrogen-ion concentration in (WARBURG), A., i, 788.
 effect of potassium cyanide on catalase in (WELKER and BOLLMAN), A., i, 192.
 distribution of chlorides in (MUKAI; VAN CREVELD; NORGAARD and GRAN), A., i, 287; (MUKESANU), A., i, 260; (DISCHER), A., i, 1086.
 distribution of chlorine in (SIEBECK and HACKMACK), A., i, 1208.
 chloroform in, after death (GIBSON and LAIDLAW), A., i, 1218.
 cholesterol derivatives in (LIFSCHÜTZ), A., i, 392.
 coagulation of (FUNCK), A., i, 290; (HELLER), A., i, 291; (STUBER and FUNCK), A., i, 393; (PICKERING and HEWITT), A., i, 393, 1208.
 effect of carbon monoxide, illuminating gas and benzene on (FORBES and HOMPE), A., i, 483.
 effect of alkaline earth chlorides on (LEHMANN), A., i, 892.
 by euphylline (MEISSNER), A., i, 193.

Blood, action of nucleic acids on (DORON), A., i, 82, 1087.
 action of phosphatides on (ZUNZ and LA BARRÉ), A., i, 697.
 presence of cyanic acid in (NICLOUX and WELTER), A., i, 789.
 enzymes in (COMPTON), A., i, 392, 1087; (BACH and ZUBKOWA), A., i, 393; (FINCUSSEN), A., i, 393.
 transformation of the fat in (LIFSCHÜTZ), A., i, 392.
 action of germanium oxide on (HAMNETT, NOWREY, and MÜLLER), A., i, 1086.
 physical chemistry of hæmoglobin in (BROWN and HILL), A., i, 1189.
 preparation of histidine from (DEM-JANOVSKI), A., i, 1052.
 distribution of hormones and drugs in (SCHAEFFER), A., i, 290.
 reaction of, with nascent hydrogen peroxide (PATZAUER), A., ii, 172.
 inorganic ions in, in tetany (GROSS and UNDERHILL), A., i, 1210.
 lipoids of in tuberculosis (HENNING), A., i, 963.
 nitrogen constituents of, in chronic nephritis (WILLIAMS), A., i, 401.
 inorganic phosphates in (LEHMAN), A., i, 191.
 effect of pilocarpine on the composition of (BORNSTEIN and VOGEL), A., i, 80.
 proteins of, and their part in the transport of carbon dioxide (AUSSENDA), A., i, 1209.
 action of purine diuretics on the concentration of (NONNENBRUCH), A., i, 79.
 distribution of quinine in (RONA and BLOCH), A., i, 290.
 hæmolytic of, by quinine (LUGER), A., i, 81.
 action of salvarsan on the serum and corpuscles of (KRITSCHESKY), A., i, 394.
 effect of administration of sodium chloride on (SAMSON), A., i, 82.
 sugar in (CAMMIDGE, FORSYTH, and HOWARD), A., i, 81; (FALTA and RICHTER-QUITTNER), A., i, 696; (ROSENBERG), A., i, 789; (STASIAR), A., i, 1208.
 in narcosis and disease (CHANTRAINE), A., i, 192.
 action of adrenaline on (PETÉNYI and LAX), A., i, 408.
 effect of amino-acids and fatty acids on (POLLAK), A., i, 483.
 effect of carbon dioxide on (BINSWANGER), A., i, 962.
 action of phosphates on (ELIAS and WEISS), A., i, 1085.

Blood, effect of sugar injection on the sugar content and the white corpuscles of (MORACZEWSKI and LINDNER), A., i, 402.
 arterial and venous, distribution of sugar in (HENRIQUES and EGE), A., i, 191; (TURBAN), A., i, 482.
 capillary and venous, sugar distribution in (NEUWIRTH and KLEINER), A., i, 1208.
 fate of sulphides in (HAGGARD and CHARLTON), A., i, 287.
 distribution of urea in (ETIENNE and VÉRAIN), A., i, 963.
 distribution of uric acid in (THEIS and BENEDICT), A., i, 82; (CHAUFFARD, BRODIN, and GRIGAUT), A., i, 1086.
 free and bound water in (NEUHAUSEN), A., i, 696.
 morphological changes of (PENTIMALLI), A., i, 302.
 defecation of, for the estimation of carbamide (IONESCU), A., ii, 596.
 respiratory changes in (AUSTIN, CULLEN, HASTINGS, MCLEAN, PETERS, and VAN SLYKE; PETERS, CULLEN, and AUSTIN), A., i, 1207.
 calves', concentration of proteins in (HOWE), A., i, 1084.
 effect of ingestion of colostrum on (HOWE), A., i, 80.
 of children in tetany, calcium in (BORLES), A., i, 192.
 dogs', lactic acid in (HASTINGS), A., i, 81.
 human, hydrogen-ion concentration in (BARCROFT, BOCK, HILL, PARSONS, and SHOJI), A., i, 891.
 partial pressure of oxygen in (BARCROFT and NAGAHASHI), A., i, 890.
 reducing power of (COOPER and WALKER), A., i, 1084.
 uric acid in (MORRIS and MACLEOD), A., i, 892.
 shed, acid-production in (EVANS), A., i, 890.
 reaction of (DE CORRAL), A., i, 79; (CONWAY and STEPHEN), A., i, 892.
Blood detection and estimation :—
 analysis of (RICHTER-QUITTNER), A., ii, 240, 407; (PINCUSSEN and FLOROS; PINCUSSEN and MOMFERATOS-FLOROS), A., ii, 408; (FOLIN and WU), A., ii, 540; (FOLIN), A., ii, 596; (UTZ), A., ii, 800.
 analyses of the corpuscles and plasma of (WU), A., i, 483.
 analysis of gases of (MEIER and KRÖNIG), A., i, 191.

Blood detection and estimation :—
 collection and analysis of (AUSTIN, CULLEN, HASTINGS, MCLEAN, PETERS, and VAN SLYKE), A., i, 1207.
 detection and estimation of, in urine (JOHANNESSEN), A., ii, 724.
 estimation of acetone substances in (HUBBARD), A., ii, 284.
 estimation of fatty acids in (BLOOR, PELKAN and ALLEN), A., ii, 593.
 estimation of amino-acids in (PETIT-JEAN), A., ii, 536.
 estimation of ammonia in (GAD-ANDRESEN), A., ii, 523.
 estimation of bilirubin in (HOLZER and MEHNER), A., ii, 799.
 estimation of calcium in (WEISS: CLARK), A., ii, 227; (LING and BUSHILL), A., ii, 587.
 estimation of carbon monoxide in (SAYERS and YANT), A., ii, 868.
 estimation of chlorides in (FRIEND), A., ii, 389; (PETSCHACHER; ISAACS), A., ii, 716.
 estimation of creatine and creatinine in (BEHRE and BENEDICT), A., i, 789.
 estimation of ethyl alcohol in (WIDMARK), A., ii, 789.
 estimation of fibrin in (GRAM), A., ii, 240.
 estimation of fibrinogen in (GRAM), A., i, 288; (LEENDERTZ and GROMELSKI), A., ii, 798.
 estimation of gases in (VAN SLYKE and STADIE; STADIE), A., ii, 78.
 apparatus for estimation of gases in (VAN SLYKE), A., ii, 158.
 estimation of hydrogen ions in (LINDHARD), A., ii, 240.
 estimation of hydrogen carbonate and hydrogen ions in plasma and (VAN SLYKE; CULLEN), A., ii, 672.
 estimation of iron in (BROWN), A., ii, 319.
 estimation of lecithin in (RANDLES and KNUDSON), A., ii, 719.
 estimation of lipoids in (LEMELAND), A., ii, 666.
 estimation of magnesium in (DENIS), A., ii, 659.
 estimation of non-protein nitrogen in (PONDER), A., ii, 583; (RICHTER-QUITTNER and HOENLINGER), A., ii, 717.
 estimation of the oxygen-combining power of (LUNDGAARD and MÖLLER), A., ii, 724.
 estimation of phenols in (PELKAN), A., ii, 399.
 estimation of phosphoric acid in (BLOOR), A., ii, 84.

Blood detection and estimation :—

- estimation of proteins in (HOWE), A., ii, 171, 172.
- estimation of proteins and nitrogen in (BIERRY and MOQUET), A., ii, 886.
- estimation of sugar in (ROSENBERG), A., i, 482; (GUY), A., ii, 94; (TERVAERT), A., ii, 186; (REIST), A., ii, 323; (STEFF), A., ii, 592; (LABBÉ, NEPVEUX, and NOMIDIS), A., ii, 663; (ERNST and WEISS), A., ii, 724; (CSONKA and TAGGART), A., ii, 879.
- estimation of sulphates in (DENIS), A., ii, 225.
- estimation of urea in (NICLOUX and WELTER), A., ii, 170; (GAD-ANDRESEN), A., ii, 536.
- ureometer for estimation of urea in (CLOGNE), A., ii, 237.
- estimation of uric acid in (GUILLAUMIN), A., ii, 170; (GRIGAUT; BENEDICT), A., ii, 405; (PUCHER), A., ii, 668.
- estimation of uric acid and urates in (BAUMAN and KEELER; GUILLAUMIN), A., ii, 796.
- Blood-corpuscles**, effect of narcotics on the charge on (MEIER and KRÖNIG), A., i, 191.
- agglutination and sedimentation of (STARLINGER; RUNNSFROM), A., i, 289.
- volume curve of, in hypertonic solutions (TAKEL), A., i, 289.
- stability of suspensions of (BRINKMAN and WASTL), A., i, 289.
- effect of extirpation of glands on the development of (ASHER and MATSUMO), A., i, 298.
- permeability of, for dextrose (VAN CREVELD and BRINKMAN), A., i, 192.
- hemolysis of, by eosin (SCHMIDT and NORMAN), A., i, 892.
- fixation of quinine by (RONA and BLOCK), A., i, 484.
- red, colloidal structure of (HATTORI), A., i, 192.
- osmotic resistance curves of (HAMBURGER), A., i, 606.
- permeability of (WIECHMANN), A., i, 80, 289.
- influence of potassium and radioactive substances on the oxidation velocity of (ELLINGER), A., i, 288.
- human, dextrose content of (BÜNNIGER), A., i, 606.
- white, glycogen content of (DE HAAN), A., i, 484.

- Blood-corpuscles**, estimation of non-hæmoglobin nitrogen in (SCHOEN), A., i, 484.
- Blood-louse**. See *Schizoneura lanigera*.
- Blood-pigments** (KÜSTER and GERLACH), A., i, 596; (KUSTER), A., i, 884.
- Blood-plasma**, combination of, with carbon dioxide (TAISTRA; CHANUTIN), A., i, 288.
- sugar in, in relation to sugar in cerebrospinal fluid and aqueous humour (DE HAAN and VAN CREVELD), A., i, 295.
- human, flocculation of (STARLINGER), A., i, 288.
- estimation of phosphates in (DENIS and v. MEYSENBUG), A., ii, 584.
- estimation of inorganic phosphorus in (MYERS and SHEVSKY), A., ii, 455.
- Blood pressure**, effect of adrenaline on (RICHAUD), A., i, 891.
- Blood-serum**, ultra-violet absorption spectra and optical rotation of the proteins of (LEWIS), A., ii, 245.
- buffer systems in (DOISY, EATON, and CHOUKE), A., i, 963.
- calcium in, in pregnancy (MAZZOCCO and MORON), A., i, 892.
- Blood-serum**, cholesterol in (STRAUSS and SCHUBARDT), A., i, 1209.
- isolation of the coagulating enzyme from (BLEIBTREV), A., i, 1084.
- colloidal equilibrium of (FISCHER), A., i, 1209.
- action of soaps with (JARISCH), A., i, 1087.
- estimation of atropine in (VAN DER HEYDE), A., ii, 669.
- estimation of globulins in (HENLEY), A., ii, 671.
- estimation of indican in (SNAPPER and VAN VLOTEN), A., ii, 793.
- estimation of salicylic acid in (HERISSEY), A., ii, 880.
- Blood-vessels**, action of calcium and potassium ions on (HAMBURGER), A., i, 606.
- Blumea Malcomii*, essential oil from (SIMONSEN and RAU), T., 876.
- Body fluids**, physical chemistry of (RUSZNYÁK and HETÉNYI), A., i, 291.
- sulphates and sulphuric acid in (HEUBNER and MEYER-BISCH), A., i, 291.
- Boiling point**, relation of melting point to (LORENZ and HERZ), A., ii, 739.
- relation between melting point, critical temperature and (PRUD'HOMME), A., ii, 349.
- Boldo leaf oil**, constituents of (ROURE-BERTRAND FILS), A., i, 846.

- Bone fat**, analytical constants of (ECKART), A., i, 967.
- Borneol**, preparation of (FABRIQUES DE PRODUITS CHIMIQUES DE THANN ET DE MULHOUSE), A., i, 560.
from pinene hydrochloride (VAVON and BEITON), A., i, 943.
- Borneolcarboxylic acids** (BREDT), A., i, 937.
- n*- and *iso*-Bornyl chloride, equilibrium of camphene hydrochloride with (MEERWEIN, VAN EMSTER, and JOUSSEN), A., ii, 751.
- Bornylene**, preparation of (MEERWEIN and JOUSSEN), A., i, 1042.
- Bornylsemicarbazide** (NEIGHBORS, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.
- Boric acid**. See under Boron.
- Boroacetic acid**, hydroxyanthraquinone esters of (DIMROTH and FAUST), A., i, 156.
- Boron**, atomic weight of (BAXTER and SCOTT), A., ii, 285; (HÖNIGSCHMID and BIRCKENBACH), A., ii, 641.
New Zealand, isotopes of (MONRO), T., 986.
Röntgen rays from (HUGHES), A., ii, 184; (HOLTMARK), A., ii, 543.
- Boron alloys** with carbon and iron (VOGEL and TAMMANN), A., ii, 852.
- Boron nitride**, reaction of, with metallic oxides (SBORGI and NASINI), A., ii, 564.
trioxide, phosphorescence of hydrated (TIEDE and WULFF), A., ii, 245.
- Boric acid**, organic phosphors of (TIEDE, WULFF, and RAGOSS), A., ii, 8; (TOMASCHKE), A., ii, 763.
effect of hydroxy-organic compounds on the hydrogen-ion concentration of (RIMBACH and LEY), A., ii, 342.
and its salts, complex compounds of, with fructose and with mannitol (GILMOUR), T., 1333.
action of, on mannitol (DUBRISAY), A., i, 1110.
hydroxyanthraquinone esters of (DIMROTH and FAUST), A., i, 155.
estimation of (STRECKER and KANNAPPEL), A., ii, 784.
estimation of, in presence of phosphoric acid (KOLTHOFF; DEERNS), A., ii, 867.
- Borates** (SBORGI and FERRI), A., ii, 568, 764.
dissociation and hydrolysis of, in solution (MENZEL), A., ii, 345.
- Polyborates** (ROSENHEIM and LEYSER), A., ii, 50.
- Boron** :—
Hexahydrodioxidiboron, potassium salt (RAY), T., 1088.
- Boron organic compounds**, preparation of (KRAUSE and NITSCHÉ), A., i, 22.
trialkyls (KRAUSE and NITSCHÉ), A., i, 22.
mono- and di-phenyl fluorides and tri-phenyl (KRAUSE and NITSCHÉ), A., i, 694.
- Brain**, constituents of (SHIMIZU), A., i, 85.
phosphosulphatides and lecithin from (FRÄNKEL and GILBERT; FRÄNKEL and KÄSZ), A., i, 294.
zinc and copper compounds in (BODANSKY), A., i, 194.
- Brass**, protection of, from corrosion (MANLEY), A., ii, 840.
estimation of antimony in (EVANS), A., ii, 231.
red, estimation of antimony and tin in (MÜCK), A., ii, 722.
- Brassicidic acid**, ethyl ester and anhydride of (HOLDE and SCHMIDT), A., i, 982.
- iso*Brazilein, synthesis of (CRABTREE and ROBINSON), T., 1033.
- Bridge linkings**, rupture of (WAHL), A., i, 653.
- Bromates**. See under Bromine.
- Brometone**. See *tert*-Butyl alcohol, *tribromo*-.
- Bromic acid and Bromides**. See under Bromine.
- Bromination** of organic acids (WARD), T., 1161.
- Bromine**, atomic weight of (MOLES), A., ii, 141; (HAXIER), A., ii, 370.
spectrum of the vapour of (RIBAUD), A., ii, 3.
solid, vapour pressure of (HENGLEIN, v. ROSENBERG, and MUCHLINSKI), A., ii, 760.
water, action of, on indene (READ and HURST), T., 2550.
- Hydrobromic acid**, heats of neutralisation of (RICHARDS and ROWE), A., ii, 425.
gaseous, density of (MOLES), A., ii, 140.
- Bromides**, retention and excretion of, in the organism (BAUR and OPPENHEIMER), A., i, 968.
estimation of, with silver nitrate (KOLTHOFF), A., ii, 581.
estimation of chlorides and, in presence of iodides (KOLTHOFF), A., ii, 79.
- Bromic acid**, estimation of (KIKUCHI), A., ii, 519.

- Bromine:**—
Bromates, reflection spectra of (SCHAEFER and SCHUBERT), A., ii, 180.
Bromine organic compounds, preparation of, without production of hydrogen bromide (GUPTA and THORPE), T., 1896.
Bromine estimation:—
 estimation of, in animal tissues (DAMIENS), A., ii, 79.
Bromoform, vapour pressure of mixtures of ethyl ether and (DOLEZALEK and SCHULZE), A., ii, 118.
 action of, on magnesium alkyl haloids (BINAGHI), A., i, 1002.
Brownian movement, theory of (JÄGER), A., ii, 30.
Brucine hexosophosphate (NEUBERG and DALMER), A., i, 920.
Brucinesulphonic acid, quinols and quinones from, and their derivatives (LEUCHS and FRICKER), A., i, 677.
isoButaldehyde, α -bromo-, action of sodiomalonic ester with (FRANKK and GROEGER), A., i, 808.
Butane, $\alpha\beta$ -tri-amino-, salts of (WINDAUS, DORRIES, and JENSEN), A., i, 60.
 β -bromo- and β -chloro-hydroxy- (FOURNEAU and PUYAL), A., i, 639.
 α -chloro- β -hydroxy- (HELPERICH and SPEIDEL), A., i, 6.
cycloButane-1-carboxylic acid, 1-hydroxy-, pyrogenic decomposition of, and its anhydrides (DEMJanov and DOJARENKO), A., i, 1161.
cycloButane-1:1-dicarboxylamide (INGOLD, SAKO, and THORPE), T., 1191.
cycloButane-1:1-dicarboxylbromoamide (INGOLD, SAKO, and THORPE), T., 1191.
Butane- α -diol- $\alpha\beta$ -dicarboxylic acid, and its salts and derivatives (INGOLD), T., 2694.
5-cycloButanespirohydantoin (INGOLD, SAKO, and THORPE), T., 1191.
cycloButanone, preparation of (DEMJanov and DOJARENKO), A., i, 1161.
 Δ^{α} -Butene- $\alpha\beta$ -dicarboxylic- β -acetic acids, and their silver salts and derivatives (FARMER), T., 2019.
 Δ^{α} -Butene- $\alpha\beta$ -dicarboxylic- β -malonic acid, ethyl esters, and sodium salt (FARMER), T., 2017.
 Δ^{α} -Buten- γ -ol, properties of (BAUDRENGHIEN), A., i, 710.
Butenonitriles (BRUYLANTS), A., i, 817, 924.
 β -Butoxybutyronitriles (BRUYLANTS), A., i, 924.
7-n-Butoxy-7:12-dihydro- γ -benzophenarsazine (LEWIS and HAMILTON), A., i, 188.
- 6-n- and -iso-Butoxy-2-methylquinoline**, and its ethiodide (GUTEKUNST and GRAY), A., i, 951.
Butter, fatty acids of (FROG and SCHMIDT-NIELSEN), A., i, 494.
 detection of coconut oil in (MUTTELET), A., ii, 236.
isoButylacetylmalonic acid, ethyl ester, and its phenylhydrazone (GAULT and SALOMON), A., i, 428.
n-Butylalanine. See α -Methyl-n-hexonic acid, α -amino-.
n-Butyl alcohol (TSUJIMOTO and TOYAMA), A., i, 297.
 preparation of (GRÜNSTEIN), A., i, 112.
 chlorination of (GAULT and GUILLEMET), A., i, 910.
isoButyl alcohol, solubility of, in water (MICHELS), A., ii, 486.
tert.-Butyl alcohol, tribromo-, and trichloro-, benzoyl and nitrobenzoyl esters (ALDRICH and BLANNER), A., i, 910.
n- and iso-Butylammonium ruthenipentabromides (GUTHRIE and KRAUSS), A., i, 16.
dl-p-sec.-Butylaniline, resolution of (GLATTFIELD and WERTHEIM), A., i, 385.
5-n-Butylbarbituric acid, and 5-bromo-, ethyl ester (DOX and YODER), A., i, 808.
5-sec.-Butylbarbituric acid (DOX and YODER), A., i, 817.
isoButylboric acid (KRAUSE and NITSCHKE), A., i, 22.
Butyldichloroarsine (QUICK and ADAMS), A., i, 601.
Buchu-camphor, preparation of (ASHINA and MITUHOI), A., i, 667.
Buckwheat, glucenasein of (KIESEL), A., i, 412.
Buffer solutions (VAN SLYKE), A., i, 893.
 for colorimetry (ACREE, MELLON, AVERY, and SLAGLE), A., ii, 157.
 systems in blood-serum (DOIRY, EATON, and CHOUKE), A., i, 963.
Bufotalan (WIELAND), A., i, 200.
Bufotalidin (WIELAND), A., i, 200.
Bufotalin, and its acetyl derivative (WIELAND), A., i, 200.
Bufotalone (WIELAND), A., i, 200.
Bufotoxin (WIELAND and ALLES), A., i, 784.
Burner, rotary (LOCKEMANN), A., ii, 43.
Butadiene, formation of, from ethylene (ZANETTI, SUYDAM, and OFFNER), A., i, 977.

Butaldehyde, preparation of (GRÜN-STEIN), A., i, 112; (ADAM and LEGG), A., i, 222.

***n*-Butyl-*NN'*-dimethylmalonamide** (Dox and YODER), A., i, 808.

$\Delta\alpha$ -Butylene, action of selenium chloride on (BOORD and COPE), A., i, 421.

$\Delta\alpha$ -Butylene, γ -chloro-(BAUDRENGHIEN), A., i, 710.

α -iodo-(KAUFMANN and SCHWEITZER), A., i, 214.

$\Delta\beta$ -Butylene, α -bromo-, and α -chloro- (BAUDRENGHIEN), A., i, 710.

***iso*Butylene- $\alpha\gamma\gamma'$ -tetracarboxylic acid**, ethyl ester (INGOLD and NICKOLLS), T., 1643.

***iso*Butylene- $\alpha\gamma\gamma'$ -tricarboxylic acid**, and its ethyl ester (INGOLD and NICKOLLS), T., 1644.

Butylhydroxymalonic acid, β -hydroxy-, and its barium salt (HELFERICH and SPEIDEL), A., i, 7.

γ -*iso*Butylidenemalonolactone, ethyl ester (FRANKE and GROEGER), A., i, 808.

***sec*-Butylmalonamide** (Dox and YODER), A., i, 816.

***n*-Butylmalonic acid**, and bromo-, derivatives of (Dox and YODER), A., i, 808.

***sec*-Butylmalonic acid**, methyl ester (Dox and YODER), A., i, 816.

***cyclo*Butylmethyl dimethylamine**, and its salts (DEMJEANOV and DOJARENKO), A., i, 997.

***iso*Butyl methyl ketone**, α -chloro- β -hydroxy- (*menthyl oxide chlorohydrin*) (PASTUREAU and BERNARD), A., i, 717.

***dl*-*p*-*sec*-Butylphenylhydrazine**, and its derivatives (GLATFIELD and WERTHEIM), A., i, 386.

***tert*-Butyl propyl ketone** (LEROIDE), A., i, 216.

2-*n*-Butylthiol-3-phenyl-5-anilino-methylene-4-glyoxalane (DAINS, THOMPSON, and ASENDORF), A., i, 1186.

Butyranilide, tribromo- (VAN PESKI), A., i, 106.

***iso*Butyrylhydroxamic acid**, and its salts and derivatives (JONES and SCOTT), A., i, 454.

Butyric acid, preparation of (ADAM and LEGG), A., i, 222.

formation of, by fermentation (NEUBERG and ARINSTEIN), A., i, 91.

oxidation of, by hydrogen peroxide (WITZEMANN), A., i, 6.

estimation of, in mixtures with acetic acid (NOTENBAART), A., i, 791.

***n*-Butyric acid**, α -amino-, and its derivatives (CURTIUS and SIEBER), A., i, 723.

β -hydroxy-, estimation of (HUBBARD), A., ii, 234.

***dl*-Butyric acid**, $\alpha\gamma$ -*d*ihydroxy-, preparation and resolution of (GLATFIELD and SANDER), A., i, 318.

***iso*Butyric acid**, α -hydroxy-, action of thionyl chloride on (BLAISE and MONTAGNE), A., i, 715.

Butyrolactone, chloro-, and its acetal (CARRIÈRE), A., i, 319.

Butyrophenol, tribromo- (VAN PESKI), A., i, 106.

***iso*Butyryl chloride**, α -chloro-, and its anilide (BLAISE and MONTAGNE), A., i, 716.

***iso*Butyryloxyisobutyryl chloride**, α -chloro-, and its anilide (BLAISE and MONTAGNE), A., i, 716.

***n*-Butyrylsulphuric acid** (VAN PESKI), A., i, 106.

Bystropogon canus, constituents of (WHELAN), A., i, 847.

C.

Cacotheline, preparation and derivatives of (LEUCHS, OSFERNBURG, and KAEHRN), A., i, 362.

degradation of, by bromine, and its derivatives (LEUCHS, MILDBRAND, and LEUCHS), A., i, 1052.

colour reactions of (LEUCHS and KAEHRN), A., i, 463.

Cactus alkaloids (SPATH), A., i, 163, 567; (SPATH and RÖDER), A., i, 352.

Cadalene, and its synthesis and derivatives (RUZICKA, MEYER, and MINGAZZINI), A., i, 560; (RUZICKA and SEIDEL), A., i, 562.

methylisopropylaphthalenes from (RUZICKA and MINGAZZINI), A., i, 1001.

Cade oil from *Juniperus oxycedrus* (HUIERRE), A., i, 505.

Cadmium, band spectrum of (HULTHÉN and BENGTSSON), A., ii, 677.

fluorescence of the vapour of (VAN DER LINGEN), A., ii, 181.

heat of solution of, in hydrochloric acid (RICHARDS and TAMARU), A., ii, 475.

rate of solution of, in ferric alum (COLLENDERG and BODFORS), A., ii, 431.

solubility of lead in (DI CAPUA), A., ii, 576.

reductions with, in volumetric analysis (TREADWELL), A., ii, 780, 788.

Cadmium alloys with magnesium (GUILLET), A., ii, 570.

heat of formation of (BILTZ and HOHORST), A., ii, 350.

with mercury, use of, in analysis (KANÓ), A., ii, 529.

Cadmium iodide, crystal structure of (BOZORTH), A., ii, 851.

nitrate, mixed crystals of calcium

nitrate and (HASSELBLATT), A., ii, 61.

nickel nitrites, complex (CUTTICA), A., ii, 449.

alkali selenates (TUTTON), A., ii, 502.

sulphate, specific heat of (COHEN and MOESVELD), A., ii, 347.

Cadmium organic compounds :—

dimethyl, preparation of (DE MAHLER), A., i, 240.

Cadmium estimation :—

estimation of, volumetrically, with ferrocyanides (TREADWELL and CHERVET), A., ii, 786.

Cadmium cells. See Cells.

Cæsium, absorption spectrum of (LINDSAY), A., ii, 599.

induction spectrum of (DUNOYER), A., ii, 729.

photoelectric effect in the vapour of (WILLIAMS and KUNZ), A., ii, 809.

density of, at absolute zero (HERZ), A., ii, 289.

Cæsium silver gold bromides (SUSCHNIG), A., ii, 514.

chloride, use of, in microchemistry (DUCLOUX), A., ii, 77.

auric chlorides, complex (WELLS), A., ii, 449, 514.

Cæsium detection :—

detection of (MISSENDEN), A., ii, 658.

Caffeine, chloroform extraction apparatus for (SCHAAP), A., ii, 797.

in the blood and urine after parental administration (LOEN), A., i, 705.

excretion of, in urine (FRIEDBERG), A., i, 88.

in urine after tea and coffee drinking (OKUSHIMA), A., i, 703.

estimation of, with silicotungstic acid (AZADIAN), A., ii, 237.

Calcite, equilibrium of aragonite with, in aqueous solution (BACKSTROM), A., ii, 849.

Calcium, exploded-wire spectrum of (SAWYER and BROKER), A., ii, 803.
vapour pressure of (PILLING), A., ii, 291.

and its alloys, absorption of nitrogen by (RUFF and HARTMANN), A., ii, 377.

Calcium alloys with mercury, electrolytic preparation of (NEUHAUSEN), A., ii, 643.

Calcium salts, antagonism of potassium salts and (SPIRO), A., i, 85.

fixation of, by animal tissues (FREUDENBERG and GYÓRGY), A., i, 87, 291, 299, 608.

distribution of, in blood (JONES), A., i, 80; (MAZZOCCO), A., i, 788, 891; (KRAMER and TINDALL), A., i, 1087; (VAN PAASSEN), A., i, 1208.

in blood of children with tetany (ROBLES), A., i, 192.

in blood serum in pregnancy (MAZZOCCO and MORON), A., ii, 892.

metabolism of. See Metabolism.

precipitation of, in the presence of hydroxyl ions (MATTSO), A., i, 800.

Calcium carbide, preparation of (BOTOLFSSEN), A., i, 1107.

carbonate, solubility of, in water (OSAKA), A., ii, 830.

precipitation of, to resemble mother-of-pearl (CLÉMENT and RIVIERE), A., ii, 500.

estimation of alkali in, by the A.R. method (SINGLETON and WILLIAMS), A., ii, 527.

chlorite (LEVI), A., ii, 567.

haloids, ammonia compounds of (HUTTIG), A., ii, 849.

hydroxide, colloidal (v. GLASENAPP), A., ii, 848.

nitrate, crystal structure of (VEGARD), A., ii, 503.

mixed crystals of cadmium nitrate and (HASSELBLATT), A., ii, 61.

oxide, precipitation of, with ferric oxide (CHARRIOU), A., ii, 163.

phosphate, liquid crystals of (GAUBERT), A., ii, 445.

phosphates, equilibrium of formation of, and their solubility (DIECKMANN and HOUDREMONT), A., ii, 292.

silicides (WÖHLER and MÜLLER), A., ii, 293.

sulphate, equilibrium of sodium carbonate and (SYRKIN), A., ii, 699.

See also Gypsum.

tungstate (SMITH), A., ii, 774.

Calcium organic compounds :—

cyanide, pure, preparation of (PINCANS), A., i, 532.

Calcium estimation :—

estimation of (LAIDLAW and PAYNE), A., ii, 786.

estimation of, volumetrically (VÜRTHEIM and VAN BERS), A., ii, 869.

estimation of, in blood (WEISS; CLARK), A., ii, 227; (LING and BUSHILL), A., ii, 587.

estimation of, in natural phosphates (MEURIOE), A., ii, 653.

Calcium estimation :—

estimation of, in presence of magnesium and phosphates (SHOHL), A., ii, 395.

estimation of, in soils (SHEDD), A., ii, 527.

estimation of, in urine (SHOHL and PEDLEY), A., ii, 395.

Calcium ammonium (BOJOLFSEN), A., ii, 570.

Calorimeter, continuous flow (KEYES, GILLESPIE, and MITSUKURI), A., ii, 424.

Calorimetry, researches on (VERKADE, COOPS and HARTMAN), A., ii, 474, (VERKADE), A., ii, 740.

of slow processes (BARRY), A., ii, 473.

animal (TAINTNA ; CHANUIN), A., i, 288 ; (LUSK and EVENDEN), A., i, 292 ; (ATKINSON, RAPPOET, LUSK, SODERSTROM, and EVENDEN), A., i, 966.

bomb, correction for temperature in (EKGSTROM), A., ii, 27.

Calves, blood of. See Blood.

Camel, colostrum of. See Colostrum.

Camera inodorata, use of the (KOMURO), A., i, 84.

Campanula rotundifolia, constituents of (SPRINGER), A., i, 797.

Camphene, hydrochloride, preparation of (CHEMISCHE FABRIK AUF AKTIEN FORM. E. SCHERING), A., i, 943.

equilibrium of *n*- and *iso*-bornyl chlorides with (MEERWEIN, VAN EMSER and JOUSSEN), A., ii, 751.

Camphenecarboxylic acids, and their salts and derivatives (BREDT), A., i, 937.

Camphene glycol, chlorohydrin of (DEVEUF), A., i, 327.

Camphenilone group (KOMPPA and ROSCHIER), A., i, 1167.

Camphenillyxanthic acid, methyl ester (KOMPPA and ROSCHIER), A., i, 1187.

p-Camphocarboxylic acid (BREDT), A., i, 938.

trans- α -Campholanamide (KLEUCKER), A., i, 736.

dl-cis- and -trans- α -Campholanic acids, and their derivatives (KLEUCKER), A., i, 736.

Campholic acid, hydroxy- (PALFRAY), A., i, 548.

isoCampholic acid, and bromo- and their salts and derivatives (RUPE and BRIELLMANN), A., i, 1017.

dl-isoCampholic acid, derivatives of (KLEUCKER), A., i, 736.

α -Campholytic acid, intra-annular tautomerism of (CHANDRASENA, INGOLD, and THORPE), T., 1542.

Camphopropan- β -ol, and its phenylurethane (HALLER and RAMART-LUCAS), A., i, 480.

Camphor, and α -bromo-, rotation of, in various solvents (WETTERFORS), A., ii, 247.

equilibrium of ethyl alcohol, water and (SCHEKINGA), A., ii, 535.

equilibria of phenols with (KREMANN and ODELGA), A., i, 159.

analysis of preparations of (FRANÇOIS and LUCE), A., ii, 667.

d Camphor, catalytic reduction of (KOMATSU and MASUMOTO), A., i, 752.

Camphor, bromo-, estimation of (EATON), A., ii, 326.

α' -bromo- (LOWRY, STEELE, and BURGESS), T., 633.

Camphor oil, constants of (GATTEFOSSÉ), A., i, 1187.

Camphor- β -allylaminoethylimide, and its hydrobromide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camphor- β -aminoethylimide, and its hydrobromide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camphor- β -bromoethylimide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camphor- β -chloroethylimide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camphor- β -diethylaminoethylimide, and its hydrobromide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camphor- β -dimethylaminoethylimide, and its hydrobromide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

α -Camphordioxime, nickel compound (BIGIAYI), A., i, 353.

Camphoreins (SINGH, RAI, and LAL), T., 1421.

Camphorescin. See Resorcinolcamphoroid, tetrabromo-.

Camphor- β -hydroxyethylimide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camphoric acid, preparation of derivatives of (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 254 ; (FAUST), A., i, 748.

Camphoric anhydride, action of *p*-cresol with (KRISHNA), T., 255.

dl-Camphorimine nitrite (KLEUCKER), A., i, 736.

Camphor- β -piperidylethylimide, and its hydrobromide (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.

Camzellite, from British Columbia (ELLSWORTH and POITEVIN), A., ii, 304.

Cancer. See Carcinoma.

Candlenut oil. See Lumbang oil.

Caoutchouc (india-rubber), structure of (OLIVIER), A., i, 45; (KIRCHHOF), A., i, 945.

constitution and hydrogenation of (STAUDINGER and FRITSCHI), A., i, 1043.

size of molecules of (HARRIES and EVERS), A., i, 357.

problem of the synthesis of (PLOTNIKOW), A., i, 419.

action of concentrated sulphuric acid on (KIRCHHOF), A., i, 562.

vulcanisation of (BOIRY), A., i, 848.

use of mercaptothiazoles in (BRUNI and ROMANI), A., i, 755.

analysis of articles made of (DUBOSC), A., ii, 91.

vulcanised, estimation of sulphur in (DYER and WATSON), A., ii, 656, 782.

Caprifoliaceæ, constituents of the berry fruits of (NOWAK and ZELLNER), A., i, 615.

Capsaicin, constitution of (OTT and ZIMMERMANN), A., i, 137.

Capsella bursa pastoris, use of, as an ergot substitute (WASICKY), A., i, 907.

Capsularin, and its derivatives (SAHA and CHOUDHURY), T., 1044.

Carbamic acid, ammonium salt. equilibrium of formation and stability of (FAURHOLT), A., ii, 272, 747; (WEGSCHEIDER), A., ii, 376; (MATIGNON and FRÉJACQUES), A., ii, 445, 499.

Carbamic acid, dichloro-, methyl ester (DATTA and CHATTERJEE), A., i, 814.

Carbamic acids, formation of azides of (OLIVERI-MANDALÀ), A., i, 473.

Carbamic acids, thio-. See Thiocarbamic acids.

Carbamide (urea), preparation of (BADISCHE ANILIN- & SODA-FABRIK), A., i, 992.

from cyanamide (FARBWERKE VORM. MEISTER, LUCIUS & BRÜNING), A., i, 993.

synthesis of (BAILEY), A., i, 815.

from ammonia (MATIGNON and FRÉJACQUES), A., i, 723; ii, 272.

compounds of benzoic acid and OSAKA and ANDO), A., i, 1127.

estimation of, with sodium hypobromite (WERNER), T., 2318.

Carbamide (urea), estimation of, in blood (IONESCU), A., ii, 596.

See also Urea.

Carbamide, chloro- (DETGEUF), A., i, 236, 327.

Carbamides, constitution of (WERNER), T., 2318.

Carbamides, substituted, preparation of (ALBERT), A., i, 370; (TANBERG and WINKEL), A., i, 1009.

thio-. See Thiocarbamides.

3:3-Carbamidebis-6-methoxybenzoic acid (HERMANN), A., i, 1151.

o-Carbamidobenzenesulphonic acid, and its salts and acetyl derivative, potassium salt (SCOTT and COHEN), T., 2042.

3-Carbamido-1-o-carboxybenzylphthalazone (RUGGLI and MEYER), A., i, 345.

3-Carbamido-6-ethoxybenzoic acid, and its ethyl ester, and 3-thio- (HERMANN), A., i, 1151.

4-Carbamido-1-ethoxynaphthalene, and 4-thio- (HERMANN), A., i, 1151.

6-Carbamido-2-hydroxy-3-naphthoic acid (FROELICHER and COHEN), T., 1660.

3-Carbamido-4-methoxybenzoic acid (FROELICHER and COHEN), T., 1656.

3-Carbamido-6-methoxybenzoic acid, and its ethyl ester (HERMANN), A., i, 1151.

Carbamidomethoxynaphthoic acids (FROELICHER and COHEN), T., 1658.

p-Carbamidophenoxyacetone, and its derivatives (SPECKAN), A., i, 580.

p-Carbamidophenoxyacetophenone (SPECKAN), A., i, 580.

a-p-Carbamidophenoxybutyric acid, ethyl ether (SPECKAN), A., i, 580.

p-Carbamidophenyl ethyl carbonate and diphenylcarbamate (SPECKAN), A., i, 580.

p-Carbamidophenylglycidic ether (SPECKAN), A., i, 580.

1-Carbamidocyclopropane-1-carboxylic acid, and 1-nitroso- (INGOLD, SAKO, and THORPE), T., 1195.

p-Carbamido-m-sulphonic acid, and its salts, and acetyl derivative, potassium salt (SCOTT and COHEN), T., 2039.

Carbamine cyanides, preparation of (FROMM and WENZL), A., i, 487.

Carbamylazopropane (NEIGHBORS, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.

Carbanilide, thio-. See Thiocarbanilide.

Carbanilinoceinnamaldoximes (BRADY and THOMAS), T., 2104.

Carbanilinoceinnamantaldoximes (BRADY and THOMAS), T., 2108.

- Carbasinic acid**, *dithio*-, salts, decomposition of, and its thioanhydride (LOANITCH), T., 2542.
- Carbazole**, melting and boiling points of (KIRBY), A., i, 27.
- equilibria of, with nitrobenzenes and nitrotoluenes (KREMAN and STRZELDA), A., i, 176.
- Carbazole**, *N*-nitroso- (MORGAN and READ), T., 2712.
- Carbazole series**, diazo-reaction in the (MORGAN and READ), T., 2709.
- Carbazole-3-azoacetoacetic acid**, ethyl ester (MORGAN and READ), T., 2715.
- Carbazole-3-azoimide**. See 3-Triazo-carbazole.
- Carbazole-3-azo- β -naphthol** (MORGAN and READ), T., 2713.
- Carbazole-3-azo- β -naphthylamine** (MORGAN and READ), T., 2714.
- Carbazole-3-azoresorcinol** (MORGAN and READ), T., 2713.
- Carbazole- α -benzoic acid**, and its derivatives (ECKERT, SEIDEL, and ENDLER), A., i, 952.
- Carbazole-3-diazocyanide** (MORGAN and READ), T., 2714.
- Carbazole-3-diazoimine** (MORGAN and READ), T., 2712.
- Carbazole-3-diazonium salts** (MORGAN and READ), T., 2711.
- Carbethoxyaminoaceto-*p*-anisidide** (HILL and KELSEY), A., i, 1141.
- Carbethoxyaminoaceto-*p*-chloroanilide** (HILL and KELSEY), A., i, 1141.
- Carbethoxyaminoaceto-*m*-toluidide** (HILL and KELSEY), A., i, 1141.
- m*-Carbethoxyamino-*p*-hydroxyphenylarsinic acid** (BART), A., i, 1202.
- 3-Carbethoxyamino-4-methoxybenzoic acid** (FROELICHER and COHEN), T., 1658.
- Carbethoxyaminomethoxynaphthoic acids** (FROELICHER and COHEN), T., 1658.
- o*-Carbethoxyaminophenol**, *p*-amino- (BART), A., i, 1202.
- 3-Carbethoxyaminophenylphosphinic acid**, and nitro- (NIJK), A., i, 961.
- α -Carbethoxyaminoresacetophenone** (SONN and FALKENHEIM), A., i, 1164.
- o*-Carbethoxybenzylmalonic acid**, ethyl ester (DAVIES and PERKIN), T., 2210.
- Carbethoxydihydrosinapic acid**. See β -4-Ethylcarbonato-3:5-dimethoxyphenylpropionic acid.
- 3-Carbethoxy-2:4-dimethylpyrrole-5-acetic acid**, ethyl ester (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 3-Carbethoxydimethylpyrroleglyoxylic acids**, and their ethylesters (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 3-Carbethoxy-2:4-dimethylpyrrole-5-glyoxylonitrile** (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- 3-Carbethoxy-2:4-dimethylpyrroldiphenylcarbinol**, and its derivatives (FISCHER and KAN), A., i, 869.
- α -Carbethoxy- γ -hexolactone**, and α -bromo- (HELFERICH and SPEIDEL), A., i, 7.
- 4-Carbethoxy-5-methylpyrrolenyl-3-*p*-dimethylaminophenylmethane**, 3-hydroxy- (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- 4-Carbethoxy-5-methylpyrrolenyl-3-furylmethane**, 3-hydroxy- (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- 4-Carbethoxy-5-methylpyrrolenyl-3-*p*-methoxyphenylmethane**, 3-hydroxy- (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- o*-Carbethoxy- β -phenylpropionic acid**, α -cyano-, ethyl ester (DAVIES and PERKIN), T., 2210.
- Carbides of metalloids**, preparation of (DE MAHLER), A., i, 101.
- Carbinols**, and their corresponding dienes, physical constants of (V. AUWERS and WESTERMANN), A., ii, 97.
- halochromism of (SKRAUP and FREUNDLICH), A., i, 539.
- optically active, lubricating power of (DOUBLEDAY), T., 2875.
- Carboceric acid** (TROPSCH and KREUTZER), A., i, 804.
- Carboceric acid** (TROPSCH and KREUTZER), A., i, 817.
- Carbocyanines**, isomeric (BRAUNHOLTZ), T., 169.
- Carbodiimides**, condensation of hydrochlorides of, with phenols (SHORT and SMITH), T., 1803.
- Carbodilactic acids**, *trithio*-, stereoisomeric, and their salts (HOLMBERG), A., i, 1113.
- Carbohydrates** (BRIGL), A., i, 1117.
- in dead leaves (COMBES and KOHLER), A., i, 1222.
- acetone and butyl alcohol fermentation of (ROBINSON), A., i, 971.
- reduction of, with hydriodic acid and phosphorus (WILLSTÄTTER and KALB), A., i, 989.

- Carbohydrates**, assimilation of, in man (SAKAGUCHI, ASAKAWA, and MATSUYAMA), A., i, 699.
- transportation, retention and excretion of (FOLIN and BERGLUND), A., i, 487.
- metabolism of. See Metabolism.
- relations between fats and (MULLER), A., i, 486.
- degradation of, in muscle (LAQUER), A., i, 298, 1039.
- sulphates of (OHLE), A., i, 936.
- benzyl ethers of (GOMBERG and BUCHLER), A., i, 112.
- detection of (BAUDISCH and DEUEL), A., ii, 664.
- Carboligase** (NEUBERG and LIEBERMANN), A., i, 305; (NEUBERG and OHLE), A., i, 480, 540; (ROSENTHALER; NEUBERG and HIRSCH); A., i, 600; (HIRSCH), A., i, 973.
- Carbomethoxybenzoyl chlorides**, reactions of, with aromatic hydrocarbons (SMITH), A., i, 140.
- 4-Carbomethoxyphenylarsinic acid**, 3-nitro- (CHRISTIANSEN), A., i, 1203.
- Carbon**, atomic weight of (MOLES), A., ii, 51.
- hexavalency of (HANTZSCH), A., i, 24.
- pure, preparation of (WIBAUT), A., ii, 565.
- spectrum of (MULLER), A., ii, 725.
- Röntgen rays from (HUGHES), A., ii, 184; (HOLZMARK), A., ii, 543.
- atoms, linkings of, in organic compounds and in graphite (WIBAUT; BEUTLER), A., ii, 259; (V. FREIGER), A., ii, 618.
- kinetics of, open saturated chains of (WOJNICZ-SIANOZENCKI), A., i, 330.
- and the related elements in the periodic system (STOCK), A., ii, 564.
- co-ordination number of (HANTZSCH), A., i, 26.
- behaviour of, at high temperatures (SAUERWALD; RYSCHKWITSCH), A., ii, 443.
- diffusion of, in metals (TAMMANN and SCHÖNERT), A., ii, 772.
- solubility limits of, in ternary steels (DAEVES), A., ii, 70.
- colloidal, cataphoresis of (GOLDBERG), A., ii, 423.
- solutions of (THORNE), A., ii, 695.
- oxidation of (PHILIPPI; PHILIPPI and RIE; PHILIPPI and THELEN), A., i, 837.
- and its oxides, equilibrium of, with iron and its oxides (REINDERS and VAN GRONINGEN), A., ii, 153; (EASTMAN), A., ii, 506.
- Carbon**, action of nitrogen on barium carbonate and (ASKENASY and GRUDE), A., ii, 445.
- action of nitrogen on sodium carbonate and (INGOLD and WILSON), T., 2278.
- catalysis of the reaction of steam with (TAYLOR and NEVILLE), A., ii, 143.
- behaviour, of, when heated with sulphur (WIBAUT), A., ii, 373.
- Carbon alloys** with boron and iron (VOGEL and TAMMANN), A., ii, 852.
- with chromium and iron (DAEVES), A., ii, 70.
- with iron (HONDA), A., ii, 711.
- with iron and tungsten (DAEVES), A., ii, 70.
- Carbon compounds**, synthesis of, from air, by means of radium (GLEW), A., ii, 607.
- natural system of (DECKER), A., i, 417, 513.
- bivalent (GILLET), A., i, 213, 513.
- Carbon tetrachloride**, physical constants of (TIMMERMANS, VANDER HORST, and ONNES), A., ii, 258.
- thermal constants of (LATIMER), A., ii, 256.
- action of fuming sulphuric acid on (LEPIN), A., ii, 847.
- suboxide, preparation and properties of (OTT and SCHMIDT), A., ii, 641.
- monoxide, equilibrium of the formation of, from carbon dioxide and carbon (JELLINEK and DIETHELM), A., ii, 833.
- origin of the spectrum of (DUFREUX), A., ii, 597.
- electrical moment of (WEIGT), A., ii, 109.
- solubility of, in serum and plasma (O'BRIEN and PARKER), A., i, 394.
- viscosities of nitrogen and (SMITH), A., ii, 549.
- absorption of, by acid cuprous chloride (KROFF), A., ii, 657.
- action of, on diamond (FOIX), A., ii, 641.
- action of oxygen with, on platinum (LANGMUIR), A., ii, 629.
- catalytic combustion of, in hydrogen (LAMB, SCALIONE, and EDGAR), A., ii, 443.
- catalytic oxidation of (ROGERS, PIGGOT, BAHLEKE, and JENNINGS; MERRILL and SCALIONE), A., ii, 136.
- in tobacco smoke (ARMSTRONG and EVANS), A., i, 1226.

Carbon monoxide, estimation of, with an iodine pentoxide indicator (KATZ and BLOOMFIELD), A., ii, 585.
 estimation of, in blood (SAYERS and YANT), A., ii, 868.
dioxide, air-free, apparatus for preparation of (SCHOELLER), A., ii, 159.
 ultra-red absorption spectra of (BARKER), A., ii, 805.
 movement of electrons in (SKINKER), A., ii, 837.
 electrical moment of (WEIGT), A., ii, 109.
 moment of inertia and structure of molecules of (EUCKEN), A., ii, 349.
 specific heat of (PARTINGTON and CANT), A., ii, 191.
 compressibility of (GUYE and BATUECAS), A., ii, 618.
 solubility of, in various solvents (KUNERTH), A., ii, 823.
 aqueous solutions of (WILKE), A., ii, 52.
 viscosities of nitrogen monoxide and (SMITH), A., ii, 549.
 absorption of, by alkaline solutions (RIOU), A., ii, 433, 487.
 absorption of, by ammoniacal solutions (RIOU), A., ii, 742.
 rate of evolution of, from solution in presence of colloids (FINDLAY and HOWELL), T., 1046.
 reaction of benzene with, under the electric discharge (MIYAMOTO), A., i, 418.
 assimilation of, by plants (MOLISCH; MÜLLER and WARBURG), A., i, 411; (WARBURG and NEGELEIN), A., i, 1097.
 effect of anæsthetics on the output of (BODINE), A., i, 889.
 relation between oxygen and, in respiration (THUNBERG), A., i, 889.
 content of, in blood (PETERS, CULLEN, and AUSTIN), A., i, 1207.
 estimation of (SHAW), A., ii, 226.
 estimation of, in air (RAUCH), A., ii, 316; (NOYONS), A., ii, 868.
 apparatus for estimation of, in air (LUNDEGÅRDH), A., ii, 719.
 estimation of, in carbonates (SAYCE and CRAWFORD), A., ii, 316.
 oxides, reactions with, in blast furnaces and gas generators (v. JUPTNER), A., ii, 374.
Carbonic acid, velocity of the formation of (KAURHOLT), A., ii, 272.
 state of, in aqueous solution (THIEL), A., ii, 374.

Carbon :—

Carbonates, dissociation and hydrolysis of, in solution (MENZEL), A., ii, 345.
 oxychloride and oxysulphide. See Carbonyl chloride and sulphide.
disulphide, molecular structure of (RANKINE), A., ii, 704.
 physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.
 properties of (SCHULZE), A., ii, 53.
 equilibrium of ethyl alcohol, water, and (SCHOORL and REGENBOGEN), A., i, 419.
 equilibrium of, with ethyl and methyl alcohols (MCKELVY and SIMPSON), A., ii, 271.
 compound of azoimide and (OLIVERI-MANDALÀ), A., i, 1070.
 action of, on mercuric acetate (BERNARDI and ROSSI), A., i, 421.
 action of, with thalious β -diketones (MORGAN and LEDBURY), T., 2892.
 sulphides (WIBAUT), A., ii, 52.
Carbon estimation :—
 estimation of, in cast iron and steel (BATA and THYSSSEN), A., ii, 395.
 estimation of, in organic compounds (KLEIN and STREUBER), A., ii, 159; (WREDE), A., ii, 316; (SIMON and GUILLAUMIN), A., ii, 867; (SIMON), A., ii, 868.
 estimation of, volumetrically, in organic compounds (LINDNER), A., ii, 657.
Carbonatotetramminecobaltiselenate. See under Cobalt.
Carbonic acid. See under Carbon.
Carbonium colouring matters, constitution of (KAUFFMANN), A., i, 820.
Carbonium salts, constitution of (HANTZSCH), A., i, 24, 820; (KEHRMANN), A., i, 259, 831.
Carbonyl chloride, mechanism of the formation of (LEPIN), A., ii, 847.
 pressure of the saturated vapour of (NIKITIN), A., ii, 847.
 adsorption of, by beechwood charcoal (BUNBURY), T., 1525.
 action of, with benzene and *m*-xylene, in presence of aluminium chloride (WILSON and FULLER), A., i, 827.
 sulphide, molecular structure of (RANKINE), A., ii, 704.
 viscosity and molecular dimensions of (SMITH), A., ii, 686.
 pharmacology of (FISCHER), A., i, 402.

- Carbonyls, metallic** (MOND and WALLIS), T., 29.
 action of nitric oxide on (MOND and WALLIS), T., 32.
- Carbonyl compounds, metallic compounds of the enolic forms of** (SCHEINER and ZIEGLER), A., i, 426.
 additive reactions of, with phosphorus haloids (CONANT), A., i, 41.
- Carbonylens** (STAUDINGER) A., i, 239.
 action of, on phosphazines (STAUDINGER and BRAUNHOLTZ), A., i, 70.
- o*-Carboxybenzylideneanthranillo acid** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 2-*o*-Carboxybenzyl-2-methylcyclopropane-1-carboxylic acid, 8-hydroxy-, lactone, and its silver salt** (KON, STEVENSON, and THORPE), T., 662.
- α*-*o*-Carboxybenzyl-*α*-methylsuccinic acid** (KON, STEVENSON, and THORPE), T., 662.
- Carboxycamphenecamphoric acid**, (BREDT), A., i, 937.
- Carboxycinnamic acids, hydroxy-** (WAYNE and COHEN), T., 1028.
- 2-Carboxy-3:4-dimethoxyphenyl-3-carboxy-2-hydroxy-*α*-naphthylcarbinol, and its sodium salt and methyl ester** (STORIUS), A., i, 746.
- d*-3-Carboxy-1-1-dimethylcyclopropane-2-propionic acid** (SIMONSEN), T., 2297.
- α*-Carboxyglutaconolactone, *γ*-hydroxy-, diethyl ester** (FALTIS and DE ROXAS), A., i, 625.
- Carboxyhaemoglobin, dissociation of** (HILL), A., i, 193.
- cis*- and *trans*-2-Carboxycyclohexane-1-acetic acids** (WINDAUS and HÜCKEL), A., i, 658.
- trans*-2-Carboxycyclohexane-1-propionic acid, and its dianilide** (WINDAUS and HÜCKEL), A., i, 658.
- 1-Carboxycyclohexyl-1-acetic acid** (LAPWORTH and McRAE), T., 2754.
- 2-Carboxyindole-1:3-diacetic acid, and its diethyl ester** (KERMACK, PERKIN, and ROBINSON), T., 1893.
- Carboxylase of yeast, action of** (ABDERHALDEN and FODOR), A., i, 92.
- 4-Carboxy-6-methoxyquinoline-2-*β*-acrylic acid** (HALBERKANN), A., i, 175.
- β*-Carboxymethylbutan-*α**β*-dicarboxylic acid, and its derivatives** (CARRIÈRE), A., i, 319.
- 4-Carboxy-3-methylphenylacetic acid** (v. AUWERS and JÜLICHER), A., i, 842.
- β*-*o*-Carboxytrinitrophenylpropionic acid** (SCHROETER, KINDERMANN, DIRTRICH, BEYSLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 123.
- m*-Carboxyphenylarsinic acid** (BART), A., i, 1201.
- vic*-2 *p*-Carboxyphenylbenzotriazole-4:5-dicarboxylic acid, and its anhydride** (CHARRIER and CRIPPA), A., i, 771.
- m*-Carboxyphenyl-6-methyl-*ψ*-azimino-benzene, 5-amino-2-*p*-hydroxy-** (KALLE & Co.), A., i, 62.
- α*-Carboxy-*β*-phenyl-*α*-methylglutaconic acid, ethyl ester, ozonides of** (FEIST, BREUER, and LUBRICHT), A., i, 553.
- 4-Carboxy-2-phenylquinoline-6-arsinic acid, and its salts** (JOHNSON and ADAMS), A., i, 187.
- 2-Carboxyphenyldithioglycollic acid** (SMILES and McLELLAND), T., 88.
- Carcinoma, anæmia in** (ROBIN and BOURNIGAUT), A., i, 497.
 zinc content of tumours in (CRISTOL), A., i, 497.
- Carnosine, distribution of, in muscle of various animals** (CLIFFORD), A., i, 398.
 in muscle after cold storage (CLIFFORD), A., i, 791.
 estimation of, in muscle (HUNTER), A., ii, 328, 885.
- Carnotite, estimation of uranium in** (SCOTT), A., ii, 788.
- Carp.** See *Cyprinus carpio*.
- Carrageen, constituents of** (RUSSELL-WELLS), A., i, 1223.
 as a protective colloid (GUTBIER and HUBER; GUTBIER, WOLF, and KIESS), A., ii, 203.
- tert*-Carvomenthyl chloride** (MAXWELL), A., i, 753.
- d*-Carvotanacetone, and its derivatives** (SIMONSEN and RAU), T., 879.
- Caryophyllaceæ, glucosides from** (VERGELOT), A., i, 207.
- Caryophyllene, hydration of** (ASAHI and TSUKAMOTO), A., i, 845.
- Casein.** See Caseinogen.
- Caseinogen from cow's milk** (BLEYER and SEIDL), A., i, 479.
 ultramicroscopic investigation of (BLEYER and SEIDL), A., i, 886.
 effect of electrolytes on the solution and precipitation of (J. and R. F. LOEB), A., i, 182.
 hydrolysis of, with alkalis (GRIGGS), A., i, 182.
 hydrolysis of, by proteolytic enzymes (DUNN and LEWIS), A., i, 292.
 action of nitrous acid on (DUNN and LEWIS), A., i, 279.

- Caselnogen**, digestion of, with trypsin (FRANKEL and FELDSEBERG), A., i, 184; (FRANKEL and JELLINEK), A., i, 965.
- Cassiterite**, occurrence of germanium in (HADDING), A., ii, 855.
- Castela Nicholsoni*, castelin from (BOSMAN), T., 969.
- Castelagenin** (BOSMAN), T., 971.
- Castelamarin** (BOSMAN), T., 970.
- Castelin** (BOSMAN), T., 969.
- Catalase**, influence of the medium on the action of (SAMMARTINO), A., i, 391.
- effect of cobaltamines on the action of (FUNK), A., i, 481.
- in blood, effect of potassium cyanide on (WELKER and BOLLMAN), A., i, 192.
- of flour (MERL and DAIMER), A., i, 184.
- Catalase**, effect of metals on the action of, and its estimation (SANTESSON), A., i, 1077.
- estimation of, in blood (BACH and ZUBKOWA), A., i, 393.
- Catalysis** (GILFILLAN), A., i, 709; (PURKAYOSTHA and DHAR), A., ii, 362; (PERRIN; LEWIS; BAILY; LOWRY), A., ii, 628; (LANGMUIR), A., ii, 629; (DHAR), A., ii, 730; (BOESEKEN), A., ii, 755; (BANERJI and DHAR), A., ii, 756.
- studies in (TAYLOR and LEWIS), T., 665; (MORAN and LEWIS), T., 1613.
- ionic activity in (HARNED and SELTZ), A., ii, 631.
- by ions, effect of a magnetic field on, in presence of a paramagnetic salt (GARNER and JACKMAN), T., 1298.
- effect of adsorption on (GUICHARD), A., ii, 630.
- rôle of protective colloids in (IREDALE), T., 1536.
- in fuller's earth (RIDEAL and THOMAS), T., 2119.
- in homogeneous gas reactions (HERZFELD), A., ii, 136.
- hydrogen-oxygen, by means of platinum (HOFMANN), A., ii, 490.
- negative (DHAR and MITTRA), A., ii, 630.
- by platinum black (VAVON and HUSON), A., ii, 631.
- at solid surfaces (ARMSTRONG and HILDITCH), A., ii, 41, 756, 757.
- Catalysts**, influencing of the activity of (ROSENEMUND and ZETZSCHE), A., ii, 41, 834; (ABEL), A., ii, 276.
- influence of, on the thermodynamics of velocity of reaction (VAN TUIJL), A., ii, 754.
- Catalysts**, influence of, on equilibrium (CLARENS), A., ii, 436; (DURAND), A., ii, 701.
- behaviour of metals as (SANDONNINI), A., ii, 557.
- Catalytic analysis**. See **Analysis**.
- hydrogenation (SKITA), A., ii, 207.
- in presence of nickel salts (v. BRAUN and KIRSCHBAUM), A., i, 727.
- with reduced nickel (RIDEAL), T., 309.
- of liquids in presence of nickel (ARMSTRONG and HILDITCH), A., ii, 41.
- with exclusion of oxygen (NORMANN), A., ii, 631.
- of phenols (SENDERENS and ABOULENO), A., i, 136.
- oxidation (KARCZAG), A., ii, 42, 137.
- reactions, course of (ZALKIND), A., ii, 835.
- Catechin** (v. EULER), A., i, 45; (FREUDENBERG), A., i, 756.
- constitution of (NIERENSTEIN), T., 604.
- Catechins**, optical activity of (FEIST and FUTTERMENGER), A., i, 565.
- stereoisomeric, and their derivatives (FREUDENBERG, BOHME, and PURMANN), A., i, 756.
- Catechol** (*pyrocatechol*), compound of diketopiperazine and (POVARNIN and TICHOMIROV), A., i, 1185.
- Catecholcamphorein** (SINGH, RAI, and LAL), T., 1427.
- Catecholsulphamphthalein** (DUTT), T., 2392.
- Catechu** (v. EULER), A., i, 45.
- Catechutannins** (NIERENSTEIN), T., 23.
- Cathetus fusciculata*, oil from (GATTEFOSSE), A., i, 1167.
- Cathodes**, deposits on, from electrolysis of mixed salts (CREUTZFELDT), A., ii, 347.
- mercury, overvoltage of (NEWBERRY), T., 7.
- Cell or Cells**, electrochemical, intensification of the current in (REICHINSTEIN and KLEMENT), A., ii, 112; (REICHINSTEIN and ROHSCHILD), A., ii, 736.
- electromotive force of (WASASTJERNA), A., ii, 815.
- temperature coefficients of electromotive force of (GERKE), A., ii, 682.
- cadmium low-voltage (OBLATA), A., ii, 343.
- carbon monoxide-oxygen (KALLMANN), A., ii, 467.

- Cell or Cells, electrochemical**, hydrogen-calomel, effect of hydrogen pressure on electromotive force of (HAINSWORTH and MACINNES), A., ii, 467.
- Cell or Cells, photo-electric**, measurement of light absorption by (v. HALBAN and SIEDENTOPF), A., ii, 332.
- alkali metal, fatigue of (ROSENBERG), A., ii, 189.
- selenium (SPATH), A., ii, 189.
- Cell or Cells, physiological**, permeability of (SCHAEFFI), A., i, 290.
- permeability of, to acids (CROZIER), A., i, 897, 1090.
- influence of optical activity on the permeability of (KOTAKE and OKAGAWA), A., i, 696.
- living, surface reactions in (WARBURG), A., i, 787.
- respiration of (WARBURG), A., i, 190.
- pharmacology of the respiration of (ELLINGER), A., i, 402.
- role of inorganic salts in (ZONDEK), A., i, 399.
- sensitiveness to poisons (HANDOVSKY), A., i, 697.
- animal, absorption of potassium by (MITCHELL, WILSON, and STANTON), A., i, 196.
- function of vitamins in (HESS), A., i, 399, 788; (ABDERHALDEN), A., i, 607.
- detection of potassium in (MACALLUM), A., ii, 869.
- estimation of the reducing power of (LIPSCHITZ and GOTTSCHALK), A., i, 298.
- estimation of substances which injure (SCHNABEL), A., i, 304.
- Cellobiose** in yeast (GROENEWEGE), A., i, 903.
- Celloisobiose** (OST and KNOTH), A., i, 526.
- Cellosan**, constitution of (KARRER and SMIRNOV), A., i, 435.
- Cellulose** (HESS), A., i, 12; (HESS and WITTELSBACH), A., i, 116.
- preparation and properties of solutions of (WAENTIG), A., i, 988.
- fluorescence of (LEWIS), A., ii, 334.
- constitution and nitration of (DE BRUIN), A., i, 13.
- structure of, and its compound with alkali (KARRER), A., i, 231.
- action of heat on (JUSTIN-MUELLER), A., i, 11.
- solubility of, in alkaline copper hydroxide and copper oxide-ammine solutions (TRAUBE), A., i, 115.
- viscosity of (JOYNER), T., 1511, 2395.
- Cellulose**, adsorption of alumina by (TINGLE), A., i, 434.
- degradation products of (KARRER), A., i, 231.
- complex copper compounds of (HESS, MESSMER, and JAGLA), A., i, 988.
- action of formaldehyde on (SAMEC and FERJANČIČ), A., i, 115.
- action of iodine on (HUEBNER and SINHA), A., i, 434.
- oxidation of (KNECHT and THOMPSON), A., i, 637.
- reactions of, with sodium chloride and neutral salt solutions (MASTERS), T., 2026.
- saccharification of (WOHLAND KRULL), A., i, 232; (BUDNIKOV and ZOLO-TAREV), A., i, 922.
- effect of water and organic salts on (HUEBNER and KAYE), A., i, 435.
- fermentation of, in the paunch of the ox (KLEIN), A., i, 82.
- alkali-soluble (KNOEVENAGEL and BUSCH), A., i, 636.
- benzyl derivatives of (GOMBERG and BUCHLER), A., i, 112.
- cotton, properties of (COLLINS), A., i, 1120.
- glucose from (IRVINE and HIRST), T., 1585.
- estimation of, in wood (HEUSER and CASSEUS), A., ii, 664.
- Cellulose esters**, preparation of (GRÜN and WITTKA), A., i, 114.
- acetate, preparation of, from wood cellulose (HAGGLUND, LÖFMANN, and FARBER), A., i, 323.
- solubility of, in alkali and alkaline-earth salts (SCHWEIGER), A., i, 324.
- viscosity of solutions of (BARR and BIRCUMSHAW), A., i, 232.
- nitrate (ANGELI), A., i, 922.
- as an emulsifying agent (HOLMES and CAMERON), A., ii, 268.
- α -, β -, and γ -Celluloses (OPFERMANN), A., ii, 665.
- Celtium**, *L*-series spectrum of, and its atomic number (DAUVILLIER), A., ii, 463.
- atomic number of (URBAIN), A., ii, 505.
- Cement**, equilibrium between lime-water and (LORENZ and HÄGERMANN), A., ii, 59.
- Cenchrus tribuloides* (sand spur), constituents of the fruit of (YOUNG KEN and LA WALL), A., i, 1100.
- Cephalin**, brain, composition and properties of (FENGER), A., i, 315.

- Cereals**, effect of application of nitrogen on the protein content of (GERICKE), A., i, 1226.
- Cerebronic acid**, constitution of (LEVENE and TAYLOR), A., i, 714.
- dl-Cerebronic acid** (LEVENE and TAYLOR), A., i, 715.
- Cerebrospinal fluid**, chemistry of (EGGERER-SEHAM and NIXON), A., i, 395.
- acetone in (KOOPMAN), A., i, 194.
- combined carbon dioxide in (TOKUOKA and OGASAWARA), A., i, 895.
- cholesterol in (FABRIS), A., i, 969.
- sugar in (COOPER; DE HAAN and VAN CREVELD; STEVENSON), A., i, 295.
- testing of, with colloidal gold solutions (GETTLER and JACKSON), A., ii, 240.
- estimation of albumin in (AYER and FOSTER), A., ii, 798.
- Cerium**, arc spectra of (KIESS, HOPKINS, and KREMER), A., ii, 244.
- L-series spectrum of (DAUVILLIER), A., ii, 101.
- Cerium alloys** (GUILLET), A., ii, 572.
- Cerium compounds**, pure, preparation of (MANSKE), A., ii, 646.
- Cerium dioxide**, action of light on (RENTZ), A., ii, 61.
- Cerous salts**, electrolysis of aqueous solutions of (SCHLOTZ), A., ii, 22.
- Cerium separation**:—
separation of, from the other earths (PRANDTL and LÖSCH), A., ii, 770.
- Ceruleofibrite** (HOLDEN), A., ii, 516.
- Cetene**, compounds of, with zinc and nitrogen tetroxide (SCHAARSCHMIDT, VEIDT, and SCHLOSSER), A., i, 646.
- ψ -nitrosite (SCHAARSCHMIDT, VEIDT, and SCHLOSSER), A., i, 646.
- Cetyl**. See Hexadecyl.
- Cetylsulphonic acid**. See Hexadecanesulphonic acid.
- Cevadine**, and its o-nitrobenzoyl derivative (MACBETH and ROBINSON), T., 1571.
- Cevine**, preparation of, and its di-o-nitrobenzoyl derivative (MACBETH and ROBINSON), T., 1574.
- Chetopterus variopedatus*, manganese in the tube of (BERKELEY), A., i, 493.
- Chalcedony**, comparison of, with other forms of silica (WASHBURN and NAVIAS), A., ii, 804.
- Charcoal**, adsorption by (HERBST), A., ii, 30; (FODOR and SCHÖNFELD), A., ii, 688.
- adsorption of gases by (MCLEAN), A., ii, 352; (GUSTAFER), A., ii, 479.
- adsorption of saturated vapours by (DRIVER and FIRTH), T., 2409.
- Charcoal**, pressure and volume of liquids adsorbed by (HARKINS and EWING), A., ii, 123.
- oxidising properties of (FEIGL), A., ii, 51.
- effect of heating sulphur with (WIBAUT), A., ii, 52.
- beechwood, adsorption of carbonyl chloride by (BUNBURY), T., 1525.
- sugar, adsorption by (BARTELL and MILLER), A., ii, 741.
- is Chavieic acid* (OTT, EICHLER, LÜDEMANN, and HEIMANN), A., i, 1027.
- Chavicine** from pepper-resin (OTT, EICHLER, LÜDEMANN, and HEIMANN), A., i, 1026.
- Chemical compounds**, classification of (LANGMUIR), A., ii, 138.
- graphic representation of the composition of (HUBSCHER), A., ii, 156.
- space formulae for (ZWIKKER), A., i, 567.
- constitution and antiseptic action (BROWNING, COHRN, GAUNT, and GULBRANSEN), A., i, 612.
- and colour (MOIR; KEHRMANN), A., ii, 333.
- in relation to lubricating power (DOUBLEDAY), T., 2875.
- and odour (ZWAARDEMAKER), A., i, 607.
- relation between pharmacological action and (V. BRAUN, BRAUNSDORF, and RATH), A., i, 759.
- and formation of resins (HERZOG and KREIDL), A., i, 1168.
- and rotatory power (RUPE and JAGGI), A., i, 840; (RUPE), A., ii, 602.
- relation between peppery taste and (OTT and ZIMMERMANN), A., i, 137.
- reactions and radiation (TAYLOR; LEWIS), A., ii, 336.
- influence of substitution on (FRANZEN and STAUBLE), A., i, 450; (BODFORSS), A., ii, 698.
- occurrence of free radicals in (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- valency in relation to the mechanism of (PERKINS), A., ii, 438.
- induced (MITTHA and DHAR), A., i, 1210; (DHAR and MITTHA), A., ii, 630.
- theory of (WINTHER), A., ii, 336.
- simultaneous, course of (WEGSCHEIDER), A., ii, 489.
- theory, rôle of the physicist in development of (WALKER), T., 735.

- Chemistry**, Arabic (HOLMYARD), A., ii, 536.
- Chemoluminescence** (GRINBERG), A., ii, 806.
- Cherry**. See *Prunus avium*.
- Cherry laurel**, constituents of distillate from (PECKER), A., ii, 537.
- Chetoptera*, chetopterin from, and its spectrum (ROMIEU and OBATON), A., i, 793.
- Chickens**, enzymes in the alimentary canal of (PLIMMER and ROSEDALE), A., i, 485.
- Children**, underweight, basal metabolism in (BLUNT, NELSON, and OLESON), A., i, 83.
- China wood oil**, constituents of (BAUER and HERBERTS), A., i, 806.
- Chloranil**, compound of acenaphthene with (PFEIFFER and FLATER), A., i, 342.
- Chlorates**. See under Chlorine.
- Chlorella vulgaris*, assimilation of carbon by (MÜLLER and WARBURG), A., i, 411.
- Chloric acid**. See under Chlorine.
- Chlorides**. See under Chlorine.
- Chlorination**, by a mixture of carbon monoxide and chlorine (VENABLE and JACKSON), A., ii, 73.
by means of nitrogen trichloride (COLEMAN and NOYES), A., i, 133.
with sulphuryl chloride (SILBERRAD), T., 1015.
- Chlorine** from apatite, atomic weight of (GLEDITSCH and SAMDAHL), A., ii, 281.
Röntgen ray absorption spectrum of (LINDH), A., ii, 177.
isotopes of (BRONSTED, HARKINS; HARTLEY, PONDER, BOWEN, and MERTON), A., ii, 280; (LUDLAM), A., ii, 497; (ISHINO), A., ii, 760.
separation of isotopes of (HARKINS and HAYES), A., ii, 140.
distribution of electrons in the atom of (BRAGG, JAMES, and BOSANQUET), A., ii, 703.
decomposition of, into atoms (TRAUTZ and STÄCKEL), A., ii, 760.
dissociation of (v. WARTENBERG and HENGLEIN), A., ii, 441.
thermal decomposition of (HENGLEIN), A., ii, 823.
solid vapour pressure of (HENGLEIN, v. ROSENBERG, and MUCHLINSKI), A., ii, 760.
photochemical reactions between hydrogen and (GOHRING), A., ii, 9; (WEIGERT and KELLERMANN), A., ii, 808.
- Chlorine**, distribution of, between water and air (TITOV), A., ii, 269.
distribution and estimation of, in plants (JUNG), A., i, 1098.
distribution of, in blood (SIEBECK and HACKMACK), A., i, 1208.
metabolism. See Metabolism.
- Chlorine dioxide**, structure of (PERKINS), A., ii, 441.
action of, on organic compounds (SCHMIDT and BRAUNSDORF), A., i, 781.
- Hydrochloric acid**, synthesis of, from its elements without explosion (NEUMANN, BERGDAHL, BROY, and KARWAT), A., ii, 44.
absorption spectrum of (COLBY and MEYER), A., ii, 5; (RANDALL, COLBY, and PATON), A., ii, 805.
electrical conductivity of, in presence of sucrose (KIFRAN), A., ii, 812.
heats of neutralisation of (KEYES, GILLESPIE, and MITSUKURI), A., ii, 424; (RICHARDS and ROWE), A., ii, 425.
formation of complex ions in solutions of (HIXON), A., ii, 815.
- Chlorides**, contact potentials of solutions of (MACINNES and YEH), A., ii, 252.
inorganic, dissociation of, in aqueous solution (GÜNTHER-SCHULZE), A., ii, 765.
effect of sucrose on the ionic activity of (CORRAN and LEWIS), A., ii, 691.
distribution of, in blood (MUKAI; VAN CREVELD; NORGAARD and GRAM), A., i, 287; (MUREBANU), A., i, 290; (DISCHE), A., i, 1086.
metabolism of. See Metabolism.
estimation of, by Bang's micro-method (PRIGGE), A., ii, 715; (PETSCHACHER), A., ii, 716.
estimation of, with silver nitrate (KOLTHOFF), A., ii, 581.
estimation of, in blood (FRIEND), A., ii, 389; (ISAACS), A., ii, 716.
estimation of, in presence of iodides (BONINO), A., ii, 78.
estimation of bromides and, in presence of iodides (KOLTHOFF), A., ii, 79.
- Chloric acid**, estimation of (HAKONORI), A., ii, 389; (KIKUCHI), A., ii, 519.
- Chlorates**, reflection spectra of (SCHAEFER and SCHUBERT), A., ii, 180.
effect of chromates on the electrolysis of (PAMFILOV), A., ii, 712.

Chlorine :—

Chlorates, estimation of, by reduction (TREADWELL, HIRSTIE, EGGER, and STURZENEGGER), A., ii, 780.

Perchloric acid, and its salts, velocity of reduction of (BREDIG and MICHEL), A., ii, 359.

estimation of (KÖNIG), A., ii, 310 ; (LENNER and TOSTERUD), A., ii, 395.

Hypochlorous acid, ionisation constant of (NOYES and WILSON), A., ii, 692.

estimation of, in concentrated salt solutions (TAYLOR and GAMMAL), A., ii, 581.

Chlorites (LEVI), A., ii, 567.

Chlorine estimation :—

estimation of, in benzaldehyde (SCHIMMEL & Co.), A., ii, 78.

estimation of, in organic compounds (KLIMONT), A., ii, 580 ; (VOTOCEK), A., ii, 863.

estimation of, in organic matter (v. BOGDÁNDY), A., ii, 519.

estimation of, in concentrated salt solutions (TAYLOR and GAMMAL), A., ii, 581.

estimation of, in gastric juice (MALGOYRE), A., ii, 780.

estimation of, in tissues (PICO and MURTAGH), A., ii, 716.

Chlorites, white (SHANNON and WHEBRY), A., ii, 517.

See also under Chlorine.

Chloro-acids, preparation of (IKEDA and KODAMA), A., i, 219.

Chloroamines, preparation of (COLEMAN and NOYES), A., i, 183.

aromatic (GOLDSCHMIDT and STROHMENGER), A., i, 1004.

Chloroform, physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.

equilibrium of ethyl alcohol, water and (SCHOORL and REGENBOGEN), A., i, 215.

action of, on magnesium alkyl haloids (BINAGHI), A., i, 1002.

derivatives as narcotics (JOACHIMOGLU), A., i, 199.

in blood after death (GIBSON and LAIDLAW), A., i, 1218.

effect of, on peptic digestion (ASTRUC and RENAUD), A., i, 281.

Chlorohydrins, preparation of (DETIGUÉ), A., i, 286, 327.

sensory stimulation by (IRWIN), A., i, 899.

β -Chlorohydrins, preparation of (SMITH), A., i, 3.

Chloropentamminocobaltic 2:4-dinitro- α -naphthoxide, and its 7-sulphonic acid (MORGAN and KING), T., 1727.

Chloropentamminocobaltic dipicrylamine (MORGAN and KING), T., 1726.

Chlorophyll, chemistry of (COSTANTIN), A., i, 162.

photoelectric properties of (DIXON and BALL), A., ii, 248.

effect of sucrose on the formation of, in etiolated cotyledons (MANSKY), A., i, 1222.

Chloropicroin, action of, with mercaptans (RAY and DAB), T., 323.

Chloroplatinic acid. See under Platinum.

Chlorosulphonic acid, mechanism of the formation of (LEPIN), A., ii, 847.

analysis of (WEISSENBERGER and ZODER), A., ii, 390.

Cholesterol (WINDAUS), A., i, 541.

action of benzoyl peroxide on (WINDAUS and LUDERS), A., i, 453.

iodine value of (HOLDE, WERNER, TACKE, and WILKE), A., ii, 728.

origin and destiny of, in the animal organism (GARDNER and FOX), A., i, 89, 790.

distribution of, in blood (ROUZAUD and THIÉRY), A., i, 394.

in blood serum (SRIKAUS and SCHUBART), A., i, 1209.

derivatives of, in blood (LIFSCHUTZ), A., i, 392.

*di*bromide (LIFSCHÜTZ), A., i, 251.

detection of, colorimetrically (KAHLENBERG), A., ii, 591.

α -**Cholesterol oxide**, chlorohydrin of (WINDAUS and LUDERS), A., i, 541.

Cholic acid, oxidation of (WIELAND and SCHLICHTING), A., i, 554.

preparations of derivatives of (RIEDEL), A., i, 553 ; (FARBENFABRIKEN VORM. F. BAYER & Co), A., i, 554.

*apo***Cholic acid**, additive compounds of (RIEDEL), A., i, 554.

Choline, production of peristalsis by (ARAI), A., i, 970.

effect of, on movements of the intestines and stomach (LE HEUX), A., i, 85.

Cholines, preparation of, from amino-acids (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 813.

proteinogenous, decomposition of (KARRER and HORLACHER), A., i, 825.

*iso***Chondodendrine** (FALTIS and NEUMANN), A., i, 569.

Chondrus crispus. See Carrageen.

Chromioxalic acid, potassium salt, auto-racemisation of (RIDEAL and THOMAS), T., 196.

Chromiselenates. See under Chromium.
Chromithiocyanates (BJERRUM), A., i, 18.

Chromium, preparation of (FUJIBAYASHI), A., ii, 647.

Röntgen-ray spectrum of (DUANE and FRICKE), A., ii, 804.

cathodic deposition of metals on (KYROPOULOS), A., ii, 22.

expansion of, and of its alloys with nickel (CHEVENARD), A., ii, 153.

Chromium alloys with carbon and iron (DAEVES), A., ii, 70.

with nickel, estimation of chromium in (LOFFELBEIN; HILD), A., ii, 660.

Chromium bases (*chromiumammynes*), solubilities of salts of (BRONSTED and PETERSEN), A., ii, 199.

complex chloroiridates of (BENRATH, BÜCHER, and ECKSTEIN), A., ii, 515.

Chromium azide, and its complex derivatives (OLIVERI-MANDALÀ and COMELLA), A., ii, 382.

sesquioxide (*chromic oxide*) as a mordant (BANCROFT), A., ii, 822.

oxidation of organic compounds by (SIMON), A., ii, 868.

trioxide, equilibrium of sulphur trioxide, water and (GILBERT, BUCKLEY, and MASSON), T., 1934.

oxides, hydrated (WEISER), A., ii, 575.

Chromic acid, constitution of (DHAR), A., ii, 382.

Chromates, reflection spectra of (SCHARFER and SCHUBERT), A., ii, 179.

magnetic susceptibility of solutions of, and sulphuric acid (CABREIRA and PIÑA DE RUBIES), A., ii, 612.

electrolysis of, with diaphragm (LOTTERMOSER and FALK), A., ii, 736.

effect of, on the electrolysis of chlorates (PAMFILOV), A., ii, 712.

estimation of, iodometrically (VOSBURGH), A., ii, 863.

Dichromates, estimation of, acidimetrically (KOLTHOFF and VOGELLENZANG), A., ii, 88.

estimation of, electro-volumetrically (ETTFLEY and VOSBURGH), A., ii, 876.

Chromium sulphate, green (RECOURA), A., ii, 508.

Chromic chloride, electrolytic reduction of (TAYLOR, GERSDORFF, and TOVREA), A., ii, 382.

Chromium:—

Hexamminechromiselenate (MEYER and SPEICH), A., ii, 71.

Pentamminechromiselenate, chloro- (MEYER and SPEICH), A., ii, 71.

Pentaquochromiselenate, chloro- (MEYER and SPEICH), A., ii, 71.

Tetraquochromihexaquochromiselenate, dichloro- (MEYER and SPEICH), A., ii, 71.

Chromium organic compounds:—

penta-, tetra-, and tri-phenyl salts (HEIN), A., i, 77.

tetraphenylchromitetrathiocyanatodiammine (HEIN), A., i, 76.

triphenylchromitetrathiocyanatodiammine (HEIN), A., i, 78.

Chromic salts with azosalicylic acids (MORGAN and SMITH), T., 2866.

Chromiselenates of carbamide and triethylenediamine (MEYER and SPEICH), A., ii, 71.

Chromium estimation:—

estimation of, in metals (LOFFELBEIN; HILD), A., ii, 660.

estimation of, in ferrochromium (KELLEY and WILEY), A., ii, 164.

estimation of, in ferrovanadium (KELLEY, WILEY, BOHN, and WRIGHT), A., ii, 89.

estimation of, in steel (SIMION), A., ii, 529; (LOSANA and CAROZZI), A., ii, 589.

Chromo-isomerism, theory of (LUCAS and KEMP), A., i, 30.

Chrysazin boroacetate, and its acetyl derivative (DIMROTH and FAUST), A., i, 157.

Chrysoidine, action of mercuric acetate on (VECCHIOTTI), A., i, 478.

Chrysophanic acid. See 3-Methyl-anthraquinone, 1:8-dihydroxy-.

Chymosin. See Rennin.

Ciloidanic acid (WIELAND and SCHLICHTING), A., i, 838.

Cinchona alkaloids (RABE), A., i, 360; (RABE, KINDLER, and WAGNER), A., i, 361; (ACTON), A., i, 610.

stereochemistry of (KING and PALMER), T., 2577.

action of hydrogen peroxide on (SPEYER and BECKER), A., i, 674.

pharmacology of (ACTON), A., i, 899.

distribution of, in the organism (SCHNABEL), A., i, 290.

syntheses of derivatives of (JACOBS and HEIDELBERGER), A., i, 671, 672, 673.

hydrogenated, preparation of amino-derivatives of (HOWARD & SONS, BLADEN, and NIERENSTEIN), A., i, 853.

- Cinchona** alkaloids, hydrogenated, aminosulphonic acids from (BOEHRINGER & SÖHNE), A., i, 46.
 estimation of, volumetrically (SCHOORL), A., ii, 538.
- Cinchonidine** hexosephosphate (NEUBERG and DALMER), A., i, 920.
 salt of tropic acid with (KING and PALMER), T., 2585.
- Cinchonine** ferrioxalate (BURROWS and TURNER), A., i, 916.
- Cinchotinine**, ethyl and methyl ethers (HEIDELBERGER and JACOBS), A., i, 673.
- Cinchotoxine**, salt of tropic acid with (KING and PALMER), T., 2584.
- Cinnamaldehyde**, condensation of *p*-nitrobenzyl chloride with (KLEUCKER), A., i, 734.
 and nitro, oximes of (BRADY and THOMAS), T., 2098.
 picrate, compound of thiocarbamide and (TAYLOR), T., 2270.
- Cinnamaldehyde**, *m*-nitro-, diethylhydrazone (BRADY and McHUGH), T., 1651.
- Cinnamaldoxime** *O*- and *N*-benzyl ethers (BRADY and THOMAS), T., 2106.
- Cinnamantaldoxime** 2:4-dinitrophenyl ether (BRADY and THOMAS), T., 2106.
- Cinnamantaldoxime**, *o*-, *m*- and *p*-nitro- (BRADY and THOMAS), T., 2106.
- Cinnamic acid**, use of, in cryoscopy (FALCIOLA), A., ii, 421.
 sulphonation of (MOORE and THOMAS), A., i, 454.
 condensation of resorcinol with (SHORT and SMITH), T., 1808.
 chlorohydrin, preparation and reactions of (FORSTER and SAVILLE), T., 2595.
 4-chlorosulphonyl derivative, preparation of (STEWART), T., 2561.
 cholesteryl ester, inversion of rotation of (ROYER), A., ii, 415.
 vanillylamide of (OTT and ZIMMERMANN), A., i, 137.
- trans-Cinnamic acid**, *p*-amino-, α -chloro-*p*-amino-, α -chloro-*p*-nitro-, and *p*-nitro-, and their salts and derivatives (PFEIFFER and HAEFELIN), A., i, 738.
- Cinnamic acids**, action of light on (DE JONG), A., i, 339.
 photo-reactions of (STOBBE and STEINBERGER), A., i, 1018.
- Cinnamic acids**, *trans*-substituted, addition of bromine to (VAN DUIN), A., i, 737.
 nitro-, estimation of nitrogen in, by the Kjeldahl method (MARGOSCHES and VOGEL), A., ii, 522.
- Cinnamon bark oil** (ROURE-BERTRAND FILS), A., i, 847.
- Cinnamonitrile**, *m*- and *p*-nitro- (BRADY and THOMAS), T., 2107.
- Cinnamoyldiazoacetic acid**, methyl ester, interaction of triphenylphosphine with (STAUDINGER and LUSCHER), A., i, 238.
- Cinnamoylformic acids**, nitro- (HELLER, LAUTH, and BUCHWALDT), A., i, 1024.
- Cinnamylidenemethyl α -hydroxyisopropyl ketone**. See α -Phenyl- ζ -methyl- $\Delta^{\alpha\gamma}$ -heptadien- ζ -ol- ϵ -one.
- Cinnamylidenequinaldine** ethohaloids (MILLS and HAMER), T., 2012.
- Citraconic acid**, *p*-chlorophenylhydrazine ester and phenyl- and tolyl-hydrazides (CHATTAWAY and PARKES), T., 285.
- Citraconyl-mono- and -di-bromophenylhydrazides** (CHATTAWAY and PARKES), T., 286.
- Citraconyl-mono- and -di-chlorophenylhydrazides** (CHATTAWAY and PARKES), T., 286.
- Citraconylhydrazides**, isomeric (CHATTAWAY and PARKES), T., 283.
- Citraconyltolylhydrazides** (CHATTAWAY and PARKES), T., 288.
- Citric acid**, dissociation constants of (HASTINGS and VAN SLYKE), A., i, 985.
 formation of, in *Citromyces* cultures, and its estimation (BUTKEWITSCH), A., i, 973.
 in milk and milk products (SUPPLEE and BELLIS), A., i, 197.
 of cows' milk, decomposition of, by bacteria (KICKINGER), A., i, 1219.
 ammonium salt, preparation of a neutral solution of (ROBINSON), A., ii, 84.
 analysis of (ROBINSON and BANDEMAR), A., ii, 460.
 cobalt salt, tetrammine (CLARK and BUCKNER), A., ii, 300.
 zirconyl salt (VENABLE and LINEBERRY), A., i, 917.
 estimation of, in urine (McCLURE), A., ii, 791.
- Citromyces**, formation of oxalic acid and ammonia in growth of, on peptone (BUTKEWITSCH), A., i, 707.
 cultivation of, on sugar, and the formation and estimation of citric and oxalic acids (BUTKEWITSCH), A., i, 973.
- Citronella oil**, estimation of geraniol content of (DE JONG and RECLAIRE), A., ii, 790.
- Clays**, formation of layers in (UNGERER), A., ii, 96.
 amphoteric action of (ARRHENIUS), A., i, 707.

Clupanodonis acid (TSUJIMOTO), A., i, 917.
Clupea alosa (shad), chemistry of the spermatozoa of (STEUDEL), A., i, 294.
Cnidic acid (MURAYAMA), A., i, 310.
Cnidiolactone (MURAYAMA), A., i, 310.
Coagulation, kinetics of the process of (MUKHERJEE and PAPACONSTANTINOU), A., ii, 694.
 sensitisation of the processes of (PESKOV), A., ii, 828.
 of colloids (MURRAY; v. HAHN), A., ii, 37.
Coal, structure and formation of (MARCUSSON), A., i, 437.
 composition of (TIDESWELL and WHEELER), T., 2345.
 chemistry of (PEARSON, SINKINSON, and STOCKINGS), A., ii, 286.
 absorption of oxygen by (SCHRADER), A., i, 638.
 from Algerian mines, composition and calorific power of (FOIX), A., ii, 713.
 British, composition and analysis of (DRAKELEY), T., 221.
 estimation of volatile matter in (DESSEMOND), A., ii, 456.
Coal fire, blue flame produced by common salt on a (SMITHILLS), A., ii, 645.
Coal-tar colouring matters, chemical constitution and antiseptic action of (FAIRBROTHER and RENSHAW), A., i, 612.
Cobæa scandens, formation of anthocyanin in (JONESCO), A., i, 97.
Cobalt, occurrence and estimation of, in plants (BERTRAND and MOKRAGNATZ), A., i, 1099.
 in arable soils (BERTRAND and MOKRAGNATZ), A., i, 975.
Cobalt alloys with manganese, electrochemistry of (TAMMANN and VADERS), A., ii, 380.
 with tungsten (KREITZ), A., ii, 381.
Cobalt bases (cobaltamines), heats of solution and of transformation of (LAMB and SIMMONS), A., ii, 121.
 relative strengths of (LAMB and YNGVE), A., ii, 217.
 solubilities of salts of (BRÜNSTED and PETERSEN), A., ii, 199.
 hydrolysis of, by barium hydroxide (JOB), A., ii, 301.
 effect of, on the action of enzymes (FUNK), A., i, 481.
 complex chloroiodides of (BENRATH, BUCHER, and ECKSTEIN), A., ii, 515.
Aquopentamminecobaltiselenate (MEYER and MOLDENHAUER), A., ii, 72.
Aquatetramminecobaltichloride selenate, chloro- (MEYER and MOLDENHAUER), A., ii, 72.

Cobalt bases :—
Carbonatotetramminecobaltiselenate (MEYER and MOLDENHAUER), A., ii, 73.
Diaquotetramminecobaltiselenate (MEYER and MOLDENHAUER), A., ii, 72.
Hexamminecobaltic chloride, preparation of (MORGAN and SMITH), T., 1970.
 salts (EPHRAIM and MOSIMANN), A., ii, 575.
Hexamine(luteo)cobaltiselenate (MEYER and MOLDENHAUER), A., ii, 72.
Pentamminecobaltiselenates, chloro- and nitro- (MEYER and MOLDENHAUER), A., ii, 72.
Roseocobaltic salts, hydrolysis of (JOB), A., ii, 382.
Selenatopentamminecobalti-salts (MEYER and MOLDENHAUER), A., ii, 72.
Sulphatopentamminecobaltiselenate (MEYER and MOLDENHAUER), A., ii, 72.
Tetramminecobaltiselenates, dichloro- and dinitro- (MEYER and MOLDENHAUER), A., ii, 73.
 See also Eldmann's salt.
Cobalt salts, preparation and properties of hydrates and amines of (CLARK and BUCKNER), A., ii, 300.
Cobalt chloride, equilibrium in the system, ammonium chloride, water and (CLENDINNNEN), T., 801.
 fluoride, hydrate and amines of (CLARK and BUCKNER), A., ii, 300.
 hydroxide, preparation of colloidal solutions of (TOWER and COOKE), 853.
 nitrosotricarbonyl (MOND and WALLEN), T., 34.
 tungstate (SMITH), A., ii, 774.
Cobalt organic compounds :—
 with α -oximinoketones (PONZIO), A., i, 661.
Cobalt bases (cobaltamines), salts of (DUFF), T., 450; (MORGAN and SMITH), T., 1956, 2366.
 of nitro-colouring matters (MORGAN and KING), T., 1723.
Cobalt detection and estimation :—
 detection of (DITZ), A., ii, 229.
 estimation of, in silicate rocks (HACKL), A., ii, 458.
 estimation of, in steel (EDER), A., ii, 458.
 estimation and separation of (WILLARD and HALL), A., ii, 874, 875.
Cobalt lakes, of the alizarin series (MORGAN and SMITH), T., 160.

- Cobalt lakes**, with quinoneoximes (MORGAN and MOSS), T., 2857.
- Cocaine**, hydrolysis of (TROGER and SCHWARZENBERG), A., i, 167.
detection of, by the Vitali reaction (HARDY), A., ii, 95.
- Coconut**, nitrogen distribution of proteins from (FRIEDERMANN), A., i, 505.
- Coconut oil**, detection of, in butter (MUTTELET), A., ii, 236.
- Codeine**, salts, preparation of solutions of, for injection (MARTIN), A., i, 948.
estimation of, in opium (RAKSHIT), A., ii, 96.
- isoCodeine**, hydrogenation of (SPEYER and WIETERS), A., i, 47.
- ψ -Codeine**, hydrogenation of (SPEYER and WIETERS), A., i, 47.
- ψ -Codeine N-oxide**, and its picrate and sulphonic acid (SPEYER and WIETERS), A., i, 170.
- Codeineoxidesulphonic acids** (SPEYER and WIETERS), A., i, 169.
- Cod-liver oil**, sulphuric acid reaction for (RICHMOND and ENGLAND), A., ii, 792.
- Coke ash**, estimation of phosphorus in, colorimetrically (MISSON), A., ii, 718.
- Cola nuts**, detection of dextrose in extract of (ARNOLD), A., i, 311.
- Collagen**, isoelectric point of (THOMAS and KELLY), A., i, 387.
- Collidine**, condensation of acetaldehyde with (KONDO and TAKAHASHI), A., i, 1177.
- Collodion particles**, charge of (LOEB), A., ii, 742, 746.
thimbles, preparation of, and their use in osmosis (BROWN), A., ii, 690.
- Colloids**, fundamental conceptions of (ZSIGMONDY), A., ii, 129.
theory of (PESKOV), A., ii, 825.
theory of the mechanical synthesis of (SEKERA), A., ii, 693.
electrical synthesis of (BODFORS and FRÖLICH), A., ii, 826.
action of penetrating radium rays on (FERNAT and PAULI), A., ii, 202.
as indicators for photo-electric effects (PESKOV), A., ii, 828.
chemistry of (ZSIGMONDY), A., ii, 484.
adsorption of electrolytes by (WEISER), A., ii, 262.
coagulation of (WESTGREN and REITSTÖTTER), A., ii, 625; (MUKHERJEE and PAPACONSTANTINOU), A., ii, 694.
by electrolytes (MURRAY), A., ii, 37.
- Colloids**, coagulation of, by sunlight (GANGULY and DHAR), A., ii, 604.
energy of the coagulation of (FRICKE), A., ii, 484.
precipitation and peptisation of (CHATTERJI and DHAR; OSTWALD and v. HAHN), A., ii, 205.
electrical precipitation of (HALL), A., ii, 555.
precipitation of, by electrolytes (WEISER and NICHOLAS), A., ii, 266.
effect of hydrogen-ion concentration on the precipitation of (TARTAR and GALEY), A., ii, 829.
pseudo-extraction of (CHARITSCHKOV), A., ii, 826.
swelling and osmosis of (BARTELL and SIMS), A., ii, 267.
transition from crystalloids to (WITZEMANN), A., ii, 428.
effect of salts on the ultrafiltration of (RICHTER-QUITNER), A., ii, 204.
effect of, on the rate of evolution of carbon dioxide from solution (FINLAY and HOWELL), T., 1046.
hydrophilic, protective and flocculating action of (FREUNDLICH and LOENING), A., ii, 356.
lyophile, electrical behaviour of (KRUYT and DE JONG), A., ii, 357.
protective (GUTHRIE and HUBER; GUTHRIE, WOLF, and KRESS), A., ii, 203; (GUTHRIE, HUBER, and ZWEIFLE), A., ii, 485; (GUTHRIE and EMSLANDER), A., ii, 625.
lecture experiment on (FRIEND), A., ii, 267.
 rôle of, in catalysis (IREDALE), T., 1536.
effect of, on the corrosion of metals (FRIEND and VALLANCE), T., 466.
in plants (SAMEC and FERJANČIĆ), A., i, 115; (SAMEC and MAYER), A., i, 921.
action of muscle poisons on (RIESSER and NEUSCHLOSZ), A., i, 1212.
estimation of, in urine (OTTENSTEIN), A., i, 609.
- Colloidal particles**, electrical charges on (MUKHERJEE), A., ii, 198; (LOEB), A., ii, 354, 467, 746.
solutions, double refraction of (ZOCHEK), A., ii, 102.
dielectric constants of (ERRERA), A., ii, 694.
cell for the observation of (FAIRBROTHER), A., ii, 485.
coagulation of (WIEGNER, MAGASANIK, and GESSNER), A., ii, 356.
organic, effect of salts on the dispersion of (TADOKORO), A., ii, 204.

- Colloidal state**, theory of (CHARIT-SCHKOV), A., ii, 827.
 systems, nephelometry of (BECHHOLD and HEBLER), A., ii, 693.
 analytical chemistry of (GUTHRIE, HUBER, and KUPPINGER), A., ii, 396.
- Colophenic acid** (ASCHAN), A., i, 221.
- Colophenic acids** (FAHRION), A., i, 423.
- Colorimeters** (GILLESPIE), A., ii, 157.
 (MANNERBACH), A., ii, 158; (MOREAU and BONIS), A., ii, 862.
- Colorimetric analysis**. See Analysis.
- Colorimetry** (DOSNE; FLEURY; GARREAU), A., ii, 518.
 standard measurement of (IVES), A., ii, 221.
 use of coloured glasses as standards in (SONDÉN), A., ii, 862.
- Colostrum**, estimation of proteins in A., ii, 670.
 camel's, analysis of (FALES), A., i, 1090.
- Colour**, theory of (MOIR), T., 1808; (WELLS), A., ii, 464
 and chemical constitution (MOIR; KERRMANN), A., ii, 333.
 calculation of, of azo-colouring matters (MOIR), T., 1555.
- Colour vision**, physical chemistry of (WEIGERT), A., i, 607.
- Colouring matter**, $C_{11}H_{11}O_3N_3$, from ethyl 3-hydroxy-5-methylpyrrole-4-carboxylate and diazobenzene chloride (FISCHER and HERRMANN), A., i, 1055.
- $C_{21}H_{16}N_2$, and its salts, from condensation of quinoline and epichlorohydrin (GIUA), A., i, 681.
- Colouring matters**, preparation of, and their intermediate products (SOCIETY FOR CHEMICAL INDUSTRY IN BASLE), A., i, 579.
 absorption of light by (LASAREV), A., ii, 332.
 bleaching of, by light (LASAREV), A., ii, 103.
 fluorescence of, in relation to wavelength of the exciting light (VAVILOV), A., ii, 181.
 effect of hydrogen-ion concentration on the diffusion of, through membranes or protein sols (BETHE), A., ii, 352.
 adsorption of, by crystals (PANETH and VORWERK), A., ii, 619.
 preparation of intermediates for (ATAK and SOUTAR), A., i, 259; (THOMAS, DAVIES, and SCOTTISH DYES, LTD), A., i, 260.
 of the alizarin series (MORGAN and SMITH), T., 160.
- Colouring matters** from camphoric anhydride (SIRCAR and DUTT), T., 1283.
 coal tar, chemical constitution and antiseptic action of (FAIRBROTHER and KENSHAW), A., i, 612.
 cyanine (MILLS), T., 455; (MILLS and BRAUNHOLTZ), T., 1489; (BRAUNHOLTZ and MILLS), T., 2004.
 antiseptic properties of (BROWNING, COHEN, and GULBRANSEN), A., i, 1097.
 from $\alpha\alpha'$ -dicyanodibenzyl diketone (DUTT and SEN), T., 2663.
 flavone, absorption spectra of (SHIBATA and NAGAI), A., ii, 413.
 containing the furan ring (KENSHAW and NAYLOR), A., i, 566.
 indigoid (FRIEDLANDER, HERZOG, and V. VOSS), A., i, 764.
 of the isatin-yellow series (MARTINET), A., i, 278.
 mono- and di-chelate mordant (MORGAN and SMITH), T., 162.
 monocyclic, calculation of the colour of (MOIR), A., ii, 333.
 optically active (SINGH, RAI, and LAL), T., 1421.
 from phenanthraquinone (SIRCAR and DUTT), T., 1944; (DUTT), T., 1951.
 from "saccharin" (DUTT), T., 2389.
 red sulphide (WATSON and DUTI), T., 1939, 2414.
 from 1:2:4:5-tetrahydroxybenzene (MUKERJI), T., 545.
 triarylmethane (BRITISH DYE STUFFS CORPORATION, LTD., GREEN, SAUNDERS and BATE), A., i, 1068.
 vat, containing sulphur and nitrogen (REISSERT), A., i, 583.
 estimation of, in wines (FRESENIUS and GRÜNHUT), A., ii, 96.
 nitro-, cobaltamines of (MORGAN and KING), T., 1723.
- See also:—
 Bilirubin.
 Biliverdin.
 Hemocyanin.
 Hemoglobin.
 Lignocyanin.
- Columbite** from Ishikawa (SHIBATA and KIMURA), A., ii, 220.
 from South Dakota (HEADEN), A., ii, 387.
- Columbium pentachloride**, preparation of, and its insulating properties (BILTZ and VOIGT), A., ii, 302.
- Columbium estimation and separation**:—
 estimation of, by reduction (TREADWELL, FREULER, and WEBER), A., ii, 781.

Columbium estimation and separation:—

separation of, from tantalum (MERRILL), A., ii, 230

Colza oil, fatty acids of (RAYMOND), A., i, 798.

Combustion, temperature of (BRONN), A., ii, 548.

of mixed gases (PAYMAN and WHEELER), T., 363.

Compounds, complex (MAGNUS), A., ii, 559

spiro-Compounds, formation and stability of (KON), T., 513; (DICKENS, KON, and THORPE), T., 1496; (BIRCH and THORPE), T., 1821.

Conifers, resin acids of (ASCHAN, FONTELLI, and SIMOLA), A., i, 1152.

Co-ordination and residual affinity (MORGAN and SMITH), T., 160, 1956, 2866; (MORGAN and DREW), T., 922; (MORGAN and KING), T., 1723; (MORGAN, DREW, and BARKER), T., 2432; (MORGAN and MOSS), T., 2857; (MORGAN and LEDBURY), T., 2892.

Copal oil, constants of (PIERAERTS), A., i, 908.

Copper, emissivity of (LUBOWSKY), A., ii, 108.

solubility of, in dilute solutions of benzaldehyde (BERNOULLI and SCHAAF), A., i, 1029.

rate of solution of, in ferric alum (COLLENBERG and BODFORS), A., ii, 431.

catalytic, preparation of (PICCARD), A., ii, 216.

catalytic action of (KOPETSHNI and WIENLER), A., i, 844; (PALMER), A., ii, 437; (BROWN and HENKE), A., ii, 833.

cementation of, by means of chromomanganese (SROVICH and CARTOCETI), A., ii, 571.

with ferro-manganese (SROVICH and CARTOCETI), A., ii, 68

colloidal, coloured forms of (PAAL and STEYER), A., ii, 294.

corrosion of, by salt solutions (MÜLLER), A., ii, 645.

action of ammonium nitrate and aqueous ammonia on (BASSETT and DURRANT), T., 2630.

action of ammonium nitrite on (DHAR), A., ii, 49.

combination of phosphorus with (EDWARDS and MURPHY), A., ii, 645.

action of, with trichloroacetic acid (DOUGHTY and FREEMAN), A., i, 427.

Copper alloys with gold, electrochemistry of (LORENZ, FRAENKEL, and WORMSER), A., ii, 21.

reaction limit of (TAMMANN), A., ii, 75.

with manganese, electrochemistry of (TAMMANN and VADERS), A., ii, 380.

with mercury (PAAL and STEYER), A., ii, 446.

with nickel and zinc (VOIGT), A., ii, 295.

Copper salts, investigation of the complex formation in solutions of (GUNTHER-SCHULZE), A., ii, 504.

reduction of ammoniacal solutions of (KOHN), A., ii, 149.

use of solutions of, as sprays (G. and (MME) G. VILLEDIEU), A., i, 415.

in tissues and in tumours (WHITE), A., i, 399.

Copper arsenite. See Scherle's green.

hydroxide, rate of dehydration of (H. and V. VON EULER), A., ii, 832.

solubility of, in sodium hydroxide solutions (MELBYE), A., ii, 851.

colloidal, preparation of (PAAL and STEYER), A., ii, 216.

oxide, action of, with molten alkali chlorides (HEDVALL and LÖNNBERG), A., ii, 68.

reduction of, by hydrogen (PEASE and TAYLOR), A., ii, 148.

trivalent oxide of (SCAGLIARI and TORELLI), A., ii, 68.

peroxides (ALDRIDGE and APPLEBEY), T., 238.

sulphate, equilibrium of ammonium sulphate, potassium sulphate, water and (HAYAMI), A., ii, 748.

sulphide, structural formula of (GLUUD), A., ii, 572.

oxidation of (GLUUD), A., ii, 446.

Cupric chloride, spectrophotometry of solutions of (GETMAN), A., ii, 411.

reaction of hypophosphorous acid with (MITCHELL), T., 1624.

action of, on magnesium organic compounds (TURNER), A., i, 23.

lithium chloride, absorption of light by (GETMAN), A., ii, 412.

tetrammine nitrate and nitrite, properties of (BASSETT and DURRANT), T., 2630.

alkali sulphates, dissociation pressures of (CAVEN and FERGUSON), T., 1406.

Cuprous bromide and chloride, phototropy of (SINGH), T., 782.

Copper :—

- Cuprous** chloride, preparation of, and its oxidation by sulphur dioxide (WARDLAW and PINKARD), T., 210.
- absorption of carbon monoxide by acid solutions of (KROFF), A., ii, 657.
- action of sulphur on (PINKARD and WARDLAW), T., 1300.
- haloids, crystallography of (WYCKOFF and POSNJAK), A., ii, 295.
- compounds of ammonia with (BILTZ and STOLLENWERK), A., ii, 67.
- sulphide, solubility of (TRUMPLER), A., ii, 111.
- Cuprates** (ALDRIDGE and APPLEBY), T., 242.
- Copper organic compounds :—**
- Copper bases** (*cuprammines*), complex salts of (HALLE and HIRSCHKO), A., ii, 768.
- compounds of, with polyhydroxy-compounds (TRAUBE), A., i, 718.
- Copper detection, estimation, and separation :—**
- detection of (THOMAS and CARPENTIER), A., ii, 86; (CARLETTI), A., ii, 787.
- detection and estimation of traces of (HAHN and LEIMBACH), A., ii, 870.
- estimation of, electrovolumetrically (ZINTL and WATTENBERG), A., ii, 871.
- estimation of, iodometrically (LANG), A., ii, 318.
- estimation of, iodometrically, in presence of arsenic (KOLTHOFF and CREMER), A., ii, 86.
- estimation of, iodometrically, in presence of iron (WARK), T., 358; (WOBER), A., ii, 588.
- estimation of, volumetrically (MINOVICI and IONESCU), A., ii, 162.
- estimation of, volumetrically, in presence of iron (THORNTON), A., ii, 528.
- estimation of, volumetrically, with sodium nitroprusside (JORET), A., ii, 872.
- estimation of, and its separation from antimony, lead and tin (KLING and LASSIEUR), A., ii, 86.
- estimation of antimony in (EVANS), A., ii, 231.
- separation of, from the group metals (WILLARD and HALL), A., ii, 872.
- Corchorus capsularis* (jute), glucoside from the leaves of (SAHA and CHOUDHURY), T., 1044.
- Cornetite**, absence of cobalt in (SCHOEP), A., ii, 219.

- Cornus sanguinea*, absence of hydrocyanic acid from the leaves of (ROSENTHALER), A., i, 412.
- Corpses**, constituents of the wax of (GOY and WENDE), A., i, 969.
- Corresponding states**, law of (HERZ), A., ii, 821.
- Corrosion** of metals, retardation of, by protective colloids (FRIEND and VALLANCE), T., 466.
- Corydaline**, constitution of (SPATH and LANG), A., i, 168; (GADAMER and V. BRUCHHAUSEN), A., i, 675.
- Coto-bark**, constituents of (SPATH and FUCHS), A., i, 558.
- Cotoin**, synthesis of (SPATH and FUCHS), A., i, 558.
- Cotton**, structure of the hair of (DENHAM), A., i, 615.
- heat developed during the mercerisation of (BARRATT and LEWIS), A., i, 526.
- effect of salts on dyeing of (AUERBACH), A., ii, 353.
- raw, constituents of (FARGHER and WITHERS), A., i, 615.
- detection and estimation of acidity and alkalinity in (COWARD and WIGLEY), A., ii, 531.
- Cotton-seed meal**, nitrogen distribution of proteins from (FRIEDEMANN), A., i, 505.
- Coumaran-2-ones**, 4- and 6-chloro-, and their derivatives (MINON and STEPHEN), T., 1601.
- Coumaranone series**, studies in the (MINON and STEPHEN), T., 1598.
- Coumarins**, preparation of (PONNDORF), A., i, 565.
- Creatine**, origin of (STEUDEL and FREISE), A., i, 793.
- in muscle (RIESSER), A., i, 791; (HAMMETT), A., i, 1090.
- formation of, in muscular dystrophy (GIBSON and MARTIN), A., i, 300.
- estimation of, in blood (BEHRE and BENEDICT), A., i, 789.
- Creatinine**, origin of (STEUDEL and FREIRE), A., i, 793.
- in muscle (HAMMETT), A., i, 1090.
- metabolism. See Metabolism.
- estimation of (PFIZENMAIER and GALANOS), A., ii, 797.
- estimation of, in blood (BEHRE and BENEDICT), A., i, 789.
- o*-Cresol, bromoamino-derivatives, and their derivatives (RAIFORD), A., i, 335.
- p*-Cresol, oxidation of (PUMMERER, MELAMED, and PUTTFARCKEN), A., i, 1161.
- action of camphoric anhydride with (KRISHNA), T., 255.

- Cresols** and their derivatives, reciprocal polarity effects in (LAPWORTH and SHOESMITH), T, 1391.
- equilibrium and properties of mixtures of phenol with (KENDALL and BEAVER), A., i, 186.
- compounds of, with acetone, benzene, ethyl alcohol and ethyl ether (BERL and SCHWEBEL; C. and W. VON RECHENBERG), A., i, 932.
- Cresols**, bromoamino-, and their derivatives (RAIFORD and COUTURE), A., i, 931.
- nitro-, lead derivatives of (GODDARD and WARD), T., 265.
- o*-Cresolcamphoroin**, and its derivatives, and *di*bromo- (SINGH, RAI, and LAL), T., 1426.
- o*-Cresol-5-sulphonic acid**, 3-nitro-, salts of (DAVIES), T., 791.
- o*-Cresolsulphonphthalein**, absorption spectra of (ORNDORFF, GIBBS, SCOTT, and JACKSON), A., ii, 806.
- Cristobalite** from Washington (SHANNON), A., ii, 450.
- Critical temperature**, relation between boiling and melting points and (PRUD'HOMME), A., ii, 349.
- Croceine**, action of phenylhydrazine and sulphite on (BUCHERER and ZIMMERMANN), A., i, 466.
- Crocidolite** from Pennsylvania (WHERRY and SHANNON), A., ii, 518.
- Crocodile oil**, constituents of (KOBAYASHI), A., i, 792.
- Croconic acid**, constitution of (HOMOLKA), A., i, 630.
- Crops**, effect of silicic acid on production of (LEMMERMANN and WIESSMANN), A., i, 1103.
- Crotonaldehyde**, preparation of (CONSORTIUM FÜR ELEKTROCHEMISCHE INDUSTRIE), A., i, 1115.
- catalytic hydrogenation of (GRUNSTEIN), A., i, 112.
- Crotonic acid**, transformations of, in ultra-violet light (STOEBMER and ROBERT), A., i, 519.
- vanillylamide of (OTT and ZIMMERMANN), A., i, 187.
- Crotonic acid**, β -amino-, ethyl ester, compound of methyl azodicarboxylate and (DIELS), A., i, 774.
- Crucibles**, silica, use of, in estimation of potassium (JONES and REEDER), A., ii, 85.
- Cryoscopy**, use of cinnamic acid in (FALCIOLA), A., ii, 421.
- Crystal** from eucalyptus oil (PENFOLD), T., 266.
- Crystals**, structure of (BRAGG), T., 2766.
- Crystals**, structure of, and constitution of atoms (NIGGLI), A., ii, 36.
- by means of Röntgen rays (HULL and DAVEY; HULL), A., ii, 624.
- of complex haloids, by means of Röntgen rays (SCHERRER and STOLL), A., ii, 514.
- electronic (HUGGINS), A., ii, 744.
- absorption of Röntgen rays in (AURÉN), A., ii, 810.
- reflexion of Röntgen rays from (DARWIN), A., ii, 416; (DARDORD), A., ii, 673.
- scattering of Röntgen rays by (BRAGG, JAMES, and BOSANQUET), A., ii, 184.
- Röntgenographic determination of the arrangement of (POLÁNYI), A., ii, 623.
- Röntgen ray investigation of the lattice of (GERLACH), A., ii, 36, 265.
- use of Röntgen rays in analysis of (CLARK and DUANE), A., ii, 483; (QUARDER), A., ii, 677.
- luminous path of α -rays in (GEIGER and WERNER), A., ii, 183.
- photography of (FRANÇOIS and LORMAND), A., ii, 127.
- rotatory power of (LONGCHAMON), A., ii, 603.
- ionic properties of (GRIMM), A., ii, 127.
- size of atoms in (PEASE), A., ii, 428.
- radius of hydrogen atoms in (AMINOFF), A., ii, 496.
- loosening of the lattice in (v. HEVESY), A., ii, 623.
- hardness of (REIS and ZIMMERMANN), A., ii, 745.
- adsorption of colouring matters by (PANETH and VORWERK), A., ii, 619.
- absorption of gases by (SEELIGER and LAPKAMP), A., ii, 30.
- velocity of decomposition of (SIEVERTS), A., ii, 700.
- diamond type, size of atoms in (PEASE), A., ii, 634.
- liquid, existence of (LEHMANN), A., ii, 692.
- formation of, and the symmetry of molecules (VORLÄNDER), A., ii, 554.
- structure of drops of (LEHMANN), A., ii, 201.
- mixed, resistance limits in (MASING), A., ii, 37.
- binary mixed, chemical behaviour of (TAMMANN), A., ii, 63.
- organic, structure of (BRAGG; BECKER and JANCKE), A., ii, 128.

- Crystalline** condition, distinction between the amorphous and the (HABER), A., ii, 553.
 substances, space formulæ for (RINKE), A., ii, 368.
- Crystallisation**, process of (SCHAUM), A., ii, 264.
 produced by annealing (GAUBERT), A., ii, 35.
 starting of, by friction of a glass rod against the wall of the vessel (DEDE; FRICKE), A., ii, 744.
 prevention of, by addition of proteins (CAVAZZANI), A., ii, 745.
 spontaneous, influence of pressure on (HASSELBLATT), A., ii, 35.
 of supercooled liquids (HINSHELWOOD and HARTLEY), A., ii, 200.
- Crystalloids**, transition from, to colloids (WITZEMANN), A., ii, 428.
- Crystal-violet-leucosulphonic acid** (WIELAND and SCHEUING), A., i, 58.
- Cubebin** (MAMELI), A., i, 347.
- Cubebinic acid**, dibromohydroxy-, and hydroxy-, amides of (MAMELI), A., i, 348.
- Cubebinolide**, derivatives of (MAMELI), A., i, 347.
- Culture solutions**, estimation of alkalinity of (MICHAELIS), A., i, 405.
- iso- ψ -Cumenol** (3:5:6-trimethylphenol) (v. AUWERS and SAURWEIN), A., i, 1033.
- α - ψ -Cumidine**, di- ω -chloro-. See 2:3:5-Trimethylbenzene, 1-dichloroamino-
- Cuminaldehyde** from eucalyptus oil (PENFOLD), T., 266.
- Cumylcoumquinoned-*i*-imide**, chloro- (GOLDSCHMIDT and STROHMENGER), A., i, 1005.
- Cuprates**. See under Copper.
- Cuprene**, constitution of (KAUFMANN and SCHNEIDER), A., i, 245.
- Cupretenine methyl ether**, hydrochloride of (HEIDELBERGER and JACOBS), A., i, 673.
- Cureanolic acid** (GRIMME), A., i, 98.
- Curcine** (GRIMME), A., i, 98.
- Curite** (SCHOEP), A., ii, 77.
- Current**. See *Ribes rubrum*.
- Cuttlefish**, action of inky secretion of, on pyrrole (RONDONI), A., i, 64.
- Cyanamide**, preparation of carbamide from (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 993.
 physiological action of (HESSE), A., i, 1093.
- Cyanamides**, condensation of hydrochlorides of, with phenols (SHORT and SMITH), T., 1803.
- Cyanamidoethyl alcohol**, and its derivatives (FROMM and HOWOLD), A., i, 529.
- Cyanates**, thio-. See Thiocyanates.
- Cyanoic acid** and **Cyanides**. See under Cyanogen.
- Cyanines**, thio-, tautomerism of (MILLS and BRAUNHOLTZ), T., 1489.
- isoCyanines**, brominated (MOUDGILL), T., 1509.
 thio- (BRAUNHOLTZ and MILLS), T., 2004.
- Cyanine colouring matters** (MILLS), T., 455; (MILLS and BRAUNHOLTZ), T., 1489; (BRAUNHOLTZ and MILLS), T., 2004.
 antiseptic properties of (BROWNING, COHEN, and GULBRANSEN), A., ii, 1097.
- Cyanogen**, spectrum of (KRATZER), A., ii, 4.
 band spectrum of (OKUBO), A., ii, 601.
- Cyanogen bromide**, action of, on tertiary arsines (STEINKOPF, DONAT, and JAEGER), A., i, 994.
 haloids, action of, on phenylhydrazine (PELLIZZARI), A., i, 585.
- Hydrocyanic acid**, formation of, from nitrogen and hydrocarbons (KOEING and HUBBUOH), A., i, 642
 preparation of, in the laboratory (FRITZMANN), A., i, 1128.
 production of, by *Bacillus pyocyaneus* (PATTY), A., i, 407.
 synthesis of, by oxidation of various organic compounds (FOSSE; FOSSE and HIEULLE), A., i, 117.
 tautomerism of (USHERWOOD), T., 1604.
 polyamide of, and its derivatives (GRISCHKEVITSCH-TROCHIMOVSKI), A., i, 723.
 oxidation of formaldehyde to (FOSSE and HIEULLE), A., i, 523.
 in plants (ROSENTHALER), A., i, 412, 614; (SIEGFRIED), A., i, 614.
 calcium salt, preparation of, pure (PINCASS), A., i, 532.
 magnesium salt, and its complex silver compounds (FICHTER and SUTER), A., i, 532.
 mercuric salt, compounds of, ammonia with (BRINKLEY), A., i, 724.
 potassium salt, crystal structure of (BOZORTH), A., i, 441; (COOPER), A., i, 1128.
 potassium cadmium, potassium mercury, and potassium zinc salts, crystal structure of (DICKINSON), A., i, 532.
 detection of (SUNDBERG), A., ii, 404.
- Cyanides**, estimation of, electrometrically, in presence of haloids (MÜLLER and LAUTERBACH), A., ii, 403.

Cyanogen :—

Cyanic acid in blood (NIDOUX and WELTER), A., i, 789.

isoCyanic acid, condensation of, with aromatic o-aminosulphonic acids (SCOTT and COHEN), T., 2034.

Cyanogen, estimation of (YANAGISAWA), A., ii, 667.

Cyanohydrins, synthesis of, by means of emulsion (NORDEFELDT), A., i, 66.

Cyanuric dihydrazidomonoazide, monohydrazidodiazide and triazide (OTT), A., i, 1128.

apocyclene (KOMPPA and ROSCHIER), A., i, 1167.

Cyclic compounds, formation of, from halogenated open-chain derivatives (CHANDRASENA and INGOLD), T., 1306; (INGOLD), T., 2676.

formation of, from hydroaromatic dicarboxylic acids (WINDAUS and HUCKEL), A., i, 658.

space configuration of (DERX), A., i, 651.

tautomerism of, with open-chain isomerides (KON, STEVENSON, and THORPE), T., 650.

halogen derivatives, absorption spectra of (GRAHAM and MACBETH), T., 2601.

Cymbopogon cæsius (inch grass), essential oil from (MOUDGILL and IYER), A., i, 945.

p-Cymene, bromo-2-amino-, and its derivatives (WHEELER and SMITHEY), A., i, 832.

γ-2-Cymylbutyric acid (RUZICKA and MINGAZZINI), A., i, 1001.

β-2-Cymylethyl alcohol, and its bromide (RUZICKA and SEIDEL), A., i, 562.

β-2-Cymylethylmalonic acid, ethyl ester (RUZICKA and MINGAZZINI), A., i, 1001.

methyl ester (RUZICKA and SEIDEL), A., i, 562.

β-2-Cymyl-α-methylbutyric acid, and its chloride (RUZICKA and SEIDEL), A., i, 562.

Cyprine, from New York (LEWIS and BAUER), A., ii, 714.

Cyprinus carpio (carp), constituents of the ovaries of (FAURE-FREMIET and GARRAULT), A., i, 700.

Cystine (MERRILL), A., i, 326.

preparation of (SCHMIDT), A., i, 992.

acid hydrolysis of (HOFFMAN and GORTNER), A., i, 429.

oxidation of, in the animal organism (LEWIS and ROOT), A., i, 487.

excretion of, as cysteine (LEWIS, MCGINTY, and ROOT), A., i, 1088.

estimation of, in proteins (FOLIN and LOONEY), A., ii, 539.

Cysts, ovarian-dermoid, unsaturated alcohols from (MUCK), A., i, 1092.

Cytopectans (CLAYSON, NORRIS, and SCHRYVER), A., i, 206.

Cytopectic acid (CLAYSON, NORRIS, and SCHRYVER), A., i, 206.

D.

Dakin's hypochlorite solution, action of, on organic compounds (ENGFELDT), A., i, 812.

Deaminoproteins (HERZIG and LIEB), A., i, 386.

Decahydronaphthalene (*decalin*), physical properties of (HERZ and SCHULTAN), A., i, 647.

Decahydronaphthalenes, isomerism of (MOHR), A., i, 248.

Decalactone (GRÜN and WIRTH), A., i, 805.

Decalin. See Decahydronaphthalene.

Decanedial, and its derivatives (ROSENMUND, ZETZSCHE, and ENDERLIN), A., i, 431.

Decarboxylation in biochemistry (SPIRO), A., i, 1213.

Δ⁹-Decenoic acid, synthesis of (GRÜN and WIRTH), A., i, 804.

from butter, and *dibromo-*, and their methyl esters (GRÜN and WIRTH), A., i, 806.

Decoisobutylamide (ASAHINA and ASANO), A., i, 506.

Decoic acid, *i*-chloro-, methyl ester (GRÜN and WIRTH), A., i, 805.

i-hydroxy-, and its methyl ester (GRÜN and WIRTH), A., i, 805.

n-Decyl fluoride (SWARTS), A., i, 101.

Dehydrocholic acid, trisemicarbazone of (BORSCHKE and HALLWASS), A., i, 1159.

Dehydrolithocholic acid, methyl ester, and its oxime (BORSCHKE and HALLWASS), A., i, 1159.

Dehydroxidation (MÜLLER), A., i, 110.

Delphinidae, oil from the heads of (NAKATOGAWA and KOBAYASHI), A., i, 701.

Demethoxy-yanogonine (MURAYAMA and MAYEDA), A., i, 265.

Demethylscopolinone, and its salts and derivatives (HESS and WAHL), A., i, 857.

Denitrification with formates (GROENEWEGER), A., i, 971.

Density (*specific gravity*), determination of, of liquids (WIEDBRAUK), A., ii, 740.

contraction of flasks, used in the determination of, of gases (MOLES and MIRAVALLES), A., ii, 617.

- Deoxybenzoin** acetal and anil, and their derivatives (KNOEVENAGEL and GOOS), A., i, 752.
- α - and β -**Deoxybenzoinpinacones** (GARCÍA BANÚS and PASCUAL VILA), A., i, 734.
- Deoxybillionic acid**, semicarbazone of (BORSCH and HALLWASS), A., i, 1160.
- iso***Deoxybillionic acid**, formula of, and its derivatives (WIELAND and ADICKES), A., i, 838.
- 2-Deoxyglucose**, and its derivatives (BERGMANN, SCHOTTE, and LECHINSKY), A., i, 227.
- Deoxyindigotin**, and its sulphonic acid and sodium salt (BORSCH and MEYER), A., i, 55.
- Deoxymatrine**, and its salts and derivatives (KONDÓ, KISHI, and ARAKI), A., i, 269.
- Deoxyscopolamine**, and its salts (HESS and WAHL), A., i, 856.
- Deoxytropol bromide** (HESS and WAHL), A., i, 855.
- α - and β -**Deoxytropylscopolein**, and their salts (HESS and WAHL), A., i, 855.
- Detectors**, new, for wireless telegraphy (HEIKE), A., ii, 19.
- Detonators**, α -particles as (HENDERSON), A., ii, 606.
- Dewindtite**, a radioactive mineral (SCHOEP), A., ii, 305.
- Dextrin**, adsorption in solutions of, and their emulsifying action (CLARK and MANN), A., ii, 550.
- Dextrose** (*d-glucose*; *grape-sugar*), mutarotation of (MURSCHAUSER), A., i, 322, 432; ii, 92.
- solubility of, in water (JACKSON and SILSBEE), A., i, 986.
- decomposition of, in alkaline solution (WATERMAN and VAN TUSSEN-BROEK), A., i, 433.
- decomposition of, by pyocyanic bacilli (AUBEL), A., i, 201.
- lactic acid fermentation of, by peptone (SCHLATTER), A., i, 1096; (BAUR and HRRZFELD), A., i, 1097.
- function of phosphates in oxidation of, by hydrogen peroxide (HARDEN and HENLEY), A., i, 433.
- oxidation of, in the body (HIRSCH), A., i, 87.
- catalytic reduction of (CAKE), A., i, 523.
- electrolytic reduction of (FINDLAY and WILLIAMS), A., i, 523.
- acetone derivatives of (IRVINE and PATTERSON), T., 2146.
- action of ammonia on (LING and NANJI), A., i, 631.
- action of ozone on solutions of (SCHONEBAUM), A., i, 223.
- Dextrose**, distribution of, in arterial and venous blood (HENRIQUES and EGE), A., i, 191.
- permeability of blood-corpuscles for (VAN CREVELD and BRINKMAN), A., i, 192.
- action of injections of, on diuresis (CORT), A., i, 1091.
- sulphate (NEUBERG and LIEBERMANN), A., i, 222.
- and its tetra-acetyl derivative, sulphate compounds of, and their salts (OHLE), A., i, 986.
- detection of, biochemically (ARNOLD), A., i, 311.
- detection of, in urine by means of *o*-nitrophenylpropionic acid (RODIL-LON), A., ii, 166.
- estimation of (GREINER), A., ii, 400; (LEMKE and LANDSBERG), A., ii, 724; (KRULL; VAN DER HAAR), A., ii, 790; (FLEURY and BOUTOT), A., ii, 879.
- separation of sucrose from, by dialysis (CONGDON and INGERSOLL), A., i, 322.
- Diabetes** (*glycosuria*), chemistry of (IONESCU), A., i, 198.
- acetone substances in blood in (FITZ), A., i, 192.
- acetonuria in (HUBBARD and NICHOLSON), A., i, 969.
- aldehyde in urine in (STEPP and FEULGEN), A., i, 300.
- aldol and acetaldehyde in urine in (FRICKE), A., i, 300.
- carbohydrate metabolism in (ALLEN and WISHART), A., i, 893; (WAGNER and PARNAS), A., i, 965.
- utilisation of fat in (BLATHERWICK), A., i, 89.
- administration of levulose in (DESGREZ, BERRY, and RATHERY), A., i, 1215.
- adrenaline (ELIAS and SAMMARTINO), A., i, 86.
- phloridzin, sugar in the kidneys in (NASH), A., i, 497.
- action of desiccated thyroid in (ASHER and HERRISBERGER), A., i, 293.
- Di-5-acenaphthylcarbamide** (FLEISCHER and SCHRANZ), A., i, 1143.
- Diacetonedeoxyglucose** (FREUDENBERG and BRAUNS), A., i, 1117.
- Diacetoneglucosene** (FREUDENBERG and BRAUNS), A., i, 1117.
- Diacetoneglucose**, transformations of (FREUDENBERG and BRAUNS), A., i, 1117.
- sulphuric ester of (LEVENE, MEYER, and WEBER), A., i, 987.

- Diacetonexylose** (FREUDENBERG and SVANBERG), A., i, 1116.
- pp'-Diacetoxybenzene** (FICHTER and JAECK), A., i, 62.
- 7:12-Diacetoxydibenzothianthren**, 5:14-dihydroxy- (BRASS and KÖHLER), A., i, 1051.
- 5:14-Diacetoxydibenzothianthrenquinone** (BRASS and KÖHLER), A., i, 1051.
- Diacetoxydiethyl sulphide** (FARBWERNER VOM MEISTER, LUCIUS, & BRÜNING), A., i, 420.
- 3:5-Diacetoxydimercuri-4-hydroxybenzaldehyde** (HENRY and SHARP), T., 1058.
- 3:5-Diacetoxydimercurialicylaldehyde** (HENRY and SHARP), T., 1056.
- Diacetoxy-7:7-dimethyl- α -dinaphthaxanthens** (SEN-GUPTA and TUCKER), T., 560, 562.
- Diacetoxymercurinroanilines** (KHARASCH, LOMMEN, and JACOBSON), A., i, 604.
- Diacetoxymercuri-5-nitrosalicylic acid** (PAIZISS and PROSKOURIAKOFF), A., i, 604.
- Diacetoxy-7-methyl-7-ethyl- α -dinaphthaxanthens** (SEN-GUPTA and TUCKER), T., 563.
- 1:5-Diacetoxynaphthalene**, and *d*-nitro- (HEMELMAYR), A., i, 836.
- 6:11-Diacetoxy- $\beta\beta$ -naphthaphenthiazine**, 3-amino-, acetyl derivative (FRIES and KERKOW), A., i, 578.
- 2:8-Diacetoxyphenyl- n -octaldehyde** (MAJIMA and TAHARA), A., i, 263.
- 3:4-Diacetoxystyrene**, ω -nitro- (ROSENMUND and PFANNKUCH), A., i, 1030.
- Diacetylacetone**, preparation of, and its salts and derivatives (COLLIE and REILLY), T., 1984.
- 1:1'-Diacetyldihydro-4:4'-dipyridyl** (DIMROTH and FRISTER), A., i, 678.
- Diacetylurpurin 2-methyl ether** (DIMROTH, SCHULTZE, and HEINZE), A., i, 157.
- Diacetylquinizarin**, *mono*- and *di*-bromo- (DIMROTH, SCHULTZE, and HEINZE), A., i, 157.
- Diacetylsuccinic acid**, ethyl ester, tautomerism of (KAUFMANN), A., i, 985.
- isomerism of, and its estimation and iron salts (KNORR and KAUFMANN), A., i, 220.
- Diacetylsuccinic acid**, *di*bromo-, ethyl ester, preparation of (HIRST and MACBETH), T., 2175.
- 1:1'-Diacetyltetrahydro-4:4'-dipyridyl** (DIMROTH and HEENE), A., i, 48.
- 1:1'-Diacetyltetrahydrodiquinolyl** (DIMROTH and HEENE), A., i, 50.
- $\alpha\alpha'$ -Diacetyl- $\alpha\alpha'\beta\beta'$ -tetraphenyltetrazan** (GOLDSCHMIDT and EULER), A., i, 475.
- $\alpha\alpha'$ -Diacetyl- $\alpha\alpha'\beta\beta'$ -tetra-*p*-tolyltetrazan** (GOLDSCHMIDT and EULER), A., i, 476.
- Dialkyl sulphides**, chloro- (POPE and SMITH), T., 1166.
- 7:9-Dialkyldeoxyuric acids** (BILTZ), A., i, 380; (BILTZ and BÜLOW), A., i, 381, 382, 383, 384; (BILTZ and HEIDRICH), A., i, 382, 383.
- Diallylanthranilic acid** (FÖLDI), A., i, 732.
- Diallylbenzidine**, 3:3':5:5'-*tetra*nitro- (VAN ROMBURGH), A., i, 276.
- NN'-Di(*p*-allyloxyphenyl)acetamidine** (SCHULER), A., i, 179.
- Dialyser**, rapid (GUTBIER, HUBER, and SCHIEBER), A., ii, 551.
- simple (GUTBIER and MAYER), A., ii, 353.
- Diamminodinitro-oxalatocobaltates**, crystallography and optical properties of (KNAGGS), T., 2069.
- Diamond**, energy of atomic linkings in (FAJANS), A., ii, 818.
- forces between atoms in, and in aliphatic hydrocarbons (FAJANS), A., ii, 194.
- action of, on carbon monoxide (FOIX), A., ii, 641.
- Diamyl selenide**, $\beta\beta'$ -*dichloro*-, and its dichloride (BOORD and COPE), A., i, 421.
- Diamylose**, constitution of (KARRER and SMIRNOV), A., i, 435.
- Diisoamyloxydiphenylmethane** (MACKENZIE), T., 1696.
- 2:7-Dianilinonaphthylavinduline** (DUTT), T., 1953.
- 2:7-Dianilinophenanthranaphthazine**, and its nitro- derivatives (SIRCAR and DUTT), T., 1948.
- 3:5-Dianilino-1:2:4-triazole**, and its dibenzoyl derivative (FROMM, KAYSER, BRIEGLEB, and FOHRENBACH), A., i, 378.
- Dianilino vinyl ketone**, and its derivatives (BORSCHKE and BONACKER), A., i, 50.
- o*-Dianisidine hydroferrocyanide** (CUMMING), T., 1296.
- Di-*p'*-anisyl-*p*-benzoquinone** (PUMMERER and PRELL), A., i, 1165.
- Di-*p*-anisyl-($\beta\beta$ -diphenylvinyl)methyl perchlorate** (ZIEGLER, OCHS, BREMER, and THIEL), A., i, 1049.
- NN'-Di-*p*-anisylhydrazine**, and its derivatives (WIELAND, WECKER, and ALBERT), A., i, 780.
- 2:5-Di-*p*-anisylquinol** (PUMMERER and PRELL), A., i, 1165.

- Di-*p*-anisylstyryl carbinol**, and its derivatives (ZIEGLER and OCHS), A., i, 151, 1048.
- Dianthraquinone**, 2:2'-*d*ihydroxy- (BRADSHAW and PERKIN), T., 921.
- Dianthraquinonylcarbamides**, isomeric (BATTEGAY and BERNHARDT), A., i, 1041.
- Dianthrone**, 3:8'-*d*ihydroxy-, and its derivatives (PERKIN and WHATTAM), T., 294.
- Dianthryl**, 2:2'-*d*ihydroxy-, and its derivatives (PERKIN and WHATTAM), T., 292.
- Diantipyril selenide**, *di-p* nitro- (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 1066.
- Diaquotetrammincobaltiselenate**. See under Cobalt.
- Diaquotetrapyridinemagnesium bromide** (SPACU), A., i, 859.
- Diarylatyrylcarbinols**, halochromy of (ZIEGLER), A., i, 151.
- Diatase**, inactivation of, by metals (LUGER), A., i, 65.
regeneration of (BIEDERMANN), A., i, 698.
action of pepsin and trypsin on (BIEDERMANN), A., i, 480.
malt, thermostability of (EUNSTROM), A., i, 599.
- 2-Diazoamino-3-bromo-*p*-cymene** (WHEELER and SMITHEY), A., i, 333.
- Diazobenzene(4-arsinic acid)-4'-aminoacetanilide**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-aminoacetophenone**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-amino-6'-acetophenoxyacetic acid**, and its sodium salt and methyl ester (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-3'-aminoanisic acid**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)aminobenzoic acids**, and their sodium salts (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-amino-cinnamic acid**, and its salts and ethyl ester (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-amino-2'-5'-dimethylphenoxyacetic acid**, sodium salt and methyl ester (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene-4-arsinic acid-3'-amino-6'-methoxybenzoic acid**, methyl ester, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-amino-2'-methylphenoxyacetic acid**, and its methyl ester and sodium salts (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)amino-methylisopropylphenoxyacetic acids**, sodium salts and methyl esters (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-4'-aminophenol**, and its sodium salt and benzoic ester (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-aminophenoxyacetic acid**, and its salts and derivatives (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-4'-aminophenylarsinic acid**, and its salts (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-6'-aminopiperonylic acid**, methyl ester, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)aniline**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)anisidines**, and their sodium salts (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)benzylglycine** (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-6-bromo-4-amino-2-methylphenoxyacetic acid**, and its sodium salt and methyl ester (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-8'-bromo-4'-aminophenoxyacetic acid**, and its sodium salt and methyl ester (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-4-chloroaniline**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)diethylamine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)dimethylamine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-4'-ethoxyphenylglycine**, and its sodium salts and ethyl ester (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-4'-methoxyphenylglycine**, and its sodium salts and ethyl ester (JACOBS and HEIDELBERGER), A., i, 74.

- Diazobenzene(4-arsinic acid)-3'-methyl-4'-aminophenoxyacetic acid**, and its methyl ester and sodium salts (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)-4'-methyl-aminophenoxyacetic acid** (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)methyl-aniline**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- bis***Diazobenzene(4-arsinic acid)pentamethylenetetramine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)phenylglycine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 74.
- Diazobenzene(4-arsinic acid)piperidine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-p-toluidine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 73.
- Diazobenzene(4-arsinic acid)-p-tolylglycine**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 74.
- Diazo-2-bromobenzene(4-arsinic acid)-p-aminophenoxyacetic acid**, and its methyl ester and sodium salts (JACOBS and HEIDELBERGER), A., i, 74.
- Diazo-compounds** (ANGELI, A., i, 774 ; (PIEBONI), A., i, 1071.
- aliphatic (STAUDINGER and LÜSCHER), A., i, 237 ; (STAUDINGER), A., i, 238, 240.
- asymmetry of (LEVENE and MIKESKA), A., i, 818.
- mechanism of reaction of (OLIVERI-MANDALÀ), A., i, 994.
- optically active (CHILES and NOYES), A., i, 924.
- α -Diazoglutaric acid**, esters (CHILES and NOYES), A., i, 924.
- Diazo-groups**, replacement of nitro-groups by (VESELÝ and DVOŘÁK), A., i, 690.
- α -Diazo-*n*-hexole acid**, *d*- and *l*-ethyl esters (CHILES and NOYES), A., i, 925.
- Diazomalonic acid** ethyl and methyl esters, interaction of triphenylphosphine with (STAUDINGER and LÜSCHER), A., i, 238.
- Diazomethane**, action of, on uracil (JOHNSON, HILL, and CASE), A., i, 471.
- action of, on ureides and uric acid (HERZIG), A., i, 373.
- Diazonium salts**, interaction of phenols and (CHATTAWAY and HILL), T., 2756.
- Diazo-reaction** in the carbazole series (MORGAN and READ), T., 2709.
- Diazotisation**, velocity of reaction of (SCHOUTIJSSEN), A., i, 181.
- 6-Diazo-*o*-toluene(3-arsinic acid)-p-aminophenoxyacetic acid**, and its sodium salt and methyl ester (JACOBS and HEIDELBERGER), A., i, 74.
- 5-Dibenzenesulphonylaminoacenesaphthene** (FLEISCHER and SCHRANZ), A., i, 1144.
- Dibenzhydrazide** chloride, action of sodium azide with (STOLLÉ and NETZ), A., i, 690.
- Dibenzofulvene** (*diphenylene-ethylene*), and its derivatives (WIELAND, REINDEL, and FERRER), A., i, 1137.
- and its picrate and polymeride, and 2:7-*di*bromo- (SIEGLITZ and JASSOY), A., i, 820.
- Dibenzothianthren**, 5:7:12:14-*tetra*hydroxy (BRASS and KOHLER), A., i, 1051.
- Dibenzothianthrendiquinone**, preparation of, and its sulphoxide and 1:14-*di*hydroxy- (BRASS and KÖHLER), A., i, 1051.
- Dibenzothiazolylmethane**, and its ethiodide and methiodide (MILLS), T., 464.
- Dibenzoylalizarin**, preparation of (BARNETT and COOK), T., 1389.
- $\alpha\delta$ -Dibenzoylamino- β -ketobutane**, and its semicarbazone (WINDAUS, DORRIES, and JENSEN), A., i, 60.
- $\alpha\delta$ -Dibenzoylamino- γ -ketovaleric acid**, methyl ester, and its phenylhydrazone (WINDAUS, DÖRRIES, and JENSEN), A., i, 61.
- Dibenzoylcytisine**, properties of (WOLF and RIDEAL), A., i, 1114.
- colloidal gels of (GORTNER and HOFFMAN), A., ii, 130.
- Dibenzoyl- β -hydroxyethylguanidine** (FROMM and HONOLD), A., i, 530.
- Dibenzoylmethane** of Wislicenus, true nature of (DUFRAISSE and GÉRALD), A., i, 89.
- selenium compounds of, and chloro- (MORGAN, DREW, and BARKER), T., 2463.
- Dibenzoylmethylethylglyoxime** (PONZIO), A., i, 18.
- Dibenzoylmethylglyoxime** (PONZIO), A., i, 17.
- 1:4- and 1:5-Dibenzoylnaphthalenes** (SCHOLL and NEUMANN), A., i, 261.
- 5:14-Dibenzoyloxydibenzothianthrendiquinone** (BRASS and KOHLER), A., i, 1051.
- 5:9-Dibenzoyloxy-7:7-dimethyl- α -dinaphthaxanthen** (SEN-GUPTA and TUCKER), T., 562.

- 5:9-Dibenzoyloxy-7-methyl 7-ethyl- α -dinaphthaxanthen** (SEN-GUPTA and TUCKER), T., 564.
- 1:12 Dibenzoyloxyperylene** (ZINCKE and DENG), A., i, 1013.
- Dibenzoylphenylmethane** (MEISENHEIMER and WEIBEZAHN), A., i, 178.
- Dibenzoylpyrasoleanthrone-yellow** (MAYER and HEHL), A., i, 878.
- $\alpha\alpha'$ -Dibenzoyl- $\alpha\alpha'\beta\beta'$ -tetraphenyltetra-**zan, and its derivatives (GOLDSCHMIDT and EULER), A., i, 475.
- $\alpha\alpha'$ -Dibenzoyl- $\alpha\alpha'\beta\beta'$ tetra-*p*-tolyltetra-**zan, and its compound with triphenyl methyl (GOLDSCHMIDT and EULER), A., i, 476.
- Dibenzopyrene-5:8 quinone** (SCHOLL and NEUMANN), A., i, 261.
- Dibenzyl**. See *s*-Diphenylethane.
- Dibenzyl ether** (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 934.
- disulphide, di-*o*-hydroxy-** (ZEMPLEN and HOFFMANN), A., i, 564.
- Dibenzylacetohydroxamic acid**, and its salts and derivatives (JONES and SCOTT), A., i, 454.
- Dibenzylacetyl chloride** (JONES and SCOTT), A., i, 454.
- Dibenzylamine**, acetyl derivative (SHORT and SMITH), T., 1806.
- Dibenzylbenzamidine**, and its salts (CURTIUS and EHRHART), A., i, 775.
- Dibenzylcyanamide**, dihydrochloride (SHORT and SMITH), T., 1806.
- 2:3-Dibenzyl-5:6-dihydropyrazine**, $\alpha\alpha'$ -*dicyano*- (DUIT and SEN), T., 2664.
- Dibenzyl diketone**, $\alpha\alpha'$ -*dicyano*-, colouring matters from (DUIT and SEN), T., 2663.
- 9:12-Dibenzylidiphensuccindane** (BRAND and MÜLLER), A., i, 445.
- 1:1'-Dibenzyl-4:4'-dipyridinium** (WEITZ and LUDWIG), A., i, 366.
- hydrogen carbonate (WEITZ and KÖNIG), A., i, 1186.
- Dibenzylidenacetone**. See Distyryl ketone.
- Dibenzylidenecyclobutanone** (DEMJANOV and DOJARENKO), A., i, 1161.
- 9:12-Dibenzylidenediphensuccindane** (BRAND and MÜLLER), A., i, 445.
- 9:12-Dibenzylidene- Δ^{10} -diphensuccindene** (BRAND and MÜLLER), A., i, 445.
- Dibenzylmalonic acid**, di-*o*-nitro-, methyl ester (LEUCHS, v. KATINSKY, and CONRAD), A., i, 472.
- Dibenzylmethylglucoside** (GOMBERG and BUCHLER), A., i, 112.
- 2:3-Dibenzyl-8-methylquinoxaline**, $\alpha\alpha'$ -*dicyano*- (DUIT and SEN), T., 2665.
- 2:3-Dibenzyl-1:4-naphthaquinoxaline**, $\alpha\alpha'$ -*dicyano*- (DUIT and SEN), T., 2665.
- 2:3-Dibenzyl-1:4-naphthaquinoxaline-7-sulphonic acid**, $\alpha\alpha'$ -*dicyano*- (DUIT and SEN), T., 2665.
- 2:3-Dibenzyl-1:4-naphthaquinoxaline-8-sulphonic acid**, $\alpha\alpha'$ -*dicyano*-10-hydroxy- (DUIT and SEN), T., 2666.
- 2:3-Dibenzylquinoxaline**, $\alpha\alpha'$ -*dicyano*-, and $\alpha\alpha'$ -*dicyano*-7-amino- (DUIT and SEN), T., 2664.
- Dibenzylsulphine-*p*-toluenesulphonylimine** (MANN and POPE), T., 1053.
- 1:1'-Dibenzyltetrahydro-4:4'-dipyridyl**, preparation and reactions of (EMMERT and VARENKAMP), A., i, 1065.
- dissociation of (WEITZ), A., i, 470.
- 3:5-Dibenzylthiol-1:2:4-triazole**, and its hydrochloride (FROMM, KATNER, BRIEGLEB, and FOHRENBAUCH), A., i, 379.
- Dibutyl selenide**, $\beta\beta'$ -dichloro-, and its dichloride (BOORD and COPE), A., i, 421.
- Dibutylarsinic acid**, and its copper salt (QUICK and ADAMS), A., i, 600.
- Diisobutylbenzidine**, 3:3':5:5'-*tetranitro*- (VAN ROMBURGH), A., i, 276.
- Diisobutylcyanosarsine** (SPINKOFF, DONAT, and JARGER), A., i, 995.
- Diisobutyl ketone** semicarbazone (LEROIIDE), A., i, 217.
- $\alpha\alpha'$ -Dibutyrylethane**, and its copper derivative (LEROIIDE), A., i, 217.
- Dibutyrylmethane**, and its copper derivative (LEROIIDE), A., i, 217.
- $\beta\beta'$ -Dibutyrylpropane**, and its disemicarbazone (LEROIIDE), A., i, 217.
- Dicamphomethylchloroamine** (RUPE and SCHMID), A., i, 1042.
- Dicamphomethylhydroxylamine**, and its salts (RUPE and SCHMID), A., i, 1041.
- $\alpha\alpha'$ -Dicarbamyl- $\alpha\alpha'$ -dimethylcyclohexane-1:1-diacetic acid**, ω -imide of (KON and THORPE), T., 1803.
- α -*NN'*-Dicarbethoxyhydrazinoacetoacetic acid**, ethyl ester (DIELS and FULDNER), A., i, 1196.
- β -Dicarbethoxyhydrazinoacetylacetone** (DIELS and FULDNER), A., i, 1196.
- γ -*NN'*-Dicarbethoxyhydrazinoacetylacetone** (DIELS), A., i, 774.
- 10- $\alpha\beta$ -Dicarbethoxyhydrazinoanthracene**, 9-amino-, and its derivatives (DIELS and MÖHL), A., i, 1194.
- 2- and 4- $\alpha\beta$ -Dicarbethoxyhydrazino-naphthalenes**, 1-amino- (DIELS and SÖRENSEN), A., i, 1194.
- 2- $\alpha\beta$ -Dicarbethoxyhydrazino-naphthalene-4-sulphonic acid**, sodium salt (DIELS and SÖRENSEN), A., i, 1194.
- 6- $\alpha\beta$ -Dicarbethoxyhydrazino-*p*-xylene**, 3-amino-, and its derivatives (DIELS and ECKELMANN), A., i, 1195.

- α -*NN'*-Dicarbomethoxyhydrazinoacetoacetic acid, ethyl ester (DIELS and FULDNER), A., i, 1196.
- β -Dicarbomethoxyhydrazinoacetylacetone (DIELS and FULDNER), A., i, 1196.
- α -*NN'*-Dicarbomethoxyhydrazinoacetononic acid, β -amino-, ethyl ester (DIELS and FULDNER), A., i, 1196.
- 1- $\alpha\beta$ -Dicarbomethoxyhydrazinonaphthalene, 2-amino-, and its derivatives (DIELS and WACKERMANN), A., i, 1196.
- 4- $\alpha\beta$ -Dicarbomethoxyhydrazinonaphthalene, 1-amino- (DIELS and KLEINFELLER), A., i, 1195.
- 2:4-Dicarboxycyclobutane-1:3-diacetic acids, and the α -anhydride (INGOLD, PERREN, and THORPE), T., 1789.
- cis*- and *trans*-2:4-Dicarboxycyclobutane 1:3-di- α -propionic acids, and 2:4-dicyano-, ethyl ester (INGOLD, PERREN, and THORPE), T., 1787.
- 4- $\alpha\beta$ -Dicarboxyethylamidohydrazinonaphthalene, 1-amino- (DIELS and SØRENSEN), A., i, 1195.
- $\alpha\gamma$ -Dicarboxyglutaconic acid, γ -bromo-, ethyl ester, transformations of (FALTIS, and DE ROXAS), A., i, 624.
- $\alpha\gamma$ -Dicarboxyglutaconolactone, γ -hydroxy-, triethyl ester (FALTIS and DE ROXAS), A., i, 625.
- $\omega'\omega''$ -Dicarboxymethanetriacetic acid, ω -cyano-, ethyl ester (INGOLD and PERREN), T., 1418.
- $\alpha\beta$ -Dicarboxymethylamidohydrazinonaphthalenes, amino- (DIELS and SØRENSEN), A., i, 1195.
- Dichromates. See under Chromium.
- Dicumylamine, chloroamino- (GOLDSCHMIDT and STROHMENGER), A., i, 1005.
- Dicyanamide, and its salts (MADELUNG and KERN), A., i, 438.
- Dicyanimide (FRANKLIN), A., i, 440.
- Dicyanodiamide, action of aqueous ammonia on (DAVIS), A., i, 118.
formation of guanidine from (BLAIR and BRAHAM), A., i, 1123.
action of, with methyl- and dimethylammonium chlorides (WERNER and BELL), T., 1790.
- Di-*p*-dimethylaminostyryl ketone, α -chloro- (BAUER and WERNER), A., i, 1035.
- Didymium, absorption spectrum of (WEIGERT), A., ii, 413.
- Dieckmann reaction (KON), T., 513.
- Dieckmann-Komppa reaction (DICKENS, KON, and THORPE), T., 1496.
- Dielectric constants, measurement of (KING and PATRICK), A., ii, 109.
of colloidal solutions (ERREKA), A., ii, 694.
- Diet, effect of, on fixation of nitrogen (DIENES), A., i, 293.
replacement of protein by urea in (MORGEN, SCHÖLER, WINDHEUSER, and OHLMER), A., i, 293.
- γ -Diethoxyacetoacetic acid, ethyl ester, use of, in the synthesis of glyoxaline derivatives (PUCHER and JOHNSON), A., i, 549.
- 2:9-Diethoxyacridine (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 469.
- 4:4'-Diethoxybenzil, 3:3'-*d*-nitro- (SCHÖNBERG and KRAEMER), A., i, 664.
- Diethoxy-1:1'-diethylcarbocyanine iodides (BRAUNHOLTZ), T., 171.
- 4:4'-Diethoxy-1:1'-dinaphthyl (SCHOLL and SEER), A., i, 336.
- 4:4'-Diethoxy-1:1'-di-*ar*-tetrahydronaphthyl (SCHROETER, KINDERMANN, DIEFRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 124.
- $\alpha\epsilon$ -Diethoxy- β -methylpentan- β -ol- δ -one (DAUDEL), A., i, 432.
- Di-*p*-ethoxyphenylformazyl methyl ketone (JACOBSEN), A., i, 596.
- 2:3-Di-*p*-ethoxyphenyl- $\alpha\beta$ -naphthaquinoxaline, and 2:3-di-*m*-nitro- (SCHÖNBERG and KRAEMER), A., i, 664.
- Diethyl ether. See Ethyl ether.
sulphide, $\beta\beta'$ -dichloro-, preparation and reactions of (MANN and POPE), T., 594.
hydrolysis of, and preparation of a non-vesicant isomeride (BALES and NICKELSON), T., 2137.
reaction of, with sodium silicate, and the treatment of burns produced by it (KRAMER), A., i, 914.
- d*-hydroxy-, preparation of esters of (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 420.
sulphoxides, $\alpha\beta\beta'$ -tri- and $\alpha\beta\beta\beta'$ -tetra- chloro- (MANN and POPE), T., 603.
- Diethylacetal, $\beta'\beta''$ -dichloro- (GRIGNARD and PURDY), A., i, 803.
- Diethylacetylallophanic acid, bromo-, ethyl ester (CALISEN), A., i, 1128.
- Diethylallylamine, and its oxide, and their picrates (MEISENHEIMER and WILLMERSDORF), A., i, 334.
- 5-Diethylaminocacenaphthene, and its derivatives (FLEISCHER and SCHRANZ), A., i, 1143.
- p*-Diethylaminobenzylideneanthranilic acid (EKELEY, ROGERS, and SWISHER), A., i, 935.
- β -Diethylaminobutan- γ -ol (FOURNEAU and PUYAL), A., i, 639.

- Diethylaminobutylacetoacetic acid**, ethyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- 2-Diethylamino-2-ethoxyacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 469.
- α -Diethylaminoethylacetoacetic acid**, ethyl and methyl esters (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- Diethylaminoethylacetonedicarboxylic acid**, ethyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- Diethylaminoethylacetylacetone** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- α -Diethylaminoethylbutyric acid**, β -hydroxy-, ethyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 640.
- Diethylaminoethylcamphorcarboxylic acid**, methyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- Diethylaminoethylcyanoacetic acid**, ethyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- Diethylaminoethylmalonic acid**, ethyl ester (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- α -Diethylaminopropan- β -ol** (FOURNEAU and PUYAL), A., i, 639.
- ϵ -Diethylamino- $\beta\beta\beta$ -trimethylheptan- α -one** (BILLON), A., i, 717.
- Diethylammonium ruthenipentabromide** (GUTHRIE and KRAUSS), A., i, 16.
- Diethylaniline**, compound of 3:5-dinitrobenzoic acid and (LEY and GRAU), A., i, 536.
- 1:1'-Diethylazurine iodide**, bromo-derivatives (MOUDGILL), T., 1510.
- Diethylbarbituric acid**, action of diazomethane on (HERZIG), A., i, 873.
- Diethylbarbituric acid**, calcium and magnesium salts (WÜLFING), A., i, 1066.
- Diethylbarbituric acid**, compound of 4-dimethylamino-1-phenyl-2:3-dimethyl-5-pyrazolone and (CHEMISCHE FABRIK AUF AKTIEN VORM. E. SCHERING), A., i, 582.
- Diethylbenzidine**, 3:3':5:5'-tetranitro- (VAN ROMBURGH), A., i, 275.
- Diethylbromocetylamine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- 1:1-Diethylcyclobutane-3-one-2-carboxylic acid**, methyl ester, semicarbazone (DICKENS, KON, and THORPE), T., 1504.
- Diethylcarbodi-imide** (STAUDINGER and HAUSER), A., i, 70.
- 2:2'-Diethylcarbothiocyanine iodide** (MILLS and BRAUNHOLTZ), T., 1493.
- Diethylcarbothiocyanine iodide**, crystallography of (MILLS), T., 462.
- Diethylchlorobutylamine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- Diethylchloroethylamine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- 7:9-Diethyldeoxyuric acid**, and its salts (BILTZ and BULOW), A., i, 384.
- α -N.O-Diethyldihydrocupreicimol** (HEIDELBERGER and JACOBS), A., i, 674.
- 7:7-Diethyl- α -dinaphthazanthene**, oxidation of, and 5:9-dihydroxy- (SENGUPTA and TUCKER), T., 565.
- Diethylene disulphide tetraiodide**, as an antiseptic (BACHEM), A., i, 611.
- Diethylenediaminecobaltic bromides**, conductivities of (DUFF), T., 450.
- $\alpha\alpha$ -Diethylethylenediamine** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 255.
- $\beta\beta$ -Diethylglutaric acid**, methyl ester (DICKENS, KON, and THORPE), T., 1504.
- $\beta\beta$ -Diethylglutaric acid**, $\alpha\alpha'$ -dibromo-, and its ethyl ester (DESHAPANDE and THORPE), T., 1436.
- gem*-Diethyl groups**, effect of, on the carbon tetrahedral angle (DESHAPANDE and THORPE), T., 1430.
- 5:5-Diethylhydantoin** (INGOLD, SAKO, and THORPE), T., 1192.
- Diethyliminovioluric acid**, and its salts (LIFSCHITZ and HEPNER), A., i, 768.
- 2:2-Diethylindan-1-one** (HALLER and BAUER), A., i, 259.
- Diethylmalonbromamide** (INGOLD, SAKO, and THORPE), T., 1192.
- Diethylmalonic acid**, and its ethyl hydrogen ester, esterification of (DUMESNIL), A., i, 622.
- Diethylmalonic acid**, ethyl esters, hydrolysis of (DUMESNIL), A., i, 520.
- Diethylmalonic acid**, derivatives of (DUMESNIL), A., i, 807.
- Diethylmalonodi-5-acenaphthylamide** (FLEISCHER and SCHRANZ), A., i, 1143.
- Diethylnitrosoamine** (SCHMIDT and FISCHER), A., i, 822.
- $\gamma\gamma$ -Diethyloctane- $\gamma\gamma$ -diol**, and its oxide (CARRIÈRE), A., i, 320.
- Diethylcyclopentamethylenearsonium iodide** (STEINKOPF, DONAT, and JAEGER), A., i, 996.
- 1:1-Diethylcyclopentane-3:4-dione**, and its disemicarbazone (DICKENS, KON, and THORPE), T., 1506.

- 7:8-Diethylphenalene** (MAYER and SIEGLITZ), A., i, 741.
- Diethylpiperazine** dichloroethylate (FARBWERKEFORM. MEISTER, LUCIUS, & BRUNING), A., i, 529.
- 1:1-Diethylcyclopropane-2-carboxylic acid**, 2:3-dicyano-, and its amide (BIRCH and THORPE), T., 1825.
- 1:1-Diethylcyclopropane 2:3-dicarboxylic acid**, and its dinitrile (BIRCH and THORPE), T., 1826.
- 1:1-Diethylcyclopropane-2:3-dicarboxylic acid**, 2:3-dicyano-, imide (BIRCH and THORPE), T., 1825.
- cis*- and *trans*-**3:3-Diethylcyclopropane-1:2-dicarboxylic acids**, and their derivatives (DESHAPANDE and THORPE), T., 1437.
- 3:3-Diethylcyclopropan-1-ol-1:2-dicarboxylic acid**, and its derivatives (DESHAPANDE and THORPE), T., 1440.
- 2:2-Diethylresorcinol** 3-ethyl ether, 4-chloro- (FABRE), A., i, 1148.
- Diethylrhodamine** (LEONARD), A., i, 1182.
- Diethylsulphine-(bis)-*p*-toluenesulphonylimine** (MANN and POPE), T., 1054.
- Diethylsulphine-*p*-toluenesulphonylimine**, and *ββ'*-dichloro- (MANN and POPE), T., 1053.
- 2:2'-Diethylthiocyanine iodide** (MILLS), T., 461.
- 2:1'-Diethylthioisocyanine** iodide (BRAUNHOLTZ and MILLS), T., 2006.
- 7:9-Diethyl-8-thiouric acids** (BILTZ and BÜLOW), A., i, 834.
- 4:6-Diethyl-*m*-xylene** (PHILIPPI, SEKA, and FROESCHL), A., i, 837.
- 2:5-Diethyl-*p*-xylene**, 3:6-dibromo-, and 3:6-dinitro- (PHILIPPI, SEKA, and ROBINSON), A., i, 837.
- Diffusion**, effect of temperature on (ÖHOLM), A., ii, 621.
of electrolytes into gels (STILES and ADAIR; STILES), A., ii, 125.
in porous vessels (HERREIRA), A., ii, 427.
formation of Liesegang's rings in (BRADFORD), A., ii, 358.
and intertraction (SCHONEBOOM), A., ii, 690.
- Difluorenylene**, and its picrate, and *di*-hydroxy-, and *dinitro*- (DZIEWOŃSKI and SUSZKO), A., i, 730.
- 1:3-Difurfurylidene-cyclohexan 2-one** (WOLFF), A., i, 668.
- Digalloylævoglucozan**, and its potassium salt (KARRER and SALOMON), A., i, 267.
- Digallylamine**, and its salts (ROSEN-MUND and PFANNKUCH), A., i, 1080.
- Digestion**, disappearance of nitrogen in (GOUIN and ANDOUARD), A., i, 698.
of proteins by trypsin (NORTHERP), A., i, 693.
effect of splenectomy on (MOLLOW), A., i, 397.
peptic, rôle of acids in (OSTWALD and KUHN), A., i, 598.
- Digestive tract**, activities of the secretions of (HARAMAKI), A., i, 698.
- Digitalis**, assay of the active principles of (KNUDSON and DRESBACH), A., ii, 882.
toxicity of extracts of (v. DAFERT), A., i, 97.
- Digitalis purpurea**, extraction of digitoxin from (BINAGHI), A., i, 412.
- Digitalonic acid**, constitution of (KILIANI), A., i, 224.
- Digitan**, estimation of digitoxin in (WARREN), A., ii, 403.
- Digitic acid**, dimethyl ester (WINDAUS and WEIL), A., i, 849.
- β*-Digitogenic acid**, dimethyl ester (WINDAUS and WEIL), A., i, 849.
- Digitolic acid**, dimethyl ester (WINDAUS and WEIL), A., i, 849.
- Digitonin**, constitution and derivatives of (WINDAUS and WEIL), A., i, 848.
- Digitoxin**, estimation of, in digitan (WARREN), A., ii, 403.
- Digitoxonic acid**, constitution of (KILIANI), A., i, 224.
- Digitoxose**, constitution of, and its carboxylic acid (KILIANI), A., i, 224.
- Diglucon**, constitution of (KARRER and SMIRNOV), A., i, 229.
- Diglycolamidic acid**, di-*p*-anisylamide, di-*p*-chlorophenylamide, and di-*m*-tolylamide (HILL and KELSEY), A., i, 1141.
- Diglycoll-1:2:4:5-*ψ*-cumidic acid** (ANSCHÜTZ and JAEGER), A., i, 428.
- Diglycollic acid**, and its derivatives (ANSCHÜTZ and JAEGER), A., i, 428.
- Diglycollnaphthalidic acids** (ANSCHÜTZ and JAEGER), A., i, 428.
- Diglycollnaphthils** (ANSCHÜTZ and JAEGER), A., i, 428.
- Diglycolltoluidic acids** (ANSCHÜTZ and JAEGER), A., i, 428.
- Diglycollxylic acids** (ANSCHÜTZ and JAEGER), A., i, 428.
- Diglycollxylls** (ANSCHÜTZ and JAEGER), A., i, 428.
- Di-*cis*- and *trans*-heptadecylenylcarbamides** (NICOLET and PELO), A., i, 644.
- Di-*n*-hexaldehyde**, *di*-*β*-hydroxy-. See Bis-6-methyltetrahydro-2-*α*-pyryl ether.
- Disixhexoylpentamethylenediamine**, *di-α*-bromo- (BERGELL), A., i, 720.

- 4:4'-Dihydrazino-1:1'-di-*ar*-tetrahydronaphthyl** (SCHROETER, KINDERMANN, DIETRICH, BEYCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 124.
- Dihydrocabinol** (RUZICKA and MEYER), A., i, 830.
- 5:8-Dihydroaceto- α -naphthalide**, 4-bromo- (ROWE and DAVIES), T., 1006.
- Dihydrobenzofurans**, mercury compounds of (ADAMS, ROMAN, and SPERRY), A., i, 946.
- 7:12-Dihydro-*p*-benzophenarsazine**, oxide and sulphide, and 7-bromo-, 7-chloro-, and 7-iodo- (LEWIS and HAMILTON), A., i, 187.
- dl*-Dihydrocampholenitrile** (KLEUCKER), A., i, 736.
- Dihydro- α -campholytic acid**, γ -*di*-hydroxy-, and its diacetyl derivative (CHANDRAMENA, INGOLD, and THORPE), T., 1548.
- Dihydrocarbostyrl-3:3'-spiran**, *di*-bromo-, *dichloro*-, and *dinitro*-derivatives (LEUCHS, CONRAD, and v. KATINSZKY), A., i, 873.
- d*- and *l*-Dihydrocinchonine** (HEIDELBERGER and JACOBS), A., i, 674.
- Dihydrocinchonine**, chloro-, and its hydrochloride (JACOBS and HEIDELBERGER), A., i, 672.
- Dihydrocinidic acid** (MURAYAMA), A., i, 310.
- Dihydrocodeine**, *a*-nitro- (SPEYER and WIETERS), A., i, 169.
- Dihydrocodeine**, and its salts (SPEYER and WIETERS), A., i, 47.
- Dihydrocodeine-*N*-oxide**, and its salts and sulphonic acid (SPEYER and WIETERS), A., i, 169.
- Dihydrocodeinesulphonic acids** (SPEYER and WIETERS), A., i, 169.
- Dihydrocodide-*N*-oxidesulphonic acid**, chloro- (SPEYER and WIETERS), A., i, 169.
- Dihydrocodidesulphonic acid**, chloro- (SPEYER and WIETERS), A., i, 170.
- Dihydrocupreicine** hydrobromide (HEIDELBERGER and JACOBS), A., i, 673.
- Dihydrocupreine oxide** (SPEYER and BROKER), A., i, 675.
- Dihydrodehydrocorydaline** (GADAMER and v. BRUCHHAUSEN), A., i, 675.
- Dihydrodemethylscopolines**, and their salts and derivatives (HESS and WAHL), A., i, 856.
- Dihydrodeoxycinchonine**, and its hydrochloride (JACOBS and HEIDELBERGER), A., i, 672.
- Dihydrofencholamine**, and its salts and derivatives (MAXWELL), A., i, 753.
- Dihydroisindole**, preparation of (v. BRAUN and NELKEN), A., i, 863.
- Dihydromethystic acids** (GOEBEL), A., i, 657.
- Dihydromethysticin** (GOEBEL), A., i, 657.
- Dihydromethysticoles**, and their semicarbazones (GOEBEL), A., i, 657.
- Dihydromorphine-*N*-oxide** (SPEYER and BECKER), A., i, 676.
- Dihydromorphinesulphonic acids** (SPEYER and BECKER), A., i, 676.
- Δ^{α} -Dihydromuconic acid**, δ -bromo- γ -hydroxy-, and γ -chloro- δ -bromo-, methyl ester (CHANDRASENA and INGOLD), T., 1818.
- Dihydronaphthalenealkamine**. See Tetrahydro- β -naphthylamine, β -hydroxy-.
- Dihydronaphthalene-2-imino-oxazolidine**, and its salts (TAKEDA and KURODA), A., i, 275.
- Dihydronaphthalene series** (ROWE and DAVIES), T., 1000.
- 5:8-Dihydro- α -naphthylamine**, oxidation and bromination of (ROWE and DAVIES), T., 1000.
- Dihydronaphthylethanol**, and its phenylurethane (DE POMMEREAU), A., i, 828.
- 7:8-Dihydrophenalene-9**, and its oxime (MAYER and SIEGLITZ), A., i, 741.
- 9:16-Dihydrophenanthranaphthazine**, 2:7-*di*-amino-, and 2:7-*di*-hydroxy- (SIRCAR and DUTT), T., 1950.
- d*- and *l*-Dihydroquinine**, and their salts (HEIDELBERGER and JACOBS), A., i, 674.
- Dihydroquinine ethyl bromide** (HEIDELBERGER and JACOBS), A., i, 673.
- Dihydroquinine**, 5:8-*di*-amino-, 8-amino-5-hydroxy-, and 5:8-*di*-hydroxy-, and their salts (JACOBS and HEIDELBERGER), A., i, 671.
- Dihydroquinine oxide**, preparation of, and its dinitrate (SPEYER and BECKER), A., i, 675.
- Dihydrrubescene**, and its picrate (DZIEWONSKI and SUSZKO), A., i, 730.
- 4:5-Dihydrothiazole**, 2-thiol-, compound of chloropicrin and (RAY and DAS), T., 325.
- 1:4-Di(5'-hydroxy-7'-sulpho-2'-naphthylamino)benzene** (KALLÉ & Co.), A., i, 824.
- Di*-isatinazine**, and its acetyl derivative (BORSCHKE and MEYER), A., i, 54.
- 5:5'-*Di*-isatylmethane** (BORSCHKE and MEYER), A., i, 53.
- 3:5-Diketo-1-benzyl-1:4-hexahydro-diazine** (DUBSKY and DINGEMANSE), A., i, 57.

- Diketocholeonic acid**, and its ethyl ester, and its dioxime (BORSCHKE and HALLWASS), A., i, 1158.
- 2:5-Diketo-1:4-dibenzoylmethylpiperazine** (CHERBULIEZ and FEER), A., i, 1065.
- 2:5-Diketo-1:4-dibenzoylpiperazine**, preparation of (SASAKI and HASHIMOTO), A., i, 56.
- 2:5-Diketo-3:6-dibenzylidene-1-methylpiperazine**. See Xanthorocellin.
- 2:6-Diketo-3:5-dibenzylidene-2:3:5:6-tetrahydro-m- α -benzobispyrrole** (DAVIES and HICKOX), T., 2653.
- 2:5-Diketo-1:4-dibenzylpiperazine**, preparation of (SASAKI and HASHIMOTO), A., i, 56.
- 2:5-Diketo-1:4-dichloromethylpiperazine** (CHERBULIEZ and FEER), A., i, 1065.
- 2:2'-Diketo- $\Delta^{1,4}$ -dicoumaran** (*oxindigo*), synthesis of (FRIES and HASENJÄGER), A., i, 46.
- 2:5-Diketo-1:4-diethoxymethylpiperazine** (CHERBULIEZ and FEER), A., i, 1066.
- 5:6-Diketo-7:7-diethyl-5:6-dihydro- α -dinaphthaxanthen**, and its derivatives (SEN-GUPTA and TUCKER), T., 565.
- 2:5-Diketo-1:4-dihydroxymethylpiperazine** (CHERBULIEZ and FEER), A., i, 1065.
- 2:5-Diketo-1:4-dimethoxymethyl-2:5-piperazine** (CHERBULIEZ and FEER), A., i, 1065.
- 5:6-Diketo-7:7-dimethyl-5:6-dihydro- α -dinaphthaxanthen**, and its derivatives (SEN-GUPTA and TUCKER), T., 569.
- 2:5-Diketo-1:4-dinaphthylmethylpiperazine** (CHERBULIEZ and FEER), A., i, 1066.
- 2:5-Diketo-1:4-dipiperidinomethylpiperazine** (CHERBULIEZ and FEER), A., i, 1065.
- 2:2'-Diketo-6-methyl- $\Delta^{1,4}$ -dicoumaran** (FRIES and HASENJÄGER), A., i, 46.
- 5:6-Diketo-7-methyl-7-ethyl-5:6-dihydro- α -dinaphthaxanthen**, and its derivatives (SEN-GUPTA and TUCKER), T., 563.
- 5:6-Diketo-7-methyl-7-n-propyl-5:6-dihydro- α -dinaphthaxanthen** (SEN-GUPTA and TUCKER), T., 566.
- 1:4-Diketo-2-methyltetrahydronaphthalene**, 2:3-dibromo-, and 2:3-dichloro- (FRIES and LOHMANN), A., i, 30.
- Diketone**, $C_{22}H_{14}O_6$, from the methyl ether of the substance $C_{22}H_{12}O_4$ (FRIES and LEUE), A., i, 462.
- $C_{26}H_{40}O_4$ and its dioxime, from the oxidation product of cholestan-4:7-dione (WINDAUS and GRIMMER), A., i, 541.
- 1:2-Diketones** (SCHÖNBERG), A., i, 27; (SCHÖNBERG and KRAEMER), A., i, 663.
- β -Diketones**, action of carbon disulphide on thallous compounds of (MORGAN and LEDBURY), T., 2892.
- interaction of selenium tetrachloride and (MORGAN, DREW, and BARKER), T., 2433.
- $\alpha\epsilon$ -Diketo- $\alpha\beta\gamma\delta\epsilon$ -pentaphenyl- Δ^8 -pentene** (DILTHEY and KAFFER), A., i, 669.
- 2:4-Diketo-5-phenyl-5-ethyltetrahydro-oxazole**, preparation of (ALTWEGG and EBIN), A., i, 53.
- 2:4-Diketo-5-phenyl-5-methyltetrahydro-oxazole** (ALTWEGG and EBIN), A., i, 53.
- Diketopiperazine**, molecular compounds of phenols and (POVARNIN and TICHOMIROV), A., i, 1185.
- 2:5-Diketopiperazine**, formaldehyde derivatives of (CHERBULIEZ and FEER), A., i, 1065.
- Diketopiperazines** (DUBSKY and DINGEMANSE), A., i, 56; (DUBSKY and HOHER; DUBSKY, HOHER, and DINGEMANSE), A., i, 57.
- 2:5-Diketopiperazine-1:4-diacetodi-naphthalide** (DUBSKY, HOHER, and DINGEMANSE), A., i, 57.
- θ -Diketostearic acid**, fusion of, with potassium hydroxide, and its ethyl and methyl esters (NICOLET and JURIST), A., i, 622.
- Dimatridine**, and its salts and derivatives (KONDŌ, KISHI, and ARAKI), A., i, 269.
- Dimercuridiaminoazobenzene diacetate** (VECCHIOTTI), A., i, 478.
- Dimercuri-4-hydroxybenzaldehyde**, 3:5-dichloro-, and 3:5-dihydroxy- (HENRY and SHARP), T., 1058.
- Dimercurialcylaldehyde**, 3:5-dichloro- and 3:5-dihydroxy- (HENRY and SHARP), T., 1057.
- 1:3-Dimethindiazidine**, synthesis and division of derivatives of (INGOLD and PIGGOTT), T., 2793.
- 2:4-Dimethoxyacetophenone-3-hydroxy-**, and its derivatives (BRAND and COLLISCHONN), A., i, 452.
- 4:4'-Dimethoxyarsenobenzene**, 3:3'-diamino-, dihydrochloride (CHRISTIANSEN), A., i, 1203.
- 3:4-Dimethoxybenzaldehyde**, 5-hydroxy- (SPÄTH and RÖDER), A., i, 853.
- 2:2'-Dimethoxybenzhydrol** (ZIEGLER and OCHS), A., i, 1049.
- 2:2'-Dimethoxybenzyl**, and its derivatives (SCHÖNBERG and KRAEMER), A., i, 664.

- 3:4-Dimethoxybenzylideneanthranilic acid (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 9:3'-4'-Dimethoxybenzylidene-fluorene-2'-carboxylic acid, 2:7-dibromo-, and its derivatives (SIEGLITZ and JASOY), A., i, 821.
- Dimethoxy-carboxybenzene-azobenzene-4'-arsinic acids, amino-, and their salts (JACOBS and HEIDELBERGER), A., i, 75.
- 2:2'-Dimethoxy-5:5'-dimethylbenzil, and its osazone (SCHÖNBERG and KRAEMER), A., i, 663.
- 5:9-Dimethoxy-7:7-dimethyl- α -dinaphthaxanthen (SEN-GUPTA and TUCKER), T., 562.
- 6:7-Dimethoxy-2:3-dimethylquinoline, and its salts (RILLIET), A., i, 840.
- 4:4'-Dimethoxy-1:1'-dinaphthyl (SCHOLL and SEER), A., i, 336.
- 4:4'-Dimethoxydi-*m*-tolyl disulphide (STEWART), T., 2558.
- 3':4'-Dimethoxy-2-hydroxydistyryl ketone (BUCK and HEILBRON), T., 1099.
- Dimethoxy-*p*-menthane, dibromo- (SCHMIDT, BARTHOLOMÉ, and LUBKE), A., i, 827.
- 5:9-Dimethoxy-7-methyl-7-ethyl- α -dinaphthaxanthen (SEN-GUPTA and TUCKER), T., 564.
- 6:7-Dimethoxy-2-methylquinazoline, and its picrate (RILLIET), A., i, 840.
- 4:6-Dimethoxyphenylarsinic acid, 3-nitro- (CHRISTIANSEN), A., i, 1203.
- 2:4-Dimethoxyphenyl bromomethyl ketone (SONN and FALKENHEIM), A., i, 1163.
- 3:4-Dimethoxyphenyl α -chloro-4'-dimethylanilomostyryl ketone (BAUER and WERNER), A., i, 1035.
- 3:4-Dimethoxyphenyl α -chlorostyryl ketone (BAUER and WERNER), A., i, 1035.
- 3:4-Dimethoxyphenylethylamine, β -5-hydroxy- (SPATH and RÖDER), A., i, 853.
- 3:5-Dimethoxyphenyl ethyl ketone (MAUTHNER), A., i, 457.
- α -3:4-Dimethoxyphenyl- β -hydroxypropylamine. See Methylisoeugenolalkamine.
- α -3:4-Dimethoxyphenyl- β -hydroxypropyldimethylamine. See Trimethylisoeugenolalkamine.
- α -3:4-Dimethoxyphenylmethyl- β -hydroxypropylamine. See Dimethylisoeugenolalkamine.
- 2:3-Di-*p*-methoxyphenyl- α - β -naphthaxinoxaline (SCHÖNBERG and KRAEMER), A., i, 665.
- 3:5-Dimethoxyphenylpropionamide, β -4-hydroxy- (SPATH and RÖDER), A., i, 852.
- 6:7-Dimethoxy-2-phenylquinazoline, and its salts (RILLIET), A., i, 840.
- 6:7-Dimethoxy-2-phenylquinoline, and its salts (RILLIET), A., i, 840.
- 2:4-Dimethoxyphenyl styryl ketone, 3-hydroxy-, and its dibromide (BRAND and COLLISCHONN), A., i, 452.
- 3:4-Dimethoxyphenyl styryl ketone dichloride (BAUER and WERNER), A., i, 1035.
- 3:4-Dimethoxyphenyl β -veratrylethyl ketone, 2-hydroxy- (CRABTREE and ROBINSON), T., 1038.
- 3:5-Dimethoxy-1-*n*-propylbenzene (MAUTHNER), A., i, 457.
- 6:7-Dimethoxyquinoline, and its salts (RILLIET), A., i, 839.
- 4:4'-Dimethoxystilbene, α -nitro- (MEISENHEIMER and WEIBEZAHN), A., i, 177.
- 3':4'-Dimethoxy-2-styrylbenzopyrylium chloride (BUCK and HEILBRON), T., 1210.
- 5:3'-4'-Dimethoxy-2'-styrylbenzylidene-aminoacene-naphthene (FLEISCHER and SCHRANZ), A., i, 1143.
- 6:8-Dimethoxy-1:2:3:4-tetrahydroisoquinoline, 7-hydroxy- (SPATH and RÖDER), A., i, 852.
- 6:7-Dimethoxy-1:3':4':5'-trimethoxybenzyl-1:2:3:4-tetrahydroisoquinoline, and its derivatives (SPATH and BOHM), A., i, 1175.
- Dimethoxyvinylidihydrophenanthrene (FALTIS and KRAUSZ), A., i, 677.
- Dimethyl ether, *s*-dichloro-, reaction of ethyl malonate with (DOX and YODER), A., i, 429.
- sulphide, *s*-dichloro- (BLOCH and HOHN), A., i, 218.
- telluride dihaloids, physiological action of (COW and DIXON), A., i, 402.
- Dimethylacethydroxamic acid. See isobutyrylhydroxamic acid.
- α -Dimethylaconitic acid (FARMER, INGOLD, and THORPE), T., 149.
- N*-Dimethylalanine, ethyl ester (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.
- 2:4-Dimethyl-5-aldehydopyrrole (FISCHER and ZERWECK), A., i, 758.
- Dimethyl-5-aldehydopyrrole-3-carboxylic acids, derivatives of (FISCHER and ZERWECK), A., i, 758.
- Dimethylallyl bromide. See γ -Methyl- Δ β -butene, α -bromo-.
- Dimethylallylacetacetic acid. See δ -Methyl- $\Delta\gamma$ -hexenoic acid.

- Dimethylallylamine oxide, picrate** (MEISENHEIMER and WILLMERSDORF), A., i, 334.
- 3:7 Dimethyl-allylxanthine** (HOFFMANN-LA ROCHE & Co.), A., i, 585.
- Dimethylamine chloride** (LEVI), A., i, 528.
compound of tellurium tetrabromide with (LOWY and DUNBROOK), A., i, 446.
- 5-Dimethylaminoacenaphthene and its derivatives** (FLEISCHER and SCHRANZ), A., i, 1143.
- 5-Dimethylaminoacetyl-2:4 dimethylpyrrole-3-carboxylic acid, ethyl ester** (FISCHER, SCHNELLER, and ZERWECK), A., i, 1056.
- p-Dimethylaminonitro-6-methyl-2:2'-dicoumarones, isomeric** (FRIES and HASENJÄGER), A., i, 45.
- p-Dimethylaminobenzaldehyde, action of, on urine** (HÄBI), A., i, 88.
- 4-Dimethylaminobenzonitrile, 3-nitro-** (MATTAAR), A., i, 251.
- p-Dimethylaminobenzophenoneanil hydrochloride** (REDELLEN and DANIOF), A., i, 148.
- 2-p Dimethylaminobenzylidene-5-methylpyrrole-4 carboxylic acid, 3-hydroxy-, ethyl ester** (FISCHER and HERMANN), A., i, 1054.
- β-Dimethylaminobutan-γ-ol** (FOURNEAU and PUYAL), A., i, 639.
- trans-p-Dimethylaminocinnamic acid, and α-bromo-, esters of, and their derivatives** (PFEIFFER and HAEFFELIN), A., i, 738.
- 2-Dimethylamino-1-α-dicarbethoxyhydrazinonaphthalene, and its perchlorate** (DIELS and KLEINFELLER), A., i, 1195.
- 1-Dimethylamino-4-α-dicarbomethoxyhydrazinonaphthalene, and its hydrochloride** (DIELS and KLEINFELLER), A., i, 1195.
- 5:6-Dimethylamino-1:3 dimethyluracil, and its hydrochloride** (BILTZ and HEIDRICH), A., i, 384.
- 4-Dimethylaminodiphenyl, and 3-amino-, and 3-mono- and 3:5:2':4'-tetra-nitro-** (GARCÍA BANÚS and FERBER TOMÁS), A., i, 333.
- 4'-Dimethylaminodiphenylamine, 5-chloro-2-nitro-** (JACOBSEN), A., i, 594.
- 4'-Dimethylamino-4-ethoxyazobenzene** (JACOBSEN), A., i, 593.
- 4'-Dimethylamino-5-ethoxydiphenylamine, 2-amino-, and 2-nitro-** (JACOBSEN), A., i, 593.
- 4'-Dimethylamino-4-ethoxyhydrazobenzene, isomerisation of** (JACOBSEN), A., i, 593.
- α-Dimethylaminoethylacetoacetic acid, ethyl ester** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 529.
- 4'-Dimethylamino-trihydroxy-9-phenylfluorones** (MUKERJI), T., 550.
- 7-Dimethylamino-3-hydroxy-4-thiolphenazine** (WATSON and DUTT), T., 1942.
- α-Dimethylamino-β-methylbutan-β-ol** (FOURNEAU and PUYAL), A., i, 639.
- 2-Dimethylamino-6-methylphenazine, methiodide, and 7-amino-, methochloride** (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.
- m-Dimethylaminophenolcamphorcin hydrochloride** (SIRCAR and DUTT), T., 1284.
- 4-Dimethylaminophenyl 2'-nitrophenyl disulphide** (LECHER and SIMON), A., i, 1013.
- 6-Dimethylamino-2-phenylbenzthiazole** (BOGERT and ABRAHAMSON), A., i, 576.
- p-Dimethylaminophenyl-bis(3-acetyl-2:4-dimethylpyrrol)methane, and its ferrichloride** (FISCHER and LUCKMANN), A., i, 276.
- Dimethylaminophenyl-bis-(3-carbethoxydimethylpyrrol)methanes, and their ferrichlorides** (FISCHER and LUCKMANN), A., i, 276.
- 4-Dimethylamino-1-phenyl-2:3-dimethyl-5-pyrazolone, compound of diethylbarbituric acid with** (CHEMISCHE FABRIK AUF AKTIEN VORM. E. SCHERING), A., i, 582.
- 1-p-Dimethylaminophenyl-6-ethoxybenzimidazole, and its derivatives** (JACOBSEN), A., i, 593.
- 4'-Dimethylamino-9-phenylfluorone, 2:4:5:7-tetra-amino-, trichloride** (MUKERJI), T., 550.
- α-Dimethylaminopropan-β-ol** (FOURNEAU and PUYAL), A., i, 639.
- α-Dimethylaminopropionic acid. See N-Dimethylalanine.**
- ω-Dimethylaminopropiophenone, and its hydrochloride and oxime** (MANNICH and HEILNER), A., i, 351.
- α-Dimethylaminopyridine methiodide** (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.
- α-p-Dimethylaminostyryl methylenedioxytyrlic ketone, α-chloro-** (BAUER and WERNER), A., i, 1085.
- 4-Dimethylaminostyryl methyl ketone, α-chloro-, and its derivatives** (BAUER and WERNER), A., i, 1085.
- 2-p-Dimethylaminostyrylpyridine methiodide** (MILLS and POPE), T., 946.

- 1-*p*-Dimethylaminostyrylisoquinoline methiodide (MILLS and SMITH), T., 2733.
- Dimethylaminosulphonic acid, copper and nickel salts (DELEPINE and DEMARS), A., i, 923.
- ε-Dimethylamino-βββ-tetramethylheptan-α-one (BILLON), A., i, 717.
- γ-Dimethylamino-γγε-tetramethyloctan-δ-one (BILLON), A., i, 717.
- p*-Dimethylaminothiobenzophenone (REDELLEN and DANILOF), A., i, 148.
- 7-Dimethylamino-4-thiophenazine, 3-amino- (WATSON and DUTT), T., 1941.
- 9-Dimethylamino-2-thiol-3-phenoxazone (WATSON and DUTT), T., 1942.
- ε-Dimethylamino-βββ-trimethylheptan-α-one (BILLON), A., i, 717.
- 5:6-Dimethyldiaminouracil, and its derivatives (BILTZ and BÜLOW), A., i, 382.
- α-Dimethylaminoisovaleric acid, ethyl ester (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.
- Dimethylammonium chloride, action of dicyanodiamide with (WERNER and BELL), T., 1790.
- ruthenipentabromide (GUTBIER and KRAUSS), A., i, 16.
- Dimethylanhalamine methiodide (SPATH), A., i, 165.
- Dimethylaniline, electrochemical oxidation of (FICHER and ROTHENBERGER), A., i, 447.
- action of 1-chloro-2-aminoanthraquinone with, in presence of copper (KOPETSHNI and WIESLER), A., i, 844.
- and *p*-bromo- and *p*-nitroso-, hydroferrocyanides (CUMMING), T., 1297.
- Dimethylaniline, *o*-chloro-*p*-iododichloride hydrochloride and *o*-chloro-*p*-iodo- (SACHS and LEOPOLD), A., i, 822.
- m*-nitro-, nitro-derivatives of (FORSTER and COULSON), T., 1988.
- 5:8-Dimethylanthraquinone, 1-chloro- (MAYER and HEIL), A., i, 878.
- 2:6-Dimethylapophyllenic acid, and its ethyl and methyl esters (MUMM and GOTTSCHALDT), A., i, 263.
- 5:5'-Dimethylarsenobenzene, 3:3'-diamino-4:4'-dihydroxy-, dihydrochloride (CHRISTIANSEN), A., i, 1203.
- Dimethylarsinecarboxylic acid (JOB and GUINOT), A., i, 75.
- 2:4-Dimethylatropic acid, β-chloro- (v. AUWERS and ZIEGLER), A., i, 140.
- 2:3-Dimethylazobenzene, 4-amino-, derivatives of (JACOBSEN), A., i, 594.
- 3:3'-Dimethylazobenzene, 4:4'-(*f*)-dichloro- (GOLDSCHMIDT and STROMMINGER), A., i, 1005.
- 3:5-Dimethylazobenzene (JACOBSEN), A., i, 595.
- 4:6-Dimethylbenzaldehyde, 2-hydroxy-, and its derivatives (v. AUWERS and SAURWEIN), A., i, 1032.
- Dimethylbenzobisthiazoles, and their salts (EDGE), T., 774.
- 1:4-Dimethylbenzenesulphonophthalein, 5-hydroxy-. See Xylenol-blue.
- 4:6-Dimethylbenzonitrile, 2-hydroxy- (v. AUWERS and SAURWEIN), A., i, 1032.
- o-2':5'-Dimethylbenzoylbenzoic acid, β-chloro- (MAYER and HEIL), A., i, 878.
- 2:4-Dimethylbenzoylbenzoic acids, and their silver salts and methyl esters (SMITH), A., i, 141.
- 1:5-Dimethylbenzthiazole ethiodide (MILLS), T., 463.
- βγ-Dimethyl-Δγ-butadiene (*disopropenyl*), combination of hydrogen bromide with (BERGMANN), A., i, 1106.
- ββ-Dimethylbutane, synthesis of (VAN RISSEGHEN), A., i, 813.
- βγ-Dimethylbutane, αγ-dibromo- (BERGMANN), A., i, 1106.
- β-chloro γ-hydroxy- (DETÈUF), A., i, 327.
- βγ-Dimethyl-Δα-butene, γ-bromo- (BERGMANN), A., i, 1106.
- β-Dimethylchloroacamin (KÜSTER and GERLACH), A., i, 597.
- α-1:3-Dimethyl-1-dichloromethyl-Δ²5-cyclohexadien-Δ⁴-acetic acid, and its ethyl ester (v. AUWERS and ZIEGLER), A., i, 140.
- 1:4-Dimethyl-1-dichloromethyl-Δ²5-cyclohexadien-4-ol, 5-mono-, and 3:5-di-bromo (v. AUWERS and ZIEGLER), A., i, 145.
- 1:4-Dimethyl-1-trichloromethyl-Δ²5-cyclohexadien-4-ol, 3:5-di-bromo- (v. AUWERS and JÜLICHER), A., i, 843.
- 1:3-Dimethyl-1-dichloromethyl-Δ²5-cyclohexadien-4-one, 5-bromo- (v. AUWERS and ZIEGLER), A., i, 145.
- 1:2-Dimethyl-1-dichloromethyl-Δ²5-cyclohexadien-4-ones, 5-mono- and 3:5 di-chloro- (v. AUWERS and ZIEGLER), A., i, 144.
- 1:2-Dimethyl-1-dichloromethyl-Δ⁴-cyclohexen-4-one, 5:6-di-chloro- (v. AUWERS and ZIEGLER), A., i, 144.
- 2:6-Dimethyleinchomeronic acid, esters of, and their methiodides (MUMM and GOTTSCHALDT), A., i, 861.
- 1:1'-Dimethylisocyanine, 6-bromo-, iodide (MOUDGILL), T., 1510.

- 7:9-Dimethyldeoxyuric acid**, and its salts and derivatives (BILTZ and BÜLOW), A., i, 382.
- $\alpha\alpha'$ -Dimethyldisobutenyl disulphide** (POPE and SMITH), T., 1167.
- $\alpha\alpha'$ -Dimethyldisobutyl disulphide**, $\beta\beta'$ -dichloro- (POPE and SMITH), T., 1167.
- 5:5'-Dimethyl-2:2'-diethylcarbothiocyanine iodide** (MILLS), T., 464.
- Dimethyldihydroapomorphimethine**, and its methiodide (FALTIS and KRAUSZ), A., i, 676.
- 1:6-Dimethyl- Δ^6 ,7-5:8-dihydronaphthalene**, and its dibromide (MAYER and SCHULTE), A., i, 820.
- 2:6-Dimethyl- Δ^2 ,3-dihydronaphthalene** (MAYER and ALKEN), A., i, 999.
- Dimethyldihydroresorcinol**, chloro-, action of phosphorus chlorides on (HINKEL and WILLIAMS), T., 2498.
- 7:7-Dimethyl- α -dinaphthaxanthen**, oxidation of (SEN-GUPTA and TUCKER), T., 559.
- 7:7-Dimethyl- α -dinaphthaxanthen**, 5:9-dihydroxy- (SEN-GUPTA and TUCKER), T., 561.
- 3:3'-Dimethyldinaphthyl sulphate**, 4:4'-diamino-, and its derivatives (FRIES and LOHMANN), A., i, 30.
- 2:4-Dimethyldiphenyl** (2,4-ditolyl), and tetranitro- (JACOBSEN), A., i, 595.
- 2:6-Dimethyldiphenyl** (2:6-ditolyl), and 4:4'-diamino-, and its benzylidene derivative, and tri- and tetra-nitro- (JACOBSEN), A., i, 595.
- 3':4-Dimethyldiphenyl** (3':4-ditolyl), 2:4':5-triamino-, acetyl derivatives, and their derivatives (JACOBSEN), A., i, 595.
- 4:4'-Dimethyldiphenyl** (4:4'-ditolyl), 2,2'-dichloro-6:6'-dinitro- (BURTON and KENNER), T., 495.
- 2:4'-Dimethyldiphenylamine**, 4-amino-, and its salicylaldehyde derivative (JACOBSEN), A., i, 593.
- 3:3'-Dimethyldiphenylamine**, 4:4'-diamino-, 4'-acetyl derivative, and its salicylidene derivative (JACOBSEN), A., i, 595.
- 3':6-Dimethyldiphenylamine**, 2-amino-4'-hydroxy- (JACOBSEN), A., i, 593.
- $\alpha\alpha'$ -Dimethyl- $\beta\beta'$ -dipropylvalerolactone**, and its barium salt (LEROIUE), A., i, 218.
- 8:5-Dimethyldisulphon-1-phenyl-1:2:4-triazole** (FROMM, KAYSER, BRIEGLER, and FÖHRENBACH), A., i, 873.
- Dimethyldithioethylene**, reactions of metallic salts with (MORGAN and LEBURY), T., 2882.
- 2:3-Di-*mp*-methylenedioxyphenyl- $\alpha\beta$ -naphthaquinoxaline** (SCHÖNBERG and KRAEMER), A., i, 665.
- 3:4:3':4'-Dimethylenetetraoxybenzoylbenzoin**, 6'-nitro- (GREENE and ROBINSON), T., 2192.
- $\beta\beta'$ -Dimethyl- α -ethylglutaric acid**, $\alpha\alpha'$ -dicyano-, ω -imide and *N*-ethyl- ω -imide of (KON and THORPE), T., 1800.
- 1:1-Dimethyl-4-ethyl- $\Delta^{2,5}$ -cyclohexadien-4-ol** (V. AUWERS and ZIEGLER), A., i, 120.
- 1:4-Dimethyl-1-ethylcyclohexane-3:5-dione** (BECKER and THORPE), T., 1305.
- 1:1-Dimethyl-4-ethylidene- $\Delta^{2,5}$ -cyclohexadiene** (V. AUWERS and ZIEGLER), A., i, 120.
- 2:3-Dimethyl-4-ethylpyrrole** (*hæmopyrrole*), derivatives of (FISCHER, SCHNELLER, ZERWECK, and SCHUBERT), A., i, 1057.
- 2:6-Dimethyl-4-ethylquinoline**, and its tartrate (KNOEVENAGEL and BARR), A., i, 751.
- 1':6'-Dimethyl-2-ethylthioisocyanine iodide** (BRAUNHOLTZ and MILLS), T., 2007.
- Dimethylisoeugenolalkamine**, and its salts and derivatives (TAKEDA and KURODA), A., i, 275.
- O*-*N*-Dimethylisoeugenol-2-imino-oxazolidine**, and its hydrochloride (TAKEDA and KURODA), A., i, 274.
- 3:6-Dimethylfluoran**, 3'- (or 6')-nitro- (EDER and WIDMER), A., i, 260.
- Dimethyl γ -fructose** (IRVINE, STEELE, and SHANNON), T., 1074.
- $\alpha\beta$ -Dimethylglutaconic acids**, and their ethyl esters and ozonides (FEIST and BREUER), A., i, 521.
- $\beta\beta'$ -Dimethylglutaric acid**, $\alpha\alpha'$ -dihydroxy-, condensation of, with *o*-phenylenediamine (KON, STEVENSON, and THORPE), T., 665.
- 1:4- and 1:5-Dimethylglyoxalines**, orientation of, and their amino- and nitro-derivatives (PYMAN), T., 2616.
- Dimethylglyoxime**, preparation of, and its cobaltous salt (PONZIO), A., i, 17, 1039.
- $\beta\beta'$ -Dimethylguanidine**, preparation of (WERNER and BELL), T., 1790.
- Dimethylhematin** (KUSTER), A., i, 884.
- Dimethylhamin**, bromo- and hydroxy-, and their derivatives (KUSTER), A., i, 884.
- Dimethylharminesulphonic acid** (KER MACK, PERKIN, and ROBINSON), T., 1896.
- $\beta\epsilon$ -Dimethyl- $\Delta^{\alpha\gamma}$ - and - $\Delta^{\beta\gamma}$ -hexadienes** (KRETSINSKI), A., i, 1180.

- Di- γ -methylhexaldehyde**, *di- γ -hydroxy-*. See Bis-5-ethyl-6-methyltetrahydro-2-furyl ether.
- cis-2-trans-4-Dimethylcyclohexane**, *cis*-1-amino-, and its salts and derivatives (SKITA), A., i, 534.
- Dimethylcyclohexanes** (CHAVANNE and BECKER), A., i, 442.
- stereoisomeric (SKITA and SCHENCK), A., i, 241.
- aa'-Dimethylcyclohexane-1:1-diacetic acid**, *aa'-diacyano-*, ω -imide of (KON and THORPE), T., 1802.
- 1:1-Dimethylcyclohexane-3:5-dione**, bromo- and chloro-derivatives, preparation of (HIRST and MACBETH), T., 2176.
- 1:4-Dimethyl-1-cyclohexane-3:5-dione-2-carboxylic acid**, ethyl ester (BECKER and THORPE), T., 1304.
- $\beta\beta$ -Dimethylhexan- γ -ol**, and its phenylurethane (LEROIDE), A., i, 216.
- 1:4-Dimethylcyclohexan-2-ol** (SKITA), A., ii, 208.
- 1:1-Dimethyl- Δ^4 -cyclohexen-3-one**, 4:5-dichloro- (HINKEL and WILLIAMS), T., 2500.
- Di-methylcyclohexylamines**, and their hydrochlorides (MAILHE), A., i, 332.
- 5:5-Dimethylhydantoin** (INGOLD, SAKO, and THORPE), T., 1192.
- 3:5-Dimethylhydrazobenzene** (JACOBSEN), A., i, 595.
- 2:4- and 3:4-Dimethyl-1-hydrindones**, 7-hydroxy-, derivatives of (V. AUWERS, HILLIGER, and WULF), A., i, 1194.
- 2:2-Dimethylindan-1-one**, and its semicarbazone (HALLER and BAUER), A., i, 258.
- 2:2'-Dimethyl-3:3'-indil(2:2'-dimethyl-3:3'-indolyl)**, and its derivatives (ODDO and SANNA), A., i, 371.
- 5:7-Dimethylisatin O-methyl ether**, and its phenylhydrazone (HELLER, BENADE, and HOCHMUTH), A., i, 1060.
- Di-1-methylisatinazaine** (BORSCHKE and MEYER), A., i, 55.
- Dimethylitaconic acid**. See Teraconic acid.
- Dimethylketenphenylimine** (STAUDINGER and HAUSER), A., i, 29.
- d-1:1-Dimethyl-2- γ -ketobutylcyclopropane-3-carboxylic acid**, and its derivatives (SIMONSEN), T., 2295.
- l-aa-Dimethyl-leucinol**, and its salts (KARRER, GISLER, HORLAHER, LOCHER, MÄDER, and THOMANN), A., i, 814.
- Dimethylmalonbromoamide** (INGOLD, SAKO, and THORPE), T., 1192.
- 1:1-Dimethyl-6-methylene(1:2:2)-dicycloheptane-3-carboxylic acid** (BREDT), A., i, 937.
- 1:1'-Dimethylstreptomovinylen-2:2'-quinocyanine iodide** (KÖNIG), A., i, 1190.
- 1:6-Dimethylnaphthalene**, reduction of (MAYER and SCHULTE), A., i, 819.
- 2:6-Dimethylnaphthalene**, 1-amino-, 1-mono-, and tri-nitro-, and their derivatives (MAYER and ALKIN), A., i, 1000.
- ee-Dimethylnonan-3 ζ -diol** (LEROIDE), A., i, 217.
- $\beta\eta$ -Dimethyl- $\Delta\beta\zeta$ -octadiene** (STAUDINGER, KREIS, and SCHILT), A., i, 978.
- $\beta\delta$ -Dimethylpentane**, γ -chloro- $\beta\delta$ -dihydroxy-(tetramethylglycerol) (PASTUREAU and BERNARD), A., i, 717.
- 1:1-Dimethylcyclopentane-3:4-dione**, preparation and derivatives of (KON), T., 524.
- $\beta\beta$ -Dimethylpentan- γ -ol**, and its phenylurethane (LEROIDE), A., i, 216.
- 5:5-Dimethylcyclopentan-3-ol-1:2-dicarboxylic acid** (FARMER, INGOLD, and THORPE), T., 152.
- 5:5-Dimethylcyclopentan-3-one-2-carboxylic acid**, bromo-derivatives (FARMER, INGOLD, and THORPE), T., 158.
- 5:5-Dimethyldicyclopentan-3-one-1-carboxylic acid**, bromo- and chloro-derivatives (FARMER, INGOLD, and THORPE), T., 153.
- 5:5-Dimethyl- Δ^1 -cyclopentene-3:4-dione-1-carboxylic acid** (FARMER, INGOLD, and THORPE), T., 148.
- 5:5-Dimethyl- Δ^4 -cyclopenten-1-ol-3-one**, and its chloro-derivatives (FARMER, INGOLD, and THORPE), T., 156.
- 5:5-Dimethyl- Δ^4 -cyclopenten-3-one-1-carboxylic acid**, 4-oximino- (FARMER, INGOLD, and THORPE), T., 148.
- N-Dimethyl-s-phthalylidithiourethane** (V. BRAUN and KAISER), A., i, 659.
- Dimethylpicrocellin** (FORSTER and SAVILLE), T., 820.
- 2:4-Dimethyl-6-propenylpyridine**, and its salts (KONDÓ and TAKAHASHI), A., i, 1177.
- aa-Dimethyl- γ -isopropylglycerols**, and their derivatives (KRETSINSKI), A., i, 1130.
- 1:1-Dimethyl-4-n-propyl- $\Delta^{2:5}$ -cyclohexadien-4-ol** (V. AUWERS and ZIEGLER), A., i, 120.
- $\beta\beta$ -Dimethyl- γ -propylhexan- γ -ol** (LEROIDE), A., i, 216.
- 1:5-Dimethyl-2-isopropylcyclohexanol**. See Methylmenthol.
- aa-Dimethyl- β -propylhexoic acid**, β -hydroxy-, ethyl ester (LEROIDE), A., i, 217.

- 1:1-Dimethyl-4-*n*-propylidene- $\Delta^{3,5}$ -cyclohexadiene (v. AUWERS and ZIEGLER), A., i, 120.
- 1:6-Dimethyl-4-isopropyl-naphthalene. See Cadalene.
- 2:4-Dimethyl-6-propylpiperidine, and its salts (KONDŌ and TAKAHASHI), A., i, 1177.
- 1:5-Dimethylpyrazole, 3-chloro-, 3-chlorobromo-, and 3-chloronitroso- (ROJAHN), A., i, 1183.
- 3:4-Dimethylpyrazole, 5-chloro- (ROJAHN), A., i, 1184.
- 5:8-Dimethylpyrazole-anthrone (MAYER and HEIL), A., i, 878.
- 2:2'- and 4:4'-Dimethylpyrazole-anthrone-yellow, and their derivatives (MAYER and HEIL), A., i, 878.
- 1:5-Dimethylpyrazol-3-one, 4-bromo- (ROJAHN), A., i, 1184.
- 3:5-Dimethylpyrrole-4-carboxylic acid, 2-hydroxy-, acetyl derivative (FISCHER and HERRMANN), A., i, 1054.
- 2:4-Dimethylpyrrole-3- and -5-carboxylic acids, nitro-, and their ethyl esters (FISCHER and ZERWECK), A., i, 759.
- 2:4-Dimethylpyrrole-3:5-dicarboxylic acid, esters (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 857.
- 2:4-Dimethylquinoline, and its salts (KNOEVENAGEL and BAHR), A., i, 751.
- Dimethylisaccharic acid, lead salt (IRVINE and HIRST), T., 1223.
- N*-Dimethylisosafrrolealkamine, and its salts and derivatives (TAKEDA and KURODA), A., i, 274.
- Dimethylisosafrrole-2-imino-oxazolidine, and its hydriodide (TAKEDA and KURODA), A., i, 274.
- α -Dimethylsuccinic hydroxyanhydride, chloro- (FARMER, INGOLD, and THORPE), T., 158.
- 3:5-Dimethylsulphone-1-nitrophenyl-1:2:4-triazole (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 378.
- 1:1'-Dimethyltetrahydro-4:4'-dicollidyl. See 1:2:4:6:1':2':4':6'-Octamethyltetrahydro-4:4'-dipyridyl.
- 1:6-Dimethyl-5:6:7:8-tetrahydronaphthalene (MAYER and SCHULTE), A., i, 820.
- 3:7-Dimethyl-7:8:9:10-tetrahydro- ψ -phenanthroline, and its salts (LINDNER), A., i, 688.
- 2:3'-Dimethylthiocyanine iodide (MILLS), T., 465.
- NS*-Dimethylthiodi- β -naphthylamine-sulphonium perchlorate (KEHRMANN and DARDEL), A., i, 1064.
- NS*-Dimethylthiodiphenylaminesulphonium hydroxide, and its perchlorate (KEHRMANN and DARDEL), A., i, 1063.
- 3:5-Dimethylthiol-1-phenyl-1:2:4-triazole (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 378.
- NS*-Dimethylthiophenyl- β -naphthylaminesulphonium perchlorate (KEHRMANN and DARDEL), A., i, 1064.
- 7:9-Dimethyl-8-thiouric acids (BILTZ and BULOW), A., i, 381.
- $\alpha\beta$ -Dimethyltricarballic acid (HOPE and SHELDON), T., 2230.
- Dimethyltriketone-trioxime, and its derivatives (PONZIO and RUGGERI), A., i, 630.
- 3:3-Dimethyl-2:5:5-tripropyltetrahydrofuran (LEROIDE), A., i, 217.
- Dimethylvalinol (KARRER, GISLER, HORLACHER, LOCHER, MADER and THOMANN), A., i, 814.
- Dimethylxylan (HEUSER and RUPPEL), A., i, 811.
- Di- $\alpha\beta$ -naphthacarbazole (KALLE & Co.), A., i, 867.
- Di- $\alpha\beta$ -naphthacarbazole-5:8-disulphonic acid (KALLE & Co.), A., i, 867.
- Dinaphthamine. See Di- $\alpha\beta$ -naphthacarbazole.
- Di- β -naphthalenesulphonylpentamethylenediamine (BERGELL), A., i, 720.
- Dinaphthalenesulphonylpiperazine (BERGELL), A., i, 720.
- $\beta\beta$ -Di-2- α -naphthaquinonylbutane (SEN-GUPTA and TUCKER), T., 564.
- $\beta\beta$ - and $\gamma\gamma$ -Di-2- α -naphthaquinonylpentanes (SEN-GUPTA and TUCKER), T., 565.
- $\beta\beta$ -Di-2- α -naphthaquinonylpropane (SEN-GUPTA and TUCKER), T., 560.
- Dinaphthathiophen, 5:6:11:13-tetrahydroxy-, and its derivatives (BRASS and KÖHLER), A., i, 1052.
- Dinaphthathiophendiquinone (BRASS and KÖHLER), A., i, 1052.
- Dinaphthathiophen-6:11-quinone, 5:13-dihydroxy-, and its acetyl derivatives (BRASS and KÖHLER), A., i, 1052.
- α -Dinaphthaxanthens, oxidation of (SEN-GUPTA and TUCKER), T., 557.
- Di- α -naphthol trisulphide (WATSON and DUTT), T., 2418.
- $\beta\beta'$ -Di-2-naphthoxydiethylamine (CLEMO and PERKIN), T., 646.
- Dinaphthyl 1:1'-disulphide, 5:5'-dichloro-, and 8:8'-dinitro- (REISSERT), A., i, 584.
- 5:5'-disulphide, 1:1'-dihydroxy- (WATSON and DUTT), T., 2416.
- 4:4'-dinitro-1:1'-diamino-, diacetyl derivative (REISSERT), A., i, 584.

- Dinaphthyl 8:8'-disulphide, 1:1'-diamino-** (REISSERT), A., i, 584.
- 1:1'-Dinaphthyl, 5:5'-dinitro-** (SCHOLL, SEER, and WEITZENBÖCK), A., i, 651.
- 2:2'-Dinaphthyl, 1:1'-diamino-** (KALLE & Co.), A., i, 867.
- Dinaphthylazobenzene** (PUMMERER, BINAFFL, BITTNER, and SCHURGRAF), A., i, 1197.
- Dinaphthylsulphonic acids, diamino-** (KALLE & Co.), A., i, 867.
- Di- α -naphthylketoneanil** (REDELLEN), A., i, 146.
- Di- α -naphthylketone- α -naphthyl** (REDELLEN), A., i, 146.
- Dinorcamphocyclohexanedione, and its disemicarbazone** (WEDEKIND and WEINAND), A., i, 549.
- Diolefines, preparation of, and their polymerides and derivatives** (TRAUN'S FORSCHUNGS-LABORATORIUM), A., i, 514, 515.
- Dioxalodipyridineiridic acid, potassium salt, and its derivatives** (DELEPINE), A., i, 860.
- Dioximes** (PONZIO), A., i, 17, 1037; (PONZIO and RUGGERI), A., i, 627.
- 4:4'-Di(oximinoacetyl-amino)-diphenylmethane** (BORSCHKE and MEYER), A., i, 53.
- 1:2-Dioxo-3-aminoanthraquinone, 2-tetraamino-1:2-dicobaltic salt** (MORGAN and SMITH), T., 168.
- 1:2-Dioxyanthraquinone, 2-pentammino-1:2-dicobaltic salt** (MORGAN and SMITH), T., 166.
- 1:2-Dioxyanthraquinone-3- and 4-quinolines, aminocobaltic salts** (MORGAN and SMITH), T., 168.
- 1:2-Dioxyanthraquinone-3-sulphonic acid, 2:3-dipentammino-1:2:3-tricobaltic salt** (MORGAN and SMITH), T., 167.
- Dioxycholadienic acid, and its salts and esters** (BOEDECKER and VOLK), A., i, 1027.
- 1:2-Dioxo-5:8-dihydroxyanthraquinone, 2-pentammino-1:2-dicobaltic salt** (MORGAN and SMITH), T., 167.
- 1:2-Dioxhydroxyanthraquinones, cobalt and aminocobalt salts** (MORGAN and SMITH), T., 166.
- 1:2-Dioxo-3-nitroanthraquinone, 2-pentammino-1:2-dicobaltic salt** (MORGAN and SMITH), T., 166.
- $\alpha\gamma$ -Dipalmitin** (AMBERGER and BROMIG), A., i, 804.
- $\beta\gamma$ -Dipalmitin- α -iodohydrin** (AMBERGER and BROMIG), A., i, 804.
- Diphenanthrylamine, di-9:10-mono-hydroxy-, as a reagent for nitrates** (FORESTI), A., ii, 524.
- Diphenic acid, γ -6:6'-dinitro-, 4:6:4':6'-tetranitro-, resolution of, and their salts and esters** (CHRISTIE and KENNER), T., 614.
- Diphenonaphthadiazole** (BUCHERER and WAHL), A., i, 465.
- 4:4'-Diphenoxybenzil** (SCHÖNBERG and KRAEMER), A., i, 664.
- $\beta\beta'$ -Diphenoxy- $\alpha\alpha'$ -dimethyldisobutyl disulphide** (POPE and SMITH), T., 1168.
- 2:3-Di- p -phenoxyphenyl-6-methylquinoxaline** (SCHONBERG and KRAEMER), A., i, 664.
- 2:3-Di- p -phenoxyphenyl- $\alpha\beta$ -naphth-quinoxaline** (SCHÖNBERG and KRAEMER), A., i, 664.
- $\Delta^9, 11$ -Diphenauccindadiene, 9 12-dichloro-** (BRAND and MÜLLER), A., i, 444.
- Diphenauccindane, 9:9:12:12-tetrachloro-** (BRAND and MÜLLER), A., i, 444.
- Δ^{10} -Diphenauccindene** (BRAND and MÜLLER), A., i, 445.
- Diphenauccindene series** (BRAND and MÜLLER), A., i, 444.
- Diphenyl hydrogen phosphate (diphenyl-phosphoric acid), compounds of, with amino-compounds** (BERNTON), A., i, 1146.
- disulphide, 2:5:2':5'-tetrachloro-, preparation of** (STEWART), T., 2558.
- di- m -hydroxy-** (WATSON and DUTT), T., 2415.
- Diphenylacetamidine, p -dihydroxy-, diallyl ether, and its hydrochloride** (SOCIETY FOR CHEMICAL INDUSTRY IN BASLE), A., i, 953.
- Diphenylacethydroxamic acid, and its salts and derivatives** (JONES and HUND), A., i, 249.
- Diphenylacetomethylamide** (STAUDINGER and HAUSER), A., i, 29.
- Diphenylacetoxystibine** (SCHMIDT), A., i, 1204.
- 4:6-Diphenyl-2-(6-acetoxy- m -tolyl)pyridine, and its picrate** (DILTHER, NUSSLEIN, MEYER, and KAFFER), A., i, 949.
- Diphenylallylamine** (FÖLDI), A., i, 732.
- Diphenylamine, constitution of the blue oxidation product of** (KEHRMANN and ROY), A., i, 467.
- action of nitric and sulphuric acids on** (KEHRMANN and MICEWICZ), A., i, 28.
- compound of tellurium tetrabromide with** (LOWY and DUNBROOK), A., i, 446.
- Diphenylamine, bromo-** (GOLDSCHMIDT and RENN), A., i, 477.

- Diphenylamine**, 4-bromo-4'-amino-, and its derivatives (JACOBSEN), A., i, 593.
 5-bromo-2:4-dinitro-2', -3', and -4'-hydroxy- (GIUA and ANGELETTI), A., i, 649.
 chloronitro- and chloronitroamino-derivatives (BLOM), A., i, 27.
d-nitro-, preparation of (MORAN), A., i, 648.
 thio-, sulphonium bases from (KEHRMANN and DARDEL), A., i, 1063.
- Diphenylaminocarboxylic acids**, 5-bromo-2:4-dinitro- (GIUA and ANGELETTI), A., i, 650.
- 1:3-Diphenyl-5-anilinomethylene-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- $\gamma\gamma$ -Diphenyl- $\alpha\alpha$ -di-*p*-anisylallene** (ZIGLER, OCHS, BREMER, and THIEL), A., i, 1049.
- Diphenyldi-*p*-anisylallyl alcohol** (ZIGLER, OCHS, BREMER, and THIEL), A., i, 1049.
- 3:5-Diphenyl-4-anisylisoxazole** (MEISENHEIMER and WEIBEZAHN), A., i, 177.
- Diphenylarsinecarboxylic acid** (JOB and GUINOT), A., i, 76.
- Diphenylarsinic acid**, and *di-p*-nitro- (BART), A., i, 1202
- Diphenylarsinoformamide** (JOB and GUINOT), A., i, 76.
- 2:2-Diphenylbenzobisthiazole**, 4-amino-, and its acetyl derivative, and 4-nitro- (BOGERT and ABRAHAMSON), A., i, 576.
- NN'*-Diphenylbenzidine**, *di-p*-chloro- and its derivatives (WIELAND, WECKER, and ALBERT), A., i, 780.
- 2:3-Diphenylbenzopyran-2-ol**, and its ethers (DECKER and BECKER), A., i, 358.
- 2:5-Diphenyl-*p*-benzoquinone**, *p-p'*-di-hydroxy-, and its acetate (PUMMERER and PRELL), A., i, 1165.
- Diphenylbenzylpyrrodiazole** (CURTIUS and EHREHART), A., i, 775.
- Diphenylbisacidi- α -naphthol trisulphide** (WATSON and DUTT), T., 2418.
- Diphenylbisacetyldihydroxynaphthyl mercaptans** (WATSON and DUTT), T., 2416.
- Diphenyl-4:4'-bisacetyldihydroxy-2-phenyl mercaptan** (WATSON and DUTT), T., 2415.
- Diphenyl-*pp'*-bisacophenol**, and its acetate (FICHTER and JAECK), A., i, 62.
- 1:1'-Diphenylbis-5:5'-tetrazole** (STOLLÉ), A., i, 689.
- Diphenylbromobismuthine**, *di-p*-chloro- (CHALLENGER and RIDGWAY), T., 110.
- Diphenylbromomethane**, action of phenylhydroxylamine with (RUPE and WITTEW), A., i, 449.
- 1:4-Diphenyl-3-*p*-bromophenyl-2-*p*-hydroxyphenyl-1:3-dimethindiazidine** (INGOLD and PIGGOTT), T., 2798.
- 4:6-Diphenyl-2-*p*-bromophenylpyridine** (GASTALDI), A., i, 368.
- Diphenylbromophthalides** (STEPHENS), A., i, 141.
- Diphenyltetrabromophthalide** (LAWRANCE and ODDY), A., i, 455.
- Diphenylcarbamic acid**, *p*-carbamidophenyl ester (SPECKAN), A., i, 580.
- 4:6-Diphenyl-2-*p*-chlorophenylpyridine** (GASTALDI), A., i, 367.
- 4:6-Diphenyl-2-*p*-chlorophenylpyridinium nitrate** (GASTALDI), A., i, 367.
- Diphenylchlorostibine**, *di-p*-amino-, acetyl derivative (SCHMIDT), A., i, 1204.
- Diphenyleanoarsine** (STEINKOPF and SCHWEN), A., i, 72; (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- Diphenyl-4:4'-diallyldinitroamine**, 3:3':5:5'-tetranitro- (VAN ROMBURGH), A., i, 276.
- Diphenyl-4:4'-disobutylidinitroamine** (VAN ROMBURGH), A., i, 276.
- 2:5-Diphenyl-3:6-disobutyltetrahydro-1:4:2:5-dioxadiarsine**, and its salts (PALMER and ADAMS), A., i, 787.
- Diphenyl-3:4-dicarboxylic anhydride** (V. AUWERS and JÜLICHER), A., i, 842.
- BB'*-Diphenyldiethyl sulphide**, *BB'*-di-chloro- (POPE and SMITH), T., 1168.
- Diphenyl-4:4'-diethyldinitroamine**, 3:3':5:5'-tetranitro- (VAN ROMBURGH), A., i, 275.
- 2:5-Diphenyl-3:6-diethyltetrahydro-1:4:2:5-dioxadiarsine** (PALMER and ADAMS), A., i, 787.
- 2:5-Diphenyl-3:6-difuryltetrahydro-1:4:2:5-dioxadiarsine** (PALMER and ADAMS), A., i, 787.
- 4:4-Diphenyl-4:5-dihydro-5-glyoxalone-2-carboxylic acid** (BILTZ, SEYDEL, and HAMBURGER-GLAZER), A., i, 872.
- s*-Diphenyldimethylcarbamide** (TANBERG and WINKEL), A., i, 1009.
- $\alpha\alpha$ -Diphenyl-*BB*-dimethylpropan- α -ol**, dehydration of (RAMART), A., i, 34.
- BB*-Diphenyl- $\alpha\alpha$ -dimethylpropiofenone** (RAMART and ALBESCO), A., i, 663.
- 2:5-Diphenyl-3:6-dimethyltetrahydro-1:4:2:5-dioxadiarsine**, and its salts (PALMER and ADAMS), A., i, 787.
- 5:5-Diphenyl-1:2-dimethyltetrahydro-4-glyoxalone-1':2-oxide**, and its derivatives (BILTZ, SEYDEL, and HAMBURGER-GLAZER), A., i, 871.

- Diphenyl-4:4'-di-*n*- and -*iso* propyldinitroamine, 3:3':5:5'-*tetra*nitro- (VAN ROMBURGH), A., i, 276.
- 2:5-Diphenyl-3:6-di-*n*-propyltetrahydro-1:4:2:5-dioxadiarsine (PALMER and ADAMS), A., i, 1787.
- Diphenylene-ethylene. See Dibenzotulvene.
- Diphenyleneoxide, 5-bromo- (MAYER and KRIEGER), A., i, 747.
- γ -5-Diphenyleneoxide-*n*-butyric acid, and its derivatives (MAYER and KRIEGER), A., i, 747.
- Diphenyleneoxide-5-carboxylic acid (MAYER and KRIEGER), A., i, 746.
- γ -5-Diphenyleneoxide- γ -keto-*n*-butyric acid, ethyl ester (MAYER and KRIEGER), A., i, 747.
- γ -5-Diphenyleneoxide-*n*-propylamine, and its hydrochloride and acetyl derivative (MAYER and KRIEGER), A., i, 747.
- γ -5-Diphenyleneoxide-*n*-propylurethane (MAYER and KRIEGER), A., i, 747.
- 3-5-Diphenyleneoxidepyridazin-6-one (MAYER and KRIEGER), A., i, 747.
- 3-Diphenylethane (*dibenzyl*) disulphoxide (WAHL), A., i, 653.
- β -Diphenylethylamine picrate (SIEGLITZ and SPITZER), A., i, 829.
- Diphenylethylarsine, and its salts (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- Diphenylethylcarbamide (SIEGLITZ and SPITZER), A., i, 829.
- 5:5-Diphenyl-1-ethyltetrahydro-4-glyoxalone 1':2'-oxide (BILTZ, SEYDEL, and HAMBURGER-GLAZER), A., i, 872.
- aa*-Diphenylethyl- β -urethane (SIEGLITZ and SPITZER), A., i, 829.
- Diphenylformylacetic acid, ethyl ester, acetal (STAUDINGER and RATHSAM), A., i, 1015.
- N*-Diphenylglycine-*o*-carboxylic acid, and *p*-chloro-, and their esters (FRIEDLANDER and KUNZ), A., i, 765.
- s*-Diphenylguanidine, use of, in acidimetry and alkalimetry (CARLTON), A., ii, 654.
- 1:4-Diphenylhexahydro-1:2:4:5-tetrazine, and its derivatives (THIELEPAPE and SPRECKELSEN), A., i, 1192.
- Diphenylhydroxytibine, *di-p*-amino-, acetyl derivative (SCHMIDT), A., i, 1204.
- Diphenyldihydroxy-*m*- and -*p*-tolylmethanes (MACKENZIE), T., 1698.
- 4:6-Diphenyl-2-(6-hydroxy-*m*-tolyl)pyridine, and its picrate (DILTHEY, NÜSSLEIN, MEYER, and KAFFER), A., i, 949.
- Diphenylimidoacetic acid, ethyl ester, and its hydrochloride (STAUDINGER and RATHSAM), A., i, 1015.
- 1:2-Diphenylindene, formation of (ORÉKHOFF and TIFFENEAU), A., i, 458.
- 1:1'-Diphenylindigotin, and 1:1'-*di-p*-chloro- (FRIEDLANDER and KUNZ), A., i, 766.
- Diphenyliodobismuthine, *di-p*-chloro- (CHALLENGER and RIDGWAY), T., 110.
- Diphenyltetraiodophthalide (LAWRANCE and ODDY), A., i, 455.
- Diphenylketenmethylimine (STAUDINGER and HAUSER), A., i, 29.
- Diphenylketimine, 2:4:6:2'-tetrahydroxy-, and its hydrochloride (NISHIKAWA and ROBINSON), T., 840.
- Diphenylmethane, action of aluminium chloride with (SCHOLL, SEER, and (NEUMANN), A., i, 336.
- Diphenylmethane, 2:2':4:4'-tetraamino-5:5'-dithiol- (WATSON and DUTT), T., 1943.
- 4:6-Diphenyl-2-(4-methoxy-3-methylphenyl)pyridine, and its hydrochloride (DILTHEY, NÜSSLEIN, MEYER, and KAFFER), A., i, 949.
- Diphenylmethylamine ferrocyanide (FÖLDI), A., i, 732.
- Diphenylmethylarsine, salts of (STEINKOPF and SCHWEN), A., i, 72.
- β -Diphenyl- γ -methylbutane, γ -chloro- (RAMART), A., i, 35.
- β -Diphenyl- γ -methylbutan- γ -ol (RAMART), A., i, 35.
- Diphenylmethylcarbonyl chloride (JONES and HURD), A., i, 249.
- Diphenylmethylcarbamide (JONES and HURD), A., i, 249.
- 4:4-Diphenyl-2-methyl-4:5-dihydro-5-glyoxalone, and its derivatives (BILTZ, SEYDEL, and HAMBURGER-GLAZER), A., i, 871.
- Diphenylmethylphosphinic acid, α -hydroxy- (CONANT, MACDONALD, and KINNEY), A., i, 186.
- $\gamma\gamma$ -Diphenyl- β -methylpropane (RAMART and ALBESCO), A., i, 663.
- 4:6-Diphenyl-2-methylpyridine (GAS-TALDI), A., i, 573.
- 2:6-Diphenyl-4-methylpyrylium salts (SCHNEIDER and ROSS), A., i, 1172.
- 5:5-Diphenylmethyltetrahydro-4-glyoxalones, and their derivatives (BILTZ, SEYDEL, and HAMBURGER-GLAZER), A., i, 871.
- Diphenyl- α -methylthionaphthylacetic acid, and its derivatives (BISTRZYCKI and KUBA), A., i, 34.

- Diphenyl- α -methylthiolnaphthylmethane** (BISTRZYCKI and KUBA), A., i, 84.
- Diphenylmethyllurethane** (JONES and HURD), A., i, 249.
- Diphenyl- α -naphthylbismuthine dichloride** (CHALLENGER and WILKINSON), T., 102.
- Diphenyl- β -naphthylmethane** (McMULLEN), A., i, 1026.
- Diphenyl- β -naphthylmethyl**, and its derivatives (GOMBERG and SULLIVAN), A., i, 929.
- $\alpha\alpha$ -Diphenyl- β -trinitrophenylhydrazine** (GOLDSCHMIDT and RENN), A., i, 477.
- $\alpha\alpha$ -Diphenyl- β -trinitrophenylhydrazyl**, and β -hydroxy-, and their derivatives (GOLDSCHMIDT and RENN), A., i, 477.
- 1:4-Diphenyl-3-*m*-nitrophenyl-2-*p*-hydroxyphenyl-1:3-dimethindiazidine** (INGOLD and PIGGOTT), T., 2798.
- 1:4-Diphenyl-3-*o*-nitrophenyl-2-*p*-nitrophenyl-1:3-dimethindiazidine** (INGOLD and PIGGOTT), T., 2799.
- Diphenylnitrosoamine** (SCHMIDT and FISCHER), A., i, 822.
- 3:5-Diphenylisoxazole-4-carboxylic acids**, derivatives of (BETTI and BERLINGOZZI), A., i, 52.
- $\beta\beta$ -Diphenylpentane**, γ -chloro- (RAMART), A., i, 35.
- $\beta\beta$ -Diphenylpentan- γ -ol** (RAMART), A., i, 35.
- Diphenylphenylethynylcarbinol**, preparation and reactions of (HESS and WELTZEN), A., i, 35.
- Diphenylphenylethynylcarbinol**, *di-p*-chloro- (MEYER and SCHUSTER), A., i, 556.
- Diphenylphosphoric acid**. See Diphenyl hydrogen phosphate.
- $\beta\beta$ -Diphenylpropionic acid**, ethyl ester, azide and hydrazide of (SIEGLITZ and SPITZER), A., i, 828.
- Diphenylpropylcarbinol** (SKRAUP and FREUNDLICH), A., i, 540.
- $\alpha\gamma$ -Diphenylpropylphosphinic acid**, α -hydroxy- (CONANT, MACDONALD, and KINNEY), A., i, 186.
- 2:4-Diphenylpyridine**, and its derivatives (GASTALDI), A., i, 574, 679.
- 2:4-Diphenylpyridine**, 6-hydroxy-, and its 5-carboxylic acid, ethyl ester (KOHLEK), A., i, 461.
- 2:4-Diphenylpyridine-6-carboxylic acid**, and its sodium salt (GASTALDI), A., i, 574.
- 2:4-Diphenyl-6-pyrone** (KOHLEK), A., i, 461.
- 2:4-Diphenyl-6-pyrone-5-carboxylic acid**, and its sodium salt and esters (KOHLEK), A., i, 461.
- Diphenylpyrrylcarbinol**, and its derivatives (FISCHER and KAAK), A., i, 869.
- 2:5-Diphenylquinol**, *pp'*-*di*hydroxy-, and its tetra-acetate (PUMMERER and PRELL), A., i, 1165.
- 2:3-Diphenylquinoline-4-carboxylic acid** (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 867.
- Diphenylstibinic acid**, *di-p*-amino-, acetyl derivative (SCHMIDT), A., i, 1204.
- Diphenylstyrylcarbinol** (MEYER and SCHUSTER), A., i, 540.
- S*-Diphenylsuccinic acid**, preparation of (LAPWORTH and MCKAE), T., 1709.
- S*-Diphenylsuccinimide** (LAPWORTH and MCKAE), T., 1712.
- Diphenylsulphone**, *di*bromo-, *di*chloro-, and *di*iodo-derivatives (GRANDMOUGIN), A., i, 251.
- 3:3'-*di*iodo- and 2:2' *di*nitro-** (GRANDMOUGIN), A., i, 331.
- 2:5-Diphenyltetrahydro-1:4:2:5-*di*oxarsine** (PALMER and ADAMS), A., i, 786.
- Diphenylthiocyanobismuthine** (CHALLENGER and WILKINSON), T., 98.
- Diphenylthiolbenzene**, derivatives of (SMILES and GRAHAM), T., 2506.
- 1:3-Diphenylthiophthalan** (BISTRZYCKI and BRENNEN), A., i, 269.
- 2:4-Diphenyl-6-*p*-tolylpyridine** (GASTALDI), A., i, 368.
- 2:4-Diphenyl-6-*p*-tolylpyrylium nitrate** (GASTALDI), A., i, 368.
- 9- $\beta\beta$ -Diphenylvinylidenexanthene** (ZIEGLER, OCHS, BREMER, and THIEL), A., i, 1049.
- $\beta\beta$ -Diphenylvinylxanthyl**, and its derivatives (ZIEGLER, OCHS, BREMER, and THIEL), A., i, 1049.
- 2:4-Diphthaliminophenol** (MUKERJI), T., 549.
- 4:6-Diphthaliminoresorcinol** (MUKERJI), T., 549.
- Dipthalyl-lactonic acid**, preparation of (RUGGLI and MEYER), A., i, 343.
- Diphtheria toxin** (WALBUM), A., i, 902.
- Dipicrylamine**, cobaltamines of (MORGAN and KING), T., 1726.
- Dipiperidino-4-phenylmethane**, *di*-4-hydroxy- (BORSCHKE and BONACKER), A., i, 51.
- 2:5-Dipiperonyloxazole** (GREENE and ROBINSON), T., 2188.
- 2:3-Dipiperonylquinoxaline** (GREENE and ROBINSON), T., 2194.
- Diisopropenyl***. See $\beta\gamma$ -Dimethyl- Δ^{γ} butadiene.
- Dipropionylpentamethylenediamine**, *di*- α -bromo- (BERGELL), A., i, 720.

- Dipropyl ethers**, $\gamma\gamma'$ -dibromo-, and $\gamma\gamma'$ -dichloro- (KAMM and NEWCOMB), A., i, 105.
- selenide, $\beta\beta'$ -dichloro-, and its dichloride (BOORD and COPE), A., i, 421.
- $\beta\beta'$ -Dipropyl sulphides, $\gamma\gamma'$ -tetrachloro- (POPE and SMITH), T., 1168.
- $\beta\beta'$ -Dipropylacetaldehyde, and its semicarbazone (LOCQUIN and WOUSENG), A., i, 711.
- $\gamma\gamma'$ -Dipropylallyl alcohol, and its allophanate (LOCQUIN and WOUSENG), A., i, 711.
- Dipropylammonium** ruthenibromides (GUTHRIE and KRAUSS), A., i, 16.
- Dipropylbarbituric acid**, action of diazomethane on (HERZIG), A., i, 574.
- Di-*n*- and iso-propylbenzidines**, 3:3':5:5'-tetranitro- (VAN ROMBURGH), A., i, 275.
- Dipropylcampholactone**, and its copper salt (LEROIDE), A., i, 218.
- Di-*n*-propyleyanosamine** (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- $\delta\eta$ -Dipropyl- Δ^c -decinene- $\delta\eta$ -diol (LOCQUIN and WOUSENG), A., i, 617.
- Dipropylethynylcarbinol**, and its allophanate (LOCQUIN and WOUSENG), A., i, 617.
- Dipropylethylcarbinol**, and its allophanate (LOCQUIN and WOUSENG), A., i, 617.
- s-Diisopropylhydrazine**, and its derivatives (LOCHTE, BAILEY, and NOYES), A., i, 329.
- $\delta\epsilon$ -Dipropylactane (PICCARD and BREWSTER), A., i, 313.
- Diisopropylphenylthiosemicarbaside** (LOCHTE, BAILEY, and NOYES), A., i, 329.
- Dipropylisopropylcarbinol** phenylurethane (LEROIDE), A., i, 217.
- Dipropylvinylcarbinol**, and its allophanate (LOCQUIN and WOUSENG), A., i, 710.
- Dipyridineiridic acid**, tetrachloro-, potassium salt (DELÉPINE), A., i, 859.
- Dipyridineiridium**, tetrachloro-, and its derivatives (DELÉPINE), A., i, 859.
- Dipyridinium** μ -dichloro-hexachloropyridinediferrate and μ -dichloro-tetrachlorodinitratodipyridinediferrate (WEINLAND and KISSLING), A., i, 364.
- 4:4'-Dipyridyl diisomyl hydroxide** and its salts (EMMERT and PARR), A., i, 179.
- diisobutyl iodide and platinichloride (EMMERT and PARR), A., i, 180.
- Dipyrryl**, heptaiodo- (PIERONI), A., i, 763.
- Dipyrrylphenylmethane colouring matters**, iron salts of (FISCHER and LUCKMANN), A., i, 276.
- 6-Diquinaldylmethane**, and its picrate (BORSCHKE and MEYER), A., i, 53.
- Di-2-quinolyl ketone**, and its derivatives (SCHEIBE and SCHMIDT), A., i, 1190.
- Diquinolylmethane** (BORSCHKE and MEYER), A., i, 53.
- Disaccharides**, constitution of (HAWORTH and LEITCH), T., 1921.
- containing selenium and sulphur, synthesis of (WREDE), A., i, 226.
- Disalcin disulphide**, and its octa-acetate (ZEMPLÉN and HOFFMANN), A., i, 563.
- Disalcinamine** (ZEMPLÉN and KUNZ), A., i, 564.
- Disalcylaldehyde**, constitution of (ADAMS, FOGLER, and KREGER), A., i, 660.
- Disalcyllic acid** (*anhydrosalcyllic acid*), and its salts and derivatives (AUSCHUTZ and CLAASEN), A., i, 456.
- Disinfection**, theory of (TRAUBE and SOMOGYI; TRAUBE), A., i, 204; (VERMAST), A., i, 406.
- Dispersion of colloidal solutions** (TADOKORO), A., ii, 204.
- rotatory (LOWRY and AUSTIN), A., ii, 414.
- calculation of (RUPE), A., ii, 602.
- anomalous (RUPE and SCHMID), A., ii, 603.
- of organic compounds (LOWRY and CUTTER), T., 532.
- Dispersity** (OSTWALD), A., ii, 131.
- Dispersoid analysis**. See Analysis.
- Diss**, ergot of. See Elgot.
- Dissociation constants** of dibasic acids (DUBOUX), A., ii, 346.
- Distillation** (GAY), A., ii, 120.
- of foaming liquids (KLANHARDT), A., ii, 472.
- separation of miscible liquids by (DUFTON), T., 306.
- Distillation apparatus**, receiver for fractionation (WILLIAMS), A., ii, 26; (WHEELER and BLAIR), A., ii, 258.
- stillhead for prevention of spray (HAHN), A., ii, 368.
- Distyryl sulphide** (POPE and SMITH), T., 1168.
- Distyryl ketone** (*dibenzylideneacetone*), halochromism of (HANTZSCH), A., i, 556.
- Distyryl ketone**, unsaturated halogen derivatives of, and their conductivity (STRAUS and DÜTZMANN), A., i, 148.
- 3':2-dihydroxy-, and its dibenzoyl derivative (BUCK and HEILBRON), T., 1097.

- Distyryl ketones**, benzopyrylium salts of (BUCK and HEILBRON), T., 1198.
- Di-*o*-styrylxanthyl ether** (ZIEGLER and OCHS), A., i, 1048.
- Disulphides** (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 877.
aromatic, action of sulphuric acid on (SMILES and McCLELLAND), T., 86.
- p*-Disulphidobenzoic acid** (SMILES and HARRISON), T., 2022.
- 5-Disulphidobis-2:4-dimethylpyrrole-3-carboxylic acid**, ethyl ester (FISCHER and HERMANN), A., i, 1055.
- Disulphidodiphenylbisazo- α -naphthol-2 sulphonic acid** (WATSON and DUTT), T., 2418.
- Disulphosulcylaldehyde**, and its salts and derivatives (WEILL and BRIMMER), A., i, 349.
- Di-*ar*-tetrahydronaphthyls**, diamino-, and their salts (SCHROETER, KINDERMANN, DIERICH, BEYSLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 123.
- Dithienoylhydroxamic acid**, and its salts (JONES and HURD), A., i, 250.
- 2-Dithiobenzoyl** (SMILES and McCLELLAND), T., 89.
- Dithiocarbamic acid**, derivatives of, as accelerators of vulcanisation (TWISS, BRAZIER, and THOMAS), A., i, 460.
- Dithionates**. See under Sulphur.
- 3:5-Dithion-1:2:4-tribenzo-1:2:4-triazole** (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 379.
- Di(toluene-*p*-sulpho)diphenylethylene-diamine** (CLEMO and PERKIN), T., 648.
- 5-Ditoluene-*p*-sulphonylaminoacene-naphthene** (FLEISCHER and SCHRANZ), A., i, 1144.
- Di-*p*-toluenesulphonylethylene** (FROMM and SIEBERT), A., i, 543.
- Ditoly**. See Dimethyldiphenyl.
- Ditoly dimethyl ether** (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 934.
- Ditolyamine**, amino- (GOLDSCHMIDT and STROHMENGER), A., i, 1005.
- Di-*p*-tolylantipyrrol selenide** (FARRWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 1066.
- NN'*-Di-*p*-tolylbenzidine** (WIELAND, WECKER, and ALBERT), A., i, 780.
- 2:5-Di-*p'*-tolyl-*p*-benzoquinone** (PUMMERER and PRELL), A., i, 1165.
- Ditolytetra-bromophthalide** (LAWRANCE and ODDY), A., i, 455.
- 8-*γ*-Di-*m*-tolylbutane** (v. AUWERS and KOLLIGS), A., ii, 176.
- s*-Di-*p*-tolylearbamide-2:2'-disulphonic acid**, potassium salt (SCOTT and COHEN), T., 2041.
- Di-*p*-tolylechlorobismuthine** (CHALLENGER and RIDGWAY), T., 113.
- Di-*p*-tolylhydrazine dinitrosoamine** (WIELAND, WECKER, and ALBERT), A., i, 780.
- Ditolytetraiodophthalide** (LAWRANCE and ODDY), A., i, 455.
- 2:3-Di-*p*-tolyl- $\alpha\beta$ -naphthaquinoxaline** (SCHONBERG and KRAEMER), A., i, 665.
- 2:5-Di-*p*-tolylquinol**, and its derivatives (PUMMERER and PRELL), A., i, 1165.
- 3:4'-Ditolylsulphone**, 4-chloro- (STEINKOPF and BUCHHEIM), A., i, 122.
- s*-Ditolythiocarbamide**, compound of chloropicrin and (RAY and DAS), T., 327.
- Di-*p*-tolylthiolacetylene**, and its dibromide (FROMM and SIEBERT), A., i, 542.
- Di-*p*-tolylthioethylene**, and its derivatives (FROMM and SIEBERT), A., i, 542.
- Di-trimethylene glycol ether**, and its derivatives (ROJAHN), A., i, 103.
- Diuresis**, action of injections of dextrose and gum arabic on (CORT), A., i, 1091.
phosphate excretion in (BOCK and IVERSEN), A., i, 1090.
- Diuretics**, distribution coefficients of (AIELLO), A., i, 301.
- Divaleraldehyde**, *di-γ*-hydroxy-. See Bis-5-methyltetrahydro-2-furyl ether.
- $\alpha\beta$ -Disovalerylaminopropane**, and its dipicrate (WINDAUS, DORRIES, and JENSEN), A., i, 60.
- Divinyl sulphide**, synthesis of (BALES and NICKELSON), T., 2137.
- Divinylarsinic acid**, $\beta\beta'$ -dichloro- and its salts (MANN and POPK), T., 1756.
- 1:1'-Dixanthonyl**, 4:4'-dichloro- (ECKERT and ENDLER), A., i, 941.
- 4:5-Dixanthylpyrocatechol** (FABRE), A., i, 1148.
- 2:3-Dixanthylquinol** (FABRE), A., i, 1148.
- 2:4-Dixanthylresorcinol**, and its diacetyl derivative (FABRE), A., i, 1148.
- Di-*o*-xylyleneimine**, *p*-nitrobenzoyl derivative (v. BRAUN and NELKEN), A., i, 864.
- Dodecamethyltetra-*m*-voglucosan** (PRINGSHEIM and SCHMALZ), A., i, 1118.
- Doebner's reaction** (CIUSA), A., i, 1062.
- Doebner's-violet-leucosulphonic acid** (WIELAND and SCHEUING), A., i, 58.

- Dogs**, effect of injection of sugars on respiration of (BÜRGER), A., i, 286.
- Dopplerite** (TIDESWELL and WHEELER), T., 2345.
- Doryphora sassafras***, oil from the leaves of (PENFOLD), A., i, 847.
- Drosophila***, vitamins required by (BACOT and HARDEN), A., i, 493.
- Drugs**, distribution of, between immiscible solvents (EMERY and WRIGHT), A., ii, 205.
distribution of, in blood (SCHAEFFI), A., i, 290.
influence of the reaction of the solution on the action of, on bacteria or tadpoles (LABES), A., i, 901.
- Drying**, change of properties of substances on (BAKER), T., 568.
influence of, on internal change (SMITS), A., ii, 358.
- Dugaldin** (MARSH, CLAWSON, COUCH, and MARSH), A., i, 413.
- Dulcin**. See *p*-Phenethylcarbamide.
- Duodenal fluid**, analysis of (DAMADE), A., ii, 887.
- Duralumin**, analysis of (COSTA-VET), A., ii, 528.
- Dyeing**, use of mordants in (BANCROFF), A., ii, 551, 822; (PADDON), A., ii, 822.
of cotton, effect of salts on (AUERBACH), A., ii, 353.
of hides and leather (RAKUZIN), A., i, 702.
of deaminated wool (PADDON), A., i, 608.
- Dypnonanil** (KNORVENAGEL and GOOS), A., i, 751.

E.

- Earths, rare** (YNTEMA and HOPKINS), A., ii, 462; (HOPKINS and DRIGGS), A., ii, 770.
preparation of, pure, and their spectra (KIESS, HOPKINS and KREMERs), A., ii, 244.
preparation of pure compounds of (MANSKE), A., ii, 646.
place of, in the periodic system (NORRISH), A., ii, 211; (RENZ), A., ii, 758.
minerals containing (SHIBATA and KIMURA), A., ii, 220; (HENRICH and HILLER), A., ii, 860.
absorption spectra of (HERZFELD; MEYER and GREULICH), A., ii, 6.
structure of the ions of the (GRIMM), A., ii, 635.
rates of hydrolysis of carbonates of (BRINTON and JAMES), A., ii, 39.
- Earths, rare**, density and molecular volume of oxides of the (PRANDTL), A., ii, 379.
magnesium sulphide phosphors of (TIEDE and SCHLEDE), A., ii, 769.
detection of, spectroscopically (MEYER and GREULICH), A., ii, 6.
concentration and separation of (BRINTON and JAMES), A., ii, 62.
separation of (PRANDT and RAUCHENBERGER), A., ii, 298, 769; (PRANDTL and LÖSCH), A., ii, 770.
- Earthworm**, Japanese, constituents of (MURAYAMA and AOYAMA), A., i, 898.
- Ebulliscopy** of binary mixtures (BECKMANN and LIESCHE), A., ii, 117.
- Ecgonine** (GADAMER and JOHN), A., i, 167.
- Edestin**, colloidal properties of (HITCHCOCK), A., i, 693.
- Eggs**, hens', sugar contents of (HEPBURN and ST. JOHN), A., i, 294.
- Egg-albumin**. See Albumin.
- Egg-shells**, utilisation of calcium salts in formation of (BUCKNER, MARTIN, PIERCE and PETER), A., i, 488.
- Egg-yolk**, absorption spectrum of (LEWIN and STENGER), A., ii, 414.
- Einstein's law** in relation to photochemical action and fluorescence (WEIGERT), A., ii, 681.
- α -Eläostearic acid**, methyl ester, conversion of, into methyl β -eläostearate (MORRELL), A., i, 982.
- Elaidanilide dibromide** (NICOLET), A., i, 106.
- Elaidoacetylhydroxamic acid** (NICOLET and PELC), A., i, 644.
- Elaidodiacetylhydroxamic acid** (NICOLET and PELC), A., i, 644.
- Elaidohydroxamic acid** (NICOLET and PELC), A., i, 644.
- Electric discharge**, absorption of gases in the (NEWMAN), A., ii, 546.
disappearance of gases in the (GENERAL ELECTRIC Co.), A., ii, 417.
chemical reactions induced by the (MIYAMOTO), A., i, 418.
corona, chemical reactions induced by the (MONTEMARTINI), A., ii, 734.
action of, on fatty acids and their glycerides (EICHWALD), A., i, 982.
- furnace, ionisation experiments with the (KING), A., ii, 810.
tungsten (COMPTON), A., ii, 822.
- Electrical conductivity** of mixtures of acids with acids, esters and ketones (KENDALL and GROSS), A., ii, 33.

- Electrical conductivity** of electrolytes, measurement of (ABRIBAT), A., ii, 344; (ROLLA and MAZZA), A., ii, 544.
- of electrolytes in solution (KRAUS), A., ii, 256.
- of ternary electrolytes in propyl alcohol (KRAUS and BISHOP), A., ii, 19.
- of solid electrolytes (VAILLANT), A., ii, 418.
- of formates in formic acid and of sulphates in sulphuric acid (KENDALL, ADLER and DAVIDSON), A., ii, 126.
- of photographic emulsions and films (RABINOVICH), A., ii, 605.
- of molten salts and their mixtures (BENRATH and DREKOFF), A., ii, 109.
- Electrical endosmosis** (STRICKLER and MATHEWS), A., ii, 688.
- Electro-adsorption** (KOLTHOFF), A., ii, 197; (OSTWALD), A., ii, 427.
- Electrochemistry** of non-aqueous solutions (MÜLLER; MÜLLER and DUSCHKE), A., ii, 612.
- Electrodes**, overvoltage of (ISGARISCHEV and BERKMANN), A., ii, 253, 254.
- reproducible, for acidimetry and alkalimetry (SHARP and MACDOUGALL), A., ii, 579.
- hydrogen, quick acting (MOLONEY), A., ii, 253.
- use of phthalate solutions for measuring (OAKES and SALISBURY; CLARK), A., ii, 468; (WOOD and MURDICK), A., ii, 735.
- electro-volumetric analysis with, in alcoholic solution (BISHOP, KITTREDGE and HILDEBRAND), A., ii, 308.
- Clark hydrogen (HEALY and KARAKER), A., ii, 519; (CULLEN), A., ii, 611.
- iodine, potential of (McKEOWN), A., ii, 417.
- potentiometric estimations with (KOLTHOFF), A., ii, 388.
- mercury drop (HEYROVSKY), A., ii, 816.
- non-attackable (KLEMENC), A., ii, 253.
- platinum film, construction and use of (EILERT), A., ii, 610.
- quinhydrone (S. P. L. and M. SÖRENSEN and LINDERSTRÖM-LANG); (BIILMANN and LUND), A., ii, 111.
- sodium amalgam (NEUHAUSEN), A., ii, 610.
- zinc, effect of hydrogen-ion concentration on the potential of (RICHARDS and DUNHAM), A., ii, 418.
- Electrolysis**, rapid, without rotating electrodes (EDGAR and PURDUM), A., ii, 547.
- with mercury drops as electrode (HEYROVSKY), A., ii, 816.
- Electrolytes**, recent conception of (SCHREINER), A., ii, 736.
- constitution and function of (MADELUNG), A., ii, 344.
- structure of molecules of (WASASTJERNA), A., ii, 2.
- measurement of the conductivity of (KRAUS), A., ii, 258; (ABRIBAT), A., ii, 344; (ROLLA and MAZZA), A., ii, 544.
- activity coefficients and colligative properties of (HARNED), A., ii, 255.
- theory of solutions of (GHOSH), A., ii, 125.
- effect of, on the solubility of non-electrolytes (McKEOWN), A., ii, 552.
- diffusion of, into gels (STILES and ADAIR; STILES), A., ii, 125.
- adsorption and precipitation of, by colloids (WEISER), A., ii, 262; (WEISER and NICHOLAS), A., ii, 266.
- coagulation of colloids by (MURRAY), A., ii, 37; (FREUNDLICH and SCHOLZ), A., ii, 828.
- equilibrium law for (JABLONCZYŃSKI and WIŚNIEWSKI), A., ii, 190.
- hydration of (ASCHKENASI), A., ii, 482.
- condition of, in blood (NEUHAUSEN and MARSHALL), A., ii, 1085.
- amphoteric, viscosity of solutions of (HEDESTRAND), A., ii, 821.
- precipitation conditions for (PRINS), A., ii, 77.
- binary, physical properties of non-aqueous solutions of (WALDEN), A., ii, 345.
- binary mixed, equilibrium in (KRAUS), A., ii, 269.
- solid, changes in the conductivity of (VAILLANT), A., ii, 418.
- strong, theory of (KALLMANN), A., ii, 125; (ARRHENIUS), A., ii, 345.
- ionic equilibria of (STEARNS), A., ii, 420.
- molecular conductivity of (LORENZ), A., ii, 19.
- anomalous behaviour of (EBERT), A., ii, 113; (KENDALL), A., ii, 419.
- application of the gas laws to (BRONSTEN), A., ii, 113.
- ternary, conductivity of, in propyl alcohol (KRAUS and BISHOP), A., ii, 19.

- Electrolytes**, weak, ionisation of (SCHOORL), A., ii, 31.
- Electrolytic detector**, new (REICHSTEIN and ROTHCHILD), A., ii, 736.
- dissociation, relation between adsorption and (RAKUZIN), A., ii, 619.
- of salts (RABINOWITSCH), A., ii, 186, 187, 188.
- ions, theory of (LORENZ), A., ii, 19.
- oxidation of organic compounds (MÜLLER), A., ii, 469.
- reduction of organic compounds (CONANT, KAHN, FIESER, and KURTZ), A., ii, 547.
- Electromotive force of electrolytic cells** (GERKE), A., ii, 682; (WASASTJERNA), A., ii, 815.
- from the movement of an electrode in an electrolyte (PROCOPIU), A., ii, 112.
- at the junction of two liquids (FREUNDLICH), A., ii, 19.
- metallic conductivity (TAMMANN), A., ii, 255.
- of metallic compounds with electron conductivity (TRUMPLER), A., ii, 110.
- Electrons and atoms** (PEASE), A., ii, 757.
- binding of, by atoms (NICHOLSON), A., ii, 544.
- motion of, in gases (TOWNSEND and BAILEY), A., ii, 43, 836; (SKINKER), A., ii, 887.
- vibration of, without radiation resistance (FOKKER), A., ii, 106.
- free, behaviour of, towards gas molecules (WAHLIN), A., ii, 608.
- positive (LODGE), A., ii, 836.
- Electron gas**, entropy of (TOLMAN), A., ii, 18.
- Electron theory and photosynthesis** (DIXON and BALL), A., ii, 248.
- application of, to the structure of solids (THOMSON), A., ii, 355.
- Electro-osmosis** (ATEN; PRAUSNITZ), A., ii, 114.
- potential difference in (FREUNDLICH), A., ii, 189.
- Elements, genesis of** (NEUBURGER), A., ii, 365.
- structure of (ALLEN), A., ii, 758; (WACHTER), A., ii, 837.
- electronic structure of (COSTER), A., ii, 677; (DAUVILLIER), A., ii, 678.
- periodic classification of (BALAREFF), A., ii, 365.
- periodic system of the ions of (GRIMM), A., ii, 635.
- physical and chemical properties of (BOHR), A., ii, 363.
- Elements, properties of, and their compounds**, in relation to the dimensions of the component atoms and ions (CLARK), A., ii, 634.
- Röntgen spectra of (HJALMAR; COSTER), A., ii, 180; (SMEKAL), A., ii, 181.
- ionisation of the vapours of, at high temperatures (NOYES and WILSON), A., ii, 811.
- disintegration of (WÄCHTER), A., ii, 838.
- artificial disintegration of (RUTHERFORD), T., 400.
- disintegration of, by α -particles (RUTHERFORD and CHADWICK), A., ii, 682.
- entropy of, in relation to the periodic system (HEYZ), A., ii, 548.
- linear relationships in the periodic system of (BILTZ), A., ii, 494.
- relative volumes of (COLLINS), A., ii, 28.
- light, ultra-violet spectra of (MILLIKAN), A., ii, 100.
- radioactive. See Radioactive elements.
- Ellagic acid** in raspberry juice (KUNZ-KRAUSE), A., i, 210.
- Elsholtzic acid**, constitution of (ASHINA and KUWADA), A., i, 1047.
- Emilium** (LOISEL), A., ii, 15.
- Emulsin** (WILLSTÄTTER and OPPENHEIMER), A., i, 959.
- preparation and activity of (HELPERICH; WILLSTÄTTER and CSÁNYI), A., i, 390.
- decomposition of amygdalin by (GIAJA), A., i, 185.
- synthesis of cyanohydrins by means of (NORDEFELDT), A., i, 66, 1077.
- δ - and σ -Emulsins (ROSENTHALER), A., i, 600.
- Emulsions**, inhibition period in the separation of (NUGENT), A., ii, 625.
- aniline-water, coagulation of (BHATNAGAR), A., ii, 268.
- chromatic (HOLMES and CAMERON), A., ii, 269.
- oil-water, reversal of phases in (BHATNAGAR), A., ii, 204.
- Enantiomorphism of matter** (CONDELLI), A., i, 410.
- Energy of gaseous molecules** (PARTINGTON), A., ii, 614; (PLANCK), A., ii, 689.
- Enols**, azo-ester reaction with (DIELS), A., i, 774, 1194.
- Entropy**, calculation of (PLANCK), A., ii, 191.
- relation of probability to (LINHART), A., ii, 257, 738.

Entropy of elements (LEWIS and LATIMER), A., ii, 471.

in relation to the periodic system (HERZ), A., ii, 548.

of metals (LATIMER), A., ii, 814.

Enzymes (BIEDERMANN and RUEHA), A., i, 66.

chemistry of (v. EULER and MYRBÄCK), A., i, 1201.

periodicity of (SLUITER), A., i, 887.

combined action of (VANDEVELDE), A., i, 958.

adsorption of (JACOBY and SHIMIZU), A., i, 481; (RAKUZIN), A., i, 1199.

effect of acids, bases and salts on (BOKORNY), A., i, 93.

action of poisons on (SANTESSON), A., i, 1077.

effect of yeast extracts on the action of (ABDERHALDEN and WERTHEIMER), A., i, 796.

of blood (COMPTON), A., i, 392, 1087; (BACH and ZWIKOWA), A., i, 392; (PINCUSSEN), A., i, 393.

autolytic (BRADLEY), A., i, 896.

digestive, during foetal life (PORCHER and TAFERNOUX), A., i, 1088.

hydrolytic, action of (VAN LAER), A., i, 64.

oxidising (MARINESCO), A., i, 790, 1211.

oxidising and reducing, mechanism of the action of (BATELLI and STERN), A., i, 1077.

proteolytic, of malt (LUNDIN), A., i, 959. in albuminous urine (HEDIN), A., i, 609.

estimation of (PINCUSSEN), A., ii, 884.

of plants, velocity of reaction of (MAESTRINI), A., i, 507, 508.

Enzymes. See also:—

Amylase.

Carboligase.

Carboxylase.

Catalase.

Cellobiase.

Diastase.

Emulsin.

Esterase.

α -Glucosidase.

Hexosephosphatase.

Invertase.

Lactase.

Levidulinase.

Luciferase.

Maltase.

Mannanase.

α -Methylglucosidase.

Oxydase.

Pectinase.

Pepsin.

Enzymes. See also:—

Peroxydase.

Phosphatase-Phosphatase.

Protease.

Ptyalin.

Raffinase.

Reductase.

Rennin.

Saccharase.

Saccharophosphatase.

Succinodehydrogenase.

Takadiastase.

Tannase.

Tributyrynase.

Trypsin.

Urease.

Zymase.

Enzyme action (ARMSTRONG), A., i, 1078.

hypothesis of (EHRENBERG), A., i, 597.

asymmetric syntheses by means of (ROSENTHALER), A., i, 480.

Epicatechins, and their derivatives (FREUDENBURG, BOHME, and PURRMANN), A., i, 757.

Epichlorohydrin, equilibrium of acetic acid, water and (LEONE and BENELLI), A., ii, 744.

action of, on sodium phosphate (BAILLY), A., i, 980.

Equation, Dushman's, for unimolecular reactions (GARNER), A., ii, 358.

of state (SCHAMES), A., ii, 24;

(WOHL), A., ii, 117; (WEGSCHEIDER), A., ii, 192.

theory of (WAGNER), A., ii, 117.

van der Waals's, in relation to space-filling numbers (VAN LAAR), A., ii, 279.

Equilibria, the Le-Chatelier-Braun principle for (BENEDICKS), A., ii, 358.

and velocity of reaction (BRANDSMA), A., ii, 699.

of reactions, calculation of (CREIGHTON), A., ii, 426.

effect of catalysts on (CLARENS), A., ii, 436; (DURAND), A., ii, 701.

in mixed binary electrolytes (KRAUS), A., ii, 269.

in binary systems, influence of substitution on (KREMANN and MÜLLER; KREMANN, ODELGA, and ZAWODSKY), A., i, 131; (KREMANN, HOHL, and MÜLLER), A., i, 138;

(KREMANN and ODELGA), A., i, 159; (KREMANN and STRZELBA), A., i, 176; (KREMANN, HEMMELMAYR and RIEMER), A., i, 1010.

in heterogeneous systems (SYRKIN), A., ii, 699.

non-, uni- and bi-variant (SCHREINEMAKERS), A., ii, 430.

- Equilibria** of salt solutions (LE CHATELIER), A., ii, 555.
in systems of water and alcohols (PUSHIN and GLAGOLEVA), T., 2813.
chemical, statistics applied to (HERZFELD), A., ii, 269.
- Equilibrium** law for electrolytes (JABLZYŃSKI and WIŚNIEWSKI), A., ii, 190.
- Erbium** oxide, emission spectra of (NICHOLS and HOWES), A., ii, 411.
- Erdmann's salt**, constitution of (RIESENFELD and KLEMENT), A., ii, 853.
- Ergot**, active constituent of (SPIRO and STOLL), A., i, 47.
of diss and of oats, chemical composition of (TANRET), A., i, 504.
- Ergotamine** (SPIRO and STOLL), A., i, 47.
- Erucic acid**, iodine value of (HOLDE, WERNER, TACKE, and WILKE), A., ii, 723.
and its anhydride (HOLDE and WILKE), A., i, 317, 519, 713.
anilide and esters of (TOYAMA), A., i, 1111.
- Erythritol**, equilibrium in the system, water and (PUSHIN and GLAGOLEVA), T., 2820.
- Esters**, conductivity of mixtures of acids and (KENDALL and GROSS), A., ii, 33.
dielectric constants of (JACKSON), A., ii, 252.
velocity of saponification of (MC-COMBIE, SCARBOROUGH, and SETTLE), T., 2308.
influence of the constitution of the alcohol on the velocity of alkaline hydrolysis of (SMITH and OLSSON), A., ii, 701.
interchange of alcohol radicals in (SHIMOMURA and COHEN), T., 883, 2051.
action of hydrazine on the halogen derivatives of (HIRST and MACBETH), T., 2169.
aromatic, preparation of, in the presence of organic bases (SUZUKI and KAISHA), A., i, 543.
 α -disubstituted, reactions of, with magnesium alkyl haloids (LEBOIDE), A., i, 215.
- Esterase**, estimation of, in blood (BACH and ZUBKOWA), A., i, 392.
- Esterification** of organic acids (ZOLLINGER-JENNY), A., i, 914.
- Ethane**, chloro-derivatives, antiseptic action of (JOACHIMOGLU), A., i, 804.
hexachloro-, formation of, from chloropicrin (SILBERRAD), A., i, 2.
- Ethanesulphonic acid**, β -chloro-, lead salt (MANN and POPE), T., 600.
- Ethenylcarbinols** (SCHEIBLER and FISCHER), A., i, 1108.
- Ethers**, dithio-, formation and properties of (NAIK and AVASARE), T., 2592.
- Ethinylcarbinols**, tertiary, conversion of, into unsaturated ketones (MEYER and SCHUSTER), A., i, 556.
- α -Ethoxalyl-*N*-methylacetanilide (THIELEPAPE), A., i, 271.
- Ethoxyacetone**, spontaneous condensation of (DAUDEL), A., i, 432.
- 2-Ethoxyacridine**, 3:9-diamino-, 9-chloro-3-nitro-, and 3-nitro-9-amino- (FARBWERKE vorm. MEISTER, LUCIUS, & BRÜNING), A., i, 469.
- 9-Ethoxyacridine** (FARBWERKE vorm. MEISTER, LUCIUS, & BRÜNING), A., i, 468.
- 2-Ethoxy-9-antipyrineaminoacridine** (FARBWERKE vorm. MEISTER, LUCIUS, & BRÜNING), A., i, 468.
- 4'-Ethoxyazobenzene**, 4-bromo- (JACOBSEN), A., i, 591.
- Ethoxybenzanthrone**, and its salts (PERKIN and SPENCER), T., 477.
- 2-p-Ethoxybenzenesazoglyoxaline** (PUMMERER, BINAFFL, BITTNER, and SCHUEGRAF), A., i, 1198.
- 4-Ethoxybenzonitrile**, 3-nitro- (MATTAAR), A., i, 251.
- 4-Ethoxybenzophenone-4'-arsinic acid** (LEWIS and CHEETHAM), A., i, 187.
- p-Ethoxybenzylideneanthranilic acid** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- Ethoxybutyrolactone** (CABRIERE), A., i, 320.
- 8-Ethoxybutyronitrile** (BRUTLANTS), A., i, 924.
- 5-Ethoxy-2:3-diethyltetrahydrofuran** (CABRIERE), A., i, 319.
- 6'-Ethoxy-2:1'-diethylthioisocyanine iodide** (BRAUNHOLZ and MILLS), T., 2007.
- α -Ethoxydihydro-*o*- and -*p*-anethole, 8-nitro- (SCHMIDT, SCHUMACHER, BÄJEN and WAGNER), A., i, 733.
- 7-Ethoxy-7:12-dihydro- γ -benzophenarsazine** (LEWIS and HAMILTON), A., i, 188.
- 6-Ethoxy-1-p-dimethylaminophenyl-2-o-hydroxyphenyl-6-benziminasole** (JACOBSEN), A., i, 594.
- 6-Ethoxy-2:4-dimethylquinoline**, preparation of (PALKIN and HARRIS), A., i, 951.
- 4'-Ethoxydiphenylamine**, 5-bromo-2-amino-, and 5-bromo-2-nitro- (JACOBSEN), A., i, 591.
4-nitro-2-amino-, and 2:4-dinitro- (BLUM), A., i, 28.

- 5-Ethoxydiphenylamine**, 4'-bromo-2-amino-, and its derivatives (JACOBSEN), A., i, 591.
- 2-Ethoxy-9-ethanolaminoacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 468.
- 4'-Ethoxyhydrazobenzene**, 4-bromo- (JACOBSEN), A., i, 591.
- 2-Ethoxy-9-p-hydroxyphenylethylaminoacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 468.
- 6'-Ethoxy-5-methyl-2:1'-diethylthioisocyanine iodide** (BRAUNHOLTZ and MILLS), T., 2008.
- Ethoxymethylenemalononic acid**, menthyl ester (SHIMOMURA and COHEN), T., 2055.
- 5-Ethoxy-9-methyl-7-ethyl-4:5-dihydrouric acid**, 4-hydroxy- (BILTZ and HEIDRICH), A., i, 383.
- 4'-Ethoxy-4-methylhydrazobenzene**, isomerisation of (JACOBSEN), A., i, 591.
- β -Ethoxy- β -methyl- γ -propylhexan- γ -ol** (LEROIDE), A., i, 218.
- 6-Ethoxy-5-methyl-1-p-tolylbenzimidazole**, and its salts (JACOBSEN), A., i, 593.
- 5-Ethoxy-1:4-naphthaquinone**, 2:3-dibromo-8-hydroxy, and its sodium salt (WHEELER and ANDREWS), A., i, 354.
- 6-Ethoxy-2-p-nitrobenzylthiol-4-methylpyrimidine** (HORN), A., i, 375.
- β -Ethoxypentane**, α -chloro- δ -dibromo- (LESPIEAU), A., i, 620.
- 5-Ethoxy- Δ -pentene**, ϵ -chloro- (LESPIEAU), A., i, 619.
- 9-Ethoxy-10-phenanthryl peroxide** (GOLDSCHMIDT and SCHMIDT), A., i, 1150.
- 3-Ethoxyphenyl-5-anilinomethylene-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- 4'-Ethoxyphenylaziminobenzene**, 4-nitro- (BLOM), A., i, 28.
- 3-Ethoxy-2-phenylbenzopyrylium salts**, and 7-hydroxy- (PRATT and ROBINSON), T., 1580.
- 1-p-Ethoxyphenylbenzotriazole**, 6-bromo- (JACOBSEN), A., i, 592.
- 2-Ethoxy-9-phenylhydrazinoacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 469.
- α -p-Ethoxyphenylhydrazinopropionic acid** (JACOBSEN), A., i, 596.
- 3-p-Ethoxyphenyl-5-hydroxymethyl-oxazolidine** (BERGMANN, ULPTS, and CAMACHO), A., i, 1181.
- 1-p-Ethoxyphenyl-6-methylbenzotriazole** (JACOBSEN), A., i, 592.
- 3-p-Ethoxyphenyloxazolidine** (BERGMANN, ULPTS, and CAMACHO), A., i, 1182.
- p-Ethoxyphenylstibinic acid** (SCHMIDT), A., i, 1204.
- α -Ethoxypropylbenzene**, β -bromo- (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 826.
- Ethoxyquinaldines**, preparation and derivatives of (BRAUNHOLTZ), T., 170.
- d-Ethoxysuccinamide** (MCKENZIE and SMITH), T., 1361.
- 1:6-Ethoxy-m-tolyl-6-methylbenzimidazole**, and its salts (JACOBSEN), A., i, 593.
- Ethyl alcohol**, preparation of (PASCAL), A., i, 3; (DAMIENS), A., i, 1105. history of (v. LIPPMANN), A., i, 102. density and refractive index of mixtures of ether, water, and (SANFOURCHE and BOUTIN), A., i, 709. vapour pressure of dilute solutions of (THOMAS), A., ii, 321. equilibrium of camphor, water, and (SCHERINGA), A., ii, 535. equilibrium of carbon disulphide with (MCKELVY and SIMPSON), A., ii, 271. equilibrium of carbon disulphide, water, and (SCHOORL and REGENBOGEN), A., i, 419. equilibrium of chloroform, water, and (SCHOORL and REGENBOGEN), A., i, 215. equilibrium of water and, with aromatic hydrocarbons (ORMANDY and CRAVEN), A., i, 215. solubility of substances in mixtures of water and (WRIGHT), T., 2251. surface tension of mixtures of water and (BIRCUMSHAW), T., 887. catalytic dehydration and esterification of (GILFILLAN), A., i, 709. inflammability of mixtures of petrol and (BOUSSU), A., ii, 557. compounds of cresols and (BREL and SCHWEBEL; C. and W. v. RECHENBERG), A., i, 932. detection of acetone in (HARRISSON), A., ii, 667. estimation of, in blood (WIDMARK), A., ii, 789.
- Ethyl ether**, physical constants of (TIMMERMANS, VAN DER HORST and ONNES), A., ii, 253. density and refractive index of mixtures of ethyl alcohol, water, and (SANFOURCHE and BOUTIN), A., i, 709. heat of solution and solubility of, in sodium chloride (McKEOWN), A., ii, 552.

- Ethyl ether**, vapour pressure of mixtures of bromoform and (DOLEZALEK and SCHULZE), A., ii, 118.
 properties of mixtures of sulphuric acid, water, and (POUND), T., 941.
 partition of compounds between water and (SMITH), A., ii, 270.
 autoxidation of (CLOVER), A., i, 619.
 compounds of cresols and (BERL and SCHWEBEL; C. and W. V. RECHENBERG), A., i, 932.
- Ethyl ether**, dichloro-, action of, on magnesium allyl bromide (LESPERAU), A., i, 619.
αα'-dichloro- (GRIGNARD and PURDY), A., i, 802.
ββ'-dichloro- (KAMM and WALDO), A., i, 105.
- Ethyl β-hydroxy-β-phenylethyl ether** (DETÈUF), A., i, 327.
 iodide, compounds of, with triethylene tetrasulphide and with ethylene mercaptide nitrite (RÅY), T., 1282.
 mercaptan, β-chloro- (BENNETT), T., 2145.
 selenate (MEYER and WAGNER), A., i, 620.
 sulphate, preparation of (DREYFUS), A., i, 517; (DAMIENS), A., i, 1105.
- Ethylacetonylmalonic acid**, and its ethyl ester, and their phenylhydrazones (GAULT and SALOMON), A., i, 428.
- C-Ethylacetylacetones**, selenium (MORGAN, DREW, and BARKER), T., 2460.
- α-Ethyl β-aldehydopropionic acid**, and its derivatives (CARRIERE), A., i, 320.
- Ethylallylaniline N-oxide**, picrate of (MEISENHEIMER and GREESKE), A., i, 334.
- Ethylaminesulphonitrile**, and its salts (OLIVERI-MANDALÀ), A., i, 1008.
- 9 Ethylaminocacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 469.
- p-Ethylaminobenzoic acid**, *p*-β-hydroxy- (ALTWEGG and LANDRIVON), A., i, 1022.
- 4-Ethylaminobenzonitrile**, 3-nitro- (MATTAR), A., i, 251.
- 4-Ethylamine-5-carboxybenzeneazobenzene-4'-arsinic acid**, and its sodium salt (JACOBS and HEIDELBERGER), A., i, 75.
- 6-Ethylamino-5-formylethylaminouracil** (BLITZ and BÜLOW), A., i, 384.
- 2-Ethylaminophenanthraquinone** (BRASS and FIEBER), A., i, 356.
- o'-Ethylaminophenyl-p-tolylsulphone** (HALBERKANN), A., i, 1133.
- 2-Ethylamino-p-toluic acid**, 3:5-dinitro-, methyl ester (GIUA), A., ii, 534.
- Ethylammonium**, ruthenipentabromide (GUTBIER and KRAUSS), A., i, 16.
- Ethylaniline**, 5-bromo-2:4-dinitro- (GIUA and ANGELETTI), A., i, 649.
 2-mono- and 2:6-di-bromo-4-nitro- (KHARASCH and JACOBSON), A., i, 189.
- Ethylazide** (STAUDINGER and HAUSER), A., i, 69.
- 2-Ethyl-4:5-benzointhane** (MAYER and SIFGLITZ), A., i, 743.
- 2-Ethyl-4:5-benzoindan-1-one** (MAYER and SIFGLITZ), A., i, 743.
- 2-Ethylbenzothiazolonylbenzothiazolymethane** (MILLS), T., 464.
- 2-Ethylbenzoxazole** (SKRAUP and MOSER), A., i, 575.
- O-Ethylisoburet** (MADELUNG and KERN), A., i, 439.
- 5-Ethyl-5-n-butylbarbituric acid** (DOX and YODER), A., i, 808.
 hypnotic power of (CARNOT and TIFENEAU), A., i, 900.
- α-Ethylbutylcarbamide**, bromo- (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 993.
- α-Ethylbutylphosphinic acid**, α-hydroxy-, lead salt (CONANT, MACDONALD, and KINNEY), A., i, 186.
- α-Ethyl-β-butyrolactone** (JOHANSSON and HAGMAN), A., i, 426.
- N-Ethylcarbazole-3-azoacetoacetic acid**, ethyl ester (MORGAN and READ), T., 2716.
- N-Ethylcarbazole-3-diazonium salts** (MORGAN and READ), T., 2715.
- Ethylcarbonatoacetoneitrile**, and its derivatives (SONN and FALKENHEIM), A., i, 1164.
- 5-Ethylcarbonato-3:4-dimethoxybenzaldehyde** (SPATH and RÖDER), A., i, 853.
- 5-Ethylcarbonato-3:4-dimethoxybenzoic acid** (SPATH and RÖDER), A., i, 853.
- 4-Ethylcarbonato-3:5-dimethoxybenzylmalonic acid** (SPATH and RÖDER), A., i, 852.
- 4-Ethylcarbonato-3:5-dimethoxyphenylacetaldoxime** (SPATH and RÖDER), A., i, 852.
- β-4-Ethylcarbonato-3:5-dimethoxyphenylethylamine**, and its salts (SPATH and RÖDER), A., i, 852.
- β-4-Ethylcarbonato-3:5-dimethoxyphenylpropionic acid**, and its derivatives (SPATH and RÖDER), A., i, 852.
- 4- and 5-Ethylcarbonatodimethoxystyrenes**, α-nitro- (SPATH and RÖDER), A., i, 852.

- 7-Ethylcarbonato-6:8 dimethoxy-1:3:3:4-tetrahydroisoquinoline** and its hydrochloride (SPATH and RÖDER), A., i, 852.
- 2-Ethylcarbonato-1:3-dimethylcyclobuten-4-one-3-carboxylic acid**, methyl ester (DIECKMANN and WITTMANN), A., i, 1156.
- Ethylcarbonatophenylalanine** (CURTIUS and SIEBER), A., i, 722.
- α -Ethylcarbonatoresacetophenone** (SONN and FALKENHEIM), A., i, 1164.
- 1-[Ethylcarbonatoisovanillyl]-6:7-dimethoxy-3:4-dihydroisoquinoline**, and its hydrochloride (SPATH and LANGE), A., i, 569.
- Ethyl-cellulose**, de-polymerisation of (HESS and WITTELSBACH), A., i, 116.
- Ethylconiine**, hydroxy-, and its derivatives (V. BRAUN, BRAUNSDORF, and RATH), A., i, 760.
- O-Ethylcyanoisocarbamide**, and its sodium salt (MADELUNG and KERN), A., i, 439.
- Ethylidibutylarsine**, and its salts (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- Ethylidihydrocupreicine** (*optotoxin*) sulphate (HEIDELBERGER and JACOBS), A., i, 678.
- d-O-Ethylidihydrocupreicinol**, and its hydrochlorides (HEIDELBERGER and JACOBS), A., i, 674.
- Ethylidihydrocupreine ethyl bromide** (HEIDELBERGER and JACOBS), A., i, 673.
- oxide, and its dinitrate (SPEYER and BECKER), A., i, 675.
- N-Ethylidihydroisoidole**, β -hydroxy-, and its picrate (V. BRAUN, BRAUNSDORF, and RATH), A., i, 760.
- Ethylidihydronaphthalene** (DE POMMEREAU), A., i, 828.
- 8-Ethyl-7:8 dihydrophenalene** (MAYER and SIEGLITZ), A., i, 741.
- 8-Ethyl-7:8-dihydrophenalene-9** (MAYER and SIEGLITZ), A., i, 741.
- N-Ethylidihydroquinicine hydrochloride** (HEIDELBERGER and JACOBS), A., i, 673.
- d- and l-N-Ethylidihydroquinicinol**, and their derivatives (HEIDELBERGER and JACOBS), A., i, 674.
- α -Ethyl- $\beta\beta$ -dimethylglutaric acid**, and its derivatives (KON and THORPE), T., 1800.
- 4-Ethylidiphenyl** (V. AUWERS and JÜLICHER), A., i, 842.
- Ethylid-n-propylarsine**, and its salts (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- Ethylene**, preparation of, by reduction of acetylene (CHEVALIER and BOURCET), A., i, 801.
- compressibility of (BATUECAS), A., ii, 618.
- absorption of, by sulphuric acid (DA-MIENS), A., i, 1105.
- catalytic hydrogenation of, with nickel (RIDEAL), T., 309.
- formation of butadiene from (ZANETTI, SUYDAM, and OFFNER), A., i, 977.
- reaction of nitrogen and, under the electric discharge (MIYAMOTO), A., i, 418.
- cyanohydrin. See Propionitrile, β -hydroxy-.
- dicarbamidophenyl ether (SPECKAN), A., i, 580.
- halohydrins, esters of (ALTWEGG and LANDRIVON), A., i, 315.
- chloro-derivatives, antiseptic action of (JOACHIMOGLU), A., i, 304.
- di-, tri-, and tetra-chloro-*, iodine numbers of (MARGOSCHES and BARU), A., ii, 235.
- diiodo-*derivatives (KAUFMANN), A., i, 214; (LATIERS), A., i, 314.
- Ethylenediamine**, compounds of, with metallic salts (PETERS), A., i, 48.
- Ethylenediphenyldiarsinous acid** (QUICK and ADAMS), A., i, 601.
- Ethylene glycol**, *monothio-*, and its metallic derivatives (BENNETT), T., 2139.
- derivatives of (ROSEN and REID), A., i, 420.
- dithio-*, potassium salt, compound of chloropierin and (RAY and DAS), T., 326.
- Ethylene mercaptan**, compound of mercuric nitrite and (RAY), T., 1282.
- Ethylene oxide**, physical properties of (MAASS and BOOMER), A., i, 912.
- hydrate of (MAZZUCHELLI and ARMENANTE), A., i, 620.
- 9-Ethylfluorene**, 2:7-*di*bromo-9-hydroxy-, and 9-chloro-2:7-*di*bromo- (SIEGLITZ and JASSOV), A., i, 821.
- γ -Ethylglutaconic acid**, α -cyano-, ethyl ester (INGOLD, PERRIN, and THORPE), T., 1782.
- 4-Ethylglyoxaline**, β -amino- (*histamine*), physiological action of (SCHENCK), A., i, 498.
- in the intestines (MEAKINS and HARRINGTON), A., i, 396; (GERARD), A., i, 790.
- 1-Ethylhexahydropyridine-3-carboxylic acid**, ethyl ester (WOLFFENSTEIN), A., i, 950.

- α -Ethylcyclohexane-1:1-diacetic acid**, and $\alpha\alpha$ -dicyano-, ω -imides of (KON and THORPE), T., 1800.
- γ -Ethylhexane- γ -diol**, and its derivatives (CARRIÈRE), A., i, 320.
- Ethylhydrocupreidine**, salts of tropic acid with (KING and PALMER), T., 2583.
- Ethylhydrocupreine**, and its oxide sulphate, and chloro- (VEREINIGTE CHININFABRIKEN ZIMMER & Co.), A., i, 948.
- salt of tropic acid with (KING and PALMER), T., 2583.
- Ethylhydrocupreine**, nitro-, and its sulphonic acid (BOEHRINGER and SÖHNE), A., i, 46.
- Ethylidenemalonie acid**, ethyl ester, preparation of γ -alkylidene derivatives of (HIGGINBOTHAM and LAFWORTH), T., 2823.
- 2-Ethylindole-3-carboxylic acid**, 2- β -amino-, and its picrate (ASAHINA and FUJITA), A., i, 47.
- Ethylmalonasic acid** (CURTIUS and SIEBER), A., i, 723.
- Ethylmenthol** (OGATA and MIYASHITA), A., i, 844.
- Ethylmethylaniline**, β -chloro- (CLEMO and PERKIN), T., 648.
- α -Ethyl- β -naphthacinehoninylpropionic acid**, and its derivatives (CARRIÈRE), A., i, 320.
- 4-Ethylnitroaminobenzonitrile**, 3:5-dinitro- (MATTAAR), A., i, 251.
- γ -Ethyl- γ -octandiol**, and its derivatives (CARRIÈRE), A., i, 319.
- Ethylcyclopentamethylenearsine**, and its salts (STEINKOPF, DONAT, and JAEGER), A., i, 996.
- γ -Ethylpentane- $\alpha\beta$ -dicarboxylic acid**, $\alpha\beta$ -dicyano- γ -hydroxy-, and γ -hydroxy-, lactone (BIRCH and THORPE), T., 1826.
- γ -Ethylpentane- $\alpha\alpha\beta$ -tricarboxylic acid**, γ -hydroxy-, lactone (BIRCH and THORPE), T., 1826.
- γ -Ethylpentan- α -ol**, γ -chloro- (MOUREU and BARRETT), A., i, 5.
- Ethyl pentyl ketone**, α -chloro-, and its semicarbazone (DETÈUR), A., i, 327.
- Ethylphenetidide**, β -hydroxy-, and its hydrogen oxalic ester (BERGMANN, ULFTS, and CAMACHO), A., i, 1182.
- Ethylphthalimide**, hydroxy- (DERSIN), A., i, 142.
- N-Ethylpiperidine**, β -hydroxy-, and their derivatives (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- O-Ethylpivalylacetone** (MORGAN and DREW), T., 938.
- Ethyl-n-propylcyanarsine** (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- N-Ethylpyrrolidine**, β -hydroxy-, and its derivatives (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 760.
- N-Ethylquinisine** hydrochloride (HEIDELBERGER and JACOBS), A., i, 673.
- 8'-Ethylisoquinoline-red** (HARRIS and POPE), T., 1032.
- Ethylsuccinic acid**, derivatives of (CARRIÈRE), A., i, 819.
- β -Ethylsulphonylpropionic acid**, β -bromo-, and its salts (FITGER), A., i, 108.
- Ethylsulphonylsuccinic acid**, and α -bromo-, and its salts (FITGER), A., i, 108.
- Ethylsulphoxysuccinic acid**, and its salts (FITGER), A., i, 108.
- N-Ethyltetrahydroisoquinoline**, β -hydroxy-, and its salts (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 761.
- 2-Ethyl-5-thienyl methyl ketone** semicarbazone (STEINKOPF and HEROLD), A., i, 851.
- β -Ethylthielaerylic acid** (FITGER), A., i, 108.
- β -Ethylthiolmaleic acid**, α -bromo-, and its barium salt and anhydride (FITGER), A., i, 108.
- 2-Ethylthiol-3-phenyl-5-anilinomethylene-4-glyoxalone** (DAINS, THOMPSON, and ASENDORF), A., i, 1186.
- Ethylthiolsuccinic acids**, stereoisomeric, and their salts (FITGER), A., i, 107.
- 3-Ethylthiophen** mercurichlorides (STEINKOPF and HEROLD), A., i, 850.
- Ethyltrigonelline** hydrochloride (MEERCK), A., i, 949.
- Eucalyptus** oil, aromatic aldehydes in (PENFOLD), T., 266.
- thymol, menthone and menthol from (SMITH and PENFOLD), A., i, 31.
- Eudalene**, and its salts (RUZICKA, MEYER, and MINGAZZINI), A., i, 560.
- isoEugenol**, derivatives of (RASTELLI), A., i, 1010.
- n- and iso-Eugenols**, ultra-violet absorption spectra of (DURRANS), A., ii, 6.
- isoEugenyl methyl ether**, β -nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.
- Euphylline**, coagulation of blood by (MEISSNER), A., i, 193.
- Europium oxide**, density and molecular volume of (FRANDTL), A., ii, 379.
- Euxanthogen**. See Mangiferin.
- Explosion** of mixed gases (CAMPBELL), T., 2483.
- Explosions**, thermodynamical theory of (HENDERSON and HASSE), A., ii, 207.
- of organic dust (BEYERSDORFER), A., ii, 749.

Explosives, sensitiveness of (EGGERT), A., ii, 133.

high, velocity of decomposition of, in a vacuum (FARMER), T., 174.

Extraction of colloids by Winkelblech's method (CHARITSCHKOV), A., ii, 826.

Extraction apparatus, chloroform (SCHAAP), A., ii, 797.

funnel (WOLFFRAM), A., ii, 221.

micro-chemical (LAQUER), A., ii, 325.

perforation, for extraction of non-miscible liquids (FAYOLLE and LOR-MAND), A., ii, 759.

F.

Fæces, human, hydrogen-ion concentration of (ROBINSON), A., i, 899.

Fasting, effect of, on intestines (DOM-BROWSKI and KOZLOWSKI), A., i, 606.

Fat, Illipé. See Illipé fat.

Fats, production of, from proteins (ATKINSON, RAPPORT, LUSK, SÖDERSTROM, and EVENDEN), A., i, 966.
enzymic synthesis of (SPIEGEL), A., i, 694.

chemistry of (ANDRÉ), A., i, 713.

effect of radiations on the hydrolysis of (PINCUSSEN and ANAGNOSTU), A., i, 485.

catalysis in the hydrolysis of (BRINER and TRAMPLER), A., ii, 208.

saponification of (NORRIS and McBAIN), T., 1362; (LANGTON), A., i, 982.

nutritive value of (TAKAHASHI), A., i, 699.

relations between carbohydrates and (MULLER), A., i, 486.

degradation of, in the body (BAER), A., i, 490.

excretion of (HILL and BLOOR), A., i, 968.

analytical constants of, from beef bone and neat's foot oil (ECKART), A., i, 967.

partly hydrolysed, analysis of (FAHRION), A., ii, 401.

animal, detection of vegetable oils in (MUTTELET), A., ii, 168.

estimation of the acetyl value of (LEYS), A., ii, 167.

estimation of the iodine-bromine number of, without using potassium iodide (WINKLER), A., ii, 534.

Feeding stuffs, estimation of amino-acids in (HAMILTON, NEVENS and GRINDLEY), A., ii, 92.

Fehling's solution, behaviour of, in light (BENRATH and OBERBACH), A., i, 108.

Felspar, constitution of (DENAEYER), A., ii, 451.

potash, melting of (MOREY and BOWEN), A., ii, 577.

β -Fenchosamphorol, and its acetate (KOMPPA and ROSCHIER), A., i, 1167.

Fenchone, derivatives of (MAXWELL), A., i, 753.

Fermentation, activators of (v. EULER and KARLSSON), A., i, 972; (LINDBERG), A., i, 1219.

acetic, effect of radioactive substances on (LABORDE, JALOUSTRE, and LEULLIER), A., i, 1219.

acetone and butyl alcohol, of carbohydrates (ROBINSON), A., i, 971.

alcoholic, nature of (WARDEN), A., i, 307.

rôle of acetaldehyde in (FERNBACH and SCHOEN), A., i, 203.

acceleration of, by animal and plant extracts (FRÄNKEL and HAGER; FRÄNKEL and SCHARF), A., i, 409.

in presence of calcium carbonate (KERR and ZECKENDORF), A., i, 305.

co-enzyme of, and its separation from vitamin-B (THOLIN), A., i, 305.

stimulants for (NEUBERG, REINFURTH, and SANDBERG), A., i, 306; (NEUBERG and SANDBERG), A., i, 408.

influence of thyroxin on (TOMITA), A., i, 972.

in presence of urea (SANDBERG), A., i, 502.

by means of yeast-cells (ABDERHALDEN), A., i, 92.

butyric acid, nature of (NEUBERG and ARINSTEIN), A., i, 91.

lactic acid, effect of lactic acid on (HOLWERDA), A., i, 611.

by micro-organisms, formation of acetaldehyde in (NEUBERG and COHEN), A., i, 304.

yeast. See Yeast.

without yeast (BAUR and HERZFELD), A., i, 93; (HAU), A., i, 307.

Ferric salts. See under Iron.

Ferrioxalic acid, salts, preparation of (BURROWS and TURNER), A., i, 916.

potassium salt, crystallography and optical properties of (KNAOGS), T., 2070.

Ferrochromium, estimation of chromium in (KELLEY and WILEY), A., ii, 164.

Ferrocyanides, titration of metals with (TREADWELL and CHERVET), A., ii, 786.

- Ferrocyanides**, estimation of electro-volumetrically (MULLER and LAUTERBACH), A., ii, 795.
 estimation of, volumetrically, with permanganate (KOLTHOFF), A., ii, 587.
- Ferroflicon**, toxic properties of (KURNAKOV and URASOV), A., ii, 845.
- Ferrotungsten**, analysis of (LOSANA and CAROZZI), A., ii, 661.
- Ferrous salts**. See under Iron.
- Ferrovandium**, estimation of chromium and vanadium in (KELLEY, WILEY, BOHN, and WRIGHT), A., ii, 89.
- Fertilisers**, estimation of nitrogen in (MACH and SINDLINGER), A., ii, 783.
- isoFerulic acid**. See 4-Methoxycinnamic acid, 3-hydroxy-.
- Fibres**, textile, behaviour and structure of (NODDER), A., ii, 791.
- Fibrin**, swelling of, by acids (SOMOGYI), A., i, 182.
 estimation of, in blood (GRAM), A., ii, 240.
- Fibrinogen**, estimation of, in blood (GRAM), A., i, 288; (LEENDERTZ and GROMELSKI), A., ii, 798.
- Fibrinolysis** (ROSENMAN), A., i, 596.
- Ficus fulva**, stearic acid from the latex of (ULTER), A., i, 1100.
- Films**, thin, properties and molecular structure of (ADAM), A., ii, 687.
- Filters**, membrane, use of, in analysis (JANDER), A., ii, 457.
- Filter paper**, apparatus for testing the permeability of (GRIFFIN and PARISH), A., ii, 309.
- Fisetole**. See Resacetophenone, ω -hydroxy-.
- Fish**, physiology of the respiration of (POWERS), A., i, 286.
 fresh water, respiratory exchange in (GARDNER, KING, and POWERS), A., i, 1084.
 sperm, phosphatides of (SANO), A., i, 699.
- Flagstaffite**, identity of, with terpin hydrate (GUILD), A., ii, 76.
- Flame**, propagation of, in mixtures of air with gases (PAYMAN and WHEELER), T., 363; (WHITE), T., 1244, 1688, 2561.
 luminescence of ions of, in the air spark (GREINACHER), A., ii, 250.
 inverted air, gaseous products from (HOFMANN and WIL), A., i, 1107.
- Flash-point**, apparatus for determination of (WRIGHT), T., 2247.
- Flavanol**, and its derivatives (FREUDENBERG and ORTHNER), A., i, 758.
- Flavanone**, reduction of (FREUDENBERG and ORTHNER), A., i, 757.
- Flavonols**, 2- and 4-chloro-, and their acetyl derivatives (MINTON and STEPHEN), T., 1602.
- Flax**, distinction between hemp and (NODDER), A., ii, 791.
- Flocculation** by electrolytes (FREUNDLICH and SCHOLZ), A., ii, 828.
- Flour**, relation between the catalytic activity of, and its grade of extraction (FERNANDEZ and PIZARROSO), A., ii, 94.
 catalase of (MERL and DAIMER), A., i, 184.
- Flowers**, manganese in (WESTER), A., i, 309.
- Fluoran**, 2:4:5:7-tetra-amino-1:8-dihydroxy-, diphthalyl derivative of (MUKERJI), T., 549.
- Fluorene**, pyrogenic transformation of (DZIEWONSKI and SUSZKO), A., i, 730.
 catalytic oxidation of (WEISS and DOWNS), A., i, 27.
- Fluorene**, 2:7-dibromo-9-hydroxy- (SIEGLITZ and JASSOY), A., i, 821.
- Fluorene series** (SIEGLITZ and JASSOY), A., i, 820.
- Fluorene-9-acethydrazide**, 2:7-dibromo-, and its hydrazones (SIEGLITZ and JASSOY), A., i, 821.
- 9-Fluorene-9-acetic acid**, 2:7-dibromo-9-hydroxy-, and its ethyl ester (SIEGLITZ and JASSOY), A., i, 820.
- Fluorenoneanil-p'-carboxylic acid** (REDELLEN), A., i, 146.
 methyl ester (REDELLEN and DANILOF), A., i, 148.
- Fluorenonebenzoylhydrazone** (REDELLEN), A., i, 146.
- Fluorenonebenzoylimide** (REDELLEN and DANILOF), A., i, 148.
- Fluorenonemethylimide** (REDELLEN), A., i, 146.
- Fluorenylidene-p-aminodiphenyl** (REDELLEN), A., i, 146.
- Fluorenyl-9-methylurethane**, 2:7-dibromo- (SIEGLITZ and JASSOY), A., i, 821.
- Fluorescein**, preparation and derivatives of (FISCHER and BOLLMANN), A., i, 936.
- Fluorescein**, dithiol- (WATSON and DUTT), T., 1942.
- Fluorescence** and photochemistry (WOOD), A., ii, 334; (WEIGERT), A., ii, 681.
 disturbance of the power of, by light (PRINGSHEIM), A., ii, 602.
 by flame excitation (NICHOLS and WILBER), A., ii, 806.
 estimation of (LEWIS), A., ii, 334.

- Fluorescent substances**, action of, on cells (JODLBAUER and HAFNER), A., i, 94.
- Fluorides**. See under Fluorine.
- Fluorine**, spectrum of (SMYTHE), A., ii, 99.
- detection and estimation of (SMITT), A., ii, 716.
- Hydrofluosilicic acid**, estimation of, by conductometric titration (KANÓ), A., ii, 719.
- Fluorine organic compounds** :—
- Fluorides**, organic (KRAUSE and POHLAND), A., i, 644.
- aliphatic (SWARTS), A., i, 101.
- Fluorine detection** :—
- detection of (FETKENHEUER), A., ii, 655.
- detection of, in Spanish grapes (PONDAL), A., i, 1100.
- Fluorites**, colour of (SAKAO and HIROSE), A., ii, 779.
- Foods**, phytin in (ARBENZ), A., i, 1102.
- tryptophan content of (IDE), A., i, 414.
- microchemical analysis of (LÜHRIG), A., ii, 534.
- detection of β -naphthol in (KINUGASA and TATSUNO), A., ii, 400.
- detection of sulphites in (CHAPMAN), A., ii, 520.
- detection of vitamins in (DRUMMOND and WATSON), A., ii, 596.
- estimation of the alkalinity of the ash of (PFYL), A., ii, 586.
- See also Feeding stuffs
- Formaldehyde**, preparation of (TRAUN'S FORSCHUNGSLABORATORIUM), A., i, 522, 528; (BLAIR and WHEELER), A., i, 917.
- and its polymerides (LUSH), A., i, 625.
- production of, by the action of ozone on hydrocarbons (WHEELER and BLAIR), A., i, 1105.
- water-soluble form of (COHN), A., i, 918.
- alcoholic fermentation of (MÜLLER), A., i, 110, 809.
- dehydroxidation of (MÜLLER), A., i, 109.
- formation of hydrocyanic acid by (FOSSE and HIEULLE), A., i, 523.
- action of phosphorus trihydride on (HOFFMAN), A., i, 8.
- as an intermediate product of assimilation (JACOBY), A., i, 502.
- effect of, on bacteria and on spores (HALLER), A., i, 408.
- in urine (VOIT), A., i, 1218.
- Formaldehyde hyposulphite**, preparation of, and its zinc compound (MALVEZIN, RIVALLAND and GRAND-CHAMP), A., i, 8.
- detection of, with phenols (PFYL, REIF, and HANNER), A., ii, 94.
- detection of, in urine (STEFF), A., ii, 793.
- Formaldehydesulphoxylic acid**, sodium salt, as an antianaphylactic (BRODIN and HUCHET), A., i, 90.
- Formamide**, synthesis of (MEYER and ORTHNER), A., i, 529.
- Formamides**, reactions of (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- Formic acid**, anhydrous, preparation of (BOSWELL and CORMAN), A., i, 712.
- formation of, during the decomposition of dextrose (WATERMAN and VAN TUSSEN BROEK), A., i, 433.
- and its salts, electrical conductivity of mixtures of (KENDALL, ADLER, and DAVIDSON), A., ii, 126.
- partition of, between water and organic solvents (GORDON and REID), A., ii, 824.
- catalytic decomposition of, on platinum and silver (TINGEY and HINSHELWOOD), T., 1668.
- catalytic dehydroxidation of (MÜLLER and SPONSEL), A., ii, 558.
- thermal decomposition of (HINSHELWOOD, HARTLEY, and TOPLEY), A., ii, 274.
- action of, on hexamethylenetetramine salts (SOMMELET and GUIOTH), A., ii, 333.
- velocity of reaction between manganates and (HOLLUTA), A., ii, 700.
- mechanism of the reduction of permanganate by (HOLLUTA), A., ii, 448; (HOLLUTA and WEISER), A., ii, 628.
- compounds of molybdic acid and (JAKÓB), A., i, 712.
- condensation of, with 3-oxythionaphthen and with phloroglucinol (SCHWENK), A., i, 153.
- salts, denitrification with (GROENEWEGE), A., i, 971.
- alkali salts, solubility of, in water (SINGWICK and GENTLE), T., 1837.
- metallic salts, freezing point curves of, in formic acid solution (ADLER), A., i, 5.
- lithium salt, preparation of, and its use in the production of alcohols and ketones (BADISCHE ANILIN- & SODA-FABRIK), A., i, 213.

- Gallaldehyde**, preparation of, and its derivatives (ROSENMUND), A., i, 1030.
- Gallium**, arc spectrum of (UHLER and TANCH), A., ii, 677.
- Gallotannin**, constitution of (NIERENSTEIN), A., i, 266.
- Galloylæovoglucosan** (KARRER and SALOMON), A., i, 267.
- Gallylamine hydrochloride** (ROSENMUND and PFANNKUCH), A., i, 1031.
- Gambier - catechin - carboxylic acids** (NIERENSTEIN), T., 26.
- Gas**, detonating, ignition of (MITSCHERLICH), A., ii, 358.
illuminating, estimation of sulphur in, with perhydrol (KLEMMER), A., ii, 224.
- Gases**, properties of (WÄCHTER), A., ii, 476.
ultra-red spectra of (SCHAEFFER), A., ii, 727.
motion of electrons in (TOWNSEND and BAILEY), A., ii, 43.
behaviour of free electrons towards the molecules of (WAHLIN), A., ii, 608.
resonance and ionisation potentials of (BOUCHER), A., ii, 608.
disappearance of, in the electric discharge (GENERAL ELECTRIC CO.), A., ii, 417; (NEWMAN), A., ii, 546.
specific heat of (INGOLD and USHERWOOD), T., 2286; (MEWES), A., ii, 737.
determination of high temperatures by effusion of (YAMAGUCHI), A., ii, 470.
thermo-compression of (DELEUVER), A., ii, 684.
compressibility of, at low temperature (GUYE and BATUECAS), A., ii, 617; (BATUECAS), A., ii, 618.
contraction of flasks in the determination of density of (MOLES and MIRAVALLS), A., ii, 617.
solubility of, in liquids (NEUHAUSEN and PATRICK), A., ii, 264; (NEUHAUSEN), A., ii, 621.
adsorption of, calculation of (BERÉNYI), A., ii, 426.
kinetic theory of adsorption of (HENRY), A., ii, 740.
adsorption of, by charcoal (GUSTAFER), A., ii, 479.
adsorption of, by charcoal and lignite (MCLEAN), A., ii, 352.
adsorption of, by crystals (SEELIGER and LAPKAMP), A., ii, 30.
application of the quantum theory to (SCHAPOSCHNIKOV), A., ii, 839.
relation between the internal friction and chemical constitution of (SCHMIDT), A., ii, 278.
- Gases**, energy of the molecules of (PARTINGTON), A., ii, 614.
containing hydrogen, molecular structure of (RANKINE), A., ii, 635.
ignition of (MASON and WHEELER), T., 2079.
of blood, analysis of (MEIER and KRONIG), A., i, 191.
diatomic, chemical constants of (PARTINGTON), A., ii, 889.
mixed, explosion of (CAMPBELL), T., 2488.
ignition of (PAYMAN and WHEELER), T., 363; (WHITE), T., 1244, 1688, 2561; (MASON and WHEELER), T., 2079.
monatomic, calculation of the chemical constants of (BRODY), A., ii, 191.
natural, collection and examination of (HENRICH and PRELL), A., ii, 858.
volcanic, analyses of (ALLEN), A., ii, 219.
estimation of, in metals (SIMONS), A., ii, 719.
separation of volatile substances from (BERL and SCHWEBEL), A., i, 932.
- Gas analysis**, absorption by pyrogallol in (HOFFMANN), A., ii, 582.
receiver for (GUTHRIE), A., ii, 158.
- Gas analysis apparatus** (VAN SLYKE), A., ii, 158; (MOSER and BRUKL), A., ii, 393; (BLAIR and WHEELER; MOSER), A., ii, 519; (SCHALLER and BERNDT), A., ii, 862.
- Gas constants**, numerical values of (HENNING), A., ii, 191.
- Gas generator** with washing apparatus (MÜLLER), A., ii, 706.
carbon oxides in (V. JÜPTNER), A., ii, 374.
- Gas reactions**, homogeneous, catalysis in (HERZFELD), A., ii, 136.
- Gas washing apparatus** (ERNST), A., ii, 839.
- Gastric juice** (TAKATA), A., i, 698.
relation of saliva to (NAKAGAWA), A., i, 789.
detection of bile pigments in (UTZ), A., ii, 799.
estimation of chlorine in (MALGOTRE), A., ii, 780.
- Gelatin**, preparation of, pure (LOEB), A., i, 387.
isoelectric condition of (RAWLING and CLARK), T., 2530.
solubility of (FAIRBROTHER and SWAN), T., 1237.
physical properties of solutions of (DAVIS and OAKES), A., i, 597.
effect of electrolytes on the solution and precipitation of (J. and R. F. LOEB), A., i, 182.

- Gelatin** gels, thermal expansion of (TAFTEL), T., 1971.
 elasticity of, as a function of the hydrogen-ion concentration (SHEPPARD, SWEET, and BENEDICT), A., ii, 747.
- Gelatin** as a protective colloid (GUTHRIE, HUBER, and ZWEIGLE), A. ii, 485; (GUTHRIE and EMSLANDER), A., ii, 625.
 dialysed, properties of (LLOYD), A., i, 1075.
 swelling of (MACDOUGALL), A., i, 205.
 drying and swelling of (SHEPPARD and ELLIOT), A., i, 387; (BOGUE), A., i, 388.
 swelling of, in organic acids (KUHN), A., i, 183.
 swelling and hydrolysis of, by acids (MOELLER), A., ii, 689.
 action of sulphosalicylic acid on the swelling of (OSTWALD and KUHN), A., i, 598.
 products of hydrolysis of (VAN SLYKE and HILLER), A., i, 63.
 viscosity of solutions of (DAVIS, OAKES, and BROWNE), A., i, 63.
 viscosity and plasticity of solutions of (BOGUE), A., i, 782; ii, 122.
 combination of acids and alkalis with (LLOYD and MAYES), A., i, 280.
 combination of hydrochloric acid with (WINTGEN and VOGEL), A., i, 280; (HITCHCOCK), A., i, 882.
 mixtures of, with asaric acid (TIEBACKX), A., i, 638.
 coagulation of mixtures of, with gum-arabic (TIEBACKX), A., ii, 698.
 nitrogenous value of, in the diet of man (ROBISON), A., i, 488.
- Gels**, action of ultra-violet light on (HOLMES and PATRICK), A., ii, 338.
 diffusion of electrolytes into (STILES and ADAIR; STILES), A., ii, 125.
 elastic, structure of (BOGUE), A., i, 782.
- Generator gas**, equilibrium of (JELLINEK and DIETHELM), A., ii, 838.
- Germanium**, occurrence and extraction of (DENNIS and PAPISH), A., ii, 150.
 occurrence of, in cassiterite (HADDING), A., ii, 855.
 crystal structure of (KOLKMEIJER), A., ii, 713.
- Germanium tetrabromide** and tetrachloride (DENNIS and HANCE), A., ii, 302.
 hydride (MÜLLER and SMITH), A., ii, 775; (PANETH and SCHMIDT-HEBBEL), A., ii, 776; (SCHENCK and IMKER), A., ii, 855.
- Germanium dioxide**, effect of, on blood (HAMMETT, NOWREY, and MÜLLER), A., i, 1086.
- Germanium detection and separation** :—
 detection of, by the arc spectrum (PAPISH), A., ii, 163.
 separation of arsenic and (MÜLLER), A., ii, 320.
- German silver**, equilibrium of the formation of (VOIGT), A., ii, 295.
- Germination**, action of nitrogenous compounds on (BOKORNY), A., i, 93.
- Glands**, physiology of (ASHER and KODA; ASHER and DOUBLER), A., i, 286; (ASHER and MATSUNO), A., i, 298; (ASHER and BERNET), A., i, 491.
- Glass**, molecular structure of (RAMAN), A., ii, 201.
 determination of the durability of (TURNER), A., ii, 719.
 effect of lime and magnesia on the durability of (DIMBLEBY, MUIRHEAD, and TURNER), A., ii, 709.
 preparation of suspensions of (FRICKE), A., ii, 692.
 diffusion of hydrogen and helium through (WILLIAMS and FERGUSON), A., ii, 841.
 migration of sodium ions through (PIRANI and LAX), A., ii, 817.
 colourless, use of selenium compounds in production of (COUSEN and TURNER), A., ii, 708.
 natural, properties and composition of (TILLEY), A., ii, 220.
 containing didymium, absorption spectra of (WEIGERT), A., ii, 413.
- Glassware**, chemical, action of analytical reagents on (TURNER and WILSON), A., ii, 501.
- Glauconite** from Sussex (HALLIMOND and RADLEY), A., ii, 861.
- Gliadin**, wheat, rate of hydrolysis of (VICKERY), A., ii, 754.
- Globulin**, carbohydrate as a constituent of (LANGSTEIN), A., i, 479.
 serum, colloidal behaviour of (HITCHCOCK), A., i, 1074.
- Globulins**, estimation of, in blood-serum (HENLEY), A., ii, 671.
 separation of, from serum (VILA), A., i, 1209.
- Glucamine**, production of (LING and NANJ1), T., 1686.
- Glucimidazole**, and 2-thiol- (PAULY and LUDWIG), A., i, 953.
- Glucimidazolone** (PAULY and LUDWIG), A., i, 953.
- Glucinum** (*beryllium*), atomic weight of (HONGSCHMID and BIRCKENBACH), A., ii, 214.

Glucinum (*beryllium*), spectrum of, and its relation to the spectrum of aluminium (GLASER), A., ii, 675.

anode rays of (THOMSON), A., ii, 106.

crystal structure of, and its oxide (McKEEHAN), A., ii, 766.

Glucinum fluoride, spectrum of (DATTÀ), A., ii, 411.

sulphate, dissociation of (MARCHEL), A., ii, 620.

equilibrium of ammonium sulphate, water and (BRITTON), T., 2612.

Glucinum separation :—

separation of aluminium and (BRITTON), A., ii, 228.

Glucosacose. See 2-Deoxyglucose.

α -Glucoseptol, synthesis of (PICTET and BARBIER), A., i, 4.

Glucoseogeny (MÜLLER; JUNG and MÜLLER), A., i, 486.

Gluconic acid, preparation of (LING and NANJÌ), A., i, 221.

Glucosamine, and its derivatives, mutarotation of (IRVINE and EARL), T., 2370.

formation of, heterocyclic compounds from (PAULY and LUDWIG), A., i, 953.

isoglucosamine (LING and NANJÌ), T., 1682.

d-Glucosamine, salicylidene derivatives of (IRVINE and EARL), T., 2376.

Glucose, thio-, and its silver salt and tetra-acetyl derivative (WREDE), A., i, 525.

d-Glucose. See Dextrose.

Glucoses, substituted, phosphoric and sulphuric esters of (LEVENE, MEYER, and WEBER), A., i, 987.

Glucose-ammonia (LING and NANJÌ), T., 1682; A., i, 631.

Glucose- α -galactoside (PICTET and VERNET), A., i, 811.

α -Glucosidase, non-identity of maltase and (WILLSTÄTTER and STEIBELT), A., i, 282.

Glucosides (KARRER, NÄGELI, and SMIRNOV), A., i, 253.

studies on (MACBETH and PRYDE), T., 1660.

new, synthesis of (DE FAZI), A., i, 755.

coumarin, in orchids (HÉRISSEY and DELAUNÉY), A., i, 210.

detection and extraction of, from plants (VERGÉLOT), A., i, 207.

Glucosides. See also :—

Amygdalin.

Anthochlor.

Aucubin.

Glucosides. See also :—

Capsularin.

Castelin.

Digitonin.

Dugaldin.

Hyssopin.

Indican.

Pelargononin.

Pelargonin.

Glue, animal, action of alum on (GUTHRIER, SAUER, and SCHELLING), A., i, 783.

Glutaconic acid (VERKADE), A., i, 520.

action of muscle enzymes on (DAKIN), A., i, 792.

derivatives of, addition of hydrogen cyanide to (HOPE and SHELTON), T., 2223.

Glutaconic acid, β -chloro-, preparation of, and its ethyl ester (INGOLD and NICKOLLS), T., 1643.

Glutaconic acids, isomerism of (FEIST), A., i, 521.

esters, self-condensation of (INGOLD, PERREN, and THORPE), T., 1783.

d-Glutamic acid, esters of, and their hydrochlorides (CHILES and NOYES), A., i, 924.

Glutamine, constitution of (THIERFELDER), A., i, 326.

synthesis of, in the human organism (SHIPLE and SHERWIN), A., i, 492.

Glutaric acid, ethyl ester, oxalyl derivative, and its semicarbazone (DICKENS, KON, and THORPE), T., 1501.

$\alpha\beta$ -trialkylated derivatives, preparation of (KON and THORPE), T., 1795.

Glutaric acid, α -hydroxy-, isopropyl ester (CHILES and NOYES), A., i, 925.

Glutaric acid series, esters of the, Dieckmann reaction with (KON), T., 513.

Glutaronitrile, action of magnesium organic compounds on (BRUYLANTS), A., i, 531.

Glycemia, respiratory metabolism in (BORNSTEIN and MÜLLER), A., i, 392; (BORNSTEIN and HOLM), A., i, 890.

d-Glyceraldehyde, conversion of, into active glyceric acid (WOHL and SCHELLENBERG), A., i, 626.

Glycerides, synthesis of (AMBERGER and BROMIG), A., i, 804.

constitution of (GRÜN), A., i, 420, 517, 621.

thin films of binary mixtures of (COLLET), A., ii, 477.

Glycerol, preparation of, from sugar (VEREINIGTE CHEMISCHE WERKE), A., i, 980.

synthesis of (PICTET and BARBIER), A., i, 4.

- Glycerol**, partition of compounds between acetone and (SMITH), A., ii, 270.
 equilibrium in the system, water and (PUSHIN and GLAGOLEVA), T., 2818.
 fermentation of, in presence of sulphur (H. and L. MÜLLER), A., i, 904.
 action of copper ethylenediamine hydroxide with (TRAUBE) A., i, 718.
 velocity of saponification of the acetins of (SMITH), A., i, 915.
 crude, distillation of (LEWIS), A., i, 419; (RAYNER), A., i, 711.
 action of *Aspergillus glaucus* on (TRAETTA-MOSCA and PBETI), A., i, 91.
 decomposition of, by *Bacillus lactis aerogenes* (KUMAGAWA), A., i, 972.
 estimation of (KELLNER), A., ii, 389.
 estimation of, by the dichromate method (KELLNER), A., ii, 723.
 estimation of, in presence of sugars (HOYT and PEMBERTON), A., ii, 321.
 estimation of trimethylene glycol in (COOKS and SALWAY), A., ii, 232.
Glycerophosphoric acid, ionisation constants of, and their use as buffers (MELLONS, ACREE, AVERY, and SLAGLE), A., i, 405.
Glyceryl chloro- and bromo-hydrins, preparation of, from allyl alcohol (READ and HURST), T., 989.
Glycine (aminoacetic acid), synthesis of, from formaldehyde (LING and NANJ1), A., i, 1124.
 synthesis of, in the human organism (SHIPLE and SHERWIN), A., i, 492.
 modifications of, and their action with diazomethane (BILTZ and PAETZOLD), A., i, 528.
Glycocholic acid, sodium salt, hæmolytic action of (PONDER), A., i, 292.
Glycogen (KARRER), A., i, 11.
 effect of sugar concentration on the synthesis of (EDERER), A., i, 895.
 change of, when exposed to light (BAYER), A., i, 323.
 content of white blood-corpuscles (DE HAAN), A., i, 484.
 formation in the embryo liver (ARON), A., i, 702.
 action of pilocarpine on the content of, in organs (HORNEMANN), A., i, 296.
 acetate (FRINGSHEIM and LASSMANN), A., i, 684.
Glycols, amino-, and nitro- (SCHMIDT and WILKENDORF), A., i, 314.
Glycollanilide sulphite (BLAISE and MONTAGNE), A., i, 520.
Glycollic acid, action of thionyl chloride on (BLAISE and MONTAGNE), A., i, 520.
Glycolysis, influence of the dextrose concentration on (MAURIAU and SERVANTIE), A., i, 1211.
Glycosuria. See Diabetes.
Glycylglycine, hydrolysis of, by hydrochloric acid (JAITSCHNIKOV), A., i, 1124.
Glyoxaline derivatives, synthesis of (PUCHER and JOHNSON), A., i, 549.
Glyoxaline, bromo-, and bromonitro-derivatives, and their salts (BALABAN and PYMAN), T., 947.
Glyoxaline-4-carboxylic acid, and 2:5-dibromo-, and its ethylester (BALABAN and PYMAN), T., 954.
Glyoxalinedicarboxylic acid, salts of, with organic bases (PAULY and LUDWIG), A., i, 953.
Glyoxaline-5-sulphonic acid, 4-bromo-, and its barium salt (BALABAN and PYMAN), T., 954.
Glyoxalones, thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
Glyoximes, structure of (BIGIAYI), A., i, 352.
 bisulphite compounds of (GASTALDI and BRAUNIZER), A., i, 626.
Gold, L-series spectrum of (DAUVILLIER), A., ii, 101; (DUANE and PATTERSON), A., ii, 463.
 Röntgen ray spectrum of (DE BROGLIE), A., ii, 330.
 preparation of colloidal solutions of (GETTLER and JACKSON), A., ii, 240.
 colloidal, effect of temperature on the coagulation of (LACHS and GOLDBERG), A., ii, 697.
 re-crystallisation of pure unworked (FRAENKEL), A., ii, 778.
 anodic oxidation of, in sulphuric acid (JIRSA and BURYÁNEK), A., ii, 713.
 action of selenium on (PÉLABON), A., ii, 303.
Gold alloys with copper, electrochemistry of (LORENZ, FRAENKEL, and WORMSER), A., ii, 21.
 reaction limit of (TAMMANN), A., ii, 75.
Gold silver bromides (SUSCHNIG), A., ii, 514.
 chlorides, complex (WELLS), A., ii, 449.
 Auric cesium chloride (WELLS), A., ii, 514.
Gold haloids, compounds of benzyl sulphide with (SMITH), A., i, 933.
 sulphides (GUTHRIE and DÜRRWACHTER), A., ii, 513.

Gold estimation:—

estimation of, colorimetrically (MULLER and FOIX), A., ii, 662.

Gouft oil, constituents of (ROURE-BERTRAND FILS), A., i, 845.

Grapes, Spanish, fluorine in (PONDAL), A., i, 1100.

Grape juice, methyl anthranilate in (POWER and CHESNUT), A., i, 97.

Grape seed oil, solid fatty acids from (ANDRÉ), A., i, 908.

Graphite, linkings of carbon atoms in (WIBAUT), A., ii, 259.

"Graphites" from pyrrole and from thiophen (CIUSA), A., i, 1057.

Grass, inchi. See *Cynbopogon coccineus*.

Grignard reaction (OSTLING), A., i, 843.

Grignard reagents, constitution of (V. BRAUN), A., i, 22.

use of, in organic syntheses (HERWORTH), A., i, 118.

action of, on thiocyanates (ADAMS), A., i, 531.

Guaiacol, 5-iodo-, behaviour of, in the organism (SIMON), A., i, 970.

Guaiacoldimercuri-compounds

(MAMELI), A., i, 1082.

Guaiacoyl- γ -methyl- $\Delta\beta$ -butenyl ether (STAUDINGER, KREIS, and SCHILT), A., i, 978.

Guanidine, preparation and derivatives of (MARCKWALD and STRUWE), A., i, 328.

mechanism of formation of, from dicyanodiamide (BLAIR and BRAHAM), A., i, 1123.

nitrate, preparation of (DAVIS), A., i, 117.

estimation of (DODD), A., ii, 536.

Guanidine, nitro-, action of sulphuric acid on (DAVIS), A., i, 530.

Guanidinophenylthiourea, amino-, and its benzylidene derivative (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 378.

Guanidoethyl alcohol, and its derivatives (FROMM and HONOLD), A., i, 530.

Guanidonium salts (MARCKWALD and STRUWE), A., i, 328.

***l*-Guleonic acid**. See *δ -epi-*l*-Rhamnonic acid*.

Gum arabic, adsorption in solutions of (CLARK and MANN), A., ii, 550.

effect of injections of, on diuresis (CORI), A., i, 1091.

coagulation of mixtures of, with gelatin (TIEBACKX), A., ii, 698.

Gypsum, crystalline, preparation of (BOURGEOIS), A., ii, 292.

dispersoid chemistry of (NEUGEBAUER), A., ii, 643.

Gypsum, transformation of, into ammonium sulphate (MATIGNON and FRÉJACQUES), A., ii, 570.

burnt, setting and velocity of solution of (BUDNIKOV and SYBKIN), A., ii, 849.

H.

*iso***Hæmatein**, and its ferrichloride tetramethyl ether (CRABTREE and ROBINSO), T., 1039.

Hæmatopyrinuria, experimental (BINDA), A., i, 89.

Hæmaturia, effect of hexamethylene-tetramine in the production of (BLOEDORN and HOUGHTON), A., i, 1091.

Hæmochrome, nature of (PARTOS), A., i, 597.

Hæmocyanin (DHÉRE and SCHNEIDER), A., i, 884.

Hæmoglobin, electric charge of (MICHAELIS and AIRILA), A., i, 63.

effect of acids and alkalis on the chemical and physical properties of (QUAGLIARIELLO), A., i, 1074.

heat of reaction of oxygen with (ADOLPH and HENDERSON), A., ii, 350.

crystallisation of (AMANTEA and KRZYSZKOWSKY), A., i, 387.

darkening of solutions of (GRAM), A., ii, 886.

physical chemistry of, in blood (BROWN and HILL), A., i, 1199.

combinations of, with carbon monoxide and oxygen, in blood (HILL), A., i, 193, 696.

estimation of (MEULENGRACHT), A., ii, 798; (TERRILL), A., ii, 799.

Hæmolysis, effect of temperature on (JARISCH), A., i, 394.

effect of metallic salts on (PURDY and WALBUM), A., i, 1087.

by morphine and its derivatives (RHODE), A., i, 1088.

by quinine (LUGER), A., i, 81.

Hæmotoxins from parasitic worms (SCHWARTZ), A., i, 493.

Halochromism (SKRAUP and FREUNDLICH), A., i, 539; (HANTZSCH), A., i, 556.

Halogens, physical properties and molecular structure of (HENGLEIN), A., ii, 44.

velocity of the action of, on metals (TAMMANN and KOSTER), A., ii, 831.

velocity of reaction of liberation of (FREUNDLICH and BARTELS), A., ii, 489.

- Halogens**, elimination of, in the reduction of halogenated nitro-compounds (BURTON and KENNER), T., 675.
 estimation of, by means of mercuric nitrate (KOLTHOFF and BAK), A., ii, 159.
 estimation of, microchemically (HOLTZ), A., ii, 521.
- Halogen atoms**, lability of, in organic compounds (HENDERSON and MACBETH), T., 892; (HIRST and MACBETH), T., 904, 2169; (GRAHAM and MACBETH), T., 1109, 2601; (MACBETH), T., 1116; (BLACK, HIRST, and MACBETH), T., 2527.
- hydrides**, formation and dissociation of, in aqueous solution (RAY and SARKAR), T., 1449.
 ionisation tension of (KNIPPING), A., ii, 186.
 viscosity of (HARLE), A., ii, 140.
- Halogenation** (DATTA and CHATTERJEE), A., i, 815.
- Halogen organic compounds**, action of ammonia and of titanous chloride on (BLACK, HIRST, and MACBETH), T., 2527.
 containing nitrogen, action of hydrazine on (HIRST and MACBETH), T., 904.
- Haloids**, constitution of (BILTZ), A., ii, 370.
 relative toxicity of ions of (CAMERON and HOLLENBERG), A., i, 499.
 inorganic, interaction of organo-metallic compounds with (CHALLENGER and RIDGWAY), T., 107.
- Hardness of solids** (REIS and ZIMMERMANN), A., ii, 745.
- Harmaline** (KERMACK, PERKIN, and ROBINSON), T., 1872.
- Harmine** (KERMACK, PERKIN, and ROBINSON), T., 1872.
- Harrison Memorial**, T., 2894.
 trust-deed establishing, T., 2917.
- Hasel nut oil** (PRITZKER and JUNGKUNG), A., i, 208.
- Heart**, action of metallic ions on the (CATE; ZONDEK), A., i, 296.
 muscle, creatine in (CONSTABEL), A., i, 296.
 action of drugs and alkaline earth salts on (BRAUN), A., i, 1213.
 frog's, effect of various ions on the (SPIRO), A., i, 489.
- Heat**, mechanical equivalent of (SWIENTOSLAWSKI), A., ii, 194.
 animal. See Animal heat.
 specific (PADOA), A., ii, 348.
 determination of (COHEN and MOESVELD), A., ii, 347.
- Heat**, specific, determination of, at low temperatures (SIMON), A., ii, 684.
 of gases (INGOLD and USHERWOOD), T., 2286; (MEWES), A., ii, 787.
 of metals (LATIMER), A., ii, 814.
 of solids (KRASE), A., ii, 421.
 of sulphides used in metallurgy (BORNEMANN and HENGSTENBERG), A., ii, 613.
- Heat of combustion** (HÜCKEL), A., ii, 195, 818.
 in relation to constitution (BINDER), A., ii, 27.
- Heat of dissociation** (HÜCKEL), A., ii, 818.
- Heat of evaporation** (HERZ), A., ii, 818.
- Heat of formation**, relation of, to atomic weight and volume change (COLLINS), A., ii, 685.
 in relation to ionic properties (GRIMM), A., ii, 690.
- Heat of fusion**, latent (STRATTON and PARTINGTON), A., ii, 258; (MORTIMER), A., ii, 622.
- Heat of solution** of organic substances, and its relation to the heat of fusion (GEHLHOFF), A., ii, 121.
- Heat of vaporisation** of non-associated liquids (MORTIMER), A., ii, 615.
- Hederagenin**, and its derivatives (VAN DER HAAR and TAMBURELLO), A., i, 160.
- α-Hederin**, and its salts and derivatives (VAN DER HAAR), A., i, 160.
- Helenium hoopesii** (sneezeweed), constituents of (MARSH, CLAWSON, COUCH, and MARSH), A., i, 413.
- Helianthus annuus**, oxidation of proteins by the green pigment in seeds of (OPARIN), A., i, 308.
- Heliotherapy** (VALLOT), A., i, 87.
- Helium**, thermodynamics of the formation of, from hydrogen (TOLMAN), A., ii, 738.
 spectrum of (LYMAN; SILBERSTEIN), A., ii, 674; (HICKS), A., ii, 675.
 band-spectrum of (CURTIS), A., ii, 330.
 series spectrum of (SEELIGER), A., ii, 17.
 ultra-violet spark spectrum of (MCLENNAN and PETRIE), A., ii, 330.
 neutral, spectrum of (RAMAN), A., ii, 803.
- Kehle model** of the atoms of (VAN VLECK), A., ii, 838.
- ionisation and resonance potentials** of (FRANCK), A., ii, 811.
- isotherms** of (HOLBORN and OTTO), A., ii, 737.

- Helium**, solubility of, in water (Cady, ELSEY, and BERGER), A., ii, 642.
diffusion of, through glass (WILLIAMS and FERGUSON), A., ii, 841.
- iso*Helium, atomic weight of (NEUBURGER), A., ii, 365.
- Hemicellulose**, digestion of (SHIMIZU), A., i, 82.
- Hemimellitene** (3:4:5-*trimethylphenol*), 2:6-dibromo-, and its derivatives (v. AUWERS and SAURWEIN), A., i, 1032.
- Hemimellitylquinitol**, dibromo-, and dinitro- (v. AUWERS and SAURWEIN), A., i, 1032.
- Hemimellityl- ψ -quinol**, dinitro- (v. AUWERS and SAURWEIN), A., i, 1033.
- Hemp**, distinction between flax and (NODDER), A., ii, 791.
- cis* and *trans*-**Heptadecenylurethanes** (NICOLET and PELO), A., i, 644.
- Heptadecenylcarbamide**, dihydroxy- (NICOLET and PELO), A., i, 644.
- Heptaldehydesemioxamazone** (LAPWORTH and McRAE), T., 2753.
- Heptamethylamygdalinic acid**, methyl ester (HAWORTH and LEITCH), T., 1924.
- n*-**Heptane**, action of nitrosyl chloride on (LYNN and HILTON), A., i, 417.
- cyclo***Heptane**, models for (MOHR), A., i, 441.
- cyclo***Heptane-1:2-diols**, space configuration of (DERX), A., i, 651.
- n*-**Heptenecarboxylic acid**, ethyl ester (CARMICHAEL), T., 2549.
- n*-**Heptyl fluoride** (SWARTS), A., i, 101.
- cyclo***Heptylcycloheptanol**, and its allophanic ester (GODCHOT and BRUN), A., i, 350.
- cyclo***Heptylidene-cycloheptanone** (GODCHOT and BRUN), A., i, 350.
- Herring roe**, fatty constituents of (TAKAHASHI), A., i, 700.
- Hesperidine**, chemistry, properties, and detection of (NILSSON), A., i, 211.
- Heterocyclic compounds**, effect of substitution on the formation and stability of (INGOLD, SAKO, and THORPE), T., 1177.
formation of, from glucosamine (PAULY and LUDWIG), A., i, 953.
- Heteropoly-acids** (ROSENHEIM and LEYSER), A., ii, 50.
of elements of the sulphur group (MEYER and STACECZY), A., ii, 773.
- 1:2:4:5:7:8-Hexa-acetoxyfluoran** (MUKERJI), T., 548.
- Hexa-amylose dodeca-acetate** (PRINGSHEIM and PERSCH), A., i, 632.
- Hexadecanesulphonic acid**, preparation and properties of (NORRIS), T., 2162.
- Hexadecyl fluoride** (SWARTS), A., i, 101.
- Hexahydrocinchonine dihydrobromide** (JACOBS and HEIDELBERGER), A., i, 672.
- α - and β -**Hexahydrodeoxycinchonines**, and their derivatives (JACOBS and HEIDELBERGER), A., i, 672.
- Hexahydrodioxidiboron**. See under Boron.
- Hexahydro- α -hydrindone**, and its derivatives (WINDAUS and HÜCKEL), A., i, 658.
- Hexahydroquinine**, and nitroso-, and their derivatives (JACOBS and HEIDELBERGER), A., i, 678.
- Hexahydrosalicylaldehyde**, and its hydrochloride (WEIL, TRAUN, and MARCEL), A., i, 1023.
- n*-**Hexaldehyde**, δ -hydroxy-, and its *p*-bromophenylhydrazone (HELFFERICH and MALKOMES), A., i, 431.
- Hexamethylenes**, stereoisomeric (SKITA), A., i, 534.
- Hexamethylenedicarbimide** (SCHMIDT), A., i, 761.
- Hexamethyleneimine**, and its salts and acetyl derivatives (SCHMIDT), A., i, 761.
- Hexamethylenetetramine**, preparation of (TRAUN's FORSCHUNGS-LABORATORIUM), A., i, 528; (RIEDEL), A., i, 991.
catalytic synthesis of (ROMBAUT and NIEUWLAND), A., i, 990.
action of alkyl haloids with (DELÉPINE and JAFFEUX), A., i, 234.
action of sodammonium on (PICON), A., i, 1123.
preparation of compounds of esters of fatty acids with (RIEDEL), A., i, 991.
role of, in production of hæmatumia (BLOEDORN and HOUGHTON), A., i, 1091.
salts of, action of formic acid with (SOMMELET and GUIOTH), A., i, 338.
- Hexaminechromiselenate**. See under Chromium.
- Hexaminecobaltic salts**. See under Cobalt.
- Hexaminecobaltic 2:4-dinitro- α -naphthoxide**, and its 7-sulphonic acid (MORGAN and KING), T., 1727.
- Hexaminecobaltic dipicrylamine** (MORGAN and KING), T., 1727.
- Hexane**, β -chloro- γ -hydroxy- (DETREUF), A., i, 327.
- cyclo***Hexane**, 1-chloro-2-hydroxy- (DETREUF), A., i, 327.
- cyclo***Hexanes**, substituted, action of hydrazine on halogen derivatives of (HIRST and MACBETH), T. 2169.

- cycloHexanespirocyclobutan-3-one*, and its derivatives (KON), T., 520.
- cycloHexane-1:1-diacetic acid*, methyl ester (DICKENS, KON, and THORPE), T., 1505.
- cycloHexane-1:2-diols*, space configuration of (DERX), A., i, 651.
- cycloHexanespirocyclopentane-3:4-dione*, and its derivatives (KON), T., 522.
- cycloHexanetriols*, catalytic preparation of (SENDERENS and ABOULENC), A., i, 337.
- cycloHexanol*, preparation of (BROCHET), A., i, 1144.
- Hexaphenylethane*, mechanism of the dissociation of (COLE), A., ii, 438.
- Hexapyridine μ -dichlorodiferridi(tetrachloroferrate)* (WEINLAND and KISSLING), A., i, 364.
- Hexapyridinemagnesium salts* (SPACU), A., i, 859.
- Hexen- β -diol*, $\alpha\zeta$ -dichloro- $\gamma\delta$ -dibromo- (LESPIEAU), A., i, 103.
- cycloHexene*, compounds of, with nitrogen tetroxide and zinc (SCHAAR-SCHMIDT, VEIDT, and SCHLOSSER), A., i, 646.
- Δ^8 -*Hexenoic acid*, vanillylamide of (OTT and ZIMMERMANN), A., i, 137.
- Δ^8 -*Hexenoyl chloride* (HELFERICH and MALKOMES), A., i, 431.
- Hexinene- $\alpha\beta\zeta$ -tetrol*, dioxide from (LESPIEAU), A., i, 103.
- n-Hexoic acid*, $\alpha\gamma$ -dihydroxy-, barium salt (HELFERICH and SPEIDEL), A., i, 7.
- γ *isoHexolactone*, $\alpha\beta$ -dihydroxy- (FRANKE and GROEGER), A., i, 808.
- Hexolactones*, bromo- and α -hydroxy- (HELFERICH and SPEIDEL), A., i, 7.
- 2-Hexosamic acids**, synthesis of (LEVENE), A., i, 327.
- 2-Hexosamines**, synthesis of (LEVENE), A., i, 327.
- Hexosediphosphoric acid*, strychnine salt (NEUBERG and DALMER), A., i, 920.
- Hexosephosphatase* (TOMITA), A., i, 960.
- Hexosephosphoric acid*, brucine, cinchonidine, and strychnine salts (NEUBERG and DALMER), A., i, 920.
- cycloHexyl allyl*, ethyl and methyl esters, bromo- (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- 5-cycloHexylbarbituric acid** (DOX and YODER), A., i, 817.
- cycloHexylcarbinyl alcohol*, and its urethane (DE POMMEREAU), A., i, 339.
- cycloHexylmalonamide* (DOX and YODER), A., i, 817.
- cycloHexylmalonic acid*, methyl ester (DOX and YODER), A., i, 816.
- cycloHexyl methyl ketone*, 2-chloro-, and its semicarbazone (WIELAND and BETTAG), A., i, 1034.
- cycloHexyl phenyl ketone*, 2-chloro-, and its semicarbazone (WIELAND and BETTAG), A., i, 1034.
- Hexylsuccinamic acid** (HIGGINBOTHAM and LAPWORTH), T., 52.
- n-Hexylsuccinimide* (LAPWORTH and McRAE), T., 2750.
- 5-cycloHexyl-2-thiobarbituric acid** (DOX and YODER), A., i, 817.
- Hibernium**, a possible new radioactive element (JOLY), A., ii, 651.
- Hides**, dyeing and tanning of (RAKUZIN), A., i, 702.
- Hippuric acid**, synthesis of, in the liver (DELPRAT and WHIPPLE), A., i, 86.
- action of Dakin's hypochlorite solution on (ENGELDT), A., i, 813.
- estimation of, in urine (SNAPPER and LAQUEUR), A., ii, 92.
- Hirudo medicinalis* (leech), physiology and pharmacology of (TESCHENDORF), A., i, 970.
- Histamine**. See 4-Ethylglyoxaline, β -amino-.
- Histidine**, preparation of, from blood (DEMJANOVSKI), A., i, 1052.
- production of urocanic acid from (KOTAKE and KONISHI), A., i, 1217.
- detection of (HUNTER), A., ii, 885.
- Histidine anhydride** dihydrochloride (FRANKEL and JELLINEK), A., i, 965.
- Histone sulphate**, digestion of, by pepsin (FELIX), A., i, 693.
- Histopeptone** (FELIX), A., i, 479.
- Homocamphoric acid**, phenyl ethyl ester of (PALFRAY), A., i, 548.
- Homocamphoric glycol**, and its derivatives (PALFRAY), A., i, 548.
- Homogeneity** (OSTWALD), A., ii, 131.
- Homogentisic acid** (MÖRNER), A., i, 341.
- estimation of, in urine (BRIGGS), A., ii, 534.
- Homologous series**, variation of physical properties in (PAULY), A., i, 1; (STARCH), A., i, 2.
- Homophthalic acid**, ethyl ester, preparation of (DAVIES and PERKIN), T., 2208.
- Homopiperonyl methyl ester**, and 6-nitro- (GREENE and ROBINSON), T., 2194.
- p-Homosalicylideneanthranilic acid** (EKELRY, ROGERS, and SWISHER), A., i, 935.
- Homoterpenylic acid** (SIMONSEN), T., 2298.
- Honey**, formation and ripening of (SARIN), A., i, 197.

- Hormones**, distribution of, in blood (SCHAEFFI), A., i, 290.
- Horse**, sedimentation of blood corpuscles of the (RUNNSTROM), A., i, 289.
- "Ksiung Ch'uang,"** a Chinese drug, constituents of (MURAYAMA), A., i, 810.
- Humic acids** (ELLER; JONAS), A., i, 326.
structure and formation of (MARCUS-SON), A., i, 487.
- Humoeric acid** (ASCHAN), A., i, 714.
- Humus**, spontaneous oxidation of (SCHRADER), A., i, 637.
estimation of, by oxidation with chromic acid (GEHRING), A., ii, 592.
- Hydantoin** (READ), A., i, 954.
effect of substitution on the formation and stability of (INGOLD, SAKO, and THORPE), T., 1177.
- Hydrargyrum oxycyanatum**. See Mercury oxycyanide.
- Hydration**, static method of study of (GUICHARD), A., ii, 555.
- Hydratropic acid**, fate of, in the body (KAY and RAPER), A., i, 1093.
- Hydrazide**, $C_6H_5ON_2S_2$, from methyl chlorothioformate and thiosemicarbazide (ARNDT, MILDE, and TSCHENSCHER), A., i, 376.
 $C_6H_5N_4S_2$, from thiosemicarbazide and phenylthiocarbimide (ARNDT, MILDE, and TSCHENSCHER), A., i, 377.
 $C_6H_5ON_4S$, from phenylthiocarbimide and semicarbazide hydrochloride (ARNDT, MILDE, and TSCHENSCHER), A., i, 376.
- Hydrazine stannichloride** (DRUCE), A., i, 639.
estimation of, volumetrically (KURTE-NACKER and WAGNER), A., ii, 312.
- Hydrazines** (WIELAND and WECKER), A., i, 779.
optically active, preparation of (GLATTFELD and WERTHEIM), A., i, 385.
- Hydrazinediacetoneamides** containing sulphur, ring closure with (FROMM), A., i, 62; (ARNDT, MILDE, and TSCHENSCHER), A., i, 375.
- 9-Hydrazinoacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 469.
- Hydrazino-9:9-bisacridine** (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 469.
- Hydrazinodiacetoneglucose** (FREUDENBERG and BRAUNS), A., i, 1117.
- 1-Hydrazino-4-methylanthraquinone** (MAYER and HEIL), A., i, 878.
- o-Hydrazinophenylacetic acid** (NEBER), A., i, 546.
- Hydrazinopyrine picrate** (THIELEPAPE and SIRRECKELSEN), A., i, 1191.
- 2-Hydrazinoquinoline-4-carboxylic acid**, and its hydrazide (THIELEPAPE), A., i, 271.
- Hydrazobenzene**, 5-bromo-2:4-dinitro- (GIUA), A., i, 691.
- Hydrazo-compounds** (CHARRIER), A., i, 769.
isomeric changes of (JACOBSEN), A., i, 589, 596.
- Hydrazoic acid**. See Azoimide.
- Hydrazones** (BRADY and McHUGH), T., 1648.
basic properties of (CIUSA and VECCHIOTTI), A., i, 474.
- 1:1-Hydrazo-ar-tetrahydronaphthalene** (SCHROETER, KINDERMANN, DIETRICH, BEYSLAG, FLEISCHHAUER, RIEBENSAMM, and OESTERLIN), A., i, 123.
- Hydrindene derivatives**, dehydrogenation of (V. BRAUN, HAHN, and SEEMANN), A., i, 728.
- Hydrindene**, bromohydroxy- (READ and HURST), T., 2554.
- 1-Hydrindone**, 5-hydroxy-, derivatives (V. AUWERS, HILLIGER, and WULF), A., i, 1194.
- Hydrindones** (V. KONEK and SZAMAK), A., i, 458.
- 2-Hydrindone-1-carboxylic acid**, ethyl ester, and its phenylhydrazones (PERKIN and TITLEY), T., 1566.
- 2-Hydrindone-3-carboxylic acid**, ethyl ester (DIECKMANN), A., i, 1022.
- Hydriodic acid**. See under Iodine.
- Hydrobenzoin**, 2:4-dinitro- (BISHOP and BRADY), T., 2367.
- Hydrobromic acid**. See under Bromine.
- Hydrobromoeupreine** dihydrobromide (HEIDELBERGER and JACOBS), A., i, 673.
- Hydrobutoxin** (WIELAND and ALLES), A., i, 784.
- Hydrocarbon**, $C_{10}H_{18}$, from dehydration of dihydrofencholenyl alcohol (MAXWELL), A., i, 753.
- $C_{10}H_{18}$, from reduction of 1:2:2:3-tetramethylcyclopentyl phenyl ketone (RUPE and JÄGGI), A., i, 840.
- $C_{18}H_{36}$, from condensation of *p*-tolyl methyl ketone (GASTALDI), A., i, 367.
- Hydrocarbons**, formation of, from carbon dioxide and from formates (LEVI), A., i, 801.
preparation of (PLAUSON and V. TISCHENKO), A., i, 818.
relation between molecular weights and iodine values of (WOO), A., i, 101.

- Hydrocarbons**, viscosity of (CHAVANNE and VAN RISEGHEM), A., ii, 352.
- nitration of (BATTESAY and BRANDT), A., i, 1001.
- oxidation of (BLAIR and WHEELER), A., i, 917.
- action of ozone on (WHEELER and BLAIR), A., i, 1105.
- action of sodammonium on (LEBEAU and PICON), A., i, 801.
- aliphatic, forces between carbon atoms in, and in diamond (FAJANS), A., ii, 194, 818.
- oxidation of, with nitrogen peroxide (GRANACHER and SCHAUFELBERGER), A., i, 513.
- aromatic, molecular refraction of (v. AUWERS), A., ii, 98; (v. AUWERS and KOLLIGS), A., ii, 174.
- action of, with azo-compounds and aluminium chloride (PUMMERER and BINAPFL), A., i, 24.
- reaction of azobenzene hydrochloride with (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1196.
- estimation of, in mineral oils (WATERMAN and PERQUIN), A., ii, 399.
- cyclic, stereoisomerism of (SKITA and SCHENCK), A., i, 240.
- ethylenic, action of chlorocarbamide on (DETGEUF), A., i, 236, 327.
- liquid, preparation of, from ethylene and sulphuric acid (DAMIENS), A., i, 1105.
- paraffin, catalytic oxidation of (SALWAY and WILLIAMS), T., 1348.
- saturated and unsaturated, energy of atomic linkings in (WIBAUT), A., i, 909.
- semibenzene (v. AUWERS and ZIEGLER), A., i, 119.
- unsaturated, hydrogenation of (CHEMISCHE FABRIK GRIESHEIM-ELEKTRON), A., i, 977.
- aliphatic, replacement of halogens in (KAUFMANN), A., i, 213.
- Hydrocarbons**, chloro-derivatives, elimination of hydrogen chloride from (FARAGHER and GARNER), A., i, 3.
- Hydrocarbostyryldihydrocoumarin-3:3-spiro** (LEUCHS, v. KATINSKY, and CONRAD), A., i, 473.
- Hydrocellulose**, action of heat on (JUSTIN-MUELLER), A., i, 11.
- Hydrochloric acid**. See under Chlorine.
- Hydrocinehonidine**, salts of tropic acid with (KING and PALMER), T., 2585.
- Hydrocinehonine**, salts of tropic acid with (KING and PALMER), T., 2584.
- Hydrocinehonine**, nitro-, and its sulphonic acid (BOEHRINGER & SÖHNE), A., i, 46.
- Hydrocupreine oxide** (VEREINIGTE CHININFABRIKEN ZIMMER & Co.), A., i, 948.
- Hydrodiphthal-lactonic acid**, and bromo-, and their derivatives (RUGGLI and MEYER), A., i, 344.
- Hydroesgonidine hydrochloride** and aurichlorides (GADAMER and JOHN), A., i, 168.
- Hydroferrieyanic acid**, zirconium salt (VENABLE and MOEHLMANN), A., ii, 712.
- Hydroferrieyanides of organic bases** (CUMMING), T., 1287.
- Hydroferroeyanic acid**, potassium salt, solubility of, in water (FABRIS), A., i, 329.
- decomposition of (BAUDISCH and BASS), A., i, 993.
- zirconium salt (VENABLE and MOEHLMANN), A., ii, 712.
- Hydroferroeyanides of organic bases** (CUMMING), T., 1287.
- Hydrofluosilicic acid**. See under Fluorine.
- Hydrogen**, model of the molecule of (CREHORE), A., ii, 494.
- model of the ionised molecule of (PAULI), A., ii, 703.
- spectrum of (WOOD), A., ii, 273; (MERTON and BARRATT), A., ii, 461.
- continuous spectrum of (LEWIS), A., ii, 802.
- line spectrum of (OLDENBURG), A., ii, 725; (GEHRCKE and LAU), A., ii, 726; (KIMURA and FUKUDA; HULBERT), A., ii, 801.
- series spectrum of (SEELIGER), A., ii, 17; (OLDENBURG), A., ii, 410.
- stability of the luminous discharge in (GIBSON and NOYES), A., ii, 812.
- atomic, incandescence of substances in (WOOD), A., ii, 759.
- atoms, radius of, in crystals (AMINOFF), A., ii, 496.
- triatomic (WENDT and LANDAUER), A., ii, 369.
- nuclei, collision of α -particles with (CHADWICK and BIELER; MCAULAY), A., ii, 12.
- structure and magnetic susceptibility of molecules of (HONDA), A., ii, 838.
- movement of electrons in (TOWNSEND and BAILEY), A., ii, 836.
- rectilinear diameter of (MATHIAS, CROMMELIN, and ONNES), A., ii, 440, 561.

- Hydrogen**, effect of a glow discharge in, between tin electrodes (PANETH), A., ii, 363.
 absorption of, in the electric discharge (NEWMAN), A., ii, 279, 546, 639; (WENDT), A., ii, 639.
 specific heat of (INGOLD and USHERWOOD), T., 2286.
 heat of vaporisation and specific heat of (MATHIAS, CROMMELIN, and ONNES), A., ii, 472.
 occlusion of, by palladium (OXLEY), A., ii, 469.
 properties of, when desorbed from platinum and palladium (ANDERSON), T., 1158.
 compressibility of (GUYE and BATURECAS), A., ii, 618.
 vapour pressure of (MARTINEZ and ONNES), A., ii, 472; (ONNES and MARTINEZ), A., ii, 759.
 velocity of sound in (GRÜNEISEN and MERKEL), A., ii, 190.
 thermodynamics of the formation of helium from (TOLMAN), A., ii, 738.
 diffusion of, through glass (WILLIAMS and FERGUSON), A., ii, 841.
 through iron and platinum (SCHMIDT and LÜCKE), A., ii, 198.
 dissociation of, in a tungsten furnace (DUFFENBACK), A., ii, 620.
 by excited mercury atoms (CARIO and FRANCK), A., ii, 809.
 formation and dissociation of polyhalogen compounds of, in aqueous solution (RAY and SARKAR), T., 1449.
 equation of state for (SCHAMES), A., ii, 24.
 equilibrium of the reduction of nickel chloride by (BERGER and CRUT), A., ii, 38.
 combination of, with acetylene derivatives (ZALKIND), A., ii, 835.
 photochemical reactions between chlorine and (GÖHRING), A., ii, 9; (WEIGERT and KELLERMANN), A., ii, 808.
 action of oxygen with, on platinum (LANGMUIR), A., ii, 629.
 combination of oxygen and, in presence of colloidal palladium (SANDONINI and QUAGLIA), A., ii, 556.
 catalytic formation of water from oxygen and (PEASE and TAYLOR), A., ii, 701.
 ignition of mixtures of oxygen and (MITCHELLICH), A., ii, 358.
 flame, luminescence excited by the (HOWES), A., ii, 335.
 simple generator for (CATHCART), A., ii, 440.
- Hydrogen compounds**, gaseous, molecular structure of (RANKINE), A., ii, 635.
Hydrogen bromide. See Hydrobromic acid under Bromine.
 chloride. See Hydrochloric acid under Chlorine.
 cyanide. See Hydrocyanic acid under Cyanogen.
 haloids. See Halogen hydrides.
 iodide. See Hydriodic acid under Iodine.
 peroxide, constitution of (ODDO and BINAGHI), A., i, 314.
 production of, by bacteria (MCLEOD and GORDON), A., i, 1095.
 catalytic formation and decomposition of, by platinum (HOFMANN), A., ii, 490.
 effect of radium rays on the rate of formation and decomposition of (KAILAN), A., ii, 106, 466, 758.
 catalytic decomposition of, by finely-divided platinum (MAXTED), T., 1760.
 effect of Rontgen rays on the catalytic decomposition of (SCHWARZ, FRIEDRICH, and WUNNERLICH), A., ii, 436.
 oxidation of butyric acid by (WITZEMANN), A., i, 6.
 influence of ferrous oxide on the action of, on iodides (PAMFILOV and PETIN), A., ii, 835.
 velocity of the action of, on metals (TAMMANN and KOSTER), A., ii, 831.
 velocity of reaction of potassium permanganate with (ZAWIDZKI), A., ii, 207.
 action of, on solutions of sugars (SCHONEBAUM), A., i, 920.
 bactericidal action of (MÜLLER), A., i, 903.
 estimation of, colorimetrically (ISAACS), A., ii, 715.
 phosphide, estimation of (MOSEER and BRUKL), A., ii, 393.
 selenide, preparation of (MOSEER and DOCTOR), A., ii, 46.
 sulphide, effect of, on respiration (HAGGARD and HENDERSON), A., i, 1206.
 hindrance of precipitation with (DEDE and BONIN), A., ii, 766.
 di- and tri-sulphides, preparation and properties of (WALTON and PARSONS), A., ii, 281.
 telluride, preparation of (MOSEER and ERTL), A., ii, 48.

Hydrogen estimation and separation:—
estimation of, in organic compounds (WREDE), A., ii, 316; (LINDNER), A., ii, 657; (SIMON and GUILLAUMIN), A., ii, 867; (SIMON), A., ii, 868.
estimation of traces of oxygen in (LARSON and WHITE), A., ii, 311.
estimation of, and its separation from gaseous paraffins (MULLER and FOIX), A., ii, 655.

Hydrogen-calomel cell. See under Cells, electrochemical.

Hydrogen electrode. See Electrode.

Hydrogen ions, determination of concentration of (MICHAELIS and KRÜGER; ACREE, MELLON, AVERY, and SLAGLE; GALLIESPIE), A., ii, 157.
colorimetrically (KOLTHOFF), A., ii, 222
colorimetrically, and electrometrically (CULLEN and HASTINGS), A., ii, 657.
in blood (LINDHARD), A., ii, 240.
apparatus for (KLING, LASSIEUR, and LASSIEUR; HEALY and KARAKER), A., ii, 519; (KLING and LASSIEUR), A., ii, 234, 715.
measurement of, and their hydration (SCHREINER), A., ii, 466.
effect of sucrose on the activity of (CORRAN and LEWIS), A., ii, 691.

Hydrogenation apparatus (KLIMONT), A., ii, 369.

Hydroalcohol dimethyl ether, and 5-mono- and 5:6-di-nitro (MAJIMA and CHIBA), A., i, 264.

Hydrolysis of salts, theory of (TIAN), A., ii, 362.
of weak acids and weak bases (GRIFFITH), A., ii, 420.

Hydronitric acid. See Azoimide.

Hydrophthalyl-lactonic acid, chloro-, esters of (RUGGLI and MEYER), A., i, 344.

Hydroquinidine, salt of tropic acid with (KING and PALMER), T., 2582.

Hydroquinine, salts of tropic acid with (KING and PALMER), T., 2582.

Hydroquinine, amino-, and its sulphonic acid (BOEHRINGER & SÖHNE), A., i, 46.
nitro-, and its sulphonic acid (BOEHRINGER & SÖHNE), A., i, 46.

Hydrosols, freezing of (LOTTERMOSER), A., ii, 286.
coloured, of nephelometry (BECHOLD and HEBLER), A., ii, 652.

Hydrothitsiol, and its dimethyl ether and 6-nitro-, dimethyl ether (MAJIMA and CHIBA), A., i, 264.

Hydroxamic acids (JONES and SCOTT), A., i, 453; (OLIVERI-MANDALA), A., i, 544.
preparation and rearrangement of (JONES and HURD), A., i, 248.

Hydroxy-acids, action of potassium iodide and iodate on (KRISHNA and POPE), T., 798.
degradation of, in the animal organism (WORINGER), A., i, 90.
esters, oxidation of, by oxygen or air (SIMON), A., i, 984.
aromatic, in urine (DE SANCTIS and FIORI), A., i, 609.

α -Hydroxy-acids, configuration of (FREUDENBERG and BRAUNS), A., i, 623.
action of thionyl chloride on (BLAISE and MONTAGNE), A., i, 520, 715.

Hydroxyaldehydes, synthesis of substances like disaccharides from (HELFERICH and WEIDENHAGEN), A., i, 1115.
aromatic, manufacture of (SOCIÉTÉ CHIMIQUE DES USINES DU RHÔNE), A., i, 257.

γ -Hydroxyaldehydes (HELFERICH and GEHRKE), A., i, 9.

Hydroxy-amines, aldehyde derivatives of (BERGMANN, ULPTS, and CAMACHO), A., i, 1180.

Hydroxycarbonyl compounds (KARRER), A., i, 40.

Hydroxy-compounds, aromatic, action of sulphites on (BUCHERER and WAHL), A., i, 464; (BUCHERER and ZIMMERMANN), A., i, 465.

Hydroxy-ketones, migration of acyl residues in acyl derivatives of the phenylhydrazones of (V. AUWERS, HILLIGER, and WULF), A., i, 1192.

Hydroxyl groups, auxiliary valency of (REIHLEN), A., i, 1146; (REIHLEN and SAPPER), A., i, 1147.

α -Hydroxy-lactones (HELFERICH and SPEIDEL), A., i, 6.

Hydroxylactonic acid, $C_6H_4O_6$, and its ethyl ester, from hydrolysis of ethyl $\alpha\alpha'$ -dibromo- $\beta\beta$ -diethylglutarate (DESHAPANDE and THORPE), T., 1438.

Hydroxylamine, preparation of (LECHER and HOFMANN), A., ii, 442.
reactions of (KJELLIN), A., ii, 640.
action of, and its derivatives on sodium arsenite (GUTMANN), A., ii, 844.
stannichloride (DRUCE), A., i, 1206.
estimation of, volumetrically (KURTENACKER and WAGNER), A., ii, 312.

Hydroxymethyleneacetophenone. See Phenyl β -hydroxyvinyl ketone.

- α -Hydroxymethylene- α -benzylacetone.**
See β -Phenyl- α -hydroxymethylene-ethyl methyl ketone.
- Hydroxymethylenedecybenzoin,** action of phenylhydroxylamine with (RUPE and WITTWER), A., i, 448.
- Hygroscopy** of soluble substances (EDGAR and SWAN), A., ii, 349.
- Hyoscyamine,** preparation and racemisation of and its sulphate (GORIS and COSTY), A., i, 1174.
- Hyperglycemia,** adrenaline (ACHARD, RIBOT, and BINET), A., i, 963; (BRÜSAMLEN), A., i, 1092.
- Hypochlorous acid.** See under Chlorine.
- Hypocloudous acid.** See under Iodine.
- Hypophosphorous acid.** See under Phosphorus.
- Hyposulphurous acid.** See under Sulphur.
- Hypoxanthine,** oxidation of, by tissues and by milk (MORGAN, STEWART, and HOPKINS), A., i, 1078.
- Hyasopin** (OESTERLE), A., i, 849.
- Hyascinglycone,** and its acetate (OESTERLE), A., i, 849.
- Hytasarindipyrindinium dibromide,** and its phenobetaine (BARNETT and COOK), T., 1388.

I.

- Ice,** crystal structure of (BRAGG), A., ii, 440.
- Ichthyol,** standardisation of commercial preparations of (PÉPIN and REAUBOURG), A., ii, 784.
- Ignition** of gases (PAYMAN and WHEELER), T., 363; (WHITE), T., 1244, 2561; (MASON and WHEELER), T., 2079.
- of mixtures of ammonia with air and oxygen (WHITE), T., 1688.
- Illipé fat,** constituents of (KOBAYASHI), A., i, 1113.
- Illipene,** and its hydrochloride (KOBAYASHI), A., i, 1113.
- Ilsemanite,** occurrence of (COOK), A., ii, 576.
- Imidasolones,** thio-. See Glyoxalones, thiol-.
- Imidoacetyl-di-*N,N*-rhodanine- α -acenaphthenequinone** (GRANACHER, REIS, and POOL), A., i, 577.
- "Iminasoleisopiperidin."** See 1:3:5-Benzotriazole.
- Iminoacetiopropionic acid,** ethyl ester (RUZICKA and SEIDEL), A., i, 1058.
- Imiscanilourazole,** and its derivatives (CASSELLA & Co.), A., i, 277.
- Imino-aryl ethers** (CHAPMAN), T., 1676.
- Iminodi-*N,N*-rhodanine- α -acenaphthenequinone** (GRANACHER, REIS, and POOL), A., i, 577.
- Imino-ethers,** spontaneous decomposition of (JOHNSON and BASS), A., i, 736.
- Incandescence** of substances in hydrogen gas (WOOD), A., ii, 759.
- Indazole, 5-amino-, and 5-iodo-, and its acetyl derivative** (v. AUWERS and LANGE), A., i, 687.
- bromo- and chloro-, and their derivatives (v. AUWERS and LANGE), A., i, 686.
- Indazoles,** stereoisomerism of, and of their halogen derivatives (v. AUWERS and LANGE), A., i, 684.
- Indazole-3-carboxylic acid,** and 5-bromo-, ethyl esters and derivatives (v. AUWERS and LANGE), A., i, 687.
- Indene,** catalytic hydrogenation of (v. BRAUN and KIRSCHBAUM), A., i, 727.
- action of bromine water on (READ and HURST), T., 2550.
- 2:3-Indeno(2:1)-indole** (ARMIT and ROBINSON), T., 838.
- 2:3-Indeno(2:1)-indolecarboxylic acid** ethyl ester (PERKIN and TITLEY), T., 1568.
- Indian yellow,** precursor of (GORTER), A., i, 932.
- Indican,** constitution of (MACBETH and PRYDE), T., 1660.
- estimation of, in blood serum (SNAPPER and VAN VLOTEN), A., ii, 793.
- Indicators,** sensitiveness of (KOLTHOFF), A., ii, 156.
- use of, without buffer solutions (MICHAELIS and KRUGER), A., ii, 157.
- coloured, salt error of (KOLTHOFF), A., ii, 222.
- modified methyl-orange (HICKMAN and LINSTED), T., 2502.
- mixed, use of (COHEN), A., ii, 780.
- radioactive (PANETH), A., ii, 785.
- Indigotin,** synthesis of (BAILEY and POTTER), A., i, 370.
- acylation and alkylation of leuco-derivatives of (GRANDMOUGIN), A., i, 470.
- Indigotin, octabromo-** (GRANDMOUGIN), A., i, 55.
- Indigotins,** halogenated (GRANDMOUGIN), A., i, 180.
- Indigotinsulphonic acid,** estimation of, by reduction (TREADWELL, DREIFUSS, and BOSSI), A., ii, 789.
- Indilidi-indoyls** (ODDO and SANNA), A., i, 371.

- Indium**, arc spectrum of (UHLER and TANCH), A., ii, 677.
- Indole**, formation of, by bacteria, and its reactions (FRIEBER), A., i, 901.
effect of sugar on the production of (APPELMANS), A., i, 794.
mechanism of the Fischer synthesis of (HOLLINS), A., i, 863.
detection of, by the cotton wool plug test (GORE), A., ii, 535.
- Indole group**, syntheses in the (ODDO and SANNA), A., i, 371.
- Indole-2-carboxydimethylacetalyamide** (KERMACK, PERKIN, and ROBINSON), T., 1892.
- Indole-2-carboxydimethylacetaly-methylamide** (KERMACK, PERKIN, and ROBINSON), T., 1886.
- 2-Indoxyl-1'-indaneindigo**. See 2-Indoxyl-1-indan-2-one.
- Indolinone**, derivatives of (TOMIČEK), A., i, 679.
- Indophenol** and its oxide, monomethyl ethers of (KEHRMANN, DECKER, and SCHMAJEWSKI), A., i, 32.
- Indophenols**, preparation of (SCHOUTISEN), A., i, 135.
- 2-Indoxyl-1-indan-2-one**, and dibromo- (FRIEDLÄNDER, HERZOG, and v. VOSS), A., i, 764.
- Inflammation** (WOLF), A., i, 401.
- Inorganic compounds**, heat of formation, lattice energy and work of ionisation of (GRIMM), A., ii, 690.
regularities in the molecular volumes of (HENGLEIN), A., ii, 260.
- Inositol**, decomposition of, by *Bacillus lactis aerogenes* (KUMAGAWA), A., i, 972.
- i-Inositol**, fermentation of (HEWITT and STEABEN), A., i, 406.
- Interferometer**, water, use of, in analysis of non-aqueous solutions (COHEN and BRUNS), A., ii, 77.
use of, as a pressure gauge (VAN DOREN, PARKER, and LOTZ), A., ii, 263.
- Intertraction and diffusion** (SCHONEBOOM), A., ii, 690.
- Intestines**, sterilisation of, during fasting (DOMBROWSKI and KOZŁOWSKI), A., i, 606
intoxication in (GERARD), A., i, 790.
effect of choline on movements of (LE HEUX), A., i, 85.
histamine in (MEAKINS and HARINGTON), A., i, 396.
saccharase in (v. EULER and SVANBERG), A., i, 296; (KNAFFL-LENZ), A., i, 485.
permeability of, to sucrose (WORINGER), A., i, 1214.
- Inulin** (PRINGSHEIM and LASSMANN), A., i, 634; (PRINGSHEIM and ARONOWSKY), A., i, 635.
preparation of, from artichoke tubers (WILLAMAN), A., i, 434.
constitution of, and its methylation (IRVINE, STEELE, and SHANNON), T., 1060.
digestion of (SHIMIZU), A., i, 82.
compounds of potassium and sodium hydroxides with (KARRER, STAUB, and WÄLTJ), A., i, 229.
detection of (CASTELLANI and TAYLOR), A., ii, 879.
- Invertase** (WILLSTÄTTER and RACKE), A., i, 598; (WILLSTÄTTER, GRABER, and KUHN), A., i, 1200.
action of (VOSBURGH), A., i, 64; (NELSON and HITCHCOCK), A., i, 388.
activity of, when adsorbed (NELSON and HITCHCOCK), A., i, 184.
action of nitrophenols on (RONA and BACH), A., i, 65.
action of quinine and narcotics on (RONA and BLOCH), A., i, 65; (RONA, ATRILA, and LASNITZKI), A., i, 959.
activity of, in yeast (MILLER), A., i, 202.
estimation of the activity of (HARDING), A., ii, 800.
- Iodates**. See under Iodine.
- Iodimetry**, potassium dichromate as a standard in (VOSBURGH), A., ii, 863.
- Iodine** in species of *Laminaria* (FREUNDLER, MENAGER, and LAURENT), A., i, 98.
absorption spectrum of (LINDSAY), A., ii, 599.
absorption and fluorescence spectra of (PRINGSHEIM), A., ii, 178.
emission spectrum of the vapour of (LANDAU-ZIERNECKI), A., ii, 674.
resonance spectrum of (MECKE), A., ii, 177; (PRINGSHEIM), A., ii, 178.
electrodeless discharge in the vapour of (ROBERTSON), A., ii, 610.
fractional diffusion of the vapour of (KOBLEWEILER), A., ii, 497.
absorption of active hydrogen and nitrogen by (WENDT; NEWMAN), A., ii, 639.
adsorption of, by silver iodide (GERMANN and TRAXLER), A., ii, 371.
colour of solutions of, at low temperatures (PICCARD and HERRMANN), A., ii, 655.
kinetics of alkaline solutions of (LIÉVIN), A., ii, 859.
equilibrium between sulphur and, in solution (AMADORI), A., ii, 561.

- Iodine**, comparative purity of various specimens of (FOULK and MORRIS), A., ii, 810.
- action of potassium azide with, in presence of carbon disulphide (BROWNE and HOEL), A., ii, 847.
- compounds of starch with (v. EULER and MYRBÄCK), A., i, 527, 1120; (v. EULER and BERGMAN; v. EULER and LANDERGHEN), A., i, 921.
- action of sulphurous acid with (MACAULAY), T., 552.
- introduction of, into the human system, and its elimination (BOURGUIGNON and CONDUOHÉ), A., i, 704.
- Iodine pentoxide** indicator for estimation of carbon monoxide (KATZ and BLOOMFIELD), A., ii, 585.
- Hydriodic acid**, heats of neutralisation of (RICHARDS and ROWE), A., ii, 425.
- photochemical oxidation of (WINTHER), A., ii, 808.
- Iodides**, photochemistry of the reaction between ferric salts and (SASAKI), A., ii, 772.
- influence of ferrous oxide on the action of hydrogen peroxide on (PAMFILOV and PETIN), A., ii, 835.
- action of nitrous acid on, in presence of oxygen (LOMBARD), A., ii, 313.
- estimation of, in presence of bromides and chlorides (KOLTHOFF), A., ii, 581.
- estimation of, by the formation of cyanogen iodide (LANG), A., ii, 782.
- Iodic acid**, reaction between sulphurous acid and (SKRABAL), A., ii, 488.
- use of, in detection of organic bases (ROSENTHALER), A., ii, 327.
- estimation of (KIKUCHI), A., ii, 519.
- Iodates**, reflection spectra of (SCHAEFER and SCHUBERT), A., ii, 180.
- detection of, in potassium iodide (LAGHARTRE), A., ii, 713.
- Hypiodous acid**, dissociation constant of (FURTH), A., ii, 281.
- Iodine estimation** :—
- estimation of, in animal tissues (DAMIENS), A., ii, 79.
- Iodine electrode**. See **Electrode**.
- Iodoform**, action of light on (BUTLER), A., ii, 604.
- action of, on magnesium alkyl haloids (BINAGHI), A., i, 1002.
- action of, on magnesium alkyl and aryl bromides (ODNO and BINAGHI), A., i, 313.
- Iodoform**, compounds of, with salts of organic bases (STEINKOPF and SCHWEN), A., i, 118.
- Ions**, structure of, and isomorphism (GRIMM), A., ii, 483.
- properties of, and chemical facts (GRIMM), A., ii, 690.
- existence of homogeneous groups of large (NOLAN), A., ii, 251.
- adsorption of (KOLTHOFF), A., ii, 197; (MUKHERJEE), A., ii, 689.
- hydration of (ASCHKENASI), A., ii, 258.
- antagonism of (BRENNER), A., i, 907.
- specific interaction of (BRÖNSTED), A., ii, 481.
- relative toxicity of (CAMERON and HOLLENBERG), A., i, 499.
- electrolytic. See **Electrolytic ions**.
- Ionisation**, theories of (FREDENHAGEN), A., ii, 112.
- by α -rays (WILKINS), A., ii, 607.
- and lattice energy, in relation to ionic properties (GRIMM), A., ii, 690.
- influence of the solvent on (NEALE), A., ii, 420.
- in electric furnace experiments (KING), A., ii, 810.
- experiments on, with a tungsten furnace (COMPTON), A., ii, 822.
- amphoteric (NOYES and WILSON), A., ii, 692.
- thermal, of metallic vapours (TOLMAN), A., ii, 18.
- of aliphatic acids (WINDISCH and OSSWALD), A., ii, 123.
- of gaseous elements at high temperatures (NOYES and WILSON), A., ii, 810.
- of weak electrolytes (SCHOORL), A., ii, 31.
- Ionisation potentials**, measurement of, in thermionic valves (PALMER), A., ii, 108.
- of gases (BOUCHER), A., ii, 608.
- Ionium** in radium residues (RONA), A., ii, 250.
- Ionogen linking**, nature of the (KAUFFMANN), A., ii, 366.
- Ipyrolic acid**, constitution of (ASAHINA and SHIMIDZU), A., i, 506.
- Iridium** :—
- Chloroiridates of cobalt- and chromium-ammines (BENRATH, BUCHER, and ECKSTEIN), A., ii, 515.
- Iridium organic compounds** with pyridine (DELEPINE), A., i, 859.
- Iron**, structure of (MÜGGE), A., ii, 381.
- spectrum of (MÜLLER), A., ii, 725.
- mass spectrum of (ASTON), A., ii, 710.
- Röntgen-ray spectrum of (DUANE and FRICKE), A., ii, 804.

Iron, emissivity of (LUBOWSKY), A., ii, 108.
 as a photochemical catalyst (BAUDISCH and BASS), A., i, 993.
 diffusion of carbon in mixed crystals of (TAMMANN and SCHÖNERT), A., ii, 772.
 diffusion of hydrogen through (SCHMIDT and LÜCKE), A., ii, 198.
 crystalline structure of (WESTGREN and LINDH), A., ii, 152; (WESTGREN and PHRAGMÉN), A., ii, 711.
 rate of solution of, in ferric alum (COLLENBERG and BODFORSS), A., ii, 431.
 rate of solution of, in dilute sulphuric acid (FRIEND and DENNETT), T., 41.
 and its oxides, equilibrium of, with carbon and its oxides (REINDERS and VAN GRONINGEN), A., ii, 153; (EASTMAN), A., ii, 506.
 meteoric. See Meteoric iron.
 "saccharated," colloidal nature of (MANNICH and ROJAHN), A., i, 718.
Iron alloys with carbon (HONDA), A., ii, 711.
 with carbon and boron (VOGEL and TAMMANN), A., ii, 852.
 with carbon and chromium (DAEVES), A., ii, 70.
 with carbon and tungsten (DAEVES), A., ii, 70.
 with manganese, electrochemistry of (TAMMANN and VADERS), A., ii, 380.
 with uranium (POLUSHKIN), A., ii, 152.
Iron salts, excretion of, in urine (KISCH), A., i, 898; (EHRENBERG and KARSTEN), A., i, 967.
Iron pentacarbonyl (MOND and WALLIS), T., 30.
 hydroxide, adsorption of, by precipitated manganese dioxide (GELOSO), A., ii, 589.
 hydroxides, naturally-occurring (WILLMANN), A., ii, 76.
 nitrosocarbonyl (MOND and WALLIS), T., 35.
 oxide, separation of, from calcium oxide (CHARRIOU), A., ii, 319.
Ferric salts, photochemistry of the reaction between iodides and (SASAKI), A., ii, 772.
 reduction of, by mercury (McCAY and ANDERSON), A., ii, 217.
 chloride, photochemical dissociation of (PUXEDDU and VODRET), A., ii, 415.
 chloride, reduction of (PICKLES), A., ii, 299.

Iron:—

Ferric hydroxide, colloidal, preparation of (BRADFIELD), A., ii, 507.
 adsorption of sodium alizarin by (BULL and ADAMS), A., i, 355.
 adsorption of radium-B and -C by (CRANSTON and HUTTON), T., 2843.
 oxide, heat of coagulation of, by electrolytes (BROWNE and MATHEWS), A., ii, 196.
 equilibrium of sulphur trioxide, water and (POSNJAK and MERWIN), A., ii, 772.
 colour of (HEDVALL), A., ii, 381.
 precipitation of calcium oxide with (CHARRIOU), A., ii, 163.
 separation of aluminium oxide and, from magnesium oxide (CHARRIOU), A., ii, 878.
 sulphates, equilibrium of formation of (APPLEBEY and WILKES), T., 337.
Ferrous chloride, equilibrium in the system ammonium chloride, water, and (CLENDDINEN), T., 801.
 hydroxide, reducing action of (MIYAMOTO), A., ii, 647.
 oxide, mechanism of the catalytic action of (PANFILOV and PETIN), A., ii, 835.
 sulphate, compound of chlorine and (RÖHM), A., ii, 648.
Iron organic compounds with phosphoric acids (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 631.
Ferric salts of 1:2:3:4-benzodiquinone-1:3-dioxime (MORGAN and MOSS), T., 2857.
 with pyridine or quinoline (WEINLAND and KISSLING), A., i, 863.
Ferrous ethyl iodide (JOB and REICH), A., i, 645.
Iron, cast, estimation of carbon in (BATA and THYSEN), A., ii, 395.
 estimation of sulphur in (MARINOT), A., ii, 224.
Steel, crystalline structure of (WESTGREN and LINDH), A., ii, 152; (WESTGREN and PHRAGMÉN), A., ii, 711.
 tempering of (POUCHOLLE), A., ii, 299; (CHARPY and GRENET), A., ii, 507.
 A¹ transformation and quenching of (CHIKASHIGE), A., ii, 852.
 solubility limits of carbon in (DAEVES), A., ii, 70.
 solid sodium hydroxide as solvent for carbon dioxide in analysis of (KELLEY and EVERS), A., ii, 160.

Iron, cast:—

- Steel, detection and estimation of vanadium in (MISSON), A., ii, 459.
- estimation of carbon in (BATA and THYSEN), A., ii, 395.
- estimation of chromium in (LOSANA and CAROZZI), A., ii, 589.
- estimation of chromium and nickel in (SIMION), A., ii, 529.
- estimation of cobalt in (EDER), A., ii, 458.
- estimation of manganese in colorimetrically (HESLINGA), A., ii, 660.
- estimation of nickel in (RUBRICIUS), A., ii, 183.
- estimation of sulphur in (MARINOT), A., ii, 224.
- estimation of vanadium in (CAIN and HOSTETTER), A., ii, 320; (KROPF), A., ii, 590.

Iron estimation and separation:—

- estimation of (KIKUCHI; KANÔ), A., ii, 721.
- estimation of, after reduction with hydrogen sulphide (LUNDELL and KNOWLES), A., ii, 88.
- estimation of, in presence of copper (WARK), T., 358; (THORNTON), A., ii, 528.
- estimation of, in blood (BROWN), A., ii, 319.
- estimation of phosphorus in (MURMANN), A., ii, 226.
- separation of aluminium and (CHALUPNY and BREISCH), A., ii, 588.
- separation of manganese from (CARUS), A., ii, 88.

Iron pyrites. See Pyrites.

Isatin, and its derivatives (BORSCHÉ and MEYER), A., i, 53.

Isatins, silver salts and ethers of (HANTZSCH and MEYER), A., i, 1178.

Isatins, bromo- and chloro-, and their derivatives (GRANDMOUGIN), A., i, 368.
 5:7-dibromo-, 5:7-dichloro-, and 4-chloro-5-bromo-, and their silver compounds and derivatives (HELLER, BENADE, and HOCHMUTH), A., i, 1059.

Isatin series, isomerism in the (HELLER), A., i, 1058; (HANTZSCH), A., i, 1177.

Isatin- α -benzoyloxime, and its alkali salts (HANTZSCH and MEYER), A., i, 1179.

Isatinhydrazone (BORSCHÉ and MEYER), A., i, 54.

Isatin-5-sulphonic acid, and its potassium and sodium salts, hydrazones of (MARTINER), A., i, 279.

Isatin-6-sulphonic acid, sodium salt, hydrazones of (MARTINER), A., i, 278.

Isatin-yellow series, colouring matters of the (MARTINER), A., i, 278.

Isatogenic acid, ethyl ester, oxime (HELLER and BOESSNECK), A., i, 370.

Isatogens (HELLER and BOESSNECK), A., i, 369.

Isatoic acid, 5:7-dibromo-, and 5:7-dichloro-, and 4-chloro-5-bromo-, and their salts and derivatives (HELLER, BENADE, and HOCHMUTH), A., i, 1059.

Isatoids (HELLER and BENADE), A., i, 582.

Isatole, 5:7-dichloro- (HELLER, BENADE, and HOCHMUTH), A., i, 1060.

Ishikawaite (SHIBATA and KIMURA), A., ii, 361.

Isomerism, geometric, in relation to the formation of films (ADAM), A., ii, 424.

ψ -Isomerism, Hantzsch's theory of (MADELUNG), A., ii, 344.

Isomorphism, ionic structure and (GRIMM), A., ii, 483.

Isoprene in relation to caoutchouc (STAUDINGER and FRITSCHI), A., i, 1043.

bromination of (BERGMANN), A., i, 1106.

addition of hydrogen haloids to (STAUDINGER, KREIS, and SCHILT), A., i, 978.

dibromide and dibromo- (STAUDINGER, MUNTWYLER, and KUPFER), A., i, 979.

Isotopes, structure of (NEUBURGER), A., ii, 107.

band spectra of (GREBE and KONEN), A., ii, 4.

difference between series spectra of (EHRENFEST; BOHR), A., ii, 598; (NICHOLSON), A., ii, 599.

and the transformation of atoms (MEITNER), A., ii, 15.

separation of (MULLIKEN and HARKINS), A., ii, 295; (MULLIKEN), A., ii, 492; (HARKINS), A., ii, 702.

J.

***Jatropha curcas*, oil from seeds of (GRIMME), A., i, 98.**

Jaundice, bile pigments in (ROSENTHAL and MEIER), A., i, 198.

Jejunum, human, sucrose-splitting enzyme in the (v. EULER and MYRBÄCK), A., i, 295.

Juglone. See 1:4-Naphthaquinone, 5-hydroxy-.

***Juncus effusus*, constituents of (ZELLNER), A., i, 1098.**

Juniper oil from Cyrenaica (MASSERA), A., i, 945.

Juniperus oxycedrus, constituents of the wood of (HUERRE), A., i, 505.
Jute. See *Corchorus capsularis*.

K.

Kagurassame oil, constituents of (TSUJIMOTO and TOYAMA), A., i, 297.

Kaolinite (*kaolin*), decomposition of, by micro-organisms (VERNADSKY), A., i, 1096.

Kasolite, composition and radioactivity of (SCHOEP), A., ii, 154.

Kataphoresis of metallic compounds of proteins (BENEDICENTI and REBELLO-ALVES), A., ii, 683.
 of water in organic liquids (GYEMANT), A., ii, 684.

Kawaic acid (MURAYAMA and MAYEDA), A., i, 265.

β -Kawaic acid (MURAYAMA and MAYEDA), A., i, 265.

Kawa-kawa resin (MURAYAMA and MAYEDA), A., i, 265.

"**Kazunoko**." See Herring roe.

Keeleyite from Bolivia (GORDON), A., ii, 859.

Keratin (HEIDUSCHKA and KOMM), A., i, 967.

Ketens (STAUDINGER and HAUSER), A., i, 28; (STAUDINGER), A., i, 238, 240; (STAUDINGER and RATHSAM), A., i, 1014; (STAUDINGER and MEYER), A., i, 1015.

dimeric (DIECKMANN and WITTMANN), A., i, 1156.

Ketenacetals (SCHEIBLER and ZIEGNER), A., i, 426; (STAUDINGER and RATHSAM), A., i, 1014; (STAUDINGER and MEYER), A., i, 1015.

Ketenethyimine (STAUDINGER and HAUSER), A., i, 29.

Ketenimine derivatives (STAUDINGER and HAUSER), A., i, 28.

Ketenium compounds (WEDERKIND and WEINAND), A., i, 234.

Ketenphenylimine (STAUDINGER and HAUSER), A., i, 29.

Ketentriethylum, bromo-, and mono- and di-chloro- (WEDERKIND and WEINAND), A., i, 235.

Keto-acids, degradation of, in the animal organism (WORINGER), A., i, 90.

1-Keto 2 aldehydoacenaphthene, 5(or 6)-bromo-, and its derivatives (MAYER and SCHÖNFELDER), A., i, 1167.

1-Keto-5-aminotetrahydronaphthalene, 5-acetyl derivative (SCHROETER and TETRALIN G. M. B. H.), A., i, 1036.

Keto-anils (REDELLEN), A., i, 145; (KNOEVENAGEL and BAHR), A., i, 750; (KNOEVENAGEL and GOOS), A., i, 751; (KNOEVENAGEL), A., i, 1060.

β -Keto-bases, synthesis of (MANNICH and HEILNER), A., i, 351.

δ -Ketocamphane- ϵ -pi-carboxylic acid. See *p*-Camphocarboxylic acid.

α -Keto- α -campholenic acid, and its semicarbazone (CHANDRASENA and INGOLD), T., 1554.

δ -Keto- β -carboxy- $\gamma\gamma$ -dimethylhexoic acid (CHANDRASENA, INGOLD, and THORPE), T., 1549.

1-Keto-2-cinnamylidenetetrahydronaphthalene (HERZOG and KREIDL), A., i, 1168.

α -Keto- $\beta\beta$ -diethylglutaric acid, and its calcium salt and semicarbazone (DESHAPANDE and THORPE), T., 1439.

5-Keto-4:7-dimethyl-4:5-dihydroindole-diazine(1:4) (KERMACK, PERKIN, and ROBINSON), T., 1888.

5-Keto-1:6-dimethyl-4-isopropyltetrahydronaphthalene (RUZICKA and SEIDEL), A., i, 562.

Ketodinitrones, formation and properties of (ALESSANDRI), A., i, 558.

Keto-enolic mixtures, spectrochemical and volumetric analysis of (v. AUWERS and JACOBSEN), A., ii, 168.

3-Keto-2-ethoxydihydroindole-2-carboxylic acid, 1-hydroxy-, ethyl and methyl esters (HELLER and BOESNECK), A., i, 370.

Ketogenic balance, calculation of, from respiratory quotients (SHAFFER), A., i, 83.

Ketogenesis (WILDER and WINTER), A., i, 893.

5-Keto- d -gluconic acid semicarbazone-semicarbazide (KILIANI), A., i, 1112.

α -Ketoglutaric acid, condensation of, with *o*-phenylenediamine (KON, STEVENSON, and THORPE), T., 664.

α -1-Keto-2-hydroxy-3-methyltetrahydronaphthalene-3-acetic acid, lactone, and its silver salt and derivatives (KON, STEVENSON, and THORPE), T., 660.

1-Keto-5- and -8-hydroxytetrahydronaphthalenes, and their derivatives (SCHROETER and TETRALIN G. M. B. H.), A., i, 1036.

3-Keto-2-methoxydihydroindole-2-carboxylic acid, 1-hydroxy-, ethyl and methyl esters (HELLER and BOESNECK), A., i, 369.

1-Keto-2-methyl-2:3:4:4-tetrachlorotetrahydronaphthalene (FRIES and LOHMANN), A., i, 29.

- 3-Keto-4-methyl-3:4-dihydro-4-carboline** (KERMACK, PERKIN, and ROBINSON), T., 1887.
- 5-Keto-1-methyl-4-isopropyltetrahydronaphthalene**, and its semicarbazone (RUZICKA and MINGAZZINI), A., i, 1001.
- 2-Keto-1-methyltetrahydronaphthalene**, 1:3:4:4-tetrachloro- (FRIES), A., i, 43.
- 1-Ketopernaphthindane**, 3:3-dichloro- (FLEISCHER and RETZE), A., i, 1138.
- 2-Keto-1:3:4-naphthoisotriazine-4-carboxylic acid**, ethyl ester (DIELS and SÖRENSEN), A., i, 1194.
- 2-Keto-1:3:4-naphthoisotriazine-6-sulphonic acid** (DIELS and SÖRENSEN), A., i, 1194.
- Ketones**, preparation of (BADISCHE ANILIN- & SODA FABRIK), A., i, 218; (HUNT), A., i, 810.
- preparation of, from mineral and tar oils (HARRIES), A., i, 514.
- conductivity of mixtures of acids with (KENDALL and GROSS), A., ii, 33.
- halochromism of (HANTZSCH), A., i, 556.
- action of phosphorus trichloride on (CONANT, MACDONALD, and KINNEY), A., i, 186.
- interaction of thiocarbamides and (TAYLOR), T., 2267.
- sodium derivatives, action of acetylene on (LOCQUIN and WOUSENG), A., i, 617.
- aliphatic, decomposition of (MAILHE), A., i, 985.
- action of trimethylene chlorobromide on (BILLON), A., i, 717.
- aliphatic-aromatic, melting points of (MAJIMA, NAGAOKA, and TAMADA), A., i, 257.
- aromatic, synthesis of, by means of zinc organic compounds (MAUTHNER), A., i, 457.
- cyclic, of the quinoline series, preparation of (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 867.
- unsaturated, reactivity of (BUCK and HEILBRON), T., 1095.
- formation of, from tertiary ethinyl-carbinols (MEYER and SCHUSTER), A., i, 556.
- detection of (BOUGAULT and GROS), A., ii, 666.
- estimation of, with hydroxylamine (BENNETT and DONOVAN), A., ii, 535.
- Ketones, α -halogenated**, action of amines with (MAYER and FREUND), A., i, 865.
- Ketones, α -halogenated**, condensation of aldehydes with (BAUER and WERNER), A., i, 1084.
- oximino-, bisulphite compounds of (GASTALDI and BRAUNIZER), A., i, 626.
- dithio-, formation and properties of (NAIK and AVASARE), T., 2592.
- Ketone-alcohol**, $C_6H_5O_2Cl$, and its derivatives, from *o*-chlorobenzaldehyde and acetaldehyde (NEUBERG and LIEBERMANN), A., i, 305.
- Ketonic compounds**, rôle of, in the formation of sugars (GERLMUYDEN), A., i, 607.
- β -Ketonic esters**, action of bromine on (KÖHLER), A., i, 552.
- 3-Keto-2-*o*-nitrophenyl-3:4-dihydroquinoline** (NEBER), A., i, 545.
- 4-Keto-*s*-octahydroanthracene**, and its semicarbazone (SCHROETER and TETRALIN G. M. B. H.), A., i, 1136.
- 4-Keto-*s*-octahydrophenanthrene**, and its semicarbazones (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- 2-Keto-4-phenyl-1-methyl-1:2-dihydropyrimidine**, and its 5:6-dibromide, and 6(or 5)-bromo- (MANNICH and HEILNER), A., i, 372.
- 2-Keto-4-phenyl-1-methylhexahydropyrimidine** (MANNICH and HEILNER), A., i, 372.
- 2-Keto-4-phenyl-1-methyl-1:2:5:6-tetrahydropyrimidine**, and its bromo-derivative (MANNICH and HEILNER), A., i, 372.
- Ketopinic anhydride** (WEDEKIND and WEINAND), A., i, 549.
- Ketopinyl chloride**, and its action with triethylamine (WEDEKIND and WEINAND), A., i, 548.
- 4-Ketopiperidinepropionic acid**, ethyl ester (RUZICKA and SWEDEL), A., i, 1057.
- Ketorhammonic acid**, constitution of (KILIANI), A., i, 1112.
- α -Ketorhammonic acid**, lactone of, and its *p*-nitrophenylhydrazone (KILIANI), A., i, 224.
- 2-Keto-6-sulpho-1:3:4-naphthoisotriazine-4-carboxylic acid**, ethyl ester (DIELS and SÖRENSEN), A., i, 1194.
- α -Ketotetrahydronaphthalene** (SCHROETER and TETRALIN G. M. B. H.), A., i, 1036.
- ac*-1-Ketotetrahydronaphthalene-3-acetic acid**, and its semicarbazone (STEVENSON and THORPE), T., 1721.
- 1-Ketotetrahydronaphthalene-3-sulphonic acid**, 5-hydroxy-, and its phenylhydrazone with its metallic salts (FUCHS and STIX), A., i, 451.

Ketotricarboxylic acids, $C_{12}H_{14}O_7$, from pyrocholoidanic acid and alkali hydroxide (WIELAND and SCHULENBURG), A., i, 848.

α -Keto- $\beta\beta$ -trimethylsuccinic acid, and its condensation products with phenylenediamines (KON, STEVENSON, and THORPE), T., 663.

Ketoximes, thermochemistry of (DORABIALSKA), A., ii, 548.

Kidneys, permeability of the glomerulus membrane of, for sugars (HAMBURGER), A., i, 490, 491, 790, 919. proteolytic enzymes of (HEDIN), A., i, 1212.

Kinetics of unimolecular reactions (GARNER), A., ii, 358.

Kojic acid, and bromo-, and hydroxy-, and their salts and derivatives (YABUTA), A., i, 939.

Krypton, estimation of (MOUREU and LEPAPE), A., ii, 394.

Kynurenic acid, constitution and derivatives of (SPÄTH), A., i, 173.

L.

Lac, Japanese, constituents of (MAJIMA), A., i, 262, 263.

Lacöl dimethyl ether ozonide (MAJIMA and KUDO), A., i, 264.

α -Lactanilide, *S*-trichloro- (PASSERINI), A., i, 731.

Lactase in yeast (WILLSTÄTTER and OPPENHEIMER), A., i, 203.

Lactic acid, formation of, from dextrose by peptone (SCHLATTER), A., i, 1096. synthesis of inactive (PIRAK), A., i, 810. heat of combustion of (MEYERHOF), A., ii, 475.

action of thionyl chloride on (BLAISE and MONTAGNE), A., i, 715.

in plants (FRANZEN and KEYSSNER), A., i, 310; (FRANZEN and STERN), A., i, 311.

destruction of, by yeast (LIEBEN), A., i, 796; (v. FÜRTH and LIEBEN), A., i, 1219.

by yeast and by blood-cells (v. FÜRTH and LIEBEN), A., i, 502. in blood (HASTINGS), A., i, 81.

formation of, in muscle (MEYERHOF), A., i, 86.

function of, in muscle (FOSTER and MOYLE), A., i, 398.

effect of sugar injections on the excretion of (MORACZEWSKI and LINDNER), A., i, 402.

detection of, in organic liquids (D'ARBELA), A., ii, 480.

detection of, in tissues (LAQUER), A., ii, 325.

estimation of (CLAUSEN), A., ii, 593.

Lactone, $C_6H_{10}O_2$, from cyclohexanone oxime (HELFEHICH and MALIKOMES), A., i, 431.

Lactones, velocity of formation and hydrolysis of (KAILAN and NEUMANN), A., ii, 433.

β -Lactones (JOHANSSON and HAGMAN), A., i, 425.

Lactose (*milk-sugar*), action of ozone on (SCHONEBAUM), A., i, 717.

detection of, in urine (HERZBERG), A., ii, 167.

micro-estimation of, in milk (FONTES and THIVOLLE), A., ii, 323.

Lævidulinase (MAYEDA), A., i, 694.

Lævoglucoan, formation of, by distillation of carbohydrates (KARRER and ROSENBERG), A., i, 812.

polymerides of (PICTET and ROSS), A., i, 525.

Lævulic acid α -alkyl derivatives of (GAULT and SALOMON), A., i, 427.

Lævulose (*d-fructose*; *fruit sugar*), preparation of (HARDING), A., i, 919.

action of ammonia on (LING and NANJJI), A., i, 631.

action of ozone on solutions of (SCHONEBAUM), A., i, 223.

acetone derivatives of (IRVINE and PATTERSON), T., 2146.

constitution and rotation of compounds of, with boric acid and sodium hydroxide (GILMOUR), T., 1333.

decomposition of, by pyrocyanic bacilli (AUBEL), A., i, 201.

effect of, in diabetes (DESGREZ, BERRY, and RATHERY), A., i, 1215.

detection of, in presence of aldoses (KOLTHOFF), A., ii, 166.

estimation of, in straw (COLLINS), A., ii, 323.

Lævulosecarboxylic acid, *l*-phenylhydrazide of (KILIANI), A., i, 1112.

Lævulosephosphoric acids, iron salts (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 631.

Lævuloseuria (BARRENSCHEEN), A., i, 496.

Laminaria, iodine in species of (FREUNDLER, MENAGER, and LAURENT), A., i, 98.

Laminarin, action of ferments on (GRUZEWSKA), A., i, 11.

Lamp, sodium, for polarimetry (FALES and MORRELL), A., ii, 43.

Landolt's reaction (SREBAL), A., ii, 488.

Lantana camara, oil from the leaves of (MOUDGILL and VRIDHACHALAM), A., i, 754.

Lanthanum, atomic weight of (BAXTER), A., ii, 298; (HOPKINS and DRIGGS), A., ii, 770.

spectrum of (ÖKUBO), A., ii, 676.

- Lanthanum**, arc spectra of (KIESS, HOPKINS, and KREMERS), A., ii, 244.
- Lanthanum oxide**, density and molecular volume of (PRANDTL), A., ii, 379.
- Laudanine**, synthesis of (SPATH and LANG), A., i, 568.
- Laudanum**, detection of saffron in (GUERBET), A., ii, 793.
- Lauric acid**, potassium salt, ultra-filtration of solutions of (MCBAIN and JENKINS), T., 2325.
- sodium salt, equilibrium of sodium chloride, water and (MCBAIN and BURNETT), T., 1320.
- cellulose ester (GRUN and WITKA), A., i, 114.
- Lavender oil**, constituents of (ROURE-BERTRAND FILS), A., i, 847.
- Lead** from uranium minerals of Madagascar (MUGUET), A., ii, 216.
- arc spectrum of (HEMSALECH and DE GRAMONT), A., ii, 176.
- band spectra of, and of uranium-lead (GREBE and KONEN), A., ii, 4.
- electro-deposition of (HUGHES), A., ii, 446.
- and of its dioxide (MELLON and REINHARD), A., ii, 817.
- separation of the isotopes of (DILLON, CLARKE, and HINCHY), A., ii, 710.
- vapour pressure curve of (INGOLD), T., 2419.
- liquid, density and surface tension of (HOGNESS), A., ii, 29.
- solubility of bismuth in (DI CAPUA), A., ii, 576.
- solubility of cadmium in (DI CAPUA), A., ii, 576.
- test-papers, preparation of (OLIVIER), A., ii, 396.
- reductions with, in volumetric analysis (TREADWELL), A., ii, 780.
- poisoning. See Poisoning.
- Lead alloys** with mercury, estimation of lead in (MELLON and REINHARD), A., ii, 787; (MELLON), A., ii, 870.
- with strontium, constitution of (PIWOWARSKY), A., ii, 644.
- with tellurium (DREIFUSS), A., ii, 508.
- Lead salts**, basic, constitution of (WEINLAND and STROH), A., ii, 767.
- soluble, action of, on plant growth (BONNET), A., i, 412.
- Lead azide**, abnormal crystallisation of (LOWNDEN), A., ii, 216.
- chromate, solvolysis of (HICKS and CRAIG), A., ii, 622.
- chromates, basic (HICKS), A., ii, 147.
- equilibria of (JAEGER and GERMS), A., ii, 65.
- Lead haloids**, compounds of ammonia with (BILTZ and FISCHER), A., ii, 851.
- molybdates, equilibria of (JAEGER and GERMS), A., ii, 65.
- nitrate, crystal structure of (VEGARD), A., ii, 503.
- electrometric titrations with (KOLTHOFF), A., ii, 781.
- thallous nitrite (CUTTICA and PACIELLO), A., ii, 378.
- oxides, physical chemistry of (GLASSTONE), T., 58, 1456, 1469, 2091.
- monoxide, isomerism of (APPLEBEY and REID), T., 2129.
- hydrates of (GLASSTONE), T., 58.
- density of (BROWN, COOK, and WARNER), A., ii, 571.
- fusion diagram of mixtures of bismuth oxide and (BELLADEN), A., ii, 777.
- sesquioxide (GLASSTONE), T., 1456.
- dioxide, electromotive properties of (GLASSTONE), T., 1469.
- peroxide, estimation of, in minium (BONIS), A., ii, 587.
- Triplumbic tetroxide** (GLASSTONE), T., 1456.
- Lead sulphates**, equilibria of (JAEGER and GERMS), A., ii, 65.
- sulphide, solubility of (TRUMPLER), A., ii, 111.
- tungstates, equilibria of (JAEGER and GERMS), A., ii, 65.
- Plumbates**, preparation and electrochemistry of (GRUBE), A., ii, 570.
- Lead organic compounds**, action of thallic chloride on (D. and A. E. GODDARD), T., 260.
- diphenyl, salts of, with organic acids (GODDARD, ASHLEY, and EVANS), T., 981.
- di-*p*-tolyl, triphenyl, tritolyls, and tri-*p*-xylyl, and their haloids (KRAUSE and REISSAUS), A., i, 602.
- tetraphenyl, use of, in preparation of organo-metallic compounds (GODDARD, ASHLEY, and EVANS), T., 978.
- trialkyl and triaryl fluorides (KRAUSE and FOHLAND), A., i, 644.
- Lead detection, estimation and separation** :—
- detection of small quantities of, in urine (SCHUMM), A., ii, 317.
- estimation of (FAIRHALL), A., ii, 659.
- estimation of, electro-volumetrically (MACINNES and TOWNSEND), A., ii, 456.
- estimation of, volumetrically (JELLINEK and ENS), A., ii, 864.

- Lead detection, estimation, and separation:**—
 estimation of, volumetrically, with ferrocyanides (TREADWELL and CHERVER), A., ii, 786.
 estimation of, in lead amalgams (MELLON and REINHARD), A., ii, 787; (MELLON), A., ii, 870.
 estimation of, in water (AVERY, HEMINGWAY, ANDERSON, and READ), A., ii, 161.
 estimation of, and its separation from antimony, copper and tin (KLING and LASSIEUR), A., ii, 86.
- Lead anode.** See Anode.
- Lead tree, formation and disglomeration of** (THIEL), A., ii, 63.
- Leather, dyeing, tanning and preservation of** (RAKUZIN), A., i, 702.
- Leaves, distribution of carbohydrates in** (SCHROEDER and HORN), A., i, 906.
 disappearance of starch from (MOLISCH), A., i, 309.
 dead, carbohydrates in (COMBES and KOHLER), A., i, 1222.
 old and young, manganese content of (JADIN and ASTRUC), A., i, 1098.
 young, oxalic acid in (BAU), A., i, 309.
- Lecithin, precipitation of hydrosols of,** by electrolytes (KAKIUCHI), A., i, 711.
 from eggs, unsaturated fatty acids from (LEVENE and ROLF), A., i, 621.
 from liver, unsaturated fatty acids of (LEVENE and SIMMS), A., i, 424.
 estimation of, as ether-soluble phosphorus (LE BRETON), A., ii, 168.
 estimation of, in blood (RANDLES and KNUDSON), A., ii, 719.
- Lectures, delivered before the Chemical Society** (RUTHERFORD). T., 400; (GREGORY), T., 750; (BRAGG), T., 2766.
- Lecture experiments with atomic models** (KING), A., ii, 705.
 showing a time reaction (MEYER), A., ii, 636.
 on protective colloids (FRIEND), A., ii, 267.
 to show the detection and estimation of traces of copper (HAHN and LEIMBACH), A., ii, 870.
 with hydrazoic acid and azides (BROWNE and HOEL), A., ii, 840.
 to show the antiseptic action of mercury ions (JOACHIMOGLU), A., ii, 211.
- Leech.** See *Hirudo medicinalis*.
- Leguminosae, assimilation of carbon dioxide by** (KOSTYTSCHEV), A., i, 618.
- Lemon juice, growth-promoting factor of** (LEICHTENTRITT and ZIELASKOWSKI), A., i, 1094.
- Lens, crystalline, lipoids of the** (GOLD-SCHMIDT), A., i, 489.
 cleavage products from proteins of the (HIJIKATA), A., i, 489; (JESS), A., i, 1199.
- Leptospermum, essential oils from species of** (PENFOLD), A., i, 44.
- Letsche's acid.** See Biloidanic acid.
- L-Leucine, degradation of, by bacteria** (ARAI), A., i, 303.
- L-Leucinecholine, salts of** (KARRER, GISLER, HORLACHER, LOCHER, MÄDER, and THOMANN), A., i, 815.
- Leucocyanoselenopyronin** (BATTEGAY and HUGEL), A., i, 669.
- Leucocytes, effect of chemicals on the chemotaxis of** (WOLF), A., i, 401.
 glycolytic action of (FUKUSHIMA), A., i, 697.
- Leuconic acid, constitution of** (HOMOLKA), A., i, 630.
- Leucosin, isoelectric point of** (LÜERS and LANDAUER), A., i, 881.
- Lichenin, digestion of** (SHIMIZU), A., i, 82.
- Liesegang's rings, formation of** (McGUIGAN), A., ii, 38; (FISCHER and McLAUGHLIN), A., ii, 206.
- Light, emission of, by atoms** (FOSTERLING), A., ii, 729; (SEELIGER), A., ii, 804.
 scattering of, by dust-free liquids (MARTIN and LEHRMAN), A., ii, 335.
 molecular scattering of, in *n*-pentane (VENKATESWARAN), T., 2655.
 absorption of, by colouring matters (LABAREV), A., ii, 103, 332.
 by solutions (LUNELUND), A., ii, 8; (v. HALBAN and SIEDENTOFF), A., ii, 332.
- blue, from the Röntgen tube, polarisation of** (SEEMAN), A., ii, 8.
- ultra-violet, measurement of absorption of** (WINTHER, BAGGESGAARD-RASMUSSEN, and SCHREINER), A., ii, 729.
 action of, on gels (HOLMES and PATRICK), A., ii, 338.
 action of, on aqueous solutions of organic acids and their salts (JAEGER and BERGER), A., i, 316.
 action of, on colloidal platinum (SPEAR, JONES, NEAVE, and SHLAGER), A., ii, 11.

- Lignin** (MELANDER), A., i, 325.
 from straw (PASCHKE), A., i, 325.
 from winter rye straw (BECKMANN, LIESCHE, and LEHMANN), A., i, 233.
 methylation of (HEUSER, SCHMITT, and GUNKEL), A., i, 325.
 oxidation of (HEUSER and SAMUELSEN), A., i, 812.
 spontaneous oxidation of (SCHRADER), A., i, 637.
 reduction of, with hydriodic acid and phosphorus (WILLSTÄTTER and KALB), A., i, 939.
 pine (KLASON), A., i, 324, 413.
 detection of, in cells (VAN ZIJF), A., ii, 94.
 α -Lignin, constitution of (KLASON), A., i, 324.
Ligninsulphonic acids (McKEE and BARSKY), A., i, 1121.
Lignite, constituents of (CIUSA and CROCE), A., ii, 385.
 heat developed in adsorption of gases by (McLEAN), A., ii, 352.
 producer tar from (RUHEMANN), A., i, 22; (FISCHER), A., i, 330.
Lignoceric acid, and its derivatives (BRIGL and FUCHS), A., i, 712.
Lignocerin (BRIGL and FUCHS), A., i, 713.
Lignoceryl cyanide (LEVENE and TAYLOR), A., i, 715.
Lignoceryl alcohol, and its acetate (BRIGL and FUCHS), A., i, 713.
Lignocyanin (PASCHKE), A., i, 326.
Lignosulphonic acid (KLASON), A., i, 14.
 oxidation of (HEUSER and SAMUELSEN), A., i, 812.
 α -Lignosulphonic acid, and its salts (KLASON), A., i, 324.
Lilium candidum and *martagon*. See Lily.
 Lily, formation of anthocyanin in the bulb scales of (MIRANDE), A., i, 1100, 1224.
Lime-water, equilibrium between cement and (LORENZ and HAEGERMANN), A., ii, 59.
Linolohydroxamic acid (NICOLET and PELO), A., i, 644.
Linseed oil, drying of (SLANSKY), A., i, 1113.
 oxidation of (COFFEY), T., 18.
Linolic acid, methyl ester (BAUER and HARDEGG), A., i, 984.
Lipæmia (BLOOR), A., i, 89.
Lipase from castor beans, preparation and properties of (HALEY and LYMAN), A., i, 390.
 action of antiseptics on the activity of (PALMER), A., i, 886.
Lipase, action of quinine on (RONA and REINICKE), A., i, 67.
 liver, action of quinine and atoxyl on (RONA and PAVLOVIĆ), A., i, 887.
 stomach, activity of (SLUITER), A., i, 887.
Lipoids (FRÄNKEL and GILBERT; FRÄNKEL and KASZ), A., i, 294.
 physical chemistry of (LOEWY), A., ii, 354.
 ultramicroscopy of (HATTORI), A., i, 192.
 action of, on metabolism in plant cells (BOAS), A., i, 94, 613.
 nutritive value of (TAKAHASHI), A., i, 699.
 estimation of, in blood (LEMELAND), A., i, 666.
Liquids, physical properties of, in equilibrium with their vapour (SWIEN-TOSLAWSKI), A., ii, 192.
 electromotive force and concentration at the junction of (FREUNDLICH), A., ii, 19; (FREUNDLICH and GYEMANT), A., ii, 342.
 density of (WIEDEBRAUK), A., ii, 740.
 equation of compression of (CARL), A., ii, 477.
 with intersecting vapour pressure curves (v. RECHENBERG), A., ii, 120.
 solubility of, at high pressures (TIMMERMANS), A., ii, 482.
 solubility of gases in (NEUHAUSEN), A., ii, 621.
 solubility of liquids in (GORDON and REID), A., ii, 824.
 partial miscibility of (DUBRISAY), A., ii, 428.
 surface pressure of thin layers of (MARCELIN), A., ii, 686.
 viscosity of (HERZ), A., ii, 821.
 calculation of viscosity and surface tension of (CREIGHTON), A., ii, 426.
 laws of internal friction of (HERZ), A., ii, 352.
 adsorption of, by charcoal (DRIVER and FIRTH), T., 2409; (HARKINS and EWING), A., ii, 123.
 change of properties of, on drying (BAKER), T., 568.
 molecular anisotropy in (RAMAN), A., ii, 603.
 extraction of, with the Soxhlet apparatus (HANDORF), A., ii, 496.
 anisotropic, structure of (FRIEDEL and ROYER), A., ii, 129.
 scattering of Röntgen rays by (HÜCKEL), A., ii, 14.
 inversion of rotation in (ROYER), A., ii, 415.
 dust-free, scattering of light by (MARTIN and LEHRMAN), A., ii, 335.

- Liquids**, foaming, distillation of (KLÄNHARDT), A., ii, 472.
 slightly miscible, determination of reciprocal solubility of (CHÉNEVEAU), A., ii, 355, 427.
 mixed, separation of, by centrifuging (MULLIKEN), A., ii, 686.
 separation of, by distillation (DUF-TON), T., 306.
 binary mixed, theory of (SCHMIDT), A., ii, 472.
 vapour pressure of (SCHMIDT), A., ii, 119; (FAUST), A., ii, 423; (CASSEL; SCHULZE), A., ii, 424.
 non-associated, vapour pressure and heat of vaporisation of (MORTIMER), A., ii, 615.
 non-miscible, apparatus for extraction of (FAYOLLE and LORMAND), A., ii, 759.
 organic, properties of (HERZ), A., ii, 23.
 refractive index of (HIMSTEDT and WERTHEIMER), A., ii, 725.
 freezing points of (TIMMERMAN, VAN DER HORST, and ONNES), A., ii, 258.
 partition coefficients between water and (SMITH), A., ii, 430, 431.
 internal friction of (HERZ), A., ii, 30.
 kataphoresis of water in (GYEMANT), A., ii, 684.
 velocity of hydrogenation of, in presence of nickel (ARMSTRONG and HILDITCH), A., ii, 41.
 binary organic, classification of (TIMMERMAN), A., ii, 25.
 supercooled, spontaneous crystallisation of (HINSHLWOOD and HARTLEY), A., ii, 200.
 superheated, pressure volume relation of (WISMER), A., ii, 425.
 undercooled, melting interval of (FERGUSON), A., ii, 614.
Lithium, isotopes of (DEMPSTER), A., ii, 417.
 arc spectra of isotopes of (McLENNAN and AINSLIE), A., ii, 541.
 line spectrum of (MERTON), A., ii, 803.
 crystal structure of (BIJVOET and KARSEN), A., ii, 499.
 attempt at transmutation of (WYCKOFF), A., ii, 642.
 ions, hydration of (BABOROVSKY and VELISEK), A., ii, 816.
 vapour, electrodeless discharge in (ROBERTSON), A., ii, 609.
Lithium salts of weak inorganic acids (ROSENHEIM and REGLIN), A., ii, 289.
Lithium antimoniodobromide (VOURNAZOS), A., ii, 651.
Lithium chloride, solubility of, in quinoline (WALTON and WISE), A., i, 370.
 cupric chloride, absorption of light by (GETMAN), A., ii, 412.
 chlorite (LEVI), A., ii, 567.
 hydride (EPHRAIM and MICHEL), A., ii, 58.
 structure of (BIJVOET and KARSEN), A., ii, 569.
 hydroxide, heats of neutralisation of (RICHARDS and ROWE), A., ii, 425.
 neutralisation of phenylstibinic acids by (SCHMIDT), A., i, 482.
 nitrate, solubility of, in ammonia (DAVIS, OLMSTEAD, and LUNDSTRUM), A., ii, 56.
n- and *iso*-Lithobillanic acids, and their derivatives (BORSCHÉ and HALLWASS), A., i, 1160.
Lithocholic acid, methyl ester (BORSCHÉ and HALLWASS), A., i, 1159.
Lithospermum erythrorhizon, colouring matter from the roots of (MAJIMA and KURODA), A., i, 946.
Liver, function of the (DELPRAT and WHIPPLE), A., i, 86; (PELKAN and WHIPPLE), A., i, 490.
 fat exchange in the (LOMBROSO), A., i, 702.
 spacial separation of glycogen and diastase in the cells of (LESSER), A., i, 195.
 basal metabolism and utilisation of protein in diseases of (AUB and MEANS), A., i, 193.
 yellow atrophy of (KAHN and BARSKY), A., i, 969.
 embryonic, glycogen formation in (ARON), A., i, 702.
Liver oil, detection of, with sulphuric acid (DRUMMOND and WATSON), A., ii, 665.
Loellingite, crystal structure of (HUGGINS), A., ii, 651.
Loranthaceæ, constituents of some (WESTER), A., i, 208.
Lubricating oils. See under Oils.
Lubrication, boundary, in relation to chemical constitution (DOUBLEDAY), T., 2875.
Lucerne (alfalfa), proteins of (OSBORNE, WAKEMAN, and LEAVENWORTH), A., i, 99.
 water-soluble constituents of (OSBORNE, WAKEMAN, and LEAVENWORTH), A., i, 1104.
 hay, nitrogen compounds in (MILLER), A., i, 414.
Luciferase, specificity of (HARVEY), A., i, 299.

- Luciferin**, specificity of (HARVEY), A., i, 299.
- Lumbang oil** (WEST and MONTES), A., i, 95.
- Luminescence** by flame excitation (HOWES), A., ii, 335; (NICHOLS and WILBER), A., ii, 806.
- inorganic (TIEDE and RICHTER), A., ii, 215.
- of metallic oxides (NICHOLS and WILBER), A., ii, 105.
- of incandescent solids (NICHOLS and HOWES), A., ii, 597.
- Lungs**, chemistry of the (SAMMARTINO), A., i, 298, 1089.
- Lung stones** (SCHERER), A., i, 966.
- Lupcol**, identity of xanthosterol with (ULTÉE), A., i, 826.
- Lupines**, effect of nitrogenous fertilisers on the alkaloids in (VOGEL and WEBER), A., i, 798.
- Lupinus albus**, effect of cocaine on the growth of (MACHT and LIVINGSTON), A., i, 798.
- Lutecium**, atomic number of (URBAIN), A., ii, 505.
- L*-series spectrum of (DAUVILLIER), A., ii, 463.
- Lutecobaltic chloride**. See Hexaminocobaltic chloride under Cobalt.
- Lythium salicaria**, constituents of (ZELLNER), A., i, 799.

M.

- Magnesium**, arc spectrum of (FOOTE, MEGGERS, and MOHLER), A., ii, 4; (HEMSALECH and DE GRAMONT), A., ii, 411.
- effect of electric fields on the spectrum of (DE GRAMONT and HEMSALCH), A., ii, 243.
- electromotive properties of (BECK), A., ii, 545.
- isotopes of (DEMPSTER), A., ii, 417.
- Magnesium alloys**, heat of formation of (BILTZ and HONORST), A., ii, 850.
- with cadmium (GUILLET), A., ii, 570.
- with mercury, thermal analysis of (BECK), A., ii, 545.
- equilibrium of ammonia with (LOOMIS), A., ii, 294.
- Magnesium salts**, distribution of, in blood (KRAMER and TISDALL), A., i, 1087.
- Magnesium dihydrogen orthantimonate** (OBAJDIN), A., ii, 501.
- carbonate, formation of, from the oxide (HEDVALL and HEUBERGER), A., ii, 766.
- Magnesium perchlorate**, preparation and properties of, and its use as a drying agent (WILLARD and SMITH), A., ii, 850.
- haloids, compounds of ammonia with (BILTZ and HÜTTIG), A., ii, 59.
- nitrate, equilibrium of, with sodium nitrate and water, and magnesium sulphate and water (JACKMAN and BROWNE), T., 694.
- silicide, preparation of, and its decomposition (SCHWARZ and KONRAD), A., ii, 846.
- sulphate, equilibrium of ammonium and potassium sulphates, water and (WESTON), T., 1223.
- equilibrium of magnesium nitrate, water and (JACKMAN and BROWNE), T., 694.
- pharmacology of (FERRARA), A., i, 500.
- sulphide, preparation of, pure, and its phosphorescence (TIEDE and RICHTER), A., ii, 215, 247.
- rare earth phosphors of (TIEDE and SCHLEEDÉ), A., ii, 769.
- Magnesium organic compounds**, use of, in syntheses (HEPWORTH), A., i, 118.
- electrolysis of (KONDYREV), A., i, 1128.
- action of cupric chloride with (TURNER), A., i, 23.
- action of, on glutaronitrile (BRUYLANTS), A., i, 531.
- action of, on nitriles (BAERTS), A., i, 817.
- with olefines (KRESTINSKI), A., i, 1128.
- with pyridine (SPAC[†]), A., i, 859.
- reactions of (GARCÍA BANÚS and PASCUAL VILA), A., i, 734.
- alkyl haloids, action of halogen derivatives of ethane and methane on (BINAGHI), A., i, 1002.
- alkyl and aryl haloids, action of iodoform on (ODDO and BINAGHI), A., i, 318.
- allyl bromide, action of dichloroethyl ether on (LESPIEAU), A., i, 619.
- cyanide, and its complex silver salts (FICHTER and SUTER), A., i, 532.
- ethyl bromide, action of, with ethyl chloropropionates (MOUREU and BAERRETT), A., i, 4.
- methyl bromide, action of, with phenylacetoneitrile (RONDOU), A., i, 934.
- phenyl bromide, action of thallous bromide on (D. and A. E. GODDARD), T., 259.

Magnesium detection, estimation, and separation:—

detection of, with 8-hydroxyquinone (MÖRNER), A., ii, 659.

detection of, in presence of manganese and phosphoric acid (PURGOTTI), A., ii, 85.

estimation of, colorimetrically (HAMMETT and ADAMS), A., ii, 587; (BRIGGS; DENIS), A., ii, 659.

estimation of, in nickel (CHALUPNY and BREISCH), A., ii, 227.

estimation of, volumetrically, in potassium salts (VÜRTHEIM), A., ii, 870.

estimation of, and separation from alkalis (BORGSTRÖM), A., ii, 659.

Magnetic field, effect of π , on catalysis by ions (GARNER and JACKMAN), T., 1298.

Magnetism and constitution of atoms (WEISS), A., ii, 23.

relation between atomic structure and (CABRERA; OXLEY), A., ii, 469.

Maize, constituents of the pollen of (ANDERSON and KULP), A., i, 508.

effect of aluminium and iron salts on growth of (ARNDT), A., i, 1103.

absorption of calcium chloride by the roots of (REDFERN), A., i, 614.

proteins in (SHOWALTER and CARR), A., i, 1103.

Maize oil, chemical composition of (BAUGHMAN and JAMIESON), A., i, 414.

Malachite-green-leucosulphonic acid, pararosaniline salt (WIELAND and SCHEUING), A., i, 58.

Maleic acid, preparation of (DOWNS), A., i, 984.

action of muscle enzymes on (DAKIN), A., i, 792.

calcium salt, crystallography and optical properties of (WHERRY and HANN), A., i, 716.

methyl ester, alkaline hydrolysis of (SKRABAL and RAITH), A., ii, 434.

Malic acid, formation of (WEISS and DOWNS), A., i, 622.

impure form of, in Crassulacæ (FRANZEN and OSTERTAG), A., i, 1223.

excretion of (TOMITA), A., i, 301.

d-**Malic acid** (FREUDENBERG and BRAUNA), A., i, 623.

l-**Malic acid**, activation of racemic acid by (MCKENZIE and WALKER), T., 349.

Malic acid, hydroxy-. See Succinic acid, aminohydroxy-.
nitro-, and its salts (LACHMAN), A., i, 107.

Malonamide, *isonitroso*-, derivatives of (RENDALL and WHITELEY), T., 2110.

Malonanilide, amino- (RENDALL and WHITELEY), T., 2119.

isonitroso-, ethers of (RENDALL and WHITELEY), T., 2115.

Malon-*p*-bromonanilide, *mono*-, and *di*-chloro-, and chlorobromo- (WEST), T., 2201.

Malondibenzylamide, *isonitroso*-, ethers of (RENDALL and WHITELEY), T., 2118.

Malondithylamide, chlorobromo- (WEST), T., 2201.

Malondimethylamide, *dichloro*-, and chlorobromo- (WEST), T., 2202.

isonitroso-, ethers of (RENDALL and WHITELEY), T., 2116.

Malonhydrazidic acid, potassium ethyl ester and benzylidene derivative (CURTIUS and SIEBER), A., i, 723.

Malonic acid, diethyl ester, reaction of *s*-.*dichlorodimethyl* ether with (DOX and YODER), A., i, 429.

and its substituted derivatives, decomposition velocity of (BERNOULLI and JAKUBOWICZ), A., ii, 40; (JAKUBOWICZ), A., ii, 361.

sodium derivative, condensation of, with ethyl muconate (FARMER), T., 2015.

alkyl derivatives, conversion of, into α -amino-acids (CURTIUS and SIEBER), A., i, 721.

esters, condensation of, with aromatic amidines (DOX and YODER), A., i, 374.

Malonic acid, halogen derivatives, quantitative reduction of, by hydriodic acid (WEST), T., 2196.

bromo-derivatives, absorption spectra of derivatives of (GRAHAM and MACBETH), T., 1109.

esters, action of hydrazine on (HIRST and MACBETH), T., 904.

mono-, and *di*-chloro-, methyl esters (HIRST and MACBETH), T., 2177.

ethyl esters, preparation of (MACBETH), T., 1120.

Malonmethylanilide, bromo-, and chloro- (WEST), T., 2199.

Malonmethylbromonanilide, bromo- (WEST), T., 2199.

Malonodi-5-acenaphthylamide (FLEISCHER and SCHRANZ), A., i, 1143.

Malononitrile, *dibromo*-, *dichloro*-, and *diiodo*- (OTT and LÖPMANN), A., i, 643.

Malontetrabenzylamide, bromo-, and *mono*- and *di*-chloro- (WEST), T., 2200.

Malontetrabromophenylamide, bromo- (WEST), T., 2200.

- Malontetraphenylamide**, bromo- and chloro- (WEST), T., 2199.
- Malt**, diastatic action of, and its preparations (LECOQ), A., i, 312; (DUCHÁČEK), A., i, 974.
- proteolytic enzymes of (LUNDIN), A., i, 959.
- Maltase** (WILLSTÄTTER and STEIBELT), A., i, 306.
- removal of, from adsorbed substances (WILLSTÄTTER and KUHN), A., i, 283.
- action of arsenic compounds on (RONA, AIRILA, and LASNITZKI), A., i, 959.
- non-identity of α -glucosidase and (WILLSTÄTTER and STEIBELT), A., i, 282.
- genesis of, in plants (PALLADIN and POPOV), A., i, 614.
- in blood (COMPTON), A., i, 392.
- of dog's serum (COMPTON), A., i, 1087.
- Maltose**, action of ozone on pure solutions of (SCHÖNEBAUM), A., i, 921.
- l-Mandelamide**, preparation and racemisation of (McKENZIE and SMITH), T., 1353.
- Mandelic acid**, tetra-acetyl glucose esters (KARRER, NÄGELI, and SMIRNOV), A., i, 253.
- Manganates**. See under Manganese.
- Manganese**, preparation of (FUJIMAYASHI), A., ii, 647.
- spectra of (CATALÁN), A., ii, 726.
- Röntgen-ray spectrum of (DUANE and FRICKE), A., ii, 804.
- Manganese alloys**, electrochemistry of (TAMMANN and VADERS), A., ii, 380.
- Manganese compounds** in flowers (WESTER), A., i, 309.
- in the leaves of plants (BERTRAND and ROSENBLATT), A., i, 411.
- in young and old leaves (JADIN and ASTRUC), A., i, 1098.
- role of, in plants (McHARGUE), A., i, 906.
- in higher plants (BERTRAND and ROSENBLATT), A., i, 95.
- in seeds (WRITER), A., i, 93.
- Manganese dioxide**, coagulation of sols of (GANGULY and DHAR), A., ii, 829.
- adsorption of iron hydroxide by precipitates of (GELOSO), A., ii, 589.
- use of, in the catalytic oxidation of ammonia (PIGGOT), A., ii, 142.
- alkali selenates (TUTTON), A., ii, 505.
- tungstate (SMITH), A., ii, 774.
- Manganous sulphide**, green, preparation of (HAHN), A., ii, 380.
- Manganese**:—
- Manganates**, velocity of reaction between formic acid and (HOLLUTA), A., ii, 700.
- Permanganates**, mechanism of the reduction of (HOLLUTA), A., ii, 700, 771.
- Manganites**, preparation and properties of (SARKAR and DHAR), A., ii, 398.
- Manganese organic compounds**:—
- Manganic-acetic and -benzoic acids**, complex salts of (WEINLAND and FISCHER), A., i, 421.
- Manganese detection, estimation, and separation**:—
- detection of (DITZ), A., ii, 229.
- detection of, by oxidation to permanganate (HESLINGA), A., ii, 589.
- estimation of (HUBER; SARKAR and DHAR; NICOLARDOT, GELOSO, and REGLADE), A., ii, 398; (WINKLER), A., ii, 589; (MINOVICI and KOLLO), A., ii, 787.
- estimation of, colorimetrically, in steel and alloys (HESLINGA), A., ii, 660.
- separation of, from iron (CARUS), A., ii, 88.
- Manganites**. See under Manganese.
- Mangiferin** (*cucurbitogen*), and its derivatives (GORTER), A., i, 933.
- Mannanase** (MAYEDA), A., i, 694.
- Mannitol**, action of boric acid on (DUBREY), A., i, 1110.
- constitution and rotation of compounds of, with boric acid and sodium hydroxide (GILMOUR), T., 1333.
- Mannitolphosphoric acid**, iron salt (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 631.
- l-Mannohepturonic acid**, and its lactone, derivatives of (KILIANI and WINGLER), A., i, 321.
- Mannose**, preparation of (CLARK), A., i, 433.
- Manurial experiments**, with aluminium and iron salts (ARNDT), A., i, 1103.
- with silicic acid (LEMMERMANN and WIRSSMANN), A., i, 1108.
- Maple**, Korean. See *Acer ginnala*.
- Marble**, corrosion of, by plant roots (CHEMIN), A., i, 94.
- Marcasite**, crystalline structure of (HUGGINS), A., ii, 651.
- Martensite**, heat of transformation of, into pearlite (YAMADA), A., ii, 475.
- Matrine**, constitution of (KONDŌ, KISHI, and ARAKI), A., i, 289.
- Matter**, enantiomorphism of (CONDELLI), A., i, 410.

- Mechanics**, statistical, relation between, and thermodynamics (TOLMAN), A., ii, 257.
- Meconic acid**, estimation of, in opium (ANNETT and BOSE), A., ii, 791.
- Medicago falcata**, yellow pigment from flowers of, and its transformation by oxydases (JONESCO), A., i, 1224.
- Melamine**, tricyano-, and its salts (MADELUNG and KERN), A., i, 440.
- Melampyrum arvense**, aucubin and melampyritol from (BRIDEL and BRAECKE), A., i, 799.
aucubin and sucrose in the seeds of (BRIDEL and BRAECKE), A., i, 209.
- Melanin** from pyrrole (RONDONI), A., i, 64.
- Melanin acids**, formation of, from organic compounds (ADLER and WIECHOWSKI), A., i, 1170.
physiological action of (ADLER and WIECHOWSKI), A., i, 498.
- Melanovanadite** (LINDGREN), A., ii, 155; (LINDGREN, HAMILTON and PALACHE), A., ii, 305.
- Meleagris gallinapavo** (turkey), enzymes of the fat of (HEPBURN), A., i, 196.
- Mellilites**, natural and synthetic (BUDINGTON), A., ii, 155.
- Mellitic acid** (PHILIPPI; PHILIPPI and RIX; PHILIPPI and THELEN; PHILIPPI, SEKA, and FROESCHL; PHILIPPI, SEKA, and ROBINSON), A., i, 837.
- Melting point**, relation of boiling point to (LORENZ and HERZ), A., ii, 739.
relation between boiling point, critical temperature, and (PRUD'HOMME), A., ii, 349.
of aliphatic-aromatic ketones (MAJIMA, NAGAOKA, and TAMADA), A., i, 257.
- Melting point determinations** (HEIKE), A., ii, 25.
apparatus for (UYTENBOGAART), A., ii, 471.
- Melting point tubes**, use of, as reaction vessels (FUOHS), A., ii, 759.
- Membranes**, collodion, anomalous osmosis through (LOEB), A., ii, 742.
de Haas's, for electro-osmosis (PRAUSNITZ), A., ii, 114.
- Mentha aquatica**, oil from (KREMERS), A., i, 848.
- Menthane**, cyano- (LÜCK), A., i, 44.
- Menthanealdehyde** (LÜCK), A., i, 44.
- Methanecarboxylic acid**, and its potassium salt (LÜCK), A., i, 44.
- Menthene glycol**, chlorohydrin of (DETUËF), A., i, 327.
- Menthol**, preparation of (SMITH and PENFOLD), A., i, 31.
- l-Menthol**, derivatives of (LÜCK), A., i, 44.
- Menthone**, preparation of (SMITH and PENFOLD), A., i, 31.
oxidation of, by ferric chloride (ASHINA and MITUHOI), A., i, 667.
- Menthylsemicarbazide** (NEIGHBORS, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.
- Mercaptans**, action of chloropierin with (RAY and DAS), T., 323.
- 2-Mercapto-1-p-iodophenyl-6-methyl-benzimidazole**, and its derivatives (JACOBSEN), A., i, 591.
- 2-Mercapto-5-methylthiazole**, and its metallic salts and their use in vulcanisation (BRUNI and ROMANI), A., i, 755.
- 3-Mercapto-1:4-naphthaquinone**, 2-amino-, acetyl derivative, and its S-methyl ether (FRIES and KERKOW), A., i, 578.
- Mercapturic acid**, formation of, during the ingestion of proteins (KAPFHAMMER), A., i, 293.
- Mercurisation**, heat developed in (BARRATT and LEWIS), A., i, 528.
- Mercuriale vivace**. See *Mercurialis perennis*.
- Mercurialis perennis*, content and distribution of sugars in (GILLOT), A., i, 1101.
- Mercury**, atomic weight of (BRÖNSTED and v. HEVESY), A., ii, 645.
absorption and spectrum of the vapour of (MCLENNAN, AINSLIE, and CALE; FRANCK and GHOTRIAN), A., ii, 728.
arc spectrum of (PROCOPIŮ), A., ii, 600.
continuous spectrum of the vapour of (CHILD), A., ii, 676.
fluorescence spectrum of (VAN DER LINGEN and WOOD), A., ii, 245; (VAN DER LINGEN), A., ii, 543.
series spectrum of (SEELIGER), A., ii, 17.
stability of the luminous discharge in (GIBSON and NOYES), A., ii, 812.
vapour, electrodeless discharge in (ROBERTSON), A., ii, 609.
separation of isotopes of (BRÖNSTED, and v. HEVESY), A., ii, 149; (MULLIKEN and HARKINS), A., ii, 295.
critical constants of (WEBER), A., ii, 645, 769.
density and surface tension of (HOGNESS), A., ii, 29.
velocity of vaporisation of (VOLMER and ESTERMANN), A., ii, 193.
decomposition of hydrogen molecules by (CARIO and FRANCK), A., ii, 809.

- Mercury**, solution of metals in (SKAUPP), A., ii, 297; (TAMMANN and JANDER), A., ii, 825.
 reduction of ferric salts by (McCAY and ANDERSON), A., ii, 217.
 reduction of vanadic acid by (McCAY and ANDERSON), A., ii, 530.
- Mercury alloys (amalgams)**, use of in volumetric analysis (KIKUCHI), A., ii, 519, 721; (KANÓ), A., ii, 529, 721.
 with alkali metals, relative densities of (BAIN and WITHROW), A., ii, 145.
 with aluminium, electromotive properties of (MÜLLER and HÖLZL), A., ii, 341.
 with cadmium, use of, in analysis (KANÓ), A., ii, 529.
 with calcium, electrolytic preparation of (NEUHAUSEN), A., ii, 643.
 with copper (PAAL and STEYER), A., ii, 446.
 with lead, estimation of lead in (MELLON and REINHARD), A., ii, 787; (MELLON), A., ii, 870.
 with magnesium, thermal analysis of (BECK), A., ii, 545.
 equilibrium of ammonia with (LOOMIS), A., ii, 294.
 with silver, preparation of (MÜLLER and HÖNIG), A., ii, 500.
 with sodium, electrochemistry of (RICHARDS and CONANT), A., ii, 340.
 with thallium (RICHARDS and SMYTH), A., ii, 341.
- Mercury compounds**, lecture experiment to show the antiseptic action of (JOACHIMOGLU), A., ii, 211.
- Mercury salts**, physiological action of (SALANT and KRITMAN), A., i, 794.
- Mercury perchlorate**, electrometric titrations with (KOLTHOFF), A., ii, 655.
 chlorides, action of sulphur dioxide on (STEWART and WARDLAW), T., 1481.
 oxychlorides (TODA), A., ii, 769.
- Mercuric chloride**, compound of pyridinebetaine and (CASSELLA & Co.), A., i, 860.
 poisoning. See Poisoning.
 oxide, catalytic influence of oxides on the decomposition of (KENDALL and FUCHS), A., ii, 147.
 sulphide, complex compound of mercuric acetate and (MIOLATI), A., i, 982.
- Mercury organic compounds** (ROSSI), A., i, 605; (MAMELI), A., i, 695, 1082; (MAMELI and MAMELI-MANNESSIER), A., i, 1080; (MAMELI and COCCONI), A., i, 1083.
- Mercury organic compounds with dihydrobenzofurans** (ADAMS, ROMAN, and SPERRY), A., i, 946.
 with hydroxybenzaldehydes (HENRY and SHARP), T., 1055.
 with nitroanilines (KHARASCH, LOMMEN, and JACOBSON), A., i, 603.
 dialkyls, preparation of (MARVEL and GOULD), A., i, 329.
 3:3'-diethyldithienyl (STEINKOPF and HEROLD), A., i, 850.
 mercaptide nitrites, and their derivatives (RAY), T., 1281.
 2:4:2':4'-tetranitrodibenzyl (KHARASCH), A., i, 190.
*o*o'-dinitrodiphenyl (WIBAUT and JÜRGENS), A., i, 694.
 2:4:6:2':4':6'-hexanitrodiphenyl (KHARASCH), A., i, 190.
 oxycyanide, explosion of (MERCK), A., i, 640.
*o*o'-Mercury-bis-*p*-nitroaniline (KHARASCH, LOMMEN, and JACOBSON), A., i, 603.
- Mercuric cyanide**, equilibrium of ammonia with (BRINKLEY), A., i, 724.
- Mercuri-compounds** (KHARASCH), A., i, 188, 189; (KHARASCH and JACOBSON), A., i, 189.
- Mercuriacetic acids**, phenol derivatives (MAMELI), A., i, 695.
- Mercuridiaminoazobenzene acetate** (VECCHIOTTI), A., i, 478.
- Mercuribenzoic acid**, *o*-chloro-, *n*-butyl ester (WHITMORE and MIDDLETON), A., i, 889.
- o*-Mercuribis-*p*-nitrobenzoic acid**, esters of (WHITMORE and MIDDLETON), A., i, 889.
- Mercuridicarboxylic acids**, esters, preparation and hydrolysis of (SCHOELLER and SCHBAUTH), A., i, 605.
- Mercuridimethylaniline**, *o* nitro-*p*-chloro- (KHARASCH and JACOBSON), A., i, 189.
- 1:1-Mercuridimethylenebis-1:2-dihydrobenzofuran** (ADAMS, ROMAN, and SPERRY), A., i, 947.
- Mercuriethylaniline**, *p*-nitro-*o*-chloro- (KHARASCH and JACOBSON), A., i, 189.
- Mercuri-iodides of alkaloids** (FRANÇOIS and BLANC), A., i, 851.
- Mercurimethylaniline**, nitrochloro-derivatives (KHARASCH and JACOBSON), A., i, 189.
- Mercurimethyl-1:2-dihydrobenzofuran**, 1-bromo-, 1-chloro-, and 1-iodo- (ADAMS, ROMAN, and SPERRY), A., i, 947.

Mercury organic compounds:—

Mercurinitroanilines, chloro- (KHARASCH, LOMMEN, and JACOBSON), A., i, 603.

Mercuri-*p*-nitrobenzoic acid, *o*-chloro- and *o*-hydroxy-, derivatives of (WHITMORE and MIDDLETON), A., i, 888.

Mercurinitrophenols, chloro- (RAIZISS and PROSKOURIAKOFF), A., i, 604.

Mercuri-*o*-nitrophenoxide, hydroxy-, sodium (RAIZISS and PROSKOURIAKOFF), A., i, 604.

Mercuri-2-nitroresorcinol, 4-chloro- (RAIZISS and PROSKOURIAKOFF), A., i, 604.

Mercuri-5-nitrosalicylic acid, 3-chloro- (RAIZISS and PROSKOURIAKOFF), A., i, 604.

Mercuriphenoxide, 2-nitro-4-hydroxy-, sodium (SCHAMBERG, RAIZISS, and KOLMER), A., i, 391.

Mercury estimation and separation:—
estimation of, in ores (HEINZELMANN), A., ii, 162.

estimation of, in pills (FRANÇOIS), A., ii, 87.

separation of, from silver salts (KOLTHOFF), A., ii, 160.

Mercury cathode. See Cathode.

Mercury electrode. See Electrodes.

Mesityl oxide, chlorohydrin. See *iso*-Butyl methyl ketone, α -chloro-*B*-hydroxy-.

Mesitylene, additive compound of tetrachlorophthalic anhydride and (PFEIFER and FLATER), A., i, 341.

Mesochatopterus Taylori, constituents of the tube of (BERKELEY), A., i, 400, 493.

Mesochydry (ODDO), A., ii, 367.

Mesothorium, chemical properties of (YOVANOVITCH), A., ii, 712.

Mesoxalic acid, sodium salt and hydrazones of (CHATTAWAY and HARRIS), T., 2706.

Mesoxamide, oxime of. See Malonamide, *isonitroso*-.

Metabolism, effect of radiothorium on (MIYADERA), A., i, 966.

effect of putrefaction products on (HIJIKATA), A., i, 495.

effect of injection of sulphur on (MEYER-BISCH and BASCH), A., i, 83.

in alcaptonuria (GIBSON and HOWARD), A., i, 401.

basal, in underweight children (BLUNT, NELSON, and OLESON), A., i, 83.

in menstruation (WILTSHIRE), A., i, 395.

calcium (ROSEMAN), A., i, 1210.

Metabolism of calcium and phosphoric acid (BLUHDORN), A., i, 83.

carbohydrate (ELIAS and SAMMARINO), A., i, 86; (HEWITT and DE SOUZA), A., i, 395; (ELIAS and WEISS; NEUWIRTH), A., i, 485.

effect of light on (PINCUSSEN), A., i, 1088.

in diabetes (ALLEN and WISHART), A., i, 893; (WAGNER and PAENAS), A., i, 965.

effect of simultaneous injection of adrenaline and pilocarpine on (VOGEL and BORNSTEIN), A., i, 395.

chloride, during the menstrual period (EISENHARDT and SCHAEFER), A., i, 83.

chlorine, in tuberculosis (BOENHEIM), A., i, 1092.

creatinine, effect of the thyroid on (SCHENK), A., i, 1212.

endogenous, in tissues (MITCHELL, NEVENS, and KENDALL), A., i, 897.

nitrogen, of higher plants (CHIBNALL), A., i, 1225.

nucleic acid, effect of Röntgen rays on (PINCUSSEN and MOMFERRATUS-FLORES), A., i, 395.

nuclein (ROTHER), A., i, 292.

of phosphorus in tetany (ELIAS and SPIEGEL), A., i, 401.

of pigments (FROMHOLDT and NERSESSOV), A., i, 387, 404.

purine (KIKUCHI), A., i, 698.

influence of food ingestion on (ROSE), A., i, 194.

respiratory (BURGER; PEDOTTI; ASHER and KODA; ASHER and DOUBLER), A., i, 286.

in glycaemia (BORNSTEIN and MULLER), A., i, 392; (BORNSTEIN and HOLM), A., i, 890.

of inorganic salts (GROSS and UNDERHILL), A., i, 1210.

of sugars, intermediary (STAUB), A., i, 83.

sulphur (LEWIS and ROOT), A., i, 487; (LEWIS, MCGINTY, and ROOT), A., i, 1088.

Metacholesterol, preparation and properties of (LIFSCHUTZ), A., i, 541.

Metallic alloys. See Alloys.

carbonyls (MOND and WALLIS), T., 29.
action of nitric oxide on (MOND and WALLIS), T., 32.

chlorides, dissociation of, in aqueous solutions (GÜNTHER-SCHULZE), A., ii, 765.

compounds, possible existence of, as vapour (EUCKEN and NEUMANN), A., ii, 567.

Metallic conductivity in relation to electromotive force (TAMMANN), A., ii, 255.

haloids, complex, determination of crystal structure of, by means of Röntgen rays (SCHERZER and STOLL), A., ii, 514.

hydrides (EPHRAIM and MICHEL), A., ii, 58.

gaseous, preparation of (PANETH, JOHANNSEN, and MATTHIES; PANETH, MATTHIES, and SCHMIDT-HEBBEL), A., ii, 383.

hydroxides, constitution of (TIRMANN), A., ii, 53.

peptisation of (CHATTERJI and DHAR), A., ii, 205, 627.

oxides, formation of, from hydroxides (H. and U. v. EULER), A., ii, 831.

Röntgen-ray spectra of (HEDVAL), A., ii, 306.

luminescence of, sublimed in the electric arc (NICHOLS and WILBER), A., ii, 105.

isomerism of (APPLEBEY and REID), T., 2129

hydrated (WEISER), A., ii, 575, 853.

catalytic decomposition of (TAYLOR and HULETT; KENDALL and FUCHS), A., ii, 646.

salts, refractive index of (LIMANN), A., ii, 173.

molten, molecular refraction of (MEYER and HECK), A., ii, 329.

absorption of light by (GEORGE), A., ii, 806.

crystalline particles in solutions of, visible in the Tyndall cone (TRAUBE and KLEIN), A., ii, 201.

electrolytic dissociation of, in solution and in fused and solid states (RABINOWITSCH), A., ii, 186, 187, 188.

mixed, electrolysis of (CREUTZFELDT), A., ii, 347.

and their mixtures, electrical conductivity of, in molten condition (BENRATH and DREKOPF), A., ii, 109.

temperature of the vapour from boiling solutions of (HARKER), A., ii, 26.

vapour pressure of (v. WARTENBERG and SCHULZ), A., ii, 146; (v. WARTENBERG and BOSSE), A., ii, 739.

vapour pressure of saturated solutions of (EDGAR and SWAN), A., ii, 349.

vapour pressure curves of ammonia with solutions of (DAVIS, OLMSTEAD, and LUNDSTRUM), A., ii, 49.

Metallic salts, solubility of, in aqueous ethyl alcohol and water (TREADWELL), A., ii, 31.

fused (HERZ), A., ii, 116, 739.

thermal expansion of solutions of (MEYER), A., ii, 787.

adsorption of, by metal surfaces (v. EULER and ZIMMERLUND), A., ii, 822.

preparation of gels of (CHARITSCHKOV), A., ii, 827.

complex coloured, grouping of atoms in (WELLS), A., ii, 464.

sparingly soluble solid, reaction affinity in systems of (BRÖNSTED), A., ii, 132.

double decomposition of, and its geometrical representation (JAN-ECKE), A., ii, 427.

reactions of, with dimethyldithiolethylene (MORGAN and LEDBURY), T., 2882.

additive compounds of organic bases with (PETERS), A., i, 48.

action of solutions of, on the heart (CATE; ZONDEK), A., i, 296.

selenides, preparation of (MOSER and DOCTOR), A., ii, 46.

sulphides, solubility of (WEIGEL), A., ii, 644.

used in metallurgy, specific heat of (BORNEMANN and HENGSTENBERG), A., ii, 613.

tellurides, preparation of (MOSER and ERTL), A., ii, 48.

vapours, thermal ionisation of (TOLMAN), A., ii, 18.

Metallo-albumins, toxicity of (ARIOLA), A., i, 500.

Metals, constitution of (KRAUS), A., ii, 566.

vacuum spark spectra of (CARTER), A., ii, 599.

electromotive behaviour of (DHAR), A., ii, 20.

allotropy and electromotive behaviour of (SMITS), A., ii, 110.

electrolytic deposition of (KLING and LAESSIEUR), A., ii, 587.

cathodic deposition of, on aluminium and chromium (KYROPOULOS), A., ii, 22.

thermo-electric force, entropy, and specific heat of, at high temperatures (LATIMER), A., ii, 814.

crystal structure of (HULL), A., ii, 624; (HAUGHTON and FORD), A., ii, 825.

impurities between the crystallites of (TAMMANN), A., ii, 502.

mechanical properties of (GUILLET and COURNOT), A., ii, 261.

- Metals**, increase in the size of grain of, formed from powdered material (SAUERWALD), A., ii, 746.
solubility of, in acids (PRINS), A., ii, 488.
diffusion in, in the solid state (SIROVICH and CARTOCETI), A., ii, 571.
diffusion and cementation of (SIROVICH and CARTOCETI), A., ii, 68.
ionic equilibria on the surfaces of (v. EULER, HEDELIUS, and ZIMMERLUND), A., ii, 251.
adsorption of metallic salts on the surface of (v. EULER and ZIMMERLUND), A., ii, 822.
reactions on the surfaces of (RUFF), A., ii, 363.
corrosion of, influence of protective colloids on (FRIEND and VALLANCE), T., 466.
passivity of (EVANS; GILLIS), A., ii, 814.
behaviour of, as catalysts (SANDONINI), A., ii, 557.
catalytic oxidation by means of (KARCZAG), A., ii, 42.
velocity of the action of oxygen, hydrogen sulphide, and the halogens on (TAMMANN and KÖSTER), A., ii, 831.
action of nitric acid on (BANERJI and DHAR), A., ii, 756.
compounds of proteins with (BENEDICENTI and REBELLO-ALVES), A., ii, 693.
action of, on toxins (ERDSTEIN and FÜRTH), A., i, 90.
deflagrated, positive ray analysis of the gases given off by (THOMSON), A., ii, 565.
dissolved in mercury, molecular condition of (SKAUPP), A., ii, 297.
affinity of, for each other (TAMMANN and JANDER), A., ii, 825.
powdered, toxicity of (ARIOLA), A., i, 500.
bearing, estimation of tin in (NÄGEL), A., ii, 721.
precipitation of, by hydrogen sulphide (SMITH), A., ii, 626.
estimation of gases in (SIMONS), A., ii, 719.
Metal wires, hard drawn, structure of (ETTSCH, POLÁNYI, and WEISSENBERG), A., ii, 201.
Meteoric iron from Texas and California (MERRILL), A., ii, 451, 452.
Meteorite, the Annaheim (JOHNSTON and ELLSWORTH), A., ii, 306.
the Pitts (MCCALLIE), A., ii, 306.
the Strathmore, composition of (MC-LINTOCK and ENNOS), A., ii, 861.
Methæmoglobin, absorption spectra of, and its transformation into oxyhæmoglobin (QUAGLIARIELLO), A., i, 883.
Methanalsulphurous acid. See Sulphiformin.
Methane, preparation of (OHMANN), A., i, 2.
action of ozone on (WHEELER and BLAIR), A., i, 1105.
ignition of mixtures of atmospheric air and (MASON and WHEELER), T., 2079.
synthesis of the polyacetic acids of (INGOLD and PERREN), T., 1414; (INGOLD and NICKOLLS), T., 1638.
estimation of, in small quantities (MURMANN), A., ii, 591.
Methane, bromotrinitro- (SCHMIDT, BARTHOLOMÉ, and LUBKE), A., i, 826.
chloro-derivatives, antiseptic action of (JOACHIMOGLU), A., i, 304.
halogen derivatives, action of, on magnesium alkyl compounds (BINAGHI), A., i, 1002.
tetranitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.
action of triphenylhydrazine on (GOLDSCHMIDT and RENN), A., i, 478.
Methanedisulphonic acid, amino-, copper and nickel salts (DELÉPINE and DEMARS), A., i, 923.
Methanetetra-acetic acid, and its salts and derivatives, and α -cyano-, ethyl ester (INGOLD and NICKOLLS), T., 1645.
Methanetriacetic acid, preparation of, and its esters (INGOLD and PERREN), T., 1414.
Methenylbis-2:4-dimethylpyrrole-3-carboxylic acid, ethyl ester (KUSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
hydrochloride (FISCHER and ZERWECK), A., i, 758.
Methoxide, sodium, action of, on benzophenone chloride and benzyldene chloride (MACKENZIE), T., 1695.
4-Methoxyacetophenone, 3-hydroxy-. See *iso*Acetovanillone.
2-Methoxy-5-aldehydobenzoic acid, and its derivatives (WAYNE and COHEN), T., 1022.
2-Methoxyanthraquinone-1-carboxylic acid (BRADSHAW and PERKIN), T., 917.
2-Methoxyanthraquinone-1-glyoxylic acid, methyl ester (BRADSHAW and PERKIN), T., 917.

- 2-o-Methoxybenzeneasoglyoxaline** (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1198.
- 2-o'-Methoxybenzeneazo-1:8-naphthasultam-4-sulphonic acid**, and its potassium salt (KONIG and KEIL), A., i, 823.
- 2'-Methoxybenzil**, 2-hydroxy- (SCHONBERG and KRAEMER), A., i, 664.
- Methoxybenzoic acids**, amino-, and nitro- (FROELICHER and COHEN), T., 1652.
- 4-Methoxybenzonitrile**, 3-nitro- (MATTAAR), A., i, 251.
- 4-Methoxybenzophenone-4'-arsinic acid** (LEWIS and CHEETHAM), A., i, 187.
- o-4'-Methoxybenzoylbenzoic acid**, preparation of, and its salts (ORNDORFF and KELLEY), A., i, 834.
- 4'-Methoxybenzoylbenzoic acid** (GREENE and ROBINSON), T., 2189.
- γ-4-Methoxybenzoyl-β-phenylethylmethylmalonic acid**, and γ-bromo-, methyl esters (KOHLEH), A., i, 552.
- Methoxybenzyl bromides**, isomeric, preparation and properties of (LAPWORTH and SHOESMITH), T., 1391.
- p-Methoxybenzylcinnamic acids**, and their anilides (INGOLD and PERREN), T., 2386.
- 5-p-Methoxybenzylidenesaminoacenaphthene** (FLEISCHER and SCHRANZ), A., i, 1143.
- o-Methoxybenzylidenesanthranilic acid** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- p-Methoxybenzyl methyl ketone**, derivatives of (LE BRAZIDEZ), A., i, 456.
- 5-Methoxy-2-benzyl-2-methyltetrahydrofuran** (HELFERICH and GRIKKE), A., i, 10.
- o- and p-Methoxy-β-bromo-α-allylpropylbenzene** (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- p-Methoxy-β-bromo-α-ethoxypropylbenzene** (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- β-Methoxy-α-bromomercuri-β-p-methoxyphenylpropionic acid**, ethyl ester (SCHRAUTH and GELLER), A., i, 1126.
- β-Methoxy-α-bromomercuri-β-phenylpropionic acid** (SCHRAUTH and GELLER), A., i, 1125.
- β-Methoxy-α-bromomercuriisovaleric acid**, ethyl ester (SCHRAUTH and GELLER), A., i, 1125.
- o- and p-Methoxy-β-bromo-α-methoxypropylbenzene** (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- α-p-Methoxybromophenyl-β-hydroxypropylamine**. See Anetholealkamine, bromo-.
- β-Methoxybutyronitrile** (BRUYLANTS), A., i, 924.
- 6-Methoxy-2-γγγ-trichloro-Δ^α-propenylquinoline-4-carboxylic acid** (HALBERKANN), A., i, 175.
- 4-Methoxycinnamic acid**, 3-amino-, and 3-hydroxy- (MAUTHNER), A., i, 936.
- α-amino-3-hydroxy-, α-benzoyl derivatives (SPATH and LANG), A., i, 568.
- p-Methoxycinnamyl alcohol** (KARER and HORLACHER), A., i, 825.
- p-Methoxycinnamylidenemalonic acid** (HIGGINBOTHAM and LAPWORTH), T., 2828.
- 5-Methoxydihydrindone**, 6-hydroxy-, and its phenylhydrazones (v. KONEK and SZAMÁK), A., i, 459.
- α-Methoxydihydro-o-anethole**, β-nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.
- 7-Methoxy-7:12-dihydro-γ-benzophenarsazine** (LEWIS and HAMILTON), A., i, 188.
- α-Methoxydihydroisosaftrole**, β-nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 738.
- 2'-Methoxy-5:5'-dimethylbenzil**, 2-hydroxy- (SCHONBERG and KRAEMER), A., i, 663.
- 2-Methoxy-4:6-dimethylbenzoic acid** (v. AUWERS and SAURWEIN), A., i, 1032.
- 5-Methoxydioxindole**, and its diacetyl derivative (HALBERKANN), A., i, 172.
- 4'-Methoxydiphenylamine**, 2:4-dinitro- (BLUM), A., i, 28.
- 6-Methoxy-2:3-diphenylquinoline-4-carboxylic acid** (HALBERKANN), A., i, 175.
- 6-Methoxy-2-(2')-furylquinoline-4-carboxylic acid** (HALBERKANN), A., i, 175.
- Methoxy-2-hydroxydiatryl ketones**, and their derivatives (BUCK and HEILBRON), T., 1097.
- 3'-Methoxy-4'-hydroxy-2-styrylbenzopyrylium chloride** (BUCK and HEILBRON), T., 1208.
- 6-Methoxyindole**, and its picrate (KERMACK, PERKIN, and ROBINSON), T., 1879.
- 6-Methoxyindole-3-aldehyde** (KERMACK, PERKIN, and ROBINSON), T., 1882.
- 6-Methoxyindole-2-carboxydimethylacetalylmethylamide** (KERMACK, PERKIN, and ROBINSON), T., 1889.
- 10-Methoxyindophenazine** (HALBERKANN), A., i, 173.
- β-Methoxy-α-iodomercuri-β-p-methoxyphenylpropionic acid**, ethyl ester (SCHRAUTH and GELLER), A., i, 1126.

- 5-Methoxyisatin, and its derivatives (HALBERKANN), A., i, 172.
- 5-Methoxy-2-ketodihydroindole-3-carboxylic acid, 3-hydroxy-, ethyl ester, and its diacetyl derivative (HALBERKANN), A., i, 172.
- 6-Methoxy-2-keto-1:2-dihydroquinoline-4-carboxylic acid (HALBERKANN), A., i, 175.
- 11-Methoxy-3-keto-4-methyl-3:4-dihydro-4-carbeline (KERMAK, PERKIN, and ROBINSON), T., 1889.
- 11-Methoxy-5-keto-4-methyl-4:5-dihydroindole-diazine(1:4) (KERMAK, PERKIN, and ROBINSON), T., 1890.
- 5-Methoxy-1-ketotetrahydronaphthalene-3-sulphonic acid, sodium salt (FUCHS and STIX), A., i, 451.
- Methoxyl groups, apparatus for estimation of (CUMMING), A., ii, 232.
- 5-Methoxymandelic acid, 2-amino- (HALBERKANN), A., i, 172.
- Methoxy-*p*-menthan-8-ol, bromo- (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- β -Methoxy- β -*p*-methoxybromophenylpropionic acid, α -amino- (SCHRAUTH and GELLER), A., 1126.
- β -Methoxy- β -*p*-methoxyphenylpropionic acid, α -amino-, and α -iodo-, and ethyl ester of the latter (SCHRAUTH and GELLER), A., i, 1126.
- 4-Methoxy-3-methylbenzylideneanthranilic acid (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 6-Methoxy-4-methyl-2:3-diketocoumaran (SCHONBERG and KRAEMER), A., i, 664.
- 5-Methoxy 9-methyl-7-ethyl-4:5-dihydric acid, 4-hydroxy- (BILTZ and HEIDRICH), A., i, 383.
- 7-Methoxy-4-methyl-1-hydrindone, hydrazones of (V. AUWERS, HILLIGER, and WULF), A., i, 1193.
- (61)-Methoxy-1-methylindole-2-carboxylic acid (KERMAK, PERKIN, and ROBINSON), T., 1881.
- 4-Methoxy-5-methylphenylarsinic acid, and 3-nitro- (CHRISTIANSEN), A., i, 1203.
- 1-Methoxy-3-methylcyclopropane-1:2-dicarboxylic acid (INGOLD), T., 2694.
- 5-Methoxy-2-methyl-2-propyltetrahydrofuran (HELFERICH and GEHRKE), A., i, 9.
- 6-Methoxy-2-methylquinoline-4-carboxylic acid (HALBERKANN), A., i, 175.
- 5-Methoxy-3-methylsulphone-1-phenyl-1:2:4-triazole (FROMM, KAYSER, BRIGLEB, and FÜHRENBACH), A., i, 378.
- 6-Methoxy-2-methyltetrahydropyran (HELFERICH and MALKOMES), A., i, 431.
- 2-Methoxynaphthalene, bromo- and chloro-derivatives (FRANZEN and STÄUBLE), A., i, 450.
- Methoxynaphthaleneazobenzene-4'-arsinic acids, 1-amino- (JACOBS and HEIDELBERGER), A., i, 74.
- 5-Methoxy-1:4-naphthaquinone, 2:3-dibromo-8-hydroxy- (WHEELER and ANDREWS), A., i, 354.
- Methoxynaphthoic acids, and amino-, and nitro- (FROELICHER and COHEN), T., 1856.
- 9-Methoxy-10-phenanthryl peroxide (GOLDSCHMIDT and SCHMIDT), A., i, 1150.
- 2-Methoxy-*p*-phenetidine, 3:5-dinitro- (REVERDIN and KOETHLISBERGER), A., i, 538.
- 4-Methoxyphenylacetic acid, 3-hydroxy- (SPATH and LANGK), A., i, 569.
- 4-Methoxyphenylarsinic acid, 3-amino-, acetyl derivative, and 3-nitro- (CHRISTIANSEN), A., i, 1203.
- 6-Methoxy-3-[phenyl-(4'-arsinic acid)-azo]-phenoxyacetic acid, 4-amino-, and its salts (JACOBS and HEIDELBERGER), A., i, 75.
- 4-*m*-Methoxyphenylglyoxaline, 2:4-diamino-, and its salts (PUMMERER, BINAFFL, BITTNER, and SCHUEGRAF), A., i, 1198.
- 2-Methoxyphenyl iodomethyl ketone, 4-hydroxy- (SONN and FALKENHEIM), A., i, 1164.
- m*-Methoxyphenylmethylhydrazine, preparation of (KERMAK, PERKIN, and ROBINSON), T., 1880.
- 5-Methoxy-2-phenyl-2-methyltetrahydrofuran (HELFERICH and GEHRKE), A., i, 9.
- 3-Methoxyphenyloctoleic acid, 2-hydroxy- (MAJIMA and TAKAYAMA), A., i, 263.
- Methoxyphenylpropanone. See *p*-Methoxybenzyl methyl ketone.
- β -Methoxy- β -phenylpropionic acid, α -amino-, and α -bromo-, and their derivatives (SCHRAUTH and GELLER), A., i, 1125.
- β -2-hydroxy-, and its derivatives (LANGLEY and ADAMS), A., i, 1164.
- 4-Methoxyphenylpyruvic acid, 3-hydroxy- (SPATH and LANG), A., i, 568.
- 6-Methoxy-2-phenylquinoline-4-carboxylic acid, 5-amino-, and 5-nitro- (HALBERKANN), A., i, 174.
- 8-Methoxy-2-phenyltetrahydroquinoline-4-carboxylic acid (ZUCKMAYER), A., i, 574.

- 6-Methoxyquinoline**, 5:8-diamino-, 8-amino-5-hydroxy-, and 5:8-dihydroxy- (JACOBS and HEIDELBERGER), A., i, 671.
- 7-Methoxyquinoline**, 3-chloro- (KERMACK, PERKIN, and ROBINSON), T., 1883.
- 6-Methoxyquinoline-2:4-dicarboxylic acid**, and its salts (HALBERKANN), A., i, 174.
- 6-Methoxyquinoline 2:3:4-tricarboxylic acid** (HALBERKANN), A., i, 174.
- 6'-Methoxy-9-rubranone**, and its salts (RABE, KINDLER, and WAGNER), A., i, 362.
- 6'-Methoxy-9-rubatoxanone**, and its salts (RABE, KINDLER, and WAGNER), A., i, 361.
- Methoxystearic acid**, bromo-, methyl ester (SCHMIDT, BARTHOLOMÉ, and LUBKE), A., i, 827.
- 3'- and 4'-Methoxy-2-styrylbenzopyrylium chlorides** (BUCK and HEILBRON), T., 1206.
- Methoxytetrahydronaphthalenes**, bromo- (SCHMIDT, BARTHOLOMÉ, and LUBKE), A., i, 827.
- 2-Methoxy-*ar*-tetrahydronaphthalenes**, amino-, bromoamino-, nitro-, and nitroamino-, and their derivatives (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSABH), A., i, 128.
- 2-Methoxy-*ar*-tetrahydronaphthalene-3-sulphonic acid** (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSABH), A., i, 127.
- Methoxytetrapyridineferritetetrachloroferrate**, hydroxy- (WEINLAND and KISSLING), A., i, 364.
- Methoxytetrapyridineferritetetrathiocyanatoferrate**, hydroxy- (WEINLAND and KISSLING), A., i, 365.
- 2- and 4-Methoxytoluoyl chlorides** (MAUTHNER), A., i, 457.
- 2- and 4-Methoxytolyl ethyl ketones** (MAUTHNER), A., i, 457.
- α -(*o*-Methoxy-*m*-tolyl)-naphthylamine**, β -hydroxy- (WAHL and LANTZ), A., i, 823.
- β -Methoxyisovaleric acid**, α -amino-, and α -bromo-, and ethyl ester of the latter (SCHRAUTH and GELLER), A., i, 1125.
- 1-Methoxyxanthone**, 4-chloro- (ECKERT and ENDLER), A., i, 941.
- Methyl alcohol**, preparation of (TRAUN'S FORSCHUNGS-LABORATORIUM), A., i, 522.
purification of, with sodium hypochlorite (MENZIES), T., 2787.
equilibrium of carbon disulphide with (MCKELVY and SIMPSON), A., ii, 271.
- Methyl alcohol**, equilibrium in the system, water and (PUSHIN and GLAGOLEVA), T., 2813.
reaction of water and, with heated copper (CHRISTIANSEN), A., i, 3.
poisoning. See Poisoning.
toxicity of (v. EULER), A., i, 90.
fate of, in the animal organism (POHL), A., i, 498.
detection of (HAMALAINEN), A., ii, 723.
estimation of, and its occurrence in the body (JANSCH), A., ii, 232.
- Methyl groups**, reactivity of, in heterocyclic bases (MILLS and SMITH), T., 2724.
- Methyl iodide**, interaction of potassium plumbite and (DRUCE), A., i, 516.
selenate (MEYER and WAGNER), A., i, 620.
- Methylabietin** (RUZICKA and MEYER), A., i, 830.
- α -Methyl- β -acetylsuccinic acid**, phenylhydrazone (FISCHER and HERRMANN), A., i, 1054.
- 10-Methylacridinium chloride**, 3:6-diamino- (*tryptofurin*) (THIEME), A., i, 59.
and its condensation with formaldehyde (CASSELLA & Co.), A., i, 276.
arsenical compounds of (BENDA), A., i, 888.
- Methylal**, dibromo- (FEIST), A., i, 912.
- α -Methylallyl alcohol**. See Δ^{α} -Buten- γ -ol.
- Methylamide**, dicyano- (MADELUNG and KERN), A., i, 439.
- Methylamine chlorite** (LEVI), A., i, 527.
- 4-Methylaminobenzenitrile**, 3-nitro- (MATTAAR), A., i, 251.
- Methylaminobisdimethylacetal** (KERMACK, PERKIN, and ROBINSON), T., 1886.
- 4-Methylamino-5-carboxybenzeneseleno-benzene-4'-arsinic acid**, and its sodium salts (JACOBS and HEIDELBERGER), A., i, 75.
- 1-Methylamino-4- α -dicarbomethoxyhydrazinomethane** (DIELS and KLEINFELLER), A., i, 1195.
- 6-Methylamino-5-ethylaminouracil**, and its derivatives (BILTZ and HEIDRICH), A., i, 383.
- 4- β -Methylaminoethylglyoxaline**, physiological action of (DALE and DUDLEY), A., i, 403.
- 6-Methylamino-5-formylethylaminouracil** (BILTZ and HEIDRICH), A., i, 383.
- 6-Methylamino-5-formylmethylamino-1:3-dimethyluracil** (BILTZ and HEIDRICH), A., i, 384.

- 6-Methylamino-5-formylmethylamino-3-methyluracil (BILTZ and BÜLOW), A., i, 383.
- 6-Methylamino-5-formylmethylamino-uracil (BILTZ and BÜLOW), A., i, 382.
- p*-Methylaminophenol (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 31.
- ω -Methylaminopropiophenone, and its hydrochloride and derivatives (MANICH and HEILNER), A., i, 351, 371.
- Methylammonium chloride, action of dicyanodiamide with (WERNER and BELL), T., 1790.
- ruthenipentabromide (GUTHIER and KRAUSS), A., i, 16.
- Methyl- Δ^8 -amylenylcarbinol (HELPERICH and MALKOMES), A., i, 431.
- Methyl Δ^8 -amylenyl ketone, and its semicarbazone (HELPERICH and MALKOMES), A., i, 431.
- α -Methylamylpiperidinium iodide (LEONARD), A., i, 363.
- Methylanhydriopierocellin (FORSTER and SAVILLE), T., 821.
- Methylaniline hydrochloride, conversion of, into toluidine hydrochloride (BECKMANN, CORRENS, and LIESCHE), A., i, 535.
- hydroferrocyanide (CUMMING), T., 1296.
- 4-Methylanilinobenzonitrile, 3-nitro- (MATTAAR), A., i, 252.
- o*-4'-Methylanilino-*p*-methylbenzophenone, *o*-2'-nitro- (MAYER and FREUND), A., i, 867.
- Methyl-*m*-anisidine, preparation of (KERMACK, PERKIN, and ROBINSON), T., 1880.
- 2-Methylantraquinone, 5:6:7:8-tetrachloro-, and 5:6:7:8-tetrachloro-1-nitro- (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 179.
- 3-Methylantraquinone, 1:5- and 1:8-dihydroxy- (EDER and WIDMER), A., i, 260.
- 8-Methylantraquinone, 1:4-dihydroxy-. See Shikazarin.
- o*-2'-Methylantraquinonyl-1'-aminobenzoylebenzene (MAYER and FREUND), A., i, 866.
- 3-Methyl- $\alpha\beta$ -anthraquinonylisooxazole (FARBWERKE, VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 179.
- p*-Methylatropic acid, $\beta\beta$ -dichloro- (v. AUWERS and JÜLICHER), A., i, 842.
- 4-Methylazobenzene, 4'-iodo- (JACOBSEN), A., i, 590.
- Methylbenzanthrones (MAYER and SIEGLITZ), A., i, 740, 743.
- 4-Methylbenzophenone-4'-arsenious oxide and -4'-arsinic acid (LEWIS and CHEETHAM), A., i, 187.
- Methylbenzoxazole, 2-hydroxy- (SKRAUP and MOSER), A., i, 575.
- 1-Methylbenzthiazole ethiodide (MILLS), T., 460.
- 5-Methylbenzthiazole, and its ethiodide (MILLS and BRAUNHOLTZ), T., 1492.
- Methylbenzthiazoles, 1-thio- (ROMANI), A., i, 467.
- 2-Methylbenzthiazolenylnbenzothiazolylmethane (MILLS), T., 465.
- 6-(5-Methylbenzthiazolyl)quinoline (LOGERT and MEYER), A., i, 869.
- Methylbenzylglyoxime, nickel compound (PONZIO), A., i, 18.
- 1-Methyl- α -bromoethylbenzenes (v. AUWERS and KOLLIGS), A., ii, 176.
- 1-Methyl- α -bromopropylbenzenes (v. AUWERS and KOLLIGS), A., ii, 176.
- Methylcyclobutane (DEMANOV and DOJARENKO), A., i, 996.
- β -Methylbutane, $\alpha\delta$ -dibromo- $\beta\gamma$ -dihydroxy- (STAUDINGER, MUNTWYLER, and KUPFER), A., i, 979.
- β -Methylbutane- $\alpha\gamma$ -dicarboxylic acid, $\alpha\delta$ -dicyano-, ethyl ester (HOPE and SHELTON), T., 2229.
- γ -Methylbutane- $\beta\delta$ -dicarboxylic acid, $\beta\gamma$ -dicyano-, ethyl ester (HOPE and SHELTON), T., 2232.
- β -Methylbutan- β -ol, α -bromo- (FOURNEAU and PUYAL), A., i, 639.
- 3-Methylcyclobutan-1-one-3-acetic acid, ethyl ester, and its *p*-nitrophenylhydrazones (INGOLD), T., 1151.
- γ -Methyl- $\Delta\beta$ -butene, α -bromo- (STAUDINGER, KREIS, and SCHILT), A., i, 978.
- p*- γ -Methyl- $\Delta\beta$ -butenylguaiacol (STAUDINGER, KREIS, and SCHILT), A., i, 978.
- γ -Methyl- $\Delta\beta$ -butenylmalonic acid, and its ethyl ester (STAUDINGER, KREIS, and SCHILT), A., i, 978.
- p*- γ -Methyl- $\Delta\beta$ -butenylveratrole (STAUDINGER, KREIS, and SCHILT), A., i, 978.
- γ -Methyl- $\Delta\alpha$ -butinene (isopropenylacetylene), preparation and derivatives of (SCHEIBLER and FISCHER), A., i, 1109.
- γ -Methyl- $\Delta\alpha$ -butinen- γ -ol, preparation and derivatives of (SCHEIBLER and FISCHER), A., i, 1108.
- β -Methyl- β -tert.-butylacetaldehyde, and its semicarbazone (LOCQUIN and WOUSSENG), A., i, 711.
- β -Methyl- β -tert.-butylacrylic acid (LOCQUIN and WOUSSENG), A., i, 711.
- γ -Methyl- γ -tert.-butylallyl alcohol, and its allophanate (LOCQUIN and WOUSSENG), A., i, 711.

- β -Methyl- $\Delta\alpha$ -butylene**, formation of, from isoamyl alcohol (PUYAL, A., i, 617).
- β -Methyl- $\Delta\beta$ -butylene sulphide** (CALINGAERT, A., i, 421).
- Methyltert.-butylethynylcarbinol**, and its allophanate (LOCQUIN and WOUSENG), A., i, 618.
- Methyltert.-butylethylcarbinol**, and its allophanate (LOCQUIN and WOUSENG), A., i, 618.
- Methyl- ψ -butylvinylcarbinol**, and its allophanate (LOCQUIN and WOUSENG), A., i, 710.
- α -Methylbutyric acid**, β -bromo-, preparation of (JOHANSSON and HAGMAN), A., i, 425.
- α -Methyl- β -butyrolactone**, preparation of (JOHANSSON and HAGMAN), A., i, 425.
- 1-Methyl-4-carboline**, 3-chloro- (KERMACK, PERKIN, and ROBINSON), T., 1885.
- Methylcarbonatoacetoneitrile** (SONN and WALKENHEIM), A., i, 1164.
- Methylcarbonato- α -amino- n -butyric acid** (CURTIUS and SIEBER), A., i, 723.
- 2-Methylcarbonato-1:3-dimethylcyclobuten-4-one-3-carboxylic acid**, ethyl ester (DIECKMANN and WITTMANN), A., i, 1156.
- Methylcarbonatophenylalanine** (CURTIUS and SIEBER), A., i, 722.
- ω -Methylcarbonatoresacetophenone** (SONN and FALKENHEIM), A., i, 1164.
- Methylchavicol**, ultra-violet absorption spectrum of (DURREANS), A., ii, 6.
- α -1-Methyl-1-dichloromethyl- $\Delta^{2,5}$ -cyclohexadiene- Δ^4 -propionic acid**, and 4-hydroxy-, and their ethyl esters (V. AUWERS and ZIEGLER), A., i, 139.
- 1-Methyl-1-trichloromethyl- $\Delta^{2,5}$ -cyclohexadien-4-ol-4-acetic acid**, and its ethylester (V. AUWERS and JÜLICHER), A., i, 842.
- 1-Methyl-1-dichloromethyl- $\Delta^{2,5}$ -cyclohexadien-4-one**, 5-mono- and 3:5-dibromo- (V. AUWERS and ZIEGLER), A., i, 145.
- 1-Methyl-1-trichloromethyl- $\Delta^{2,5}$ -cyclohexadien-4-one**, and 3-mono- and 3:5-dichloro-, and their derivatives (V. AUWERS and JÜLICHER), A., i, 841.
- 1-Methyl-1-dichloromethylcyclohexan-4-one**, 2:3:5:6-tetrabromo- (V. AUWERS and ZIEGLER), A., i, 145.
- 1-Methyl-1-trichloromethylcyclohexan-4-one**, 2:3:5:6-tetrachloro- (V. AUWERS and JÜLICHER), A., i, 843.
- 1-Methyl-1-dichloromethyl- $\Delta^{2,5}$ -cyclohexen-4-one**, 5:6-dibromo- (V. AUWERS and ZIEGLER), A., i, 145.
- 1-Methyl-1-trichloromethyl-4-methylene- $\Delta^{2,5}$ -cyclohexadiene**, and 3:5-dibromo- (V. AUWERS and JÜLICHER), A., i, 842.
- Methylisochondodendrine**, and its derivatives (FALTIS and NEUMANN), A., i, 570.
- α -Methylcinnamaldehyde**, *m*- and *p*-nitro- (HELLER, LAUTH, and BUCHWALDT), A., i, 348.
- 6-Methylcoumaran-2-one**, synthesis of, and its derivatives (MAMELI), A., i, 669.
- 6-Methylcoumarin** (PONNDORF), A., i, 566.
- Methylisocrotylcarbinol**, and its acetyl derivative (KRESTINSKI), A., i, 1129.
- Methyldeoxybenzoin-2:2'-dicarboxylic acid** (RUGGLI and MEYER), A., i, 344.
- ω -Methyldibenzofulvene**, 2:7-dibromo- (SIEGLITZ and JASSOY), A., i, 821.
- 5-(or 5')-Methyl-2:3'-diethylthioeyanine iodide** (MILLS), T., 463.
- 5-Methyl-2:1'-diethylthioeyanine iodide** (BRAUNHOLTZ and MILLS), T., 2007.
- 1-Methyl-1:2-dihydrobenzofuran**, and 1-iodo- (ADAMS, ROMAN, and SPERRY), A., i, 948.
- O-Methyliso- α -dihydrodemethylscopoline**, and its salts (HESS and WAHL), A., i, 856.
- Methyldihydronaphthalene-2-imino-oxazolidine**, and its salts (TAKEDA and KURODA), A., i, 275.
- α -Methyldihydropalmatine** (SPÄTH and LANG), A., i, 168.
- N-Methyldihydroquinoline hydrochloride** (HEIDELBERGER and JACOBS), A., i, 673.
- d*- and *l*-N-Methyldihydroquinolineol**, and their derivatives (HEIDELBERGER and JACOBS), A., i, 674.
- Methyl- $\alpha\beta$ -dimethyl- α -propenylcarbinol** (KRESTINSKI), A., i, 1131.
- O-Methyl-N-dimethyltyrosine**, ethyl ester (KARRER, GISLER, HORLACHER, LOCHER, MÄDER, and THOMANN), A., i, 814.
- O-Methyl-N-dimethyltyrosinol** (KARRER, GISLER, HORLACHER, LOCHER, MÄDER, and THOMANN), A., i, 814.
- 3-Methyldiphenyl**, 4-amino-, and its derivatives (PUMMERER, BINAPPI, BITTNER, and SCHUEGRAF), A., i, 1197.
- 4-Methyldiphenyl ether**, 2-amino-, and 2-nitro- (MAYER and KRIEGER), A., i, 747.

- 4'-Methyldiphenylamine**, 5-iodo-2-amino-, and 5-iodo-2-nitro- (JACOBSEN), A., i, 591.
- 5-Methyldiphenylamine**, 2-amino-4'-hydroxy- (JACOBSEN), A., i, 592.
- 4'-iodo-2-amino-, and its derivatives (JACOBSEN), A., i, 590.
- 5-Methyldiphenyleneoxide** (MAYER and KRIEGER), A., i, 747.
- 2-Methyldixanthone**, and its derivatives (v. DEM KNESEBECK and ULLMANN), A., i, 360.
- Methylene iodide**, action of aluminium with (THOMAS), A., i, 330.
- Methylenbis-2:3:5:6-tetrahydroxybenzene** (MUKERJI), T., 549.
- Methylene-blue**, diffusion of in organic solvents (LOEWY), A., ii, 354.
- bleaching of, by yeast (KUMAGAWA), A., i, 306.
- reduction of, by the enzymes of muscle (AHLGREN), A., i, 792.
- estimation of (KIKUCHI), A., ii, 721.
- Methylenecyclobutane**, derivatives of (DEM'JANOV and DOJARENKO), A., i, 1009.
- Methylenecamphor**, hydroxy-, reduction products of (RUPE and SCHMIDT), A., i, 1041.
- Methylenecamphor- α -aminocamphor** (RUPE and SCHMIDT), A., ii, 604.
- Methylenedi-*p*-chlorodiphenyldiurethane** (DATTA and CHATTERJEE), A., i, 816.
- Methylenecitric acid**, preparation of (GASTALDI), A., i, 809.
- Methylenediamine** stannichloride (DRUCE), A., i, 639.
- Methylenedicarbamic acid**, esters (DATTA and CHATTERJEE), A., i, 816.
- Methyleneditetrachloro- α -naphthyldiurethane** (DATTA and CHATTERJEE), A., i, 816.
- 6:6'-Methylene-di-(2-methylquinoline-4-carboxylic acid)** (BORSCHKE and MEYER), A., i, 53.
- 3:4'-Methylenedioxybenzoylbenzoin** (GREENE and ROBINSON), T., 2190.
- 3':4'-Methylenedioxybenzoylbenzoin**, and 6'-nitro- (GREENE and ROBINSON), T., 2190.
- 3:4-Methylenedioxybenzylideneglycine** (SCHEIBLER and BAUMGARTEN), A., i, 656.
- 3':4'-Methylenedioxy-2-benzylidene-1-hydrindone**, 6' amino-, and 6'-nitro- (ARMIT and ROBINSON), T., 830.
- 3:4-Methylenedioxy- β -bromo- α -ethoxypropylbenzene** (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- 3:4-Methylenedioxy- β -bromo- α -methoxypropylbenzene** (SCHMIDT, BARTHOLOMÉ, and LÜBKE), A., i, 827.
- 3':4'-Methylenedioxy-2-hydroxydistyryl ketone** (BUCK and HEILBRON), T., 1099.
- 6:7-Methylenedioxy-2:3-indenoquinolines**, and their derivatives (ARMIT and ROBINSON), T., 832.
- 6:7-Methylenedioxy-2:3(3-ketoindeno) (1:2)-quinoline**, and its methosulphate (ARMIT and ROBINSON), T., 835.
- α -Methylenedioxyphenylmethyl- β -hydroxypropylamine**. See *N*-Methylisosalrolealkamine.
- 4-Methylenedioxyphenyl-5-methyl-oxazolidone** (TAKEDA and KURODA), A., i, 273.
- α -3:4-Methylenedioxyphenyl- δ -methyl- Δ^{α} -penten- δ -ol- γ -one**, and its acetate (SCHEIBLER and FISCHER), A., i, 1110.
- 3':4'-Methylenedioxy-2-styrylbenzopyrylium chloride** (BUCK and HEILBRON), T., 1209.
- 6:6'-Methylenedi-(2-phenylquinoline-4-carboxylic acid)** (BORSCHKE and MEYER), A., i, 53.
- 1 1'-Methylenedipyrrolidine** (PUTOCHIN), A., i, 1176.
- Methylethylallylamine**, and its oxide, and their salts (MEISENHEIMER and LOHSNER), A., i, 813.
- 7-Methyl-9-ethyldeoxyuric acid**, and its salts (BILTZ and BÜLOW), A., i, 382.
- 9-Methyl-7-ethyldeoxyuric acid**, and its salts (BILTZ and HEIDRICH), A., i, 383.
- d*-N-Methyl-*O*-ethylidihydrocupreicinal** (HEIDELBERGER and JACOBS), A., i, 674.
- 9-Methyl-7-ethyl-4:5-dihydrouric acid**, 4:5-dihydroxy- (BILTZ and HEIDRICH), A., i, 383.
- 7-Methyl-7-ethyl- α -dinaphthaxanthen**, oxidation of, and 5:9-dihydroxy- (SEN-GUPTA and TUCKER), T., 563.
- Methylethyl-di-*n*-propylarsonium iodide** (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- β -Methyl- β -ethylglutaric acid**, methyl ester (DICKENS, KON, and THORPE), T., 1503.
- Methylethylisohexylcarbinol**, and its allophanate (LOCQUIN and WOUSSENG), A., i, 617.
- Methyl ethyl ketone**, condensation of, in presence of calcium carbide (BECKER and THORPE), T., 1303.
- Methylethylmonylcarbinol** (LOCQUIN and WOUSSENG), A., i, 617.
- Methylethylcyclopentamethylenearsonium iodide** (STEINKOPF, DONAT, and JAEGER), A., i, 996.

- 1-Methyl-1-ethylcyclopentane-3:4-dione** (DICKENS, KON, and THORPE), T., 1508.
- 1-Methyl-1-ethylcyclopentane-3:4-dione-2:5-dicarboxylic acid**, methyl ester, and its derivatives (DICKENS, KON, and THORPE), T., 1508.
- γ -Methyl- γ -ethylpentan- β -one**, and its semicarbazone (NYBERGH), A., i, 802.
- Methylethylpropylamine**, and its oxide, and their salts (MEISENHEIMER and BERNHARD), A., i, 813.
- 5-Methyl-1-ethyl-2-isopropylcyclohexanol**. See Ethylmenthol.
- 5-Methyl-1-ethylpyrazole**, 3-chloro-, and its ethobromide, and 3-chloro-4-bromo- (ROJAHN), A., i, 1184.
- 5-Methyl-1-ethylpyrazol-3-one**, and bromo- (ROJAHN), A., i, 1184.
- 3-Methyl-5-ethylpyridazinone-5-carboxylic acid**, and its ethyl ester (GAULT and SALOMON), A., i, 873.
- 2-Methyl-4-ethylquinoline**, and its salts (KNOEVENAGEL and BAHR), A., i, 751.
- Methylethylsulphine-*p*-toluenesulphonylimine** (MANN and POPE), T., 1053.
- 2-Methyl-2-ethyltetrahydrofuran**, 5-hydroxy- (HELPERICH and GEHRKE), A., i, 9.
- 7-Methyl-9-ethyl-8-thiouric acids** (BILTZ and BULOW), A., i, 382.
- 9-Methyl-7-ethyl-8-thiouric acids** (BILTZ and HEIDRICH), A., i, 382.
- 9-Methyl-7-ethyluric acid** (BILTZ and HEIDRICH), A., i, 383.
- Methylisoeugenol**, ultra-violet absorption spectrum of (DURRANS), A., ii, 7.
- Methylisoeugenolalkamine**, and its salts (TAKEDA and KURODA), A., i, 274.
- Methylisoeugenol-2-imino-oxazolidine**, and its salts (TAKEDA and KURODA), A., i, 274.
- 9-Methylfluorene**, 2,7-dibromo-9-hydroxy-, and 9-chloro-2:7-dibromo- (SIEGLITZ and JASSOY), A., i, 821.
- Methylformoxyhæmin** (KÜSTER and GERLACH), A., i, 597.
- γ -Methylfructoside** (MENZIES), T., 2238.
- Methylfuran-3-carboxylic acid**, bromo-2-hydroxy-, and its acetate (ASAHINA and KUWADA), A., i, 1047.
- Methylfurfuraldehyde**, and hydroxy-, detection of, as phloroglucides (TADOKORO), A., ii, 236.
- α -Methylglucosidase**, action of arsenic compounds on (RONA, AIRILA, and LASNITZKI), A., i, 959.
- β -Methylglucoside**, and its triacetyl derivative, sulphate compounds of (OHLE), A., i, 987.
- Methylglucoside- ζ -bromohydrin**, triacetate of (WREDE), A., i, 226.
- β -Methylglutaconic acid**, α -cyano-, ethyl ester, and its methyl homologues, and their action with ammonia and with water (HOPE), T., 2216.
- β -Methylglutaconic acids**, ethyl esters, and their derivatives (FRIST and BREUER), A., i, 521.
- β -Methylglutaric acid**, α -mono- and α -di-bromo-, and α -mono- and α -di-hydroxy-, and α -iodo-, and their salts and derivatives (INGOLD), T., 2684.
- $\alpha\beta$ -dicyano-**, ethyl ester, preparation of (HOPE and SHELDON), T., 2226.
- 2-Methylglyoxaline**, bromo-derivatives, and their salts, and 4-bromo-5-nitro- (LIGHT and PYMAN), T., 2626.
- 2-Methylglyoxaline-5-sulphonic acid**, 4-bromo- (LIGHT and PYMAN), T., 2629.
- Methylglyoxime**, compounds of potassium and sodium hydrogen sulphites with (GASTALDI and BRAUNIZER), A., i, 627.
- Methylglyoxime**, amino-, and its salts and derivatives (PONZIO and RUGGERI), A., i, 629.
- chloro-**, nickel compound (PONZIO), A., i, 18.
- Methylguanidine**, preparation of (WERNER and BELL), T., 1790.
- Methylhæmatin** (KÜSTER), A., i, 884.
- γ -Methylheptaldehyde**, γ -hydroxy- (HELPERICH and GEHRKE), A., i, 9.
- 1-Methylhexahydropyridine-3-carboxylic acid**, methyl ester (WOLFFENSTEIN), A., i, 365.
- and its methiodide (MERCK), A., i, 949.
- methohaloids of (WOLFFENSTEIN), A., i, 950.
- γ -Methyl-*n*-hexaldehyde**, γ -hydroxy-, and its methylsemiactal (HELPERICH and GEHRKE), A., i, 9.
- Methylcyclohexane**, physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.
- ϵ -Methylhexane**, α chloro- β -hydroxy- (DETGEUF), A., i, 327.
- α -Methylcyclohexane-1:1-diacetic acid**, and $\alpha\alpha$ -dicyano-, *N*-methyl- ω -imide of (KON and THORPE), T., 1801.
- ω -Methylcyclohexanols**, isomeric, and their derivatives (GODCHOT and BÉDOS), A., i, 334.
- Methylcyclohexanones**, derivatives of (MAILHE), A., i, 332.
- δ -Methyl- $\Delta\gamma$ -hexenoic acid** (STAUDINGER, KREIS, and SCHILT), A., i, 978.

- α -Methyl-*n*-hexoic acid**, α -amino- ϵ -hydroxy-, and its copper salt (ZELINSKY and DENGIN), A., i, 1127.
- β -Methyl- β -isohexylacetaldehyde**, and its semicarbazone (LOCQUIN and WOUSSENG), A., i, 711.
- γ -Methyl- γ -isohexylallyl alcohol** (LOCQUIN and WOUSSENG), A., i, 711.
- Methylcyclohexylamines**, and their derivatives (MAILHE), A., i, 832.
- Methyl-*n*-hexylcarbinol**, resolution of (KENYON), T., 2540.
- 3- and 4-Methylcyclohexylethane- $\alpha\beta$ -dicarboxylic acids**, $\alpha\beta$ -dicyano-1-hydroxy- (BIRCH and THORPE), T., 1836.
- Methylisohexylethynylcarbinol**, and its allophanate (LOCQUIN and WOUSSENG), A., i, 617.
- 1-Methyl-1-*n*-hexylcyclopropane-2-carboxylic acid**, 2:3-dicyano-, and its silver salt and amide (BIRCH and THORPE), T., 1827.
- cis*- and *trans*-1-Methyl-1-*n*-hexylcyclopropane-2:3-dicarboxylic acids, and their derivatives (BIRCH and THORPE), T., 1828.
- Methylisohexylvinylcarbinol** (LOCQUIN and WOUSSENG), A., i, 710.
- N*-Methylhistamine**. See 4- β -Methylaminoethylglyoxaline.
- Methylhydrazobenzene**, 5-bromo-2:4-dinitro- (GIUA), A., i, 691.
- 4-Methylhydrazobenzene**, 4'-iodo- (JACOBSEN), A., i, 590.
- 4-Methyl-1-hydrindamine**, 7-hydroxy-, derivatives of (V. AUWERS, HILLIGER, and WULF), A., i, 1193.
- 4-Methyl-1-hydrindone**, 7-hydroxy-, derivatives of (V. AUWERS, HILLIGER, and WULF), A., i, 1193.
- 1-Methyl-2-hydrindone-1-carboxylic acid**, ethyl ester, and its derivatives (PERKIN and TINGLEY), T., 1569.
- Methylhydrodiphtalyl** (RUGGLI and MEYER), A., i, 344.
- 2-Methyl-5- $\alpha\beta\gamma\delta$ -tetrahydroxybutyl pyrrole-3-carboxylic acid**, ethyl ester (PAULY and LUDWIG), A., i, 954.
- N*-Methyl- β -hydroxyethylphenetidine**, and its benzoic ester (BERGMANN, ULITS, and CAMACHO), A., i, 1182.
- N*-Methyl-*N*- $\beta\gamma$ -dihydroxypropylphenetidine** (BERGMANN, ULITS, and CAMACHO), A., i, 1182.
- 5-Methylimino-2-thio-3- and -4-naphthyl-2:3:4:5-tetrahydro-1:3:4-thiodiazoles** (GUHA), A., i, 877.
- 5-Methylimino-2-thio-4-phenyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole** (GUHA), A., i, 877.
- 5-Methylimino-2-thio-4-tolyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole** (GUHA), A., i, 877.
- Methyliminoisoviouric acid**, and its salts (LIFSCHITZ and HERNER), A., i, 768.
- Methylindazoles**, bromo-, and their derivatives (V. AUWERS and LANGE), A., i, 686.
- 3-Methylindazole-2-sulphonic acid**, and its dihydrate and sodium salt (V. AUWERS and HULTENES), A., i, 684.
- 1-Methyl-2:3-indenindoles** (ARMIT and ROBINSON), T., 838.
- 3-Methylindole-2-carboxydimethyl-acetalymethylamide** (KERMACK, PERKIN, and ROBINSON), T., 1888.
- 1-Methylisatinhydrazone**, and its salts and acetyl derivative (BORSCHKE and MEYER), A., i, 55.
- Methylisatoid**, tetrachloro- (HELLER, BENADE, and HOCHMUTH), A., i, 1060.
- Methylitamic acid**, and its barium salt (INGOLD), T., 2689.
- α -Methylignosulphonic acid**, β -naphthylamine salt (KLASON), A., i, 324.
- α -Methyl-*d*-mannoside**, biochemical synthesis of (HÉRISSEY), A., i, 112.
- Methylmenthol** (OGATA and MIYASHITA), A., i, 844.
- β -Methylmethanetriacetic acid**, preparation of, and its ethyl ester (INGOLD), T., 1148.
- Methylcyclomethylenehydrazineimide-1:3-dicarboxylic acid**, 4-chloro-, ethyl ester and its derivatives (DATTA and CHATTERJEE), A., i, 816.
- 2-Methylnaphthalene**, 3-bromo-1:4-dihydroxy-, 3-chloro-1:4-dihydroxy-, and 1:4-dihydroxy- (FRIES and LOHMANN), A., i, 30.
- Methylnaphthalenes**, ring closure in derivatives of (MAYER and SIEGLITZ), A., i, 999.
- 1-Methylnaphthalene-4-carboxylic acid**, and its derivatives (MAYER and SIEGLITZ), A., i, 740.
- 2-Methylnaphthalene-1-carboxylic acid**, and its derivatives (MAYER and SIEGLITZ), A., i, 742.
- 6-Methylnaphthalene-2-carboxylic acid**, 1-amino-, and 1-mito- (MAYER and ALKEN), A., i, 1000.
- 12-Methyl- $\beta\beta$ -naphthaphenthiazine-6:11-quinone** (FRIES and KERKOW), A., i, 578.
- 2-Methyl-1:4-naphthaquinone**, and 2-bromo-, and 3-chloro- (FRIES and LOHMANN), A., i, 30.

- 2-Methyl- α -naphthiminazole**, 1-chloro- (FISCHER and KRACKER), A., i, 957.
- Methyl-1- α -naphthiminazolonecarbamide** (DIELS and SØRENSEN), A., i, 1195.
- 7-Methyl- α -naphthindole** (MAYER and ALKEN), A., i, 1000.
- 7-Methyl- α -naphthindole-2-carboxylic acid**, and its barium salt (MAYER and ALKEN), A., i, 1000.
- 7-Methyl- α -naphthisatin**, and its phenylhydrazone (MAYER and ALKEN), A., i, 1000.
- 1-Methyl- β -naphthol**, 4-chloro-, and 4-chloro-6-bromo- (FRIES), A., i, 42.
- 2-Methyl- α -naphthol**, 4-chloro-, and its acetate (FRIES and LOHMANN), A., i, 29.
- 7-Methyl- α -naphthoxindole** (MAYER and ALKEN), A., i, 1000.
- 6-Methylnaphthyl-2-acetaldehyde**, 1-nitro-, and its phenylhydrazone (MAYER and ALKEN), A., i, 1000.
- 6-Methylnaphthyl-2-acetic acid**, 1-nitro-, and its derivatives (MAYER and ALKEN), A., i, 1000.
- 6-Methylnaphthyl-2-pyruvic acid**, 1-nitro-, and its derivatives (MAYER and ALKEN), A., i, 1000.
- p-Methylnitrosoaminobenzonitrile** (SCHMIDT, SCHUMACHER, BAJEN, and WAGNER), A., i, 734.
- γ -Methylnonane- $\alpha\beta$ -dicarboxylic acid**, $\alpha\beta$ -dicyano- γ -hydroxy-, and γ -hydroxy-, lactone (BIRCH and THORPE), T., 1830.
- γ -Methylnonane- $\alpha\alpha\beta$ -tricarboxylic acid**, γ -hydroxy-, lactone (BIRCH and THORPE), T., 1829.
- Methylnonylcarbinol**, and its allophanate (LOCQUIN and WOUSSENG), A., i, 617.
- 8-Methyloctahydro- ψ -phenanthroline**, and its dihydrochloride (LINDNER), A., i, 688.
- Methylglycoen** (KARRER), A., i, 11.
- Methyl-orange**, improved modification of, as an indicator (HICKMAN and LINSTAD), T., 2502.
- 1-Methyloxindole**, 3-dichloro- (STOLLÉ), A., i, 762.
- 8-Methyl- Δ^{ψ} -pentadiene** (KRESTINSKI), A., i, 1129.
- 8-Methyl- $\Delta^{\alpha\delta}$ -pentadiene- $\alpha\alpha$ -dicarboxylic acid**, ethyl ester (STAUDINGER, MUNTWYLER, and KUPFER), A., i, 979.
- γ -Methylpentane- $\beta\beta$ -dicarboxylic acid**, $\beta\gamma$ -dicyano-, ethyl ester (HOPE and SHELTON), T., 2234.
- γ -Methyl- $\Delta\gamma$ -pentenes**, isomeric (VAN RISSEGHEM), A., i, 909.
- 8-Methyl- Δ^6 -pentenecarboxylic acid**, γ -hydroxy-, lactone derivatives of (STAUDINGER, MUNTWYLER, and KUPFER), A., i, 979.
- 8-Methyl- $\Delta\gamma$ -pentenyl-2:5:8-trihydroxy-1:4-naphthaquinone**. See Shikonin.
- 3-Methyl- ψ -phenanthroline**, preparation and constitution of (LINDNER), A., i, 688.
- 6-Methylphenylacetacetic acid**, 5-chloro-2:4-dinitro-, ethyl ester (DAVIES and HICKOX), T., 2647.
- 5-Methylphenylarsinic acid**, 3-nitro-4-hydroxy- (CHRISTIANSEN), A., i, 1203.
- 1- α -N-Methylpiperidylethan- α -ols** (HESS and CORLEIS), A., i, 170.
- 1- α -N-Methylpiperidylethan- α -one**, and its salts and derivatives (HESS and CORLEIS), A., i, 170.
- β -Methylpropane**, α -amino- β -hydroxy-, and β -chloro- α -amino-, and their salts (DERSIN), A., i, 142.
- 1-Methylcyclopropane-1-carboxylic acid**, methyl ester (KOHN and MENDELEWITSCH), A., i, 518.
- 3-Methylcyclopropane-1:2-dicarboxylic acids**, and 1-bromo-, and their salts (INGOLD), T., 2687.
- 3-Methylcyclopropan-1-ol-1:2-dicarboxylic acid**, and its silver salt (INGOLD), T., 2693.
- Methylisopropenylaniline**, and its derivatives (KNOEVENAGEL and BAHR), A., i, 750.
- Methylisopropenyl ketone** (SCHEIBLER and FISCHER), A., i, 1110.
- α -Methylpropionic acid**, α -amino- β -hydroxy-, and its copper salt (ZRLINSKY and DENGIN), A., i, 1129.
- Methylcyclopropylcarbinyl iodide** (DEM-JANOV and DOJARENKO), A., i, 1014.
- 3-Methyl-6-isopropylcoumaran-1:2-dione** (MAMELI), A., i, 671.
- 3-Methyl-6-isopropylcoumaran-8-one**, and its derivatives (MAMELI), A., i, 669.
- Methylisopropylcoumarophenazine** (MAMELI), A., i, 671.
- 7-Methyl-7-n-propyl- α -dinaphthaxanthene**, oxidation of, and 5:9-dihydroxy- (SEN-GUPTA and TUCKER), T., 563.
- 5-Methyl-2-isopropyl-1-ethylidenecyclohexane** (OGATA and MIYASHITA), A., i, 844.
- Methyl-*n*- and -*iso*-propylglyoximes**, and their salts and dibenzoyl derivatives (PONZIO), A., i, 18.
- Py-N-Methyl-*ind*-N-propylharmine chloride** (KERMAK, PERKIN, and ROBINSON), T., 1893.

- cis*-1-Methyl-4-isopropylcyclohexane (SKITA and SCHENCK), A., i, 241.
- 5-Methyl-2-isopropylidenepyrrolenine-4-carboxylic acid, 8-hydroxy-, ethyl ester (KUSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIO), A., i, 858.
- 5-Methyl-2-isopropyl-1-methylenecyclohexane (OGATA and MIYASHITA), A., i, 844.
- Methylisopropyl-naphthalenes and their salts (RUZICKA and MINGAZZINI), A., i, 1001.
- 6-Methyl-4-isopropyl-1-naphthoic acid, (RUZICKA, MEYER, and MINGAZZINI), A., i, 561.
- 5-Methyl-6-propyloctan- ϵ -ol, and its phenylurethane (LEROIDE), A., i, 216.
- α -Methylpropylphosphinic acid, α -hydroxy-, lead salt (CONANT, MACDONALD, and KINNEY), A., i, 186.
- 5-Methyl-1-propyl-2-isopropylcyclohexanol. See Propylmenthol.
- 1-Methyl-4-isopropyl-3-propylidenecyclohexane (OGATA and MIYASHITA), A., i, 844.
- 2-Methyl-2-propyltetrahydrofuran, 5-hydroxy- (HELFERICH and GEHRKE), A., i, 9.
- 1-Methyl-4-isopropyl-5:6:7:8-tetrahydronaphthalene (RUZICKA and MINGAZZINI), A., i, 1001.
- α -Methyl- α -propylvaleric acid, and its ethyl ester and amide (LEROIDE), A., i, 216.
- 3-Methylpyrazole, 5-chloro-, sodium salt (ROJAHN), A., i, 1183.
- Methylpyrazoleanthrones (MAYER and HEIL), A., i, 878.
- 3-Methyl-6-pyridazinone (GAULT and SALOMON), A., i, 873.
- 3-Methyl-6-pyridazinone-5-carboxylic acid, ethyl ester and hydrazide (GAULT and SALOMON), A., i, 873.
- 1-Methylpyridine-3-carboxylic acid, methyl ester, chloride (MERCK), A., i, 950.
- methosulphate (WOLFFENSTEIN), A., i, 861.
- 1-Methyl-2-pyridone-2-benzoylimide, and its picrate (TSCHITSCHIBABIN and BYLINKIN), A., i, 578.
- 2-Methylpyrimidine, 6-chloro-5-bromo-, and 6-hydroxy- (CHERBULIEZ and STAVRITCH), A., i, 581.
- 2-Methylpyrimidine-4-carboxylic acid, 5-bromo-6-hydroxy-, and 6-hydroxy-, and their copper salts (CHERBULIEZ and STAVRITCH), A., i, 581.
- 5-Methylpyrrole-4-carboxylic acid, 5-amino-2-hydroxy-, ethyl ester (FISCHER and HERMANN), A., i, 1054.
- 5-Methylpyrrole-4-carboxylic acid, 2- and 3-hydroxy-, ethyl esters, derivatives of (FISCHER and HERMANN), A., i, 1054.
- Methylpyrrolidine, 1-hydroxy- (PUTOCHIN), A., i, 1176.
- 3-Methylquinoline, and its acetyl derivative (GRANDMOUGIN), A., i, 585.
- N-Methylquinicine dihydrochloride (HEIDELBERGER and JACOBS), A., i, 673.
- 2-Methylquinoline, 6-amino-, stannic- and stannous-chlorides (DRECE), A., i, 1206.
- bromo-, and their salts (MOUDGILL), A., i, 173.
- Methylisquinolines, and their methiodides (MILLS and SMITH), T., 2732.
- 2-Methylquinoline-3-carboxylic acid, 4-hydroxy-, oxide and its ethyl ester and derivatives (MCCLUSKEY), A., i, 864.
- 2-Methylquinoline-4-carboxylic acid, 6-hydroxy- (HALBERKANN), A., i, 175.
- 6'-Methylisquinoline-red (HARRIS and POPE), T., 1032.
- 1-Methyl-2-quinolone-4-carboxylic acid, ethyl ester (THIELEPAPE), A., i, 271.
- Methylquinoxalinocyclopentane-2:3-dione-1-carboxylic acid, ethyl ester (DIRCKMANN), A., i, 1022.
- Methylretene, formation of, from abietic acid (RUZICKA and MEYER), A., i, 829.
- Methylretenequinone, and its quinoxaline derivative (RUZICKA and MEYER), A., i, 830.
- N-Methylisosaftrolealkamine, and its salts and derivatives (TAKEDA and KURODA), A., i, 274.
- Methylisosaftrole-2-imino-oxazolidine (TAKEDA and KURODA), A., i, 274.
- α -Methylserine. See α -Methylpropionic acid, α -amino- β -hydroxy-.
- Methylstannonic acid, constitution and derivatives of (LAMBOURNE), T., 2533.
- β -Methylsticine acid (MURAYAMA and MAYEDA), A., i, 265.
- α -Methylstyryl methyl ketone, amino- and nitro-derivatives (HELLER, LAUTH, and BUCHWALDT), A., i, 349.
- 3-Methylsulphone-1-nitrophenyl-1:3:4-triazole, 5-hydroxy- (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 378.
- 3-Methylsulphone-1-phenyl-1:3:4-triazole, 5-hydroxy- (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 378.

- p*-Methylsulphonetriphenylacetic acid (BISTRZYCKI and KUBA), A., i, 33.
- p*-Methylsulphonetriphenylcarbinol (BISTRZYCKI and KUBA), A., i, 34.
- p*-Methylsulphoxidetriphenylacetic acid (BISTRZYCKI and KUBA), A., i, 33.
- 3-Methyltetradecahydro- ψ -phenanthroline (LINDNER), A., i, 688.
- 2-Methyltetrahydronaphthalene (TETRALIN G. M. B. H.), A., i, 1040.
- 3-Methyltetrahydro-2:1-naphthanthraquinone (TETRALIN G. M. B. H.), A., i, 1040.
- 2-Methyl- $\beta\beta$ -*or*-tetrahydronaphthiminazole (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSABM, and OESTERLIN), A., i, 125.
- 3 Methyl-2-tetrahydronaphthoyl-*o*-benzoic acid (TETRALIN G. M. B. H.), A., i, 1040.
- N*-Methyltetrahydronorharmine (KERMACK, PERKIN, and ROBINSON), T, 1890.
- α -Methyltetrahydropalmatines (SPATH and LANG), A., i, 169.
- 3-Methyltetrahydro- ψ -phenanthroline, and its hydrochlorides (LINDNER), A., i, 688.
- 2-Methyltetrahydropyran, 6-hydroxy. See *n*-Hexaldehyde, 6-hydroxy.
- 1-Methyltetrahydroquinoline, 6-amino- (LINDNER), A., i, 688.
- N*-Methyltetrahydroquinoline-2-amino-phenazine methochloride, zinc salt (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.
- 5-Methylthiazole sulphide (BRUNI and ROMANI), A., i, 755.
- S*-Methylthiodi- α - and β -naphthylamine-sulphonium salts (KEHRMANN and DARDEL), A., i, 1064.
- Methylthiol groups, estimation of, in methylated. thiolbenzenes (POLLAK and SPITZER), A., ii, 789.
- 8-Methylthiol-9-methyl- Δ^7 -isoxanthine (BILTZ and BULOW), A., i, 384.
- 2-Methylthiol-3-phenyl-5-anilino-methylene-4-glyoxalone (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- 9-*p*-Methylthiophenylfluorene, and its 9-carboxylic acid (BISTRZYCKI and KUBA), A., i, 34.
- 5-Methylthiol-1-phenyl-1:2:4-triazole, 3-amino-, and its benzoyl derivatives (FROMM, KAYSER, BRIEGLEB, and FÖHRNBACH), A., i, 379.
- 5-Methylthiolisalicylic acid (SIEWART), T, 2560.
- p*-Methylthioltriphenylacetic acid, and its derivatives (BISTRZYCKI and KUBA), A., i, 33.
- p*-Methylthioltriphenylcarbinol (BISTRZYCKI and KUBA), A., i, 33.
- p*-Methylthioltriphenylchloromethane (BISTRZYCKI and KUBA), A., i, 33.
- p*-Methylthioltriphenylmethane (BISTRZYCKI and KUBA), A., i, 33.
- S*-Methylthiophenyl- α - and - β -naphthylamine-sulphonium salts (KEHRMANN and DARDEL), A., i, 1063.
- O*-Methyl-*l*-tyrosine, and its ethyl ester (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.
- O*-Methyltyrosine-choline, and its salts (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.
- α -Methyl-*n*-valeric acid, α -amino- β - and - δ -hydroxy-, and their copper salts (ZELINSKY and DENGIN), A., i, 1126.
- Methyl-violet, preparation of (CREIGHTON), A., i, 473.
- 9-Methyl- Δ^7 -*S*-isoxanthine, and 8-iodo-, and their salts (BILTZ and BULOW), A., i, 384.
- 3-Methylxanthinepropionic acid (MERCK, WOLFES, and KORNICK), A., i, 1071.
- 2-Methylxanthone, 7-chloro- (v. DEM KNESEBECK and ULLMANN), A., i, 360.
- 2-Methylxanthone-8-carboxylic acid, 5- and 7-chloro- (v. DEM KNESEBECK and ULLMANN), A., i, 359.
- Methylxanthorocellin (FORSTER and SAVILLE), T, 822.
- Methysticin, catalytic hydrogenation of (GOEBEL), A., i, 657.
- Meyersite (ELSCHNER), A., ii, 714.
- Mica, black, of Ytterby, haloes in (JOLY), A., ii, 651.
- Michael's addition and separation rule (SMITH), A., ii, 751.
- Microbes (*micro-organisms*), formation of acetaldehyde in fermentation by (NEUBERG and COHEN), A., i, 304.
- action of, on amino-acids (HANKE and KOESSLER), A., i, 406.
- decomposition of kaolin by (VERNADSKY), A., i, 1096.
- Microcolorimeter (BAUDOUIN and BERNARD), A., ii, 862.
- Micro-incineration (SCHOELLER), A., ii, 654.
- Micro-organisms. See Microbes.
- Microspira desulfuricans* in soils (KÜHR), A., i, 1228.
- Milfoil. See *Achillea millefolium*.
- Milk, amino-acids in (VIALE), A., i, 898.
- coagulation of, by rennet (BAUR and HERZFELD), A., i, 284.

- Milk**, surface tension phenomena in the separation of cream from (RAHN), A., ii, 478.
 effect of administration of inorganic and organic substances on the composition of (DENIS, SISSON, and ALDRICH), A., i, 494.
 chlorides in (SISSON and DENIS), A., i, 87.
 and its products, citric acid in (SUPPLEE and BELLIS), A., i, 197.
 origin of the fat of (SHEEHY), A., i, 400.
 oxidation of xanthine by (MORGAN, STEWART, and HOPKINS; ARMSTRONG), A., i, 1078.
 toxicity of, and its derivatives (PENTIMALLI), A., i, 302.
 cow's, amino-acids in (HIJIKATA), A., i, 494.
 decomposition of citric acid in, by bacteria (KICKINGER), A., i, 1219.
 diastatic action of (WELZMÜLLER), A., i, 400.
 estimation of urea in (MORIMOTO), A., i, 703.
 goat's, non-protein nitrogen in (TAYLOR), A., i, 1213.
 sour, analysis of (KLING and LASSEUR), A., ii, 402.
 micro-estimation of lactose in (FONTÈS and THIVOLLE), A., ii, 323.
 estimation of peroxylase in (RICE and HANZAWA), A., ii, 407.
- Mineral carbonates**, estimation of carbon dioxide in (SAYCE and CRAWFORD), A., ii, 317.
 phosphates, solubility of (TOCHER), A., ii, 525.
- Minerals** from Bosnia (KOECHLIN), A., ii, 860.
 from Franklin, New York (LEWIS and BAUER), A., ii, 714.
 Japanese, analyses of (SHIBATA and KIMURA), A., ii, 220, 516; (SHIBATA and UEMURA), A., ii, 305.
 from Morocco (BARTHOUX), A., ii, 651.
 action of bacteria on (HELBRONNER and RUDOLFS), A., i, 706.
 heavy liquids for the mechanical separation of (CLERICI), A., ii, 578.
 spectrographic analysis of (HADDING), A., ii, 780.
 estimation of phosphorus in, colorimetrically (MISSON), A., ii, 718.
- Mixtures**, binary, theory of (DOLEZALEK and SCHULZE), A., ii, 118.
 ebullioscopy of (BECKMANN and LIESCHE), A., ii, 117.
 vapour pressure of (SCHMIDT), A., ii, 119.
- Mobility** of symmetrical tautomeric triad systems (INGOLD and PIGGOTT), T., 2881.
- Molecular elotropy** in liquids (RAMAN), A., ii, 603.
 conductivity of strong electrolytes (LORENZ), A., ii, 19.
 constitution, methods for determination of (WAGNER), A., ii, 116.
 refraction. See Refraction.
 structure, relation of odour to (WELANGE), A., i, 894.
 weights. See Weights, molecular.
- Molecules**, structure of (CREHORE), A., ii, 438.
 in the light of the dispersion theory (WASASTJERNA), A., ii, 491.
 arrangement of electrons in (BURY), A., ii, 43.
 determination of the diameter of (GUYE and RUDY), A., ii, 278.
 symmetry of, and the formation of liquid crystals (VORLÄNDER), A., ii, 554.
 collisions between atoms and (FRANCK), A., ii, 464.
 motion of, in space (BÖESEKEN, VAN LOON, DERX, and HERMANS), A., ii, 210.
 anisotropy of (RAMAN), A., ii, 102.
 gaseous, free energy of (PLANCK), A., ii, 689.
 unsaturated, electron structure of (EASTMAN), A., ii, 367.
- Molybdenum**, K-series spectrum of (DUANE and PATTERSON), A., ii, 463.
 Röntgen-ray spectrum of (WINGÅRDH), A., ii, 249.
 valency of, in the octacyanides (COLLENBERG), A., ii, 508.
- Molybdenum carbonyl** (MOND and WALLIS), T., 29.
 dichloride (LINDNER), A., ii, 509.
 trioxide, crystal structure of, and its hydrate (BURGER), A., ii, 508.
- Molybdic acid**, effect of hydroxy-organic compounds on the hydrogen-ion concentration of (RIMBACH and LEY), A., ii, 342.
- Molybdoselenites** (ROSENHEIM and KRAUSE), A., ii, 48.
- Molybdenum organic compounds**:—
 Molybdic acid, compounds of, with formic acid (JAKÓB), A., i, 712.
- Molybdenum estimation and separation**:—
 estimation of, in tungsten (HALL), A., ii, 660.
 separation of, from tungsten (MERRILL), A., ii, 229.
- Molybdomalic acid**, ammonium salts (DARMOIS), A., i, 220, 522.

Molybdoselenites. See under Molybdenum.

Monazite from Ishikawa (SHIBATA and KIMURA), A., ii, 220.

decomposition and analysis of (WENGER and CHRISTIN), A., ii, 722.

Montan wax. See Wax.

Mordants (BANCROFT), A., ii, 551, 646, 822; (PADDON), A., ii, 822.

Morphine and its derivatives, hæmolytic by (RHODE), A., i, 1088.

allyl ether, and its hydrochloride (V. KERESZTY and WOLF), A., i, 854.

derivatives of (SPEYER and BECKER), A., i, 675.

detection of (COLLEDGE), A., ii, 327.

detection and estimation of, in animal tissues (WACHTEL), A., ii, 171.

estimation of, in opium (RAKSHIT), A., ii, 96.

*apo*Morphine, reduction of (FALTIS and KRAUSZ), A., i, 676.

Moss, Irish, extraction of colloidal solutions from (GUTHIER and HUBER; GUTHIER, WOLF, and KIESS), A., ii, 205.

Mother-of-pearl, precipitation of calcium carbonate in the form of (CLÉMENT and RIVIÈRE), A., ii, 500.

Mould fungi. See Fungi.

Moulds, fermentation of pentoses by (PETERSON, FRED, and SCHMIDT), A., i, 1220.

action of, on racemic acids (CONDELLI), A., i, 410.

Mountain ash, constituents of the berries of (v. LIPPMANN), A., i, 1224.

acids from the berries of (FRANZEN and OSTERTAG), A., i, 616.

Muconic acid, constitution of, and α -bromo, methyl ester (CHANDRASENA and INGOLD), T., 1306.

ethyl ester, addition of ethyl sodiocyanoacetate and ethyl sodiomalonate to (FARMER), T., 2015.

Mucor racemosus, invertase of (KOSTYTSCHEV and ELIASBERG), A., i, 410.

Muscarine, isolation of, from *Amanita muscaria* (KING), T., 1743.

Muscle, chemical composition and structure of, with reference to function (BEHRENDT), A., i, 197.

chemical and physical properties of (QUAGLIARIELLO), A., i, 608.

diffusion of hydrogen and hydroxyl ions in relation to the action of (ADAMS), A., i, 1211.

extraction of, by solutions of calcium and sodium chlorides (WIDMARK and LINDAHL), A., ii, 607.

cause of contraction of (v. FURTH; DUX and LOW), A., i, 398.

Muscle, tension and extensibility of, during contraction (VERZÁR, BÖGEL, and SZÁNYI), A., i, 1212.

permeability of (EMBDEN and ADLER; VOGEL; SIMON), A., i, 196.

effect of adrenaline on the permeability of (LANGE), A., i, 791.

effect of alkaloids and organic electrolytes on the permeability of (VOR-SCHUTZ), A., i, 791.

amino-acids in (ROSEDALE), A., i, 492.

production of ammonia in (LEE and TASHIRO), A., i, 1089.

degradation of carbohydrates in (LAQUER), A., i, 298, 1089.

carnosine in, after cold storage (CLIFFORD), A., i, 791.

of various animals (CLIFFORD), A., i, 398.

creatine in (RIESSER), A., i, 791.

formation of creatine in dystrophy of (GIBSON and MARTIN), A., i, 300.

creatine and creatinine in (HAMMFT), A., i, 1090.

action of enzymes in, on fumaric, glutaconic, and maleic acids (DAKIN), A., i, 792.

reducing action of the enzymes of, on methylene-blue (AHLGREN), A., i, 792.

relative toxicity of haloid ions towards (CAMERON and HOLLENBERG), A., i, 499.

energy changes and formation of lactic acid in (MEYERHOFF), A., i, 86, 897.

function of lactic acid in (FOSTER and MOYLER), A., i, 398.

potassium paralysis in (VOGEL), A., i, 196.

cross-striped, effect of potassium and of uranium salts on movements of (VERZÁR and SZÁNYI), A., i, 1213.

frog's, change of choline content of, on electrical stimulation (GEIGER and LOEWI), A., i, 492.

formation of phosphoric acid during contraction of (EMBDEN and LAWACZEK), A., i, 492.

heart. See Heart.

pig's, organic bases from (SMORODIN-CEV), A., i, 1213.

smooth, action of acid and alkali on (HEYMANN), A., i, 791.

striated, action of poisons on (RIESSER and NEUSCHLOSZ), A., i, 1212.

estimation of carnosine in (HUNTER), A., ii, 328, 885.

Mustard flour, estimation of allylthiocarbimide in (LUCE and DOUCER), A., ii, 537.

Mutarotation (RIIBER), A., ii, 807.

Mya arenaria, respiration of (COLLIP), A., i, 286.

Myrica gale, essential oils from (SCHOOF), A., i, 1043.

Myristicin, β -nitro- (SCHMIDT, SCHUMACHER, BÄJEN, and WAGNER), A., i, 733.

N.

Nalita (SAHA and CHOUDHURY), T., 1044.

1:5-Naphthabisdioxindole (MARTINET and VACHER), A., i, 689.

$\beta\beta$ -Naphthabisdioxindole-3:8-dicarboxylic acid, ethyl ester, and its acetyl derivatives (MARTINET and VACHER), A., i, 689.

Naphthacinchoninylpropionic acid, ethyl ester (CARRIÈRE), A., i, 319.

β -Naphthacoumarin, nitro- (MORGAN and REEVES), T, 7.

$\alpha\beta$ -Naphthadihydroxoxazine (CLEMO and PERKIN), T., 647.

1:5-Naphthadi-isatin, and its derivatives (MARTINET and VACHER), A., i, 689.

Naphthadivinulines, amino-, bromo-, hydroxy-, and nitro- (DUTT), T., 1951.

α -Naphthaldehyde, 2- and 4-hydroxy-, azomethine derivatives of (MORGAN and REEVES), T., 1.

nitro-2-hydroxy- (MORGAN and REEVES), T., 6.

β -Naphthaldehyde, phenylhydrazone of (WEIL and OSTERMEIER), A., i, 139.

α - and β -Naphthaldehydes, bromo- (MAYER and SIEGLITZ), A., i, 742, 744.

Naphthalene, structure of, and its derivatives (VOROSHOV), A., i, 1135.

absorption of ultra-violet rays by (HENRY and STEINER), A., i, 928.

heat of combustion of (VERKADE, COOPS, and HARTMAN), A., ii, 475 ; (SWIENTOSLAWSKI and STARCZEWSKA), A., ii, 616.

vapour pressure of (NELSON and SENSEMAN), A., i, 245.

ethylation of (MILLIGAN and REID), A., i, 380.

nitration of (DAVIS), A., i, 818.

catalytic oxidation of (DOWNS), A., i, 27 ; (CONOVER and GIBBS), A., i, 454.

and its derivatives, electrolytic oxidation of (ONO), A., i, 1008.

hydrogenated derivatives (SCHROETER), A., i, 122, 123, 126 ; (SCHROETER and TETRALIN G. M. B. H.), A., i, 1036.

Naphthalene, 5-(or 8)-bromo-2-amino-, -2-hydroxy-, and -2-nitro-, and their derivatives (v. BRAUN, HAHN, and SEEMANN), A., i, 730.

bromonitro-1:5-dihydroxy- (HEMMELMAYR), A., i, 836.

1:8-dibromo-2:7-dihydroxy-, and 5-iodo-1-nitro-, and their derivatives (SCHOLL, SEER, and WEITZENBÖCK), A., ii, 650.

2:3-dibromo-1:4:5:8-tetrahydroxy-, 2:3:8-tribromo-1:4 5-trihydroxy-, and 1:4:5:8-tetrahydroxy- (WHEELER and ANDREWS), A., i, 354.

chloro-, chloroamino-, chlorohydroxy-, and chloronitro-derivatives (KALLE & Co.), A., i, 819.

dichloronitro- and trichloro-derivatives (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 244.

1:4-dicyano- (SCHOLL and NEUMANN), A., i, 261.

1:5-dihydroxy-, effect of heating, with potassium hydrogen carbonate (HEMMELMAYR), A., i, 836.

1:5- and 2:7-dihydroxy-, tautomerism of (FUCHS and STIX), A., i, 451.

Naphthalenes, dihydroxy-, equilibria of amines with (KREMAN, HEMMELMAYR, and RIEMER), A., i, 1010.

Naphthalene-2-aldehyde, 4-bromo-1-hydroxy-, 4-chloro-1-hydroxy-, and their derivatives (WEIL and HEERDT), A., i, 256.

Naphthalene-1-azo-*p*-benzylcamphor, β -hydroxy- (HALLER and BOUDIN), A., i, 857.

Naphthalene-1-azo-*p*-benzylidenecamphor, β -hydroxy- (HALLER and BOUDIN), A., i, 356.

5-Naphthaleneazo-6-methoxy-2-phenylquinoline-4-carboxylic acid, 5-hydroxy- (HALBERKANN), A., i, 175.

Naphthaleneazo-4-nitro- α -naphthol, 4-nitro- (MUKERJI), T., 2881.

Naphthalene-1-carboxylic acid, 4-bromo-, and its methyl ester (MAYER and SIEGLITZ), A., i, 740.

Naphthalene-2-carboxylic acid, 4-chloro-1-hydroxy- (WEIL and HEERDT), A., i, 257.

Naphthalene-1:4-dicarboxyl chloride (SCHOLL and NEUMANN), A., i, 261.

Naphthalene-1:4-dicarboxylic acid, and its methyl ester (MAYER and SIEGLITZ), A., i, 741.

Naphthalene-2:6-dicarboxylic acid, 1:5-dihydroxy-, and its barium salts and dinitro-derivative (HEMMELMAYR), A., i, 836.

- Naphthalene-1:5-disulphonic acid**, salts of (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 244.
- Naphthalene-3:6-disulphonic acid**, 1:8-diamino-, and 1:8-dinitro-, salts of (LYNCH), A., ii, 881.
- Naphthalenedi-*m*-thiobenzoic acid**, (SMILES and GRAHAM), T., 2509.
- Naphthalene-1-sulphonic acid**, 5-amino-, acetyl derivative, and 5-chloro- (REISSERT), A., i, 584.
- 8-amino-, and 5- and 8-nitro-, and their salts (REISSERT), A., i, 583.
- Naphthalenesulphonamides**, cyano- (KAUFMANN and ZOBEL), A., i, 744.
- Naphthalene-1-sulphonic acid**, 2:5-diamino- (BUCHERER and WAHL), A., i, 464.
- Naphthalene-2-sulphonic acid**, hexaminechromic and hexaminecobaltic salts (EPHRAIM and MOSIMANN), A., ii, 575.
- Naphthalene-4-sulphonic acid**, 1:8-dinitro-, sodium salt and sulphonyl chloride of (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 245.
- Naphthalene-5-sulphonic acid**, 1:6-dihydroxy-, disulphurous acid ester (BUCHERER and WAHL), A., i, 464.
- Naphthalene-8-sulphonic acid**, 4-chloro-1-hydroxy- (KALLE & Co.), A., i, 819.
- 4-chloro-1-nitro-, and its sodium salt (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 244.
- Naphthalenesulphonic acids**, amine salts, solubilities of (WALES), A., i, 727.
- substituted, identification of (LYNCH), A., ii, 881.
- Naphthalene-1-sulphonyl chloride**, 2-cyano- (KAUFMANN and ZOBEL), A., i, 744.
- Naphthalene-5-sulphonyl chloride**, 4-nitro-1-amino-, acetyl derivative (REISSERT), A., i, 584.
- 3-β-Naphthalenesulphonyl-2-methylquinoline**, and its platinichloride (TROGER and MENZEL), A., i, 270.
- β-Naphthalenesulphonylpiperidine** (BERGELL), A., i, 720.
- ββ-Naphthaphenthiazine**, and 3-amino-, 6:11-dihydroxy-, and 3-imino-, and their derivatives (FRIES and KERKOW), A., i, 578.
- ββ-Naphthaphenthiazine-6:11-dicarboxylic acid**, ethyl ester, 11:12-lactam (FRIES and KERKOW), A., i, 578.
- ββ-Naphthaphenthiazine-6:11-quinone**, and its derivatives, and 3-amino-, 3-bromo-, and 3-chloro- (FRIES and KERKOW), A., i, 578.
- 1:2-Naphthaquinols** (FRIES), A., i, 42.
- β-Naphthaquinoline**, diamino-, and dinitro- (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.
- Naphthaquinone**, hydroxy-derivatives (WHEELER and ANDREWS), A., i, 354; (WHEELER and NAIMAN), A., i, 1165.
- 1:4-Naphthaquinone**, mono- and dibromo-5-hydroxy-, 2-bromo-2:3-dihydroxy-, and mono- and di-chloro-5-hydroxy- (WHEELER and NAIMAN), A., i, 1165.
- Naphthaquinoneanil**, hydrochloride and methyl ether of (KALLE & Co.), A., i, 824.
- Naphthasaccharin I. and II.** See Naphthoicsulphonimides.
- 4:1:8-Naphthasultamazo-1'-naphthalene-4'-sulphonic acid** (KÖNIG and KÖHLER), A., i, 880.
- 1:8-Naphthasultam-4-sulphonic acid**, and its salts and derivatives (KÖNIG and KEIL), A., i, 822.
- Naphthasultone**, preparation of sulphonyl chlorides of (KALLE & Co.), A., i, 819.
- Naphthathiam**, and nitroso- (REISSERT), A., i, 583.
- Naphthathiam-blue**, constitution of, and diamino- (REISSERT), A., i, 583.
- α-Naphthatriazole-2:3-dicarboxylic acid**, 3-hydroxy-, methyl ester (DIELS and WACKERMANN), A., i, 1195.
- α-Naphthiminazole**, 1-chloro- (FISCHER and KRACKER), A., i, 957.
- αβ-Naphthiminazole-6-sulphonic acid** (MORGAN and ROOKE), A., i, 134.
- α-Naphthiminoazolone** (DIELS and SØRENSEN), A., i, 1195.
- 2:2'-ββ-Naphthindigotin** (FIERZ and TORLER), A., i, 869.
- Naphthisatin**, β-thio- (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1179.
- meso-Naphthodixanthylene** (ECKERT and ENDLER), A., i, 941.
- 2-Naphthoic acid**, 1-bromo- (MAYER and SIEGLITZ), A., i, 744.
- 3-Naphthoic acid**, 6-amino-2-hydroxy- (FROELICHER and COHEN), T., 1659.
- Naphthoic acids**, reduction of (WEIL and OSTERMEIER), A., i, 139.
- Naphthoicsulphonimides**, and their derivatives (KAUFMANN and ZOBEL), A., i, 744.
- α-Naphthol**, 3-chloro- (FRANZEN and STAUBLE), A., i, 450.
- 5-iodo- (SCHOLL, SEER, and WEITZENBÖCK), A., i, 651.
- 2:4-dinitro-, and its 7-sulphonic acid, cobaltammines of (MORGAN and KING), T., 1727.

- β*-Naphthol**, preparation of (MAY), A., i, 449.
 detection of, in foods, spices, and beverages (KINUGASA and TATSUNO), A., ii, 400.
- β*-Naphthol**, bromo- and bromonitro-derivatives (FRANZEN and STÄUBLE), A., i, 450.
- 4:8-dichloro-, and its methyl ether (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 244.
- Naphthols**, substitution in (FRANZEN and STAUBLE), A., i, 450.
- o*-***β*-Naphtholazophenylacetic acid**, and its sodium salt and esters (NEBER), A., i, 545
- β*-Naphthol-3-carboxylic acid**, methyl ester, condensation of methyl *n*-opionate with (STOSIUS), A., i, 746.
- Naphtholcarboxylic acids**, reduction of (WEIL and OSTERMEIER), A., i, 139; (WEIL and HEERDT), A., i, 256.
- α*-Naphthol-3:6-disulphonic acid**, 8-amino- (*H-acid*), cobaltammine salt of (MORGAN and MOSS), T., 2864.
 estimation of (LEE), A., ii, 235.
- α*-Naphthol-2:5- and 4:5-disulphonic acids**, 6-amino- (BUCHERER and WAHL), A., i, 248.
- α*-Naphthol-2-sulphonic acid**, sodium salt, use of in estimation of amino-compounds (MATHEWSON), A., ii, 882.
- α*-Naphthol-5-sulphonic acid**, 6-amino-, and its derivatives (BUCHERER and WAHL), A., i, 247.
 action of sulphites on (BUCHERER and WAHL), A., i, 464.
- Naphtholsulphonic acids**, amino-, action of phenylhydrazine-bisulphite mixture on (BUCHERER and ZIMMERMANN), A., i, 465.
- β*-Naphthoxyethyl aniline** (CLEMO and PERKIN), T., 649.
- α*- and *β*-Naphthoxyparaffins** (ELEKTRO-CHEMISCHE WERKE, BOSSHARD, and STRAUSS), A., i, 913.
- α*- and *β*-Naphthylbenzoic acids**, acetyl derivatives (MCMULLEN), A., i, 1025.
- α*-Naphthyl methyl ether**, compound of tetrachlorophthalimide with (PFEIFFER and FLATER), A., i, 342.
 5-iodo- (SCHOLL, SEER, and WEITZENBÖCK), A., i, 651.
 thiocyanate (CHALLENGER and WILKINSON), T., 100.
- β*-Naphthyl ethyl disulphide** (LECHER and WITTMER), A., i, 642.
- α*- and *β*-Naphthyl *β*-chloroethyl ethers**, and 1-amino-, and 1-nitro- (CLEMO and PERKIN), T., 646.
- α*- and *β*-Naphthyl mercaptans**, hydroxy- (WATSON and DUTT), T., 2416.
- α*- and *β*-Naphthylacetaldoximes** (MAYER and SIEGLITZ), A., i, 742, 743.
- α*- and *β*-Naphthylacetic acid**, bromo-, and their derivatives (MAYER and SIEGLITZ), A., i, 742, 744.
- d*-Naphthylalanine** (SASAKI and KINOSE), A., i, 303.
- dl*-Naphthylalanine**, action of *Bacillus proteus* on (SASAKI and KINOSE), A., i, 303.
- α*-Naphthylamine**, electrolytic oxidation of (ONO), A., i, 1008.
 substitution in (FRANZEN and STAUBLE), A., i, 450.
- α*-Naphthylamine**, 5-iodo-, and its salts (SCHOLL, SEER, and WEITZENBÖCK), A., i, 650.
 8-thiol-, and its dibenzoyl derivative (REISSERT), A., i, 584.
- β*-Naphthylamine**, compound of tellurium tetrabromide with (LOWY and DUNBROOK), A., i, 446.
 additive compounds of, with azo-esters, and their oxidation products (DIELS and WACKERMANN), A., i, 1068.
 analysis of (LEE and JONES), A., ii, 883.
- α*- and *β*-Naphthylamines**, equilibria of, with dihydroxynaphthalenes (KREMAN, HEMMELMAYR, and RIEMER), A., i, 1010.
 hydroferrocyanides of (CUMMING), T., 1295.
- α*- and *β*-Naphthylamines**, chloro-derivatives, and their derivatives (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 244.
- α*-Naphthylamine-8-sulphonic acid**, 4-chloro-, sodium salt (FRIEDLANDER, KARAMESSINIS, and SCHENK), A., i, 244.
- o*-*α*-Naphthylaminobenzoylbenzene**, *o*-2' nitro- (MAYER and FREUND), A., i, 866.
- Naphthylaminodiphenyl**, 4-amino-1:4'-dihydroxy- (KALLE & Co.), A., i, 824.
- N*-*α*-Naphthyl-*β*-aminoethyl hydrogen sulphate** (SAUNDERS), T., 2675.
- p*-*α*-Naphthylaniline**, and its benzylidene derivative (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1197.
- α*-Naphthyl 1-anthraquinonyl ketone** (SCHOLL, SEER, and DATMER), A., i, 258.
- α*-Naphthylarsinic acid**, sulphonic acid of, and its potassium salts (HILL and BALLS), A., i, 1080.

- Naphthylazobenzene** (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1197.
- 2- β -Naphthylbenzoxazole** (SKRAUP and MOSER), A., i, 575.
- α -1- and -2-Naphthylbutanes** (MAYER and SIEGLITZ), A., i, 741, 743.
- α -1- and -2-Naphthylbutan- γ -ones, and their derivatives** (MAYER and SIEGLITZ), A., i, 741.
- 1- α -Naphthyl-2:3-dibenzylquinoxalinium chloride, $\alpha\alpha'$ -dicyano-** (DUTT and SEN), T., 2666.
- β -Naphthylidimethylamine-6-sulphonic acid, potassium salt** (MORGAN and ROOKE), A., i, 135.
- β -Naphthyl- β -dimethylaminoethyl ether, hydrochloride** (CLEMO and PERKIN), T., 646.
- 1:2-Naphthylencarbamide.** See α -Naphthiminoazalone.
- 1:4-Naphthylenediamine-2-*p*-toluene-6(7)sulphonic acid, monoacetyl derivative, and its sodium salt** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 579.
- α -1-Naphthylethylamine** (SAMUELSON), A., i, 823.
- α - and β -Naphthylethylamines, and their derivatives** (MAYER and SIEGLITZ), A., i, 742, 743.
- β -Naphthylethylmalonic acid** (MAYER and SIEGLITZ), A., i, 743.
- α -Naphthylglycine-8-carboxylic acid, sodium salt** (FIERZ and SALLMANN), A., i, 870.
- β -Naphthylglycine-3-carboxylic acid, and its sodium salt** (FIERZ and TOBLER), A., i, 870.
- Naphthylglycinesulphonic acids** (FIERZ and SALLMANN), A., i, 870.
- Naphthylideneanthranilic acid, β -hydroxy-** (EKELEY, ROGERS, and SWISHER), A., i, 935.
- α -Naphthylidenenitroanilines, 2- and 4-hydroxy-** (MORGAN and REEVES), T., 4.
- 5-Naphthylimino-2-thio-3-naphthyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole** (GUHA), A., i, 877.
- 5-Naphthylimino-2-thio-3-phenyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole** (GUHA), A., i, 877.
- 5-Naphthylimino-2-thio-3-tolyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole** (GUHA), A., i, 877.
- α -Naphthylmethyl ethyl ether, 4-bromo-** (MAYER and SIEGLITZ), A., i, 742.
- α - and β -Naphthylmethyl bromides, bromo-** (MAYER and SIEGLITZ), A., i, 742, 743.
- α - and β -1-Naphthylmethylacetacetic acids, ethyl esters** (MAYER and SIEGLITZ), A., i, 741, 743.
- β -Naphthylmethylamine-6-sulphonic acid, and its salts and derivatives, and 1-nitroso-** (MORGAN and ROOKE), A., i, 134.
- α -1- and -2-Naphthylmethylbutyric acids, and their derivatives** (MAYER and SIEGLITZ), A., i, 741, 743.
- α - and β -Naphthylmethylmalonic acids, and their ethyl esters** (MAYER and SIEGLITZ), A., i, 741, 743.
- β -Naphthyl methyl ketone, and its derivatives** (V. BRAUN, HAHN, and SEEMANN), A., i, 729.
- α - and β -Naphthylmethylmalonic acids, and bromo- and their derivatives** (MAYER and SIEGLITZ), A., i, 741, 743.
- β -Naphthylmethylnitrosoamine-6-sulphonic acid** (MORGAN and ROOKE), A., i, 134.
- α -1- and -2-Naphthyl- β -nitroethylenes** (MAYER and SIEGLITZ), A., i, 742, 743.
- Naphthylpropionic acids, bromo-, and their derivatives** (MAYER and SIEGLITZ), A., i, 742.
- 2- β -Naphthylquinoline, methiodide of** (V. BRAUN, HAHN, and SEEMANN), A., i, 729.
- α -Naphthylthioselenium acetylacetone** (MORGAN, DREW, and BARKER), T., 2453.
- Narcosis, theory of** (TRAUBE and KLEIN), A., i, 198; (JOACHIMOGLU), A., i, 199.
- Narcotics, distribution coefficients of** (AIELLO), A., i, 301.
- mode of action of** (WIELAND), A., i, 497.
- effective strengths of** (FÜHNER), A., i, 199.
- Narcotine, estimation of, in opium** (RAKSHIT), A., ii, 96.
- Naurite** (ELMCHNER), A., ii, 714.
- Neat's foot oil, analytical constants of** (EUKART), A., i, 967.
- Neosarphenamine.** See Neosalvarsan.
- Neodymium oxide, density and molecular volume of** (PRANDTL), A., ii, 379.
- Neon, spectrum of** (BROUGHALL), A., ii, 541.
- Röntgen spectrum and ionisation potential of** (GROTRIAN), A., ii, 179.
- series spectrum of** (SEELIGER), A., ii, 17.
- excitation and ionisation potentials of** (HERTZ), A., ii, 733.
- frequency of electrons in the atom of** (BROUGHALL), A., ii, 213.

- Neosalvarsan**, toxicity of (HART and PAYNE), A., i, 704.
- Neotruxinic acids**, and their derivatives (STOERMER and BACHÉ), A., i, 832.
- Neocyttaridium**, atomic number of (URBAIN), A., ii, 505.
- Nepeta japonica*, essential oil from (MURAYAMA and ITAGAKI), A., i, 44.
- Nephelometer** (KUGELMASS), A., ii, 692.
- Nephelometers**, theory and use of (WELLS), A., ii, 308; (WEINBERG), A., ii, 309; (BAUDOUIN and BÉNAUD), A., ii, 862.
- Nephelometry** of colloidal solutions (BECHHOLD and HEBLER), A., ii, 693. of coloured hydrosols (BECHHOLD and HEBLER), A., ii, 652.
- Nephritis** from chromic acid (MAJOR), A., i, 610. nitrogen constituents of blood in (WILLIAMS), A., i, 401. nitrogen retention in (WEISS and GARNER), A., i, 704. dropsical, excretion of chlorides in (VIOLE), A., i, 968.
- Nerves**, diffusion of hydrogen and hydroxyl ions in relation to the action of (ADAMS), A., i, 1211. sensory, stimulation of, by alcohols and chlorohydrins (IRWIN), A., i, 899.
- Nervous system**, biochemistry of the (PIGHINI), A., i, 295.
- Neutral salt action** at high concentrations (AKERLÖF), A., ii, 134.
- Nickel**, spectrum of (MÜLLER), A., ii, 725. electrolytic, structure of, and its deposition and solution potential (KOLLSCHÜTTER and SCHÖDL), A., ii, 648. scattering of electrons by (DAVISSON and KUNSMAN), A., ii, 251. catalytic hydrogenation with (RIDEAL), T., 309. effect of copper on the catalytic activity of (ARMSTRONG and HILDITCH), A., ii, 757. occurrence and estimation of, in plants (BERTRAND and MOKRAGNATZ), A., i, 1099. in arable soils (BERTRAND and MOKRAGNATZ), A., i, 975.
- Nickel alloys** with chromium, expansion of (CHENEVARD), A., ii, 153. estimation of chromium in (LÖFFELBEIN; HILD), A., ii, 660. with copper and zinc (VOIGT), A., ii, 295. with manganese, electrochemistry of (TAMMANN and VADERS), A., ii, 380.
- Nickel salts**, catalytic hydrogenation by means of (v. BRAUN and KIRSCHBAUM), A., i, 727.
- Nickel chloride**, equilibrium in the system, ammonium chloride, water, and (CLENDINNEN), T., 801. equilibrium of the reduction of, by hydrogen (BERGER and CRUT), A., ii, 38. haloids, crystal structure of the ammoniates of (WYCKOFF), A., ii, 573. hydroxide, preparation of colloidal solutions of (TOWER and COOKE), A., ii, 853. nitrate, composition and crystal structure of (WYCKOFF), A., ii, 573. nitrites, complex (CUTTICA), A., ii, 448. oxide, reduction of, by hydrogen (BERGER), A., ii, 508. tungstate (SMITH), A., ii, 774.
- Nickel organic compounds** :— with dioximes (BJERRUM), A., i, 18, 19, 20, 21.
- Nickel estimation** :— estimation of, electrometrically (MÜLLER and LAUTERBACH), A., ii, 875. estimation of, gravimetrically, as dioxide (VAUBEL), A., ii, 875. estimation of, in silicate rocks (HACKL), A., ii, 458. estimation of, in steel (RUBRICIUS), A., ii, 163; (SIMON), A., ii, 529. estimation of magnesium in (CHALUPNY and BREISCH), A., ii, 227.
- Nicotine**, estimation of, in tobacco (LIOTTA), A., ii, 404.
- Nicotinic acid** (*pyridine-3-carboxylic acid*), amyl and ethyl esters (WOLFFENSTEIN), A., i, 861. ethyl ester, *N*-ethiodide (WOLFFENSTEIN), A., i, 365. methyl ester, preparation of *N*-alkylhaloids of (WOLFFENSTEIN), A., i, 365. dipropylamide and piperidide of (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1058.
- Nicotinodilethylamide** (HARTMANN and SEIBERTH), A., i, 679.
- Nicotinodipropylamide** (HARTMANN and SEIBERTH), A., i, 679.
- Nicotinopiperidide** (HARTMANN and SEIBERTH), A., i, 679.
- Night-blue**, surface tension of aqueous solutions of (DE LAQUIERRE), A., ii, 262. effect of anions on the colloidal state of (AKAMATSU), A., ii, 830.
- Niobium pentoxide**, action of light on (RENZ), A., ii, 61.

- Nirvanol** (4-phenyl-4-ethylhydantoin), synthesis of (READ), A., i, 954.
- Niton** (*radium emanation*), spectrum of (NYSWANDER, LIND, and MOORE), A., ii, 803.
- activity of γ -rays from (SLATER), A., ii, 681.
- effect of, on blood and urine (HAUENSTEIN), A., i, 80.
- Nitration** of hydrocarbons (BATTEGAY and BRANDT), A., i, 1001.
- Nitrides**, rate of formation of (TAMMANN), A., ii, 852.
- Nitrification** (MIYAKE and SOMA), A., i, 1096.
- in soils (STEPHENSON), A., i, 416.
- Nitriles**, dynamics of the formation of, from anhydrides and amides (KREMANN, ZOFF, and OSWALD; KREMANN, ROSLER, and PENKNER), A., ii, 748.
- action of hydrogen peroxide on (OLIVERI-MANDALA), A., i, 544.
- action of organomagnesium compounds on (BAERTS), A., i, 817.
- aromatic, electrochemical oxidation of (FICHTER and GRISARD), A., i, 37.
- iso* **Nitriles** (PASSERINI), A., i, 731.
- Nitro-compounds**, reduction of, by stannous chloride (GOLDSCHMIDT, STORM, and HASSEL), A., ii, 361.
- reduction of, by titanium trichloride (RATHSBURG), A., i, 242.
- aromatic, formation of salts from (LIFSCHITZ), A., i, 737.
- catalytic reduction of (BRAND and STEINER), A., i, 536.
- removal of halogen atoms from (MACBETH), T., 1116.
- action of sodium hydrogen sulphite on (WEIL and MOSER), A., i, 443.
- halogenated, reduction of (BURTON and KENNER), T., 675.
- detection of (PRINS), A., ii, 877.
- Nitrogen atom**, structure of the (MOIR), T., 1808.
- active modification of (NEWMAN), A., ii, 279, 639; (WENDT), A., ii, 639.
- origin of the spectrum of (DUFFIEUX), A., ii, 597.
- ionisation and excitation tension of (BRANDT), A., ii, 186.
- heat of vaporisation and specific heat of (MATHIAS, CROMMELIN, and ONNES), A., ii, 472.
- pure, density of (MOLES), A., ii, 562.
- atmospheric, from Madrid, density of (PAYÁ and MOLES), A., ii, 762.
- equation of state for (BARTELS and EUCKEN), A., ii, 117.
- isotherms of (HOLBORN and OTTO), A., ii, 737.
- Nitrogen**, rectilinear diameter of (MATHIAS, ONNES, and CROMMELIN), A., ii, 562.
- absorption of, by calcium and its alloys (RUFF and HARTMANN), A., ii, 377.
- viscosities of carbon monoxide and (SMITH), A., ii, 549.
- electrolytic reduction of (FICHTER and SUTER), A., ii, 372.
- explosion of acetylene and (GARNER and MATSUNO), T., 1729.
- reaction of ethylene and, under the electric discharge (MIYAMOTO), A., i, 418.
- reaction between carbon, barium carbonate and (ASKENASY and GRUDE), A., ii, 445.
- reaction between carbon, sodium carbonate and (INGOLD and WILSON), T., 2278.
- quadrivalent, radicals containing (WIELAND and KOGL), A., i, 726.
- fixation of, in soils (GRAVES, CARTER, and LUND), A., i, 976.
- effect of diet on fixation of, in the organism (DIENES), A., i, 293.
- Nitrogen compounds**, photosynthesis of (BALY, HEILBRON, and HUDSON), T., 1078.
- Nitrogen trichloride**, attempted preparation of (NOYES), A., ii, 143.
- chlorination by means of (COLEMAN and NOYES), A., i, 133.
- monoxide* (*nitrous oxide*), viscosities of carbon dioxide and (SMITH), A., ii, 549.
- solubility of, in various solvents (KUNERTH), A., ii, 823.
- narcotic action of (WIELAND), A., i, 497.
- dioxide* (*nitric oxide*), preparation of, from boron nitride and metallic oxides (SBOGI and NASINI), A., ii, 564.
- oxidation of (BURDICK), A., ii, 272; (BRINER, NIEWIAZSKI, and WISWALD), A., ii, 563.
- action of, on metallic carbonyls (MOND and WALLIS), T., 32.
- action of, and its derivatives on sodium arsenite (GUTMANN), A., ii, 844.
- estimation of (KLEMENC and BUNZL), A., ii, 783.
- trioxide* (*nitrous anhydride*) (FOERSTER), A., ii, 284.
- per-* or *tetr-*oxide, liquid, analysis of (SANFOURCHE), A., ii, 454.
- oxides, formation and decomposition of (BODENSTEIN, BOES, LINDNER, and RAMSTETTER), A., ii, 372.

Nitrogen oxides, formation of, in gas analysis (JONES and PARKER), A., ii, 228.

reactions of alkalis with (SAN-FOURCHE), A., ii, 762.

compounds of, with olefines and zinc (SCHAARSCHMIDT, VEIDT, and SCHLOSSER), A., i, 645.

analysis of mixtures of (BURDICK), A., ii, 583.

estimation of, in air (ALLISON, PARKER, and JONES), A., ii, 313.

Nitric acid, oxidation reactions in the synthesis of (MATIGNON), A., ii, 563.

electrolytic concentration of aqueous solutions of (CREIGHTON), A., ii, 212.

heats of neutralisation of (RICHARDS and ROWE), A., ii, 425.

heat developed on mixing sulphuric acid, water and (McDAVID), A., ii, 617.

vapour pressure of aqueous solutions of (SPROESSER and TAYLOR), A., ii, 143.

action of, on metals (BANERJI and DHAR), A., ii, 756.

estimation of nitrogen in esters of (KESSELER, ROHM, and LUTZ), A., ii, 392.

Nitrates, organic compounds giving precipitates with (GIRAL PEREIRA), A., ii, 523.

assimilation of (BAUDISCH), A., i, 194.

detection of (FORESTI), A., ii, 524.

detection and estimation of, in serum and in urine (NOLIE), A., ii, 583.

estimation of, in drinking water (REUSS), A., ii, 454.

Nitrous acid, velocity of decomposition of (KLEMENC and POLLAK), A., ii, 432.

rate of reaction of, with aliphatic amino-acids (DUNN and SCHMIDT), A., i, 1024.

action of, on iodides, in presence of oxygen (LOMBARD), A., ii, 313.

and its salts, detection of (FALCIOLA), A., ii, 783.

estimation of, in presence of arsenious acid (KLEMENC), A., ii, 865.

Nitrites, reaction of, with thiosulphates (FALCIOLA), A., ii, 453.

assimilation of (BAUDISCH), A., i, 194.

estimation of, in presence of nitrates (MACH and SINDLINGER), A., ii, 783.

Nitrogen :—

Pernitric acid, properties and structure of (TRIFONOV), A., ii, 845.

as an analytical reagent (MATHEWSON), A., ii, 883.

Nitrogen estimation :—

estimation of, by Dumas's method (MOHR), A., ii, 82.

estimation of, by the Kjeldahl method (PAUL and BERRY), A., ii, 82; (PARNAS and WAGNER), A., ii, 312; (M. and I. SBOROWSKY), A., ii, 783.

estimation of, by a modified Kjeldahl method (MEARS and HUSSEY), A., ii, 159; (SKUTIL), A., ii, 582.

estimation of, by a micro-Kjeldahl method (LING and PRICE), A., ii, 522.

estimation of, microchemically (SCHOELLER), A., ii, 159; (ACÉL), A., ii, 225; (VALLÉE and POLONOVSKI), A., ii, 312; (HOLTZ), A., ii, 521.

estimation of, in ammonium salts (MEURICE), A., ii, 225.

estimation of, in fertilisers (MACH and SINDLINGER), A., ii, 783.

estimation of, in nitric esters (KESSELER, ROHM, and LUTZ), A., ii, 392.

amino-acid, estimation of, by the ninhydrin reaction (RIFFART), A., ii, 718.

non-protein, estimation of, in blood (PONDER), A., ii, 583; (RICHTER-QUITZNER and HOENLINGER), A., ii, 717.

estimation of, in serum (CRISTOL), A., ii, 583.

Nitro-groups, replacement of diazo-groups by (VESELY and DVOŘÁK), A., i, 690.

aromatic, reduction of, by tissues (LIPSCHITZ and GOTTSCHALK), A., i, 298.

estimation of, in aromatic organic compounds (CALLAN and HENDERSON), A., ii, 524.

***β*-Nitrole- γ -ketopentane**, α -oximino- (PONZIO and RUGGERI), A., i, 629.

***β*-Nitrole- γ -ketophenylbutane**, α -oximino- (PONZIO), A., i, 1038.

Nitrones, catalytic reduction of (CUSMANO), A., i, 143.

Nitrosyl chloride, action of, on *n*-heptane (LYNN and HILTON), A., i, 417.

Nitrosylselenic acid (MEYER and WAGNER), A., ii, 372.

Nitrous acid. See under Nitrogen.

Nitrous anhydride. See Nitrogen trioxide.

- Nitzschia closterium*, synthesis of vitamin-A by (JAMESON, DRUMMOND, and COWARD), A., i, 1099.
- Nonadecan- δ -ol**, and its phenylurethane (LEROIIDE), A., i, 217.
- Δ^{α} -Nonenoic acid**, vanillylamide and *p*-hydroxybenzylamide of (OTT and ZIMMERMANN), A., i, 137.
- Nonoisobutylamide** (ASAHINA and ASANO), A., i, 506.
- n*-Nonoic acid**, preparation of (ASAHINA and ISHIDA), A., i, 520.
sodium salt, properties of solutions of (FLECKER and TAYLOR), T., 1101.
- Norcholoidanic acid**, and its dimethyl ester (WIELAND and ADICKES), A., i, 338.
- Norpinic acid**, structure and synthesis of (GANGULY), A., i, 1153.
- Norsolannellic acid** (WIELAND and SCHULENBURG), A., i, 346.
- Nortropinonecarboxylic acid**, ethyl ester, and its derivatives (MERCK, WOLFES, and MAEDER), A., i, 1173.
- Novocaine**, estimation of (HANSON), A., ii, 405.
- Nucleic acid**, compounds of proteins with (STEUDEL and PEISER), A., i, 1200.
metabolism of. See Metabolism.
animal, preparation and analysis of (LEVENE), A., i, 1075.
in bacteria (SCHAEFFER, FOLKOFF, and JONES), A., i, 1095.
pancreatic (HAMMARSTEN and JORPES), A., i, 387.
- Nucleic acids**, anticoagulating action of (DOYON), A., i, 82, 1087.
from spleen of cattle (STEUDEL), A., i, 297.
- Nuclein metabolism**. See Metabolism.
- Nutmeg butter** (B.), A., i, 975.
- Nuts**, Barbassu, constituents of (JUMELLE), A., i, 207.
- Nut tree oil**, purging (L.), A., i, 908.
- Nystagmus** (PENTIMALLI), A., i, 302.
- O.
- Oak**, black powder from a hollow (v. LIPPMANN), A., i, 311.
tannin from the (FREUDENBERG and VOLBRECHT), A., i, 1046.
- Oats**, ergot of. See Ergot.
- Obituary notices** :—
Ganesh Sakharan Agashe, T., 745.
James Robert Appleyard, T., 2898.
Adrian Brown, T., 2899.
William Gowland, T., 2907.
Philippe Auguste Guye, T., 2909.
Edward William Taylor Jones, T., 746.
- Obituary Notices** :—
William Kellner, T., 2912.
George William Macdonald, T., 2913.
John Spiller, T., 748.
Lionel William Stansell, T., 2916.
- Octa-acetyldisalleinethylamine** (ZEMPLEN and KUNZ), A., i, 564.
- Octa-acetyldisalleinmethylamine** (ZEMPLEN and KUNZ), A., i, 564.
- Δ^{γ} -Octadiene**, and its ozonide (SCHWEITZER), A., i, 215.
- $\Delta^{\alpha\gamma}$ -Octadiene- $\alpha\delta\theta$ -tetracarboxylic acid**, δ -bromo- $\gamma\delta$ -dihydroxy- (CHANDRASENA and INGOLD), T., 1318.
- s*-Octahydroanthracene**, and its sulphonic acid (SCHROETER and TETRALIN G. M. B. H.), A., i, 1136.
- s*-Octahydrophenanthrene**, and its derivatives (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- Octamethyltetra-amylose** (PRINGSHEIM and PERSCH), A., i, 113.
tetra-acetate (PRINGSHEIM and PERSCH), A., i, 632.
- 1:2:4:6:1':2':4':6'-Octamethyltetrahydro-4:4'-dipyridyl** (EMMERT and WERR), A., i, 680.
- Octane**, β -chloro- γ -hydroxy- (DETREUF), A., i, 327.
- 0:3:3-dicycloOctane** (SCHROETER and VOSSEN), A., i, 122.
- 0:3:3-dicycloOctane-3:7-diol** (SCHROETER and VOSSEN), A., i, 122.
- 0:3:3-dicycloOctane-3:7-dione** (SCHROETER and VOSSEN), A., i, 122.
- Octapyridine- μ -dichlorodiferriehloride** (WEINLAND and KISSLING), A., i, 365.
- $\Delta^{\alpha\delta\theta}$ -Octatriene- $\alpha\delta\theta$ -tetracarboxylic acid**, $\gamma\delta$ -dihydroxy- (CHANDRASENA and INGOLD), T., 1319.
- B* γ -Octene oxide** (DETREUF), A., i, 327.
- 0:3:3-dicyclo- Δ^1 -Octene-3:7-dione-2:4:6:8-tetracarboxylic acid**, methyl ester (SCHROETER and VOSSEN), A., i, 122.
- Octopus**, muscle extract of (QUAGLIARIELLO), A., i, 608.
- n*- and *sec*-Octyl fluorides** (SWARTS), A., i, 101.
- Odour**, relation of, to chemical constitution (ZWAARDEMAKER), A., i, 607.
and molecular structure (DELANGE), A., i, 894.
perception of (KOMURO), A., i, 84.
value of (UNGERER and STODDARD), A., i, 894.
- Odours**, aromatic, classification of (OHMA), A., i, 1090.
- Edema**, nitrogen content of the fluid in (STRISOWER), A., i, 964.

- Edema**, artificial, effect of calcium and potassium ions on the production of (HAMBURGER), A., i, 606.
- Oils**, relation between the refractive index and chemical characteristics of (PICKERING and COWLISHAW), A., ii, 325.
- velocity of extension of, on the surface of water (WOOE), A., ii, 197.
- catalysts in the hydrogenation of (UENO), A., i, 983.
- saponification of (NORRIS and MCBAIN), T., 1362; (LANGTON), A., i, 982.
- drying, oxidation of (COFFEY), T., 17.
- essential. See Oils, vegetable.
- fatty, drying of (SLANSKY), A., i, 1113.
- lubricating mineral, chemistry of (DUNSTAN and THOLE), A., i, 213.
- mineral and tar, oxidation of (HARRIES), A., i, 514.
- See also Petroleum.
- vegetable (ROURE-BERTRAND FILS), A., i, 845, 846, 847.
- from the Sudan (JOSEPH and WHITFIELD), A., i, 754.
- from Tonkin and Annam (GATTEFOSSÉ), A., i, 1167.
- effect of sunlight on the formation of, in plants (CANALS), A., i, 907.
- detection of, in animal fats (MUTTELET), A., ii, 168.
- estimation of the acetyl value of (LEYS), A., ii, 167; (COOK), A., ii, 324.
- estimation of the iodine-bromine number of, without using potassium iodide (WINKLER), A., ii, 534.
- Oil-water** emulsions, reversal of phases in (BHATNAGAR), A., ii, 204.
- Oleannilide dibromide** (NICOLET), A., i, 106.
- Olefines**, magnesium compounds of (KRESTINSKI), A., i, 1128.
- Oleic acid**, measurement of the surface pressure of thin layers of (MARCELIN), A., ii, 687.
- catalytic decomposition of (MAILHE), A., i, 423.
- relation of, to its halogen additive products (NICOLET), A., i, 106.
- sodium salt, ultra-filtration of solutions of (MCBAIN and JENKINS), T., 2325.
- vanillylamide of (OTT and ZIMMERMANN), A., i, 137.
- Oleic anhydride**, iodine value of (HOLDE, WERNER, TACKE, and WILKE), A., ii, 723.
- α -Olein** (AMBERGER and BROMIG), A., i, 804.
- Oleacetylhydroxamic acid** (NICOLET and PELC), A., i, 644.
- Oleodiacetylhydroxamic acid** (NICOLET and PELC), A., i, 644.
- α -Oleo- β -distearin** (AMBERGER and BROMIG), A., i, 804.
- Oleum fagi sylvaticæ*. See Beechnut oil.
- Oligodynamy** of silver (DOERR and BERGER), A., i, 1097.
- Olive oil**, coloration of, with the Villavecchia reagent (PRAX), A., ii, 595.
- Olivine** from lithosiderites (CHIRVINSKII), A., ii, 714.
- n*-Oplanic acid**, methyl ester, condensation of methyl β -naphthol-3-carboxylate with (STOSIUS), A., i, 746.
- Opium**, estimation of meconic acid in (ANNETT and BOSE), A., ii, 791.
- Indian, estimation of alkaloids in (RAKSHIT), A., ii, 96.
- Optical anomalies**, negative, effect of substitution on (LE BAS), A., ii, 9.
- properties of solutions (WASASTJERNA), A., ii, 2.
- rotation. See Rotation.
- Optically active compounds**, influence of constitution on the rotatory power of (RUPE and JAGGI), A., i, 840; (RUPE), A., ii, 602.
- Optotoxin**. See Ethyldihydrocupreicine.
- Oranges**, changes in, on keeping (ANDRÉ), A., i, 209.
- Orchids**, coumarin glucosides in (HÉRISSEY and DELAUNRY), A., i, 210.
- Orcinolsulphamphthalein** (DUTT), T., 2391.
- Ores**, genesis of, in relation to geographical distribution (GREGORY), T., 750.
- Organic chemistry**, problems of (TRAUTZ and WINKLER), A., i, 926.
- Organic compounds**, atomic and molecular refraction in (SWIENTOSLAWSKI), A., ii, 173; (V. AUWERS and KOLLIGS), A., ii, 174.
- rotatory dispersion of (LOWRY and CUTTER), T., 532.
- crystalline, structure of (BRAGG: BECKER and JANCKE), A., ii, 128.
- valency of (HENRICH), A., ii, 704, (STIEGLITZ), A., ii, 705.
- polar and non-polar valency in (GARNER), A., ii, 758.
- linkings of carbon atoms in (WIBAUT), A., ii, 259; (BEUTLER), A., ii, 259, 348.

- Organic compounds**, lability of halogen atoms in (HENDERSON and MACBETH), T., 892; (HIRST and MACBETH), T., 904, 2169; (GRAHAM and MACBETH), T., 1109, 2601; (MACBETH), T., 1116; (BLACK, HIRST, and MACBETH), T., 2527.
- numerical relationships between carbon atoms in (DECKER), A., i, 417.
- thermochemistry of (SWIENTOSLAWSKI), A., ii, 195; (FAJANS), A., ii, 818; (THIEL), A., ii, 819.
- relation between heats of fusion and of solution of (GEHLHOFF), A., ii, 121.
- wet combustion of, by means of chromic acid (SIMON), A., ii, 593.
- biochemical and electrochemical oxidation of (FICHTER), A., ii, 23; (NATHANSON), A., ii, 421; (MULLER), A., ii, 469.
- electrochemistry of the reversible reduction of (CONANT, KAHN, FIESER, and KURTZ), A., ii, 547.
- catalytic reduction of, by platinum oxide (VOORHEES and ADAMS), A., ii, 558.
- phosphorescent mixtures of boric acid with (TIEDE, WULFF, and RAGOSS), A., ii, 8.
- action of Dakin's hypochlorite solution on (ENGFELDT), A., i, 812.
- preparation of bromo-derivatives of, without production of hydrogen bromide (GUPTA and THORPE), T., 1896.
- elimination of carbon dioxide from (KUNZ-KRAUSE and MANICKE), A., ii, 731.
- behaviour of, in plants (CIAMICIAN and GALIZZI), A., i, 503.
- aromatic, estimation of nitro-groups in (CALLAN and HENDERSON), A., ii, 524.
- homologous, variation of physical properties of (PAULY), A., i, 1; (STARK), A., i, 2.
- giving precipitates with nitrates (GIRAL PEREIRA), A., ii, 523.
- micro-analysis of (WELTER), A., ii, 399.
- detection of oxygen in (PICCARD), A., ii, 389.
- estimation of carbon and hydrogen in (WREDE), A., ii, 316; (LINDNER), A., ii, 657; (SIMON and GUILLAUMIN), A., ii, 867; (SIMON), A., ii, 868.
- estimation of carbon and oxygen in (KLEIN and STUBER), A., ii, 159.
- estimation of chlorine in (KLIMONT), A., ii, 580; (VOTOCÉK), A., ii, 863.
- Organic compounds**, estimation of oxygen in (MEULEN), A., ii, 717.
- estimation of phosphorus in (MANCINI), A., ii, 88.
- estimation of phosphorus in, iodometrically (SVANBERG, SjöBERG, and ZIMMERLUND), A., ii, 867.
- estimation of sulphur in (MEULEN), A., ii, 311.
- estimation of thallium in (A. E. and D. GODDARD), T., 488.
- Organic dust**, explosions of (BEYERSDORFER), A., ii, 749.
- Organism**, animal, synthesis of amino-acids in the (SHIPLE and SHERWIN), A., i, 492.
- deamination of amino-acids in the (KOTAKE), A., i, 1218.
- oxidation in the (MITTRA and DHAR), A., i, 1210.
- Organo-metallic compounds** (SCHLUBACH and GOES), A., i, 1204.
- preparation of (GODDARD, ASHLEY, and EVANS), T., 978.
- interaction of, with inorganic haloids (CHALLENGER and RIDGWAY), T., 107.
- Organum vulgare*, constituents of oils from (ANGELESCU), A., i, 460.
- Ornithine**, presence of, in plants (KIESEL), A., i, 412.
- Osazones**, formation of (VAN LAER and LOMBAERS), A., i, 113.
- Osmium**, detection of, with potassium thiocyanate (HIRSCH), A., ii, 459.
- Osmometers**, collodion (BROWN), A., ii, 690.
- Osmosis**, anomalous (LOEB), A., ii, 124, 354, 467, 742.
- See also Electrical endosmosis.
- Osmotic pressure**, measurement of, with the water interferometer (LOTZ and FRAZER), A., ii, 264.
- regulation of (PALMER, ATCHLEY, and LOEB), A., i, 692.
- determination of molecular weight by means of (FOIX), A., ii, 621.
- of protein solutions (LOEB), A., ii, 742.
- Ovalbumin**. See under Albumin.
- Overvoltage** (NEWBURY), T., 7.
- measurement of (TARTAR and KEYES), A., ii, 343.
- on electrodes (ISGARISCHEV and BERKMANN), A., ii, 253, 254.
- Oxalic acid**, and its potassium salt, action of radium rays on (KAILAN), A., ii, 543.
- action of electric arc light on the catalytic decomposition of solutions of (KUNZ-KRAUSE and MANICKE), A., ii, 731.

- Oxalic acid**, crystallised, as a standard in oxidation processes (HILL and SMITH), A., ii, 388.
 formation of, in growth of *Aspergillus* and *Citromyces* (BUTKEWITSCH), A., i, 707, 978.
 in young leaves (BAU), A., i, 309.
 behaviour of, in the animal organism (PINCUSSEN), A., i, 403.
 calcium salt, monohydrate, optical and crystallographic properties of (WHERRY), A., ii, 450.
 equilibrium of potassium carbonate with (RAMANN and SALLINGER), A., ii, 131.
 lead diphenyl salt (GODDARD, ASHLEY, and EVANS), T., 981.
 potassium salt, preparation of, from potassium acetate (EVANS and HINES), A., i, 803.
 sodium salt, formation of, from sodium formate (MATIGNON and MARCHAL), A., i, 915.
 uranyl salt, photolysis of (BAUE and REBMANN), A., ii, 337; (BAUR and HAGENMACHER), A., ii, 338.
 methyl ester, decomposition of, by acetic acid (TURNER and WILSON), A., i, 916.
 detection and estimation of, and its use in standardisation (ROSENTHALER), A., ii, 594.
 estimation of, with permanganate without addition of acid (WITT), A., ii, 594.
 estimation of, in urine (SALKOWSKI), A., ii, 92; (MISLOWITZER), A., ii, 325.
- Oxalodi-5-acenaphthylamide** (FLEISCHER and SCHRANZ), A., i, 1143.
- β -Oxalyl-n-butyric acid**, ethyl ester (FEIST and BREUER), A., i, 521.
- α -Oxalecarboxylic acids**, isomerism of (BETTI and BERLINGOZZI), A., i, 52.
- Oxidation**, simultaneous reduction and (KOTZ and RATHERT), A., i, 236.
 catalysis of (KARCZAG), A., ii, 137.
 biochemical and electrochemical, of organic compounds (FICHTER), A., ii, 23; (NATHANSON), A., ii, 421; (MÜLLER), A., ii, 469.
 electrolytic. See Electrolytic oxidation.
 with potassium permanganate (BIGELOW), A., i, 998.
- Oxidation-reduction reactions**, effect of substitution on the free energy of (LAMER and BAKER), A., ii, 735.
- NN'*-Oxidobisdihydromorphinesulphonic acid** (SPREYER and BECKER), A., i, 676.
- Oximes**, thermochemistry of (DORABIALSKA), A., i, 725; ii, 548.
 isomerism of (BRADY and THOMAS), T., 2098.
- Oximinoacetoacetic acid**, ethyl ester, stereoisomerism of (DORABIALSKA), A., i, 725.
- Oximinoacetophenone**, cobaltous salt (PONZIO), A., i, 662.
- Oximinoacetylacetone**, cupric and cobaltous salts (PONZIO), A., i, 662.
- Oximino benzoylacetone**, cobaltous salt (PONZIO), A., i, 662.
- N*-Oximino-ethers** (BARROW, GRIFITHS, and BLOOM), T., 1713.
- α -Oximinoketones**, cobalt compounds of (PONZIO), A., i, 661.
- Oxindigo**, synthesis of (FRIES and HASENJAGER), A., i, 45.
- Oxindole**, 1-amino-, and its *o*-nitrobenzylidene derivative (NEBER), A., i, 546.
- Oxonium salts** from phenols and their ethers (KEHRMANN, DECKER, and SOLOMINA), A., i, 32; (KEHRMANN, DECKER, and SCHMAJEWSKI), A., i, 33.
- Oxyallobetulin**, and its acetate (SCHULZE and PIEROH), A., i, 1046.
- Oxycellulose** (HEUSER and STOCKIGT), A., i, 719.
 formation and reactions of (CLIFFORD and FARGHER), A., i, 1121.
 action of heat on (JUSTIN-MUELLER), A., i, 11.
 reactions of, with alkalis (SCHWALBE and BECKER), A., i, 232.
 detection of (BECKER), A., ii, 94.
- Oxydases**, chemical nature of (VAN DER HAAR), A., i, 284.
- Oxydehydrocorydaline** (GADAMER and v. BRUCHHAUSEN), A., i, 675.
- Oxydigitogenic acid**, trimethyl ester (WINDAUS and WEIL), A., i, 849.
- Oxygen**, absorption spectra of liquid and gaseous (SHAVER), A., ii, 331.
 band spectrum of (RUNGE), A., ii, 329.
 heat of reaction of, with hæmoglobin (ADOLPH and HENDERSON), A., ii, 350.
 heat of vaporisation and specific heat of (MATHIAS, CROMMELIN, and ONNES), A., ii, 472.
 rectilinear diameter of (MATHIAS and ONNES), A., ii, 561.
 isotherms of (HOLBORN and OTTO), A., ii, 737.
 univalent (GOLDSCHMIDT), A., i, 1148; (GOLDSCHMIDT and SCHMIDT), A., i, 1149.
 compressibility of (GUYE and BATHÉCAS), A., ii, 617.

Oxygen, density of (MOLES), A., ii, 141;
(MOLES and GONZÁLEZ), A., ii, 497;
(MOLES and CRESPI), A., ii, 636.
solubility of, in organic solvents
(FISCHER and PFLEIDERER), A., ii,
841.
velocity of the action of, on metals
(TAMMANN and KOSTER), A., ii, 881.
equilibrium of silver oxide and (KEYES
and HARA), A., ii, 353.
ignition of mixtures of ammonia and
(WHITE), T., 1688.
ignition of mixtures of hydrogen and
(MITSCHEHLICH), A., ii, 358.
combination of hydrogen and, in
presence of colloidal palladium
(SANDONNINI and QUAGLIA), A., ii,
556.
catalytic formation of water from
hydrogen and (PEASE and TAYLOR),
A., ii, 701.
partial pressure of, in human blood
(BARCROFT and NAGAHASHI), A., i,
890.
relation between carbon dioxide and,
in respiration (THUNBERG), A., i,
889.
subcutaneous absorption of, at high
altitudes (BAYEUX), A., i, 79.
Oxygen detection and estimation:—
detection of, in organic compounds
(PICCARD), A., ii, 389.
estimation of small quantities of
(SHEAFF), A., ii, 582.
estimation of, in hydrogen (LARSON
and WHITE), A., ii, 311.
estimation of, in organic compounds
(KLEIN and STEUBER), A., ii, 159;
(MEULEN), A., ii, 717.
Oxyhemoglobin, crystalline, preparation
of (HEIDELBERGER), A., i, 962.
dissociation of (HILL), A., i, 193.
Oxynitrilase. See β -Emulsin.
Oxynitrilase. See σ -Emulsin.
10-Oxy-9-oxythionaphthyl-6-oxy-
phenanthrene(FRIEDLÄNDER,HERZOG,
and v. Voss), A., i, 764.
3-Oxythionaphthen, condensation of
formic acid with (SCHWENK), A., i,
153.
2-Oxythionaphthyl-2-indane-1:3-dione
(FRIEDLÄNDER, HERZOG, and v.
Voss), A., i, 764.
Ozone, preparation and properties of
(RIESENFELD and SCHWAB), A., ii,
637.
luminescence of, when decomposing
(TRAUTZ and SEIDEL), A., ii, 730.
physical constants of (RIESENFELD
and SCHWAB), A., ii, 761.
action of, on amines (STRECKER and
BALTES), A., i, 14.

Ozone, action of, on hydrocarbons
(WHEELER and BLAIR), A., i
1105.
action of, on lactose (SCHONEBAUM),
A., i, 717.
action of, on sugars (SCHONEBAUM),
A., i, 223.
Ozoniser, new (NĚMEČEK), A., ii, 841.

P.

Pagoda corn oil, constituents of (ROURE-
BERTRAND FILS), A., i, 845, 846.
Palladium, occlusion of hydrogen by
(OXLEY), A., ii, 469.
properties of hydrogen desorbed from
(ANDERSON), T., 1153.
separation of, from platinum, by
means of dimethylglyoxime (DAVIS),
A., ii, 662.
Palmitine, conversion of berberine into
(SPATH and LANG), A., i, 166.
Palmitic acid, sodium salt, hydrolysis of
solutions of (MCBAIN, TAYLOR, and
LAING), T., 621.
cetyl ester (BRIGL and FUCHS), A., i,
713.
vanillylamide of (OTT and ZIMMER-
MANN), A., i, 137.
 α -Palmito- β -diolein (AMBERGER and
BROMIG), A., i, 804.
Palmitodistearins, isomeric (AMBERGER
and BROMIG), A., i, 804.
Pancreas, function of the (ADLER), A.,
i, 195.
action of amino-acid hydrochlorides on
the secretion of (ARAI), A., i, 297.
amylolytic activity of, in polyneuritis
(TIGER and SIMONNET), A., i, 195.
nucleic acid from (HAMMARSTEN and
JORPES), A., i, 387.
hydrolysis of yeast-nucleic acid by
boiled extracts of (JONES), A., i,
479.
Papilionaceæ, glucosides from (VERGE-
LOT), A., i, 207.
Paracetaldehyde, preparation of (VOGT
and NIEUWLAND), A., i, 111.
Paracetaldehyde, bromo- (HELFERICH
and SPEIDEL), A., i, 6.
Paraffins, oxidation of, with nitrogen per-
oxide (GRÄNACHER and SCHAUFEL-
BERGER), A., i, 513.
Paraffins, mono- and di-chloro- (ELEK-
TROCHEMISCHE WERKE, BOSSHARD,
and STRAUSS), A., i, 913.
nitro-, absorption spectra of (GRAHAM
and MACBETH), T., 1109.
substituted, reduction of, with
titanium salts (HENDERSON and
MACBETH), T., 892.

- Paraffin wax**, composition of (FRANCIS), T., 496; (FRANCIS, WATKINS, and WALLINGTON), T., 1529, 2804.
 action of, with oxygen, sulphur, selenium or tellurium (SIEBENECK), A., i, 417.
- Paralysis agitans**, sugar regulation in (DRESEL and LEWY), A., i, 969.
- Paramacium**, toxicity of alkoids for (CRANE), A., i, 404.
- Pararosanilineleucosulphonic acid**, salts and hydrates of (WIELAND and SCHEUING), A., i, 58.
- Paris green**, estimation of copper and arsenic in (KOLTHOFF and CREMER), A., ii, 86.
- α -Particles**, disintegration of elements by (RUTHERFORD and CHADWICK), A., ii, 682.
 excitation of γ -rays by (SLATER), A., ii, 13.
 collisions of, with hydrogen nuclei (CHADWICK and BIELER; MCAULAY), A., ii, 12.
 as detonators (HENDERSON), A., ii, 606.
- Passivity** (EVANS; GILLIS), A., ii, 814.
- Patchouli oil** (ROURE-BERTRAND FILS), A., i, 846.
- Paullinia tannin** (NIERENSTEIN), T., 23.
- Pea**, absorption of calcium chloride by the roots of (REDFERN), A., i, 614.
- Peaches**, odorous constituents of (POWER and CHESNUT), A., i, 99.
- Pectin** in stored fruit (CARRÉ), A., i, 1222.
 estimation of, in apples (CARRÉ and HAYNES), A., ii, 401.
- Pectinase** produced by *Rhizopus* (HARTER and WEIMER), A., i, 507.
- Pelargonenin** (CURREY), T., 322.
- Pelargonin** (CURREY), T., 319.
- Pelargonium**, scarlet, colouring matter of the (CURREY), T., 319.
- Pellagra**, deficiency of amino-acids as a cause of (GOLDBERGER and TANNER), A., i, 1092.
 amino-acids and hippuric acid in the urine in (MURLIN), A., i, 965.
- Pellotine**, constitution of (SPÄTH), A., i, 163.
- Penta-acetylsalicylaminethylamine** (ZEMPLEN and KUNZ), A., i, 564.
- Penta-acetylsalicylaminmethylamine** (ZEMPLEN and KUNZ), A., i, 564.
- isoPentacosane** (LEVENE and TAYLOR), A., i, 715.
- isoPentacosic acid**, and its ethyl ester, and α -bromo-, and α -hydroxy- (LEVENE and TAYLOR), A., i, 714.
- isoPentacosyl iodide** (LEVENE and TAYLOR), A., i, 715.
- isoPentacosyl alcohol** (LEVENE and TAYLOR), A., i, 715.
- Pentamethoxy- β -gambier-catechin carboxylic acid** (NIERENSTEIN), T., 28.
- 3:4:6:3':4'-Pentamethoxy-3-phenylchroman**, and 2-chloro-, and 2-hydroxy-, and its acetyl derivative (NIERENSTEIN), T., 613.
- Pentamethyl- Δ^2 cyclohexadien-4-ols** (v. AUWERS and ZIEGLER), A., i, 121.
- 2:2:3:3:5-Pentamethylindoline**, and its salts and derivatives (KNOEVENAGEL), A., i, 1061.
- Pentamminechromiselenates**. See under Chromium.
- Pentamminecobaltiselenates**. See under Cobalt.
- n-Pentane**, molecular scattering of light in (VENKATESWARAN), T., 2655.
- n-Pentane**, $\beta\gamma\delta$ -trioximino-. See Dimethyl triketone trioxime.
- isoPentane**, physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.
- cycloPentane**, effect of attached groups on ring formation in (CHANDRASENA and INGOLD), T., 1552.
- cycloPentane-1:1-acetic acid**, methyl ester (DICKENS, KON, and THORPE), T., 1502.
- cycloPentanespiro-2:3-dicyanocyclopropane-2-carboxylic acid**, and its amide (BIRCH and THORPE), T., 1834.
- cycloPentanespiro-2:3-dicyanocyclopropane-2:3-dicarboxylic acid**, imide (BIRCH and THORPE), T., 1834.
- Pentane- $\beta\delta$ -diol**, β -amino-, oxalate, and γ -chloro- γ -nitro-, and γ -nitro- (SCHMIDT and WILKENDORF), A., i, 314.
- cycloPentane-2:3-dione-1-carboxylic acid**, ethyl ester, diphenylhydrazone (DIECKMANN), A., i, 1022.
- ae-Pentanedialphenylidihydraside** (CLUTTERBUCK and COHEN), T., 128.
- cycloPentanespirocyclopentane-3:4-dione**, and its derivatives (KON), T., 526.
- cycloPentanespirocyclopentane-3:4-dione-2:5-dicarboxylic acid**, methyl ester, and its quinoxaline derivative (DICKENS, KON, and THORPE), T., 1502.
- cycloPentanespirocyclopropane-2:3-dicarboxylic acid**, *cis*- and *trans*-forms, and their dinitrile (BIRCH and THORPE), T., 1835.
- isoPentane- $\alpha\beta\delta\delta$ -tetracarboxylic acid**, and β -cyano-, ethyl ester (INGOLD and NICKOLLS), T., 1647.
- Pentaphenylethane**, chloro- (SCHLENK and MARK), A., i, 1003.

- Pentaphenylethyl**, (SCHLENK and MARK), A., i, 1002.
- Pentaphenylpyridine**, salts of (DILTHEY, NUSSLEIN, MEYER, and KAFFER), A., i, 949.
- Pentaphenylpyrylium** salts (DILTHEY and KAFFER), A., i, 668.
- Pentapyridinium** μ -chlorodecachlorodiferrate (WEINLAND and KISSLING), A., i, 864.
- Pentaquochromiselenates**. See under Chromium.
- cyclo*- Δ^2 - α -Penthiophen-5-carboxylic acid**, ethyl ester, and its derivatives (STEINKOPF and AUGESTAD-JENSEN), A., i, 851.
- Pentosans** (HEUSER), A., i, 113, 921.
- Pentoses**, fermentation of, by moulds (PETERSON, FRED, and SCHMIDT), A., i, 1220.
- Pentosuria**, *r*-arabinose in (WRZENIEWSKI), A., i, 1215.
- cyclo*Pentylethane- $\alpha\beta$ -dicarboxylic acid**, $\alpha\beta$ -dicyano-1-hydroxy-, and 1-hydroxy-, lactone (BIRCH and THORPE), T., 1836.
- cyclo*Pentylethane- $\alpha\alpha\beta$ -tricarboxylic acid**, 1-hydroxy-, lactone (BIRCH and THORPE), T., 1835.
- Pepper**, artificial and natural, constituents of (OTT, EICHLER, LÜDEMANN, and HEIMANN), A., i, 1026.
- Peppermint oil** (KREMER), A., i, 357, 848.
- Pepsin**, action of (HAMMARSTEN), A., i, 958.
action of, on diastase (BIEDERMANN), A., i, 480.
effect of chloroform on the activity of (ASTRUC and RENAUD), A., i, 281.
diffusion of, in gels (PEKELHARING), A., i, 388.
digestion of histone sulphate by (FELIX), A., i, 698.
estimation of (VALTEICH and GLOVER; GLASSNER), A., ii, 406; (TAKATA), A., ii, 886.
- Peptides**, estimation of, by alkalimetry (WILLSTÄTTER and WALDSCHMIDT-LEITZ), A., ii, 169.
- Peptisation** of metallic hydroxides (CHATTERJI and DHAR), A., ii, 205, 627.
- Peptone**, fermentation of dextrose by (SCHLATTER), A., i, 1096; (BAUR and HERZFELD), A., i, 1097.
- Peptones**, toxicity of (PENTIMALLI), A., i, 302.
- Perchlorates** and **Perchloric acid**. See under Chlorine.
- Perilla oil** (BAUER and HARDEGG), A., i, 983.
- Periodic system**, linear relationships in the (BILTZ), A., ii, 494.
of the ions of elements (GRIMM), A., ii, 635.
- Peristalsis**, production of, by choline (ARAI), A., i, 970.
- Permeability** of cells to acids (CROZIER), A., i, 897.
influence of optical activity on (KOTAKE and OKAGAWA), A., i, 696.
of plant cells (OSTERHOUT; BROOKS; KAHHO), A., i, 308.
- Permutite**, equilibrium of bases in (SCHULZE), A., ii, 486.
volumes of kations in (LORENZ), A., ii, 196.
- Pernitric acid**. See under Nitrogen.
- Peroxides** and **glyoximes**, structure of (BIGIAMI), A., i, 352.
- Peroxydase**, estimation of, in blood (BACH and ZUBKOWA), A., i, 392.
estimation of, in milk (RICE and HANZAWA), A., ii, 407.
- Persulphates**. See under Sulphur.
- Perylene**, preparation of (ZINKE), A., i, 132; (HANSGIRG), A., i, 245.
- Perylene**, 1:12-dihydroxy- (ZINKE), A., i, 132.
- Petrol**, inflammability of mixtures of alcohol and (BOUSSU), A., ii, 557.
- Petroleum**, determination of the iodine number of (HLADKÝ), A., ii, 722.
iodine number of, in the Bergius process (WATERMAN and PERQUIN), A., ii, 90.
ozonides of (KOETSCHAU), A., i, 977.
light, detection of benzene in (SCHWARZ), A., ii, 531.
estimation of, the vapour of, in air (FRITZMANN and MACULEVITSCH), A., ii, 877.
estimation of aromatic hydrocarbons in (WATERMAN and PERQUIN), A., ii, 399.
- Pharbitis nil chois***, constituents of the seeds of (ASAHINA and SHIMIDZU), A., i, 506.
- Pharmacological action**, relation between chemical constitution and (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 769.
- Phaseolus angularis*** (adzuki bean), proteins of (JONES, FINKS, and GERSDORFF), A., i, 504.
- Phaseolus lunatus*** (lima bean), proteins of (JONES, GERSDORFF, JOHNS, and FINKS), A., i, 1101.
- Phellandral** from eucalyptus oil (PENFOLD), T., 266.
- Phellodendron amurense***, constituents of (SHIMO), A., i, 99.
- Phenacetic acid**, detection of, colorimetrically (EKKERT), A., ii, 169.

α -Phenacylmethyl- α' -antipyrino-4-methylmethylaniline, and its hydrochloride (MANNICH and HEILNER), A., i, 373.

α s-Phenacylmethylmethylcarbamide (MANNICH and HEILNER), A., i, 372.

Phenanthranaphthazines, amino-, bromo-, hydroxy-, and nitro- (SIRCAR and DUTT), T., 1945.

Phenanthranaphthazine-12-sulphonic acid, 2-*mono*- and 2:7-*di*-amino-, and 2:7-*di*hydroxy- (SIRCAR and DUTT), T., 1950.

Phenanthraphenazines, 2-amino-, 2-*mono*- and 2:3:4-*tri*-hydroxy-, and 3:4-*di*nitro-2-hydroxy-, and their derivatives (SCHMIDT and SPOUN), A., i, 665.

Phenanthraquinol methyl ethers (GOLDSCHMIDT and SCHMIDT), A., i, 1149.

Phenanthraquinol, 3:4-*di*nitro-2-hydroxy-, diacetate (SCHMIDT and SPOUN), A., i, 666.

Phenanthraquinone, 2-amino-, acetyl derivative, and its diacetate (BRASS and FERBER), A., i, 356.

***di*bromohydroxy-**, 3:4-*di*nitro-2-hydroxy-, and 2:3:4-*tri*hydroxy-, and their derivatives (SCHMIDT and SPOUN), A., i, 666.

Phenanthraquinone colouring matters (SIRCAR and DUTT), T., 1944; (DUTT), T., 1951.

Phenanthrene, melting and boiling points (KIRBY), A., i, 27.

vapour pressure of (NELSON and SENSEMAN), A., i, 245.

solubility of, in organic solvents (HENSTOCK), T., 2124.

action of aluminium chloride with (SCHOLL and SCHWARZER), A., i, 331.

estimation of (WILLIAMS), A., ii, 90.

Phenanthrene, 2-amino-9- and -10-hydroxy-, and their derivatives (SCHMIDT and SPOUN), A., i, 666.

Phenanthrene series (SCHMIDT and SPOUN), A., i, 665.

Phenanthrene, dichloronitro-, derivatives (SCHMIDT and SPOUN), A., i, 666.

Phenanthroxazine, preparation of (FORESTI), A., i, 1062.

detection of nitrates by means of (FORESTI), A., ii, 524.

Phenanthroxyls (GOLDSCHMIDT and SCHMIDT), A., i, 1149.

Phenazarsine oxide (SCHMIDT), A., i, 285.

Phenazarsinic acid, and *di*nitro-, and its sodium salt (SCHMIDT), A., i, 285.

***o*-Phenetidine**, use of, in separation of aluminium and iron (CHALUPNY and BREISCH), A., ii, 538.

***p*-Phenetidine**, 2:5-*di*nitro-, and its acetyl derivative (REVERDIN and ROETHELISBERGER), A., i, 538.

2-*p*-Phenetidinobenzoic acid, 4-nitro- (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 469.

β -*p*-Phenetidine-*p*-phenetolecarbamide (SPECKAN), A., i, 580.

Phenetole, iodonitro-derivatives (APOSTOLO), A., i, 335.

***p*-Phenytylcarbamide** (*dulcin*), effect of substitution on the sweetness of, and β -bromo- (SPECKAN), A., i, 579.

derivatives of (HERMANN), A., i, 1151.

detection of, microchemically (DENIGES and TOURROU), A., ii, 95.

***p*-Phenytylcarbamine cyanide** (FROMM and WENZL), A., i, 438.

***p*-Phenytylsulphamic acid**, sodium salt (WEIL and MOSER), A., i, 444.

Phenotallene (ASCHAN, FONTELL, and SIMOLA), A., i, 1152.

Phenol, formation of, by bacteria (SIEKE), A., i, 902.

ultra-violet absorption spectrum of (KLINGSTEDT), A., ii, 332.

latent heat of fusion of (STRATTON and PARTINGTON), A., ii, 258.

equilibrium of water and (LEONE and ANGELESCU), A., ii, 743.

equilibrium of the freezing point of, in mixtures with water (RHODES and MARKLEY), A., i, 135.

equilibrium and properties of mixtures of cresols with (KENDALL and BEAVER), A., i, 136.

mechanism of the bromination of (BAINES), T., 2810.

catalysis of the hydrogenation of, with sodium carbonate (ARMSTRONG and HILDITCH), A., ii, 756.

ethers, action of nitric acid on (MEYER and GOTTLIEB-BILLROTH), A., i, 539.

detection of, colorimetrically (BELLUSSI), A., ii, 723.

estimation of, in mixtures of tar acids (HOFFERT), A., ii, 879.

Phenol, 2:4:5-*tri*bromo- (AKTIEN-GESELLSCHAFT FÜR ANILINFABRIKATION), A., i, 1145.

pentabromo-, silver salts, chromo-isomeric (LUCAS and KEMP), A., i, 30.

3:5-*di*nitro-, solubility and volatility of (SIDGWICK and TAYLOR), T., 1853.

Phenols, tautomerism of (FUCHS and STIX), A., i, 451.

equilibria of camphor with (KREMANN and ODELGA), A., i, 159.

- Phenols**, equilibria of triphenylmethane with (KREMANN, ODELGA, and ZAWODSKY), A., i, 131.
 equilibria of water and (LEONE and ANGELESOU), A., ii, 743.
 introduction of the chloroethyl group into (CLEMO and PERKIN), T., 642.
 condensation of, with hydrochlorides of cyanamides and carbodi-imides (SHORT and SMITH), T., 1803.
 interaction of diazonium salts and (CHATTAWAY and HILL), T., 2756.
 catalytic hydrogenation of (SENDERENS and ABOULENC), A., i, 136.
 action of nitrous acid on (SCHOUTISEN), A., i, 135.
 oxidation of (PUMMERER, MELAMED, and PUTTFARCKEN), A., i, 1161.
 and their derivatives, oxidation of (GOLDSCHMIDT), A., i, 1148; (GOLDSCHMIDT and SCHMIDT), A., i, 1149.
 and their ethers, oxonium salts from (KEHRMANN, DECKER, and SOLOMINA), A., i, 32; (KEHRMANN, DECKER, and SCHMAJEWSKI), A., i, 32.
 sulphonation of (CAMPBELL), T., 847.
 physiology of (PELKAN and WHIPPLE), A., i, 499; (PELKAN), A., ii, 399.
 monohydric, preparation of halogen substitution products of (AKTIEN-GESELLSCHAFT FÜR ANILINFABRIKATION), A., i, 1145.
 dihydric, ultra-violet absorption spectra of (KLINGSTEDT), A., ii, 680.
 detection of (MOIR), A., ii, 321.
 estimation of, in the blood (PELKAN), A., ii, 399.
 estimation and separation of (HANCHE and KOESSLER), A., ii, 322.
- Phenols**, amino-, preparation of (LEWCOCK, ADAM, SIDERFIN, and GALBRAITH), A., i, 650.
 chloro-, solubility of (SIDGWICK and TURNER), T., 2256.
 halogenated, nitration of (RAIFORD), A., i, 335.
 triiodo-, preparation of, from hydroxy-acids (KRISHNA and POPE), T., 798.
 nitro-, solubility and cooling curves of (CARRICK), A., i, 334.
 equilibria of, with triphenylcarbinol (KREMANN, HOHL, and MÜLLER), A., i, 138.
 metallic derivatives of (D. and A. E. GODDARD), T., 54; (GODDARD and WARD), T., 262.
 estimation of nitrogen in, by the Kjeldahl method (MARGOSCHES and VOGEL), A., ii, 522.
- Phenolcamphorsin**, and its derivatives (KRISHNA), T., 253; (SINGH, RAI, and LAL), T., 1425.
Phenolcamphorsin, *m*-amino- (SIRCAR and DUTT), T., 1286.
Phenolcarboxylic acids, bismuth salts, hydrolysis of, by water (PERLING), A., i, 252.
Phenol-4:6-disulphonic acid, *o*-amino-, potassium hydrogen salt (SAKELLARIOS), A., i, 1145.
Phenol-*o*- and -*p*-mercuri-salts (MAMELI), A., i, 695.
Phenolphthalein, influence of ethyl alcohol on the colour change of (WEGSCHEIDER), A., ii, 387.
Phenolphthalein, bromo- and chloro-derivatives (THIEL and MÜLLER), A., i, 659.
Phenolphthaleins, colour changes of (THIEL), A., i, 455.
Phenol-red, use of, as an indicator (MASSINK), A., ii, 307.
Phenolsulphamphthalein, and its derivatives, and 2:2'-imino- (DUTT), T., 2390.
Phenolsulphocarboxylic acids, nitro-, action of bromine on (SAKELLARIOS), A., i, 1144.
Phenol-4-sulphonic acid, 2-amino-, cobaltammine salt (MORGAN and MOSE), T., 2865.
 2-bromo-6-amino-, and 2-bromo-6-nitro-, potassium salt (SAKELLARIOS), A., i, 1145.
Phenolsulphonic acids, nitro-, action of bromine on (SAKELLARIOS), A., i, 1144.
Pheno- β -naphthacarbazole, 4-hydroxy- (BUCHERER and WAHL), A., i, 464.
Pheno- $\beta\beta$ -naphthacarbazole-1:3-disulphonic acid, 5-hydroxy- (BUCHERER and WAHL), A., i, 465.
Pheno- β -naphthacarbazole-1:7-disulphonic acid, sodium salt (BUCHERER and ZIMMERMANN), A., i, 465.
Pheno- $\beta\beta$ -naphthacarbazole-1-sulphonic acid, 5-hydroxy- (BUCHERER and WAHL), A., i, 464.
Pheno- α -naphthacarbazole-2-sulphonic acid, and 4-hydroxy-, sodium salts (BUCHERER and ZIMMERMANN), A., i, 465.
Pheno- β -naphthacarbazole-2-sulphonic acid, 4-hydroxy- (BUCHERER and WAHL), A., i, 465.
Phenoxides, nitro-, alkali (D. and A. E. GODDARD), T., 54.
 lead, magnesium and silver (GODDARD and WARD), T., 262.
Phenoxyacetic acid, *m*-amino-, and its acetyl derivative (MINTON and STEPHEN), T., 1597.

- Phenoxyacetic acids**, chloro-, and their derivatives (MINTON and STEPHEN), T., 1599.
 nitro-, preparation of, and their derivatives (MINTON and STEPHEN), T., 1591.
- Phenoxyacetone**, *p*-amino-, and *p*-nitro-, and their derivatives (SPECKAN), A., i, 580.
- o*-Phenoxybenzoic acid, 2:5-dichloro- (ECKERT and ENDLER), A., i, 941.
- p*-Phenoxybenzoic acid (SCHÖNBERG and KRAEMER), A., i, 664.
- 4-Phenoxybenzonitrile**, 3-nitro- (MATTAAR), A., i, 251.
- 4-Phenoxybenzophenone-4'-arsinic acid** (LEWIS and CHEETHAM), A., i, 187.
- 5-Phenoxy-5-*n*-butylbarbituric acid** (DOX and YODER), A., i, 808.
- Phenoxy-*n*-butylmalonic acid**, ethyl ester (DOX and YODER), A., i, 808.
- Phenoxychloroparaffin** (ELEKTROCHEMISCHE WERKE, BOSSHARD, and STRAUSS), A., i, 913.
- β-Phenoxydiethylaniline**, *p*-nitroso- (CLEMO and PERKIN), T., 645.
- 7-Phenoxy-7:12-dihydro-γ-benzophenarsazine** (LEWIS and HAMILTON), A., i, 188.
- β-Phenoxyethylaniline** (CLEMO and PERKIN), T., 645.
- Phenoxyethylphenylarsinous acid** (QUICK and ADAMS), A., i, 601.
- Phenoxy-malonic acid**, menthyl ester (SHIMOMURA and COHEN), T., 2054.
- 7-Phenoxy-2-methylxanthone-8-carboxylic acid**, and its sodium salt (v. DEM KNESEBECK and ULLMANN), A., i, 360.
- Phenoxyparaffin**, and its sulphonic acid (ELEKTROCHEMISCHE WERKE, BOSSHARD, and STRAUSS), A., i, 913.
- 3-Phenoxy-2-phenylbenzopyrylium ferrichloride**, and 7-hydroxy- (PRATT and ROBINSON), T., 1582.
- Phenyl allyl ether**, *p*-amino-, and its derivatives (GUTEKUNST and GRAY), A., i, 951.
- iso*amyl ethers, *p*-amino-, and *p*-nitro-, and their derivatives (GUTEKUNST and GRAY), A., i, 951.
- benzyl ether*, *p*-amino-, acetyl derivative (GUTEKUNST and GRAY), A., i, 951.
- butyl ethers*, *p*-amino- and *p*-nitro-, and their derivatives (GUTEKUNST and GRAY), A., i, 950.
- β*-chloroethyl ether, *o*- and *p*-amino-, acetyl derivatives (CLEMO and PERKIN), T., 645.
- dithiochloride*, *o*-nitro- (LECHER and SIMON), A., i, 1013.
- Phenyl ethyl and β-naphthyl disulphides** (LECHER and WITTEBER), A., i, 642.
- mercaptan*, *m*-hydroxy- (WATSON and DUTT), T., 2415.
- mercurichloride*, 2:4:6-trinitro- (KHARASCH), A., i, 190.
- Phenylacetaldehyde**, preparation of (KODAMA), A., i, 349.
 fate of, in the body (KAY and RAFFER), A., i, 1093.
- Phenylacetic acid**, and *o*-nitro-, sodium salt, behaviour of, in the animal organism (SOHEMPF), A., i, 403.
 sodium uranyl salt (BARLOT and BRENET), A., ii, 167.
 chlorosulphonyl derivative, preparation of (STEWART), T., 2560.
- Phenylacetic acid**, *o*-amino-, and its salts and derivatives (NEBER), A., i, 545.
 2:4-dinitro-, mercuric salt (KHARASCH), A., i, 190.
- Phenylacetic-*o*-sulphinic anhydride** (NEBER), A., i, 546.
- Phenylacetoneitrile**, action of, with magnesium methyl bromide (RONDOU), A., i, 934.
- Phenylacetoneitriles**, dialkylated (BLONDEAU), A., i, 654.
- Phenyl-*o*-acetoxystyrylsulphone** and *p*-chloro- (TROGER and BOLTE), A., i, 267.
- Phenylacetyl-*dl*-alanine** (SHIPLE and SHERWIN), A., i, 1093.
- Phenylacetylene**, preparation of (HESSELER), A., i, 442.
 hydrogenation of (ZALKIND), A., i, 1134.
- Phenylacetyl-*dl*-leucine** (SHIPLE and SHERWIN), A., i, 1093.
- 5-Phenylacridine**, 2':4'-chloronitro-, 1-mono-, and 1:8-di-nitro- (MAYER and FREUND), A., i, 866.
- β-Phenylacrylic acid**, α-cyano-, preparation of (LAPWORTH and McRAE), T., 1700.
- Phenylalanine**, behaviour of, in the animal organism (KOTAKE, MASAI, and MORI), A., i, 1217.
 hydrochloride and anilide (CURTIUS and SIEBER), A., i, 722.
- dl*-Phenylalaninecholine**, salts of (KARBER, GISLER, HORLACHER, LOCHER, MÄDER, and THOMANN), A., i, 815.
- α-Phenylallyl alcohol**, and its benzoate (RUPE and MÜLLER), A., i, 40.
- 3-Phenyl-1- and -3-allylindazoles**, and their picrates (v. AUWERS and HÜLTENES), A., i, 683.

- Phenylaminoanthraquinone**, *diamino-* (BADISCHE ANILIN- & SODA-FABRIK), A., i, 942.
- N*-Phenyl- β -aminoethyl, and *p*-chloro-, and *m*-nitro-, hydrogen sulphates (SAUNDERS), T., 2671.
- Phenylamino-2- and -4-hydroxy- α -naphthylacetoneitriles**, and nitro- (MORGAN and REEVES), T., 4.
- Phenyl-1-amino- β -naphthylamine**, preparation of (DUTT), T., 1952.
- 1-Phenyl-4-*p*-aminophenylbenzene**, and its diazonium sulphate (PUMMERER, BINAPFL, BITTNER, and SCHUEGRAF), A., i, 1187.
- 3-Phenyl-5-anilinomethylene-4-glyoxal-one**, 2-thiol-, and 2-thiol-3-*p*-chloro- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- γ -Phenyl- α -anisyl- β -methylpropan- β -ol (LE BRAZIDEC), A., i, 457.
- γ -Phenyl- α -anisyl- β -methyl- Δ^{α} -propene (LE BRAZIDEC), A., i, 457.
- N*-Phenylanthranilic acid, and *p*-chloro-, formalides and nitriles from (FRIEDLÄNDER and KUNZ), A., i, 765.
- Phenylarsenious acid**, *p*-nitro- (BART), A., i, 1202.
- Phenylarsine**, *o*- and *p*-chloro- (PALMER and ADAMS), A., i, 786.
- Phenylarsinic acid**, *p*-amino-, sodium salt. See Atoxyl.
- o*-chloro- (PALMER and ADAMS), A., i, 786.
- chloro-, hydroxy-, and nitro-derivatives (BART), A., i, 1201.
- p*-hydroxy-*m*-amino-, sodium salt (FOURNIER, GUÉNOT, and SCHWARTZ), A., i, 300.
- Phenylarsinoacetanilide**, and *p*-amino-, and its derivatives (QUICK and ADAMS), A., i, 601.
- Phenylarsinoacetic acid** (QUICK and ADAMS), A., i, 601.
- Phenylarsinoacetarsanilic acid**, and *p*-amino-, and its acetyl derivative (QUICK and ADAMS), A., i, 601.
- Phenylarsinoacetophenetidine**, and *p*-amino-, and its acetyl derivative (QUICK and ADAMS), A., i, 601.
- Phenylarsino-*o*- and -*p*-acetylaminobenzoic acids** (QUICK and ADAMS), A., i, 601.
- Phenylaziminobenzene**. See 2-Phenyl-1:2:3-benzotriazole.
- 2-Phenyl- ψ -aziminobenzene**. See 2-Phenyl- ψ -1:2:3-benzotriazole.
- Phenylazobis-*p*-dimethylaminotriphenylmethane** (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazobis-*p*'-dimethylaminotriphenylmethane**, *p*-chloro- (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazocarbonamide**, *p*-mono- and 2:4-*di*-bromo (PIERONI), A., i, 1072.
- Phenylazotri-*p*-aminophenylmethane** (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazotri-*p*-anisylmethane** (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazotri-*p*-dimethylaminophenylmethane** (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazotri-*p*'-dimethylaminophenylmethane**, *p*-chloro- (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazotri-*p*-tolylmethane**, *p*-chloro- (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylazoxycarbonamide**, and *p*-bromo-, and *p*-nitro- (PIERONI), A., i, 1072.
- 3-Phenylbarbituric acid**, 4-imino- (LIFSCHITZ and HEPNER), A., i, 768.
- N*-Phenylbenziminom-*m*-hydroxyphenyl ether, and its hydrochloride (CHAPMAN), T., 1676.
- 5-Phenyl-2:1-benzoacridine** (MAYER and FREUND), A., i, 866.
- 2-Phenyl-1:3-benzodioxan** (ADAMS, FOGELER, and KREGER), A., i, 681.
- 2-Phenylbenzoselenazole**, 6-*mono*- and 4:6-*di*-amino-, 6-*mono*-, and 4:6-*di*-nitro-, and their derivatives (BOGERT and CHEN), A., i, 1183.
- 2-Phenylbenzoselenazole-azo- β -naphthol** (BOGERT and CHEN), A., i, 1183.
- 2-Phenylbenzoxazole**, 2-*o*-hydroxy-, and its salts (SKRAUP and MOSER), A., i, 575.
- 2 *p*-nitro- (SKRAUP and MOSER), A., i, 575.
- Phenyl-*o*-benzoyloxystyrylsulphone** and *p*-chloro- (TROGER and BOLTE), A., i, 287.
- Phenyl- β -benzoyl- α -phenylethylphosphinic acid**, and its anhydride (CONANT and POLLACK), A., i, 67.
- cis*- and *trans*- β -Phenyl- β -(α -benzoyl)-phenylvinylhydroxylamine, and their derivatives (RUPE and WITTEW), A., i, 448.
- Phenyl- β -benzoyl- α -styrylethylphosphinic acid** (CONANT, BUMP, and HOLT), A., i, 68.
- 2-Phenylbenzthiazole**, synthesis of, and 6-amino-, 6-bromo-, and 6-cyano-, and their derivatives (BOGERT and ABRAHAMSON), A., i, 576.
- 2-Phenylbenzthiazole-6-carboxylic acid** (BOGERT and ABRAHAMSON), A., i, 576.
- 2-Phenyl-1:2:3-benzotriazole**, 4'-chloro-4-amino-, acetyl derivative, and 4'-chloro-4-nitro- (BLUM), A., i, 28.
- 2-Phenyl- ψ -1:2:3-benzotriazole**, 5-amino- (KALLÉ & Co.), A., i, 61.

- β*-Phenyl-*β*-benzylamine**, and its hydrochloride (BLONDEAU), A., i, 655.
- N*-Phenyl-*N*-benzyl-*β*-aminoethyl sodium sulphate** (SAUNDERS), T., 2674.
- β*-Phenyl-*β*-benzylbutanol**, and its derivatives (BLONDEAU), A., i, 655.
- α*-Phenyl-*α*-benzylbutyric acid**, and its amide, ester, and nitrile (BLONDEAU), A., i, 654.
- β*-Phenyl-*α*-benzylethylphosphinic acid**, *α*-hydroxy- (CONANT, MACDONALD, and KINNEY), A., i, 186.
- Phenyl *α*-benzyl-*α*-ethylpropyl ketone** (HALLER and BAUER), A., i, 259.
- 1-Phenyl-4-benzylidenehydantoin**, 1-*p*-chloro-, and 2-thio-1-*p*-chloro- (HILL and KELSEY), A., i, 1141.
- Phenylbenzylmethylarsine**, and its salts (STEINKOPF, DONAT, and JAEGER), A., i, 996.
- N*-Phenyl-*O*-benzyl-*N*-methylhydroxylamine** (MEISENHEIMER and WILLMERSDORF), A., i, 334.
- Phenyl *α*-benzylpropyl ketone**, and its oxime (HALLER and BAUER), A., i, 259.
- 6-Phenyl-4-benzylpyridine-3-carboxylic acid**, 2-hydroxy-6-*p*-chloro-, ethyl ester (KOHLER and SMITH), A., i, 458.
- 5-Phenyl-3-benzylpyrone**, *p*-chloro- (KOHLER and SMITH), A., i, 458.
- 5-Phenyl-3-benzylpyrone-2-carboxylic acid**, methyl ester, monochloro-derivatives (KOHLER and SMITH), A., i, 458.
- Phenylbiuret-*o*-sulphonic acid**, *p*-bromo-, *p*-chloro-, and *p*-iodo-, and their salts and derivatives (SCOTT and COHEN), T., 2045.
- Phenylboric acid** (KRAUSE and NITSCHKE), A., i, 694.
- Phenylbromoarsineacetanilide** (QUICK and ADAMS), A., i, 601.
- Phenylbromoarsineacetic acid** (QUICK and ADAMS), A., i, 601.
- Phenyl-*β*-bromo-*β*-benzoyl-*α*-phenyl-ethylphosphinic acids** (CONANT and POLLACK), A., i, 67.
- Phenyl*δ*-bromobismuthine**, *p*-chloro- (CHALLENGER and RIDGWAY), T., 110.
- α*-Phenyl-*β*-(*α'*-bromocinnamoyl)-ethylphosphinic acid** (CONANT, BUMP, and HOLT), A., i, 67.
- Phenyl bromomethyl ketone**, 2:4-dihydroxy-, and its derivatives (SONN and FALKENHEIM), A., i, 1163.
- Phenyl bromonitro-*p*-chlorophenylpropyl ketones** (KOHLER and SMITH), A., i, 458.
- Phenyl *α*-*γ*-dibromo-*β*-nitro-*γ*-*p*-chlorophenylpropyl ketone** (KOHLER and SMITH), A., i, 458.
- Phenyl *α*-bromo-*γ*-nitro-*β*-phenylpropyl ketone**, *p*-chloro- (KOHLER and SMITH), A., i, 457.
- 3-Phenyl-5-*p*-bromophenylaminomethylene-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- α*-Phenyl-*β*-(*α'*-*δ*-dibromo-*β'*-phenylpropionyl)-ethylphosphinic acid** (CONANT, BUMP, and HOLT), A., i, 67.
- β*-Phenyl-bromo-*β*-phenylvinylhydroxylamine** (RUPE and WITTWER), A., i, 449.
- Phenylisobutylsuccinic acid** (UPSON and THOMPSON), A., i, 343.
- γ*-Phenylbutyric acid**, *α*-amino-, excretion of, as its *N*-acetyl derivative (KNOOP), A., i, 486.
- γ*-2:4-dihydroxy-** (LANGLEY and ADAMS), A., i, 1154.
- Phenylcamphenol**, and its anhydride (NORDSTROM), A., i, 944.
- Phenylcarbamide**, *p*-hydroxy-, preparation of ethers of (RIEDEL), A., i, 579.
- Phenylcarbamide-2:4-disulphonic acid**, potassium salt (SCOTT and COHEN), T., 2043.
- Phenylcarbamide-*o*-sulphonic acid**, *p*-bromo-, and its salts, and acetyl derivative, potassium salt (SCOTT and COHEN), T., 2047.
- Phenylcarbamine cyanide** (FROMM and WENZL), A., i, 438.
- Phenylcarbasole**, preparation of (ECKERT, SEIDEL, and ENDLER), A., i, 952.
- 3-Phenyl-1-*o*-carboxybenzylphthalazone** (RUGGLI and MEYER), A., i, 345.
- α*-Phenyl-*β*-carboxyethylphosphinic acid** (CONANT, BUMP, and HOLT), A., i, 67.
- Phenylchloroarsineacetic acid** (QUICK and ADAMS), A., i, 601.
- Phenyl *p*-chlorophenylacetyl methyl ketone**, and its copper derivative (KOHLER and SMITH), A., i, 458.
- 1-Phenyl-4-*p*-chlorophenylfuran** (KOHLER and SMITH), A., i, 458.
- Phenyl*δ*-dichlorostibine**, *p*-amino-, acetyl derivative, and *m*-amino-*p*-hydroxy-, and their hydrochlorides (SCHMIDT), A., i, 1203.
- Phenyl*tetra*-chlorostibine**, *p*-amino-, hydrochloride (SCHMIDT), A., i, 1204.
- α*-Phenylcinchoninic acid**. See 2-Phenylquinoline-4-carboxylic acid.
- cis*- and *trans*-*α*-Phenyl-*o*-cesumarephenones**, and their derivatives (DECKER and BECKER), A., i, 358.
- γ*-Phenylcrotonic acid**, *α*-cyano-, amide of (STEVENSON and THORPE), T., 1720.

- Phenylisocrotylcarbinol** (KRETSINSKI), A., i, 1180.
- Phenylcyanourethane** (SCOTT and COHEN), T., 2050.
- Phenyl-3:4-diacetoxystyrylsulphone**, and *p*-chloro- (TROGER and BOLTE), A., i, 268.
- Phenyl-di-*p*-anisylisooxazoles** (MEISENHEIMER and WEIBEZAHN), A., i, 177.
- γ -Phenyl- α -di-*p*-anisylpropan- α -one** (ZIEGLER and OCHS), A., i, 1049.
- Phenyl-2:4-dibenzoyloxystyrylsulphone**, and *p*-chloro- (TROGER and BOLTE), A., i, 268.
- 1-Phenyl-2:3-dibenzyl-1,4-naphthoquinoxalium chloride**, *aa'*-dicyano- (DUTT and SEN), T., 2666.
- Phenyl *aa*-dibenzylpropyl ketone** (HALLER and BAUER), A., i, 259.
- 1-Phenyl-2:3-dibenzylquinoxalium chloride**, *aa'*-dicyano- (DUTT and SEN), T., 2666.
- 4-Phenyl-2:6-di-(*p*-bromophenyl)pyridine**, and its salts (DILTHEY, NUSSLEIN, MEYER, and KAFFER), A., i, 949.
- Phenyl di-*p*-chloro- β -phenylstyryl ketone** (MEYER and SCHUSTER), A., i, 556.
- Phenyl-diethylphenylphosphazide** (STAUDINGER and HAUSER), A., i, 69.
- Phenyl-diethylphosphinebenzoylimine** (STAUDINGER and HAUSER), A., i, 70.
- Phenyl-diethylphosphinephenylimine** (STAUDINGER and HAUSER), A., i, 69.
- β -Phenyl- $\alpha\beta$ -diethylpropiofenone** (RAMART and ALBESCO), A., i, 663.
- 2-Phenyl-dihydro-1:2:4-isonaphthatriazine-3-carboxylic acid**, 2-*p*-nitro-ethyl ester (FIERZ and SALLMANN), A., i, 870.
- 9-Phenyl-9:10-dihydro-*meso*thioanthracene**, and 2-chloro- (BISTRZYCKI and BRENNEN), A., i, 268.
- 4-Phenyl-4:5-dihydro-1:3:4-thiadiazole**, 2-thiol-5-thio-, potassium salt, compound of chloropicrin and (RAY and DAS), T., 325.
- 2-Phenyl-1:3-dihydro-1:2:3-triazole-4-carboxylic acid**, and its bromo-derivative (GASTALDI and BRAUNIZER), A., i, 626.
- Phenyl-di- α -hydroxybenzylarsine**, *o*- and *p*-chloro- (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxy-*o*-carbomethoxybenzylarsine** (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxy-*p*-chlorobenzylarsine** (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxyethylarsine**, and its platinichloride, and *p*-chloro- (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxy-*n*-heptylarsine** (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxy-*p*-methoxybenzylarsine** (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxy-*n*-propylarsine**, and its platinichloride (PALMER and ADAMS), A., i, 786.
- Phenyl-di- α -hydroxyisovalerylarsine**, and its platinichloride (PALMER and ADAMS), A., i, 786.
- 5-Phenyl-2:7-dimethylacridine**, 9-nitro- (MAYER and FREUND), A., i, 867.
- α -Phenyl- γ -dimethylaminopropan- α -ol**, and its derivatives (MANNICH and HEILNER), A., i, 351.
- 4-Phenyl-2-*p*-dimethylaminostyrylthiazole methiodide** (MILLS and SMITH), T., 2735.
- Phenyl-dimethylarsine dihydroxide**, and its salts (STEINKOPF and SCHWEN), A., i, 72.
- 3-Phenyl-1:2-dimethylindazolium iodide** (v. AUWERS and HULTENES), A., i, 683.
- 2-Phenyl-5:5-dimethylloxazoline**, and its salts (DERSIN), A., i, 142.
- 1-Phenyl-2:3-dimethyl-5-pyrazolone**, 4-amino- and 4-chloroamino-, acetyl derivative, and their derivatives (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 954.
- Phenyl-di- α -naphthylbromomethane**, and its metallic derivatives (SCHOEFFLE), A., i, 338.
- Phenyl-di- α -naphthylcarbinol**, preparation and derivatives of (SCHOEFFLE), A., i, 337.
- Phenyl-di- α -naphthylchloromethane**, and its metallic derivatives (SCHOEFFLE), A., i, 338.
- Phenyl-di- α -naphthylmethyl** (SCHOEFFLE), A., i, 338.
- Phenyl-diphenylethynylcarbinol**, preparation and reactions of (HESS and WELTZEN), A., i, 35.
- 4-Phenyl-dithiourazole**, and 3-imino-, and their derivatives (ARNDT, MILDE, and TSCHENSCHER), A., i, 377.
- 2-Phenyl-4:6-di-*p*-tolylpyridine**, and 2-*p*-chloro- (GASTALDI), A., i, 368.
- 2-Phenyl-4:6-di-*p*-tolylpyrylium salts**, and 2-*p*-chloro- (GASTALDI), A., i, 368.
- o*-Phenyleneammethyl chloride** (PELLIZZARI), A., i, 585.

- o*-Phenylenediacetic acid, diethyl ester, preparation of (PERKIN and TITLEY), T., 1565.
- m*-Phenylenediacetic acid, 4:6-dinitro-, and its lead salt (DAVIES and HICKOX), T., 2651.
- m*-Phenylenediallyldiamine, 2:4-dinitro- (GIUA and ANGELETTI), A., i, 649.
- o*-Phenylenediamine, 3:6-dichloro- (MACLEOD, PFUND, and KILPATRICK), A., i, 1181.
- p*-Phenylenediamine, compound of tellurium tetrabromide with (LOWY and DUNBROOK), A., i, 446.
- Phenylenediamines, equilibria of, with dihydroxynaphthalenes (KREMANN, HEMMELMAYR, and RIEMER), A., i, 1011.
- equilibria of triphenylcarbinol with (KREMANN, HOHL, and MÜLLER), A., i, 138.
- hydroferrocyanides (CUMMING), T., 1295.
- m*-Phenylenediaminecamphorcin (SIRCAR and DUTT), T., 1286.
- p*-Phenylenedimethyldiamine hydroferrocyanide (CUMMING), T., 1296.
- m*-Phenylenedisulphamic acid, sodium salt (WEIL and WASSERMANN), A., i, 1007.
- o*-Phenylenemelanuric acid (PELLIZARI), A., i, 586.
- β*-Phenylethane, α -chloro-*β*-hydroxy- (DETGEUF), A., i, 327.
- 1-Phenyl-6-ethoxybenzimidazole, 1-*p*-bromo- (JACOBSEN), A., i, 591.
- 1-Phenyl-6-ethoxybenztriazole, 1-*p*-bromo- (JACOBSEN), A., i, 591.
- 2-Phenyl-3-*p*-ethoxyphenyloxazolidine (BERGMANN, ULPTS, and CAMACHO), A., i, 1182.
- Phenylethylacetylaminooacetic acid, and its nitrile (READ), A., i, 955.
- N*-Phenyl-*N*-ethyl-*O*-allylhydroxylamine (MEISENHEIMER and GRESKE), A., i, 334.
- α -Phenylethylamine hydroferrocyanide (CUMMING), T., 1294.
- N*-Phenyl-*N*-ethyl- β -aminoethyl, and *p*-nitroso-, hydrogen sulphates (SAUNDERS), T., 2678.
- Phenylethylbarbituric acid, action of diazomethane on (HERZIG), A., i, 374.
- calcium and magnesium salts (WÜLFING), A., i, 1066.
- β*-Phenyl- β -ethylbutanol, and its derivatives (BLONDEAU), A., i, 655.
- β*-Phenyl- β -ethylbutylamine, and its hydrochloride (BLONDEAU), A., i, 655.
- Phenylethylcarbamine cyanide (FROMM and WENZL), A., i, 438.
- Phenylethylcyanosarin (STEINKOFF, DONAT, and JAEGER), A., i, 996.
- 3-Phenyl- γ -ethylhexane (RAMART and ALBESCO), A., i, 663.
- 4-Phenyl-4-ethylhydantoin. See Nirvanol.
- β -Phenylethyl β -hydroxyvinyl ketone, and its salts (RUPE and MÜLLER), A., i, 41.
- 3-Phenyl-1- and -2-ethylindazoles, and their picrates (v. AUWERS and HÜLTENES), A., i, 682.
- 1-Phenyl-5-ethylpyridazine-6-one-5-carboxylic acid, and its ethyl ester (CARRIÈRE), A., i, 319.
- 1-Phenyl-4-ethylpyrrolidine-5-one-3-carboxylic acid, and its ethyl ester (CARRIÈRE), A., i, 319.
- Phenylethylsemicarbazide (NEIGHBORS, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.
- δ - α -Phenylethylsemicarbazides, and their hydrochlorides (WILSON, HOPPER, and CRAWFORD), T., 869.
- Phenylethylsuccinic acid (UPSON and THOMPSON), A., i, 343.
- Phenylfurylethylene oxides, *p*-nitro- (KLEUCKER), A., i, 735.
- Phenylfuryl-2-keto-3-methylcyclohexylmethane (WOLFF), A., i, 668.
- γ -Phenylglutaconic acid, α -cyano-, ethyl ester (INGOLD, PERREN, and THORPE), T., 1782.
- β -Phenylglyceric acids, isomerism of phenylglycidic acid and (BOESEKEN and DE GRAAFF), A., i, 551.
- α -Phenylglycerol, and its triacetyl derivative (MOUREU and GALLAGHER), A., i, 34.
- Phenylglycidic acid, *cis-p*-nitro- (KLEUCKER), A., i, 735.
- β -Phenylglycidic acid, isomerism of β -phenylglyceric acids and (BOESEKEN and DE GRAAFF), A., i, 551.
- Phenylglycine, preparation of compounds of (BRITISH DYESTUFFS CORPORATION, LTD., LEVINSTEIN, and IMBERT), A., i, 252.
- Phenylglycine anhydride, preparation and reactions of (FUCHS), A., i, 1152.
- p*-Phenylglycinearsinic acid, and its sodium salt (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- p*-Phenylglycylamidoarsinic acid, and its sodium salt (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- Phenylglycyl-*n'*-aminophenol-*p*-arsenoxide (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 962.

- p*-Phenylglycyl-*m*'-aminophenolarsinic acid (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- p*-Phenylglycylanilinoarsinic acid (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- Phenylglycylanthrannilic acid *p*-arsenoxide (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 962.
- p*-Phenylglycylcarbamidoarsinic acid (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- p*-Phenylglycylmethylcarbamidoarsinic acid (ROCKEFELLER INSTITUTE FOR MEDICAL RESEARCH), A., i, 961.
- 3-Phenyl-4-glyoxalone, 2-thiol-3-*p*-chloro- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- α -Phenyl- β -glyoxyethylphosphinic acid (CONANT, BUMP, and HOLT), A., i, 67.
- Phenylglyoxylic acid, 2:3:4-*tri*hydroxy-, and its salts and derivatives (FINGER and EIRICH), A., i, 254.
- Phenyl hexadecyl ketone, 3:4-*di*hydroxy- (MAJIMA and CHIBA), A., i, 265.
- 1-Phenylhydantoin, 1-*p*-chloro-, and 2-thio-1-*p*-chloro- (HILL and KELSEY), A., i, 1141.
- Phenylhydrazine, equilibrium of acetone or acetaldehyde with (HAMMERSTEN), A., ii, 830.
action of cyanogen haloids on (PELLIZARI), A., i, 585.
- Phenylhydrazine, 5-bromo-2:4-*di*nitro-, and its acetyl derivative (GIUA), A., i, 691.
p-nitro-, preparation of (DAVIES), T., 715.
- 1-Phenylhydrazine-3-carboxylic acid, 5-chloro-2-hydroxy- (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 385.
- Phenylhydrazinedisulphonic acid, *p*-nitro-, ammonium salt (DAVIES), T., 720.
- 9-Phenylhydrazinocridine (FARRWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 469.
- Phenylhydrazinoethanesulphonylphenylhydrazide hydrochloride (CLUTTERBUCK and COHEN), T., 127.
- 5-Phenylhydrazino-1-phenyl-1:2:3:4-tetrazole (STOLLE), A., i, 689.
- Phenylhydrazobis-*p*-dimethylaminotriphenylmethane (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylhydrazobis-*p*'-dimethylaminotriphenylmethane, *p*-chloro- (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylhydrazones, *o*- and *p*-nitro-, quinonoid salts of (CIUSA and RASTELLI), A., i, 1073.
- Phenylhydrazothiodicarbonylamine *S*-methyl ether (ARNDT, MILDE, and TSCHENSCHER), A., i, 376.
- Phenylhydrazotri-*p*-aminophenylmethane (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylhydrazotri-*p*'-aminophenylmethane, *p*-chloro- (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylhydrazotri-*p*-anisylmethane (WIELAND, POPPER and SEEFRIED), A., i, 773.
- Phenylhydrazotri-*p*-dimethylaminophenylmethane (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- Phenylhydrazotri-*p*'-dimethylaminophenylmethane, *p*-chloro- (WIELAND, POPPER, and SEEFRIED), A., i, 773.
- 1-Phenyl-6-hydroxybenzimidazole, 1-*p*-bromo- (JACOBSEN), A., i, 591.
- N*-Phenyl-*N*'-hydroxyisobutylthiocarbonylamine (DERSIN), A., i, 142.
- Phenyl 2-hydroxy- β -ethoxystyryl ketone (PRATT and ROBINSON), T., 1580.
- Phenyl-*mono*- and -*di*-hydroxyethyl carbamides, *p*-hydroxy- (KIEDEL), A., i, 579.
- Phenylhydroxylamine, condensation of, with hydroxymethylene compounds and carbinols (RUPE and WITTEWIT), A., i, 448, 449.
- β -Phenyl- α -hydroxymethylene-ethyl methyl ketone, and its salts and derivatives (RUPE and MÜLLER), A., i, 41.
- Phenyl 2-hydroxy- β -phenoxystyryl ketone (PRATT and ROBINSON), T., 1581.
- 3-Phenyl-2-*p*-hydroxyphenyl-1:4-*di*-phenyl-1:3-*di*methindiazidine, 3-*p*-bromo-, and 3-*m*-nitro- (INGOLD and PIGGOTT), T., 2798.
- α -Phenyl- β -hydroxypropylamine, α -*p*-hydroxy-, and its salts (TAKEDA and KURODA), A., i, 273.
- Phenyl-*o*-hydroxystyrylsulphone, and *p*-chloro- (TROGER and BOLTE), A., i, 267.
- Phenyl-2:4-*di*hydroxystyrylsulphone, and *p*-chloro-, and their dimethyl ethers (TROGER and BOLTE), A., i, 268.
- Phenyl β -hydroxyvinyl ketone, reduction of (RUPE and MÜLLER), A., i, 40.
- Phenylimineketendicarboxylic acid, ethyl ester (STAUDINGER and HAUSER), A., i, 29.
- 5-Phenylimino-2-thio-3-naphthyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole (GUHA), A., i, 877.
- 5-Phenylimino-2-thio-3-phenyl-2:3:4:5-tetrahydro-1:3:4-thiodiazole (GUHA), A., i, 876.

- 5-Phenylimino-2-thio-2:3:4:5-tetrahydro-1:3:4-thiadiazole (GUHA), A., i, 876.
- 5-Phenylimino-2-thio-3-tolyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole (GUHA), A., i, 877.
- 2-Phenylindazole, 3-hydroxy- (HELLER), A., i, 1066.
- 3-Phenylindazole-2-carboxylic acid, methyl ester (v. AUWERS and HÜLTENES), A., i, 683.
- 3-Phenylindazole-2-sulphonic acid, metallic salts (v. AUWERS and HÜLTENES), A., i, 683.
- 3-Phenylindazole-2-sulphonoxide (v. AUWERS and HÜLTENES), A., i, 684.
- N-Phenylindirubin (FRIEDLANDER and KUNZ), A., i, 766.
- 1-Phenylindoxyl, and its sodium salt, and 4-chloro- (FRIEDLANDER and KUNZ), A., i, 765.
- Phenylindoxyllic acid, ethyl and methyl esters (FRIEDLÄNDER and KUNZ), A., i, 765.
- 1-Phenylindoxyl-2-thionaphthenones (FRIEDLANDER and KUNZ), A., i, 766.
- Phenyl iodomethyl ketone, 2:4-dihydroxy-, and its diacetate (SONN and FALKENHEIM), A., i, 1163.
- Phenylisatin, 4-chloro- (FRIEDLANDER and KUNZ), A., i, 766.
- 1-Phenylisatin-2-p-dimethylaminoanil (FRIEDLANDER and KUNZ), A., i, 765.
- Phenylisatinic acid, 4-chloro-, sodium salt (FRIEDLÄNDER and KUNZ), A., i, 766.
- Phenylketenacetal (STAUDINGER and RATHSAM), A., i, 1015.
- Phenyl-lactic acid, behaviour of, in the animal organism (KOTAKE and MORI; MORI), A., i, 1216.
- Phenyl-lactic acid, hydroxy-, excretion of, after administration of tyrosine (KOTAKE and OKAGAWA), A., i, 1217.
- d- and l-Phenyl-lactic acids, and hydroxy-, production of acetoacetic acid from, in the liver (MORI), A., i, 1217.
- Phenylmalonamide (DOX and YODER), A., i, 817.
- 1-Phenyl-2-mercapto-6-ethoxybenzimidazole, 1-p-bromo- (JACOBSEN), A., i, 591.
- Phenyl-p-methoxyphenylpropanes, $\beta\gamma$ -dibromo- (INGOLD and PIGGOTT), T., 2386.
- Phenyl-p-methoxyphenyl- Δ^2 -propenes (INGOLD and PIGGOTT), T., 2386.
- 5-Phenyl-3-methylacridine, 1-amino- and 1-nitro- (MAYER and FREUND), A., i, 866.
- 7:9-diamino- and 7:9-dinitro- (MAYER and FREUND), A., i, 867.
- Phenylmethylaminoethanol, dihydroxy-, hydrochloride, preparation of (NAGAI), A., i, 652.
- N-Phenyl-N-methyl- β -aminoethyl hydrogen sulphate (DUTT and SEN), T., 2674.
- Phenylmethylarsine, cyano- (STEINKOPF and SCHWEN), A., i, 72.
- 4-Phenyl-2-methylbenzoic acid, calcium salt (v. AUWERS and JÜLICHER), A., i, 842.
- 1-Phenyl-6-methylbenzotriazole, 1-p-hydroxy- (JACOBSEN), A., i, 592.
- β -Phenyl- β -methylbutanol, and its derivatives (BLONDEAU), A., i, 655.
- β -Phenyl- β -methylbutylamine, and its hydrochloride (BLONDEAU), A., i, 655.
- Phenyl α -methylbutylketone (LEROIDE), A., i, 216.
- α -Phenyl- α -methylbutyric acid, and its amide, esters, and nitrile (BLONDEAU), A., i, 654.
- Phenyl 3':4'-methylenedioxytyreryl ketone dibromide (BAUER and WERNER), A., i, 1035.
- Phenylmethylthylarsine, and its salts (STEINKOPF, DONAT, and JAEGER), A., i, 995.
- 1-Phenyl-3-methyl-2-ethyl-5-pyrazolone, 4-amino- and 4-chloro-amino-, acetyl derivative, (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 954.
- β -Phenyl- α methylglutaconic acids, isomeric, and their esters and ozonides (FEIST, BREUER, and LUBRICHT), A., i, 553.
- Phenylmethylglyoxime, nickel compound (PONZIO), A., i, 18.
- α -Phenyl- ζ -methyl- Δ - α -heptadien- ζ -ol-one (SCHEIBLER and FISCHER), A., i, 1110.
- β -Phenylmethylheptenol (HELFERICH and GEHRKE), A., i, 9.
- as-Phenylmethylhydrazine, p-nitro-, and its derivatives (CIUSA and RASTELLI), A., i, 1073.
- 4-Phenylmethylhydrazinobenzonitrile, 3-nitro- (MATTAAR), A., i, 252.
- α -Phenylmethyl- β -hydroxypropylamine, α -p-hydroxy-, and its hydrochloride (TAKEDA and KURODA), A., i, 273.
- 3-Phenyl-1- and -2-methylindazoles, and their picrates (v. AUWERS and HÜLTENES), A., i, 683.
- Phenyl-2'-methyl-3'-indolydene-methane, 3:4:5-trihydroxy-, and its derivatives (ROSENKUND and FRANK-KUCH), A., i, 1031.
- 9-Phenyl-3-methylnaphthacridine, 2-amino- (BROWNING, COHEN, GAUNT, and GULBRANSEN), A., i, 612.

- 5 Phenyl-3-methylisoxazole-4-carboxylic acids, derivatives of (BETTI and BERLINGOZZI), A., i, 52.
- α -Phenyl-5-methyl- Δ^2 -penten-5-ol-one, and its derivatives (SCHEIBLER and FISCHER), A., i, 1109.
- Phenylmethyl- α -propylarsine, and its salts (STEINKOFF, DONAT, and JAEGER), A., i, 998.
- Phenyl α -methyl- α -propylbutyl ketone (LEROIDE), A., i, 216.
- Phenyl-1-methylpyrazole, chloro-4-bromo- (ROJAHN), A., i, 1184.
- 3-Phenyl-1-methylpyrazoline, and its hydrochloride (MANNICH and HEILNER), A., i, 372.
- 1-Phenyl-3-methyl-5-pyrazolone, 1-5'-nitro-2'-hydroxy- (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 385.
- Phenyl-1-methylpyrazolones (ROJAHN), A., i, 1184.
- 1-Phenyl 3-methyl-5-pyrazolone-3'-carboxylic acid, 1-5'-chloro-2'-hydroxy-, and 1-4'-hydroxy- (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 385.
- N-Phenyl-2-methylpyridazonexanthone, 5- and 7-chloro- (V. DEM KNESEBECK and ULLMANN), A., i, 360.
- 2-Phenyl-6-methylquinoline-4-carboxylic acid, allyl ester (ROSENMUND), A., i, 52.
- Phenylmethylsuccinic acid (UPSON and THOMPSON), A., i, 343.
- Phenylmethylthiazole methiodides (MILLS and SMITH), T., 2735.
- 4-Phenyl-3-methyl-2-thiazolonyl-4-quinolylmethane methiodide (MILLS and SMITH), T., 2736.
- Phenyl methyl triketone- α - and - β -trioximes, and their nickel salts and triacetyl derivatives (PONZIO), A., i, 1038.
- 2-Phenyl- α -8-naphthatriazole, 2- p -amino- (CHARRIER and VIOLA), A., i, 771.
- 6:2- and 9:2-diamino-, and 9:2-dinitro- (MORGAN and GILMOUR), A., i, 380.
- Phenyl- α -naphthylamine, β -hydroxy- (WAHL and LANTZ), A., i, 823.
- 3-Phenyl-5-naphthylaminomethylene-4-glyoxalone, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- Phenyl α -naphthyl ketones, chloro- (SCHOLL, SEER, and DAIMER), A., i, 258.
- 2-Phenyl- α - and - β -naphthylmethylbenzoic acids, and their salts (McMULLEN), A., i, 1025.
- Phenyl- α - and - β -naphthylphthalides (McMULLEN), A., i, 1025.
- Phenyl- β -naphthylsulphone, 2:4-dinitro- (GRANDMOUGIN), A., i, 331.
- Phenyl γ -nitro- β - p -chlorophenylpropyl ketone (KOHLE and SMITH), A., i, 458.
- Phenyl β -nitro- γ -hydroxy- γ - p -chlorophenylpropyl ketone (KOHLE and SMITH), A., i, 458.
- cis*- and *trans*- α -Phenyl- δ - p -nitrophenyl- Δ^2 -butadiene γ - α -chlorohydrius and oxides (KLEUCKER), A., i, 734, 735.
- 3-Phenyl-2- p -nitrophenyl-1:4-diphenyl-1:3-dimethindiazidine, 3- o -nitro- (INGOLD and PIGGOTT), T., 2799.
- Phenyl γ -nitro- p -phenylpropyl ketone, p -chloro- (KOHLE and SMITH), A., i, 457.
- o -Phenylnitrosoaminophenyl- p -tolylsulphone. See o - p -Toluenesulphonyldiphenylnitrosoamine.
- 1-Phenyloxindole, 3-*di*chloro- (STOLLÉ), A., i, 762.
- 5-Phenylcyclopentane-2:3-dione-1-carboxylic acid, ethyl ester, diphenylhydrazone (DIECKMANN), A., i, 1022.
- 2-Phenylperithiazine (REISSERT), A., i, 584.
- Phenyl phenacetyl methyl ketone, p -chloro-, and its copper derivative (KOHLE and SMITH), A., i, 458.
- 10-Phenylphenazonium hydroxide, 2-amino- δ -thiol- (WATSON and DUTT), T., 1941.
- Phenyl- α -phenyl- β -(α' -bromocinnamoyl)-ethylphosphinic acid (CONANT, BUMP, and HOLT), A., i, 68.
- Phenyl- α -phenyl- β -($\alpha'\beta'$ -*di*bromophenylpropionyl)-ethylphosphinic acid (CONANT, BUMP, and HOLT), A., i, 68.
- Phenyl- α -phenyl- β -carboxyethylphosphinic acid (CONANT, BUMP, and HOLT), A., i, 68.
- Phenyl- α -phenyl- β -cinnamoyl ethylphosphinic acid (CONANT, BUMP, and HOLT), A., i, 68.
- γ -Phenyl- α -(β -phenylethyl)propylphosphinic acid, α -hydroxy- (CONANT, MACDONALD, and KINNET), A., i, 186.
- β -Phenyl- β -phenylvinylhydroxylamine (RUPE and WITTEBER), A., i, 449.
- Phenylphosphinic acid, nitroamino- (NIJK), A., i, 961.
- 3-Phenyl-1-phthalidophthalazone (RUGGLI and MEYER), A., i, 343.
- 1-Phenylpiperidine, 4-hydroxy-, 4-hydroxy-1- p -amino-, and their derivatives (BORSCHKE and BONACKER), A., i, 51.
- α -Phenylpropane, α -*di*hydroxy-, and its dibenzoate (RUPE and MÜLLER), A., i, 40.

- γ -Phenylpropan- α - and - β -ols, β - and α -iodo-** (PORCHER), A., i, 539.
- l - β -Phenylpropionamide, l - β -hydroxy-** (MCKENZIE and SMITH), T., 1359.
- Phenylpropionic acid, cis - β (?)chloro- p -nitro-** (KLEUCKER), A., i, 735.
- β -Phenylpropionic acid, α -thiol-** (GRÄNACHER), A., i, 850.
- β -Phenylpropionolactone, β -2:4- di - and β -2:4:6- tri -hydroxy-, and β -5-nitroso-2:4- di hydroxy-** (LANGLEY and ADAMS), A., i, 1153.
- Phenylpropionylphenylmethylhydrazine** (TOMIČEK), A., i, 679.
- Phenylpropionyltolylhydrazines** (TOMIČEK), A., i, 679.
- γ -Phenylpropyl alcohol, β -amino-, and its hydrochloride** (KARRER), A., i, 991.
- Phenyl α -propylbutyl ketone** (LEROIIDE), A., i, 216.
- Phenyl- n -propyleycarsine** (STEINKOPF, DONAT, and JAEGER), A., i, 996.
- α -Phenylisopropylethyl methyl ketone, α -semicarbazono of** (RUPE and MÜLLER), A., i, 41.
- α -Phenylisopropyl β -hydroxyvinyl ketone, and its salts** (RUPE and MÜLLER), A., i, 41.
- Phenyl- n -propylsuccinic acid** (UPSON and THOMPSON), A., i, 343.
- 5-Phenyl-isopropyltriazole, 3-hydroxy-** (NEIGHBORS, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.
- α -Phenylpropan- β -dicarboxylic acid, ethyl ester** (DIELS and AUBART), A., i, 1194.
- Phenylpyridinium bromide, 2:4:5- tri -nitro-** (GIUA), A., i, 691.
- 1-Phenyl-4-pyridone** (SMIRNOV), A., i, 464.
preparation and derivatives of (BORSCHÉ and BONACKER), A., i, 50.
- 2-Phenylpyrimidine, and 5-bromo-6-hydroxy-, 6-chloro-5-bromo-, and 6-hydroxy-, and their salts** (CHERBULIEZ and STAVRITCH), A., i, 581.
- 2-Phenylpyrimidine-4-carboxylic acid, 5-bromo-6-hydroxy-, and 6-hydroxy-, and their derivatives** (CHERBULIEZ and STAVRITCH), A., i, 581.
- 1-Phenylpyrrolidine-5-one-3-carboxylic acid, ethyl ester** (CARRIERE), A., i, 318.
- Phenylpyruvic acid, behaviour of, in the animal organism** (KOTAKE and MORI), A., i, 1216.
and hydroxy-, asymmetrical reduction of, in organs (MORI and KANAI), A., i, 1217.
- Phenylpyruvic acid, ethyl ester, isomerism of** (GAULT and WEICK), A., i, 1024.
- Phenylquinoline, 3-hydroxy-2- o -nitro-** (NEBER), A., i, 545.
- 2-Phenylquinoline-4-carboxylic chloride** (ROSENMUND), A., i, 52.
- 2-Phenylquinoline-4-carboxylic acid (*atophan*), esters of** (ROSENMUND), A., i, 52.
benzyl ester (GAMS and KAISER), A., i, 51.
- 2-Phenylquinoline-4-carboxylic acid, 5-amino-6-hydroxy-, and 6-hydroxy-** (HALBERKANN), A., i, 174.
6-hydroxy- (ZUCKMAYER), A., i, 952.
- 3-Phenylquinoline-4-carboxylic acid, 2-hydroxy-** (FARBWERKE VORM. MEISTER, LUCIUS, & BRÜNING), A., i, 867.
- 2-Phenylquinoline-4-formylacetic acid, ethyl ester** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolylaminoethanol dihydrochloride** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolyl bromomethyl ketone, and its hydrobromide** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolyl diethylaminomethyl ketone, and its hydrobromide** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolyl dimethylaminoethanol dihydrochloride** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolyl dimethylaminomethyl ketone, and its hydrochloride** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolyl methyl ketone, and its hydrobromide** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- 2-Phenyl-4-quinolyl piperidinomethyl ketone, and its hydrochloride** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1180.
- N -Phenylrhodanine, α -oximino-, and its derivatives** (GRÄNACHER, REIS, and POOL), A., i, 577.
- N -Phenylrhodanine- α -acenaphthenequinone, and its d -nitro-derivative** (GRÄNACHER, REIS, and POOL), A., i, 577.
- Phenylstibinic acid, p -amino-, and p -hydroxy-** (SCHMIDT), A., i, 1204.
 p -nitro- (CHARRIER), A., i, 1080.
- Phenylstibinic acids, neutralisation of, with lithium hydroxide** (SCHMIDT), A., i, 482.

- Phenylstibinous oxide**, *p*-amino-, acetyl derivative (SCHMIDT), A., i, 1208.
- Phenyl styryl ketone** (*benzylidenacetophenone*), and its derivatives, absorption spectra of (SHIBATA and NAGAI), A., ii, 413.
- Phenyl styryl ketone**, α -bromo-, action of alcohols on (DUFRAISSE and GÉRALD), A., i, 843.
- 4-Phenyl-2-styrylthiazole** (MILLS and SMITH), T., 2784.
- 9-Phenyl-9-styrylxanthene** (ZIEGLER and OCHS), A., i, 1048.
- Phenylsuccinic acid**, preparation of (LAPWORTH and McRAE), T., 1704.
- Phenylsuccinyleosin** (LAPWORTH and McRAE), T., 2723.
- Phenylsulphamic acid**, and *o*- and *p*-chloro-, sodium salts (WEIL and MOSER), A., i, 444.
- Phenylsulphamic acid**, *p*-amino-, and its acetyl and benzoyl derivatives, and *m*-nitro-, and their salts (WEIL and WASSERMANN), A., i, 1006.
- Phenylsulphamphthalein**, 4:4'-diamino-2:2'-imino- (DUTT), T., 2392.
- 2-Phenyltetrahydroquinoline-4-carboxylic acid**, and its lithium salt (ZUCKMAYER), A., i, 574.
- 2-Phenyltetrahydroquinoline-4-carboxylic acid**, 7-amino-, acetyl derivative, and its potassium salt, and 6-hydroxy- (ZUCKMAYER), A., i, 952.
- 4-Phenyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole**, 5-imino-2-thio- (GUHA), A., i, 876.
- 1-Phenyl-1:2:3:4-tetrazole**, 5-amino-, preparation and derivatives of (STOLLÉ), A., i, 689.
- 4-Phenylthiazole-2-aldehyde** *p*-dimethylaminoanil methiodide (MILLS and SMITH), T., 2735.
- 4-Phenyl-2-thiazolylphthalonylmethane** (MILLS and SMITH), T., 2734.
- Phenylthiocarbamide**, *p*-chloro-, compounds of, with chloropicrin and mercuric nitrite (RAY and DAS), T., 326.
- Phenyl- η - and - ψ -thiocarbaminecyanides**, salts of (FROMM and WENZEL), A., i, 438.
- Phenylthiolacetyl-2:4-dimethylpyrroles**, *o*-nitro- (FISCHER and HERRMANN), A., i, 1055.
- Phenylthioldimethylpyrrole-3-carboxylic acids**, *o*-nitro-, ethyl esters (FISCHER and HERRMANN), A., i, 1055.
- 4-Phenylthiol-2:3:5-trimethylpyrrole**, 4-*o*-nitro- (FISCHER and HERRMANN), A., i, 1055.
- Phenylthioxanthyl** (GOMBERG and MINNIS), A., i, 163.
- r*- and *l*-**Phenyl-*p*-tolylacetamides** (McKENZIE and SMITH), T., 1357.
- Phenyl-*m*- and -*p*-tolylamines**, 5-bromo-2:4-dinitro- (GIUA and ANGELETTI), A., i, 649.
- 1-Phenyl-3-*p*-tolyl-5-anilinomethylene-4-glyoxalone**, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- N*-Phenyl-*N*-*p*-tolylhydroxylamine** (WIELAND and KÖGL), A., i, 727.
- 4-Phenyl-1-*p*-tolyl-3-methylpyrazole** (ROJAHN), A., i, 373.
- Phenyl-*p*-tolylnitrogen oxide** (WIELAND and KÖGL), A., i, 727.
- Phenyl-*p*-tolylsulphone**, *o'*- and *p'*-amino- (HALBERKANN), A., i, 1133.
- Phenyl-1-*p*-tolylsulphone**, 2:5-diamino-, derivatives of (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 579.
- β -Phenyl- $\alpha\beta$ -triethylpropionic acid** (RAMART and ALBESCO), A., i, 663.
- β -Phenyl- $\alpha\beta$ -triethylpropiofenone** (RAMART and ALBESCO), A., i, 663.
- Phenyltrimethylarsonium iodide**, compound of iodoform and (STEINKOPF and SCHWEN), A., i, 118.
- Phenyltrimethylphenylenediamine**, and its derivatives (FICHTER and ROTHENBERGER), A., i, 447.
- β -Phenyl-8:1:3:4 (or -1:3:5)-triphenyl- Δ^2 -5 (or -4)-pyrrolinylhydroxylamine**, and its picrate and chloro-derivative (RUPE and WITTWER), A., i, 449.
- 4-Phenylurazole**, and its derivatives and 3-imino- (ARNDT, MILDE, and TSCHENSCHER), A., i, 376.
- γ -Phenylvaleraldehyde**, γ -hydroxy- (HELFERICH and GEHRKE), A., i, 9.
- α -Phenylvinylphosphinic acid** (CONANT, MACDONALD, and KINNEY), A., i, 186.
- 3-Phenylvioluric acid**, 4-imino- (LIFSCHITZ and HEPNER), A., i, 768.
- Phloroglucinol**, phytochemical synthesis of (NIERENSTEIN), A., i, 136.
- condensation of formic acid with (SCHWENK), A., i, 153.
- Phloroglucinolcamphorsin** (SIRCAR and DUTT), T., 1285; (SINGH, RAI, and LAL), T., 1428.
- Phloroglucinolsulphamphthalein** (DUTT), T., 2391.
- Phosphatases** (TOMITA), A., i, 960.
- Phosphatase-phosphatase** (BARENDRECHT), A., i, 67.
- Phosphatides**, solubility of (STUCKERT), A., i, 1089.
- biochemistry of (BRINKMAN and WASTL), A., i, 289.
- of blood (BRINKMAN), A., i, 1086.
- action of, on coagulation of blood (ZUNZ and LA BARRE), A., i, 697.

Phosphasines, preparation and reactions of (STAUDINGER and LÜSCHER), A., i, 237.

reactions of (BRAUNHOLTZ), T., 300.

action of carbonylene derivatives with (STAUDINGER and BRAUNHOLTZ), A., i, 70.

Phosphineimines (STAUDINGER and HAUSER), A., i, 68.

Phospholipin, physico-chemical properties of (KAKIUCHI), A., i, 711.

Phosphonium iodide, crystal structure of (DICKINSON), A., ii, 640.

Phosphorescence of mixtures of boric acid with organic compounds (TIEDE, WULF, and RAGOSA), A., ii, 8.

Phosphorescent sulphides. See Phosphors and Sulphides.

Phosphors of the alkaline earths, photoelectric effect of (GÖGGER), A., ii, 732.

rare earth magnesium sulphide (TIEDE and SCHLEDE), A., ii, 769.

organic, of boric acid (TOMASCHEK), A., ii, 763.

Phosphorus, allotropy of (MARCKWALD and HELMHOLTZ), A., ii, 845.

absorption of active hydrogen and nitrogen by (WENDT; NEWMAN), A., ii, 639.

combination of copper with (EDWARDS and MURPHY), A., ii, 645.

Phosphorus compounds, molecular susceptibility of (PASCAL), A., ii, 285.

biochemistry of (ROGOZINSKI), A., i, 1226.

estimation of, in blood plasma (MYERS and SHEVKY), A., ii, 455.

Phosphorus carbide (DE MAHLER), A., i, 101.

chlorides, action of, on chlorodimethyldihydroresorcinol (HINKEL and WILLIAMS), T., 2498.

haloids, additive compounds of (CONANT), A., i, 41; (CONANT and POLLACK; CONANT, BUMP, and HOLT), A., i, 67; (CONANT, MACDONALD, and KINNEY), A., i, 186.

trihydride, action of formaldehyde on (HOFFMAN), A., i, 8.

pentoxide (*phosphoric oxide*), purification of (FINCH and PETO), T., 692.

treated with ozone, use of, as a drying agent (MANLEY), T., 331.

Phosphoric acid in soils (FRAPS), A., i, 616.

metabolism of. See Metabolism.

detection of, by means of benzidine (FEIGL), A., ii, 865.

estimation of (CLARK and KEELER), A., ii, 84.

Phosphorus :—

Phosphoric acid, estimation of, iodometrically (SVANBERG, SJÖBERG, and ZIMMERLUND), A., ii, 867.

estimation of, and its salts, volumetrically (KOLTHOFF), A., ii, 314; (MOERK; MOERK and HUGHES), A., ii, 866.

estimation of, in blood (BLOOR), A., ii, 84.

separation of, in analysis (BALAREFF), A., ii, 525; (TARUGI), A., ii, 784.

Phosphates, colloidal, formation of (ELSCHNER), A., ii, 714.

inorganic, in blood (LEHMAN), A., i, 191.

mineral, solubility of, in citric acid (TOCHER), A., ii, 525.

effect of, on plant growth (v. WRANGELL), A., i, 1098.

excretion of, in urine (FISKE), A., i, 88; (BOCK and IVERSEN), A., i, 1090.

adsorption of, by stannic sulphide in estimations (KIKUCHI), A., ii, 525.

estimation of, by the Bell-Doisy method (BRIGGS), A., ii, 718.

estimation of, in blood plasma (DENIS and VON MEYSENBUG), A., ii, 584.

Pyrophosphoric acid, structure of (BALAREFF), A., ii, 49.

Pyrophosphates, estimation of, volumetrically (MOERK), A., ii, 866.

Hypophosphorous acid, studies on (MITCHELL), T., 1624.

Phosphorus organic compounds

(BRAUNHOLTZ), T., 300; (CONANT), A., i, 41; (CONANT and POLLACK;

CONANT, BUMP, and HOLT), A., i, 67; (STAUDINGER and HAUSER),

A., i, 68; (STAUDINGER and BRAUNHOLTZ), A., i, 70; (CONANT, MAC-

DONALD, and KINNEY), A., i, 186; (STAUDINGER and LUSCHER), A., i,

237; (NIJK), A., i, 960.

Phosphoric acid, glucose esters of (LEVENE, MEYER, and WEBER), A., i, 987.

Phosphorus estimation :—

estimation of, colorimetrically (LOSANA), A., ii, 392.

estimation of, in iron (MURMANN), A., ii, 226.

estimation of, in minerals and in coke ash, colorimetrically (MISSON), A., ii, 718.

estimation of, in organic compounds (MANCINI), A., ii, 83; (SVANBERG, SJÖBERG, and ZIMMERLUND), A., ii, 867.

Phosphorus estimation :—

inorganic, estimation of, in serum (TISDALL), A., ii, 392.

Phosphosulphatide from the lungs (SAMMARTINO), A., i, 1089.

Photocatalysis (BALY, HEILBRON, and HUDSON), T., 1078.

Photochemical action in relation to fluorescence (WEIGERT), A., ii, 681. equivalent law (PRINGSHEIM), A., ii, 602.

verification of, with photographic dry plates (EGGERT and NODDACK), A., ii, 9.

processes, conversion of energy in (BYK), A., ii, 415.

velocity of reaction in relation to concentration in (VOLMER and RIGGERT), A., ii, 336.

studies (PLOTNIKOW), A., i, 419; ii, 248.

Photochemistry, laws of (LASAREV), A., ii, 103, 332.

and fluorescence (WOOD), A., ii, 334.

Photo-electric effects, colloids as indicators of (PESKOV), A., ii, 828.

Photographic copying process, mechanism of (WEIGERT and SCHÖLLER), A., ii, 10.

emulsions and films, electrical conductivity of (RABINOVICH), A., ii, 605.

plates, dry, verification of the photochemical equivalent law with (EGGERT and NODDACK), A., ii, 9.

Photography, new sensitiser for (MILLS and POPE), T., 946.

of crystals (FRANÇOIS and LORMAND), A., ii, 127.

Photo-haloids, effect of an electric current on (TUBANDT and ESCHENHAGEN), A., ii, 346.

Photosynthesis (BALY), A., i, 307; (KOSTYTSHEV), A., i, 307, 308, 613.

and the electron theory (DIXON and BALL), A., ii, 248.

Phototropy and photoelectric effects (GALLAGHER), A., ii, 11.

of metallic salts (SINGH), T., 782.

of sulphides, effect of heat on (MOURELO), A., ii, 604.

Phrenosine, behaviour of, in the body (SHIMIZU), A., i, 91.

Phthalaldehydic acids, preparation of (DAVIES, PERKIN, and CLAYTON), T., 2214.

Phthaleins, colour changes of (THIEL), A., i, 453.

Phthalic acid, salts, potentials of, against the hydrogen electrode (WOOD and MURDICK), A., ii, 735.

Phthalic acid, tetrachlorides, constitution and tautomerism of (OTT), A., i, 832.

amaryl esters, and their silver and strychnine salts (PAOLINI), A., i, 755.

*iso*Phthalic acid, 2:4:6-trinitro- (GIUA), A., i, 534.

Phthalic anhydride, preparation of, from naphthalene (CONOVER and GIBBS), A., i, 454.

Friedel and Crafts' reaction with (McMULLEN), A., i, 1025.

substituted derivatives, condensation of toluene and aluminium chloride with (LAWRANCE), A., i, 340.

Phthalic anhydride, tetrabromo-, di- and tetra-chloro-, additive compounds of (PFEIFFER and FLATER), A., i, 341.

Phthalimide, preparation of (BRITISH DYESUFFS CORPORATION, LTD., GREEN and GREEN), A., i, 836.

potassium and sodium derivatives, preparation of (HAMMICK and LOCKET), T., 2362.

Phthalimide, tetrachloro-, compound of α -naphthyl methyl ether and (PFEIFFER and FLATER), A., i, 342.

Phthalimidoacetaldehyde, and its derivatives (RADDE), A., i, 1155.

Phthalimidoacetoneitrile (SONN and FALKENHEIM), A., i, 1164.

α -Phthalimidoethyl- γ -diethoxyacetoacetic acid, ethyl ester (PUCHER and JOHNSON), A., i, 550.

β -Phthalimidoethylmalonic acid, and its ammonium salt and methyl ester (RADDE), A., i, 1155.

α -Phthalimidopropaldehyde, and its semicarbazone (RADDE), A., i, 1155.

γ -Phthalimido-*n*-propylmalonic acid (RADDE), A., i, 1155.

α -Phthalimino- β -hydroxy- β -methylpropane (DERSIN), A., i, 142.

α -Phthalimino- β -methylpropane, β -bromo-, β -chloro-, and β -iodo- (DERSIN), A., i, 142.

Phthalyl chlorides, isomeric (v. BRAUN and KAISER), A., i, 658.

Phthalyl- α -alanylamide, and its derivatives (RADDE), A., i, 1155.

1:2-Phthalylanthraquinone and its azine (SCHOLL and NEUMANN), A., i, 262.

Phthalyl-*NN'*-dipteridide (v. BRAUN and KAISER), A., i, 659.

Phthalylglycylamide (RADDE), A., i, 1155.

Phthalylglycylanilide (RADDE), A., i, 1154.

Phthalylglycylonitrile, and its derivatives (RADDE), A., i, 1155.

- Phthalylphenylcarbamide** (ELLIOTT), T., 206.
- Physicist**, rôle of the, in development of chemical theory (WALKER), T., 735.
- Physiological fluids**, equilibrated sterilisable (FLEISCH), A., i, 964.
- precipitation of proteins from (GRIGAUT and ZIZINE), A., ii, 886.
- Phytin** in foods (ARBENZ), A., i, 1102.
- Phytosterols**, chemistry of (VESTERBERG; VESTERBERG and WESTERLIND), A., i, 825.
- iodine value of (HOLDE, WERNER, TACKE, and WILKE), A., ii, 723.
- Pichurin bean oil**, constituents of (ROURE-BERTRAND FILS), A., i, 846.
- β -Picolinium** ruthenibromides (GUTHRIE and KRAUSS), A., i, 16.
- Picolinoylaminoanthraquinones** (BARNETT), A., i, 943.
- Picric acid**, cobaltamines of (MORGAN and KING), T., 1725.
- hexamminechromic and hexamminecobaltic salts (EPHRAIM and MOSMANN), A., ii, 575.
- Picrocrocin**, and its derivatives (WINTERSTEIN and TELECZKY), A., i, 563.
- Picrocellin**, constitution of (FORSTER and SAVILLE), T., 816.
- Picrotin ketone**, and α -bromo-, and oximino-, and their derivatives (HORMANN and BISCHOF), A., i, 161.
- Picrotoxin** (HORMANN and BISCHOF), A., i, 161.
- Pigments**, metabolism of. See Metabolism.
- anthocyanin, formation of (COMBES), A., i, 412.
- related to anthocyanins, tinctorial properties of (EVEREST and HALL), A., i, 265.
- Pike**, constituents and toxicity of the roes of the (MCCRUDDEN), A., i, 194.
- Pills**, estimation of mercury in (FRANÇOIS), A., ii, 87.
- Pilocarpine**, action of, on blood (BORNSTEIN and VOGEL), A., i, 80.
- d*-Pimaric acid**, preparation of (KNECHT and HIBBERT), A., i, 1020.
- Pinelic acid**, *pentahydroxy*-, brucine and quinine salts (KILIANI), A., i, 1112.
- Pinablotic acid**, nitrosate, nitrosite and nitroschloride of, and oximino- (ASCHAN, FONTELL, and SIMOLA), A., i, 1152.
- Pinacolin transformation**, reversed (NYBERGH), A., i, 918.
- mobility of ethyl groups in the (NYBERGH), A., i, 802.
- Pinacolylthiocarbamide**, compound of chloropicrin and (RÂY and DAS), T., 326.
- Pinacone**, equilibrium in the system, water and (PUSHIN and GLAGOLEVA), T., 2816.
- Pinacyanols**, constitution of (KÖNIG), A., i, 1188.
- Pine**, Swedish, constituents and properties of (WAHLBERG), A., i, 1101.
- Pine leaves**, constituents of (v. EULER), A., i, 100.
- Pine lignin**. See Lignin.
- Pine needles**, wax from (KAUFFMANN and FRIEDEBACH), A., i, 740.
- Pine wood**, cellulose content of (KLASON), A., i, 99.
- Pinene hydrochloride**, preparation of borneol from (VAVON and BERTON), A., i, 943.
- Pinonic acid**, ethyl esters, semicarbazones (SIMONSEN), T., 2297.
- Pinus sylvestris*, constituents of the pollen of (KIESEL), A., i, 799.
- iso*Piperic acid** (OTT, EICHLER, and HEIMANN), A., i, 1027.
- Piperidine**, compounds of, with metallic salts (PETERS), A., i, 48.
- action of formaldehyde on (PUTOCHIN), A., i, 1176.
- haloid compounds of (LEONARD), A., i, 362.
- Piperidine**, 1-chloro-, preparation of (HIRST and MACBETH), T., 2173.
- Piperidinepropionic acid**, 4-hydroxy-, ethyl ester (RUZICKA and SEIDEL), A., i, 1057.
- 9 Piperidino-2-ethoxyacridine**, and its hydrochloride (FARBWERKE VORM. MEISTER, LUCIUS, & BRUNING), A., i, 469.
- Piperidinomethylenemalonie acid**, ethyl ester (INGOLD, PERREN, and THORPE), T., 1785.
- 4-Piperidone**, polymerisation of (RUZICKA and SEIDEL), A., i, 1057.
- 1- α -Piperidylethan-1-ol**, and its derivatives (HESS and CORLEIS), A., i, 170.
- N*-Piperidyl-*s*-phthalylidithiourethane** (v. BRAUN and KAISER), A., i, 659.
- β -4-Piperidylpropionic acid**, ethyl ester, benzoyl derivative (RABE, KINDLER, and WAGNER), A., i, 361.
- Piperil**, 6:6'-dinitro- (GREENE and ROBINSON), T., 2193.
- Piperitone** (READ and SMITH), T., 574, 1863; (READ, SMITH, and BENTIVOLIO), T., 582.
- constitution and oxidation of, and its semicarbazones (PENFOLD), A., i, 259, 1035.
- dl*-Piperitone**, and its semicarbazones (READ and SMITH), T., 1863.

- dl*-Piperitoneoximes, and their derivatives (READ, SMITH, and BENTIVOGLIO), T., 582.
- Piperolin, 6'-*mono*- and 6:6'-*di*-nitro- (GREENE and ROBINSON), T., 2192.
- Piperonaldehyde, preparation of, from isosafrole (NAGAI), A., i, 839.
- Piperonyl chloride, *dichloro*- (HAMBURGER), A., i, 556.
- Piperonylacetic acid, 6-nitro-, methyl ester (GREENE and ROBINSON), T., 2195.
- β*-Piperonylacrylic acid, α -cyano-, preparation of (LAPWORTH and McRAE), T., 1701.
- N*-Piperonylaminoacetic acid, α -cyano-, ethyl ester, hydrochloride (SCHEIBLER and BAUMGARTEN), A., i, 656.
- N*-Piperonylglycine hydrochloride (SCHEIBLER and BAUMGARTEN), A., i, 656.
- Piperonylideneaminoacetic acid, sodium salt (SCHEIBLER and BAUMGARTEN), A., i, 656.
- 3-Piperonylideneamino-2-methoxy-*ar*-tetrahydronaphthalene (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAM), A., i, 128.
- Piperonylideneanthranilic acids, nitro- (EKELEY, ROGERS, and SWISHER), A., i, 935.
- Piperonylideneisatinazine (BORSCH and MEYER), A., i, 54.
- Piperonylideneethyl α -hydroxyisopropyl ketone. See α -3:4-Methylene-dioxiphenyl- β -methyl- Δ^{α} -penten- β -ol- γ -one.
- Piperonylideneresorcinol, and its acetyl derivative (FABRE), A., i, 1148.
- 2-Piperonylquinoline-4-carboxylic acid, esters (ROSENEMUND), A., i, 52.
- Piperonylsuccinic acid, preparation of (LAPWORTH and McRAE), T., 1707.
- Piperylmethylglyoxime, nickel compound (BIGIARI), A., i, 353.
- Pipettes, testing and graduation of (STOTT), A., ii, 388.
- standard dropping (HALPHEN), A., ii, 221.
- Pituitary body, tethelin from the (DRUMMOND and CANNAN), A., i, 491.
- activity of extracts of (DALE and DUDLEY), A., i, 397.
- Pivalic acid (*aa*-dimethylpropionic acid), dibromo-, and dihydroxy-, and their methyl esters (KOH and MENDELWITSCH), A., i, 518.
- Plants, chemistry of (NOWAK and ZELLNER), A., i, 615; (SPRINGER), A., i, 797; (ZELLNER), A., i, 799, 1098.
- Plants, acidity of juices of, and their growth and feeding power (BAUER and HAAS), A., i, 975.
- growth and nutrition of, in relation to hydrogen-ion concentration (ARRHENIUS), A., i, 1097.
- growth of the organs of, in absence of air (MAQUENNE and DEMOUSSY), A., i, 707.
- biological significance of alkaloids in (CIAMICIAN and RAVENNA), A., i, 797.
- absorption of aluminium by the roots of (STOKLASA, ŠEBOR, TÝMICH, and UWACHA), A., i, 502.
- effect of aluminium compounds on the growth of (CONNER and SEARS), A., i, 613.
- genesis of amylase and maltase in (PALLADIN and POPOV), A., i, 614.
- distribution of anthocyanidins in (JONESCO), A., i, 797.
- effect of arsenic compounds on the growth of (STEWART), A., i, 1221; (STEWART and SMITH), A., i, 1222.
- assimilation in (BALY), A., i, 307; (KOSTYTSHEV), A., i, 307, 308; (MOLISCH; MÜLLER and WARBURG), A., i, 411; (WARBURG and NEGELEIN), A., i, 1097.
- formation of chlorophyll in (MANSKY), A., i, 1222.
- action of coal gas on (WEHMER), A., i, 211.
- corrosion of marble by the roots of (CHEMIN), A., i, 94.
- colloids in (SAMEC and FERJANČIČ), A., i, 115; (SAMEC and MAYER), A., i, 921.
- effect of hippuric acid and uric acid on the nutrition of (BOKORNY), A., i, 1222.
- hydrocyanic acid in (SIEGFRIED; ROSENTHALER), A., i, 614.
- incrusted tissues of (SCHMIDT and DUYSEN), A., i, 206.
- absorption of ions by the roots of (REDFERN), A., i, 614.
- action of soluble lead salts on (BONNET), A., i, 412.
- role of manganese salts in (McHARGUE), A., i, 906.
- manganese content of the leaves of, and its variation with age (BERTRAND and ROSENBLATT), A., i, 411.
- effect of neutral salts on heat coagulation of the protoplasm of (KAHO), A., i, 94, 205.
- effect of the reaction of nutritive solutions on the germination and growth of (HIXON), A., i, 1221.
- absorption of nutritive substances from soils (CASALE), A., i, 508.

- Plants**, effect of sunlight on the formation of essential oils in (CANALS), A., i, 907.
- behaviour of organic compounds in (CIAMICIAN and GALIZZI), A., i, 503.
- pectic substances of (CLAYSON, NORRIS, and SCHRYVER), A., i, 206.
- nutrition of, with mineral phosphates (V. WRANGELL), A., i, 1098.
- saccharophosphatase in (NĚMEC and DUCHOŇ), A., i, 206.
- action of selenium compounds on the growth of (STOKLASA), A., i, 614.
- action of selenium on the metabolism of, in presence of radioactive air and soil (STOKLASA, KŘÍŽKA, PĚNKAVA, ZELENKA, CHMELÁK, and JÁNSKÝ), A., i, 974.
- action of selenium, sulphur and tellurium salts on growth of (TUBINA), A., i, 707.
- effect of reaction of soils on the distribution of (ATKINS), A., i, 415, 509; (COMBER), A., i, 416; (FISHER), A., i, 510.
- germinating, effect of soil acidity on (LEMMERMANN and FRESERIUS), A., i, 510.
- chlorophyll-free, vitamin-B in (ORTON, McCOLLUM, and SIMMONDS), A., i, 974.
- green, constituents of (FRANZEN and SCHUMACHER; FRANZEN and KEYSNER; FRANZEN and STERN), A., i, 311, 975; (FRANZEN and OVERTAG), A., i, 616, 1223; (FRANZEN and HELWERT), A., i, 1102.
- higher, distribution of manganese in (BERTRAND and ROSENBLATT), A., i, 95.
- nitrogen metabolism of (CHIBNALL), A., i, 908, 1225.
- detection of the ψ -bases of anthocyanidins in (COMBES), A., i, 206.
- detection of tryptophan in (KRETZ), A., ii, 668.
- estimation and distribution of chlorine in (JUNG), A., i, 1098.
- occurrence and estimation of cobalt and nickel in (BERTRAND and MOKRAGNATZ), A., i, 1099.
- Plant cells**, hydrogen-ion concentration of (ATKINS), A., i, 411.
- permeability of (OSTERHOUT; BROOKS; KAHHO), A., i, 308.
- rhythmic precipitation in the membranes of (MÖLLER), A., i, 94.
- action of bases and salts on masses of (MACDOUGALL), A., i, 205.
- action of lipoids on (BOAS), A., i, 94, 613.
- Plant plasma**, toxic action of salts on (KAHHO), A., i, 311.
- Platinum**, L-series spectrum of (DUANE and PATTERSON), A., ii, 463; (WEBSTER), A., ii, 805.
- photoelectric wave-length of (STUHL-MANN), A., ii, 809.
- heated, emission of electrons from (LOCKROW), A., ii, 608.
- loss of weight of, after ignition (BALAREFF), A., ii, 579.
- diffusion of hydrogen through (SCHMIDT and LÜCKE), A., ii, 198.
- properties of hydrogen desorbed from (ANDERSON), T., 1153.
- catalytic action of (LANGMUIR), A., ii, 629.
- catalytic decomposition of hydrogen peroxide by (MAXTED), T., 1760.
- oxygen-hydrogen catalysis by (HOFMANN), A., ii, 490.
- colloidal, action of ultra-violet light on (SPEAB, JONES, NEAVE, and SHLAGER), A., ii, 11.
- Platinum salts**, hydrolysis of (ARCHIBALD and GALE), T., 2849.
- Platinic chloride**, compound of triethylene trisulphide and (RÄY), T., 1283.
- Platinum oxide**, use of, in catalytic reduction of organic compounds (VOORHEES and ADAMS), A., ii, 558.
- Chloroplatinic acid**, preparation of (RUDNICK), A., ii, 303.
- Platinum separation** :—
- separation of palladium from, by means of dimethylglyoxime (DAVIS), A., ii, 662.
- Platinum bases (platiniammines)** (TSCHUGAEV and PSCHENIOVYN), A., ii, 856.
- electrical conductivity of nitrites of (TSCHUGAEV and VLADIMIROV), A., ii, 814.
- Platinum black**, catalysis by (VAVON and HUSSON), A., ii, 631.
- Platinum electrodes**. See **Electrodes**.
- Platinum metals**, catalysis by (HOFMANN), A., ii, 276.
- Plotohæmolysin** (KABESHIMA), A., i, 608.
- Plotospasmin** (KABESHIMA), A., i, 608.
- Plotosus anguillaris*, toxin of (KABESHIMA), A., i, 608.
- Plototoxin** (KABESHIMA), A., i, 608.
- Poison** of scorpions (WILSON), A., i, 969.
- tetrodon (ISHIHARA), A., i, 85.
- toad's, constituents of (WIELAND), A., i, 199; (WIELAND and ALLEN), A., i, 784.
- Poisons**, specificity of (BERGELL), A., i, 970.

- Poisons**, sensitiveness of cells to (HANOVSKY), A., i, 697.
 influence of the reaction of the solution on the action of, on bacteria or tadpoles (LABES), A., i, 901.
 action of, on enzymes (SANTESSON), A., i, 1077.
 action of, on striated muscle (RIESSER and NEUSCHLOSZ), A., i, 1212.
- Poisoning** by arsenic (SCHELLBACH), A., i, 900.
 by barium salts (IONESCU), A., i, 199.
 (KRAFFT), A., i, 301; (BAUMANN), A., i, 898.
 by lead compounds, hæmatoporphyrin in urine in (SCHUMM), A., i, 609.
 with mercuric chloride (BORNSTEIN and KERB), A., i, 404.
 with methyl alcohol, biochemistry of (RABINOVITCH), A., i, 963.
 by strychnine, effect of magnesium sulphate on (FERRARA), A., i, 500.
- Polarity**, induced, of atoms (LAPWORTH), T., 416; (KERMAK and ROBINSON), T., 427.
- Polecium**, velocity of emission of the α -rays of (CURIE), A., ii, 606.
- Polonium** hydride (PANETH and JOHANNSEN), A., ii, 777.
- isoPoly-acids** (ROSENHEIM and LRYSEY), A., ii, 50.
- Polyamylases**, and their derivatives (PRINGSHEIM and PERSCH; PRINGSHEIM and DERNIKOS), A., i, 632; (PRINGSHEIM and GOLDSTEIN), A., i, 638.
 physiology of (PRINGSHEIM and MÜLLER), A., i, 411.
- Polyborates**. See under Boron.
- Polycyclic compounds**, chemistry of, in relation to their homocyclic unsaturated isomerides (FARMER, INGOLD, and THORPE), T., 128; (CHANDRASANA, INGOLD, and THORPE), T., 1542.
- Polymerisation** (FIELDING), A., ii, 23, 421.
 under critical conditions (MULLER), A., ii, 820.
- Polynuritis** in birds, amyolytic action of the liver and pancreas in (TIGER and SIMONNER), A., i, 195.
- Polyoxymethylene** (HAMMICK and BORRER), T., 2738.
- Polypeptides**, refraction of (HIRSCH and KUNZ), A., i, 781.
- Polysaccharides** (KARRER), A., i, 11; (ABDERHALDEN), A., i, 82; (KARRER, STAUB, and WÄLTJ), A., i, 229; (KARRER and BÜRKLIN; KARRER and SMIRNOV), A., i, 435; (KARRER and FLORONI), A., i, 1119.
- Polysaccharides**, constitution of (IRVINE, STEELE, and SHANNON), T., 1060; (IRVINE and HIRST), T., 1585; (ZWIKKER), A., i, 230, 432; (KOMATSU and KASHIMA), A., i, 811.
 digestion of (SHIMIZU), A., i, 82.
- Polythionic acids**. See under Sulphur.
- Pompelmuus** oil, constants of (GATTEFOSSÉ), A., i, 1167.
- Potassium**, bright metallic, preparation of, in nitrogen (BORNEMANN), A., ii, 444.
 absorption spectrum of the vapour of (DATTA), A., ii, 678; (NARAYANA and GUNNAIYA; HARRISON), A., ii, 679.
 arc spectrum of (FOOTE, MEGGERS, and MOHLER), A., ii, 598.
 electrical conductivity of, in liquid ammonia (KRAUS and LUCASSE), A., ii, 252.
 vapour, ionisation of, by light (WILLIAMSON), A., ii, 681.
 electrodeless discharge in (ROBERTSON), A., ii, 609.
 chemical constants of (LADENBURG and MINKOWSKI), A., ii, 191.
 crystal structure of (MCKEEHAN), A., ii, 709.
- Potassium salts**, absorption of, by animal cells (MITCHELL, WILSON, and STANTON), A., i, 196.
 action of, on isolated tissues, and its substitutes (CLARK), A., i, 399.
 antagonism of calcium salts and (SPIRO), A., i, 85.
 replacement of, by uranium and thorium in Ringer's solution (ZWAARDEMAKER), A., i, 1213.
 distribution of, in blood (KRAMER and TISDALL), A., i, 1087.
 estimation of magnesium in (VÜRTHEIM), A., ii, 870.
- Potassium azide**, action of iodine with, in presence of carbon disulphide (BROWNE and HOEL), A., ii, 847.
 azidodithiocarbonate (BROWNE and HOEL), A., ii, 848.
 bromide, ionisation of, in aqueous solutions (PEARCE and HART), A., ii, 254.
 antimoniodobromide (VOURNAZOS), A., ii, 651.
 platinibromide, hydrolysis of (ARCHIBALD and GALE), T., 2849.
 carbonate, equilibrium of, with barium sulphate and with calcium oxalate (RAMANN and SALLINGER), A., ii, 181.
 chlorate, action of radium rays on (KAILAN), A., ii, 543.

Potassium chlorate, equilibrium of potassium nitrate, water, and (TODA), A., ii, 748.
perchlorate, preparation of, from potassium chlorate (LENHER, STONE, and SKINNER), A., ii, 287.
 chloride, electrical conductivity of solutions of, in presence of sucrose (KIERAN), A., ii, 812.
 platinichloride, refractive index of (RAITERI), A., ii, 541.
 stannichloride, crystal structure of (DICKINSON), A., ii, 287.
 chromate, equilibrium of silver iodate with (RAMANN and SALLINGER), A., ii, 131.
dichromate as a standard in iodimetry (VOSBURGH), A., ii, 863.
 hydroxide, preparation of, in alcoholic solution for volumetric work (MCALLUM), A., ii, 54.
 heats of neutralisation of (RICHARDS and ROWE), A., ii, 425.
 iodate and iodide, action of, on hydroxy-acids (KRISHNA and POPE), T., 798.
 iodide, compound of pyridinebetaine and (CASSELLA & Co.), A., i, 860.
 deflection of iodates in (LACHARTRE), A., ii, 716.
permanganate, properties of (FESTER and BRUDE), A., ii, 771.
 absorption spectrum of (HAGENBACH and PERCY), A., ii, 600.
 solubility of, in solutions of potassium and sodium sulphates (TRIMBLE), A., ii, 374.
 thermal decomposition of (MOLES and CRESPI), A., ii, 374.
 velocity of the reduction of, by formic acid (HOLLUTA), A., ii, 448; (HOLLUTA and WEISER), A., ii, 628.
 velocity of reaction of hydrogen peroxide with (ZAWIDZKI), A., ii, 207.
 oxidation of side-chains with (BIGELOW), A., i, 998.
 effect of filter paper on titrations with (SIMPSON), A., ii, 234.
 nitrate, equilibrium of potassium chlorate, water, and (TODA), A., ii, 748.
 cadmium nickel nitrite (CUTTICA), A., ii, 449.
 ammonium phosphates (CORELLI), A., ii, 287.
 plumbite, interaction of methyl iodide and (DRUCE), A., i, 516.
 silicate, equilibrium of potassium tungstate with (VAN LIEMPT), A., ii, 775.

Potassium silicofluoride, refractive index of (RAITERI), A., ii, 541.
 sulphate, equilibrium of aluminium sulphate, water, and (BRITTON), T., 982.
 equilibrium of ammonium and magnesium sulphates, water, and (WESTON), T., 1223.
 equilibrium of copper sulphate, ammonium sulphate, water, and (HAYAMI), A., ii, 748.
 hydrogen sulphate, preparation of (DOMINIK), A., ii, 145.
 alum, mordanting of wool with (PADDON), A., ii, 822.
 tungstate, equilibrium of, with potassium silicate and with sodium tungstate (VAN LIEMPT), A., ii, 775.
Potassium organic compounds :—
 cyanide, crystal structure of (BOZORTH), A., i, 441; (COOPER), A., i, 1128.
 cadmium, mercury, and zinc cyanides, crystal structure of (DICKINSON), A., i, 532.
 ferricyanide, use of, in iodimetry (KOLTHOFF), A., ii, 224.
 decomposition of, by heat (CUTTICA), A., i, 441.
 ferrocyanide, decomposition of, in daylight (BAUDISCH and BASS), A., i, 993.
 solubility of, in water (FABRIS), A., i, 329; (VALLANCE), A., i, 724.
 action of, on silver haloids (BONINO), A., ii, 78.
 use of, in electrometric titration (KOLTHOFF), A., ii, 580.
 estimation of, with permanganate (KOLTHOFF), A., ii, 537.
 phenyl selenate (MEYER and WAGNER), A., i, 620.
Potassium detection and estimation :—
 detection of, in animal and vegetable cells (MACALLUM), A., ii, 869.
 estimation of (HAZEN), A., ii, 658.
 estimation of, volumetrically (MACHELEIDT), A., ii, 227.
 estimation of, in presence of magnesium, sodium, phosphates, and sulphates (WATSON), A., ii, 586.
 estimation of, in soils (JONES and REEDER), A., ii, 85.
Potassoxyethoxydiphenylketen (STAUDINGER and MEYER), A., i, 1015.
Potatoes, manganese content of (BODE and HEMBD), A., i, 415.
Potential, differences of, at the surface of two liquids (FREUNDLICH and GYEMANT), A., ii, 342.

- Powders**, adsorbing, determination of the surface of (PANETH and VORWERK), A., ii, 618.
- Praseodymium** oxide, density and molecular volume of (PRANDTL), A., ii, 879.
- Precipitates**, carrying down of soluble salts by (DUTOIT and GROBET), A., ii, 550.
- Pregl's solution**, composition of (BACHSTEZ), A., ii, 792.
- Pregnancy**, chemistry of (KILLIAN and SHERWIN; CALDWELL and LYLE), A., i, 489.
- Probability**, correlation of entropy and (LINHART), A., ii, 257, 738.
- Procaine**. See Novocaine.
- Propane**, amino- and hydroxy-chloro-derivatives, constitution of (SMITH and PLATON), A., i, 1121.
- cyclo***Propane**, preparation of, pure (TRAUTZ and WINKLER), A., i, 926.
- isomerisation of (TRAUTZ and WINKLER), A., i, 926.
- cyclo***Propane**, nitro-, action of alkalis on (KOHLE and SMITH), A., i, 457.
- cyclo***Propane-1-carboxylamide-1-carboxylic acid**, ethyl ester (INGOLD, SAKO, and THORPE), T., 1189.
- cyclo***Propanecarboxylhydroxamic acid**, and its salts and derivatives (JONES and SCOTT), A., i, 453.
- cyclo***Propane-1-carboxylic acid**, 1-amino-, and its derivatives, and 1-hydroxy- (INGOLD, SATO, and THORPE), T., 1196.
- cyclo***Propane-1:1-dicarboxylamide** (INGOLD, SAKO, and THORPE), T., 1189.
- cyclo***Propane-1:1-dicarboxylbromoamide** (INGOLD, SAKO, and THORPE), T., 1190.
- cyclo***Propane-1:1-dicarboxylic acid**, ethyl ester, and amides, and condensation products with carbamides (DOX and YODER), A., i, 180.
- Propane- α -diol**, β -chloro- β -nitro-, and its dibenzoate (SCHMIDT and WILKENDORF), A., i, 314.
- α -**Propanedisulphonyldihydraside** (CLUTTERBUCK and COHEN), T., 128.
- α -**Propanedisulphonyldiphenylhydrazide** (CLUTTERBUCK and COHEN), T., 127.
- 5-cyclo****Propanespirohydantoin**, and dibromo- (INGOLD, SAKO, and THORPE), T., 1190.
- Propanesulphonic acid**, β - γ -dichloro-, ammonium and barium salts (POPE and SMITH), T., 1169.
- Propane- α - γ -tricarboxylic acid** (DICKENS, KON, and THORPE), T., 1502.
- Propan- β -oleamphorecarboxylic acids**, isomeric (HALLER and RAMART-LUCAS), A., i, 460.
- Propan- β -oleamphorecarboxylolides**, isomeric (HALLER and RAMART-LUCAS), A., i, 460.
- iso***Propenylacetylene**. See γ -Methyl- Δ^2 -butenine.
- Propionic acid**, lead diphenyl salt (GODDARD, ASHLEY, and EVANS), T., 981.
- Propionic acid**, bromo-, sodium salt, separation of bromine from (v. EULER and FAHLANDER), A., ii, 360.
- β -bromo- and β -chloro-, preparation of (ROJAHN), A., i, 105.
- α - and β -chloro-, ethyl esters, action of magnesium ethyl bromide with (MOUREU and BARRETT), A., i, 4.
- β -cyano-, ethyl ester, and its oxime (CARRIERE), A., i, 318.
- Propionitrile**, β -hydroxy- (RÖHM and HAAS), A., i, 531.
- Propionylbutyric acid** (BRUYLANTS), A., i, 531.
- 3-Propionyl-*p*-cresol**, derivatives of (v. AUWERS, HILLIGER, and WULF), A., i, 1192.
- Propionylxypropionyl chloride**, α -chloro- and its anilide (BLAISE and MONTAGNE), A., i, 716.
- Propiophenones**, $\alpha\alpha\beta\beta$ -substituted, decomposition of, with sodamide (RAMART and ALBESCO), A., i, 662.
- β -Propoxybutyronitriles** (BRUYLANTS), A., i, 924.
- 7-*n*-Propoxy-7:12-dihydro- γ -benzophenarsazine** (LEWIS and HAMILTON), A., i, 188.
- 6-Propoxy-2-methylquinoline**, and its ethiodide (GUTKUNST and GRAY), A., i, 951.
- n*-Propyl *p*-carbamidophenyl ether**, β -hydroxy- (SPECKAN), A., i, 580.
- hydrogen selenate, and its metallic salts (MEYER and WAGNER), A., i, 620.
- iso***Propyl alcohol**, pharmacology of (MACHT), A., i, 1093.
- oxidation of, by potassium permanganate (EVANS and SEFTON), A., i, 1108.
- fate of, in the animal organism (POHL), A., i, 498.
- iso***Propyl alcohol**, α -chloro- α -nitro-, and α -nitro-, sodium derivative (SCHMIDT and WILKENDORF), A., i, 314.
- n*-Propylalanine**. See α -Methyl-*n*-valeric acid, α -amino-.
- α -cyclo***Propylalanine**. See *α -cyclo***Propylpropionic acid**, α -amino-.

- isoPropylallylbarbituric acid**, preparation of (HOFFMANN-LA ROCHE & Co.), A., i, 372.
- N-isoPropylaminoacetic acid**, α -cyano-, ethyl ester, and its hydrochloride (SCHEIBLER and BAUMGARTEN), A., i, 657.
- n- and isoPropylammonium** ruthenipentabromides (GUTHRIE and KRAUSS), A., i, 16.
- isoPropylbenzoic acid**, *o*-amino-*p*- α -hydroxy-, and *o*-nitro-*p*- α -hydroxy- (PHILLIPS), A., i, 955.
- n-Propylboric acid** (KRAUSE and NITSCHKE), A., i, 22.
- Propyl-n-butylamine** (v. BRAUN and WEISMANTEL), A., i, 1151.
- Propylbutylarsinic acid** (QUICK and ADAMS), A., i, 600.
- isoPropyl isobutyl ketone semicarbazone** (LEROIDE), A., i, 217.
- N-Propylconine**, γ -hydroxy-, hydrochloride (v. BRAUN, BRAUNSDORF, and RATH), A., i, 760.
- 3-Propyl-p-cresol**, α -amino-, benzoyl derivative (v. AUWERS, HILLIGER, and WULF), A., i, 1193.
- isoPropylisocrotylcarbinol**, and its acetyl derivative (KRESTINSKI), A., i, 1129.
- Propylene**, preparation of, pure (TRAUTZ and WINKLER), A., i, 909.
production of, from cyclopropane (TRAUTZ and WINKLER), A., i, 927.
action of selenium chloride on (BOORD and COPE), A., i, 421.
- Propylenediamine stannichloride** (DRUCE), A., i, 639.
- a*-cycloPropylethyldimethylamine**, salts of (DEMJanov and DOJARENKO), A., i, 1014.
- cycloPropylethylene glycol** (DEMJanov and DOJARENKO), A., i, 1014.
- 9-n-Propylfluorene**, 2:7-dibromo-9-hydroxy- (SIEGLITZ and JASSOY), A., i, 821.
- N-isoPropylglycine**, and its hydrochloride (SCHEIBLER and BAUMGARTEN), A., i, 657.
- N-isoPropylideneaminoacetic acid**, sodium salt (SCHEIBLER and BAUMGARTEN), A., i, 657.
- 4-isoPropylidene-cyclohexanone**, separation and reduction of (HENDERSON, ROBERTSON, and BROWN), T., 2721.
- isoPropylindigotin**, 6:6-di- α -hydroxy- (PHILLIPS), A., i, 955.
- n-Propylisatoid** (HELLER and BENADE), A., i, 582.
- isoPropylmalonamide** (DOX and YODER), A., i, 816.
- Propylmenthol** (OGATA and MIYASHITA) A., i, 844.
- Py-N-Propyl-Ind-N-methylarsine** chloride (KERMAK, PERKIN, and ROBINSON), T., 1894.
- β -Propylmethylheptenol** (HELFERICH and GEHRKE), A., i, 9.
- Propylphenetidine**, β - γ -dihydroxy-, and its oxalic ester (BERGMANN, ULPTS, and CAMACHO), A., i, 1181.
- N-Propylpiperidine**, γ -hydroxy-, and its derivatives (v. BRAUN, BRAUNSDORF, and RATH), A., i, 760.
- a*-cycloPropylpropionic acid**, α -amino-, and its salts (ZELINSKY and DENGIN), A., i, 1127.
- 2-isoPropylquinoline**, 2- α -hydroxy- (SCHEIBLER and FISCHER), A., i, 1110.
- isoPropylsemicarbazide**, and its salts and derivatives (NEIGHBORS, FOSTER, CLARK, MILLER, and BAILEY), A., i, 881.
- isoPropylstannonic acid**, and its salts and derivatives (DRUCE), T., 1859.
- N-Propyltetrahydroisoquinoline**, γ -hydroxy-, and its salts (v. BRAUN, BRAUNSDORF, and RATH), A., i, 761.
- Prosolannellie acid** (WIELAND and SCHULENBURG), A., i, 346.
- Protamines** (GROSS), A., i, 784.
- Protease**, estimation of, in blood (BACH and ZUBKOWA), A., i, 392.
- Proteic acids**, hydroxy- (EDLBACHER), A., i, 692, 883.
- Proteins**, physical chemistry of (COHN), A., i, 882.
mobility of the ions of (PAULI), A., i, 478.
potential differences between solutions of (LOEB), A., i, 692.
heat of coagulation of (HIRSCH-POGANY), A., i, 596.
colloid chemistry of (FODOR), A., i, 691.
coagulation of, by sunlight (YOUNG), A., i, 386.
coagulation of, by heat (LEPESCHKIN), A., i, 1198; (LÜERS and LANDAUER), A., ii, 697.
influence of acids on the osmotic pressure of solutions of (LOEB), A., ii, 742.
influence of aggregates on the potential and osmotic pressure of (LOEB), A., i, 883.
determination of the isoelectric point of (LÜERS and LANDAUER), A., 881.
effect of salts on the isoelectric point of (LOEB), A., i, 882.
solubility of, at the isoelectric point (COHN), A., i, 882.
precipitation of (CRISTOL), A., ii, 583.

- Proteins and their derivatives, fractionation of** (RAKUZIN), A., i, 957.
adsorption of, by aluminium hydroxide (RAKUZIN), A., i, 1199.
free amino-groups in (ENGELAND; EDLBACHER), A., i, 279.
amino-acids from products of hydrolysis of (BUSTON and SCHRYVER; JONES and JOHNS), A., i, 182.
basic derivatives of (FELIX), A., i, 295.
electric kataphoresis of metallic compounds of (BENEDICENTI and REBELLO-ALVES), A., ii, 683.
action of β -naphthalenesulphonyl chloride on (EDLBACHER and FUCHS), A., i, 279.
compounds of nucleic acid and (STEUDEL and PEISER), A., i, 1200.
sulphur in (HOFFMAN and GORTNER), A., i, 429.
"double-nitrogen" as a measure of degradation of (HAHN), A., i, 291.
toxicity of (PENTIMALLI), A., i, 302.
action of, in the organism (LUTTICHAU), A., i, 892.
effect of polysaccharides on sparing of (SHIMIZU), A., i, 83.
replacement of, by urea in diet (MORGEN, SCHÖLER, WINDHEUSER, and OHLMER), A., i, 293.
absorption and utilisation of, in liver diseases (AUB and MEANS), A., i, 193.
digestibility of (JONES and WATERMAN), A., i, 893.
digestion of, by trypsin (NORTHROP), A., i, 693.
Bence Jones', distribution of nitrogen in (LUSCHER), A., i, 1199.
of blood sera, ultra-violet absorption spectra and rotation of (LEWIS), A., ii, 245.
natural (SCHMIDT and BRAUNSDORF), A., i, 781.
of organs, decomposition of (THOMAS), A., i, 790.
of serum (PIETTRE and VILA), A., i, 63.
in yeast, decomposition of, during fermentation (IVANOV), A., i, 202.
- Proteins and soaps** (FISCHER, McLAUGHLIN, and HOOKER), A., ii, 480, 829.
precipitation of (HILLER and VAN SLYKE), A., i, 1074.
from physiological fluids (GRIGAUT and ZIZINE), A., ii, 886.
detection of, and their derivatives, colorimetrically (RAKUZIN), A., ii, 670.
- Proteins, estimation of, in blood** (HOWE), A., ii, 171, 172; (BIERRY and MOQUET), A., ii, 886.
estimation of, in colostrum (HOWE), A., ii, 670.
estimation of, in plasma (WU), A., ii, 406.
estimation of amino-acids in (FOLIN and LOONEY), A., ii, 539.
estimation of ammonia in (FROIDEVAUX), A., ii, 454.
estimation of tryptophan in (LÜSCHER), A., i, 1199.
estimation of tyrosine in (v. FÜRTH and FLEISCHMANN), A., ii, 406.
- Protocatechualdehyde, preparation of** (HAMBURGER), A., i, 556.
- Protocatechuic acid, compound of diketopiperazine and** (POVARNIN and TICHOMIROV), A., i, 1185.
- Protoplasm, chemical constitution of** (WALTER), A., i, 308.
relation between the colloidal state and the physiological functions of (WURMSER and JACQUOT), A., i, 1221.
- Prunus avium* (cherry), acids of** (FRANZEN and HELWERT), A., i, 1102.
- Pseudowavellite** (HENRICH and HILLER), A., ii, 860.
- Psoriasis, composition of the scales in** (ABDERHALDEN and ZOERN), A., ii, 794.
- Ptyalin, thermostability of** (ERNSTRÖM), A., i, 599.
- Puerperal fever, changes in blood and urine in** (COUINAUD and CLOGNE), A., i, 401.
- Pump, water** (WETZEL), A., ii, 139.
- Purine metabolism. See Metabolism.**
- Purinecarboxylic acids, preparation of** (MERCK, WOLFES, and KORNICK), A., i, 1071.
- Purpura aperta* and *lapillus*, colouring matters from** (FRIEDLANDER), A., i, 793.
- Purpurin diboroacetate, and its acetyl derivative** (DIMROTH and FAUST), A., i, 157.
2-methyl ether, 3-bromo- (DIMROTH, SCHULTZE, and HEINZE), A., i, 158.
- Pycnometry** (SAAE), A., ii, 549.
- Pyranhydrones** (SCHNEIDER and ROSS), A., i, 1171.
- Pyrazole derivatives, electrochemical oxidation of** (FICHTER and DE MONTMOLLIN), A., i, 470.
- Pyrazoleanthrone-yellow, constitution of** (MAYER and HEIL), A., i, 877.
- Pyrene, synthesis of** (FLEISCHER and RETZE), A., i, 1138.

- Pyridine**, reduction of (LEBEAU and PICON), A., ii, 48; (DIMROTH and FRISTER), A., i, 678.
 derivatives, reduction of (THIELEPAPE and SPRECKEISEN), A., i, 1191.
 electrolysis of metallic salts in (MÜLLER; MÜLLER and DUSCHEK), A., ii, 612.
 compounds of, with metallic salts (PETERS), A., i, 48.
 compound of sodammonium with (LEBEAU and PICON), A., i, 48.
 hydroferrocyanide (CUMMING), T., 1297.
 aryl derivatives of (DILTHEY), A., i, 272.
 diisopropyl- and isopropyl-stannisalts (DRUCK), T., 1861.
 additive compounds of, with metallic alkyl and aryl xanthates (DUBSKY and APTEKMAN), A., i, 104.
 additive compounds of *s*-trinitroanisole with (GIUA), A., i, 572.
 detection of (LEHNER; GORIS and LARSONNEAU), A., ii, 795.
 separation of quinoline and (CUMMING), T., 1298.
- Pyridine**, 2-amino-, benzoyl derivative, and its salts (TSCHITSCHIBABIN and BYLINKIN), A., i, 573.
- Pyridines**, physical properties of aqueous solutions of (JONES and SPEAKMAN), A., i, 171.
 methylated, preparation of, pure (HEAP, JONES, and SPEAKMAN), A., i, 171.
 arylated (DILTHEY, NÜSSLEIN, MEYER, and KAFFER), A., i, 949.
- Pyridine-4-acetamide**, 2:6-dihydroxy- (INGOLD and NICKOLLS), T., 1644.
- Pyridinebetaine**, compounds of, with metallic salts (CASSELLA & Co.), A., i, 860.
- Pyridine-3-carboxylic acid**. See Nicotinic acid.
- Pyridine-3-carboxyldiethylamide** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 860.
- Pyridine-2:3-dicarboxylic acid**, methyl ester (MERCK), A., i, 950.
- Pyridineiridide acid**, pentachloro-, and its potassium salt (DELÉPINE), A., i, 859.
- Pyridine-4-propionic acid**, 2:6-dihydroxy-, and its salts (FARMER), T., 2018.
- Pyridinium salts** (BARNETT and COOK), T., 792.
 tetrachloroferrate (WEINLAND and KISLING), A., i, 363.
 ruthenipentabromide (GUTHIER and KRAUSS), A., i, 16.
- Pyrimidine**, 5-bromo-6-hydroxy-, and 6-chloro-5-bromo-, and their salts (CHERBULIEZ and STAVRITCH), A., i, 581.
- Pyrimidines** (DOX and YODER; HORN), A., i, 374; (CHERBULIEZ and STAVRITCH), A., i, 581; (JOHNSON and BAUDISCH), A., i, 328.
- spiro***Pyrimidines** (DOX and YODER), A., i, 180.
- Pyrimidines**, 2-thiol-, alkylation of (HORN), A., i, 374.
- Pyrimidine-4-carboxylic acid**, 5-bromo-6-hydroxy-, and 6-hydroxy-, and their derivatives (CHERBULIEZ and STAVRITCH), A., i, 581.
- Pyrites**, use of, as a catalyst in Friedel and Crafts' reaction (SMYTHE), T., 1270.
 action of acetylene on (STEINKOPF and HEROLD), A., i, 850.
 estimation of sulphur in (GADAIS), A., ii, 79; (CHAUDRON and JUREBOIRARD), A., i, 311.
- Pyrocatechol**, complex salts of, and its derivatives (REIHLEN), A., i, 1146; (REIHLEN and SAPPER), A., i, 1147.
- Pyrofulmin** (LANGHANS), A., i, 328.
- Pyrogallol**, absorption of oxygen by, in gas analysis (HOFFMANN), A., ii, 582.
 compound of diketopiperazine and (POVARNIN and TICHOMIROV), A., i, 1185.
 equilibrium of triphenylcarbinol with (KREMANN, HOHL, and MÜLLER), A., i, 138.
 1:3-dimethyl ether, and its acetyl derivative, amino- and nitro-derivatives of (BRAND and COLLISCHONN), A., i, 452.
- Pyrogallolcamphorein**, and its tetraacetyl derivative (SINGH, RAI, and LAL), T., 1428.
- Pyromellitic acid** (PHILIPPI; PHILIPPI and RIE; PHILIPPI and THELEN; PHILIPPI, SEKA, and FROESCHL; PHILIPPI, SEKA, and ROBINSON), A., i, 837.
- Pyromellitic acid**, dibromo- (PHILIPPI, SEKA, and ROBINSON), A., i, 837.
- Pyromucic acid**, bactericidal action of (KAUFMANN), A., i, 804.
- Pyromucylhydroxamic acid**, and its acetyl ester, and their salts (JONES and HURD), A., i, 249.
- γ-Pyrone**, action of aniline on (BORSCHKE and BONACKER), A., i, 50.
- Pyrophosphoric acid**. See under Phosphorus.
- Pyroprosolanellie acid** (WIELAND and SCHULENBURG), A., i, 346.

Pyrocolannellie acid (WIELAND and SCHULENBURG), A., i, 846.

Pyrosulphuryl chloride. See under Sulphur.

Pyrrrole, catalytic hydrogenation of (PUTOCHIN), A., i, 1176.

behaviour of, in the body (SHIMIZU), A., i, 91.

melanin from (RONDONI), A., i, 64.

derivatives (KUSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 857.

iodo-derivatives (PIERONI), A., i, 763.

Pyrrroles (FISCHER and ZERWECK), A., i, 758; (FISCHER and HERRMANN), A., i, 1054; (FISCHER, SCHNELLER, and ZERWECK), A., i, 1055.

constitution of polymeric (PIERONI and MOGGI), A., i, 766.

Pyrrrolealdehydes (FISCHER and ZERWECK), A., i, 758.

Pyrrrole-3:5-diacetic-2:4-dicarboxylic acid, methyl ester (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 859.

Pyrrrole-3:5-dicarboxylic acid, 2:4-dinitro-, ethyl ester (FISCHER and ZERWECK), A., i, 759.

Pyrrrolidine, preparation of (PUTOCHIN), A., i, 1176.

action of formaldehyde on (PUTOCHIN), A., i, 1176.

5-Pyrrrolidine-2-carboxylic acid, *n*-butyl ester (CHILES and NOYES), A., i, 925.

α -Pyrrroleinohonic acid, and its sodium salt (CIUSA), A., i, 1062.

Pyrus aucuparia. See Mountain ash.

Pyrus coronaria, organic acids from (SANDO and BARTLETT), A., i, 100.

Pyruvaldehyde, *p*-ethoxyphenylhydrazone (JACOBSEN), A., i, 596.

Pyruvic acid, as a fermentation product (v. GRAB), A., i, 306.

decarboxylation of, by yeast (HANK and KOEHLER), A., i, 409.

as the acetylating agent of the body (KNOOP), A., i, 487.

in urine (FRICKE), A., i, 495.

p-ethoxyphenylhydrazone (JACOBSEN), A., i, 596.

m-methoxyphenylmethylhydrazone (KERMAK, PERKIN, and ROBINSON), T., 1881.

o- and *p*-nitrophenylhydrazones, alkali salts of (CIUSA and RASTELLI), A., i, 1073.

as-p-nitrophenylmethylhydrazone (CIUSA and RASTELLI), A., i, 1074.

Pyrylium compounds (DILTHEY and KAFFER), A., i, 668.

Pyrylium salts, synthesis of (PRATT and ROBINSON), T., 1577.

Q.

Quantum theory, application of, to gases (SCHAPOSCHNIKOV), A., ii, 889.

Quartz, heat of crystallisation of (RAY), A., ii, 685.

crystal structure of (HUGGINS), A., ii, 641.

Quinaldine synthesis (LINDNER), A., i, 687.

Quinaldinic acid, esters and quaternary salts of (MILLS and HAMER), T., 2008.

Quince tree, exudation from the stems of (v. LIPPMANN), A., i, 1223.

Quindoline methochloride and methosulphate (ARMIT and ROBINSON), T., 836.

Quindolines (GRANDMOUGIN), A., i, 584.

Quinhydrone electrode. See Electrode.

Quinic acid, synthesis of (KAUFMANN), A., i, 464.

derivatives of (HALBERKANN), A., i, 174.

Quinicoylglycineanilide dihydrochloride (HEIDELBERGER and JACOBS), A., i, 673.

Quinicoylglycine-*p*-hydroxyanilide (HEIDELBERGER and JACOBS), A., i, 673.

Quinidine oxide, picrate of (SPEYER and BECKER), A., i, 675.

Quinine, effect of, on blood (LUGER), A., i, 81.

fixation and distribution of, in blood (RONA and BLOCH), A., i, 290, 484.

salts, ionisation and luminescence of (SCHMIDT), A., ii, 182.

detection of quinotoxine in (GANASINI), A., ii, 539.

estimation of (LIVERSEDGE and ANDREWS), A., ii, 669.

acetoxybenzoyl derivatives of (MERCK, DIEHL, and MAYEN), A., i, 46.

oxide, salts of (SPEYER and BECKER), A., i, 675.

separation of strychnine and (WARREN and CLARK), A., ii, 171; (EVERS), A., ii, 669.

Quinine alkaloids, distribution of, in the organism (BOECKER), A., ii, 896.

Quinizarin, action of bromine on (DIMROTH, SCHULTZE, and HEINZE), A., i, 157.

Quinizarin, 2-fluoro-, and its diacetyl derivative (DIMROTH and HILCKEN), A., i, 158.

Quinizarin-2:3-dipyridinium dibromide, and its phenobetaïne (BARNETT and COOK), T., 1384.

Quinizarin-2-phenylsulphone, and its diacetyl derivative (DIMROTH and HILCKEN), A., i, 158.

Quinizarinquinone dibromide and methobromide (DIMROTH, SCHULTZE, and HEINZE), A., i, 157.

Quinizarinquinone, 5- and 6-hydroxy- (DIMROTH and HILCKEN), A., i, 158.

Quinobensidine, and its bishydrazine (WIELAND, WECKER, and HAAS), A., i, 780.

Quinocyanines, chemistry of (KÖNIG), A., i, 1188.

Quinol, electromotive equilibria of quinone with (GRANGER and NELSON), A., i, 43.
chlorination of (ECKERT and ENDLER), A., i, 392.
oxidation of, in presence of aliphatic amines (HARGER), A., i, 539.
compound of diketopiperazine and (POVARNIN and TICHOMIROV), A., i, 1185.

Quinol, nitro-derivatives, and their benzoates (KEHRMANN, SANDOZ, and MONNIER), A., i, 33.

Quinolecamphorcin (SINGH, RAI, and LAL), T., 1427.

Quinoline, solubility of lithium chloride in (WALTON and WISE), A., i, 370.
hydroferrocyanides (CUMMING), T., 1298.
separation of pyridine and (CUMMING), T., 1298.

Quinoline, 5-amino-, stanni- and stannochloride (DRUCE), A., i, 1206.
8-hydroxy-, magnesium compound (MÖRNER), A., ii, 659.

*iso***Quinoline** (HARRIS and POPE), T., 1029.
stannichloride (DRUCE), A., i, 1206.

Quinolines (PALKIN and HARRIS), A., i, 951.

Quinoline series (THIELEPAPE), A., i, 271.
preparation of amino-acids of the (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 1179.

Quinoline-4-carboxylic acid, and its hydrazide, and 2-chloro-, ethyl ester, and 2-iodo- (THIELEPAPE), A., i, 271.

6-Quinoline-2:4-dicarboxylic acid, 6-hydroxy-, and its ammonium hydrogen salt (HALBERKANN), A., i, 174.

*iso***Quinoline-red** (HARRIS and POPE), T., 1029.

Quinolinium ruthenibromides (GUTHIER and KRAUSS), A., i, 16.

Quinolsulphamphthalein, hydroxy- (DUTT), T., 2391.

Quinone. See Benzoquinone.

Quinones, action of, on bacteria (MORGAN and COOPER), A., i, 204.

o-**Quinones** (SCHONBERG), A., i, 27; (SCHÖNBERG and KRAEMER), A., i, 663.

Quinone-1-imide-aci-2(or 4)-nitro-4(or 2)-mercury (KHARASCH, LOMMEN, and JACOBSSON), A., i, 604.

Quinotoxine, salts of tropic acid with (KING and PALMER), T., 2584.
detection of, in quinine salts (GANASINI), A., ii, 539.

R.

Rabbits, zinc content of the organs of (BERTRAND and VLADESCO), A., i, 493.

Racemic acid, optical activation of, by *l*-malic acid (MCKENZIE and WALKER), T., 349.
dissociation of, in solution (PENNYCUICK), A., i, 624.

Racemic acids, action of bacteria and moulds on (CONDELLI), A., i, 410.

Racemic compounds, active (LANDRIEU), A., i, 808.
transformations during the fission of (WEGSCHEIDER), A., i, 441.

Racemisation of optically active acid amides (MCKENZIE and SMITH), T., 1348.

Radiation, absorption and dispersion of (WEIGERT), A., ii, 605.
and chemical reactions (TAYLOR; LEWIS), A., ii, 386.

Radiation theory, and its applications (PERRIN; LEWIS; BALY; LOWRY), A., ii, 628; (DHAR), A., ii, 730.

Radioactive elements (HOFFMANN), A., ii, 184.
Meitner nuclear model of (NEUBURGER), A., ii, 183.
atomic disintegration of (MEITNER), A., ii, 15.
isotopy of (NEUBURGER), A., ii, 107, 250.
oxidising properties of (LEWIS and JALOUSTRÉ), A., ii, 13.

minerals, new (SCHOEP), A., ii, 77, 450, 451.

substances, β -ray spectra from (MEITNER), A., ii, 416.
disintegration of (HEVESTY), A., ii, 608.
adsorption of (EBLER and VAN RHYN), A., ii, 13.

Radioactive substances, distribution of, in solutions (LACHS and WERTENSTEIN), A., ii, 681.

use of, as indicators (PANETH), A., ii, 785.

Radioactivity of water. See under Water.

Radio-elements. See Radioactive elements.

Radiothorium, effect of, on metabolism (MIYADERA), A., i, 966.

Radium content of British coal (DRAKELEY and SMITH), T., 237.

adsorption of, by barium sulphate (GERMANN), A., ii, 16.

chemical action of the penetrating rays of (KAILAN), A., ii, 466, 543.

action of the penetrating rays of, on colloids (FERNAU and PAULI), A., ii, 202.

synthesis of carbon compounds from air by means of (GLEW), A., ii, 607.

rays, production of hydrogen peroxide from water by (KAILAN), A., ii, 106.

γ -rays from, absorption of, by luminous compounds (OWEN and PAGE), A., ii, 108.

effect of, on the germination of seeds (STOKLASA), A., i, 613.

emanation. See Niton.

chloride, positive ray analysis of the gases given off by (THOMSON), A., ii, 565.

residues, ionium in (RONA), A., ii, 250.

solutions, permanency of (BECKER), A., ii, 810.

estimation of small quantities of (SZILARD), A., ii, 586.

Radium-Band-C, β -ray spectra of (CHADWICK and ELLIS), A., ii, 802.

adsorption of, by ferric hydroxide (CRANSTON and HUTTON), T., 2843.

Radium-C, emission of α -particles by (SHENSTONE and SCHLUNDT), A., ii, 465.

Raffinase, specific nature of (WILLSTÄTTER and KUHN), A., i, 284.

Raffinose, preparation of (CLARK), A., i, 323.

Ragweed, constituents of the pollen of (HEYL), A., i, 1224.

Rain-water. See under Water.

Rana temporaria (common frog), constituents of eggs of (FAURÉ-FREMIER and DE STREEL), A., i, 85.

Rape oil, fatty acids of (TOYAMA), A., i, 1113.

Rapic acid (GRABNER), A., i, 519.

Raspberry. See *Rubus idæus*.

Raspberry juice, cause of the clouding of (KUNZ-KRAUSE), A., i, 210.

Rayfish liver oil, constituents of (TSUJIMOTO and TOYAMA), A., i, 297; (TOYAMA), A., i, 895.

Rays, absorbed, photochemical efficiency of (WINTHER), A., ii, 808.

Becquerel, coloration and luminescence produced by (MEYER and PRZIBRAM), A., ii, 339.

positive, analysis by (THOMSON), A., ii, 565.

Röntgen, reflexion of, from crystals (DARWIN), A., ii, 416; (DARDORN), A., ii, 673.

determination of crystal structure by means of (CLARK and DUANE), A., ii, 483; (BIJVOET and KARSSEN; POSNJAK and WYCKOFF), A., ii, 499; (HULL and DAVEY; HULL), A., ii, 624.

investigation of crystal lattices with (GERLACH), A., ii, 36.

and thermodynamic equilibrium (DE BROGLIE), A., ii, 249.

absorption of (DE BROGLIE; DUANE), A., ii, 104; (RICHTMYER), A., ii, 105, 804; (WINGÅRDH), A., ii, 249.

in crystals (AURÉN), A., ii, 810.

scattering of, by anisotropic liquids (HÜCKEL), A., ii, 14.

scattering of, by crystals (BRAGG, JAMES, and BOSANQUET), A., ii, 184.

soft (KURTH), A., ii, 809.

excitation of (RICHARDSON and BAZZONI), A., ii, 14.

biological action of (PETRY), A., i, 204.

spectra of. See Spectra.

α -Rays, radioactivity resulting from bombardment by (SHENSTONE), A., ii, 415.

ionisation by (WILKINS), A., ii, 607.

luminous path of, in crystals (GEIGER and WERNER), A., ii, 183.

range of, in liquids and gases (TRAUBENBERG and PHILIPP), A., ii, 12.

β -Rays, scattering of, by thin metal sheets (GEIGER and BOTHE), A., ii, 13.

relation between the scattering of, and atomic number (GLASSON), A., ii, 183.

relation between γ -rays and (MEITNER), A., ii, 416, 732.

spectra of. See Spectra.

γ -Rays, absorption of, by luminous compounds (OWEN and PAGE), A., ii, 108.

- γ -Rays**, excitation of, by α -particles (SLATER), A., ii, 13.
 wave-lengths of (MEITNER), A., ii, 733.
 relation between β -rays and (MEITNER), A., ii, 416, 732.
 spectra of. See Spectra.
- Recrystallisation**, theory of (ALTERTHUM), A., ii, 623.
- Rectification** (GAY), A., ii, 120.
- Reductases** (SMORODINCEV), A., i, 1201.
- Reduction** by the Kishner-Wolff method (THIELEPAPE), A., i, 271; (THIELEPAPE and SPRECKELSEN), A., i, 1191.
 simultaneous oxidation and (KÖTZ and RATHERT), A., i, 236.
- Reductodehydrocholic acid**, and its esters, and derivatives (BORSCH and HALLWASS), A., i, 1158.
- Reeds**, sucrose from the roots of (V. LIPPMANN), A., i, 311.
- Refraction** and atomic structure (WASASTJERNA), A., ii, 1.
 atomic and molecular, of organic compounds (SWIENTOSLAWSKI), A., ii, 173.
 double, of colloidal solutions (ZACHER), A., ii, 102.
 molecular, atomic constants of (HUCKEL), A., ii, 195.
 coefficients of (V. AUWERS and KOLLIGS), A., ii, 174.
 relation of, to other properties (HERZ), A., ii, 409.
 of aromatic hydrocarbons (V. AUWERS), A., ii, 98.
 of molten salts (MEYER and HECK), A., ii, 241, 329.
 of substances with higher melting points (EISENLOHR), A., ii, 1.
- Refractive index** at corresponding temperatures (HERZ), A., ii, 97.
 of metallic salts in aqueous solution (LIMANN), A., ii, 173.
 of organic liquids (HIMSTEDT and WERTHEIMER), A., ii, 725.
 of phosphorescent sulphides (CURIE), A., ii, 801.
- Refractivity**, relation between atomic volume and (LE BAS), A., ii, 241.
- Refractometric analysis**. See Analysis.
- Refractories**, expansion of, at high temperatures (BOGITCH), A., ii, 115.
- Relativity**, laboratory test of the theory of (KING), A., ii, 439.
- Rennin** (*chymosin*), action of (HAMMARSTEN), A., i, 958.
 chemical action of (INICHOV), A., i, 960.
 coagulation of milk by (BAUR and HERZFELD), A., i, 284.
- Resacetophenone**, amino-, ω -benzoyl derivative (SONN and FALKENHEIM), A., i, 1164.
 ω -hydroxy- (*fisetole*), synthesis of (SONN and FALKENHEIM), A., i, 1163.
- Resins**, constituents of (ZINKE, FRIEDRICH, JOHANNSEN, and RICHTER), A., i, 667; (ZINKE, HANSELMAYER, and EHMER), A., i, 668.
 from spruce needles (V. EULER), A., i, 283.
 chemistry of (VESTERBERG; VESTERBERG and WESTERLIND), A., i, 825.
 relation between chemical constitution and ability to form (HERZOG and KREIDL), A., i, 1168.
- Resin acids** of the Coniferae (ASCHAN, FONTELL and SIMOLA), A., i, 1152.
- Resonance potentials** of gases (BOUCHER), A., ii, 608.
- Resorcinol**, constitution of, and its derivatives (FABRE), A., i, 1147.
 tautomerism of (FUCHS), A., i, 336.
 condensation of cinnamic acid with (SHORT and SMITH), T., 1808.
 compound of diketopiperazine and (POVARNIN and TICHOMIROV), A., i, 1185.
 use of, in qualitative inorganic analysis (LAVOYE), A., ii, 779.
- Resorcinol**, 4-chloro-2-nitroso- (FABRE), A., i, 1148.
 dinitroso-. See 1:2:3:4-Benzodiquinoxone-1:3-dioxime.
 dinitrosothiol- (WATSON and DUTT), T., 1943.
- Resorcinolcamphorein** (KRISHNA), T., 255; (SIRCAR and DUTT), T., 1284.
- Resorcinolcamphorein**, *tetrabromocamphoreosin* (SIRCAR and DUTT), T., 1284; (SINGH, RAI, and LAL), T., 1429.
- Resorcinolphthalphthalein**, hydrochloride and acetyl derivative of (FISCHER and BOLLMANN), A., i, 936.
- Resorcinolphenylosuccinein** (LAPWORTH and McRAE), T., 2722.
- Resorcinolsulphamphthalein** (DUTT), T., 2390.
- β -Resoreylaldehyde methyl ethers**, constitution of (ORT and NAUEN), A., i, 555.
- Respiration**, effect of anaesthetics on (BODINE), A., i, 889.
 effect of extirpation of the spleen on (ASHER and KODA; ASHER and DOUBLER), A., i, 286.
 effect of lack of calcium on basal metabolism in (PEDOTTI), A., i, 286.

- Respiration**, influence of hydrogen sulphide on (HAGGARD and HENDERSON), A., i, 1206.
 relation between oxygen and carbon dioxide in (THUNBERG), A., i, 889.
 of cells, physical chemistry of (WARBURG), A., i, 190.
 of dogs, effect of injection of sugars on (BÜRGER), A., i, 286.
 of fishes, physiology of (POWERS), A., i, 286.
- Respiratory exchange** in fresh-water fish (GARDNER, KING, and POWERS), A., i, 1084.
 metabolism. See Metabolism.
- Retina**, photochemistry of the (WEIGERT), A., ii, 11.
 liberation of phosphoric acid by the (LANGE and SIMON), A., i, 701.
- Rhammonic acid**, preparation of (KILIANI), A., i, 223.
- δ*-epi-l-Rhammonic acid** (*guleonic acid*), and its salts and derivatives (KILIANI), A., i, 1112.
- Rhamnus frangula*, constituents of the bark of (OESTERLE), A., i, 100.
- Rhinanthin**, comparison of aucubin and (BRIDEL and BRAECKE), A., i, 1168.
- Rhinanthus crista-galli*, aucubin and sucrose from seeds of (BRIDEL and BRAECKE), A., i, 1225.
- Rhinolith**, constituents of a (DEBUCQUET), A., i, 497.
- Rhizopus*, production of pectinase by (HARTER and WEIMER), A., i, 507.
- Rhizostoma cuvieri*, constituents of the gonads of (HAUROWITZ), A., i, 1210.
- Rhodanine** (GRÄNACHER, REIS, and POOL), A., i, 576; (GRANACHER), A., i, 849.
- Rhodanine**, α -oximino-, and its silver salt (GRÄNACHER, REIS, and POOL), A., i, 577.
- Rhodanine- α -acenaphthenequinone** (GRANACHER, REIS, and POOL), A., i, 577.
- Rhodium bases** (*rhodiumammynes*), solubilities of salts of (BRÖNSTED and PETERSEN), A., ii, 199.
- Rhodymenia palmata*, pentosan from (SAUVAGEAU and DENIGES), A., i, 507.
- Rhus glabra*, organic acids from (SANDO and BARTLETT), A., i, 100.
- Ribes rubrum* (currant), acids in (FRANZEN and SCHUMACHER), A., i, 310.
- Rice**, vitamin content of (FLEMING), A., i, 93.
- Ricinine**, constitution of (SPÄTH and TSCHERNITZ), A., i, 571.
- Ricnoleohydroxamic acid** (NICOLET and PELC), A., i, 644.
- Rings**, four-membered, additive formation of (INGOLD and PIGGOTT), T., 2793.
- Ringer's solution**, replacement of potassium salts in (ZWAARDEMAKER), A., i, 1213.
- Rocella fuciformis*, picrorocellin from (FORSTER and SAVILLE), T., 816.
- Rochelle salt** See Tartaric acid, potassium sodium salt.
- Rosaniline-sulphurous acid**, and its reactions with aldehydes (WIELAND and SCHEUING), A., i, 58.
- Roses**, red, colouring matter of (CURREY), A., i, 413.
- Roseocobaltic salts**. See under Cobalt.
- Rotation**, inversion of, in anisotropic liquids (ROYER), A., ii, 415.
 optical, experimental proof of theories of (WETTERFORS), A., ii, 247.
 of sugars (MALBY), T., 2608.
- Rotatory dispersion**. See Dispersion.
 power and chemical constitution (RUPE and JAGGI), A., i, 840; (RUPE), A., ii, 602
 of crystals (LONGCHAMON), A., ii, 603.
- Ruban** (RABE), A., i, 360.
- 9-Rubanone**, and its picrate (RABE, KINDLER, and WAGNER), A., i, 361.
- Rubatoxan** (RABE), A., i, 361.
- 9-Rubatoxanone**, and 8-bromo-, and their salts (RABE, KINDLER, and WAGNER), A., i, 361.
- Rubicene**, and its picrate, and dibromo-, and dinitro- (DZIEWOŃSKI and SUSZKO), A., i, 730.
- Rubicenedisulphonic acid** (DZIEWOŃSKI and SUSZKO), A., i, 731.
- Rubidium**, induction spectrum of (DUNOYER), A., ii, 729.
- Rubidium bromate**, preparation of (BUELL and McCROSKY), A., ii, 146.
 silver gold bromides (SUSCHNIG), A., ii, 514.
- Rubidium detection**:—
 detection of (MISSENDEN), A., ii, 658.
- Rubilic acid**, hexachloro-, and pentachlorohydroxy-, and their derivatives (KÜSTER and HERRMANN), A., i, 886.
- Rubus fruticosus* (blackberry), lactic acid in the leaves of (FRANZEN and KEYSNER), A., i, 310.
- Rubus idæus* (raspberry), ellagic acid in (KUNZ-KRAUSE), A., i, 210.
 occurrence of lactic acid and succinic acid in the leaves of (FRANZEN and STERN), A., i, 311, 975.
- Rutacarpine**, constitution of (ASAHI and FUJITA), A., i, 47.
- Ruthenium carbonyls** (MOND and WALLIS), T., 30.

- Ruthenium tetroxide** (KRAUSS), A., ii, 75.
 alkali sulphites (REMY), A., ii, 857.
Ruthenibromides (GUTHRIE and KRAUSS), A., i, 16.
Rye straw, lignin from (BECKMANN, LIESCHE, and LEHMANN), A., i, 233.

S.

- Sabellaria alveolata**, constituents of eggs of (FAURÉ-FREMIET), A., i, 85.
Sabinene, oxidation of, with chromyl chloride (HENDERSON, ROBERTSON, and BROWN), T., 2717.
Sabinenilaldehyde, and its semicarbazone (HENDERSON, ROBERTSON, and BROWN), T., 2720.
Sabinenilanic acid, and its salts (HENDERSON, ROBERTSON, and BROWN), T., 2720.
Saccharase, diffusion of solutions of (v. EULER and ERICSON), A., i, 886.
 acidity and thermostability of (v. EULER and MYRBACK), A., i, 693.
 removal of, from adsorbed substances (WILLSTATTER and KUHN), A., i, 283.
 specific nature of (WILLSTATTER and KUHN), A., i, 284.
 reactivation of, by dialysis (v. EULER and SVANBERG), A., i, 284.
 inactivation of, by iodine (v. EULER and LANDERGREEN), A., i, 1076.
 phosphorus content of preparations of (v. EULER and SVANBERG), A., i, 1200.
 inactivation of, by silver salts (v. EULER and MYRBACK), A., i, 959.
 silver compound of (v. EULER and JOSEPHSON), A., i, 1076.
 inversion of sucrose by (COLIN and CHAUDUN; CHAUDUN), A., i, 389.
 in blood and yeast (KNAFFL-LENZ), A., i, 694.
 in the intestine (v. EULER and SVANBERG), A., i, 296; (KNAFFL-LENZ), A., i, 435.
 analysis of (CANALS), A., i, 1075.
d-Saccharic acid, degradation of (BERGMANN), A., i, 7.
Saccharimeter, testing of the (KRAISY), A., ii, 233.
"Saccharin" (*o-benzoic sulphinide*), colouring matters from (DUTT), T., 2389.
 estimation of *p*-sulphaminobenzoic acid in (HERZOG and KREIDL), A., ii, 237.
C₁-Saccharinic acids (GLATTFIELD and SANDER), A., i, 318.
Saccharomyces cerevisiae, action of ultra-violet light on (DE FAZI), A., i, 1219.
Saccharomyces Marxianus, fermentation by (v. EULER and JOSEPHSON), A., i, 706.
Saccharophosphatase (TOMITA), A., i, 960.
 occurrence and action of, in plants (NĚMEC and DUCHOŇ), A., i, 206.
Saffron, constituents of (WINTERSTEIN and TELECZKY), A., i, 563.
 detection of the colouring matter of (GUERBET), A., ii, 793.
^{iso}**Safrole ozonide** (NAGAI), A., i, 839.
^{iso}**Safrolealkamines**, and their salts and derivatives, and bromo- (TAKEDA and KURODA), A., i, 273.
^{iso}**Safrole-2-imino-oxazolidine**, and bromo-, and their derivatives (TAKEDA and KURODA), A., i, 273.
Sakoe oil from Madagascar (JUMELLE), A., i, 210.
Salicin thiocyanate, tetra-acetyl derivative (ZEMPLĚN and HOFFMANN), A., i, 563.
Salicindithylamine (ZEMPLĚN and KUNZ), A., i, 564.
Salicinphenylmethylamine (ZEMPLĚN and KUNZ), A., i, 565.
Salicylaldehyde picrate, compound of thiocarbamide and (TAYLOR), T., 2269.
Salicylaldehyde, 5-amino-, and its derivatives (WEIL, TRAUN, and MARCEL), A., i, 1023.
Salicylatotetramminecobaltic salts (MORGAN and SMITH), T., 1956.
Salicylic acid, antipyrine salt, hydrolysis of (KOLTHOFF), A., i, 471.
 cobaltamine salt of (MORGAN and SMITH), T., 2874.
 sodium salt, compound of pyridine-betaine and (CASSELLA & Co.), A., i, 860.
 substituted derivatives of (H. P. and W. KAUFMANN), A., i, 252.
 5-chlorosulphonyl derivative (STEWART), T., 2259.
 estimation of, in blood-serum (HERISSEY), A., ii, 880.
Salicylic acid, 4-nitro- (KONDŌ, NAKAJIMA, and MURAKAWA), A., i, 745.
 5:5'-dithio-, and 5-thiol- (STEWART), T., 2259.
Salicylic acids, substituted, reduction of (WEIL and BRIMMER), A., i, 349; (WEIL, TRAUN, and MARCEL), A., i, 1023.
Salicylides, polymeric (PAOLINI and SCALBA), A., i, 746.
Salicylideneanthranilic acid, 3- and 5-nitro- (EKELEY, ROGERS, and SWISHER), A., i, 985.

- Salicylidene glucosamine** (IRVINE and EARL), T., 2378.
- 2-Salicylidene-1-methyl glucosamine** (IRVINE and EARL), T., 2379.
- 2-Salicylidene-3:5:6-triacetyl-1-ethyl glucosamine** (IRVINE and EARL), T., 2380.
- 2-Salicylidene-3:5:6-triacetyl glucosamine, 1-bromo-** (IRVINE and EARL), T., 2375.
- 2-Salicylidene-3:5:6-triacetyl-1-methyl glucosamine** (IRVINE and EARL), T., 2379.
- α -Salicylylidenedeoxybenzoïn.** See *trans*- α -Phenyl-*o*-coumarophenone.
- Saligenin, derivatives of** (HART and HIRSCHFELDER), A., i, 38.
- Saliva, relation of, to gastric juice** (NAKAGAWA), A., i, 739.
- human, hydrogen-ion concentration of (STARR), A., i, 1209.
- detection of bismuth in (GANASSINI), A., ii, 590.
- Salts, relations between temperature and the solubility, activity, and osmotic coefficients of** (BRONSTED), A., ii, 854.
- slow hydrolysis of (TIAN), A., ii, 362.
- complex, crystallography and optical properties of (KNAGGS), T., 2069.
- fused (HERZ), A., ii, 739.
- molecular refraction and dissociation of (MEYER and HECK), A., ii, 241.
- reactions in mixtures of (HICKS and CRAIG), A., ii, 622.
- triple (WELLS), A., ii, 560.
- of weak acids and weak bases, hydrolysis of (GRIFFITH), A., ii, 420.
- See also Metallic salts.
- Salt solutions, osmotic and activity functions in** (BRONSTED), A., ii, 482.
- equilibria of (LE CHATELIER), A., ii, 555.
- redissolution of a salt during evaporation of (RENGADE), A., ii, 31.
- Salvarsan (arsphenamine), relation between mode of synthesis and toxicity of** (CHRISTIANSEN), A., i, 186.
- colloidal properties of (RAIZISS and GAYRON), A., i, 1202.
- compounds of aldehydes with (RAIZISS and BLATT), A., i, 1079.
- sulphur content of, and its relation to toxicity (CHRISTIANSEN), A., i, 601, 1202.
- Samarium oxide, density and molecular volume of** (FRANDTL), A., ii, 379.
- Samarakite** (SHIPATA and KIMURA), A., ii, 516.
- Sandal wood oil, Dutch East Indies, preparation and properties of** (ROJDESTWENSKY), A., ii, 792.
- Sand spur.** See *Cenchrus tribuloides*.
- Santal oil, estimation of santalol in** (HARRISON), A., ii, 400.
- Santalol, estimation of, in santal oil** (HARRISON), A., ii, 400.
- Saponins** (VAN DER HAAR), A., i, 160, 565, 1168; (VAN DER HAAR and TAMBURELLO), A., i, 160.
- and their derivatives, effect of alkali or bromine on the biological activity of (SIEBURG and BACHMANN), A., i, 404.
- surface activity and toxic action of (KOFLER), A., i, 610.
- differentiation and estimation of (KOFLER), A., ii, 595.
- Sativic acids** (NICOLET and COX), A., i, 320; (REINGER), A., i, 623.
- Saturra montana*, constituents of oil from (LEONE and ANGELESCU), A., i, 357.
- Sauerkraut, fermentation of** (BRUNKOW, PETERSON, and FRED), A., i, 312.
- Scandium, aic spectrum of** (KING), A., ii, 100.
- extraction and purification of, from thorveitite (P. and G. URBAIN), A., ii, 504.
- Scatole, toxicity of** (SALANT and KLEITMAN), A., i, 794.
- Scheele's green, composition of** (BORNE-MANN), A., ii, 852.
- Scheih oil, constituents of** (ROURE-BERTRAND FILS), A., i, 845.
- Schiff's reagent, modified** (WERTHEIM), A., ii, 793.
- Schizoneura lanigera* (blood-louse), colouring matter and wax of (SCHULZ), A., i, 494.
- Schizophycæ, water-soluble colouring matters from** (BOKESCH), A., i, 210.
- Schweinfurth green, estimation of arsenic and copper in** (KOLTHOFF and CREMER), A., ii, 86.
- Scopolamine, constitution of** (HESS and WAHL), A., i, 854.
- Scopoline** (HESS and WAHL), A., i, 854.
- Scorpion venom.** See Poison.
- Scrophulariaceæ, sugars and glucosides in the** (BRAECKE), A., i, 1225.
- Sea urohin, constituents of** (TAKAHASHI), A., i, 609.
- Sea water.** See under Water.
- Sea-wolf liver oil** (LEXOW), A., i, 195.
- Sebacic acid, ethyl and ethyl hydrogen esters** (GRÜN and WIRTH), A., i, 805.
- Sebacic dialdehyde.** See Decanedial.
- Secretin** (VAN EWEYK and TENNENBAUM), A., i, 397.

- Sedimentation analysis, apparatus for** (F. V. and D. V. HAHN), A., ii, 705.
- Sedum telephium*, glucoside in the stems and roots of (BRIDEL), A., i, 799, 1225.
- Seeds, evaluation of the vitality of** (NĚMEC and DUCHOŇ), A., i, 94, 411.
- influence of calcium on the germination of (MAQUENNE and DEMOUSSY), A., i, 905.
- effect of selenium and radium on the germination of (STOKLASA), A., i, 618.
- Dutch, manganese in (WESTER), A., i, 93.
- "Seid" oil, constituents of (JOSEPH and WHITEFIELD), A., i, 754.
- Selachyl alcohol** (TSUJIMOTO and TOYAMA), A., i, 297.
- Selenatopentamminecobalti-salts.** See under Cobalt.
- Selenious acid.** See under Selenium.
- Selenium, constitution of** (PĚLABON), A., ii, 141.
- isotopes of (ASTON), A., ii, 842.
- colloidal, and the action of gelatin on its solutions (GUTBIER and EMSLANDER), A., ii, 625.
- effect of freezing on (GUTBIER, HEINRICH, and HUBER), A., ii, 142; (GUTBIER and EMSLANDER), A., ii, 283.
- action of, on gold (PĚLABON), A., ii, 303.
- Selenium compounds, action of, on the growth of plants** (STOKLASA), A., i, 614.
- effect of, on the germination of seeds (STOKLASA), A., i, 613.
- pharmacology of (JOACHIMOGLU and HIROSE), A., i, 396, 406.
- Selenium salts, action of, on growth of plants** (TURINA), A., i, 707.
- action of, on plant metabolism in presence of radioactive air and soil (STOKLASA, KŘIČKA, PĚNKAVA, ZELENKA, CHMELÁŘ, and JÁNSKÝ), A., i, 974.
- Selenium tetrachloride, action of, on amylene, butylene, and propylene** (BOORD and COPE), A., i, 421.
- action of, with β -diketones (MORGAN, DREW, and BARKER), T., 2433.
- dioxide, preparation of (MEYER), A., ii, 639.
- hydrates of (MANCHOT and ORTNER), A., ii, 283.
- oxybromide (LENHER), A., ii, 707.
- oxychloride, properties of (LENHER), A., ii, 706.
- Selenium oxychloride, vapour pressure of** (LENHER, SMITH, and TOWN), A., ii, 371.
- Selenates, reflection spectra of** (SCHAEFER and SCHUBERT), A., ii, 179.
- complex (MAYER), A., ii, 70.
- double (TUTTON), A., ii, 502, 505.
- Selenious acid, and its compounds with molybdenum and vanadium** (ROSENHEIM and KRAUSE), A., ii, 47.
- Selenides, preparation of** (MOSER and DOCTOR), A., ii, 46.
- Selenium organic compounds** (BOGERT and CHEN), A., i, 1182.
- aromatic (FARBWERKE FORM. MEISTER, LUGIUS, & BRÜNING), A., i, 1066.
- acetylacetones, and cyano- (MORGAN, DREW, and BARKER), T., 2443.
- benzoylacetones, and cyano- (MORGAN, DREW, and BARKER), T., 2453.
- dibenzoylmethanes, and cyano- (MORGAN, DREW, and BARKER), T., 2464.
- C-ethylacetylacetones (MORGAN, DREW, and BARKER), T., 2460.
- Selenic acid, salts and esters of** (MEYER and WAGNER), A., i, 620.
- Selenium detection and estimation:—**
detection and estimation of (MÜLLER, MENZEL, and SCHUBERT), A., ii, 390.
- estimation of (LOSANA), A., ii, 865.
- Selenium cells, mode of action and equilibrium of** (SPÄTH), A., ii, 189.
- 4-Seleno-2-methylquinazalone** (BOGERT and CHEN), A., i, 1183.
- Semicarbazide hydrochloride** (OCHI), A., i, 723.
- Semicarbazidedithiocarboxylic acid, thio-, potassium salt** (GUHA), A., i, 877.
- Semicarbasido-*dl*-piperitone, and its hydrochloride** (READ and SMITH), T., 1869.
- Semicarbazones, action of amines on** (WILSON, HOPPER, and CRAWFORD), T., 366.
- thio-, reactions of (WILSON and BURNS), T., 870.
- Semina cardui Maris, tyramine from** (ULLMANN), A., i, 616.
- Serum, ionic distribution in** (REICHEI), A., i, 485.
- viscosity and velocity of ultrafiltration of (ELLINGER and NEUSCHLOSZ), A., i, 484.
- proteins of (PIETTRE and VILA), A., i, 63.

- Serum**, condition of sugar in (RUSZYŃSKI and HETÉNYI), A., i, 291.
 action of kaolin on the tributyrinase in (OLSEN), A., i, 292.
 horse, separation of globulins of (VILA), A., i, 1209.
 human, estimation of bilirubin in (THANNHAUSER and ANDERSEN), A., ii, 671.
 detection and estimation of nitrates in (NOLTE), A., ii, 583.
 estimation of non-protein nitrogen in (CRISTOL), A., ii, 583; (CRISTOL and SIMONNET), A., ii, 887.
 estimation of inorganic phosphorus in (TISDALL), A., ii, 392.
- Serum albumin**. See under Albumin.
Serum globulin. See under Globulin.
- Sesquiterpenes**, chemistry of (RUZICKA, MEYER, and MINGAZZINI), A., i, 560.
- Shad**. See *Clupea alosa*.
- Shaking machine**, for large quantities of precipitates (FRANZEN), A., ii, 759.
- Shark liver oil**, constituents of (TSUJIMOTO and TOYAMA), A., i, 297; (TOYAMA), A., i, 895.
- Shark oil**, catalytic decomposition of (MAILHE), A., i, 424.
- Shikazarin** (MAJIMA and KURODA), A., i, 946.
- Shikon**. See *Lithospermum erythrorhizon*.
- Shikonin**, constitution and derivatives of (MAJIMA and KURODA), A., i, 946.
- o*-**Sialosinolic acid**, oxidation of (ZINKE, HANSELMAYER, and EHMER), A., i, 668.
- Silica**. See Silicon dioxide.
- Silicane**. See Silicon tetrahydride.
- Silicanetetra-1-piperidinium chloride** (LEONARD), A., i, 363.
- Silicon**, modifications of (MANCHOT), A., ii, 144; (MANCHOT and FUNK), A., ii, 286, 764.
- Silicon compounds**, unsaturated, luminescence of (KAUTSKY and ZOCHER), A., ii, 464.
- Silicon tetrachloride**, thermal constants of (LATIMER), A., ii, 256.
- tetrahydride (silicane)**, preparation of (SCHWARZ and KONRAD), A., ii, 846.
 viscosity and molecular dimensions of (RANKINE and SMITH), A., ii, 709.
- dioxide (silica)**, reflection spectra of (SCHAEFER and SCHUBERT), A., ii, 179.
 heat of wetting of gels of (PATRICK and GRIM), A., ii, 122.
 catalytic activity of (GILFILLAN), A., i, 709.
- Silicon dioxide (silica)**, adsorption of ammonia by gels of (DAVIDHEISER and PATRICK), A., ii, 262.
 action of sodium chloride with (CLEWS and THOMPSON), T., 1442.
 crucibles. See Crucibles.
 estimation of (MURMANN), A., ii, 226.
 estimation of, in filtered sea-water (WELLS), A., ii, 868.
- Silicic acid**, colloidal, preparation of (KRÖGER), A., ii, 212, 213; (BRADFELD), A., ii, 507.
 electrical properties of (LÖSENBECK), A., ii, 695.
 analysis of gels of (NEUHAUSEN and PATRICK), A., ii, 144.
- Silicates**, ultra-red spectra of (SCHAEFER and SCHUBERT), A., ii, 727.
 colloidal, in soils, effect of alum on (SCOFIELD), A., i, 212.
- Silicate rocks**, estimation of arsenic in (HACKL), A., ii, 159.
 estimation of cobalt and nickel in (HACKL), A., ii, 458.
- Silk**, action of iodine on (HUEBNER and SINHA), A., i, 434.
 artificial, manufacture of (HESS), A., i, 922.
- Silk fibroin**, composition and structure of (ABDERHALDEN), A., i, 784.
- Silver**, Röntgen-ray spectrum of (DE BROGLIE), A., ii, 330.
 absorption of K-radiation from, in gases (BURBIDGE), A., ii, 184.
 photoelectric wave-length of (STUHLMAN), A., ii, 809.
 electrolytic potential of (KOLTHOFF), A., ii, 20.
 purity of, for use in atomic weight determinations (BAXTER and WOODMAN), A., ii, 376; (BAXTER), A., ii, 377.
 oligodynamy of (DOERR and BERGER), A., i, 1097.
- colloidal** (GUTBIER, HUBER, and ZWEIFLER), A., ii, 485.
 colours of (SCHAUM and MARX), A., ii, 696.
 carrageen as a protective colloid for (GUTBIER, WOLF, and KIESS), A., ii, 203.
 sols of Carey Lea, behaviour of, towards electrolytes and hydrophilic colloids (FREUNDLICH and LOENING), A., ii, 696.
- Silver alloys with mercury**, preparation of (MÜLLER and HÖNIG), A., ii, 500.
- Silver compounds**, photochemistry of (WEIGERT and SCHÖLLER), A., ii, 10.

Silver salts, use of ammoniacal solutions of, in valency determinations (COLLENBERG), A., ii, 495.

Silver autimonide, velocity of solution of, in silver (WEISS and HENRY), A., ii, 207.

arsenate, compound of acetylene with (OBERDOERFER and NIEUWLAND), A., i, 515.

bromate, preparation and properties of (REEDY), A., ii, 56.

bromide, action of light on (HARTUNG), T., 682.

effect of colloids on the photochemical decomposition of (SCHWARZ and STOCK), A., ii, 731.

gold bromides (SUSCHNIG), A., ii, 514.

chloride, solubility of, in solutions of chlorides (FORBES and COLE), A., ii, 291.

ammonium auric chloride (WELLS), A., ii, 449.

perchlorate, equilibrium in the system, benzene, water, and (HILL), A., ii, 555.

chromate, formation of, in Liesegang's rings (McGUIGAN), A., ii, 38.

solvolytic (HICKS and CRAIG), A., ii, 622.

haloids, action of light on (KOCH and SCHRADER), A., ii, 182.

chlorination of, in Gooch crucibles (MELLON and SIEGISMUND), A., ii, 781.

action of, on potassium ferrocyanide (BONINO), A., ii, 78.

iodate, equilibrium of potassium chromate with (RAMANN and SALLINGER), A., ii, 131.

iodide, heat of formation of (TAYLOR and ANDERSON), A., ii, 121.

adsorption of iodine by (GERMANN and TRAXLER), A., ii, 371.

permanganate, velocity of decomposition of (SIEVERTS and THEBERATH), A., ii, 360.

molybdate, crystal structure of (WYCKOFF), A., ii, 765.

nitrate, electrolysis of pyridine solutions of (MÜLLER; MÜLLER and DUSCHEK), A., ii, 612.

oxide, crystal structure of (WYCKOFF), A., ii, 291.

equilibrium of oxygen and (KEYES and HARA), A., ii, 353.

catalytic influence of oxides on the decomposition of (KENDALL and FUCHS), A., ii, 147.

phosphate, compound of acetylene with (OBERDOERFER and NIEUWLAND), A., i, 515.

Silver phosphide (MOSE and BRUKL), A., ii, 393.

sulphate, action of, on alkyl haloids, in sulphuric acid solution (SCHILOV), A., i, 918.

thiosulphates, alkaline, compounds of ammonia with (JONSSON), A., ii, 57.

Silver organic compounds:—

Silver-albumose, estimation of (HERZOG), A., ii, 798.

Silver-thioglycollic acid, and its sodium salt (CHEMISCHE FABRIK FLORA), A., i, 425.

Silver estimation and separation:—

estimation of, in presence of colloidal silver (GUTHRIE, HUBER, and KUPPINGER), A., ii, 396.

separation of, from mercurous salts (KOLTHOFF), A., ii, 160.

Silver anode. See Anode.

Silver ores from Nevada (SHANNON), A., ii, 859.

Sincosite (SCHALLER), A., ii, 459.

Slags, dephosphoration, constituents of (DEMOLON), A., ii, 564.

Sneezeweed. See *Helenium hoopesii*.

Snow, analyses of (SCHAFER), A., i, 512.

Soaps, colloid chemistry and manufacture of (FISCHER, McLAUGHLIN, and HOOKER), A., ii, 430, 829.

effect of addition of salt on the viscosity of solutions of (KING), A., i, 621.

adsorption of, by wool (BRITISH RESEARCH ASSOCIATION FOR THE WOOLLEN AND WORSTED INDUSTRIES), A., ii, 551.

action of blood-serum with (JARISCH), A., i, 1087.

estimation of, in wool (BRITISH RESEARCH ASSOCIATION FOR THE WOOLLEN AND WORSTED INDUSTRIES), A., ii, 594.

Soap solutions, preparation and hydro-metric precision of (JUSTIN-MUELLER), A., ii, 658.

constitution of (McBAIN; TAYLOR and LAING), T., 621; (FLECKER and TAYLOR), T., 1101; (NORRIS), T., 2161.

effect of electrolytes on the constitution of (SALMON), T., 711; (McBAIN and BURNETT), T., 1320.

detergent power of (MEES), A., ii, 268.

ultra-filtration of (McBAIN and JENKINS), T., 2325.

Sodamide, syntheses by means of (HALLER and BAUER), A., i, 258; (HALLER and BENOIST), A., i, 350; (HALLER and BOUDIN), A., i, 356.

Sodammonium, action of, on hydrocarbons (LEBEAU and PICON), A., i, 801.

action of, with hexamethylenetetramine, tetramethyldiaminomethane, and ethylidene-ethylamine (PICON), A., i, 1128.

compound of pyridine with (LEBEAU and PICON), A., i, 48.

Soddlite, radioactive (SCHOEP), A., ii, 451.

Sodium, bright metallic, preparation of, in nitrogen (BORNEMANN), A., ii, 444.

absorption spectrum of (LADENBURG), A., ii, 6; (HARRISON), A., ii, 679.

arc spectrum of (FOOTE, MEGGERS, and MOHLER), A., ii, 598.

influence of the pressure of foreign gases on the spectrum of (MINKOWSKI), A., ii, 242.

distribution of electrons in the atom of (BRAGG, JAMES, and BOSANQUET), A., ii, 703.

chemical constants of (LADENBURG and MINKOWSKI), A., ii, 191.

vapour, electrodeless discharge in (ROBERTSON), A., ii, 609.

electrical conductivity of, in liquid ammonia (KRAUS and LUCASSE), A., ii, 252.

ions, electrolytic migration of, through glass (PIRANI and LAX), A., ii, 817.

heat of vaporisation of (LADENBURG and MINKOWSKI), A., ii, 194.

resistance temperature coefficients of solutions of, in liquid ammonia (KRAUS and LUCASSE), A., ii, 734.

equilibrium of mixtures of ammonia and (KRAUS and LUCASSE), A., ii, 764.

equilibrium of mixtures of tellurium and, in liquid ammonia (KRAUS and CHIU), A., ii, 765.

action of, with bromo-derivatives of benzene and toluene (FUCHS and METZL), A., i, 442.

Sodium alloys with mercury, electrochemistry of (RICHARDS and CONANT), A., ii, 840.

Sodium salts, distribution of, in blood (KRAMER and TISDALL), A., i, 1087.

Sodium antimoniodobromide (VOURNAZOS), A., ii, 651.

arsenite, action of derivatives of nitric oxide and hydroxyl on (GUTMANN), A., ii, 844.

azide, action of, with benzylidene-benzhydrazide chloride and dibenzhydrazide chloride (STOLLÉ and NETZ), A., i, 690.

Sodium perborate, electrolytic preparation of (ALSGAARD), A., ii, 375; (ABENDT and HANTGE), A., ii, 569.

bromate and chlorate, crystal structures of (DICKENSON and GOODHUE), A., ii, 145.

carbonate, manufacture of, by the ammonia process (LE CHATELIER), A., ii, 375.

velocity of absorption of carbon dioxide by (RIOU), A., ii, 438.

equilibrium of calcium sulphate and (SYRKIN), A., ii, 699.

equilibrium in the system, sodium chloride, water, and (FREETH), A., ii, 627.

equilibrium of sodium sulphate, water, and (DAWKINS), T., 776.

reaction between nitrogen, carbon and (INGOLD and WILSON), T., 2278.

minerals from Kenya Colony (WALTHER), A., ii, 859.

hydrogen carbonate, preparation of (TOPORESCU), A., ii, 375, 642.

chloride, effect of albumin on the conductivity of solutions of (PALMER, ATCHLEY, and LOEB), A., i, 692.

melting and freezing points of (FERGUSON), A., ii, 845.

diffusion of (STILES and ADAIR), A., ii, 125.

equilibrium of ammonium nitrate and (PERMAN), T., 2473.

equilibrium in the system, sodium carbonate, water, and (FREETH), A., ii, 627.

equilibrium of sodium laurate, water, and (MCBAIN and BURNETT), T., 1320.

blue flame produced by, on a coal fire (SMITHELLS), A., ii, 645.

reactions of cellulose with (MASTERS), T., 2026.

compounds of pyridinebetaine and (CASSELLA & Co.), A., i, 860.

action of silica with (CLEWS and THOMPSON), T., 1442.

chloride and sulphate, equilibrium of, with ammonium chloride and sulphate and water (RIVETT), T., 379.

chlorite (LEVI), A., ii, 567.

chromate, mixed crystals of sodium sulphate and (RICHARDS and MELDRUM), A., ii, 54.

hydroxide, preparation of, free from carbon dioxide (CORNOG), A., ii, 288.

heats of neutralisation of (RICHARDS and ROWE), A., ii, 425.

Sodium hydroxide, heat of neutralisation of, with hydrochloric acid (KEYES, GILLESPIE, and MITSUKURI), A., ii, 424.
 reactions of, with aluminium salts (GROBER), A., ii, 573.
 hypochlorite, red colour of solutions of (MARIO), A., ii, 457.
 hyposulphite (HEYL and GREER), A., ii, 288.
 iodide, conductivity of, in amyl alcohol (KRAUS and BISHOP), A., ii, 813.
 nitrate, equilibrium of ammonium nitrate with (EARLY and LOWRY), T., 963.
 equilibrium of magnesium nitrate, water and (JACKMAN and BROWNE), T., 694.
 peroxide, estimation of the active oxygen in (MILBAUER), A., ii, 521.
 phosphate, action of epichlorohydrin on (BAILLY), A., i, 980.
 dihydrogen phosphate, effect of, on body power (HERZHEIMER), A., i, 970.
 silicate (ERDENBRECHER), A., ii, 444.
 electrolysis of solutions of (SPENCER and PROUD), A., ii, 611.
 equilibrium of sodium tungstate with (VAN LIEMPT), A., ii, 775.
 silicofluoride, refractive index of (RATTEI), A., ii, 541.
 sulphate, equilibrium of sodium carbonate, water and (DAWKINS), T., 776.
 mixed crystals of sodium chromate and (RICHARDS and MELDRUM), A., ii, 54.
 sulphide, use of, instead of hydrogen sulphide in qualitative analysis (VORTMANN), A., ii, 653.
 sulphite, alkylation of (BAGGESGAARD-KASMUSSEN and WERNER), A., i, 104.
 hydrogen sulphite, action of, on nitro-compounds (WEIL and MOSER), A., i, 443.
 tungstates (SMITH), A., ii, 774.
 equilibrium of, with sodium silicate and with potassium tungstate (VAN LIEMPT), A., ii, 775.
 for use in blood analysis (FOLIN), A., ii, 596.
 paratungstate, preparation of (LOT-TERMOSER), A., ii, 510.
Sodium estimation:—
 estimation of, in aluminium and alumina (GRITH), A., ii, 720.
 estimation of, in tinned dishes (WILSON), A., ii, 395.

Sodium amalgam electrode. See Electrodes.
Sodium lamp. See Lamp.
Sodium press, laboratory (BECKMANN), A., ii, 369.
Soils, absorbent power of (CASALE), A., i, 508.
 absorption and basic exchange in (v. NOSTITZ), A., i, 511.
 absorption of ammonium salts from solutions by (AARNIO), A., i, 1227.
 absorption of water by colloids in (ROBINSON), A., i, 1228.
 flocculation of (COMBER), A., i, 212.
 precipitation of the constituents of, by calcium salts in presence of hydroxyl ions (MATTSON), A., i, 800.
 acidity of (KÖNIG, HASENBÄUMER and KRÖGER; LEMMERMAN and FRESSENIUS), A., i, 510.
 effect of lime, phosphates, etc., on (BAUER and HAAS), A., i, 975.
 reaction of, and its relation to plant distribution (ATKINS), A., i, 415, 509; (COMBER), A., i, 416; (FISHER), A., i, 510; (SALISBURY), A., i, 1104.
 relation of hydrogen-ion concentration in, to their lime requirements (JOHNSON), A., i, 708.
 effect of aluminium salts on ammonification and nitrification in (DENISON), A., i, 512.
 carbon-nitrogen ratio in (READ), A., i, 416.
 origin of colloids in (WHITNEY), A., i, 708; (GORDON), A., i, 1227.
 effect of salts on nitrogen fixation in (GREAVES, CARTER, and LUND), A., i, 976.
 organic nitrogen in, availability of (ROBINSON, WINTER, and MILLER), A., i, 212.
 phosphoric acid in (FRAPS), A., i, 616.
 effect of colloids on the availability of salts in (GORDON and STARKEY), A., i, 1104.
 effect of alum on colloidal silicates in (SCOFFIELD), A., i, 212.
 oxidation of sulphur compounds in (LIPMAN, WAKSMAN, and JOFFE), A., i, 303; (DEMOLON), A., i, 312; (WAKSMAN), A., i, 706.
 oxidation of sulphur and transformation of phosphates in (WAKSMAN and JOFFE), A., i, 416.
 sulphate reducing organism in (KÜHR), A., i, 1228.
 effect of drying, on the water-soluble constituents of (GUSTAFSON), A., i, 708.

- Soils**, classification of water in (PARKER), A., i, 616.
 toxicity, acidity and basicity of (CARR), A., ii, 172.
 acid, influence of calcium salts on (ROBINSON and BULLIS), A., i, 976.
 nitrification in (STEPHENSON), A., i, 416.
 arable, cobalt and nickel in (BERTRAND and MOKRAGNATZ), A., i, 975.
 humus, mechanical analysis of (ROBINSON), A., ii, 888.
 mechanical analysis of (ROBINSON), A., ii, 888.
 analysis of, from the formation of layers (UNGEREK), A., ii, 96.
 detection of acidity of (HUDIG and HETTERSCHIJ), A., i, 1104.
 estimation of moisture in (DEIGHTON), A., i, 1227.
 estimation of calcium in (SHEDD), A., ii, 527.
 estimation of hydrogen-ion concentration in (HEALY and KARRAKER), A., ii, 519.
 estimation of organic matter in (READ and RIDGELL), A., ii, 540.
 estimation of potassium in (JONES and REEDER), A., ii, 85.
 estimation of sulphates in (HIRST and GRAVES), A., ii, 521.
Soil solutions, method of obtaining (PARKER), A., i, 511.
Soja beans, culture of, and the occurrence of urease in these plants (WESTER), A., i, 311.
 nitrogen distribution of proteins from (FRIEDEMANN), A., i, 505.
Solannellie acid (WIELAND and SCHULENBURG), A., i, 346.
Solanum esculentum. See Tomato.
Solar atmosphere, ionisation in the (RUSSELL), A., ii, 675.
Solids, structure of (THOMSON), A., ii, 355, 745.
 incandescent, luminescence of (NICHOLS and HOWES), A., ii, 597.
 specific heat of (KRASE), A., ii, 421.
 hardness of, and its relation to constitution (REIS and ZIMMERMANN), A., ii, 745.
 interpenetration of, by chemical reaction (WEISS and HENRY), A., ii, 207, 487.
 velocity of chemical reactions in (HINSHELWOOD and BOWEN), A., ii, 628.
 extraction of (CHARITSCHKOV), A., ii, 826.
- Sols**, hydrophobic, action of, with hydrophilic colloids (FREUNDLICH and LOENING), A., ii, 356.
 lyophile, capillary-electric phenomena in (KRUYT and DE JONG), A., ii, 357.
 mastic, precipitation of (MICHAELIS and HIRABAYASHI), A., ii, 429.
Solubility (HILDEBRAND and JENKS), A., ii, 141; (BRÖNSTED and PETERSEN), A., ii, 199; (BRÖNSTED), A., ii, 481; (EPHRAIM and MOSIMANN), A., ii, 574.
 determination of (DUNDON and HENDERSON), A., ii, 552.
 calculation of (MORTIMER), A., ii, 621.
 partition in (SMITH), A., ii, 430, 431.
 prediction of, in polar solutions (KENDALL, DAVIDSON, and ADLER), A., ii, 34.
 of a compound in presence of other compounds (LEONE and ANGEL-ESCU), A., ii, 743; (LEONE and BENELLI), A., ii, 744.
 of gases in liquids (NEUHAUSEN and PATRICK), A., ii, 264.
 reciprocal, of slightly miscible liquids (CHÉNEVEAU), A., ii, 355, 427.
 in mixed solvents (PUCHER and DEHN), A., ii, 126.
 of substances in two solvents (SCHILOV, LEPIN, and JANTSCHAK), A., ii, 824.
 of substances in mixtures of alcohol and water (WRIGHT), T., 2251.
Solutions, theory of (CASSEL), A., ii, 481; (HERRMANN), A., ii, 552.
 optical properties of (WABASTJERNA), A., ii, 2.
 absorption of light by (LUNELUND), A., ii, 8; (v. HALBAN and SIEDEN-TOFF), A., ii, 332.
 additive compounds in, and ionisation (KENDALL and GROSS), A., ii, 32.
 thermochemistry of (LEVALT-EZER-SKY), A., ii, 819.
 adhesion in (SCHILOV and others), A., ii, 350; (SCHILOV, LEPIN, JANTSCHAK, and DUBNIN), A., ii, 626.
 adsorption of (OSTWALD and DE IZAGUIRRE), A., ii, 480.
 concentrated, theory of (TIMMERMANS), A., ii, 25.
 dilute aqueous, application of the ideal equation to (KENDALL), A., ii, 32.
 of metallic salts, crystalline particles in (TRAUBE and KLEIN), A., ii, 201.
 non-aqueous, electrochemistry of (MÜLLER; MÜLLER and DUSCHKE), A., ii, 612.
 of binary electrolytes, physical properties of (WALDEN), A., ii, 345.

- Solutions, non-aqueous, analysis of, with the interferometer** (COHEN and BRUNS), A., ii, 77.
 dispersoid analysis of (OSTWALD), A., ii, 198.
- Solvates, theory of** (SYRKIN), A., ii, 823.
- Solvatochromism** (HANTZSCH), A., i, 556.
- Solvents, partition of substances between** (SCHILOV, LEPIN, JANTSCHAK, and DUBININ), A., ii, 626; (SCHILOV, LEPIN, and JANTSCHAK), A., ii, 824.
 recovery of, in extraction processes (PICHLER), A., ii, 715.
 influence of, on the velocity of formation of quaternary ammonium compounds (HAWKINS), T., 1170.
 mixed, solubility in (PUCHER and DEHN), A., ii, 126.
 velocity of reaction in (CASHMORE, McCOMBIE, and SCARBOROUGH), T., 243; (McCOMBIE, SCARBOROUGH, and SETTLE), T., 2308.
- Solvolytic mixtures of fused salts** (HICKS and CRAIG), A., ii, 622.
- Sorbic acid, piperidide of** (OTT and ZIMMERMANN), A., i, 138.
- Sorghum vulgare*, protein from the caryopsis of (VISCO), A., i, 211.
- Sound, velocity of, in air and hydrogen** (GRÜNEISEN and MERKEL), A., ii, 190.
- Specific gravity.** See Density.
- Spectra in relation to atomic structure** (KING), A., ii, 277.
 produced by collision with electrons (SEELIGER), A., ii, 17.
 broadening of lines in (FRANCK), A., ii, 241; (FÜCHTBAUER and JOOS), A., ii, 242.
 of gases, excitation of, during chemical reactions (HABER and ZISCH), A., ii, 461.
 absorption, of elements barium to antimony (LINDSAY), A., ii, 599.
 of bromomalonics derivatives and nitroparaffins (GRAHAM and MACBETH), T., 1109.
 of halogen derivatives of cyclic compounds (GRAHAM and MACBETH), T., 2601.
 of metallic salts (GEORGE), A., ii, 806.
 arc (ST. JOHN and BABCOCK), A., ii, 4.
 intensity-current curves of (MOORE), A., ii, 242.
 occurrence of spark lines in (HEMSALECH and DE GRAMONT), A., ii, 410.
- Spectra, arc, values of constants in** (FOOTE and MOHLER), A., ii, 410.
 wave-lengths in (WALTERS), A., ii, 100.
 arc-cathode (DUNSTAN and WOOTEN), A., ii, 99.
 band, origin of (TAKAHASHI), A., ii, 3.
 regularities in (KRATZER), A., ii, 409.
 of isotopes (GREEE and KONEN), A., ii, 4.
 line, production of enhanced (SAWYER and BECKER), A., ii, 242.
 quantum theory of (BOHR), A., ii, 801.
 β -ray (ELLIS), A., ii, 339.
 and their connexion with γ -rays (MEITNER), A., ii, 732.
 from radioactive substances (MEITNER), A., ii, 416.
 β - and γ -ray (ELLIS), A., ii, 466.
 Röntgen ray (DUANE), A., ii, 104; (RICHTMYER), A., ii, 105, 804; (HJALMAR; COSTER), A., ii, 180; (SMEKAL), A., ii, 15, 181, 600, 607; (DOLEJSEK; DUANE and PATTERSON), A., ii, 463; (DUANE and FRICKE), A., ii, 804.
 precise measurements of (SIEGBAHN), A., ii, 104.
 spark lines in (WENTZEL), A., ii, 249.
 and the structure of atoms (COSTER), A., ii, 491, 677; (DAUVILLIER), A., ii, 678.
 in relation to valency (WENTZEL), A., ii, 607.
 K-series, of the light elements (DOLEJSEK; DAUVILLIER), A., ii, 243.
 L-series (COSTER), A., ii, 244, 462; (DAUVILLIER), A., ii, 463.
 N-series in (DOLEJSEK), A., ii, 411.
 of metallic oxides (HEDVAL), A., ii, 300.
 of organic crystalline compounds (BRAGG; BECKER and JANCKE), A., ii, 128.
 ultra-violet (KURTH), A., ii, 410.
 series, difference between, of isotopes (EHRENFEST; BOHR), A., ii, 598; (NICHOLSON), A., ii, 599.
 Balmer series, structure of (GEHRCKE and LAU), A., ii, 3.
 spark, identification of air lines in (MERRILL, HOPPER, and KEITH), A., ii, 802.
 ultra-red (SCHAEFER), A., ii, 727.
 ultra-violet, absorption (WINTHER, BAGGESGAARD-RASMUSSEN, and SCHREINER), A., ii, 729.
 of dihydric phenols (KLINGSTEDT), A., ii, 680.

- Spectra** vacuum spark, of metals (CARTER), A., ii, 599.
 visible, doublets in (GOUDSMIT), A., ii, 462.
- Spectrograph**, mass (ASTON and FOWLER), A., ii, 241.
- Röntgen ray** (SEEMANN), A., ii, 15.
- Spectrography** (v. ANGERER), A., ii, 3.
- Spectrophotometer**, double slit (NARAYAN and SUBRAHMANYAN), A., ii, 329.
- Spectroscopy**, new apparatus for affording source of light for (GERLACH and KOCH), A., ii, 330.
- Spectrum**, ultra-violet, extension of the (MILLIKAN), A., ii, 100.
 continuous, emission and absorption of (DUCLAUX), A., ii, 99.
- Spectrum lines**, errors in the measurement of (MERTON and HARRISON), A., ii, 673.
- Spermatozoa**, histochemistry of (STEUDER), A., i, 294.
- Sphingomyelin**, optical properties of (SANO), A., i, 701.
- Splianthol** from paracress (ASAHINA and ASANO), A., i, 505.
- Spirans** (LEUCHS, v. KATINSKY, and CONRAD), A., i, 471, 873.
- Spleen**, function of the (ASHER and BERNET), A., i, 491.
 effect of extirpation of, on blood-corpuscles (ASHER and MATSUMO), A., i, 298.
 formation of bilirubin in the (ERNST and SZAPPANYOS), A., i, 1089.
 of cattle, nucleic acids from (STEUDEL), A., i, 297.
- Spruce**, Swedish, constituents and properties of (WAHLBERG), A., i, 1101.
- Spruce needles**, resins and tannins from (v. EULER), A., i, 233.
- Stains**, detection of urine in (JEMMA), A., ii, 460.
- Stannated-1-piperidinium chloride**, *di-chloro* (LEONARD), A., i, 363.
- Stannic acid and salts**. See under Tin.
- Stannous salts**. See under Tin.
- Staphylolysin**, effect of metallic salts on the formation of (WALBUM), A., i, 795.
- Star anise oil**, constants of (GATTE-FOSSE), A., i, 1167.
- Starch** (ZWIKKER), A., i, 10.
 constitution of (DE HOOP), A., i, 434.
 measurement of the liquefaction of (OLSSON), A., ii, 401.
 adsorption in solutions of, and their emulsifying action (CLARK and MANN), A., ii, 550.
- Starch**, preservation of solutions of (KANÖ), A., i, 280.
 chemistry of (PRINGSHEIM and PERSCH), A., i, 113, 632; (PRINGSHEIM and DERNIKOS), A., i, 632; (PRINGSHEIM and GOLDSTEIN), A., i, 633.
 decomposition products of (PICTET and JAHN), A., i, 987.
 temperature coefficients in the degradation of (ERNSTRÖM), A., i, 598.
 effect of amino-acids on the hydrolysis of, by enzymes (SHERMAN and WALKER; SHERMAN and CALDWELL), A., i, 283.
 benzyl derivatives of (GOMBERG and BUCHLER), A., i, 112.
 iodide, constitution of (LOTTERMOSE and STEUDE), A., i, 10.
 compounds of iodine with (v. EULER and MYRRÄCK), A., i, 527, 1120; (v. EULER and BERGMANN; v. EULER and LANDERGREN), A., i, 921.
 activity of reductase from (SMORODINCEV), A., i, 1201.
 disappearance of, from leaves (MOLISCH), A., i, 309.
 estimation of (LING, CALLOW, and PRICE), A., ii, 879.
- Starfish**, constituents of (HINARD and FILLON), A., i, 87.
- Stasite** (SCHOEP), A., ii, 386.
- Stearic acid**, oxidation of (ASAHINA and IWHIDA), A., i, 520.
 cellulose ester (GRÜN and WITTKA), A., i, 114.
 vanillylamide of (OTT and ZIMMERMANN), A., i, 137.
- Stearic acid**, *dibromodihydroxy-* and *dichlorodihydroxy-* (NICOLET and COX), A., i, 320.
tetrahydroxy-. See Sativic acid.
- Stearodipalmitins**, isomeric (AMBERGER and BROMIG), A., i, 804.
- Stearoxydecoic acid**, methyl ester (GRÜN and WIRTH), A., i, 805.
- Steel**. See under Iron.
- Stereochemistry**, recent advances in (SINGH), A., ii, 103.
 studies in (HOLMBERG), A., i, 1113.
- Stereoisomerides**, biological difference of (JUNG and MÜLLER), A., i, 486.
- Stereoisomerism**, ethylenic (DUFRAISSE), A., i, 534.
- Sterols**, biochemistry of (BRINKMAN and WASTL), A., i, 239.
- Stibinetri-1-piperidinium chloride** (LEONARD), A., i, 363.
- Stibiobenzene**, *mm'*-diamino-*pp'*-*di*hydroxy- (SCHMIDT), A., i, 1204.

- Stilbene**, action of aluminium chloride with (SCHOLL and SCHWARZER), A., i, 331.
- Stilbene**, amino- and nitro-derivatives, stereoisomeric (STOERMER and OEHLERT), A., i, 647.
- nitro-derivatives, preparation of (BISHOP and BRADY), T., 2366.
- 4-nitro-2-cyano-, dibromide (RUGGLI and MEYER), A., i, 346.
- Stilbene-4-carboxylic acids**, 2-amino- and 2-nitro-, stereoisomeric, and their esters (STOERMER and OEHLERT), A., i, 648.
- Stilbene-2:3'-dicarboxyl chloride**, and its derivatives and 7:7'-dichloro-, methyl ester (RUGGLI and MEYER), A., i, 345.
- Stilbene-2:2'-dicarboxylic acid**, esters of (RUGGLI and MEYER), A., i, 344.
- Stomach**, effect of choline on the movements of the (LE HEUX), A., i, 85.
- enzymes of the (HAMMARSEN), A., i, 958.
- human, action of calcium and potassium ions on the (TEZNER and TUROLT), A., i, 396.
- Straw**, lignin from (PASCHKE), A., i, 325.
- estimation of levulose in (COLLINS), A., ii, 323.
- Streptococcus**, hæmolytic, extraction of enzymes from (STEVENS and WEST), A., i, 903.
- Strontium alloys** with lead, constitution of (PIWOWARSKY), A., ii, 644.
- Strontium chlorite** (LEVI), A., ii, 567.
- hydroxide, crystalline, solubility of (SIDERSKY), A., ii, 501.
- nitrate, crystal structure of (VEGARD), A., ii, 503.
- silicate (ESKOLA), A., ii, 849.
- tungstate (SMITH), A., ii, 774.
- Strontium detection**:—
- detection of (LUIZ), A., ii, 227.
- Strychnine**, physiology of (WEISS and HATCHER), A., i, 900.
- poisoning. See Poisoning.
- ferrioxalate (BURROWS and TURNER), A., i, 916.
- hexose- and sucrose-phosphates (NEUBERG and DALMER), A., i, 920.
- separation of quinine from (WARREN and CLARK), A., ii, 171; (EVERS), A., ii, 669.
- iso***Strychnine**, preparation of (LEUCHS and NITSCHKE), A., i, 1175.
- Strychnos alkaloids** (LEUCHS, OSTERBURG, and KAEHN), A., i, 362; (LEUCHS and KAEHN), A., i, 463; (LEUCHS and FRICKER), A., i, 677; (LEUCHS, MILDBRAND, and LEUCHS), A., i, 1052; (LEUCHS and NITSCHKE), A., i, 1175.
- Styphnic acid**, retene ester (RUZICKA and MEYER), A., i, 830.
- Styrene**, chloro-, and its chlorohydrin (FORSTER and SAVILLE), T., 2600.
- o*-nitro-3:4:5-trihydroxy- (ROSEN-MUND and PFANNKUCH), A., i, 1030.
- 2-Styrylbenzopyrylium chloride**, hydroxy-derivatives (BUCK and HEILBRON), T., 1204.
- Styryl α -chloro-*p*-dimethylaminostyryl ketone** (BAUER and WERNER), A., i, 1035.
- Styryl dichloromethyl ketone**, nitro-derivatives (HELLER, LAUTH, and BUCHWALDT), A., i, 348.
- 9-Styryldi- $\beta\beta'$ -naphthoxanthenyl perchlorate** (ZIEGLER), A., i, 152.
- Styryl furfurylidene-methyl ketone**, 2-hydroxy- (BUCK and HEILBRON), T., 1100.
- 9-Styryl-1-methoxyxanthenyl perchlorate** (ZIEGLER), A., i, 152.
- Styryl methyl ketone**, *o*- and *p*-hydroxy- (BUCK and HEILBRON), T., 1100.
- 1-Styrylisoquinoline** (MILLS and SMITH), T., 2732.
- 9-Styrylxanthanol** (ZIEGLER and OCHS), A., i, 1048.
- 9-Styrylxanthenyl perchlorate** (ZIEGLER), A., i, 151.
- 9-Styrylxanthyl bromide hydrobromide** and chloride hydrochloride, and their perchlorides (ZIEGLER and OCHS), A., i, 1047.
- ethyl ether (ZIEGLER and OCHS), A., i, 1048.
- Suberdialdehyde**, and its diphenylhydrazone (ROSEN-MUND, ZETZSCHE, and FLÜTSCH), A., i, 39.
- Suberone**, (?)2:7-dibromo- (GODOHOT and BRUN), A., i, 350.
- Substance**, $C_8H_8N_8$, from azoimide and methylthiocarbimide (OLIVERI-MANDALA), A., i, 473.
- $C_6H_5O_4$, and its methyl ether, from glycerol and *Aspergillus glaucus* (TRAETTA-MOSCA and PRETI), A., i, 91.
- C_6H_5N , from *p*-xylene and sulphuryl azide (CURTIUS and SCHMIDT), A., i, 777.
- $C_6H_5O_3N_2S$, from *p*-carbamidotoluene-*m*-sulphonic acid and phosphoryl chloride (SCOTT and COHEN), T., 2041.
- $C_6H_{13}O_4N$, from hydrolysis of ethyl $\beta\gamma$ -dicyano- γ -methylbutane- $\beta\beta$ -dicarboxylate (HOPE and SHELTON), T., 2232.
- $C_{10}H_{18}O_4$, from α -bromoiso-butanaldehyde and sodiomalonate ester (FRANKE and GROßER), A., i, 808.

- Substance, $C_{10}H_{18}O_5$, from β -methylbutan- β -ol- γ -one and phosphoric oxide (SCHEIBLER and FISCHER), A., i, 1110.
- $C_{12}H_{10}O_6SNa$, and its amino-derivative, from sodium hydrogen sulphite and 2:7-dihydroxynaphthalene (FUCHS and STIX), A., i, 451.
- $C_{16}H_{11}O_2N_2S_2$, from oxidation of 3:5-dimethylthiol-1-phenyl-1:2:4-triazole with permanganate (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 378.
- $C_{11}H_{20}O$, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
- $C_{12}H_{22}O$, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
- $C_{15}H_{16}O_5$, from oxidation of the lactone of 3-hydroxy-2-*o*-carboxybenzyl-2-methylcyclopropane-1-carboxylic acid (KON, STEVENSON, and THORPE), T., 663.
- $C_{15}H_{10}O_6$, from formic acid and phloroglucinol (SCHWENK), A., i, 153.
- $C_{13}H_{12}O_7$, from formic acid and phloroglucinol (SCHWENK), A., i, 153.
- $C_{12}H_{24}O$, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
- $C_{14}H_{14}O_5$, from ethyl 5-aldehydo-salicylate and acetylacetone (WAYNE and COHEN), T., 1027.
- $C_{12}H_{16}O_3$, from kawaic acid and alcoholic potash (MURAYAMA and MAYEDA), A., i, 265.
- $C_{14}H_{16}O_4$, and its platinumchloride, from hydrolysis of $C_{22}H_{26}N_4$ (CIUSA and VECCHIOTTI), A., i, 474.
- $C_{14}H_{16}O_4N_2Cl_2$, from urine and dichlorobenzenediazonium chloride (HERMANS), A., i, 1091.
- $C_{15}H_9O_5$, from *s-o*-phthalyl chloride and sodium salicylate (H. P. and W. KAUFMANN and GÖTTING), A., i, 253.
- $C_{15}H_{12}O_4$, from the bark of *Rhamnus frangula* (OESTERLE), A., i, 100.
- $C_{15}H_{22}O$, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
- $C_{15}H_{22}O_3$, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
- $C_{15}H_{16}O_2N_2$, from hydrolysis of aldehyde phenylmethylhydrazine hydrochloride (CIUSA and VECCHIOTTI), A., i, 474.
- $C_{15}H_{20}ON_2$, from cyclohexanone and ethyl α -cyanopropionate (KON and THORPE), T., 1802.
- $C_{16}H_{16}O_2N_2S$, from monothioethylene glycol and phenylcarbimide (BENNETT), T., 2146.
- $C_{12}H_{24}O$, from ethylene and nitrogen (MIYAMOTO), A., i, 418.
- Substance, $C_{11}H_{14}O$, from 9-styrylxanthenol and acetic acid (ZIEGLER and OCHS), A., i, 1048.
- $C_{21}H_{12}O_4$, and its derivatives, from autoxidation of 2-acetyl- α -naphthol (FRIES and LEUE), A., i, 462.
- $C_{22}H_{16}O_6$, from *m*-cresol and α -nitrophthalic anhydride (EDER and WIDMER), A., i, 260.
- $C_{22}H_{14}O_5N_2$, from β -ketohydrindene and *p*-nitrobenzaldehyde (FRIEDLANDER, HERZOG, and v. VOSS), A., i, 764.
- $C_{23}H_{25}NPI$, from triphenylphosphine-ethylimine and ethyl iodide (STAUDINGER and HAUSER), A., i, 69.
- $C_{23}H_{16}O_5$, from β -ketohydrindene and *p*-hydroxybenzaldehyde (FRIEDLANDER, HERZOG, and v. VOSS), A., i, 764.
- $C_{23}H_{25}O_2NPI$, from ethyl azidoacetate, triphenylphosphine and methyl iodide (STAUDINGER and HAUSER), A., i, 69.
- $C_{25}H_{30}O$, from magnesium phenylethynyl bromide and distyryl ketone (HESS and WELTZIEN), A., i, 35.
- $C_{25}H_{23}O_2N_3$, and its salts, from azodibenzyl and dimethyl- β -naphthylamine (DIELS and KLEINFELLER), A., i, 1195.
- $C_{27}H_{18}O$, from γ -diphenyl- α -di-*p*-anisylallene and acetic acid (ZIEGLER, OCHS, BREMER, and THIEL), A., i, 1049.
- $C_{28}H_{14}O_4$, from 2-hydroxyanthraquinone and dextrose (BRADSHAW and PERKIN), T., 921.
- $C_{28}H_{28}N_4$, from hydrolysis of benzaldehyde phenylhydrazine hydrochloride (CIUSA and VECCHIOTTI), A., i, 474.
- $C_{28}H_{24}ON_4$, from *N*-ethylcarbazole-3-diazonium chloride and ammonia (MORGAN and READ), T., 2717.
- $C_{32}H_{22}O_6$, from oxidation of phenanthraquinyl acetate (GOLDSCHMIDT and SCHMIDT), A., i, 1150.
- $C_{33}H_{22}O_6N_2S_2$, from "saccharin" and *m*-phenylenediamine hydrochloride (DUTT), T., 2393.
- $C_{33}H_{22}O_6N_2S_2$, from "saccharin" and 4:6-diaminoresorcinol hydrochloride (DUTT), T., 2394.
- $C_{34}H_{22}O_3$, from magnesium phenylethynyl bromide and oxalyl chloride (HESS and WELTZIEN), A., i, 36.
- $C_{40}H_{36}O_4$, from diphenylketen and phenylketenacetal (STAUDINGER and RATHSAM), A., i, 1015.

- Substitution**, influence of, on equilibria in binary systems (KREMANN and MÜLLER; KREMANN, ODELGA, and ZAWODSKY, A., i, 181; (KREMANN, HOHL, and MÜLLER), A., i, 138; (KREMANN and ODELGA), A., i, 159; (KREMANN and STRZELBA), A., i, 176.
influence of, on chemical reactions (FRANSEN and STÄUBLE), A., i, 450; (BODFORSS), A., ii, 698.
effect of, on the free energy of oxidation-reduction reactions (LAMER and BAKER), A., ii, 735.
- Substitution reactions** (v. AUWERS), A., ii, 98.
- Succinohloroimide**, preparation of (HIRST and MACBETH), T., 2174.
- Succinic acid**, preparation of (SUZUKI and MATSUYAMA), A., i, 716.
aldehyde acids derived from (CARRIÈRE), A., i, 318.
benzyl ester, enzymic hydrolysis of (HOWARD), A., i, 960.
menthyl hydrogen ester (SHIMOMURA and COHEN), T., 2056.
- Succinic acid**, tetrachloro-, diethyl ester (DOUGHTY and FREEMAN), A., i, 427.
- Succinic acids**, substituted, formation of, from esters of $\alpha\beta$ -unsaturated acids (HIGGINBOTHAM and LAPWORTH), T., 49.
syntheses of (LAPWORTH and McRAE), T., 1699, 2741.
aminohydroxy-, synthesis of, and their salts and derivatives (DAKIN), A., i, 143.
resolution of, and their alkaloidal salts (DAKIN), A., i, 430.
- Succinodehydrogenase** (WIDMARK), A., i, 600.
effect of hydrogen-ion concentration on the action of (OHLSSON), A., i, 785.
- Succinodi-5-acenaphthylamide** (FLEISCHER and SCHRANZ), A., i, 1143.
- Succinonitrile**, dichloro- (OTT and LÖPMANN), A., i, 643.
- Sucrose** (*saccharose*, *cane-sugar*), preparation of, chemically pure (KRAISY), A., ii, 233.
spectrum of the triboluminescence of (LONGCHAMON), A., ii, 542.
polarisation of normal solutions of (STANĚK), A., ii, 167.
heat of combustion of (SWIENTO-SŁAWSKI and STARCZEWSKA), A., ii, 616.
osmotic pressure of (LOTZ and FRAZER), A., ii, 264.
inversion of (MORAN and LEWIS), T., 1613; (SCATCHARD), A., i, 230.
- Sucrose** (*saccharose*, *cane-sugar*), velocity of inversion of (DHAR), A., ii, 89; (FALES and MORRELL), A., ii, 832.
inversion of, by alkaline copper solution (MAQUENNE), A., i, 920; (CANAIS), A., ii, 592.
inversion of, by saccharase (COLIN and CHAUDUN; CHAUDUN), A., i, 389.
hydration of, in water solution (SCATCHARD), A., i, 230.
velocity of hydrolysis of (CLARK), A., ii, 135.
oxidation of, by nitric acid (CHATTAWAY and HARRIS), T., 2703.
action of ozone on solutions of (SCHONEBAUM), A., i, 223.
explosion of finely divided (BEYERSDORFFER), A., ii, 749.
sulphate (NEUBERG and LIEBERMANN), A., i, 222.
benzyl derivatives of (GOMBERG and BUCHLER), A., i, 112.
permeability of the intestines to (WORINGER), A., i, 1214.
detection of, colorimetrically (KRYZ), A., ii, 233.
detection of, in presence of dextrose (CONGDON and STEWART), A., ii, 233.
estimation of, with invertase (HARDING), A., ii, 167.
estimation of, in presence of other sugars (BEHRE and DURING), A., ii, 790.
separation of dextrose from, by dialysis (CONGDON and INGERSOLL), A., i, 322.
- Sucrosephosphoric acid**, strychnine salt (NEUBERG and DALMER), A., i, 920.
- Sugars**, effect of ketonic compounds on the formation of (GEELMUYDEN), A., i, 607.
synthesis of (WAGNER and PARNAS), A., i, 965.
from formaldehyde (EWART), A., i, 10.
in the body (PARNAS and WAGNER), A., i, 487.
fluorescence of (LEWIS), A., ii, 334.
optical rotation of (MALTEY), T., 2608.
adsorption in solutions of, and their emulsifying action (CLARK and MANN), A., ii, 550.
chemistry of (KILIANI), A., i, 223, 321, 1111.
action of amino-acids on (GRÜNHUT and WEBER), A., i, 235.
fermentation of, by pentose-fermenting bacteria (PETERSON, FRED, and ANDERSON), A., i, 971.

- Sugars**, fermentation of, preparation of glycerol from (VEREINIGTE CHEMISCHE WERKE), A., i, 980.
- in blood (CAMMIDGE, FORSYTH, and HOWARD), A., i, 81; (ROSENBERG), A., i, 789; (NEUWIRTH and KLEINER; STASIAK), A., i, 1208.
- distribution of (FALTA and RICHTER-QUITTNER), A., i, 696.
- effect of amino-acids and fatty acids on (POLLAK), A., i, 483.
- effect of carbon dioxide on (BIN-SWANGER), A., i, 962.
- in narcosis and disease (CHANTRAINE), A., i, 192.
- in arterial and venous blood (TURBAN), A., i, 482.
- relation between the content of, in urine and in blood (TERVAERT), A., i, 1215.
- in blood and urine, action of phosphates on (ELIAS and WEISS), A., i, 1085.
- in cerebrospinal fluid, aqueous humour, and blood plasma (COOPE; DE HAAN and VAN CREVELD; STEVENSON), A., i, 295.
- steric transformation of, in tissues (ISAAC and ADLER), A., i, 297.
- stereoisomeric, permeability of the glomerulus membrane for (HAMBURGER), A., i, 490, 491, 790, 919.
- synthesis of acylated halogen derivatives of (FREUDENBERG and IVERS), A., i, 523.
- action of hydrogen peroxide on solutions of (SCHONEBAUM), A., i, 920.
- metabolism of. See Metabolism.
- oxidation of acids derived from (GREINERT), A., i, 1111.
- unsaturated reduction products of (BERGMANN, SCHOTTE, and LECHINSKY), A., i, 227; (BERGMANN and MIEKELF), A., i, 618.
- transportation, retention and excretion of (FOLIN and BERGLUND), A., i, 487.
- excretion of, in urine (MURSCHEHAUSER), A., i, 198; (NEUWIRTH), A., i, 485.
- reducing, action of ammonia on (LING and NANJI), A., i, 631.
- clarification of solutions containing, by means of lead acetate (ENGLIS and TSANG), A., ii, 459.
- estimation of (QUISUMBING and THOMAS), A., ii, 92; (HANAK), A., ii, 166; (FLEURY and BOUTOT), A., ii, 879.
- estimation of, in blood (LABBÉ, NERVEUX, and NOMIDIS), A., ii, 663.
- Sugars**, reducing, estimation of, in urine (SMITH), A., ii, 663.
- estimation of, by fermentation methods (LANGE), A., ii, 98.
- estimation of, iodometrically (AUERBACH and BODLANDER), A., ii, 878.
- estimation of, in blood (ROSENBERG), A., i, 482; (GUY), A., ii, 94; (REIST), A., ii, 323; (STEFF), A., ii, 592; (ERNST and WESSE), A., ii, 724; (CSONKA and TAGGART), A., ii, 879.
- estimation of, in blood and urine (TERVAERT), A., ii, 166.
- estimation of, in urine (FOLIN and BERGLUND), A., ii, 400.
- Sulphamidophthalic acid**, constitution of, and its salts and esters (ZINCKE and GREUNE), A., i, 550.
- p*-**Sulphaminobenzoic acid**, sodium salt (WEIL and MOSER), A., i, 444.
- estimation of, in commercial saccharin (HERZOG and KREIDL), A., ii, 237.
- Sulphamphthaleins** (DUTT), T., 2389.
- cis*-**Sulphatodiethylenediaminecobaltic bromide** (DUFF), T., 452.
- "**Sulphatoethylamine**." See *N*-Phenyl- β -aminoethyl hydrogen sulphate.
- 1:1-Sulphatomercuridimethylene-bis-1:2-dihydrobenzofuran** (ADAMS, ROMAN, and SPERRY), A., i, 947.
- Sulphatopentamminecobaltiselenate**. See under Cobalt.
- Sulphides**. See under Sulphur.
- Sulphide colouring matters**, red (WATSON and DUTT), T., 1939, 2414.
- Sulphidobisdimethylpyrrole-3-carboxylic acids**, ethyl esters (FISCHER and HERRMANN), A., i, 1055.
- Sulphiformin** (*methanalsulphurous acid*) (MALVEZIN), A., i, 222.
- Sulphilimines** (MANN and POPE), T., 1052.
- Sulphimidophthalic acid**, constitution of, and its salts and derivatives (ZINCKE and GREUNE), A., i, 551.
- p*-**Sulphinobenzoic acid** (SMILES and HARRISON), T., 2024.
- 5-Sulphinosalicylic acid** (STEWART), T., 2559.
- Sulphite liquor lactone** (HINTIKKA), A., i, 347.
- cis*-**Sulphitodiethylenediaminecobaltic bromide** (DUFF), T., 452.
- Sulphoacetic acid** as a condensing agent (SCHNEIDER and KRAFT), A., i, 749.
- salts, detection and estimation of, in cellulose acetates (ENTAT and VULQUIN), A., ii, 583.
- cobalt and copper salts (BACKER and DUBSKY), A., i, 428.

- Sulpho-acid groups, steric hindrance of** (VAN DUIN), A., i, 189.
- 8-*p*-Sulphobenzeneazo-5-amino-6-methoxyquinoline** (JACOBS and HEIDELBERGER), A., i, 671.
- 5-Sulphobenzeneazo-2:4-dimethylpyrrole-3-carboxylic acid, ethyl and methyl esters** (KÜSTER, WEBER, MAURER, NIEMANN, SCHLACK, SCHLAYERBACH, and WILLIG), A., i, 858.
- 8-*p*-Sulphobenzeneazo-5-hydroxy-6-methoxy-2-phenylquinoline-4-carboxylic acid** (HALBERKANN), A., i, 175.
- 8-*p*-Sulphobenzeneazo-5-hydroxy-6-methoxyquinoline** (JACOBS and HEIDELBERGER), A., i, 672.
- 5-*p*-Sulphobenzeneazo-6-hydroxy-2-phenylquinoline-4-carboxylic acid** (HALBERKANN), A., i, 174.
- Sulphobenzide.** See Diphenylsulphone.
- Sulphocarboxylic acids, salts of** (BACKER and DUBSKY), A., i, 423.
- m*-Sulphocinnamic acid, and its aniline hydrogen salt** (MOORE and THOMAS), A., i, 454.
- Sulphohæmoglobinæmia** (VAN DEN BERGH), A., i, 798.
- 1-3'-Sulpho-2-hydroxyphenyl-3-methyl-5-pyrazolone, 5'-chloro-** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 385.
- 1-5'-Sulpho-2'-hydroxyphenyl-3-methyl-5-pyrazolone-3'-carboxylic acid** (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 385.
- Sulphonamides, aliphatic** (CLUTTERBUCK and COHEN), T., 120.
substituted, preparation of (FARBEN-FABRIKEN VORM. F. BAYER & Co.), A., i, 999.
- Sulphonation of phenols** (CAMPBELL), T., 847.
- α -Sulphonedibutyric acid, resolution of** (AHLBERG), A., i, 625.
- Sulphonic acids, esters, action of, with amines** (FOLDI), A., i, 782.
- Sulphonic acids, *o*-amino-, aromatic, condensation of isocyanic acid with** (SCOTT and COHEN), T., 2034.
- 2:2'-Sulphonidetriphenylcarbinol** (GOMBERG and BRITTON), A., i, 163.
- 2:2'-Sulphonidetriphenylcarbinyl chloride** (GOMBERG and BRITTON), A., i, 163.
- 2:2'-Sulphonidetriphenylmethane** (GOMBERG and BRITTON), A., i, 163.
- 2:2'-Sulphonidetriphenylmethyl, and its peroxide** (GOMBERG and BRITTON), A., i, 163.
- Sulphonyl chlorides, aromatic** (STEWART), T., 2555.
- 2-*p*-Sulphophenyl-6-methyl- ψ -asimino-benzene, 5-amino-** (KALLE & Co.), A., i, 61.
- α -Sulphopropionic acid, cobalt and copper salts** (BACKER and DUBSKY), A., i, 423.
- Sulphosalicylaldehyde, preparation and derivatives of** (WEIL and BRIMMER), A., i, 849.
- Sulphosalicylic acid, action of, on the swelling of gelatin and albumin** (OSTWALD and KUHN), A., i, 598.
- 5-Sulphosalicylic acid, 3-nitro-, potassium hydrogen salt** (SAKELLARIOS), A., i, 1145.
- Sulphoxyllic acid, estimation of, volumetrically** (DE BACHO), A., ii, 311.
- Sulphur, absorption spectrum of** (FUKUDA), A., ii, 728.
- Röntgen-ray spectrum of** (LINDH), A., ii, 542.
- latent heat of fusion of** (STRATTON and PARTINGTON), A., ii, 258.
- rhombic, solubility of** (HILDEBRAND and JENKS), A., ii, 141.
- solubility of, in organic liquids** (DELAPLACE), A., ii, 706.
- valency of** (LECHER and SIMON), A., i, 1018.
- absorption of active hydrogen and nitrogen by** (WENDT; NEWMAN), A., ii, 639.
- colloidal, physico-chemical investigation of** (ROSSI), A., ii, 485.
- hydrophobic and hydrophilic sols of** (FREUNDLICH and SCHOLZ), A., ii, 841.
- effect of heating charcoal with** (WIBAUT), A., ii, 52, 373.
- action of, on cuprous chloride** (PINKARD and WARDLAW), T., 1300.
- equilibrium between iodine and, in solution** (AMADORI), A., ii, 561.
- and its compounds, action of, on terpenes** (BUDNIKOV and SCHILOV), A., i, 944.
- metabolism.** See Metabolism.
- in proteins** (HOFFMAN and GORTNER), A., i, 429.
- Sulphur compounds in agar** (NEUBERG and OHLE), A., i, 828.
- action of, on growth of plants** (TURINA), A., i, 707.
- oxidation of, by bacteria in soils** (LIPMAN, WAKSMAN, and JOFFE), A., i, 803; (WAKSMAN), A., i, 706.
- fate of, in the animal organism** (SCHMIDT and CLARK), A., i, 968.
- Sulphur monochloride, action of aniline with** (COFFEY), A., i, 132.

Sulphur:—

Sulphuryl chloride, action of, on aromatic amines (ELLER and KLEMM), A., i, 448.

chlorination of benzene with (SILBERRAD), T., 1015.

action of, on organic compounds (DURRANS), T., 44.

Pyrosulphuryl chloride, action of toluene on (STEINKOPF and BUCHHEIM), A., i, 122.

Thionyl chloride, action of, on glycollic acid (BLAISE and MONTAGNE), A., i, 520.

action of, on α -hydroxy-acids (BLAISE and MONTAGNE), A., i, 715.

Sulphides, phosphorescent, refractive index of (CURIE), A., ii, 801.

action of red rays on (CURIE), A., ii, 246.

effect of heat on phototropy of (MOURELO), A., ii, 604.

preparation of sols of (v. HAHN), A., ii, 866.

fate of, in the blood (HAGGARD and CHARLTON), A., i, 287.

estimation of (BUDNIKOV and KRAUSE), A., ii, 717, 782.

estimation of, volumetrically (WILLARD and CAKE), A., ii, 80.

Sulphur dioxide, specific heat of (PARTINGTON and CANTR), A., ii, 191.

solubility of, in suspensions of calcium and magnesium hydroxides (SMITH and PARKHURST), A., ii, 761.

vapour pressure of (BERGSTROM), A., ii, 423.

viscosity and molecular dimensions of (SMITH), A., ii, 686.

equilibrium between water and (BICHOWSKY), A., ii, 270.

oxidising properties of (WARDLAW and PINKARD), T., 210.

oxidising and reducing properties of (STEWART and WARDLAW), T., 1481.

estimation of (BARHAM), A., ii, 520.

trioxide, physical properties of (BERTHOUD), A., ii, 638.

equilibrium of chromium trioxide, water and (GILBERT, BUCKLEY, and MASSON), T., 1934.

equilibrium of ferric oxide, water and (POENJAK and MERWIN), A., ii, 772.

tetroxide (MEYER, BAILLEUL, and HENKEL), A., ii, 843.

Sulphurous acid, autoreduction of (BENNETT) T., 1794.

Sulphur:—

Sulphurous acid, reaction between iodic acid and (SKRABAL), A., ii, 488.

action of iodine with (MACAULAY), T., 552.

estimation of (COPPETTI), A., ii, 80.

Sulphites, preparation of, pure (SHENEFIELD, VILBRANDT, and WITHROW), A., ii, 45.

alkali hydrogen, equilibria in aqueous solutions of (BALY and BAILEY), T., 1813.

detection of, in foods (CHAPMAN), A., ii, 520.

Sulphuric acid, lead chamber process for (FORRER), A., ii, 761.

and its salts, electrical conductivity of mixtures of (KENDALL, ADLER, and DAVIDSON), A., ii, 126.

determination of the transport numbers for (FERGUSON and FRANCE), A., ii, 113, 114.

properties of mixtures of ethyl ether, water and (POUND), T., 941.

heat developed on mixing nitric acid, water and (McDAVID), A., ii, 617.

absorption of ethylene by (DAMIENS), A., i, 1105.

action of, with aromatic disulphides (SMILES and McCLELLAND), T., 86.

detection and estimation of, in cellulose acetate (ENTAT and VULQUIN), A., ii, 533.

estimation of, as barium sulphate (BALAREFF), A., ii, 864.

estimation of, in presence of aluminium (MOSER and KOHN), A., ii, 782.

Sulphates, diffusion of (STILES), A., ii, 125.

double hydrated, dissociation pressures of (CAVEN and FERGUSON), T., 1406.

estimation of (KOLTHOFF), A., ii, 81.

estimation of, as barium sulphate (CHATTERJEE), A., ii, 390.

estimation of, volumetrically (JELLINEK and ENS), A., ii, 864.

estimation of, in soils (HIRST and GREAVES), A., ii, 521.

estimation of, in blood (DENIS), A., ii, 225.

Persulphates, estimation of (SCAGLIARINI and TORELLI), A., ii, 81.

Hyposulphurous acid, estimation of, volumetrically (DE BACHO), A., ii, 311.

Sulphur:—

Thiosulphuric acid, occurrence and estimation of, in the animal organism (DEZANI), A., i, 968.

Thiosulphates, periodicity of reactions between arsenites or arsenates and (FORBES, ESTILL, and WALKER), A., ii, 271.

reaction of, with nitrites (FALCIOLA), A., ii, 453.

ageing of standard solutions of (HAHN and WINDISCH), A., ii, 873.

Dithionates, estimation of, volumetrically (FISCHER and CLASSEN), A., ii, 453.

Polythionic acids, and their salts (RIESENFELD and FELD), A., ii, 45.

Polythionates, analysis of (KURTENACKER and FRITSCH), A., ii, 521.

Sulphur organic compounds (MANN and POPE), T., 1052.

autoxidation of (DELÉPINE), A., i, 621, 914.

thiocyanates (LECHER and WITTEW; LECHER and GOEBEL), A., i, 640.

Sulphuric acid, glucose esters of (LEVENE, MEYER, and WEBER), A., i, 987.

mixed, anhydrides of, with carboxylic acids (VAN PESKI), A., i, 106.

Sulphur estimation:—

estimation of (LOSANA), A., ii, 582, 656.

estimation of, microchemically (HOLTZ), A., ii, 521.

estimation of, in antimony sulphide (LUFF and PORRITT), A., ii, 225.

estimation of, in illuminating gas, by means of perhydrol (KLEMMER), A., ii, 224.

estimation of, in commercial ichthyol preparations (PÉPIN and REAUBOURG), A., ii, 784.

estimation of, in cast iron and steel (MARINOT), A., ii, 224.

estimation of, in organic compounds (MEULEN), A., ii, 311.

estimation of, in pyrites (GADAIS), A., ii, 79; (CHAUDRON and JURE-BOIRARD), A., ii, 311.

estimation of, in vulcanised rubber (DYER and WATSON), A., ii, 656, 782.

estimation of, in urine (ROBISON), A., ii, 389.

Sulphur group, heteropoly-acids of the elements of the (MEYER and STANECZY), A., ii, 773.

Sulphuric and Sulphurous acids. See under Sulphur.

Sulphuryl chloride. See under Sulphur.

Sulphurylaside, action of, on benzene (SCHMIDT), A., i, 777.

action of, on *p*-xylene (CURTIUS and SCHMIDT), A., i, 776.

Sunlight, effect of various rays of, on the formation of essential oils in plants (CANALS), A., i, 907.

coagulation of colloids by (GANGULY and DHAR), A., ii, 604.

Surfaces, orientation of molecules in (HARKINS and ROBERTS), A., ii, 422.

Surface tension, measurement of (SUGDEN), T., 858; (ELDER), A., ii, 618.

calculation of (CREIGHTON), A., ii, 426.

effect of addition of electrolytes on (IZAGUIRRE), A., ii, 262.

effect of, on rate of solution (GANGULY and BANERJI), A., ii, 821.

of corresponding states (LORENZ and HERZ), A., ii, 261.

of thin layers of liquids (MARCELIN), A., ii, 686.

Suspensoids, coagulation of (v. HAHN), A., ii, 37.

Symphoricarpos racemosus, sugars from the fruit of (v. LIPPMANN), A., i, 311.

Synergism, ionic (MICHAELIS and HIRABAYASHI), A., ii, 429.

Synovial fluid, analyses of (MALMÉJAC), A., i, 703.

Syphilis, action of bismuth compounds on (SAZERAC and LEVADITI), A., i, 89; (FOURNIER and GUÉNOT), A., i, 301.

action of sodium *p*-hydroxy-*m*-aminophenylarsinate on (FOURNIER, GUÉNOT, and SCHWARTZ), A., i, 300.

Systems, binary. See Binary systems.

conjugated, of double linkages, combination in (BERGMANN), A., i, 1106.

disperse, optical properties of (LIFSCHITZ and BECK), A., ii, 597.

heterogeneous, equilibria in (SYEKKIN), A., ii, 699.

velocity of reaction in (COLLENBERG and BODFORSS), A., ii, 431.

symmetrical tautomeric triad, mobility of (INGOLD and PIGGOTT), T., 2381.

T.

Tadpoles, influence of the reaction of the solution on the action of poisons or drugs on (LABES), A., i, 901.

- Takadiastase**, inactivation of (JACOBY and SHIMIZU), A., i, 481.
- Tallow**, Mafurra (RINDL), A., i, 1102.
- Tannase** (FREUDENBERG and VOLLBRECHT), A., ii, 285.
estimation of (RHIND and SMITH), A., ii, 407.
- Tannin**, detection of (ATKINSON and HAZLETON), A., ii, 793.
estimation of, in wines (MALVEZIN), A., ii, 172.
- Tannins** (FREUDENBERG, BÖHME, and PURRMANN), A., i, 756; (FREUDENBERG and VOLLBRECHT), A., i, 1046.
from *Acer ginnala* (PERKIN and UYEDA), T., 66.
catechu (NIERENSTEIN), T., 23.
Chinese (FREUDENBERG and SCILASI), A., i, 1169.
crystalline synthetic (KARRER and SALOMON), A., i, 266.
from spruce needles (v. EULEK), A., i, 233.
estimation of, in wines (FRESENIUS and GRUNHUT), A., ii, 96.
- Tanning** of hides and leather (RAKUZIN), A., i, 703.
- Tantalum dichloride** (LINDNER), A., ii, 509.
pentachloride, preparation of, and its insulating properties (BILTZ and VOIGT), A., ii, 303.
reduction of (RUFF and THOMAS), A., ii, 513.
pentoxide, action of light on (RENZ), A., ii, 51.
- Tantalum separation**:—
separation of, from columbium (MERRILL), A., ii, 230.
- Tapiolite** from South Dakota (HEADDEN), A., ii, 387.
- Tar acids**, estimation of phenol in (HOFFERT), A., ii, 878.
- l-Tartardialdehyde**, and its derivatives (BERGMANN), A., i, 8.
- Tartaric acid**, and its salts, rotatory dispersion of (LOWRY and AUSTIN), A., ii, 414.
active racemic compound of (LANDRIEU), A., i, 808.
and its salts, action of uranyl acetate on (KOPATSCHEK), A., i, 984.
potassium sodium salt, piezo-electricity of crystals of (SCOTT), A., ii, 609.
ethyl ester, preparation and properties of (LOWRY and CUTTER), T., 532.
green coloration of (PATTERSON), T., 1042.
methyl ester, acetyl derivative (FREUDENBERG and BRAUNS), A., i, 628.
- Tartaric acid**, estimation of, volumetrically, in presence of alkaline earth chlorides (SIMON and ZIVY), A., ii, 880.
- Tartaric acid, dihydroxy-** (LACHMAN), A., i, 109.
- Taste**, peppery, relation between chemical constitution and (OTT and ZIMMERMANN), A., i, 137.
- Tautomerism**, complex (LEY and GRAU), A., i, 536.
of dyads (USHERWOOD), T., 1604.
intra-annular (FARMER, INGOLD, and THORPE), T., 128.
keto-enolic (KAUFFMANN), A., i, 985.
application of Claisen's rule to (DIECKMANN), A., i, 1020.
ring-chain (KON, STEVENSON, and THORPE), T., 650; (DESHAPANDE and THORPE), T., 1490; (INGOLD, PERREN, and THORPE), T., 1765.
- Taxine**, and its derivatives (WINTERSTEIN and IATRIDES), A., i, 572.
- Taxus baccata*** (yew), taxine from (WINTERSTEIN and IATRIDES), A., i, 572.
- Telegraphy**, wireless, new detectors for (HEIKE), A., ii, 19.
- Telluric acid**. See under Tellurium.
- Tellurium**, absorption spectrum of (LINDSAY), A., ii, 599.
allotropy of (DAMIENS), A., ii, 498.
amorphous, crystallisation of (DAMIENS), A., ii, 562.
metallurgy of (HULOT), A., ii, 142.
equilibrium of mixtures of sodium and, in liquid ammonia (KRAUS and CHIU), A., ii, 765.
- Tellurium alloys** with antimony and with lead (DREIFUSS), A., ii, 503.
- Tellurium compounds**, pharmacology of (JOACHIMOGLU and HIROSE), A., i, 396, 406.
- Tellurium salts**, action of, on growth of plants (TURINA), A., i, 707.
action of, on bacteria (JOACHIMOGLU), A., i, 1095.
- Telluric acid**, preparation of (MEYER and MOLDENHAUER), A., ii, 49.
- Tellurides**, preparation of (MOSER and ERTL), A., ii, 48.
- Tellurium organic compounds**:—
tetrabromide, compounds of amines with (LOWY and DUNBROOK), A., i, 446.
bisacetylacetone dichloride (MORGAN and DREW), T., 929.
bisbenzoylacetone dichloride (MORGAN and DREW), T., 930.
bischloroacetylacetone dichloride (MORGAN and DREW), T., 936.
bispivalylacetone dichloride (MORGAN and DREW), T., 938.

- Tellurium organic compounds** :—
 chloroacetylacetone, and its dichloride (MORGAN and DREW), T., 985.
 dibenzoylmethane trichloride (MORGAN and DREW), T., 939.
 3-ethylacetylacetone and its dihaloids (MORGAN and DREW), T., 932.
 O-ethylpivalylacetone trichloride (MORGAN and DREW), T., 937.
 3-methylacetylacetone, and its dihaloids (MORGAN and DREW), T., 931.
- Tellurium detection and estimation** :—
 detection and estimation of (MULLER, MENZEL, and SCHUBERT), A., ii, 390.
- Temperature curves** (ARIANO), A., ii, 470.
 of combustion (BRONN), A., ii, 548.
 high, determination of, by effusion of gases (YAMAGUCHI), A., ii, 470.
 investigations at (RUFF, SCHMIDT, and MUGDAN), A., ii, 818.
 constant high, attainment of (MOSEK), A., ii, 115.
 low, maintenance of constant (CIOFFI and TAYLOR), A., ii, 817.
 of the body, influence of the thyroid on regulation of (GRAFE and v. REDWITZ; SCHENCK), A., i, 491.
- Teraconic acid**, preparation of (FARMER, INGOLD, and THORPE), T., 150.
- Terephthalaldehyde**, phenylhydrazones of (ROSENMUND, ZETZSCHE, and FLÜTSCH), A., i, 39.
- Terephthalylbis(diphenylamide)** (ROSENMUND, ZETZSCHE, and FLÜTSCH), A., i, 39.
- Terephthalylidiane** (ROSENMUND, ZETZSCHE, and FLÜTSCH), A., i, 39.
- Terephthalylidene- α -naphthalide** (ROSENMUND, ZETZSCHE, and FLÜTSCH), A., i, 39.
- Terephthalylidene- α -naphthylamine** (ROSENMUND, ZETZSCHE, and FLÜTSCH), A., i, 39.
- Terpenes**, chemistry of (VESTERBERG; VESTERBERG and WESTERLIND), A., i, 825.
 action of sulphur and sulphur compounds on (BUDNIKOV and SCHILOV), A., i, 944.
 dicyclic, formulæ of (DUPONT), A., i, 1042.
- Terpene compounds** (RUZICKA and MEYER), A., i, 547, 829; (RUZICKA, MEYER, and MINGAZZINI), A., i, 560; (RUZICKA and SEIDEL), A., i, 562.
- Terpene compounds**, solubility, capillary activity, and hæmolytic activity of (RHODE), A., i, 964.
- Terpin**, commercial, melting point of (CLAVERA), A., i, 1042.
 hydrate, identity of flagstaffite with (GUILD), A., i, 76.
- Terpineol**, preparation of (MARCHANT), A., i, 262.
 hydrate, detection of (DENIGES), A., ii, 789.
- Tetany**, effect of sodium salts on the production of (TISDALL), A., i, 1216.
 calcium in blood in (ROBLES), A., i, 192.
 phosphorus metabolism in (ELIAS and SPIEGEL), A., i, 401.
- Tethelin** (DRUMMOND and CANNAN), A., i, 491.
- 5:7:12:14-Tetra-acetoxylbenzothianthren** (BRASS and KOHLER), A., i, 1051.
- Tetra-acetoxystearic acid** (NICOLET and COX), A., i, 320.
- Tetra-acetyl- γ -bromoglucose** (WREDE), A., i, 226.
- Tetra-acetyltrichloroacetylglucoses** (BRIGL), A., i, 225.
- Tetra-acetyl-gallaldehyde-cyanohydrin** (ROSENMUND and PFANNKUCH), A., i, 1031.
- Tetra-acetylglucose**, mandelic acid derivatives of (KARBER, NAGELI, and SMIRNOV), A., i, 253.
- Tetra-acetylmannose**, chloro- (BRAUNS), A., i, 433.
- Tetra-acetylsalicylphenylmethylamine** (ZEMPLÉN and KUNZ), A., i, 564.
- Tetra-acetylsalicyltrimethylammonium bromide** (ZEMPLÉN and KUNZ), A., i, 565.
- α -Tetra-amylose dodeca-acetate** (PRINGSHEIM and DERNIKOS), A., i, 632.
- Tetra-aquodipyridinemagnesium salts** (SPACU), A., i, 859.
- 5:7:12:14-Tetrabenzoxyloxydibenzothianthren** (BRASS and KÖHLER), A., i, 1051.
- Tetrabenzyl-benzylglucoside** (GOMBERG and BUCHLER), A., i, 112.
- 2:2:4:4-Tetracarboxycyclobutane-1-malonic-3-acetic acid**, ethyl ester (INGOLD, PERREN, and THORPE), T., 1788.
- Tetracosic acid**, esters of (BRIGL and FUCHS), A., i, 712.
- n -Tetracosyl alcohol**, and its derivatives (BRIGL and FUCHS), A., i, 713.
- Tetraethylammonium**, and its derivatives (SCHLUBACH and BALLAUF), A., i, 15.
 ruthenipentabromide (GUTHRIE and KRAUSS), A., i, 16.

- Tetraethyldiphenylquinoxalinium perchlorate** (KEHRMANN, ROY, and RAMM), A., i, 468.
- Tetraethylphosphonium bromide and iodide**, compounds of iodoform with (STEINKOPF and SCHWEN), A., i, 118.
- Tetraethylstibinium iodide**, compound of iodoform and (STEINKOPF and SCHWEN), A., i, 118.
- Tetraglucosan**, and its derivatives (PRINGSHEIM and SCHMALZ), A., i, 1118.
- Tetrahydroacenaphthene derivatives** (V. BRAUN, HAHN, and SEEMANN), A., i, 728.
- Tetrahydroacenaphthene**, 4-hydroxy-, and 4-thiol-, and their derivatives (V. BRAUN and KIRSCHBAUM), A., i, 728.
- Tetrahydroacenaphthene-4-sulphinic acid** (V. BRAUN and KIRSCHBAUM), A., i, 728.
- Tetrahydroacenaphthene-4-sulphonic acid**, and its salts and derivatives (V. BRAUN and KIRSCHBAUM), A., i, 728.
- Tetrahydroaceto- α -naphthalide**, 6:7-di- and 4:6:7-tri-bromo- (ROWE and DAVIES), T., 1005.
- Tetrahydrobenzyl alcohol** (DE POMMEREAU), A., i, 825.
- Tetrahydrobufotalin** (WIELAND), A., i, 200.
- l-Tetrahydrocarvone** (SIMONSEN and RAU), T., 881.
- Tetrahydrocinchonine** (JACOBS and HEIDELBERGER), A., i, 672.
- Tetrahydro- ψ -codeine**, and its salts (SPEYER and WIETERS), A., i, 47.
- α - and β -Tetrahydrodeoxyquinonines** (JACOBS and HEIDELBERGER), A., i, 672.
- γ -5-Tetrahydrodiphenyleneoxide-*n*-butyric acid**, and its derivatives (MAYER and KRIEGER), A., i, 747.
- γ -5-Tetrahydrodiphenyleneoxide- γ -keto-*n*-butyric acid**, ethyl ester (MAYER and KRIEGER), A., i, 747.
- γ -5-Tetrahydrodiphenyleneoxide-*n*-propylamine**, and its hydrochloride and acetyl derivative (MAYER and KRIEGER), A., i, 748.
- Tetrahydrodipyridyl hydrate**, and its salts (PETERS), A., i, 48.
- Tetrahydrodipyridyls**, dialkyl derivatives, action of iodine on (EMMERT and PARR), A., i, 179.
- Tetrahydrodurene** (SKITA and SCHENCK), A., i, 241.
- Tetrahydromethystic acid** (GOEBEL), A., i, 657.
- Tetrahydromethysticols**, and its semicarbazone (GOEBEL), A., i, 657.
- Tetrahydro- β -naphthaldehyde**, and its phenylhydrazones (WEIL and OSTERMEIER), A., i, 139.
- av*- α - and - β -Tetrahydronaphthaldehydes**, and their derivatives (V. BRAUN, MOLDAENKE, DIRLAM, and GRUBER), A., i, 749.
- Tetrahydronaphthalene** (*tetralin*), physical properties of (HERZ and SCHUFTAN), A., i, 647.
- preparation of derivatives of (STEVENSON and THORPE), T., 1717.
- derivatives, dehydrogenation of (V. BRAUN, HAHN, and SEEMANN), A., i, 728.
- fate of, in the organism (RÖCKEMANN), A., i, 499.
- Tetrahydronaphthalene**, dibromonitro- (V. BRAUN, HAHN, and SEEMANN), A., i, 729.
- nitro-derivatives (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 128.
- ar*-Tetrahydronaphthalenes**, triamino-, and their derivatives (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 126.
- 5:6:7:8-Tetrahydronaphthalene-1- and -2-aldehydes**, and their derivatives (FLEISCHER and FELDMEIER), A., i, 1160.
- ar*-Tetrahydronaphthalenecarboxylic acids**, aminohydroxy-, hydroxy- and nitrohydroxy-, and their derivatives (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 129.
- ar*-Tetrahydronaphthalenesulphonic acids**, and nitro-, and their salts and derivatives (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 126.
- ar*-Tetrahydronaphthalene-1-thiolacetic acid** (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 127.
- α - and β -Tetrahydronaphthanthraquinones** (TETRALIN G. M. B. H.), A., i, 1040.
- 7:8:9:10-Tetrahydro- β -naphthaquinoline**, and its salts (V. BRAUN and GRUBER), A., i, 762.
- Tetrahydronaphthasines**, and its salts and derivatives (V. BRAUN and GRUBER), A., i, 762.
- ar*-Tetrahydro- α - and - β -naphthols** (SCHROETER and TETRALIN G. M. B. H.), A., i, 1036.
- and amino-, bromo-, chloro- and nitro-derivatives, and their derivatives (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 127.

- ar*-Tetrahydro- β -naphthol-4-azobenzene-*p*-sulphonic acid (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 130.
- ar*-Tetrahydro- β -naphthol-3-sulphonic acid, and 1-bromo-, and their salts and derivatives (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 127.
- ar*-Tetrahydronaphthonitrile (v. BRAUN, MOLDAENKE, DIRLAM, and GRUBER), A., i, 749.
- α - and β -Tetrahydronaphthoylacetyl chlorides (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- α - and β -Tetrahydronaphthoylacetylmalonie acids, ethyl esters (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- β -Tetrahydronaphthoyl-*o*-benzoie acid, and its derivatives (TETRALIN G. M. B. H.), A., i, 1039.
- α - and β -Tetrahydronaphthoylpropionic acids (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- Tetrahydronaphthylacetic acid, and its ethyl ester (DE POMMERAU), A., i, 828.
- ar*-Tetrahydro- α -naphthylamine, electrolytic oxidation of (ONO), A., i, 1008.
- Tetrahydro- β -naphthylamine, β -hydroxy-, and its salts (TAKEDA and KURODA), A., i, 275.
- ar*-Tetrahydro- α - and - β -naphthylamines, and their derivatives and nitro- (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 124.
- ar*-Tetrahydronaphthylaminesulphonic acids (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 129.
- α - and β -Tetrahydronaphthylbutyric acids (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- β -Tetrahydronaphthylbutyryl chloride (SCHROETER and TETRALIN G. M. B. H.), A., i, 1137.
- 2- β -Tetrahydronaphthylcinchoninic acid, and its salts and ethyl ester (v. BRAUN, HAHN, and SEEMANN), A., i, 728.
- Tetrahydro- β -naphthyl dimethylamine, β -hydroxy-, and its salts (TAKEDA and KURODA), A., i, 275.
- ar*-Tetrahydronaphthylenediamines, and their derivatives (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 125.
- 2:3-*ar*-Tetrahydronaphthylphenanthrazine (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 125.
- Tetrahydronaphthylethanol, and its phenylurethane (DE POMMERAU), A., i, 828.
- Tetrahydro β -naphthylmethylamine, β -hydroxy-, and its salts (TAKEDA and KURODA), A., i, 275.
- Tetrahydronaphthylmethylamines, and their salts and derivatives (v. BRAUN, MOLDAENKE, DIRLAM, and GRUBER), A., i, 749.
- ar*-Tetrahydro- α - and - β -naphthylmethylamines, acetyl derivatives (SCHROETER, KINDERMANN, DIETRICH, BEYSCHLAG, FLEISCHHAUER, RIEBENSAHM, and OESTERLIN), A., i, 124.
- o*- β Tetrahydronaphthylmethylbenzoie acid (TETRALIN G. M. B. H.), A., i, 1040.
- β -Tetrahydronaphthylphthalide (TETRALIN G. M. B. H.), A., i, 1039.
- 2- β -Tetrahydronaphthylquinoline, and its derivatives (v. BRAUN, HAHN, and SEEMANN), A., i, 729.
- 2-*ar*-Tetrahydronaphthylquinoline-4-carboxylic acid, and 7-bromo- (CHEMISCHE FABRIK AUF AKTIEN VORM. E. SCHEERING), A., i, 952.
- 1- and 2-Tetrahydronaphthylthiols (TETRALIN G. M. B. H.), A., i, 340.
- ar*-Tetrahydronaphthylthiols (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 126.
- ar*-Tetrahydronaphthyl-2-thiolacetic acid (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 127.
- 1- and 2-Tetrahydronaphthylthiolacetic acids (TETRALIN G. M. B. H.), A., i, 340.
- Tetrahydropyran-4:4-dicarboxylic acid, ethyl ester (KAMM and WALDO), A., i, 105.
- Tetrahydropyran-4:5-*spiro*-3:4:6-triketo-hexahydropyrimidine (KAMM and WALDO), A., i, 105.
- Tetrahydropyrido-3:4-iminasole. See 1:3:5-Benzotriazole.
- Tetrahydrothiodiazole, *dithio*- (GUHA), A., i, 877.
- bis*-*ar*-Tetrahydrothionaphthendisulphonic acid, sodium salt (SCHROETER, SVANOE, EINBECK, GELLER, and RIEBENSAHM), A., i, 127.
- 2:3:5:6-Tetraketo-1:4-dimethylpiperazine (FORSTER and SAVILLE), T., 824.
- Tetralaeoglucosan, and its derivatives (PRINGSHEIM and SCHMALZ), A., i, 1118.

Tetralin. See Tetrahydronaphthalene.

1:3:5:7-Tetramethoxyanthraquinone, and 2:4:6:8-tetranitro- (HELLER and LINDNER), A., i, 1040.

4:4':6:6'-Tetramethoxyarsenobenzene, 3:3'-diamino-, dihydrochloride (CHRISTIANSEN), A., i, 1203.

4:6:3':4'-Tetramethoxy-3-ethoxy-3-phenylchroman, 2-hydroxy- (NIERENSTEIN), T., 610.

7:8:5':6'-Tetramethoxy-2-methyl-3:4-indeno(2':1')-benzopyrylium ferri-chloride (CRABTREE and ROBINSON), T., 1041.

4:6:3':4'-Tetramethoxy-3-phenylchroman, and 3-bromo-, 3-bromo-2-hydroxy-, and 2-chloro- (NIERENSTEIN), T., 609.

2:8-Tetramethyldiaminoacridine (MOUDGILL), T., 1506.

3:6-Tetramethyldiaminocyanoselenopyronin, and its salts (BATTEGAY and HUGEL), A., i, 669.

Tetramethyldiaminodiphenylmethane, compound of tellurium tetrabromide with (LOWRY and DUNBROOK), A., i, 446.

4:4'-Tetramethyldiaminodiphenylmethane, 3:3'-din Nitro- (MOUDGILL), T., 1507.

4:4'-Tetramethyldiaminodihydroxy-fuchsones (MUKERJI), T., 551.

4:4'-Tetramethyldiamino-2:2'-oxido-phenylsulphamphthalein (DUTT), T., 2394.

3:6-Tetramethyldiaminoselenoxanthone (BATTEGAY and HUGEL), A., i, 669.

Tetramethylammonium aluminate, condition of, in solution (BĀEZINA), A., i, 638.

ruthenipentabromide (GUTHRIE and KRAUSS), A., i, 16.

chlorite (LEVI), A., i, 528.

iodide, compound of iodoform and (STEINKOPF and SCHWEN), A., i, 118.

Tetramethylarsonium iodide, compound of iodoform and (STEINKOPF and SCHWEN), A., i, 118.

1:2:4:6-Tetramethyl-1-dichloromethyl- $\Delta^{2,5}$ -cyclohexadien-4-ol (V. AUWERS and ZIEGLER), A., i, 121.

1:3:7:9-Tetramethyldeoxyuric acid, and its salts (BILTZ and HEIDRICH), A., i, 384.

Tetramethyldiallyl. See $\beta\eta$ -Dimethyl- Δ^8 -octadiene.

Tetramethyldiamylose (PRINGSHEIM and PERSCH), A., i, 632.

Tetramethyldiethanoltriaminotriphenylcarbinol hydrochloride (BRITISH DYE-STEUFFS CORPORATION, LTD., GREEN, SAUNDERS, and BATE), A., i, 1068.

Tetramethyldiphenocuinonolmenium perchlorate (KEHRMANN, ROY, and RAMM), A., i, 467.

Tetramethylenediguanidine, and its salts (KIESEL), A., i, 531.

action of arginase on (KIESEL), A., i, 413.

Tetramethylethanoltriaminodiphenyl-naphthylcarbinol hydrochloride (BRITISH DYE-STEUFFS CORPORATION, LTD., GREEN, SAUNDERS, and BATE), A., i, 1068.

1:1:3:6-Tetramethyl-4-ethyl- $\Delta^{2,5}$ -cyclohexadien-4-ol (V. AUWERS and ZIEGLER), A., i, 121.

1:1:3:6-Tetramethyl-4-ethylidene- $\Delta^{2,5}$ -cyclohexadiene (V. AUWERS and ZIEGLER), A., i, 121.

1:3:4:6-Tetramethyl fructose (IRVINE and PATTERSON), T., 2697.

Tetramethylglycerol chlorohydrin. See $\beta\delta$ -Dimethylpentane, γ -chloro- $\beta\delta$ -dihydroxy-.

$\beta\beta\delta$ -Tetramethylheptan- γ -one, η -chloro- (BILLON), A., i, 717.

Tetramethyl- $\Delta^{2,5}$ -cyclohexadien-4-ols (V. AUWERS and ZIEGLER), A., i, 120.

1:2:4-Trimethylcyclohexanes, stereoisomeric (SKITA and SCHENCK), A., i, 241.

2:2:3:3-Tetramethylindoline, and 5-nitroso- and their salts (KNOVENAGEL), A., i, 1061.

Tetramethyl-4-methylene- $\Delta^{2,5}$ -cyclohexadienes (V. AUWERS and ZIEGLER), A., i, 121.

Tetramethylmucic acid, amide and methyl ester of (KARRER and PEYER), A., i, 809.

$\gamma\gamma\epsilon\epsilon$ -Tetramethyloctan- δ -one, η -chloro- (BILLON), A., i, 717.

Tetramethylolphosphonium chloride, and its derivatives (HOFFMAN), A., i, 8.

1:2:2:3-Tetramethylcyclopentyl benzyl ketone (RUPE and JÄGGI), A., i, 840.

1:2:2:3-Tetramethylcyclopentyl α -naphthyl ketone (RUPE and JÄGGI), A., i, 840.

1:2:2:3-Tetramethylcyclopentyl- β phenylethyl ketone (RUPE and JÄGGI), A., i, 840.

1:2:2:3-Tetramethylcyclopentyl phenyl ketone (RUPE and JÄGGI), A., i, 840.

1:2:2:3-Tetramethylcyclopentylpropylcarbinol (LEROIDE), A., i, 217.

NV' -Tetramethylphthalamide (V. BRAUN and KAISER), A., i, 659.

2:4:2':4'-Tetramethylpyrocoll-3:8'-dicarboxylic acid, ethyl ester (KÜSTER, SCHLACK, SCHLAYERRACH, and WIL-LIG), A., i, 858.

Tetramethylsaccharic acid, salts, amide and methyl ester of (KARRER and PEYER), A., i, 809.

2:2:3:8-Tetramethyl-*ar*-tetrahydro-*a*-naphthindoline, and its salts (KNOEVENAGEL), A., i, 1061.

1:3:7:9-Tetramethylthiouric acids (BILTZ and HEIDRICH), A., i, 383.

Tetrammincobaltiselenates. See under Cobalt.

2:4:5:6-Tetraphenylpyridine picrate (DILTHEY, NUSSLEIN, MEYER, and KAFFER), A., i, 949.

Tetraphthene. See Tetrahydroacenaphthene.

***s*-Tetrapropylethane.** See *δ**c*-Dipropyloctane.

Tetrapyridinecobaltiselenate, dichloro- (MEYER and MOLDENHAUER), A., ii, 73.

Tetrapyridineferritetetrachloroferrate, dihydroxy- (WEINLAND and KISSLING), A., i, 364.

Tetrapyridinium μ -dichlorodinitratohexachlorodiferrate (WEINLAND and KISSLING), A., i, 364.

Tetraquinolinium μ -dichlorooctachlorodiferrate (WEINLAND and KISSLING), A., i, 364.

Tetraquochromihexaquochromiselenate. See under Chromium.

Tetrazoles, preparation of (STOLLÉ), A., i, 689.

Tetrodon poison. See Poison.

Textile fibres. See Fibres.

Thallium, electrode potential of, and its amalgams (RICHARDS and SMYTH), A., ii, 341.

flame reactions of (PAPISH), A., ii, 528.

Thallium compounds, studies on (BERRY), T., 394.

Thallium chloride, photochemistry of (RENZ), A., ii, 66.

chlorides, action of, on organo-metallic compounds (D. and A. E. GODDARD), T., 256.

chromates (CANNERI), A., ii, 297.

cadmium nickel nitrite (CUTTICA), A., ii, 449.

arseno-, bismutho-, and stibio-thio-sulphates (CANNERI), A., ii, 378.

Thallic-thalious sulphate (BENRATH and ESPENSCHIED), A., ii, 504.

Thalious borates (CANNERI and MORELLI), A., ii, 571.

bismuth bromide and iodide (CANNERI and PERINA), A., ii, 512.

chlorite (LEVI), A., ii, 567.

Thallium:—

Thalious haloids, compounds of ammonia with (BILTZ and STOLLENWERK), A., ii, 67.

nitrite, double salts of metallic nitrites and (CUTTICA and PACIELLO), A., ii, 377.

Thallium organic compounds:—

dialkyl salts and diaryl haloids (GODDARD), T., 36; (D. and A. E. GODDARD), T., 256.

diphenyl oxide and salts (A. E. and D. GODDARD), T., 486.

Thallium estimation:—

estimation of, in organic compounds (A. E. and D. GODDARD), T., 488.

Thapsic acid, electro-synthesis of, and its ethyl ester (CARMICHAEL), T., 2545.

Thaumasite, constitution of (HOLDEN), A., ii, 860.

Theobromine, chloroform extraction apparatus for (SCHAAP), A., ii, 797.
soluble compounds of, with alkali benzoates and salicylates (KNOLL & Co.), A., i, 464.

estimation of, in urine (GUNZBERG), A., i, 703.

Theobromine-1-acetic acid (MERCK, WOLFES, and KORNICK), A., i, 1071.

Theophylline, soluble compounds of, with alkali benzoates and salicylates (KNOLL & Co.), A., i, 464.

Theophylline-7-acetic acid (MERCK, WOLFES, and KORNICK), A., i, 1071.

Thermochemistry, unit of measurement in (SWIENTOSLAWSKI), A., ii, 548.

of organic compounds (SWIENTOSLAWSKI), A., ii, 195; (FAJANS), A., ii, 818; (THIEL), A., ii, 819.

of solutions (LEVALT-EZERSKY), A., ii, 819.

Thermo-compression (DELEUVER), A., ii, 684.

Thermodynamics, relation between statistical mechanics and (TOLMAN), A., ii, 257.

of mixtures (WAGNER), A., ii, 116.

Thermo-electric force of metals (LATIMER), A., ii, 814.

Thermo-regulator (STARKEY and GORDON), A., ii, 547.

Thermos flask, use of, in the laboratory (BOURLET and THOMAS), A., ii, 139.

Thermostat to determine the effect of temperature on diffusion (ÖHOLM), A., ii, 621.

adjustable (LEWIS and WOOD), A., ii, 612.

electrical heating apparatus for (RAWLING), A., ii, 612.

- Thiazoles** (BOGERT and ABRAHAMSON), A., i, 576; (BOGERT and MEYER), A., i, 868.
- Thienoylhydroxamic acid**, and its salts and esters (JONES and HURD), A., i, 249.
- Thiobacillus thio-oxidans*, growth of (LIPMAN, WAKSMAN, and JOFFE), A., i, 303.
- Thiocarbamic acids**, decomposition of azides of (OLIVERI-MANDALÀ), A., i, 1007.
- Thiocarbamides**, preparation of (GOOD-YEAR TIRE AND RUBBER CO.), A., i, 286.
interaction of aldehydes or ketones with (TAYLOR), T., 2267.
- 3-Thiocarbamido-6-methoxybenzoic acid** (HERMANN), A., i, 1151.
- Thiocarbamine cyanides**, preparation of (FROMM and WENZL), A., i, 437.
- Thiocarbanilide**, compound of chloropierin and (RAY and DAS), T., 327.
- Thiocarbimides** (HILL and KELSEY), A., i, 1141.
- Thiocyanates** (HILL and KELSEY), A., i, 1141.
action of Grignard reagents on (ADAMS), A., i, 531.
- Thiocyanic acid**, tetra-acetylsalicylic salt (ZEMPLÉN and HOFFMANN), A., i, 563.
ethyl-, β -naphthyl- and phenyl-thiol esters (LECHER and WITTEW), A., i, 641.
phthalimidomethyl ester (PUCHER and JOHNSON), A., i, 549.
sulphur compounds of (LECHER and WITTEW; LECHER and GOEBEL), A., i, 640.
- 1:3:4-Thiodiazole**, 5-amino-2-thiol-, and its salts and derivatives (GUHA), A., i, 875.
2:5-dithiol- (LOSANITCH), T., 2544.
potassium salt, compound of chloropierin and (RAY and DAS), T., 325.
- Thiodiazoles**, synthesis of (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 377.
- 1:3:4-Thiodiazolyl sulphides**, 2-thiol- (LOSANITCH), T., 2544.
- Thionaphthen-2-carboxylic acid**, and its methyl ester (GESELLSCHAFT FÜR TEERVERWERTUNG M. B. H., WEISSGERBER and KRUBER), A., i, 567.
- Thionaphthen-2:3-dicarboxylic acid**, and its methyl ester (GESELLSCHAFT FÜR TEERVERWERTUNG M. B. H., WEISSGERBER and KRUBER), A., i, 567.
- 2-Thionaphthen-2'-indoneindigo**. See 2-Oxythionaphthenyl-2-indane-1:3-dione.
- 2-Thionaphthen-9-phenanthreneindigo**. See 10-Oxy-9-oxythionaphthenyl-6-oxyphenanthrene.
- Thionaphthensulphonic acid**, preparation of (GESELLSCHAFT FÜR TEERVERWERTUNG M. B. H., and WEISSGERBER), A., i, 1172.
- 3-Thion-2:4-dibenzoyl-1:2:4-triazole**, 5-imino- (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 379.
- 3-Thion-4:1:2-thiodiazole**, 5-amino-, lead salt and its dibenzoyl derivative (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 379.
- Thiophen series** (STEINKOPF and HEROLD), A., i, 850; (STEINKOPF and AUGSTAD-JENSEN), A., i, 851.
- Thiosulphates**. See under Sulphur.
- cis-Thiosulphatodiethylenediamine-cobaltic bromide** (DUFF), T., 453.
- Thiosulphuric acid**. See under Sulphur.
- Thioliol dimethyl ether** (MAJIMA and CHIBA), A., i, 264.
- Thorium**, *N*-series spectrum of (DOLEJSEK, DUANE, and PATTERSON), A., ii, 463.
adsorption of, by basic ferric acetate (BROWN), T., 1736.
- Thorium oxide** (*thoria*), catalytic activity of (GILFILLAN), A., i, 709.
catalytic action of, on acetates (ADKINS and KRAUSE), A., i, 422.
reduction of, by metallic tungsten (GENERAL ELECTRIC CO.), T., 2236.
- Thorium estimation** :—
estimation of, in monazite sand (HELMICK), A., ii, 164.
- Thorium-C**, emission of α -particles by (SHENSTONE and SCHLUNDT), A., ii, 465.
- Thorium-X**, oxidising properties of (LEMAI and JALOUSTRÉ), A., ii, 186.
- Thortveitite** (SCHEFFELIG), A., ii, 306.
from Madagascar (BOULANGER and URBAIN), A., ii, 517.
extraction and purification of scandium from (P. and G. URBAIN), A., ii, 504.
- Thrombin and antithrombin** (PICKERING and HEWITT), A., i, 1208.
- Thrombolytic** (ROSENMAN), A., i, 596.
- Thymine**, detection of (BAUDISCH and JOHNSON), A., ii, 238; (JOHNSON and BAUDISCH), A., ii, 328.
detection of, in the presence of sugar (DEUEL and BAUDISCH), A., ii, 670.

Thymol, preparation of (COLE; SMITH and PENFOLD), A., i, 31.

Thymolmercuri-compounds, and 6-iodo- (MAMELI and MAMELI-MANNESSIER), A., i, 1080; (MAMELI), A., i, 1082.

Thymus, effect of extirpation of, on blood-corpuscles (ASHER and MATSUMO), A., i, 298.

basic peptone in (FELIX), A., i, 702.

Thymus striatus, constituents of oil from (LEONE and ANGELESCU), A., i, 460.

Thymus vulgaris, constituents of oil from (LEONE and ANGELESCU), A., i, 357.

Thyroid, chemical constituents of (ZUNZ), A., i, 397.

new constituent of the (SAMMARTINO), A., i, 1212.

proteinogenic amines in the (SAMMARTINO), A., i, 966.

effect of, on metabolism and on the regulation of body temperature (GRAFE and v. REDWITZ; SCHENCK), A., i, 491.

effect of, on creatinine metabolism (SCHENCK), A., i, 1212.

effect of adrenal feeding on the iodine content of the (BLACK, HUPPER, and ROGERS), A., i, 966.

action of, in phloridzin diabetes (ASHER and HERRISBERGER), A., i, 293.

Thyroxin, influence of, on alcoholic fermentation (TOMITA), A., i, 972.

Tin, arc spectrum of (HEMSALECH and DE GRAMONT), A., ii, 176.

Röntgen-ray spectrum of (DE BROGLIE), A., ii, 330.

isotopes of (ASTON), A., ii, 650.

liquid, density and surface tension of (HOGNESS), A., ii, 29.

rate of solution of, in ferric alum (COLLENERG and BODFORS), A., ii, 431.

Tin tetrachloride, thermal constants of (LATIMER), A., ii, 256.

catalysis of the formation of, in the glow discharge (PANETH), A., ii, 363.

hydride (PANETH, JOHANNSEN, and MATHIES; PANETH, MATTHIAS, and SCHMIDT-HEBBEL), A., ii, 383.

oxides, hydrated (WEISER), A., ii, 853.

Stannous chloride, reduction of nitro-compounds by (GOLDSCHMIDT, STORM, and HASSEL), A., ii, 361.

oxide and its hydrates, preparation and reactions of (BURY and PARTINGTON), T., 1998.

Stannic hydroxide, amphoteric properties of (COLLINS and WOOD), T., 441.

Tin :—

Stannic acids, isomerism of (COLLINS and WOOD), T., 441.

action of hydrochloric acid on (COLLINS and WOOD), T., 1122.

action of alkali hydroxides on (COLLINS and WOOD), T., 2760.

Tin organic compounds (DRUCE), A., i, 639, 1206.

diisopropylxide and haloids and isopropyltribromide (DRUCE), T., 1861.

mercaptides (WUYTS and VANGIN-DERTAELEN), A., i, 250.

tetraethyl and tetraphenyl, action of thallic chloride on (D. and A. E. GODDARD), T., 259.

tetraphenyl, use of, in preparation of organo-metallic compounds (GODDARD, ASHLEY, and EVANS), T., 978.

Tin detection, estimation and separation :—

detection of (HELLER), A., ii, 458.

estimation of, iodometrically (KOLLER), A., ii, 530.

estimation of, volumetrically (SMITH), A., ii, 398.

estimation of, in bearing metals (NÄGEL), A., ii, 721.

estimation of, volumetrically, in red brass (MÜCK), A., ii, 722.

estimation of, and its separation from antimony, copper, and lead (KLING and LASSIEUR), A., ii, 86.

Tissues, permeability of (SCHAEFFI), A., i, 290.

non-protein constituents of, and endogenous katabolism (MITCHELL, NEVENS, and KENDALL), A., i, 897.

action of potassium salts on (CLARK), A., i, 399.

oxidation of xanthine by (MORGAN, STEWART, and HOPKINS), A., i, 1078.

animal. See Animal tissues.

animal, vegetable, and tumour, copper salts in (WHITE), A., i, 399.

subcutaneous, swelling in (MORAWITZ and DENECKE), A., i, 492.

detection of urea in (BONNET and HAUSHALTER), A., ii, 794.

estimation of chlorine in (PICO and MURTAGH), A., ii, 716.

estimation of the reducing power of (LIPSCHITZ and GOTTSCHALK), A., i, 298.

estimation of urea in (GAD-ANDRESEN), A., ii, 536.

Tissue extracts, estimation of uric acid in (STEUDER and SUZUKI), A., ii, 538.

Titanazetetra-1-piperidinium chloride (LEONARD), A., i, 863.

Toluene compounds, Me=1.
Titanium salts, photochemical reduction of (BENRATH and OBLADEN), A., ii, 781.
 reduction of nitroparaffins with (HENDERSON and MACBETH), T., 892.
Titanium trichloride, reduction with (RATHSBURG), A., i, 242.
 action of, on halogen organic compounds (BLACK, HIRST, and MACBETH), T., 2527.
tetrachloride, thermal constants of (LATIMER), A., ii, 256.
sesquioxide, catalytic activity of (GILLILLAN), A., i, 709.
 catalytic action of, on acetates ADKINS and KRAUSE), A., i, 422.
dioxide, action of light on (RENZ), A., ii, 61.
 estimation of, volumetrically, in bauxite (WINCH and CHANDRATREYA), A., ii, 459.
 sulphate, use of solutions of, in controlling the course of catalysis (HOFMANN), A., ii, 490.
 use of, in estimation of iron and copper (THORNTON), A., ii, 528.
Titanium estimation:—
 estimation of (KIKUCHI; KANÔ), A., ii, 721.
 estimation of, by reduction (TREADWELL, BLUMENTHAL, and STARKLE), A., ii, 788.
Toads, poison from. See Poison.
Tobacco, biochemistry of (PARIS), A., i, 211.
 carbon monoxide in the smoke of (ARMSTRONG and EVANS), A., i, 1226.
 estimation of nicotine in (LIOTTA), A., ii, 404.
Tobaku oil, composition and properties of (IWAMOTO), A., i, 98.
o-Tolidine hydroferrocyanide (CUMMING), T., 1296.
Toluene, physical constants of (TIMMERMANS, VAN DER HORST, and ONNES), A., ii, 258.
 compressibility of (RICHARDS, BARTLETT, and HODGES), A., ii, 28.
 equilibrium of ethyl alcohol, water, and (ORMANDY and CRAVEN), A., i, 215.
 nitration of (GIBSON, DUCKHAM, and FAIRBAIRN), T., 270.
 action of pyrosulphuryl chloride on (STEINKOPF and BUCHHEIM), A., i, 122.
 estimation of (ZABOROWSKI), A., ii, 877.
Toluene, 3:5-dibromo-, action of sodium with (FUCHS and METEL), A., i, 442.

Toluene compounds, Me=1.
Toluene, chloro-derivatives, isomeric (WAHL, NORMAND, and VERMEYLEN), A., i, 442.
 6-chloro-3-amino- (GOLDSCHMIDT and STROHMENGER), i, 1006.
dichloronitro-derivatives (DAVIES), T., 812.
 nitro-derivatives, equilibrium of, with carbazole and acenaphthene (KREMMANN and STRZELBA), A., i, 176.
m-nitro-, nitration of (BRADY), T., 328.
Toluenes, bromo-, oxidation of, with potassium permanganate (BIGELOW), A., i, 998.
 nitro-, equilibria in binary systems of (BELL and McEWEN), A., i, 726.
p-Toluenesulphuric acid-β-naphthylamine (FISCHER and MEIER), A., i, 956.
Toluenesulphuric acid-β-naphthylamine (FISCHER and MEIER), A., i, 956.
Toluene-4-azo-β-naphthol, 2:6-dichloro- (DAVIES), T., 813.
o-Toluenesulphuric acid-β-naphthylamine, and its acetyl derivative (FISCHER and SCHWAPPACHER), A., i, 957.
p-Toluene-1-azo-5-nitro-β-naphthylamine (MORGAN and CHAZAN), A., i, 181.
1-p-Toluenesulphuric acid-8-nitro-β-naphthylamine (MORGAN and GILMOUR), A., i, 380.
2-o-Toluenesulphuric acid-phenol (CHATTAWAY and HILL), T., 2758.
p-Toluenediazonium chloride, compound of bismuth trichloride and (CHALLENGER and WILKINSON), T., 103.
o-Toluenesulphonamide, electrochemical oxidation of (FICHTER and LÖWE), A., i, 242.
 oxidation of (PAMFILOV), A., i, 928.
p-Toluenesulphonamide, compound of triphenylarsine-*p*-toluenesulphonylimine and (MANN and POPE), T., 1054.
o- and p-Toluenesulphonamides, thermal analysis of the system (DOBRJANSKI), A., i, 1182.
p-Toluene-p-sulphonbenzylamide (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 999.
Toluene-p-sulphondi-p-tolylamide, and its derivatives (HALBERKANN), A., i, 1134.
3-p-Toluenesulphonecenmarin, 7-hydroxy- (TRÖGER and BOLTE), A., i, 268.
p-Toluene-p-sulphonethylamide (FARBENFABRIKEN VORM. F. BAYER & Co.), A., i, 999.

Toluene compounds, Me = 1.

- Toluenesulphonic acid**, 4-toluenesulphonmethylamidophenyl ester (SOCIETY OF CHEMICAL INDUSTRY IN BASLE), A., i, 31.
- Toluene-*m*-sulphonic acid**, 6-chloro-5-nitro-, metallic salts and amide of (DAVIES), T., 789.
- Toluene-*p*-sulphonic acid**, 1-methyl-papaverinium, 1-methylpyridinium and phenyltrimethylammonium salts (FÖLDI), A., i, 732.
- β -chloroethyl ester (CLEMO and PERKIN), T., 644.
- transformations of diarylamides of (HALBERKANN), A., i, 1132.
- Toluene-*p*-sulphonphenyl-*p'*-tolylamide**, and its derivatives (HALBERKANN), A., i, 1133.
- m*-Toluenesulphonyl chloride**, 6-chloro-, and 6-chloro-5-nitro- (DAVIES), T., 787.
- 5-Toluene-*p*-sulphonylaminosce naphthene** (FLEISCHER and SOBRANZ), A., i, 1144.
- p*-Toluenesulphonyl-*o''*-anilino-*p'*-benzeneazo- β -naphthol** (HALBERKANN), A., i, 1133.
- p*-Toluenesulphonylbenzeneazo- β -naphthol** (HALBERKANN), A., i, 1133.
- 3-*p*-Toluenesulphonylcoumarin** (TRÖGER and BOLTE), A., i, 267.
- 3-Toluene-*p*-sulphonyldextrose** (FREUDENBERG and IVERS), A., i, 524.
- Toluene-*p*-sulphonyldextrosediacetone** (FREUDENBERG and IVERS), A., i, 524.
- o*'-*p*-Toluenesulphonyldiphenylnitrosoamine** (HALBERKANN), A., i, 1132.
- 3-*p*-Toluenesulphonyl-2-methylquinoline**, and its salts and derivatives (TRÖGER and MENZEL), A., i, 269.
- p*-Toluenesulphonyl-2-nitro-3-anilino-*p*-phenetidine** (REVERDIN and ROETHLISBERGER), A., i, 538.
- p*-Toluenesulphonyl-3:5-dinitro-3-anilino-*p*-phenetidine** (REVERDIN and ROETHLISBERGER), A., i, 538.
- p*-Toluenesulphonyl-2:3-dinitro-*N*-methyl-*p*-phenetidine** (REVERDIN and ROETHLISBERGER), A., i, 538.
- p*-Toluenesulphonyl-mono-, di- and tri-nitro-*p*-phenetidines**, and their derivatives (REVERDIN and ROETHLISBERGER), A., i, 538.
- 3-*p*-Toluenesulphonyl-1:3-oxazolid-2-one** (FROMM and HONOLD), A., i, 580.
- p*-Toluenesulphonyl-*p*-phenetidine**, nitro-derivatives (REVERDIN and ROETHLISBERGER), A., i, 538.
- β -*p*-Toluenesulphonylquinophthalone** (TROGER and MENZEL), A., i, 270.

Toluene compounds, Me = 1.

- Toluene-*p*-sulphonyltetra-acetylglucose** (FREUDENBERG and IVERS), A., i, 524.
- p*-Toluenesulphonyl-*p*-tolylsulphoxid-ethylene** (FROMM and SIEBERT), A., i, 543.
- p*-Toluenesulphonyl-*p* tolylthioethane** (FROMM and SIEBERT), A., i, 543.
- Toluene-*p*-sulphonyltriacytylbromoglucose** (FREUDENBERG and IVERS), A., i, 524.
- Toluene-*p*-sulphonyltriacytylmethylglucoside** (FREUDENBERG and IVERS), A., i, 524.
- o*-Toluic acid**, 3-hydroxy- (ASAHINA and KONDO), A., i, 657.
- p*-Toluic acid**, bromo-2-amino-, and its derivatives (WHEELER and SMITHER), A., i, 332.
- Toluic acids**, solubility of, in xylenes (CHAPAS), A., i, 339.
- Toluic acids**, bromo- and chloro-derivatives, and their ethyl esters, amides, bromides, and chlorides (DAVIES and PERKIN), T., 2202.
- dinitroamino-**, methyl ester, and tri-nitro-, and their silver salts and ethyl ester (GIUA), A., i, 534.
- Toluidine hydrochloride**, formation of, from methylaniline hydrochloride (BECKMANN, CORRENS, and LIESCHE), A., i, 535.
- hydroferrocyanides** (CUMMING), T., 1293.
- o*-Toluidine**, separation of, from *p*-toluidine (CUMMING), T., 1293.
- m*-Toluidine**, di-*o*-chloro- (GOLDSCHMIDT and STROHMENGER), A., i, 1005.
- 6-iodo-**, and its hydrochloride and acetyl derivative (BURTON and KENNER), T., 680.
- 5-nitro-**, preparation of (BRADY, DAY, and ROLT), T., 527.
- p*-Toluidine**, 2:6-dichloro-, and 2:6-dichloro-3-nitro-, and their derivatives (DAVIES), T., 813.
- 2:3- and 2:5-dinitro-** (SCOTT and ROBINSON), T., 844.
- Toluidines**, dinitro- (BRADY, DAY, and ROLT), T., 526.
- Toluidine-blue**, metachromism of (SCHWARZ and HEHMANN), A., i, 930.
- p*-Toluidine-*m*-sulphonic acid**, salts of (SCOTT and COHEN), T., 2039.
- o*- and *p*-Toluidinesulphonitriles**, and their salts (OLIVERI-MANDALA), A., i, 1008.
- Toluidinoanthraquinones**, amino- (BADISCHE ANILIN- & SODA-FABRIK), A., i, 942.

Toluene compounds, Me = 1.

- 4-Toluidinobenzonitriles, 3-nitro- (MATTAAR), A., i, 251.
- o*-*p'*-Toluidinobenzoylbenzene, *o*-2'-nitro- (MAYER and FREUND), A., i, 866.
- β -*p*-Toluidinobutyric acid, ethyl ester, and its hydrochloride (STOERMER and ROBERT), A., i, 519.
- p*-Toluidinoethylitaconic acid, ethyl ester (CARRIÈRE), A., i, 319.
- 1-*p*-Toluidino-2-furyl-5-methylbenziminazole (FISCHER and MEIER), A., i, 957.
- 1-*o*-Toluidino-2-furylnaphthiminazole (FISCHER and SCHWAPPACHER), A., i, 957.
- 1-*p*-Toluidino-2-furylnaphthiminazole, and its nitrosoamine (FISCHER and MEIER), A., i, 956.
- 1-*p*-Toluidino-4-hydroxynaphthalene (KALLE & Co.), A., i, 824.
- 1-*p*-Toluidino-2-*o*-hydroxyphenyl-5-methylbenziminazole, and its diacetyl derivative (FISCHER and MEIER), A., i, 957.
- 1-*p*-Toluidino-2-*o*-hydroxyphenyl-naphthiminazole, and its diacetyl derivative (FISCHER and MEIER), A., i, 956.
- 1-*o*-Toluidino-2-hydroxyphenylnaphthiminazoles, and their derivatives (FISCHER and SCHWAPPACHER), A., i, 957.
- 3-*p*-Toluidino-5-ketoisoxazole, and its derivatives (WORRALL), A., i, 874.
- 3-*p*-Toluidino-6-ketopyrazole, and its hydrochloride and acetyl derivative (WORRALL), A., i, 875.
- 4-*p*-Toluidino-2-methylanthraquinone, 1-amino- (BADISCHE ANILIN- & SODA-FABRIK), A., i, 942.
- 1-*o*-Toluidino-2-methylnaphthiminazole, and its acetyl derivative (FISCHER and SCHWAPPACHER), A., i, 957.
- 8-*o*- and -*p*-Toluidino-1:4-naphthaquinones, 2:3-dibromo-5-hydroxy- (WHEELER and ANDREWS), A., i, 354.
- 1-*o*-Toluidino- α -naphthiminazole, and its nitrosoamine (FISCHER and SCHWAPPACHER), A., i, 957.
- β -Toluidino-*p*-phenetolecarbamides (SPECKAN), A., i, 580.
- 1-*o*-Toluidino-2-phenyl- α -naphthiminazole, and its picrate (FISCHER and SCHWAPPACHER), A., i, 957.
- 1-*p*-Toluidino-2-phenyl- α -naphthiminazole, and its nitrosoamine (FISCHER and MEIER), A., i, 957.
- 2-*p*-Toluidinophenyl-*p*-tolylsulphone, and its derivatives (HALBERKANN), A., i, 1133.

Toluene compounds, Me = 1.

- 5-*m*-Toluidino-1-*m*-tolyl-1:2:3:4-tetraazole (STOLLE), A., i, 689.
- p*-Toluoilaminobenzoic acids (LAW-RANCE), A., i, 340.
- p*-Toluoil-*o*-benzoic acid, preparation of (MCMULLEN), A., i, 140.
- p*-Toluoil-*m*-benzoic acid, and its silver salt and methyl ester (SMITH), A., i, 141.
- Toluoilbenzoic acids, aminohydroxy-, hydroxy-, and nitrohydroxy- (EDER and WIDMER), A., i, 261.
- tetrabromo-, tetraiodo- and nitro-, and their methyl esters (LAWRANCE), A., i, 340.
- m*-Toluoilbenzoic acids, 3:4- and 3:6-dichloro-2-*p*-hydroxy- (v. DEM KNESEBECK and ULLMANN), A., i, 359.
- o*-Tolyl- β -chloroethyl ether, and *p*-nitro- (CLEMO and PERKIN), T., 645.
- p*-Tolyl *p'*-amino- and *p'*-nitro-benzyl ethers (PUMMERER, MELAMED, and PUTTFARCKEN), A., i, 1163.
- methyl ether, 3-chlorosulphonyl derivative, preparation of (STEWART), T., 2558.
- p*-Tolyl-*o*-acetoxystyrylsulphone (TROGER and BOLTE), A., i, 268.
- N*-*o*-Tolyl- β -aminoethyl hydrogen sulphate (SAUNDERS), T., 2673.
- 3-Tolyl-5-anilinomethylene-4-glyoxalones, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- p*-Tolyl-1-anilino-*m*-tolylsulphone, and its derivatives (HALBERKANN), A., i, 1134.
- o*- and *p*-Tolylarsines (PALMER and ADAMS), A., i, 786.
- 1-*p*-Tolylbenzothiazole, and its derivatives (BOGERT and MEYER), A., i, 868.
- p*-Tolyl-*o*-benzoyloxystyrylsulphone (TROGER and BOLTE), A., i, 268.
- 1-*m*-Tolyl-4-benzylidenehydantoin, and 2-thio- (HILL and KELSEY), A., i, 1142.
- p*-Tolylbiuret (FROMM and WENZL), A., i, 438.
- α -*p*-Tolylisobutyric acid, $\beta\beta$ -dichloro- (v. AUWERS and ZIEGLER), A., i, 140.
- p*-Tolylcarbamine cyanide (FROMM and WENZL), A., i, 438.
- Tolyl-2:4-diacetoxystyrylsulphone (TROGER and BOLTE), A., i, 268.
- Tolyl-2:4-dibenzoyloxystyrylsulphone (TROGER and BOLTE), A., i, 268.
- 1-*p*-Tolyl-2:3-dibenzyl-6-methylquin-oxaline, *aa'*-dicyano- (DUTT and SEN), T., 2666.

Toluene compounds, Me = 1.

- o*- and *p*-Tolyldi- α -hydroxybenzylarsines (PALMER and ADAMS), A., i, 786.
- o*- and *p*-Tolyldi- α -hydroxyethylarsines (PALMER and ADAMS), A., i, 786.
- m*- and *p*-Tolyldiphenylmethane, dihydroxy- (MACKENZIE), T., 1698.
- 2:4-Tolylene-diamine hydroferrocyanide (CUMMING), T., 1295.
- compound of tellurium tetrabromide with (LOWY and DUNBROOK), A., i, 446.
- p*-Tolyethylcarbinol (v. AUWERS and KOLLIGS), A., ii, 176.
- 1-*p*-Tolyl-4-ethylpyrroline-5-one-3-carboxylic acid, and its ethyl ester (CARRIERE), A., i, 319.
- p*-Tolylfuryl-2-keto-3-methylcyclohexylmethane (WOLFF), A., i, 668.
- 3-*m*-Tolyl-4-glyoxalone, 2-thiol- (DAINS, THOMPSON, and ASENDORF), A., i, 1185.
- 1-*m*-Tolyldihantoin, and 2-thio- (HILL and KELSEY), A., i, 1142.
- p*-Tolyldiazine, *m*-nitro-, preparation of (DAVIES), T., 720.
- p*-Tolyldiazinesulphonic acid, *m*-nitro-, ammonium salt (DAVIES), T., 720.
- p*-Tolyl-*o*-hydroxystyrylsulphone (TROGER and BOLTE), A., i, 267.
- Tolyl-2:4-dihydroxystyrylsulphone, and its dimethyl ether (TROGER and BOLTE), A., i, 268.
- m*-Tolyldieneanthranilic acid (EKELEY, ROGERS, and SWISHER), A., i, 935.
- 5-Tolylimino-2-thio-3-naphthyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole (GUHA), A., i, 877.
- 5-Tolylimino-2-thio-3-phenyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole (GUHA), A., i, 877.
- 5-Tolylimino-2-thio-3-tolyl-2:3:4:5-tetrahydro-1:3:4-thiadiazole (GUHA), A., i, 876.
- m*-Tolyl-6-methylbenzimidazole 1:6-hydroxy-, and its salts (JACOBSEN), A., i, 593.
- 2-*m*-Tolyl-3-methylindole, 2-*p*-hydroxy- (v. AUWERS, HILLIGER, and WULF), A., i, 1193.
- o*-Tolylmethylnitrosamine (SCHMIDT and FISCHER), A., i, 822.
- 2-*p*-Tolyl- α -8-naphthatriazole, 6-amino-, and its derivatives and 6-nitro- (MORGAN and CHAZAN), A., i, 181.
- 9-amino-, and 9-nitro- (MORGAN and GILMOUR), A., i, 380.
- p*-Tolyl- α -8-naphthatriazole- α - β -naphthol (MORGAN and CHAZAN), A., i, 181.

Toluene compounds, Me = 1.

- α -*o*- and *p*-Tolynaphthylamines, β -hydroxy- (WAHL and LANTZ), A., i, 823.
- Tolyloxides, nitro-, lead (GODDARD and WARD), T., 265.
- Tolyloxyacetic acids, amino-, nitro-, and their derivatives (MINTON and STEPHEN), T., 1594.
- 2- and 3-*p*'-Tolyloxy-*p*-cresols, and their derivatives (PUMMERER, MELAMED, and PUTTFARCKEN), A., i, 1163.
- 2-*p*-Tolyloxy-2:3-dihydro-*p*-benzoquinomethane, and its derivatives (PUMMERER, MELAMED, and PUTTFARCKEN), A., i, 1162.
- α -*p*-Tolylpropionic acid, $\beta\beta\beta$ -trichloro- (v. AUWERS and JULICHER), A., i, 842.
- β -*o*-Tolylpropionolactone, β :3:5-dihydroxy- (LANGLEY and ADAMS), A., i, 1154.
- α -*p*-Tolylprozan- β -dicarboxylic acid, ethyl ester (DIELS and AUBART), A., i, 1194.
- 1-*p*-Tolylpyrroline-5-one-3-carboxylic acid, ethyl ester (CARRIERE), A., i, 318.
- o*-Tolylsulphamic acid, *p*-amino-, benzoyl derivative, sodium salt (WEIL and WASSERMANN), A., i, 1006.
- p*-Tolylsulphamic acid, sodium salt (WEIL and MOSER), A., i, 444.
- o*-Tolyl-*p*-sulphamsulphonic acid, amino-, sodium salt (WEIL and WASSERMANN), A., i, 1007.
- p*-Tolylsulphonyl-*p*-tolylthioethylene (FROMM and SIEBERT), A., i, 543.
- 1-*p*-Tolyl-1:2:3:4-tetrazole, 5-amino- (STOLLÉ), A., i, 689.
- p*-Tolylthioacetaldehyde, phenylhydrazones and tolylmercaptal derivatives of (FROMM and SIEBERT), A., i, 542.
- p*-Tolylthiothioacetic acid, *p*-tolyl ester (FROMM and SIEBERT), A., i, 543.
- m*-Tolyltoluquinonedimide (GOLDSCHMIDT and STROHMENGER), A., i, 1006.
- Tomato (*Solanum esculentum*), proteins of the seeds of (JOHNS and GERSDORFF), A., i, 800.
- Toxicity (RONA and BLOCH; RONA and BACH), A., i, 65; (RONA and REINICKE), A., i, 67; (RONA, AIRILA, and LASNITZKI), A., i, 959.
- Toxins, formation of, by bacteria (WALBUM), A., i, 902.
- chemical nature of (SALKOWSKI), A., i, 1216.
- adsorption of, by aluminium hydroxide (RAKUZIN), A., i, 1199.
- action of metals on (ERDSTEIN and FÜRTH), A., i, 90.

- Trees**, Indian forest, oils and fats from the seeds of (RAU and SIMONSEN), A., i, 1100.
- Trehalose**, fermentation of, by bacilli (KOSER), A., i, 407.
- Triacetin**, hydrolysis of (YAMAZAKI), A., i, 422.
- 3:4:5-Triacetoxystyrene**, ω -nitro- (ROSENMUND and PFANNKUCH), A., i, 1080.
- 7 ϵ -Triacetyl- β -chloroacetylglucose**, α -chloro- (BRIGL), A., i, 225.
- 7 ϵ -Triacetyl- β -trichloroacetylglucose**, α -chloro- (BRIGL), A., i, 225.
- Triacetyl-ethylglucoside- ζ -bromohydrin** (WREDE), A., i, 226.
- Triacetyl-gallaldehyde**, and its derivatives (ROSENMUND and PFANNKUCH), A., i, 1080.
- 7 ϵ -Triacetylglucose**, α - and β -forms of, and its $\alpha\beta$ -anhydride (BRIGL), A., i, 1117.
- 7 ϵ -Triacetylglucose**, α -chloro-, and its β -chlorosulphinite and $\alpha\beta$ -dichloro- (BRIGL), A., i, 225.
- 3:5:8-Triacetyl-1-methyl glucosamine** (IRVINE and EARL), T., 2381.
- 7 ϵ -Triacetyl- α -methylglucoside** (BRIGL), A., i, 1117.
- Trisoamylphosphine-ethylimine** (STAUDINGER and HAUSER), A., i, 70.
- Trisoamylphosphinephenylimine** (STAUDINGER and HAUSER), A., i, 70.
- Trisoamylphosphinephenylphosphazide** (STAUDINGER and HAUSER), A., i, 70.
- Trianilinearsine** hydrochloride (SCHMIDT), A., i, 285.
- 3:4:5-Tri-*p*-anisylisooxazole** (MEISENHEIMER and WEIBEZAHN), A., i, 177.
- Triaquotripyridinemagnesium chloride** (SPACU), A., i, 859.
- Triazo-group** (FORSTER and SAVILLE), T., 2595.
- 3-Triazocarbazole** (MORGAN and READ), T., 2714.
- α -Triazo- β -hydroxy- β -phenylpropionic acid** (FORSTER and SAVILLE), T., 2595.
- 1:2:3-Triazole**, 4:5-dicyano-, and its derivatives (GRISCHKEVITSCH-TROCHIMOVSKI), A., i, 724.
- 1:2:4-Triazole**, 5-amino-3-thiol-, and 3:5-dithiol-, and their salts (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 379.
- Triazoles** (ARNDT, MILDE, TSCHENSCHER, BIELICH, and ECKERT), A., i, 277.
- synthesis of (FROMM, KAYSER, BRIEGLEB, and FÖHRENBACH), A., i, 377.
- Triazoles**, aromatic, preparation of (KALLE & Co.), A., i, 61.
- 1:2:3-Triazole-5-carboxylic acid**, 4-cyano-, ethyl ester (GRISCHKEVITSCH-TROCHIMOVSKI), A., i, 724.
- α -Triazo- β -methoxy- β -phenylpropionic acid** (FORSTER and SAVILLE), T., 2601.
- Triazosulphole**, and its compound with mercuric oxide (OLIVERI-MANDALA), A., i, 1008.
- Tribenzoylgallaldehyde**, and its derivatives (ROSENMUND and PFANNKUCH), A., i, 1030.
- Tribenzoylphenylmethane** (MEISENHEIMER and WEIBEZAHN), A., i, 178.
- Tribenzylamine**, *tri-o*-hydroxy-, hydrochloride (ZEMPLÉN and KUNZ), A., i, 564.
- Tribenzylmethyl chloride** (JONES and SCOTT), A., i, 454.
- Trisobutylcarbinol** (SKRAUP and FREUNDLICH), A., i, 540.
- Tributyrlinase**, action of kaolin on the activity of, in serum (OLSEN), A., i, 292.
- 2:4:4-Tricarboxycyclobutane-1-acetic-3- α -propionic acid**, 2-cyano-, ethyl ester (INGOLD, PERREN, and THORPE), T., 1788.
- 2:4:4-Tricarboxycyclobutane-1-malonic-3- α -propionic acid**, 2-cyano-, ethyl ester (INGOLD, PERREN, and THORPE), T., 1788.
- Trichilia emetica**, mafurra tallow from the nuts of (RINDL), A., i, 1102.
- isoTricosole acid**, and its ethyl ester (LEVENE and TAYLOR), A., i, 715.
- isoTricosyl iodide** (LEVENE and TAYLOR), A., i, 715.
- isoTricosyl alcohol** (LEVENE and TAYLOR), A., i, 715.
- isoTricosylmalonic acid**, and its diethyl ester (LEVENE and TAYLOR), A., i, 715.
- Tridymite** from Italy (BILLOWS), A., ii, 578.
- 2:4:5-Tri-*p*-ethoxyphenyloxazole** (SCHÖNBERG and KRAEMER), A., i, 664.
- Triethylamine**, action of, on ketopinyl chloride (WEDEKIND and WEINAND), A., i, 549.
- Triethylammonium ruthenipentabromide** (GUTBIEB and KRAUSS), A., i, 16.
- Triethylene tri- and tetra-sulphides**, and their compounds with mercuric salts, and with ethyl iodide (RAY), T., 1279.
- Triethylphosphinebenzoylimine** (STAUDINGER and HAUSER), A., i, 70.
- Triethylphosphine-ethylimine** (STAUDINGER and HAUSER), A., i, 70.

Triethylphosphineimines (STAUDINGER and HAUSER), A., i, 70.
Triethylphosphinemethylimine (STAUDINGER and HAUSER), A., i, 70.
Triethylphosphinephenylimine (STAUDINGER and HAUSER), A., i, 70.
Triethylsulphonium bromide, absorption spectra of, in various solvents (TAYLOR and LEWIS), T., 665.
 β -Trigalloyllavoglucosan, and its potassium salt (KARRER and SALOMON), A., i, 267.
Triglycolamidic acid, tri-*p*-anisylamide of (HILL and KELSEY), A., i, 1141.
Trihexosan, and its nona-acetyl derivative (PICTET and JAHN), A., i, 937.
2:3:5-Triketo-6- α -methoxybenzyl-1:4-dimethylpiperazine (FORSTER and SAVILLE), T., 824.
2:3:5-Triketo-6- α -methoxybenzyl-1(or 4)-methylpiperazine (FORSTER and SAVILLE), T., 824.
2:4:5-Trimercuriacetanilide, acetate of (RONNI), A., i, 605.
2:3:5-Trimethoxybenzoic acid (FALTIS and NEUMANN), A., i, 570.
2:4:4'-Trimethoxybenzophenone (ZIEGLER and OCHS), A., i, 1049.
2:4:5-Trimethoxybenzylideneanthranilic acid (EKELEY, ROGERS, and SWISHER), A., i, 935.
6:7:8-Trimethoxy-1:2-dimethyl-1:2:3:4-tetrahydroisoquinoline, salts of (SPATH), A., i, 165.
7:5:6'-Trimethoxy-2-ethyl-3:4-indeno-(2:1')-benzopyrylium ferrichloride (CRABTREE and ROBINSON), T., 1037.
Trimethoxy- β -gambier-catechinacarbonylic acid (NIERENSTEIN), T., 28.
6:7:8-Trimethoxy-1-methyl-3:4-dihydroisoquinoline, and its salts (SPATH), A., i, 165.
7:5:6'-Trimethoxy-2-methyl-3:4-indeno-(2:1)-benzopyrylium ferrichloride (CRABTREE and ROBINSON), T., 1036.
6:7:8-Trimethoxy-1-methyl-1:2:3:4-tetrahydroisoquinoline, salts of (SPATH), A., i, 165.
2:3:4-Trimethoxyphenyl styryl ketone (BRAND and COLLISCHONN), A., i, 452.
6:7:8-Trimethoxy-1:2:3:4-tetrahydroisoquinoline, and its salts (SPATH), A., i, 165.
Trimethylamine chlorite (LEVI), A., i, 528.
 hydroferrocyanide (CUMMING), T., 1297.
Trimethylammonium ruthenipentabromide (GUTHRIE and KRAUSS), A., i, 16.

2:3:5-Trimethylaniline, 4(?)-chloro- (GOLDSCHMIDT and STROHMENGER), A., i, 1005.
2:3:4-Trimethylbenzaldehyde, 6-hydroxy-, and its semicarbazone (V. AUWERS and ZIEGLER), A., i, 121.
2:3:5-Trimethylbenzene, 1-dichloro-amino- (GOLDSCHMIDT and STROHMENGER), A., i, 1005.
 $\beta\beta\gamma$ -Trimethylbutane (CHAVANNE and LEJEUNE), A., i, 417.
 $\beta\gamma\gamma$ -Trimethyl- $\Delta\alpha$ -butylene (CHAVANNE and LEJEUNE), A., i, 417.
 $\alpha\alpha\beta$ -Trimethylbutyric acid, β -hydroxy-, ethyl ester (LEROIDE), A., i, 218.
Trimethylcamphorylmethylammonium bromide, preparation of (RUPE), A., i, 666.
1:2:3-Trimethyl-5-di- ω -chloroethylbenzene (V. AUWERS and ZIEGLER), A., i, 121.
1:2:6-Trimethyl-1-dichloromethyl- $\Delta^{2,5}$ -cyclohexadien-4-one, and its derivatives (V. AUWERS and ZIEGLER), A., i, 121.
1:3:6-Trimethyl-1-dichloromethyl- $\Delta^{2,5}$ -cyclohexadien-4-one, 5-chloro- (V. AUWERS and ZIEGLER), A., i, 145.
1:7:9-Trimethyldeoxyuric acid, and its salts (BILLIZ and BULOW), A., i, 383.
3:4:5-Trimethyl-2:6-dibromomethylphenol (V. AUWERS and SAURWEIN), A., i, 1033.
3:4:5-Trimethyl-2:6-dihydroxymethylphenol (V. AUWERS and SAURWEIN), A., i, 1033.
Trimethylene chlorobromide, action of, on aliphatic ketones (BILLON), A., i, 717.
 dinitrate (BLECHTA), A., i, 518.
2:4:5-Tri-*mp*-methylenedioxyphenyl-oxazole (SCHONBERG and KRAEMER), A., i, 665.
Trimethylene glycol, poly-ethers of (ROJAHN), A., i, 103.
 preparation of, β -bromo- and β -chloropropionic acids from (ROJAHN), A., i, 105.
 estimation of, in crude glycerol (COOKS and SALWAY), A., ii, 232.
***peri*-Trimethylenenaphthalene**, and its derivatives (FLEISCHER and RETZE), A., i, 1188.
Trimethylethylene. See β -Methyl- $\Delta\beta$ -butylene.
1:1:3-Trimethyl-4-ethyl- $\Delta^{2,5}$ -cyclohexadiene (V. AUWERS and ZIEGLER), A., i, 120.
1:1:3-Trimethyl-4-ethyl- $\Delta^{2,5}$ -cyclohexadien-4-ol (V. AUWERS and ZIEGLER), A., i, 120.

- Trimethylethylmethane.** See $\beta\beta$ -Dimethylbutane.
- 5:1':6'-Trimethyl-2-ethylthioisocyanine iodide** (BRAUNHOLTZ and MILLS), T., 2007.
- Trimethylisoeugenolalkamine**, and its hydrochloride (TAKEDA and KURODA), A., i, 275.
- 2:3:5-Trimethylglucose**, oxidation of (HAWORTH and LEITCH), T., 1929.
- 2:3:6-Trimethylglucose**, and its derivatives (IRVINE and HIRST), T., 1213.
- $\alpha\beta\beta$ -Trimethylglutaric acid**, $\alpha\alpha'$ -dicyano-, *N*-methyl- ω -imide (KON and THORPE), T., 1799.
- $\beta\beta\beta$ -Trimethylheptan γ -one**, η -chloro- (BILLON), A., i, 717.
- Trimethylhomogalloylhomoveratrylamine**, and its derivatives (SPATH and BÖHM), A., i, 1175.
- Trimethylmelamine**, *tricyano-* (MADELUNG and KERN), A., i, 440.
- 1:1:2-Trimethyl-4-methylene- $\Delta^{2,5}$ -cyclohexadiene** (v. AUWERS and ZIEGLER), A., i, 120.
- 2:3:6-Trimethyl methylglucoside** (IRVINE and HIRST), T., 1221.
- Trimethyl methylxyloside**, preparation of (CARRUTHERS and HIRST), T., 2304.
- Trimethylmuic acid**, methyl ester (KARRER and PEYER), A., i, 809.
- 3:4:5-Trimethylphenol**. See Hemimelitenol.
- 3:5:6-Trimethylphenol**. See *iso*- ψ -Cumenol.
- $\beta\gamma\gamma$ -Trimethyl- δ -propylheptan- $\beta\delta$ -diol** (LEROIDE), A., i, 218.
- Trimethylisopropylmethane**. See $\beta\beta\gamma$ -Trimethylbutane.
- $\alpha\beta\beta$ -Trimethylpropylphosphinic acid**, α -hydroxy-, lead salt (CONANT, MACDONALD, and KINNEY), A., i, 186.
- 2:4:6-Trimethylpyridine** platinichloride (EMMERT and WEBB), A., i, 680.
- 2:4:6-Trimethylpyridinium benzyl iodide** and perchlorate (WEITZ and KONIG), A., i, 1188.
- 2:4:6-Trimethylquinoline**, and its salts (KNOEVENAGEL and BARR), A., i, 751.
- 1:7:9-Trimethyl-8-thiouric acids** (BILTZ and BÜLOW), A., i, 383.
- Trimethyltricarballic acid**, methyl ester and anhydride of (HOPE and SHELTON), T., 2235.
- Trinitrides**. See Azides.
- 1:2:4-Trioxo-5:8-dihydroxyanthraquinone**, aminocobaltic salt (MORGAN and SMITH), T., 168.
- α -Trioxymethylene**, preparation of (HAMMICK and BOERER), T., 2738.
- Triphenylacethydroxamic acid**, and its acetyl ester, and their salts (JONES and HURD), A., i, 249.
- Triphenyl-3-acetyl-2:4-dimethylpyrrylmethane** (FISCHER and LUCKMANN), A., i, 276.
- Triphenylallene** (MEYER and SCHUSTER), A., i, 540.
- Triphenylamines**, *o*- and *m*-amino-, and their derivatives (PICCARD and BREWSTER), A., i, 370.
- Triphenylarsine bromocyanide** (STEINKOPF and SCHWEN), A., i, 72.
- Triphenylarsine-*p*-toluenesulphonylimine**, compound of *p*-toluenesulphonamide with (MANN and POPE), T., 1054.
- Triphenylbiphenylene-ethyl** and its chloride (SCHLENK and MARK), A., i, 1004.
- Triphenylbismuthine dichloride** and difluoride (CHALLENGER and WILKINSON), T., 96.
- Triphenylbismuthine**, *tri-p*-bromo- and *tri-p*-chloro-, and their dibromides and dichlorides (CHALLENGER and RIDGWAY), T., 108.
- 4:5:6-Triphenyl-2-(*p*-bromophenyl)pyridine**, and its picrate (DILTHEY, NÜSSLEIN, MEYER, and KAFFER), A., i, 949.
- Triphenyl-3-carbethoxy-2:4-dimethylpyrrylmethane** (FISCHER and LUCKMANN), A., i, 276.
- Triphenylcarbinol**, equilibria of, with pyrogallol, nitrophenols, nitrobenzenes, and phenylenediamines (KREMANN, HOHL, and MÜLLER), A., i, 138.
- Triphenylcarbinol**, *pp''*-*tri*thiol- (WATSON and DUTT), T., 1940.
- Triphenyl-*p*-dimethylaminophenylmethane** (FISCHER and LUCKMANN), A., i, 276.
- 1:3:4-Triphenyldipyrazole** (ROJAHN), A., i, 373.
- Triphenylethynylcarbinol**, preparation and reactions of (HESS and WELTZIEN), A., i, 35.
- Tri-*p*-phenylethylcarbinol** (HESS and WELTZIEN), A., i, 36.
- 1:3:5-Triphenyl- $\Delta^{1,4}$ -cycloheptadiene** (GASTALDI), A., i, 367.
- Triphenylhydrazine**, action of tetranitromethane on (GOLDSCHMIDT and RENN), A., i, 478.
- 1:2:6-Triphenyl-4-*p*-hydroxyphenylpyridinium chloride** (DILTHEY), A., i, 272.
- Triphenylmethane**, coloured alkali salts of (HANTZSCH), A., i, 26.
- halochromism of derivatives of (HANTZSCH), A., i, 26.

- Triphenylmethane**, equilibria of amines and phenols with (KREMANN, ODELGA, and ZAWODSKY), A., i, 181.
- unsaturated halogen derivatives of, and their conductivity (STRAUS and DÜTZMANN), A., i, 148.
- Triphenylmethane colouring matters**, absorption spectra of (ADINOLFI), A., ii, 601.
- free radicals of (WIELAND, POPPER, and SEEFRIED), A., i, 772.
- peroxidation products of the leucobases of (KEHRMANN, ROY, and RAMM), A., i, 467.
- Triphenylmethanesulphonic acids**, photochemistry of (HOLMES), A., ii, 465.
- Triphenylmethyl** (GOMBERG and SULLIVAN), A., i, 929.
- α -Triphenylmethylaminocamphor** (RUPE and SCHMID), A., ii, 604.
- Triphenylmethylarsonium bromide**, and its compound with iodoform (SREIN-KOPF and SCHWEN), A., i, 118.
- Triphenylsacoxazole**, *tri-p*-nitro- (MEI-SSENHEIMER and WEIBEZAHN), A., i, 177.
- 2:4:4-Triphenyl-1:3-oxthiophan-5-one**, action of sulphuric acid on (BIS-TRYCKI and BRENNEN), A., i, 268.
- Triphenylpararosaniline hydrochloride**, preparation of (HORIUCHI), A., i, 1067.
- Triphenylphosphine**, phosphazines from (STAUDINGER and LÜSCHER), A., i, 237.
- Triphenylphosphineacetylglxylic acid**, ethyl ester, azine of (STAUDINGER and LÜSCHER), A., i, 238.
- Triphenylphosphinebenzaldehydeazine**, preparation and reactions of (BRAUN-HOLTZ), T., 305.
- Triphenylphosphinebenzilazine** (STAUDINGER and LÜSCHER), A., i, 237.
- Triphenylphosphinebenzophenoneazine**, reactions of (BRAUNHOLTZ), T., 301.
- Triphenylphosphinebenzoylimine** (STAUDINGER and HAUSER), A., i, 69.
- Triphenylphosphine-ethylimine** (STAUDINGER and HAUSER), A., i, 69.
- Triphenylphosphineglyoxylic acid-azine**, ethyl ester, reactions of (BRAUN-HOLTZ), T., 304.
- Triphenylphosphineimineazide** (STAUDINGER and HAUSER), A., i, 69.
- Triphenylphosphinemethylimine** (STAUDINGER and HAUSER), A., i, 69.
- Triphenylphosphine- α -naphthylazide** (STAUDINGER and HAUSER), A., i, 69.
- Triphenylphosphine- α -naphthylimine** (STAUDINGER and HAUSER), A., i, 69.
- Triphenylphosphinedinitroquinoneazine** (STAUDINGER and LÜSCHER), A., i, 238.
- $\alpha\beta\gamma$ -Triphenylpropane- $\alpha\beta$ -diol**, dehydration of (ORÉKHOFF and TIFFENEAU), A., i, 438.
- 2:4:6-Triphenylpyrylium nitrate** (GAST-ALDI), A., i, 367.
- 1:2:6-Triphenyl-4-quinopyridan** (DIL-THY), A., i, 272.
- Triphenylstibine**, and *tri-p*-amino-, derivatives of (SCHMIDT), A., i, 1203.
- Triple salts**. See Salts.
- Tripropylcarbinol phenylurethane** (LE-ROIDE), A., i, 217.
- Tripyridineiridium**, trichloro- (DELÉ-PINE), A., i, 860.
- Tripyridinium hexachloroferrate** and μ -dichloro-heptachloropyridinediferate (WEINLAND and KISSLING), A., i, 364.
- Trisalicinamine**, dodeca-acetyl deriv-ative (ZEMPLEN and KUNZ), A., i, 564.
- 2:4:6-Tri-*p*-tolylpyridine** (GASTALDI), A., i, 368.
- Tri-(triacetylalloyl)lævoglucosan** and its potassium salt (KARRER and SALOMON), A., i, 266.
- Tri-trimethylene glycol ether diacetate** (ROJAHN), A., i, 103.
- $\alpha\beta$ -Trisovalerylaminobutane** (WIND-AUS, DÖRRER, and JENSEN), A., i, 60.
- Trivinylarsenic oxide**, $\beta\beta'\beta''$ -trichloro- (MANN and POPE), T., 1757.
- Trivinylarsine**, $\beta\beta'\beta''$ -trichloro-, salts (MANN and POPE), T., 1757.
- Trivinylarsine-*p*-toluenesulphonylimine**, $\beta\beta'\beta''$ -trichloro- (MANN and POPE), T., 1758.
- Trivinylhydroxyarsonium nitrate**, $\beta\beta'\beta''$ -trichloro- (MANN and POPE), T., 1757.
- Trivinylmethylarsonium iodide**, $\beta\beta'\beta''$ -trichloro- (MANN and POPE), T., 1758.
- Trochic acid** (YANAGISAWA and TAKA-SHIMA), A., i, 652.
- Trochodiol** and its phenylcarbamate (NISHIZAWA), A., i, 653.
- Trochophytosterol**, and its acetyl deriv-ative (YANAGISAWA and TAKA-SHIMA), A., i, 652.
- Trollite** from California (EAKLE), A., ii, 858.
- Tropic acid**, resolution of, and its salts (KING and PALMER), T., 2577.
- esters of (v. BRAUN, BRAUNSDORF, and RÄTH), A., i, 769.
- fate of, in the body (KAY and RAPER), A., i, 1093.

- Tropinecarboxylic acid**, and its methyl ester (MERCK, WOLFES, and MAEDER), A., i, 1173.
- Tropinone** (MERCK, WOLFES, and MAEDER), A., i, 1174.
- Tropinonecarboxylic acid**, esters, preparation of (MERCK and WOLFES), A., i, 567; (MERCK, WOLFES, and MAEDER), A., i, 568, 1173; (WILLSTATTER, WOLFES, and MAEDER; WOLFES and MAEDER), A., i, 938.
- Tropinonedicarboxylic acid**, ethyl ester (MERCK, WOLFES, and MAEDER), A., i, 1173.
- N- β -Tropoxyethylpiperidine**, and its acetyl derivative, and their hydrochlorides (v. BRAUN, BRAUNSDORF, and RATH), A., i, 760.
- N- γ -Tropoxypropylpiperidine** hydrochloride (v. BRAUN, BRAUNSDORF, and RATH), A., i, 760.
- Trout**. See *Trutta fario*.
- Trust-deed** establishing the Harrison Memorial Fund, T., 2917.
- Trutta fario* (trout), constituents of the eggs of (FAURÉ-FREMIET and GARBAULT), A., i, 700.
- Truxillic acids**, constitution of (DE JONG), A., i, 839.
- configuration of (STOERMER and BACHER), A., i, 830.
- Truxinic acids**, configuration of (STOERMER and BACHER), A., i, 830.
- δ -Truxinic acids**, and their salts and derivatives (STOERMER and BACHER), A., i, 832.
- Trypflavin**. See 1-Methylacridinium chloride, 3:6-diamino-.
- Trypsin**, influence of reaction on the action of (RINGER), A., i, 282.
- inactivation of (NORTHROP), A., i, 282.
- kinetics of digestion by (NORTHROP), A., i, 693.
- digestion of casein with (FRÄNKEL and FELDSEBERG), A., i, 184.
- action of, on diastase (BIEDERMANN), A., i, 480.
- estimation of (KAI), A., ii, 672.
- Tryptophan**, content of, in foods (v. FURTH and LIEBEN), A., i, 293; (IDE), A., i, 414.
- detection of, microchemically, in plants (KRETZ), A., ii, 668.
- estimation of, in proteins (FOLIN and LOONEY), A., ii, 539; (LÜSCHER), A., i, 1199.
- d*-Tryptophan anhydride** (FRÄNKEL and FELDSEBERG), A., i, 184.
- l*-Tryptophan**, action of *Bacillus proteus* on (SASAKI and OTSUKA), A., i, 302.
- Tschernigite**, from Wyoming (ERICKSON), A., ii, 304.
- Tuberculosis**, chlorine metabolism in (BOENHEIM), A., i, 1092.
- lipoids of blood in (HENNING), A., i, 963.
- protein requirements in (McCANN), A., i, 497.
- Tumours**, amyloid, constituents of (EPFINGER), A., i, 497.
- Tungsten**, Röntgen-ray spectrum of (YOSHIDA and TANAKA), A., ii, 805.
- valency of, in the octacyanide (COLLEMBERG), A., ii, 508.
- attempts to decompose, at high temperatures (WENDT and IRION), A., ii, 773.
- and its oxides, equilibria of, with hydrogen and water vapour and with carbon oxides and oxygen (v. LIEMPT), A., ii, 301.
- reduction of thorium oxide by (GENERAL ELECTRIC Co.), T., 2236.
- Tungsten alloys**, analysis of (SEEL), A., ii, 876.
- with carbon and iron (DAEVES), A., ii, 70.
- with cobalt (KREITZ), A., ii, 381.
- Tungsten dichloride** (LINDNER), A., ii, 509.
- hydroxide, preparation of hydrosols of (KROGER), A., ii, 212, 213.
- oxide, catalytic activity of (GILLFILLAN), A., i, 709.
- trioxide**, crystal structure of, and its hydrates (BURGER), A., ii, 508.
- green colour of (VAN LIEMPT), A., ii, 73.
- hydrates of (HUTTIG and KURRE), A., ii, 773.
- Tungstates** (SMITH), A., ii, 774.
- complex, preparation of (LOTTERMOSER), A., ii, 510.
- Tungsten detection, estimation and separation**:—
- detection of (VAN LIEMPT), A., ii, 787.
- estimation of (LAVERS), A., ii, 164.
- colloidal, estimation of, in tungsten powder (LOTTERMOSER), A., ii, 230.
- estimation of aluminium in (V. and K. FROBOESE), A., ii, 397.
- estimation of molybdenum in (HALL), A., ii, 660.
- separation of, from molybdenum (MERRILL), A., ii, 229.
- Tungsten furnace**. See Electric furnace.
- Turbidity**, standard for (BECHHOLD and HEBLER), A., ii, 693.
- reversible (LIESEGANG), A., ii, 369.
- Turkey**. See *Meleagris gallinapavo*.
- Turpentine oil**, Aleppo, composition of (DUPONT), A., i, 857; (VÉZES and DUPONT), A., i, 1043.

- Tyramine**, estimation of (HANKE and KOESSLER), A., ii, 322.
- Tyramine**, *dischloro*-, hydrochloride, preparation of (ZEYNEK), A., ii, 254.
- Tyrosine**, deaminating of, in the organism (KOTAKE, MATSUOKA, and OKAGAWA), A., i, 1218.
- detection of, in urine (SCHUMM and PAPENDIECK), A., i, 899.
- estimation of (HANKE and KOESSLER), A., ii, 322.
- estimation of, in proteins (v. FURTH and FLEISCHMANN), A., ii, 406; (FOLIN and LOONEY), A., ii, 539.
- Tyrosine**, 3:5-*di*-bromo-, and -chloro-, preparation of (ZEYNEK), A., i, 254.
- d*-**Tyrosine anhydride** (FRANKEL and FELDSEBERG), A., i, 184.
- Tyrosine-choline**, and its salts (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.

U.

- Ultra-filtration** (McBAIN and JENKINS), T., 2325.
- Ulva**, spectrum of chlorophyll from (ROMIEU and OBATON), A., i, 793.
- Umbelliferæ**, hesperidine in (NILSSON), A., i, 211.
- Undecenoic acid**, arylamides of (OTT and ZIMMERMANN), A., i, 137.
- Undecenol**, and its hydrogen sulphate (GRUN and WIRTH), A., i, 805.
- Unsaturated compounds**, formation of, from halogenated open-chain derivatives (CHANDRASENA and INGOLD), T., 1306; (INGOLD), T., 2626.
- homocyclic, chemistry of polycyclic compounds in relation to isomeric (FARMER, INGOLD, and THORPE), T., 128; (CHANDRASENA, INGOLD, and THORPE), T., 1542.
- determination of the iodine value of (HOLDE, WERNER, TACKE, and WILKE), A., ii, 533; (HOLDF), A., ii, 665.
- Uracil**, action of diazomethane on (JOHNSON, HILL, and CASE), A., i, 471.
- Uranium**, Rontgen ray spectrum of (DE BROGLIE), A., ii, 330.
- N*-series spectrum of (DOLEJŠEK), A., ii, 463.
- disintegration of, and its isotopes (NEUBURGER), A., ii, 185.
- Uranium alloys** with iron (POLUSHKIN), A., ii, 152.
- Uranium salts**, action of light on (ALOY and RODIER; BAUR and REBMANN), A., ii, 337; (BAUR and HAGGENMACHER), A., ii, 338.
- Uranium salts**, triboluminescence of (SIEMSEN), A., ii, 511.
- Uranium oxides** (JOLIBOIS and BOSUET), A., ii, 301; (LEBEAU), A., ii, 302.
- radioactivity of (STAHLING), A., ii, 106.
- tri*- and *tetr*-oxides, hydrates of (HUTTIG and v. SCHROEDER), A., ii, 510.
- Uranyl nitrate**, hydrate of (GERMANN), A., ii, 649.
- radioactivity of the precipitate from sodium hydroxide and (JOLIBOIS and BOSSUET), A., ii, 575.
- hypophosphites (ROSENHEIM and TREWENDT), A., ii, 650.
- Uranium organic compounds** :—
- Uranylacetic acid**, sodium salt, refractive index of (RAITERI), A., ii, 541.
- Uranium detection and estimation** :—
- detection of (BUELL), A., ii, 590.
- detection of, spectroscopically (MEYER and GREULICH), A., ii, 6.
- estimation of (KIKUCHI; KANO), A., ii, 721.
- estimation of, electrometrically (EWING and ELDRIDGE), A., ii, 661.
- estimation of, by reduction (TREADWELL and BLUMENTHAL), A., ii, 788.
- estimation of, in carnotite (SCOTT), A., ii, 788.
- estimation of, in presence of phosphoric acid (SCHOPF and STEINKUHLER), A., ii, 530.
- Uranium-V** (PICARD and STAHEL), A., ii, 185; (HAHN), A., ii, 340.
- Uranium-X**, adsorption of, by basic ferric acetate (BROWN), T., 1736.
- Uranium-lead**, band spectra of, and of lead (GREBE and KONEN), A., ii, 4.
- Urazole**, thio-, and its derivatives (ARNDT, MILDE, and TSCHENSCHER), A., i, 375.
- dithio*-, constitution of, and its derivatives (GUHA), A., i, 875.
- dithio*-, and iminothio-, preparation of (FROMM), A., i, 62.
- Urea (carbamide)**, influence of amino-acids on formation of (FOLIN and BERGLUND), A., i, 702.
- replacement of proteins by, in diet (MOIGEN, SCHÖLER, WINDHEUSER, and OHLMER), A., i, 293.
- excretion of, according to Ambard's laws (LUBLIN), A., i, 400.
- as a plant nutrient (BOKORNY), A., i, 1096.
- in fungi (GORIS and COSTY), A., i, 1220.

- Urea** (*carbamide*), distribution of, in blood (ETIENNE and VÉRAIN), A., i, 963.
- detection of (BONNET and HAUSHALTER; BARRENSCHEEN and WELTMANN), A., ii, 794; (PINCUSSEN), A., ii, 884.
- detection of, in tissues, by means of xanthidol (STUBEL), A., i, 387.
- estimation of (ASZÓDI), A., ii, 536; (CARRA), A., ii, 668.
- estimation of, gasometrically (MEZGER), A., ii, 170.
- estimation of, by the hypobromite method (MÉNAUL; STEHLÉ), A., ii, 403.
- estimation of, by the xanthidol method, effect of sodium fluoride on (POLONOVSKI and AUGUSTE), A., ii, 668.
- estimation of, microchemically, in blood (NICLOUX and WELTER), A., ii, 170.
- estimation of, in blood, tissues, etc. (GAD-ANDRESEN), A., ii, 536.
- estimation of, in cow's milk (MORIMOTO), A., i, 703.
- estimation of, in wine (JANET), A., ii, 794.
- See also Carbamide.
- Urease** (LÖVGREN), A., i, 185.
- inactivation of, by metals (JACOBY), A., i, 480; (JACOBY and SHIMIZU), A., i, 481.
- in soja beans (WESTER), A., i, 311.
- distribution of, in plants (KIESEL and TROITZKI), A., i, 410.
- in fungi (GORIS and COSTY), A., i, 1220.
- effect of chemicals on the ureolitic power of (WESTER), A., i, 391.
- Ureometers** (CLOGNE), A., ii, 237; (SCIORTINO), A., ii, 794.
- Urethane**, equilibrium in the system, *m*-dinitrobenzene and (PUSHIN and FIOLETOVA), T., 2822.
- Uric acid**, synthesis of, in the organism (KOLLMANN), A., i, 293.
- dissociation of, and its salts (KANITZ), A., i, 277.
- influence of hydrogen-ion concentration on the solubility of (JUNG), A., i, 1070.
- colloidal and supersaturated solutions of (SCHADE), A., i, 1192.
- in human blood (MORRIS and MACLEOD), A., i, 392.
- distribution of, in blood (THEIS and BENEDICT), A., i, 82; (ROUZAUD and THIÉRY), A., i, 394; (CHAUFARD, BRODIN, and GRIGAUT), A., i, 1086.
- Uric acid**, estimation of (MORRIS and MACLEOD), A., ii, 328; (JACKSON and PALMER), A., ii, 328, 795.
- estimation of, in blood (GUILLAUMIN), A., ii, 170, 796; (GRIGAUT; BENEDICT), A., ii, 405; (PUCHER), A., ii, 668; (BAUMAN and KEELER), A., ii, 796.
- estimation of, in tissue extracts (FREUDEL and SUZUKI), A., ii, 538.
- estimation of, in urine (THIÉRY), A., ii, 238; (VAILLANT), A., ii, 668; (BENEDICT and FRANKKE), A., ii, 669; (KHOURI), A., ii, 885.
- Urine**, effect of radium emanation on (HAUENSTEIN), A., i, 80.
- viscosity of (JOEL), A., i, 198.
- composition of, under various conditions (CAMPBELL and WEBSTER), A., i, 197, 495.
- effect of loss of carbon dioxide on the hydrogen-ion concentration of (MARSHALL), A., i, 494.
- acidity of (BLATHERWICK and LONG), A., i, 967.
- relation between the reaction of, and the alveolar tension of carbon dioxide (ENDRES), A., i, 1214.
- acetaldehyde in (STEPP and FEULGEN), A., i, 495.
- organic acids in (GOIFFON and NEPVEUX), A., i, 1215.
- aldehyde in, in diabetes (STEPP and FEULGEN), A., i, 300.
- alkalinity of, after meals (FISKE), A., i, 88.
- amino-acids and hippuric acid in, in pellagra (MURLIN), A., i, 965.
- amino-nitrogen in (CIACCIO), A., i, 88.
- excretion of ammonia in, following administration of acids (KEETON), A., i, 300.
- reactions of, after treatment with arsenobenzene (GAVIATI and PAVOLINI), A., i, 496.
- excretion of caffeine in (FRIEDBERG), A., i, 88; (OKUSHIMA), A., i, 703.
- carbonates and bicarbonates in (GAMBLE), A., i, 494.
- substance in, giving the diazo-reaction (HERMANN), A., i, 1091.
- action of *p*-dimethylaminobenzaldehyde on (HARR), A., i, 88.
- formaldehyde in, after administration of hexamethylenetetramine (VOIT), A., i, 1218.
- in lead poisoning, hæmatoporphyrin in (SCHUMM), A., i, 609.
- aromatic hydroxy-acids in (DE SANC-TIS and FIORI), A., i, 609.

Urine, iodine number of (WELTMANN), A., i, 88.
 excretion of iron in (KISCH), A., i, 898; (EHRENBERG and KARSTEN), A., i, 967.
 effect of diet on the distribution of nitrogen in (ROBISON), A., i, 495.
 effect of phenylacetic and phenylpropionic acids on the excretion of nitrogen in (HJIKATA), A., i, 495.
 excretion of phosphates in (FISKE), A., i, 88.
 during water diuresis and purine diuresis (BOCK and IVERSEN), A., i, 1090.
 proteic acids of (EDLBACHER), A., i, 692.
 pyruvic acid in (FRICKE), A., i, 495.
 association of radicles in (WHITE), A., i, 1214.
 action of phosphates on sugar in (ELIAS and WEISS), A., i, 1085.
 relation between the sugar in, with that in blood (TERVAERT), A., i, 1215.
 excretion of sugars in (MURACH-HAUSER), A., i, 198; (NEUWIRTH), A., i, 485.
 albuminous, proteolytic enzymes in (HEDIN), A., i, 609.
 whales', composition of (SCHMIDT-NIELSEN and HOLMSEN), A., i, 704.
Urine, analytical methods relating to:—
 analysis of (PINCUSSEN and FLOROS; PINCUSSEN and MOMFERRATOS-FLOROS), A., ii, 408.
 detection of, in stains, by indican reactions (JEMMA), A., ii, 460.
 detection of acetaldehyde in (STEFF), A., ii, 403.
 detection of acetone in (TROISE), A., ii, 595.
 detection of albumin in (RENOUX), A., ii, 797.
 detection of aldol in, in diabetes (FRICKE), A., ii, 326.
 detection of aldol and acetaldehyde in (FRICKE), A., i, 300.
 alkapton, detection of (KATSCH and NEMET), A., i, 198.
 detection of bile pigments in (SILBERSTERN), A., ii, 799.
 detection of bismuth in (AUBRY), A., ii, 165; (GANASSINI), A., ii, 590.
 detection and estimation of blood in (JOHANNESSEN), A., ii, 724.
 detection of dextrose in (RODILLON), A., ii, 166.
 detection of formaldehyde in (STEFF), A., ii, 793.
 detection of lactose in (HERZBERG), A., ii, 167.

Urine, analytical methods relating to:—
 detection of small quantities of lead in (SCHUMM), A., ii, 317.
 detection and estimation of nitrates in (NOLTE), A., ii, 583.
 detection of tyrosine in (SCHUMM and PAPENDIECK), A., i, 899.
 detection of urobilin in (RODILLON), A., ii, 888.
 estimation of acetone in (LAX), A., ii, 326; (KADING), A., ii, 793.
 estimation of acetone substances in (HUBBARD), A., ii, 234.
 estimation of albumin in (GÉRARD), A., ii, 670.
 estimation of the amino-acid nitrogen in, colorimetrically (FOLIN), A., ii, 536.
 estimation of ammonia in (MESTREZAT and JANET), A., ii, 453.
 estimation of total bases in (FISKE), A., ii, 408.
 estimation of calcium in (SHOHL and PEDLEY), A., ii, 395.
 estimation of colloids in (OTTENSTEIN), A., i, 609.
 estimation of citric acid in (McCLURE), A., ii, 791.
 use of zinc ferrocyanide in estimation of dextrose in (CARREZ), A., ii, 233.
 estimation of hippuric acid in (SNAPPER and LAQUEUR), A., ii, 92.
 estimation of homogentisic acid in (BRIGGS), A., ii, 534.
 estimation of hydrogen-ion concentration in (SILBERSTEIN), A., ii, 452.
 estimation of oxalic acid in (SALKOWSKI), A., ii, 92; (MISLOWITZER), A., ii, 325.
 estimation of purine bases in (STEUDEL and CHOU), A., ii, 239; (SALKOWSKI), A., ii, 405.
 estimation of sugar in (TERVAERT), A., ii, 166; (FOLIN and BERGLUND), A., ii, 400.
 estimation of reducing sugars in (SMITH), A., ii, 663.
 estimation of sulphur in (ROBISON), A., ii, 389.
 estimation of theobromine in (GÜNZBERG), A., i, 703.
 estimation of thiosulphuric acid in (DEZANI), A., i, 968.
 estimation of urea in (JANET), A., ii, 794.
 estimation of uric acid in (THIÉRY), A., ii, 238; (VAILLANT), A., ii, 668; (BENEDICT and FRANKE), A., ii, 669; (KHOURI), A., ii, 885.

- Urobilin**, absorption spectrum of (LEWIN and STENGER), A., ii, 414.
 detection of, in urine (RODILLON), A., ii, 888.
- Urocanic acid**, production of, from histidine in dogs (KOTAKE and KONISHI), A., i, 1217.
 production of acetoacetic acid from, in the liver (KONISHI), A., i, 1211.
- Urochrome** as a chlorophyll derivative (ROAF), A., i, 401.
- Urusene** (MAJIMA and TAKAYAMA), A., i, 263.
- Urushiol**, and its derivatives (MAJIMA), A., i, 262.
- Uterus**, anticoagulating substances from the mucous membrane of the (KING), A., i, 701.

V.

- Vacciniin**, constitution of (OHLE), A., i, 1018.
- Valency** (CAVEN; SMITH), A., ii, 279; (HUTTIG), A., ii, 849.
 determination of, by means of ammoniacal silver solutions (COLLEMBERG), A., ii, 495.
 theories of (LAPWORTH), T., 416; (KERMAK and ROBINSON), T., 427.
 electron theory of, applied to organic compounds (STIEGLITZ), A., ii, 560.
 octet theory of (PERKINS), A., ii, 133.
 in relation to Röntgen-ray spectra (WENTZEL), A., ii, 607.
 in relation to chemical reaction (PERKINS), A., ii, 438.
 neutralisation of (DUBSKY and APTEK-MANN), A., i, 104.
 types of (LANGMUIR), A., ii, 137.
 negative, and co-ordination number (SCHILOV), A., ii, 839.
 of organic compounds (HENRICH), A., ii, 704; (STIEGLITZ), A., ii, 705.
 polar and non-polar, in organic compounds (GARNER), A., ii, 758.
 secondary (SMITH), A., ii, 270, 430, 431.
 subsidiary (CLARK and BUCKNER), A., ii, 300.
- Valerianilide**, $\beta\beta\gamma$ -trichloro- α -hydroxy- (PASSEKINI), A., i, 732.
- Valeric acid**, lead diphenyl salt (GODDARD, ASHLEY, and EVANS), T., 981.
- isoValeric acid**, α -amino- β -hydroxy-, preparation and derivatives of (SCHRAUTH and GELLER), A., i, 1125.
- Valine-choline**, and its salts (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.

- Valinol**, and its hydrochloride (KARRER, GISLER, HORLACHER, LOCHER, MADER, and THOMANN), A., i, 814.
- Vanadic acid**. See under Vanadium.
- Vanadioselenites**. See under Vanadium.
- Vanadium pentoxide**, coagulation of sols of (WIEGNER, MAGASANIK, and GESSNER), A., ii, 356.
- Vanadic acid**, estimation of, by reduction by mercury (McCAY and ANDERSON), A., ii, 530.
- Vanadioselenites** (ROSENHEIM and KRAUSE), A., ii, 48.
- Vanadium detection and estimation**:—
 detection and estimation of, in steel (MISSION), A., ii, 459.
 estimation of (KANÔ), A., ii, 721.
 estimation of, after reduction with hydrogen sulphide (LUNDELL and KNOWLES), A., ii, 88.
 estimation of, in ferrovanadium (KELLEY, WILEY, BOHN, and WRIGHT), A., ii, 89.
 estimation of, colorimetrically, in steel (KROPP), A., ii, 590.
 estimation of, in steel, by precipitation with ammonium phosphomolybdate (CAIN and HOSTETTER), A., ii, 520.
- Vanillin**, preparation of (SIEVERS and GIVAUDAN & Co.), A., i, 38.
 estimation of (TSCHIRSCH), A., ii, 403.
- Vanillin glyceride** (DODGE), A., i, 748.
- Vanillylamides**, (OTT and ZIMMERMANN), A., i, 137.
- Vanillylidenerescorinol**, and its acetyl derivative (FABRE), A., i, 1148.
- Vaporisation**, process of (HARKINS and ROBERTS), A., ii, 422.
 coefficients of (VOLMER and ESTERMANN), A., ii, 193.
- Vapours**, precipitated, rhythmic deposition of (KARRER), A., ii, 496.
- Vapour density**, apparatus for measurement of (MAGNUS and SCHMID), A., ii, 260.
 at low pressures and extended range of temperature (MAASS and BOOMER), A., i, 912.
- Vapour pressure**, formula for (HENGLEIN), A., ii, 118.
 at high temperatures (INGOLD), T., 2419.
 intersecting curves of (v. RECHENBERG), A., ii, 120.
 of binary mixed liquids (SCHMIDT), A., ii, 119; (FAUST), A., ii, 423; (CASSEL; SCHULZE), A., ii, 424.
 of non-associated liquids (MORTIMER), A., ii, 615.

- Vapour pressure** of metallic salts (v. WARTENBERG and SCHULZ), A., ii, 146; (v. WARTENBERG and BOSSE), A., ii, 739.
of saturated solutions (EDGAR and SWAN), A., ii, 349.
- Vascular system**, action of organic bases on the (TESCHENDORF), A., i, 90.
- Velardeite** from California (SHANNON), A., ii, 517.
- Velocity of chemical reaction** in solids (HINSHELWOOD and BOWEN), A., ii, 628.
- Velocity of crystallisation** under pressure (HASSELBLATT), A., ii, 35.
- Velocity of decomposition** of crystals (SIEVERTS), A., ii, 700.
- Velocity of hydrolysis** of esters (SMITH and OLSSON), A., ii, 701.
- Velocity of reaction**, theory of (DHAR), A., ii, 39; (BRONSTED), A., ii, 699. Michael's rule for (SMITH), A., ii, 751.
lecture experiment to show (MEYER), A., ii, 636.
influence of catalysts on the thermodynamics of (VAN THIEL), A., ii, 754.
and equilibria (BRANDSMA), A., ii, 699.
in heterogeneous systems (COLLEBERG and BOEFORS), A., ii, 431.
of liberation of halogens (FREUNDLICH and BARTELS), A., ii, 489.
in mixed solvents (CASHMORE, MCCOMBIE, and SCARBOROUGH), T., 243; (MCCOMBIE, SCARBOROUGH, and SETTLE), T., 2308.
- Velocity of saponification** of esters (MCCOMBIE, SCARBOROUGH, and SETTLE), T., 2308.
of oils and fats (NORRIS and MCBAIN), T., 1362.
- Veratraldehyde**, 5-amino- (SPATH and RÖDER), A., i, 853.
6-amino- and its derivatives, and 6-bromo-, 6-chloro-, and 6-iodo- (RILLIET), A., i, 839.
- Veratrylideneaniline**, 6-amino- and 6-nitro- (RILLIET), A., i, 839.
- Veratrylidene-p-anisidine**, 6-amino- and 6-nitro- (RILLIET), A., i, 839.
- Veratrylidene-gallacetophenone** dimethyl ether, preparation of (CRABTREE and ROBINSON), T., 1038.
- Veratrylidene-toluidines**, 6-amino- and 6-nitro- (RILLIET), A., i, 839.
- Vernin**, detection of (STEUDEL and FREISE), A., ii, 535.
- Veronal**, effect of lecithin on the excretion of (BACHEM), A., i, 400.
detection of (FABRE), A., ii, 795.
- Veronal derivatives**, detection of (HANDORF), A., ii, 884.
- Viburnum opulus**, detection of dextrose in fruits of (ARNOLD), A., i, 311.
- Vine**, Canadian. See *Vitis hederacea*.
- Vinegar**, detection of mineral acids in (KLING and LASSIEUR), A., ii, 234; (KLING, LASSIEUR, and LASSIEUR), A., ii, 519.
- Vinyl chloride**, polymerisation of, on exposure to ultra-violet light (PLOTNIKOW), A., i, 419.
compounds, preparation and polymerisation of (TRAUN'S FORSCHUNGS-LABORATORIUM), A., i, 516, 517.
haloids, preparation of (TRAUN'S FORSCHUNGS-LABORATORIUM), A., i, 517.
- Vinyl alcohol**, formation of, from acet-aldehyde, in presence of potassium hydroxide (EVANS and LOOKER), A., i, 102.
- Vinylarsenious sulphide**, β -chloro- (MANN and POPE), T., 1756.
- Vinylarsines**, β -chloro-, and their salts (MANN and POPE), T., 1754.
- Vinylarsinic oxide**, β -chloro- (MANN and POPE), T., 1755.
- Vinylarsinic acid**, β -chloro-, and its ammonium salt (MANN and POPE), T., 1755.
- Vinyl carbinols**, poly-arylated (ZIEGLER and OCHS), A., i, 1047.
- Vinylidene glycol** diethyl ether. See Ketenacetel.
- Vinylcyclopropane**, and its dibromide (DEMJEANOV and DOJARENKO), A., i, 1014.
- Vinylsulphuric acid**, preparation of (TRAUN'S FORSCHUNGS-LABORATORIUM), A., i, 517.
- Viola odorata** (violet), oil from the roots of (GORIS and VISCHNIAC), A., i, 848.
- Violet**. See *Viola odorata*.
- Violuric acids**, imino-, chromoisomerism of (LIFSCHITZ and HEPNER), A., i, 767.
- Viscera**, extraction of alkaloids from (COLLEDGE), A., ii, 327.
- Viscosity**, calculation of (CREIGHTON), A., ii, 426.
and compound formation in binary mixtures of acids with ketones and with esters (KENDALL and BRAKELEY), A., ii, 126.
of amphoteric electrolytes (HEDESTRAND), A., ii, 821.
of gases (SMITH), A., ii, 549.
of hydrocarbons (CHAVANNE and VAN RISSEGHEM), A., ii, 352.
of liquids (HERZ), A., ii, 821.
negative (RABINOWITSCH), A., ii, 478.

Viscostalagmometer, new (TRAUBE), A., ii, 122.

Vitamins (GRALKA and ARON), A., i, 395; (FRANKEL and HAGER; FRÄNKEL and SCHARF), A., i, 409. structure of (WILLIAMS), A., i, 280. physical chemistry of (LAMER), A., i, 281.

isolation of (SEIDELL), A., i, 887.

inactivation of (ZILVA), A., i, 488.

rôle of, in the chemistry of cells (HESS), A., i, 399, 788.

relation of, to cell oxidation processes (ABDERHALDEN), A., i, 607.

in lemon juice (LEICHTENTRITT and ZIELASKOWSKI), A., i, 1094.

in rice, and their stimulation of yeast (FLEMING), A., i, 93.

in yeast (FUNK and DUBIN), A., i, 203.

antineuritic. See Vitamin-B.

fat-soluble, estimation of (ZILVA and MIURA), A., i, 194.

detection of, in foodstuffs (DRUMMOND and WATSON), A., ii, 596.

Vitamin-A, synthesis of, by marine diatoms (JAMESON, DRUMMOND, and COWARD), A., i, 1099.

Vitamin-B, preparation and properties of (TSUKIYE), A., i, 974.

isolation of (SEIDELL), A., i, 1078.

extraction of, with glacial acetic acid (LEVINE, MCCOLLUM, and SIMMONDS), A., i, 975.

presence of, in chlorophyll-free plants (ORTON, MCCOLLUM, and SIMMONDS), A., i, 974.

bacteria as a source of (DAMON), A., i, 201.

physiological action of (BLOHM, SANTESSON, and v. EULER), A., i, 1218.

estimation of (v. EULER and MYRBÄCK), A., i, 293.

effect of temperature and hydrogen-ion concentration on the destruction of (SHERMAN, LAMER, and CAMPBELL), A., i, 207; ii, 225.

estimation of (SHERMAN, LAMER, and CAMPBELL), A., ii, 407.

Vitis hedernacea (Canadian vine), grape oil from (ANON.), A., i, 97.

Volcanic gases. See Gases.

Volcanism, chemistry of (ALLEN), A., ii, 219.

Volume, atomic, relation between refractivity and (LE BAS), A., ii, 241.

atomic and molecular, relation between, at the absolute zero (HERZ), A., ii, 29.

molecular, regularities of, of inorganic compounds (HENGLEIN), A., ii, 280.

Volumes, relative, of chemical elements (COLLINS), A., ii, 28.

Volumenometer for microanalysis (BOLLAND), A., ii, 221.

Vulcanisation, accelerators of (TWISS BRAZIER, and THOMAS), A., i, 460.

W.

War memorial, unveiling of, T., 2895.

Water, catalytic formation of, from hydrogen and oxygen (PEASE and TAYLOR), A., ii, 701.

band spectrum of (WEICHMANN), A., ii, 5, 331.

specific volume of molecules of fixed (TROMP), A., ii, 476.

molecular structure of (PICCARD), A., ii, 212.

surface tension of mixtures of alcohol and (BIRCUMSHAW), T., 887.

velocity of extension of oils on the surface of (WOOG), A., ii, 197.

kataphoresis of, in organic liquids (GYEMANT), A., ii, 684.

vapour, catalysis of the reaction of carbon with (TAYLOR and NEVILLE), A., ii, 143.

chemical reactions involved in purification of (GREENFIELD and BUSWELL), A., ii, 653.

NATURAL WATER:—

Potable or drinking water, estimation of nitrates in (REUSS), A., ii, 454.

Rain-water, analyses of (SCHAFER), A., i, 512.

Sea-water, chemical constituents of (BERTRAND, FREUNDLER, and MÉNAGER), A., ii, 441.

filtered, estimation of silica in (WELLS), A., ii, 868.

Spring and mineral waters, radioactivity of (EBLER and VAN RHYN), A., ii, 16.

of Bagnoles-de-l'Orne, radioactivity of (LOISEL), A., ii, 15.

Water analysis:—

lactose-fermenting bacteria in relation to (LEVINE), A., i, 901.

estimation of, in fuel (MARINOT), A., ii, 223.

estimation of lead in (AVERY, HEMINGWAY, ANDERSON, and READ), A., ii, 161.

Water pump. See Pump.

Wattle, gidgee. See *Acacia cambagei*.

Wax, bees', Indian, constants of (ROBERTS and ISLIP), A., ii, 534.

fossil, of Monte Falo (CIUSA and VOIS), A., ii, 386.

- Wax**, montan, acids of (TROPSCH and KREUTZER), A., i, 317, 804.
 colouring constituents of (MARCUSON and SMELKUS), A., i, 807.
 paraffin. See Paraffin wax.
 from pine needles (KAUFMANN and FRIEDEBAUGH), A., i, 740.
- Weights**, brass, protection of, from rust (MANLEY), A., ii, 840.
- Weights**, molecular, determination of (KAST), A., ii, 421.
 determination of, by means of osmotic pressure (FOIX), A., ii, 621.
 determination of, in alcoholic solution by means of the flash-point elevation (WRIGHT), T., 2247.
- Weinschenkite** (HENRICH and HILLER), A., ii, 860.
- Whale**, urine of. See Urine.
- Wheat**, hydrolysis of gliadin from (VICKERY), A., ii, 754.
 estimation of starch in (LING, CALLOW, and PRICE), A., ii, 879.
- White metal**, analysis of (KLING and LASSIEUR), A., ii, 87; (BERTIAUX), A., ii, 396.
- Willia javanica*, occurrence of emulsin and of cellobiase in (GROENEWEGE), A., i, 903.
- Wines**, odour and chemical composition of (MALVEZIN), A., i, 1220.
 effect of radioactive substances on the acetic fermentation of (LABORDE, JALOUSTRÉ, and LEULIER), A., i, 1219.
 detection of formic acid in (FRESENIUS and GRUNHUT), A., ii, 234.
 estimation of the acids in (v. FELENBERG), A., ii, 534; (MALVEZIN), A., ii, 880.
 estimation of tannin in (MALVEZIN), A., ii, 172.
 estimation of tannin and colouring matters in (FRESENIUS and GRUNHUT), A., ii, 96.
- Wood**, detection of, by a colour reaction (ADLER), A., ii, 401.
 estimation of cellulose in (HEUSER and CASSEUR), A., ii, 664.
- Wool**, adsorption of neutral soap by (BRITISH RESEARCH ASSOCIATION FOR THE WOOLLEN AND WORSTED INDUSTRIES), A., ii, 551.
 action of iodine on (HUEBNER and SINHA), A., i, 434.
 mordanting of, with potash alum (PADDON), A., ii, 822.
 deaminated, dyeing of (PADDON), A., i, 608.
 estimation of soap in (BRITISH RESEARCH ASSOCIATION FOR THE WOOLLEN AND WORSTED INDUSTRIES), A., ii, 594.
- Worms**, parasitic, hæmatotoxins from (SCHWARTZ), A., i, 493.
- Wurts-Fittig synthesis** (FUCHS and METZL), A., i, 442; (SCHLUBACH and GOES), A., i, 1204.
- X**
- X-rays**. See Rays, Röntgen.
- Xanthic acid**, cobalt, nickel, potassium and sodium alkyl and aryl salts, and their compounds with pyridine (DUBSKY and APTEKMAN), A., i, 104.
 diphenylbisazo-2-hydroxy-7-naphthyl ester (WATSON and DUTT, T., 2417).
- Xanthine**, oxidation of, by tissues and by milk (MORGAN, STEWART, and HOPKINS; ARMSTRONG), A., i, 1078.
- Xanthone**, condensed derivatives of (ECKERT and ENDLER), A., i, 941.
- Xanthone**, 1:4-dichloro-, and 4-chloro-1-hydroxy- (ECKERT and ENDLER), A., i, 941.
 1:3-dihydroxy- (NISHIKAWA and ROBINSON), T., 841.
- Xanthone series** (v. DEM KNESEBECK and ULLMANN), A., i, 359.
- Xanthone-4-carboxylic acid**, and its salts and derivatives (ANSCHÜTZ and CLAASEN), A., i, 456.
- Xanthorocellin** (FORSTER and SAVILLE), T., 821.
- Xanthosterol** (DIETERLE), A., i, 652.
 identity of lupeol with (ULTÉE), A., i, 826.
- Xanthydrol**, use of, in detection of urea in tissues (STÜBEL), A., i, 397.
- 3-Xanthylbenzoic acid**, 2:4-dihydroxy- (FABRE), A., i, 1148.
- 4-Xanthyl-1:2-diethylresorcinol** 3-ethyl ether (FABRE), A., i, 1148.
- 3-Xanthylpyrocatechol**, and 4-chloro- (FABRE), A., i, 1148.
- 2-Xanthylquinol**, and 3-chloro- (FABRE), A., i, 1148.
- 2-Xanthylresorcinol**, 4-chloro-, and 4-nitroso- (FABRE), A., i, 1148.
- 4-Xanthylresorcinol**, and its diacetyl derivative, and 2-nitroso- (FABRE), A., i, 1148.
- Xenon**, isotopes of (ASTON), A., ii, 844.
 estimation of (MOUREU and LÉPAGE), A., ii, 394.
- Xylan** (SALKOWSKI), A., i, 323.
 preparation and hydrolysis of (HEUSER, BRADEN, and KÜRSCHNER), A., i, 113.
 and its acetyl derivatives (KOMATSU and KASHIMA), A., i, 811.
 methyl ethers of (HEUSER and RUPPEL), A., i, 810.

(*o*-Xylene, *Me:Me* = 1:2; *m*-xylene, *Me:Me* = 1:3; *p*-xylene, *Me:Me* = 1:4.)

Xylene, equilibrium of ethyl alcohol, water and (ORMANDY and CRAVEN), A., i, 215.
 estimation of (ZABOROWSKI), A., ii, 877.
m-Xylene, vapour pressure of liquid mixtures of ammonia and (KRAUS and ZEITFUCHS), A., i, 725.
 action of carbonyl chloride on, in presence of aluminium chloride (WILSON and FULLER), A., i, 827.
p-Xylene, action of sulphuryl azide on (CURTIUS and SCHMIDT), A., i, 776.
Xylenes, trinitro-, oxidation of (GIUA), A., i, 534.
Xylene cyanole FF as an indicator (HICKMAN and LINSTED), T., 2502.
p-Xylenedi-*p*-thiodimethylaniline (SMILES and GRAHAM), T., 2510.
Xylenol, tribromo-, action of, on tubercle bacilli (DUBOC), A., i, 972.
Xylenol-blue, use of, as an indicator (COHEN), A., ii, 387.
p-Xylidine hydroferrocyanide (CUMMING), T., 1294.
 ψ -Xylidine, and its salts (CURTIUS and SCHMIDT), A., i, 776.
 β -*m*-Xylidino-*p*-phenetolecarbamide (SPECKAN), A., i, 580.
Xylose, action of lactic bacteria on (FRID, PETERSON, and ANDERSON), A., i, 201.
 methylation of (CARRUTHERS and HIRSH), T., 2299.
p-Xylylene-2:5-diacetic acid, 3:6-di-bromo- (PHILIPPI, SEKA, and ROBINSON), A., i, 837.
p-Xylylene-2:5-diglyoxylic acid, 3:6-di-bromo- (PHILIPPI, SEKA, and ROBINSON), A., i, 837.
 4-*o*-Xylyl ethyl ketone, and its semicarbazone (v. AUWERS and ZIEGLER), A., i, 120.
 α -*m*-4-Xylylpropionic acid, and $\beta\beta$ -dichloro-, ethyl ester (v. AUWERS and ZIEGLER), A., i, 140.
m-Xylylpyruvic acid, *d*-nitro- (DAVIES and HICKOX), T., 2646.

Y.

Yeast, longevity of (LING and NANJ), A., i, 92.
 changes in nitrogenous substances during autolysis by (IVANOV), A., i, 202.
 action of, on acid amides (DIETER), A., i, 795.
 action of, on aldehydes (KUMAGAWA), A., i, 305.

Yeast, influence of mineral water on the carbohydrate exchange in (MAYER), A., i, 972.
 toxicity of chlorine derivatives of methane, ethane and ethylene towards (PLAGGE), A., i, 93.
 extracts, effect of, on the action of enzymes (ABDERHALDEN and WERTHEIMER), A., i, 796.
 fat of (MACLEAN), A., i, 304, 795.
 fermentation by (ABDERHALDEN; ABDERHALDEN and FODOR), A., i, 92; (WILLSTATTER and STEIBELT), A., i, 306.
 effect of acids on (SOMOGYI), A., i, 201.
 stimulants for (NEUBERG and SANDBERG), A., i, 408.
 in presence of urea (SANDBERG), A., i, 502.
 gum in (SALKOWSKI), A., i, 304.
 invertase activity of (MILLER), A., i, 202.
 lactase content of (WILLSTATTER and OPPENHEIMER), A., i, 203.
 destruction of lactic acid by cells of (LIEBEN), A., i, 796; (v. FÜRTH and LIEBEN), A., i, 1219.
 bleaching of methylene-blue by (KUMAGAWA), A., i, 306.
 nitrogenous constituents of (MEISENHEIMER), A., i, 304; (SCHENCK), A., i, 305.
 nitrogenous nutrition of (SWOBODA), A., i, 795.
 hydrolysis of the protein of (KIESEL), A., i, 408.
 decomposition of proteins in, during fermentation (IVANOV), A., i, 202.
 effect of quinine, phenol, and mercuric chloride on the growth of (JOACHIMOGLU), A., i, 903.
 action of silver salts on (ZERNER and HAMBURGER), A., i, 307.
 urea as a nutrient for (BOKORNY), A., i, 1096.
 vitamins in (FUNK and DUBIN), A., i, 203.
 stimulation of, by vitamins (FLEMING), A., i, 93.
 rate of growth of, in beer wort (CLARK), A., i, 501.
 zymase formation in (HAYDUCK and HAEHN), A., i, 611.
 Japanese, fermentation by species of (KUMAGAWA), A., i, 972.
 oxygenated, action of amino-acids on (LIEBEN), A., i, 1219.
R, fermentation by (v. EULER and JOSEPHSON), A., i, 706.
 See also *Wulha javanica*.

Yeast-cells, carbohydrate metabolism of (ELIAS and WEISS), A., i, 485.
 decomposition of lactic acid by (v. FÜRTH and LIEBEN), A., i, 502.
 action of saponins on (BOAS), A., i, 502.
Yeast-nucleic acid (STEUDEL and PEISER), A., i, 279, 782.
 action of human feces on (ROTHER), A., i, 292.
 hydrolysis of, by boiled pancreatic extracts (JONES), A., i, 479.
Yew. See *Taxus baccata*.
Yohimba bark, estimation of yohimbine in (SCHOMER), A., ii, 797.
Yohimbine, estimation of, in yohimba bark (SCHOMER), A., ii, 797.
Ytterbium, *L*-series spectrum of (DAUVILLIER), A., ii, 463.
Yttrium, atomic weight of (FOGG and JAMES), A., ii, 297.
 arc spectra of (KIESS, HOPKINS, and KREMER), A., ii, 244; (YNTEMA and HOPKINS), A., ii, 462.

Z.

Zeeman effect (WOLTJER), A., ii, 102.
Zinc, heat of solution of, in hydrochloric acid (RICHARDS and THORVALDSEN), A., ii, 475.
 liquid, density and surface tension of (HOGNESS), A., ii, 29.
 rate of solution of, in ferric alum (COLLENBERG and BODFORS), A., ii, 431.
Zinc alloys with arsenic (HEIKE), A., ii, 60.
 with copper and nickel (VOIGT), A., ii, 295.
 with magnesium, heat of formation of (BILTZ and HÖRST), A., ii, 350.
Zinc compounds, content of, in animal organs (BERFRAND and VLADESCO), A., i, 493.
 in the animal and human organism (ROST), A., i, 87.
 in carcinoma (CRISTOL), A., i, 497.
Zinc salts, feeding of animals with (BERTRAND and BENZON), A., i, 893.
Zinc arsenide as detector for wireless telegraphy (HEIKE), A., ii, 19.
 borate (BROEKSMIT), A., ii, 377.

Zinc bromate hexahydrate, crystal structure of (WYCKOFF), A., ii, 710.
antimoniodibromide (VOURNAZOS), A., ii, 651.
 sulphate, electrical conductivity of solutions (TARTAR and KEYES), A., ii, 186.
 sulphide, phosphorescent (GUNTZ), A., ii, 502.
 containing copper (GUDDEN and POHL), A., ii, 680.
 sulphides, phosphorescent, action of ultra-violet light on (LOEB and SCHMIDESKAMP), A., ii, 8.
Zinc organic compounds, action of, on acid chlorides (MAUTHNER), A., i, 457.
 with olefines and nitrogen oxides (SCHAARSCHMIDT, VEIDT, and SCHLOSNER), A., i, 645.
Zinc estimation and separation:—
 estimation of (URBASCH), A., ii, 317.
 estimation of, by electrometric titration (KOLTHOFF), A., ii, 580.
 estimation of, gasometrically (BEYNE), A., ii, 85.
 estimation of, as pyrophosphate (BALAREFF), A., ii, 228.
 estimation of, as sulphate (GUTHRIE and STAIB), A., ii, 396.
 estimation of, volumetrically (MONASCH), A., ii, 160.
 estimation of, volumetrically, with ferrocyanides (TREADWELL and CHERVET), A., ii, 786.
 estimation of, volumetrically, with sodium sulphide (OLIVIER), A., ii, 396.
 separation of, by means of ammonium phosphate (LUFF), A., ii, 456.
 separation of, from nickel (LUDWIG), A., ii, 786.
Zinc blende. See Blende.
Zirconium (MARDEN and RICH), A., ii, 153.
Zirconium organic compounds:—
 ferri- and ferro-cyanides (VENABLE and MOEHLMANN), A., ii, 712.
Zymase, formation of, in yeast (HAYDUCK and HAEHN), A., i, 611.
 of yeast, action of (ABDERHALDEN and FODOR), A., i, 92.
Zymogens, adsorption of (JACOBY and SHIMIZU), A., i, 481.
 artificial (JACOBY), A., i, 480.
 (JACOBY and SHIMIZU), A., i, 481.

INDEX TO PATENTS.

Austrian Patent.

83136, A., i, 954

British Patents.

144681, A., i, 1009

145060, A., i, 992

147001, A., i, 178

147118, A., i, 112

147119, A., i, 111

148419, A., i, 340

153605, A., i, 262

153917, A., i, 567

154579, A., i, 430

154907, A., i, 420

155577, A., i, 481

156103, A., i, 425

156117, A., i, 516

156121, A., i, 517

156116, A., i, 514

156122, A., i, 515

156136, A., i, 528

156148, A., i, 522

156215, A., i, 559

156538, A., i, 559

158558, A., i, 582

160433, A., i, 178

161539, A., i, 639

164302, A., i, 560

164326, A., i, 236

164715, A., i, 257

165446, A., i, 585

165770, A., i, 132

165771, A., i, 132

167157, A., i, 926

167781, A., i, 529

171292, A., i, 942

172682, A., i, 259

173004, A., i, 222

173006, A., i, 260

173063, A., i, 254

173097, A., i, 218

173166, A., i, 260

173540, A., i, 252

174101, A., i, 354

175019, A., i, 448

176038, A., i, 468

176925, A., i, 560

177189, A., i, 517

177807, A., i, 1172

179753, A., i, 650

180016, A., i, 625

181247, A., i, 872

182986, A., i, 853

183044, A., i, 836

183351, A., i, 941

183897, A., i, 914

184625, A., i, 1058

185612, A., i, 1068

185913, A., i, 1179

186859, A., i, 1179

Canadian Patent.

217486, A., i, 828

Dutch Patents.

6352, A., i, 962

6581, A., i, 961

French Patents.

22347, A., i, 385

521119, A., i, 75

521281, A., i, 579

521469, A., i, 75

523108, A., i, 810

525539, A., i, 531

526129, A., i, 801

526686, A., i, 942

German Patents (D.R.-P.).

301278, A., i, 993

301279, A., i, 992

338735, A., i, 631

338737, A., i, 565

338738, A., i, 46

338853, A., i, 46

338926, A., i, 61

339101, A., i, 579

339350, A., i, 553

339494, A., i, 605

339561, A., i, 554

339562, A., i, 514

339945, A., i, 556

339947, A., i, 46

340744, A., i, 464

340872, A., i, 815

340873, A., i, 365

340874, A., i, 365

341112, A., i, 762

341837, A., i, 567

342048, A., i, 574

342733, A., i, 824

343054, A., i, 861

343055, A., i, 854

343056, A., i, 819

343057, A., i, 823

343147, A., i, 819

343148, A., i, 860

343149, A., i, 867

343249, A., i, 822

343321, A., i, 980

343322, A., i, 867

343930, A., i, 934

344027, A., i, 952

344028, A., i, 949

344029, A., i, 950

344030, A., i, 950

344140, A., i, 948

344501, A., i, 952

344878, A., i, 913

345145, A., i, 918

345361, A., i, 1066

346065, A., i, 818

346188, A., i, 942

346236, A., i, 915

346325, A., i, 706

346383, A., i, 991

346461, A., i, 950

346462, A., i, 991

346673, A., i, 1039

346810, A., i, 999

346888, A., i, 949

347377, A., i, 991

347604, A., i, 980

347609, A., i, 992

348379, A., i, 950

348484, A., i, 943

348906, A., i, 1066

349794, A., i, 1145

349915, A., i, 1115

350376, A., i, 1066

350429, A., i, 977

352129, A., i, 1160

352719, A., i, 1137

352720, A., i, 1036

352721, A., i, 1136

352980, A., i, 1071

353195, A., i, 1131

353932, A., i, 1172

354696, A., i, 1173

354698, A., i, 1161

354950, A., i, 1173

355926, A., i, 1128

Japanese Patents.

37211, A., i, 219
37212, A., i, 349
38647, A., i, 543
38752, A., i, 523
39210, A., i, 716

Swiss Patents.

73683, A., i, 942
88188, A., i, 3
88561, A., i, 31
89053, A., i, 38
89055, A., i, 818
89241, A., i, 276
90480, A., i, 942
90587, A., i, 828

90590, A., i, 827
90807, A., i, 860
91728, A., i, 953
92296, A., i, 992

United States Patents.

1874695, A., i, 27
1874720, A., i, 984
1874721, A., i, 1039
1874722, A., i, 27
1875949, A., i, 53
1878343, A., i, 51
1878939, A., i, 31
1884615, A., i, 245
1884637, A., i, 179
1888573, A., i, 654

1390972, A., i, 391
1393191, A., i, 315
1398597, A., i, 370
1394851, A., i, 355
1399082, A., i, 666
1399144, A., i, 652
1401631, A., i, 648
1403117, A., i, 679
1406547, A., i, 748
1408974, A., i, 888
1417875, A., i, 943
1418900, A., i, 1022
1419091, A., i, 938
1419092, A., i, 938
1424236, A., i, 1128

ERRATA.

VOL. 104 (ABSTR., 1913).

Page Line

- i. 681 11 for "4-β-iminoethylglyoxaline" read "4-β-aminoethylglyoxaline."

VOL. 106 (ABSTR., 1914).

- ii. 135 10* for "Thorium" read "Thallium."
ii. 874 8* col. ii. for "thorium" read "thallium."

VOL. 108 (ABSTR., 1915).

- i. 719 8* delete "VERNIER."
ii. 891 17 col. ii. " Vernier "
ii. 965 29 col. ii. " Vernier see Léon Givaudin."
ii. 1103 { 18 col. i. } " VERNIER."
26

VOL. 120 (ABSTR., 1921).

- i. 165 25 for "ZEIGLER" read "ZIEGLER."
i. 796 16* " ethyl r-pinate" read "ethyl r-pinonate."

VOL. 122 (ABSTR., 1922).

- i. 224 24 for "Digitoxone" read "Digitoxose."
i. 319 1* " furfuran" read "furan."
i. 371 18 " Bis-1:3'-indyl" read "Bis-1:3'-indil."
22 " 3:3'-indyl" read "3:3'-indil."
27 " 1:1'-indyl" read "1:1'-indil."
31 " 3:3'-Methylketyl (3:3'-dimethylketyl)" read "2:2'-Dimethyl-3:3'-indil (2:2'-dimethyl-3:3'-diindyl)."
i. 554 17 " SCHLICHTUNG" read "SCHLICHTING."
i. 563 16 " TELECKY" read "TELECZKY."
i. 758 25 " bis-3-carbethoxy-2:4-dimethylpyrrolmethene" read "ethyl m-thenylbis-2:4-dimethylpyrrole-3-carboxylate."
" CO₂Et·C:Me" read " CO₂Et·C:Me" $\begin{matrix} \text{MeC}=\text{N} \end{matrix} \begin{matrix} \text{C}=\text{etc.} \end{matrix}$ read $\begin{matrix} \text{MeC}=\text{N} \end{matrix} \begin{matrix} \text{C}=\text{etc.} \end{matrix}$
i. 758 17* " $\begin{matrix} \text{MeC}=\text{N} \end{matrix} \begin{matrix} \text{C}=\text{etc.} \end{matrix}$ read $\begin{matrix} \text{MeC}=\text{N} \end{matrix} \begin{matrix} \text{C}=\text{etc.} \end{matrix}$ "
i. 848 18* insert "[cf. Stevens, A., 1921, i, 735]."
i. 945 13 for "Cynpopogan" read "Cynbopogan."
ii. 77 8 " PbO₅UO₃·4H₂O" read "2PbO₅UO₃·4H₂O."
ii. 415 20 " 122" read "121."
ii. 519 18* " KANAKER" read "KARRAKER."
ii. 582 bottom " 30% of hydrogen peroxide" read "30% hydrogen peroxide."
ii. 685 3 " christobalite" read "cristobalite."
ii. 710 16* " lead tetraethyl" read "magnesium ethyl iodide."
ii. 714 5 " naurite" read "navrute."
ii. 714 12 " variszite" read "variscite."
ii. 768 2* " HALLE" read "HALLA."
ii. 830 13* " HAMMERSTEN" read "HAMMARSTEN."

* From bottom.

UNGERER, E., ii, 96.

Uyeda, Y. See A. G. Perkin.

VALLOT, J., i, 87.

Venable, F. P., and D. H. Jackson,
ii, 73.

Vila, A. See M. Piettre.

Vilbrandt, F. C. See S. L. Shene-
field.

William Ring See S. L. Shenefield

Willmann, K., ii, 76.

Windaus, A., W. Dörries, and H.
Jensen, i, 60.

Withrow, J. R. See S. L. Shenefield.

Witzemann, E. J., i, 6.

Woringer, P., i, 90.

Wormall, A. See P. K. Dutt.

Wormser, M. See R. Lorenz.

Wright, W. C. See G. L. Kelley.

June 21st, 1917 Nov. 7th, 1921

VOL. 120 (ABSTR., 1921).

Page	Line
i, 388	2 for " <i>Soil. Sci.</i> , 172," read " <i>Soil Sci.</i> , 11."
ii, 573	3 ,, " <i>BARBOROVSKÝ</i> " read " <i>BABOROVSKÝ</i> ."
	4 ,, " <i>HANÁK</i> " read " <i>HANAKOVÁ</i> ."
ii, 621	21* ,, " <i>103</i> " read " <i>100</i> ."

* From bottom.

PROCEEDINGS

OF THE

CHEMICAL SOCIETY.

Ordinary Scientific Meeting, Thursday, January 19th, 1922, at 8 p.m., SIR JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

The PRESIDENT referred to the loss sustained by the Society, through death, of the following Fellows :

	Elected.	Died.
Giacomo Ciamician (Honorary Fellow) ...	March 2nd, 1911	
Ganesh Sakharan Agashe	June 15th, 1916	Dec. 2nd, 1921
James Albert Allison	Dec. 7th, 1905	Dec. 9th, 1921
Charles Edward Cassal	Feb. 6th, 1899	Dec. 22nd, 1921
William King	June 21st, 1917	Nov. 7th, 1921

The PRESIDENT announced :

(1) That in accordance with the Bye-laws the following vacant places fall due to be filled at the Annual General Meeting on March 30th, 1922 :

Vice-Presidents who have filled the office of

President Two vacancies.

Vice-Presidents who have not filled the office

of President Two vacancies.

Secretary One vacancy.

Ordinary Members of Council :

(a) Town Members (that is, those living within a radius of 50 miles from London) Three vacancies.

(b) Country Members (that is, those living beyond a radius of 50 miles of London) Three vacancies.

The Council has nominated Dr. Irvine Masson to the office of Secretary.

The names of the Members of Council who retire at the Annual General Meeting on March 30th, 1922, and who are ineligible for re-election to the same office until the lapse of one year, are as follows :

Vice-Presidents who have filled the office of President :

Sir James J. Dobbie and Sir William J. Pope.

Vice-Presidents who have not filled the office of President :

Professor J. B. Cohen and Dr. H. J. H. Fenton.

Ordinary Members of Council :

(a) Town Members :

Mr. J. A. Gardner, Dr. C. A. Keane, and Sir Robert Robertson.

(b) Country Members :

Dr. E. F. Armstrong, Professor F. E. Francis, and Principal J. C. Irvine.

(2) That arrangements have been made with the Society of Chemical Industry whereby provincial Local Sections of that Society may give opportunity to Fellows of the Chemical Society to read papers on pure chemistry and to attend the meetings at which such papers are read. Further information in regard to the scheme will be circulated at an early date.

(3) That a pamphlet entitled "Notes on the Furnishing and Equipment of Chemical Laboratories" has just been issued by the Society. Arising from a Conference held in July, 1920, to consider whether economies could be effected in the equipment of chemical laboratories, a small Committee was appointed by the Council to look into and report upon the matter. These "Notes" represent the results of the inquiry and can be obtained free of charge on application to the Assistant Secretary.

Certificates were read for the first time in favour of :

Jnanendra Kumar Adhya, M.Sc., 30, Judge's Court Road, Alipore, Calcutta.

Gilbert James Avey, Northern Polytechnic Institute, Holloway, N. 7.

Bhagwan Das Batta, B.Sc., F. C. College, Lahore, India.

Darrell Webb Beesley, A.I.C., 29, King Edward Road, Dentons Green, St. Helens.

William Alfred Bishop, B.A., 33, Banstead Road, Purley.

Leslie James Burrage, B.Sc., 32, Emerson Road, Ilford.

Gabriel Cofman-Nicoresiti, 15, Ludlow Road, Ealing, W. 5.

Archibald Neil Currie, M.A., B.Sc., A.I.C., 44, Fisher Street, Dennistoun, Glasgow.

James Owen Chesshyre Dalton, B.A., The Plot, Adams Road, Cambridge.

Arthur William Edwards, 4, Dunvegan Road, Eltham, S.E. 9.

Samuel Ellingworth, M.Sc., 74, Millfield Road, York.

Kenneth Fisher, M.A., Ph.D., Eton College, Windsor.

Miri Lal Goel, M.Sc., Royal Artillery Bazar, Ambala, India.

George Grimshaw Hammond, B.Sc., 254, Mill Street, Toxteth, Liverpool.

William Allen Hamor, M.A., Mellon Institute of Industrial Research, The University, Pittsburg, Pa., U.S.A.

Frank Mouat Irvine, B.Sc., A.I.C., 30, Clayton Road, Jesmond, Newcastle-on-Tyne.

John Albert Jenkinson, B.Sc., School House, Little Sutton, Birkenhead.
 Rowland Nicholas Johnson, B.Sc., Greenstede, West Hill, East Grinstead.
 David John Lake, B.Sc., 36, Gerald Street, Hafod, Swansea.
 Herbert Langwell, Wentworth, Denham Road, Epsom.
 Vernon Howes Legg, B.Sc., 95, Carter Knowle Road, Sheffield.
 John Alexander McRae, M.A., Ph.D., F.I.C., Queen's University, Kingston,
 Ont., Canada.
 Walter Thomas James Morgan, 171, Kensington Avenue, Manor Park, E. 12.
 John Sydney Morrison, 23, Seymour Road, Cheadle Hulme, Manchester.
 Partridge Leslie Podd, B.Sc., 8, St. Peter's Street, Ipswich.
 Krishnarajpet Venkata Rau, B.A., B.Sc., Ananda College, Colombo, Ceylon.
 Frank Crafer Ray, B.A., 47, Alfriston Road, Clapham Common, S.W. 11.
 Jessie Stewart, B.Sc., School House, Helmington Row, Crook, Co. Durham.
 George Taylor, Frigorifico, Campana F.C.C.A., Argentine.
 James Rawson Walmsley, A.I.C., Dean Brook Cottage, Dean Lane, Moston,
 Manchester.
 Walter Arthur White, 144, Shelley Avenue, Manor Park, E. 12.
 Robert Norman Wright, B.Sc., A.I.C., 16, Redcliffe Street, S. Kensington,
 S.W. 10.

Prof. A. SMITHELLS then gave an account of the Lewis-Langmuir theory of atomic structure, the exposition being illustrated with a large number of models. On the motion of Prof. Sir W. H. Bragg, supported by Prof. Lowry and the President, a cordial vote of thanks to the lecturer was passed with acclamation, and Prof. Smithells made suitable acknowledgment.

ORDINARY SCIENTIFIC MEETING.

The next Ordinary Scientific Meeting will be held on Thursday, February 2nd, 1922, at 8 p.m.

LECTURE.

At the Ordinary Scientific Meeting to be held on Thursday, February 9th, 1922, at 8 p.m., in the Lecture Hall of the Institution of Mechanical Engineers, Storey's Gate, S.W.1., Sir Ernest Rutherford, M.A., D.Sc., F.R.S., will deliver his Lecture, entitled "Artificial Disintegration of Elements."

SETS OF THE *BERICHTE* AND OF THE *ZENTRALBLATT*.

The Council has secured a few sets of the *Berichte* and of the *Zentralblatt* from August, 1914, to December, 1920, inclusive. These can be obtained, *in complete sets only*, from the Assistant Secretary, price £10 a set of the *Berichte*, or of the *Zentralblatt*.

The number of sets is strictly limited, and applications will be dealt with in the order in which they are received.

ADDITIONS TO THE LIBRARY.

I. Donations.

DEUTSCHE CHEMISCHE GESELLSCHAFT. *Chemisches Zentralblatt*. Generalregister IV, über die Jahrgänge 1912–1916. Edited by IGNAZ BLOCH. Leipzig 1921. pp. vi + 2016. (*Reference.*)

From the Deutsche Chemische Gesellschaft.

FÉDÉRATION DES INDUSTRIES CHIMIQUES DE BELGIQUE. *Bulletin*. Vol. I., etc. Gand 1921 +. [Issued with the *Bull. Soc. Chim. Belg.*] (*Reference.*)

From the Société Chimique de Belgique.

FOURNEAU, ERNEST. *Préparation des médicaments organiques*. Paris 1921. pp. viii + 350. 25 fr. (*Recd.* 13/12/21.)

From the Author.

PARKES, SAMUEL. *The rudiments of chemistry; illustrated by experiments, and eight copper-plate engravings of chemical apparatus*. London 1810. pp. xii + 294. ill. (*Reference.*)

From Mr. W. Livingstone Gunn.

II. By Purchase.

FISCHER, FERDINAND. *Kraftgas: Theorie und Praxis der Vergasung fester Brennstoffe*. (*Chem. Tech. in Einzeldarstellungen.*) 2nd edition. Edited by J. GWOSDZ. Leipzig 1921. pp. viii + 428. ill. (*Recd.* 21/12/21.)

MINISTRY OF MUNITIONS and DEPARTMENT OF SCIENTIFIC AND INDUSTRIAL RESEARCH. *Technical records of explosives supply, 1915–1918*. No. IV. *The theory and practice of acid mixing*. London 1921. pp. vi + 94. ill. 12s. net. (*Recd.* 8/12/21.)

VANINO, LUDWIG. *Handbuch der präparativen Chemie*. 2nd edition. Vol. I. *Anorganischer Teil*. Stuttgart 1921. pp. xxiv + 821. ill. (*Recd.* 13/12/21.)

III. Pamphlets.

MEES, CHARLES EDWARD KENNETH, and CLARKE, HANS THATCHER. *The preparation of synthetic organic chemicals at Rochester*. [A lecture delivered before the Society of Chemical Industry, Montreal, August 31, 1921.] Rochester, N.Y. 1921. pp. 40. ill.

MOIR, JAMES. (1) *The permanganate absorption-spectrum: a claim for priority*. (2) *A formula for calculating the uranium spectrum*. (From the *Trans. Roy. Soc. S. Africa*, 1921. 10.)

SCIENTIFIC AND INDUSTRIAL RESEARCH, DEPARTMENT OF.
Bulletin No. 5. The cleaning and restoration of museum exhibits :
report upon investigations conducted at the British Museum.
London 1921. pp. 12. ill.

— First report of the Gas Cylinders Research Committee.
London 1921. pp. 126. ill.

— *Food Investigation Board*. Special Report No. 5. Report
on heat insulators by the Engineering Committee of the Food
Investigation Board. Experiments by EZER GRIFFITHS. London
1921. pp. 61. ill.

— *Fuel Research Board*. Technical Paper No. 3. The efficiency
of low temperature coke in domestic appliances. By MARGARET
WHITE FISHENDEN. London 1921. pp. 35. ill.

WRIGHT, ALLISTER MACLEAN. The anticomplementary properties
observed in certain serum reactions. (From the *Trans. New Zealand
Inst.*, 1921, 53.)

WRIGHT, ALLISTER MACLEAN, BEVIS, MISS J. F., and the late
PERCY S. NELSON. The chemistry of flesh foods. No. 5. The
nitrogenous constituents of meat-extracts. (From the *Trans. New
Zealand Inst.*, 1921, 53.)

CERTIFICATES OF CANDIDATES FOR ELECTION AT THE BALLOT TO BE HELD AT THE ORDINARY SCIENTIFIC MEETING ON THURSDAY, FEBRUARY 16TH, 1922.

N.B.—The names of those who sign from “General Knowledge”
are printed in *italics*.

ADHYA, JNANENDRA KUMAR, M.Sc. (Cal.), 30, Judge's Court Road, Alipore,
Calcutta. Chemical Laboratory Assistant, Govt. Test House, Alipore,
Calcutta. Took the M.Sc. degree of the Calcutta University in 1920. Served
as an asst. chemist in the Research Dept. of Messrs. Bird & Co., Calcutta.
At present employed as a Chemical Laboratory Assistant, Govt. Test House,
Alipore, Calcutta. Late Hon. Secretary, Presidency College Chemical
Society, Calcutta. (*Signed by*) N. Brodie, J. Bhaduri, B. B. Dey.

APPLEBEE, HARRY CHARLES, 509, Edge Lane, Droylsden, nr. Manchester.
Chief Works Chemist, Manchester Corporation Gas Works, Bradford Road,
Manc. Finsbury Technical College (Chemical Department), 1902-1905
(Certificate). Associate of the Institute of Chemistry (1918). Lecturer,
Chemistry of Gas Manufacture, Manchester College of Technology. (*Signed
by*) E. F. Morris, F. M. Rowe, Frank Lee Pyman, F. S. Sinnatt, E. Williams.

AVEY, GILBERT JAMES, Chemical Department, Northern Polytechnic
Institute, Holloway, London, N.7. Lecturer in Chemistry. Wish to receive
the Journals of the Society. (*Signed by*) T. J. Drakeley, Raymond R. Butler,
Philip Schidrowitz.

BAKER, JOHN WILLIAM, 20, Redfield Lane, Earls Court Rd., S.W.5. Research Student in Organic Chemistry at the Imperial Coll. of Science and Technology. Royal Scholarship in Chemistry, 1916. A.R.C.S. Chem., First Class, 1921. B.Sc. Hons. Chem., First Class, 1921 (London). (*Signed by*) Jocelyn Thorpe, James C. Philip, M. A. Whiteley, C. K. Ingold.

BATTA, BHAGWAN DAS, Nabha State, Panjab (India). Demonstrator in Chemistry, F. C. College, Lahore. Appearing for the Panjab University M.Sc. Degree. B.Sc., Panjab University. Demonstrator in chemistry at Forman Christian College, Lahore. Working on the "Active principle of *Melia Azideracta* (India) leaves." (*Signed by*) N. A. Yajnik, Edwin Theodore Dean, S. R. Khosla.

BEESELEY, DARRELL WEBB, 29, King Edward Rd., Dentons Green, St. Helens. Technical Chemist. Associate of Institute of Chemistry. Asst. works manager, Messrs. Tennants (Lancs.), Ltd., Manchester; previously head chemist for same; prior to this, asst. chemist with United Alkali Co. (*Signed by*) Chas. W. Carpenter, E. L. Rhead, Wm. E. Kay.

BISHOP, WILLIAM ALFRED, 33, Banstead Rd., Purley, Surrey. Chemist. B.A. Cantab. (*Signed by*) H. J. H. Fenton, C. T. Heycock, F. W. Dootson.

BRISLEY, CHARLES WILLIAM, 105, Thurlow Park Road, London, S.E.21. Student at University of Oxford. B.A. Honours (Chemistry). At present doing research for Part II. School of Natural Science (Chemistry). (*Signed by*) F. D. Chattaway, E. Hope, S. G. P. Plant, M. P. Applebey, L. Gowing-Scopes.

BURKE, KATHARINE ALICE, 17, Lion Gate Gardens, Richmond, Surrey. Assistant Lecturer in Chemistry, University College, W.C.1. B.Sc. London. Published papers in the *Journal of the Chemical Society*, and *Zeitschrift fur Physikalische Chemie*. Translated "Thermochemistry," by Julius Thomsen, in Sir Wm. Ramsay's series. (*Signed by*) J. N. Collie, Irvine Masson, F. G. Donnan.

BURRAGE, LESLIE JAMES, 32, Emerson Road, Ilford, Essex. Undergraduate at King's College, London. Taking Hons. B.Sc., Chemistry (Oct. 1921). B.Sc. degree obtained Oct. 1921. Being now engaged on research work at King's College I wish to keep in touch with fresh chemical developments. (*Signed by*) Arthur Fairbourn, A. J. Allmand, Harold Toms.

BYLES, JOHN EDWARD, 87, Newlands Park, Sydenham, S.E.26. Chemist, Class II, Government Laboratory. B.Sc. (Manchester), A.I.C. Nine years at Government Laboratory. (*Signed by*) A. G. Francis, F. S. Aumonier, J. R. Nicholls.

CHILD, REGINALD, Yateley Lodge Cottage, Yateley, Camberley. Honours Student of King's College, London. (*Signed by*) Samuel Smiles, H. W. Cremer, C. S. Salmon.

CHRISTIE, GEORGE HALLATT, 155, Sharrow Vale Rd., Sheffield. Engaged in Research in Organic Chemistry with Dr. J. Kenner, Sheffield University. Master of Science (1st Class Hons. Chemistry, Sheffield University). Organic Chemical Research under Dr. Kenner since June 1920. I wish to have access to the Society's Library and receive the Society's Journal. (*Signed by*) W. P. Wynne, J. Kenner, F. G. Tryhorn, A. W. Chapman, Thos. B. Smith.

COFMAN-NICORESTI, GABRIEL, 15, Ludlow Road, Ealing, W.5. Pharmacist Manufacturing Chemist, Member of the Pharmaceutical Society. Director of the Solidol Chemical Co., Mill Street, Dockhead, S.E.1. Inventor and patentee of improvements in or connected with stoves for burning solidified spirit or the like, No. 101569/16, and also improved liniment, No. 18060/15. I desire to make use of the Library and keep in touch with current chemical

literature. (*Signed by*) Jules Cofman-Nicoressti, Hartley Shaw, S. B. Tallantyre.

COURTNEY, WILLIAM BARRY, 12, Caithness Road, Mitcham, Surrey. Student, engaged on research work. I have been a student of the Chemical Department of the South-Western Polytechnic Institute for five years, and have passed the B.Sc. (Honours) examination of the London University in Oct. 1921. (*Signed by*) J. B. Coleman, F. H. Lowe, F. Arnall, J. C. Crocker.

CRUTCHLON, BERNARD WILLIAM ALFRED, 65, Durlston Rd., Upper Clapton, E.5. Honours Student of King's College, London. (*Signed by*) Samuel Smiles, H. W. Cremer, C. S. Salmon.

CURRIE, ARCHIBALD NEIL, M.A., B.Sc. A.I.C., 44, Fisher St., Dennistoun, Glasgow. (The applicant holds the Glasgow University B.Sc. degree with special distinction in his principal subject—Chemistry.) Biochemist to the Glasgow Royal Cancer Hospital. The applicant has two papers ready for publication. The first is "The Fat of Adipose Tissue in Malignant Disease," and will be published shortly either as a Hospital Bulletin or in the *Journal of Exptl. Pathology and Bacteriology*. The second, which is one of a series of "Studies in Fat Pigment," is entitled: "A Preliminary Note on the Separation of Pigment from Accompanying Fat." (*Signed by*) G. G. Henderson, T. S. Patterson, J. Roberts.

DALTON, JAMES OWEN CHESSHYRE, The Plot, Adams Road, Cambridge. Engaged in Research work at Cambridge University. Honours Degree in Nat. Science Tripos Part II., June 1921. (*Signed by*) W. J. Pope, T. M. Lowry, W. H. Mills.

EDWARDS, ARTHUR WILLIAM, 4, Dunvegan Road, Eltham, S.E.9. Chemist. (1) Early education at Dorchester Grammar School; winner of the science prize in 1902. (2) Studied Analytical Chemistry and Physics at Duncan's School of Pharmacy under Mr. William Duncan, F.C.S., Ph.C., also at Darlington Technical College, and under private tutors. (3) Matriculated at Edinburgh University in 1909, for the study of chemistry. (4) Passed the minor examination of the Pharmaceutical Society of Great Britain in 1909; Member of the British Pharmaceutical Conference. (5) Engaged in bacteriological and pharmaceutical chemistry under Mr. J. E. Langford Moore, F.C.S., at St. Bartholomew's Hospital. (6) For five years with the English Company of Farbenfabriken vorm. Friedrich Bayer & Co., Elberfeld and Leverkusen, under Dr. C. R. Henning, Ph.D., F.C.S. (7) Served in the Royal Engineers—gas services—during the War. (8) At present with Messrs. R. W. Greeff & Co., Ltd., of Thames House, Queen Street Place, E.C.4, in an advisory and technical capacity. (9) Keenly desirous of keeping in touch with up-to-date Chemical Research and receiving the publications of the Chemical Society. (*Signed by*) J. E. Langford Moore, Percy Barrs, Frank B. Arnold, William Duncan, Charles Cockle.

ELLINGWORTH, SAMUEL, 74, Millfield Road, York. Research Chemist, Dept. of Organic Chemistry, University of Leeds. B.Sc. (Leeds), 1st Class Honours, 1916. M.Sc. 1918. Chemist to the British Dyestuffs Corporation (Huddersfield), Ltd. (1916–1921). (*Signed by*) J. B. Cohen, P. K. Dutt, Arthur Smithells, Akira Shimomura.

FISHER, KENNETH, Eton College. Schoolmaster. M.A. (Oxon), Ph.D. (Jena). (*Signed by*) W. H. Perkin, D. Ll. Hammick, R. H. Thomas, T. W. J. Taylor, E. Hope, N. V. Sidgwick.

FREDERIKSE, WILLIAM ALEXANDER, Amsterdam, Honthorststraat 2. Chemical Student, University of Amsterdam. Qualifications none. Desires to become a fellow of the Chemical Society, in order to profit by the Journal

and other publications. (*Signed by*) H. J. Backer, J. C. Irvine, R. Robinson.

GOEL, MIRI LAL, Royal Artillery Bazar, Ambala, Punjab (India). 2nd Professor of Chemistry, Meerut College, Meerut (U.P.). B.Sc. with Honours in Chemistry; M.Sc. of the Punjab University; Research Scholar of the Punjab University for some time; Demonstrator of the B.Sc. IV year class Govt. College Chem. Lab. while student of the M.Sc. class; Professor of the Divisional College, Meerut (U.P.), since November 1919, taking both Intermediate and B.Sc. classes. Joint author of the paper published in J.C.S. (Trans., 210; 1921 Feb.); worked as Consulting Chemist of the Goel Scientific and Technical Works, Meerut; Organised "Iswar" Soap Factory, R.A. Bazar, Ambala; doing research work on essential oils and dyeing extracts from vegetable dyestuffs in the College at present. (*Signed by*) Horace B. Dunicliff, N. A. Yajnik, Edwin Theodore Dean, Shanti Swarupa Bhatnagar, Bawa Kartar Singh.

GOSS, FRANK ROBERT, 4, Abbotsford Rd., Goodmayes, Ilford, Essex. Neil Arnott Research Scholar of the University of London. Student from 1918 to 1921 at the Royal College of Science, obtaining on completion of course the Associateship (A.R.C.S.) in Chemistry, and B.Sc. (Lond.) with Honours in Chemistry. I have been awarded the Neil Arnott Medal in Chemistry of the Univ. of London, and am at present working in the Organic Chemistry Research Laboratories of the Royal Coll. of Science. (*Signed by*) Jocelyn Thorpe, M. A. Whiteley, C. K. Ingold.

GREY, EGERTON CHARLES, Medical School, Cairo, Egypt. Professor of Chemistry. D.Sc., M.B. Author of several papers on the Chemical Action of Bacteria, published in the Proc. of the Royal Society. (*Signed by*) Arthur Harden, F. R. Henley, Robert Robison, Ida Smedley Maclean.

GRIMWOOD, ROBERT CHARLES, 11, Billiter Square, E.C.3. Research Student, Royal College of Science. Associate City and Guilds, Finsbury Institute. Registered Student at Institute of Chemistry. (*Signed by*) Jocelyn Thorpe, Robt. G. Grimwood, M. A. Whiteley, C. K. Ingold.

HAMMOND, GEORGE GRIMSHAW, 254, Mill Street, Toxteth, Liverpool. Analyst and Chemist. Bachelor of Science (Manchester). Member of the Pharmaceutical Society. (*Signed by*) Arthur W. Duncan, Frederick Owen, Walter Salmon.

HAMOR, WILLIAM ALI EN, Mellon Institute of Industrial Research, University of Pittsburgh, Pittsburgh, Pa., U.S.A. Assistant Director of the Mellon Institute of Industrial Research. Educated, University of Pittsburgh, 1904-1907; Master of Arts, 1913. Research Chemist, College of the City of New York, 1907-1914. Assistant to the Director, Mellon Institute, Pittsburgh, Pa., 1914-1916. Assistant Director, Mellon Institute, since 1916. Shale Oil Expert, Department of Mines of Canada, 1908. Editorial Staff, *Journal of Industrial and Engineering Chemistry*, 1912-1914. Editorial Board Member, "The Science History of the Universe," 1909. Commissioned a Major in the Chemical Warfare Service, United States Army, November 28, 1917; served in France ten months as assistant chief of the Technical Division, C.W.S., A.E.F.; honourably discharged, December 6, 1918. Member of the National Research Council on the Division of Research Extension, 1921-1924. Member of the National Research Council's Sub-Committee on Airplane Fuels, 1917. Member of various committees of the American Chemical Society, 1913-1921. Specialist in hydrocarbon chemistry. Author of the following books: "A History of Chemistry," 1909; "The American Petroleum Industry" (with R. F. Bacon), 2 vols., 1916; "The Examination

of Petroleum" (with F. W. Padgett), 1920; and "American Fuels" (with R. F. Bacon), 2 vols., 1922. Also author of a series of papers and reports of researches on oil-shales, petroleum, fuels, and anaesthetics. Contributor to scientific journals on educational subjects, especially on the value of industrial research and chemistry. Member of the American Chemical Society, American Electrochemical Society, and Alpha Chi Sigma and Phi Lambda Upsilon fraternities. Residence: 1030, Victoria Avenue, New Kensington, Pa. Business Address, Mellon Institute of Industrial Research, Pittsburgh, Pa., U.S.A. (*Signed by*) Marston Taylor Bogert, Raymond F. Bacon, Chas. Baskerville.

HART, LESLIE RALPH, 'Beaulieu,' St. Albans, Herts. Research Student. London University B.Sc. Examination in Honours Chemistry, 1st Class, November 1921. (*Signed by*) Samuel Smiles, A. J. Allmand, A. Fairbourn, D. T. Gibson.

HUME-ROTHERY, WILLIAM, Mendip House, Headington Hill, Oxford Research Student in Chemistry. I am a Natural Science Demy of Magdalen College, Oxford, and have taken the Oxford B.A. Degree in Chemistry (Part I of the Final Honours Course) and am now engaged in Research Work in Inorganic Chemistry (for Part II of the Final Honours Course). I intend to devote my life to Research Work and wish to join the Society to keep in touch with the developments of the Science. (*Signed by*) Frederick Soddy, B. Lambert, M. P. Applebey, A. F. Walden, W. H. Perkin, E. Hope.

IRVINE, FRANK MOUAT, 30, Clayton Road, Jesmond, Newcastle-on-Tyne. Lecturer in Chemistry at Armstrong College, Newcastle. B.Sc. (1st Class Hons. Chemistry) (Dunelm), A.I.C. Formerly Research Chemist with Messrs. Cookson & Co., Ltd., Lead and Antimony Manufacturers, Willington, Quay-on-Tyne. (*Signed by*) W. N. Haworth, Grace C. Leitch, J. A. Smythe, C. F. Allpress.

JENKINSON, JOHN ALBERT, School House, Little Sutton, nr. Birkenhead. Science Master at the Plympton Grammar School, South Devon. Student at Liverpool University (Final Examination 1921). B.Sc. (Hons. Chem.). Teacher of Science at Plympton Grammar School. (*Signed by*) E. C. C. Baly, W. C. McC. Lewis, I. M. Heilbron.

JOHNSON, ROWLAND NICHOLAS, 'Greenstede,' West Hill, East Grinstead, Sussex. Research Student at King's College, London. Honours B.Sc. (London), November 1921. First Class in Chemistry with Physics. (*Signed by*) A. J. Allmand, Samuel Smiles, A. Fairbourn.

KOHLI, SARAL JANG, M.Sc., Sat Updesh, Lahore (Punjab), India. Professor of Chemistry, Prince of Wales College (Kashmir State College), Jammu (Kashmir), India. (1) A First Class B.Sc. of the Punjab University, standing first in my combination (Physics and Chemistry) of subjects. (2) A First Class M.Sc. in Chemistry, standing first in the Punjab University. (3) Have done some work on the Radioactivity of Indian Minerals. (4) Have great desire to continue my studies and research work. (5) Am working as a Professor of Chemistry, teaching B.Sc. classes. (*Signed by*) N. A. Yajnik, E. T. Dean, S. R. Khosla.

LAKE, DAVID JOHN, 36, Gerald St., Hafod, Swansea. Science Master, Daventry Grammar School. Passed B.Sc. (London), 1920. Student Swansea Technical College, 1916-1920; Student Swansea University, 1920-Jan. 1921; since Jan. 1921, Science Master, Daventry Grammar School, Northants. (*Signed by*) E. A. Tyler, T. H. Hooper, D. G. Hopkins.

LANGWELL, HUBERT, Wentworth, Denham Road, Epsom, Surrey. Research Chemist. Ten years research on the fermentation of celluloses. (*Signed by*) C. F. Cross, Robt. Ensoll, E. J. Charlton, William Bacon.

LEGG, VERNON HOWES, 95, Carter Knowle Road, Sheffield. Student, Sheffield University. Took the three years' day course in Chemistry at Sheffield University and obtained the B.Sc. Honours Degree (Chemistry). Am at present engaged in Research Work at the Applied Science Dept. (*Signed by*) W. P. Wynne, R. V. Wheeler, J. Kenner, Arthur W. Chapman.

MCRÆ, JOHN ALEXANDER, Queen's University, Kingston, Ont., Canada. Assistant Professor of Chemistry, Queen's University. M.A. (Queen's Univ., Kingston), Ph.D. (Manchester), F.I.C. 1912-1915, Lecturer on Chemistry, Queen's University, Kingston, Canada. 1915-1919, Asst. Professor of Chemistry, Queen's University, Kingston, Canada. 1919-1921, Research Student at Univ. of Manchester as holder of Special Fellowship from the Honorary Advisory Council for Scientific and Industrial Research of Canada. (*Signed by*) Arthur Lapworth, F. P. Burt, Henry Stephen, F. Fairbrother, Frederick Challenger.

MATTHEWS, MARCUS AURELIUS, 'Burghley,' 47, Spratthall Rd., Wanstead, E.11. Student for M.Sc. Student at East London College for three years. B.Sc., Second Class Honours in Chemistry. (*Signed by*) F. G. Pope, F. W. Turner, J. R. Partington.

MAXTED, DURBAN ROY, 7, New Road, Reading. Lecturer, University College, Reading. A.R.C.Sc. (1918); B.Sc. (London). 1st Class Hons, Chemistry, 1919. Assistant chemist to Bow Bridge Dye Works, London, E., June 1919-Jan. 1920. Lecturer in Organic Chemistry, University College, Reading, since Feb. 1920. (*Signed by*) H. Bassett, Reginald G. Durrant, J. W. Dodgson.

MORGAN, WALTER THOMAS JAMES, 171, Kensington Avenue, Manor Park, London, E.12. Student, East London College (University of London), Entering for B.Sc. (Hons. Chem.) in Oct. 1922. Inter. B.Sc., 1920. Assistant chemist in the Research Laboratory of the South Metropolitan Gas Company from June 1917-Oct. 1918. (*Signed by*) E. V. Evans, J. R. Partington, F. G. Pope.

MORRISON, JOHN SYDNEY, 23, Seymour Rd., Cheadle Hulme, Manchester. Technical Chemist and English Representative of Messrs. Perry & Hope, Ltd., Firth and Clyde Chemical Works, Nitshill, nr. Glasgow. 1908-1913, Manc. School of Technology, awarded diploma for pure and applied chemistry, also courses in gas engineering, gas manufacturing, bleaching and dyeing, dairy analysis. Ministry of Munitions, D.E.S., London, 1916-1919. Superintendent Chemist and Assistant Manager at Folly explosive gas factories, Gretna, Litherland, West Gorton, and Sutton Oak. Awarded O.B.E. July 1920. Messrs. Perry & Hope, Ltd., Glasgow, 1920-1921, Technical Chemist and Representative. (*Signed by*) James Booth, Herbert H. H. Lorenz, John H. Jeffery, John Turnbull.

OWEN, EDWARD JOHN, Porthdafarch Farm, Holyhead, N. Wales. Organic Chemist, employed by Messrs. The British Drug Houses, Ltd., City Road, London, N.1, in the manufacture of complex organic compounds since Oct. 3, 1921. B.Sc. (Wales) with Second Class Honours in Chemistry. (*Signed by*) Kennedy Orton, Owen Owen, L. Hunter, H. B. Watson, Frederick Soper.

PARSONS, ALFRED TENNYSON, 25, Westwell Rd., Streatham, S.W.16. Chemist, Class 2, Government Laboratory. B.Sc. (Dunelm), A.I.C. (*Signed by*) Robert Robertson, Geo. Stubbs, J. Carmichael.

PATTERSON, WILLIAM HAMILTON, University of London Club, 21, Gower St., W.C.1. Lecturer in Chemistry, East London College, University of London, M.Sc., University of Manchester, A.I.C. Assistant Versuchsanstalt, Karlsruhe, Germany. Carnegie Research Scholar, Iron and Steel Institute,

1913. Various Research Publications. (*Signed by*) F. G. Pope, O. L. Brady, W. E. Garner.

PODD, PARTIDGE LESLIE, 8, St. Peter's St., Ipswich. Post-graduate Research Student at University College, London. Passed B.Sc. Honours Exam. of London University, 1921. As a research student, I desire to have the privilege of using the library of the Society, and I also desire to keep in touch with the advance in knowledge of all branches of chemistry through the medium of the Chemical Society. (*Signed by*) Irvine Masson, Henry Terrey, J. N. E. Day.

RAU, KRISHNARAJPET VENKATA, Ananda College, Colombo, Ceylon. Assistant Master, Ananda College. B.A. (Madras), B.Sc. (London). Ten years' teaching experience. (*Signed by*) J. P. C. Chandrasena, W. N. Rae, M. Parthasarathy.

RAY, FRANK CRAFER, 47, Alfriston Rd., West Side, Clapham Common, S.W.11. Assistant Lecturer, Chemical Department, Battersea Polytechnic, S.W.11. B.A. (Cantab.) (*Signed by*) Harold Hunter, A. Houssa, S. Greenberg, J. L. White, Robert H. Pickard, J. Kenyon.

STEVENSON, FRANCIS MALCOLM, 26, Albert St., Belper, nr. Derby. Dyers' Chemist. Graduated at Leeds University in Colour Chemistry (Hons.), and now desirous of keeping in touch with chemical advances and research. (*Signed by*) A. G. Perkin, R. B. Forster, J. B. Cohen.

STEWART, JESSIE, School House, Helmington Row, Crook, Co. Durham. Research Student. B.Sc. Durham University. First Class Honours in Chemistry. (*Signed by*) Samuel Smiles, A. J. Allmand, David T. Gibson.

TATNER, GILBERT HENRY, 52, Melrose Street, Hull. Student, University of Leeds. Studying for Honours B.Sc. (Chemistry). Date of Honours Examination, 1922. (*Signed by*) Arthur Smithells, J. B. Cohen, H. M. Dawson, M. Sybil Leslie, W. Lowson.

TANSLEY, LEONARD BEAUMONT, M.C., M.A., 'Espérance,' Purley, Surrey. Science Coach. Honours Graduate in Chemistry, etc., of Trinity College, Cambridge University. Lecturer and Examination Coach in Science at the General Engineering College, S.W.5. (*Signed by*) Francis Arnall, Frank Matthews, J. C. Crocker.

TAYLOR, GEORGE, Frigorifico, Campana, F.C.C.A., Argentine, S. America. Chief Chemist. Member of British Assn. of Chemists. Late Asst. Chemist to Liverpool Gas Co. (*Signed by*) Harold A. Auden, Frank B. Small, J. C. Frodsham.

THOMSON, ROBERT WILLIAM MURRAY, 117, Highbury Hill, London, N.5. Lecturer. B.Sc. of St. Andrews University. Assistant Lecturer in Physical Chemistry at King's College, University of London. (*Signed by*) Samuel Smiles, H. W. Cremer, John H. Foulger.

WALMSLEY, JAMES RAWSON, Dean Brook Cottage, Dean Lane, Moston, Manchester. Analyst and Pharmaceutical Chemist. Member of the Pharmaceutical Society of Gt. Britain. Associate of the Institute of Chemistry. Associate of the Manc. College of Technology (Genl. Chemical Technology). Lecturer (part-time) College of Technology, Manc. (*Signed by*) Frederick Owen, Samuel E. Hodgkinson, Walter Salmon, William Jackson Lund, Jas. Grant.

WHITE, WALTER ARTHUR, 144, Shelley Avenue, Manor Park, E.12. Assistant Analytical Chemist. Assistant to Dr. G. W. Clough in his analytical practice at the Royal Veterinary College, Camden Town. Formerly assistant to Dr. G. D. Lander. Student at the Sir John Cass Institute for the Final B.Sc. (Honours Chemistry) Examination of the University of London. (*Signed by*) G. W. Clough, W. H. Barker, E. D. Griffiths, W. T. Clough.

WIGHTMAN, WILLIAM PERSEHOUSE DELISLE, 59, Holmewood Gardens, Streatham Hill, London, S.W.2. Student. Student for degree in Chemistry. (*Signed by*) James C. Philip, M. A. Whiteley, H. F. Harwood, H. Ellingham.

WISEMAN, JULIUS, B.Sc., 102, Cambridge Rd., Mile End, E.1. Demonstrator Chemistry Dept., Guy's Hospital, and Student in Medicine. (*Signed by*) C. S. Gibson, A. J. Allmand, Samuel Smiles.

WOOSLEY, DUNCAN PAX, 96, Roxborough Road, Harrow, Middlesex. Chemical Laboratory Assistant. With The British Drug Houses, Ltd., Graham St., City Rd., N.1., since Aug. 1919. Student at Birkbeck College Univ. of London, Sept. 1919 to date, reading for B.Sc. Hons. Chem. I desire to have access to the Library and to keep in touch with modern developments. (*Signed by*) Francis H. Carr, O. A. Le Beau, James Taylor.

WRIGHT, ROBERT NORMAN, 16, Redcliffe St., South Kensington, S.W.10. Assistant Chemist to the Dried Milk Dairy Products Company, Hemyock, Devon. A.R.C.S. (1st Class in Chemistry), B.Sc. (Lond.). (1st class Honours), A.I.C. Dec. 1920, appointed assistant chemist in the Research Laboratory of Messrs. J. Lyons & Co., Ltd. (*Signed by*) H. B. Baker, James C. Philip, A. E. Dunstan, F. B. Thole, Philip C. Bull.

Certificates have been accepted by the Council under Bye-Law I (2) in favour of the following :

BEARPARK, ARTHUR FREDERICK, 'Kingsfold,' Mill Street, Capetown. Postal Address, P.O. Box 1159, Capetown. Postal address in U.K. c/o Henry Tyrer & Co., Ltd., 360, Royal Liver Building, Liverpool. General Manager, Whaling Companies. Three years pupil with H. Thompson, Esq., F.C.S., Hull. Five years chemist to British Oil and Cake Mills, Limited. Member of the Royal Society of South Africa. Member of the South African Chemical Institute. Member of the Chemical, Metallurgical and Mining Society of South Africa. (*Signed by*) Harry Thompson, *Edward B. Maxted*.

SEN, SURESH CHANDRA, Mungpoo P.O., Riyang D. H. Ry., India. Assistant Quinologist, Government of Bengal. B.Sc. (Calcutta), B.A. (Cantab.). Desires to receive the Journal. (*Signed by*) G. E. Shaw.

PROCEEDINGS

OF THE

CHEMICAL SOCIETY.

Ordinary Scientific Meeting, Thursday, February 2nd, 1922, at 8 p.m., Professor H. E. ARMSTRONG, LL.D., F.R.S., Vice-President, in the Chair.

The Chairman referred to the loss sustained by the Society, through death, of the following Fellows :

	Elected.	Died.
Thomas Duxbury	Dec. 15th, 1892.	Nov. 26th, 1921.
James Hill Fairweather	June 18th, 1908.	
Herbert Garland	May 15th, 1913.	April, 1921.
David Emrys Williams	Dec. 6th, 1917.	Jan. 9th, 1922.

The following were formally admitted Fellows of the Chemical Society: H. C. Bell, S. F. Birch, W. A. P. Challenor, P. C. Chaumeton, L. G. F. Dolley, D. T. Gibson, E. J. Hartung, D. N. Jackman, J. H. Middleditch, J. R. Pickering, E. H. Usherwood, I. W. Wark, and L. D. Williams.

Certificates were read for the first time in favour of :

George Macdonald Bennett, M.A., M.Sc., F.I.C., 94, Mercers Road, Tufnell Park, N.19.

Eric Sykes Brooks, Verandah Buildings, Dalton-in-Furness, Lancs.

Fred Brumfitt, 123, Tennyson Place, Bradford.

Carlton Butler, B.Sc., A.I.C., 23, Winifred Road, Urmston, Manchester.

Francis Arthur Freeth, O.B.E., M.Sc., F.I.C., Heather Croft, Hartford.

Thomas Weatherill Harrison, B.Sc., F.I.C., 18, Elfindale Road, S.E.24.

Eusebius Holmes, B.Sc., Eastern Green, Coventry.

Lingaiah Basavalingaiah Manjunath, B.A., University College, Gower Street, W.C.2.

George Flaxman Underhay, B.Sc., Allerton, Woodside Avenue, Muswell Hill, N.10.

William Andrew Shakespeare Vardy, Winksley Vicarage, Ripon.

A Certificate has been authorised by the Council for presentation to ballot under Bye-Law I (2) in favour of:

Suresh Chandra Sen, B.A., B.Sc., Mungpoo P.O., Riyang, D.H.Ry., India.

The following papers were read :

"The action of light on silver bromide." By E. J. HARTUNG.

"The structure of the benzene nucleus. Part I. Intra-nuclear tautomerism." By C. K. INGOLD.

"The structure of the benzene nucleus. Part II. Synthetic formation of the bridged modification of the nucleus." By C. K. INGOLD.

"The structure of the benzene nucleus. Part III. The suppression of intra-nuclear change." By C. K. INGOLD and H. A. PIGGOTT.

"Platinum catalyst for automatic gas lighters." By L. D. WILLIAMS.

Ordinary Scientific Meeting, held in the Lecture Hall of the Institution of Mechanical Engineers, on Thursday, February 9th, 1922, at 8 p.m., SIR JAMES WALKER, D.Sc., LL.D., F.R.S., President, in the Chair.

SIR ERNEST RUTHERFORD, D.Sc., F.R.S., delivered his Lecture entitled "Artificial Disintegration of Elements." A vote of thanks to the Lecturer, proposed by Professor H. E. Armstrong, seconded by Professor F. G. Donnan, and supported by the Chairman, was carried with acclamation, and acknowledged by Sir Ernest Rutherford.

Ordinary Scientific Meeting, Thursday, February 16th, 1922, at 8 p.m., SIR JAMES WALKER, D.Sc., LL.D., F.R.S., President, in the Chair.

The PRESIDENT announced that the Society had lost, through death, the following Fellows :

	Elected.	Died.
Charles Baskerville.....	April 21st, 1898	Jan. 28th, 1922
George Gilbert Pond	April 22nd, 1903	May 20th, 1920

The following were formally admitted Fellows of the Chemical Society : A. N. Campbell, W. Davies, A. A. Davis, F. Dickens, A. S. Russell, P. M. Tooke, and M. Williams.

Certificates were read for the first time in favour of :

Frederick Mason Brewer, Lincoln College, Oxford.

Alfred Duncan Gay, 49, Thornlaw Road, West Norwood, S.E.27.

Douglas Cecil Henry, M.A., Hulme Hall, Victoria Park, Manchester.

William Hurst, 2, Edditch Grove, Tonge, Bolton.
 Thomas Iredale, B.Sc., 94, Tyrwhitt Road, Brockley, S.E.4.
 George Lewis Kennaby, B.Sc., Anglo-Persian Oil Co., Abadan, Persian Gulf.
 George Allan Russell, B.Sc., A.I.C., Barnsheen, Ditton, Widnes.
 Ernest Arthur Seeley, B.Sc., 10, Hamilton Road, N.5.
 Brian Duncan Shaw, Ferns' Hollow, Station Road, Ilkeston.
 Arthur Frederick Watson, M.Sc., A.I.C., Brockwell Hall, Herne Hill, S.E.23.
 Llywelyn Williams, B.Sc., Herne Bay College, Kent.

The following Certificate has been authorised by the Council for presentation to ballot under Bye-Law I (2) in favour of :

Walter Albert Lawrance, M.A., Ph.D., Bates College, Lewiston, Maine, U.S.A.

Messrs. R. Whytlaw Gray and F. G. Pope were elected Scrutators, and a ballot for the election of Fellows was held. The following were subsequently declared elected as Fellows :

Jnanendra Kumar Adhya, M.Sc.	Rowland Nicholas Johnson, B.Sc.
Harry Charles Applebee, A.I.C.	Saral Jang Kohli, M.Sc.
Gilbert James Avey.	David John Lake, B.Sc.
John William Baker, B.Sc.	Hubert Langwell.
Bhagwan Das Batta, B.Sc.	Vernon Howes Legg, B.Sc.
Arthur Frederick Bearpark.	John Alexander McRae, M.A., Ph.D.,
Darrell Webb Beesley, A.I.C.	F.I.C.
William Alfred Bishop, B.A.	Marcus Aurelius Matthews, B.Sc.
Charles William Brisley, B.A.	Durban Roy Maxted, B.Sc.
Katharine Alice Burke, B.Sc.	Walter Thomas James Morgan.
Leslie James Burrage, B.Sc.	John Sydney Morrison.
John Edward Byles, B.Sc., A.I.C.	Edward John Owen, B.Sc.
Reginald Child.	Alfred Tennyson Parsons, B.Sc., A.I.C.
George Hallatt Christie, M.Sc.	William Hamilton Patterson, M.Sc.,
Gabriel Cofman-Nicoresti.	A.I.C.
William Barry Courtney, B.Sc.	Partridge Leslie Podd, B.Sc.
Bernard William Alfred Crutchlow.	Krishnarajpet Venkata Rau, B.A.,
Archibald Neil Currie, M.A., B.Sc.,	B.Sc.
A.I.C.	Frank Crafer Ray, B.A.
James Owen Chesshyre Dalton, B.A.	Suresh Chandra Sen, B.A., B.Sc.
Arthur William Edwards.	Francis Malcolm Stevenson, B.Sc.
Samuel Ellingworth, M.Sc.	Jessie Stewart, B.Sc.
Kenneth Fisher, M.A., Ph.D.	Gilbert Henry Tatner.
William Alexander Frederikse.	Leonard Beaumont Tansley, M.A.
Miri Lal Goel, M.Sc.	George Taylor.
Frank Robert Goss, B.Sc.	Robert William Murray Thomson,
Egerton Charles Grey, D.Sc., M.B.	B.Sc.
Robert Charles Grimwood.	James Rawson Walmsley, A.I.C.
George Grimshaw Hammond, B.Sc.	Walter Arthur White.
William Allen Hamor, M.A.	William Persehouse Delisle Wightman.
Leslie Ralph Hart, B.Sc.	Julius Wiseman, B.Sc.
William Hume-Rothery, B.A.	Duncan Pax Woosley.
Frank Mouat Irvine, B.Sc., A.I.C.	Robert Norman Wright, B.Sc., A.I.C.
John Albert Jenkinson, B.Sc.	

The following papers were read :

- " A theoretical derivation of the principle of induced alternate polarities." By A. LAPWORTH.
- " An explanation of the property of induced polarity of atoms and an interpretation of the theory of partial valencies on an electronic basis." By W. O. KERMACK and R. ROBINSON.
- " Some reactions of benzanthrone." By A. G. PERKIN and G. D. SPENCER.
- " A rapid iodimetric estimation of copper and iron in mixtures of their salts." By I. W. WARK.

ANNUAL GENERAL MEETING.

The Annual General Meeting will take place on Thursday, March 30th, at 4.30 p.m. An Informal Meeting will be held the same evening at 8 p.m.

REPORTS ON THE PROGRESS OF APPLIED CHEMISTRY.

(Issued by the Society of Chemical Industry.)

The Society of Chemical Industry has generously offered to supply Volume VI of these Reports to Fellows of the Chemical Society, who are not Members of the Society of Chemical Industry, at the reduced price of 10/-, post free.

Orders, with remittance, should be sent to Dr. J. P. Longstaff, Secretary, Society of Chemical Industry, 46 and 47, Finsbury Square, London, E.C. 2.

ADDITIONS TO THE LIBRARY.

I. *Donations.*

KINGZETT, CHARLES THOMAS. The popular chemical dictionary. 2nd edition. London 1921. pp. viii + 540. (*Reference.*) 21s. net.

From the Author.

RUSSELL, T. H. The planning and fitting-up of chemical and physical laboratories. London 1903. pp. xx + 178. ill. (Second Copy.) (*Recd.* 10/1/22.)

From the Author.

II. *By Purchase.*

CHAMOT, ÉMILE MONNIN. Elementary chemical microscopy. 2nd edition. New York 1921. pp. xvi + 480. ill. 25s. net. (*Recd.* 21/1/22.)

P R O C E E D I N G S

O F T H E

C H E M I C A L S O C I E T Y.

Ordinary Scientific Meeting, Thursday, March 2nd, 1922, at 8 p.m., Dr. M. O. FORSTER, F.R.S., Treasurer, in the Chair.

It was announced that the Society had lost, through death, the following Fellows :

	Elected.	Died.
James Robert Appleyard.....	May 15th, 1890.	Oct. 1921.
Thomas Lynton Briggs	April 19th, 1883.	April 3rd, 1921.
Edward William Taylor Jones	April 6th, 1871.	Feb. 19th, 1922.

The following were formally admitted Fellows of the Chemical Society : F. B. Arnold, J. W. Baker, A. W. Edwards, F. R. Goss, H. Hunter, W. H. Patterson, F. C. Ray, and S. P. Schotz.

The following List of Nominations for vacant places on the Council, the names of those nominated by the Council being indicated by an asterisk, was read from the Chair :

Vice-Presidents who have filled the office of President (two vacancies).—*H. B. Dixon; *P. F. Frankland.

Vice-Presidents who have not filled the office of President (two vacancies).—*E. C. C. Baly; *T. M. Lowry.

Secretary (one vacancy).—*Irvine Masson.

Ordinary Members of Council : (a) Town Members, that is, those resident within 50 miles of Charing Cross (three vacancies).—C. Dorée; J. J. Fox; J. R. Partington. (b) Country Members, that is, those resident beyond 50 miles from Charing Cross (three vacancies).—*I. M. Heilbron; J. W. McBain; *W. H. Mills.

Certificates were read for the first time in favour of :

Brojo Kishore Banerjee, B.Sc., Jamshedpur Technical Institute, Tata Nagar, India.

Benoy Bhusan Basu, B.Sc., M.B., 1/1, Nepal Bhattacharji's Street, Kalighat P.O., Calcutta.

Leslie Ashburn Dunn, 9, Pembroke Terrace, Bridlington.
 Richard Fisher, Hudshaw House, Hexham, Northumberland.
 Herbert Hodgson, 35, Ryan Street, Bradford.
 Arthur Cyril Percy Lunn, King's College, Cambridge.
 Robert Craig McDowall, 6, Ascog Street, Crosshill, Glasgow.
 Akella Suryanarayana Rao, B.Sc., Vizagapatam, India.
 Harry Lister Riley, 7, Trafalgar Square, Chelsea, S.W.3.
 James Robinson, B.Sc., Milton High School, Bulawayo, S. Rhodesia.
 Suhrmania Sastry, B.A., 5, Krishnappa Naik Agrabaram, Georgetown,
 Madras.

The following papers were read :

- "New halogen derivatives of camphor. Part II. α' -Bromo-camphor." By T. M. LOWRY, V. STEELE, and H. BURGESS.
 "The rotatory dispersive power of organic compounds. Part X. The preparation and properties of pure ethyl tartrate." By T. M. LOWRY and J. O. CUTTER.
 "The molecular configuration of polynuclear aromatic compounds. Part I. The resolution of γ -6:6'-dinitro- and 4:6:4':6'-tetranitrodiphenic acids into optically active components." By G. H. CHRISTIE and J. KENNER.

Ordinary Scientific Meeting, Thursday, March 16th, 1922, at 8 p.m., SIR JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

A. T. Parsons was formally admitted a Fellow of the Chemical Society.

Certificates were read for the first time in favour of :

John Wilham Blagden, M.A., Ph.D., F.I.C., Apple Tree House, Grove Road, Woodford, E.18.
 Charlie Lyons Carter, M.Sc., 9, Carr Street, N.E. Valley, Dunedin, N.Z.
 Arthur Rockett Chadwick, Corporation Sewage Works, Marley, Keighley.
 Douglas Creese Harrison, B.Sc., 18, Church Crescent, Muswell Hill, N. 10.
 Henry Unwin Harrow, 13, Southwark Bridge Road, S.E. 1.
 Ethel Mary James, B.Sc., 1, Rusham Road, Balham, S.W. 12.
 Edward Ingleson Lawson, 65, Warrington Road, Newcastle-on-Tyne.
 John Mylne Mullaly, 14, Bardwell Road, Oxford.
 Harold Edward Charles Powers, B.Sc., 4, Grove Crescent, Woodford, E. 18.
 Albert W. Smith, B.S., Ph.D., Case School of Applied Science, Cleveland, Ohio, U.S.A.
 Clarence Stott, B.Sc., A.I.C., Benroyd, Holywell Green, Halifax.
 William Thomas, B.A., M.Sc., A.I.C., Penfro, Stoneywood, Aberdeen.
 John George Walters, B.Sc., 46, Durham Road, East Finchley, N. 2.
 William Wallace Watt, 29, Woodstock Road, Golders Green, N.W. 4.
 Harry Alexander Wood, Santa Elena, Entre Rios, Argentine.

The following Certificate has been authorised by the Council for presentation to ballot under Bye-Law I (2) :

Harilal Basu, P.O. Box 84, Rangoon, Burma.

The following papers were read :

- Change of properties of substances on drying." By H. B. BAKER.
 The influence of nitro-groups on the reactivity of substituents in the benzene nucleus. Part VI. The elimination of halogen during the reduction of halogenated nitro-compounds." By H. BURTON and J. KENNER.

REPORTS ON THE PROGRESS OF APPLIED CHEMISTRY.

(Issued by the Society of Chemical Industry.)

The Society of Chemical Industry has generously offered to supply Volume VI of these Reports to Fellows of the Chemical Society, who are not Members of the Society of Chemical Industry, at the reduced price of 10s. post free.

Orders, with remittance, should be sent to Dr. J. P. Longstaff, Secretary, Society of Chemical Industry, 46 and 47, Finsbury Square, London, E.C.2.

ADDITIONS TO THE LIBRARY.

I. *Donations.*

DUDLEY, DUD, and others. *Mettallum Martis*: or iron made with pit-coale, sea-coale, etc., 1665. A treatise of metallica. By JOHN ROVENZON, 1613. *Metallica*, or the treatise of metallica. By SIMON STURTEVANT, 1612. [Edited by JOHN N. BAGNALL. Wolverhampton 1854.] pp. xix + 43, ii + 30, ii + 120. ill. (*Reference.*) From Professor P. Phillips Bedson.

FISCHER, EMIL. *Aus meinem Leben*. Berlin 1922. pp. vi + 202. ill. (*Recd.* 18/2/22.) From the Publisher: Julius Springer.

HADFIELD, SIR ROBERT ABBOTT. The work and position of the metallurgical chemist, also references to Sheffield and its place in metallurgy. An address. [London 1921.] pp. 97. ill. (*Recd.* 2/3/22.) From the Author.

HILDITCH, THOMAS PERCY. A concise history of chemistry. 2nd edition. London [1922]. pp. xii + 276. 6s. net. (*Recd.* 24/2/22.) From the Publishers: Messrs. Methuen & Co.

JOHNSTONE, SYDNEY JAMES. Potash. (Imperial Institute Monograph on Mineral Resources.) New edition. London 1922. pp. x + 122. 6s. net. (*Recd.* 2/3/22.) From the Director.

KANTHACK, R. (Compiler.) Tables of refractive indices. Vol. II. Oils, fats, and waxes. Edited by JOHN NAISH GOLDSMITH. London 1921. pp. 296. 25s. net. (*Recd.* 3/3/22.)

From the Publishers : Messrs. Adam Hilger Ltd.

MINISTRY OF MUNITIONS and DEPARTMENT OF SCIENTIFIC AND INDUSTRIAL RESEARCH. Technical records of explosives supply 1915-1918. No. V. Manufacture of sulphuric acid by contact process. London 1921. pp. vi + 128. ill. 25s. net. (*Recd.* 3/3/22.)

— — — No. VI. Synthetic phenol and picric acid. London 1921. pp. vi + 98. ill. 15s. net. (*Recd.* 14/2/22.)

From the Department.

PAULI, WOLFGANG. Colloid chemistry of the proteins. Part I. Translated by PERCY CYRIL LESLEY THORNE. London 1922. pp. xii + 140. ill. 8s. 6d. net. (*Recd.* 28/2/22.)

From the Publishers : Messrs. J. & A. Churchill.

II. By Purchase.

ARCHIVIO DI FARMACOLOGIA SPERIMENTALE E SCIENZE AFFINI. Year XVIII, Vol. XXVII, etc. Siena 1919 +. (*Reference.*)

BAUER, K. H. Analytische Chemie der Alkaloide. Berlin 1921. pp. vi + 425. ill. (*Recd.* 18/2/22)

BEILBY, SIR GEORGE. Aggregation and flow of solids : being the records of an experimental study of the micro-structure and physical properties of solids in various states of aggregation. 1900-1921. London 1921. pp. xvi + 256. + xxxiv plates. 20s. net. (*Recd.* 21/2/22.)

CHEMIKER-KALENDER 1922 Edited by WALTHER ROTH. Year 34. 2 vols. Berlin 1922. (*Reference*)

DEERR, NOËL. Cane sugar : a textbook on the agriculture of the sugar cane, the manufacture of cane sugar, and the analysis of sugar-house products. 2nd edition. London 1921. pp. viii + 644. ill. 42s. net. (*Recd.* 17/1/22.)

FOERSTER, FRITZ. Elektrochemie wässriger Lösungen. 3rd edition. Leipzig 1922. pp. xx + 900. ill. (*Recd.* 18/2/22.)

FORTSCHRITTE IN DER ANORGANISCH-CHEMISCHEN INDUSTRIE : an Hand der Deutschen Reichs-Patente dargestellt. Edited by ADOLF BRÄUER and JOHANNES D'ANS. Vol. I, 1877-1917, etc. Berlin 1921 +. ill. (*Reference.*)

GATTEFOSSÉ, R. M., and GATTEFOSSÉ, J. Nouveaux parfums synthétiques. Lyon 1921. pp. 208. ill. 5s. 6d. net. (*Recd.* 6/2/22.)

GOW, COLIN CLARENCE. The electro-metallurgy of steel. London 1921. pp. xvi + 351. ill. 27s. 6d. net. (*Recd.* 17/1/22.)

GRASSER, GEORG. Synthetic tannins: their synthesis, industrial production, and application. Translated by FINI G. A. ENNA. London 1922. pp. viii + 143. 12s. net. (*Recd.* 17/1/22.)

HIGGINS, SYDNEY HERBERT. Bleaching: being a résumé of the important researches on the industry published during the years 1908—1920. Manchester 1921. pp. viii + 138. 10s. 6d. net. (*Recd.* 21/1/22.)

HIRSCH, PAUL. Die Einwirkung von Mikroorganismen auf die Eiweisskörper. (Biochemie in Einzeldarstellungen. IV.) Berlin 1918. pp. x + 256. ill. (*Recd.* 3/3/22.)

HÖFER, JOHANNES. Die Fabrikation künstlicher plastischer Massen sowie der künstlichen Steine, Kunststeine, Stein- und Zementgüsse. 4th edition. Wien 1921. pp. xii + 366. ill. (*Recd.* 25/1/22.)

ILLINGWORTH, STEWART ROY, assisted by JENKYN GRIFFITHS. The analysis of coal and its by-products. London [1921]. pp. [xii] + 380. ill. 21s. net. (*Recd.* 17/1/22.)

KANITZ, ARISTIDES. Temperatur und Lebensvorgänge. (Biochemie in Einzeldarstellungen. I.) Berlin 1915. pp. x + 175. ill. (*Recd.* 3/3/22.)

KAUFFMANN, HUGO. Beziehungen zwischen physikalischen Eigenschaften und chemischer Konstitution. (Chemie in Einzeldarstellungen, Vol. X.) Stuttgart 1920. pp. xii + 422. (*Recd.* 13/2/22.)

KLIMONT, I. M. Der technisch-synthetische Campher. (Chem. Tech. in Einzeldarstellungen.) Leipzig 1921. pp. viii + 132. ill. (*Recd.* 3/3/22.)

KOLTHOFF, I. M. Der Gebrauch von Farbenindikatoren. Ihre Anwendung in der Neutralisationsanalyse und bei der colorimetrischen Bestimmung der Wasserstoffionenkonzentration. Berlin 1921. pp. iv + 144. ill. (*Recd.* 3/3/22.)

ORGANIC SYNTHESES. An annual publication of satisfactory methods for the preparation of organic chemicals. Edited by ROGER ADAMS, JAMES BRYANT CONANT, HANS THACHER CLARKE, and OLIVER KAMM. Vol. I, etc. New York 1921 +. ill. (*Reference.*) 8s. 6d. net.

PRINGSHEIM, PETER. Fluoreszenz und Phosphoreszenz im Lichte der neueren Atomtheorie. Berlin 1921. pp. viii + 202. ill. (*Recd.* 13/2/22.)

RICHTER, VICTOR VON. Organic chemistry, or chemistry of the carbon compounds. Edited by RICHARD ANSCHÜTZ and HANS MEERWEIN. II. Carbocyclic compounds. Translated from the 11th German edition by EDMUND EDWARD FOURNIER D'ALBE. London 1922. pp. xvi + 760. (*Reference.*) 35s. net.

RÖHMANN, FRANZ. Über künstliche Ernährung und Vitamine. (Biochemie in Einzeldarstellungen. II.) Berlin 1916. pp. vi + 150. ill. (*Recd.* 3/3/22.)

SIEGFRIED, MAX. Über partielle Eiweisshydrolyse. (Biochemie in Einzeldarstellungen. III.) Berlin 1916. pp. iv + 64. (*Recd.* 3/3/22.)

SOCIÉTÉ DE CHIMIE BIOLOGIQUE. Bulletin. Vol. I, etc. Paris 1914 +.

STEPHENSON, CHARLES H. Some micro-chemical tests for alkaloïds. Including "Chemical tests of the alkaloids used." By CHARLES EDWIN PARKER. London 1921. pp. iv + 110 + xxvii plates. 21s. net. (*Recd.* 17/1/22.)

TAYLOR, HUGH STOTT. Industrial hydrogen. New York 1921. pp. 210. ill. 20s. net. (*Recd.* 21/1/22.)

THOMSON, SIR JOSEPH JOHN. Rays of positive electricity and their application to chemical analyses. 2nd edition. London 1921. pp. x + 238. ill. 16s. net. (*Recd.* 21/1/22.)

TREADWELL, FREDERICK PEARSON. Analytical chemistry. Vol. I. Qualitative analysis. 5th English edition. Translated by WILLIAM T. HALL. New York 1921. pp. xviii + 598. ill. 23s. net. (*Recd.* 12/1/22.)

III. Pamphlets.

ARRHENIUS, SVANTE. Aus meiner Jugendzeit. Leipzig 1913. pp. 18.

ASSOCIATION OF BRITISH CHEMICAL MANUFACTURERS and BRITISH CHEMICAL PLANT MANUFACTURERS' ASSOCIATION. *Joint Research Committee on Chemical Plant*. Report of sub-committee for standardisation of jacketted pans. [London 1922.] pp. 14. ill.

CHEMICAL SOCIETY. Notes on the furnishing and equipment of chemical laboratories. London 1921. pp. 11.

CROMPTON, HOLLAND. Nebular zones and the formation of planets. [London 1921.] pp. 19.

IMPERIAL MINERAL RESOURCES BUREAU. The Mineral Industry of the British Empire and Foreign Countries. War Period. Mica (1913—1919). London 1922. pp. 32.

—— — Sulphur and iron-pyrites. (1913—1919.) London 1922. pp. 67.

SCIENTIFIC AND INDUSTRIAL RESEARCH, DEPARTMENT OF. *Building Research Board*. Special Report No. 4. The transmission of heat and gases through, and the condensation of moisture on the surface of, wall materials : an extract from the report of the Building Materials Research Committee. London 1921. pp. 19. ill.

SCIENTIFIC AND INDUSTRIAL RESEARCH, DEPARTMENT OF. *Fuel Research Board*. Fuel for motor transport. Second Memorandum. London 1921. pp. iv + 16.

— — — Special Report No. 1. Pulverised coal systems in America. By LEONARD C. HARVEY. Revised edition. London 1922. pp. iv + 117. ill.

SINGH, BAWA KARTAR. Recent advances in stereochemistry. (From the *J. Proc. Asiatic Soc. Bengal*, 1921, N.S. 17.)

CERTIFICATES OF CANDIDATES FOR ELECTION AT
THE BALLOT TO BE HELD AT THE ORDINARY
SCIENTIFIC MEETING ON THURSDAY, MAY 4TH,
1922.

N.B.—The names of those who sign from "General Knowledge" are printed in *italics*.

BANERJEE, BROJO KISHORE, Jamshedpur Technical Institute, G. Town, Jamshedpur, Tatanagar, India. Assistant, Chemical and Analytical Department, Jamshedpur Technical Institute, Tatanagar, India. B.Sc. (Calcutta University). Read up to M.Sc. Chemistry with Inorganic Chemistry as special subject in the post-graduate classes of the Calcutta University. Had been assistant in the Chemical and Metallurgical Dept. of the Bengal Engineering College, Sibpur, India, for two and a half years nearly. Now assistant in the Chemical and Analytical Dept. of the Jamshedpur Technical Institute since August 1921. Lecturing on General Inorganic Chemistry since March 1919. (*Signed by*) R. N. Sen, P. C. Chattopadhyay, J. Bhaduri.

BASU, BENOY BHUSAN, 1/1, Nepal Bhattacharji's Street, Kalighat P.O., Calcutta, India. B.Sc., M.B. Assistant Chemical Analyst to the Calcutta Corporation, intending to keep in touch with the modern advancement in Chemistry and the chemical world. (*Signed by*) B. N. Ghosh, T. K. Ghose, K. B. Sen.

BENNETT, GEORGE MACDONALD, 94, Mercers Road, Tufnell Park, N.19. Demonstrator, Chemical Department, Guy's Hospital Medical School. B.A., M.Sc. (London), M.A. (Cantab.), F.I.C. Fellow of St. John's College, Cambridge. Author of several papers in the Transactions since 1913. (*Signed by*) C. S. Gibson, D. C. Vining, Eustace E. Turner, H. C. Bell.

BLAGDEN, JOHN WILLIAM, Apple Tree House, Grove Road, Woodford, E.18. Head of Research Dept., Messrs. Howards & Sons, Ilford. M.A. (Cantab.), Ph.D. (Würzburg), F.I.C. 1899–1900 Demonstrator at the Electrochemical Institute of the Polytechnic, Darmstadt. 1900–1914 Research Chemist with Messrs. Boehringer & Sons, Mannheim. (1914–1919 Civilian Prisoner of War at Ruhleben.) 1919–1922 with Messrs. Howards & Sons as above stated. (*Signed by*) H. B. Baker, J. T. Hewitt, M. O. Forster.

BREWER, FREDERICK MASON, 42, Aberdeen Park, London, N.5. Student. Science Scholar of Lincoln College, Oxford. (*Signed by*) W. H. Perkin, T. W. J. Taylor, D. Ll. Hammick, E. Hope, S. G. P. Plant, N. V. Sidgwick.

BROOKS, ERIC SYKES, M.P.S., Verandah Buildings, Dalton-in-Furness,

Lancs. Chemist. Member of Pharmaceutical Society of Great Britain. Student of Chemistry for the past eight years, also as private coach for elementary chemistry. (*Signed by*) Alex. W. McLaren, Chas. Morton, Alfred Thomas.

BRUMFITT, FRED, 123, Tennyson Place, Bradford. Director and Manager, Wholesale Druggists. Diploma Chemistry and Dyeing, Bradford Technical College. Wholesale Druggist. (*Signed by*) V. Steele, F. W. Richardson, Barker North.

BUTLER, CARLTON, B.Sc., A.I.C., 23, Winifred Road, Urmston, nr. Manchester. Research Student in the Victoria University of Manchester. Lanc. C.C. Scholar 1917. Served Sp. Brig. R.E. until Jan. 1919. First Class Honours Chemistry 1921. Research, and part-time demonstrator, session 1921-22. (*Signed by*) H. B. Dixon, Colin Campbell, Frederick Challenger, Henry Stephen.

CARTER, CHARLIE LYONS, 9, Carr Street, N.E. Valley, Dunedin. Lecturer on Chemistry, Otago University, Dunedin, New Zealand. M.Sc. with 1st Class Honours in Chemistry, N.Z., 1919. A.I.C. (London), 1921. Demonstrator in chemistry, Otago University, 1917-19. Lecturer on Chemistry, 1919. Publication: "Mutton-bird Oil," *Jour. Soc. Chem. Ind.*, Sept. 1920, p. 220. (*Signed by*) J. K. H. Inglis, J. Williams, C. S. Hicks.

CHADWICK, ARTHUR ROCKETT, Corporation Sewage Works, Marley, Keighley. Chemist and Manager, Keighley Corporation Sewage Works. Eighteen and a half years' experience in the Chemistry of Sewage Disposals. Chemist and Manager, Keighley Corporation Sewage Works for two years. Chemist and Manager, Bingley U.D.C. Sewage Works for five years. Chemist and Manager, Prestwich U.D.C. Sewage Works for one and a half years. Chemist and under-Manager at the Thornhill Sewage Works of the County Borough of Dewsbury for ten years. Eight years student at the Dewsbury Technical School in Chemistry and Allied Subjects, and hold Board of Education certificates for all stages. I desire to join your Society so that I can keep in touch with modern methods of water, sewage and trades-waste purification, to obtain your publications and to have the use of your library. (*Signed by*) James H. Kershaw, Herbert J. Taylor, John Senior.

DUNN, LESLIE ASHBURN, 9, Pembroke Terrace, Bridlington. Assistant Works Chemist, employed by Messrs. Raines & Porter, Ltd., Hull. Was educated at Bridlington School (by scholarship). Studied Chemistry (inorganic, organic, and industrial) at Hull Technical College. Served with 1st E. Yorks. Regt. during the war. Finally assistant to T. A. Nightscapes, Esq., F.C.S. (*Signed by*) Thos. A. Nightscapes, Harry Thompson, W. Geary.

FISHER, RICHARD, Hudshaw House, Hexham, Northumberland. Student. During war I was employed in the Hydrophone Department of the Navy (Special Technical Branch). After the war I took up Medical Studies and have always wished for up-to-date chemical information. (*Signed by*) F. C. Garrett, C. R. Gent, John Gibson.

FREETH, FRANCIS ARTHUR, Heather Croft, Hartford. Chief Chemist, Messrs. Brunner, Mond & Co., Ltd. M.Sc. (Liv.), F.I.C., F.Inst.P., O.B.E. (*Signed by*) F. G. Donnan, W. E. Garner, Henry Terrey, Irvine Masson.

GAY, ALFRED DUNCAN, 49, Thornlaw Road, West Norwood, S.E.27. Analytical Chemist. Three years Junior Assistant to Dr. Bernard Dyer. Fifteen months Assistant Chemist to The Magadi Soda Co., Ltd., B.E.A. Recently engaged as Analytical Chemist to Schweppes, Ltd., Mineral Water Manufacturers. (*Signed by*) R. Lessing, Bernard Dyer, Wm. H. Woodcock, George Taylor.

HARRISON, DOUGLAS CREESE, 18, Church Crescent, Muswell Hill, London, N.10. Research Student in Organic Chemistry under Prof. Smiles at King's College, London. Passed Exam. for B.Sc. (London) with 1st Class Honours, with Chemistry as principal subject and Physics subsidiary. (*Signed by*) Samuel Smiles, A. J. Allmand, A. Fairbourne.

HARRISON, THOMAS WEATHERILL, 18, Elfindale Road, S.E.24. Chemist, Government Laboratory, London. B.Sc. (London). F.I.C. "Titration with Permanganate in Presence of Hydrochloric Acid." "The Action of Dimethyl Sulphate upon Oils of the Aromatic and Aliphatic Series." (*Signed by*) E. Grant Hooper, Geo. Stubbs, J. J. Fox.

HARROW, HENRY UNWIN, 13, Southwark Bridge Road, S.E.1. Senior Assistant Chemist, Barclay, Perkins' Brewery. Inter. B.Sc. (London), 1906. Chemist, South African Breweries, Johannesburg, from 1913 to 1920. Sitting for B.Sc. Hons. (London) in 1923 (October). (*Signed by*) J. Kenyon, J. L. White, S. Greenberg.

HENRY, DOUGLAS CECIL, M.A., Hulme Hall, Victoria Park, Manchester. Lecturer in Chemistry at the University of Manchester; Fellow of Trinity College, Cambridge; late Coutts-Trotter Scholar. Two years' research work, not at present published, on Adsorption from Solution, the results embodied in a thesis; won a Fellowship of Trinity College, Cambridge (Oct. 1921). Teaching experience—two years' coaching and tutorial work for Trinity, Cambridge. Lecturing and demonstrating at Manchester. (*Signed by*) Harold B. Dixon, Arthur Lapworth, F. P. Burt, J. E. Myers, Henry Stephen, Angus F. Core, Colin Campbell.

HODGSON, HERBERT, 35, Ryan Street, Bradford, Yorks. Analytical Chemist. Over twenty years assistant to Mr. F. W. Richardson, City Analyst of Bradford, and County Analyst of W.R. of Yorks. Also with the firm of Messrs. Richardson & Jaffé, Industrial Chemists. Experience in most branches of Analytical Chemistry. Also with R.E. "Anti-Gas," University College, London. (*Signed by*) F. W. Richardson, A. Jaffé, Geoffrey A. Bracewell.

HOLMES, EUSEBIUS, Eastern Green, Coventry. Student of Chemistry. Qualified for B.Sc. (Birm.). Desiring access to recent chemical literature. (*Signed by*) C. E. Wood, Wm. Wardlaw, F. Ashley Cooper.

HURST, WILLIAM, 2, Edditch Grove, Tonge, Bolton. Analytical and Manufacturing Chemist, c/o Thos. Moscrop & Co., Bolton, Oil Blenders and Refiners, Lubricants, etc. 1st Class Certificates, Lancs. and Cheshire Inst.: Mathematics 3rd year, and Chemistry, Inorganic and Organic, 4th year. Final Certificate of The City and Guilds Inst., London, in Oils, Fats, and Waxes. Two and a half years' Senior Chemistry Course (full time), Mining and Technical College, Wigan, 1919–1921. At present attending the College of Technology, Manchester. (*Signed by*) John Allan, Frank Lee Pyman, E. B. Naylor, T. H. Winstanley.

IREDALE, THOMAS, 94, Tyrwhitt Road, St. Johns, S.E.4. Scholar of the 1851 Exhibition. B.Sc. (Sydney, Australia), 1919. Have since been engaged in research. Papers published in Transactions of Chem. Soc. (1921). (*Signed by*) F. G. Donnan, I. W. Wark, E. J. Hartung, D. C. Jones.

JAMES, ETHEL MARY, 1, Rusham Road, Balham, S.W.12. Assistant Science Mistress, Clapham County Secondary School, S.W.11. B.Sc. (Lond.); Hons. Chem. 1920, after studying at Bedford College from 1916–20. At present teacher of chemistry at the above school. (*Signed by*) Holland Crompton, J. F. Spencer, M. S. Crewdson.

KENNABY, GEORGE LEWIS, Abadan, Persian Gulf. Chemist. B.Sc.

(London), 3rd Class Honours in Chemistry, 1920, since which I have been working for the Anglo-Persian Oil Co. at their refinery in Persia. I am desirous of keeping in touch with the recent developments in chemistry with the object of taking my A.I.C. at some future date. (*Signed by*) F. G. Donnan, W. E. Garner, Irvine Masson, Henry Terrey.

LAWSON, EDWARD INGLESON, 65, Warrington Road, Newcastle-on-Tyne. Teacher of Chemistry at Dame Allan's School, Newcastle-on-Tyne. Student of Chemistry for four years at Leeds University. Teacher of Chemistry for four years at Newcastle. (*Signed by*) J. R. Scott, R. M. Woodman, John Stanley Carter, Edgar Rhodes.

LUNN, ARTHUR CYRIC PERCY, King's College, Cambridge. Student. Part I Nat. Science Tripos, 1921. Taking Part II (Chemistry, May 1922). (*Signed by*) Hamilton McCombie, Harold A. Scarborough, A. E. Cashmore, C. T. Heycock, W. H. Mills, Douglas F. Harrison.

MCDOWALL, ROBERT CRAIG, 6, Ascog Street, Crosshill, Glasgow. Analytical Chemist to Messrs. Perry & Hope, Ltd., Forth & Clyde Chemical Works, Nitshill. Cream of Tartar Subs., Malt Extract, Food Products generally 1904-1911 (7 years). Technical and Chemical Adviser to Messrs. Craig, McDowall & Co., Thornliebank, Glasgow, Cream of Tartar Subs., Baking Powders, and other Food Products, Mag. Sulph., Ferro Sulph., Soda Sulph., etc., 1911-1915 (4½ years). Glasgow Highlanders, H.M. Forces, 1915-1918 (3½ years). Technical Chemist and Representative, also Process Superintendent to Messrs. Perry & Hope, Ltd., Forth & Clyde Chemical Works, Nitshill, 1918-1922 (4 years). Having been connected for the past sixteen years with the manufacture and analysis of foodstuffs I am desirous of keeping in touch with all latest books and papers appertaining to same. (*Signed by*) James Booth, J. Turnbull, Herbert Lorenz, W. McHutchison.

MANJUNATH, LINGAIAH BASAVALINGAIAH, London. Student of the London University. B.A. degree in Chemistry of the Madras University. At present a student for the London M.Sc. degree in Chemistry in the University College. (*Signed by*) O. L. Brady, E. H. Thierry, J. N. E. Day.

MULLALY, JOHN MYLNE, 14, Bardwell Road, Oxford. Student of Chemistry. Part author of paper *Trans. Chem. Soc. CXIX*, p. 1802 (1921). Commoner of Balliol College. (*Signed by*) H. Hartley, C. N. Hinshelwood, E. J. Bowen, H. R. Raikes.

POWERS, HAROLD EDWARD CHARLES, 4, Grove Crescent, Woodford, E.18. Analytical Chemist. B.Sc. Hons. (2nd Class). About six years' practical experience in technical laboratory. (*Signed by*) Hugh Main, George Senter, F. Barrow, S. Sugden, Alfred J. White.

RAO, AKELLA SURYANARAYANA (Junior), Vizagapatam, India. B.Sc. in Chemistry, Botany (National University, Adyar, Madras). I intend joining the Society fully believing that by doing so I shall be in the know of the recent work in chemistry, which would be of immense use to me as a prospective research chemist. (*Signed by*) T. S. Natrajan, S. N. Dhar, K. C. Viraraghavan.

RILEY, HARRY LISTER, 7, Trafalgar Square, Chelsea, S.W.3. Student (Research) at the Royal College of Science. A.R.C.Sc. and B.Sc. (Hons.) London. (*Signed by*) H. B. Baker, James C. Philip, J. N. Sugden, H. J. T. Ellingham, A. A. Eldridge.

ROBINSON, JAMES, Bulawayo, S. Rhodesia. Science Master, Milton High School, Bulawayo. B.Sc., Univ. of Wales. Completed 1st year London Final Sc. Course in Mechanical Engineering (Matriculated 1st Div. Inter. Sc., London). Completed advanced practical courses on Fibres, Detergents,

Alkaloids, Assays. One session's research work on "The Action of SO_2 , Sulphites, Meta-bisulphate on various Peroxides (BaO_2 , H_2O_2 , CeO_2 , PbO_2) in Comparison with MnO_2 and with the action on $\text{K}_2\text{Mn}_2\text{O}_8$, CrO_3 , and Per-sulphates as Oxidising Agents," under Prof. Herbert Jackson, King's College, London. (*Signed by*) Geo. A. Pingstone, Arthur Slator, A. L. Stern.

RUSSELL, GEORGE ALLAN, Barnsheen, Ditton, Widnes. Chemist to Messrs. T. Vickers & Sons, Ltd., Phosphate Works, Widnes. B.Sc. (Dun.). A.I.C. Five years' laboratory and works experience with the United Alkali Co., Ltd., Widnes, in the manufacture of monochlorobenzene, chloroacetic acid, and concentration of sulphuric acid, and subsequently two and a half years' (to date) experience with Messrs. T. Vickers & Sons, Ltd., Widnes, in the manufacture of sulphuric acid, Fertilisers and Phosphate of Soda, and in the corresponding analytical work. (*Signed by*) F. C. Garrett, J. A. Smythe, P. L. Robinson, John W. Towers, *John Hargreaves*.

SASTRY, M. SUHRAMANIA, 5, Krishnappa Naik Aghraharam, Georgetown, Madras, India. Chemical Assistant, Government Leather Trades Institute, Madras. B.A. (Madras). Post Graduate Course for two and a half years (Madras University). Chemical Assistant. (*Signed by*) Kalyan C. Srinivasan, T. S. Natrajan, K. C. Viraraghavan, *S. N. Dhar*.

SEELEY, ERNEST ARTHUR, 10, Hamilton Road, London, N.5. Chemical Research Student. B.Sc. 1st Class Hons. Chem. (Lond.), A.R.C.S. (1st Chem.). D.I.C. (Organic Chemistry Research). (*Signed by*) Jocelyn Thorpe, C. K. Ingold, L. C. Nickolls.

SHAW, BRIAN DUNCAN, Ferns' Hollow, Station Road, Ilkeston. Student. Second Year Hon. Chem. Student (for B.Sc. Lond.) at University College, Nottingham. (*Signed by*) F. S. Kipping, F. C. Laxton, G. Green.

SMITH, ALBERT W., Case School of Applied Science, Cleveland, Ohio, U.S.A. Professor of Chemistry. Graduate of Chemical Dept. of Case School of Applied Science (B.S.). University of Zürich, Switzerland (Ph.D.). (*Signed by*) Olin F. Tower, Chas. F. Guhlmann, Chas. H. Herty.

STOTT, CLARENCE, Benroyd, Holywell Green, nr. Halifax. Works Chemist. (a) B.Sc. (Leeds); (b) A.I.C.; (c) Diploma in Gas Engineering (Leeds); (d) War Service: Army, $4\frac{1}{2}$ years; B.E.F. France, $3\frac{1}{2}$ years; rank: Capt. The last two years of service overseas were spent in Chemical Warfare as Gas Officer to the 49th (West Riding) Division. Honours: (1) Mentioned in Despatches; (2) Military Cross. (e) Works experience: $1\frac{1}{2}$ years' laboratory and plant experience with the Farnley Iron Co., nr. Leeds. (*Signed by*) John W. Cobb, Arthur Smithells, Herbert Ingle.

THOMAS, WILLIAM, Penfro, Stoneywood, Aberdeen. Lecturer in Chemistry, University, Aberdeen. M.Sc. (Wales). B.A. (Cantab.). A.I.C. (*Signed by*) Alex. Findlay, T. Harold Reade, Edward Bloom.

UNDERHAY, GEORGE FLAXMAN, Allerton, Woodside Avenue, Muswell Hill, N.10. Research at University College, London. 2nd Class Honours B.Sc. (Chemistry). Now engaged on research under Prof. F. G. Donnan, C.B.E., F.R.S., at University College. (*Signed by*) F. G. Donnan, J. N. Collie, Irvine Masson, O. L. Brady.

VARDY, WILLIAM ANDREW SHAKESPEARE, Winksley Vicarage, Ripon. Analytical Chemist, relinquished on account of ill-health. Late Student in Science and Member of Ripon G.S. Scientific Society. Analytical Chemist. (*Signed by*) A. Mortimer, Barker North, T. W. Price.

WALTERS, JOHN GEORGE, 46, Durham Road, East Finchley, N.2. Teacher. B.Sc. (London). Senior Chemistry Master, County Secondary School, Hendon, N.W.4. Late Commandant XVII Corps Gas School, B.E.F., and Chemical

Adviser XVII Corps, B.E.F. (*Signed by*) T. J. Drakeley, P. Schidrowitz, Raymond R. Butler.

WATT, WILLIAM WALLACE, 29, Woodstock Road, Golders Green, N.W.4. Consulting Petroleum Technologist. Chief Chemist, Burmah Oil Co., Rangoon (13½ years). Technical Staff (London), Anglo-Persian Oil Co. (1½ years). Deputy Controller of Oil to H.M. Govt. of India (1 year). Technical Adviser Caltex Oil Co., London (1½ years). (*Signed by*) F. B. Thole, A. E. Dunstan, J. C. Crocker, F. Arnall.

WATSON, ARTHUR FREDERICK, Brockwell Hall, Herne Hill, S.E.23. Research Chemist. University College, London, 1913-15, 1919-21; B.Sc. (War) degree, 1917. B.Sc. (Honours Chemistry), 1920. A.I.C. 1921. Research in Biological Chemistry, 1920-1921. M.Sc. (Physiological Chemistry), 1921, London. (*Signed by*) J. C. Drummond, Irvine Masson, F. G. Donnan.

WILLIAMS, LLYWELYN, Herne Bay College, Kent. Assistant Master. B.Sc. (Wales) in Chemistry and Physics. My present appointment is that of Senior Science and Mathematics Master at Herne Bay College, Kent. (*Signed by*) T. Campbell James, C. R. Bury, L. J. Hudleston.

WOOD, HARRY ALEXANDER, Santa Elena, Entre Rios, Argentine Republic. Industrial Chemist. (a) Analytical Assistant to Prof. James Hendrick, B.Sc., F.I.C., Professor of Agriculture, Aberdeen University (3 years); (b) Analytical Assistant to Dr. J. F. Tocher, F.I.C., Public Analyst and Official Agricultural Analyst (3 years); (c) Works Chemist: Establecimientos Argentinos de Bovril Santa Elena, Entre Rios, Argentine Republic (1½ years). (*Signed by*) James Hendrick, J. F. Tocher, A. Searl.

The following Certificates have been authorised by the Council for presentation to ballot under Bye-law I (2) :

BASU, HARILAL, Care of the Inspector-General of Civil Hospitals, Burma, Rangoon, Post Box No. 84. Civil Surgeon and District Public Health Officer of Yamethin District and Superintendent of Yamethin District Jail (Prison). L.M. & S. (Calcutta University). A good and successful student in the chemistry classes and always takes special interest in practical chemical work. Registered Medical Practitioner of the British Medical Council (London). Attended pulp making from bamboo for making of paper. Wants to do research work in chemistry. (*Signed by*) L. B. Seal, M. Hunter.

LAWRANCE, WALTER ALBERT, Lewiston, Maine, U.S.A. Professor of Chemistry, Bates College. Degrees : B.A., M.A. (McMaster); M.A., Ph.D. (Toronto). "Poly-nitro-compounds," *Proc. Royal Soc. Can.*, May 1918. "Friedel-Crafts' Reaction," *J. Am. Chem. Soc.*, Sept. 1920. "Friedel-Crafts' Reaction," *J. Am. Chem. Soc.*, Dec. 1921. "Friedel-Crafts' Reaction," *J. Am. Chem. Soc.* (in course of publication). (*Signed by*) W. Lash Miller.

P R O C E E D I N G S

O F T H E

C H E M I C A L S O C I E T Y .

Ordinary Scientific Meeting, Thursday, April 6th, 1922, at 8 p.m., PROFESSOR H. E. ARMSTRONG, LL.D., F.R.S., Vice-President, in the Chair.

Reference was made to the loss sustained by the Society, through death, of :

	Elected.	Died.
Philippe Auguste Guye (Honorary Fellow)	Mar. 7th, 1912.	Mar. 27th, 1922.
Seward Whiting Williams	June 18th, 1891.	Aug. 25th, 1921.

It was announced—

(1) That in order to commemorate the centenary of the birth of Pasteur the University and town of Strasbourg have decided to erect a statue facing the University where, as a professor, Pasteur commenced his career. The inauguration ceremonies, under the patronage of the President of the Republic and other distinguished men, will be held on May 1st, 1923, and will include the unveiling of the statue and the opening of an Exhibition of Hygiene and Bacteriology.

With the object of evincing the sympathy of this country with the project in view, a British Committee has been formed of the following : Chairman, Sir Charles Sherrington, G.B.E. (President of the Royal Society), Treasurer, Mr. A. Chaston Chapman, F.R.S. (President of the Institute of Chemistry), Mr. H. E. Field (President of the Institute of Brewing), Professor Percy F. Frankland, C.B.E., F.R.S., Sir John M'Fadyean, M.R.C.V.S., LL.D. (Principal of the Royal Veterinary College), Professor C. J. Martin, C.M.G., F.R.S. (Director of the Lister Institute), Sir William J. Pope, K.B.E., F.R.S., Sir James Walker, F.R.S. (President of the Chemical Society), and Sir Almroth Wright, K.B.E., F.R.S. (Principal of the Institute of Pathology and Research).

It is hoped there may be many British chemists desirous of contributing to the Memorial Fund. Contributions may be sent

to Monsieur Th. Héring, 6, Rue des Veaux, Strasbourg, or to Mr. A. Chaston Chapman, Institute of Chemistry, 30, Russell Square, W.C. 1. The subscription list will close about the end of next June.

Manufacturers and business firms in this country to whom the Exhibition may be of interest are cordially invited to offer assistance, and those able and willing to do so should communicate with Professor Borrel, 3, Rue Koeberlé, Strasbourg.

(2) That the following Committees for 1922—1923 had been appointed by the Council :

Finance Committee.—Messrs. E. F. Armstrong, A. Chaston Chapman, C. A. Hill, G. T. Moody, and the Officers.

House Committee.—Messrs. J. L. Baker, Sir Herbert Jackson, Alexander Scott, Sir William A. Tilden, E. W. Voelker, and the Officers.

Library Committee.—Messrs. P. P. Bedson, O. L. Brady, Sir Thomas K. Rose, E. J. Russell, J. M. Thomson (Chairman), Sir William A. Tilden, the Editor, the Librarian, and Officers, together with the following representatives of contributing Societies : Messrs. J. L. Baker, C. Dorée, J. C. Drummond, A. Harden, C. A. Keane, R. L. Mond, and G. Senter.

Publication Committee.—Messrs. A. J. Allmand, O. L. Brady, C. H. Desch, I. M. Heilbron, J. T. Hewitt, J. C. Irvine, H. King, T. M. Lowry, J. W. McBain, R. H. Pickard, N. V. Sidgwick, J. F. Thorpe, and the Officers.

Research Fund Committee.—Messrs. H. B. Baker, D. L. Chapman, Sir James J. Dobbie, F. G. Donnan, G. G. Henderson, F. S. Kipping, R. H. Pickard, Sir William J. Pope, R. Robinson, S. Smiles, and the Officers.

The following were formally admitted Fellows of the Chemical Society : H. B. Courtney and G. Thompson.

Certificates were read for the first time in favour of :

George Baborovský, Ph.D., Falkensteinerova ul. 7, Brno, Czecho-Slovakia.

Stanley John Bailey, B.A., The Braes, South Nutfield, Redhill.

Robert Stanley Colborne, B.Sc., A.R.C.S., 57, County Road, Swindon.

James Brierley Firth, D.Sc., F.I.C., 44, Lucknow Avenue, Mapperley Park, Nottingham.

Dorothea Annie Hoffert, 50, Hampstead Way, N.W. 11.

Ralph Henry Parker, Radio House, Wilson Road, Smethwick, Staffs.

Edmond Samuels, 40, Castlereagh Street, Sydney, N.S.W.

Ronald William Stevenson, M.C., M.A., Eaglehurst, Ditton Hill, Surbiton.

A Certificate has been authorised by the Council for presentation to ballot under Bye-Law I (2) in favour of :

Robert Swainson-Hall, Government Plantations Department, Victoria, British Cameroons, West Africa.

The following papers were read :

- " Constitution of picrorocellin, a nitrogenous constituent of *Roccella fuciformis*." By M. O. FORSTER and W. B. SAVILLE.
 - " The determination of surface tension from the maximum pressure in bubbles." By S. SUGDEN.
 - " The variation of refractive index and specific gravity of benzene with temperature." By W. B. PARKER and G. THOMPSON.
 - " The influence of substituents on the formation and stability of heterocyclic compounds. Part I. Hydantoins." By C. K. INGOLD, S. SAKO, and J. F. THORPE.
-

ORDINARY SCIENTIFIC MEETING.

The next Ordinary Scientific Meeting will be held on Thursday, May 4th, 1922, at 8 p.m., when a ballot for the election of Fellows will be held.

INFORMAL MEETING.

The next Informal Meeting will be held on Thursday, May 18th, 1922, after the conclusion of the formal business of the Ordinary Scientific Meeting, to be held at 8 p.m.

RESEARCH FUND COMMITTEE.

A meeting of the Research Fund Committee will be held in June next. Applications for Grants, to be made on forms which can be obtained from the Assistant Secretary, Chemical Society, Burlington House, W. 1., must be received on or before Thursday, June 1st, 1922.

All persons who receive grants in June, 1921, or in June of any previous year, whose accounts have not been declared closed by the Council, are reminded that reports must be returned by Thursday, June 1st.

SETS OF THE *BERICHTE* AND OF THE *ZENTRALBLATT*.

The Society still has a few remaining sets of the *Berichte* and of the *Zentralblatt* from August, 1914, to December, 1920, inclusive, for disposal. These can be obtained, in complete sets only, from the Assistant Secretary, price £10 per set of the *Berichte*, and of the *Zentralblatt*.

The number of sets is strictly limited and application should be made without delay.

LIST OF FELLOWS, 1922.

The Council has decided to supply the List of Fellows for 1922 to those Fellows only who make application to the Assistant Secretary before August 31st, 1922.

ADDITIONS TO THE LIBRARY.

I. *Donations.*

CORNUBERT, R. Dictionnaire Anglais-Français-Allemand : de mots et locutions intéressant la physique et la chimie. Paris 1922. pp. xxxii + 298. 42 fr. (*Reference.*)

From the Publisher : M. Dunod.

HALE, ARTHUR JAMES. Modern chemistry, pure and applied. Vol. IV. London [1922]. pp. viii + 280. ill. 16s. net. (*Recd.* 31/3/22.)

From the Publishers : Messrs. Virtue & Co.

IMPERIAL INSTITUTE. Indian Trade Enquiry. Reports on lac, turpentine, and rosin. London 1922. pp. x + 74. 5s. net. (*Recd.* 1/4/22.)

From the Director.

IMPERIAL MINERAL RESOURCES BUREAU. Laws and regulations relating to lead poisoning : being an analysis with texts of the laws and regulations made in the chief industrial countries to prevent plumbism. By GILBERT STONE. London 1922. pp. 247. (Two copies.) 5s. net.

From the Bureau.

INDIAN SCIENCE CONGRESS. Handbook for the use of members attending the ninth meeting to be held at Madras . . . 1922. Madras 1921. pp. x + 166. (*Recd.* 21/3/22.)

From Captain Clive Newcomb.

LOWRY, THOMAS MARTIN. Inorganic chemistry. London 1922. pp. xii + 944. ill. 28s. net. (*Recd.* 30/3/22.)

From the Author.

MINISTRY OF MUNITIONS and DEPARTMENT OF SCIENTIFIC AND INDUSTRIAL RESEARCH. Technical records of explosives supply 1915-1918. No. 8. Solvent recovery. London 1921. pp. iv + 22. ill. 3s. net. (*Recd.* 10/4/22.)

From the Department.

MONIER-WILLIAMS, GORDON WICKHAM. Power alcohol : its production and utilisation. London [1922]. pp. xii + 324. ill. 21s. net. (*Recd.* 10/4/22.)

From the Publishers : Messrs. Henry Frowde and
Hodder & Stoughton.

SCIENTIFIC AND INDUSTRIAL RESEARCH, Department of. *Fuel Research Board*. A handbook on the winning and utilisation of peat. By A. HAUSDING. Translated from the 3rd German edition by HUGH RYAN. London 1921. pp. xxiv + 506. ill. 30s. net. (*Recd.* 16/3/22.) From the Department.

SHERMAN, HENRY CLAPP, and SMITH, SYBIL L. The vitamins. New York 1922. pp. iv + 274. ill. \$4 net. (*Recd.* 21/3/22.)

From the Publishers: The Chemical Catalogue Company, Inc.

VENABLE, FRANCIS P. Zirconium and its compounds. New York 1922. pp. 1-10, 15-174. \$2.50 net. (*Recd.* 30/3/22.)

From the Publishers: The Chemical Catalogue Company, Inc.

II. By Purchase.

ASTON, FRANCIS WILLIAM. Isotopes. London 1922. pp. viii + 152. ill. 9s. net. (*Recd.* 14/3/22.)

BODFORSS, SVEN. Die Aethylenoxyde. Ihre Darstellung und Eigenschaften. (Sammlung, Vol. XXVI.) Stuttgart 1920.

BÖCKMANN, FRIEDRICH. Celluloid: its raw material, manufacture, properties and uses. 2nd English edition, translated from the 3rd German edition by HERBERT BIRTWHISTLE STOCKS. London 1921. pp. viii + 188. ill. 8s. 6d. net. (*Recd.* 16/3/22.)

CHAPMAN, ALFRED CHASTON. Einige Fortschritte auf dem Gebiete der analytischen Chemie. Translated by OSKAR BAUDISCH. (Sammlung, Vol. XXVI.) Stuttgart 1921.

DONATH, EDUARD, and LISSNER, A. Kohle und Erdöl. (Sammlung, Vol. XXVI.) Stuttgart 1920.

FUNK, CASIMIR. The vitamins. Translated from the 2nd German edition by HARRY E. DUBIN. Baltimore 1922. pp. 502. ill. 31s. 6d. net. (*Recd.* 9/3/22.)

GROSSMANN, HERMANN, and WRESCHNER, MARIE. Die anomale Rotationsdispersion. (Sammlung, Vol. XXVI.) Stuttgart 1921.

HAHN, FRIEDRICH-VINCENZ VON. Ueber die Herstellung und Stabilität kolloider Lösungen anorganischer Stoffe. (Mit besonderer Berücksichtigung der Sulfidsole.) (Sammlung, Vol. XXVI.) Stuttgart 1922.

HATSCHKE, EMIL. An introduction to the physics and chemistry of colloids. 4th edition. London 1922. pp. xiv + 172. ill. 7s. 6d. net. (*Recd.* 16/3/22.)

HURST, GEORGE H. Soaps: a practical manual on the manufacture of domestic, toilet, and other soaps. 3rd edition. By HERBERT BIRTWHISTLE STOCKS. London 1922. pp. viii + 432. ill. 21s. net. (*Recd.* 14/3/22.)

JONES, LLEWELLYN, and SCARD, FREDERIC ISENBART. The manufacture of cane sugar. 2nd edition. London [1921]. pp. xx + 482. ill. 25s. net. (*Recd.* 16/3/22.)

JULIAN, H. FORBES, and SMART, EDGAR. Cyaniding gold and silver ores. 3rd edition. By A. W. ALLEN. London 1921. pp. xxiv + 418. ill. 36s. net. (*Recd.* 16/3/22.)

MEADE, ALWYNE. Modern gasworks practice. 2nd edition. London 1921. pp. xii + 816. 55s. net. (*Recd.* 14/3/22.)

MELLOR, JOSEPH WILLIAM. A comprehensive treatise on inorganic and theoretical chemistry. Vols. I and II. London 1922. pp. xvi + 1066, viii + 894. ill. (*Reference.*) £6 6s. net.

NEUBURGER, MAXIMILIAN CAMILLO. Das Problem der Genesis des Actiniums. (Sammlung, Vol. XXVI.) Stuttgart 1921.

— Neuere Ergebnisse der Forschung über die Radioaktivität des Kaliums und Rubidiums im letzten Dezennium. (Sammlung, Vol. XXVI.) Stuttgart 1921.

PFEIFFER, PAUL. Organische Molekülverbindungen. (Chemie in Einzeldarstellungen, Vol. XI.) Stuttgart 1922. pp. xiv + 328. (*Recd.* 14/3/22.)

REDWOOD, SIR BOVERTON. Petroleum. 3 vols. 4th edition. London 1922. pp. xxx + 1353. ill. £5 5s. net. (*Recd.* 10/3/22.)

SIMMONS, WILLIAM HERBERT, and MITCHELL, CHARLES AINSWORTH. Edible fats and oils: their composition, manufacture, and analysis. 2nd edition. London 1921. pp. viii + 190. ill. 10s. 6d. net. (*Recd.* 16/3/22.)

YOUNG, SYDNEY, and others. Distillation principles and processes. London 1922. pp. xiv + 510. ill. 40s. net. (*Recd.* 14/3/22.)

ZEITSCHRIFT FÜR LEDER- UND GERBEREICHIMIE. Vol. I., etc. Hamburg 1922 +. ill. (*Reference.*)

III. Pamphlets.

ALEXANDER, ELIZABETH H. A bibliography of John Ferguson. (From the *Trans. Glasgow Bibliographical Soc.*, with alterations.) 1920. ill.

IMPERIAL MINERAL RESOURCES BUREAU. The Mineral Industry of the British Empire and Foreign Countries. War Period. Lead. (1913-1919.) London 1922. pp. 95.

— — Platinum and allied metals. (1913-1919.) London 1922. pp. 84.

— — Tin. (1913-1919.) London 1922. pp. 98.

INSTITUTE OF CHEMISTRY OF GREAT BRITAIN AND IRELAND. Lecture on modern applications of chemistry to crop production. By EDWARD JOHN RUSSELL. London 1922. pp. 23. ill.

MIALL, STEPHEN. The structure of the atom : notes on some recent theories. London 1922. pp. vi + 26. ill.

TADOKORO, TETSUTARO. On the spectrochemical reaction of methylfurfurol- and oxymethylfurfurol phloroglucid. (From the *J. Coll. Agric., Hokkaido Imp. Univ.*, 1921, 10.) ill.

—— Ultraviolet spectroscopical studies on the antagonistic action of salts in organic colloidal solution. (From the *J. Coll. Agric., Hokkaido Imp. Univ.*, 1921, 10.) ill.

P R O C E E D I N G S

O F T H E

C H E M I C A L S O C I E T Y .

Ordinary Scientific Meeting, Thursday, May 4th, 1922, at 8 p.m.,
Professor J. F. THORPE, C.B.E., F.R.S., Vice-President, in the
Chair.

The CHAIRMAN referred to the loss sustained by the Society,
through death, of the following Fellows :

	Elected.	Died.
Sir William Phipson Beale	Dec. 5th, 1867.	April 13th.
Joseph Albert Hughes	Dec. 3rd, 1908.	Jan. 9th.
Charles Mann Luxmoore	Feb. 16th, 1893.	April 16th.
James Morris	May 1st, 1919.	Mar. 31st.

The CHAIRMAN announced :

1. That the Council had sealed the following Address of
Congratulation to be presented to the Royal University of Padua
on the celebration of the Seven Hundredth Anniversary of its
Foundation to be held on May 14th to 17th, 1922 :

T H E C H E M I C A L S O C I E T Y

T O

T H E R E C T O R , S E N A T E , P R O F E S S O R S A N D A L U M N I

O F T H E

R O Y A L U N I V E R S I T Y O F P A D U A .

On behalf of the Chemical Society, we have the honour to
convey our warmest felicitations on the celebration of the
Seven Hundredth Foundation Anniversary of the Royal
University of Padua.

The knowledge that your University flourished in the 13th
century, that six hundred years ago it pointed the way of
learning to the peoples—not only of Italy, but of Europe—
and that inscribed on the roll of its learned professors in

science are to be found the immortal names of Galileo and Morgagni, is indeed an inspiration to the whole world. It is our earnest wish that the lamp of learning, kindled so long ago in your ancient University, may continue to burn with increasing radiance.

In the name of the Fellows and Council of the Chemical Society we welcome this happy opportunity of recording our deep appreciation of the commanding position achieved by your University in the advancement of knowledge, uniting therewith our most sincere wishes for the continued prosperity of this illustrious and venerable seat of learning.

Signed on behalf of the Council of
The Chemical Society.

JAMES WALKER, President.

M. O. FORSTER, Treasurer.

IRVINE MASSON, } Secretaries.
JAMES C. PHILIP, }

ARTHUR W. CROSSLEY, Foreign Secretary.

(L.S.)

Sealed in Council this Fourth day of May, One Thousand Nine Hundred and Twenty-Two.

2. That Mr. G. H. Gabb had presented to the Society's collection a medal of Joseph Priestley, executed by Halliday about 1804. The inscription on the reverse of the medal is taken from the tablet erected in the new Meeting House, Birmingham, to the memory of Priestley and was written by Dr. Parr.

The following were formally admitted Fellows of the Chemical Society : P. Appleyard, K. A. Burke, M. H. Cocks, K. J. Isaac, L. O. Kekwick, and H. L. Saunders.

Certificates were read for the first time in favour of :

Frank Belton, B.Sc., Athlone, Bush Hill Park, Enfield.

Robert Keith Cannan, M.Sc., 4, Antrim Mansions, N.W. 3.

Bibhu Charan Chatterjee, M.Sc., University College of Science, 92, Upper Circular Road, Calcutta, India.

Shiavaksha Hormasji, Contractor, Karanjia Building, opp. Moos Building, Grant Road, Bombay, India.

Margaret White Fishenden, D.Sc., 19, Shepherd's Hill, Highgate, N. 6.

Frédéric Walter Goddard, M.A., 12, College Street, Winchester.

Edmund Langley Hirst, M.A., B.Sc., Ph.D., Ainslie Villa, Largo Road, St. Andrews.

William Job Jenkins, 37, Marston Road, Knowle, Bristol.

Peppo Joseph Levy, 42, Granville Gardens, Shepherd's Bush, W. 12.

David McCall, B.Sc., 4, Lime Street, Dundee.

Richard Edmund Anthony Mallet, B.A., Redford, Ockley, Surrey.

Ronald George Wreyford Norrish, B.A., Emmanuel College, Cambridge.
 John Goodrich Oats, 3, Montpelier Terrace, Ilfracombe.
 Claude Bernard Meister Platt, M.B.E., Middleton House, Beckett Road,
 Doncaster.
 Arthur Plowman, B.Sc., A.R.C.S., A.I.C., 8, Rowantree Road, Bycullah
 Park, Enfield.
 John Pryde, B.Sc., 4, Gowrie Street, Blackness Avenue, Dundee.
 Juda Hirsch Quastel, B.Sc., A.R.C.S., A.I.C., Trinity College, Cambridge.
 John Charles Smith, B.Sc., 31, Bidwell Street, Wellington, N.Z.
 Valerie Emeline Sophie Swainson, B.Sc., 21, Hungerford Road, N. 7.
 Nowroji Jamshedji Vazifdar, 17, Queens Road, opp. Charin Road Station,
 Bombay, India.
 George William Wigg, 72nd Northumbrian Bde., R.F.A., Barrack Road,
 Newcastle-on-Tyne.

Messrs. C. Gilling and J. Reilly were elected Scrutators, and a
 ballot for the election of Fellows was held. The following were
 subsequently declared elected as Fellows :

Brojo Kishore Banerjee, B.Sc.	Walter Albert Lawrance, M.A., Ph.D.
Benoy Bhusan Basu, B.Sc., M.B.	Edward Ingleson Lawson.
Harilal Basu.	Arthur Cyrie Percy Lunn, B.A.
George Macdonald Bennett, M.A., M.Sc., F.I.C.	Robert Craig McDowall.
John William Blagden, M.A., Ph.D., F.I.C.	Lingaiah Basavalingaiah Manju- nath, B.A.
Frederick Mason Brewer.	John Mylne Mullaly.
Eric Sykes Brooks.	Harold Edward Charles Powers, B.Sc.
Fred Brumfitt.	Akella Suryanarayana Rao, B.Sc.
Carlton Butler, B.Sc., A.I.C.	Harry Lister Riley, B.Sc.
Charlie Lyons Carter, M.Sc., A.I.C.	James Robinson, B.Sc.
Arthur Rockett Chadwick.	George Allan Russell, B.Sc., A.I.C.
Leslie Ashburn Dunn.	M. Suhramania Sastry, B.A.
Richard Fisher.	Ernest Arthur Seeley, B.Sc.
Francis Arthur Freeth, M.Sc., F.I.C.	Brian Duncan Shaw.
Alfred Duncan Gay.	Albert William Smith, B.Sc., Ph.D.
Douglas Creese Harrison, B.Sc.	Clarence Stott, B.Sc., A.I.C.
Thomas Weatherill Harrison, B.Sc., F.I.C.	William Thomas, B.A., M.Sc., A.I.C.
Henry Unwin Harrow.	George Flaxman Underhay, B.Sc.
Douglas Cecil Henry, M.A.	William Andrew Shakespeare Vardy.
Herbert Hodgson.	John George Walters, B.Sc.
Eusebius Holmes, B.Sc.	Arthur Frederick Watson, M.Sc., A.I.C.
William Hurst.	William Wallace Watt.
Thomas Iredale, B.Sc.	Llywelyn Williams, B.Sc.
Ethel Mary James, B.Sc.	Harry Alexander Wood.
George Lewis Kennaby, B.Sc.	

The following papers were read :

- "On dopplerite. Studies in the composition of coal." By F. V.
 TIDESWELL and R. V. WHEELER.
 "Bromo-derivatives of glyoxaline." By I. E. BALABAN and
 F. L. PYMAN.

- "The properties of ammonium nitrate. Part IV. The reciprocal salt-pair $\text{NH}_4\text{NO}_3 + \text{NaCl} \rightleftharpoons \text{NH}_4\text{Cl} + \text{NaNO}_3$." By E. P. PERMAN.
- "The reactivity of ammonia." By E. C. C. BALY and H. M. DUNCAN.
- "Photocatalysis. Part II. The photosynthesis of nitrogen compounds from nitrates and carbon dioxide." By E. C. C. BALY, I. M. HEILBRON, and D. P. HUDSON.

Ordinary Scientific Meeting, Thursday, May 18th, 1922, at 8 p.m., SIR JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

Reference was made to the loss sustained by the Society, through death, of :

	Elected.	Died.
Adolphus Collenette	March 1st, 1883.	May 7th, 1922.

The following were formally admitted Fellows of the Chemical Society : G. M. Bennett and J. G. Mitchell.

Certificates were read for the first time in favour of :

Harry Baines, B.Sc., 22, Henry Road, West Bridgford, Nottingham.
 Victor Cecil Branson, B.Sc., A.R.C.S., A.I.C., Messrs. Naparima Oilfields, Ltd., San Fernando, Trinidad, B.W.I.
 Albert Hassell, 47, Stanley Street, Tunstall, Stoke-on-Trent.
 Edith Annie Peach, 43, St. Gabriel's Road, Cricklewood, N.W.2.
 Clifford Dane Thomas, B.Sc., 10, Lansdowne Place, Blackheath, S.E.10.

The following Certificate has been authorised by the Council for presentation to ballot under Bye-law I (2) :

Walter Herbert Fulweiler, B.Sc., c/o The United Gas Improvement Co., Philadelphia, Pa., U.S.A.

The meeting was then adjourned and the Informal Meeting declared open. A selection of autograph letters from the Roscoe collection and of books of historical interest from the Library was exhibited.

LECTURE.

At the Ordinary Scientific Meeting to be held on Thursday, June 8th, 1922, at 8 p.m., in the Lecture Hall of the Institution of Mechanical Engineers, Storey's Gate, S.W.1, Dr. H. H. Dale, C.B.E., F.R.S., will deliver his Lecture entitled : "Chemical and Physiological Properties."

LIST OF FELLOWS, 1922.

The Council has decided to supply the List of Fellows for 1922 to those Fellows only who make application to the Assistant Secretary before August 31st, 1922.

List of Papers received between April 7th and May 18th, 1922 :

- " A synthesis of *isobrazilein* and certain related anhydropyranol salts. Part II. Synthesis of *isohæmatein*." By H. G. CRABTREE and R. ROBINSON.
- " Studies in catalysis. Part XVI. The inversion of sucrose by hydrogen ion." By T. MORAN and W. C. MCC. LEWIS.
- " 2-*p*-Dimethylaminostyrylpyridine methiodide, a new photographic sensitiser." By W. H. MILLS and W. J. POPE.
- " On dopplerite. Studies in the composition of coal." By F. V. TIDESWELL and R. V. WHEELER.
- " Bromo-derivatives of glyoxaline." By I. E. BALABAN and F. L. PYMAN.
- " The oxidation of hexadecane and saturated fatty acids by air or oxygen." By A. H. SALWAY and P. N. WILLIAMS.
- " Estimation of lead peroxide." By C. RAMASOMAYAJULU.
- " Dyes derived from phenanthraquinone (naphthaflavindulines)." By S. C. DATTA.
- " Researches on sulphuryl chloride. Part II. A new chlorinating agent; preparation of poly-chloro-derivatives of benzene." By O. SILBERRAD.
- " The decomposition of benzyl disulphoxide." By J. A. SMYTHE.
- " The use of iron pyrites in a Friedel-Crafts' reaction." By J. A. SMYTHE.
- " The properties of ammonium nitrate. Part IV. The reciprocal salt-pair, $\text{NH}_4\text{NO}_3 + \text{NaCl} \rightleftharpoons \text{NH}_4\text{Cl} + \text{NaNO}_3$." By E. P. PERMAN.
- " The reactivity of ammonia." By E. C. C. BALY and H. M. DUNCAN.
- " Photocatalysis. Part II. The photosynthesis of nitrogen compounds from nitrates and carbon dioxide." By E. C. C. BALY, I. M. HEILBRON, and D. P. HUDSON.
- " Attempts to prepare red sulphide dyes. Part I." By E. R. WATSON and S. DUTT.
- " Limits for the propagation of flame in vapour-air mixtures. Part I. Mixtures of air and one vapour at the ordinary temperature and pressure." By A. G. WHITE.
- " Studies on hypophosphorous acid. Part IV. Its reaction with cupric chloride." By A. D. MITCHELL.

- "The labile nature of the halogen atom in organic compounds. Part III. The absorption spectra of bromomalonic derivatives and nitroparaffins, and their bearing on the question of an oxygen-halogen linking." By H. GRAHAM and A. K. MACBETH.
- "The labile nature of the halogen atom in organic compounds. Part IV. The tautomeric hydrogen hypothesis and the removal of the halogen atom from aromatic nitro-compounds." By A. K. MACBETH.
- "The action of sulphur on cuprous chloride." By F. W. PINKARD and W. WARDLAW.
- "Dyes derived from phenanthraquinone. Part I. Phenanthranaphthazines." By A. C. SIRCAR and S. B. DUTT.
- "Physical chemistry of the oxides of lead. Part IV. Red lead and lead sesquioxide." By S. GLASSTONE.
- "The sulphilimines, a new class of organic compounds containing quadrivalent sulphur." By F. G. MANN and W. J. POPE.
- "The aldehydo-salicyclic acids and their derivatives." By E. J. WAYNE and J. B. COHEN.
- "The effect of an electrolyte on solutions of pure soap. Phase rule equilibria in the system sodium laurate-sodium chloride-water." By J. W. MCBAIN and A. J. BURNETT.
- "Investigations into the analytical chemistry of tantalum, columbium, and their mineral associates. Part III. A critical study of the volumetric estimation of columbium." By W. R. SCHOELLER and E. F. WATERHOUSE.
- "Mercury compounds of hydroxybenzaldehydes." By T. A. HENRY and T. M. SHARP.
- "*iso*Quinoline and the *iso*quinoline-reds." By J. E. G. HARRIS and W. J. POPE.
- "Studies in optically active dyes. Part I. Camphoreins." By B. K. SINGH, R. RAI, and R. LAL.
- "The hydroferrocyanides and hydroferricyanides of the organic bases." Part I. By W. M. CUMMING.
- "The viscosity of cellulose in cuprammonium hydroxide solution. Part I. The determination of the viscosity." By R. A. JOYNER.
- "The oxidising and reducing properties of sulphur dioxide. Part IV. Mercury chlorides." By L. M. STEWART and W. WARDLAW.
- "The conditions underlying the formation of unsaturated and cyclic compounds from halogenated open-chain derivatives. Part IV. Products formed from halogen derivatives of muconic acid. The constitution of muconic acid." By J. P. C. CHANDRASENA and C. K. INGOLD.

- "Studies in the coumaranone series. Part II. The preparation of 6- and 4-chlorocoumaran-2-ones and their conversion into 4- and 2-chloroflavanols respectively and some derivatives of *o*- and *p*-chlorophenoxyacetic acids." By T. H. MINTON and H. STEPHEN.
- "The preparation of *o*-, *m*-, and *p*-nitrophenoxyacetic acids and various nitrotolxyoxyacetic acids and their derivatives." By T. H. MINTON and H. STEPHEN.
- "Some properties of hydrogen desorbed from platinum and palladium." By P. ANDERSON.
- "The reactivity of doubly-conjugated unsaturated ketones. Part III. Unsymmetrical hydroxy- and methoxy-derivatives." By J. S. BUCK and I. M. HEILBRON.
- "Phenopyryllium salts of distyryl ketones. Part I." By J. S. BUCK and I. M. HEILBRON.
- "The mathematics of the dicyclic colour theory and a new theory of the nitrogen atom." By J. MOIR.
- "Physical chemistry of the oxides of lead. Part V. The electro-motive behaviour of lead dioxide." By S. GLASSTONE.
- "Studies of the constitution of soap solutions. Sodium behenate and sodium nonylate." M. TAYLOR.
- "Studies in the anthracene series. Part II." By E. DE B. BARNETT and J. W. COOK.

ADDITIONS TO THE LIBRARY.

I. *Donations.*

ACTUALITÉS DE CHIMIE CONTEMPORAINE. Publiées sous la direction de A. HALLER. 1st series. Paris 1922. pp. viii + 326. ill. 12 fr. (*Recd.* 11/4/22.)

From the Publisher : M. Gaston Doin.

IMPERIAL MINERAL RESOURCES BUREAU. The Mineral Industry of the British Empire and Foreign Countries. War Period. Copper (1913—1919). London 1922. pp. 204. 4s. net. (*Recd.* 1/5/22.)

— — — Gold (1913—1919). London 1922. pp. 366. 6s. net. (*Recd.* 11/4/22.)

From the Bureau.

PURVIS, JOHN EDWARD, and HODGSON, THOMAS REGINALD. The chemical examination of water, sewage, foods and other substances. 2nd edition. Cambridge 1922. pp. viii + 346. 20s. net. (*Recd.* 3/5/22.)

From the Syndics, Cambridge University Press.

STALEY, RALPH. Town gas manufacture : a practical introductory treatment of the equipment and processes of an average gas works. London 1922. pp. xii + 108. ill. 2s. 6d. net. (*Recd.* 1/5/22.)

From the Publishers : Sir Isaac Pitman & Sons.

II. By Purchase.

SAINTE-CLAIRE DEVILLE, ÉMILE. Manuel de chimie gazière : méthodes et procédés des essais et analyses en usage à l'usine expérimentale de la villette de la Société du Gaz de Paris. Paris 1921. pp. viii + 208. ill. (*Recd.* 2/5/22.)

TRIVELLI, ADRIAAN PETER HERMAN, and SHEPPARD, SAMUEL EDWARD. The silver bromide grain of photographic emulsions. Rochester, N.Y. 1921. pp. 144. ill. 16s. net. (*Recd.* 1/5/22.)

VRIES, OTTO DE. Estate rubber : its preparation, properties and testing. Batavia 1920. pp. xii + 650. ill. 40s. net. (*Recd.* 2/5/22.)

WILLGERODT, CONRAD. Die organischen Verbindungen mit mehrwertigem Jod. (Chemie in Einzeldarstellungen, Vol. VII.) Stuttgart 1914. pp. xii + 266. (*Recd.* 2/5/22.)

ZEITSCHRIFT FÜR LEDER- UND GERBEREI-CHEMIE. Vol. I, etc. Hamburg 1921 +. ill. (*Reference.*)

ZEITSCHRIFT FÜR PFLANZENERNÄHRUNG UND DÜNGUNG. A. Wissenschaftlicher Teil. B. Wirtschaftlich-Praktischer Teil. Vol. I, etc. Leipzig 1922 +. (*Reference.*)

CERTIFICATES OF CANDIDATES FOR ELECTION AT THE BALLOT TO BE HELD AT THE ORDINARY SCIENTIFIC MEETING ON THURSDAY, JUNE 15TH, 1922.

N.B.—The names of those who sign from "General Knowledge" are printed in *italics*.

BABOROVSKÝ, GEORGE, Ph.D. (Prague), Brno, Falkensteinerova, ul. 7. Professor in the Technical High School of Brno, Techeoslowackia. Member of the Société chimique de France, Techeoslowack Chemical Society, formerly member of the American Electrochemical Society, formerly Professor in the University of Prague. 1900–1905 assistant, 1905–1911 lecturer in physical chemistry; 1911–1912 professor of physical chemistry in the Czech University of Prague; since 1912 professor of physical chemistry in the Czech Technical High School of Brno. Publications on different physico-chemical and electrochemical subjects in following periodicals : *Bulletin of the Czech Acad. of Science*, *Zeit. f. phys. Chem.*, *Zeit. f. Elektrochem.*, *Phys. Zeitschr.*, *Chem. Listy*. Books : "Electrochemistry" (with Prof. F. Plzák), "Theoretical and Physical Chemistry" (1920) (boh.). (*Signed by*) Gerald Druce, A. Šimek, J. Heyrovský.

BAILEY, STANLEY JOHN, The Braes, South Nutfield, Surrey. Student (Cambridge University), Natural Science Tripos, Part I, in 1921, including Chemistry. (*Signed by*) T. Martin Lowry, Alfred Greeves, Hamilton McCombie, William J. Pope.

BAINES, HARRY, 22, Henry Rd., West Bridgford, Notts. British. Research Student. B.Sc. (Lond.), external, 1st Class Hons. in Chemistry. Now engaged on research course. (*Signed by*) F. Stanley Kipping, S. J. Pentecost, F. C. Laxton.

BELTON, FRANK, "Athlone," Bush Hill Park, Enfield. Headmaster, County Secondary School, Edmonton, N.9. (1) B.Sc. (London), Chemistry at Final Exam. (2) Studied chemistry at the Royal Technical College, Salford; with a short course at the Royal College of Science, London. (3) Head of Science Dept. and teacher of chemistry, Central Higher Grade School, Salford. (4) Author of text-books on physics and chemistry. (*Signed by*) B. Prentice, E. Clark, E. Green.

BRANSON, VICTOR CECIL, Naparima Oilfields, Ltd., San Fernando, Trinidad, B.W.I. Analytical Chemist. A.R.C.Sc. (1st class chemistry), B.Sc. Lond. (1st class honours in Chemistry), A.I.C. Chemist to the Naparima Oilfields, Ltd., since Jan. 1920. (*Signed by*) L. H. Parker, James C. Philip, H. B. Baker.

CANNAN, ROBERT KEITH, 4, Antrim Mansions, N.W.3. Assistant in Biochemistry, University College, London. Several papers in *Biochemical Journal* and *Journal of Physiology*. M.Sc. (Lond.) in Biochemistry. (*Signed by*) J. C. Drummond, R. H. A. Plimmer, Irvine Masson.

CHATTERJEE, BIBHU CHARAN, Ariadah, 24, Perganas, Bengal. Research Scholar in Chemistry. B.Sc. (Hons. in Chemistry), 1919; M.Sc. in Chemistry, 1921; first in Class I; conducted a piece of research work in organic chem. entitled, "On carbamic esters; chlorine as a simultaneous oxidising and chlorinating agent"; Sir T. N. Palit Research Scholar in Chemistry, Calcutta University. (*Signed by*) P. C. Rây, R. L. Datta, J. N. Mukherjee.

COLBORNE, ROBERT STANLEY, 57, County Road, Swindon. Research Student in Physical Chemistry (Imperial College, London). Student at Imperial College since 1919 (Oct.); A.R.C.S. (1st Class) 1921 (July); B.Sc. (Hons. 1st Class) 1921 (Nov.). (*Signed by*) H. B. Baker, M. A. Whiteley, James C. Philip, H. J. T. Ellingham.

CONTRACTOR, SHIAVAKSHA HORMASJI, F.N.B.A., F.S.Sc. (Lond.), Karanjia Building, opp. Moos Building, Grant Rd., Bombay. Scientist, Artist and Craftsman. Inventor of gold mirror glasses; Permanent blotters; Sizing Composition, etc. Recipient of 50 gold, silver, and bronze medals, and hundreds of diplomas at the world's various exhibitions. (*Signed by*) J. C. Mascarenhas, N. M. Bhathena, D. D. Kanga.

FIRTH, JAMES BRIERLEY, 44, Lucknow Avenue, Mapperley Park, Nottingham. Lecturer in Chemistry, University College, Nottingham. D.Sc. (Manc.), M.Sc. (Dunelm), F.I.C. Author or joint-author of some sixteen published researches, most of which have appeared in the *Journal of the Chemical Society*. (*Signed by*) F. S. Kipping, Henry Stephen, J. E. Myers, Colin Campbell, Norman Smith, Arnold Weston.

FISHENDEN, MARGARET WHITE, 19, Shepherds Hill, Highgate, N.6. Scientific Investigator for Fuel Research Board. 1909, B.Sc. (Manchester), 1st Class Honours Physics; 1919, D.Sc. Have published a number of papers on Meteorology (upper air) and conditions in weaving factories and spinning mills. Also the following: "The Coal Fire" (Special Report, No. 3, of Fuel Research Board). "The Efficiency of Low Temperature Coke in Domestic Appliances" (Technical Paper, No. 3, of Fuel Research Board). (*Signed by*) George Beilby, Harold B. Dixon, Frank Lee Pyman.

GODDARD, FREDERICK WALTER, 12, College St., Winchester. Assistant Master, Winchester College. M.A. (Cantab.); 1st Class Nat. Science Tripos

(1st Class Chemistry, Physics, Mineralogy); 1st Class Maths. Tripos, Pt. 1. War Service. Five years teaching Chemistry at Win. Coll. to Scholarship Candidates and post scholarship. Head of Chem. Dept., Win. Coll. (*Signed by*) Llewellyn L. Garbutt, D. Ll. Hammick, A. J. Berry.

HASSELL, ALBERT, 47, Stanley Street, Tunstall, Stoke-on-Trent. Science Master, Denstone College, Staffs. Chemistry Master, Denstone College, Staffs. I wish to obtain the Transactions so as to keep in touch with current chem. literature. (*Signed by*) Joseph West, H. V. Thompson, F. H. Clews.

HIRST, EDMUND LANGLEY, Ainslie Villa, Largo Rd., St. Andrews. Assistant in Chemistry Department, University of St. Andrews. Degrees: M.A., B.Sc., Ph.D. Assistant in Chemistry at St. Andrews University from October 1920. Publication: "The constitution of the disaccharides, Part V (Cellobiose)," by Walter Norman Haworth and Edmund Langley Hirst (T., 1921, 119, 193). (*Signed by*) J. C. Irvine, R. Robinson, R. C. Menzies, A. Killen Macbeth.

HOFFERT, DOROTHEA ANNIE, 50, Hampstead Way, London, N.W.11. Research worker under the Food Investigation Board. 1916-1920, four years' experience in analytical and research work on dopes and varnishes for aeroplanes (war work). One year, 1920-1921, student in training for research, Cambridge University, with grant from Scientific and Industrial Research Department. (*Signed by*) William J. Pope, W. H. Mills, Charles T. Heycock, Frances M. Hamer, Ida Smedley Maclean, J. N. Goldsmith.

JENKINS, WILLIAM JOB, 37, Marston Road, Knowle, Bristol. Research Chemist. Plant chemist at H.M. Factory, Pembrey, 1915-1918. Research Chemist at Messrs. Nobels Ardeer Factory, 1920-1921. Research worker under Department of Scientific and Industrial Research, 1921-1922. (*Signed by*) Wm. Rintoul, N. Pictou, R. A. Joyner, Wm. Barbour.

LEVY, PEPPO JOSEPH, 42, Granville Gardens, Shepherds Bush, London W.12. Spanish. Chemist (M.F.G.) intending. I am a student at the day school of Chemistry of the Polytechnic and wish to follow the latest developments in chemical science, and to use the Society's library. (*Signed by*) Frank E. Weston, W. H. Collier, Frank Matthews.

MCCALL, DAVID, 4, Lime Street, Dundee. Head of Pharmacy Dept., Technical College, Dundee. Degree of B.Sc. (St. Andrews), July 1919. Diploma of Pharmaceutical Chemist, April 1922. Member of Pharmaceutical Society of Gt. Britain. Lecturer on Botany, Pharmacy, and Materia Medica. Head of Pharmacy Dept., Technical College, Dundee, 1919. (*Signed by*) Alex. McKenzie, John S. Lumsden, John S. W. Boyle, John Foggie, H. J. Plenderleith.

MALLET, RICHARD EDMUND ANTONY, "Redford," Ockley, Surrey. Research Chemist. B.A. of Oxford University. Honours degree passed with distinction. Now employed as chemist at the British Museum. Have been engaged in research work at Oxford with Prof. T. R. Merton, F.R.S. (*Signed by*) Alexander Scott, D. Ll. Hammick, C. N. Hinshelwood, H. R. Raikes, H. Hartley.

NORRISH, RONALD GEORGE WREYFORD, Emmanuel College, Cambridge, and 48, Kimberley Rd., Cambridge. Research Student in Photochemistry. Scholar of Emmanuel College, Cambridge. Natural Sciences Tripos, Part I, Class I. Natural Sciences Tripos, Part II Chemistry, Class I. Paper: "The Transition Elements and the Octet Theory," *Chem. News*, 1922, 124, 16-22. My reason for seeking admission is that I hope to make the study of Chemistry my chief occupation and interest in life. (*Signed by*) W. J. Pope, W. H. Mills, Eric K. Rideal, T. M. Lowry.

OATS, JOHN GOODRICH (junr.), 3, Montpelier Terrace, Ilfracombe. Engineering Student at R. A. Lister & Co., Dursley. Also chemical student (private laboratory). Student of metallurgy, chemistry, and analysis; holding a diploma from the International Correspondence Schools, including the subjects: mathematics, physics, inorganic chemistry, qualit. analysis, quant. analysis, and the manufacture of iron and steel. (*Signed by*) F. Southerden, Alfred L. Grey, R. Barton, G. F. Hall, Ernest F. Hooper, Jas. Walker.

PARKER, RALPH HENRY, "Radio House," Wilson Rd., Smethwick, Staffs. Student in Electrical and Mechanical Engineering and allied subjects. Research Student. (1) Took the 3 years' course in physics (mechanics, heat, electricity, and magnetism) at the (full time) Junior Day Technical School, Smethwick, years 1915-1918. (2) For three years continued with part day classes, 2 years' evening tuition, also matriculation class in chemistry, Smethwick Municipal Tech. (3) Studying Organic Chemistry at present time (1922). I will endeavour to forward the interests of the Society to the best of my ability and endeavour to keep abreast with modern chemical science. (*Signed by*) Arthur Adams, Joseph Lones, J. H. Wood.

PEACH, EDITH ANNIE, 43, St. Gabriel's Road, Cricklewood, N.W.2. Student of Chemistry at University College, London. Desirous of receiving the *Journal* and of using the Library. (*Signed by*) Irvine Masson, F. G. Donnan, J. C. Drummond.

PLATT, CLAUDE BERNARD MEISTER, Captain, M.B.E., R.E., Middleton House, Beckett Rd., Doncaster. Colliery Chemist, and Fuel Technologist. Two years as assistant to Prof. R. V. Wheeler, D.Sc., F.I.C., F.C.S.; engaged on research on spontaneous combustion of coal, composition of coal, research on explosions in mines and general laboratory work at Home Office Experimental Station. 2nd year Imperial College of Science and Technology (interrupted by the war). During war engaged on Army Wireless Telegraphy. At present chemist to Doncaster Collieries Association, Ltd., and assistant to Hot and Deep Mines Research Committee of Dept. of Scientific and Industrial Research. Published in *Trans. Chem. Soc.*, 1913, 108, 1713. M.Soc.C.I., A.M.I.Min.E. Twelve months' experience on technical side of boiler practice—especially as regards fuel combustion. (*Signed by*) R. Lessing, W. Payman, R. V. Wheeler.

PLOWMAN, ARTHUR, 8, Rowantree Rd., Bycullah Park, Enfield. Research Student. B.Sc. (Lond.), A.R.C.S., A.I.C. (*Signed by*) R. Winton West, G. A. R. Kon, E. H. Farmer.

PRYDE, JOHN, 4, Gowrie St., Blackness Ave., Dundee. Research Chemist, St. Andrews University, N.B. Degree: B.Sc. (Pure Science). 1851 Exhibition Research Scholar, 1921. Publication (with J. A. Hewitt): "Metabolism of Carbohydrates, Pt. I," *Biochem. J.*, XIV, 3 and 4, 395. (*Signed by*) J. C. Irvine, R. Robinson, R. C. Menzies, A. Killen Macbeth.

QUASTEL, JUDA HIRSCH, Trinity College, Cambridge. Research Student, Biochemical Dept., Cambridge. Obtained A.R.C.S. (Lond. First Cl.), June 1921; elected Associate of Institute of Chemistry, June 1921; obtained B.Sc. Hons. (Lond. First Cl.) in October 1921. Accepted as candidate for Ph.D. (Cambridge). I wish to receive the Society's publications and to take part in its activities. (*Signed by*) F. G. Hopkins, H. B. Baker, James C. Philip.

SAMUELS, EDMOND, 40, Castlereagh St., Sydney, N.S.W. Pharmaceutical Chemist. Chemistry course, Sydney University. Interested in preparation of pharmaceutical products for physicians' use. Originator of several

medicinal products. Member of Pharmaceutical Society of N.S.W.; member of Millions Club, N.S.W.; member of Justices Association, N.S.W. (*Signed by*) J. Laker Macmillan, G. Spurge, Francis L. Watt.

SMITH, JOHN CHARLES, 31, Bidwell Street, Wellington, N.Z. Student of Chemistry. B.Sc. (N.Z.) 1919-20. Thesis on "The Condensation of Phenolic Substances with Basic Imino-haloids," presented for the degree of M.Sc. (N.Z.), Nov. 1921. (*Signed by*) F. P. Worley, W. F. Short, A. H. Bowell.

STEVENSON, RONALD WILLIAM, M.C., M.A., Major, Eaglehurst, Ditton Hill, Surbiton, Surrey. Manufacturing Chemist, Analytical and Research work. Director, Messrs. Stevenson and Howell, Ltd., Manufacturing Chemists. M.A., Cambridge (Science). (*Signed by*) Arthur Hutchinson, William Stevenson, Vernon J. Tilley.

SWAINSON, VALERIE EMELINE SOPHIE, 21, Hungerford Rd., N.7. Secondary School teacher (teaching Chemistry up to Intermediate Standard). B.Sc. (Hons.) Lond. in Chemistry. Demonstrator (temporary), mainly in Inorganic Chemistry, for two terms, at University College, Dundee (Jan.-June, 1921). (*Signed by*) H. Crompton, M. S. Crewdson, J. F. Spencer.

THOMAS, CLIFFORD DANE, 10, Lansdowne Place, Blackheath, S.E.10. British. Research Chemist. Second Class Honours, B.Sc., 1920. One year's research under Dr. O. L. Brady at University College, 1920-1921. Junior Research Assistant at Research Dept., Woolwich. (*Signed by*) Eustace E. Turner, Sydney A. Pearman, W. J. Powell.

VAZIFDAR, NOWROJI JAMSHEDJI, 17, Queens Road, Opp. Charin Road Station, Bombay. First Assistant Chemical Analyser to Government. L.M. & S. Bombay. Teacher in Chemistry, Grant Medical College, Bombay. Chemical analyser to Govt. for Sind, Karachi, 1914-15; 1916-17. Author of "Physiology of the Nervous System and Special Senses." Coach in physiology for the last 15 years. (*Signed by*) D. D. Kanga, M. J. Gajjar, K. G. Naik.

WIGG, GEORGE WILLIAM, 72nd Northumbrian Bde., R.F.A., Barrack Rd., Newcastle-on-Tyne. Lt., R.F.A. 4th-5th year medical student, University of Durham, College of Medicine. 3rd year honours chemistry student, Armstrong college, Newcastle-on-Tyne. (*Signed by*) W. N. Haworth, Frank M. Irvine, D. Arthur Ruell.

The following Certificates have been authorised by the Council for presentation to ballot under Bye-law I (2) :

HALL, ROBERT SWAINSON, F.L.S., F.A.M.S., c/o Govt. Plantations Dept., Victoria, British Cameroons, West Africa. Supt. of agriculture to Nigerian Government. Three years' training in chemistry, Waterloo Secondary School, Oldham, Lancs. Four years' practical experience in agricultural chemistry in Portuguese Congo, S.W. Africa. Contribution to science mostly on mycological and agricultural matter; see *Kew Bulletin*, *Bull. of Imperial Institute*, etc. (*Signed by*) John Taylor.

FULWEILER, WALTER HERBERT, Wallingford, Pa., U.S.A. Chemical Engineer with The United Gas Improvement Co., Phila., Pa. B.S. in Chemistry, University of Pennsylvania, 1901. Chemist with Phila. Gas Works, 1901-1902. Chief chemist, The United Gas Improvement Co., 1913-1920. Chemical engineer, The United Gas Improvement Co., 1920-date. (*Signed by*) H. Jermain Creighton.

PROCEEDINGS

OF THE

CHEMICAL SOCIETY.

Ordinary Scientific Meeting, Thursday, June 1st, 1922, at 8 p.m.,
Professor T. M. LOWRY, C.B.E., F.R.S., Vice-President, in the
Chair.

The CHAIRMAN referred to the loss sustained by the Society,
through death, of :

	Elected.	Died.
George Robertson Hislop	Jan. 20th, 1870.	May 18th, 1922.

The following were formally admitted Fellows of the Chemical
Society : A. D. Gay and E. A. Seeley.

Certificates were read for the first time in favour of :

Allin Cottrell, 3, Oxford Terrace, Edinburgh.
Reginald William Griffiths, B.Sc., 137, Holly Road, Handsworth, Birmingham.
John Watson Jenkin, B.Sc., A.I.C., Tregardock, St. Austell.
Ghulam Saffdar Khan, B.Sc., Veterinary College, Lahore, India.
Hiroshi Nomura, Chemical Institute, Imperial University, Sendai, Japan.
Harold Pickles, The Gasworks, Kildwick, Keighley.
Stephen John Watson, B.Sc., 35, Oakfield Terrace, Gosforth, Newcastle-on-
Tyne.
Alexander Mitchell Williams, M.A., D.Sc., Shirley Institute, Didsbury,
Manchester.

The following papers were read :

- "The reactivity of doubly-conjugated unsaturated ketones.
Part III. Unsymmetrical hydroxy- and methoxy-derivatives."
By J. S. BUCK and I. M. HEILBRON.
- "Phenopyryllium salts of distyryl ketones. Part I." By J. S.
BUCK and I. M. HEILBRON.
- "Ring-chain tautomerism. Part II. The effect of the gem-
diethyl group on the carbon tetrahedral angle." By S. S.
DESHAPANDE and J. F. THORPE.

Ordinary Scientific Meeting, held in the Lecture Hall of the Institution of Mechanical Engineers, on Thursday, June 8th, 1922, at 8 p.m., Professor H. E. ARMSTRONG, LL.D., F.R.S., Vice-President, in the Chair.

Dr. H. H. Dale, C.B.E., F.R.S., delivered his Lecture entitled : "Chemical and Physiological Properties." A vote of thanks to the Lecturer, proposed from the Chair, and seconded by Mr. Francis H. Carr, was carried with acclamation, and acknowledged by Dr. Dale.

Ordinary Scientific Meeting, Thursday, June 15th, 1922, at 8 p.m., Sir JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

Reference was made to the loss sustained by the Society, through death, of :

	Elected.	Died.
William Gowland	Feb. 2nd, 1871.	June 10th.
Thomas Henry Johnson	May 5th, 1921.	Feb. 19th.

The following were formally admitted Fellows of the Chemical Society : R. A. Fitzsimons, H. U. Harrow, E. M. James, and H. L. Riley.

Certificates were read for the first time in favour of :

William Leslie Carter, Ellesmere, Bloxcliffe Street, Langley Green, Birmingham.

Wilfred Thomas Grey Davies, B.Sc., Plas y Coed, Derllwyn Road, Tondur, Bridgend.

Roy Gardner, B.Sc., Technical High School, Dunedin, New Zealand.

John Ernest Jewell, 34, Grove End Road, St. John's Wood, N.W. 8.

Charles William Moorhead, 23, Newbridge Road, Bath.

Albert Bramah Pearce Page, A.R.C.S., 27, Tanza Road, Hampstead, N.W. 3.

Arthur Scholefield, Poulton Hey, Bromborough, Birkenhead.

Stanley James Corral Snedker, The Rowans, Wood Hey, Rock Ferry, Birkenhead.

Dr. H. M. Atkinson and Dr. P. C. Austin were elected Scrutators, and a ballot for the election of Fellows was held. The following were subsequently declared elected as Fellows :

George Baborovský, Ph.D.

Stanley John Bailey, B.A.

Harry Baines, B.Sc.

Frank Belton, B.Sc.

Victor Cecil Branson, B.Sc., A.R.C.S.,
A.I.C.

Robert Keith Cannan, M.Sc.

Bibhu Charan Chatterjee, M.Sc.

Robert Stanley Colborne, B.Sc.,
A.R.C.S.

James Brierley Firth, D.Sc., F.I.C.

Margaret White Fishenden, D.Sc.

Walter Herbert Fulweiler, B.S.

Frederick Walter Goddard, M.A.

Robert Swainson Hall.

Albert Hassell.

Edmund Langley Hirst, M.A., B.Sc., Ph.D.	Arthur Plowman, B.Sc., A.R.C.S., A.I.C.
Dorothea Annie Hoffert.	John Pryde, B.Sc.
William Job Jenkins.	Juda Hirsch Quastel, B.Sc., A.R.C.S.
Peppo Joseph Levy.	Edmond Samuels.
David McCall, B.Sc.	John Charles Smith, B.Sc.
Richard Edmund Antony Mallet, B.A.	Ronald William Stevenson, M.C., M.A.
Ronald George Wreyford Norrish, B.A.	Valerie Emeline Sophie Swainson, B.Sc.
John Goodrich Oats (Junr.).	Clifford Dane Thomas, B.Sc.
Ralph Henry Parker.	Nowroji Jamshedji Vazifdar.
Edith Annie Peach.	George William Wigg.
Claude Bernard Meister Platt, M.B.E.	

The following papers were read :

- "Ring-chain tautomerism. Part III. The occurrence of tautomerism of the three-carbon (glutaconic) type between a homocyclic compound and its unsaturated open-chain isomeride." By C. K. INGOLD, E. A. PERREN, and J. F. THORPE.
- "The tautomerism of dyads. Part I. Experiments on the detection of tautomeric equilibria in hydrocyanic acid." By E. H. USHERWOOD.

LIST OF FELLOWS, 1922.

The List of Fellows for 1922 is being prepared, and changes of address received after July 31st, 1922, cannot be included therein. Fellows whose Christian names and degrees do not appear in full are particularly requested to notify the Assistant Secretary.

Owing to the heavy cost of printing, the List will be sent only to those Fellows who make application prior to August 31st, 1922.

THE LIBRARY.

The Library will be closed for Stocktaking from Monday, August 7th, until Saturday, August 19th, 1922, inclusive.

During the four weeks August 21st to September 16th the Library will close daily at 5 p.m.

List of Papers received between May 19th and June 15th, 1922 :

- "The interaction of sodium chloride and silica." By F. H. CLEWS and H. V. THOMPSON.
- "Ring-chain tautomerism. Part II. The effect of the gem-diethyl group on the carbon tetrahedral angle." By S. S. DESHAPANDE and J. F. THORPE.

- "Dyes derived from saccharine. The sulphim-phthaleins." By S. DATTA.
- "Dyes derived from diphenyl-ketipic-dinitrile." By S. DATTA and N. K. SEN.
- "Thermal expansion of gelatin gels." By A. TAFFEL.
- "The rôle of protective colloids in catalysis. Part II." By T. IREDALE.
- "The chlorinated dialkyl sulphides." By W. J. POPE and J. L. B. SMITH.
- "Gallotannin. Part XIII. The identity of digallic acid from gallotannin with synthetic *m*-digallic acid." By M. NIERENSTEIN and C. W. SPIERS.
- "Experiments on the synthesis of the polyacetic acids of methane. Part VI. Conditions of formation by the cyanoacetic ester method of unstable methanetriacetic esters and a new synthesis of methanetriacetic acid." By C. K. INGOLD and E. A. PERREN.
- "Tables of average specific and atomic heat capacities of chemical elements in the order of the increment of the atomic weights and their interpretations. The law of Dulong and Petit." By P. N. TCHIRVINSKY.
- "The dissociation pressures of hydrated double sulphates. Part I. Hydrated cupric alkali sulphates." By R. M. CAVEN and J. FERGUSON.
- "The action of sodium sulphite on lead iodide." By G. SINGH.
- "2 : 3 : 6-Trimethyl glucose." By J. C. IRVINE and E. S. STEELE.
- "Researches on the constitution of coal." By S. R. ILLINGWORTH.
- "The composition of paraffin wax. Part II." By F. FRANCIS, C. M. WATKINS, and R. W. WALLINGTON.
- "Ring-chain tautomerism. Part III. The occurrence of tautomerism of the three-carbon (glutaconic) type between a homocyclic compound and its unsaturated isomeride." By C. K. INGOLD, E. A. PERREN, and J. F. THORPE.
- "Note on the effect of a magnetic field on catalysis by ions in the presence of a para-magnetic salt." By W. E. GARNER and D. N. JACKMAN.
- "Reciprocal induced polarity effects in cresols and their derivatives. Properties of the isomeric methoxybenzyl bromides." By A. LAPWORTH and J. B. SHOESMITH.
- "Intermetallic actions. The system aluminium-arsenic." By Q. A. MANSURI.
- "The tautomerism of dyads. Part I. Experiments in the detection of tautomeric equilibria in hydrocyanic acid." By E. H. USHERWOOD.

- "The nitro-derivatives of metanitrodimethylaniline." By A. FORSTER and W. COULSON.
- "Catalytic racemisation of optically active acid amides." By A. MCKENZIE and I. A. SMITH.
- "The sorption of phosgene by beech wood charcoal." By H. M. BUNBURY.
- "The adsorption of uranium-X and its isotope thorium by basic ferric acetate." By A. C. BROWN.
- "The reactions of cellulose with sodium chloride and other neutral salt solutions. Part I. Preliminary survey." By H. MASTERS.
- "The imino-aryl ethers. Part I. *N*-Phenylbenziminom-*m*-hydroxyphenyl ether and the synthesis of 2:4-dihydroxybenzophenone." By A. W. CHAPMAN.
- "The intermolecular condensation of methylethyl ketone in the presence of carbide." By O. BECKER and J. F. THORPE.
- "The equilibrium in systems composed of water and alcohols. The system methyl alcohol-water." By N. A. PUSHIN and A. A. GLAGOLEVA.
- "The equilibrium in the system dinitrobenzene-urethane." By N. A. PUSHIN and A. FIOLETOVA.
- "The chemistry of polycyclic structures in relation to their homocyclic unsaturated isomerides. Part III. Intra-annular tautomerism of α -campholytic acid." By J. P. C. CHANDRASENA, C. K. INGOLD, and J. F. THORPE.
- "Adsorption and catalysis in Fuller's earth." By E. K. RIDEAL and W. THOMAS.
- "The formation and stability of *spiro*-compounds. Part VIII. The Dieckmann-Komppa reaction." By F. DICKENS, G. A. R. KON, and J. F. THORPE.
- "The cyanine dyes. Part V. The virtual tautomerism of the thio-cyanines." By W. H. MILLS and W. T. K. BRAUNHOLTZ.

ADDITIONS TO THE LIBRARY.

I. *Donations.*

BERNTHSEN, AUGUST. A textbook of organic chemistry. New edition, revised to date. By JOHN JOSEPH SUDBOROUGH. London 1922. pp. xvi + 908. 12s. 6d. net. (*Recd.* 16/5/22.)

From the Publishers: Messrs. Blackie & Son.

BROOKS, BENJAMIN TALBOTT. The chemistry of the non-benzenoid hydrocarbons and their simple derivatives. New York 1922. pp. 612. ill. \$7 net. (*Recd.* 11/5/22.)

From the Publishers: The Chemical Catalog Company, Inc.

DUCLAUX, JACQUES. Les colloïdes. 2nd edition. Paris 1922. pp. x + 306. 10 fr. (*Recd.* 13/6/22.)

From the Publishers : MM. Gauthier-Villars et Cie.

FALK, KAUFMAN GEORGE. Catalytic action. New York 1922. pp. 172. \$2.50 net. (*Recd.* 8/6/22.)

From the Publishers : The Chemical Catalog Company, Inc.

FRIEND, JOHN NEWTON. The chemistry of combustion. London 1922. pp. viii + 110. ill. 4s. net. (*Recd.* 15/5/22.)

From the Publishers : Messrs. Gurney & Jackson.

HICKS, WILLIAM MITCHINSON. A treatise on the analysis of spectra : based on an essay to which the Adams prize was awarded in 1921. Cambridge 1922. pp. viii + 326. ill. 35s. net. (*Recd.* 11/5/22.)

From the Syndics, Cambridge University Press.

IMPERIAL MINERAL RESOURCES BUREAU. Iron ore. (Summary of information as to the present and prospective iron-ore supplies of the world.) Part I. United Kingdom. London 1922. pp. 238. ill. 6s. net. (*Recd.* 7/6/22.)

— The Mineral Industry of the British Empire and Foreign Countries. War Period. Coal, coke, and by-products (1913—1919). Part II. London 1922. pp. 160. ill. 6s. 6d. net. (*Recd.* 29/5/22.)

From the Bureau.

KEMPTON, PERCIVAL HENRY SYDNEY. Industrial nitrogen : the principles and methods of nitrogen fixation and the industrial applications of nitrogen products in the manufacture of explosives, fertilizers, dyes, etc. London 1922. pp. xii + 104. ill. 2s. 6d. net. (*Recd.* 2/6/22.)

From the Publishers : Sir Isaac Pitman & Sons.

KNOX, JOSEPH. The fixation of atmospheric nitrogen. 2nd edition. London 1921. pp. viii + 124. ill. 4s. net. (*Recd.* 15/5/22.)

From the Publishers : Messrs. Gurney & Jackson.

LAURIE, ARTHUR PILLANS. Building materials : an introduction to the study of the principal materials used in building construction. Edinburgh 1922. pp. viii + 188. ill. 6s. net. (*Recd.* 15/5/22.)

From the Publishers : Messrs. Oliver & Boyd.

PLIMMER, VIOLET G., and PLIMMER, ROBERT HENRY ADERS. Vitamins and the choice of food. London 1922. pp. xii + 164. ill. 7s. 6d. net. (*Recd.* 22/5/22.)

From the Publishers : Messrs. Longmans, Green & Co.

SHREVE, RANDOLPH NORRIS, WATSON, WARREN N., and WILLIS, A. R. Dyes classified by intermediates. Dyes tabularly arranged under each intermediate, with statistical and other data for both dyes and intermediates. Glossary of dye and intermediate names alphabetically arranged. New York 1922. pp. ii + 632. \$10 net. (*Recd.* 31/5/22.)

From the Publishers : The Chemical Catalog Company, Inc.

STEWART, ALFRED WALTER. Some physico-chemical themes. London 1922. pp. xii + 420. ill. (*Reference.*) 21s. net.

From the Publishers : Messrs. Longmans, Green & Co.

II. *By Purchase.*

BUDOWSKI, I. Die Naphthensäuren. Berlin 1922. pp. vi + 116. ill. (*Recd.* 31/5/22.)

FODOR, ANDOR. Das Fermentproblem. Dresden 1922. pp. x + 280. ill. (*Recd.* 23/5/22.)

FUCHS, WALTER. Der gegenwärtige Stand des Gärungsproblems. (Sammlung, Vol. XXVII.) Stuttgart 1922. (*Reference.*)

GLAZEBROOK, SIR RICHARD. [Editor.] A dictionary of applied physics. Vol. I., etc. London 1922 +. ill. (*Reference.*)

GUILLET, LÉON, and PORTEVIN, ALBERT. An introduction to the study of metallography and macrography. Translated by LEONARD TARVENER. London 1922. pp. xvi + 290. ill. 30s. net. (*Recd.* 13/6/22.)

KEGHEL, MAURICE DE. Les encres, les cirages, les colles et leur préparation. Paris 1922. pp. 384. ill. (*Recd.* 10/6/22.)

LEWKOWITSCH, JULIUS. Chemical technology and analysis of oils, fats, and waxes. 6th edition, revised by GEORGE H. WARBURTON. Vol. II. London 1922. pp. xii + 960. ill. 42s. net. (*Recd.* 25/5/22.)

MANN, JOHN DIXON. Forensic medicine and toxicology. 6th edition. Revised by WILLIAM A. BREND. London 1922. pp. xii + 574. ill. 30s. net. (*Recd.* 13/6/22.)

PROCTER, HENRY RICHARDSON. The principles of leather manufacture. 2nd edition. London 1922. pp. viii + 688. ill. 32s. net. (*Recd.* 13/6/22.)

REVUE DE METALLURGIE. Fondée avec concours de la Société d'Encouragement pour l'Industrie Nationale, du Comité des Forges de France et de différents établissements industriels. Vol. 16, etc. Paris 1919 +. (*Reference.*)

TUTTON, ALFRED EDWIN HOWARD. Crystallography and practical crystal measurement. 2 vols. [2nd edition.] London 1922. pp. xviii + 746 + xiv, viii + 747 to 1446. ill. £5 net. (*Recd.* 13/6/22.)

VALENTA, EDUARD. Photographische Chemie und Chemikalienkunde mit Berücksichtigung der Bedürfnisse der graphischen Druckgewerbe. Part I. Anorganische Chemie. 2nd edition. Halle (Saale) 1921. pp. xvi + 282. ill. (*Recd.* 31/5/22.)

ZERR, GEORG, and RÜBENCAMP, ROB. Handbuch der Farbenfabrikation. 3rd edition. Berlin 1922. pp. xvi + 908. ill. (*Recd.* 23/5/22.)

III. *Pamphlets.*

BAILEY, E. H. S. A dietary study of some Kansas institutions under the control of the State Board of Health. Lawrence, Kansas 1921. pp. 70.

BRITISH COTTON INDUSTRY RESEARCH ASSOCIATION. The Shirley Institute. [Didsbury] 1922. pp. 22. ill.

CANADIAN INSTITUTE OF CHEMISTRY. [History, etc.] Montreal [1922]. pp. 56. ill.

INSTITUTE OF CHEMISTRY OF GREAT BRITAIN AND IRELAND. Lecture on chemists and the patent laws. By HORATIO BALLANTYNE. London 1922. pp. 31.

MOIR, JAMES. Colour and chemical constitution. Part XIII. The calculation of the colour of the monocyclic dyes. Part XIV. The calculation of the colour of the dicyclic dyes. (From the *Trans. Roy. Soc. S. Africa*, 1921, 10.)

SCIENTIFIC AND INDUSTRIAL RESEARCH, Department of. Bulletin No. 6. On the electro-deposition of iron. With an appendix containing a bibliography on the subject. By WILLIAM EARL HUGHES. London 1922. pp. iv + 50. ill.

— Deterioration of structures of timber, metal, and concrete exposed to the action of sea-water. Second (interim) report of the Committee of the Institution of Civil Engineers. Edited by P. M. CROSTHWAITE and GILBERT R. REDGRAVE. London 1922. pp. 57.

PROCEEDINGS

OF THE

CHEMICAL SOCIETY.

Ordinary Scientific Meeting, Thursday, October 5th, 1922, at 8 p.m., Sir JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

Reference was made to the loss sustained by the Society, through death, of the following Fellows :

	Elected.	Died.
Frederick Bedford	June 20th, 1901.	July 13th.
Frederick Douglas Brown	Mar. 20th, 1876.	Aug. 2nd.
William Kellner	Mar. 19th, 1874.	Sep. 12th.
George William MacDonald ...	Dec. 6th, 1894.	Aug. 7th.
Alfred Ernest Moore	June 17th, 1903.	Mar. 1st.
Arthur Faraday Price	May 4th, 1882.	Mar. 17th.
Alexander Smith	April 17th, 1890.	Sep. 8th.
Lionel William Stansell	Feb. 3rd, 1887.	May 22nd.
James Stenhouse	May 5th, 1892.	Mar. 18th.
Matthew Carrington Sykes	Feb. 19th, 1891.	Aug. 25th.
Jokichi Takaminé	Feb. 3rd, 1887.	July 22nd.
Stanley Goulden Thompson ...	May 1st, 1919.	Feb. 23rd.
James Waterhouse	Feb. 7th, 1901.	Sep. 28th.

In announcing the resignation of Dr. M. O. Forster from the office of Treasurer, the President referred to the many valuable services Dr. Forster had rendered to the Society during the past twenty years. His resignation was received with profound regret and his presence at the meetings would be greatly missed, and the President wished him, in the name of the Society, every success in the important work he was about to undertake as Director of the Institute of Science at Bangalore. Dr. Forster briefly responded.

It was stated that the Council had nominated Professor J. F. Thorpe, C.B.E., F.R.S., to fill the office of Treasurer until the date of the next Annual General Meeting, and that Dr. J. T. Hewitt, F.R.S., had been nominated to fill, for the same period, the vacancy amongst the Vice-Presidents caused by Professor Thorpe's appointment.

The President further announced that some time ago the American Chemical Society submitted to the Council certain proposals according to which we should be associated with them in the future conduct and control of the *Journal of Physical Chemistry*. Advantage was taken of the recent presence in this country of Dr. Parsons, Secretary of the American Chemical Society, to carry through the negotiations. The Council has to report that a satisfactory issue has been reached, the scheme having been widened to admit of the association of the Faraday Society with the enterprise. From some future date, probably January 1924, the *Journal of Physical Chemistry* will be accordingly issued under the auspices of the American Chemical Society, the Chemical Society, and the Faraday Society. The editorial board will consist of four American and four British representatives, together with an Editor-in-Chief appointed by these eight.

The following were formally admitted Fellows of the Chemical Society: J. P. Levy, F. M. Brewer, V. E. S. Swainson, and R. S. Colborne.

Certificates were read for the first time in favour of :

Robert Allanson, 10, Grosvenor Drive, New Brighton, Wallasey.
 Thomas Cuthbert Allchin, M.B.E., E. C. Powder Co., Ltd., Longfield, Kent.
 James Wilson Armit, M.A., The Manse, Monikie, Dundee.
 Menghraj Choithram Bathija, B.A., Fitzwilliam Hall, Cambridge.
 Donald Charles Broome, Disgwylfa, Penmaenmawr, Carnarvonshire.
 Walter Ferguson Buist, B.Sc., 24, New Hall Lane, Clubmoor, Liverpool.
 Stanley Robert Bult, 63, Addiscombe Road, Croydon.
 Arthur Josiah Burford, 419, Edgware Road, Maida Vale, W. 2.
 Katragadda Seshachalam Choudary, M.A., M.Sc., 71, Coral Merchant Street, George Town, Madras, India.
 Elliott Martin Christie, M.Sc., 37, Lynwood Avenue, Dunedin, New Zealand.
 Frederick Stanley Clarke, 64, Golders Manor Drive, N.W. 11.
 Sydney George Cavigan, 22, Walker Street, Burton-on-Trent.
 Edward Dance, B.Sc., Castle Inn, Sandal, Wakefield.
 • Milton Dewhurst, B.Sc., A.I.C., 1280, Warwick Road, Acocks Green, Birmingham.
 Samuel Frederick Ford, 65, Abbots Road, Southall.
 John Masson Gulland, B.Sc., 11, Chester Street, Edinburgh.
 Herbert Haigh, B.Sc., 211, Park Road, Barnsley.
 Thomas Edward Hubbard, 9, Cunliffe Street, Pendleton, Manchester.
 James Ernest Humphries, B.A., M.Sc., 29, St. John's Road, Newport, Mon.
 John Huntley, The Villa, Burstwick, nr. Hull.
 Sydney Alfred Hurren, M.C., 39, Upper Marylebone Street, W. 1.
 Habib Iskander, Heliopolis, Cairo, Egypt.
 Edgar Jones, 167, Upper Dale Road, Derby.
 Jan Kalf, D.Chem., Nieuwe Prinsengracht 126, Amsterdam.
 Shankar Nath Kapur, B.Sc., 11, George Square, Edinburgh.
 William Dawson Lloyd, 42, Tavistock Road, Westbourne Park, W. 11.
 Philip Patrick Lynch, B.Sc., Church Street, Timaru, New Zealand.

- Francis Geoffrey Macalpine, B.Sc., Broad Oak, Accrington.
 Frederick Henry McDowall, M.Sc., Knox College, Dunedin, New Zealand.
 Kenneth McKenzie, Endrik, Hamilton Road, Cambuslang, Glasgow.
 Archibald Roy Hammond Martin, B.A., 8, Kingsland Road, Worthing.
 Dina Nath, c/o P. Lal, Esq., Vakil, Jullundhur City, Punjab, India.
 Torkild Haastrup Nielsen, B.Sc., Siam Cement Co., Ltd., Bangsue, Siam.
 Hidejiro Nishikawa, 3, Hachimancho, Asakusa, Tokyo, Japan.
 Joseph Edwards Pooley, 1, Margaretting Road, Manor Park, E. 12.
 Nagendra Prasad, M.Sc., Ranchi, Behar & Orissa, India.
 Edmund Brydges Rudhall Prideaux, M.A., D.Sc., F.I.C., 6, Arboretum Street, Nottingham.
 James Robert Pound, M.Sc., 601a, Dana Street, Ballarat, Victoria, Australia.
 Alfred Norman Douglas Pullen, The Larches, Greenham Common, Newbury.
 Thomas Abraham Robertson, Sassafras, Rickard Road, Bankstown, N.S.W.
 Burke Rushworth, 4, West Park Terrace, Healey, Batley, Yorks.
 Gopal Singh, M.Sc., The Brij, Gawalmaudi, Lahore, India.
 George Victor Smith, A.R.C.S., Westleigh, Eggington Road, Wollaston, Wellingborough.
 Francis Wilbert Stoye, M.Sc., 37, Kansas Avenue, Belfast.
 Leslie Stuart Theobald, A.R.C.S., 3, Ennismore Avenue, Chiswick, W. 4.
 William Thompson, B.Sc., Ravensworth, Cockton Hill, Bishop Auckland.
 Laurence Barnett Timmis, M.Sc.Tech., A.I.C., 19, Trinity Marsh Lane, Waltham Cross.
 Richard Truszkowski, B.Sc., Koszykowa 21, m-3, Warsaw, Poland.
 Humphrey Terence Viney, B.Sc., Bramerton, Blakebrook, Kidderminster.
 Eric Richard Boraman Woodward, B.Sc., The Vicarage, Herne Bay, Kent.
 William James Uglow Woolcock, C.B.E., M.P., 166, Piccadilly, W. 1.

The following papers were read :

- "Cupric tetrammine nitrite and the corrosion of copper by aqueous solutions of ammonia and of ammonium nitrate."
 By H. BASSETT and R. G. DURRANT.
 "The additive formation of four-membered rings. Part I. The synthesis and resolution of some derivatives of 1 : 3-diazidine."
 By C. K. INGOLD and H. A. PIGGOTT.
 "Rates of reduction of nitro-groups by hydrogen in the presence of colloidal palladium, especially of nitrotoluene isomerides."
 By W. H. PATTERSON.
 "A determination of the relative proportions of the isomerides in nitrotoluene or toluidine." By W. H. PATTERSON.

Ordinary Scientific Meeting, Thursday, October 19th, 1922, at 8 p.m. Professor H. E. Armstrong, LL.D., F.R.S., Vice-President, in the Chair.

Reference was made to the loss sustained by the Society, through death, of :

	Elected.	Died.
Frederick Thomas Harry	June 20th, 1908.	Sept. 14th, 1922.
Edmond Herbert Grove-Hills	Feb. 15th, 1894.	Oct. 2nd, 1922.

Mr. C. D. Thomas was formally admitted a Fellow of the Chemical Society.

Certificates were read for the first time in favour of :

Leslie Bains, B.A., B.Sc., A.I.C., 5, Carlton Road, Stroud Green, N. 4.

George Joseph Burrows, B.Sc., Chemical Department, The University, Sydney, N.S.W.

John Hardwick Long Champion, B.Sc., 41, Triangle West, Clifton, Bristol.

Norman Evers, B.Sc., F.I.C., The Cottage, Salisbury Avenue, Cheam, Sutton.

Joshua William Farmery, M.A., A.I.C., 56, Sheen Road, Richmond, Surrey.

Philip Henry Gifford, B.Sc., Boys' High School, Salisbury, Rhodesia.

Joseph Keith Matthews, M.A., Cedric House, Rainhill, Liverpool.

John Ewart Moss, M.Sc., 298, Franklin Road, Kings Norton, Birmingham.

Harry Ward, M.Sc., A.I.C., 28, Commercial Street, E. 1.

The following papers were read :

"The polarity of double bonds. An extension of the theories of Lapworth and Robinson." By T. M. LOWRY.

"The theories of polarity in organic compounds." By J. KENNER.

"A model to illustrate the orbits of the electrons above the carbon atom." By W. E. GARNER.

NOTICES.

The next Ordinary Scientific Meeting takes place on November 2nd, at 8 p.m., and an Informal Meeting will be held on November 16th, after the conclusion of the formal business of the Ordinary Scientific Meeting at 8 p.m.

RESEARCH FUND.

A meeting of the Research Fund Committee will be held in December next. Application for Grants, to be made on forms obtainable from the Assistant Secretary, Chemical Society, Burlington House, W. 1, must be received on or before Friday, December 1st, 1922.

All persons who received Grants in December, 1921, or December of any previous year, whose accounts have not been closed by the Council, are reminded that reports must be returned by December 1st.

COLLECTIVE INDEX, VOL. VI (1913-1922).

Volume VI of the Collective Index will be published in two parts. Part I (Index to Authors) will be issued in 1923 and Part II (Index to Subjects) in 1924.

The price to Fellows who apply before *Christmas Day, 1922*, is £2 17s. post free. A remittance, made payable to Mr. S. E. Carr, must accompany the application in each case and should be sent to the Assistant Secretary, Chemical Society, Burlington House, W. 1. The price to Fellows who apply after December 25th, 1922, will be £3 7s.

Arrangements have been made by which cases in blue or maroon cloth for binding this Index in three volumes (Authors Index 1 vol. and Subject Index, 2 vols.) will be supplied for the sum of Seven Shillings. The binding will involve an additional charge of 2s. 3d. per volume. The total cost, therefore, of binding the Index in cloth will be 13s. 9d.

Fellows desiring to avail themselves of this offer should apply to Messrs. Gurney & Jackson, 33, Paternoster Row, London, E.C. 4, and also inform the Assistant Secretary, so that the necessary instructions may be given for the Index to be posted direct to the Publishers.

RUSSIAN FUND.

A further appeal has been issued on behalf of Russian Men of Science. In addition to donations of money to be devoted to purchasing requisite articles, gifts of clothing (which, if used, must be in good condition) and recent chemical literature will be welcome.

The Chemical Society will act as the receiving depôt, and cheques (made payable to Mr. S. E. Carr and crossed "Russian Fund"), together with parcels of clothing, boots, books, etc., should be addressed to the Assistant Secretary, Chemical Society, Burlington House, W. 1.

List of Papers received between June 16th and October 19th, 1922 :

- " Experiments on the synthesis of the polyacetic acids of methane. Part VII. The preparation of methylidenetriacetic acid and of methanetetra-acetic acid." By C. K. INGOLD and L. C. NICKOLLS.
- " An instance illustrating the effect of attached groups on the ease of formation of the *cyclopentanone* ring." By J. P. C. CHANDRASENA and C. K. INGOLD.
- " The formation and stability of *spiro-compounds*. Part IX. The influence on stability of groups of high molecular weight." By S. F. BIRCH and J. F. THORPE.

- "The viscosity of cellulose. Part II. The lowering of the viscosity of cellulose by various reagents." By R. A. JOYNER.
- "The constitution of polysaccharides. Part V. The yield of glucose from cotton cellulose." By J. C. IRVINE and E. L. HIRST.
- "The action of solvents on coal. Part I." By S. R. ILLINGWORTH.
- "Studies in the glucosides. Part I. The constitution of indican." By A. K. MACBETH and J. PRYDE.
- "The alkylhydrazones." By O. L. BRADY and G. P. MCHUGH.
- "The formation of bromine derivatives of carbon without the production of hydrogen bromide." By B. M. GUPTA and J. F. THORPE.
- "Limits for the propagation of flame at various temperatures in mixtures of ammonia with air and oxygen." By A. G. WHITE.
- "The molecular scattering of light in normal pentane." By R. VENKATESWARAN.
- "Preparation and reactions of stannous oxide and stannous hydroxides." By F. W. BURY and J. R. PARTINGTON.
- "Investigation of antimony metal from different sources. Indication of isotopes." By S. D. MUZAFFAR.
- "The catalytic decomposition of formic acid on surfaces of platinum and silver." By H. C. TINGEY and C. N. HINSHELWOOD.
- "Inorganic complex salts. Crystallographic and optical study. Part I." By I. E. KNAGGS.
- "The formation of derivatives of tetrahydronaphthalene from γ -phenyl fatty acids. Part II." By A. STEVENSON and J. F. THORPE.
- "N-Oximino-ethers. Part II. N-Aryl ethers of 2:4- and 2:6-dinitrobenzaloxime." By F. BARROW, E. D. GRIFFITHS, and E. BLOOM.
- "The nitro- and amino-derivatives of *o*- and *p*-methoxybenzoic acid and of α - and β -methoxynaphthoic acid." By V. FROELICHER and J. B. COHEN.
- "The addition of ethyl sodiocyanoacetate and ethyl sodiomalonate to ethyl muconate." By E. H. FARMER.
- "A synthesis of pyrylium salts of anthocyanidin type." By D. D. PRATT and R. ROBINSON.
- "The action of sodium methoxide and its homologues on benzophenone chloride and benzylidene chloride. Part III." By J. E. MACKENZIE.
- "On diacetylacetone." By J. N. COLLIE and A. A. B. REILLY.
- "The explosion of acetylene and nitrogen. Part II." By W. E. GARNER and K. MATSUNO.

- "The applicability of a linear poisoning law to the catalytic decomposition of hydrogen peroxide." By E. B. MAXTED.
- "Molecular weights in alcoholic solutions from the elevation of the flash point." By R. WRIGHT.
- "Selective solubility shown by aqueous alcohol when acting as a solvent." By R. WRIGHT.
- "Attempts to prepare red sulphide dyes. Part II. Mercaptan derivatives of azo-dyes." By E. R. WATSON and S. DUTT.
- "The isolation of muscarine, the potent principle of *Amanita muscaria*." By H. KING.
- "Researches on residual affinity and co-ordination. Part IX. Cobaltammine salts of the nitro-dyes." By G. T. MORGAN and H. J. S. KING.
- "The mobility of symmetrical triad systems. Part I. The conditions which confer mobility on symmetrical open-chain systems terminated by phenyl groups." By C. K. INGOLD and H. A. PIGGOTT.
- "Cevadine. Part I." By A. K. MACBETH and R. ROBINSON.
- "Cyclic addition. Part I. The occurrence and reversibility of reactions in which cyclic structures are formed by the direct union of unsaturated linkages. Part II. The reaction between azomethines." By C. K. INGOLD and H. A. PIGGOTT.
- "*iso*Propylstannonic acid and some of its derivatives." By J. G. F. DRUCE.
- "Syntheses of alkylidenecyanoacetic acids and of substituted succinic acids. Part I. Acids containing aromatic residues." By A. LAPWORTH and J. A. McRAE.
- "On crystalline glucose-ammonia and *isoglucosamine*." By A. R. LING and D. R. NANJ1.
- "The autoreduction of sulphurous acid." By G. M. BENNETT.
- "Researches on the conditions of formation of rings attached to the *o*-, *m*-, and *p*-positions of the benzene nucleus. Part I. The action of sodium on *o*-phenylenediacetic ester." By W. H. PERKIN and A. F. TITLEY.
- "A method for the preparation of $\alpha\beta$ -trialkylated glutaric acids." By G. A. R. KON and J. F. THORPE.
- "The bromine compounds of phenanthrene. Part II." By H. HENSTOCK.
- "The solubility of phenanthrene in various organic solvents." By H. HENSTOCK.
- "The sorption of liquids by charcoal." By J. DRIVER and J. B. FIRTH.
- "The equilibria in aqueous solutions of the alkali metal bisulphites." By E. C. C. BALY and R. A. BAILEY.

- "Studies in the anthracene series. Part III." By E. DE B. BARNETT, J. W. COOK, and H. H. GRAINGER.
- "The solubilities of the alkaline formates and acetates in water." By N. V. SIDGWICK and J. A. H. R. GENTLE.
- "The solubility of the alkaline salts of benzoic and the hydroxybenzoic acids in water." By N. V. SIDGWICK and E. K. EWBANK.
- "The mechanism of the formation of benzoylbenzoin by treatment of benzoylmandelonitrile with an alcoholic solution of sodium ethoxide." By H. GREENE and R. ROBINSON.
- "The oxime of mesoxamide (isonitrosomalonomide) and some allied compounds. Part IV. The ethers of isonitrosomalonanilide, and isonitrosomalondimethylamide, and isonitrosomalondibenzylamide." By A. G. RENDALL and M. A. WHITELEY.
- "Researches on residual affinity and co-ordination. Part X. Salicylatotetramminocobaltic salts and the constitution of oxonium compounds." By G. T. MORGAN and J. D. MAIN SMITH.
- "The cyanine dyes. Part VI. Dyes containing a quinoline and a benzothiazole nucleus. The thioisocyanines." By W. T. K. BRAUNHOLTZ and W. H. MILLS.
- "The isomerism of the oximes. Part X. Cinnam- and nitrocinnam-aldoximes." By O. L. BRADY and C. D. THOMAS.
- "The problem of substitution in the benzene nucleus and the Thomson-Lewis-Langmuir theory of co-valence." By R. FRASER and J. E. HUMPHRIES.
- "The constitution of soap solutions. Cetylsulphonic acid and other sulphonates." By M. H. NORRIS.
- "Physical chemistry of the oxides of lead. Part VI. The anodic behaviour of lead and lead peroxide." By S. GLASSTONE.
- "Use of paraformaldehyde (gaseous formaldehyde) and paraldehyde (gaseous acetaldehyde) in the synthesis of alcohols." By C. E. WOOD and F. SCARF.
- "Rotatory dispersion of the esters of lactic acid. Normal esters. Part I." By C. E. WOOD, J. E. SUCH, and F. SCARF.
- "The solubility and volatility of 1:3:5-dinitrophenol." By N. V. SIDGWICK and T. W. J. TAYLOR.
- "The labile nature of the halogen atom in organic compounds. Part V. The action of hydrazine on the halogen derivatives of some esters and cyclohexanes." By E. L. HIRST and A. K. MACBETH.
- "Esters of the hydroxyalkylarylamines. Part I. Acid sulphuric esters of the simple monohydroxyethylarylamines." By K. H. SAUNDERS.

- "Harmine and harmaline. Part VI. The synthesis of *n*-methyl-tetrahydroharmine and the constitution of harmaline and of the alkylated harmines." By W. O. KERMACK, W. H. PERKIN, and R. ROBINSON.
- "Note on the decomposition of the dithiocarbazinates." By S. M. LOSANITCH.
- "The interaction of aldehydes and ketones with thiocarbamides in the presence of acids. Part II." By J. TAYLOR.
- "The preparation of methylguanidine, and of $\beta\beta$ -dimethylguanidine by the interaction of dicyanodiamide, and methylammonium and dimethylammonium chlorides respectively." By E. A. WERNER and J. BELL.
- "Adsorption and catalysis in Fuller's earth." By E. K. RIDEAL and W. THOMAS.
- "Quantitative reduction by hydriodic acid of halogenated malonyl derivatives. Part II. The *s*-tetra-substituted amides of bromo- and chloro-malonic acid." By R. W. WEST.
- "Amylases of the cereal grains—the 'insoluble' amylase of barley." By J. L. BAKER and H. F. E. HULTON.
- "Reactivity of hydrogen atoms in methylene groups." By T. H. READE.
- "The condensation of the aromatic *o*-amino-sulphonic acids with isocyanic acid." By J. R. SCOTT and J. B. COHEN.
- "Monothioethylene glycol." By G. M. BENNETT.
- "The β -chlorovinylarsines." By F. G. MANN and W. J. POPE.
- "Piperitone. Part IV. The interaction of *dl*-piperitone and semicarbazide, and the isolation of pure *dl*-piperitone." By J. READ and H. G. SMITH.
- "The chlorination and bromination of the toluic acids and the preparation of the phthalaldehydic acids." By W. DAVIES and W. H. PERKIN.
- "The isomerism of metallic oxides. Part I. Lead monoxide." By M. P. APPLEBEY and R. D. REID.
- "The action of the chlorides of phosphorous on chlorodimethyldihydroresorcin." By L. E. HINKEL and W. D. WILLIAMS.
- "On the movements of explosion waves in tubes of different shapes." By C. CAMPBELL.
- "The chlorination of benzoyl chloride." By E. HOPE and G. C. RILEY.
- "Ethyl α -cyano- β -methyl glutaconate and its methyl homologues." By E. HOPE.
- "The addition of hydrogen cyanide to derivatives of glutaconic acid. Part I. The addition of hydrogen cyanide to ethyl- α -

- cyano- β -methyl glutaconate and its homologues." By E. HOPE and W. SHELDON.
- "The velocity of reaction in mixed solvents. Part III. The influence of temperature on the velocity of saponification of esters." By H. MCCOMBIE, H. A. SCARBOROUGH, and R. H. SETTLE.
- "The action of ferrous chloride on some organic hydrochlorides." By W. M. CUMMING.
- "The form of the vapour-pressure curve at high temperature. Part I. The curve for lead." By C. K. INGOLD.
- "The optical rotations of the sugars." By J. G. MALTBY.
- "Heteromorphism of calcium sulphate. Alabaster and its synthesis." By M. COPISAROW.
- "Heteromorphism of calcium carbonate. Marble: synthetic and metamorphic." By M. COPISAROW.
- "Researches on residual affinity and co-ordination. Part IX. Interaction of selenium tetrachloride and β -diketones." By G. T. MORGAN, H. D. K. DREW, and T. V. BARKER.
- "The ignition of gases. Part II. Ignition by a heated surface. Mixtures of methane and air." By W. MASON and R. V. WHEELER.
- "On the interchange of alcohol radicles in esters. Part II." By A. SHIMOMURA and J. B. COHEN.
- "Note on the reduction of thorium oxide by metallic tungsten." THE RESEARCH STAFF OF THE GENERAL ELECTRIC CO., LTD., LONDON (C. J. SMITHELLS).
- " γ -Methylfructoside." By R. C. MENZIES.
- "The ultrafiltration of soap solutions: sodium oleate and potassium laurate." By J. W. MCBAIN and W. J. JENKINS.
- "Some new derivatives of methylstannonic acid; their bearing upon its constitution." By H. LAMBOURNE.
- "Note on the hydrolysis of $\beta\beta'$ -dichloroethyl sulphide. Part I. Synthesis of divinyl sulphide and the preparation of a non-vesicant isomer of $\beta\beta'$ -dichloroethyl sulphide." By S. H. BALES and S. A. NICKELSON.
- "The quaternary salts of quinaldinic acid." By W. H. MILLS and F. M. HAMER.
- "A modified methyl-orange indicator." By K. C. D. HICKMAN and R. P. LINSTEAD.
- "The labile nature of the halogen atom in organic compounds. Part VI. The action of titanous chloride and of ammonia on representative halogen compounds." By I. A. BLACK, E. L. HIRST, and A. K. MACBETH.
- "The interaction of aniline and acrolein." By F. G. MANN.
- "*p*-Dithiobenzoic acid." By S. SMILES and D. C. HARRISON.

- "The reversibility of the reaction between nitrogen, carbon, and sodium carbonate." By C. K. INGOLD and D. WILSON.
- "Limits for the propagation of flame in vapour-air mixtures. Part II. Mixtures of more than one vapour and air at the ordinary temperature and pressure." By A. G. WHITE.
- "The constitution of carbamides. Part XIV. The decomposition of urea by sodium hypobromite in alkaline solution, and an improved procedure for the estimation of urea by this means." By E. A. WERNER.
- "The system chromium trioxide-sulphur trioxide-water." By L. F. GILBERT, H. BUCKLEY, and I. MASSON.
- "The constitution of acetone derivatives of glucose and fructose." By J. C. IRVINE and J. PATTERSON.
- "The formation and properties of dithio-ketones ($R_2C:S:S$) and dithio-ether ($R_2S:S$). Part III." By K. G. NAIK and M. D. AVASARE.
- "The constitution of the terpene present in the essential oil from *Andropogon Jwarancusa*, Jones." By J. L. SIMONSEN.
- "The action of bromine water on indene." By J. READ and E. HURST.
- "Note on the dilution of ethylenebromohydrin with water." By J. READ and G. J. BURROWS.
- "The solubilities of the aldehydo-benzoic acids." By N. V. SIDGWICK and H. CLAYTON.
- "The solubility of the chlorophenols." By N. V. SIDGWICK and S. L. TURNER.
- "The resolution of tropic acid and the stereochemical configuration of the cinchona alkaloids." By H. KING and A. D. PARKER.
- "The methylation of xylose." By E. L. HIRST and A. CARRUTHERS.
- "The specific heats of gases with special reference to hydrogen." By C. K. INGOLD and E. H. USHERWOOD.
- "Purification of methyl alcohol by means of sodium hypochlorite." By R. C. MENZIES.
- "Cupric tetrammine nitrite and the corrosion of copper by aqueous solutions of ammonia and of ammonium nitrate." By H. BASSETT and R. G. DURRANT.
- "The speed of the uniform movement of flame in mixtures of the paraffins with air." By W. MASON.
- "Mutarotation and pseudo-mutarotation in the glucosamine series." By J. C. IRVINE and J. C. EARL.
- "Salicylidene derivatives of *d*-glucosamine." By J. C. IRVINE and J. C. EARL.
- "Perhalides of the phenyltrimethylammonium series." By H. McCOMBIE and T. H. READE.

- "Note on the preparation of the sodium and potassium derivatives of phthalimide." By D. L. HAMMICK and G. H. LOCKET.
- "The additive formation of four-membered rings. Part I. The synthesis and resolution of some derivatives of tetrahydro-1:3-diazine." By C. K. INGOLD and H. A. PIGGOTT.
- "Derivatives of dithiophenyl-benzene." By S. SMILES and H. GRAHAM.
- "The products of fermentation of cellulose." By H. LANGWELL.
- "The labile nature of the halogen atom in organic compounds. Part VII. The absorption spectra of the halogen derivatives of some cyclic compounds and their bearing on the question of an oxygen-halogen linking." By H. GRAHAM and A. K. MACBETH.
- "The polarity of double bonds. An extension of the theories of Lapworth and Robinson." By T. M. LOWRY.
- "2:4-Dinitrobenzil." By G. B. BISHOP and O. L. BRADY.
- "Rates of reduction of nitro-groups by hydrogen in the presence of colloidal palladium, especially of nitrotoluene isomerides." By W. H. PATTERSON.
- "A determination of the relative proportions of the isomerides in nitrotoluene or toluidine." By W. H. PATTERSON.
- "The synthesis of *m*- α -benzbispyrrole derivatives." By W. DAVIES and E. H. C. HICKOX.
- "Viscosity of organic liquid mixtures and its relationship to chemical composition and vapour pressure and certain other physical properties." By M. A. SOOFI.
- "The system tin-arsenic." By Q. A. MANSURI.
- "On the preparation of glucinum sulphate and the system: ammonium sulphate-glucinum sulphate-water at 25° C." By H. T. S. BRITTON.
- "Aromatic sulphonic chlorides." By J. STEWART.
- "The oleo-resin of *Canarium muelieri*." By T. G. H. JONES and F. SMITH.
- "The conditions underlying the formation of unsaturated and cyclic compounds from halogenated open-chain derivatives. Part V. Products derived from α -halogenated β -methylglutaric acid." By C. K. INGOLD.
- "The behaviour of the stannic acids towards solutions of alkaline hydroxides." By G. E. COLLINS and J. K. WOOD.
- "A specific cellulose-fermenting organism." By H. LANGWELL.
- "The mechanism of the bromination of phenol in aqueous solution." By H. BAINES.
- "The theories of polarity in organic compounds." By J. KENNER.

- "The solubility and volatility of the nitrobenzaldehydes." By N. V. SIDGWICK and W. M. DASH.
- "The action of magnesium on symmetrical tetrachloro- and tetrabromo-tetramethylmethane." By A. FAIRBOURNE.
- "The triazo-group. Part XXII. Cinnamic acid chlorohydrin and its conversion into α -triazole- β -hydroxy- β -phenylpropionic acid." By M. O. FORSTER and W. B. SAVILLE.
- "A model to illustrate the orbits of the electrons about a carbon atom." By W. E. GARNER.
- "The labile nature of the halogen atom in organic compounds. Part VIII. The reduction of chloro- and bromo-trinitromethane and of tetranitromethane." By T. HENDERSON and A. K. MACBETH.
- "The condensation of 9:10-diaminophenanthrene with phenanthraquinone and other *o*-quinones." By S. DUTT and E. R. WATSON.
- "Studies in the β -phenylglutaric acid series. Part I. A possible basis of a new method for the determination of the relative distribution of the valencies of polyvalent elements." By J. G. JACKSON and J. KENNER.
- "A simplified method for the resolution into its optically active components of *sec*-octyl alcohol (methyl-*n*-hexylcarbinol)." By J. KENYON.
- "Electro-synthesis of azelaic and thapsic acids." By M. CARMICHAEL.
- "Photocatalysis. Part III. The photosynthesis of naturally occurring nitrogen compounds from carbon dioxide and ammonia." By E. C. C. BALY, I. M. HEILBRON, and H. J. STERN.
- "The preparation of potassium and sodium arylsulphoniodoamides." By E. ROBERTS.

ADDITIONS TO THE LIBRARY.

I. Donations.

BARKER, THOMAS VIPOND. Graphic and tabular methods in crystallography as the foundation of a new system of practice: with a multiple tangent table and a 5-figure table of natural cotangents. London [1922]. pp. xvi+152. ill. 14s. net. (*Recd.* 3/10/22.) From the Author.

BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE. A retrospect 1831—1921. By O. J. R. HOWARTH. London 1922. pp. viii+318. ill. (*Reference.*) From the British Association.

BRITISH ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE. Fourth report on colloid chemistry and its general and industrial applications. London 1922. pp. 382. 5s. 6d. net. (*Recd.* 15/9/22.)

From the Dept. of Scientific and Industrial Research.

CAEN, JOHN CANNELL. The manufacture of dyes. London 1922. pp. x+274. 12s. 6d. net. (*Recd.* 3/10/22.)

From Mrs. J. C. Cain.

DENHAM, HENRY GEORGE. An inorganic chemistry. London 1922. pp. viii+684. ill. 12s. 6d. net. (*Recd.* 5/7/22.)

From the Author.

FOOTE, PAUL D., and MOHLER, F. L. The origin of spectra. New York 1922. pp. 250. ill. \$4.50. (*Recd.* 30/9/22.)

From the Publishers: The Chemical Catalog Company, Inc.

[FRITZ, JOHN.] The John Fritz Medal. [2nd edition. New York 1922.] pp. 128. ill. (*Recd.* 23/6/22.)

From the Board of Award.

GRIFFITHS, HUGH. Materials of chemical plant construction—non-metals. London 1922. pp. 64. ill. 3s. net. (*Recd.* 4/9/22.)

—. The general principles of chemical engineering design. London 1922. pp. 64. ill. 3s. net. (*Recd.* 4/9/22.)

From the Publishers: Messrs. Benn Brothers.

HALE, ARTHUR JAMES. Modern chemistry, pure and applied. Vol. V. London [1922]. pp. viii+280. ill. 16s. net. (*Recd.* 7/9/22.)

From the Publishers: Messrs. Virtue & Co.

HENRICH, FERDINAND. Theories of organic chemistry. Translated from the revised 4th German edition of 1921 by **TREAT BALDWIN JOHNSON** and **DOROTHY A. HAHN**. New York 1922. pp. xvi+604. ill. 30s. net. (*Recd.* 9/10/22.)

From the London Publishers: Messrs. Chapman & Hall.

HOLMES, HARRY N. Laboratory manual of colloid chemistry. New York 1922. pp. xii+128. 10s. net. (*Recd.* 21/8/22.)

From the London Publishers: Messrs. Chapman & Hall.

IMPERIAL INSTITUTE. Indian Trade Enquiry. Drugs and tanning materials. Reports on cinchona bark and myrobalans. London 1922. pp. x+60. 4s. net. (*Recd.* 28/7/22.)

From the Director.

INTERNATIONAL BREWERS' CONGRESS. Report of the proceedings of the Second Congress held at Chicago, October 18—21, 1911: with an account of the Second International Barley and Hop Prize Exhibition, and brief history of the American Exposition of brewing machinery, materials, and products. Vol. I. Chicago 1912. pp. 642. (*Reference.*)

From Mr. J. C. N. Eastick.

JOURNAL DE PHYSIQUE ET LE RADIUM. Vol. III., etc. Paris 1922+. (*For Circulation.*)

From the Faraday Society.

KOLBE, HERMANN [Editor]. Das chemische Laboratorium der Universität Marburg und die seit 1859 darin ausgeführten chemischen Untersuchungen nebst Ansichten und Erfahrungen über die Methode des chemischen Unterrichts. Braunschweig 1865. pp. 524. ill. (Recd. 21/8/22.) From Professor P. Phillips Bedson.

—. Das chemische Laboratorium der Universität Leipzig und die seit 1866 darin ausgeführten chemischen Untersuchungen. Braunschweig 1872. pp. xii+668. ill. (Recd. 21/8/22.)

From Professor P. Phillips Bedson.

KRAUS, CHARLES AUGUST. The properties of electrically conducting systems including electrolytes and metals. New York 1922. pp. 416. ill. \$4.50 net. (Recd. 21/8/22.)

From the Publishers: The Chemical Catalog Company, Inc.

LEWIS, WILLIAM. Commmercium philosophico-technicum; or, the philosophical commerce of arts: designed as an attempt to improve arts, trades, and manufactures. London 1763. pp. iv+xviii+x+646+xiv. ill. (Recd. 30/9/22.)

From Mr. E. Gardner.

LIND, SAMUEL COLVILLE. The chemical effects of alpha particles and electrons. New York 1921. pp. 182. ill. \$3 net. (Recd. 21/8/22.)

From the Publishers: The Chemical Catalog Company, Inc.

LOEB, JACQUES. Proteins and the theory of colloidal behaviour. New York 1922. pp. xii+292. ill. 15s. net. (Recd. 6/7/22.)

From the Publishers: The McGraw Hill Book Company.

MALAN, H. L., and ROBINSON, A. I. The weighing and measuring of chemical substances. London 1922. pp. 64. ill. 3s. net. (Recd. 4/9/22.) From the Publishers: Messrs. Benn Brothers.

MINISTRY OF MUNITIONS and DEPARTMENT OF SCIENTIFIC AND INDUSTRIAL RESEARCH. Technical records of explosives supply. No. 7. Manufacture of nitric acid from nitre and sulphuric acid. London 1922. pp. vi+86. ill. 10s. 6d. net. (Recd. 20/7/22.)

From the Department.

MITCHELL, CHARLES AINSWORTH. Documents and their scientific examination, with especial reference to the chemistry involved in cases of suspected forgery, investigation of disputed documents, handwriting, etc. London 1922. pp. xii+216. ill. 10s. 6d. net. (Recd. 21/8/22.) From the Publishers: Messrs. C. Griffin & Co.

OIL AND COLOUR CHEMISTS' ASSOCIATION. Journal. Vol. V. etc. London 1922+. (Reference.) From the Association.

PARTINGTON, JAMES RIDDICK, and PARKER, LESLIE HENRY. The nitrogen industry. London 1922. pp. xvi+336. ill. 21s. net. (Recd. 4/10/22.)

From the Publishers: Messrs. Constable & Co.

POLSKIEGO TOWARZYSTWA CHEMICZNEGO. *Roczniki Chemji*. Vol. I. Warszawa 1921. (*Reference*.) From the Society.

SCHLESISCHEN KOHLENFORSCHUNGSINSTITUT DER KAISER-WILHELM-GESELLSCHAFT IN Breslau. *Mitteilungen*. Vol. I. Berlin 1922. pp. viii+180. ill. (*Recd.* 21/8/22.)

From the Publishers: Gebrüder Borntraeger.

SCIENTIFIC AND INDUSTRIAL RESEARCH, Department of. *Food Investigation Board*. Special Report No. 7. The preservation of food by freezing with special reference to fish and meat: a study in general physiology. By WALTER STILES. London 1922. pp. 186. ill. (*Reference*.) 10s. net. From the Department.

SEARS, GEORGE W. A systematic qualitative chemical analysis: a theoretical and practical study of analytical reactions of the more common ions of inorganic substances. New York 1922. pp. vi+120. 8s. 6d. net. (*Recd.* 21/8/22.)

From the London Publishers: Messrs. Chapman & Hall.

SIMMONDS, CHARLES. Alcohol in commerce and industry. London 1922. pp. xii+120. ill. 3s. net. (*Recd.* 15/9/22.)

From the Publishers: Sir Isaac Pitman & Sons.

SINNATT, FRANK STURDY, and others. Coal and allied subjects: a compendium of the first ten bulletins issued by the Lancashire and Cheshire Coal Research Association. [1918—1921.] London [1922]. pp. viii+205. ill. 15s. net. (*Recd.* 9/10/22.)

From the Publishers: Messrs. H. F. & G. Witherby.

SWARTS, FRED. Cours de chimie organique. 3rd edition. Bruxelles 1921. pp. vi+674. ill. (*Recd.* 3/10/22.)

——. Cours de chimie inorganique. 3rd edition. Bruxelles 1922. pp. vi+734. ill. (*Recd.* 3/10/22.)

From the Publisher: M. Maurice Lamertin.

SWINDIN, NORMAN. Pumping in the chemical works. London 1922. pp. 80. ill. 3s. net. (*Recd.* 4/9/22.)

——. The flow of liquids in pipes. London 1922. pp. 64. ill. 3s. net. (*Recd.* 4/9/22.)

From the Publishers: Messrs. Benh Brothers.

TAYLOR, HUGH STOTT. Industrial hydrogen. New York 1921. pp. 210. ill. \$3.50 net. (*Recd.* 21/8/22.)

From the Publishers: The Chemical Catalog Company, Inc.

TÔHOKU IMPERIAL UNIVERSITY. Technology reports. Vol. I., etc. Sendai 1920+. ill. (*Reference*.) From the University.

TSAO, HUI CHUN, and ADOLPH, WILLIAM H. A text-book of general chemistry. Based on McPherson and Henderson: A course in general chemistry. Shanghai 1922. pp. xxvi+653. [In Chinese.] From the Authors.

PROCEEDINGS

OF THE

CHEMICAL SOCIETY.

Ordinary Scientific Meeting, held in the Lecture Hall of the Institution of Mechanical Engineers, on Thursday, October 26th, 1922, at 8 p.m., Sir JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

Sir William H. Bragg, K.B.E., F.R.S., delivered his Lecture entitled: "The Significance of Crystal Structure." A vote of thanks to the Lecturer, proposed by Professor J. Norman Collie, and seconded by Dr. A. E. H. Tutton, was carried with acclamation, acknowledgment being made by Sir William Bragg.

Ordinary Scientific Meeting, Thursday, November 2nd, 1922, at 8 p.m., Professor J. F. THORPE, C.B.E., D.Sc., F.R.S., Treasurer, in the Chair.

Reference was made to the great loss the Society had sustained by the death, on October 28th, of Professor Alexander Crum Brown, who was elected a Fellow on December 5th, 1867. The Chairman referred to Professor Crum Brown's long connexion with the Society, of which he had served as President for two years and as Vice-President for thirty-three years, and to his many valuable contributions to the science of chemistry.

The following were formally admitted Fellows of the Chemical Society: R. A. Mallet and Owen Owen.

Certificates were read for the first time in favour of:

Leopold Louis Bacarese, 2, Stirling Road, Clapham Road, S.W. 9.
Nicholas Timothy Belaiew, C.B., 33, Elvaston Place, Queen's Gate, S.W. 7.
James Bell, 30, Trinity College, Dublin.
John Arnold Cranston, D.Sc., 14, Fernleigh Road, Newlands, Glasgow.
Hugh John Curnow, 22, Halfway Street, Sidcup.
Richard Victor Eaton, B.Sc., A.R.C.S., Lakeside, Seldown, Poole.
Roland Hall Griffith, B.A., 14, Bigwood Avenue, Hove.
Lucy Higginbotham, M.Sc., 57, Wilbraham Road, Chorlton-cum-Hardy, Manchester.
Svend Hald Lassen, Briarwood, Manorgate Road, Kingston-on-Thames.

Reginald Arthur McNicol, M.Sc., Birnam, Rotherfield Road, Carshalton.
 Thomas Malkin, B.Sc., A.I.C., Golden Square, Warrington.
 Thomas Trevor Potts, 60, Christ Church Square, Macclesfield.
 George Clifford Riley, B.A., B.Sc., 15, Wellington Square, Oxford.
 Oscar Walter Snow, B.A., Graeme's Bank, Hatch End.
 Thomas John Wall, B.A., Northcote, Alexandra Road, Malvern.

The following papers were read :

- "The solubility and volatility of the nitrobenzaldehydes." By N. V. SIDGWICK and W. M. DASH.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XIII. The spatial configuration of the unbranched aliphatic chain." By R. H. PICKARD, J. KENYON, and H. HUNTER.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XIV. The normal aliphatic ethers of *d*- β -octanol." By J. KENYON and R. A. McNICOL.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XV. The normal aliphatic ethers of *d*-methylbenzylcarbinol." By H. PHILLIPS.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XVI. A new type of Walden inversion." By H. PHILLIPS.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XVII. The di-*d*- β -octyl esters of the acids of the general formula $(CH_2)_n(CO_2H)_2$." By L. HALL.
- "Orientation of the 1 : 4- and 1 : 5-dimethylglyoxalines. Mode of fission of 5-aminoglyoxalines." By F. L. PYMAN.
- "Bromo-derivatives of 2-methylglyoxaline." By L. LIGHT and F. L. PYMAN.
- "2 : 5-Iminodihydro-1 : 2 : 3-triazole. Part I. Constitution of Dimroth's 5-anilidotriazole." By P. K. DUTT.

Unveiling Ceremony of the War Memorial to Lt.-Colonel E. F. Harrison, C.M.G., and other Fellows of the Chemical Society on Thursday, November 16th, 1922, at 3 p.m.

A report of the proceedings of the meeting will appear in the December issue of the Journal.

Ordinary Scientific Meeting, Thursday, November 16th, 1922, at 8 p.m., Professor J. C. PHILIP, O.B.E., D.Sc., F.R.S., Secretary, in the Chair.

Reference was made to the loss sustained by the Society, through death, of the following Fellows :

	Elected.	Died.
Edward Henry Grossmann	Dec. 5th, 1895.	Aug. 29th.
Leo Tschugaeff	Dec. 1st, 1910.	Sept. 26th.

Messrs. J. R. Green and T. Iredale were formally admitted Fellows of the Chemical Society.

Certificates were read for the first time in favour of :

Kenneth Claude Bailey, M.A., 137, South Circular Road, Dublin.
 Katherine Hope Coward, M.Sc., 14, Parliament Hill, Hampstead, N.W. 3.
 Allan Robert Earle, B.A., 90, Shadwell Road, Portsmouth.
 Samuel Gray Fenton, 27, Somerset Road, Harrow.
 Phillip Marcus Fisk, 92, Aldborough Road, Seven Kings, Ilford.
 Thomas Graham, 47, Canada Road, Cardiff.
 Leslie Frank Hewitt, B.Sc., A.I.C., 8, Maze Road, Kew, Surrey.
 John Ingham, 19, Greyhound Road, Tottenham, N. 17.
 Wilfred Herbert Linnell, B.Sc., A.I.C., 12, Highbury, Monkseaton, Whitley Bay.
 Ramprasad Pitamberdas Mahta, M.A., Organic Chemistry Dept., University College, Gower Street, W.C. 1.
 Jocelyn Patterson, Ph.D., M.Sc., 106, Market Street, St. Andrews.
 Tara Chand Rana, M.Sc., III Bridge, Srinagar, Kashmir, India.
 John Baldwin Shoesmith, M.Sc., 10, Spottiswoode Road, Edinburgh.
 Reginald Ernest Tebbit, 120, Lordship Lane, E. Dulwich, S.E. 22.
 Eric Wilfred Tong, B.Sc., A.R.C.S., Hollingbourne, Maidstone.
 Thomas Arthur Williams, Ridgeway House, Mill Hill, N.W. 7.

A Certificate has been authorised by the Council for presentation to ballot under Bye-Law I (2) in favour of: Babu Lall Govila, Central Polytechnic Institute, Gwalior.

The meeting was then adjourned and the Informal Meeting declared open.

List of Papers received between October 20th and November 16th, 1922 :

- "Investigations on the dependence of rotatory power on chemical constitution. Part XIII. The spatial configuration of the unbranched aliphatic chain." By R. H. PICKARD, J. KENYON, and H. HUNTER.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XIV. The normal aliphatic ethers of *d*- β -octanol." By J. KENYON and R. A. McNICOL.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XV. The normal aliphatic ethers of *d*-methyl benzylcarbinol." By H. PHILLIPS.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XVI. A new type of Walden inversion." By H. PHILLIPS.

- "Investigations on the dependence of rotatory power on chemical constitution. Part XVII. The di-*d*- β -octyl esters of the acids of the general formula $(\text{CH}_2)_n(\text{CO}_2\text{H})_2$." By L. HALL.
- "Orientation of the 1:4- and 1:5-dimethylglyoxalines. Mode of fission of 5-aminoglyoxalines." By F. L. PYMAN.
- "Bromo-derivatives of 2-methylglyoxaline." By L. LIGHT and F. L. PYMAN.
- "The oxidation of cane sugar by nitric acid." By F. D. CHATTAWAY and H. J. HARRIS.
- "Castelamarin—a bitter principle from *Castela Nicholsoni*." By L. P. BOSMAN.
- "Catalytic hydrogenation with nickel. Factors determining catalytic activity." By R. THOMAS.
- "2:5-Iminodihydro-1:2:3-triazole. Part I. Constitution of Dimroth's 5-anilidotriazole." By P. K. DUTT.
- "The action of alcohol on the sulphates of ammonium." By H. B. DUNNICLIFF.
- "Refraction and dispersion of potassium vapour." By A. L. NARAYAN and G. SUBRAHMANYAM.
- "The interaction of diazonium salts and phenols." By F. D. CHATTAWAY and H. R. HILL.
- "The relation between the crystal structure and constitution of carbon compounds. Part I. Compounds of the type CX_4 ." By I. E. KNAGGS.
- "Binary critical solution temperatures as criteria of the purity of acetic acid." By D. C. JONES.
- "Ternary critical solution temperatures as a criterion of liquid purity." By D. C. JONES.
- "Syntheses of alkylidenecyanoacetic acids and of substituted succinic acids. Part II. Preparation of acids containing saturated aliphatic residues and the constitution of the aliphatic alkylidenecyanoacetic esters." By A. LAPWORTH and J. A. McRAE.
- "Formation of γ -alkylidene derivatives from ethylidenemalonic ester." By A. LAPWORTH and L. HIGGINBOTHAM.
- "Resorcinolphenylsuccinein." By A. LAPWORTH and J. A. McRAE.
- "The preparation of α -trioxymethylene and a new polymer of formaldehyde." By D. L. HAMMICK and A. R. BOEREE.
- "The isoelectric condition of gelatin." By S. O. RAWLING and W. CLARK.
- "The nitration of 3-chloroacenaphthene." By G. FARNELL.
- "The diazo-reaction in the carbazole series. Carbazole-3-diazoimine and -3-diazonium salts." By G. T. MORGAN and H. N. READ.

- "The oxidation of sabinene with chromyl chloride." By G. G. HENDERSON, J. M. ROBERTSON, and D. C. BROWN.
- "Scattering of light by CO_2 and N_2O , and by some organic vapours." By A. L. NARAYAN.
- "1:3:4:6-Tetramethyl fructose." By J. C. IRVINE and J. PATTERSON.
- "Researches on residual affinity and co-ordination. Part XII. Interactions of metallic salts and dimethyldithiolethylene." By G. T. MORGAN and W. LEDBURY.
- "Researches on residual affinity and co-ordination. Part XIII. Cobaltammine and ferric lakes of dinitrosoresorcinol." By G. T. MORGAN and J. E. MOSS.
- "Researches on residual affinity and co-ordination. Part XIV. Cobaltammine and cobaltic lakes of the azo-salicylic acids." By G. T. MORGAN and J. D. M. SMITH.
- "Boundary lubrication and chemical constitution. The optically active carbinols of the formula $\text{C}_2\text{H}_5\cdot\text{CH}(\text{OH})\cdot\text{C}_n\text{H}_{2n+1}$." By I. DOUBLEDAY.
- "The velocity of reaction in mixed solvents. Part IV. The influence of the base on the velocity of saponification of esters." By A. CASHMORE, H. McCOMBIE, and H. A. SCARBOROUGH.
- "The adsorption of radium-B and radium-C by ferric hydroxide." By J. A. CRANSTON.
- "The hydrolysis of platinum salts. Part II. Potassium platini-chloride." By E. H. ARCHIBALD and W. A. GALE.
- "On the reactivity of methyl groups in heterocyclic bases." By W. H. MILLS and J. L. B. SMITH.
- "An examination of the reputed antimonious hydroxides." By C. LEA and J. K. WOOD.
- "The adsorption of stannous chloride by stannic acid." By G. E. COLLINS and J. K. WOOD.
- "The composition of paraffin wax. Part III." By F. FRANCIS, C. M. WATKINS, and R. W. WALLINGTON.
- "The hydrogen sulphates of the alkali metals and ammonium." By H. B. DUNNICLIFF.

LECTURE.

On Thursday, December 14th, at 8 p.m., at the Institution of Mechanical Engineers, Storey's Gate, Westminster, S.W. 1, Professor Cecil H. Desch, D.Sc., Ph.D., will deliver a Lecture entitled "The Metallurgical Applications of Physical Chemistry."

COLLECTIVE INDEX, VOL. VI (1913-1922).

Volume VI of the Collective Index, to be published in two parts, will be issued Part I in 1923 and Part II in 1924.

The price to Fellows who apply before *Christmas Day, 1922*, is £2 17s. post free. A remittance, made payable to Mr. S. E. Carr, must accompany the application in each case and should be sent to the Assistant Secretary, Chemical Society, Burlington House, W. 1. The price to Fellows who apply after December 25th, 1922, will be £3 7s.

Arrangements have been made by which cases in blue or maroon cloth for binding this Index in three volumes (Authors' Index, 1 vol. and Subject Index, 2 vols.) will be supplied for the sum of Seven Shillings. The binding will involve an additional charge of 2s. 3d. per volume. The total cost, therefore, of binding the Index in cloth will be 13s. 9d.

Fellows desiring to avail themselves of this offer should apply to Messrs. Gurney & Jackson, 33, Paternoster Row, London, E.C. 4.

CERTIFICATES OF CANDIDATES FOR ELECTION AT THE BALLOT TO BE HELD AT THE ORDINARY SCIENTIFIC MEETING ON THURSDAY, DECEMBER 7TH, 1922.

N.B.—The names of those who sign from "General Knowledge" are printed in *italics*.

ALLANSON, ROBERT, "Oaklands," 10, Grosvenor Drive, New Brighton, Wallasey, Cheshire. British. Director of Powell & Scholefield, Ltd., Chemical Manufacturers, Liverpool, and interested in research on behalf of the chemical industry and in connexion with manufacturing chemistry. (*Signed by*) R. L. Siau, Arnold R. Tankard, Douglas Henville.

ALLCHIN, THOMAS CUTHBERT, (M.B.E.), Longfield, Kent. Works manager (chemical), E. C. Powder Coy., Ltd. Manufacturing chemistry. To receive publications and attend meetings. (*Signed by*) Edgar Neumann, Matthew J. Cannon, Herbert E. Burgess.

ARMIT, JAMES WILSON, The Manse, Monikie, nr. Dundee. Chemical research student at St. Andrew's University. M.A., St. Andrews (1920). Research student in chemistry since October 1921. (*Signed by*) Robert Robinson, J. C. Irvine, John Pryde, E. L. Hirst.

BACARESE, LEOPOLD LOUIS, 2, Stirling Road, Clapham Road, S.W.9. British. Managing Chemist. (1) Two years student at the Westminster School of Pharmacy, London, S.W. (2) Passed the qualifying exam., Gibraltar. (3) Member of the Pharmaceutical Society. (4) Member of the British Optical Association and the Institute of Ophthalmic Opticians. (5) Four years Managing Chemist to Heppell & Co., London. Keenly desirous of becoming a Fellow to keep in touch and attend lectures, receive Society's publications, with a view to improving experience in Organic Chemistry.

(Signed by) Robert S. Glennie, W. Bennison, P. H. Woodnoth, H. Sampson Wills.

BAINS, LESLIE, 5, Carlton Road, Stroud Green, N.4. British. Research Student. Natural Sciences Tripos, Cambridge: Part I (1st Class), Part II (2nd Class). Two years' research in Organic Chemistry at Cambridge; now at the Imperial College of Science, South Kensington. B.A. (Cantab.), B.Sc. (London). Associate of Institute of Chemistry. (Signed by) Jocelyn Thorpe, C. K. Ingold, Arnold Stevenson, G. A. R. Kon.

BATHJA, MENGHAJ CHOITHRAM, Fitzwilliam Hall, Cambridge. British subject. Student. I have remained as a student of chemistry for the last six years at the universities of Bombay and Cambridge, respectively, and have obtained Honours degree of Bombay in chemistry and physics, and 2nd Class Honours degree of Cambridge in chemistry, physics, and botany. (Signed by) H. J. H. Fenton, A. J. Berry, T. M. Lowry.

BELAJEW, NICHOLAS TIMOTHY, Colonel, C.B., 33, Elvaston Place, Queen's Gate, London, S.W.7. Russian. Metallurgist. Formerly Lecturer in Chemistry and Metallurgy at the Michael Artillery Academy, Petrograd. Colonel, C.B., M.I.Met., M.I.Min.E., Hon. M.I.Ae.E. Honorary Corresponding Fellow of the Royal Society of Arts. Honorary Corresponding Member of the Council of the Institute of Metals. Member of the Council of the I. Aeronautical Engineers. Member of the Iron and Steel Institute, the Faraday Society, the Society of Chemical Industry, and of the Mineralogical Society. Member of the Société de Chimie Physique in Paris. In Russia member of Society of Physics and Chemistry, Imp. R. Mineralogists and others. Contributor of several papers on chemistry, physics, and crystallography. (Signed by) H. C. H. Carpenter, W. J. Pope, R. A. Hadfield.

BELL, JAMES, 30, Trinity College, Dublin. British. University Demonstrator of Chemistry. Senior Moderator in Experimental Science (with large gold medal for first place and high marks), Dublin University, 1920. Demonstrator of chemistry, Dublin University, 1919-1922. Part author of two papers on original research, viz. *J. Chem. Soc.*, 117, 1133, (1920), and *J. Chem. Soc.*, 121, 1790, (1922). (Signed by) Sydney Young, Emil A. Werner, A. G. G. Leonard.

BROOME, DONALD CHARLES, "Disgwylyfa," Penmaenmawr, North Wales. British. Technical Chemist. Educated at Emanuel School, Wandsworth Common, and later at Woolwich and Battersea Polytechnics. Have passed Inter. B.Sc. (London) and am studying for final B.Sc., having completed one year's work for this examination. Late chief assistant chemist at the South Met. Gas Co.'s Tar Works, East Greenwich. Now holding the post of works chemist at the Penmaenmawr works of the Penmaenmawr and Trinidad Lake Asphalt Co., Ltd. (Signed by) M. O. Forster, J. Kenyon, J. L. White, Herbert E. Nickels, Wallace C. Nickels.

BUIST, WALTER FERGUSON, 24, New Hall Lane, Clubmoor, Liverpool. Scotch. Graduate. Awaiting teaching post. Northern Universities Matriculation 1913. Inter.B.Sc., University of Liverpool, 1914. (War service 1914-1920.) Honours chemistry, B.Sc., 1922. Desirous of keeping in touch with current literature. (Signed by) E. C. C. Baly, Herbert H. Froyssell, I. M. Heilbron, J. S. Buck.

BULT, STANLEY ROBERT, 63, Addiscombe Rd., Croydon. British. Chemist. At present time Curator, Institute of Hygiene, lecturing and demonstrating upon dietetic and hygienic problems. Twenty-two years' experience of chemistry (with break during the war). Seventeen years' laboratory work, oils, waxes, soaps, paints, metals, rubber, foodstuffs. Some

small contributions to technical journals. (*Signed by*) Herbert J. Bult, Lawrence Briant, Julian L. Baker.

BURFORD, ARTHUR JOSIAH, c/o 419, Edgware Rd., Maida Vale, W.2. British. Chemist and Druggist. (1) Two years student in chemistry at the School of Pharmacy of the Pharmaceutical Society, Bloomsbury Square, W.C. (2) One year student in pharmaceutical and analytical chemistry at the London College of Chemistry and Botany, S.W. (3) Member of the Pharmaceutical Society. (4) Three years laboratories of Wands, Manufacturing Chemists, Leicester. (5) Three years chemist to Heppell & Co., London. (6) Keenly desirous of becoming a Fellow to keep in touch and attend lectures, receive Society's publications, with a view to improving experience in Pharmaceutical Organic Chemistry. (*Signed by*) William Bennisson, Gabriel Cofman-Nicoresti, Robert S. Glennie.

BURROWS, GEORGE JOSEPH, Chemical Department, University, Sydney, N.S.W. B.Sc. Sydney (1910). Lecturer in Chemistry, Sydney University. Author of various publications in *J.C.S.* since 1914. (*Signed by*) Eustace E. Turner, Charles E. Fawsitt, J. A. Schofield.

CARTER, WILLIAM LESLIE, "Ellesmere," Bloxidge St., Langley Green, nr. Birmingham. British. Works Chemist (in charge). Educated at the Secondary School, Oldbury (by county scholarship). Oxford Univ. local exam. honours and distinction in chemistry. 1915, Assistant chemist, Messrs. Chance & Hunt, Ltd., Oldbury, Chemical Mfrs. 1916, Assistant chemist, Messrs. J. Hardman, Ltd., Birmingham, tar distillers. 1917, Assistant chemist, H.M. factory site "B," Oldbury. 2½ years' military service (France). 1921, Board of Education higher education course in advanced chemistry. 1922, Chemist-in-charge, Messrs. Birmingham Starch & Mfg. Co., Ltd., Birmingham. Paper in preparation on "The Manufacture of Starch." (*Signed by*) J. G. Howarth, Arthur Adams, Joseph Lones.

CAVIGAN, SYDNEY GEORGE, 22, Walker St., Burton-on-Trent. Analytical Chemist. I have had five years' experience in the analysis of metals, foods, and drugs, and general experience in biological chemistry and toxicology, and am at present in charge of the analytical department of Messrs. Evans, Sons, Lescher & Webb, Research Laboratories, Runcorn. I wish to become a member in order to obtain the *Journal* to assist me in my researches, and to keep up my knowledge of modern chemical literature. (*Signed by*) Wm. P. Skerchly, James Booth, J. Turnbull, Herbert H. H. Lorenz.

CHAMPION, JOHN HARDWICK LONG, 41, Triangle West, Clifton, Bristol. Student. Bachelor of Science (Bristol). Have done eight months' research in physical chemistry under Prof. J. W. McBain. Am at present engaged as a part-time teacher of chemistry and physics. (*Signed by*) J. W. McBain, M. Nierenstein, Millicent Taylor, B. Fullman.

CHOUDARY, KATRAGADDA SESHACHALAM, 71, Coral Merchant St., George Town, Madras, India. Indian (Hindu). Leather Trades Chemist. Master of Arts of the University of Madras, Master of Science of the University of Leeds; lecturer in chemistry in the Christian College, Madras (1917-18); lecturer in chemistry in the Government Arts College, Rajahmundry, Madras; (1918-19); chemist in the Leather Trades Institute, Madras (1919-20). (*Signed by*) P. K. Dutt, Akira Shimomura, H. M. Dawson.

CHRISTIE, ELLIOT MARTIN, 37, Lynwood Avenue, Dunedin, N.Z. Secondary School Teacher. Graduated B.Sc., Victoria University College, 1920. M.Sc. (2nd Class Honours in Chemistry), 1922. Thesis on "Oxidation of Unsaturated Organic Compounds by Persulphate." Appointed Science Master, Gore High School, 1922. (*Signed by*) J. K. H. Inglis, John Williams, C. S. Hicks.

CLARKE, FREDERICK STANLEY, 64, Golders Manor Drive, N.W.11. British. Pharmaceutical Chemist. 1899, Demonstrator (chemistry and pharmacy), Manchester. From 1900, 2½ years with the late Wm. Martindale, editor, *Extra Pharmacopoeia*. 1902 to 1920 on technical experimental staff, Burroughs, Wellcome & Co., London. At present works manager and chemist, Anglo-American Oil Co.'s Speciality Dept. for petroleum products. (Signed by) Harold King, Harold C. Sayer, Frank Lee Pyman.

COTTRELL, ALLIN, 3, Oxford Terrace, Edinburgh. British. Lecturer in Technical Chemistry, Edinburgh University. Lately chief chemist, H.M. Factory, Gretna. (Signed by) James Walker, Sydney A. Kay, Leonard Dobbin.

CRANSTON, JOHN ARNOLD, 14, Fernleigh Road, Newlands, Glasgow. British. Lecturer on Physical Chemistry, Royal Technical College, Glasgow. Doctor of Science, Glasgow University. Lecturer in Royal Technical College for three years. (Signed by) F. J. Wilson, Thomas Gray, R. M. Caven.

CURNOW, HUGH JOHN, 22, Halfway St., Sidcup. English. Undergraduate. Inter. Science (London). Student of the Institute of Chemistry. (Signed by) H. Pochin, J. R. Partington, W. H. Patterson.

DANCE, EDWARD, Castle Inn, Sandal, Wakefield. British. Student. B.Sc. Hons. in Col. Chem., Leeds, and Diploma in Dyeing, Leeds University. Desire Fellowship in order that I may receive the *Journal* and have use of Library. (Signed by) R. M. Woodman, J. R. Scott, P. W. Clutterbuck, John S. Carter.

DAVIES, WILFRED THOMAS GREY, Plas-y-Coed, Derllwyn Rd., Tondû, Glamorgan. British. Assistant, Tondû By-product Works. B.Sc., Bristol, 1920. Inter.B.Sc., 1914. Carried out research under Dr. M. Nierenstein at Bristol University. (Signed by) M. Nierenstein, M. Taylor, F. Francis, J. W. McBain, B. Fullman.

DEWHURST, MILTON, 1280, Warwick Rd., Acocoks Green, Birmingham. British. Works Chemist. B.Sc. Hons. Chemistry (Vict.). A.I.C. (Signed by) H. B. Dixon, Arthur Lapworth, F. P. Burt, Frederick Challenger, John Harwood.

EATON, RICHARD VICTOR, "Lakeside," Seldown, Poole. British. Chemist, R.N. Cordite Factory. A.R.C.Sc., B.Sc. (Lond.). (Signed by) H. B. Baker, M. A. Whiteley, James C. Philip.

EVERS, NORMAN, The Cottage, Salisbury Avenue, Cheam, Surrey. British. Chemist. B.Sc. (London), F.I.C. Chief chemist to Messrs. Allen & Hanbury's, Ltd., Bethnal Green, E.2. Author of numerous papers on chemical subjects published in the *Analyst* and elsewhere. (Signed by) F. W. Gamble, G. J. Woods, Thos. McLachlan.

FARMERY, JOSHUA WILLIAM, 56, Sheen Rd., Richmond, Surrey. British. Analytical and Research Chemist. M.A. (Cantab.). A.I.C. Research paper, "Decomposition of Nitric Esters," by Lowry, Browning, and Farmery, *J.C.S.* (Signed by) T. M. Lowry, A. F. Joseph, Cecil H. Cribb.

FORD, SAMUEL FREDERICK, 65, Abbots Road, Southall, Middlesex. British. Assistant Chemist. (Analysis of oils, essences, and foods.) Assistant for three years in laboratory of rubber and antimony manufacturer. Now assistant chemist in laboratory of food manufacturers. Ex-day and now evening student at Battersea Polytechnic. Studying for Science degree, Rutlish Secondary School, Merton, Old Boy. (Signed by) F. J. Wilson, G. S. W. Marlow, J. Beauchamp Salter.

GARDNER, ROY, 91, Lincoln Rd., Masterton, N.Z. Honours Student (Chemistry), University College, Otago, N.Z. Desirous of keeping in touch with recent developments. B.Sc. (N.Z.), now doing research work in chemistry

at Univ. of Otago, N.Z. (*Signed by*) Fredk. P. Worley, W. F. Short, A. H. Howell, J. K. H. Inglis, Thomas H. Easterfield.

GIFFORD, PHILIP HENRY, Boys' High School, Salisbury, S. Rhodesia, S. Africa. Science Master. B.Sc. (Vict.) (1st Class Chemistry and Physics). Assistant Science Master, Ballymena Academy, Co. Antrim (1913-14). Senior Science and Maths. Master, Edinburgh Institution (1914-17 and part of 1919). Senior Science Master and Vice-Principal of the Boys' High School, Salisbury, S. Rhodesia, from Jan. 1st, 1920. (*Signed by*) Charles A. Fogg, T. M. Nightingale, E. Boardman.

GRIFFITH, ROLAND HALL, 14, Bigwood Avenue, Hove, Sussex. British. B.A. (Chemistry), Oxford. Engaged on research in organic chemistry at the University laboratories. (*Signed by*) Edward Hope, W. H. Perkin, S. G. P. Plant, Allan F. Walden, N. V. Sidgwick, M. P. Applebey, B. Lambert.

GRIFFITHS, REGINALD WILLIAM, 137, Holly Road, Handsworth, Birmingham. Member of Research Staff, Research Laboratory, Dunlop Rubber Co., Ltd., Fort Dunlop, Birmingham. Research student at Birmingham Municipal Technical School. B.Sc. (Birmingham), 1920. (*Signed by*) J. Newton Friend, Frederick Challenger, J. E. Coates, S. R. Carter, G. T. Morgan.

GULLAND, JOHN MASSON, 11, Chester Street, Edinburgh. Chemical Research Student. B.Sc. in pure science, Edinburgh University, July 1921. Research student, Chemistry Department, University of St. Andrews. (*Signed by*) Robert Robinson, J. C. Irvine, John Pryde, E. L. Hirst.

HAIGH, HERBERT, 211, Park Road, Barnsley, Yorkshire. British. Student (Graduate). Alternative Matriculation for ex-service men, 1919. Inter. B.Sc., Liverpool, 1920, Maths., Physics, and Chemistry. B.Sc. Liverpool, 1922, Honours Chemistry, Class I. Desirous of keeping in touch with current literature. (*Signed by*) I. M. Heilbron, E. C. C. Baly, Herbert H. Froyssell, Harold Rogerson, J. S. Buck.

HIGGINBOTHAM, LUCY, 57, Wilbraham Rd., Chorlton-cum-Hardy, Manchester. British. Research Assistant in Organic Chemistry at the British Cotton Research Association, Didsbury. B.Sc., Honours Chemistry, 1st Class, 1919, at Manchester University. M.Sc., 1920. Working for Dr. Stephens at Manchester University. 1921-22 Private assistant to Professor Lapworth. Publications: "The Formation of Substituted Succinic Acid from Esters of $\alpha\beta$ -Unsaturated Esters," by Arthur Lapworth and Lucy Higginbotham; "Studies in the Coumaranone Series. Part I," by Lucy Higginbotham and Henry Stephen. (*Signed by*) Arthur Lapworth, Frederick Challenger, Leslie R. Ridgway, R. G. Fargher.

HUBBARD, THOMAS EDWARD, 9, Cunliffe Street, Pendleton, Manchester. Senior Assistant Chemist to the Salford Corporation Gas Works. As a student at the College of Technology, Salford, gained numerous prizes and scholarships in chemistry. Deputised for Mr. C. E. Teasdale, lecturer on gas manufacture. Lectured on calorimetry in modern gas works practice, etc. (*Signed by*) Jas. Grant, E. Bayley, Frank B. Small.

HUMPHRIES, JAMES ERNEST, 29, St. John's Rd., Newport, Mon. British. University Assistant in Chemistry, Aberdeen. B.A. (Lond. and Cantab.). M.Sc. (Wales). (*Signed by*) W. Thomas, E. Bloom, F. G. Mann.

HUNTLEY, JOHN, The Villa, Burstwick, nr. Hull. Works Chemist. Chief chemist, Humber Portland Cement Co., Ltd., Melton, East Yorks. (*Signed by*) W. H. Coates, Harry Thompson, Thos. A. Nightscapes, E. B. Atkinson, T. Luxton.

HURREN, SYDNEY ALFRED, M.C., 39, Upper Marylebone St., W.1. British.

Lecturer at Northern Polytechnic Institute on Technology of Pianoforte Construction, and technical editor of *The Pianomaker*, the monthly journal of the trade. (*Signed by*) Thomas J. Drakeley, Philip Schidrowitz. Raymond R. Butler.

ISKANDER, HABIB, Heliopolis, Cairo. Egyptian. Senior Science Master of the American University, Cairo, Intermediate Section. Graduate of Royal Training College, Cairo, with top mark in chemistry, 1915. Late Senior Chemistry Master at Victoria College, Alexandria, 1915-21. (Victoria College is an English College qualifying for the Oxford and Cambridge Schools Examinations.) Author of "Khoulasit Al Chemia," an Arabic text-book in chemistry adopted by many schools and colleges. Contributor of some articles on "Recent Advances in Chemical Science" published in *Al Moktataf*, the biggest scientific magazine in the Arabic-speaking world. (*Signed by*) A. I. Kahil, A. H. Ahmad, W. Morris Colles, R. Fleming.

JENKIN, JOHN WATSON, "Tregardock," St. Austell, Cornwall. Metallurgical Research Chemist. B.Sc. (Hons.) London (2nd Class Hons. in Chemistry). A.I.C. (by examination). Studied chemistry at King's College, London, under Professors Crossley, Smiles, and Allmand. At present engaged in research work under Professor C. H. Desch in the Dept. of Metallurgy, University of Sheffield. (*Signed by*) Cecil H. Desch, A. J. Allmand, Samuel Smiles.

JEWELL, JOHN ERNEST, 34, Grove End Road, St. John's Wood, N.W.8. English. Pharmaceutical Chemist. Proprietor of Messrs. Heppell, West-end chemists. Member of the Pharmaceutical Society. Director of many well-known chemical firms. (*Signed by*) Frank Radcliffe, Stanley J. Peacock, C. S. Gibson.

JONES, EDGAR, Milverton House, 167, Upper Dale Road, Derby. Technical Chemist. Chief chemist, Derby Gas Light & Coke, Co. Late chemist Gas Helmet Redipping Station, R.A.O.D., Alexandria. Inter. B.Sc. (London). Lecturer, Gas Engineering and Manufacture, Derby Technical College. (*Signed by*) Jos. Yates, Cyril H. Birkitt, Stanley Dixon.

KALFF, JAN, Amsterdam, Nieuwe Prinsengracht 126. Dutch. Assistant, University, Amsterdam. Chem. Doct. Engaged in organic chemical work. Desire to keep up with the latest English investigations in chemistry. (*Signed by*) H. J. Backer, W. A. Frederikse, F. G. Donnan.

KAPUR, SHANKAR NATH, Amritsar, India (at present, 5, Grosvenor Crescent, Edinburgh). Distillery Chemist, Amritsar Distillery Co., Amritsar, India. B.Sc. 1st Class Honours in Chemistry from Government College, Lahore (Punjab University). Chemist to the Amritsar Distillery Co., Amritsar, India (from May 1917). Now sent by the Government of India to study "Industrial Alcohol" in England and on the Continent. (*Signed by*) John E. Mackenzie, Thomas H. Pope, B. H. Wilsdon, Bawa Kartar Singh, Horace B. Dunnickif.

KHAN, GHULAM SAFDAR, Veterinary College, Lahore. Lecturer in Chemistry, Veterinary College, Lahore. Passed B.Sc. examination in 1920 from Punjab University. Appearing in the M.Sc. examination 1922. Special work: (1) Separation of different qualities of turpentine oil. (2) Solubility of sulphur dioxide in sulphuric acid of different concentration at constant temperature. (*Signed by*) S. R. Khosla, N. A. Yajnik, E. T. Dean.

LASSEN, SVEND HALD, "Briarwood," Manorgate Rd., Kingston-on-Thames. Danish. Works Manager. Graduated at the University of Copenhagen. Assistant lecturer at the Royal Mineralogical Museum, Copenhagen, 1913-14. Participated in an Arctic expedition to North Greenland as mineralogist. At

present works manager to Water Softeners, Ltd. (synthetic aluminium silicates and dyestuffs). (*Signed by*) Harald Faber, Bernard Dyer, Robt. G. Grimwood.

LLOYD, WILLIAM DAWSON, 42, Tavistock Road, Westbourne Park, W.11. Medical, Veterinary, and Pharmaceutical Representative. Fifteen years' analytical and pharmaceutical experience—manufacture of bismuth salts, urine analysis, etc. (*Signed by*) A. E. Mills, Geo. A. Pemberton Wright, Herbert A. Mills.

LYNCH, PHILIP PATRICK, Church Street, Timaru, N.Z. Medical Student. B.Sc. (N.Z.). First and second professional examinations for degrees M.B., Ch.B. (N.Z.). 1918 Asst. Chemist, Department of Agriculture, Wellington. At present assistant to C. S. Hicks, M.Sc., A.I.C., Public Analyst, Dunedin (Fellow). Engaged in biochemical research on goitre, nephritis, and diabetes under Professor of Pathology at University of Otago and in collaboration with C. S. Hicks, M.Sc. (Fellow). (*Signed by*) C. S. Hicks, Thomas H. Easterfield, Fredk. P. Worley, J. K. H. Inglis.

MACALPINE, FRANCIS GEOFFREY, Broad Oak, Accrington, Lancs. British. Managing Director of Coal-Tar Distillation and Ammonia Works. Degree of B.Sc. of Manchester University. Twelve months' study of coal carbonisation and working up of by-products at Birchenwood Collieries, Kids Grove. I desire to join the Chemical Society so as to be kept informed of the progress of chemistry through the Transactions of the Society and its Library. (*Signed by*) B. Prentice, E. Clark, D. V. Hollingworth.

MCDOWALL, FREDERICK HENRY, Knox College, Dunedin. Research Student, Otago University. M.Sc. with 1st Class Honours. Thesis on "Constitution and Synthesis of Menthol." Smeaton Research Scholar of Otago University, 1921-22. Student demonstrator and half-time assistant to Dr. Inglis in Chemistry Department, Otago University, 1919, 1921-22. (*Signed by*) J. K. H. Inglis, C. L. Carter, C. S. Hicks.

MCKENZIE, KENNETH, "Endrick," Hamilton Rd., Cambuslang. Scotch. Analytical Chemist. For the past ten years engaged in the study and practice of analytical chemistry. Training: at Royal Technical College, Glasgow, and with Messrs. R. R. Tatlock & Thomson, Public Analysts, Glasgow. At present assistant chemist with Lantero Nitrate Co., Ltd., Chile, with complete control of all analytical work. Desirous of joining Chemical Society for purpose of keeping in touch with advances in analytical chemistry. Originator of method for determination of perchlorates in small quantities in caliche, etc. Refinement of ferrous sulphate assay for nitrate in caliche, whereby experimental error is reduced from $\pm 4\%$ to $\pm 0.6\%$. (*Signed by*) R. T. Thomson, A. R. Campbell, George Grant, Alex. W. McLaren.

McNICOL, REGINALD ARTHUR, "Birnam," Rotherfield Road, Carshalton, Surrey. British. Bio-chemist to the Jersey Tomato and Fruit Preserves, Ltd., St. Heliers, Jersey (J. Lyons & Co.). Pupil Sutton (Surrey) County Secondary School, Sept. 1912-July 1916. Student Battersea Polytechnic, Sept. 1916-June 1922. B.Sc. Lond. (3rd Class Hons. in Chemistry), Oct. 1920. M.Sc. Lond., May 1922. Thesis: "The Preparation and Optical Properties of the Aliphatic Ethers of Secondary Octyl Alcohol." (*Signed by*) Leslie Hall, J. Kenyon, Harold Hunter, J. L. White.

MALKIN, THOMAS, Golden Square, Warrington. British. Research Student in Chemistry, Victoria University, Manchester (Ph.D. candidate). B.Sc. (London), Hons. Chemistry. A.I.C. Two years' post-graduate research at College of Technology, Manchester. Part-time lecturer in chemistry at Warrington Technical Institute, 1920-22. (*Signed by*) Arthur Lapworth, Frank Lee Pyman, Lealie Randal Ridgway.

MARTIN, ARCHIBALD ROY HAMMOND, 8, Kingsland Rd., Worthing, Sussex. British. Student. At St. John's College, Cambridge, read for Nat. Sci. Tripos, Part I, for 2 years, taking the B.A. degree in June, 1921. Read for Nat. Sci. Tripos, Part II, in chemistry for 1 year, passing in Class 3. (*Signed by*) W. G. Palmer, T. M. Lowry, Charles T. Heycock.

MATTHEWS, JOSEPH KNITH, M.A. (Cambridge), Cedrio House, Rainhill, Lancashire. British. Fuel Technologist (Messrs. Pearson & Knowles Coal & Iron Co., Warrington, Lanes.). 1908, 1st Class Honours in Natural Sciences Tripos, Camb. (including chemistry). 1909, 2nd Class Honours in Part II Chemistry Tripos, Camb. From 1909 the applicant has been in a responsible position in chemical industry, first with Messrs. Pilkington Bros. and later with the above-named firm. (*Signed by*) A. J. Berry, Charles T. Heycock, S. R. Adcock, W. E. S. Turner.

MOORHEAD, CHARLES WILLIAM, 23, Newbridge Rd., Bath. Chemist to Meldon Valleys Co., Ltd. Twenty-five years' experience in drug and chemical trades, glass technology, manufacturing chemistry. (*Signed by*) Alan E. Gardiner, Alfred Mander, A. E. Mills.

MOSS, JOHN EWART, 298, Franklin Road, Kings Norton, Birmingham. Research Student. B.Sc. (1st Class Honours in Chemistry) and M.Sc. (Birm.). (*Signed by*) G. T. Morgan, S. R. Carter, Wm. Wardlaw, E. Ashley Cooper, C. J. V. Bews, J. D. Main Smith.

NATH, DINA, Jullundhur City (Punjab), India. At present assistant chemist to Bhatni Sugar Works, Gorakhpur (District), United Provinces, India. Graduate in Science of the Punjab University. Specialist in soap manufacture. Written a paper on soapmaking with special reference to Indian conditions. At present working on "The Action of Alkalies on Rosine under Different Conditions." Certificate from the Professor-in-Charge with regard to the ability in applied chemistry is attached herewith. (*Signed by*) Bhagwan Das Batta, N. A. Yajnik, E. T. Dean.

NIELSEN, TORKILD HAASTRUP, Knockhall Lodge, Greenhithe, Kent. Danish. Chemist at Kent Cement Works of the Associated Portland Cement Manufacturers, Limited. Bachelor of Science in chemical engineering from the Royal Technical College in Copenhagen. (*Signed by*) H. Baggesgaard Rasmussen, Chr. Winther, J. J. Fox, S. P. L. Sørensen.

NISHIKAWA, HIDEJIRO, 3, Hachimanchō, Asakusa, Tokyo, Japan. Japanese Government Research Scholar. Graduate of the section of agricultural chemistry of the Agricultural Department of the Imperial University of Tokyo, Japan. Research student of the Chemistry Department, St. Andrews University. (*Signed by*) Robert Robinson, J. C. Irvine, John Pryde, E. L. Hirst.

NOMURA, HIROSHI, Chemical Institute, Imperial University, Sendai, Japan. Japanese. Assistant Professor of Chemistry. Graduated at the Tokio Imperial University of Japan, engaged in research for one and a half years under Professor W. H. Perkin. Assistant Professor of Chemistry, Tohoku University, since 1914. (*Signed by*) W. H. Perkin, E. Hope, S. G. F. Plant, E. N. Allott, W. Davies, A. F. Titley, Thos. H. Durrans.

PAGE, ALBERT BRAMAH PEARCE, 27, Tanza Road, Hampstead, N.W.3. Research Student. Associate Royal College of Science. I desire to keep in touch with chemical progress through the *Journal of the Society*. (*Signed by*) James C. Philip, H. J. T. Ellingham, Olive Workman.

PICKLES, HAROLD, Gasworks, Kildwick, nr. Keighley. Chemist and Assistant Engineer, Keighley Corporation Gasworks. Studied chemistry (theoretical, practical, and industrial) at Keighley Trade and Grammar

School, Halifax Technical College, and Leeds University. Nine years' laboratory experience as gasworks chemist. Contributed papers to the *Gas World* on the properties of gaseous mixtures. (*Signed by*) H. Harper, Wm. Cranfield, John Wilkinson.

POOLEY, JOSEPH EDWARDS, 1, Margaretting Road, Manor Park, E.12. British. Principal, "Sandringham" Central School, Forest Gate, E.7. Trained Certificated Teacher. Inter. B.Sc. (London), chemistry as a subject. Inter. B.Sc. (Econ.), London. L.C.P. (chemistry as a subject). B. of E. Certs. in P. and T. Chemistry, advanced stage. Science lecturer at Woolmore St. L.C.C. Evening Institute, Sessions 1914-18. (*Signed by*) Evan D. Griffiths, Léon E. Walling, Alfred W. J. Denney.

POTTS, THOMAS TREVOR, 60, Christ Church Square, Macclesfield. British. Chemist to Messrs. Robert Fletcher & Son, Ltd., Kearsley Paper Works, Stoneclough, and Greenfield Paper Mill, Greenfield, Yorks. Educated at Macclesfield Grammar School (science student under Fisher Advanced Courses Scheme). Matriculated 1917. Part-time student, Manchester Municipal College of Technology in Chemistry and Technology of Paper Manufacture. Held above post from August 1919, since when I have been engaged in general chemical control of processes and private research work in chemistry as applied to the manufacture of high-grade tissues, including original work on new processes of manufacture and methods of analysis. (*Signed by*) William Bacon, Alfred Holt, Julius Hübner.

POUND, JAMES ROBERT, 601A, Dana St., Ballarat, Victoria, Australia. Lecturer in Chemistry, the School of Mines, Ballarat. M.Sc. (Melb., Chemistry). Author of papers on Ether and Sulphuric Acid, *Trans.*, 1911 and 1922, and on Treatment of Wolfram Ores, *Proc. Australian Inst. Min. Met.*, 1919. Formerly research chemist at Amalgamated Zinc (De Bovay's), Ltd., Broken Hill, N.S.W., and superintendent of Magnetic Separation Works, S. & M. Syndicate, Ltd., Launceston, Tas. (*Signed by*) Orme Masson, A. C. D. Rivett, D. Avery.

PRASAD, NAGENDRA, Ranchi, Behar and Orissa, India. Deputy Collector. B.Sc. with Honours in Chemistry. M.Sc. with Thesis and Organic Chemistry as the special subject. Joint author of "Iodination by means of Nitrogen Iodide or by means of Iodine in presence of Ammonia," published in the *Journal of the American Chemical Society*, Vol. XXXIX, 1917. Worked as a research student for some time at the Agricultural Research Institute at Pusa; has a private laboratory of his own; wants to be in touch with current literature on chemistry. (*Signed by*) Phuldeo Sahay Varma, A. S. Khan, Chandra Bhushan Roy, R. L. Datta.

PRIDEAUX, EDMUND BRYDGES RUDHALL, 6, Arboretum Street, Nottingham. British. M.A. and B.Sc. (N.Z.), D.Sc. (Lond.), F.I.C. Lecturer in Chemistry and Research Chemist at University College, Nottingham. Research falls mainly under the following heads: The compounds of fluorine; the atomic and molecular volumes of liquid elements and their liquid compounds; electrochemical investigations of solutions by the conductivity and electromotive force methods, with special reference to acidity and alkalinity; electrolytic corrosion; the evaporation and deliquescence of salt solutions; the chemical factors controlling the alkalinity of sea water. My books deal with the arithmetical problems of physical chemistry and with the theory and control of acidity and alkalinity, with special reference to the use of indicators. (*Signed by*) F. Stanley Kipping, F. G. Donnan, A. J. Allmand.

PULLEN, ALFRED NORMAN DOUGLAS, The Larches, Greenham Common,

Newbury. English. Late Analyst to China Mining and Metal Co., Ltd., Hong Kong. In charge of smelting and refining works to above Co., 1918-22. Pharmacist to Hong Kong Government, 1914-17, and assistant analyst, 1915-16. Assistant to T. Tickle, B.Sc., F.I.C., Boro' Analyst, Exeter, 1912-13. Major Diploma, Pharmaceutical Society and silver medallist (chem.). 1912. Paper on "Iodine Ointment," *Pharm. Journ.*, Nov. 16th, 1912. (*Signed by*) A. W. Crossley, Thomas Tickle, Frank Comyns.

RILEY, GEORGE CLIFFORD, B.A., B.Sc., 15, Wellington Square, Oxford. British. Research Chemist. Works chemist on production of explosives, 1915-19. Student at Oxford University, 1914-15 and 1919-21, taking 2nd Class Honours, Final Honours Chemistry School, and B.A. 1920. One year's research work in organic chemistry and B.Sc. degree, 1921. Research assistant to Professor W. H. Perkin, M.A., F.R.S., 1921 to present date. Publication: "The Chlorination of Benzoyl Chloride" (with Edward Hope), accepted for publication in the *Journal of the Society*. (*Signed by*) W. H. Perkin, N. V. Sidgwick, E. Hope, W. Davies, A. F. Tittley, D. Ll. Hammick, S. G. P. Plant.

ROBERTSON, THOMAS ABRAHAM, "Sassafras," Rickard Road, Bankstown, N.S.W. Chemist and Manager, Austral Starch Co., Nelligen, N.S.W. Student at Sydney Technical College in Inorganic Chem. complete course, Theory and Practical. Organic Chemistry 3 years, 1907-13. Matriculated for Sydney University, 1914. Assist. in Physiological Research Lab., Sydney University, for 2 years, 1910-12. Member of Sydney Technical College Chemical Society for 8 years. Special representative of commercial firms for 2 years. At present Organic Chemist and Manager of Austral Starch Co., Nelligen, N.S.W. Paper and exhibits before Sydney Tech. Coll. Chem. Society: "Chemical Substances in Human Body" and "Glandular Secretion of Iodine in Blood Stream;" other papers in course of preparation. (*Signed by*) Richard W. Challinor, Arthur R. Penfold, Archibald D. Ollé, Robt. Grant, George Z. Du Pain.

RUSHWORTH, BURKE, 4, West Park Terrace, Healey, Batley, Yorks. English. Manager, Chemist Department, Batley Co-operative Society, Ltd. Certificate winner Army Science College, Bonn University, Germany, having studied there after the armistice. Ex-student S.W. Polytechnic Institute, Chelsea, having taken the Minor Pharmaceutical Exam. Now working for the Major and F.S.M.C. diplomas. (*Signed by*) C. Morton, F. H. Lowe, F. Arnall.

SCHOLEFIELD, ARTHUR, Poulton Hey, Bromborough, Cheshire. British. Director of Powell & Scholefield, Ltd., Chemical Manufacturers, Edge Hill, Liverpool. Interested in the chemical industry, and desirous of keeping in touch with scientific research. (*Signed by*) Harold Rogerson, E. C. C. Baly, I. M. Heilbron.

SINGH, GOPAL, "The Brij," Gawalmandi, Lahore, India. Indian. Professor of Chemistry, Dyal Singh College, Lahore. B.Sc. with Honours in Chemistry, M.Sc. 1st Class First. For some time research student of the Punjab University. For about 2½ years Professor of Chemistry at the Dyal Singh College, Lahore. Joint author of the paper (*J.C.S.*, T., 1920, p. 980). Author of the paper (*J.C.S.*, T., 1922, May, p. 782) "Phototropy of Inorganic Salts, Cuprous Chloride, and Bromide." (*Signed by*) N. A. Yajnik, Horace B. Dunncliff, Tara Singh Mahal.

SMITH, GEORGE VICTOR, Westleigh, Eggington Rd., Wollaston, Stourbridge. British. Student. A.R.C.S. (Chemistry). Formerly Assistant Master, King Edward VI. Grammar School, Stourbridge. (*Signed by*) James C. Philip, M. A. Whiteley, H. F. Harwood, J. N. Sugden, J. G. Howarth.

SNEDKER, STANLEY JAMES CORRAL, "The Rowans," Wood Hey, Rock

Ferry, Cheshire. Senior Assistant, Department of Organic Chemistry, University of Liverpool. Carrying out original research. I wish to keep in touch with current chemical literature. (*Signed by*) E. C. C. Baly, I. M. Heilbron, Harold Rogerson, J. S. Buck.

SNOW, OSCAR WALTER, Graeme's Bank, Hatch End, Middlesex. British. Research Student. B.A. (Oxon.) [Part I, Final Honours School of Chemistry]. Scholar of New College, Oxford. At present engaged in research for Part II. (*Signed by*) W. H. Perkin, S. G. P. Plant, N. V. Sidgwick, Allan F. Walden, E. Hope.

STOYLE, FRANCIS WILBERT, 37, Kansas Avenue, Belfast. British. Student. B.Sc. (1st Class Hons.), Queen's University, Belfast, 1921. M.Sc. Queen's University, Belfast, 1922. (*Signed by*) Robert Robertson, A. W. Stewart, H. Graham.

THEOBALD, LESLIE STUART, 3, Ennismore Avenue, Chiswick, W.4. Student. A.R.C.S. course in chemistry just completed. (*Signed by*) James C. Philip, H. F. Harwood, J. N. Sugden.

THOMPSON, WILLIAM, "Ravensworth," Cockton Hill, Bishop Auckland. Student. Bachelor of Science, University of Durham. Degree obtained June 1922. (*Signed by*) P. Phillips Bedson, W. N. Haworth, Grace C. Leitch.

TIMMIS, LAURENCE BARNETT, 19, Trinity Marsh Lane, Waltham Cross, Herts. British. Chemist. M.Sc.Tech. (Manchester), A.I.C. Employed at Messrs. Allen & Hanbury's, London, as assistant chemist in the analytical and research departments. (*Signed by*) F. W. Gamble, Frank Lee Pyman, John K. Wood.

TRUSZKOWSKI, RICHARD, Koszykowa 21, m. 3, Warsaw, Poland. British. Research Student. B.Sc. (*Signed by*) Stanley A. Mumford, O. L. Brady, E. H. Thierry.

VINEY, HUMPHREY TERENCE, Bramerton, Blakebrook, Kidderminster. British. Carpet Industry, Application of Chemical Science to. Honours B.Sc., Birmingham University. (*Signed by*) G. T. Morgan, S. R. Carter, E. A. Cooper, C. E. Wood.

WALL, THOMAS JOHN, Northcote, Alexandra Road, Malvern. British. Research Student. Took Part I of Final Honour School of Natural Science at Oxford in June last and am now doing research in Organic Chemistry for Part II. Exhibitioner of Christ Church, Oxford. (*Signed by*) S. G. P. Plant, N. V. Sidgwick, E. Hope.

WARD, HARRY, Toynbee Hall, 28, Commercial St., E.1. British. Research Chemist. B.Sc. (Hons.) (Leeds), M.Sc., A.I.C. 2½ years' experience as research chemist with the General Electric Company. (*Signed by*) R. Le Rossignol, L. D. Goldsmith, William Singleton.

WATSON, STEPHEN JOHN, School of Agriculture, Reaseheath Hall, Nantwich, Cheshire. British. Demonstrator in Agricultural Chemistry, Armstrong College (in the University of Durham). Degree of B.Sc. in agriculture, Armstrong College, Durham University. Registered student of the Institute of Chemistry. Research on potash. Paper accepted for publication by the Society of Public Analysts. Eighteen months' experience in coal analysis. Delivered 50 lectures to Elementary Class in Agricultural Chemistry. (*Signed by*) S. Hoare Collins, Philip Buckle, Lionel Orange, Harold Scragg.

WILLIAMS, ALEXANDER MITCHELL, Shirley Institute, Didsbury, Manchester. British. Research Chemist. M.A., D.Sc. Author or joint author of over a dozen scientific papers on physical chemistry (including T., 1920, 117, p. 844). (*Signed by*) Arthur W. Crossley, D. Clibbens, R. G. Fargher, J. C. Withers.

WOODWARD, ERIC RICHARD BORAMAN, The Vicarage, Herne Bay, Kent. Chemist. Graduate (B.Sc. in Chemistry and Physics) of the University, Liverpool. (*Signed by*) I. M. Heilbron, E. C. C. Baly, Herbert H. Froyssell.

WOOLCOCK, WILLIAM JAMES UGLOW, C.B.E., M.P., 166, Piccadilly, W.1. British. General Manager, Association of British Chemical Manufacturers. Pharmaceutical Chemist. Barrister-at-law. (*Signed by*) M. O. Forster, N. Garrod Thomas, E. V. Evans, F. F. Armstrong.

The following Certificate has been authorised by the Council for presentation to ballot under Bye-law I (2) :

GOVILA, BABU LALL, Gwalior. Hindu. Principal of Central Polytechnic Institute, and Director of Industrial and Technical Education, Gwalior State. (1) Studied chemistry in Indian University College and Japan. (2) Graduated in Tokyo (Japan) in colour chemistry. (3) Studied weaving and spinning in Japan. (4) Worked as Professor and Demonstrator of applied chemistry. (5) Worked in dyeing, calico printing, and soap works in Japan and India. (6) Worked as Principal of the Institute of Textile Technology, Chanderi. (7) Worked in colour research and coal-tar products. (8) Working as Principal of Central Polytechnic Inst. with a department for applied chemistry, and training students in dyeing, calico printing, soap-making, coal-tar distillation, intermediate products, and coal-tar dyes, for the City and Guilds of London Examinations, for the last four years. (*Signed by*) Puran Singh.

ADDITIONS TO THE LIBRARY.

I. Presented by the Society of Chemical Industry.

AGRICULTURAL LEDGER. Calcutta 1892—1912. (*Reference.*)

CHEMICAL ENGINEERING AND THE WORKS CHEMIST. Vols. I—V. Dover 1911—1915. (*Reference.*)

DINGLERS POLYTECHNISCHES JOURNAL. Vols. CCIX—CCLVI. Stuttgart 1876—1885. (*For Circulation.*) [Wanting Vol. CCXXXV.]

INDIA, *Imperial Department of Agriculture*. Agricultural Journal of India. Vols. III—XIV. Calcutta 1908—1919. (*Reference.*) [Wanting Vols. V, VI, VIII, IX. Vols. III, VII, XI, XIII incomplete.]

INDIA RUBBER JOURNAL. Vols. XLIX—LVIII. London 1915—1919. (*Reference.*)

KÖGYÖ-KWAGAKU-KWAI. (Society of Chemical Industry.) Zasshi. (Journal.) Vols. XVIII—XXII. Tokyo 1915—1919. (*Reference.*) [Vols. XVIII, XXI, XXII incomplete.]

PERFUMERY AND ESSENTIAL OIL RECORD. Vols. VI—XI. London 1914—1920. (*Reference.*) [Vols. IX, X incomplete.]

PHILOSOPHICAL MAGAZINE. Vols. XXIX—XL. London 1915—1920. (*For Circulation.*) [Wanting Nos. 193, 233—234, 236.]

REVUE DE METALLURGIE. Vols. XI—XVI. Paris 1914—1919. (*Reference.*) [Vols. XI, XIV, XV incomplete.]

RUSSIAN PHYSICAL and CHEMICAL SOCIETY. *Journal*. Vols. XXV—XLVIII. St. Petersburg 1893—1916. (*For Circulation*.) [Wanting Vols. XXXV, XXXVI. Vols. XXV, XXVII, XXXII—XXXIV, XXXVII incomplete.]

U.S. DEPARTMENT OF AGRICULTURE. *Office of Experiment Stations*. Experiment Station Record. Vols. XI—XLVIII. Washington 1899—1919. (*Reference*.) [Wanting Vols. XVIII, XXXIII—XXXVI. Vols. XI, XXVI—XXX, XXXVIII, XXXIX, XLI incomplete.]

U.S. DEPARTMENT OF COMMERCE. *Bureau of Standards*. Bulletin. Vols. I—XIV. Washington 1904—1918. (*Reference*.)

Also odd volumes and parts of the following :

Circulars, Scientific and Technologic Papers of the Bureau of Standards. Biochemical Journal. Bulletins and Circulars of the U.S. Bureau of Chemistry.

II. *Miscellaneous Donations.*

ABDERHALDEN, EMIL. Handbuch der biologischen Arbeitsmethoden. Abt. I. Chemische Methoden. Abt. IV. Angewandte chemische und physikalische Methoden. Berlin 1921+.

From the Publishers : Herrn Urban & Schwarzenberg.

FUEL IN SCIENCE AND PRACTICE. A journal of the scientific and economic use of fuels. (Being the Record of the Coal Research Club.) Vol. I, etc. London 1922+. ill. (*Reference*.) [Monthly supplement to The Colliery Guardian.]

From the Publishers : The Colliery Guardian Co., Ltd.

HACKH, INGO W. D. Chemical reactions and their equations. A guide and reference book for students of chemistry. Philadelphia 1922. pp. viii+138. ill. 6s. net. (*Recd.* 13/11/22.)

From the London Publishers : Messrs. Chapman & Hall.

MINISTRY OF MUNITIONS and DEPARTMENT OF SCIENTIFIC AND INDUSTRIAL RESEARCH. Technical records of explosives supply. No. 9. Heat transmission. London 1922. pp. iv+48. ill. 5s. net. (*Recd.* 1/11/22.)

From the Department.

U.S. WAR DEPARTMENT. Report on the fixation and utilization of nitrogen. Washington 1922. pp. xii+354. ill. (*Recd.* 15/7/22.)

From the Chief of Ordnance.

III. *By Purchase.*

ARNAUD, D., and FRANÇHE, G. Manuel de céramique industrielle. 2nd edition. Paris 1922. pp. x+642. ill. (*Recd.* 4/9/22.)

ASSOCIATION OF OFFICIAL AGRICULTURAL CHEMISTS. Official and tentative methods of analysis, as compiled by the Committee on

Revision of Methods. 2nd edition, revised to November 1, 1919. Washington, D.C. 1921. pp. xii+418. ill. (*Recd.* 21/8/22.)

BARKER, ALDRED FARRER, [and others]. Textiles. Revised edition. London 1922. pp. xii+386. ill. 15s. net. (*Recd.* 16/11/22.)

BINGHAM, EUGENE C. Fluidity and plasticity. New York 1922. pp. xii+440. ill. 20s. net. (*Recd.* 16/11/22.)

BOGUE, ROBERT HERMAN. The chemistry and technology of gelatin and glue. New York 1922. pp. xii+644. ill. 30s. net. (*Recd.* 16/11/22.)

BOLL, MARCEL, and BENNEJEANT, CHARLES. Chimie des métaux et métallurgie dentaires. Paris 1922. pp. xii+252. ill. (*Recd.* 1/7/22.)

CHEMISCH-TECHNISCHE UNTERSUCHUNGSMETHODEN. 7th edition. Edited by ERNST BERL. Vol. I. etc., Berlin 1921+. ill. (*Reference.*)

DAVIS, ARTHUR CHARLES. Manufacture of Portland cement. 3rd edition. Dublin 1922. pp. xiv+416. ill. 25s. net. (*Recd.* 6/7/22.)

DAY, DAVID T., and others. A handbook of the petroleum industry. 2 vols. New York 1922. pp. x+964, vi+1006. ill. 75s. net. (*Recd.* 16/11/22.)

EINSTEIN, ALBERT. Untersuchungen über die Theorie der "Brownschen Bewegung." Edited by R. FÜRTH. (Ostwald's *Klassiker*, No. 199.) Leipzig 1922. pp. 72. ill. (*Recd.* 14/9/22.)

EPHRAIM, FRITZ. Anorganische Chemie: ein Lehrbuch zum Weiterstudium und zum Handgebrauch. Dresden 1922. pp. viii+727. (*Recd.* 3/10/22.)

FISCHER, EMIL. Untersuchungen über Kohlenhydrate und Fermente. II. (1908—1919.) Berlin 1922. pp. x+534. (*Recd.* 14/9/22.)

GILDEMEISTER, EDUARD, and HOFFMANN, FRIEDRICH. The volatile oils. 2nd edition, translated by EDWARD KREMERS. Vol. III. London 1922. pp. xx+778. ill. 32s. net. (*Recd.* 6/11/22.)

HAAS, PAUL, and HILL, THOMAS GEORGE. An introduction to the chemistry of plant products. Vol. II. London 1922. pp. viii+140. ill. 7s. 6d. net. (*Recd.* 10/10/22.)

HOFMAN, HEINRICH OSCAR. Metallurgy of zinc and cadmium. New York 1922. pp. xii+342. ill. 20s. net. (*Recd.* 20/6/22.)

INSTITUTION OF PETROLEUM TECHNOLOGISTS. The petroleum industry. A brief survey of the technology of petroleum. London [1922]. pp. viii+346. ill. 14s. net. (*Recd.* 6/7/22.)

KAYSER, EDMOND. Microbiologie appliquée à la fertilisation du sol. 4th edition. Paris 1921. pp. 325. ill. (*Recd.* 1/7/22.)

— Microbiologie appliquée à la transformation des produits agricoles. 4th edition. Paris 1921. pp. 390. ill. (*Recd.* 1/7/22.)

MEYER, RICHARD. Victor Meyer: Leben und Wirken eines deutschen Chemikers und Naturforschers 1848—1917. Leipzig 1917. pp. xvi+472. ill. (*Recd.* 21/8/22.)

— Vorlesungen über die Geschichte der Chemie. Leipzig 1922. pp. viii+468. (*Recd.* 10/10/22.)

ROBERTSON, GEORGE SCOTT. Basic slags and rock phosphates. Cambridge 1922. pp. xvi+120. ill. 14s. net. (*Recd.* 20/6/22.)

ROBINSON, CLARK SHOVE. The elements of fractional distillation. New York 1922. pp. x+206. ill. 12s. 6d. net. (*Recd.* 21/6/22.)

ROGERS, ALLEN. Practical tanning: a handbook of modern practice and processes as applied in the manufacture of leather and allied products. London 1922. pp. xxvi+700. ill. 45s. net. (*Recd.* 6/7/22.)

RYDBERG, JOHANNES ROBERT. Untersuchungen über die Beschaffenheit der Emissionsspektren der chemischen Elemente. Translated from the French by A. VON OETTINGEN. Edited by MANNE SIEGBAHN and A. VON OETTINGEN. (Ostwald's *Klassiker*, No. 196.) Leipzig 1922. pp. xvi+214. ill. (*Recd.* 14/9/22.)

SCOTT, WILFRID W. [Editor.] Standard methods of chemical analysis: a manual of analytical methods and general reference. 3rd edition. 2 vols. New York 1922. pp. xxviii+726, xviii, 1001-1570+50. ill. 52s. 6d. net. (*Recd.* 16/11/22.)

TAMMANN, GUSTAV. Aggregatzustände: die Zustandsänderungen der Materie in Abhängigkeit von Druck und Temperatur. Leipzig 1922. pp. viii+294. ill. (*Recd.* 27/6/22.)

IV. Pamphlets.

AIR MINISTRY. *Meteorological Office*. Advisory Committee on Atmospheric Pollution. Report on observations in the year ending March 31st, 1921. Forming the seventh report of the Committee for the Investigation of Atmospheric Pollution. London 1922. pp. 31. ill.

ALBERTA. *Scientific and Research Council*. Second annual report. Edmonton 1922. pp. 86. ill.

AMERICAN INSTITUTE OF BAKING. Bulletin No. 8. The diastatic enzymes of wheat flour and their relation to flour strength. By LOUYS A. RUMSEY. 1922. pp. 86. ill.

— Bulletin No. 9. Flour strength as influenced by the addition of diastatic ferments. By FERDINAND A. COLLATZ. 1922. pp. 74. ill.

BELAIEV, NICHOLAS TIMOTHY. Heat treatment of steel. (From the *Russian Economist*, 1922, 2.)

— Henry Marion Howe (1848—1922). (From the *Russian Economist*, 1922, 2.) ill.

PROCEEDINGS

OF THE

CHEMICAL SOCIETY.

Ordinary Scientific Meeting, Thursday, December 7th, 1922, at 8 p.m., Professor SAMUEL SMILES, O.B.E., D.Sc., F.R.S., Vice-President, in the Chair.

The CHAIRMAN referred to the loss sustained by the Society, through death, of :

	Elected.	Died.
Thomas William Dukes	Dec. 5th, 1889.	Nov. 8th, 1921.
David Thomas Williams	Feb. 20th, 1908.	Jan. 29th, 1922.

Mr. G. F. Underhay was formally admitted a Fellow of the Chemical Society.

Certificates were read for the first time in favour of :

- Arthur John Attwood, A.R.C.S., 11, Frederick Crescent, S.W. 9.
- Frederick Bradshaw, M.A., D.Sc., 78, Rothbury Terrace, Heaton, Newcastle-on-Tyne.
- Charles Crossley Mann Broughton, B.Sc., 2, Upton Road, Bexleyheath.
- John Southwell Bousfield, A.R.C.S., 2, Lyndale, Childs Hill, N.W. 2.
- Leslie John Couzens, 101, Kensington Avenue, Manor Park, E. 12.
- Ernest Cecil Craven, 59, Southern Road, Plaistow, E. 13.
- James Cooper Duff, M.Sc., F.I.C., 13, Queen's Road, Erdington, Birmingham.
- John Kenneth Giles, B.Sc., Hillcrest, Arkley, Barnet.
- John Preedy Griffiths, B.Sc., A.R.C.S., 15, Hazelwood Lane, Palmer's Green, N. 13.
- Leslie Julius Harris, B.Sc., 18, The Park, Ealing, W. 5.
- John Laker, B.Sc., 8, Allerton Road, N. 16.
- George Arnold Medley, 45, Hill Top Road, Dalton, Huddersfield.
- William James Osborn, 12, West Square, Southwark, S.E. 1.
- Francis James Paxon, B.Sc., A.R.C.S., 30, St. George's Road, Palmer's Green, N. 13.
- Edwin William Stanley Press, 4, Porchester Square, Hyde Park, W. 2.
- Edward Ebenezer Price, The Haven, Dormans Park, East Grinstead.
- Edgar Sanders, The White House, Blackwell, Worcestershire.
- Hedley Archibald Sloman, B.A., A.I.C., Robinhood, Undercliff Gardens, Leigh-on-Sea, Essex.
- Arthur Roy Stark, 76, De la Polo Avenue, Hull.
- Frederick Vincent Tideswell, M.Sc., Dept. of Fuel Technology, The University, Sheffield.

Frederic Jenner Tritton, B.Sc., 1F, Vicarage Mansions, West Green, N. 15.
 Paule Laure Vanderstichele, M.Sc., Bedford College for Women, Regent's
 Park, N.W. 1.

Stanley Douglas Weaver, B.Sc., A.R.C.S., 27, Hastings Road, Maidstone.
 Harold Wood, B.Sc., 39, Edmund Street, Scedley, Manchester.

The CHAIRMAN announced that a circular had been sent out to Fellows with regard to the new weekly Journal entitled "Chemistry and Industry," and that the Journal for 1923 would be supplied to Fellows for the sum of 10s.

Mr. C. S. Grace and Dr. G. A. R. Kon were elected Scrutators, and a ballot for the election of Fellows was held. The following were subsequently declared elected as Fellows :

Robert Allanson.
 Thomas Cuthbert Allchin.
 James Wilson Armit, M.A.
 Leopold Louis Bacarese.
 Leslie Bains, B.A., B.Sc., A.I.C.
 Menghraj Choithram Bathija, B.A.
 Nicholas Timothy Belaiew, C.B.
 James Bell.
 Donald Charles Broome.
 Walter Ferguson Buist, B.Sc.
 Stanley Robert Bult.
 Arthur Josiah Burford.
 George Joseph Burrows, B.Sc.
 William Leslie Carter.
 Sydney George Cavigan.
 John Hardwick Long Champion,
 B.Sc.
 Katragadda Seshachalam Choudary,
 M.A., M.Sc.
 Elliot Martin Christie, M.Sc.
 Frederick Stanley Clarke.
 Allin Cottrell.
 John Arnold Cranston, D.Sc.
 Hugh John Curnow.
 Edward Dance, B.Sc.
 Wilfred Thomas Grey Davies, B.Sc.
 Milton Dewhurst, B.Sc., A.I.C.
 Richard Victor Eaton, B.Sc., A.R.C.S.
 Norman Evers, B.Sc., F.I.C.
 Joshua William Farmery, M.A., A.I.C.
 Samuel Frederick Ford.
 Roy Gardner, B.Sc.
 Philip Henry Gifford, B.Sc.
 Babu Lal Govila.
 Roland Hall Griffith, B.A.
 Reginald William Griffiths, B.Sc.
 John Masson Gulland, B.Sc.

Herbert Haigh, B.Sc.
 Lucy Higginbotham, M.Sc.
 Thomas Edward Hubbard.
 James Ernest Humphries, B.A., M.Sc.
 John Huntley.
 Sydney Alfred Hurren, M.C.
 Habib Iskander.
 John Watson Jenkin, B.Sc., A.I.C.
 John Ernest Jewell.
 Edgar Jones.
 Jan Kalf, D.Chem.
 Shankar Nath Kapur, B.Sc.
 Ghulam Safdar Khan, B.Sc.
 Svend Hald Lassen.
 William Dawson Lloyd.
 Philip Patrick Lynch, B.Sc.
 Francis Geoffrey Macalpine, B.Sc.
 Frederick Henry McDowall, M.Sc.
 Kenneth McKenzie.
 Reginald Arthur McNicol, M.Sc.
 Thomas Malkin, B.Sc., A.I.C.
 Archibald Roy Hammond Martin, B.A.
 Joseph Keith Matthews, M.A.
 Charles William Moorhead.
 John Ewart Moss, M.Sc.
 Dina Nath.
 Torkild Haastrup Nielsen, B.Sc.
 Hidejiro Nishikawa.
 Hiroshi Nomura.
 Albert Bramah Pearco Page, A.R.C.S.
 Harold Pickles.
 Joseph Edwards Pooley.
 Thomas Trevor Potts.
 James Robert Pound, M.Sc.
 Nagendra Prasad, M.Sc.
 Edmund Brydges Rudhall Prideaux,
 M.A., D.Sc., F.I.C.

Alfred Norman Douglas Pullen.
 George Clifford Riley, B.A., B.Sc.
 Thomas Abraham Robertson.
 Burke Rushworth.
 Arthur Scholefield.
 Gopal Singh, M.Sc.
 George Victor Smith, A.R.C.S.
 Stanley James Corrall Snedker.
 Oscar Walter Snow, B.A.
 Francis Wilbert Stoyte, M.Sc.
 Leslie Stuart Theobald, A.R.C.S.

William Thompson, B.Sc.
 Laurence Barnett Timmis, M.Sc.Tech.,
 A.I.C.
 Richard Truszkowski, B.Sc.
 Humphrey Terence Viney, B.Sc.
 Thomas John Wall, B.A.
 Harry Ward, M.Sc., A.I.C.
 Stephen John Watson, B.Sc.
 Alexander Mitchell Williams, M.A., D.Sc.
 Eric Richard Boraman Woodward, B.Sc.
 William James Uglow Woolcock, C.B.E.

The following papers were read :

- "The isoelectric condition of gelatin." By S. O. RAWLING and W. CLARK.
 "Studies on metal hydrides. The electrolytic formation of stibine in sulphuric acid and caustic soda solution." By H. J. S. SAND, E. J. WEEKS, and S. W. WORRELL.
 "Three carbon tautomerism in the *cyclopropane* series." By F. GOSS, C. K. INGOLD, and J. F. THORPE.

Meeting held in the Lecture Hall of the Institution of Mechanical Engineers, on Thursday, December 14th, 1922, at 8 p.m., Professor H. B. DIXON, C.B.E., M.A., F.R.S., Vice-President, in the Chair.

Professor C. H. Desch, D.Sc., Ph.D., delivered his Lecture entitled : "The metallurgical applications of physical chemistry." A vote of thanks to the Lecturer, proposed by Professor T. M. Lowry and seconded by Dr. N. V. Sidgwick, was carried with acclamation, acknowledgment being made by Professor Desch.

Ordinary Scientific Meeting, Thursday, December 21st, 1922, at 8 p.m., SIR JAMES WALKER, D.Sc., F.R.S., President, in the Chair.

The President referred to the loss sustained by the Society, through death, of :

	Elected.	Died.
Henry Ruchet Fairweather	Dec. 4th, 1919.	Sept. 11th.

Messrs. Duncan P. Woosley and Donald C. Broome were formally admitted Fellows of the Chemical Society.

Certificates were read for the first time in favour of :

George James Alexander, B.Sc., 1, Terrace Road, Upton Manor, E. 13.
 Harold Eugene Buckley, M.Sc., 273, Wigun Road, Bolton.
 Robert Kenneth Callow, B.A., B.Sc., 17, Sutherland Street, S.W. 1.
 Alexander Coutie, 4, Priestfield Road, Edinburgh.

Thomas Arthur Harrison, 64, Eastwood Road, Goodmayes.

Herbert Frederick Miller, B.Sc., A.I.C., Griffith John College, Hankow, China.

William Owen Murch, B.Sc., National Institute for Medical Research, Hampstead, N.W. 3.

James Herbert Oliver, B.Sc., 22, Avenue Gardens, Acton, W. 3.

Thomas Workman Orr, A.I.C., Antofagasta, Chile.

Edward Arthur Perren, 70, Denton Road, N. 8.

Frank Victor Raleigh, B.A., B.Sc., 6, Cavendish Road, Harringay, N. 4.

Helena Charlotte Randall, B.Sc., A.R.C.S., 57, Farnham Road, Guildford.

Jerry Robertson, 38, Church Street, Heckmondwike.

Douglas Rosebery Snellgrove, B.Sc., 73, Millbrook Road, Southampton.

Montague Williams, B.Sc., Cleveland Cottage, The Grove, Isleworth.

Certificates have been authorised by the Council for presentation to ballot under Bye-Law I (2) in favour of :

Donald Hatch Andrews, B.A., Stirling Chemical Laboratory, New Haven, Conn., U.S.A.

Arthur D. Little, Charles River Road, Cambridge, Mass., U.S.A.

The following papers were read :

"Benzbisthiazoles. Part II." By S. R. H. EDGE.

"Ring-chain tautomerism. Part IV. The effect of the methyl ethyl grouping on the carbon tetrahedral angle." By B. SINGH and J. F. THORPE.

ORDINARY SCIENTIFIC MEETING.

The next Ordinary Scientific Meeting will be held on Thursday, January 18th, 1923, at 8 p.m.

BUREAU FOR PHYSICO-CHEMICAL STANDARDS.

Under the auspices of the Union Internationale de la Chimie Pure et Appliquée, a "Bureau des Etalons Physico-chimiques" has been established at Brussels (Université, rue des Sols). This is intended as a centre for the preparation and collection of substances of the highest purity, the physical constants of which have been determined with the utmost possible precision, and which shall be at the disposal of research workers. An appeal is accordingly made to chemists generally to collaborate in the provision of highly purified products from which the standards may be prepared.

It is further proposed to collect the scattered literature dealing with the preparation of substances in a condition of the highest purity, and with the determination of their physical constants. With this object in view, investigators are invited to furnish the Bureau with copies of any relevant papers.

List of Papers received between November 17th, 1922, and December 21st, 1922 :

- "Negative hydrogen and the electronic formula of benzene." By H. S. FRY.
- "Elimination of the amino-group of tertiary amino-alcohols. Part I." By A. McKENZIE and A. C. RICHARDSON.
- "Studies on metal hydrides. The electrolytic formation of stibine in sulphuric acid and caustic soda solution." By H. J. S. SAND, E. J. WEEKS, and S. W. WORRELL.
- "Some derivatives of methylene-diquinalide and their relationship to the carbocyanines." By F. M. HAMER.
- "Dyestuffs derived from heterocyclic bases containing reactive methyl groups." By J. L. B. SMITH.
- "1 : 8-*bis*-Naphthylene-indigo." By S. DUTT.
- "Dyes derived from diphenic anhydride." By S. DUTT.
- "The effect of metallic oxides and metals on the oxidation of vegetable oils in presence of fibrous material. Part I. The rise in temperature and the changes in iodine value." By M. B. RANE, P. N. ROY, and L. GUPTA.
- "Synthesis of substituted thianthrens. Part I. Thianthren and nitrothianthren." By S. KRISHNA.
- "The absorption of moisture by coal (and other fuels). Part I. A relation between degree of humidity in the air and moisture content of coal." By B. MOORE and F. S. SINNATT.
- "The nitration of benzaldehyde." By O. L. BRADY and S. HARRIS.
- "The investigation of *meso*-thioanthracene derivatives. Part I. Observations on the production of dithioanthraquinone, dithiodianthrone, and other closely related derivatives." By I. M. HEILBRON and J. S. HEATON.
- "The chemistry of the glutaconic acids. Part XIII. The isomerism due to retarded mobility." By J. F. THORPE and A. S. WOOD.
- "The chemistry of the glutaconic acids. Part XIV. Three carbon tautomerism in the *cyclopropane* series." By F. R. GOSS, C. K. INGOLD, and J. F. THORPE.
- "The quantitative absorption of light by simple inorganic salts. Part II. The chlorides of arsenic, antimony, and bismuth." By A. K. MACBETH and N. I. MAXWELL.
- "The production and reactions of 2-dithiobenzoyl." By M. McKIBBEN and E. W. McCLELLAND.
- "The influence of papaverine on the optical activity of narcotine in acid solution." By H. E. ANNETT.

- "The higher oxide of cobalt." By O. R. HOWELL.
- "The calculation of the colour of monocyclic substances by assigning an absorption band to each possible tautomeric formula of the substance." By J. MOIR.
- "The application of the Hoesch reaction to the preparation of some dihydroxy- and trihydroxy-deoxybenzoin." By E. CHAPMAN and H. STEPHEN.
- "Studies on thallium compounds. Part II. The reduction of thallic compounds with ferrous sulphate and with sodium arsenite." By A. J. BERRY.
- "The constitution of the disaccharides. Part VII. Sucrose." By W. N. HAWORTH and W. H. LINNELL.
- "The determination of the dissociation pressures of hydrated salts by a dynamical method. Part II." By J. R. PARTINGTON and D. B. HUNTINGFORD.
- "The constitution of the disaccharides. Part VIII. Sucrose." By W. N. HAWORTH and J. G. MITCHELL.
- "Benzobisthiazoles. Part II." By S. R. H. EDGE.
- "The upper limit of diazotisability in the benzene series. Diazo-derivatives of mesitylene." By G. T. MORGAN and G. R. DAVIES.
- "Researches on residual affinity and co-ordination. Part XV. Interactions of acetylpropionylmethane and the tetrachlorides of selenium and tellurium." By G. T. MORGAN and H. G. REEVES.
- "Studies in the normal butyl series. Part II. The four stereoisomeric β - γ -di-*p*-tolylamino-*n*-butanes." By G. T. MORGAN and W. J. HICKINBOTTOM.
- "The formation and stability of spiro-compounds. Part X. Spiro-compounds derived from *cycloheptane*." By J. W. BAKER and C. K. INGOLD.
- "The sorption by carbon prepared from pure organic compounds. Part I. The sorption of iodine by carbons prepared from carbohydrates." By J. B. FIRTH.
- "Ring-chain tautomerism. Part IV. The effect of the methyl ethyl grouping on the carbon tetrahedral angle." By B. SINGH and J. F. THORPE.
- "The transition from the colloidal to the crystalloidal state. Part I. Solutions of potassium oleate." By L. L. BIRCUMSHAW.
- "Investigations on the dependence of rotatory power on chemical constitution. Part XVIII. The di-*l*-menthyl esters of the acids of the oxalic esters." By L. HALL.

- "The propagation of flame in complex gaseous mixtures. Part V. The interpretation of the law of speeds." By W. PAYMAN.
- "The rate of detonation in complex gaseous mixtures." By W. PAYMAN and N. S. WALLS.
- "Determination of tin in wolfram. A modification of Powell's method." By O. F. LUBATTI.
- "The reactivity of doubly-conjugated unsaturated ketones. Part IV. The effect of substitution on the reactivity of 4'-dimethylamino-2-hydroxydistyryl ketone." By I. M. HEILBRON and A. B. WHITWORTH.
- "Studies in phototropy. The reversed phototropy of cinnamaldehydesemicarbazone and its methoxy derivatives." By I. M. HEILBRON, H. E. HUDSON, and D. M. HUISE.
- "Complex metallic ammines. Part VIII. The introduction of *d*- and *tri*-basic organic acid radicles into the pentamminecobaltic complex." By J. C. DUFF.

ADDITIONS TO THE LIBRARY.

I. *Donations.*

DAKIN, HENRY DRYSDALE. Oxidations and reductions in the animal body. 2nd edition. London 1922. pp. x + 176. 6s. net. (*Recd.* 1/12/22.)

From the Publishers : Messrs. Longmans, Green & Co.

HACKSPILL, LOUIS. L'azote : la fixation de l'azote atmosphérique et son avenir industriel. Paris 1922. pp. xii + 272. ill. 14 fr. net. (*Recd.* 8/12/22.) From Dr. E. B. Maxted.

KEMPTON, PERCIVAL HENRY SYDNEY. The industrial applications of X-rays : an introduction to the apparatus and methods used in the production and application of X-rays for the examination of materials and structures. London 1922. pp. xiv + 112. ill. 2s. 6d. net. (*Recd.* 6/12/22.)

From the Publishers : Sir Isaac Pitman & Sons.

II. *By Purchase.*

BACON, RAYMOND FOSS, and HAMOR, WILLIAM ALLEN. [Éditeurs.] American fuels. 2 vols. New York 1922. pp. x + 628, vi, 629—1258. ill. 60s. net. (*Recd.* 23/11/22.)

BARKER, ALDRED FARRER, and MIDGLEY, EBER. Analysis of woven fabrics. 2nd edition. London 1922. pp. xvi + 322. ill. 15s. net. (*Recd.* 16/11/22.)

CLARK, WILLIAM MANSFIELD. The determination of hydrogen ions. 2nd edition. Baltimore 1922. pp. 480. ill. \$5 net. (*Recd.* 9/12/22.)

CLARKE, ALBERT. Coal-tar colours in the decorative industries. London 1922. pp. xiv + 166. 6s. net. (*Recd.* 21/11/22.)

DESCOUR, L. Pasteur and his work. Translated by A. F. WEDD and B. H. WEDD. London [1922]. pp. 256. ill. 15s. net. (*Recd.* 21/11/22.)

GRAFE, VIKTOR. Chemie der Pflanzenzelle. Berlin 1922. pp. viii + 421. ill. (*Recd.* 28/11/22.)

HÖBER, RUDOLF. Physikalische Chemie der Zelle und der Gewerbe. 5th edition. [Part I.] Leipzig 1922. pp. xvi + 544. ill. (*Recd.* 1/12/22.)

KEWLEY, JAMES. The petroleum and allied industries: petroleum, natural gas, natural waxes, asphalts and allied substances, and shale oils. London 1922. pp. xii + 302. ill. 12s. 6d. net. (*Recd.* 21/11/22.)

MARSHALL, CHARLES EDWARD. [Editor.] Microbiology: a text-book of microorganisms general and applied. [Philadelphia 1921.] pp. xxviii + 1044. ill. 21s. net. (*Recd.* 28/11/22.)

MICHAELIS, LEONOR. Die Wasserstoffionenkonzentration: ihre Bedeutung für die Biologie und die Methoden ihrer Messung. 2nd edition. Part I. Berlin 1922. pp. xii + 262. ill. (*Recd.* 1/12/22.)

PARRY, ERNEST JOHN. The chemistry of essential oils and artificial perfumes. 4th edition. 2 vols. London 1921—22. pp. viii + 549, viii + 365. ill. 51s. net. (*Recd.* 23/11/22.)

PILE, SEYMOUR, and JOHNSTON, REGINALD. Tested methods of metallurgical analysis (non-ferrous). London 1922. pp. 128. ill. 7s. 6d. net. (*Recd.* 21/11/22.)

RUSSELL, ALEXANDER SMITH. An introduction to the chemistry of radio-active substances. London 1922. pp. xii + 174. ill. 6s. net. (*Recd.* 21/11/22.)

SOCIETY OF DYERS AND COLOURISTS. Colour index. Part I, etc. Bradford 1922 +. (*Reference.*)

SMITS, ANDREAS. The theory of allotropy. Translated by JOHN SMEATH THOMAS. London 1922. pp. xiv + 398. ill. 21s. net. (*Recd.* 9/12/22.)

SOMMERFELD, ARNOLD. Atombau und Spektrallinien. 3rd edition. Braunschweig 1922. pp. xii + 764. ill. (*Recd.* 29/11/22.)

VENERAND, WOLFGANG. Asbest und Feuerschutz: Vorkommen, Verarbeitung und Anwendung des Asbestes. Feuerschutz in Theatern, öffentlichen Gebäuden usw. 2nd edition. Wien 1920. pp. viii + 360. ill. (*Recd.* 8/12/22.)

IMPERIAL AGRICULTURAL RESEARCH
INSTITUTE LIBRARY
NEW DELHI.

[illegible]